Second leptogenesis: Unraveling the baryon-lepton asymmetry discrepancy

YeolLin ChoeJo,^a Kazuki Enomoto,^a Yechan Kim,^a Hye-Sung Lee^a

^aDepartment of Physics, Korea Advanced Institute of Science and Technology Daejeon 34141, Korea

E-mail: particlephysics@kaist.ac.kr, k_enomoto@kaist.ac.kr, cj7801@kaist.ac.kr, hyesung.lee@kaist.ac.kr

ABSTRACT: We propose a novel scenario to explain the matter-antimatter asymmetry by twofold leptogenesis, wherein heavy Majorana neutrinos exhibit temperature-dependent masses and engage in CP-violating decays. This scenario envisages two distinct phases of leptogenesis: one occurring above the electroweak scale and the other below it. The sphaleron process converts the first lepton asymmetry to baryon asymmetry, but not the second one due to its decoupling. This mechanism potentially explains the significant discrepancy between baryon and lepton asymmetries, as suggested by recent observations of Helium-4. Furthermore, our model implies that the present masses of Majorana neutrinos are lighter than the electroweak scale, offering a tangible avenue for experimental verification in various terrestrial settings.

Contents

1	Introduction	1
2	Second leptogenesis	2
3	Realization in a wave dark matter model	3
4	Parameter regions for the second leptogenesis	6
5	Quantitative result of the second leptogenesis	7
6	Summary and outlook	10

1 Introduction

The universe is baryon asymmetric as suggested by the measured baryon-to-photon ratio [1]

$$\eta_B \equiv \frac{n_B - n_{\bar{B}}}{n_{\gamma}} \simeq (6.14 \pm 0.25) \times 10^{-10}, \tag{1.1}$$

where n_B , $n_{\bar{B}}$, and n_{γ} are the number density of the baryon, antibaryon, and photon, respectively. Considering cosmic inflation, η_B must be generated after the inflation rather than being an initial condition, [2], a process known as the baryogenesis [3]. It is wellestablished that the standard model (SM) falls short in explaining this [4–9].

Leptogenesis is a plausible new physics scenario to explain η_B , where the heavy Majorana neutrino N possessing the Yukawa couplings to the lepton doublet ℓ and the Higgs doublet Φ , are added to the SM [10]. At the decoupling of N in the early universe, CPviolating decays $N \to \ell \Phi$ and $N \to \bar{\ell} \Phi^{\dagger}$ produce the lepton number (L). It is converted to the baryon number (B) via the sphaleron process [11, 12]. N also facilitates the explanation of tiny neutrino masses through the type-I seesaw mechanism [13–17].

Recently, the EMPRESS experiment reported a new result of the ⁴He abundance observation [18]. It suggests the large degeneracy parameter of the electron neutrino; $\xi_e = 0.05^{+0.03}_{-0.02}$. This implies substantial lepton-to-photon ratio given by [19]

$$\eta_L \equiv \frac{n_L - n_{\bar{L}}}{n_{\gamma}} \simeq \sum_l \frac{g_l \pi^2}{12\zeta(3)} \left(\frac{T_l}{T_{\gamma}}\right)^3 \xi_l,\tag{1.2}$$

where l represents all leptons. g_l , ξ_l , and T_l are the degree of freedom, the degeneracy parameter, and the temperature of l, respectively. The EMPRESS result suggests

$$\eta_L \simeq \frac{\pi^2 \sum_{i=e,\mu,\tau} \xi_{\nu_i}}{6\zeta(3)} \left(\frac{T_\nu}{T_\gamma}\right)^3 \simeq (7.5^{+4.5}_{-3.0}) \times 10^{-2},\tag{1.3}$$



Figure 1. (a) The standard leptogenesis with the constant mass M_0 . In the red (blue) region, N is in (out of) thermal equilibrium. Leptogenesis occurs in the blue region. (b) Leptogenesis may occur twice for the temperature-dependent mass, resulting in a larger lepton asymmetry than the baryon asymmetry.

where we have used $(T_{\nu}/T_{\gamma})^3 = 4/11$ and the flavor universality $(\xi_{\nu_e} = \xi_{\nu_{\mu}} = \xi_{\nu_{\tau}})$ due to the neutrino oscillations. Given the universe's electrical neutrality, we disregard the charged lepton asymmetry of the same magnitude as η_B , as it does not match the observed lepton asymmetry. See Refs. [20–28] for further discussions on the lepton asymmetry. While the deviation is at a 2.5 σ level and may not appear excessively large, the suggested value has a significant phenomenological impact.

The lepton asymmetry in Eq. (1.3) is much larger than the baryon asymmetry in Eq. (1.1); $\eta_L/\eta_B \simeq 10^8$. If the sphaleron process occurs frequently, the baryon and lepton asymmetries are made to have the same size, which is proportional to the initial B - L [29, 30]. Thus, the traditional baryogenesis scenario cannot explain this large discrepancy. The discrepancy seems to suggest that two asymmetries were individually generated by different new physics at different epochs in the early universe.

In this paper, we propose a novel, yet simple leptogenesis scenario to explain this discrepancy. It posits two occurrences of leptogenesis in the early universe, driven by the temperature-dependent mass of N. The first leptogenesis takes place prior to the sphaleron decoupling at the electroweak symmetry breaking $(T_{\rm sph} \simeq 100 \text{ GeV})$ [31], and the generated lepton asymmetry is converted to the baryon asymmetry. On the other hand, the second leptogenesis happens below $T_{\rm sph}$, allowing the resultant lepton asymmetry to persist into the present universe.

2 Second leptogenesis

First, we describe the scenario of the twofold leptogenesis and how it can explain the suggested large discrepancy between the baryon and lepton asymmetries. This concept is visualized in Figure 1, alongside a comparison with the standard leptogenesis scenario.

We consider the following case. In the early universe, the heavy neutrino N acquires the temperature-dependent Majorana mass by the new physics effect in addition to the bare Majorana mass M_0 . The sums of them are denoted by M(T). We assume that M_0 is smaller than the electroweak scale; $M_0 < 100$ GeV.

M(T) behaves as a constant M_* ($\gg M_0$) above the temperature T_* . At $T < T_*$, it decreases as a function of temperature. When the temperature has dropped enough below a certain temperature T_N ($\ll T_*$), the new physics effect becomes negligibly small, and $M(T) \simeq M_0$ until the current universe. A specific new physics to realize this scenario will be discussed later in this paper.

At high temperatures $T > T_*$, N behaves as Majorana fermion with the mass M_* and is thermalized via the Yukawa interaction. At $T \simeq M_*$, the production rate is exponentially suppressed, and N begins to be decoupled. The first leptogenesis occurs at this stage. L is generated, and it is converted to B, which remains as the baryon asymmetry until the current universe.

At $T < T_*$, M(T) begins to decrease as cooling of the universe; $M(T) \propto T^n$. If this decrease is faster than the temperature decrease (n > 1), M(T) can be lower than T at some point, and N can be thermalized again.

As the universe cools further, M(T) behaves as a constant again but much smaller than the mass at the high temperature; $M(T) \simeq M_0 \ll M_*$. At $T \simeq M_0$, the heavy neutrino is decoupled again. The second leptogenesis occurs at this stage, generating the extra lepton number ΔL . Since M_0 is lower than the electroweak scale, the sphaleron process has already decoupled, and ΔL remains as the additional lepton asymmetry until the current universe.

Since the baryon asymmetry is set by L, the size of ΔL has to be much larger than L to explain the large baryon-lepton asymmetry discrepancy. How can a large enhancement of ΔL be made at the second leptogenesis? It can be realized in a natural way. The size of the Yukawa coupling y is proportional to $\sqrt{M_0}$ to reproduce the neutrino mass matrix by the type-I seesaw mechanism [32]. Thus, the ratio of the production rate $\Gamma_{\text{prod}} \propto y^2 M(T)$ and the Hubble parameter $H \propto T^2$ is given by $\Gamma_{\text{prod}}/H \propto M(T)M_0/T^2$. At the first leptogenesis $(T \simeq M_*)$, the production is much suppressed because $\Gamma_{\text{prod}}/H \propto M_0/M_* \ll 1$. Such a case is referred to as the weak washout [33], and the generated lepton asymmetry is also suppressed. On the other hand, at the second leptogenesis $(T \simeq M_0)$, the production rate is not suppressed because $\Gamma_{\text{prod}}/H \propto M(T)M_0/T^2|_{T=M_0} \simeq 1$. This is the strong washout [33], and much larger lepton asymmetry can be generated via the second leptogenesis compared to the first one. It can naturally explain the large difference between the baryon and lepton asymmetries.

3 Realization in a wave dark matter model

Here, we discuss a specific realization of the temperature-dependent mass of N. The neutrino mass variation over cosmic time has been studied a lot in the context of the mass-varying neutrinos in the quintessence dark energy field, where the neutrinos may get their masses from the quintessence field [34, 35]. Lately, there have been many new studies of the mass-varying neutrinos [36–60], including those taking the scalar wave dark matter

(DM) [61] in place of the quintessence scalar field. In this paper, we focus on the case that M(T) is given by the coupling to the scalar wave DM.

The scalar wave DM ϕ obeys the following equation of motion with the assumption of spatial homogeneity;

$$\ddot{\phi} + 3H\dot{\phi} + m_{\phi}^2\phi = 0, \qquad (3.1)$$

where m_{ϕ} is the mass of ϕ and is constrained to 3×10^{-21} eV $< m_{\phi} < 30$ eV [61]. The lower bound arises from Lyman- α forest data, and the upper from the de Broglie wavelength exceeding inter-particle separation.

At high temperatures, ϕ is fixed at the nonzero initial value due to the large Hubble friction. Since the *H* decreases over time, it becomes comparable to m_{ϕ} at the temperature T_* ; $H(T_*) = m_{\phi}$. We assume the universe is radiation-dominated in the following. It leads to $T_* \simeq \left(m_{\phi} M_{\rm Pl} \sqrt{90/(8\pi^3 g_*)}\right)^{1/2}$, where $M_{\rm Pl}$ is the Planck mass and $g_* = 106.75$ is the effective degree of freedom of the energy density [38, 40, 51, 59].

At $T < T_*$, ϕ coherently oscillates by the mass term;

$$\phi(t) = \frac{\sqrt{2\rho(t)}}{m_{\phi}} \cos(m_{\phi}t), \qquad (3.2)$$

where $\rho(t) = \frac{1}{2}\dot{\phi}^2 + \frac{1}{2}m_{\phi}^2\phi^2$ is the energy density of ϕ . Since ρ behaves as the matter-like, $\rho \propto a^{-3}$ where *a* is the scale factor, the oscillation amplitude becomes smaller proportional to $a^{-3/2} \propto T^{3/2}$ as the temperature decreases [62–64].

At the current temperature $T_0 \simeq 2.73$ K [65], the oscillation energy density ρ_0 contributes to $\rho_{\rm DM}$, the relic energy density of the DM. In this paper, we assume that the oscillating ϕ contributes to the entire DM. It requires the current oscillation amplitude ϕ_0 to be $\phi_0 = \sqrt{2\rho_{\rm DM}}/m_{\phi}$.

The interaction between particles and the wave DM provides the time-dependent mass of the particles. Some references have investigated leptogenesis with the varying neutrino mass by using quintessence dark energy [66, 67], the neutrino itself as dark energy [68], or other new physics [69]. Nevertheless, none of these works discussed the possibility of the second leptogenesis.

We assume three Majorana neutrinos N_i (i = 1, 2, 3) which couple to the scalar wave DM [39, 51, 59]. The relevant part of the Lagrangian is given by

$$\mathcal{L} = -\frac{1}{2} (M_{0i} + g_i \phi) \overline{N}_i^c N_i + \text{h.c.}$$
(3.3)

where M_{0i} and g_i are the bare Majorana masses and coupling constants of N_i , respectively. We do not consider off-diagonal couplings for simplicity. The cosmic scaling of the Majorana neutrino mass in the wave DM setup was also studied in Refs. [39, 51, 59].

The second term in Eq. (3.3) generates the time-dependent mass $M_i(t) = M_{0i} + g_i \phi(t)$. If the oscillation period is much shorter than the time scale of the relevant physics (in our case, N_i decay time), the oscillating term $\phi(t)$ can be approximated by its time average.



Figure 2. The temperature-dependent mass M(T) in Eq. (3.5) using the time-average approximation (black) in comparison with temperature T (orange). Leptogenesis occurs when the heavy neutrinos decouple (two blue periods). Indices *i* are suppressed here.

m_{ϕ}	Wave dark matter mass
$\overline{M_{*i}}$	Majorana mass at T_*
M_{0i}	Bare Majorana mass
$\overline{T_*}$	Temperature when oscillation starts
T_{N_i}	Temperature when $g_i \phi$ term becomes negligible
T_0	Temperature of the current universe

Table 1. The notations of masses and temperatures. The free parameters are only m_{ϕ} , M_{*i} and M_{0i} , and others are determined by these parameters.

As a result, we obtain the temperature-dependent mass of N_i as follows;

$$M_{i}(T) = \begin{cases} M_{0i} + \frac{g_{i}\phi_{0}}{\sqrt{2}} \left(\frac{T_{*}}{T_{0}}\right)^{3/2} & T > T_{*}, \\ M_{0i} + \frac{g_{i}\phi_{0}}{\sqrt{2}} \left(\frac{T}{T_{0}}\right)^{3/2} & T_{*} > T. \end{cases}$$
(3.4)

In order to have the twofold leptogenesis scenario described earlier, we consider the case that the first term of $M_i(T)$ is dominant at low temperatures $(T \ll T_*)$; on the other hand, the second one is at high temperatures $(T \simeq T_*)$. Then, we can find the temperature T_{N_i} $(\ll T_*)$, below which the effect of the wave DM (the second term) is negligible compared with the bare mass (the first term). It is evaluated by $M_{0i} = g_i \phi_0 (T_{N_i}/T_0)^{3/2}/\sqrt{2}$.

Consequently, the behavior of $M_i(T)$ is described by

$$M_{i}(T) \simeq \begin{cases} M_{*i} & T > T_{*}, \\ M_{0i} + g_{i} \frac{\phi_{0}}{\sqrt{2}} \left(\frac{T}{T_{0}}\right)^{3/2} & T_{*} > T > T_{N_{i}}, \\ M_{0i} & T_{N_{i}} > T, \end{cases}$$
(3.5)



Figure 3. The allowed region for the second leptogenesis (blank region) for $m_{\phi} = 10^{-2}$ eV. In the blue regions, the necessary conditions for the second leptogenesis are not satisfied. The red and yellow regions are excluded by theoretical requirements to avoid a dominant quartic coupling and thermalization of ϕ , respectively. The blue star is the benchmark point for Figure 4.

where $M_{*i} \equiv M_{0i} + g_i \phi_0 (T_*/T_0)^{3/2} / \sqrt{2}$. In Figure 2, we illustrate how $M_i(T)$ varies with temperature in the second leptogenesis scenario. The masses and temperatures are summarized in Table 1.

4 Parameter regions for the second leptogenesis

Here, we consider constraints on the model parameters. The Lagrangian of the model includes three kinds of new parameters m_{ϕ} , M_{0i} , and g_i . We can also choose a more convenient set of the parameters: T_* , M_{0i} , and M_{*i} . Using these parameters, T_{N_i} is evaluated by

$$T_{N_i} = T_* \left(\frac{M_{0i}}{M_{*i} - M_{0i}}\right)^{2/3} \simeq T_* \left(\frac{M_{0i}}{M_{*i}}\right)^{2/3}.$$
(4.1)

The necessary conditions for the second leptogenesis are as follows. For the lightest N_i , (i) $M_{*i} > T_*$: The first decoupling happens earlier than T_* . (ii) $T_{N_i} > M_{0i}$: The second decoupling occurs later than T_{N_i} . (iii) $T_* > T_{N_i}$: The time interval for the temperaturedependent mass should exist, which is equivalent to $M_{*i} > 2M_{0i}$ by Eq. (4.1).

Next, we consider theoretical constraints. The quartic coupling of ϕ is radiatively induced by g_i , and it has to be smaller than the mass term at least at the matter-radiation equality $T_{mr} \simeq 1$ keV, otherwise $\rho(t)$ behaves as a^{-4} not a^{-3} [51, 57, 70]. This requires $m_{\phi}^2/\phi_{mr}^2 > g_i^4/(16\pi^2)$, where ϕ_{mr} is the oscillation amplitude at T_{mr} . In addition, in order to avoid thermalization of ϕ , the scattering rate has to be smaller than H(T) [51]. We consider two scatterings $\phi\nu \to \phi\nu$ and $\phi N_i \to \phi N_i$. The former gives a weaker constraint because ν couples to ϕ only via tiny mixing. The scattering rate of the latter is roughly given by $\Gamma_{\phi N_i \to \phi N_i} \sim g_i^4 T$ when N_i is relativistic. Thus, we obtain $g_i^4 > \sqrt{8\pi^3 g_*/90} T/M_{\rm Pl}$.

The coupling g_i is also subject to constraints from various experimental studies, such as the Majoron emitting decay [71, 72], neutrino free-streaming on the CMB [49], and neutrino oscillations [37] depending on the m_{ϕ} values. However, their constraints are weaker than others in the parameter space we are interested in.

In Fig. 3, we show the allowed parameter regions for the second leptogenesis in the case of $m_{\phi} = 10^{-2}$ eV, which corresponds to $T_* \simeq 2.7$ TeV. The blue regions do not satisfy the three conditions required for the second leptogenesis. The red and yellow regions are excluded by theoretical constraints to avoid a dominant quartic coupling at $T = T_{mr}$ and to prevent the thermalization of ϕ , respectively. The experimental constraints are too weak to be shown in the figure. Consequently, the second leptogenesis is expected to occur in the blank regions of the figure. In numerical evaluations, we use the average density of the DM $\rho_{\rm DM} = 1.2 \times 10^{-6} \text{ GeV/cm}^3$, not the local density because we investigate phenomena in the early universe.

The allowed region changes with different values of m_{ϕ} . For example, the constraint $\Gamma_{\phi N_i \to \phi N_i} > H$ becomes stronger for larger m_{ϕ} , while the quartic coupling constraint becomes more stringent for smaller m_{ϕ} . Since we consider the scenario where the oscillation of ϕ begins before the sphaleron decoupling as explained in Sec. 2, m_{ϕ} needs to be larger than 10^{-5} eV, which is derived from $T_* > T_{\rm sph} \simeq 100$ GeV. Thus, our relevant mass region is 10^{-5} eV $< m_{\phi} < 30$ eV. We have checked that the allowed region does not change significantly, and we can find a lot of parameter points to achieve the second leptogenesis in this mass region.

5 Quantitative result of the second leptogenesis

The asymmetry production in leptogenesis is evaluated by the density matrix equation including the flavor effect [73–75]. Here, we employ the formalism given in Refs. [76–78] with the addition of the temperature dependence in the masses. The equation is given by

$$\frac{\mathrm{d}N_{N_i}}{\mathrm{d}z} = -D_i(N_{N_i} - N_{N_i}^{\mathrm{eq}}),$$

$$\frac{\mathrm{d}N_{\alpha\beta}}{\mathrm{d}z} = \sum_i \left[\varepsilon_{\alpha\beta}^{(i)} D_i(N_{N_i} - N_{N_i}^{\mathrm{eq}}) - \frac{1}{2} W_i \{P_i, N\}_{\alpha\beta} \right]$$

$$- \frac{\Gamma_{\tau}}{Hz} [I_{\tau}, [I_{\tau}, N]]_{\alpha\beta} - \frac{\Gamma_{\mu}}{Hz} [I_{\mu}, [I_{\mu}, N]]_{\alpha\beta},$$
(5.1)
(5.1)

where $z = M_{01}/T$, i = 1, 2, 3, and $\alpha, \beta = e, \mu, \tau$. N_{N_i} and the diagonal terms $N_{\alpha\alpha}$ are the number of N_i and $B/3 - L_{\alpha}$, where L_{α} is the lepton number for each flavor, respectively, in a portion of the comoving volume that contains one photon at the era when N_i is relativistic and in thermal equilibrium [79, 80]. The off-diagonal terms $N_{\alpha\beta}$ ($\alpha \neq \beta$) represent the coherence between the flavors. The number of the total B-L is given by $N_{B-L} = \sum_{\alpha} N_{\alpha\alpha}$. The term D_i accounts for the decay and inverse decay of N_i . The washout effect in B-L asymmetry is described by W_i . We consider the washout effect from the inverse decay and neglect one from other lepton-number-violating processes. D_i and W_i are given by [80]

$$D_{i} = \frac{(yy^{\dagger})_{ii}}{8\pi Hz} M_{i}(T) \frac{K_{1}(M_{i}(T)/T)}{K_{2}(M_{i}(T)/T)},$$
(5.3)

$$W_i = \frac{2}{3} D_i N_{N_i}^{\text{eq}}, \tag{5.4}$$

where $N_{N_i}^{\text{eq}} = (3/8) (M_i(T)/T)^2 K_2 (M_i(T)/T)$ is the equilibrium value of N_{N_i} , and y is the Yukawa matrix for the interaction among the lepton doublet, the Higgs doublet, and the heavy neutrinos. K_n is a modified *n*-th Bessel function of the second kind. We note that the temperature dependence of the mass is included in these terms. P_i is the projection matrix constructed with the Yukawa matrix. Decoherence effects by the interchanges between the left-handed and right-handed leptons are described by the double commutator terms, where $\Gamma_{\mu(\tau)} \simeq 8 \times 10^{-3} (\sqrt{2}m_{\mu(\tau)}/v)^2 T$ [77] is the rate of the process involving $\mu(\tau)$, $I_{\mu} = \text{diag}(0, 1, 0)$, and $I_{\tau} = \text{diag}(0, 0, 1)$.

Since M_{0i} in our scenario is much lighter than the Davidson-Ibarra bound, $M_{0i} \gtrsim 10^8$ GeV [81], for the heavy neutrinos with hierarchical masses, we consider resonant leptogenesis, where the heavy neutrinos possess very close masses [82, 83]. The *CP* asymmetry parameter $\varepsilon_{\alpha\beta}^{(i)}$ is divided into contributions from vertex and self-energy diagrams [33]. The self-energy part $\varepsilon_{\alpha\beta}^{S(i)}$ is resonantly enhanced with degenerate masses and can dominate the *CP* asymmetry parameter. $\varepsilon_{\alpha\beta}^{S(i)}$ is given by

$$\varepsilon_{\alpha\beta}^{S(i)} = \frac{1}{16\pi (yy^{\dagger})_{ii}} \sum_{j\neq i} \left\{ i \left[y_{i\alpha}^* y_{j\beta} (yy^{\dagger})_{ji} - y_{i\beta} y_{j\alpha}^* (yy^{\dagger})_{ij} \right] \frac{M_j}{M_i} + i \left[y_{i\alpha}^* y_{j\beta} (yy^{\dagger})_{ij} - y_{i\beta} y_{j\alpha}^* (yy^{\dagger})_{ji} \right] \right\} \frac{(M_j^2 - M_i^2) M_i^2}{(M_j^2 - M_i^2)^2 + M_i^4 \Gamma_j^2 / M_j^2},$$
(5.5)

where M_j and Γ_j are the temperature-dependent mass and decay rate of N_j , respectively. Thus, $\varepsilon_{\alpha\beta}^{(i)}$ is significantly enhanced when the resonant condition, $|M_j - M_i| \simeq \Gamma_j/2$, is satisfied.

As a benchmark point to evaluate N_{N_1} and N_{B-L} , we assume the following input parameters: $m_{\phi} = 10^{-2}$ eV ($T_* \simeq 2.7$ TeV), $M_{01} = 0.1$ GeV, and $M_{*1} = 2.4 \times 10^5$ GeV. The current masses of N_2 and N_3 are chosen to satisfy the resonant condition at the second leptogenesis, $\Delta M_{12} \equiv M_{02} - M_{01} = 0.5 \times 10^{-19}$ GeV and $\Delta M_{13} \equiv M_{03} - M_{01} =$ 4.0×10^{-19} GeV. M_{*i} (i = 2, 3) are determined by imposing $M_{*i}/M_{*1} = M_{0i}/M_{01}$ by which the resonant condition is also satisfied at the first leptogenesis.

The Yukawa matrix y is set to satisfy the neutrino oscillation data with the normal ordering masses [65] by using the Casas-Ibarra parametrization [32], $y = \sqrt{2} \hat{M}_N^{1/2} R \hat{m}_{\nu}^{1/2} U^{\dagger} / v$, where \hat{M}_N and \hat{m}_{ν} are the diagonal mass matrices of the heavy and active neutrinos, respectively, U is the PMNS matrix [84, 85], v is the vacuum expectation value of the Higgs, and R is a complex orthogonal matrix. We note that \hat{M}_N is evaluated at the current



Figure 4. N_{N_1} and $|N_{B-L}|$ for resonant leptogenesis with the temperature-dependent masses of the neutrinos on the benchmark point described in the text. Here, $|N_{B-L}|$ is the sum of the B-Lasymmetry deposited in each flavor, and its sign is represented together. The baryon asymmetry is determined after the sphaleron decoupling ($T \simeq 100$ GeV corresponding to $z \simeq 10^{-3}$ with $M_{01} = 0.1$ GeV) with positive N_{B-L} , but the lepton asymmetry is fixed only after the second leptogenesis ($z \simeq 1$) with negative N_{B-L} . The baryogenesis fitting at the sphaleron decoupling ($z \simeq 10^{-3}$) is shown together (thin lines).

temperature. Six parameters in y are undetermined by the neutrino oscillation data: the lightest neutrino mass m_{ν_1} , two Majorana phases α_1, α_2 in the PMNS matrix U, and three complex phases $\omega_1, \omega_2, \omega_3$ in R. The notations of the Majorana phases, and the complex phases follow Refs. [65] and [76], respectively. We assume the following values for them; $m_{\nu_1} = 0 \text{ eV}, \alpha_1 = \alpha_2 = 0, \omega_1 = \omega_2 = 0$, and $\omega_3 = 0.2e^{-i\pi/4}$.

We consider thermal leptogenesis with the initial condition $N_{N_1} = N_{B-L} = 0$ at $z = 10^{-10}$ ($T = 10^9$ GeV). Figure 4 shows the behavior of N_{N_1} and N_{B-L} with the above inputs. The first leptogenesis occurs a little later than $T = M_{*1}$ ($z \simeq 10^6$) because of weak washout [80]. The produced lepton number is converted to the baryon number by the sphaleron process, and it is fixed after the sphaleron decoupling. For simplicity, we assume that the decoupling occurs instantaneously, and the final baryon number is determined by N_{B-L} at the temperature T = 100 GeV ($z = 10^3$) shown by the purple lines in Figure 4.

The baryon-to-photon ratio can be obtained by $\eta_B = a_{\rm sph} N_{B-L}/f$ at the sphaleron decoupling, where $a_{\rm sph}$ and f are the sphaleron conversion rate [29, 30] and the photon dilution factor [79] due to the increase of the photon number by annihilation of particles from the first leptogenesis till the BBN, respectively. We use $a_{\rm sph} = 28/79$, the value for the SM plasma. Since the heavy neutrinos have nearly degenerate masses, we evaluate f including the effect of N_1 , N_2 , and N_3 , not only N_1 . Then, we have f = 1232/43.¹ As a

¹With m generations of the heavy neutrinos, f = 11(427 + 7m)/172, which leads to the commonly used

result, we obtain $\eta_B \simeq 6.14 \times 10^{-10}$, which is consistent within 1σ level with the current observations [1, 86]. Also, the positive N_{B-L} at the sphaleron decoupling in our benchmark point provides the correct sign for the baryon asymmetry.

The second leptogenesis commences shortly after $T = M_{01}$ (or z = 1), due to the effects of strong washout [80]. The lepton number generated during this period persists into the current universe, as all lepton-number-violating processes have ceased. In this scenario, the photon dilution factor f' is calculated based on the change in photon numbers from the second leptogenesis to the era of BBN, rather than from the first leptogenesis. This is because the N_1 species is relativistic and returns to thermal equilibrium before the onset of the second leptogenesis. We adopt f' = 176/43, where the thermal bath consists of e^{\pm} , ν_{ℓ} , γ , and N_i before the second leptogenesis.² Consequently, we derive a lepton asymmetry value of $\eta_L = N_{B-L}/f' \simeq 5.0 \times 10^{-3}$. The lepton asymmetry is flavor-universal due to neutrino oscillation, so we consider the summation of the B - L asymmetry across all flavors, rather than focusing solely on the electron component. Since N_{B-L} is negative in the late epoch of our benchmark model, a positive lepton asymmetry is also guaranteed.

This demonstrates that the second leptogenesis can significantly amplify lepton asymmetry, increasing it by several orders of magnitude from the baryon asymmetry. In our analysis, the benchmark point, which was not fully optimized, already indicates $\eta_L \sim 10^{-3}$. This is remarkably close to the EMPRESS data, which suggests $\eta_L \sim 10^{-2}$, and represents a substantial deviation from $\eta_B \sim 10^{-10}$. A more refined analysis or the addition of more Majorana neutrinos might aid in reconciling the slight discrepancy from the observed values.

6 Summary and outlook

It is notable that a significant deviation of the lepton asymmetry from the baryon asymmetry can be explained in a rather simple framework of the second leptogenesis. This scenario allows only larger lepton asymmetry than the baryon asymmetry, not the other way around, in accordance with the measurement. A more comprehensive study will follow in the subsequent work. In the future, there will be increased CMB data from the Simons Observatory [87] and CMB-S4 [88] that can either confirm or refute the discrepancy [23].

Acknowledgments

This work was partly supported by the National Research Foundation of Korea (Grant No. NRF-2021R1A2C2009718).

References

 B.D. Fields, K.A. Olive, T.-H. Yeh and C. Young, *Big-Bang Nucleosynthesis after Planck*, JCAP 03 (2020) 010 [1912.01132].

value f = 2387/86 in the case of m = 1 [79].

²With *m* generations of the heavy neutrinos, f' = 11(43 + 7m)/172.

- [2] K. Murai, F. Takahashi, M. Yamada and W. Yin, Can baryon asymmetry be explained by a large initial value before inflation?, Phys. Rev. D 108 (2023) 083518 [2307.03049].
- [3] A.D. Sakharov, Violation of CP Invariance, C asymmetry, and baryon asymmetry of the universe, Pisma Zh. Eksp. Teor. Fiz. 5 (1967) 32.
- [4] M.B. Gavela, P. Hernandez, J. Orloff and O. Pene, Standard model CP violation and baryon asymmetry, Mod. Phys. Lett. A 9 (1994) 795 [hep-ph/9312215].
- [5] M.B. Gavela, M. Lozano, J. Orloff and O. Pene, Standard model CP violation and baryon asymmetry. Part 1: Zero temperature, Nucl. Phys. B 430 (1994) 345 [hep-ph/9406288].
- [6] M.B. Gavela, P. Hernandez, J. Orloff, O. Pene and C. Quimbay, Standard model CP violation and baryon asymmetry. Part 2: Finite temperature, Nucl. Phys. B 430 (1994) 382 [hep-ph/9406289].
- [7] P. Huet and E. Sather, Electroweak baryogenesis and standard model CP violation, Phys. Rev. D 51 (1995) 379 [hep-ph/9404302].
- [8] K. Kajantie, M. Laine, K. Rummukainen and M.E. Shaposhnikov, Is there a hot electroweak phase transition at $m_H \gtrsim m_W$?, Phys. Rev. Lett. 77 (1996) 2887 [hep-ph/9605288].
- M. D'Onofrio and K. Rummukainen, Standard model cross-over on the lattice, Phys. Rev. D 93 (2016) 025003 [1508.07161].
- [10] M. Fukugita and T. Yanagida, Baryogenesis Without Grand Unification, Phys. Lett. B 174 (1986) 45.
- [11] N.S. Manton, Topology in the Weinberg-Salam Theory, Phys. Rev. D 28 (1983) 2019.
- [12] F.R. Klinkhamer and N.S. Manton, A Saddle Point Solution in the Weinberg-Salam Theory, Phys. Rev. D 30 (1984) 2212.
- [13] P. Minkowski, $\mu \to e\gamma$ at a Rate of One Out of 10⁹ Muon Decays?, Phys. Lett. B **67** (1977) 421.
- [14] T. Yanagida, Horizontal gauge symmetry and masses of neutrinos, Conf. Proc. C 7902131 (1979) 95.
- [15] M. Gell-Mann, P. Ramond and R. Slansky, Complex Spinors and Unified Theories, Conf. Proc. C 790927 (1979) 315 [1306.4669].
- [16] R.N. Mohapatra and G. Senjanovic, Neutrino Mass and Spontaneous Parity Nonconservation, Phys. Rev. Lett. 44 (1980) 912.
- [17] J. Schechter and J.W.F. Valle, Neutrino Masses in SU(2) x U(1) Theories, Phys. Rev. D 22 (1980) 2227.
- [18] A. Matsumoto et al., EMPRESS. VIII. A New Determination of Primordial He Abundance with Extremely Metal-poor Galaxies: A Suggestion of the Lepton Asymmetry and Implications for the Hubble Tension, Astrophys. J. 941 (2022) 167 [2203.09617].
- [19] K. Kohri, M. Kawasaki and K. Sato, Big bang nucleosynthesis and lepton number asymmetry in the universe, Astrophys. J. 490 (1997) 72 [astro-ph/9612237].
- [20] M. Kawasaki and K. Murai, Lepton asymmetric universe, JCAP 08 (2022) 041 [2203.09713].
- [21] A.-K. Burns, T.M.P. Tait and M. Valli, Indications for a Nonzero Lepton Asymmetry from Extremely Metal-Poor Galaxies, Phys. Rev. Lett. 130 (2023) 131001 [2206.00693].

- [22] D. Borah and A. Dasgupta, Large neutrino asymmetry from TeV scale leptogenesis, Phys. Rev. D 108 (2023) 035015 [2206.14722].
- [23] M. Escudero, A. Ibarra and V. Maura, Primordial lepton asymmetries in the precision cosmology era: Current status and future sensitivities from BBN and the CMB, Phys. Rev. D 107 (2023) 035024 [2208.03201].
- [24] T. Takahashi and S. Yamashita, Big bang nucleosynthesis and early dark energy in light of the EMPRESS Yp results and the H0 tension, Phys. Rev. D 107 (2023) 103520 [2211.04087].
- [25] S. Kasuya, M. Kawasaki and K. Murai, Enhancement of second-order gravitational waves at Q-ball decay, JCAP 05 (2023) 053 [2212.13370].
- [26] V. Domcke, K. Kamada, K. Mukaida, K. Schmitz and M. Yamada, New Constraint on Primordial Lepton Flavor Asymmetries, Phys. Rev. Lett. 130 (2023) 261803 [2208.03237].
- [27] S. Deng and L. Bian, Constraints on new physics around the MeV scale with cosmological observations, Phys. Rev. D 108 (2023) 063516 [2304.06576].
- [28] F. Gao, J. Harz, C. Hati, Y. Lu, I.M. Oldengott and G. White, Sphaleron freeze-in baryogenesis with gravitational waves from the QCD transition, 2309.00672.
- [29] S.Y. Khlebnikov and M.E. Shaposhnikov, The Statistical Theory of Anomalous Fermion Number Nonconservation, Nucl. Phys. B 308 (1988) 885.
- [30] J.A. Harvey and M.S. Turner, Cosmological baryon and lepton number in the presence of electroweak fermion number violation, Phys. Rev. D 42 (1990) 3344.
- [31] V.A. Kuzmin, V.A. Rubakov and M.E. Shaposhnikov, On the Anomalous Electroweak Baryon Number Nonconservation in the Early Universe, Phys. Lett. B 155 (1985) 36.
- [32] J.A. Casas and A. Ibarra, Oscillating neutrinos and $\mu \rightarrow e, \gamma$, Nucl. Phys. B **618** (2001) 171 [hep-ph/0103065].
- [33] S. Davidson, E. Nardi and Y. Nir, Leptogenesis, Phys. Rept. 466 (2008) 105 [0802.2962].
- [34] R. Fardon, A.E. Nelson and N. Weiner, Dark energy from mass varying neutrinos, JCAP 10 (2004) 005 [astro-ph/0309800].
- [35] A.W. Brookfield, C. van de Bruck, D.F. Mota and D. Tocchini-Valentini, Cosmology of mass-varying neutrinos driven by quintessence: theory and observations, Phys. Rev. D 73 (2006) 083515 [astro-ph/0512367].
- [36] M.M. Reynoso and O.A. Sampayo, Propagation of high-energy neutrinos in a background of ultralight scalar dark matter, Astropart. Phys. 82 (2016) 10 [1605.09671].
- [37] A. Berlin, Neutrino Oscillations as a Probe of Light Scalar Dark Matter, Phys. Rev. Lett. 117 (2016) 231801 [1608.01307].
- [38] Y. Zhao, Cosmology and time dependent parameters induced by a misaligned light scalar, Phys. Rev. D 95 (2017) 115002 [1701.02735].
- [39] G. Krnjaic, P.A.N. Machado and L. Necib, Distorted neutrino oscillations from time varying cosmic fields, Phys. Rev. D 97 (2018) 075017 [1705.06740].
- [40] V. Brdar, J. Kopp, J. Liu, P. Prass and X.-P. Wang, Fuzzy dark matter and nonstandard neutrino interactions, Phys. Rev. D 97 (2018) 043001 [1705.09455].
- [41] H. Davoudiasl, G. Mohlabeng and M. Sullivan, Galactic Dark Matter Population as the Source of Neutrino Masses, Phys. Rev. D 98 (2018) 021301 [1803.00012].

- [42] J. Liao, D. Marfatia and K. Whisnant, Light scalar dark matter at neutrino oscillation experiments, JHEP 04 (2018) 136 [1803.01773].
- [43] F. Capozzi, I.M. Shoemaker and L. Vecchi, Neutrino Oscillations in Dark Backgrounds, JCAP 07 (2018) 004 [1804.05117].
- [44] G.-Y. Huang and N. Nath, Neutrinophilic Axion-Like Dark Matter, Eur. Phys. J. C 78 (2018) 922 [1809.01111].
- [45] Y. Farzan, Ultra-light scalar saving the 3 + 1 neutrino scheme from the cosmological bounds, Phys. Lett. B 797 (2019) 134911 [1907.04271].
- [46] J.M. Cline, Viable secret neutrino interactions with ultralight dark matter, Phys. Lett. B 802 (2020) 135182 [1908.02278].
- [47] A. Dev, P.A.N. Machado and P. Martínez-Miravé, Signatures of ultralight dark matter in neutrino oscillation experiments, JHEP 01 (2021) 094 [2007.03590].
- [48] M. Losada, Y. Nir, G. Perez and Y. Shpilman, Probing scalar dark matter oscillations with neutrino oscillations, JHEP 04 (2022) 030 [2107.10865].
- [49] G.-y. Huang and N. Nath, Neutrino meets ultralight dark matter: 0νββ decay and cosmology, JCAP 05 (2022) 034 [2111.08732].
- [50] E.J. Chun, Neutrino Transition in Dark Matter, 2112.05057.
- [51] A. Dev, G. Krnjaic, P. Machado and H. Ramani, Constraining feeble neutrino interactions with ultralight dark matter, Phys. Rev. D 107 (2023) 035006 [2205.06821].
- [52] G.-y. Huang, M. Lindner, P. Martínez-Miravé and M. Sen, Cosmology-friendly time-varying neutrino masses via the sterile neutrino portal, Phys. Rev. D 106 (2022) 033004 [2205.08431].
- [53] M. Losada, Y. Nir, G. Perez, I. Savoray and Y. Shpilman, Parametric resonance in neutrino oscillations induced by ultra-light dark matter and implications for KamLAND and JUNO, JHEP 03 (2023) 032 [2205.09769].
- [54] D. Brzeminski, S. Das, A. Hook and C. Ristow, Constraining Vector Dark Matter with neutrino experiments, JHEP 08 (2023) 181 [2212.05073].
- [55] G. Alonso-Alvarez, K. Bleau and J.M. Cline, Distortion of neutrino oscillations by dark photon dark matter, Phys. Rev. D 107 (2023) 055045 [2301.04152].
- [56] M. Losada, Y. Nir, G. Perez, I. Savoray and Y. Shpilman, Time dependent CP-even and CP-odd signatures of scalar ultralight dark matter in neutrino oscillations, Phys. Rev. D 108 (2023) 055004 [2302.00005].
- [57] H. Davoudiasl and P.B. Denton, Sterile neutrino shape shifting caused by dark matter, Phys. Rev. D 108 (2023) 035013 [2301.09651].
- [58] T. Gherghetta and A. Shkerin, Probing a local dark matter halo with neutrino oscillations, Phys. Rev. D 108 (2023) 095009 [2305.06441].
- [59] Y. ChoeJo, Y. Kim and H.-S. Lee, Dirac-Majorana neutrino type oscillation induced by a wave dark matter, Phys. Rev. D 108 (2023) 095028 [2305.16900].
- [60] Y. Chen, X. Xue and V. Cardoso, Black Holes as Neutrino Factories, 2308.00741.
- [61] L. Hui, Wave Dark Matter, Ann. Rev. Astron. Astrophys. 59 (2021) 247 [2101.11735].

- [62] J. Preskill, M.B. Wise and F. Wilczek, Cosmology of the Invisible Axion, Phys. Lett. B 120 (1983) 127.
- [63] L.F. Abbott and P. Sikivie, A Cosmological Bound on the Invisible Axion, Phys. Lett. B 120 (1983) 133.
- [64] M. Dine and W. Fischler, The Not So Harmless Axion, Phys. Lett. B 120 (1983) 137.
- [65] PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *PTEP* **2022** (2022) 083C01.
- [66] X.-J. Bi, P.-h. Gu, X.-l. Wang and X.-m. Zhang, Thermal leptogenesis in a model with mass varying neutrinos, Phys. Rev. D 69 (2004) 113007 [hep-ph/0311022].
- [67] P.-h. Gu and X.-j. Bi, Thermal leptogenesis with triplet Higgs boson and mass varying neutrinos, Phys. Rev. D 70 (2004) 063511 [hep-ph/0405092].
- [68] C. Hati and U. Sarkar, Neutrino dark energy and leptogenesis with TeV scale triplets, Eur. Phys. J. C 76 (2016) 236 [1511.02874].
- [69] K.J. Bae, H. Baer, K. Hamaguchi and K. Nakayama, Affleck-Dine Leptogenesis with Varying Peccei-Quinn Scale, JHEP 02 (2017) 017 [1612.02511].
- [70] M.S. Turner, Coherent Scalar Field Oscillations in an Expanding Universe, Phys. Rev. D 28 (1983) 1243.
- [71] M. Doi, T. Kotani and E. Takasugi, Double beta Decay and Majorana Neutrino, Prog. Theor. Phys. Suppl. 83 (1985) 1.
- [72] GERDA collaboration, Search for exotic physics in double-β decays with GERDA Phase II, JCAP 12 (2022) 012 [2209.01671].
- [73] A. De Simone and A. Riotto, On the impact of flavour oscillations in leptogenesis, JCAP 02 (2007) 005 [hep-ph/0611357].
- [74] S. Blanchet, P. Di Bari and G.G. Raffelt, Quantum Zeno effect and the impact of flavor in leptogenesis, JCAP 03 (2007) 012 [hep-ph/0611337].
- [75] S. Blanchet, P. Di Bari, D.A. Jones and L. Marzola, Leptogenesis with heavy neutrino flavours: from density matrix to Boltzmann equations, JCAP **01** (2013) 041 [1112.4528].
- [76] K. Moffat, S. Pascoli, S.T. Petcov, H. Schulz and J. Turner, Three-flavored nonresonant leptogenesis at intermediate scales, Phys. Rev. D 98 (2018) 015036 [1804.05066].
- [77] A. Granelli, K. Moffat and S.T. Petcov, Aspects of high scale leptogenesis with low-energy leptonic CP violation, JHEP 11 (2021) 149 [2107.02079].
- [78] A. Granelli, K. Hamaguchi, N. Nagata, M.E. Ramirez-Quezada and J. Wada, *Thermal leptogenesis in the minimal gauged* $U(1)_{L_{\mu}-L_{\tau}}$ model, *JHEP* **09** (2023) 079 [2305.18100].
- [79] W. Buchmuller, P. Di Bari and M. Plumacher, Cosmic microwave background, matter antimatter asymmetry and neutrino masses, Nucl. Phys. B 643 (2002) 367 [hep-ph/0205349].
- [80] W. Buchmuller, P. Di Bari and M. Plumacher, Leptogenesis for pedestrians, Annals Phys. 315 (2005) 305 [hep-ph/0401240].
- [81] S. Davidson and A. Ibarra, A Lower bound on the right-handed neutrino mass from leptogenesis, Phys. Lett. B 535 (2002) 25 [hep-ph/0202239].

- [82] A. Pilaftsis, Heavy Majorana neutrinos and baryogenesis, Int. J. Mod. Phys. A 14 (1999) 1811 [hep-ph/9812256].
- [83] A. Pilaftsis and T.E.J. Underwood, Resonant leptogenesis, Nucl. Phys. B 692 (2004) 303 [hep-ph/0309342].
- [84] B. Pontecorvo, Mesonium and anti-mesonium, Sov. Phys. JETP 6 (1957) 429.
- [85] Z. Maki, M. Nakagawa and S. Sakata, Remarks on the unified model of elementary particles, Prog. Theor. Phys. 28 (1962) 870.
- [86] PLANCK collaboration, Planck 2018 results. VI. Cosmological parameters, Astron. Astrophys. 641 (2020) A6 [1807.06209].
- [87] SIMONS OBSERVATORY collaboration, The Simons Observatory: Science goals and forecasts, JCAP 02 (2019) 056 [1808.07445].
- [88] K. Abazajian et al., CMB-S4 Science Case, Reference Design, and Project Plan, 1907.04473.