An explicit formula for the zeros of the Riemann zeta function

Eugenio P. Balanzario^{*†} and Daniel Eduardo Cárdenas Romero[†]

[†]Centro de Ciencias Matemáticas, Universidad Nacional Autónoma de México, Apartado Postal 61-3 (Xangari), Morelia Michoacán, México

December 4, 2023

Abstract

We present an explicit formula for a weighted sum over the zeros of the Riemann zeta function. This weighted sum is evaluated in terms of a sum over the prime numbers, weighted with help of the Hermite polynomials. From the explicit formula presented in this note, it follows that prime numbers determine the distribution of the zeros of the Riemann zeta function. In fact, it is possible to compute the zeros of the zeta function without actually using this function.

Introduction

Let $\Lambda(n)$ be the von Mangoldt arithmetical function and w(t) be a real valued function defined for t > 0. Let us consider the function m(y) defined by

$$m(y) = \sum_{n=1}^{\infty} \Lambda(n) w(n \beta y)$$

^{*}ebg@matmor.unam.mx

for y > 0 and where w(t) serves a weight function for $\Lambda(n)$. Assuming w(t) is well behaved, the Mellin transform of m(y) is given by

$$\widehat{m}(s) = \int_{0}^{\infty} y^{s-1} m(y) \, dy = -\frac{\zeta'(s)}{\zeta(s)} \widehat{w}(s)$$

for $s = \sigma + it$ such that $\sigma > 1$ and where $\zeta(s)$ is the Riemann zeta function. From Perron inversion formula,

$$m(y) = \lim_{q \to \infty} \frac{-1}{2\pi i} \int_{2-iq}^{2+iq} \widehat{w}(s) \beta \frac{\zeta'(s)}{\zeta(s)} \beta y^{-s} \, ds.$$
(1)

By shifting the vertical line of integration to the left, we obtain formally, for $x \ge 0$, that

$$\sum_{n=1}^{\infty} \Lambda(n) w\left(\frac{n}{x}\right) = x \beta \widehat{w}(1) - \sum_{\rho} x^{\rho} \beta \widehat{w}(\rho) + E$$
⁽²⁾

where $\rho = \beta + i\gamma$ stands for a non trivial zero of $\zeta(s)$ and E stands for contributions to the integral (1) other than those included in the first two terms on the right of equation (2).

At this point, it is desirable to choose a weight function w(t) in such a way that equation (2) is worth studying. In a previous note [2], we have chosen w(t) to be a probability density function depending on two parameters. With these two parameters we were able to place the bulk of the probability mass given by w(t) at any preassigned place in the positive real line, with the result that we could isolate a finite number of terms on the left hand side of equation (2). In the extreme case that only one term is isolated on the left hand side of equation (2) we obtain as a result a smooth version of Landau's explicit formula [5],

$$\Lambda(x) = -\frac{2\pi}{T} \sum_{0 < \lambda \le T} x^{\rho} + O\left(\frac{\log T}{T}\right).$$

In this note we choose w(t) to be such that its Mellin transform $\widehat{w}(s)$ is equal to a probability density function over the so called critical line,

which is the vertical line defined by $\sigma = 1/2$. Also, $\widehat{w}(s)$ will be such that $\widehat{w}(1/2 - it) = \widehat{w}(1/2 + it)$ for each $t \in \mathbb{R}$. With $\widehat{w}(s)$ depending on two free parameters, we will be able to locate half of the unit mass given by $\widehat{w}(1/2 + it)$ around a small neighborhood of a point on the line s = 1/2 + it with t > 0. The second half of the mass will be located symmetrically along the line segment s = 1/2 - it with t > 0. Thus, by letting this half unit mass be sufficiently concentrated around a suitable point, a single zero, together with its conjugate, will be isolated from the sum over ρ on the right hand side of equation (2).

In this note we consider a weight function w(t) such that its Mellin transform is given by

$$\widehat{w}(s) = c_1 (-1)^k \left(s - \frac{1}{2}\right)^{2k} \exp\left\{\frac{k}{\xi^2} \left(s - \frac{1}{2}\right)^2\right\} \quad \text{where} \quad c_1 = \frac{(k/\xi^2)^{k+\frac{1}{2}}}{\Gamma(k+\frac{1}{2})} \quad (3)$$

for every $s \in \mathbb{C}$. We note that $\widehat{w}(s)$ depends on $\xi > 0$ and $k \in \mathbb{N}$ as two free parameters.

Through all this note we assume the validity of the Riemann hypothesis (H_1) as well as the hypothesis affirming that all zeros of the Riemann zeta function are simple (H_2) .

Now we state the main result of this note.

Theorem 1. For $\xi > 0$ and $k \in \mathbb{N}$, let

$$S(\xi,k) = \frac{1}{\xi} \sqrt{\frac{k}{2\pi}} \sum_{n=1}^{\infty} \exp\left\{-2\frac{k}{\xi^2}(\gamma_n - \xi)^2\right\}$$
(4)

where γ_n is the imaginary part of the n-th zero of the Riemann zeta function in the upper half plane. Assuming hypotheses H_1 and H_2 , then we have

$$S(\xi,k) = \frac{1}{4\pi} \log \frac{\xi}{2\pi} - c_2 \sum_{n=1}^{\infty} \frac{\Lambda(n)}{\sqrt{n}} \exp\left\{-\frac{\xi^2}{4k} \log^2 n\right\} H_{2k}\left(\xi \frac{\log n}{2\sqrt{k}}\right) + E \quad (5)$$

with $E \ll (\sqrt{k}/\xi^2 + \exp\{-\sqrt{k}/\xi\}) \log \xi$ as $\xi \to \infty$ and where

$$c_2 = \frac{1}{2\sqrt{\pi}} \frac{(-1/4)^k}{\Gamma(k+1/2)} \quad and \quad H_k(x) = (-1)^k e^{x^2} \frac{d^k}{dx^k} e^{-x^2}$$

is the k-nth the Hermite polynomial.

Let $\beta > 1$ be a constant real number. In the next section we will find it convenient to put $k = \beta(\xi \log(\xi \log \xi))^2$, and in this case the error term E in the above theorem is given by (as $\xi \to \infty$)

$$E \ll \frac{\log^2 \xi}{\xi}.$$

Determining the zeros of $\zeta(s)$

In this section, we address the question of how many terms in the sum on the right hand side of formula (5) are necessary in order to obtain a good agreement between both sides of the equation. We do this by using the the following rough approximation to the Hermite polynomials

$$H_n(x) \approx e^{\frac{1}{2}x^2} \frac{2^n}{\sqrt{\pi}} \Gamma\left(\frac{n+1}{2}\right) \cos\left(x\sqrt{2n} - \frac{n\pi}{2}\right)$$

([1], pages 508-510, formulas 13.6.38 and 13.5.16). From this approximation to the Hermite polynomials, we obtain the following approximate expression for the weight function

$$\widetilde{w}(x) := \frac{1}{2\pi} \frac{1}{\sqrt{x}} \exp\left\{-\frac{1}{8\alpha} \log^2 x\right\} \cos(\xi \log(x) - k\beta\pi) \tag{6}$$

where here and in the remainder of this note we write $\alpha = k/\xi^2$. From equation (6) we have

$$|\widetilde{w}(x)| \le \exp\Big\{-\frac{1}{8lpha}\log^2 x\Big\}.$$

Given a small number $\epsilon > 0$, let $\tau = \tau(\epsilon)$ be the number of terms in the sum on the right hand side of formula (5) needed in order to estimate $S(\xi, k)$ within an error not greater than ϵ . For the estimation of τ , notice that

$$\sum_{n>\tau} \Lambda(n) \mathfrak{G}|\widetilde{w}(n)| \leq \int_{\tau}^{\infty} \log(x) \mathfrak{G} \exp\left\{-\frac{1}{8\alpha} \log^2 x\right\} dx$$
$$= 4\alpha \exp\left\{-\left(\frac{\log \tau}{2\sqrt{2\alpha}} - \sqrt{2\alpha}\right)^2 + 2\alpha\right\}$$



Figure 1: The graph of $S(\xi, 2\xi^2)$ where $S(\xi, k)$ is as in equation (4).

Equating this expression to ϵ , clearing for τ and writing k/ξ_0^2 in place of α , we see that for the sum $\sum_{n>\tau} \Lambda(n)\beta|\widetilde{w}(n)|$ to be less than or equal to ϵ , it suffices if

$$\tau = \exp\left\{4\left(\frac{k}{\xi_0^2} + \sqrt{\frac{k}{\xi_0^2}}\sqrt{\frac{k}{\xi_0^2} + \frac{1}{2}\log\frac{4k}{\epsilon\beta\xi_0}}\right)\right\}.$$
(7)

From equation (4) it follows that $S(\xi, k)$ places approximately half a unit mass in every sufficiently small neighborhood of ξ_0 if and only if $\xi_0 = \gamma_n$ for some $n \in \mathbb{N}$. This fact is illustrated in figure 1 where the graph was computed by using the right hand side of formula (5) but with $\tilde{w}(x)$, as given in equation (6), in place of w(x). In this figure, the imaginary part of the first three zeros of the Riemann zeta function are marked with a vertical line. We appreciate the accumulation of half a unit mass around the imaginary parts of zeros of the Riemann zeta function. This accumulation of mass allows us to decide whether or not a given number ξ_0 is equal to the imaginary part of one of the zeros of the Riemann zeta function.

Around each point $\xi_0 = \gamma_n$ with $n \in \mathbb{N}$, the local variance of $S(\xi, k)$ is equal to $\sigma^2 = \gamma_n^2/8k$. On the other hand, for the imaginary part of the *n*-th zero of the Riemann zeta function it is known that the following asymptotic relation takes place (see [4], page 20)

$$\gamma_n \sim \frac{2\pi \beta n}{\log n}$$

as $n \to \infty$. Hence, in order to place half a unit mass in a small neighborhood of γ_n , the standard deviation of $S(\xi, k)$ around such a neighborhood of $\xi_0 = \gamma_n$ must be of an order of magnitude of

$$\frac{\xi_0}{\sqrt{k}} = \frac{1}{\sqrt{\beta}\log n}$$

where $\beta > 1$ is moderately large but fixed. The larger β is, the greater the half mass will be concentrated around ξ_0 . Putting $k = \beta(\xi_0 \log n)^2$ in equation (7) we obtain

$$\tau = \exp\left\{4\left(\beta\log^2 n + \sqrt{\beta}\log n\sqrt{\beta\log^2 n + \log 2\sqrt{\beta/\epsilon}\log n}\right)\right\}.$$

If ϵ is fixed, then we have that $\tau \sim \exp\{4\beta \log^2 n\}$ as $n \to \infty$. With this expression we have that $\exp\{\beta' \log^2(\xi_0 \log \xi_0)\}$ is an estimate for the number of primes necessary in order to determine whether a number $1/2 + i\xi_0$ is a zero of the Riemann zeta function for each $\xi_0 \in \mathbb{R}$.

In this section we have seen that, assuming the Riemann hypothesis and the simplicity of the zeros of $\zeta(s)$, knowledge of the distribution of prime numbers and its integral powers is sufficient for the determination of the distribution of the nontrivial zeros of the Riemann zeta function. Thus, it is possible to compute the zeros of $\zeta(s)$ without actually using this function. However, a large number of primes are necessary for the determination of the zeros of $\zeta(s)$.

Proof of theorem 1

Lemma 1. Let $\widehat{w}(s)$ be is as in (3), c_2 as in theorem 1 and $\alpha = k/\xi^2$. Then

$$w(x) = \frac{c_2}{\sqrt{x}} \exp\left\{-\frac{1}{4\alpha}\log^2 x\right\} H_{2k}\left(\frac{\log x}{2\sqrt{\alpha}}\right).$$
(8)

Proof. Let us consider the integral

$$I = \int_{\sigma_0 - i\infty}^{\sigma_0 + i\infty} \left(s - \frac{1}{2}\right)^{2k} e^{\alpha (s - \frac{1}{2})^2} e^{-sx} \, ds \quad \text{where} \quad \sigma_0 = \frac{1}{2} \left(1 + \frac{x}{\alpha}\right).$$

We must show that $I = 2\pi i\beta c_1^{-1}(-1)^k\beta w(e^x)$ with w(x) as given in equation (8). Now,

$$I = \exp\left\{\frac{\alpha}{4} - \alpha\left(\frac{1}{2} + \frac{x}{2\alpha}\right)^2\right\} \int_{\sigma_0 - i\infty}^{\sigma_0 + i\infty} \left(s - \frac{1}{2}\right)^{2k} \exp\left\{\alpha\left(s - \frac{1}{2} - \frac{x}{2\alpha}\right)^2\right\} ds$$
$$= \exp\left\{-\frac{x}{2} - \frac{x^2}{4\alpha}\right\} \int_{-\infty}^{+\infty} \left(\frac{x}{2\alpha} + it\right)^{2k} e^{-\alpha t^2} i dt$$
$$= \exp\left\{-\frac{x}{2} - \frac{x^2}{4\alpha}\right\} \frac{i}{(2\alpha)^k} \int_{-\infty}^{+\infty} \left(\frac{x}{\sqrt{2\alpha}} + it\right)^{2k} e^{-\frac{1}{2}t^2} dt$$
$$= \exp\left\{-\frac{x}{2} - \frac{x^2}{4\alpha}\right\} i \sqrt{\frac{\pi}{\alpha}} \left(\frac{1}{4\alpha}\right)^k H_{2k} \left(\frac{x}{2\sqrt{\alpha}}\right)$$

where for the last step in the above calculation, we have made use of the following formula

$$\int_{-\infty}^{+\infty} (\sqrt{2}x + it)^{2k} e^{-\frac{1}{2}t^2} dt = \frac{\sqrt{2\pi}}{2^k} H_{2k}(x)$$

(see formula 8.951 in [3]).

In the next lemma, we will let g(s) be the logarithmic derivative of $\chi(s)$ where

$$\chi(s) = 2^s \pi^{s-1} \sin\left(\frac{\pi}{2}s\right) \Gamma(1-s) \quad \text{and} \quad \zeta(s) = \chi(s)\zeta(1-s) \quad (9)$$

is the functional equation satisfied by the Riemann zeta function (see [6], pages 234 and 270). Furthermore, the reader will notice that the proof of the lemma is an application of the saddle point method of complex analysis.

Lemma 2. Let

$$g(s) = \log 2\pi + \frac{\pi}{2} \tan\left(\frac{\pi s}{2}\right) - \psi(s) \tag{10}$$

where $\psi(s) = \Gamma'(s)/\Gamma(s)$ is the digamma function. Then, as $\xi = \sqrt{k/\alpha} \to \infty$, we have that

$$J = \int_{-\infty}^{+\infty} \widehat{w} \left(\frac{1}{2} - it\right) g\left(\frac{1}{2} - it\right) dt = \log \frac{\xi}{2\pi} + O\left(\frac{\alpha^{\frac{3}{4}} \log \xi}{\sqrt{k}} + \frac{\log \xi}{e^{\sqrt{\alpha}}}\right).$$

Proof. If we write

$$J_1^{\pm} = \int_0^\infty \widehat{w}\left(\frac{1}{2} \pm it\right) \tan\left(\frac{\pi}{4} \pm i\frac{t}{2}\right) dt \quad \text{and} \quad J_2^{\pm} = \int_0^\infty \widehat{w}\left(\frac{1}{2} \pm it\right) \beta\psi\left(\frac{1}{2} \pm it\right) dt$$

then we have

$$J = \log 2\pi + \frac{\pi}{2}(J_1^- + J_1^+) + (J_2^- + J_2^+).$$

On the other hand,

$$\tan\left(\frac{\pi}{4} \pm i\frac{t}{2}\right) = \pm i + O(e^{-t}) \quad \text{and} \quad \psi\left(\frac{1}{2} \pm it\right) = \log t \pm \frac{\pi}{2}i + O\left(\frac{1}{t}\right)$$

where the second of these formulas follows from the asymptotic expansion for the digamma function (Abranowitz formula 6.3.18)

$$\psi(z) = \log z - \frac{1}{2z} - \frac{1}{12z^2} + \frac{1}{120z^4} - \frac{1}{252z^6} + \cdots,$$

for $z \to \infty$ in $|\arg z| < \pi$.

For the estimation of J_2^{\pm} we first assume that $|t - \xi| \leq \alpha^{-\frac{1}{4}}$. Then we have

$$\widehat{w}\left(\frac{1}{2} + it\right) = \frac{\alpha^{k+\frac{1}{2}}}{\Gamma(k+\frac{1}{2})} \exp\{-\alpha t^2 + 2k \log t\} \\ = \frac{\alpha^{k+\frac{1}{2}}}{\Gamma(k+\frac{1}{2})} e^{-k} \left(\frac{k}{\alpha}\right)^k \exp\{-2\alpha(t-\xi)^2 + O\left(\sqrt{\frac{\alpha^3}{k}}(t-\xi)^3\right)\} \\ = \sqrt{\frac{\alpha}{2\pi}} \exp\{-2\alpha(t-\xi)^2\} \left(1 + O\left(\frac{1}{k}\right)\right) \left(1 + O\left(\sqrt{\frac{\alpha^3}{k}}(t-\xi)^3\right)\right).$$

Hence

$$\widehat{w}\left(\frac{1}{2}+it\right) = \sqrt{\frac{\alpha}{2\pi}} \exp\{-2\alpha(t-\xi)^2\}\left(1+O\left(\frac{\alpha^3}{\sqrt{k}}\right)\right).$$
(11)

Furthermore,

$$\begin{split} & \int_{\xi-\alpha^{-\frac{1}{4}}}^{\xi+\alpha^{-\frac{1}{4}}} \widehat{w}\Big(\frac{1}{2}+it\Big)\log(t) \, dt \\ &= \left\{1+O\Big(\frac{\alpha^{\frac{3}{4}}}{\sqrt{k}}\Big)\right\}\sqrt{\frac{\alpha}{2\pi}} \int_{\xi-\alpha^{-\frac{1}{4}}}^{\xi+\alpha^{-\frac{1}{4}}} \log(t)\beta e^{-2\alpha(t-\xi)^{2}} \, dt \\ &= \left\{1+O\Big(\frac{\alpha^{\frac{3}{4}}}{\sqrt{k}}\Big)\right\}\frac{1}{\sqrt{2\pi}} \int_{-\alpha^{\frac{1}{4}}}^{\alpha^{\frac{1}{4}}} \log\Big(\xi+\frac{t}{\sqrt{\alpha}}\Big)\beta e^{-2t^{2}} \, dt \\ &= \left\{1+O\Big(\frac{\alpha^{\frac{3}{4}}}{\sqrt{k}}\Big)\right\}\Big\{\log\xi+O\Big(\frac{\alpha^{-\frac{1}{4}}}{\xi}\Big)\Big\}\Big\{\frac{1}{2}+O(e^{-2\sqrt{\alpha}})\Big\} \\ &= \frac{1}{2}\log\xi+O\Big(\frac{\alpha^{\frac{3}{4}}\log\xi}{\sqrt{k}}\Big). \end{split}$$

For the estimation of the integral defining J_2^{\pm} over the range of t such that $|t-\xi| > \alpha^{-\frac{1}{4}}$, we first notice that

$$\begin{aligned} -\alpha t^2 + 2k\log t + k - k\log \frac{k}{\alpha} &= -\alpha (t-\xi)^2 - 2k - 2k \int_{\xi}^{t} \frac{t-y}{y^2} \, dy \\ &\leq -\alpha (t-\xi)^2. \end{aligned}$$

Hence we have

$$\left|\widehat{w}\left(\frac{1}{2}+it\right)\right| = \frac{\alpha^{k+\frac{1}{2}}}{\Gamma(k+\frac{1}{2})}\exp\{-\alpha t^2 + 2k\log t\}$$
$$\ll \sqrt{\alpha}\left(\frac{\alpha}{k}\right)^k e^k \exp\{-\alpha t^2 + 2k\log t\} \ll \sqrt{\alpha}\exp\{-\alpha(t-\xi)^2\}.$$

Thus,

$$\sqrt{\alpha} \int_{\xi+\alpha^{-\frac{1}{4}}}^{\infty} \log(t) \beta e^{-\alpha(t-\xi)^2} dt \ll \int_{\alpha^{\frac{1}{4}}}^{\infty} t \beta e^{-t^2} dt = \frac{1}{2} e^{-\sqrt{\alpha}}.$$

On the other hand

$$\sqrt{\alpha} \int_{1}^{\xi - \alpha^{-\frac{1}{4}}} \log(t) \beta e^{-\alpha(t-\xi)^2} dt \le \sqrt{\alpha} \log(\xi) \int_{-\infty}^{-\alpha^{-\frac{1}{4}}} e^{-\alpha t^2} dt \ll \log(\xi) e^{-\sqrt{\alpha}}.$$

Therefore

$$J_{2}^{-} + J_{2}^{+} = \log \xi + O\left(\frac{\alpha^{\frac{3}{4}} \log \xi}{\sqrt{k}} + \frac{\log \xi}{e^{\sqrt{\alpha}}}\right).$$

In a similar fashion we see that $J_1^- + J_1^+ = O(e^{-\xi})$. This finishes the proof of the lemma.

Lemma 3. We have that

$$c_1 \sum_{n=1}^{\infty} \gamma_n^{2k} \beta e^{-\alpha \gamma_n^2} = \sqrt{\frac{\alpha}{2\pi}} \sum_{|\gamma_n - \xi| \le \alpha^{-\frac{1}{4}}} \exp\{-2\alpha (\gamma_n - \xi)^2\} + O\left(\frac{\alpha}{\sqrt{k}} \log \xi + e^{-\sqrt{\alpha}}\right).$$

Proof. We write $s_1 + s_2$ for the sum we want to estimate, where s_1 is the sum over γ_n such that $|\gamma_n - \xi| \leq \alpha^{-\frac{1}{4}}$ and s_2 is sum over γ_n such that $|\gamma_n - \xi| > \alpha^{-\frac{1}{4}}$. Because of equation (11), for s_1 we have,

$$s_{1} = \sqrt{\frac{\alpha}{2\pi}} \left\{ 1 + O\left(\frac{\alpha^{\frac{3}{4}}}{\sqrt{k}}\right) \right\} \sum_{|\gamma_{n}-\xi| \le \alpha^{-\frac{1}{4}}} \exp\{-2\alpha(\gamma_{n}-\xi)^{2}\}$$
$$= \sqrt{\frac{\alpha}{2\pi}} \sum_{|\gamma_{n}-\xi| \le \alpha^{-\frac{1}{4}}} \exp\{-2\alpha(\gamma_{n}-\xi)^{2}\} + O\left(\frac{\alpha^{\frac{3}{4}}}{\sqrt{k}}\sqrt{\alpha}\sum_{|\gamma_{n}-\xi| \le \alpha^{-\frac{1}{4}}}1\right).$$

We recall that if N(t) is the number of zeros of $\zeta(s)$ such that $0 < \gamma_n \leq t$, then $N(t) \simeq t \log t$. Therefore,

$$s_1 = \sqrt{\frac{\alpha}{2\pi}} \sum_{|\gamma_n - \xi| \le \alpha^{-\frac{1}{4}}} \exp\{-2\alpha(\gamma_n - \xi)^2\} + O\left(\frac{\alpha}{\sqrt{k}}\log\xi\right).$$

By proceeding as in the proof of lemma 2 we see that $s_2 \ll \exp\{-\sqrt{\alpha}\}$. This finishes the proof of lemma 3.

With the help of the previous lemmas, we can proceed to the proof of theorem 1. From Perron inversion formula (1) we have

$$m(1) = -\frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \widehat{w}(s) \frac{\zeta'(s)}{\zeta(s)} \, ds.$$

By shifting the vertical line of integration from $\sigma = 2$ to $\sigma = -1$, we get

$$m(1) = c_1 \beta \frac{(-1)^{k+1}}{4^k} e^{\frac{1}{4}\alpha} - 2c_1 \sum_{n=1}^{\infty} \gamma_n^{2k} \beta e^{-\alpha \gamma_n^2} - \frac{1}{2\pi i} \int_{-1-i\infty}^{-1+i\infty} \widehat{w}(s) \frac{\zeta'(s)}{\zeta(s)} \, ds.$$

Because of lemma 3 the second term on the right hand side of this equations can be written as $2S(\xi, k) + E$. Because of the functional equation (9), we can write the last integral as

$$-\frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \widehat{w}(1-s) \left(g(s) - \frac{\zeta'(s)}{\zeta(s)}\right) ds$$

where g(s) is as in equation (10). If we set $\hat{k}(s) = \hat{w}(1-s)$, then, from the operational properties of the Mellin transform, we have that $k(x) = x^{-1}w(x^{-1})$ and therefore,

$$\frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \widehat{w}(1-s) \frac{\zeta'(s)}{\zeta(s)} \, ds = -\sum_{n=1}^{\infty} \Lambda(n) \frac{1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \widehat{k}(s) \beta n^{-s} \, ds$$
$$= -\sum_{n=1}^{\infty} \frac{\Lambda(n)}{n} w \left(\frac{1}{n}\right)$$
$$= -c_2 \sum_{n=1}^{\infty} \frac{\Lambda(n)}{\sqrt{n}} \exp\left\{\frac{-1}{4\alpha} \log^2 n\right\} H_{2k} \left(\frac{\log n}{2\sqrt{\alpha}}\right)$$
$$= -m(1).$$

Hence,

$$2\mathfrak{B}m(1) = c_1 \mathfrak{B}\frac{(-1)^{k+1}}{4^k} e^{\frac{1}{4}\alpha} - 2\mathfrak{B}S(\xi,k) + \frac{1}{2\pi}\tilde{J} + O\left(\frac{\alpha\log\xi}{\sqrt{k}} + \frac{\log\xi}{e^{\sqrt{\alpha}}}\right)$$

where

$$\frac{1}{2\pi}\tilde{J} = \frac{-1}{2\pi i} \int_{2-i\infty}^{2+i\infty} \widehat{w}(1-s) \beta g(s) \, ds = \frac{-1}{2\pi} \int_{-\infty}^{+\infty} \widehat{w}\left(\frac{1}{2}-it\right) g\left(\frac{1}{2}-it\right) \, dt$$
$$= \frac{1}{2\pi} \log \frac{\xi}{2\pi} + O\left(\frac{\alpha \log \xi}{\sqrt{k}} + \frac{\log \xi}{e^{\sqrt{\alpha}}}\right)$$

the last equality being a consequence of lemma 2. This finishes the proof.

References

- Abramowitz, M.; Stegun, I.A. Handbook of mathematical functions with formulas, graphs, and mathematical tables. National Bureau of Standards Applied Mathematics Series, No. 55, 1964.
- [2] Balanzario, E.P.; Cárdenas Romero, E.D.; Chacón Serna, R. A smooth version of Landau's explicit formula. Manuscript submitted, 2023.
- [3] Gradshteyn, I.S.; Ryzhik, I.M. Table of integrals, series, and products. Academic Press, 2007.
- [4] Ivić, A. The Riemann zeta-function. Dover Publications, Inc., 2003.
- [5] Landau, E. Uber die Nullstellen der Zetafunktion. Math. Ann. 71 (1912) 548-564.
- [6] Tenenbaum, G. Introduction to analytic and probabilistic number theory. American Mathematical Society, 2015.