Output-sensitive Complexity of Multi-Objective Integer Network Flow Problems

David Könen^{1,2} and Michael Stiglmayr^{1,3}

¹University of Wuppertal, School of Mathematics and Natural Sciences, Optimization Group, Gaußstraße 20, 42103 Wuppertal, Germany,

²Corresponding author, E-Mail: [koenen@uni-wuppertal.de,](mailto:koenen@uni-wuppertal.de) ORCID-ID: 0000-0003-1747-8791 ³E-Mail: [stiglmayr@uni-wuppertal.de,](mailto:stiglmayr@uni-wuppertal.de) ORCID-ID: 0000-0003-0926-1584

This paper addresses the *output-sensitive* complexity for linear multi-objective integer minimum cost flow (MOIMCF) problems and provides insights about the time complexity for enumerating all supported nondominated vectors. The paper shows that there can not exist an output-polynomial time algorithm for the enumeration of all supported nondominated vectors that determine the vectors in an ordered way in the outcome space unless $\mathbf{NP} = \mathbf{P}$. Moreover, novel methods for identifying supported nondominated vectors in bi-objective minimum cost flow (BOIMCF) problems are proposed, accompanied by a numerical comparison between decisionand objective-space methods. A novel, equivalent and more compact formulation of the minimum cost flow ILP formulation used in the *ε*-scalarization approach is introduced, demonstrating enhanced efficiency in the numerical tests.

Keywords: minimum cost flow, multi-objective integer linear programming, multiobjective network flow, complexity theory, output-sensitive, weakly supported, outputpolynomial algorithm

1 Introduction

The *multi-objective integer minimum cost flow* (MOIMCF) problem can be written as

$$
\min \ c(f)
$$
\n
$$
\text{s.t.} \sum_{j:(i,j)\in A} f_{ij} - \sum_{j:(j,i)\in A} f_{ji} = b_i \qquad i \in V
$$
\n
$$
l_{ij} \le f_{ij} \le u_{ij} \qquad (i,j) \in A
$$
\n
$$
f \in \mathbb{Z}_{\ge}^m
$$
\n(MOMCF)

in a directed graph $D = (V, A)$ with *n* nodes $V = \{v_1, \ldots, v_n\}$ and *m* arcs $A =$ ${e_1, \ldots, e_m}$. Let l_{ij} and u_{ij} denote the integer-valued, non-negative, finite lower and upper capacity bounds, respectively, for each arc $(i, j) \in A$. Here, b_i is the integer-valued *flow balance* of the node $i \in V$ where $b_i > 0$ are supply nodes, $b_i < 0$ are demand nodes and $b_i = 0$ are transshipment nodes.

We denote a feasible solution $f = (f_{ij})_{ij \in A} \in \mathbb{Z}_{\geq}^m$ as *flow* and define the *cost* of a flow *f* using the (vector valued) objective function $c: \mathbb{R}^m \to \mathbb{R}^d$ and the corresponding cost matrix $C \in \mathbb{R}^{d \times m}$ as

$$
c(f) := (c^1(f), \dots, c^d(f))^{\top} := \Big(\sum_{(i,j)\in A} c_{ij}^1 f_{ij}, \dots, \sum_{(i,j)\in A} c_{ij}^d f_{ij}\Big)^{\top} = C \cdot f.
$$

Note that the *(continuous) multi-objective minimum cost flow* (MOMCF) problem is the LP-relaxation of MOIMCF. From now on, we assume that *D* is connected (see [Ahuja et al.](#page-28-0) [1993](#page-28-0)) and that there is at least one feasible flow.

Then, we denote the set of feasible outcome vectors in the objective space by \mathcal{Y} := $\{C f : f \in \mathcal{X}\}\$, where X is the set of all feasible flows. We assume that the objective functions are conflicting, which implies that no solution minimizes all objectives simultaneously. Throughout this article, we will use the *Pareto concept of optimality*, which is based on the componentwise order in \mathbb{R}^d . We write $y^1 \leq y^2$ if $y^1_k \leq y^2_k$ for $k = 1, \ldots, d, y^1 \leq y^2$ if $y^1 \leq y^2$ and $y^1 \neq y^2$, and $y^1 < y^2$ if $y^1_k < y^2_k$, $k = 1, \ldots, d$. Furthermore, let $\mathbb{R}^d_{\geq} := \{x \in \mathbb{R}^d : x \geq 0\}$ denote the non-negative orthant of \mathbb{R}^d . Its interior \mathbb{R}^d_\geq is defined accordingly.

A feasible solution $f^* \in \mathcal{X}$ is *efficient* if there does not exist any other feasible solution $f \in \mathcal{X}$ such that $c(f) \leq c(f^*)$. If f^* is efficient, $c(f^*)$ is called *nondominated point*. If $f, f' \in \mathcal{X}$ are such that $c(f) \leq c(f')$ we say that *f dominates* f' and $c(f)$ *dominates* $c(f')$. Feasible solutions $f, f' \in \mathcal{X}$ are *equivalent* if $c(f) = c(f')$. The set of efficient flows is denoted by $\mathcal{X}_E \subset \mathcal{X}$ and the set of nondominated vectors by $\mathcal{Y}_\mathcal{N} \subset \mathcal{Y}$. Moreover, a feasible flow *f* is called weakly efficient if there is no other flow f' such that $c(f') < c(f)$.

For the weighted sum method, we define the set of *normalized weighting vectors* as the set $\Lambda_d = \{\lambda \in \mathbb{R}^d_{\geq} : \|\lambda\|_1 = 1\}$ or $\Lambda_d^0 = \{\lambda \in \mathbb{R}^d_{\geq} : \|\lambda\|_1 = 1\}$ if weights equal to zero are included. The *weighted-sum scalarization* with $\lambda \in \Lambda_d$ or $\lambda \in \Lambda_d^0$ is defined as the parametric program $P_{\lambda} \coloneqq \min\{\lambda^{\top} C f : f \in \mathcal{X}\}\.$ Note that the problem P_{λ} is as easy to

solve as the associated single objective problem, as long as the encoding lengths of the components of λ are not too large. Note that λ is not part of the original input. However it is shown that these lengths can be bounded by $\mathcal{O}(\text{poly}(n,m))$ [\(Bökler and Mutzel,](#page-28-1) [2015](#page-28-1)). If $\lambda \in \Lambda_d^0$ every optimal solution to P_λ is a weakly efficient solution. Moreover, every optimal solution of P_λ is efficient if $\lambda \in \Lambda_d$ [\(Ehrgott](#page-28-2), [2005\)](#page-28-2).

Then we can distinguish between several classes of efficient solutions.

- 1. *Weakly supported efficient solutions* are efficient solutions that are optimal solutions of P_λ for $\lambda \in \Lambda_d^0$, i.e., an optimal solution to a weighted sum singe-objective problem with weights strictly or equal to zero. Their images in the objective space are weakly *supported nondominated points*. All weakly supported nondominated vectors are located on the boundary of the *upper image* $P := \text{conv}(\mathcal{Y}) + \mathbb{R}^d_{\geq}$.
- 2. An efficient solution is called *supported efficient solution* if it is an optimal solution to the weighted sum scalarization P_λ for $\lambda \in \Lambda_d$, i.e., an optimal solution to a single objective weighted-sum problem where the weights are strictly positive. Its image is called *supported nondominated vector*; we use the notations \mathcal{X}_{SE} and \mathcal{Y}_{SN} . A supported nondominated vector is located on the *nondominated frontier* defined as the set $\{y \in \text{conv}(\mathcal{Y}_N): \text{ conv}(\mathcal{Y}_N) \cap (y - \mathbb{R}^d) \}$, i.e., located on the union of the *maximal nondominated faces*. Hereby, a *nondominated face* $F \subset \mathcal{Y}$ is defined as a face of the outcome space where all its points are nondominated, even in the relaxation (continuous case) of the integer problem. A face *F* is called *maximally nondominated* if there is no other nondominated face *G* such that *F* is a subset of *G* and the dimension of *G* would be greater than the dimension of *F*. Let $F_\mathcal{X}$ be the set of the preimage of a maximally nondominated face F_y of polytope \mathcal{Y} , i.e., all solutions whose image lies in F_y . We use the expression *maximally efficient face* for F_X . Even if we use the expression face for F_X , it must not be a face of the decision space since more than one feasible solution could have a vertex of $\mathcal Y$ as an image, and they may lie in different faces of X .
- 3. *Extreme supported solutions* are those solutions whose image lies on the vertex set of the upper image. Their image is called an *extreme supported nondominated vector*. We use the notation \mathcal{Y}_{EN} for the set of extreme nondominated points.
- 4. *Unsupported efficient solutions* are efficient solutions that are not optimal solutions of P_{λ} for any $\lambda \in \Lambda_d^0$. *Unsupported nondominated vectors* lie in the interior of the upper image.

Figure [1.1](#page-3-0) illustrates supported extreme, supported, and unsupported nondominated points as well as the upper image in the bi-objective case. Whereas Figure Figure [1.2](#page-4-0) illustrates extreme, supported, and weakly supported nondominated vectors in the threedimensional case.

Due to the total unimodularity of MOMCF, each extreme supported nondominated point of MOMCF has an integer preimage since *u* and *b* are integral (see [Ahuja et al.](#page-28-0) [1993](#page-28-0)). In other words, the sets of extreme supported nondominated points of MOMCF and MOIMCF, and thus the respective upper images, coincide. In the remainder of

- extreme supported nondominated
- supported nondominated \bullet
- unsupported nondominated \bullet
- \times dominated

Figure 1.1: Illustration of the upper image $P = \text{conv}(\mathcal{Y}) + \mathbb{R}^2$ and the different solution types of the BOIMCF problem presented in Figure [3.1.](#page-8-0) Note that the origin starts at (90*,* 90)[⊤] and not all dominated vectors are displayed in the figure.

this paper, only integer flows are considered, and from now on, flow always refers to an integer flow.

Since the minimum cost flow problem is a fundamental, well-studied problem in combinatorial optimization [\(Ahuja et al.,](#page-28-0) [1993](#page-28-0); [Bertsekas](#page-28-3), [1998](#page-28-3)), various polynomial algorithms exist for the single-objective version of this problem. However real-world problems often involve multiple conflicting objectives. These multi-objective minimum cost flow problems are much harder to solve than the single-objective case. MOIMCF had been reviewed in [Hamacher et al.](#page-28-4) [\(2007](#page-28-4)), where the authors comment on the lack of efficient algorithms. However, since then, there still needs to be more efficient algorithms for determining supported efficient solutions for MOIMCF problems, and we give an overview of new results below.

There are several algorithms to determine efficient solutions for bi-objective integer minimum cost flow (BOIMCF) problems, e.g., [\(Eusébio and Figueira,](#page-28-5) [2009a](#page-28-5); [Eusébio et al.](#page-28-6), [2014](#page-28-6); [Raith and Ehrgott](#page-29-0), [2009a](#page-29-0); [Sedeño-Noda](#page-29-1), [2001](#page-29-1); [Sedeño-Noda and Gonzalez-Martin](#page-29-2), [2000](#page-29-2); [Sedeño-Noda and González-Martin](#page-29-3), [2003](#page-29-3)). Raith and Sedeño-Noda introduced an enhanced parametric approach to determine all extreme efficient solutions for BOIMCF problems [\(Raith and Sedeño-Noda,](#page-29-4) [2017](#page-29-4)). However, there are only quite a few specific methods designed to determine all (or subsets) of the nondominated vectors in the objective space (nor the corresponding efficient solutions in the decision space) for MOIMCF problems [\(Eusébio and Figueira](#page-28-7), [2009c;](#page-28-7) [Fonseca et al.,](#page-28-8) [2010](#page-28-8); [Sun,](#page-29-5) [2011](#page-29-5)).

In [Könen and Siglmayr](#page-28-9) [\(2023\)](#page-28-9) an *output-polynomial* time algorithm is presented to determine all efficient supported flows for MOIMCF problems. Unfortunately, this method is not powerful enough to determine all nondominated supported vectors in outputpolynomial time, since there may be exponentially many flows mapping to the same vectors.

The analysis of the output-sensitive complexity of specific problems has gained im-

Figure 1.2: The convex hull $conv(Y)$ of a multi-objective minimum cost flow problem [\(Könen and Siglmayr,](#page-28-9) [2023](#page-28-9)) is shown in blue, its containing hyperplane $h = \{y \in$ $\mathcal{Y}: 0.5 y_1 + 0.5 y_2 = 12$ is depicted in yellow. The extreme nondominated vectors are $y^1 = (6, 16, 6)^{T}, y^2 = (12, 12, 6)^{T}, y^3 = (16, 8, 10)^{T}$. The red dots on the edge between (*y* 2 *, y*³) are supported nondominated vectors (13*,* 11*,* 7)[⊤]*,* (14*,* 10*,* 8)[⊤]*,* (11*,* 13*,* 9)[⊤]. The light red dots on the edge between the edge (y^1, d^1) are weakly supported vectors, which would be dominated in the non-integer case by the points on the edge (y^1, y^2) . The maximally nondominated faces are the edges (y^1, y^2) and (y^2, y^3) .

portance in recent years and several combinatorial problems have been studied, e.g., multi-objective shortest path [\(Bökler et al.,](#page-28-10) [2017](#page-28-10)), multi-objective spanning tree problems [\(Bökler et al.,](#page-28-10) [2017;](#page-28-10) [Okamoto and Uno](#page-29-6), [2011](#page-29-6)), mining frequent closed attribute trees [\(Arimura and Uno](#page-28-11), [2005](#page-28-11)), general multi-objective combinatorial optimization problems and multi-objective linear programs [\(Bökler and Mutzel,](#page-28-1) [2015](#page-28-1)).

The purpose of this paper is to give insights into the time complexity for the enumeration of all supported nondominated vectors for MOIMCF problems. The paper shows that there can not exist an output-polynomial time algorithm for the enumeration of all supported nondominated vectors that determine the vectors in an ordered way in the outcome space unless $NP = P$. The paper shows that the next best distinct cost flow can be determined in $\mathcal{O}(n^3)$ given an initial optimal flow. This result derives an improved algorithm to determine all supported nondominated vectors for BOIMCF problems to the algorithm presented in [Könen and Siglmayr](#page-28-9) [\(2023](#page-28-9)) if the number of branches needed is significantly smaller than the number of supported efficient solutions. The paper presents an example in which the adjusted algorithm saves an exponential amount of to considered flows to get all supported nondominated vectors.

While the algorithm in [Könen and Siglmayr](#page-28-9) [\(2023\)](#page-28-9) is a *decision-space* method, the paper presents also *objective-space* methods to determine all supported nondominated

vectors, based on the ε -scalarization method. In addition, the paper gives an equivalent more compact formulation for the ILP used in the *ε*-scalarization method, which can be solved more efficiently. Note, that the *ε*-scalarization could also be used to get all nondominated vector, even in higher dimension. The adjusted algorithm can be seen as an mixture of a decision- and objective-space method.

The remainder of the paper is structured as follows. Section [2](#page-5-0) introduces the theory of output-sensitive time complexity and summarizes the existing results from the literature on the existence of output-polynomial time algorithms for the MOIMCF problem w.r.t. different solution concepts. While Section [3](#page-7-0) provides insight about the output-sensitive complexity for MOIMCF problems. In Section [4,](#page-14-0) we derive an adjusted algorithm to determine all supported nondominated vectors. In addition, a method is presented to use objective-space methods like the *ε*-scalarization to determine all supported nondominated vector and a more compact formulation for the ILP used in the ε -scalarization is presented. Numerical results by these algorithms on different instances are reported in Section [5.](#page-24-0) Section [6](#page-25-0) summarizes the paper findings and suggests potential avenues for future research.

2 An Introduction to Output-sensitive Complexity

This section formally introduces the theory of *output-sensitive* complexity of enumeration problems and summarizes the existing results from the literature on the existence of output-polynomial time algorithms for the MOIMCF problem w.r.t. different solution concepts, for an comprehensive introduction see [Johnson et al.](#page-28-12) [\(1988\)](#page-28-12).

Definition 2.1 [\(Bökler and Mutzel 2015](#page-28-1)) *An enumeration problem is a pair* (*I, C*) *such that*

- *1.* $I \subseteq \Sigma^*$ *is the set of* instances *for some fixed alphabet* Σ *,*
- 2. $C: I \to 2^{\Sigma^*}$ maps each instance $x \in I$ to its configurations $C(x)$, and
- *3. the encoding length* |*s*| *for s in* $C(x)$ *for x in I is in* poly(|*x*|),

where Σ^* can be interpreted as the set of all finite strings over $\{0,1\}$.

We assume that *I* is decidable in polynomial time and that *C* is computable.

Definition 2.2 [\(Bökler and Mutzel 2015](#page-28-1)) *An* enumeration algorithm *for an enumeration problem* $E = (I, C)$ *is a* random access machine *that*

- *1. on input x in I outputs c in C*(*x*) *exactly once, and*
- *2. on every input terminates after a finite number of steps.*

Definition 2.3 [\(Bökler and Mutzel 2015\)](#page-28-1) *An enumeration algorithm for an enumeration problem* $E = (I, C)$ *is said to run in* output-polynomial time (is output-sensitive) *if its running time is in* poly $(|x|, |C(x)|)$ *for x in I.*

A finished decision problem E^{Fin} for an enumeration problem $E = (I, C)$ is defined as the problem: Given an instance $x \in I$ of the enumeration problem and a subset $M \subseteq C(x)$ of the configuration set, the goal is to decide if $M = C(x)$, i.e., we want to decide if we already have found all configurations. If the enumeration problem *E* can be solved in output-polynomial time then $E^{\text{Fin}} \in \mathbf{P}$ [\(Lawler et al.](#page-29-7), [1980](#page-29-7)).

Thus, MOIMCF as any discrete multi-objective optimization problem can be considered as an enumeration problem, i.e., enumerating all nondominated vectors or w.r.t. different solution concepts subsets of the nondominated vectors or solutions. Let *C* be the configuration set of all supported efficient solutions and *C* ∗ the configuration set of all nondominated vectors. An output-polynomial time algorithm $E = (I, C)$ which determines all supported efficient solutions exactly once, would also yield all nondominated vectors. However, it might not be output polynomial w.r.t. this task, as there might be an exponential number of efficient solutions mapping to a small number of nondominated points.

Thus, the algorithm for $E^* = (I, C^*)$ may output the elements (i.e., the nondominated vectors) more than once or even exponentially many times. Consider, for example, a directed graph with {1*, . . . , n*} transshipment nodes, a node *s* and *t* with flow balance *n* and $-n$, respectively. The graph contains the arcs (s, i) and (i, t) for all $i \in \{1, \ldots, n\}$ with upper capacity *n*. The cost of all arcs is equal to $(1,1)^\top \in \mathbb{R}^2$. Then we have $\binom{2n-1}{n}$ $\binom{n-1}{n}$ supported efficient solutions but all map to the same extreme nondominated point.

Table [2.1](#page-6-0) summarizes the existing results from literature on the existence of outputpolynomial time algorithms for the MOIMCF problem w.r.t. different solution concepts. In the table, a check-mark indicates existence, a cross indicates that the existence of such an algorithm can be ruled out, and the question mark indicates that this problem remains an open question.

Table 2.1: Existing results on the existence of output-polynomial time algorithms for the MOIMCF problem w.r.t. different solution concepts.

	extreme supported	supported	weakly supported	all
nondominated vectors			\mathbf{v}^2	\mathbf{v}^3
efficient solutions				

¹ [Bökler and Mutzel](#page-28-1) (2015) ; [Ehrgott et al.](#page-28-13) (2012)

² [Könen and Siglmayr](#page-28-9) [\(2023](#page-28-9))

³ [Bökler et al.](#page-28-10) [\(2017](#page-28-10))

Note that the negative results for all nondominated vectors also hold for the biobjective case and the weakly supported nondominated vectors for the three-objective case. In the bi-objective case, the set of all weakly supported nondominated vectors equals the set of all supported nondominated vectors [\(Könen and Siglmayr](#page-28-9), [2023](#page-28-9)). While

an output polynomial time algorithm exists to determine all supported efficient solutions, it remains an open question for the set of all supported nominated vectors.

3 Output-Sensitive Complexity for MOIMCF Problems

This chapter provides new insights into the time complexity for enumerating all supported nondominated vectors. We show that no output-polynomial time algorithm for enumerating all supported nondominated vectors that find the vectors in an ordered way in the outcome space exists unless $NP = P$. In order to prove this result, we first need the concept of *optimal tree solution* and *induced cycles*.

3.1 Preliminaries

Definition 3.1 *We call f and an associated* tree structure (T, L, U) *a* tree solution *if it consists of a spanning tree T of D and a disjoint edge set* $A \setminus T = L \cup U$ *with* $U \cap L = \emptyset$ *such that the flow f satisfies*

$$
f_{ij} = l_{ij} \quad \text{for all } (i,j) \in L,
$$

$$
f_{ij} = u_{ij} \quad \text{for all } (i,j) \in U.
$$

If a feasible (optimal) flow exists, then there also exists a feasible (optimal) tree solution [\(Cook et al.](#page-28-14), [1998\)](#page-28-14), respectively. The network simplex algorithm [\(Dantzig](#page-28-15), [1951](#page-28-15)) always determines an optimal tree solution [\(Ahuja et al.,](#page-28-0) [1993](#page-28-0)). Note that the network simplex algorithm does not run in polynomial time. However, using the *enhanced capacity scaling algorithm* [\(Orlin,](#page-29-8) [1993](#page-29-8)) solves the problem in time $O((m \log n)(m + n \log n))$, i.e., in strongly polynomial time would yield an optimal solution that can be transformed in a tree solution in polynomial time. W.l.o.g. we can assume that $l_{ij} = 0$ for all $(i, j) \in A$ (see [Ahuja et al. 1993](#page-28-0)).

We associate a real number y_i with each node $i \in V$. We refer to y_i as the node potential of node *i*. For a given set of node potentials *y* we define the *reduced cost* \overline{c}_{ij} of a given arc (i, j) as $\overline{c}_{ij} := c_{ij} + y_i - y_j$. Any node potential *y* that satisfies the well-known *complementary slackness optimality conditions* (see, e.g. [Ahuja et al. 1993](#page-28-0)) is *optimal*, and we have the following property of the reduced costs in the *residual graph D^f* .

Property 3.2 *Let f be an optimal flow and y an optimal node potential. Then* $\overline{c}_{ij}(f) \geq 0$ *for all* $(i, j) \in D_f$.

Hereby, $D_f = (V, A_f) \equiv A^+ \cup A^-$ is defined as the *residual graph* with respect to a feasible flow *f*, where $A^+ := \{(i, j) : (i, j) \in A, f_{ij} < u_{ij}\}\$ and $A^- := \{(j, i) : (i, j) \in A, f_{ij} < u_{ij}\}\$ *A,* $f_{ij} > l_{ij}$. In D_f , the *residual capacities* and *residual costs* are defined by $u_{ij}(f)$:= $u_{ij} - f_{ij} > 0$ and $c_{ij}(f) := c_{ij}$, respectively, if $(i, j) \in A^+$ and $u_{ij}(f) := f_{ji} - l_{ji} > 0$ and $c_{ij}(f) \coloneqq -c_{ji}$, respectively, if $(i, j) \in A^-$.

For an optimal flow f , we can determine an optimal node potential y by computing shortest path distances in D_f from an arbitrary but fixed root node r to all other nodes $i \in V$. Notice that we assume that all nodes *i* are reachable from *r* in D_f . If this is not

the case, artificial arcs with sufficiently high costs are added to D_f . In the following, *y* is considered as an optimal node potential (see [Ahuja et al. 1993](#page-28-0)).

Property 3.3 [\(Ahuja et al. 1993](#page-28-0)) *A tree solution f with an associated tree structure* (*T, L, U*) *is optimal if*

$$
(i) \quad \overline{c}_{ij} = 0 \quad \text{for all } (i, j) \in T,
$$
\n
$$
(ii) \quad \overline{c}_{ij} \ge 0 \quad \text{for all } (i, j) \in L,
$$
\n
$$
(iii) \quad \overline{c}_{ij} \le 0 \quad \text{for all } (i, j) \in U.
$$

These three conditions are equivalent to $\overline{c}_{ij}(f) \geq 0$ for all $(i, j) \in D_f$.

Definition 3.4 [\(Cook et al. 1998](#page-28-14)) *Let* $(i, j) \notin T$ *be a non-tree arc.*

- *(i) There exists a* unique *cycle* $C(T, L, U, (i, j))$ induced by (i, j) *, that is formed by* (i, j) *together with the path* P_{ji}^T *from j to i in T. This cycle is referred to as* C_{ij} *.*
- *(ii)* The arc (i, j) defines the orientation of the cycle C_{ij} . If $(i, j) \in L$, then the *orientation of the cycle is in the same direction as* (i, j) *. If otherwise* $(i, j) \in U$ *, then the cycle's orientation is in the opposite direction. We define the set of all* $arcs(u, v)$ *in* C_{ij} *that is in the same direction as the orientation of the cycle with* C_{ij}^+ and the set of all arcs (u, v) that are opposite directed with C_{ij}^- .

See Figure [3.1](#page-8-0) for an illustration.

Figure 3.1: An example of a BOIMCF problem, that contains 10 nondomianted vectors and 83 nonefficient flows. Right: an optimal tree solution for $c¹$ and a example of a unique induced cycle $C_{4,5}$ induced by the arc $(4,5)$ (non-tree arcs are dashed). Here arc $(4,5) \in U$ and therefore $C_{4,5}^+ = \{(4,5), (3,4)\}$ and $C_{4,5}^- = \{(3,5)\}$. The cost $c(C_{4,5}) = (7, -9)^\top$.

Property 3.5 [\(Cook et al. 1998\)](#page-28-14) *Given a tree structure* (T, L, U) *, the unique cycle* C_{ij} *induced by an arc* $(i, j) \notin T$ *satisfies*

• For all arcs $(u, v) \in C_{ij}$ it holds that $(u, v) \in T \cup (i, j)$.

• The cost of C_{ij} is given by $c(C_{ij}) = \overline{c}_{ij}$ if $(i, j) \in L$ and $c(C_{ij}) = -\overline{c}_{i,j}$ if $(i, j) \in U$.

Theorem 3.6 [\(Könen et al. 2022\)](#page-28-16) *Each undirected cycle C^D in D can be represented by a composition of the unique induced cycles*

$$
C_{ij} for all (i,j) \in C_D \setminus T.
$$

Any cycle $C \in D_f$ yields an undirected cycle C_D , which can be represented by a composition of the unique cycles C_{ij} for all $(i, j) \in C_D \setminus T$. We can extend this result by showing that the incidence vector and the cost are closed under this composition.

Lemma 3.7 [\(Könen et al. 2022](#page-28-16)) *For any cycle* $C \in D_f$ *let* C_D *be the undirected equivalent cycle in D. Then, it holds that:*

$$
\chi(C) = \sum_{a \in C_D \backslash T} \chi(C_a) \qquad and \qquad c(C, f) = \sum_{a \in C_D \backslash T} c(C_a),
$$

 $where \ \chi(C_a) \in \{-1, 0, 1\}^A \ \text{is defined by}$

$$
\chi_{ij}(C_a) := \begin{cases} 1, & \text{if } (i,j) \in C_a^+, \\ -1, & \text{if } (i,j) \in C_a^-, \\ 0, & \text{otherwise} \end{cases} \qquad \text{for all } (i,j) \in A.
$$

Theorem 3.8 [\(Könen et al. 2022](#page-28-16)) *Let f be an optimal tree solution with an associated tree structure* (T, L, U) *and let* f^* *be another integer flow. Then the flow* f^* *can be written as*

$$
f^* = f + \sum_{a \notin T} \lambda_a \chi(C_a)
$$

for some $\lambda \in \mathbb{Z}$ *and it holds that*

$$
c(f^*) = c(f) + \sum_{a \notin T} \lambda_a c(C_a)
$$

3.2 Output-Sensitive Complexity for BOIMCF Problems

In order to investigate the time complexity of enumerating all supported nondominated vectors for BOIMCF problems, we define first new problems in single-objective MCF problems.

Definition 3.9 *Given a single-objective MCF problem and an integer* $k \in \mathbb{Z}$ *. Then the* exact flow problem (EF) *asks whether there exists a flow f with cost* $c(f) = k$ *.*

We will prove that this decision problem is **NP**-complete by reducing it to the wellknown **NP**-complete *subset-sum* problem.

Definition 3.10 *Given a set* $N = \{1, \ldots, n\}$ *of n items with positive integer weights* w_1, \ldots, w_n *and a real value T, the* subset sum problem (SSP) *is to find a subset of N such that the corresponding total weight is exactly equal T. The formal definition is given by*

$$
\sum_{j=1}^{n} w_j x_j = T
$$

$$
x_j \in \{0, 1\} \quad \forall j \in \{1, \dots, n\}
$$

Theorem 3.11 *The exact flow problem is* **NP***-complete.*

Proof. Take an instance of the subset sum problem. Create the following instance of a exact flow problem. Create a node *i*' and *i*["] for each $i \in N$ and arcs $(i', i'') \forall i \in \{1, ..., n\}$ and $(i'',(i+1)')$ $\forall i \in \{1,\ldots,n-1\}$ all with cost equal to zero and lower and upper capacity equals to zero and one, respectively. We add arcs $(i', (i + 1)') \ \forall i \in \{1, \ldots, n-1\}$ with cost $c_{i',(i+1)'} = w_i$, lower capacity $l_{i',(i+1)'} = 0$ and upper capacity $u_{i',(i+1)'} = 1$. We define the node $1' = s$ and create an additional node t as well as the arcs (n'', t) and (n', t) with cost equal to zero and w_n , respectively. We set $b_s = -1, b_t = 1$ and $b_i = 0$ for all other nodes. A construction of this instance is illustrated in Figure [3.2.](#page-10-0)

If we can decide whether a flow $f \in \mathcal{X}$ with $c(f) = T$ exists in polynomial time, we would also solve be able to solve the subset sum problem in polynomial time. Assume the subset-sum problem is solvable, i.e., $\sum_{j=1}^{n} w_j x_j = T$. For each $x_i = 1$ we take the arc $(i', (i + 1)')$ with cost $c_{i', (i+1)'} = w_i$. For each $x_i = 0$ we take the arcs (i', i'') and $(i'',(i+1)')$ both with cost equal to zero. This is a feasible *s*-*t* path (feasible flow) with cost equal to T . On the other side, considering a path P (flow) with cost equal to T we set $x_i = 1$ if $(i', (i + 1)') \in P$ and $x_i = 0$ if not. Then *x* solves the subset problem with the same argumentation as above.

Figure 3.2: The instance of the exact flow problem corresponding to a given subset sum instance.

Theorem [3.11](#page-10-1) implies that it is also **NP**-complete to determine if a flow *f* exists with $k_1 < c(f) < k_2$ with $k_1, k_2 \in \mathbb{Z}$.

In addition, we can prove the following statement.

Theorem 3.12 *The problems to determine the k-th best flow or the k-th smallest distinct cost of a flow are* **NP***-hard.*

 \Box

Proof. Let the cost of finding the *k*-th best or *k*-th smallest distinct cost of a flow be *T*(*n*). Consider the instance of the exact flow problem for a given subset sum problem as in the proof above. Deciding whether a flow f exists with $c(f) = T$ would solve the subset sum problem. There are at most 2^n different flows (item *i* could be selected or not). Thus, a binary search for a given flow value is $\mathcal{O}(n)$. It can be concluded that the complexity of the subset sum problem, known to be \mathbf{NP} -complete, is $\mathcal{O}(nT(n))$. Therefore, the problem of determining the *k*-th best or *k*-th smallest distinct cost of a flow is **NP**-hard. \Box

Next, we will prove that we can determine all distinct costs of the flows for a singleobjective minimum cost flow problem in output-polynomial time, if we can determine all supported nondominated vectors for a bi-objective minimum cost flow problem in output-polynomial time and vice versa.

Definition 3.13 *Given a single-objective integer minimum cost flow problem, the* all distinct cost value flow problem (ACVF) *determines a minimal set of flows that includes all existing different cost values of all feasible flows. In other words, it identifies k different flows* f_1, \ldots, f_k *such that*

$$
c(f_1) < \ldots < c(f_k)
$$

and there exists no flow f_p *such that* $c(f_p) \notin \{c(f_1), \ldots, c(f_k)\}.$

Theorem 3.14 *If we can solve the ACVF problem in output-polynomial time, then we can determine all supported nondominated vectors of a BOIMCF problem in outputpolynomial time.*

Proof. Each supported nondominated vector lies on a maximal nondominated facet of the upper image since its preimage is an optimal solution to a weighted sum scalarization P_{λ} for an $\lambda \in \Lambda_d$ (i.e., weights are strictly positive). Note that in the bi-objective case, every maximally nondominated face F_V of conv(V) is a line segment connecting two adjacent extreme supported points if there is more than one nondominated point $(|\mathcal{Y}_N| > 1)$. A maximally nondominated face can only have dimension zero if there is only one extreme nondominated point, which implies that there is only one nondominated point (or in other words, the ideal point is feasible). In the following, we assume that $(|y_N| > 1)$. All N extreme points and precisely one corresponding extreme flow can be determined by using the enhanced parametric programming approach in $\mathcal{O}(M + Nn(m + 1))$ *n* log *n*)) time [\(Raith and Sedeño-Noda](#page-29-4), [2017](#page-29-4)), where *M* denotes the time required to solve a given single-objective minimum cost flow problem. Also, the algorithm stores one extreme flow for each extreme nondominated point.

In [Könen and Siglmayr](#page-28-9) [\(2023](#page-28-9)), it is shown that for each maximal nondominated facet, we can create a reduced single-objective flow problem in which each image of a feasible flow lies on the maximal nondominated face. If we assume that we can solve the (ACVF) problem in output-polynomial time, we could determine for a maximal nondominated face all supported nondominated vectors whose images lie on this facet in

output-polynomial time. Thus, we solve the ACVF problem on the reduced network with only the first objective function. Solving the ACVF problem successively for the reduced single-objective minimum cost flow problems for each *N* − 1 maximal nondominated facets would yield all supported nondominated vectors for the given BOIMCF problem. Note that some vectors may lie in more than one maximal nondominated facet, namely the extreme nondominated vectors, and therefore we have to ensure that we only store these flows once. In total, we would obtain all supported nondominated vectors in output-polynomial time. \Box

Theorem 3.15 *If we can determine all supported nondominated vectors for a bi-objective integer minimum cost flow problem in output-polynomial time, then we can solve the ACVF problem in output-polynomial time.*

Proof. Consider an instance of the ACVF problem. Let f_1 be an optimal solution of the corresponding single-objective MCF problem, with the optimal tree structure (*T, L, U*). Due to Theorem [3.8](#page-9-0) it holds that any flow \bar{f} can be written as $\bar{f} = f_1 + \sum_{a \notin T} \lambda_a \chi(C_a)$ for some $\lambda \in \mathbb{Z}$ and thus $c(\bar{f}) = c(f_1) + \sum_{a \notin T} \lambda_a c(C_a)$. We also can obtain the highest cost of a feasible flow f_k in polynomial time. Therefore, we create an MCF instance with objective function $-c(f)$ as objective and obtain the best flow in polynomial time.

We construct an BOIMCF instance by creating an artificial node *s* which has an arc (s, r) to an arbitrary but fixed node r with a lower and upper capacity equal to 1 and with costs equal to $(0, c^1(f_k))^T$. The demand of node *s* is $b_s = -1$, and we adjust the demand value of the node *r* by $b_r = b_r + 1$. For this new artificial problem, the existing cost values regarding the first objective are equal to the original problem, i.e., the set of the ACVF problem regarding the first-objective would be the same in both problems. For the second objective, we define the cost for each arc as the negative reduced cost of the first objective, i.e., $c^2(a) = -\bar{c}^1(a)$ for each $a \in A$. Then for any induced cycle C_a it holds $c^1(C_a) = -c^2(C_a)$. One extreme supported vector ist the vector $(c^1(f_1), c^2(f_1))$.

Any flow f_p can be written as $f_p = f_1 + \sum_{a \notin T} \lambda_a \chi(C_a)$ for some $\lambda \in \mathbb{Z}$ and it holds that

$$
c(f_p) = c(f_1) + \sum_{a \notin T} \lambda_a c(C_a)
$$

=
$$
(c^1(f_1) + \sum_{a \notin T} \lambda_a c^1(C_a), c^2(f_1) + \sum_{a \notin T} \lambda_a c^2(C_a))^\top
$$

=
$$
(c^1(f_1) + \sum_{a \notin T} \lambda_a c^1(C_a), c^2(f_1) - \sum_{a \notin T} \lambda_a c^1(C_a))^\top.
$$

Since $c^{1}(f_{p}) + c^{2}(f_{p}) = \text{const}$, all feasible flows f_{p} lie on the same maximally nondominated edge which is spanned by the two extreme supported nondominated vectors $(c^{1}(f_{1}), c^{2}(f_{1}))$ and $(c^{1}(f_{k}), c^{2}(f_{k}))$. If we could determine all nondominated supported vectors in output-polynomial time, we would also solve the ACVF problem for the original single-objective MCF problem. \Box

Let ACFV^{Fin} denote the decision problem of ACFV, i.e., given a set of distinct cost values $\{c(f_1), \ldots, c(f_k)\}\$ of some feasible flows, decide if there exists a feasible flow f_p with a distinct cost value to the given ones $c(f_p) \notin \{c(f_1), \ldots, c(f_k)\}\$. If we can show that $ACFV^{Fin}$ is not solvable in polynomial time, then it does not exits an output-polynomial time algorithm to determine all supported nondominated vector unless $\mathbf{NP} = \mathbf{P}$.

Using these results, we can show that if we have given two distinct supported nondominated vector y^1, y^2 with $y_1^1 < y_2^2$ for a bi-objective integer minimum cost flow (BOIMCF) problem, it is NP -hard to decide if another supported nondominated vector y^3 exists, which has higher cost in the first objective for *y* ¹ and lower cost in the first objective for y^2 .

Theorem 3.16 *Given two different supported nondominated vectors y* ¹ *and y* ² *of a BOIMCF problem with* $y_1^1 < y_1^2$, the problem of deciding whether another supported *nondominated vector* y^3 *with* $y_1^1 < y_1^3 < y_1^2$ *exists, i.e., in between the supported nondominated vector* y^1 *and* y^2 *is* **NP***-complete.*

Proof. We show the **NP**-completeness of the problem by reduction from the subset sum problem. Consider an instance of the subset sum problem. Create the singleobjective minimum cost flow problem as in the proof of Theorem [3.11,](#page-10-1) the construction is shown in Figure [3.2.](#page-10-0) An optimal flow of this instance would have a cost equal to 0 and the highest cost of a feasible flow would be $\sum_{i \in N} w_i$. Transform the single-objective MCF problem to a BOIMCF problem as in the proof of Theorem [3.15,](#page-12-0) such that all nondominated vectors are supported nondominated vectors and lie on the same edge between the two extreme supported points $(0, \sum_{i \in N} w_i)^{\top}$ and $(\sum_{i \in N} w_i, 0)^{\top}$. From the artificial node *s*, create two arcs (s, t') and (s', t') with cost (k_1, ℓ_1) and (k_2, ℓ_2) , where ℓ_1, ℓ_2 is chosen such that both points lie on the maximally nondominated face, which can be determined in polynomial time. So we have the two supported nondominated vectors $y^1 = (T - 1, \ell_1)$ and $y^2 = (T + 1, \ell_2)$. Any supported nondominated vector y^3 with the property $y_1^1 < y_1^2 < y_1^2$ would have a cost equal to *T*. If we can decide whether there exists y^3 such that $y^1_a < y^3_1 < y^2_1$ in polynomial time, then we would solve the subset sum problem in polynomial time, which yields a contradiction. \Box

Theorem [3.16](#page-13-0) implies also that it is **NP**-hard to determine the next best supported nondominated vector regarding the first or second objective for a given supported nondominated vector of a BOIMCF problem. This means it is not possible to determine all supported nondominated vectors in an ordered way regarding the first-objective (secondobjective) in output-polynomial time. However, this theorem is not enough to show that the supported nondominated vector could not be obtained in output-polynomial time and it still remains an open question.

Consider the BIOMCF problem of a subset sum instance in the proof of Theorem [3.16.](#page-13-0) We have shown that given a supported nondominated vector, it is **NP**-hard to determine the next best supported nondominated vector for this instance. However, for this instance of a BOIMCF problem, we can determine all supported-nondominated vectors

in output-polynomial time using a similar algorithm to the pseudo-polynomial labeling algorithm to solve the subset sum problem presented in [Pisinger](#page-29-9) [\(1999](#page-29-9)).

4 Determining the Supported Nondominated Vectors

In this section, we shortly describe the Algorithm [1](#page-17-0) presented in [Könen and Siglmayr](#page-28-9) [\(2023](#page-28-9)) to determine all supported efficient solutions and derive an adjusted algorithm in order to determine the supported nondominated vectors more efficiently if the number of branches needed is significantly smaller than the number of supported efficient solutions.

4.1 Determining All Supported Efficient Solutions

Algorithm [1](#page-17-0) consists of two phases and relies on the following widely-known fact for any integer supported flow, see, e.g., [Gal](#page-28-17) [\(1977\)](#page-28-17).

Theorem 4.1 *A flow f is contained in* F_X , *i.e., its image of* $c(f)$ *lies on a maximally nondominated face* F_Y *w.r.t. an associated weight vector* $\lambda \in \Lambda_d$, if *f is an optimal solution to the parametric network flow program* P_λ .

Any image of a supported flow must lie in at least one maximally nondominated face, and any integer point in a maximally nondominated face corresponds to a supported integer flow. Assuming that for each maximally nondominated face $F_i \in \{F_1, \ldots, F_t\}$, one optimal solution f^i and the corresponding weighting vectors λ^i are given the problem of determining all supported flows reduces to determining all optimal flows for each parametric single-objective problem (P_{λ^i}) . These optimal solutions can be determined by the algorithm for determining all optimum flows for single-objective minimum cost flow problems presented in [Könen et al.](#page-28-16) [\(2022](#page-28-16)), which we refer to as the all optimum flow (AOF) algorithm. The AOF algorithm successively searches for so-called *proper zero-cost* cycles efficiently using a modified depth-first search technique.

Definition 4.2 [\(Hamacher 1995](#page-28-18)) *A* proper (i, j) -cycle $C \in D_f$ *is a proper cycle that contains the arc* (i, j) *. In addition, a proper* (i, j) *-cycle* $C \in D_f$ *is called* minimal *if it has minimal cost among all proper (i, j*)*-cycles.*

In order to determine all optimal solutions for a given maximal nondominated face, i.e., an edge between two consecutive extreme vectors y^i and y^{i+1} (see Figure [4.1\)](#page-15-0), the approach uses the AOF algorithm on the network with weight vector λ corresponding to the given edge F_i . The extreme vector y_i is optimal for this network with cost function λ ^T*Cf*. In the first step, the network D_{λ} gets reduced to D' λ in the following way. We remove all arcs in $\mathcal{X} := \{(i, j) \in A : \overline{c}_{ij} \neq 0\}$, i.e., arcs with non-zero reduced cost. Note that some of these arcs in $\mathcal X$ may carry flow. Consequently, we must adjust the demand vector accordingly to preserve feasibility. Consider the reduced network $D'_{\lambda} = (V, A')$ with $A' = A \setminus \mathcal{X}$ and the new flow balance values $b'_i = b_i - \sum_{(i,j) \in \mathcal{X}} f_{ij} + \sum_{(j,i) \in \mathcal{X}} f_{ji}$. For any feasible flow f in D we can easily determine the corresponding feasible flow f' in D' where all flow values are copied for all arcs $a \in A'$.

Figure 4.1: Illustration of two consecutive extreme points, the maximally nondominated edge *Fⁱ* in blue and the cost vector of $(\lambda^i)^\top C$.

We denote by \mathcal{P}'_{λ} the set of all feasible integer flows of the reduced network D'_{λ} . The fact that there is only a finite number of flows in \mathcal{P}'_{λ} follows from the assumption that there is no arc (i, j) with $u_{ij} = \infty$.

For a feasible solution $f' \in \mathcal{P}'_{\lambda}$, we define

$$
f^*(a) = \begin{cases} f'(a), & \text{if } a \in A', \\ f(a), & \text{if } a \in \mathcal{X}. \end{cases}
$$

The next lemma demonstrates how every optimal flow in D_{λ} can be obtained from combining an initial optimal flow and some feasible flow in D'_{λ} .

Lemma 4.3 [\(Könen et al. 2022](#page-28-16)) *Let f be an optimal integer solution of the MCF problem in* D_{λ} *. If* f' *is a feasible integer solution in* D'_{λ} *and* f^* *is defined as above, then* f^* *is an optimal solution in* D_{λ} *and vice versa.*

Given a feasible flow f in D'_{λ} , we can compute another feasible flow by finding a proper cycle in the residual network of D'_{λ} . Therefore, we can determine an another optimal flow in D_{λ} by finding a proper cycle in $D'_{\lambda,f}$ instead of finding a proper 0-cycle in $D_{\lambda,f}$. This gives us the following theorem:

Theorem 4.4 [\(Könen et al. 2022](#page-28-16)) *Let f be an optimal integer flow. Then every proper cycle in* $D'_{\lambda,f}$ *induces another optimal integer flow in* D_{λ} *.*

Following this, the feasible objective space is reduced to all feasible solutions whose image lies on the edge F_i . In [Könen et al.](#page-28-16) [\(2022](#page-28-16)), it is shown that such a minimal proper cycle *C* in $D'_{\lambda,f}$ can be determined in $\mathcal{O}(mn)$, yielding a new efficient solution

whose image is on the maximal nondomianted face F_i . Since we can compute the next supported efficient solution in D'_{λ} , we can now apply the binary partition approach as used in [Hamacher](#page-28-18) [\(1995](#page-28-18)) or [Könen et al.](#page-28-16) [\(2022](#page-28-16)). Let $\mathcal{X}_{D'}$ be the set of optimal flows in D'_{λ} , which we also call the *optimal solution space*, and let f' , $f'' \in \mathcal{X}_{D'}$ as above. We then can recursively dividing $\mathcal{X}_{D'}$ into $\mathcal{X}_{D'_1}$ and $\mathcal{X}_{D'_2}$ such that f' and f'' are optimal flows in \mathcal{X}_{D_1} and \mathcal{X}_{D_2} , respectively, and then seeks more flows in $\mathcal{X}_{D'_1}$ and $\mathcal{X}_{D'_2}$. This can easily be done by identifying some arc *a* with $f'(a) \neq f''(a)$. Such an arc must exist since $c(f') \neq c(f'')$. Let $(i, j) = a$ if $a \in L$ and $(i, j) = a^{-1}$ if $a \in U$. Assume w.l.o.g. that $f'_{ij} < f''_{ij}$. We set

$$
\mathcal{X}_{D'_1} \coloneqq \{z \in \mathcal{X}_{D'} : z_{ij} \leq f'_{ij}\}
$$

and

$$
\mathcal{X}_{D_2'}\coloneqq \{z\in \mathcal{X}_{D'}: z_{ij}\geq f_{ij}'+1\}.
$$

This partitioning implies that each new minimum cost flow problem is defined in a modified flow network with an altered upper or lower capacity of a single arc (i, j) . By dividing the new solution spaces again until no more other optimal flow exists, all optimal solutions are found, and therefore, all supported efficient solutions that image lie on the maximal nondominated face *Fⁱ* . We obtain the complete set of supported efficient solutions by doing so for each maximally nondominated face and only storing newly seen flows. For each pair of consecutive extreme points y^i and y^{i+1} , we determine the weighting vector $\lambda^i \in \Lambda$ that corresponds to the normal of maximally nondominted facet F_i connecting the extreme points y^i and y^{i+1} :

$$
\lambda^i := \frac{1}{\sqrt{\lambda_1^2 + \lambda_2^2}} (\lambda_1, \lambda_2)^\top \quad \text{where} \quad \lambda_1^i := c^2(f^i) - c^2(f^{i+1})
$$

$$
\lambda_2^i := c^1(f^{i+1}) - c^1(f^i).
$$

Then f^i and f^{i+1} are both optimal flows for the single-objective weighted-sum (MCF) program $(P_{\lambda i})$ [\(Eusébio and Figueira](#page-28-5), [2009a](#page-28-5)). Hence, determining all optimal solutions for (P_{λ^i}) gives all supported efficient flows whose image lies in between F_i .

4.2 The Adjusted Algorithm

In the current version of the algorithm, we search for a proper cycle in the reduced network to obtain a second supported efficient solution. Note that in the reduced network, any cycle would have a cost equal to zero regarding the weighted sum with a weight vector λ corresponding to the current maximal nondominated face F_i . However, we will now consider the first objective. We then search for a minimal proper cycle in the reduced network, i.e., a proper cycle with minimum cost regarding the first objective under all proper cycles in D'_{λ} . A proper zero cost cycle regarding the first objective in D'_{λ} would yield a supported efficient solution whose images maps to the same vector as the initial solution (in the bi-objective case). Since we want, in the best case, only one flow per vector, we search for a minimal proper cost cycle under all proper cycles that have cost greater than zero. We then would obtain a distinct cost second-best flow regarding the

Algorithm 1: FindAllSupportedEfficientFlowsBiObjective

Data: (D, l, u, b, c^1, c^2) **Result:** The set of all supported flows \mathcal{X}_S $\mathcal{X}_S \leftarrow \varnothing$; // Determine all extreme supported points *y ⁱ* and for each one corresponding extreme flow f^i , sorted non-decreasingly in $c^1(f)$ $\{(y^i, f^i): i \in \{1, \ldots, N\}\} \leftarrow \text{EnhancedParametricNetworkAlgorithm(D)};$ **for** $i = 1, ..., N - 1$ **do** $\lambda_1^i \leftarrow c^2(f^i) - c^2(f^{i+1})$; $\lambda_2^i \leftarrow c^1(f^{i+1}) - c^1(f^i)$; $\lambda^i \leftarrow \frac{1}{\sqrt{\lambda^2_i}}$ $\frac{1}{\lambda_1^2 + \lambda_2^2}$ (λ_1, λ_2) ^T; $\mathcal{X}_S \leftarrow \mathcal{X}_S \cup \text{FindAllOptimalFlows}(P_{\lambda^i}, f^i)$; // Return only flows with $c^1(f) \neq c^1(f^i)$ to avoid repetitions.

first objective. Using the complementary slackness condition, we prove that we could obtain such a cycle in $\mathcal{O}(n^3)$. Using this fact, we could adjust the previously presented algorithm in Section [4.1](#page-14-1) to find the supported nondominated vectors more efficiently.

We want to find all supported nondominated vectors for each maximally face F_i (in this case, given by an edge between two consecutive extreme points). For that, we want to determine all supported nondominated vectors between the edge of the two extreme vectors y^i and y^{i+1} . In Figure [4.1,](#page-15-0) an example is shown.

For y_i let f' be a corresponding optimal flow. Considering the edge F_i , we know that f' is not only an optimal flow for the weight vector λ , but as well f' is an optimal flow regarding the first objective c^1 in D'_{λ} . So for f' , there can not be a negative cycle regarding the cost of c_1 in $D'_{\lambda, f'}$ and the complementary slackness conditions holds.

In order to find the next nondominated vector on the edge F_i , we determine the minimal proper cycle $C \coloneqq \operatorname{argmin} \{c^1(C) : c^1(C) > 0, C \in D'_{\lambda, f'}\}$. Using techniques from [Könen et al.](#page-28-16) [\(2022](#page-28-16)) and [Hamacher](#page-28-18) [\(1995](#page-28-18)) this can be done in $\mathcal{O}(n^3)$. Note, that this is feasible only for the first and last supported nondominated vector on a maximally nondominated face, since it requires an optimal solution w.r.t. the first or second objective function to start with. Thus, this procedure can not be extended to iteratively generate supported nondominated vectors along the face in an ordered way.

Let $\overline{D} = (\overline{d}_{ji})$ be the distance table of $D_{\lambda,f}$ concerning $\overline{c}^1(f)$, i.e., \overline{d}_{ji} is the length of the shortest path P_{ji} from *j* to *i* in D_f with length $\overline{c}^1(f, P_{ji})$. The distance table may be computed in time $O(n^3)$ using the Floyd-Warshall algorithm or in time $O(n^2 \log n + mn)$ by (essentially) repeated calls to Dijkstra's algorithm; the latter is more efficient on sparse graphs.

Hamacher shows the following property for proper minimal (i, j) -cycles:

Property 4.5 [\(Hamacher 1995\)](#page-28-18) *For any anti-parallel arc in* A_f , *i.e.*, $(i, j) \in A_f$ *with* $(j, i) \notin A_f$, the cost of a proper minimal (i, j) -cycle $C \in D_f$ is given by $c(f, C) =$ $\overline{c}_{ij}(f) + \overline{d}_{ji}$.

Algorithm 2: FindAllOptimalFlows

Data: optimal flow *f* for $P_{\lambda i}$ and the reduced costs \overline{c} **Result:** All optimal integer flows for P_{λ} *i* $f^* \leftarrow \text{FindAnotherOptimalFlow}(f, \overline{c}, D);$ **if** $f^* = \textbf{null}$ then Stop; $\text{Print}(f^*)$; // Partition solution space and find new optimal flow $a \leftarrow \text{arc } a \text{ with } f(a) \neq f^*(a);$ **if** $f(a) < f^*(a)$ **then** FindAllOptimalFlows (f, \overline{c}, D) with $u_a = f(a)$; FindAllOptimalFlows (f^*, \overline{c}, D) with $l_a = f(a) + 1$; **else** FindAllOptimalFlows(f , \overline{c} , D) with $l_a = f(a)$; FindAllOptimalFlows (f^*, \overline{c}, D) with $u_a = f(a) - 1$;

When using Hamacher's idea, the problem is that it only applies to arcs with no anti-parallel arc in A_f . For all other arcs, the cost of the corresponding proper minimal (i, j) -cycle has to be computed by finding a shortest path in $D_f \setminus \{(j, i)\}\)$. In the following result, we show that we can use the complementary slackness optimality conditions of an optimal solution to overcome this problem and restrict ourselves to consider only arcs with no anti-parallel arcs in $D_{\lambda,f}$.

Theorem 4.6 *We can determine the minimal cost cycle*

 $C \coloneqq \mathrm{argmin}_{C \in D'_{\lambda, f'}} \{c^1(C) : c^1(C) > 0\}$

over all cycles with strictly positive weight in $\mathcal{O}(n^3)$ *.*

Proof. Let *C* be a minimal proper cycle with the property $c^1(C) \neq 0$. So $\overline{c}^1(C, f) > 0$, because there cannot be a cycle with negative costs due to the negative cycle optimality conditions. Therefore, there exists an arc $(i, j) \in C$ with $\overline{c}_{ij}(f) > 0$. Since f is an optimal solution, the complementary slackness optimality conditions ensure that $\overline{c}_{ij}^1(f) > 0$ only holds if $(i, j) \in D$ and $f_{ij} = l_{ij}$ or $(j, i) \in D$ and $f_{ji} = u_{ji}$. Therefore, we have that $(i, j) \in \overline{A} := \{(i, j) \in D_f : (i, j) \in D \text{ with } f_{ij} = l_{ij} \text{ or } (j, i) \in D \text{ with } f_{ji} = u_{ji}\}\$ and $(j, i) \notin D_f$, (since the flow value of the corresponding arc of (i, j) in *D* is equal to the upper or lower capacity of this arc).

Since $(j, i) \notin D_f$ it holds that $\overline{c}^1(C, f) = \overline{c}_{ij} + \overline{d}_{ji}$ due to Property [4.5.](#page-17-1) Since $\overline{c}_{ij}^1(f) \ge 0$ any cycle C with $c^1(C)$ has at least one of such an arc. Consequently, we can determine a minimal proper cycle $C := \operatorname{argmin} \{c^1(C) : c^1(C) > 0, C \in D'_{\lambda, f'}\}$ by just choosing $C = \operatorname{argmin} \{c(f, C) = \overline{c}_{ij} + \overline{d}_{ji} : C = \{(i, j)\} \cup P_{ji} \text{ with } (i, j) \in \overline{A}\}.$ We can compute the distance table in $\mathcal{O}(n^3)$ time. Notice that this approach also provides the paths P_{ji} in addition to the cost \overline{d}_{ii} .

Given all pairwise distances \overline{d}_{ji} and the reduced costs $\overline{c}_{ij}^1(f)$ for all (i, j) in D_f^1 we can determine $c^1(f, C) = \overline{c}_{ij}^1 + \overline{d}_{ji}$ in constant time $\mathcal{O}(1)$ and can determine the argmin in $\mathcal{O}(m)$ time. As a result, we can compute a minimal proper cycle with $c^1(C) \neq 0$ in time $\mathcal{O}(n^3)$.

So let *C* be chosen as above. Let $f'' = f' + \chi(C)$. The flow f'' is supported, and its image lies on the edge F_i . We are going to prove that f'' is the next nondominated point on F_i regarding the cost of c^1 , i.e., there does not exist a flow \bar{f} whose image lies on F_i with the property $c^{1}(f') < c^{1}(\bar{f}) < c^{1}(f'')$.

Theorem 4.7 *There exists no flow* \bar{f} *whose image* $c^1(\bar{f})$ *lies on* F_1 *between the images of* f' and f'' , i.e., $c^1(f') < c^1(\bar{f}) < c^1(f'')$.

Proof. It holds that

$$
f'' = f' + \chi(C)
$$
 and thus $c^1(f'') = c^1(f') + c^1(C)$

and we know that any flow \bar{f} on F_1 can be written as

$$
\bar{f} = f' + \sum_{C_i \in D'_{\lambda, f'}} \chi(C_i) \qquad c^1(\bar{f}) = c^1(f') + \sum_{C_i \in D'_{\lambda, f'}} c^1(C_i).
$$
\n
$$
\implies c^1(f') < c^1(f') + c^1(C) = c^1(f'') \leq c^1(f') + \sum_{C_i \in D'_{\lambda, f'}} c^1(C_i) = c(\bar{f}).
$$

The inequality $\sum_{C_i \in D'_{\lambda, f'}} c^1(C_i) \geq c^1(C)$ holds since the sum contains the positive cost of at least one cycle regarding the first objective. Since *C* was chosen as the minimal cost cycle

$$
C := \mathrm{argmin}_{C \in D'_{\lambda, f'}} \{c^1(C) : c^1(C) > 0\},\
$$

 \Box

it holds that $c^1(f'') \leq c^1(\bar{f})$.

Since we can compute the next nondominated supported vector in D'_{λ} , we can now apply the binary partition approach as used in [Hamacher](#page-28-18) [\(1995](#page-28-18)) and [Könen et al.](#page-28-16) [\(2022](#page-28-16)), presented above. Thereby, we substitute FindAnotherOptimalFlow in Algorithm [2](#page-18-0) by Algorithm [3.](#page-20-0)

This adjusted algorithm can be more efficient in determining all supported nondominated vectors than by determining all supported efficient flows, as the next example will show since we might have an exponential number of supported efficient flows that we would not consider with this new technique.

Example 4.8 *Consider a graph D with nodes* $\{1, \ldots, k\}$ *, parameters* $L, M \in \mathbb{N}$ *and an optimal flow f w.r.t. the first objective.* Node 1 *has* $b_1 = 2$ *and node k has* $b_k = -2$ *. For all other nodes* $i \in \{2, \ldots, k-1\}$ *we have* $b_i = 0$. The graph *D contains arcs*

• (*i, i* + 1) *for all i* ∈ {1*, . . . , k* − 1} *with ui,i*+1 = (*k* − 3)*M* + (*L* + 2)*, fi,i*+1 = 2 *and reduced costs* $\bar{c}^1_{i,i+1} = \bar{c}^2_{i,i+1} = 0$,

Algorithm 3: FindSecondDistinctCostBestFlow

Data: optimal flow f in P_{λ} *i*

Result: A second distinct cost-best flow f' with $c(f') \neq c'(f)$, if one exists.

 $y \leftarrow$ ComputeNodePotential(*f*, *D*); $\overline{c} \leftarrow$ ComputeReducedCost (y, D) ; $(\overline{D}, P) \leftarrow$ DetermineDistanceTableAndPaths (\overline{c}, f, D) ; $C \leftarrow \text{argmin}\{c(f, C)\} = \text{argmin}_{C \in D'_{\lambda, f'}} \{c^1(C) : c^1(C) > 0\};$ **if** $C = \text{null}$ then Stop; $f' \leftarrow f + \chi(C);$ **return** *f* ′ *;*

- • *arcs* $(k-i, k-i-2)$ *for all* $i \in \{1, \ldots k-3\}$ *with* $u_{k-i,k-i-2} = M$, $f_{k-i,k-i-2} = 0$ $and \ \bar{c}_{k-i,k-i-2}^1 = \bar{c}_{k-i,k-i-2}^2 = 0, \ and$
- *one arc* $(k, k-2)$ *with* $u_{k,k-2} = L$ *,* $f_{k,k-2} = 0$ *and* $\bar{c}^1_{k,k-2} = 1$, $\bar{c}^2_{k,k-2} = -1$ *.*

Figure [4.2](#page-20-1) *illustrates this example for* $k = 5$. There are $L + 1$ *nondominated vectors which are all supported. However, we would have* $(M + 1)^{k-3}(L + 1)$ *efficient flows, which are all supported. However, using the branching technique described above, we would have only L* + 1 *leaves at the end of our branching.*

Figure 4.2: Graph *D* of the BOIMCF problem in Example [4.8](#page-19-0) with an optimal flow *f* w.r.t. the first objective. The arcs are labeled with $(u_a, f_a, \bar{c}_{ij}^1, \bar{c}_{ij}^2)$. Here $l_{ij} = 0$ for all arcs and $(u_a =$ $(k-3)M + (L+2)$, $f_a = 2$, $\bar{c}_{ij}^1 = 0$, $\bar{c}_{ij}^2 = 0$) for all arcs that are not labeled. The nodes are labeled with b_i and $b_i = 0$ for all nodes which are not labeled. $M > L$.

However, as shown in the following example, we might find supported efficient flows (even exponential many) with images corresponding to supported nondominated vectors already found in other branches.

Example 4.9 *Consider a graph D with nodes* $\{1, \ldots, k\}$ *, parameter* $L \in \mathbb{N}$ *and an optimal flow f w.r.t. the first objective. Node* 1 *has* $b_1 = 2$ *and node k has* $b_k = -2$ *. All other nodes* $i \in \{2, \ldots, k-1\}$ *are transshipment nodes,* $b_i = 0$ *. The graph D contains arcs*

• $(i, i + 1)$ *for all* $i \in \{1, ..., k - 1\}$ *with* $u_{i,i+1} = L + 2$ *,* $f_{i,i+1} = 2$ *and reduced costs* $\bar{c}^1_{i,i+1} = \bar{c}^2_{i,i+1} = 0$, and

• (*k, k* − *i*) *for all i* ∈ {2*, . . . , k* − 1} *with uk,k*−¹ = *L, fk,k*−¹ = 0 *and reduced costs* $\bar{c}^1_{k,k-1} = 1, \bar{c}^2_{k,k-i} = -1.$

Figure [4.3](#page-21-0) illustrates the graph D for k = 5*. Then, this BOIMCF has L*+1 *nondominated vectors, all of them supported. However, we have* $\sum_{i=0}^{L} \binom{(k-2)+i-1}{i}$ *i supported efficient flows, and all of them would yield a leave of our branching technique. This number can be exponential. Assume that L > k then it holds that*

$$
\sum_{i=0}^{L} \binom{(k-2)+i-1}{i} > \sum_{i=0}^{k} \binom{k-3+i}{i} > \sum_{i=0}^{k} \binom{k-3}{i} = 2^{k-3}.
$$

Since $k \in \mathcal{O}(m)$ *, we have an exponential amount of flows to consider.*

Figure 4.3: Graph *D* of the BOIMCF problem in Example [4.9](#page-20-2) with an optimal flow *f* w.r.t. the first objective. The arcs labeled with $(u_a, f_a, \bar{c}_{ij}^1, \bar{c}_{ij}^2)$. Here $l_{ij} = 0$ for all arcs and $(u_a =$ $L + 2$, $f_a = 2$, $\bar{c}_{ij}^1 = 0$, $\bar{c}_{ij}^2 = 0$) for all arcs that are not labeled. The nodes are labeled with b_i and $b_i = 0$ for all nodes which are not labeled.

4.3 *ε***-Scalarizations on the Reduced Networks**

One decision-space method for determining nondominated vectors for general multiobjective linear programs is the well-known ε -constraint method [\(Haimes et al.,](#page-28-19) [1971](#page-28-19)). In the *ε*-constraint method, there is no aggregation of criteria as in the weighted-sum method, instead, only one of the original objectives is minimized while the others are transformed into constraints.

The *ε*-constraint scalarization of [\(MOIMCF\)](#page-1-0) can be represented as:

$$
\begin{array}{ll}\n\text{min} & c^j(f) \\
\text{s.t.} & c^k(f) \le \varepsilon_k \\
& f \in \mathcal{X}\n\end{array} \quad \forall k \in \{1, \dots, d\}, \ k \neq j
$$

An illustration of the *ε*-constraint for a bi-objective minimum cost flow problem can be found in Figure [4.4.](#page-22-0)

For the BOIMCF problem, all nondominated vectors can be found by solving a sequence of ε -constraints problems. Starting with the lexicography minimal solution $\text{lexmin}\{c^1(f), c^2(f)\}\$ regarding the first objective, which can be determined in polynomial time, we can determine the next nondominated point using the ε -constraint. After each

Figure 4.4: An example of the *ε*-constraint scalarization $\{\min_{f \in \mathcal{X}} c^1(f) : c^2(f) \leq \varepsilon\}$ for a BOIMCF problem.

newly generated point, we update ε until the last nondominated point lexmin $\{c^2(f), c^1(f)\}$ is generated. In [Eusébio and Figueira](#page-28-20) [\(2009b\)](#page-28-20), an implicit enumeration algorithm for BOIMCF problems is given, which solves such a sequence of *ε*-constraint problems by computing optimal non-integer solutions with a network simplex algorithm and then determining optimal integer solutions with a branch-and-bound technique.

We also could use this *ε*-constraint method to obtain the complete set of supported nondominated vectors for the BOIMCF problems, in which we have to solve one *ε*constraint problem for each supported nondominated vector. Thereby, we solve the *ε*-constraint method on each maximally nondominated face. Let $X_{D'}$ be the set of all feasible flows for the reduced network for one maximal nondominated face, i.e., all solutions whose image lie on this maximal nondominated face. Any solution in the reduced network is a supported efficient solution. Using the *ε*-constraint method on all of these facets would determine all supported nondominated vectors. We only must solve one *ε*-constraint problem for each supported nondominated vector. Let $T(\varepsilon)$ be the longest time required to solve such an *ε*-constraint problem.

Theorem 4.10 *For a given BOIMCF problem, we can determine all S supported nondominated vectors in time* $\mathcal{O}(N n(m + nlog n) + S(T(\varepsilon) + n + m))$

Proof. The enhanced parametric network approach [\(Raith and Ehrgott](#page-29-10), [2009b](#page-29-10)) requires $\mathcal{O}(Nn(m+n\log n)+M)$ time, where M is the time required to solve a singleobjective minimum cost flow problem and *N* is the number of all extreme supported nondominated vectors. Since the algorithm determines the extreme points in decreasing order of $c_1(f)$, we do not need additional time to sort the extreme points. Defining the weight vectors λ_i for $i = 1, ..., N - 1$ and building the network with the corresponding cost function takes $\mathcal{O}(N(n+m))$ time. Determining all S_i supported nondominated vectors on the maximally nondominated face F_i we have to solve S_i-1 ε -constraint problems.

Doing so for each of the maximally nondominated faces requires $\mathcal{O}(S(T(\varepsilon) + n + m))$
time. time.

4.4 A more compact formulation for the ILP in the *ε***-method**

The *ε*-constraint problem contains *m* variables and $2m + n + 1$ constraints. However, according to Theorem [3.8,](#page-9-0) each flow *f* ∗ can be written by an initial optimal tree solution *f* and a conical combination of incidence vectors of all *induced cycles* with bounded coefficients, i.e.,

$$
f^* = f + \sum_{a \notin T} \lambda_a \chi(C_a)
$$

for some $\lambda \in \mathbb{Z}$ and it holds that

$$
c(f^*) = c(f) + \sum_{a \notin T} \lambda_a c(C_a).
$$

Therefore instead of solving a constrained minimum cost problem, we also could solve the following ILP, which searches for the best combination of induced cycles such that the capacity constraints are satisfied.

$$
\min \sum_{a \notin T} \lambda_a c^1(C_a) \qquad \qquad \text{(c-MOMCF)}
$$
\n
$$
\text{s.t.} \quad 0 - f_{ij} \leq \sum_{\substack{a:(i,j)\in C_a}} \lambda_a \chi_{ij}(C_a) \leq u_{ij} - f_{ij} \quad \forall (i,j) \in \bigcup_{a \notin T} C_a
$$
\n
$$
\sum_{a \notin T} \lambda_a c^2(C_a) \leq \varepsilon
$$

Theorem 4.11 *For ε sufficiently large the set of feasible solutions of* [\(MOIMCF\)](#page-1-0) *and* [\(c-MOIMCF\)](#page-23-0) *coincide.*

Proof. Let f be an initial optimal tree solution. Assume that f^* is a feasible solution for Equation [\(MOIMCF\)](#page-1-0). After Theorem [3.8,](#page-9-0) the solution *f* ∗ can be written as $f^* = f + \sum_{a \notin T} \lambda_a^* \chi(C_a)$ for some $\lambda \in \mathbb{Z}$. Let λ_a^* for $a \notin T$ be the solution for Equa-tion [\(c-MOIMCF\)](#page-23-0), then $0-f_{ij} \leq \sum_{a:(i,j)\in C_a} \lambda_a \chi_{ij}(C_a) \leq u_{ij}-f_{ij}$ for all $(i,j) \in \bigcup_{a \notin T} C_a$ would be satisfied, since otherwise an arc (u, v) exists where the capacity constraint $0 \leq f_{uv}^* \leq u_{uv}$ would not hold. A contradiction of the feasibility of f^* . Now assume that λ'_a for $a \notin T$ is a feasible solution of Equation [\(c-MOIMCF\)](#page-23-0). Let $f' = f + \sum_{a \notin T} \lambda'_a \chi(C_a)$. Since f was an feasible initial optimal tree solution and we only change flow on cycles, it holds that $\sum_{j:(i,j)\in A} f'_{ij} - \sum_{j:(j,i)\in A} f'_{ji} = b_i$ for all $i\in V$. The boundaries of $0 \le f_{ij} \le u_{ij}$ for all $(i, j) \in A$ are also satisfied, since otherwise $u_{ij} - f_{ij}$ for one arc $(i, j) \in \bigcup_{a \notin T} C_a$ would be not satisfies. A contradiction.

 \Box

This ILP has *n* variables and at least $n-1$ constraints less than the standard ε constraint problem. According to Theorem [3.8,](#page-9-0) the respective sets of feasible solutions coincide. In Section [5,](#page-24-0) we evaluate numerically which approach is faster in practice, the standard *ε*-constraint method or the combination approach of the induced cycles. We also compare the running times to determine all supported nondominated vectors compared to the algorithms in Section [4.1](#page-14-1) and Section [4.2.](#page-16-0)

5 Numerical experiments

In this section, we present the implementation and numerical evaluation of the four methods. The section aims to report and compare the results, providing a comprehensive understanding of their behaviors.

All computations are conducted on a computer with an Intel® Core™i8-8700U CPU 3.20 GHz processor with 32 GB RAM, using a LINUX operating system. The algorithms are implemented and run in Python (Version 3.11). In addition, for solving the *ε*scalarizations Gurobi (Version 10.3) embedded in Python is used. To ensure a fair comparisons, the ILPs in Gurobi were solved using a single thread.

For the computational experiments, we utilized test instances from Figure [4.2](#page-20-1) and Figure [4.3,](#page-21-0) as well as minimum cost flow problem classes generated by the NETGEN network generator [\(Klingman et al.](#page-28-21), [1974](#page-28-21)). The entire test comprised 10 problem classes, with each class consisting of a set of 15 randomly generated network problems. The parameters that allowed the random generation of each NETGEN instances are the number of nodes, number of arcs, the number of nodes acting as supply or sink nodes, respectively, the greatest cost, greatest capacity, and the total supply in each network. In each problem class, the number of arcs and nodes varies. The instances ranged from 50 to 2000 nodes with 100 to 8000 arcs. These variables were chosen as independent variables due to the direct influence that they have in the number of possible supported nondominated vectors or supported efficient solutions, and, therefore influencing the different number of iterations of the different methods. All the other parameters that NETGEN can accept were kept constant. All instances have 2 nodes acting as supply nodes and 2 nodes acting as sink nodes, a maximum arc cost of 10 for both objective functions, a maximum upper capacity of 50, and a total supply of 50.

Results for the NETGEN instance classes are presented in Table [5.1,](#page-26-0) while the results of the test instances from Figure [4.2](#page-20-1) and Figure [4.3](#page-21-0) are summarized in Table [5.2](#page-27-0) and Table [5.3.](#page-27-1) These tables display the number of extreme nondominated vectors, supported nondominated vectors, supported efficient solutions, and CPU-time for all four methods. We display the min, max and mean of the times and numbers of the 15 network problems. Note, that we only consider problem instances with at least two nondominated vectors.

Table [5.2](#page-27-0) indicates that, as expected, the adjusted algorithm outperforms the all optimum algorithm when the number of branches needed is significantly smaller than the number of supported efficient solutions. However, when the number of branches needed equals the number of efficient solutions, the all optimal algorithm outperform the adjusted algorithm, as evident in Table [5.3.](#page-27-1) The high computational cost of the Floyd-Warshall algorithm used in each branch contributes to the adjusted algorithm's

suboptimal performance, as highlighted in Table [5.1](#page-26-0) and indicates that this algorithm performs badly in practice.

Despite this, when the number of supported nondominated vectors and supported efficient solutions equals, and therefore the all optimal flow algorithm would run in output-polynomial time to determine the number of all supported nondominated vectors, the objective-space methods clearly outperform this algorithm.

On a positive note, the more compact formulation of the *ε*-scalarization surpasses the classic formulation in each instance, as shown in Table [5.4.](#page-27-2) The mean CPU-Time needed in the more compact formulation for the *ε*-scalarization across all instance classes is 4*.*244 times faster than the standard formulation, demonstrating its efficiency. The time gap may increase in instances including more nodes and arcs. This is a nice result since the more compact formulation can also be used in any *ε*-scalarization method for multi-objective minimum cost flow problems with $d \geq 3$ objectives or the determination of all nondominated vectors for minimum cost flow problems.

In conclusion, the computational experiments provide valuable insights into the strengths and weaknesses of the implemented algorithms. The more compact formulation of *ε*scalarization presents a promising direction for future research, offering faster computation times and potential applications in multi-objective minimum cost flow problems with three or more objectives.

6 Conclusion

This paper discusses in detail the time complexity of enumerating all supported nondominated vectors for MOIMCF problems. The paper shows that there can not exist an output-polynomial time algorithm for the enumeration of all supported nondominated vectors that determine the vectors in an ordered manner in the outcome space unless $\mathbf{NP} = \mathbf{P}$. However, it remains an open question whether an output-polynomial time algorithm exists. Future research could focus on whether an output-polynomial time algorithm exists for the bi-objective or the multi-objective case with $d \geq 3$ objectives. The numerical tests show that the outcome space methods clearly outperform the decision-space methods, even if they do not run in output-polynomial time. The compact formulation of the ILP for the *ε*-method shows a significant time improvement compared to the conventional ILP. The new formulation can also be used for the computation of all nondominated vectors in multidimensional minimum cost flow problems and could be investigated in the future.

For (MOIMCF) problems, supported nondominated vectors are often only a small part of the complete set of nondominated vectors, even for the bi-objective case. However, determining supported nondominated vectors is often needed as a first step in two-phase exacts methods and for population-based heuristics. The development of improved twophase methods that compute all nondominated vectors for (MOIMCF) problems could be explored in the future.

Table 5.1: Numerical results for the different instance classes generated with NETGEN. T- $|\mathcal{Y}_{EN}|$ displays the time needed to determine the number of extreme vectors. T-AO, T-DS, T-*ε*, T-New-*ε* refers for the CPU-time in seconds needed for Algorithm [1,](#page-17-0) the adjusted version with Algorithm [3,](#page-20-0) the *ε*-Method, and the new more compact *ε*-Method. Empty entries (–) are reflecting a CPU-time of over 500 seconds and have not been recorded.

Class		$ \mathcal{Y}_{EN} $	$T- \mathcal{Y}_{EN} $	$ \mathcal{Y}_{SN} $	$ \mathcal{X}_{SN} $	T-AO	T-DS	T- ε	T-New- ε
1	min	$\overline{2}$	0.03	$\overline{4}$	$\overline{4}$	0.0058	0.1027	0.0055	0.003
$n=50$	max	10	0.1777	41	41	0.0602	1.26	0.0346	0.0195
$m=100$	mean	4.667	0.0836	19.73	19.73	0.0283	0.5501	0.0154	0.0087
$\overline{2}$	\min	$\overline{5}$	0.1836	23	24	0.0416	0.8	0.0175	0.0107
$n=50$	max	15	0.614	97	221	0.3842	8.12	0.1069	0.0338
$m=200$	mean	9.067	0.3546	42.33	52.6	0.1016	2.009	0.0422	0.0207
3	min	$\sqrt{2}$	0.1013	3	3	0.0119	0.4053	0.0048	0.003
$n=100$	max	14	0.9123	49	96	0.3851	15.45	0.1094	0.0402
$m=200\,$	mean	6.333	0.3965	25.53	29.6	0.1213	5.103	0.0628	0.0198
$\overline{4}$	min	6	0.7169	20	20	0.1122	4.276	0.0634	0.0183
$n = 100$	max	36	4.797	128	154	0.8551	33.03	0.3975	0.1325
$m=400$	mean	11.85	1.545	52.77	63.92	0.3516	11.65	0.1652	0.0447
5	min	$\overline{4}$	0.8857	$\overline{4}$	$\overline{4}$	0.0716	4.966	0.0844	0.0155
$n = 200$	max	17	4.121	82	86	1.145	109.5	0.4663	0.0936
$m=400$	mean	8.467	$2.018\,$	31.67	33.53	0.4812	41.46	0.2308	0.0482
6	min	$\overline{2}$	0.7583	5	5	0.0655	5.481	0.0299	0.0064
$n=200$	max	24	11.39	128	311	4.528	382.6	0.708	0.1501
$m=800\,$	mean	13.6	6.506	60.2	77.8	1.3	100.7	0.4	0.0853
$\overline{7}$	min	$\overline{4}$	20.17	10	10	3.185	\equiv	0.4253	0.0926
$n = 1000$	max	18	98.72	64	64	18.76	$\overline{}$	2.271	0.5435
$m = 2000$	mean	10.73	58.84	34.4	34.93	10.75	$\qquad \qquad -$	1.331	0.3006
8	min	$\,4\,$	39.73	$\overline{\mathcal{I}}$	$\overline{7}$	2.599	$\overline{}$	0.3993	0.0854
$n = 1000$	max	32	344.4	112	121	42.37	$\overline{}$	4.186	0.8983
$m = 4000$	mean	18.13	197.4	55.4	60.93	22.44	$\overline{}$	2.387	0.5203
9 $n = 2000$ $m = 4000$	\min max mean	$\overline{4}$ 24 13.4	81.46 542.8 301.3	9 57 36.13	9 124 44.13	16.42 139.1 55.47	\equiv $\overline{}$	1.345 9.661 5.307	0.2309 1.588 0.9287
10	min	$\boldsymbol{9}$	367.5	17	17	26.64	\equiv	3.4	0.6443
$n = 2000$	max	26	1099	90	220	272.8	$\overline{}$	10.69	1.956
$m = 8000$	mean	17.4	746.2	65.2	108.4	139.8	-	7.168	1.293

Table 5.2: Numerical results for the different instances of Figure [4.2.](#page-20-1) Each instance has $|\mathcal{Y}_{EN}| = 2$ extreme supported nondominated and $|\mathcal{Y}_{SN}| = 6$ nondominated supported vectors. For the first 5 instances, it holds $M = 10$, $L = 5$. For the last, it holds $M = 1, L = 3$. In thins instances the adjusted algorithm needs only $|\mathcal{Y}_{SN}| = 6$ branches.

Class	$ \mathcal{X}_{SN} $	$T-AO$	T-DS	$T - \epsilon$	$T-New-\epsilon$
$n=5$	726	0.06906	0.001218	0.00186	0.001399
$n=6$	7986	0.6985	0.001787	0.001866	0.001458
$n=7$	87846	7.733	0.001909	0.02054	0.001622
$n=8$	966306	92.69	0.002558	0.01949	0.001775
$n=9$	10629366	1197.7119	0.002758	0.0203	0.001847
$n=20$	786432.0	162.332160	0.013405	0.023102	0.002356

Table 5.3: Numerical results for the different instances of Figure [4.3.](#page-21-0) Each instance has $|\mathcal{Y}_{EN}| = 2$ extreme supported nondominated and $|\mathcal{Y}_{SN}| = 6$ nondominated supported vectors. For the first six instances, it holds $L = 5$. For the last $L = 3$. In this instances the adjusted algorithm needs exactly $|\mathcal{X}_{SN}|$ branches.

Class	$ \mathcal{X}_{SN} $	T-AO	T-DS	$T - \varepsilon$	$T-New-\epsilon$
$n=5$	56.0	0.005528	0.011958	0.001973	0.001445
$n=6$	126.0	0.013386	0.034731	0.001971	0.001618
$n=7$	252.0	0.031600	0.089560	0.002218	0.001579
$n=8$	462.0	0.065902	0.195041	0.002353	0.001674
$n=9$	792.0	0.124758	0.399943	0.002429	0.001775
$n=10$	1287.0	0.223086	0.819848	0.020767	0.002110
$n=20$	33649.0	12.846450	84.200655	0.020696	0.002846

Table 5.4: Displays the time difference of the *ε*-epsilon method versus the more compact formulation of the mean for each randomly generated NETGEN class instances of Table [5.1.](#page-26-0)

References

- Ahuja RK, Magnanti TL, Orlin JB (1993) Network Flows: Theory, Algorithms, and Applications, 1st edn. Prentice Hall
- Arimura H, Uno T (2005) An output-polynomial time algorithm for mining frequent closed attribute trees. In: Kramer S, Pfahringer B (eds) Inductive Logic Programming, Springer Berlin Heidelberg, Berlin, Heidelberg, p 1–19
- Bertsekas DP (1998) Network optimization. Athena Scientific
- Bökler F, Mutzel P (2015) Output-sensitive algorithms for enumerating the extreme nondominated points of multiobjective combinatorial optimization problems. In: Bansal N, Finocchi I (eds) Algorithms - ESA 2015, Springer Berlin Heidelberg, Berlin, Heidelberg, p 288–299
- Bökler F, Ehrgott M, Morris C, Mutzel P (2017) Output-sensitive complexity of multiobjective combinatorial optimization. Journal of Multi-Criteria Decision Analysis 24(1-2):25–36, DOI 10.1002/mcda. 1603
- Cook WJ, Cunningham WH, Pulleyblank WR, Schrijver A (1998) Combinatorial Optimization. John Wiley & Sons, New York
- Dantzig GB (1951) Application of the simplex method to a transportation problem. In: Koopmans TC (ed) Activity Analysis and Production and Allocation, Wiley, p 359–373
- Ehrgott M (2005) Multicriteria optimization, vol 491. Springer Science & Business Media
- Ehrgott M, Löhne A, Shao L (2012) A dual variant of Benson's "outer approximation algorithm" for multiple objective linear programming. Journal of Global Optimization 52(4):757–778, DOI 10.1007/ s10898-011-9709-y
- Eusébio A, Figueira JR (2009a) Finding non-dominated solutions in bi-objective integer network flow problems. Computers & Operations Research 36:2554–2564, DOI 10.1016/j.cor.2008.11.001
- Eusébio A, Figueira JR (2009b) Finding non-dominated solutions in bi-objective integer network flow problems. Computers & Operations Research 36(9):2554–2564, DOI 10.1016/j.cor.2008.11.001
- Eusébio A, Figueira JR (2009c) On the computation of all supported efficient solutions in multi-objective integer network flow problems. European Journal of Operational Research 199(1):68–76, DOI 10.1016/ j.ejor.2008.10.031
- Eusébio A, Figueira JR, Ehrgott M (2014) On finding representative non-dominated points for biobjective integer network flow problems. Computers & Operations Research 48:1–10, DOI 10.1016/j. cor.2014.02.009
- Fonseca M, Figueira J, Resende M (2010) Solving scalarized multi-objective network flow problems using an interior point method. International Transactions in Operational Research 17:607–636, DOI 10.1111/j.1475-3995.2010.00761.x
- Gal T (1977) A general method for determining the set of all efficient solutions to a linear vectormaximum problem. European Journal of Operational Research 1:307–322, DOI 10.1016/0377-2217(77)90063-7
- Haimes Y, Lasdon LS, Wismer D (1971) On a bicriterion formation of the problems of integrated system identification and system optimization. IEEE Transactions on Systems, Man, and Cybernetics pp 296–297
- Hamacher HW (1995) A note on k best network flows. Annals OR 57(1):65–72
- Hamacher HW, Pedersen CR, Ruzika S (2007) Multiple objective minimum cost flow problems: A review. European Journal of Operational Research 176(3):1404–1422
- Johnson DS, Yannakakis M, Papadimitriou CH (1988) On generating all maximal independent sets. Information Processing Letters 27(3):119–123, DOI 10.1016/0020-0190(88)90065-8
- Klingman DD, Napier A, Stutz JD (1974) Netgen: A program for generating large scale capacitated assignment, transportation, and minimum cost flow network problems. Management Science 20:814– 821, DOI 10.1287/MNSC.20.5.814
- Könen D, Schmidt D, Spisla C (2022) Finding all minimum cost flows and a faster algorithm for the k best flow problem. Discrete Applied Mathematics 321:333–349, DOI 10.1016/j.dam.2022.07.007
- Könen D, Stiglmayr M. (2023)An output-polynomial time algorithm to determine all supported efficient solutions for multi-objective integer network flow problems. arXiv preprint. DOI 10.48550/arXiv.2305. 12867
- Lawler EL, Lenstra JK, Rinnooy Kan AHG (1980) Generating all maximal independent sets: Nphardness and polynomial-time algorithms. SIAM Journal on Computing 9(3):558–565, DOI 10.1137/ 0209042, DOI 10.1137/0209042
- Okamoto Y, Uno T (2011) A polynomial-time-delay and polynomial-space algorithm for enumeration problems in multi-criteria optimization. European Journal of Operational Research 210(1):48–56, DOI 10.1016/j.ejor.2010.10.008
- Orlin JB (1993) A faster strongly polynomial minimum cost flow algorithm. Operations Research 41(2):338–350, DOI 10.1287/opre.41.2.338
- Pisinger D (1999) Linear time algorithms for knapsack problems with bounded weights. Journal of Algorithms 33(1):1–14, DOI 10.1006/jagm.1999.1034
- Raith A, Ehrgott M (2009a) A two-phase algorithm for the biobjective integer minimum cost flow problem. http://wwwescaucklandacnz/research/tech/esc-tr-661pdf 36, DOI 10.1016/j.cor.2008.06.008
- Raith A, Ehrgott M (2009b) A two-phase algorithm for the biobjective integer minimum cost flow problem. Comput Oper Res 36(6):1945–1954
- Raith A, Sedeño-Noda A (2017) Finding extreme supported solutions of biobjective network flow problems: An enhanced parametric programming approach. Computers & Operations Research 82, DOI 10.1016/j.cor.2017.01.004
- Sedeño-Noda A (2001) An algorithm for the biobjective integer minimum cost flow problem. Computers & Operations Research 28:139–156, DOI 10.1016/S0305-0548(99)00095-7
- Sedeño-Noda A, Gonzalez-Martin C (2000) Biobjective minimum cost flow problem. European Journal of Operational Research 124:591–600, DOI 10.1016/S0377-2217(99)00191-5
- Sedeño-Noda A, González-Martin C (2003) An alternative method to solve the biobjective minimum cost flow problem. Asia-Pacific Journal of Operational Research 20
- Sun M (2011) Finding integer efficient solutions for multiple objective network programming problems. Networks 57:362–375, DOI 10.1002/net.20407