

Entanglement and Pseudo Entanglement Dynamics versus Fusion in CFT

Song He^{a,b,1}, Yu-Xuan Zhang^{a,2}, Long Zhao^{a,3}, Zi-Xuan Zhao^{a,4}

^a*Center for Theoretical Physics and College of Physics, Jilin University,
Changchun 130012, People's Republic of China*

^b*Max Planck Institute for Gravitational Physics (Albert Einstein Institute),
Am Mühlenberg 1, 14476 Golm, Germany*

Abstract

The fusion rules and operator product expansion (OPE) serve as crucial tools in the study of operator algebras within conformal field theory (CFT). Building upon the vision of using entanglement to explore the connections between fusion coefficients and OPE coefficients, we employ the replica method and Schmidt decomposition method to investigate the time evolution of entanglement entropy (EE) and pseudo entropy (PE) for linear combinations of operators in rational conformal field theory (RCFT). We obtain a formula that links fusion coefficients, quantum dimensions, and OPE coefficients. We also identify two definition schemes for linear combination operators. Under one scheme, the EE captures information solely for the heaviest operators, while the PE retains information for all operators, reflecting the phenomenon of pseudo entropy amplification. Irrespective of the scheme employed, the EE demonstrates a step-like evolution, illustrating the effectiveness of the quasiparticle propagation picture for the general superposition of locally excited states in RCFT. From the perspective of quasiparticle propagation, we observe spontaneous block-diagonalization of the reduced density matrix of a subsystem when quasiparticles enter the subsystem.

¹hesong@jlu.edu.cn

²yuxuanz20@mails.jlu.edu.cn

³zhaolong@jlu.edu.cn

⁴zzx23@mails.jlu.edu.cn

Contents

1	Introduction	1
2	Pseudo entropy amplification and fusion rules	6
2.1	The late-time formula of n -th PREE/REE	6
2.1.1	Schmidt decomposition method	6
2.1.2	Replica method	9
2.2	Pseudo entropy amplification	12
2.2.1	The REE and EE	13
2.2.2	The PREE and PE	14
2.3	A connection between Fusion numbers and OPE coefficients	16
3	The entanglement dynamics of refined linear combination operators (18)	17
3.1	Recovering information of lighter operators in the EE	18
3.2	The quasiparticle picture	19
3.3	Block diagonalization of the reduced density matrix at late times	22
4	Conclusion and prospect	24
A	The product of two semi-positive matrices is diagonalizable	25

1 Introduction

In 2D conformal field theory (CFT), fusion rules determine which operators appear in the operator product expansion (OPE) of two primary fields, originating from the BPZ equations [1, 2]. These rules are undoubtedly crucial since for 2D CFT all correlation functions can be reduced to the two-point function via the OPE. One can summarize the fusion rules in the following commutative and associative fusion algebra [3, 4]

$$[\mathcal{O}_i] \times [\mathcal{O}_j] = \sum_k N_{ij}^k [\mathcal{O}_k], \quad (1)$$

where $[\mathcal{O}_i]$ denotes the conformal family of the primary \mathcal{O}_i , and N_{ij}^k called fusion numbers are non-negative integers. Note one can rephrase eqn. (1) in terms of the quantum dimension d_i of the associated conformal family $[\mathcal{O}_i]$ [3, 5]

$$d_i d_j = \sum_k N_{ij}^k d_k. \quad (2)$$

In rational conformal field theory (RCFT), it is well-known that the fusion numbers N_{ij}^k can be computed using the Verlinde formula [3]. The counterpart in irrational conformal field theory (e.g., Liouville field theory) was studied in [6].

Interestingly, recent studies on quantum entanglement in RCFT have seemingly offered a new perspective on the connection between fusion numbers and OPE coefficients [7–12]. The related topic is called *local operator quench* [7], where one first generates a locally excited state through acting with a local operator \mathcal{O} on the vacuum state $|\Omega\rangle$ of the theory of interest, and then evolves it with the Hamiltonian of the theory,

$$|\psi(t)\rangle := \frac{1}{\sqrt{\mathcal{N}(\epsilon)}} e^{-iH(t-i\epsilon)} \mathcal{O}(x) |\Omega\rangle, \quad (3)$$

where ϵ is an infinitesimal positive parameter and $\mathcal{N}(\epsilon)$ is a normalization factor such that $\langle \psi(t) | \psi(t) \rangle = 1$. It has been shown that the local operator quench exhibits broad applicability in measuring scrambling and thermalization effects in CFTs with large central charge [13–17], as well as in probing the bulk geometry [18] and characterizing bulk dynamics [19–22] in the context of AdS/CFT correspondence [23–25]. Investigating the local operator quench in RCFT is driven by the quest to understand the intricate connections between fusion rules, entanglement properties, and the broader implications for quantum information and computing. In particular, the fusion rules and their role in entanglement in CFT have connections to topological quantum computing [26–28]. The entanglement dynamics obtained from such investigations contribute to theoretical advancements and potential practical applications.

The process of local operator quench provides an excellent platform for the study of entanglement dynamics between subsystems. In RCFT local primary or descendant operator quench, i.e., the local operator \mathcal{O} in eqn. (3) is a primary or descendant operator, it was found that the variations⁵ of entanglement entropy (EE, defined as $S(\rho_A) := -\text{tr}[\rho_A \log \rho_A]$) and Rényi entanglement entropy (REE, defined as $S^{(n)}(\rho_A) := \frac{1}{1-n} \log \text{tr}[\rho_A^n]$) saturate to a constant equal to the logarithm of quantum dimension of the local operator [8–10]. The result has been generalized to various cases [11–14, 17, 19, 21, 22, 29–36], including the multiple local excitations [11, 12],

$$|\Psi(t)\rangle = \frac{1}{\sqrt{\mathcal{N}(\epsilon)}} e^{-iH(t-i\epsilon)} \mathcal{O}_N(x_N) \dots \mathcal{O}_i(x_i) \dots \mathcal{O}_1(x_1) |\Omega\rangle. \quad (5)$$

In RCFT multiple local operator quenches, the finiteness of quantum dimension guarantees the conservation of entanglement under the scattering of quasiparticles [11]. When the subsystem A is semi-infinite, the variations of the EE and REE at late times are solely the summation of the contributions from individual local operator quenches, i.e.,

$$\lim_{t \rightarrow \infty} \Delta S^{(n)}(\rho_A^\Psi(t)) = \sum_{i=1}^N \lim_{t \rightarrow \infty} \Delta S^{(n)}(\rho_A^{\mathcal{O}_i}(t)) = \sum_{i=1}^N \log d_i, \quad (6)$$

where d_i is the quantum dimension of \mathcal{O}_i [11, 12].

This paper mainly focuses on the local quenches of linear combination operators in RCFT. Specifically, our interest centers on cases wherein local operator \mathcal{O} in eqn. (3) assumes the form

$$\mathcal{O}(x) = \sum_{k, \{\alpha\}, \{\bar{\alpha}\}} C_k^{\{\alpha, \bar{\alpha}\}} \cdot L^{2h_k + |\alpha| + |\bar{\alpha}|} \cdot \mathcal{O}_k^{\{\alpha, \bar{\alpha}\}}(x). \quad (7)$$

In the above, $C_k^{\{\alpha, \bar{\alpha}\}}$ are dimensionless combination coefficients, L is a constant with a dimension of length (such as the length of subsystem), and $\mathcal{O}_k^{\{\alpha, \bar{\alpha}\}}(x)$ are descendant operators; $\mathcal{O}_k^{\{\alpha, \bar{\alpha}\}}(x) \equiv L_{-\alpha_1} \dots \bar{L}_{-\bar{\alpha}_1} \dots \mathcal{O}_k(x)$, $|\alpha| \equiv \sum_i \alpha_i$. Generically, in RCFTs, when the subsystem A is semi-infinite, one may expect that the variation of the EE of operators taking the form of (7) at late times should be a function of $C_k^{\{\alpha, \bar{\alpha}\}}$ and d_k , denoted as $\log(f(C_k^{\{\alpha, \bar{\alpha}\}}, d_k))$.

The motivation for studying such kinds of local operator quenches will be clear from the

⁵The variations of EE and REE, denoted as $\Delta S(\rho_A^\mathcal{O}(t))$ and $\Delta S^{(n)}(\rho_A^\mathcal{O}(t))$ respectively in this paper, are defined as

$$\Delta S(\rho_A^\mathcal{O}(t)) \equiv \lim_{n \rightarrow 1} \Delta S^{(n)}(\rho_A^\mathcal{O}(t)), \quad \Delta S^{(n)}(\rho_A^\mathcal{O}(t)) := S^{(n)}(\rho_A^\mathcal{O}(t)) - S^{(n)}(\rho_A^\Omega), \quad (4)$$

where $\rho_A^\mathcal{O}(t)$ is the reduced density matrix of subsystem A when the total system is in the state $|\psi(t)\rangle$ (3), and ρ_A^Ω is the reduced density matrix for the vacuum state $|\Omega\rangle$.

perspective of OPE. The OPE of two spin-zero primary fields \mathcal{O}_i takes the following form⁶

$$\mathcal{O}_i(x_1)\mathcal{O}_j(x_2) = \sum_k C_{kij} \sum_{\{\alpha\},\{\bar{\alpha}\}} \beta_{ij}^{k\{\alpha\}} \bar{\beta}_{ij}^{k\{\bar{\alpha}\}} |x_1 - x_2|^{\alpha+|\bar{\alpha}|+2h_k-2h_i-2h_j} \mathcal{O}_k^{\{\alpha,\bar{\alpha}\}}(x_2). \quad (8)$$

One can utilize the operators on both sides of the OPE (8) to construct locally excited states. Assume that the subsystem A is semi-infinite. For multiple local excitations (5) generated by the composite operator on the l.h.s. of the OPE (8), we know from eqn. (6) that the variation of the EE (as well as REE) of the multiple locally excited state will eventually saturate to $\log d_i + \log d_j = \log d_i d_j = \log \sum_k N_{ij}^k d_k$, where the second equality comes from the fusion rule of \mathcal{O}_i and \mathcal{O}_j (see eqn. (2)). On the other hand, given that the linear combination operator on the r.h.s. of the OPE (8) is solely a specific instance of eqn. (7), and based on the discussion following eqn. (7), we conclude that the variation of the EE of the operator on the r.h.s. of the OPE (8) saturate to $\log (f(C_{kij} \beta_{ij}^{k\{\alpha\}} \bar{\beta}_{ij}^{k\{\bar{\alpha}\}}, d_k))$. Whether starting from the l.h.s. of the equation (8) or the right side, the results for the variation of the EE obtained should be the same. The function f thus characterizes a connection between the OPE coefficients and fusion numbers in RCFT since, from the above discussions, we have $f(C_{kij} \beta_{ij}^{k\{\alpha\}} \bar{\beta}_{ij}^{k\{\bar{\alpha}\}}, d_k) = \sum_k N_{ij}^k d_k$. This is the primary motivation for studying the EE and REE associated with the linear combination operator (7) in this paper.

Even in Ising CFT, however, employing the standard replica method [37, 38] to calculate the EE of a linear combination operator containing numerous descendant fields proves to be intricate (see [10] for a particular combination of descendant fields). Hence, here, we begin with a simplified scenario, excluding all descendant operators from (7)⁷,

$$|\psi\rangle = \frac{1}{\sqrt{\mathcal{N}(\epsilon)}} e^{-\epsilon H} \mathcal{O}(x) |\Omega\rangle, \quad \mathcal{O}(x) = \sum_{k \in \mathcal{S}} C_k \cdot L^{2h_k} \cdot \mathcal{O}_k(x). \quad (9)$$

where \mathcal{S} is a primary operator set $\mathcal{S} := \{\mathcal{O}_1, \mathcal{O}_2, \dots, \mathcal{O}_m\}$ ($m \in \mathbb{Z}_+$) from a RCFT. Additionally, we assume that the operators \mathcal{O}_k in \mathcal{S} are orthogonal to each other in terms of the two-point function, such that $\langle \mathcal{O}_k^\dagger \mathcal{O}_l \rangle \propto \delta_{kl}$. Although such a blunt truncation overlooks the contributions of all descendant fields to entanglement, we anticipate that our analysis of the linear combination operator (9) will provide valuable clues for the subsequent general discussions. We will employ the replica method and the Schmidt decomposition method [12] to compute the late-time EE of the linear combination operator (9). The latter method is a quantum mechanical approach that circumvents the complicated computations of all higher-point correlation functions. The consistency in results obtained from both methods instills confidence in utilizing the latter to compute the EE of operators on the r.h.s of the OPE (8).

On the other hand, we expect that our simple linear combination operators (9) can reproduce the fusion relations (2) in RCFTs in a heuristic manner. Our strategy is as follows: starting from a known fusion equation (1), we select all primary operators appearing on the r.h.s. of it to construct the linear combination operator in (9). Subsequently, we compute the EE (REE) for the state (9). It will be a function of the combination coefficients $\{C_k\}$ and operator quantum dimensions $\{d_k\}$. We look for the possibility in the parameter space of $\{C_k\}$ that operator (9) produces the same EE (i.e., $\log \sum_k N_{ij}^k d_k$) as the linear combination operator on the r.h.s. of OPE (8) corresponding to the given fusion equation. It is intriguing that, however, when attempting this, it can be proven that the EE and REE for the state (9) only capture information about the heaviest operators (those with the highest conformal dimensions) in \mathcal{S} .

⁶We assume that x_1 and x_2 lie on the real axis.

⁷Note that two operators up to a global factor yield the same state. In this paper, we partially fix this redundancy by imposing the normalization constraint $\sum_{k \in \mathcal{S}} |C_k|^2 = 1$.

For the most extreme situation, consider the case in which the set includes a unique heaviest primary operator \mathcal{O}_M , the expression for the late-time REE is

$$\lim_{t \rightarrow \infty} \Delta S^{(n)}(\rho_A^\psi(t)) = \log d_M, \quad (10)$$

where d_M is the quantum dimension of \mathcal{O}_M , and its conformal dimension is denoted by $h_M = \max_{k \in \mathcal{S}} \{h_k\}$. In this situation, information regarding all operators apart from \mathcal{O}_M is entirely lost.

To reproduce relation (2) using our simple linear combination operators (9), let us momentarily set aside the EE and replace it with a generalisation version, known as pseudo entropy (PE). PE is a two-state generalization of EE recently proposed via AdS/CFT and post-selection processes [39]. It is defined as follows,

$$S(\mathcal{T}_A^{\psi|\varphi}) = -\text{tr}[\mathcal{T}_A^{\psi|\varphi} \log \mathcal{T}_A^{\psi|\varphi}]. \quad (11)$$

The operator $\mathcal{T}_A^{\psi|\varphi}$ involving two states (i.e., the initial state ψ and the final state φ in the context of post-selection experiments [40, 41]), called reduced transition matrix, is the partial trace of the transition matrix $\mathcal{T}^{\psi|\varphi}$ [39, 42],

$$\mathcal{T}^{\psi|\varphi} = \frac{|\psi\rangle\langle\varphi|}{\langle\varphi|\psi\rangle} = \frac{\rho^\psi \rho^\varphi}{\text{tr}[\rho^\psi \rho^\varphi]}, \quad \mathcal{T}_A^{\psi|\varphi} = \text{tr}_{\bar{A}}[\mathcal{T}^{\psi|\varphi}]. \quad (12)$$

Like the calculation of EE, one usually computes the so-called pseudo-Rényi entanglement entropy (PREE) first. Subsequently, the PE is obtained by taking the limit of $n \rightarrow 1$ to the n -th PREE,

$$S^{(n)}(\mathcal{T}_A^{\psi|\varphi}) = \frac{1}{1-n} \log \text{tr}[(\mathcal{T}_A^{\psi|\varphi})^n], \quad S(\mathcal{T}_A^{\psi|\varphi}) = \lim_{n \rightarrow 1} S^{(n)}(\mathcal{T}_A^{\psi|\varphi}). \quad (13)$$

In previous studies [43, 44], it was shown that, for subsystem A being half-space, the n -th PREE under local primary or descendant operator quenches converges to the logarithm of the quantum dimension of the local operator as time t approaches infinity.⁸ In this work, to construct the transition matrix $\mathcal{T}^{\psi|\tilde{\psi}}$, we choose the state in eqn. (9) as $|\psi\rangle$ and select another state $|\tilde{\psi}\rangle$ as

$$|\tilde{\psi}\rangle = \mathcal{N}(\epsilon)^{-1/2} e^{-\epsilon H} \mathcal{O}(\tilde{x}) |\Omega\rangle. \quad (14)$$

As outlined in section 2.1, the advantage of the PE of the reduced transition matrix $\mathcal{T}_A^{\psi|\tilde{\psi}}$ compared with EE is its capability to retain information about all primary operators within \mathcal{S} . Specifically, the maximal value attainable by $S^{(n)}(\mathcal{T}_A^{\psi|\tilde{\psi}})$ but not achievable by $S^{(n)}(\rho_A^\psi)$ in general is the logarithm of the sum of the quantum dimensions of all primary operators,

$$\max_{\{C_k\}} \left\{ \lim_{t \rightarrow \infty} \Delta S^{(n)}(\mathcal{T}_A^{\psi|\tilde{\psi}}(t)) \right\} = \log \left(\sum_{k \in \mathcal{S}} d_k \right). \quad (15)$$

Comparing (15) with (10), we observe that the late-time value of PE is larger than the EE, a phenomenon identified in [53] as the *pseudo entropy amplification*.

The PE enables us to reproduce the fusion equation (1) in RCFTs under the rough truncation from eqn. (7) to eqn. (9). Consider, for instance, one of the fusion equations in the critical Ising model:

$$\sigma \times \sigma = \mathbb{I} + \varepsilon, \quad (16)$$

⁸See [45–72] for more developments on PE and related topics.

where σ is the spin operator, ε is the energy density operator and \mathbb{I} the identity. The quantum dimensions of three primaries are $d_\sigma = \sqrt{2}$, $d_\varepsilon = d_{\mathbb{I}} = 1$ respectively. We will analyze the late-time PE of the transition matrices $\mathcal{T}^{\psi|\tilde{\psi}}$ generated by both sides of eqn. (16): the composite operator $\sigma(x+L)\sigma(x)$ and the linear combination operator $C_{\mathbb{I}} \cdot \mathbb{I} + C_\varepsilon \cdot L^{2h_\varepsilon} \cdot \varepsilon(x)$. For the composite operator $\sigma(x+L)\sigma(x)$, its late-time PE equals its late-time EE [43]. Hence, according to eqn. (6), the value of the late-time PE we arrive at is $\log d_\sigma + \log d_\sigma = \log d_\sigma^2 = \log 2$. On the other hand, considering the set $\mathcal{S} = \{\mathbb{I}, \varepsilon\}$, according to eqn. (15), we find that the maximum value of the PE precisely equals $\log 2 = \log(d_{\mathbb{I}} + d_\varepsilon)$. Thus, we can always identify a set of appropriate combination coefficients $\{C_{\mathbb{I}}, C_\varepsilon\}$ to reproduce the fusion equation $d_\sigma \times d_\sigma = d_{\mathbb{I}} + d_\varepsilon$ through the PE.

As previously mentioned, if we aim not only to reproduce fusion relations (2) using quantum entanglement but also to establish a connection between fusion coefficients and OPE coefficients, the contributions of all possible descendant operators in the OPE must be added. Although counting the contributions of an infinite number of operators seems implausible, Ref. [12] presented an ingenious approach. By imposing identical conformal transformations on both sides of the OPE (8), the authors in [12] deduced the contribution to EE from each conformal family that appears to the r.h.s. of the OPE (8), i.e., the EE of the operator

$$\sum_{\{\alpha\}, \{\bar{\alpha}\}} \beta_{ij}^{k\{\alpha\}} \bar{\beta}_{ij}^{k\{\bar{\alpha}\}} |x_1 - x_2|^{\alpha + |\bar{\alpha}| + 2h_k - 2h_i - 2h_j} \mathcal{O}_k^{\{\alpha, \bar{\alpha}\}}(x_2). \quad (17)$$

For RCFTs, this contribution precisely equates to the logarithm of the corresponding quantum dimension (i.e., $\log d_k$). Based on this nontrivial result, we can utilize the Schmidt decomposition method to derive the late-time EE of the linear combination operator on the r.h.s. of the OPE (8). The result is a function of the OPE coefficients and quantum dimensions (see eqn. (76)). As we mentioned in the discussions after eqn. (8), this function characterizes a relationship between the OPE coefficients and the fusion coefficients because we have $f(C_{kij} \beta_{ij}^{k\{\alpha\}} \bar{\beta}_{ij}^{k\{\bar{\alpha}\}}, d_k) = \sum_k N_{ij}^k d_k$. By means of the technique we used in exploring the maximum PE of the simplified linear combination operator (9) (further details provided in section 2.3), we finally establish a link (see eqn. (79)) between fusion coefficients, quantum dimensions, and OPE coefficients using the function (76).

Although we reproduce the fusion relations by computing the PE of the linear combination operator (9) and establish the correlation between fusion coefficients and OPE coefficients through the linear combination operator like (7), these operators might not be suitable when considering the EE of a finite number of linear combinations of operators. As mentioned earlier, the information about light operators doesn't manifest in the EE. To find a way by which the information about light operators in \mathcal{S} can be expressed within the EE is the second objective of this paper. It turns out that by considering the following refined linear combination operator

$$\mathcal{O}(x; \epsilon) = \sum_{k \in \mathcal{S}} C_k \frac{\mathcal{O}_k(x)}{\sqrt{\langle \mathcal{O}_k^\dagger(x, \epsilon) \mathcal{O}_k(x, -\epsilon) \rangle}} = \sum_{k \in \mathcal{S}} C_k \cdot |2\epsilon|^{2h_k} \cdot \mathcal{O}_k(x) \quad (18)$$

instead of the linear combination operator in eqn. (9), all information of operators in \mathcal{S} can be encoded into the late-time EE (see section 3.1 for more details). We can delve into further details of entanglement dynamics about operator (18), particularly regarding the applicability of the quasiparticles picture. It's well-known that each primary operator \mathcal{O}_k in (18) can generate a locally excited state with a classical interpretation of quasiparticles propagation [7, 8], whether the state generated by operator (18) (i.e., the superposition of locally primary excited states) still possesses the quasiparticles picture is not immediately apparent. A common counterexample is that in quantum mechanics the superposition of coherent (quasiclassical) states usually no longer has a classical correspondence. Using the replica method, we find that the

answer is positive. The evolution of EE (and REE) corresponding to operator (18) shows a step-like behavior from zero to non-zero, which suggests that the quasiparticle propagation picture indeed holds for the general superposition of locally excited states in RCFT. Another fascinating detail in entanglement dynamics is our observation that, under the quasiparticle propagation picture, upon the entry of quasiparticles into a subsystem, the reduced density matrix of the subsystem spontaneously undergoes instantaneous block-diagonalization. Each distinct diagonal block carries information about different conformal families.

The rest of this paper is organized as follows. In section 2, we derive the late-time REE and PREE of the operator (9) through two methods, the Schmidt decomposition approach and the replica trick. We analyze the dependence of REE and PREE on the combination coefficients, showcasing the emergence of pseudo entropy amplification by demonstrating the maximum values of REE and PREE. Utilizing the Schmidt decomposition method, we establish a connection between fusion and OPE coefficients by examining the REE of operators on both sides of the OPE. In section 3, we illustrate how operator (18) facilitates the restoration of information related to light operators in the EE (REE). Additionally, we present further details of the entanglement dynamics associated with operator (18). The summary and prospect are given in section 4.

2 Pseudo entropy amplification and fusion rules

The earliest exploration of the EE (REE) of a superposition of locally primary excited states appeared in [12], followed by an extension of the PE (PREE) by the authors of [43]. According to the Schmidt decomposition method [12], the late-time limit of the n -th PREE of the reduced transition matrix generated by operator (9) is given by [43]

$$\lim_{t \rightarrow \infty} \Delta S^{(n)}(\mathcal{T}_A^{\psi|\tilde{\psi}}(t)) = \frac{1}{1-n} \log \left[\sum_{k \in \mathcal{S}} \left(\frac{d_k^{\frac{1}{n}-1} |C_k|^2 L^{4h_k} \langle \mathcal{O}_k^\dagger(\tilde{x}, \epsilon) \mathcal{O}_k(x, -\epsilon) \rangle}{\sum_{j \in \mathcal{S}} |C_j|^2 L^{4h_j} \langle \mathcal{O}_j^\dagger(\tilde{x}, \epsilon) \mathcal{O}_j(x, -\epsilon) \rangle} \right)^n \right]. \quad (19)$$

Setting $\tilde{x} = x$ allows one to obtain the late-time EE associated with operator (9). Despite the good agreement between formula (19) and numerical computations, an analytical derivation based on the standard method of the CFT replica trick remains absent to date. In this section, we bridge this gap by providing replica calculations for the cases of $n = 2$ and $n = 3$, and the computation for general n can be obtained straightforwardly. Subsequently, we aim to extract intriguing information from the late-time PREE formula. We show the pseudo entropy amplification phenomenon by exploring the maximum values of PREE and REE of locally excited states (9). These research findings point towards a potential relationship between fusion and OPE coefficients as demonstrated in subsection 2.3.

2.1 The late-time formula of n -th PREE/REE

In replica method, calculating the n -th REE/PREE for an operator \mathcal{O} , which consists of a linear combination of m primary operators, requires working with m^{2n} $2n$ -point functions on the plane. In this subsection, we first review the derivation of eqn. (19) by Schmidt composition method in Ref. [43], and then we will demonstrate the replica calculations of the second and third REE and PREE to derive the same results (19).

2.1.1 Schmidt decomposition method

As mentioned earlier, operators in \mathcal{S} are mutually orthogonal in the sense of the two-point function $\langle \mathcal{O}_k^\dagger \mathcal{O}_l \rangle$. Let us first define a series of normalized excited states with the help of

$\mathcal{O}_k \in \mathcal{S}$,

$$|\mathcal{O}_k(x, t)\rangle := \frac{1}{\sqrt{\langle \mathcal{O}_k^\dagger(x, \epsilon) \mathcal{O}_k(x, -\epsilon) \rangle}} e^{-iHt} \mathcal{O}_k(x, -\epsilon) |\Omega\rangle, \quad \langle \mathcal{O}_k(x, t) | \mathcal{O}_l(x, t) \rangle = \delta_{kl}. \quad (20)$$

Generally speaking, $|\mathcal{O}_k(x, t)\rangle$ is an entangled state living in the sub-Verma module $V_k \otimes \bar{V}_k$. It can be written in the following form by a time-dependent Schmidt decomposition

$$|\mathcal{O}_k(x, t)\rangle = \sum_u a_u^k(x, t) |k_u(x, t)\rangle \otimes |\bar{k}_u(x, t)\rangle, \quad (21)$$

where $\{|k_u(x, t)\rangle\}$ and $\{|\bar{k}_u(x, t)\rangle\}$ parameterized by (x, t) are two orthonormal bases of V_k and \bar{V}_k respectively, and $a_u^k(x, t) \geq 0$ ($\sum_u (a_u^k(x, t))^2 = 1$) are Schmidt coefficients. The REE between V_k and \bar{V}_k for the state $|\mathcal{O}_k(x, t)\rangle$ is directly given by

$$\begin{aligned} S^{(n)}(\rho_{V_k}^{\mathcal{O}_k}(t)) &= \frac{1}{1-n} \log \left\{ \text{tr}_{V_k} \left[(\text{tr}_{\bar{V}_k} |\mathcal{O}_k(x, t)\rangle \langle \mathcal{O}_k(x, t)|)^n \right] \right\} \\ &= \frac{1}{1-n} \log \sum_u (a_u^k(x, t))^{2n}. \end{aligned} \quad (22)$$

The next crucial point lies in the identification⁹ of $\Delta S^{(n)}(\rho_A^{\mathcal{O}_k}(t))$ and $S^{(n)}(\rho_{V_k}^{\mathcal{O}_k}(t))$ in the large- t limit, where

$$\rho_A^{\mathcal{O}_k}(t) \equiv \text{tr}_{\bar{A}} [|\mathcal{O}_k(x, t)\rangle \langle \mathcal{O}_k(x, t)|] \quad (23)$$

representing the reduced density matrix of subsystem $A = [0, \infty)$ when the total system is quenched by $\mathcal{O}_k(x)$ at the initial time. Consequently, we arrive at the limit as $t \rightarrow \infty$ [8]

$$\lim_{t \rightarrow \infty} S^{(n)}(\rho_{V_k}^{\mathcal{O}_k}(t)) = \log d_k, \quad (24)$$

where d_k is the quantum dimension of \mathcal{O}_k . Now we move to investigate the PREE of $e^{-iHt}|\psi\rangle$ (9) and $e^{-iHt}|\tilde{\psi}\rangle$ (14). These two states can be rewritten as two superposition states of $|\mathcal{O}_k(x, t)\rangle$

$$e^{-iHt}|\psi\rangle = \sum_{k \in \mathcal{S}} \sqrt{\lambda_k} |\mathcal{O}_k(x, t)\rangle, \quad e^{-iHt}|\tilde{\psi}\rangle = \sum_{k \in \mathcal{S}} \sqrt{\tilde{\lambda}_k} |\mathcal{O}_k(\tilde{x}, t)\rangle, \quad (25)$$

where

$$\lambda_k = \tilde{\lambda}_k = \frac{(C_k)^2 L^{4h_k} \langle \mathcal{O}_k^\dagger(x, \epsilon) \mathcal{O}_k(x, -\epsilon) \rangle}{\sum_{j \in \mathcal{S}} |C_j|^2 L^{4h_j} \langle \mathcal{O}_j^\dagger(x, \epsilon) \mathcal{O}_j(x, -\epsilon) \rangle}, \quad \sum_{k \in \mathcal{S}} |\lambda_k| = 1. \quad (26)$$

Consequently, the transition matrix becomes

$$\begin{aligned} \mathcal{T}^{\psi|\tilde{\psi}}(t) &= \frac{1}{\sum_{k \in \mathcal{S}} \sqrt{\lambda_k} \sqrt{\tilde{\lambda}_k} \langle \mathcal{O}_k(\tilde{x}, t) | \mathcal{O}_k(x, t) \rangle} \\ &\times \sum_{j, l \in \mathcal{S}} \sqrt{\lambda_j} \sqrt{\tilde{\lambda}_l} \sum_{u, v} a_u^j(x, t) a_v^l(\tilde{x}, t) |j_u(x, t)\rangle \langle \bar{j}_u(x, t)| \langle l_v(\tilde{x}, t) | \langle \bar{l}_v(\tilde{x}, t)|. \end{aligned} \quad (27)$$

⁹This identification can be understood through the following physical intuition: In the quasiparticles propagation picture, in the large- t limit, subsystem $A = [0, \infty)$ collects all the right-moving quasiparticles, which carry information about the holomorphic part of \mathcal{O}_k , while the complement \bar{A} collects all the left-moving quasiparticles, which carry information about the anti-holomorphic part of \mathcal{O}_k [7] [73]

The reduced transition matrix is obtained by tracing out the anti-holomorphic part ($V \equiv \oplus_k V_k$),

$$\begin{aligned} \mathcal{T}_V^{\psi|\tilde{\psi}}(t) &= \text{tr}_{\oplus_{\bar{k}} \bar{V}_{\bar{k}}} \mathcal{T}^{\psi|\tilde{\psi}}(t) \\ &= \sum_{k \in \mathcal{S}} \sum_{u,v,w} \frac{\sqrt{\lambda_k} \sqrt{\tilde{\lambda}_k^*} a_u^k(x,t) a_v^k(\tilde{x},t) \langle \bar{k}_v(\tilde{x},t) | \bar{k}_u(x,t) \rangle \langle k_v(\tilde{x},t) | k_w(x,t) \rangle}{\sum_{j \in \mathcal{S}} \sqrt{\lambda_j} \sqrt{\tilde{\lambda}_j^*} \langle \mathcal{O}_j(\tilde{x},t) | \mathcal{O}_j(x,t) \rangle} \cdot |k_u(x,t)\rangle \langle k_w(x,t)|, \end{aligned} \quad (28)$$

which in general is nondiagonal under the basis $\{|k_u(x,t)\rangle\}$. We can compute the trace of $(\mathcal{T}_V^{\psi|\tilde{\psi}}(t))^n$,

$$\begin{aligned} & \text{tr} \left[(\mathcal{T}_V^{\psi|\tilde{\psi}}(t))^n \right] \\ &= \sum_{k \in \mathcal{S}} \frac{\left(\sqrt{\lambda_k} \sqrt{\tilde{\lambda}_k^*} \right)^n}{\left(\sum_{j \in \mathcal{S}} \sqrt{\lambda_j} \sqrt{\tilde{\lambda}_j^*} \langle \mathcal{O}_j(\tilde{x},t) | \mathcal{O}_j(x,t) \rangle \right)^n} \\ & \quad \times \sum_{\{u\}, \{v\}} a_{u_1}^k(x,t) a_{v_1}^k(\tilde{x},t) \langle \bar{k}_{v_1}(\tilde{x},t) | \bar{k}_{u_1}(x,t) \rangle \langle k_{v_1}(\tilde{x},t) | k_{u_2}(x,t) \rangle \dots a_{u_n}^k(x,t) a_{v_n}^k(\tilde{x},t) \\ & \quad \times \langle \bar{k}_{v_n}(\tilde{x},t) | \bar{k}_{u_n}(x,t) \rangle \langle k_{v_n}(\tilde{x},t) | k_{u_1}(x,t) \rangle. \end{aligned} \quad (29)$$

To further reduce (29), let us turn to consider the n -th PREE of $|\mathcal{O}_k(x,t)\rangle$ and $|\mathcal{O}_k(\tilde{x},t)\rangle$.

$$\begin{aligned} S^{(n)}(\mathcal{T}_V^{\mathcal{O}_k(x)|\mathcal{O}_k(\tilde{x})}(t)) &= \frac{1}{1-n} \log \text{tr}_{\oplus_k V_k} \left[(\mathcal{T}_V^{\mathcal{O}_k(x)|\mathcal{O}_k(\tilde{x})}(t))^n \right] \\ &= \frac{1}{1-n} \log \left[\langle \mathcal{O}_k(\tilde{x},t) | \mathcal{O}_k(x,t) \rangle^{-n} \right. \\ & \quad \times \sum_{\{u\}, \{v\}} a_{u_1}^k(x,t) a_{v_1}^k(\tilde{x},t) \langle \bar{k}_{v_1}(\tilde{x},t) | \bar{k}_{u_1}(x,t) \rangle \langle k_{v_1}(\tilde{x},t) | k_{u_2}(x,t) \rangle \dots a_{u_n}^k(x,t) a_{v_n}^k(\tilde{x},t) \\ & \quad \left. \times \langle \bar{k}_{v_n}(\tilde{x},t) | \bar{k}_{u_n}(x,t) \rangle \langle k_{v_n}(\tilde{x},t) | k_{u_1}(x,t) \rangle \right]. \end{aligned} \quad (30)$$

Under the same spirit, we expect that

$$\lim_{t \rightarrow \infty} \Delta S^{(n)}(\mathcal{T}_A^{\mathcal{O}_k(x)|\mathcal{O}_k(\tilde{x})}(t)) = \lim_{t \rightarrow \infty} S^{(n)}(\mathcal{T}_V^{\mathcal{O}_k(x)|\mathcal{O}_k(\tilde{x})}(t)). \quad (31)$$

We have known that the late-time limit of $\Delta S^{(n)}(\mathcal{T}_A^{\mathcal{O}_k(x)|\mathcal{O}_k(\tilde{x})}(t))$ is equal to the late-time limit of $\Delta S^{(n)}(\rho_A^{\mathcal{O}_k}(t))$ [43], and the latter we already know is equal to (22).

Comparing eqn. (22) with eqn. (30), we obtain the equality

$$\begin{aligned} & \lim_{t \rightarrow \infty} \sum_{\{u\}, \{v\}} a_{u_1}^k(x,t) a_{v_1}^k(\tilde{x},t) \langle \bar{k}_{v_1}(\tilde{x},t) | \bar{k}_{u_1}(x,t) \rangle \langle k_{v_1}(\tilde{x},t) | k_{u_2}(x,t) \rangle \dots a_{u_n}^k(x,t) a_{v_n}^k(\tilde{x},t) \\ & \quad \times \langle \bar{k}_{v_n}(\tilde{x},t) | \bar{k}_{u_n}(x,t) \rangle \langle k_{v_n}(\tilde{x},t) | k_{u_1}(x,t) \rangle \\ &= \langle \mathcal{O}_k(\tilde{x}, t=0) | \mathcal{O}_k(x, t=0) \rangle^n d_k^{1-n} \end{aligned} \quad (32)$$

Substituting eqn. (32) into eqn. (29) and taking $t \rightarrow \infty$, we finally obtain the late-time PREE formula

$$\begin{aligned} \lim_{t \rightarrow \infty} \Delta S^{(n)}(\mathcal{T}_A^{\psi|\tilde{\psi}}(t)) &= \lim_{t \rightarrow \infty} \frac{1}{1-n} \log \text{tr} \left[(\mathcal{T}_V^{\psi|\tilde{\psi}}(t))^n \right] \\ &= \frac{1}{1-n} \log \left[\sum_{k \in \mathcal{S}} \left(\frac{d_k^{\frac{1}{n}-1} |C_k|^2 L^{4h_k} \langle \mathcal{O}_k^\dagger(\tilde{x}, \epsilon) | \mathcal{O}_k(x, -\epsilon) \rangle}{\sum_{j \in \mathcal{S}} |C_j|^2 L^{4h_j} \langle \mathcal{O}_j^\dagger(\tilde{x}, \epsilon) | \mathcal{O}_j(x, -\epsilon) \rangle} \right)^n \right]. \end{aligned} \quad (33)$$

The accuracy of the above formula is verified by comparing it with the results obtained through numerical replica calculations in [43]. The next section will employ the replica method and conformal mapping in 2D CFT to reproduce eqn. (33) for $n = 2$ and $n = 3$.

2.1.2 Replica method

Let us begin with the Euclidean path integral formulation of the replica calculation of the n -th PREE. Suppose that the theory of interest (with a Lagrangian $\mathcal{L}(\phi, \partial\phi)$) resides on a Euclidean plane Σ_1 with the metric $ds^2 = dwd\bar{w}$ ($(w, \bar{w}) = (x + i\tau, x - i\tau)$). The Euclidean transition matrix of interest is generated by the linear combination operator $\mathcal{O}(w, \bar{w}) \equiv e^{\tau H} \mathcal{O}(x) e^{-\epsilon H}$ (9)

$$\mathcal{T}_E^{\psi|\bar{\psi}} = \frac{\mathcal{O}(w_1, \bar{w}_1)|\Omega\rangle\langle\Omega|\mathcal{O}^\dagger(w_2, \bar{w}_2)}{\langle\Omega|\mathcal{O}^\dagger(w_2, \bar{w}_2)\mathcal{O}(w_1, \bar{w}_1)|\Omega\rangle}, \quad (34)$$

where $w_1 \equiv x - i\tau_1$, $w_2 \equiv \tilde{x} + i\tau_2$ ($\tau_{1,2} > 0$). The reduced transition matrix of subsystem A can be expressed by path integral with operators inserted at (w_1, \bar{w}_1) and (w_2, \bar{w}_2) on the w -plane with a cut on A

$$\begin{aligned} \langle\phi_{A-}|\mathcal{T}_{E,A}^{\psi|\bar{\psi}}|\phi_{A+}\rangle &= \frac{\int_{\phi(x \in A, \tau=0_+) = \phi_{A+}(x)}^{\phi(x \in A, \tau=0_-) = \phi_{A-}(x)} [d\phi] \mathcal{O}^\dagger(w_2, \bar{w}_2) \mathcal{O}(w_1, \bar{w}_1) \exp\left\{-\int_{\mathbb{R}^2} \mathcal{L}(\phi, \partial\phi)\right\}}{\int [d\phi] \mathcal{O}^\dagger(w_2, \bar{w}_2) \mathcal{O}(w_1, \bar{w}_1) \exp\left\{-\int_{\mathbb{R}^2} \mathcal{L}(\phi, \partial\phi)\right\}} \\ &= \left(\begin{array}{c} \text{---}^{+\infty} \\ \star \\ \mathcal{O}^\dagger(w_2, \bar{w}_2) \\ \text{---} \\ \mathcal{O}(w_1, \bar{w}_1) \\ \text{---}^{-\infty} \end{array} \right)^{-1} \times \left(\begin{array}{c} \text{---}^{+\infty} \\ \star \\ \mathcal{O}^\dagger(w_2, \bar{w}_2) \\ \text{---} \\ \mathcal{O}(w_1, \bar{w}_1) \\ \text{---}^{-\infty} \end{array} \right). \end{aligned} \quad (35)$$

Based on (35), $\text{tr}[(\mathcal{T}_{E,A}^{\psi|\bar{\psi}})^n]$ amounts to a $2n$ -point correlation function on a n -sheet Riemann surface Σ_n ,

$$\begin{aligned} \text{tr}[(\mathcal{T}_{E,A}^{\psi|\bar{\psi}})^n] &= \left(\begin{array}{c} \text{---}^{+\infty} \\ \star \\ \mathcal{O}^\dagger(w_2, \bar{w}_2) \\ \text{---} \\ \mathcal{O}(w_1, \bar{w}_1) \\ \text{---}^{-\infty} \end{array} \right)^{-n} \times \left(\begin{array}{c} \text{---}^{+\infty} \\ \star \\ \mathcal{O}^\dagger(w_2, \bar{w}_2) \\ \text{---} \\ \mathcal{O}(w_1, \bar{w}_1) \\ \text{---}^{-\infty} \end{array} \right)^n \\ &= \frac{Z_n}{Z_1^n} \cdot \frac{\langle\mathcal{O}(w_1, \bar{w}_1)\mathcal{O}^\dagger(w_2, \bar{w}_2)\dots\mathcal{O}(w_{2n-1}, \bar{w}_{2n-1})\mathcal{O}^\dagger(w_{2n}, \bar{w}_{2n})\rangle_{\Sigma_n}}{\langle\mathcal{O}^\dagger(w_2, \bar{w}_2)\mathcal{O}(w_1, \bar{w}_1)\rangle_{\Sigma_1}^n}, \end{aligned} \quad (36)$$

where Z_1 and Z_n are vacuum partition functions on Σ_1 and Σ_n respectively, and $\mathcal{O}(w_{2k-1}, \bar{w}_{2k-1})$ and $\mathcal{O}^\dagger(w_{2k}, \bar{w}_{2k})$ are operators inserted at the k -th sheet. The variation of the n -th PREE is defined as subtracting the contribution of the n -th REE of the vacuum from the n -th PREE,

$$\begin{aligned} \Delta S^{(n)}(\mathcal{T}_{E,A}^{\psi|\bar{\psi}}) &\equiv S^{(n)}(\mathcal{T}_{E,A}^{\psi|\bar{\psi}}) - S^{(n)}(\rho_{A, \text{vacuum state}}) \\ &= \frac{1}{1-n} \log \frac{\langle\mathcal{O}(w_1, \bar{w}_1)\mathcal{O}^\dagger(w_2, \bar{w}_2)\dots\mathcal{O}(w_{2n-1}, \bar{w}_{2n-1})\mathcal{O}^\dagger(w_{2n}, \bar{w}_{2n})\rangle_{\Sigma_n}}{\langle\mathcal{O}(w_1, \bar{w}_1)\mathcal{O}^\dagger(w_2, \bar{w}_2)\rangle_{\Sigma_1}^n}. \end{aligned} \quad (37)$$

The real-time evolution of the n -th PREE is achieved by performing a Wick rotation of the Euclidean time: $\tau_1 \rightarrow \epsilon + it$, $\tau_2 \rightarrow \epsilon - it$. For the case of $n = 2$, $\Delta S^{(2)}(\mathcal{T}_{E,A}^{\psi|\bar{\psi}})$ is linked to a series of four-point functions on Σ_2

$$e^{-\Delta S^{(2)}(\mathcal{T}_{E,A}^{\psi|\bar{\psi}})} = \frac{\sum_{ijkl \in \mathcal{S}} C_i C_j^* C_k C_l^* L^{2(h_i + h_j + h_k + h_l)} \langle\mathcal{O}_i(w_1, \bar{w}_1)\mathcal{O}_j^\dagger(w_2, \bar{w}_2)\mathcal{O}_k(w_3, \bar{w}_3)\mathcal{O}_l^\dagger(w_4, \bar{w}_4)\rangle_{\Sigma_2}}{\left(\sum_{j \in \mathcal{S}} |C_j|^2 L^{4h_j} \langle\mathcal{O}_j(w_1, \bar{w}_1)\mathcal{O}_j^\dagger(w_2, \bar{w}_2)\rangle_{\Sigma_1}\right)^2}. \quad (38)$$

These four-point functions can be evaluated by applying the conformal mapping $z = i\sqrt{-w}$ [7,8] from Σ_2 to Σ_1 . The four operators after the mapping are located at

$$\begin{aligned} z_1 = -z_3 = i\sqrt{-x-t+i\epsilon}, \quad \bar{z}_1 = -\bar{z}_3 = -i\sqrt{-x+t-i\epsilon}, \\ z_2 = -z_4 = i\sqrt{-\tilde{x}-t-i\epsilon}, \quad \bar{z}_2 = -\bar{z}_4 = -i\sqrt{-\tilde{x}+t+i\epsilon}. \end{aligned} \quad (39)$$

Let's take $i = 1, j = 2, k = 3,$ and $l = 4$ for the sake of writing convenience, the four-point function can be expressed as

$$\begin{aligned} & \langle \mathcal{O}_1(w_1, \bar{w}_1) \mathcal{O}_2^\dagger(w_2, \bar{w}_2) \mathcal{O}_3(w_3, \bar{w}_3) \mathcal{O}_4^\dagger(w_4, \bar{w}_4) \rangle_{\Sigma_2} \\ &= \prod_{i=1}^4 |2z_i|^{-2h_i} \langle \mathcal{O}_1(z_1, \bar{z}_1) \mathcal{O}_2^\dagger(z_2, \bar{z}_2) \mathcal{O}_3(z_3, \bar{z}_3) \mathcal{O}_4^\dagger(z_4, \bar{z}_4) \rangle_{\Sigma_1} \\ &= \left(\left| \frac{2z_1 z_{12} z_{14}}{z_{24}} \right|^{-2h_1} \prod_{i=2}^4 \left| \frac{2z_i z_{1i}^2 z_{24}}{z_{12} z_{14}} \right|^{-2h_i} \right) G_{34}^{21}(\eta, \bar{\eta}), \end{aligned} \quad (40)$$

where $G_{34}^{21}(\eta, \bar{\eta}) \equiv \lim_{z \rightarrow \infty} z^{2h_1} \bar{z}^{2h_1} \langle \mathcal{O}_1(z, \bar{z}) \mathcal{O}_2^\dagger(1, 1) \mathcal{O}_3(\eta, \bar{\eta}) \mathcal{O}_4^\dagger(0, 0) \rangle_{\Sigma_1}$, and $(\eta, \bar{\eta}) \equiv \left(\frac{z_{12} z_{34}}{z_{13} z_{24}}, \frac{\bar{z}_{12} \bar{z}_{34}}{\bar{z}_{13} \bar{z}_{24}} \right)$. Let us begin by coping with the G -function in terms of the conformal blocks. In general CFTs, $G_{34}^{21}(\eta, \bar{\eta})$ can be written as follows [1]

$$G_{34}^{21}(\eta, \bar{\eta}) = \sum_p C_{34}^p C_{12}^p F_{34}^{21}(p|\eta) \bar{F}_{34}^{21}(p|\bar{\eta}). \quad (41)$$

Here, C_{34}^p is the coefficient of the three-point function $\langle \mathcal{O}_p \mathcal{O}_3 \mathcal{O}_4^\dagger \rangle$, and the index p corresponds to each \mathcal{O}_p of all Virasoro primary fields. It is important to note that the conformal block has a universal behavior around $\eta = 0$,

$$F_{34}^{21}(p|\eta) = \eta^{h_p - h_3 - h_4} (1 + O(\eta)). \quad (42)$$

As shown in [43], the cross ratios $(\eta, \bar{\eta}) \simeq \left(1 + \frac{(\tilde{x}-x+2i\epsilon)^2}{16t^2}, -\frac{(\tilde{x}-x-2i\epsilon)^2}{16t^2} \right) \sim (1, 0)$ in the late-time limit. Then the fusion transformation in RCFT [74, 75]

$$F_{34}^{21}(p|1-\eta) = \sum_q F_{pq} \left[\begin{matrix} 2, 1 \\ 3, 4 \end{matrix} \right] F_{32}^{41}(q|\eta) \quad (43)$$

can be leveraged to fix the leading behavior of the conformal block in $\eta \rightarrow 1$ limit and subsequently fix the leading order of the late-time behavior of $G_{34}^{21}(\eta, \bar{\eta})$

$$\begin{aligned} \lim_{t \rightarrow \infty} G_{34}^{21}(\eta, \bar{\eta}) &\simeq \sum_{p,q} C_{12}^p C_{34}^p F_{pq} \left[\begin{matrix} 2, 1 \\ 3, 4 \end{matrix} \right] (1-\eta)^{h_q - h_2 - h_3} \bar{\eta}^{h_p - h_3 - h_4} \\ &\sim \sum_{p,q} C_{12}^p C_{34}^p F_{pq} \left[\begin{matrix} 2, 1 \\ 3, 4 \end{matrix} \right] t^{2(h_2 + 2h_3 + h_4 - h_p - h_q)}. \end{aligned} \quad (44)$$

On the other hand, we can determine the leading behavior of the prefactor for G_{34}^{21} at late times by simply applying the expressions of (z_i, \bar{z}_i) from eqn. (39). It is found that

$$\lim_{t \rightarrow \infty} \left(\left| \frac{2z_1 z_{12} z_{14}}{z_{24}} \right|^{-2h_1} \prod_{i=2}^4 \left| \frac{2z_i z_{1i}^2 z_{24}}{z_{12} z_{14}} \right|^{-2h_i} \right) \sim t^{-2(h_2 + 2h_3 + h_4)}. \quad (45)$$

Combining eqn. (44) with eqn. (45) yields a remarkable outcome: the four-point function $\langle \mathcal{O}_1 \mathcal{O}_2^\dagger \mathcal{O}_3 \mathcal{O}_4^\dagger \rangle$ does not vanish at late times if, and only if, a vacuum sector is present in the fusion of any two operators

$$\begin{aligned} \mathcal{O}_1 \times \mathcal{O}_2 = \mathbb{I} + \dots, \quad \mathcal{O}_3 \times \mathcal{O}_4 = \mathbb{I} + \dots, \\ \mathcal{O}_1 \times \mathcal{O}_4 = \mathbb{I} + \dots, \quad \mathcal{O}_2 \times \mathcal{O}_3 = \mathbb{I} + \dots \end{aligned} \quad (46)$$

The only scenario that aligns with eqn. (46) is when $\mathcal{O}_1 = \mathcal{O}_2 = \mathcal{O}_3 = \mathcal{O}_4$. Therefore, the second PREE (38) at late times is reduced to

$$\begin{aligned} \lim_{t \rightarrow \infty} e^{-\Delta S^{(2)}(\mathcal{T}_A^{\psi|\bar{\psi}}(t))} &= \frac{\sum_{k \in \mathcal{S}} |C_k|^4 L^{8h_k} \langle \mathcal{O}_k(w_1, \bar{w}_1) \mathcal{O}_k^\dagger(w_2, \bar{w}_2) \mathcal{O}_k(w_3, \bar{w}_3) \mathcal{O}_k^\dagger(w_4, \bar{w}_4) \rangle_{\Sigma_2}}{\left(\sum_{j \in \mathcal{S}} |C_j|^2 L^{4h_j} \langle \mathcal{O}_j(w_1, \bar{w}_1) \mathcal{O}_j^\dagger(w_2, \bar{w}_2) \rangle_{\Sigma_1} \right)^2} \\ &= \frac{\sum_{k \in \mathcal{S}} d_k^{-1} |C_k|^4 L^{8h_k} \langle \mathcal{O}_k(w_1, \bar{w}_1) \mathcal{O}_k^\dagger(w_2, \bar{w}_2) \rangle_{\Sigma_1}^2}{\left(\sum_{j \in \mathcal{S}} |C_j|^2 L^{4h_j} \langle \mathcal{O}_j(w_1, \bar{w}_1) \mathcal{O}_j^\dagger(w_2, \bar{w}_2) \rangle_{\Sigma_1} \right)^2}, \end{aligned} \quad (47)$$

where from the first line to the second line, we utilize the known results that the four-point function of \mathcal{O}_k on Σ_2 at late times splits into two two-point functions on Σ_1 multiplied by the inverse of the quantum dimension of \mathcal{O}_k [43]. Note eqn. (47) perfectly matches eqn. (33) for $n = 2$.

The above calculation suggests that for general n , a $2n$ -point function on Σ_n does not vanish at late times if and only if all these $2n$ operators are the same. Indeed, in the replica calculation of n -th PREE, one can readily obtain (33) by simply considering $2n$ -point functions involving only one primary. To show the remaining $2n$ -point functions do vanish at late times, we begin with the $n = 3$ case and evaluate the 6-point functions on Σ_3 .¹⁰ According to the conformal map between Σ_k and Σ_1 , a general 6-point correlation function on Σ_3 can be written as

$$\begin{aligned} &\langle \mathcal{O}_1(w_1, \bar{w}_1) \mathcal{O}_2^\dagger(w_2, \bar{w}_2) \mathcal{O}_3(w_3, \bar{w}_3) \mathcal{O}_4^\dagger(w_4, \bar{w}_4) \mathcal{O}(w_5, \bar{w}_5) \mathcal{O}_6^\dagger(w_6, \bar{w}_6) \rangle_{\Sigma_3} \\ &= \prod_{i=1}^6 |3z_i^2|^{-2h_i} \langle \mathcal{O}_1(z_1, \bar{z}_1) \mathcal{O}_2^\dagger(z_2, \bar{z}_2) \mathcal{O}_3(z_3, \bar{z}_3) \mathcal{O}_4^\dagger(z_4, \bar{z}_4) \mathcal{O}_5(z_5, \bar{z}_5) \mathcal{O}_6^\dagger(z_6, \bar{z}_6) \rangle_{\Sigma_1}. \end{aligned} \quad (48)$$

To evaluate the 6-point function on Σ_1 , we insert a complete basis into this function

$$\begin{aligned} &\langle \mathcal{O}_1(z_1, \bar{z}_1) \mathcal{O}_2^\dagger(z_2, \bar{z}_2) \mathcal{O}_3(z_3, \bar{z}_3) \mathcal{O}_4^\dagger(z_4, \bar{z}_4) \mathcal{O}_5(z_5, \bar{z}_5) \mathcal{O}_6^\dagger(z_6, \bar{z}_6) \rangle_{\Sigma_1} \\ &= \sum_p \lim_{z_0, \bar{z}_0 \rightarrow \infty} z_0^{2h_p} \bar{z}_0^{2\bar{h}_p} \\ &\quad \times \langle \mathcal{O}_1(z_1, \bar{z}_1) \mathcal{O}_2^\dagger(z_2, \bar{z}_2) \mathcal{O}_3(z_3, \bar{z}_3) \mathcal{O}_p^\dagger(0, 0) \rangle_{\Sigma_1} \langle \mathcal{O}_p(z_0, \bar{z}_0) \mathcal{O}_4^\dagger(z_4, \bar{z}_4) \mathcal{O}_5(z_5, \bar{z}_5) \mathcal{O}_6^\dagger(z_6, \bar{z}_6) \rangle_{\Sigma_1} \end{aligned} \quad (49)$$

By using Mobius transformation, the two 4-point functions in (49) can be rewritten as

$$\begin{aligned} &\langle \mathcal{O}_1(z_1, \bar{z}_1) \mathcal{O}_2^\dagger(z_2, \bar{z}_2) \mathcal{O}_3(z_3, \bar{z}_3) \mathcal{O}_p^\dagger(0, 0) \rangle_{\Sigma_1} \langle \mathcal{O}_p(z_0, \bar{z}_0) \mathcal{O}_4^\dagger(z_4, \bar{z}_4) \mathcal{O}_5(z_5, \bar{z}_5) \mathcal{O}_6^\dagger(z_6, \bar{z}_6) \rangle_{\Sigma_1} \\ &= \left| \frac{z_{12} z_1}{z_2} \right|^{-2h_1} \left| \frac{z_{12} z_1}{z_{12}^2 z_2} \right|^{2h_2} \left| \frac{z_{12} z_1}{z_{13}^2 z_2} \right|^{2h_3} \left| \frac{z_{12} z_1}{z_1^2 z_2} \right|^{2h_p} G_{3p}^{21}(\eta, \bar{\eta}) \\ &\quad \times \left| \frac{z_{04} z_{06}}{z_{46}} \right|^{-2h_p} \left| \frac{z_{04} z_{06}}{z_{04}^2 z_{46}} \right|^{2h_4} \left| \frac{z_{04} z_{06}}{z_{05}^2 z_{46}} \right|^{2h_5} \left| \frac{z_{04} z_{06}}{z_{06}^2 z_{46}} \right|^{2h_6} G_{56}^{4p}(\eta', \bar{\eta}'). \end{aligned} \quad (50)$$

with $\eta = \frac{z_{12} z_3}{z_{13} z_2}$ and $\eta' = \frac{z_{04} z_{56}}{z_{05} z_{46}}$ whose late-time behavior are $(\eta, \eta') \simeq (1 + \frac{x - \bar{x} + 2i\epsilon}{t}, 1 + \frac{x - \bar{x} + 2i\epsilon}{t}) \sim (1, 1)$.

Combining (48), (49) and (50), the late-time behavior of the holomorphic part of the 6-point function on Σ_3 is

$$\begin{aligned} &\lim_{t \rightarrow \infty} \langle \mathcal{O}_1(w_1) \mathcal{O}_2^\dagger(w_2) \mathcal{O}_3(w_3) \mathcal{O}_4^\dagger(w_4) \mathcal{O}(w_5) \mathcal{O}_6^\dagger(w_6) \rangle_{\Sigma_3} \\ &\sim \sum_{l_1, l_2, p} \left(\prod_{i=1}^6 t^{-\frac{2}{3} h_i} \right) \left(t^{-\frac{1}{3} h_1} t^{-\frac{1}{3} h_2} t^{-\frac{1}{3} h_3} t^{-\frac{1}{3} h_p} \right) t^{h_2 + h_3 - h_{l_1}} \left(t^{\frac{1}{3} h_p} t^{-\frac{1}{3} h_4} t^{-\frac{1}{3} h_5} t^{-\frac{1}{3} h_6} \right) t^{h_4 + h_5 - h_{l_2}} t^{h_1 + h_6 - h_p} \end{aligned} \quad (51)$$

¹⁰For the discussions of the higher point functions, it is more convenient to adopt the coordinate convention in [8].

where h_{l_1} and h_{l_2} are contributed from the fusion

$$\mathcal{O}_2 \times \mathcal{O}_3 = \mathcal{O}_{l_1} + \dots, \quad \mathcal{O}_4 \times \mathcal{O}_5 = \mathcal{O}_{l_2} + \dots \quad (52)$$

as shown in Fig. 1.




Figure 1: Fusion transformations to obtain $\lim_{t \rightarrow \infty} \langle \mathcal{O}_1(w_1) \mathcal{O}_2^\dagger(w_2) \mathcal{O}_3(w_3) \mathcal{O}_4^\dagger(w_4) \mathcal{O}(w_5) \mathcal{O}_6^\dagger(w_6) \rangle_{\Sigma_3}$

Using the result, we derive for the 4-point function (46), the 6-point function on Σ_3 does not vanish at late times if, and only if

$$h_{l_1} = h_{l_2} = h_p = 0 \quad (53)$$

yielding the fusion rules

$$\mathcal{O}_2 \times \mathcal{O}_3 = \mathbb{I} + \dots, \quad \mathcal{O}_4 \times \mathcal{O}_5 = \mathbb{I} + \dots, \quad \mathcal{O}_1 \times \mathcal{O}_6 = \mathbb{I} + \dots \quad (54)$$

Similarly, the late-time behavior of the anti-holomorphic part of the 6-point function yields the fusion rules

$$\mathcal{O}_1 \times \mathcal{O}_2 = \mathbb{I} + \dots, \quad \mathcal{O}_3 \times \mathcal{O}_4 = \mathbb{I} + \dots, \quad \mathcal{O}_5 \times \mathcal{O}_6 = \mathbb{I} + \dots, \quad (55)$$

which means that the 6-point function on Σ_3 does not vanish at late times only when

$$\mathcal{O}_1 = \mathcal{O}_2 = \mathcal{O}_3 = \mathcal{O}_4 = \mathcal{O}_5 = \mathcal{O}_6. \quad (56)$$

All the discussion of the 6-point function on Σ_3 can be extended to a general $2n$ -point function on Σ_n by inserting a complete basis and using the conformal block, and the results are similar, implying that the $2n$ -point function does not vanish if and only if all the operators in the $2n$ -point function belong to the same conformal family.

2.2 Pseudo entropy amplification

In this subsection, we introduce the phenomenon of pseudo entropy amplification, where PE (PREE) contains all the information about the linear combination operator whereas EE (REE) does not, through the evaluation of the maximal value of (33) by adjusting the combination coefficients C_k .

2.2.1 The REE and EE

Let us begin by focusing on the REE and EE. By simply removing the tildes in (33), we arrive at the following result

$$\begin{aligned}
\lim_{t \rightarrow \infty} \Delta S^{(n)}(\rho_A^\psi(t)) &= \lim_{t \rightarrow \infty} \Delta S^{(n)}(\mathcal{T}_A^{\psi|\psi}(t)) \\
&= \frac{1}{1-n} \log \frac{\sum_{k \in \mathcal{S}} d_k^{1-n} |C_k|^{2n} (\frac{L}{2\epsilon})^{4nh_k}}{(\sum_{j \in \mathcal{S}} |C_j|^2 (\frac{L}{2\epsilon})^{4h_j})^n} \\
&= \frac{1}{1-n} \log \frac{\sum_{k \in \mathcal{S}_M} d_k^{1-n} |C_k|^{2n}}{(\sum_{j \in \mathcal{S}_M} |C_j|^2)^n}, \tag{57}
\end{aligned}$$

where $\mathcal{S}_M := \{\mathcal{O}_k | \mathcal{O}_k \in \mathcal{S}, h_k = h_M \equiv \max_{l \in \mathcal{S}} \{h_l\}\}$. From the second equation to the third equation, we omit terms associated with lighter operators since, in the limit as $\epsilon \rightarrow 0$, they constitute sub-leading contributions within the sum. Consequently, we conclude that information about lighter operators is lost in the REE when \mathcal{S}_M is a proper subset of \mathcal{S} . This conclusion holds, especially in cases where a single heaviest primary operator with a quantum dimension of d_M exists, simplifying eqn. (57) to $\log d_M$. For cases where the conformal dimensions of primary operators in (9) are the same¹¹, the late-time behavior of n -th REE for ρ_A^ψ is

$$\lim_{t \rightarrow \infty} \Delta S^{(n)}(\rho_A^\psi(t)) = \frac{1}{1-n} \log \left(\sum_{k \in \mathcal{S}} d_k^{1-n} |C_k|^{2n} \right), \tag{58}$$

where the constraint $\sum_{k \in \mathcal{S}} |C_k|^2 = 1$ was employed to simplify the expression.

Taking the limit of $n \rightarrow 1$ to eqn. (57), we obtain the expression of the late-time EE,

$$\lim_{t \rightarrow \infty} \Delta S(\rho_A^\psi(t)) = \sum_{k \in \mathcal{S}_M} p_k \log d_k + H(p_k), \tag{59}$$

where $p_k = \frac{|C_k|^2}{\sum_{j \in \mathcal{S}_M} |C_j|^2}$ is an effective probability distribution, and $H(p_k) = -\sum_{k \in \mathcal{S}_M} p_k \log p_k$ the classical Shannon entropy of $\{p_k\}$.

As functions of the combination coefficients C_k , the maximum values of (57) and (59) can be determined using *Hölder's inequality*, a generalization of *Cauchy-Schwarz inequality*,

$$\begin{aligned}
\frac{1}{n-1} \log \frac{\left(\sum_{k \in \mathcal{S}_M} d_k^{1-\frac{1}{n}} (y_k^n)^{\frac{1}{n}} \right)^n}{\sum_{j \in \mathcal{S}_M} y_j^n} &\leq \frac{1}{n-1} \log \frac{\left((\sum_{k \in \mathcal{S}_M} d_k)^{\frac{n-1}{n}} (\sum_{k \in \mathcal{S}_M} y_k^n)^{\frac{1}{n}} \right)^n}{\sum_{j=1}^m y_j^n} \\
&= \log \left(\sum_{k \in \mathcal{S}_M} d_k \right). \tag{60}
\end{aligned}$$

Here, we redefine a series of parameters in terms of C_k

$$y_k := |C_k|^2 d_k^{\frac{1}{n}-1} > 0, \quad (k = 1, 2, \dots, m). \tag{61}$$

The equality in (60) holds if and only if

$$y_k^n \propto d_k \Rightarrow |C_k| \propto \sqrt{d_k}. \tag{62}$$

¹¹We continue to assume that the primary operators are orthogonal to each other for this case.

Thus, we conclude that the maximum late-time REE and EE saturates to the logarithm of the sum of quantum dimensions of the heaviest operators,

$$\max_{\{C_k\}} \left\{ \lim_{t \rightarrow \infty} \Delta S^{(n)}(\rho_A^\psi(t)) \right\} = \max_{\{C_k\}} \left\{ \lim_{t \rightarrow \infty} \Delta S(\rho_A^\psi(t)) \right\} = \log \left(\sum_{j \in \mathcal{S}_M} d_j \right). \quad (63)$$

The corresponding combination coefficients, denoted as C_k^* , are given by

$$|C_k^*| = \sqrt{\frac{d_k}{\sum_{j \in \mathcal{S}} d_j}}. \quad (64)$$

Note that there remains an undetermined degree of phase freedom for each coefficient C_k since the late-time formula (33) depends only on the magnitude of C_k .

We can examine our conclusions in specific RCFT models. One typical example in RCFT is the linear combination of vertex operators in 2D massless free scalar theory.¹² As demonstrated in [9,33], the variation of the second REE of the one-parameter primary operator $V_q = \sqrt{q}e^{i\alpha\phi} \pm \sqrt{1-q}e^{-i\alpha\phi}$ is maximized when the parameter q equals $\frac{1}{2}$,

$$\max_q \{ \Delta S_A^{(2)}(V_q) \} = \Delta S_A^{(2)}(V_{\frac{1}{2}}) = \log 2 = \log(1+1), \quad (65)$$

where 1 represents the quantum dimension of the vertex operators $e^{\pm i\alpha\phi}$ and $q = \frac{1}{2}$ maximizing $\Delta S_A^{(2)}(V_a)$ precisely aligns with the obtained result. A more nontrivial verification can be made in the critical Ising model. Our results indicate that the information of lighter operators in the critical Ising model will be lost in REE/EE in the limit of $\epsilon \rightarrow 0$, as the conformal dimensions of the three primary operators (i.e., the Ising spin σ , energy density ε , and identity \mathbb{I}) in the critical Ising model differ from each other.

We numerically compute the late-time second REE for two linear combination operators: $C_\sigma \cdot L^{2h_\sigma} \sigma + C_\varepsilon \cdot L^{2h_\varepsilon} \varepsilon$ and $C_\sigma \cdot L^{2h_\sigma} \sigma + C_{\mathbb{I}} \cdot \mathbb{I}$ (See Fig. 2(a) and (b) respectively). Our late-time formula (33) (solid lines) exhibits good agreement with the numerical data (hollow circles). For the linear combination of the Ising spin and the energy density (Fig. 2(a)), the late-time second REE approaches zero in the $\epsilon \rightarrow 0$ limit, as the heavier operator ε possesses a quantum dimension of 1. Conversely, for the linear combination of the Ising spin and the identity (Fig. 2(b)), the late-time second REE approaches $\log \sqrt{2}$ in the $\epsilon \rightarrow 0$ limit since the heavier operator σ has a quantum dimension of $\sqrt{2}$.

2.2.2 The PREE and PE

After exploring the late-time behavior and the maximum values of REE and EE in the previous subsection, we now focus on studying PREE and PE. This part demonstrates how the PREE and PE can preserve information about all lighter operators in \mathcal{S} . This preservation is achieved by simply shifting the excitation position of \mathcal{O} from x to \tilde{x} to create the final state $|\tilde{\psi}\rangle$ (14).

The finiteness of the difference between x and \tilde{x} prevents the divergence of two-point functions in eqn. (33) and leaves a new expression for the late-time PREE in the $\epsilon \rightarrow 0$ limit,

$$\lim_{t \rightarrow \infty} \Delta S^{(n)}(\mathcal{T}_A^{\psi|\tilde{\psi}}(t)) = \frac{1}{1-n} \log \left[\sum_{k \in \mathcal{S}} \left(\frac{d_k^{\frac{1}{n}-1} |C_k|^2 \left(\frac{L}{|x-\tilde{x}|} \right)^{4h_k}}{\sum_{j \in \mathcal{S}} |C_j|^2 \left(\frac{L}{|x-\tilde{x}|} \right)^{4h_j}} \right)^n \right]. \quad (66)$$

Since the summation is performed on all operators in \mathcal{S} , information about lighter operators will be encoded in the PREE. Compared with the results of EE, we observe the phenomenon of

¹²We express our gratitude to Tadashi Takayanagi for bringing this to our attention.




Figure 2: The variation of the late-time second REE (in panels (a) and (b)) and the late-time second PREE (in panels (c) and (d)) with respect to the combination coefficient C_σ in the critical Ising model. The solid lines are plotted using the late-time formula (33), while the hollow circles denote numerical data computed using the well-known four-point functions in the critical Ising model. The vertical dotted lines in panels (c) and (d), plotted in terms of eqn. (68), correspond to the maximum values of the second PREE. All four panels are symmetric along $C_\sigma = 0$ since the late-time formula (33) depends solely on the magnitude of the combination coefficients. Note that the UV regulator ϵ , while a sensitive parameter in REE calculations, does not impact PREE computations and can be safely set to 0.

pseudo entropy amplification [53]. It can be shown more explicitly by looking at the maximum value of the n -th PREE. Let us define a series of parameters y_k , like before,

$$y_k := d_k^{\frac{1}{n}-1} |C_k|^2 \left(\frac{L}{x - \tilde{x}} \right)^{4h_k}. \quad (67)$$

The n -th PREE (66) is then transformed into the same expression in (60). The Hölder's inequality guarantees that (66) reaches its maximum value of $\log \left(\sum_{k \in \mathcal{S}} d_k \right)$ when the combination coefficients are given by

$$|C_k^*| = \sqrt{\frac{\frac{d_k}{L^{4h_k} \langle \mathcal{O}_k^\dagger(\tilde{x}) \mathcal{O}_k(x) \rangle}}{\sum_{j \in \mathcal{S}} \frac{d_j}{L^{4h_j} \langle \mathcal{O}_j^\dagger(\tilde{x}) \mathcal{O}_j(x) \rangle}}}. \quad (68)$$

The maximum value of the late-time PREE encompasses the quantum dimension of each primary operator within the set \mathcal{S} . This outcome contrasts with the result of REE (63), thereby explicitly reflecting the pseudo entropy amplification phenomenon. On the other hand, the late-time PE of $\mathcal{T}_A^{\psi|\tilde{\psi}}(t)$, obtained by taking the limit as $n \rightarrow 1$ for (66), is given by

$$\lim_{t \rightarrow \infty} \Delta S(\mathcal{T}_A^{\psi|\tilde{\psi}}(t)) = H(p_k(x, \tilde{x})) + \sum_{k \in \mathcal{S}} p_k(x, \tilde{x}) \log d_k. \quad (69)$$

In the above, $p_k(x, \tilde{x}) := \frac{|C_k^2|L^{4h_k}\langle\mathcal{O}_k^\dagger(\tilde{x})\mathcal{O}_k(x)\rangle}{\sum_{j\in\mathcal{S}}|C_j^2|L^{4h_j}\langle\mathcal{O}_j^\dagger(\tilde{x})\mathcal{O}_j(x)\rangle}$ is a position-depended probability distribution.

Similarly to the REE, we can confirm the accuracy of our results using numerical computations in the critical Ising model. As illustrated in Fig. 2(c) and Fig. 2(d), we numerically calculate the second PREE for linear combination operators $\mathcal{O}(x)$ and $\mathcal{O}(\tilde{x})$, where \mathcal{O} represents a linear combination of the Ising spin and the energy density or the Ising spin and the identity. Our late-time formula (33) (solid lines) consistently matches the numerical data (hollow circles). Furthermore, the coefficients corresponding to the maximum of the late-time 2nd PREE are also accurately characterized by eqn. (68). By comparing the green line in Fig. 2(a) with the lines in Fig. 2(d), one can readily observe the phenomenon of pseudo entropy amplification.

2.3 A connection between Fusion numbers and OPE coefficients

The results in the above compellingly demonstrate the efficacy and simplicity of the Schmidt decomposition method in analyzing the late-time EE (PE) of linear combinations of operators, eliminating the need for computing intricate higher-point functions. This section will mainly utilize the Schmidt decomposition method to explore the relationship between OPE coefficients and fusion coefficients.

Starting from the OPE (8), one can write down an equality for the locally excited states,

$$e^{-(\epsilon+it)H}\mathcal{O}_1(x+1)\mathcal{O}_2(x)|\Omega\rangle = \sum_k \sum_{\{\alpha,\bar{\alpha}\}} C_{k12}\beta_{12}^{k\{\alpha\}}\bar{\beta}_{12}^{k\{\bar{\alpha}\}}e^{-(\epsilon+it)H}\mathcal{O}_k^{\{\alpha,\bar{\alpha}\}}(x)|\Omega\rangle. \quad (70)$$

As shown in [12], the EE of the state on the l.h.s. of (70) saturates to $\log(d_1d_2)$ at late times.¹³ To calculate the EE of the state on the r.h.s. of (70), building on the Schmidt decomposition method, we would like to recast the state into the following form

$$|\psi_{12}(t)\rangle = \frac{1}{\sqrt{\sum_k|\tilde{C}_{k12}(\epsilon)|^2}}\sum_k\tilde{C}_{k12}(\epsilon)\cdot|\psi_{k12}(t)\rangle, \quad (71)$$

with

$$|\psi_{k12}(t)\rangle := \frac{1}{\sqrt{\mathcal{N}_{k12}(\epsilon)}}\sum_{\{\alpha,\bar{\alpha}\}}\beta_{12}^{k\{\alpha\}}\bar{\beta}_{12}^{k\{\bar{\alpha}\}}e^{-(\epsilon+it)H}\mathcal{O}_k^{\{\alpha,\bar{\alpha}\}}(x)|\Omega\rangle, \quad (\langle\psi_{k12}(t)|\psi_{l12}(t)\rangle = \delta_{kl}), \quad (72)$$

$$\mathcal{N}_{k12}(\epsilon) := \sum_{\{\alpha,\bar{\alpha},\gamma,\bar{\gamma}\}}|2\epsilon|^{-(4h_k+|\alpha|+|\bar{\alpha}|+|\gamma|+|\bar{\gamma}|)}e^{\frac{i\pi}{2}(|\bar{\alpha}|+|\bar{\gamma}|-|\alpha|-|\gamma|)}\beta_{12}^{k\{\alpha\}}\beta_{12}^{k\{\gamma\}}\bar{\beta}_{12}^{k\{\bar{\alpha}\}}\bar{\beta}_{12}^{k\{\bar{\gamma}\}}c_k(\{\alpha,\bar{\alpha}\},\{\gamma,\bar{\gamma}\}), \quad (73)$$

and

$$\tilde{C}_{k12}(\epsilon) := C_{k12}\cdot\sqrt{\mathcal{N}_{k12}(\epsilon)}. \quad (74)$$

In the above, we denote by $c_k(\{\alpha,\bar{\alpha}\},\{\gamma,\bar{\gamma}\})$ the coefficient of the two-point function $\langle\mathcal{O}_k^{\{\alpha,\bar{\alpha}\}}(z,\bar{z})\mathcal{O}_k^{\{\gamma,\bar{\gamma}\}}(0,0)\rangle$ (e.g., $c_k(\{-1,0\},\{0,0\}) = -2h_k$). Since $|\psi_{k12}(t)\rangle$ are orthogonal to one another, we can apply the Schmidt decomposition method like before to derive the EE of $|\psi_{12}\rangle$ at late times, which is given by

$$\lim_{t\rightarrow\infty}\Delta S(\rho_A^{\psi_{12}}(t)) = H(p_k) + \sum_k p_k \cdot \lim_{t\rightarrow\infty}\Delta S(\rho_A^{\psi_{k12}}(t)), \quad (75)$$

¹³One might consider substituting $x+1$ with $x+2\epsilon$ to yield the second type of linear combination operators (18) on the r.h.s. of the OPE. However, it's essential to note that the EE of the composite operator on the l.h.s. of the OPE, after the substitution, does not converge to $\log(d_1d_2)$ as ϵ tends to zero and t tends to infinity [12].

where $p_k \equiv |\tilde{C}_{k12}|^2 / \sum_j |\tilde{C}_{j12}|^2$. Due to the non-orthogonality among distinct locally descendant excited states in eqn. (72), it's unattainable to obtain the EE of $\rho_A^{\psi_{k12}}$ via Schmidt decomposition method. Nevertheless, we can still compute it by employing the replica method. As demonstrated in [10, 43], for a general linear combination of descendant operators (within the same conformal family, say $[\mathcal{O}_k]$), there exists an additional correction to the subsystem's EE at late times apart from $\log d_k$ (for instance, $\lim_{t \rightarrow \infty} \Delta S_A(\partial \mathcal{O}_k + \bar{\partial} \mathcal{O}_k) = \log d_k + \log 2$). However, as shown in [12], due to the fact that the linear combination operator on the r.h.s. of the OPE need to match the composite operators on the l.h.s. under the same conformal mapping in RCFTs, this additional correction was found to be zero for the state $|\psi_{k12}\rangle$ (72). Consequently, we have

$$\lim_{t \rightarrow \infty} \Delta S(\rho_A^{\psi_{12}}(t)) = H(p_k) + \sum_k p_k \cdot \log d_k = \log \sum_k d_k, \quad (76)$$

where the second equality arises from the EE of the composite operator on the l.h.s. of the OPE and the fusion algebra (2). On the other hand, as per the conclusions from previous sections, the second equality ensures that p_k will maximize the expression in eqn. (76). We thus arrive at an equality between the quantum dimensions and OPE coefficients

$$p_k = \frac{d_k}{\sum_j d_j} = \frac{|\tilde{C}_{k12}|^2}{\sum_j |\tilde{C}_{j12}|^2} \Rightarrow \sqrt{\frac{d_k}{d_l}} = \frac{C_{k12}}{C_{l12}} \lim_{\epsilon \rightarrow 0} \sqrt{\frac{|\mathcal{N}_{k12}(\epsilon)|}{|\mathcal{N}_{l12}(\epsilon)|}}. \quad (77)$$

Notice that the r.h.s. of the above equality contains information about the operators \mathcal{O}_1 and \mathcal{O}_2 , while the l.h.s. does not. Hence the above equality is not complete. One may replace¹⁴ $d_{k(l)}$ on the l.h.s. of eqn. (77) with $N_{12}^{k(l)} d_{k(l)}$, yielding an equality relating fusion coefficients to OPE coefficients, to restore the information of \mathcal{O}_1 and \mathcal{O}_2 for the l.h.s. of eqn. (77),

$$\sqrt{\frac{N_{12}^k d_k}{N_{12}^l d_l}} = \frac{C_{k12}}{C_{l12}} \lim_{\epsilon \rightarrow 0} \sqrt{\frac{|\mathcal{N}_{k12}(\epsilon)|}{|\mathcal{N}_{l12}(\epsilon)|}}, \quad N_{12}^k \in \{0, 1\}. \quad (79)$$

While we have linked fusion coefficients and quantum dimensions to coefficients of OPE through EE, rigorously verifying the above two equations necessitates dealing with the infinite summation over secondary fields. Even in the simplest Ising model, this remains a rather challenging task. In this regard, we aim to seek relevant insights in future work.

3 The entanglement dynamics of refined linear combination operators (18)

In the previous section, we provided a connection between OPE coefficients and fusion coefficients by examining the EE of a linear combination of operators on the r.h.s. of the OPE. Such a linear combination operator is distinctive as it involves linear combinations of an infinite number of descendant operators. When we confine the number of operators to a finite set, as in

¹⁴Such a replacement may be understood in terms of the following way: we recast eqn. (76) into the following form to reveal the dependence of the result on the fusion coefficients,

$$\lim_{t \rightarrow \infty} \Delta S(\rho_A^{\psi_{12}}(t)) = H(p_k) + \sum_k p_k \cdot \log N_{12}^k d_k = \log \sum_k N_{12}^k d_k, \quad N_{12}^k \in \{0, 1\}. \quad (78)$$

Note that $p_k = 0$ if $N_{12}^k = 0$ and we define $0 \log 0 = 0$. Similar to eqn. (77), eqn. (78) holds if and only if $p_k = \frac{N_{12}^k d_k}{\sum_j N_{12}^j d_j}$.

eqn. (9), we encounter an intriguing phenomenon, i.e., the REE/EE, when applied to a general linear combination operator \mathcal{O} (9), has limitations in capturing information about lighter operators. In this section, we address this limitation¹⁵ by redefining the linear combination operator \mathcal{O} (9) as the form of (18). We then show more details about the entanglement dynamics of such refined operators.

3.1 Recovering information of lighter operators in the EE

To understand the effectiveness of this redefinition, we begin by elucidating why the REE and EE of linear combination operators in (9) fail to capture information about lighter operators. From the perspective of replica calculations, this arises due to the varying divergence of the $2n$ -point functions on the numerator at late times as ϵ approaches zero. Despite L canceling out the dimension of ϵ , allowing for the summation of correlation functions associated with different operators, only the most divergent portion is ultimately retained. While this explanation is accurate, it lacks intuitiveness because, from the definition of the linear operator in (9), all primary operators seem to be equally weighted. Approaching the issue from the perspective of quantum states rather than operators provides a more satisfactory answer. When discussing the EE of locally excited states, we should focus on normalized quantum states rather than operators. Consequently, we prefer to regard $|\psi\rangle$ (9) as a linear combination of normalized locally primary excited states in the Hilbert space rather than a state generated by a linear combination operator, i.e., we have

$$\begin{aligned} |\psi(t)\rangle &= \frac{1}{\sqrt{\mathcal{N}(\epsilon)}} \sum_{k \in \mathcal{S}} C_k L^{2h_k} e^{-iH(t-i\epsilon)} \mathcal{O}_k(x) |\Omega\rangle \\ &= \frac{1}{\sqrt{\mathcal{N}(\epsilon)}} \sum_{k \in \mathcal{S}} C_k \cdot \left(\frac{L}{2\epsilon}\right)^{2h_k} |\mathcal{O}_k(x, t)\rangle, \quad (\langle\psi(t)|\psi(t)\rangle = 1), \end{aligned} \quad (80)$$

where the state $|\mathcal{O}_k(x, t)\rangle$, as defined in (20), carries the information of entanglement among subsystems when the overall system is quenched by $\mathcal{O}_k(x)$ at the initial moment. The above expression indicates that, from the perspective of quantum states, the action of the operator $\mathcal{O}(x)$ (9) on the vacuum results in a superposition of heavy operator states, with the weight of light operator states approaching zero in the limit as ϵ tends to zero¹⁶,

$$\lim_{\epsilon \rightarrow 0} |\psi(t)\rangle = \frac{1}{\sqrt{\sum_{k \in \mathcal{S}_M} |C_k|^2}} \sum_{k \in \mathcal{S}_M} C_k \cdot |\mathcal{O}_k(x, t)\rangle. \quad (81)$$

The method to restore the contributions of light primary operators is currently evident: by following the expression in eqn. (80), and by substituting L with 2ϵ (i.e., considering linear combination operators taking the form of (18)), we can extract information about all operators from the EE. The state generated by the refined linear combination operator (18) is a superposition of all locally primary excited states,

$$|\psi(t)\rangle = \sum_{k \in \mathcal{S}} C_k \cdot |\mathcal{O}_k(x, t)\rangle, \quad \left(\sum_{k \in \mathcal{S}} |C_k|^2 = 1\right). \quad (82)$$

¹⁵We would like to thank Yuya Kusuki for valuable discussions regarding this section.

¹⁶ \mathcal{S}_M in (81) denotes the set consisting of the heaviest operators within the set \mathcal{S} , as introduced in section 2.1.




Figure 3: The variation of the late-time second REE with respect to the combination coefficients in the critical Ising model. Panels (a) and (b): A mixture of spin σ , energy density ϵ , and identity \mathbb{I} . Panels (c) and (d) correspond to linear combinations of ϵ with σ and ϵ with \mathbb{I} , respectively. Panel (b) as well as the hollow circles in panels (c) and (d) represent numerical data obtained from known correlation functions in the critical Ising model. Panel (a) as well as the solid lines in panels (c) and (d) are plotted according to eqn. (83). Note that the maximum value of the second REE is $\log(\sqrt{2}+2)$, as the quantum dimensions of three primary are $d_\sigma = \sqrt{2}$, $d_\epsilon = d_{\mathbb{I}} = 1$.

Utilizing two analytical methods introduced before, the n -th REE and EE of the above state are given by

$$\lim_{t \rightarrow \infty} \Delta S^{(n)}(\rho_A^\psi(t)) = \frac{1}{1-n} \log \left(\sum_{k \in \mathcal{S}} d_k^{1-n} |C_k|^{2n} \right), \quad (83)$$

$$\lim_{t \rightarrow \infty} \Delta S(\rho_A^\psi(t)) = H(p_k) + \sum_{k \in \mathcal{S}} p_k \log d_k, \quad (84)$$

where $p_k \equiv |C_k|^2$. Both REE and EE reach their maximum value of $\log(\sum_{k \in \mathcal{S}} d_k)$ when the combination coefficients are given by $|C_k| = |C_k^*| \equiv \sqrt{\frac{d_k}{\sum_{j \in \mathcal{S}} d_j}}$. We can also validate our results using numerical replica computations in the critical Ising model, as illustrated in Fig. 3.

3.2 The quasiparticle picture

As mentioned in the introduction, for integrable theories such as RCFTs, each time-evolving locally primary or descendant excited state (20) possesses an interpretation of the classical picture of quasiparticles propagation [7–11]: at time $t = 0$, a pair of entanglement quasiparticles

excited by the operator \mathcal{O}_k emerge at position x , and subsequently propagate at the speed of light to positive and negative infinity respectively (although, with spatially inhomogeneous Hamiltonians such as Hamiltonians under sine-square deformation [76, 77], the quasiparticles no longer travel at the speed of light [78–80]). Naturally, one would inquire whether the classical picture of quasiparticles propagation still holds for linear combinations of these locally excited states. Although it has been observed that for combinations of primary operators with the same scaling dimensions, the quasiparticles picture remains applicable [9, 33, 81]. It’s still worrisome that this classical picture may fail in general superposition states. On one hand, a combination of primary operators with differing scaling dimensions cannot be regarded as a primary operator anymore. On the other hand, within quantum mechanics, we know that superpositions of coherent (or quasiclassical) states generally no longer possess classical correspondences. We can check the validity of the classical picture by analyzing the full-time evolution of the EE of superposition states. If the EE evolution of a certain superposition state exhibits a step-like pattern (e.g., see Fig. 4), we infer that the classical interpretation of quasiparticles propagation holds for that state. Otherwise, we do not consider it to have a classical interpretation.




Figure 4: The complete time evolution of EE, which exhibits a step-like behavior.

Let us reanalyze the REE of a state $|\psi\rangle$ generated by the refined linear combination operator

$$\mathcal{O}(x; \epsilon) = \sum_{k \in \mathcal{S}} C_k \cdot (2\epsilon)^{2h_k} \mathcal{O}_k(x),$$

$$\left(\sum_{k \in \mathcal{S}} |C_k|^2 = 1, \quad |\psi(t)\rangle \equiv e^{-iHt} |\psi\rangle = \frac{1}{\sqrt{\mathcal{N}(\epsilon)}} e^{-(\epsilon+it)H} \mathcal{O}(x, \epsilon) |\Omega\rangle \right) \quad (85)$$

using the replica method. We keep subsystem A as $[0, \infty)$ and mainly focus on the case of $n = 2$ for simplicity since extending this proposition to any n is straightforward. As shown in section 2.1.2, utilizing the replica method, the second REE is reduced to a series of four-point functions on a 2-sheeted Riemann surface Σ_2

$$e^{-\Delta S^{(2)}(\rho_A^\psi(t))} = \sum_{ijkl \in \mathcal{S}} C_i C_j^* C_k C_l^* (2\epsilon)^{2(h_i+h_j+h_k+h_l)} \langle \mathcal{O}_i(w_1, \bar{w}_1) \mathcal{O}_j^\dagger(w_2, \bar{w}_2) \mathcal{O}_k(w_3, \bar{w}_3) \mathcal{O}_l^\dagger(w_4, \bar{w}_4) \rangle_{\Sigma_2}. \quad (86)$$

The four-point functions in the sum, with the help of the conformal mapping $w = z^2$, can be expressed as four-point functions on the plane Σ_1 (we take $\{i, j, k, l\} = \{1, 2, 3, 4\}$ for example),

$$\langle \mathcal{O}_1(w_1, \bar{w}_1) \mathcal{O}_2^\dagger(w_2, \bar{w}_2) \mathcal{O}_3(w_3, \bar{w}_3) \mathcal{O}_4^\dagger(w_4, \bar{w}_4) \rangle_{\Sigma_2} = \left(\left| \frac{2z_1 z_{12} z_{14}}{z_{24}} \right|^{-2h_1} \prod_{i=2}^4 \left| \frac{2z_i z_{1i}^2 z_{24}}{z_{12} z_{14}} \right|^{-2h_i} \right) G_{34}^{21}(\eta, \bar{\eta}), \quad (87)$$

where (z_i, \bar{z}_i) are defined in eqn. (39) and $(\eta, \bar{\eta}) \equiv \left(\frac{z_{12}z_{34}}{z_{13}z_{24}}, \frac{\bar{z}_{12}\bar{z}_{34}}{\bar{z}_{13}\bar{z}_{24}}\right)$ is cross ratio. $G_{34}^{21}(\eta, \bar{\eta})$ can be expanded as eqn. (41) in terms of conformal blocks $F_{34}^{21}(p|\eta)$.

At early times $0 < t < |x|$, we find that the cross ratio becomes

$$\eta \simeq \frac{\epsilon^2}{4(|x| - t)^2} \sim 0, \quad \bar{\eta} \simeq \frac{\epsilon^2}{4(|x| + t)^2} \sim 0 \quad (88)$$

in the limits of $\epsilon \rightarrow 0$. In terms of the expansion (42) of the conformal blocks at zero, we arrive at the early time behavior of the four-point function G_{34}^{21} ,

$$G_{34}^{21}(\eta, \bar{\eta}) = \sum_p \left(C_{12}^p C_{34}^p \left(\frac{\epsilon^2}{4|x|^2 - 4t^2} \right)^{2h_p - 2h_3 - 2h_4} (1 + O(\epsilon)) \right). \quad (89)$$

On the other hand, at early times, the leading order of the prefactor of G_{34}^{21} in eqn. (87) in $\epsilon \rightarrow 0$ limit reads

$$\left(\left| \frac{2z_1 z_{12} z_{14}}{z_{24}} \right|^{-2h_1} \prod_{i=2}^4 \left| \frac{2z_i z_{1i}^2 z_{24}}{z_{12} z_{14}} \right|^{-2h_i} \right) = \frac{2^{-2(h_1+h_2+h_3+h_4)} \epsilon^{-2h_1-2h_2+2h_3+2h_4}}{(4|x|^2 - 4t^2)^{2h_3+2h_4}} (1 + O(\epsilon)). \quad (90)$$

Combining (89) with (90), the four-point function on Σ_2 at early times is given by

$$\begin{aligned} & (2\epsilon)^{2(h_1+h_2+h_3+h_4)} \langle \mathcal{O}_1(w_1, \bar{w}_1) \mathcal{O}_2^\dagger(w_2, \bar{w}_2) \mathcal{O}_3(w_3, \bar{w}_3) \mathcal{O}_4^\dagger(w_4, \bar{w}_4) \rangle_{\Sigma_2} \\ &= \sum_p \left(C_{12}^p C_{34}^p \left(\frac{\epsilon^2}{4|x|^2 - 4t^2} \right)^{2h_p} (1 + O(\epsilon)) \right) \quad (0 < t < |x|). \end{aligned} \quad (91)$$

It can be observed that the above equation is non-zero if and only if there exists an identity block in G_{34}^{21} . In this scenario, we have $\mathcal{O}_3 = \mathcal{O}_4$ and $\mathcal{O}_1 = \mathcal{O}_2$. Consequently, we conclude that the four-point functions on Σ_2 at early time

$$(2\epsilon)^{2(h_i+h_j+h_k+h_l)} \langle \mathcal{O}_i(w_1, \bar{w}_1) \mathcal{O}_j^\dagger(w_2, \bar{w}_2) \mathcal{O}_k(w_3, \bar{w}_3) \mathcal{O}_l^\dagger(w_4, \bar{w}_4) \rangle_{\Sigma_2} = \delta_{ij} \delta_{kl}. \quad (92)$$

Substituting (92) into (86), we find that the 2nd REE at early times is zero,

$$\Delta S^{(2)}(\rho_A^\psi(t)) = -\log \sum_{ijkl \in \mathcal{S}} C_i C_j^* C_k C_l^* \delta_{ij} \delta_{kl} = -2 \log \sum_{i \in \mathcal{S}} |C_i|^2 = 0 \quad (0 < t < |x|). \quad (93)$$

The late-time analysis closely parallels that of section 2.1.2. It's essential to note that as we are discussing EE rather than PE (i.e., we have $\tilde{x} = x$), the cross-ratio will experience a discontinuity at $t = |x|$,

$$\eta \simeq 1 - \frac{\epsilon^2}{4(t - |x|)^2} \sim 1, \quad \bar{\eta} \simeq \frac{\epsilon^2}{4(t + |x|)^2} \sim 0, \quad (t > |x|). \quad (94)$$

This results in

$$G_{34}^{21}(\eta, \bar{\eta}) = \sum_{p,q} \left(C_{12}^p C_{34}^q F_{pq} \left[\begin{matrix} 2,1 \\ 3,4 \end{matrix} \right] \frac{(\epsilon/2)^{2(h_p+h_q-h_2-2h_3-h_4)}}{(t - |x|)^{2(h_q-h_2-h_3)} (t + |x|)^{2(h_p-h_3-h_4)}} (1 + O(\epsilon)) \right) \quad (95)$$

for $t > |x|$. On the other hand, the leading order of the prefactor at late times reads

$$\begin{aligned} & \left(\left| \frac{2z_1 z_{12} z_{14}}{z_{24}} \right|^{-2h_1} \prod_{i=2}^4 \left| \frac{2z_i z_{1i}^2 z_{24}}{z_{12} z_{14}} \right|^{-2h_i} \right) \\ &= \epsilon^{2h_3-2h_1} 4^{-h_1-2h_2-3h_3-2h_4} (t - |x|)^{-2(h_2+h_3)} (t + |x|)^{-2(h_3+h_4)} (1 + O(\epsilon)). \end{aligned} \quad (96)$$

Combining the above two equations, we find that

$$(2\epsilon)^{2(h_1+h_2+h_3+h_4)} \langle \mathcal{O}_1(w_1, \bar{w}_1) \mathcal{O}_2^\dagger(w_2, \bar{w}_2) \mathcal{O}_3(w_3, \bar{w}_3) \mathcal{O}_4^\dagger(w_4, \bar{w}_4) \rangle_{\Sigma_2} \\ = \sum_{pq} \left(C_{12}^p C_{34}^p F_{pq} \left[\begin{matrix} 2,1 \\ 3,4 \end{matrix} \right] \frac{\epsilon^{2(h_p+h_q)}}{(t-|x|)^{2h_q} (t+|x|)^{2h_p}} (1+O(\epsilon)) \right) \quad (t > |x|). \quad (97)$$

Similarly, we observe that eqn. (97) is non-zero if and only if both vacuum blocks exist in G_{34}^{21} and G_{32}^{41} . In this scenario, we have $\mathcal{O}_1 = \mathcal{O}_2 = \mathcal{O}_3 = \mathcal{O}_4$. We thus conclude that at late times

$$(2\epsilon)^{2(h_i+h_j+h_k+h_l)} \langle \mathcal{O}_i(w_1, \bar{w}_1) \mathcal{O}_j^\dagger(w_2, \bar{w}_2) \mathcal{O}_k(w_3, \bar{w}_3) \mathcal{O}_l^\dagger(w_4, \bar{w}_4) \rangle_{\Sigma_2} = \begin{cases} d_i^{-1}, & i = j = k = l, \\ 0, & \text{otherwise.} \end{cases}, \quad (98)$$

where we use the formula $F_{00} \left[\begin{matrix} \mathcal{O}, \mathcal{O} \\ \mathcal{O}, \mathcal{O} \end{matrix} \right] = d_{\mathcal{O}}^{-1}$ in RCFTs [74]. Substituting (98) into (86), one obtains the late-time value which was previously encountered in section 3.1.

We have demonstrated the 2nd REE evolution of the linear combination operator (18) exhibiting a step-like behavior. The results can be straightforwardly extended to a general n based on the methodology outlined in section 2.1.2. For the general n -th REE evolution, the $2n$ -point functions on Σ_n behave like

$$(2\epsilon)^{2\sum_{\alpha=1}^{2n} h_{i_\alpha}} \left\langle \prod_{\alpha=1}^n \mathcal{O}_{i_{2\alpha-1}}(w_{2\alpha-1}, \bar{w}_{2\alpha-1}) \mathcal{O}_{i_{2\alpha}}^\dagger(w_{2\alpha}, \bar{w}_{2\alpha}) \right\rangle_{\Sigma_2} = \begin{cases} \prod_{\alpha=1}^n \delta_{i_{2\alpha-1} i_{2\alpha}}, & 0 < t < |x|, \\ d_{i_1}^{1-n} \prod_{\alpha=2}^{2n} \delta_{i_1 i_\alpha}, & t > |x|. \end{cases} \quad (99)$$

Consequently, the complete time evolution of the n -th REE is given by

$$\Delta S^{(n)}(\rho_A^\psi(t)) = \begin{cases} 0, & 0 < t < |x|, \\ \frac{1}{1-n} \log \sum_{k \in \mathcal{S}} d_k^{1-n} |C_k|^{2n}, & t > |x|. \end{cases} \quad (100)$$

By analytically continuing n to 1, one obtains the complete time evolution of EE,

$$\Delta S(\rho_A^\psi(t)) = \begin{cases} 0, & 0 < t < |x|, \\ H(p_k) + \sum_{k \in \mathcal{S}} p_k \log d_k, & t > |x|, \end{cases} \quad (101)$$

where $p_k \equiv |C_k|^2$ and $H(p_k)$ the classical Shannon entropy of $\{p_k\}$. Eqn. (101) indicates that the classical picture of quasiparticle propagation remains applicable for the linear combination states (85) in RCFTs. It should be noted that while the evolution of EE and REE exhibits a step-like behavior, in the case of PE and PREE, the results generally do not follow a step-like pattern [43]. That is, the late-time formulae for PE (69) and PREE (66) hold only under large- t limit.

3.3 Block diagonalization of the reduced density matrix at late times

What we aim to elaborate on next is that eqn. (101) suggests that after t exceeds $|x|$, the reduced density matrix ρ_A^ψ will be block diagonal¹⁷,

$$\rho_A^\psi(t) = \bigoplus_{k \in \mathcal{S}} \rho_A^{\mathcal{O}_k}(t), \quad (t > |x|), \quad (102)$$

¹⁷We would like to thank Wu-zhong Guo for valuable discussions regarding this section.

where $\rho_A^{\mathcal{O}_k}(t)$ represents the reduced density matrix (23) generated by the primary operator \mathcal{O}_k within the set \mathcal{S} . To see this, let us first define a new reduced density matrix of A utilizing $\rho_A^{\mathcal{O}_k}$ [33],

$$\sigma_A^\psi(t) := \sum_{k \in \mathcal{S}} |C_k|^2 \rho_A^{\mathcal{O}_k}(t), \quad \text{tr}_A[\sigma_A^\psi(t)] = 1. \quad (103)$$

Despite the apparent difference between ρ_A^ψ and σ_A^ψ by definition,

$$\rho_A^\psi - \sigma_A^\psi = \sum_{i,j \in \mathcal{S}; i \neq j} C_i C_j^* \mathcal{T}^{\mathcal{O}_i | \mathcal{O}_j},$$

$$\left(\mathcal{T}^{\mathcal{O}_i | \mathcal{O}_j}(t) := |\mathcal{O}_i(x, t)\rangle \langle \mathcal{O}_j(x, t)| \equiv \frac{e^{-iHt} \mathcal{O}_i(x, -\epsilon) |\Omega\rangle \langle \Omega| \mathcal{O}_j^\dagger(x, \epsilon) e^{iHt}}{\sqrt{\langle \mathcal{O}_i^\dagger(x, \epsilon) \mathcal{O}_i(x, -\epsilon) \rangle \langle \mathcal{O}_j^\dagger(x, \epsilon) \mathcal{O}_j(x, -\epsilon) \rangle}} \right), \quad (104)$$

we demonstrate that indeed ρ_A^ψ and σ_A^ψ are equivalent throughout the entire time evolution. This can be achieved by computing the relative entropy between ρ_A^ψ and σ_A^ψ . We know that the relative entropy between two reduced density matrices is 0 if and only if these two reduced density matrices are identical. In practice, one usually compute the Rényi relative entropy¹⁸ [82]

$$S^{(n)}(\rho_A^\psi || \sigma_A^\psi) = \frac{1}{n-1} \left(\log \text{tr}_A[(\rho_A^\psi)^n] - \log \text{tr}_A[\rho_A^\psi (\sigma_A^\psi)^{n-1}] \right) \quad (105)$$

first and then take the limit of $n \rightarrow 1$ to obtain the relative entropy,

$$S(\rho_A^\psi || \sigma_A^\psi) = \lim_{n \rightarrow 1} S^{(n)}(\rho_A^\psi || \sigma_A^\psi) = \text{tr}_A[\rho_A^\psi \log \rho_A^\psi] - \text{tr}_A[\rho_A^\psi \log \sigma_A^\psi]. \quad (106)$$

Since the first part of the n -th Rényi relative entropy (105) is nothing but the n -th REE of ρ_A^ψ , we only need to focus on its second part, which amounts to the following sum of $2n$ -point functions on Σ_n ,

$$\text{tr}_A[\rho_A^\psi (\sigma_A^\psi)^{n-1}] = \frac{Z_n}{Z_1^n} \sum_{\{k_\alpha\} \in \mathcal{S}} C_{k_0} C_{k_1}^* \prod_{\alpha=2}^n |C_{k_\alpha}|^2 (2\epsilon)^{2(h_{k_0} + h_{k_1} + 2 \sum_{\alpha=2}^n h_{k_\alpha})}$$

$$\times \left\langle \mathcal{O}_{k_0}(w_1, \bar{w}_1) \mathcal{O}_{k_1}^\dagger(w_2, \bar{w}_2) \prod_{\alpha=2}^n \mathcal{O}_{k_\alpha}(w_{2\alpha-1}, \bar{w}_{2\alpha-1}) \mathcal{O}_{k_\alpha}^\dagger(w_{2\alpha}, \bar{w}_{2\alpha}) \right\rangle_{\Sigma_n}. \quad (107)$$

Leveraging eqn. (99), we can figure out the above sum and determine the evolution behavior of $\text{tr}_A[\rho_A^\psi (\sigma_A^\psi)^{n-1}]$ ¹⁹,

$$\text{tr}_A[\rho_A^\psi (\sigma_A^\psi)^{n-1}] = \begin{cases} \frac{Z_n}{Z_1^n}, & 0 < t < |x|, \\ \frac{Z_n}{Z_1^n} \sum_{k \in \mathcal{S}} d_k^{1-n} |C_k|^{2n}, & t > |x|. \end{cases} \quad (108)$$

Substituting eqns (108) and (100) into eqn. (105), it becomes evident that the Rényi relative entropy of any order remains zero throughout time evolution. Utilizing the analytic continuation of n , the relative entropy between ρ_A^ψ and σ_A^ψ stays zero consistently,

$$S(\rho_A^\psi(t) || \sigma_A^\psi(t)) = \lim_{n \rightarrow 1} S^{(n)}(\rho_A^\psi(t) || \sigma_A^\psi(t)) = 0, \quad (t > 0). \quad (109)$$

¹⁸Unlike the relative entropy, Rényi relative entropy is not non-negative.

¹⁹Recall that Z_n is the partition function of the Riemann surface Σ_n .

Therefore, we conclude that $\rho_A^\psi(t) = \sigma_A^\psi(t) = \sum_{k \in \mathcal{S}} p_k \rho_A^{\mathcal{O}_k}(t)$ throughout the evolution. Next, we demonstrate that the reduced density matrices corresponding to different primary operators are orthogonal to each other at late times. The crux of the matter is that we can find

$$\text{tr}_A \left[\sqrt{\rho_A^{\mathcal{O}_l}(t) \rho_A^{\mathcal{O}_k}(t)} \sqrt{\rho_A^{\mathcal{O}_l}(t)} \right] = \text{tr}_A [\rho_A^{\mathcal{O}_k}(t) \rho_A^{\mathcal{O}_l}(t)] = 0, \quad (t > |x|, k \neq l) \quad (110)$$

through eqn. (92). Since $\sqrt{\rho_A^{\mathcal{O}_l}(t) \rho_A^{\mathcal{O}_k}(t)} \sqrt{\rho_A^{\mathcal{O}_l}(t)}$ is a semi-positive definite operator, the above equation being valid implies that $\sqrt{\rho_A^{\mathcal{O}_l}(t) \rho_A^{\mathcal{O}_k}(t)} \sqrt{\rho_A^{\mathcal{O}_l}(t)}$ has a spectrum consisting solely of zeros. On the other hand, by the Jacobson Lemma, the operators AB and BA share the same nonzero spectrum. Hence, $\rho_A^{\mathcal{O}_k}(t) \rho_A^{\mathcal{O}_l}(t)$ also possesses a spectrum solely composed of zeros. Moreover, owing to the semi-positivity of $\rho_A^{\mathcal{O}_k}(t)$ and $\rho_A^{\mathcal{O}_l}(t)$, $\rho_A^{\mathcal{O}_k}(t) \rho_A^{\mathcal{O}_l}(t)$ is diagonalizable (see appendix A for more details), indicating that $\rho_A^{\mathcal{O}_k}(t) \rho_A^{\mathcal{O}_l}(t)$ is zero. Similarly, $\rho_A^{\mathcal{O}_l}(t) \rho_A^{\mathcal{O}_k}(t)$ is also a zero operator, so $\rho_A^{\mathcal{O}_k}(t)$ and $\rho_A^{\mathcal{O}_l}(t)$ are commutative.

When two matrices ρ_i and ρ_j are commutative, there exist a unitary matrix denoted as U such that

$$U \rho_i U^\dagger = \Lambda_i, \quad U \rho_j U^\dagger = \Lambda_j, \quad (111)$$

where Λ_i and Λ_j are diagonal matrices with non-negative eigenvalues. In this scenario, $\rho_i \rho_j = 0$ is equivalent to $\Lambda_i \Lambda_j = 0$. The latter implies that the nonzero eigenvalues of Λ_i and the nonzero eigenvalues of Λ_j are distributed at different locations along the diagonal. Consequently, $\rho_i + \rho_j$ will be a block diagonal matrix. Therefore, we conclude that $\rho_A^\psi(t) = \sum_{k \in \mathcal{S}} p_k \rho_A^{\mathcal{O}_k}(t)$ will be a block diagonal matrix (102) after $t > |x|$.

4 Conclusion and prospect

This paper investigates the late-time behavior of REE and PRE in local operator quenches of RCFT, with a particular emphasis on quenches involving linear combination operators. For a state $|\psi\rangle$ in the form of (9), the REE captures only the information of the heaviest primary (57). In contrast, when primary operators with distinct conformal weights are included, the PREE of the reduced transition matrix $\mathcal{T}_A^{\psi|\tilde{\psi}}$ preserves the information of all the primary operators (66). The maximal value attainable by $S^{(n)}(\mathcal{T}_A^{\psi|\tilde{\psi}})$ but unattainable by $S^{(n)}(\rho_A^\psi)$ is the logarithm of the sum of the quantum dimensions of all primary operators,

$$\max_{\{C_k\}} \left\{ \lim_{t \rightarrow \infty} \Delta S^{(n)}(\mathcal{T}_A^{\psi|\tilde{\psi}}(t)) \right\} = \log \left(\sum_{k \in \mathcal{S}} d_k \right). \quad (112)$$

This phenomenon, also known as ‘‘pseudo-entropy amplification’’, occurs when incorporating operators with different conformal dimensions within the linear combination operator \mathcal{O} in (9).

Furthermore, we utilized EE to explore the relationships among quantum dimension, OPE, and fusion coefficients. The computation of EE from both sides of eqn (70) should yield identical results, leading to the relationship between OPE coefficients and fusion coefficients

$$\sqrt{\frac{N_{12}^k d_k}{N_{12}^l d_l}} = \frac{C_{k12}}{C_{l12}} \lim_{\epsilon \rightarrow 0} \sqrt{\frac{|\mathcal{N}_{k12}(\epsilon)|}{|\mathcal{N}_{l12}(\epsilon)|}}, \quad N_{12}^k \in \{0, 1\}. \quad (113)$$

Finally, we discuss that to recover information about lighter operators from EE in general cases, one should consider the regularized linear combination operator (18) instead of (9), or

equivalently, from the perspective of states,

$$|\psi\rangle = \frac{1}{\sqrt{\mathcal{N}(\epsilon)}} \mathcal{O}(x; \epsilon) |\Omega\rangle = \sum_{k \in \mathcal{S}} C_k |\mathcal{O}_k(x)\rangle. \quad (114)$$

It is in combining states within the Hilbert space that we truly perform linear combinations, highlighting the nontriviality of defining the linear combination operator from the perspective of operators. The full-time evolution of EE for the regularized linear combination operator is determined to be

$$\Delta S(\rho_A^\psi(t)) = \begin{cases} 0, & 0 < t < |x|, \\ H(p_k) + \sum_{k \in \mathcal{S}} p_k \log d_k, & t > |x|, \end{cases} \quad (115)$$

indicating that the classical picture of quasiparticles propagation remains applicable for the regularized linear combination states in RCFTs. The quasiparticle picture also indicate that the reduced density matrix ρ_A^ψ will be block diagonal after t surpasses $|x|$,

$$\rho_A^\psi(t) = \bigoplus_{k \in \mathcal{S}} \rho_A^{\mathcal{O}_k}(t), \quad (t > |x|), \quad (116)$$

with each block containing information about one conformal family.

Since we are currently focusing on RCFTs, a natural question for future research is to extend the relevant investigations to Liouville CFT, holographic CFTs, $T\bar{T}$ -deformed CFTs, etc. When investigating the relationship between quantum dimension, OPE coefficients, and fusion coefficients, we must deal with a summation involving OPE coefficients β and coefficients of two point functions, which is extremely challenging at the technical level. The block diagonalization of the reduced density matrix for the linear combination operator, akin to symmetry resolved entanglement [83–87], may reveal hidden symmetries in the local linear combination operator quenches. We would like to leave them to future work.

Acknowledgements

We would like to thank Tadashi Takayanagi, Wu-zhong Guo, Yuya Kusuki, Yuan Sun, Hao Ouyang, Hong-An Zeng, and Yang Liu for their valuable discussions related to this work. We are also grateful to all the organizers of the "Quantum Information, Quantum Matter and Quantum Gravity" workshop (YITP-T-23-01) held at YITP, Kyoto University, where a part of this work was done. SH would like to appreciate the financial support from Jilin University, Max Planck Partner Group, and the Natural Science Foundation of China Grants (No.12075101, No.12235016). L.Z and Z.Z are supported by the Science and Technology Development Plan Project of Jilin Province, China (No. 20240101326JC).

A The product of two semi-positive matrices is diagonalizable

Suppose A and B are two semi-positive matrices, so there exist a invertible matrix P , such that $B = P^\dagger D P$, where

$$D = \begin{pmatrix} I_k & 0 \\ 0 & 0 \end{pmatrix}, \quad (117)$$

where I_k denotes a k by k identity matrix. Since $AB \sim PAP^\dagger(P^\dagger)^{-1}BP^{-1} = PAP^\dagger D \equiv CD$, to prove the product of two semi-positive matrices is diagonalizable, we need to show matrix CD is diagonalizable. Let matrix C be equal to

$$\begin{pmatrix} C_1 & C_2^\dagger \\ C_2 & C_3 \end{pmatrix}, \quad (118)$$

and by the definition of C , it is obviously a semi-positive matrix which can be reformulated as

$$C = M^\dagger \begin{pmatrix} I_{k'} & 0 \\ 0 & 0 \end{pmatrix} M, \quad (119)$$

where M is a invertible matrix.

Suppose x is the solution of equation $C_1 x = 0$, then we have

$$(x^\dagger \ 0) \begin{pmatrix} C_1 & C_2^\dagger \\ C_2 & C_3 \end{pmatrix} \begin{pmatrix} x \\ 0 \end{pmatrix} = (x^\dagger \ 0) M^\dagger \begin{pmatrix} I_{k'} & 0 \\ 0 & 0 \end{pmatrix} M \begin{pmatrix} x \\ 0 \end{pmatrix} = 0. \quad (120)$$

If we denote $y = M \begin{pmatrix} x \\ 0 \end{pmatrix}$, (120) can be rewritten as

$$y^\dagger \begin{pmatrix} I_{k'} & 0 \\ 0 & 0 \end{pmatrix} y = 0, \quad (121)$$

giving rise to the fact that $y = \begin{pmatrix} 0 \\ * \end{pmatrix}$ which leads to $y^\dagger \begin{pmatrix} I_{k'} & 0 \\ 0 & 0 \end{pmatrix} M = 0$, i.e.

$$(x^\dagger \ 0) \begin{pmatrix} C_1 & C_2^\dagger \\ C_2 & C_3 \end{pmatrix} = 0 \Rightarrow \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} x = 0. \quad (122)$$

Therefore, $C_1 x = 0 \iff \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} x = 0$, i.e. $\text{rank}(C_1) = \text{rank} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix}$, indicating that there exist a matrix Q such that $C_2 = QC_1$. After multiplying the matrix CD on the left by a matrix X and on the right by the inverse of X , where $X = \begin{pmatrix} I & 0 \\ -Q & I \end{pmatrix}$, CD becomes

$$\begin{pmatrix} I & 0 \\ -Q & I \end{pmatrix} CD \begin{pmatrix} I & 0 \\ Q & I \end{pmatrix} = \begin{pmatrix} I & 0 \\ -Q & I \end{pmatrix} \begin{pmatrix} C_1 & 0 \\ C_2 & 0 \end{pmatrix} \begin{pmatrix} I & 0 \\ Q & I \end{pmatrix} = \begin{pmatrix} C_1 & 0 \\ 0 & 0 \end{pmatrix}. \quad (123)$$

Since C is a semi-positive matrix, C_1 can be diagonalized, i.e. CD is diagonalizable which leads to the fact that the product of two semi-positive matrices is diagonalizable.

References

- [1] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, *Infinite Conformal Symmetry in Two-Dimensional Quantum Field Theory*, *Nucl. Phys. B* **241** (1984) 333–380.
- [2] P. Di Francesco, P. Mathieu and D. Senechal, *Conformal Field Theory*. Graduate Texts in Contemporary Physics. Springer-Verlag, New York, 1997, 10.1007/978-1-4612-2256-9.
- [3] E. P. Verlinde, *Fusion Rules and Modular Transformations in 2D Conformal Field Theory*, *Nucl. Phys. B* **300** (1988) 360–376.

- [4] J. Fuchs, *Fusion rules in conformal field theory*, *Fortsch. Phys.* **42** (1994) 1–48, [[hep-th/9306162](#)].
- [5] J. Fuchs, *Quantum dimensions*, *Commun. Theor. Phys.* **1** (1991) 59–109.
- [6] C. Jeco and J. Troost, *Notes on the Verlinde formula in non-rational conformal field theories*, *Phys. Rev. D* **74** (2006) 106002, [[hep-th/0601085](#)].
- [7] M. Nozaki, T. Numasawa and T. Takayanagi, *Quantum Entanglement of Local Operators in Conformal Field Theories*, *Phys. Rev. Lett.* **112** (2014) 111602, [[1401.0539](#)].
- [8] S. He, T. Numasawa, T. Takayanagi and K. Watanabe, *Quantum dimension as entanglement entropy in two dimensional conformal field theories*, *Phys. Rev. D* **90** (2014) 041701, [[1403.0702](#)].
- [9] P. Caputa and A. Veliz-Osorio, *Entanglement constant for conformal families*, *Phys. Rev. D* **92** (2015) 065010, [[1507.00582](#)].
- [10] B. Chen, W.-Z. Guo, S. He and J.-q. Wu, *Entanglement Entropy for Descendent Local Operators in 2D CFTs*, *JHEP* **10** (2015) 173, [[1507.01157](#)].
- [11] T. Numasawa, *Scattering effect on entanglement propagation in RCFTs*, *JHEP* **12** (2016) 061, [[1610.06181](#)].
- [12] W.-Z. Guo, S. He and Z.-X. Luo, *Entanglement entropy in (1+1)D CFTs with multiple local excitations*, *JHEP* **05** (2018) 154, [[1802.08815](#)].
- [13] P. Caputa, J. Simón, A. Štikonas and T. Takayanagi, *Quantum Entanglement of Localized Excited States at Finite Temperature*, *JHEP* **01** (2015) 102, [[1410.2287](#)].
- [14] P. Caputa, M. Nozaki and T. Takayanagi, *Entanglement of local operators in large- N conformal field theories*, *PTEP* **2014** (2014) 093B06, [[1405.5946](#)].
- [15] C. T. Asplund, A. Bernamonti, F. Galli and T. Hartman, *Holographic Entanglement Entropy from 2d CFT: Heavy States and Local Quenches*, *JHEP* **02** (2015) 171, [[1410.1392](#)].
- [16] C. T. Asplund, A. Bernamonti, F. Galli and T. Hartman, *Entanglement Scrambling in 2d Conformal Field Theory*, *JHEP* **09** (2015) 110, [[1506.03772](#)].
- [17] P. Caputa, T. Numasawa and A. Veliz-Osorio, *Out-of-time-ordered correlators and purity in rational conformal field theories*, *PTEP* **2016** (2016) 113B06, [[1602.06542](#)].
- [18] Y. Suzuki, T. Takayanagi and K. Umemoto, *Entanglement Wedges from the Information Metric in Conformal Field Theories*, *Phys. Rev. Lett.* **123** (2019) 221601, [[1908.09939](#)].
- [19] M. Nozaki, T. Numasawa and T. Takayanagi, *Holographic Local Quenches and Entanglement Density*, *JHEP* **05** (2013) 080, [[1302.5703](#)].
- [20] P. Caputa, T. Numasawa, T. Shimaji, T. Takayanagi and Z. Wei, *Double Local Quenches in 2D CFTs and Gravitational Force*, *JHEP* **09** (2019) 018, [[1905.08265](#)].
- [21] Y. Kusuki and M. Miyaji, *Entanglement Entropy after Double Excitation as an Interaction Measure*, *Phys. Rev. Lett.* **124** (2020) 061601, [[1908.03351](#)].
- [22] T. Kawamoto, T. Mori, Y.-k. Suzuki, T. Takayanagi and T. Ugajin, *Holographic local operator quenches in BCFTs*, *JHEP* **05** (2022) 060, [[2203.03851](#)].

- [23] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Adv. Theor. Math. Phys.* **2** (1998) 231–252, [[hep-th/9711200](#)].
- [24] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett. B* **428** (1998) 105–114, [[hep-th/9802109](#)].
- [25] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253–291, [[hep-th/9802150](#)].
- [26] Z. Wang, *Topological quantum computation*. No. 112. American Mathematical Soc., 2010.
- [27] V. Lahtinen and J. Pachos, *A short introduction to topological quantum computation*, *SciPost Physics* **3** (2017) 021.
- [28] E. Rowell and Z. Wang, *Mathematics of topological quantum computing*, *Bulletin of the American Mathematical Society* **55** (2018) 183–238.
- [29] M. Nozaki, *Notes on Quantum Entanglement of Local Operators*, *JHEP* **10** (2014) 147, [[1405.5875](#)].
- [30] W.-Z. Guo and S. He, *Rényi entropy of locally excited states with thermal and boundary effect in 2D CFTs*, *JHEP* **04** (2015) 099, [[1501.00757](#)].
- [31] S. He, *Conformal bootstrap to Rényi entropy in 2D Liouville and super-Liouville CFTs*, *Phys. Rev. D* **99** (2019) 026005, [[1711.00624](#)].
- [32] L. Apolo, S. He, W. Song, J. Xu and J. Zheng, *Entanglement and chaos in warped conformal field theories*, *JHEP* **04** (2019) 009, [[1812.10456](#)].
- [33] A. Bhattacharyya, T. Takayanagi and K. Umemoto, *Universal Local Operator Quenches and Entanglement Entropy*, *JHEP* **11** (2019) 107, [[1909.04680](#)].
- [34] Y. Kusuki and K. Tamaoka, *Entanglement Wedge Cross Section from CFT: Dynamics of Local Operator Quench*, *JHEP* **02** (2020) 017, [[1909.06790](#)].
- [35] K. Goto, M. Nozaki, S. Ryu, K. Tamaoka and M. T. Tan, *Scrambling and Recovery of Quantum Information in Inhomogeneous Quenches in Two-dimensional Conformal Field Theories*, [2302.08009](#).
- [36] J. Kudler-Flam, M. Nozaki, T. Numasawa, S. Ryu and M. T. Tan, *Bridging two quantum quench problems – local joining quantum quench and Möbius quench – and their holographic dual descriptions*, [2309.04665](#).
- [37] C. G. Callan, Jr. and F. Wilczek, *On geometric entropy*, *Phys. Lett. B* **333** (1994) 55–61, [[hep-th/9401072](#)].
- [38] P. Calabrese and J. L. Cardy, *Entanglement entropy and quantum field theory*, *J. Stat. Mech.* **0406** (2004) P06002, [[hep-th/0405152](#)].
- [39] Y. Nakata, T. Takayanagi, Y. Taki, K. Tamaoka and Z. Wei, *New holographic generalization of entanglement entropy*, *Phys. Rev. D* **103** (2021) 026005, [[2005.13801](#)].
- [40] Y. Aharonov and L. Vaidman, *The two-state vector formalism: an updated review*, *Time in quantum mechanics* (2008) 399–447.

- [41] J. Dressel, M. Malik, F. M. Miatto, A. N. Jordan and R. W. Boyd, *Colloquium: Understanding quantum weak values: Basics and applications*, *Rev. Mod. Phys.* **86** (Mar, 2014) 307–316.
- [42] W.-z. Guo, S. He and Y.-X. Zhang, *Constructible reality condition of pseudo entropy via pseudo-Hermiticity*, *JHEP* **05** (2023) 021, [2209.07308].
- [43] W.-z. Guo, S. He and Y.-X. Zhang, *On the real-time evolution of pseudo-entropy in 2d CFTs*, *JHEP* **09** (2022) 094, [2206.11818].
- [44] S. He, J. Yang, Y.-X. Zhang and Z.-X. Zhao, *Pseudo-entropy for descendant operators in two-dimensional conformal field theories*, 2301.04891.
- [45] P. Wang, H. Wu and H. Yang, *Fix the dual geometries of $T\bar{T}$ deformed CFT_2 and highly excited states of CFT_2* , *Eur. Phys. J. C* **80** (2020) 1117, [1811.07758].
- [46] A. Mollabashi, N. Shiba, T. Takayanagi, K. Tamaoka and Z. Wei, *Pseudo Entropy in Free Quantum Field Theories*, *Phys. Rev. Lett.* **126** (2021) 081601, [2011.09648].
- [47] A. Mollabashi, N. Shiba, T. Takayanagi, K. Tamaoka and Z. Wei, *Aspects of pseudoentropy in field theories*, *Phys. Rev. Res.* **3** (2021) 033254, [2106.03118].
- [48] K. Goto, M. Nozaki and K. Tamaoka, *Subregion spectrum form factor via pseudoentropy*, *Phys. Rev. D* **104** (2021) L121902, [2109.00372].
- [49] M. Miyaji, *Island for gravitationally prepared state and pseudo entanglement wedge*, *JHEP* **12** (2021) 013, [2109.03830].
- [50] I. Akal, T. Kawamoto, S.-M. Ruan, T. Takayanagi and Z. Wei, *Page curve under final state projection*, *Phys. Rev. D* **105** (2022) 126026, [2112.08433].
- [51] T. Nishioka, T. Takayanagi and Y. Taki, *Topological pseudo entropy*, *JHEP* **09** (2021) 015, [2107.01797].
- [52] Z. Li, Z.-Q. Xiao and R.-Q. Yang, *On holographic time-like entanglement entropy*, *JHEP* **04** (2023) 004, [2211.14883].
- [53] Y. Ishiyama, R. Kojima, S. Matsui and K. Tamaoka, *Notes on pseudo entropy amplification*, *PTEP* **2022** (2022) 093B10, [2206.14551].
- [54] K. Doi, J. Harper, A. Mollabashi, T. Takayanagi and Y. Taki, *Pseudoentropy in dS/CFT and Timelike Entanglement Entropy*, *Phys. Rev. Lett.* **130** (2023) 031601, [2210.09457].
- [55] K. Doi, J. Harper, A. Mollabashi, T. Takayanagi and Y. Taki, *Timelike entanglement entropy*, *JHEP* **05** (2023) 052, [2302.11695].
- [56] S. He, J. Yang, Y.-X. Zhang and Z.-X. Zhao, *Pseudo entropy of primary operators in $T\bar{T}/J\bar{T}$ -deformed CFTs*, *JHEP* **09** (2023) 025, [2305.10984].
- [57] Z. Chen, *Complex-valued Holographic Pseudo Entropy via Real-time AdS/CFT Correspondence*, 2302.14303.
- [58] D. Chen, X. Jiang and H. Yang, *Holographic $T\bar{T}$ deformed entanglement entropy in dS_3/CFT_2* , 2307.04673.
- [59] W.-z. Guo and J. Zhang, *Sum rule for pseudo Rényi entropy*, 2308.05261.

- [60] A. J. Parzygnat, T. Takayanagi, Y. Taki and Z. Wei, *SVD Entanglement Entropy*, 2307.06531.
- [61] P.-Z. He and H.-Q. Zhang, *Timelike Entanglement Entropy from Rindler Method*, 2307.09803.
- [62] C.-S. Chu and H. Parihar, *Time-like entanglement entropy in AdS/BCFT*, *JHEP* **06** (2023) 173, [2304.10907].
- [63] X. Jiang, P. Wang, H. Wu and H. Yang, *Timelike entanglement entropy in dS_3/CFT_2* , *JHEP* **08** (2023) 216, [2304.10376].
- [64] X. Jiang, P. Wang, H. Wu and H. Yang, *Timelike entanglement entropy and TT^- deformation*, *Phys. Rev. D* **108** (2023) 046004, [2302.13872].
- [65] K. Narayan and H. K. Saini, *Notes on time entanglement and pseudo-entropy*, 2303.01307.
- [66] T. Kawamoto, S.-M. Ruan, Y.-k. Suzuki and T. Takayanagi, *A Half de Sitter Holography*, 2306.07575.
- [67] F. Omid, *Pseudo Rényi Entanglement Entropies For an Excited State and Its Time Evolution in a 2D CFT*, 2309.04112.
- [68] W.-z. Guo and Y. Jiang, *Pseudo entropy and pseudo-Hermiticity in quantum field theories*, 2311.01045.
- [69] K. Shinmyo, T. Takayanagi and K. Tasuki, *Pseudo entropy under joining local quenches*, 2310.12542.
- [70] K. Narayan, *Comments on de Sitter space, extremal surfaces and time entanglement*, 2310.00320.
- [71] K. Narayan, *de Sitter space, extremal surfaces, and time entanglement*, *Phys. Rev. D* **107** (2023) 126004, [2210.12963].
- [72] H. Kanda, T. Kawamoto, Y.-k. Suzuki, T. Takayanagi, K. Tasuki and Z. Wei, *Entanglement Phase Transition in Holographic Pseudo Entropy*, 2311.13201.
- [73] S. He, H.-A. Zeng, Y.-X. Zhang and Z.-X. Zhao. To be appear.
- [74] G. W. Moore and N. Seiberg, *Naturality in Conformal Field Theory*, *Nucl. Phys. B* **313** (1989) 16–40.
- [75] G. W. Moore and N. Seiberg, *Polynomial Equations for Rational Conformal Field Theories*, *Phys. Lett. B* **212** (1988) 451–460.
- [76] A. Gendiar, R. Krccmar and T. Nishino, *Spherical Deformation for One-Dimensional Quantum Systems*, *Prog. Theor. Phys.* **122** (2009) 953–967, [0810.0622].
- [77] H. Katsura, *Exact ground state of the sine-square deformed XY spin chain*, *J. Phys. A* **44** (2011) 252001, [1104.1721].
- [78] X. Wen and J.-Q. Wu, *Quantum dynamics in sine-square deformed conformal field theory: Quench from uniform to nonuniform conformal field theory*, *Phys. Rev. B* **97** (2018) 184309, [1802.07765].

- [79] K. Goto, M. Nozaki, K. Tamaoka, M. T. Tan and S. Ryu, *Non-Equilibrating a Black Hole with Inhomogeneous Quantum Quench*, 2112.14388.
- [80] M. Nozaki, K. Tamaoka and M. T. Tan, *Inhomogeneous quenches as state preparation in two-dimensional conformal field theories*, 2310.19376.
- [81] J. Zhang and P. Calabrese, *Subsystem distance after a local operator quench*, *JHEP* **02** (2020) 056, [1911.04797].
- [82] N. Lashkari, *Modular Hamiltonian for Excited States in Conformal Field Theory*, *Phys. Rev. Lett.* **117** (2016) 041601, [1508.03506].
- [83] M. Goldstein and E. Sela, *Symmetry-resolved entanglement in many-body systems*, *Phys. Rev. Lett.* **120** (2018) 200602, [1711.09418].
- [84] S. Zhao, C. Northe and R. Meyer, *Symmetry-resolved entanglement in AdS_3/CFT_2 coupled to $U(1)$ Chern-Simons theory*, *JHEP* **07** (2021) 030, [2012.11274].
- [85] L. Capizzi, P. Ruggiero and P. Calabrese, *Symmetry resolved entanglement entropy of excited states in a CFT*, *J. Stat. Mech.* **2007** (2020) 073101, [2003.04670].
- [86] K. Weisenberger, S. Zhao, C. Northe and R. Meyer, *Symmetry-resolved entanglement for excited states and two entangling intervals in AdS_3/CFT_2* , *JHEP* **12** (2021) 104, [2108.09210].
- [87] Y. Kusuki, S. Murciano, H. Ooguri and S. Pal, *Symmetry-resolved entanglement entropy, spectra & boundary conformal field theory*, *JHEP* **11** (2023) 216, [2309.03287].