

Generators of the Poincaré Group for arbitrary tensors and spinor-tensors

H.V. Almeida Silva^a *; D. Dalmazi^a †; R.R. Lino dos Santos^b ‡; E.L. Mendonça^a §

a- UNESP - Campus de Guaratinguetá - DFI

b- National Centre for Nuclear Research, Pasteura 7, 02-093 Warsaw, Poland

June 7, 2024

Abstract

In this work, we systematically derive explicit expressions for the Poincaré Group generators on arbitrary-rank tensors and spinor-tensors in $D = 3 + 1$ and $D = 2 + 1$ spacetimes, thus generalizing previous works in the literature for the groups $ISO(3,1)$ and $ISO(2,1)$. From the Casimir eigenvalue equations, we demonstrate in a model-independent way the Fierz-Pauli constraints for massive particles for spins $\mathfrak{s} = 5/2, 3,$ and 4 in $D = 3 + 1$ and helicities $\alpha = 5/2, 3,$ and 4 in $D = 2 + 1$ dimensions. We also comment on the demonstration of the Fierz-Pauli constraints for the general case of arbitrary spin (helicity).

*hudson.vinicius@unesp.br

†denis.dalmazi@unesp.br

‡rafael.santos@ncbj.gov.pl

§elias.leite@unesp.br

Contents

1	Introduction	3
2	Poincaré Group $ISO(3,1)$	4
2.1	Arbitrary rank- s tensors in $D = 3 + 1$	5
2.1.1	Spacetime translations	5
2.1.2	Lorentz transformations	6
2.2	Arbitrary rank- s spinor-tensors in $D = 3 + 1$	7
2.2.1	Spacetime translations and Lorentz transformations	7
2.3	Casimir operators of $\mathfrak{iso}(3,1)$	9
3	Poincaré group $ISO(2,1)$	10
3.1	Arbitrary rank- s tensors and spinor-tensors in $D = 2 + 1$	11
3.2	Casimirs of $\mathfrak{iso}(2,1)$	12
4	Fierz-Pauli constraints from the algebra	13
4.1	Spin-5/2	13
4.1.1	Spin-5/2 in $D = 3 + 1$:	13
4.1.2	Helicity-5/2 in $D = 2 + 1$	14
4.2	Spin-3	15
4.2.1	Spin-3 in $D = 3 + 1$	15
4.2.2	Helicity-3 in $D = 2 + 1$	16
4.3	Spin-4	17
4.3.1	Spin-4 in $D = 3 + 1$	17
4.3.2	Helicity-4 in $D = 2 + 1$	18
4.4	Comments on the case of arbitrary rank- s tensors	19
5	Conclusions	20
A	Spin generators in $D = 2 + 1$	22
A.1	helicity - $\alpha = 5/2$	22
A.2	helicity - $\alpha = 3$	22
A.3	helicity - $\alpha = 4$	23
B	Rank-s algebra	24

1 Introduction

Understanding the fundamental physical laws is central to theoretical physics. At the core of this understanding lies the exploration of symmetries and symmetry groups, which are essential for characterizing the interactions and describing elementary particles. In this regard, the Poincaré Group plays a crucial role by unifying Lorentz transformations with spacetime translations.

The seminal works by Wigner and Bargmann [1, 2] have significantly shaped our understanding of the relationship between symmetries and the properties of elementary particles, such as mass and spin. These elementary particles belong to the unitary irreducible representations of the Poincaré Group labeled by the eigenvalues of the Casimir operators of the algebra, forming the mathematical basis for the description of elementary particles [3].

Additionally, for a massive particle, the so-called Fierz-Pauli constraints reduce the degrees of freedom of a tensor (or spinor-tensor) representation to match the physical degrees of freedom. These constraints were postulated long ago in [4].

In the present article we work both in $D = 3 + 1$ and $D = 2 + 1$ spacetime dimensions and focus on deriving explicit representations for the Poincaré group generators acting on tensors and spinor-tensors for arbitrary integer and half-integer spins (and helicities), respectively. Explicit expressions in the literature can only be found for specific lower spins as $s \leq 2$ in $D = 3 + 1$, see [8], and lower helicities in $D = 2 + 1$ as $\alpha = 3/2, 2$ and 3 , see [5, 6, 7].

We also demonstrate the Fierz-Pauli constraints for massive particles directly from the eigenvalue equations of the Casimir invariants in a model-independent way for the higher spin (HS) cases $\mathfrak{s} = 5/2, 3, 4$ in $D = 3 + 1$ and higher helicity cases $\alpha = 5/2, 3, 4$ in $D = 2 + 1$. The Fierz-Pauli constraints are usually obtained in a dynamical way through the manipulation of the equations of motion of a given model. Here, differently, we demonstrate that they can be obtained from the algebra, generalizing [8], where only the simpler lower spin cases $\mathfrak{s} = 1/2, 1, 3/2, 2$ have been investigated. In $D = 2 + 1$ our demonstrations for $\alpha = 5/2$ and $\alpha = 4$ are new.

Massive HS particles of arbitrary integer and half-integer spins are present in the spectrum of the superstring theory. Nowadays, motivations for HS theories are much broader, though, see, for instance, [9, 10]. We can speculate that string theory itself might be a spontaneously broken phase of a higher spin gravity (HiSGRA) theory that generalizes Einstein's spin-2 gravity, see [11] and references therein for reviewing these motivations. The powerful higher spin local symmetries of HiSGRA would contribute to a possible finiteness of the corresponding quantum field theory, see the seminal earlier works [12, 13]. Furthermore, there are also generalizations of the AdS/CFT duality to HiSGRA [14] and HS applications to $3D$ bosonization [15, 16].

Even beyond bosonization, the case of $D = 2 + 1$ dimensions is especially interesting since, differently from $D = 3 + 1$, helicity eigenstates can be described by local field theories in

$D = 2 + 1$, which are called self-dual models or parity singlets. Interestingly, the helicity $+\alpha$ and helicity $-\alpha$ parity singlets can be “soldered” into a parity doublet containing both helicities $\pm\alpha$. These soldered models are the usual massive spin- \mathfrak{s} models in any dimension $D \geq 3$ for $\mathfrak{s} = 1, 3/2, 2$. Therefore, the parity singlets in $D = 2 + 1$ work like the basic building blocks of massive particles in the real world in $D = 3 + 1$.

This article is structured as follows: Section 2 examines the Poincaré Group in $D = 3 + 1$ spacetime, $ISO(3, 1)$. We obtain general explicit expressions for the group generators within representations of arbitrary-rank tensors and spinor-tensors. Next, Section 3 extends these results to the Poincaré Group in $D = 2 + 1$ spacetime, $ISO(2, 1)$, where we then identify explicit representations of the Casimir operators of the corresponding algebraic structure $\mathfrak{iso}(2, 1)$. Notably, these results can be compared with the findings of Jackiw and Nair, as presented in [18], see also the precursor work on the Poincaré group in $D = 2 + 1$ by Binegar [19]. Then, in Section 4, we obtain the Fierz-Pauli constraints for massive spin(helicity)- $\mathfrak{s}(\alpha)$ ($\mathfrak{s}(\alpha) = 5/2, 3, 4$) particles in $D = 3 + 1$ ($D = 2 + 1$). For the convenience of our readers, in the appendices, we introduce explicit expressions of symmetric spin generators in $D = 2 + 1$ dimensions.

Conventions: In both $D = 3 + 1$ and $D = 2 + 1$, the Minkowski metric $\eta_{\mu\nu}$ is mostly plus, i.e., $\eta = \text{diag}(-, +, +, +)$ and $\eta = \text{diag}(-, +, +)$, respectively. Arbitrary tensors and spinor-tensors, with rank- s , depending on the spacetime coordinates x^μ , are represented by $T^{\mu_1 \dots \mu_s}(x)$ and $\psi^{\mu_1 \dots \mu_s}(x)$, respectively. We use the word “spin” and the symbol \mathfrak{s} for $D = 3 + 1$ dimensions and “helicity” and the symbol α for $D = 2 + 1$. Throughout the text, groups are referred to using capital letters while gothic letters are employed to represent algebras. Symmetrizations, denoted by “(…),” and anti-symmetrizations, represented by “[…]”, are normalized.

The following properties of the Dirac matrices are useful. In $D = 2 + 1$, they are 2×2 real matrices expressed in terms of Pauli matrices, as follows: $\gamma^0 = i\sigma_2$, $\gamma^1 = \sigma_2$, and $\gamma^2 = \sigma_3$. These matrices satisfy the following fundamental relations:

$$[\gamma_\alpha, \gamma_\beta] = 2\epsilon_{\alpha\beta\rho}\gamma^\rho, \quad (1)$$

$$\{\gamma_\mu, \gamma_\nu\} = 2\eta_{\mu\nu}, \quad (2)$$

$$\gamma_\mu\gamma_\nu = \eta_{\mu\nu} + \epsilon_{\mu\nu\alpha}\gamma^\alpha. \quad (3)$$

2 Poincaré Group $ISO(3, 1)$

It is well known that relativistic classical and quantum field theories must be invariant under the Poincaré transformations, which act on a four-vector x^μ spacetime coordinate as:

$$x^\mu \longrightarrow x'^\mu = \Lambda^\mu{}_\nu x^\nu + a^\mu, \quad (4)$$

i.e., such transformation is a composition of the Lorentz transformation parameterized by $\Lambda^\mu{}_\nu$, $SO(3, 1)$, plus a spacetime translation parameterized by a^μ , which together form the

inhomogeneous Lorentz group $ISO(3,1)$. Here, a^μ is a real four-vector while $\Lambda^\mu{}_\nu$ is a rank-two tensor that fulfills the relation:

$$\eta_{\mu\nu}\Lambda^\mu{}_\alpha\Lambda^\nu{}_\beta = \eta_{\alpha\beta} \quad . \quad (5)$$

Since the Poincaré group is a Lie group,¹ we can express Poincaré group transformations as

$$\Lambda^\mu{}_\nu = \delta^\mu{}_\nu - \omega^\mu{}_\nu, \quad a^\mu = \epsilon^\mu \quad , \quad (6)$$

where ϵ^μ and $\omega^\mu{}_\nu$ are continuous infinitesimal parameters. In order to satisfy the condition (5), the tensor $\omega^\mu{}_\nu$ must be antisymmetric, i.e., $\omega^\mu{}_\nu = -\omega_\mu{}^\nu$. Therefore, in $D = 3+1$ dimensions, we have six independent components corresponding to three rotation angles plus three Lorentz boosts. Using (6), the transformation (4) will be given by:

$$x^\mu \longrightarrow x'^\mu = x^\mu - \omega^\mu{}_\nu x^\nu + \epsilon^\mu \quad . \quad (7)$$

The generators of the corresponding Lie algebra satisfy the commutation relations:

$$[I^{\alpha\beta}, I^{\mu\nu}] = i(\eta^{\alpha\mu}I^{\beta\nu} + \eta^{\beta\nu}I^{\alpha\mu} - \eta^{\alpha\nu}I^{\beta\mu} - \eta^{\beta\mu}I^{\alpha\nu}), \quad (8)$$

$$[P^\mu, I^{\alpha\beta}] = i(\eta^{\mu\beta}P^\alpha - \eta^{\mu\alpha}P^\beta), \quad (9)$$

$$[P^\alpha, P^\beta] = 0, \quad (10)$$

where $I^{\alpha\beta} = -I^{\beta\alpha}$ denotes the six generators of the Lorentz group associated with the six independent components of the antisymmetric parameter $\omega_{\mu\nu}$, and P^α denotes the four generators of spacetime translations associated with the four parameters ϵ_μ .

2.1 Arbitrary rank- s tensors in $D = 3 + 1$

In this section, we obtain explicit expressions for the generators of the Poincaré group acting on arbitrary tensors of rank- s , for integer s . We consider translations and Lorentz transformations separately in order to obtain the generators, P^α and $I^{\alpha\beta}$, respectively. We start with the translation case.

2.1.1 Spacetime translations

Considering an infinitesimal spacetime translation:

$$x'^\mu = x^\mu + \epsilon^\mu, \quad (11)$$

¹A generic element of a Lie group, say $g(\theta)$, where θ is a continuous infinitesimal parameter, can always be represented by $D_R(g(\theta)) = e^{i\theta_a T_R^a}$, for a given representation D_R , where $a = 1, \dots, n$, while T_R are the generators of the group corresponding to the representation D_R . This property is independent of the representation. For a pedagogical review, see for instance [25].

Since $\partial x'^{\mu}/\partial x^{\nu} = \delta_{\nu}^{\mu}$ we have:

$$T'^{\mu_1 \dots \mu_s}(x') = T^{\mu_1 \dots \mu_s}(x) \quad . \quad (12)$$

On the one hand, expanding the left hand side of (12), we obtain:

$$T'^{\mu_1 \dots \mu_s}(x) = T^{\mu_1 \dots \mu_s}(x) - \epsilon^{\alpha} \partial_{\alpha} T^{\mu_1 \dots \mu_s}(x) + \mathcal{O}(\epsilon^2) \quad , \quad (13)$$

On the other hand, as a group element acting on the original tensor, we have:

$$T'^{\mu_1 \dots \mu_s}(x) = [\exp(-i\epsilon_{\alpha} P^{\alpha})]_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} T^{\nu_1 \dots \nu_s}(x) \quad . \quad (14)$$

By expanding the exponential on the right-hand side, we obtain:

$$T'^{\mu_1 \dots \mu_s}(x) = T^{\mu_1 \dots \mu_s}(x) - i\epsilon_{\alpha} (P^{\alpha})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} T^{\nu_1 \dots \nu_s}(x) + \mathcal{O}(\epsilon^2) \quad . \quad (15)$$

Comparing (13) with (15), we finally get an explicit expression for the generator of translations:

$$(P^{\alpha})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} = -i \left(\prod_{i=1}^s \delta_{\nu_i}^{\mu_i} \right) \partial^{\alpha} \quad , \quad (16)$$

which is valid on arbitrary rank- s tensors in $D = 3 + 1$ dimensions.

2.1.2 Lorentz transformations

In order to obtain the explicit expressions for the generators of the Lorentz transformations $I^{\alpha\beta}$, we use the same approach of the previous case, but now, notice that under the Lorentz transformation $\partial x'^{\mu}/\partial x^{\nu} = \Lambda_{\nu}^{\mu}$, so we have:

$$T'^{\mu_1 \dots \mu_s}(x') = \left(\prod_{i=1}^s \Lambda_{\nu_i}^{\mu_i} \right) T^{\nu_1 \dots \nu_s}(x) \quad . \quad (17)$$

In the infinitesimal case, considering that $\Lambda_{\nu}^{\mu} = \delta_{\nu}^{\mu} - \omega_{\nu}^{\mu}$ up until the first order in ω , we have explicitly:

$$\begin{aligned} \prod_{i=1}^s \Lambda_{\nu_i}^{\mu_i} &= (\delta_{\nu_1}^{\mu_1} \delta_{\nu_2}^{\mu_2} \dots \delta_{\nu_s}^{\mu_s} - \omega_{\nu_1}^{\mu_1} \delta_{\nu_2}^{\mu_2} \delta_{\nu_3}^{\mu_3} \dots \delta_{\nu_s}^{\mu_s} - \omega_{\nu_2}^{\mu_2} \delta_{\nu_1}^{\mu_1} \delta_{\nu_3}^{\mu_3} \dots \delta_{\nu_s}^{\mu_s} - \omega_{\nu_s}^{\mu_s} \delta_{\nu_1}^{\mu_1} \dots \delta_{\nu_{s-1}}^{\mu_{s-1}} \delta_{\nu_s}^{\mu_s}) \\ &= \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} - \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} \omega_{\nu_k}^{\mu_k} \quad . \end{aligned} \quad (18)$$

Inserting (18) into (17) and expanding the left hand side of (17) up to the first order in ω , we have:

$$T'^{\mu_1 \dots \mu_s}(x) - \omega^{\gamma}_{\alpha} x^{\alpha} \partial_{\gamma} T^{\mu_1 \dots \mu_s}(x) = \left[\prod_{i=1}^s \delta_{\nu_i}^{\mu_i} - \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} \omega_{\nu_k}^{\mu_k} \right] T^{\nu_1 \dots \nu_s}(x) \quad , \quad (19)$$

making explicit use of the antisymmetry of $\omega_{\alpha\beta} \equiv \eta_{\alpha\gamma}\omega^\gamma_\beta$ we have:

$$T'^{\mu_1 \dots \mu_s}(x) = T^{\mu_1 \dots \mu_s}(x) + \frac{\omega_{\alpha\beta}}{2} \left[(x^\beta \partial^\alpha - x^\alpha \partial^\beta) \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} - \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} (\eta^{\alpha\mu_k} \delta_{\nu_k}^\beta - \eta^{\beta\mu_k} \delta_{\nu_k}^\alpha) \right] T^{\nu_1 \dots \nu_s}(x) \quad (20)$$

As an element of the Lorentz group acting on a rank- s tensor, we have:

$$\begin{aligned} T'^{\mu_1 \dots \mu_s}(x) &= \exp\left(-\frac{i}{2}\omega_{\alpha\beta} I^{\alpha\beta}\right)_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} T^{\nu_1 \dots \nu_s}(x) \quad , \\ &= T^{\mu_1 \dots \mu_s}(x) - \frac{i}{2}\omega_{\alpha\beta} (I^{\alpha\beta})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} T^{\nu_1 \dots \nu_s}(x) + \mathcal{O}(\omega^2) \quad , \end{aligned} \quad (21)$$

where $I^{\alpha\beta}$ is the generator of Lorentz transformations which we are looking for. Comparing (21) with (20), we get an expression for the generators of Lorentz transformation:

$$(I^{\alpha\beta})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} = i \left(\prod_{i=1}^s \delta_{\nu_i}^{\mu_i} \right) (x^\beta \partial^\alpha - x^\alpha \partial^\beta) - i \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} (\eta^{\alpha\mu_k} \delta_{\nu_k}^\beta - \eta^{\beta\mu_k} \delta_{\nu_k}^\alpha) \quad . \quad (22)$$

This generator can be decomposed into an orbital angular momentum part and a spin part $I^{\alpha\beta} = L^{\alpha\beta} + S^{\alpha\beta}$, explicitly,

$$(L^{\alpha\beta})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} = i \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} (x^\beta \partial^\alpha - x^\alpha \partial^\beta) \quad , \quad (23)$$

$$(S^{\alpha\beta})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} = i \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} (\eta^{\beta\mu_k} \delta_{\nu_k}^\alpha - \eta^{\alpha\mu_k} \delta_{\nu_k}^\beta) \quad . \quad (24)$$

Now, we have all the generators of the Poincaré group on rank- s tensors, with s integer in $D = 3 + 1$ dimensions. Next, we generalize such results to the fermionic case represented by arbitrary rank- s spinor-tensors.

2.2 Arbitrary rank- s spinor-tensors in $D = 3 + 1$

In this section we obtain the generators of the Poincaré group on rank- s spinor-tensors. We denote spinor-tensors as $\psi^{\mu_1 \mu_2 \dots \mu_s}(x)$. Below, we derive the expressions of the generators proceeding as before.

2.2.1 Spacetime translations and Lorentz transformations

We start by requiring that the translated spinor-tensor at the translated spacetime coordinate is equal to the original spinor-tensor at the original spacetime coordinate:

$$\psi'^{\mu_1 \dots \mu_s}(x') = \psi^{\mu_1 \dots \mu_s}(x) \quad , \quad (25)$$

If we follow the same steps of Section 2.1.1, we end up with the following translation generator on rank- s spinor tensors:

$$(P^\alpha)_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} = -i \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} \partial^\alpha \mathbb{I} \quad . \quad (26)$$

where \mathbb{I} is a (4×4) identity matrix in the Dirac-spinor space. Except by the matrix \mathbb{I} , we have exactly the same result obtained in the pure tensor case (16).

Now, regarding the Lorentz transformations, there is a difference with respect to the bosonic case. Namely, the Lorentz transformation of a rank- s spinor tensor is given by

$$\psi'^{\mu_1 \dots \mu_s}(x') = \left(\prod_{i=1}^s \Lambda_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} \right) S \psi^{\nu_1 \dots \nu_s}(x) \quad , \quad (27)$$

where we take into account i) the tensor representation of the Lorentz group through the factors Λ acting directly on the Minkowski indices, and ii) the spinorial representation given by S , which is explicitly given by the group element:

$$S \equiv \exp\left(\frac{i}{4} \omega_{\alpha\beta} \sigma^{\alpha\beta}\right) \quad , \quad (28)$$

with the generators $\sigma_{\alpha\beta}$ given by

$$\sigma_{\alpha\beta} = -\frac{i}{2} [\gamma_\alpha, \gamma_\beta] \quad (29)$$

Notice that, in (28), the $\omega_{\alpha\beta}$ parameters are precisely the same infinitesimal antisymmetric parameters of the Lorentz transformations. However, the generators $\sigma_{\alpha\beta}$ are the corresponding Lorentz generators of the spinorial counterpart ($\sigma_{\alpha\beta}$ is explicitly antisymmetric). The symbol $[,]$ stands for commutator while the γ 's are the usual Dirac matrices.

On the one hand, expanding the left hand side of (27) and (28), using (18), we have:

$$\psi'^{\mu_1 \dots \mu_s}(x) - \omega^\gamma_\alpha x^\alpha \partial_\gamma \psi^{\mu_1 \dots \mu_s}(x) = \left[\prod_{i=1}^s \delta_{\nu_i}^{\mu_i} - \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} \omega_{\nu_k}^{\mu_k} + \dots \right] \left[\mathbb{I} + \frac{i}{4} \omega_{\alpha\beta} \sigma^{\alpha\beta} + \dots \right] \psi^{\nu_1 \dots \nu_s}(x) \dots \quad (30)$$

Recalling the antisymmetry of $\omega_{\alpha\beta}$, we have:

$$\begin{aligned} \psi'^{\mu_1 \dots \mu_s}(x) &= \psi^{\mu_1 \dots \mu_s}(x) + \frac{\omega_{\alpha\beta}}{2} \left[(x^\beta \partial^\alpha - x^\alpha \partial^\beta) \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} \mathbb{I} \right. \\ &\quad \left. - \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} (\eta^{\alpha\mu_k} \delta_{\nu_k}^\beta - \eta^{\beta\mu_k} \delta_{\nu_k}^\alpha) \mathbb{I} + \frac{i}{2} \sigma^{\alpha\beta} \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} \right] \psi^{\nu_1 \dots \nu_s}(x). \end{aligned} \quad (31)$$

On the other hand,

$$\begin{aligned} \psi'^{\mu_1 \dots \mu_s}(x) &= \exp\left(-\frac{i}{2} \omega_{\alpha\beta} I^{\alpha\beta}\right)_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} \psi^{\nu_1 \dots \nu_s}(x) \\ &= \psi^{\mu_1 \dots \mu_s}(x) - \frac{i}{2} \omega_{\alpha\beta} (I^{\alpha\beta})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} \psi^{\nu_1 \dots \nu_s}(x) \quad . \end{aligned} \quad (32)$$

Comparing (32) with (31), the generator for Lorentz transformations can be written as:

$$(I^{\alpha\beta})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} = i \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} (x^\beta \partial^\alpha - x^\alpha \partial^\beta) \mathbb{I} - i \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} (\eta^{\alpha\mu_k} \delta_{\nu_k}^\beta - \eta^{\beta\mu_k} \delta_{\nu_k}^\alpha) \mathbb{I} - \frac{1}{2} \sigma^{\alpha\beta} \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} \quad . \quad (33)$$

Again, this generator can be decomposed into an orbital angular momentum part and a spin part $I^{\alpha\beta} = L^{\alpha\beta} + S^{\alpha\beta}$ respectively given by:

$$(L^{\alpha\beta})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} = i \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} (x^\beta \partial^\alpha - x^\alpha \partial^\beta) \mathbb{I}, \quad (34)$$

$$(S^{\alpha\beta})_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} = -i \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} (\eta^{\alpha\mu_k} \delta_{\nu_k}^\beta - \eta^{\beta\mu_k} \delta_{\nu_k}^\alpha) \mathbb{I} - \frac{1}{2} \sigma^{\alpha\beta} \prod_{i=1}^s \delta_{\nu_i}^{\mu_i}, \quad (35)$$

where in the spin part $S^{\alpha\beta}$ of the total angular momentum we have an explicit separation of the integer – the first term of the right-hand side of (35)– as well as the semi-integer contributions – the second term of the right-hand side of (35).

We have, therefore, obtained representations of all Poincaré generators for arbitrary rank- s tensors and spinor-tensors. Next, we study the Casimir operators for this group.

2.3 Casimir operators of $\mathfrak{iso}(3, 1)$

The Casimir operators of a group are the operators that commute with all group generators. Consequently, these operators are invariant by the group symmetry transformations and are used to classify the irreducible representations associated with that group. According to the seminal work by Wigner in 1939 [1], mass and spin are precisely the two properties that characterize systems with invariance under the Poincaré group. Then, the irreducible representation of such a group is the basis of the description of elementary particles.

The Lie algebra $\mathfrak{iso}(3, 1)$ of the Poincaré group is composed of the generators P^μ and $I^{\alpha\beta}$. In $D = 3 + 1$, it is well known that the Poincaré algebra has two Casimirs,²

$$C_1 \equiv P^2 = P_\alpha P^\alpha, \quad (36)$$

$$C_2 \equiv W^2 = W_\alpha W^\alpha = -\frac{1}{2} I_{\alpha\beta} I^{\alpha\beta} P^2 + I_{\gamma\beta} I^{\alpha\beta} P_\alpha P^\gamma, \quad (37)$$

where W^α is the Pauli-Lubanski pseudovector, given explicitly by

$$W^\alpha = \epsilon^{\alpha\beta\gamma\delta} P_\beta I_{\gamma\delta} / 2, \quad (38)$$

and $I_{\gamma\delta}$ is given by (22) (for tensors) or (33) (for spinor-tensors). The corresponding eigenvalue equations,

$$P^2 \phi = -m^2 \phi, \quad (39)$$

$$W^2 \phi = -m^2 \mathfrak{s}(\mathfrak{s} + 1) \phi, \quad (40)$$

define the mass (m) and the spin (\mathfrak{s}) of an elementary massive particle represented (suppressing indices) by a field ϕ .

²From now on, we call a Casimir operator simply Casimir.

For the arbitrary rank- s tensor representation, we explicitly write the Casimir C_1 with the representation given by (16). Once the eigenvalue equation (39) must be satisfied, we obtain the Klein-Gordon equation in the given representation:

$$(\square - m^2)T^{\mu_1 \dots \mu_s} = 0. \quad (41)$$

Similarly, we write the eigenvalue equation of the Casimir C_2 using (16) and (22) to get

$$\begin{aligned} & 2s \square T^{\mu_1 \dots \mu_s} + 2 \sum_{\substack{k,r=1 \\ k < r}}^s \prod_{\substack{i=1 \\ i \neq k,r}}^s \delta_{\nu_i}^{\mu_i} \delta_{\nu_r}^{\mu_k} \delta_{\nu_k}^{\mu_r} \square T^{\nu_1 \dots \nu_s} - 2 \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} \partial_{\nu_k} \partial^{\mu_k} T^{\nu_1 \dots \nu_s} + \\ & + 2 \sum_{\substack{k,r=1 \\ k < r}}^s \prod_{\substack{i=1 \\ i \neq k,r}}^s \delta_{\nu_i}^{\mu_i} \left[\eta^{\mu_k \mu_r} \partial_{\nu_k} \partial_{\nu_r} - \eta_{\nu_k \nu_r} (\square \eta^{\mu_k \mu_r} - \partial^{\mu_k} \partial^{\mu_r}) - \delta_{\nu_k}^{\mu_r} \partial^{\mu_k} \partial_{\nu_r} - \delta_{\nu_r}^{\mu_k} \partial^{\mu_r} \partial_{\nu_k} \right] T^{\nu_1 \dots \nu_s} \\ & = m^2 \mathfrak{s}(\mathfrak{s} + 1) T^{\mu_1 \dots \mu_s}. \end{aligned} \quad (42)$$

Analogously, on arbitrary rank- s spinor-tensors, the eigenvalue equations are respectively:

$$(\square - m^2)\psi^{\mu_1 \dots \mu_s} = 0, \quad (43)$$

$$\begin{aligned} & \left(2s + \frac{3}{4}\right) \square \psi^{\mu_1 \dots \mu_s} + 2 \sum_{\substack{k,r=1 \\ k < r}}^s \prod_{\substack{i=1 \\ i \neq k,r}}^s \delta_{\nu_i}^{\mu_i} \delta_{\nu_r}^{\mu_k} \delta_{\nu_k}^{\mu_r} \square \psi^{\nu_1 \dots \nu_s} - 2 \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} \partial_{\nu_k} \partial^{\mu_k} \psi^{\nu_1 \dots \nu_s} + \\ & + 2 \sum_{\substack{k,r=1 \\ k < r}}^s \prod_{\substack{i=1 \\ i \neq k,r}}^s \delta_{\nu_i}^{\mu_i} \left[\eta^{\mu_k \mu_r} \partial_{\nu_k} \partial_{\nu_r} - \eta_{\nu_k \nu_r} (\square \eta^{\mu_k \mu_r} - \partial^{\mu_k} \partial^{\mu_r}) - \delta_{\nu_k}^{\mu_r} \partial^{\mu_k} \partial_{\nu_r} - \delta_{\nu_r}^{\mu_k} \partial^{\mu_r} \partial_{\nu_k} \right] \psi^{\nu_1 \dots \nu_s} \\ & - \frac{1}{2} \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta_{\nu_i}^{\mu_i} ([\not{\partial}, \gamma_{\nu_k}] \partial^{\mu_k} - [\not{\partial}, \gamma^{\mu_k}] \partial_{\nu_k} - [\gamma^{\mu_k}, \gamma_{\nu_k}] \square) \psi^{\nu_1 \dots \nu_s} \\ & = m^2 \mathfrak{s}(\mathfrak{s} + 1) \psi^{\mu_1 \dots \mu_s}, \end{aligned} \quad (44)$$

where $\not{\partial}$ denotes $\gamma_\mu \partial^\mu$.

Now that we have all the generators of the Poincaré group and the eigenvalue equations of the Casimir operators for rank- s tensors and spinor-tensors in $D = 3 + 1$, we proceed with a simple prescription that allows us to adapt our results to the lower dimension case of the $D = 2 + 1$ spacetime.

3 Poincaré group $ISO(2, 1)$

We start by taking advantage of that, in three spacetime dimensions, the 3 independent components of an antisymmetric tensor of rank two $I^{\mu\nu}$ can be written in terms of a three-component vector I_α as

$$I_\alpha = \frac{1}{2} \epsilon_{\alpha\mu\nu} I^{\mu\nu} \quad . \quad (45)$$

Then, the corresponding commutators of the Poincaré algebra $\mathfrak{iso}(2, 1)$ are

$$[I_\mu, I_\nu] = i\epsilon_{\mu\nu\rho} I^\rho \quad , \quad (46)$$

$$[I_\mu, P_\nu] = i\epsilon_{\mu\nu\rho} P^\rho \quad , \quad (47)$$

$$[P_\mu, P_\nu] = 0 \quad . \quad (48)$$

3.1 Arbitrary rank- s tensors and spinor-tensors in $D = 2 + 1$

We easily verify that the generators for spacetime translation for tensors (16) and spinor-tensors (26) in $D = 3 + 1$ preserve their expressions in $D = 2 + 1$.

The orbital angular momentum and the spin part are therefore

$$(L_\rho)^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s} = \frac{1}{2} \epsilon_{\rho\alpha\beta} (L^{\alpha\beta})^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s} \quad , \quad (49)$$

$$(S_\rho)^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s} = \frac{1}{2} \epsilon_{\rho\alpha\beta} (S^{\alpha\beta})^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s} \quad , \quad (50)$$

so that the generators (23) and (24) are generalized respectively to:

$$(L_\rho)^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s} = -i\epsilon_{\rho\alpha\beta} x^\alpha \partial^\beta \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} \quad , \quad (51)$$

$$(S_\rho)^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s} = i \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \epsilon^{\mu_k}_{\rho\nu_k} \delta_{\nu_i}^{\mu_i} \quad . \quad (52)$$

These are the representations of the generator of Lorentz transformations for rank- s tensors in $D = 2 + 1$, describing bosonic particles of mass m and helicity α .³

Proceeding in the very same way for the fermionic case, with generators (34) and (35), we obtain:

$$(L_\rho)^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s} = -i\epsilon_{\rho\alpha\beta} x^\alpha \partial^\beta \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} \mathbb{I} \quad , \quad (53)$$

$$(S_\rho)^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s} = i \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \epsilon^{\mu_k}_{\rho\nu_k} \delta_{\nu_i}^{\mu_i} \mathbb{I} + \frac{i}{2} \prod_{i=1}^s \delta_{\nu_i}^{\mu_i} \gamma_\rho \quad , \quad (54)$$

which are the representations of the corresponding generators of the Lorentz transformations on spinor-tensors in $D = 2 + 1$. Next, we explore a little more the Casimirs of the algebra $\mathfrak{iso}(2, 1)$ and their eigenvalue equations.

³Throughout this work we use the word helicity to refer to the “spin” of the particle in $D = 2 + 1$. Notice that, in $D = 3 + 1$, massive spin- s particles propagate $2s + 1$ degrees of freedom, while, in $D = 2 + 1$, they propagate helicity $+\alpha$ **and** $-\alpha$ in the case of parity preserving models (doublets of parity) and $+\alpha$ **or** $-\alpha$ in the case of parity breaking models (singlets of parity described by self-dual descriptions).

3.2 Casimirs of $\text{iso}(2, 1)$

As mentioned in the introduction, the classification of the unitary irreducible representations of the Poincaré group in three dimensions was for the very first time provided by Binégar in 1981 [19]. An interesting application of this study was given by Jackiw and Nair in 1991 [18], in which they studied one-particle states as unitary representations of the Poincaré group in three dimensions, aiming the description of *anyons*. In section II of [18], the authors provide the Poincaré algebra in $D = 2 + 1$ dimensions and its Casimirs:

- C_1 : P^2 , satisfying the eigenvalue equation $(P^2 + m^2)\phi = 0$ for a given one particle state ϕ , where m denotes the mass of a given one particle state represented by ϕ .
- C_2 : $P \cdot S$, the $D = 2+1$ version of the Pauli-Lubanski operator, satisfying $(P \cdot S - \alpha m)\phi = 0$, where α specifies the helicity of the particle.

In [18], the authors explored spin-1/2 and spin-1 representations for $ISO(2, 1)$. In the present section, we generalize such results for arbitrary rank- s tensors and spinor-tensors in the same way we have studied for $ISO(3, 1)$. The eigenvalue equations are:

$$C_1\phi \equiv P^\mu P_\mu \phi = -m^2 \phi \quad , \quad (55)$$

$$C_2\phi \equiv P^\mu S_\mu \phi = \alpha m \phi \quad , \quad (56)$$

where S_μ is given by (52) or (54). For arbitrary rank- s tensors, the explicit expressions for the eigenvalue equations (55) and (56) are respectively given by:

$$(\square - m^2)T^{\mu_1 \dots \mu_s} = 0 \quad , \quad (57)$$

$$\left(\sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s E^{\mu_k}_{\nu_k} \delta^{\mu_i}_{\nu_i} \right) T^{\nu_1 \dots \nu_s} = \alpha m T^{\mu_1 \dots \mu_s} \quad . \quad (58)$$

where we have defined the operator $E^\mu_\nu \equiv \epsilon^{\mu\rho} \nu \partial_\rho$. The equation (58) is the Pauli-Lubanski equation for an arbitrary rank- s field in the tensor representation.

Analogously, in the spinor-tensor representation, using (26) and (54), we have:

$$(\square - m^2)\psi^{\mu_1 \dots \mu_s} = 0 \quad , \quad (59)$$

$$\left(\sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s E^{\mu_k}_{\nu_k} \delta^{\mu_i}_{\nu_i} \mathbb{I} + \frac{1}{2} \prod_{i=1}^s \delta^{\mu_i}_{\nu_i} \gamma_\rho \partial^\rho \right) \psi^{\nu_1 \dots \nu_s} - \alpha m \psi^{\mu_1 \dots \mu_s} = 0 \quad . \quad (60)$$

In the next section, we demonstrate that it is possible to obtain the Fierz-Pauli constraints from the Casimir eigenvalue equations.

4 Fierz-Pauli constraints from the algebra

As we increase the spin of the massive particle, in order to propagate the correct number of physical degrees of freedom in a given representation, higher-rank tensors or spinor-tensors are required. However, higher-rank tensors introduce spurious field components that must be eliminated. This is possible by introducing constraint equations. For massive particles of integer spin- s , represented by arbitrary rank- s bosonic fields, those constraints are given by the Fierz-Pauli constraints [4]:

$$T^{\mu_1 \dots \mu_s} = T^{(\mu_1 \dots \mu_s)}, \quad (61)$$

$$\eta_{\mu_1 \mu_2} T^{\mu_1 \mu_2 \dots \mu_s} = 0, \quad (62)$$

$$\partial_{\mu_1} T^{\mu_1 \mu_2 \dots \mu_s} = 0. \quad (63)$$

For semi-integer spin, represented by arbitrary rank- s fermionic fields, they are:

$$\psi^{\mu_1 \dots \mu_s} = \psi^{(\mu_1 \dots \mu_s)}, \quad (64)$$

$$\eta_{\mu_1 \mu_2} \psi^{\mu_1 \mu_2 \dots \mu_s} = 0, \quad (65)$$

$$\partial_{\mu_1} \psi^{\mu_1 \mu_2 \dots \mu_s} = 0, \quad (66)$$

$$\gamma_{\mu_1} \psi^{\mu_1 \mu_2 \dots \mu_s} = 0. \quad (67)$$

The derivation of Fierz-Pauli constraints from a given massive higher-spin model is typically not a straightforward task, as it requires working through the equations of motion in order to demonstrate that the constraints hold and all lower-rank auxiliary fields vanish on-shell. However, in [8], the authors have achieved a model-independent derivation of the Fierz-Pauli constraints for spins $1/2$, 1 , $3/2$, and 2 in $D = 3 + 1$. In the present work, we extend these results to include bosonic cases of spin-3 and spin-4 particles, along with the fermionic example of spin-5/2 in $D = 3 + 1$. We also explore the generalization of our findings to $D = 2 + 1$ for helicities $5/2, 3, 4$ and pave the way for a demonstration for arbitrary integer spin (helicity).

4.1 Spin-5/2

In this section, we obtain the Fierz-Pauli constraints for a rank-2 spinor-tensor representation of a spin-5/2 massive particle in $D = 3+1$ and $D = 2+1$, from the eigenvalue equations provided by the Casimirs C_1 and C_2 .

4.1.1 Spin-5/2 in $D = 3 + 1$:

For a rank-2 spinor-tensor, the first Casimir operator C_1 has the Klein-Gordon equation as its eigenvalue equation:

$$\square \psi^{\mu\nu} = m^2 \psi^{\mu\nu}. \quad (68)$$

Setting $s = 2$ and $\mathfrak{s} = 5/2$ in (44) and making use of the Klein-Gordon equation (68), we find after some manipulation, the following expression for the Pauli-Lubanski equation:

$$\begin{aligned}
& m^2(-2\psi^{\mu\nu} - \eta^{\mu\nu}\psi + \psi^{\nu\mu}) + \partial^\mu\partial^\nu\psi + \eta^{\mu\nu}\partial_\alpha\partial_\beta\psi^{\alpha\beta} + \\
& - \partial^\mu\partial_\alpha\psi^{\alpha\nu} - \partial^\nu\partial_\beta\psi^{\mu\beta} - \partial^\mu\partial_\beta\psi^\nu{}_\beta - \partial^\nu\partial_\alpha\psi^{\alpha\mu} + \\
& - \frac{1}{4}([\not{\partial}, \gamma_\alpha]\partial^\mu - [\not{\partial}, \gamma^\mu]\partial_\alpha - m^2[\gamma^\mu, \gamma_\alpha])\psi^{\alpha\nu} + \\
& - \frac{1}{4}([\not{\partial}, \gamma_\beta]\partial^\nu - [\not{\partial}, \gamma^\nu]\partial_\beta - m^2[\gamma^\nu, \gamma_\beta])\psi^{\mu\beta} = 0,
\end{aligned} \tag{69}$$

where ψ denote $\eta_{\mu\nu}\psi^{\mu\nu}$.

The following contractions will result in the Fierz-Pauli constraints:

$$\partial^2(69) \Rightarrow \partial_\mu\partial_\nu\psi^{\mu\nu} = 0, \tag{70}$$

$$\gamma\partial(69) \Rightarrow \gamma_\nu\partial_\mu\psi^{\mu\nu} = \gamma_\mu\partial_\nu\psi^{\mu\nu} = 0, \tag{71}$$

$$\partial(69) \Rightarrow \partial_\mu\psi^{\mu\nu} = \partial_\nu\psi^{\mu\nu} = 0, \tag{72}$$

$$\eta(69) \Rightarrow \psi = 0, \tag{73}$$

$$\gamma\gamma(69)_a \Rightarrow \gamma_\mu\gamma_\nu\psi_a^{\mu\nu} = 0, \tag{74}$$

$$\gamma(69)_a \Rightarrow \gamma_\mu\psi_a^{\mu\nu} = \gamma_\nu\psi_a^{\mu\nu} = 0 \Rightarrow \psi_{a\mu\nu} = 0, \tag{75}$$

$$\gamma\gamma(69) \Rightarrow \gamma_\mu\gamma_\nu\psi^{\mu\nu} = 0, \tag{76}$$

$$\gamma(69) \Rightarrow \gamma_\mu\psi^{\mu\nu} = \gamma_\nu\psi^{\mu\nu} = 0. \tag{77}$$

In the equations (74) and (75) the subscript “a” means that we are taking only the anti-symmetric part of $\psi^{\mu\nu}$, i.e., $\psi_a^{\mu\nu} = (\psi^{\mu\nu} - \psi^{\nu\mu})/2$.

With the equations (72), (73), (75) and (77) in hands, we complete the demonstration of the Fierz-Pauli constraints (64) - (67) for a rank-2 spinor-tensor in $D = 3 + 1$, obtained from the Casimir equations.

4.1.2 Helicity-5/2 in $D = 2 + 1$

For a rank-2 spinor-tensor with helicity $\alpha = 5/2$, the expression (59) yields the Klein-Gordon equation while (60) provides the Pauli-Lubanski equation in $D = 2 + 1$:

$$E^\mu{}_\rho\psi^{\rho\nu} + E^\nu{}_\rho\psi^{\mu\rho} + \frac{1}{2}\not{\partial}\psi^{\mu\nu} = \frac{5}{2}m\psi^{\mu\nu}. \tag{78}$$

First, applying $\not{\partial}$ to equation (78) and using (78) itself in the terms with $\not{\partial}\psi^{\alpha\beta}$, we have:

$$9m^2\psi^{\mu\nu} + 2\partial^\mu\partial_\lambda\psi^{\lambda\nu} + 2\partial^\nu\partial_\lambda\psi^{\mu\lambda} - E^\mu{}_\rho E^\nu{}_\lambda\psi^{\rho\lambda} = 5m\not{\partial}\psi^{\mu\nu}. \tag{79}$$

Then, as we have done in $D = 3 + 1$, we suggest some steps in order to obtain the Fierz-Pauli constraints, but now it is always necessary to make use of the Klein-Gordon equation and

equations (78) and (79). Combining these equations, we have:

$$\partial^2(78); \partial^2(79) \Rightarrow \partial_\mu \partial_\nu \psi^{\mu\nu} = 0, \quad (80)$$

$$\eta(78); \eta(79) \Rightarrow \eta_{\mu\nu} \psi^{\mu\nu} = 0, \quad (81)$$

$$\partial\gamma(78); \partial\gamma(79) \Rightarrow \partial_\mu \gamma_\nu \psi^{\mu\nu} = \partial_\nu \gamma_\mu \psi^{\mu\nu} = 0, \quad (82)$$

$$\gamma^2(78); \gamma^2(79) \Rightarrow \gamma_\mu \gamma_\nu \psi^{\mu\nu} = \gamma_\nu \gamma_\mu \psi^{\mu\nu} = 0, \quad (83)$$

$$\partial(78); \partial(79) \Rightarrow \partial_\mu \psi^{\mu\nu} = \partial_\nu \psi^{\mu\nu} = 0, \quad (84)$$

$$\gamma(78); \gamma(79) \Rightarrow \gamma_\mu \psi^{\mu\nu} = \gamma_\nu \psi^{\mu\nu} = 0. \quad (85)$$

Notice that after obtaining the transversality condition and using the property $E_{\mu\nu}E_{\alpha\beta} = \square(\theta_{\mu\beta}\theta_{\nu\alpha} - \theta_{\mu\alpha}\theta_{\nu\beta})$ the equation (79) reduces to a Dirac equation:

$$(\not{\partial} - m)\psi^{\mu\nu} = 0. \quad (86)$$

In order to demonstrate that $\psi^{\mu\nu}$ is symmetric, we apply $\epsilon_{\mu\nu\alpha}$ to (78), which gives us:

$$\epsilon_{\mu\nu\alpha}\psi^{\mu\nu} = 0 \Rightarrow \psi^{\mu\nu} = \psi^{\nu\mu}. \quad (87)$$

With this result and the conditions (81), (84), and (85) the Fierz-Pauli constraints are all verified from the Casimir equations.

4.2 Spin-3

Next, we obtain the Fierz-Pauli constraints for a rank-3 tensor representation of a spin-3 massive particle in $D = 3 + 1$ and $D = 2 + 1$, from the Casimir equations.

4.2.1 Spin-3 in $D = 3 + 1$

The Casimirs C_1 and C_2 are given respectively by (36) and (37),

$$C_1 \equiv P^2 = P_\alpha P^\alpha \Rightarrow (\square - m^2)T^{\mu\nu\gamma} = 0, \quad (88)$$

$$C_2 \equiv W^2 = W_\alpha W^\alpha \Rightarrow W^2 T^{\mu\nu\gamma} = m^2 \mathfrak{s}(\mathfrak{s} + 1)T^{\mu\nu\gamma} = 12m^2 T^{\mu\nu\gamma}. \quad (89)$$

Using (38) explicitly in the expression (89) and (88), we find after some manipulation:

$$\begin{aligned} & 6m^2 T^{\mu\nu\gamma} + 2m^2 (\eta^{\nu\gamma} T^{\mu\sigma}{}_\sigma + \eta^{\mu\gamma} T^{\sigma\nu}{}_\sigma + \eta^{\mu\nu} T^{\sigma}{}_\sigma{}^\gamma) - 2m^2 (T^{\mu\gamma\nu} + T^{\gamma\nu\mu} + T^{\nu\mu\gamma}) + \\ & + 2(\partial_\alpha \partial^\gamma T^{\mu\nu\alpha} + \partial_\alpha \partial^\nu T^{\mu\alpha\gamma} + \partial_\alpha \partial^\mu T^{\alpha\nu\gamma}) - 2(\eta^{\nu\gamma} \partial_\alpha \partial_\sigma T^{\mu\alpha\sigma} + \eta^{\mu\gamma} \partial_\alpha \partial_\sigma T^{\alpha\nu\sigma} + \eta^{\mu\nu} \partial_\alpha \partial_\sigma T^{\alpha\sigma\gamma}) + \\ & - 2\partial^\nu \partial^\gamma T^{\mu\sigma}{}_\sigma - 2\partial^\mu \partial^\gamma T^{\sigma\nu}{}_\sigma - 2\partial^\mu \partial^\nu T^{\sigma}{}_\sigma{}^\gamma + 2\partial_\alpha \partial^\gamma T^{\mu\alpha\nu} + 2\partial_\alpha \partial^\gamma T^{\alpha\nu\mu} + \\ & + 2\partial_\alpha \partial^\nu T^{\mu\gamma\alpha} + 2\partial_\alpha \partial^\nu T^{\alpha\mu\gamma} + 2\partial_\alpha \partial^\mu T^{\nu\alpha\gamma} + 2\partial_\alpha \partial^\mu T^{\gamma\nu\alpha} = 0. \end{aligned} \quad (90)$$

Applying $\eta_{\nu\gamma}$, $\eta_{\mu\nu}$, and $\eta_{\mu\gamma}$, respectively, on (90), and taking one derivative of the resulting equations, we show that

$$T^{\mu\sigma}{}_\sigma = 0 \quad ; \quad T^{\sigma}{}_\sigma{}^\gamma = 0 \quad ; \quad T^{\sigma\nu}{}_\sigma = 0. \quad (91)$$

Inserting the results (91) in (90) and applying $\partial_\mu\partial_\nu\partial_\gamma$ in the obtained expression we demonstrate that $\partial_\mu\partial_\nu\partial_\gamma T^{\mu\nu\gamma} = 0$. Next, we apply $\partial_\mu\partial_\gamma$, $\partial_\mu\partial_\nu$, and $\partial_\nu\partial_\gamma$, respectively, on (90) to show that $\partial_\mu\partial_\gamma T^{\mu\nu\gamma} = 0$, $\partial_\mu\partial_\nu T^{\mu\nu\gamma} = 0$, and $\partial_\nu\partial_\gamma T^{\mu\nu\gamma} = 0$. Finally, applying ∂_μ , ∂_ν , and ∂_γ , respectively, we get the transversality conditions:

$$\partial_\mu T^{\mu\nu\gamma} = 0 \quad ; \quad \partial_\nu T^{\mu\nu\gamma} = 0 \quad ; \quad \partial_\gamma T^{\mu\nu\gamma} = 0. \quad (92)$$

Then, the expression (90) can be rewritten as:

$$3T^{\mu\nu\gamma} = T^{\mu\gamma\nu} + T^{\nu\mu\gamma} + T^{\gamma\nu\mu}. \quad (93)$$

By taking differences of (93) and relabelling some indices we obtain the following system

$$\begin{aligned} 4t_1^{\mu\nu\gamma} - t_2^{\gamma\nu\mu} - t_3^{\mu\gamma\nu} &= 0, \\ -t_1^{\mu\nu\gamma} + 4t_2^{\gamma\nu\mu} + t_3^{\mu\gamma\nu} &= 0, \\ -t_1^{\mu\nu\gamma} + t_2^{\gamma\nu\mu} + 4t_3^{\mu\gamma\nu} &= 0, \end{aligned} \quad (94)$$

where $t_1^{\mu\nu\gamma} \equiv T^{\mu\nu\gamma} - T^{\nu\mu\gamma}$, $t_2^{\gamma\nu\mu} \equiv T^{\gamma\nu\mu} - T^{\gamma\mu\nu}$ and $t_3^{\mu\gamma\nu} \equiv T^{\nu\gamma\mu} - T^{\mu\gamma\nu}$. Since the numerical matrix corresponding to the linear system (94) has a non-vanishing determinant, the only solution of (94) is the trivial one: $t_j^{\mu\nu\rho} = 0$, $j = 1, 2, 3$. Consequently, we end up with a totally symmetric tensor

$$T^{\mu\nu\gamma} = T^{(\mu\nu\gamma)}. \quad (95)$$

In this way, we deduce the Fierz-Pauli constraints for rank-3 tensors, i.e., the tensor is fully symmetric (95), traceless (91) and transverse (92). These constraints reduce the degrees of freedom of the arbitrary tensor $T^{\mu\nu\gamma}$ from 64 to 7, in agreement with the $2\mathfrak{s} + 1$ rule for the number of degrees of freedom of a massive spin- \mathfrak{s} particle in $D = 3 + 1$ at $\mathfrak{s} = 3$.

4.2.2 Helicity-3 in $D = 2 + 1$

For a rank-3 tensor with helicity $\alpha = 3$, the Pauli-Lubanski equation (58) is given by:

$$E_\gamma{}^\rho T^{\mu\nu\gamma} + E_\beta{}^\nu T^{\mu\beta\rho} + E_\alpha{}^\mu T^{\alpha\nu\rho} = 3mT^{\mu\nu\rho}. \quad (96)$$

Using the Klein-Gordon equation in (96), we can demonstrate the Fierz-Pauli constraints after some manipulations. We summarize the steps as follows:

$$\partial^3(96) \Rightarrow \partial_\mu\partial_\nu\partial_\rho T^{\mu\nu\rho} = 0 \quad (97)$$

$$\partial^2(96) \Rightarrow \partial_\mu\partial_\nu T^{\mu\nu\rho} = \partial_\mu\partial_\rho T^{\mu\nu\rho} = \partial_\rho\partial_\nu T^{\mu\nu\rho} = 0 \quad (98)$$

$$\partial(96) \Rightarrow \partial_\mu T^{\mu\nu\rho} = \partial_\nu T^{\mu\nu\rho} = \partial_\rho T^{\mu\nu\rho} = 0 \quad (99)$$

$$\eta(96) \Rightarrow T^\mu{}_\mu{}^\rho = T^{\mu\nu}{}_\mu = T^{\mu\nu}{}_\nu = 0 \quad (100)$$

where, repeated use of (96) has been done. Contracting $E^\gamma{}_\mu$, $E^\gamma{}_\nu$ and $E^\gamma{}_\rho$, one by one, with the (96) and reinserting the three resulting equations in (96), we obtain:

$$3T^{\mu\nu\gamma} = T^{\mu\gamma\nu} + T^{\nu\mu\gamma} + T^{\gamma\nu\mu} \quad (101)$$

that is the same equation (93), so performing the same steps of the previous section we conclude that the tensor is totally symmetric. This completes the demonstration of the Fierz-Pauli constraints for rank-3 tensors in $D = 2 + 1$.

4.3 Spin-4

Here we obtain the Fierz-Pauli constraints for a rank-4 tensor representation of a spin-4 massive particle in $D = 3 + 1$ and $D = 2 + 1$, from the Casimir equations.

4.3.1 Spin-4 in $D = 3 + 1$

As usual, the first Casimir operator C_1 yields the Klein-Gordon equation:

$$\square T^{\mu\nu\alpha\beta} = m^2 T^{\mu\nu\alpha\beta}. \quad (102)$$

In turn, for $s = 4$ and $\mathfrak{s} = 4$, we rewrite the Pauli-Lubanski eigenvalue equation in the form:

$$m^2 A^{\mu\nu\alpha\beta} + B^{\mu\nu\alpha\beta} = 0, \quad (103)$$

where

$$A^{\mu\nu\alpha\beta} = 6 T^{\mu\nu\alpha\beta} - (T^{\nu\mu\alpha\beta} + T^{\alpha\nu\mu\beta} + T^{\beta\nu\alpha\mu} + T^{\mu\alpha\nu\beta} + T^{\mu\beta\alpha\nu} + T^{\mu\nu\beta\alpha}), \quad (104)$$

while

$$\begin{aligned} B^{\mu\nu\alpha\beta} = & \partial^\mu \partial_\gamma T^{\gamma\nu\alpha\beta} + \partial^\mu \partial_\gamma T^{\nu\gamma\alpha\beta} + \partial^\mu \partial_\gamma T^{\alpha\nu\gamma\beta} + \partial^\mu \partial_\gamma T^{\beta\nu\alpha\gamma} + \partial^\nu \partial_\gamma T^{\gamma\mu\alpha\beta} + \partial^\nu \partial_\gamma T^{\mu\gamma\alpha\beta} \\ & + \partial^\nu \partial_\gamma T^{\mu\alpha\gamma\beta} + \partial^\nu \partial_\gamma T^{\mu\beta\alpha\gamma} + \partial^\alpha \partial_\gamma T^{\gamma\nu\mu\beta} + \partial^\alpha \partial_\gamma T^{\mu\gamma\nu\beta} + \partial^\alpha \partial_\gamma T^{\mu\nu\gamma\beta} + \partial^\alpha \partial_\gamma T^{\mu\nu\beta\gamma} \\ & + \partial^\beta \partial_\gamma T^{\gamma\nu\alpha\mu} + \partial^\beta \partial_\gamma T^{\mu\gamma\alpha\nu} + \partial^\beta \partial_\gamma T^{\mu\nu\gamma\alpha} + \partial^\beta \partial_\gamma T^{\mu\nu\alpha\gamma} \\ & - \partial_\gamma \partial_\sigma (\eta^{\mu\nu} T^{\gamma\sigma\alpha\beta} + \eta^{\mu\alpha} T^{\gamma\nu\sigma\beta} + \eta^{\mu\beta} T^{\gamma\nu\alpha\sigma} + \eta^{\nu\alpha} T^{\mu\gamma\sigma\beta} + \eta^{\nu\beta} T^{\mu\gamma\alpha\sigma} + \eta^{\alpha\beta} T^{\mu\nu\gamma\sigma}) \\ & - m^2 (\theta^{\mu\nu} T^{\gamma\alpha\beta} + \theta^{\mu\alpha} T^{\gamma\nu\beta} + \theta^{\mu\beta} T^{\gamma\nu\alpha} + \theta^{\nu\alpha} T^{\mu\gamma\beta} + \theta^{\nu\beta} T^{\mu\gamma\alpha} + \theta^{\alpha\beta} T^{\mu\nu\gamma}). \end{aligned} \quad (105)$$

In order to verify the Fierz-Pauli constraints, we perform a series of contractions and operations starting from (103), always relying on (102). Due to the large number of contraction possibilities offered by the four indices of the tensor, the required procedure for the demonstration is laborious and typically involves the use of systems of equations that are obtained from different manipulations of (103). We summarize the steps as follows:

$$\partial^4 (103) \Rightarrow \partial_\mu \partial_\nu \partial_\alpha \partial_\beta T^{\mu\nu\alpha\beta} = 0 \quad (106)$$

$$\partial^3 (103) \Rightarrow \partial_\mu \partial_\nu \partial_\alpha T^{\mu\nu\alpha\beta} = \partial_\mu \partial_\nu \partial_\beta T^{\mu\nu\alpha\beta} = \partial_\mu \partial_\alpha \partial_\beta T^{\mu\nu\alpha\beta} = \partial_\nu \partial_\alpha \partial_\beta T^{\mu\nu\alpha\beta} = 0 \quad (107)$$

$$\partial^2 \eta (103) \Rightarrow \partial_\mu \partial_\nu T^{\mu\nu\gamma} = \partial_\mu \partial_\nu T^{\mu\gamma\nu} = 0 = \dots \quad (108)$$

$$\partial^2 (103) \Rightarrow \partial_\mu \partial_\nu T^{\mu\nu\alpha\beta} = \partial_\mu \partial_\nu T^{\mu\alpha\nu\beta} = 0 = \dots \quad (109)$$

$$\partial \eta (103) \Rightarrow \partial_\mu T^{\mu\beta\gamma} = \partial_\mu T^{\mu\gamma\beta} = \partial_\mu T^{\beta\mu\gamma} = 0 = \dots \quad (110)$$

$$\eta \eta (103) \Rightarrow T^{\gamma\alpha}{}_\alpha = T^{\gamma\alpha}{}_{\gamma\alpha} = T^{\gamma\alpha}{}_{\alpha\gamma} = 0 \quad (111)$$

$$\eta (103) \Rightarrow T^{\gamma\alpha\beta}{}_\gamma = T^{\alpha\beta\gamma}{}_\gamma = 0 = \dots \quad (112)$$

$$\partial (103) \Rightarrow \partial_\mu T^{\mu\nu\alpha\beta} = \partial_\mu T^{\nu\mu\alpha\beta} = 0 = \dots, \quad (113)$$

where, in each step, we feedback equation (103) and after that it becomes:

$$6 T^{\mu\nu\alpha\beta} = T^{\nu\mu\alpha\beta} + T^{\alpha\nu\mu\beta} + T^{\beta\nu\alpha\mu} + T^{\mu\alpha\nu\beta} + T^{\mu\beta\alpha\nu} + T^{\mu\nu\beta\alpha}. \quad (114)$$

Similarly to the spin-3 case, it is convenient to define six partially antisymmetric tensors:

$$\begin{aligned} t_1^{\mu\nu\alpha\beta} &= T^{\mu\nu\alpha\beta} - T^{\nu\mu\alpha\beta} & ; & & t_4^{\alpha\mu\nu\beta} &= T^{\alpha\mu\nu\beta} - T^{\alpha\nu\mu\beta} \\ t_2^{\nu\alpha\mu\beta} &= T^{\nu\alpha\mu\beta} - T^{\mu\alpha\nu\beta} & ; & & t_5^{\beta\mu\alpha\nu} &= T^{\beta\mu\alpha\nu} - T^{\beta\nu\alpha\mu} \\ t_3^{\nu\beta\alpha\mu} &= T^{\nu\beta\alpha\mu} - T^{\mu\beta\alpha\nu} & ; & & t_6^{\beta\alpha\mu\nu} &= T^{\beta\alpha\mu\nu} - T^{\beta\alpha\nu\mu} \end{aligned} \quad (115)$$

By taking differences of (114) we derive, for instance,

$$7t_1^{\mu\nu\alpha\beta} + t_1^{\mu\nu\beta\alpha} + t_2^{\nu\alpha\mu\beta} + t_3^{\nu\beta\alpha\mu} + t_4^{\alpha\mu\nu\beta} + t_5^{\beta\mu\alpha\nu} = 0. \quad (116)$$

Notice that, in principle, $t_1^{\mu\nu\alpha\beta} \neq t_1^{\mu\nu\beta\alpha}$. Therefore, the exchange $\alpha \leftrightarrow \beta$ in (115) leads to a total of 12 partially antisymmetric rank-4 tensors. Taking differences of (114) and relabelling indices we can derive 12 equations for those 12 tensors. As in the spin-3 case, the determinant of the numerical 12×12 matrix of the corresponding linear system is different than zero, and the trivial solution $t_j^{\gamma\delta\rho\lambda} = 0$ for $j = 1, 2, \dots, 12$ is the only possible one. This immediately implies that the tensor is symmetric under the exchange of any pair of indices,

$$T^{\gamma\delta\rho\lambda} = T^{(\gamma\delta\rho\lambda)} \quad . \quad (117)$$

Thus, together with (112) and (113), we have demonstrated the Fierz-Pauli constraints for rank-4 tensors in $D = 3 + 1$, obtained from the Casimir eigenvalue equations.

4.3.2 Helicity-4 in $D = 2 + 1$

For an arbitrary rank-4 tensor, the equation (58) is given by:

$$E_\lambda^\mu T^{\lambda\nu\alpha\beta} + E_\lambda^\nu T^{\mu\lambda\alpha\beta} + E_\lambda^\alpha T^{\mu\nu\lambda\beta} + E_\lambda^\beta T^{\mu\nu\alpha\lambda} = 4 m T^{\mu\nu\alpha\beta}. \quad (118)$$

Similarly to what was done in $D = 3 + 1$, we will describe the main steps in the search for the Fierz-Pauli constraints, namely:

$$\partial^3 (118) \Rightarrow \partial_\mu \partial_\nu \partial_\alpha \partial_\beta T^{\mu\nu\alpha\beta} = 0 \quad (119)$$

$$\partial^3 (118) \Rightarrow \partial_\mu \partial_\nu \partial_\alpha T^{\mu\nu\alpha\beta} = 0 \quad (120)$$

$$\partial^2 \eta (118) \Rightarrow \partial_\mu \partial_\nu T^{\mu\nu\gamma} = 0 \quad (121)$$

$$E \partial^2 (118) \Rightarrow \partial_\mu \partial_\nu T^{\mu\nu\alpha\beta} = 0 \quad (122)$$

$$\partial \eta (118) \Rightarrow \partial_\mu T^{\mu\nu\gamma} = 0 \quad (123)$$

$$E \eta \eta (118) \Rightarrow T^\mu{}_\mu{}^\gamma{}_\gamma = 0 \quad (124)$$

$$E \eta (118) \Rightarrow T^{\mu\nu\gamma}{}_\gamma = 0 \quad (125)$$

$$E \partial (118) \Rightarrow \partial_\mu T^{\mu\nu\alpha\beta} = 0 \quad (126)$$

It is worth mentioning that we always use the Klein-Gordon equation $\square T^{\mu\nu\alpha\beta} = m^2 T^{\mu\nu\alpha\beta}$ whenever it is possible as well as the equation (118) itself is reused. We have the following steps Contracting E_μ^ρ , E_ν^ρ , E_α^ρ , and E_β^ρ with (118) and reinserting the obtained equations into (118), we have:

$$6 T^{\mu\nu\alpha\beta} = T^{\nu\mu\alpha\beta} + T^{\alpha\nu\mu\beta} + T^{\beta\nu\alpha\mu} + T^{\mu\alpha\nu\beta} + T^{\mu\beta\alpha\nu} + T^{\mu\nu\beta\alpha} \quad (127)$$

which is the same equation (114) obtained in $D = 3+1$. This last result demonstrates, therefore, that the tensor is symmetric with respect to any pair of indices. We have then demonstrated that the Fierz-Pauli constraints for a rank-4 tensor also follow from the eigenvalue equations of the Casimir operators of the Poincaré group in $D = 2 + 1$.

4.4 Comments on the case of arbitrary rank-s tensors

We have observed that in the $D = 3 + 1$, for example in the case of spin-4, only the part $A^{\mu\nu\alpha\beta}$ survived in (103) after deducing the transversality and traceless conditions. From this, we conclude that the tensor is completely symmetric. In general, for arbitrary rank-s, we would obtain:

$$m^2 A^{\mu_1 \dots \mu_s} + B^{\mu_1 \dots \mu_s} = 0. \quad (128)$$

The procedure to be followed when we are demonstrating the Fierz-Pauli constraints is then to show that $B^{\mu_1 \dots \mu_s} = 0$. To do this, we must apply s derivatives to (128) in order to demonstrate that $\partial_{\mu_1} \dots \partial_{\mu_s} T^{\mu_1 \dots \mu_s} = 0$. Then, we apply $(s-1)$ derivatives or $(s-2)$ derivatives plus the metric in order to eliminate terms of the form $\partial_{\mu_1} \dots \partial_{\mu_{s-1}} T^{\mu_1 \dots \mu_s}$ and $T^{\mu_1 \dots \gamma \gamma}$.

Notice that we first demonstrate that the scalars obtained from (128) are zero, and then the vectors and then rank-2 tensors, and so on. The idea is to gradually increase the rank of the objects. The difficulty arises because as the rank of the tensor increases, the number of possible contractions also increases, and the result is that the demonstration relies on increasingly larger systems of homogeneous equations.

The ultimate goal is to demonstrate that the trace over an arbitrary pair of indices and a contracted derivative with some index of the tensor is zero. At this stage, we will be dealing with objects of rank $(s-2)$ and $(s-1)$ respectively. In this way, (128) reduces to $A^{\mu_1 \dots \mu_s} = 0$ which can be written as:

$$\binom{s}{2} T^{\mu_1 \mu_2 \dots \mu_s} = T^{\mu_2 \mu_1 \dots \mu_s} + T^{\mu_3 \mu_2 \mu_1 \dots \mu_s} + \dots + T^{\mu_s \mu_2 \dots \mu_1} + \dots + T^{\mu_1 \mu_2 \dots \mu_s \mu_{s-1}}, \quad (129)$$

where there are exactly $\binom{s}{2}$ terms on the right-hand side of the equation, corresponding to the simple interchange of two indices of the tensor $T^{\mu_1 \dots \mu_s}$ on the left-hand side of the equation.

To demonstrate that the tensor is completely symmetric, we must permute the indices of (129), combining them to form antisymmetric tensors of the type $T^{[\mu_1 \mu_2] \dots \mu_s}$, as in equation (116). Just as it was done for spin-4, we must construct $(s-2)! \binom{s}{2} = \frac{s!}{2}$ equations to account for

the same number of variables (the number of ways to select two positions for the antisymmetric pair and then permute the remaining $(s - 2)$ positions). We then construct the matrix that represents this system⁴ of homogeneous equations and calculate the determinant. It must be verified that the determinant is nonzero, so that the solution of the system is unique and trivial, meaning that each term $T^{[\mu\nu]\dots\mu_s}$ is zero. This proves that the tensor is symmetric with respect to any pair of its indices. Therefore, it is concluded that the Pauli-Lubanski equation, together with the Klein-Gordon equation, both derived from first principles with the help of the eigenvalue equations of the Casimir operators of the Poincaré group, implies that the rank- s tensor must satisfy the Fierz-Pauli constraints and thus has precisely $2s + 1$ degrees of freedom.

Some comments on the case of $D = 2 + 1$ dimensions are also in order. We can notice that, in (96) the operator E acts on each of the indices of the tensor $T^{\mu\nu\rho}$. The same occurs in the equation for the rank-4 tensors. We conjecture that if we work with an arbitrary tensor of s indices, we will have the equation:

$$E_\alpha^{\mu_1} T^{\alpha\mu_2\dots\mu_s} + E_\alpha^{\mu_2} T^{\mu_1\alpha\mu_3\dots\mu_s} + \dots E_\alpha^{\mu_s} T^{\mu_1\mu_2\dots\mu_{(s-1)}\alpha} = \alpha m T^{\mu_1\mu_2\dots\mu_s}. \quad (130)$$

This is a necessary condition that tensors must satisfy in order to describe fields of helicity $+\alpha$ or $-\alpha$. However, it is not sufficient on its own as it does not provide an equation that is identical to the equation of motion of a massive self-dual model. It must be accompanied by the Klein-Gordon equation, which, in our work, is given by the first Casimir.

5 Conclusions

In this work, we have investigated the description of massive higher spin particles in $D = 3 + 1$ and $D = 2 + 1$ dimensions in a model-independent way. We derive the Fierz-Pauli constraints starting from algebraic properties of the Poincaré groups $ISO(3, 1)$ and $ISO(2, 1)$, respectively.

The derivation of the Fierz-Pauli constraints from the eigenvalue equations of the Casimir operators of the Poincaré algebra seems to have been employed for the first time by [8]. There, the authors have worked out specifically the spin cases $s = 1/2, 1, 3/2$ and $s = 2$ in $D = 3 + 1$. They have conjectured that their results could be extended to higher spins. It is important to stress that in the higher spin cases ($s \geq 5/2$) the corresponding massless models (Fronsdal and Fang-Fronsdal models [33, 34]) are invariant only under constrained gauge transformations where the gauge parameters must be traceless (bosons) or γ -traceless (fermions). So one might expect eventually that the model-independent analysis carried out here for the massive cases with $s = 5/2, 3$ and $s = 4$ in $D = 3 + 1$ would be affected by their massless counterparts

⁴Strictly speaking, this is valid if the tensor indices are different. Otherwise, the variables will not be independent and the system will have its order reduced.

and could present some surprises. We have shown that this is not the case and the Fierz-Pauli constraints for the higher spin models are exactly the ones postulated in the literature.

We have also extended the method of [8] to $D = 2 + 1$ dimensions. More specifically, in Sections 2 and 3, we obtain explicit expressions for the translation generators, P , and the Lorentz rotation generators, $I = L + S$, acting on arbitrary rank tensors and spinor-tensors in both aforementioned dimensions. In each section, we also provide explicit expressions for the Casimirs of the algebras $\mathfrak{iso}(3, 1)$, given by $C_1 = P^2$ and $C_2 = W^2$, and $\mathfrak{iso}(2, 1)$, given by $C_1 = P^2$ and $C_2 = P \cdot S$. With those expressions at hand and the respective eigenvalue equations defining mass and spin/helicity, in Section 4, we demonstrate that the Fierz-Pauli constraints follow from the Casimir equations, independently of any specific model. Our results also justify the expressions for the Lorentz generators in $D = 2 + 1$ used by [35] and [36], and complement previous works [18, 19]. Our specific results for helicities $5/2$ and 4 are new.

From our expressions, we can deduce the Fierz-Pauli constraints that must be present in any tensor and spinor-tensor description of massive spinning particles. Regarding the general case of arbitrary spin, we have mentioned the technical difficulties involved and paved the way for a general demonstration, see subsection 4.4. Our expressions are also useful for higher rank descriptions. In particular, it is possible to confirm the constraints appearing in the model of [37], which describes massive spin-1 particles via a symmetric rank-2 tensor $W_{\mu\nu} = W_{\nu\mu}$. Such model accommodates self-interacting vertices with super-renormalizable couplings in $D = 3 + 1$, like for example $V = g_o (\eta^{\mu\nu} W_{\mu\nu})^3$ which can not be built up via usual vector or Kalb-Ramond anti-symmetric tensor descriptions. Moreover, we can also revisit the equations of motion of first-order self-dual models describing massive particles with helicity-1 [20], $3/2$ [21], 2 [22], $5/2$ [23], 3 , and 4 [24] in $D = 2 + 1$ dimensions in order to show that, after dynamically obtaining the Fierz-Pauli constraints, they turn into the Klein-Gordon plus Pauli-Lubanski equations in terms of the generators of the Poincaré algebra $\mathfrak{iso}(2, 1)$. For convenience, in Appendix A, we provide the expressions for the spin generators for some of these cases. In appendix B we demonstrate that our original general expression for the spin operator for arbitrary helicity does satisfy the Lorentz algebra in $D = 2 + 1$.

Additionally, our readers may also find interesting the works [38, 39] and the more recent ones [40, 41] where explicit models for spinning particles at the level of first quantization (finite number of degrees of freedom) are obtained in different dimensions via the method of the coadjoint orbit of the Poincaré group.

Acknowledgements

HVAS is supported by FAPESP grant - 2022/14185-1. DD is partially supported by CNPq (grant 313559/2021-0). RRLdS is supported in part by the National Science Centre (Poland) under the research Grant No. 2020/38/E/ST2/00126. DD and RRLdS acknowledge previous support by FAPESP grant 2016/09489-0. RRLdS thanks Astrid Eichhorn and her group at

the University of Southern Denmark for their hospitality during the last stages of the work. The authors thank Prof. Julio M. Hoff da Silva for useful discussions.

A Spin generators in $D = 2 + 1$

Here, for convenience of our readers we give explicit expressions for the spin generator for the specific cases of helicities $\alpha = 5/2, 3$ and 4 . With such expressions, we can better understand the equations of motion coming from the so-called first order in derivatives self-dual descriptions. The simpler cases of helicities $\alpha = 1, 3/2$ and 2 have been used, for example, in [18], [6] and [5], respectively.

A.1 helicity - $\alpha = 5/2$

The symmetric spin operator obtained from (54), taking the rank $s = 2$, is given by:

$$(S_\rho)^{\mu\nu}{}_{\alpha\beta} = \frac{i}{2}(\delta_\alpha^\mu \epsilon^\nu{}_{\rho\beta} + \delta_\alpha^\nu \epsilon^\mu{}_{\rho\beta} + \delta_\beta^\mu \epsilon^\nu{}_{\rho\alpha} + \delta_\beta^\nu \epsilon^\mu{}_{\rho\alpha})\mathbb{I} + \frac{i}{4}(\delta_\alpha^\mu \delta_\beta^\nu + \delta_\alpha^\nu \delta_\beta^\mu)\gamma_\rho, \quad (131)$$

and satisfy the relations

$$[(S^\mu)^{\alpha\beta\rho\sigma}, (S_\nu)_{\sigma\rho\lambda\gamma}] = i\epsilon^\mu{}_{\nu\delta}(S^\delta)^{\alpha\beta\rho\sigma}, \quad (132)$$

$$(S_\rho)^{\mu\nu}{}_{\lambda\sigma}(S^\rho)^{\lambda\sigma}{}_{\alpha\beta} = -\alpha(\alpha + 1)I_{\alpha\beta}^{\mu\nu}, \quad \alpha = \frac{5}{2}, \quad (133)$$

with the identity operator $I_{\alpha\beta}^{\mu\nu}$ given by:

$$I_{\alpha\beta}^{\mu\nu} = \frac{1}{2}(\delta_\alpha^\mu \delta_\beta^\nu + \delta_\alpha^\nu \delta_\beta^\mu)\mathbb{I} - \frac{16}{35}\eta^{\mu\nu}\eta_{\alpha\beta}\mathbb{I} - \frac{4}{35}[(\delta_\beta^\nu \gamma^\mu + \delta_\beta^\mu \gamma^\nu)\gamma_\alpha + (\delta_\alpha^\nu \gamma^\mu + \delta_\alpha^\mu \gamma^\nu)\gamma_\beta]. \quad (134)$$

The self-dual description for massive spin-5/2 in $D = 2 + 1$ dimensions was introduced in the work by Aragone and Stephany in 1984 [23]. It is possible to show that, after satisfying the Fierz-Pauli constraints, the equations of motion of these models can be written in terms of (131).

A.2 helicity - $\alpha = 3$

From (52), the explicit expression for the symmetric spin generator for the rank $s = 3$ case is given by:

$$\begin{aligned} (S_\rho)^{\mu\nu\gamma}{}_{\alpha\beta\lambda} &= \frac{i}{3}(\epsilon_{\alpha\rho}{}^\mu I_{\beta\lambda}^{\nu\gamma} + \epsilon_{\beta\rho}{}^\mu I_{\alpha\lambda}^{\nu\gamma} + \epsilon_{\lambda\rho}{}^\mu I_{\beta\alpha}^{\nu\gamma} \\ &+ \epsilon_{\alpha\rho}{}^\nu I_{\beta\lambda}^{\mu\gamma} + \epsilon_{\beta\rho}{}^\nu I_{\alpha\lambda}^{\mu\gamma} + \epsilon_{\lambda\rho}{}^\nu I_{\beta\alpha}^{\mu\gamma} \\ &+ \epsilon_{\alpha\rho}{}^\gamma I_{\beta\lambda}^{\nu\mu} + \epsilon_{\beta\rho}{}^\gamma I_{\alpha\lambda}^{\nu\mu} + \epsilon_{\lambda\rho}{}^\gamma I_{\beta\alpha}^{\nu\mu}), \end{aligned} \quad (135)$$

where

$$\begin{aligned}
I_{\mu\nu\rho}^{\sigma\phi\omega} &= -\frac{1}{3} \left(\delta_{\mu}^{\sigma} I_{\nu\rho}^{\phi\omega} + \delta_{\mu}^{\phi} I_{\nu\rho}^{\sigma\omega} + \delta_{\mu}^{\omega} I_{\nu\rho}^{\phi\sigma} \right) \\
&+ \frac{1}{18} \left[\delta_{\mu\nu} (\delta_{\rho}^{\phi} \delta^{\omega\sigma} + \delta^{\sigma\phi} \delta_{\rho}^{\omega} + \delta_{\rho}^{\sigma} \delta^{\omega\rho}) + (\rho \leftrightarrow \nu) + (\rho \leftrightarrow \mu) \right].
\end{aligned} \tag{136}$$

We can check that the spin generator S_{ρ} in (135) obeys the following relations:

$$[(S^{\mu})_{\gamma\lambda\delta}^{\alpha\beta\gamma}, (S_{\nu})^{\gamma\lambda\delta}_{\rho\nu\sigma}] = i\epsilon^{\mu}_{\nu\delta} (S^{\delta})^{\alpha\beta\gamma}_{\rho\nu\sigma} \tag{137}$$

$$(S^{\mu})_{\gamma\lambda\delta}^{\alpha\beta\gamma} (S_{\mu})^{\gamma\lambda\delta}_{\rho\nu\sigma} = -\alpha(\alpha+1) I_{\rho\nu\sigma}^{\alpha\beta\gamma}, \quad \alpha = 3. \tag{138}$$

The first-order self-dual model, which describes a massive particle with helicity +3 or -3 in $D = 2 + 1$ dimensions, was introduced by Aragone and Khoudeir in 1993 in [24]. In [7], the authors use the expression (135) in order to write down the equations of motion coming from that model.

A.3 helicity - $\alpha = 4$

From (52), we can obtain the explicit expression for the symmetric spin generator for the rank $s = 4$ case, which is given by

$$\begin{aligned}
(S_{\rho})_{\lambda\sigma\gamma\varphi}^{\mu\nu\alpha\beta} &= \frac{i}{4} \left[\epsilon^{\mu}_{\rho\lambda} I_{\sigma\gamma\varphi}^{\nu\alpha\beta} + \epsilon^{\mu}_{\rho\sigma} I_{\lambda\gamma\varphi}^{\nu\alpha\beta} + \epsilon^{\mu}_{\rho\gamma} I_{\sigma\lambda\varphi}^{\nu\alpha\beta} + \epsilon^{\mu}_{\rho\varphi} I_{\sigma\gamma\lambda}^{\nu\alpha\beta} \right. \\
&+ \epsilon^{\nu}_{\rho\sigma} I_{\lambda\gamma\varphi}^{\mu\alpha\beta} + \epsilon^{\nu}_{\rho\lambda} I_{\sigma\gamma\varphi}^{\mu\alpha\beta} + \epsilon^{\nu}_{\rho\gamma} I_{\lambda\sigma\varphi}^{\mu\alpha\beta} + \epsilon^{\nu}_{\rho\varphi} I_{\lambda\gamma\sigma}^{\mu\alpha\beta} \\
&+ \epsilon^{\alpha}_{\rho\gamma} I_{\sigma\lambda\varphi}^{\nu\mu\beta} + \epsilon^{\alpha}_{\rho\sigma} I_{\gamma\lambda\varphi}^{\nu\mu\beta} + \epsilon^{\alpha}_{\rho\lambda} I_{\sigma\gamma\varphi}^{\nu\mu\beta} + \epsilon^{\alpha}_{\rho\varphi} I_{\sigma\lambda\gamma}^{\nu\mu\beta} \\
&\left. + \epsilon^{\beta}_{\rho\varphi} I_{\sigma\gamma\lambda}^{\nu\alpha\mu} + \epsilon^{\beta}_{\rho\sigma} I_{\varphi\gamma\lambda}^{\nu\alpha\mu} + \epsilon^{\beta}_{\rho\gamma} I_{\sigma\varphi\lambda}^{\nu\alpha\mu} + \epsilon^{\beta}_{\rho\lambda} I_{\sigma\gamma\varphi}^{\nu\alpha\mu} \right],
\end{aligned} \tag{139}$$

where

$$\begin{aligned}
I_{\lambda\sigma\gamma\varphi}^{\mu\nu\alpha\beta} &= \frac{1}{4} \left(\delta_{\alpha}^{\mu} I_{\sigma\gamma\varphi}^{\nu\alpha\beta} + \delta_{\alpha}^{\nu} I_{\sigma\gamma\varphi}^{\mu\alpha\beta} + \delta_{\alpha}^{\sigma} I_{\sigma\gamma\varphi}^{\nu\mu\beta} + \delta_{\alpha}^{\varphi} I_{\sigma\gamma\varphi}^{\nu\alpha\mu} \right) \\
&- \frac{1}{60} \left[\eta_{\lambda\sigma} (\delta^{\mu\nu} I_{\gamma\varphi}^{\alpha\beta} + \delta^{\mu\alpha} I_{\gamma\varphi}^{\nu\beta} + \delta^{\mu\beta} I_{\gamma\varphi}^{\alpha\nu} + \delta^{\nu\alpha} I_{\gamma\varphi}^{\mu\beta} + \delta^{\nu\beta} I_{\gamma\varphi}^{\mu\alpha} + \delta^{\alpha\beta} I_{\gamma\varphi}^{\mu\nu}) \right. \\
&\left. + (\sigma \leftrightarrow \gamma) + (\sigma \leftrightarrow \varphi) + (\lambda \leftrightarrow \gamma) + (\lambda \leftrightarrow \varphi) + (\lambda\sigma \leftrightarrow \gamma\varphi) \right].
\end{aligned} \tag{140}$$

With such expressions at hand, we are able to verify that:

$$[(S^{\mu})_{\gamma\lambda\delta\eta}^{\alpha\beta\gamma\phi}, (S_{\nu})^{\gamma\lambda\delta\eta}_{\rho\nu\sigma\xi}] = i\epsilon^{\mu}_{\nu\delta} (S^{\delta})^{\alpha\beta\gamma\phi}_{\rho\nu\sigma\xi} \tag{141}$$

$$(S_{\rho})_{\eta\omega\delta\xi}^{\mu\nu\alpha\beta} (S_{\rho})^{\eta\omega\delta\xi}_{\lambda\sigma\gamma\varphi} = -\alpha(\alpha+1) I_{\lambda\sigma\gamma\varphi}^{\mu\nu\alpha\beta}, \quad \alpha = 4. \tag{142}$$

The first-order self-dual model describing massive particles of helicity +4 or -4 in $D = 2 + 1$ dimensions was also introduced in [24]. It is possible to write the equations of motion coming from this model in terms of the symmetric generators that we have explicitly deduced here.

B Rank-s algebra

Here, we demonstrate the spin algebra for arbitrary s . From (52), we have:

$$\begin{aligned} [S_\alpha, S_\beta]_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} &= (S_\alpha)^{\mu_1 \dots \mu_s}_{\rho_1 \dots \rho_s} (S_\beta)^{\rho_1 \dots \rho_s}_{\nu_1 \dots \nu_s} - (S_\beta)^{\mu_1 \dots \mu_s}_{\rho_1 \dots \rho_s} (S_\alpha)^{\rho_1 \dots \rho_s}_{\nu_1 \dots \nu_s} \\ &= i^2 \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \epsilon^{\mu_k}_{\alpha \rho_k} \delta^{\mu_i}_{\rho_i} \sum_{l=1}^s \prod_{\substack{j=1 \\ j \neq l}}^s \epsilon^{\rho_l}_{\beta \nu_l} \delta^{\rho_j}_{\nu_j} - i^2 \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \epsilon^{\mu_k}_{\beta \rho_k} \delta^{\mu_i}_{\rho_i} \sum_{l=1}^s \prod_{\substack{j=1 \\ j \neq l}}^s \epsilon^{\rho_l}_{\alpha \nu_l} \delta^{\rho_j}_{\nu_j}. \end{aligned} \quad (143)$$

To analyze this expression, we can consider two scenarios: when $k \neq l$ and when $k = l$. Let's begin with the first case ($k \neq l$). In this scenario, k takes the value of j and l takes the value of i , resulting in contractions between Levi-Civita symbols and Kronecker deltas. Therefore, the right hand side of (143) can be expressed as follows:

$$\sum_{\substack{k,l=1 \\ k \neq l}}^s \prod_{\substack{i=1 \\ i \neq k,l}}^s \delta^{\mu_i}_{\nu_i} \epsilon^{\mu_k}_{\alpha \nu_k} \epsilon^{\mu_l}_{\beta \nu_l} - \sum_{\substack{k,l=1 \\ k \neq l}}^s \prod_{\substack{i=1 \\ i \neq k,l}}^s \delta^{\mu_i}_{\nu_i} \epsilon^{\mu_k}_{\beta \nu_k} \epsilon^{\mu_l}_{\alpha \nu_l}. \quad (144)$$

Notice that l and k are dummy indices and can be interchanged with another index. We substitute l with k and k with l :

$$\sum_{\substack{k,l=1 \\ k \neq l}}^s \prod_{\substack{i=1 \\ i \neq k,l}}^s \delta^{\mu_i}_{\nu_i} \epsilon^{\mu_k}_{\alpha \nu_k} \epsilon^{\mu_l}_{\beta \nu_l} - \sum_{\substack{k,l=1 \\ k \neq l}}^s \prod_{\substack{i=1 \\ i \neq k,l}}^s \delta^{\mu_i}_{\nu_i} \epsilon^{\mu_l}_{\beta \nu_l} \epsilon^{\mu_k}_{\alpha \nu_k} = 0. \quad (145)$$

Now, we can analyze the second case ($k = l$). In this scenario, the sums in (143) can be reduced to just one sum. Additionally, the Kronecker deltas are always contracted, which simplifies the product of symbols to

$$\begin{aligned} \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta^{\mu_i}_{\nu_i} \epsilon^{\mu_k}_{\alpha \rho_k} \epsilon^{\rho_k}_{\beta \nu_k} - \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta^{\mu_i}_{\nu_i} \epsilon^{\mu_k}_{\beta \rho_k} \epsilon^{\rho_k}_{\alpha \nu_k} &= \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta^{\mu_i}_{\nu_i} \left(-\delta^{\mu_k}_{\beta} \eta_{\alpha \nu_k} + \delta^{\mu_k}_{\nu_k} \eta_{\alpha \beta} + \delta^{\mu_k}_{\alpha} \eta_{\beta \nu_k} - \delta^{\mu_k}_{\nu_k} \eta_{\alpha \beta} \right) \\ &= \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta^{\mu_i}_{\nu_i} \left(-\delta^{\mu_k}_{\beta} \eta_{\alpha \nu_k} + \delta^{\mu_k}_{\alpha} \eta_{\beta \nu_k} \right). \end{aligned} \quad (146)$$

Notice that the term $-\delta^{\mu_k}_{\beta} \eta_{\alpha \nu_k} + \delta^{\mu_k}_{\alpha} \eta_{\beta \nu_k}$ can be rewritten as $\epsilon^{\mu_k}_{\rho \nu_k} \epsilon_{\alpha \beta}^{\rho}$, so we have

$$\epsilon_{\alpha \beta}^{\rho} \sum_{k=1}^s \prod_{\substack{i=1 \\ i \neq k}}^s \delta^{\mu_i}_{\nu_i} \epsilon^{\mu_k}_{\rho \nu_k} = -i \epsilon_{\alpha \beta}^{\rho} (S_\rho)^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s}. \quad (147)$$

Finally, by replacing this result in (143), we have

$$[S_\alpha, S_\beta]_{\nu_1 \dots \nu_s}^{\mu_1 \dots \mu_s} = -i \epsilon_{\alpha \beta}^{\rho} (S_\rho)^{\mu_1 \dots \mu_s}_{\nu_1 \dots \nu_s}, \quad (148)$$

completing the demonstration.

References

- [1] E. P. Wigner, “On Unitary Representations of the Inhomogeneous Lorentz Group,” *Annals Math.* **40**, 149-204 (1939) doi:10.2307/1968551
- [2] V. Bargmann and E. P. Wigner, “Group Theoretical Discussion of Relativistic Wave Equations,” *Proc. Nat. Acad. Sci.* **34**, 211 (1948) doi:10.1073/pnas.34.5.211
- [3] Weinberg, S. (1995). *The Quantum Theory of Fields*. Cambridge: Cambridge University Press. doi:10.1017/CBO9781139644167.
- [4] M. Fierz and W. Pauli, “On relativistic wave equations for particles of arbitrary spin in an electromagnetic field,” *Proc. Roy. Soc. Lond. A* **173**, 211-232 (1939) doi:10.1098/rspa.1939.0140
- [5] E. L. Mendonça, A. L. R. dos Santos and D. S. Lima, “Massive spin-3/2 models in $D = 2 + 1$,” *Phys. Lett. B* **775**, 147-151 (2017) doi:10.1016/j.physletb.2017.10.062
- [6] D. Dalmazi and E. L. Mendonca, “Dual descriptions of spin two massive particles in $D=2+1$ via master actions,” *Phys. Rev. D* **79**, 045025 (2009) doi:10.1103/PhysRevD.79.045025
- [7] E. L. Mendonça and D. Dalmazi, “Dual descriptions of massive spin-3 particles in $D = 2 + 1$ via Noether gauge embedment,” *Phys. Rev. D* **91**, 065037 (2015) doi:10.1103/PhysRevD.91.065037
- [8] A. Koenigstein, F. Giacosa and D. H. Rischke, “Classical and quantum theory of the massive spin-two field,” *Annals Phys.* **368**, 16-55 (2016) doi:10.1016/j.aop.2016.01.024
- [9] A. Bengtsson, De Gruiter, *Higher spin field theory v.1. free theory* (2023).
- [10] X. Bekaert, N. Boulanger, A. Campoleoni, M. Chiodaroli, D. Francia, M. Grigoriev, E. Sezgin and E. Skvortsov, “Snowmass White Paper: Higher Spin Gravity and Higher Spin Symmetry,” [arXiv:2205.01567 [hep-th]].
- [11] R. Rahman and M. Taronna, “From Higher Spins to Strings: A Primer,” [arXiv:1512.07932 [hep-th]].
- [12] E. S. Fradkin and M. A. Vasiliev, “Candidate to the Role of Higher Spin Symmetry,” *Annals Phys.* **177**, 63 (1987) doi:10.1016/S0003-4916(87)80025-8
- [13] M. A. Vasiliev, “Consistent equation for interacting gauge fields of all spins in $(3+1)$ -dimensions,” *Phys. Lett. B* **243**, 378-382 (1990) doi:10.1016/0370-2693(90)91400-6
- [14] E. Sezgin and P. Sundell, “Holography in 4D (super) higher spin theories and a test via cubic scalar couplings,” *JHEP* **07**, 044 (2005) doi:10.1088/1126-6708/2005/07/044

- [15] J. Maldacena and A. Zhiboedov, “Constraining conformal field theories with a slightly broken higher spin symmetry,” *Class. Quant. Grav.* **30**, 104003 (2013) doi:10.1088/0264-9381/30/10/104003
- [16] Z. Li, “Bootstrapping conformal four-point correlators with slightly broken higher spin symmetry and 3D bosonization,” *JHEP* **10**, 007 (2020) doi:10.1007/JHEP10(2020)007
- [17] D. Dalmazi, A. L. R. dos Santos, E. L. Mendonça and R. Schmidt Bittencourt, “Soldering spin-3 opposite helicities in $D = 2 + 1$,” *Phys. Rev. D* **100**, no.6, 065011 (2019) doi:10.1103/PhysRevD.100.065011
- [18] R. Jackiw and V. P. Nair, “Relativistic wave equations for anyons,” *Phys. Rev. D* **43**, 1933-1942 (1991) doi:10.1103/PhysRevD.43.1933
- [19] B. Binengar, “Relativistic Field Theories in Three-dimensions,” *J. Math. Phys.* **23**, 1511-1517 (1982) doi:10.1063/1.525524
- [20] P. K. Townsend, K. Pilch and P. van Nieuwenhuizen, “Selfduality in Odd Dimensions,” *Phys. Lett. B* **136**, 38 (1984) doi:10.1016/0370-2693(84)92051-3
- [21] S. Deser and J. H. Kay, “Topologically massive supergravity,” *Phys. Lett. B* **120**, 97-100 (1983) doi:10.1016/0370-2693(83)90631-7
- [22] C. Aragone and A. Khoudeir, “Selfdual massive gravity,” *Phys. Lett. B* **173**, 141-144 (1986) doi:10.1016/0370-2693(86)90234-0
- [23] C. Aragone and J. Stephany, “Dreibein massive spin 5/2,” *Class. Quant. Grav.* **1**, 265 (1984) doi:10.1088/0264-9381/1/3/005
- [24] C. Aragone and A. Khoudeir, “Selfdual spin 3 and 4 theories,” [arXiv:hep-th/9307004 [hep-th]].
- [25] Maggiore M. A Modern Introduction to Quantum Field Theory, Oxford University Press 2005.
- [26] D. Dalmazi and E. L. Mendonca, “A New spin-2 self-dual model in $D=2+1$,” *JHEP* **09**, 011 (2009) doi:10.1088/1126-6708/2009/09/011
- [27] S. Deser and J. G. McCarthy, “Selfdual formulations of $D=3$ gravity theories,” *Phys. Lett. B* **246**, 441-444 (1990) doi:10.1016/0370-2693(90)90627-I
- [28] S. Deser, R. Jackiw and S. Templeton, “Topologically Massive Gauge Theories,” *Annals Phys.* **140**, 372-411 (1982) [erratum: *Annals Phys.* **185**, 406 (1988)] doi:10.1016/0003-4916(82)90164-6

- [29] M. Gomes, L. C. Malacarne and A. J. da Silva, “On the equivalence of the selfdual and Maxwell-Chern-Simons models coupled to fermions,” *Phys. Lett. B* **439**, 137-141 (1998) doi:10.1016/S0370-2693(98)01008-9
- [30] D. Dalmazi and E. L. Mendonca, “Quantum equivalence between the self-dual and the Maxwell-Chern-Simons models nonlinearly coupled to U(1) scalar fields,” *J. Phys. A* **39**, 9355-9363 (2006) doi:10.1088/0305-4470/39/29/025
- [31] E. L. Mendonça and R. Schimidt Bittencourt, “Master Actions and Helicity Decomposition for Spin-4 Models in $3D$ ”, arXiv 2212.09688
- [32] E. L. Mendonça and H. L. Oliveira, “Reaching out a geometrical description for spin-4 self-dual models in $D=2+1$,” *Phys. Rev. D* **105**, no.2, 025002 (2022) doi:10.1103/PhysRevD.105.025002
- [33] C. Fronsdal, ‘Massless Fields with Integer Spin,’ *Phys. Rev. D* **18**, 3624 (1978) doi:10.1103/PhysRevD.18.3624
- [34] J. Fang and C. Fronsdal, “Massless Fields with Half Integral Spin,” *Phys. Rev. D* **18**, 3630 (1978) doi:10.1103/PhysRevD.18.3630
- [35] R. Gaitan, arXiv:0711.2498, “On the Coupling Problem of Higher Spin Fields in 2+1 Dimension ”, PhD thesis, in spanish
- [36] P. Arias, thesis in spanish, 0912.5106 [hep-th].
- [37] D. Dalmazi and R. C. Santos, “Spin-1 duality in D dimensions”, *Phys. Rev. D* **84**, 045027 (2011).
- [38] S. L. Lyakhovich, A. A. Sharapov and K. M. Shekhter, “Massive spinning particle in any dimension. 1. Integer spins,” *Nucl. Phys. B* **537**, 640-652 (1999) doi:10.1016/S0550-3213(98)00617-8 [arXiv:hep-th/9805020 [hep-th]].
- [39] S. L. Lyakhovich, A. A. Sharapov and K. M. Shekhter, “A Uniform model of the massive spinning particle in any dimension,” *Int. J. Mod. Phys. A* **15**, 4287-4300 (2000) doi:10.1016/S0217-751X(00)00211-X [arXiv:hep-th/0002247 [hep-th]].
- [40] T. Basile, E. Joung and T. Oh, “Manifestly Covariant Worldline Actions from Coadjoint Orbits. Part I: Generalities and Vectorial Descriptions”, *JHEP* 01 (2024) 018, e-Print:2307.13644 [hep-th].
- [41] T. Oh, “Ab Initio Construction of Poincaré and AdS Particle”, e-Print:2311.08712 [hep-th].