Massive Star Cluster Formation I.

High Star Formation Efficiency While Resolving Feedback of Individual Stars

Brooke Polak^{1, 2, *}, Mordecai-Mark Mac Low^{2, 3}, Ralf S. Klessen^{1, 4}, Jia Wei Teh¹, Claude

Cournoyer-Cloutier⁶, Eric P. Andersson², Sabrina M. Appel⁷, Aaron Tran⁸, Sean C. Lewis³, Maite J.

C. Wilhelm⁵[®], Simon Portegies Zwart⁵[®], Simon C. O. Glover¹[®], Long Wang⁹[®], and Stephen L. W.

McMillan³

- ¹ Universität Heidelberg, Zentrum für Astronomie, Institut für Theoretische Astrophysik, Heidelberg, Germany
- e-mail: brooke.polak@uni-heidelberg.de
- ² Department of Astrophysics, American Museum of Natural History, New York, NY, USA
- ³ Department of Physics, Drexel University, Philadelphia, PA, USA
- ⁴ Universität Heidelberg, Interdisziplinäres Zentrum für Wissenschaftliches Rechnen, Heidelberg, Germany
- ⁵ Sterrewacht Leiden, Leiden University, Leiden, the Netherlands
- ⁶ Department of Physics and Astronomy, McMaster University, Hamilton, ON, Canada
- ⁷ Department of Physics and Astronomy, Rutgers University, Piscataway, NJ, USA
- ⁸ Department of Physics, University of Wisconsin–Madison, Madison, WI, USA
- ⁹ School of Physics and Astronomy, Sun Yat-sen University, Daxue Road, Zhuhai, 519082, China

December 12, 2023

ABSTRACT

The mode of star formation that results in the formation of globular clusters and young massive clusters is difficult to constrain through observations. We present models of massive star cluster formation using the TORCH framework, which uses the Astrophysical MUltipurpose Software Environment (AMUSE) to couple distinct multi-physics codes that handle star formation, stellar evolution and dynamics, radiative transfer, and magnetohydrodynamics. We upgrade TORCH by implementing the N-body code PETAR, thereby enabling TORCH to handle massive clusters forming from $10^6 M_{\odot}$ clouds with $\geq 10^5$ individual stars. We present results from TORCH simulations of star clusters forming from 10^4 , 10^5 , and $10^6 M_{\odot}$ turbulent, spherical gas clouds (named M4, M5, M6) of radius $R = 11.7 \,\mathrm{pc}$. We find that star formation is highly efficient and becomes more so at higher cloud mass and surface density. For M4, M5, and M6 with initial surface densities $2.325 \times 10^{1,2,3} \,\mathrm{M_{\odot} \, pc^{-2}}$, after a free-fall time of $t_{\rm ff} = 6.7, 2.1, 0.67 \,\mathrm{Myr}$, we find that $\sim 30\%$, 40% and 60% of the cloud mass has formed into stars, respectively. The final integrated star formation efficiency is $\epsilon_{\star} = 32\%$, 65%, and 85% for M4, M5, and M6. Observations of nearby clusters similar in mass and size to M4 have similar integrated star formation efficiencies of $\epsilon_{\star} \leq 30\%$. The M5 and M6 models represent a different regime of cluster formation that is more appropriate for the conditions in starburst galaxies and gas-rich galaxies at high redshift, and that leads to a significantly higher efficiency of star formation. We argue that young massive clusters build up through short efficient bursts of star formation in regions that are sufficiently dense ($\Sigma \ge 10^2 \,\mathrm{M_{\odot} \ pc^{-2}}$) and massive $(M_c \ge 10^5 \, \mathrm{M_{\odot}})$. In such environments, the dynamical time of the cloud becomes short enough that stellar feedback cannot act quickly enough to slow star formation.

Key words. Star clusters - star formation - ISM: clouds - globular clusters: general

1. Introduction

Globular clusters (GCs), found in every massive galaxy, are some of the most ancient objects in the Universe. They serve as fossils that can reveal the elusive environment and physics of the early phases of galaxy assembly (Brodie & Strader 2006; Portegies Zwart et al. 2010; Renaud et al. 2017; Krumholz et al. 2019; Adamo et al. 2020). Yet because of their age, many aspects of cluster formation and evolution at high redshift are challenging to constrain through observation. Little is known about the efficiency and timescale at which gas is converted into stars to create such massive bound clusters.

Though the progenitors of GCs are too old to observe, younger star clusters that are as massive as GCs are forming in nearby galaxies. These young massive clusters (YMCs) have masses $M \ge 10^4 \,\mathrm{M_{\odot}}$ and ages < 100 Myr (Portegies Zwart et al. 2010). The discovery of these objects indicates that the mode of extreme star formation that forms massive star clusters still occurs today. Even more are being discovered with JWST, as many YMCs in starburst galaxies are too embedded to have been seen by *Hubble* (Whitmore et al. 2023). Although it has been suggested that YMCs are the present day analogues to young GCs, this is debated in the literature (see Renaud 2020).

^{*} Fellow of the International Max Planck Research

School for Astronomy and Cosmic Physics at the University of Heidelberg

Theory suggests that, despite the abundance of GCs, $\leq 1\%$ of clusters survive to become GCs (Fall & Zhang 2001; Fall et al. 2005; Fall 2006). The conditions that lead to bound star clusters as massive as GCs remain a mystery, and observations of forming YMCs are sparse. Star formation must be fast and efficient enough to form bound stars that can survive the epoch of stellar feedback and the dispersal of the natal gas (Lada & Lada 2003). The plethora of GCs suggests these conditions were very common in the early Universe.

The process of star formation in a cluster begins with the global gravo-turbulent collapse of giant molecular clouds (Larson 1981, GMCs;). As the collapse proceeds, fragmentation creates overdense clumps within the GMC that begin to form stars (Mac Low & Klessen 2004; McKee & Ostriker 2007; Klessen & Glover 2016). The feedback from these stars, in the form of stellar winds, jets, and radiation begins clearing out dense gas in and around the forming sub-clusters, slowing down the local (sub-cluster scale) and global (cloud scale) star formation rate (SFR; e.g. Girichidis et al. 2020; Lewis et al. 2023). Eventually, massive stars explode as SNe further dispersing gas. However, it has been argued that the efficiency at which stellar feedback slows global star formation diminishes with higher gas surface density (Grudić et al. 2018). The sub-clusters eventually merge if they are mutually gravitationally bound, forming a final cluster cleared of all natal gas (Krause et al. 2020).

Many details of star cluster formation remain poorly understood due to the difficulty of modelling such a complex process. Stellar evolution and binary dynamics need to be resolved on time scales of years and distance scales of an AU, while the magnetohydrodynamics (MHD) of the collapsing gas covers regions several parsecs across with crossing times of thousands to millions of years. Because of this, most computational star cluster formation models are limited and must make considerable approximations. Many simulations do not form individual stars but rather sink particles representing sub-clusters (e.g. Bate et al. 1995; Federrath et al. 2010) and extract the properties and feedback of individual stars from the sink particles (e.g. Sormani et al. 2017; Howard et al. 2017; Kim et al. 2017; Grudić et al. 2018; Su et al. 2018; Lahén et al. 2019). Some simulations do form single stars, but do not resolve the stellar feedback of each individual star particle (Colín et al. 2013; Li et al. 2019), instead including feedback from just the sink particles that created the stars.

Modelling individual stars is important, as this can change the efficiency and location of stellar feedback injection. Dynamical processes often eject high mass stars (Fujii & Portegies Zwart 2011); the location of massive stars directly affects how and when gas is dispersed. Gas dispersal stops star formation. Models of sub-cluster feedback may over-estimate the strength of feedback, as they do not allow spatial separation between the stars in the sub-cluster. This lack of separation also changes the morphology of the gas, affecting the number of low-density channels in the gas that can vent thermal energy from the sub-cluster. The degree to which the sub-cluster vs. star-by-star approaches differ must be constrained.

There are a few models that do evolve individual stars with both stellar feedback and higher order gravitational dynamics (Wall et al. 2020; Grudić et al. 2021). While these models include most of the relevant physics, they lack the computational efficiency to simulate star clusters forming from clouds of masses $>10^5~M_{\odot}$, and instead focus on simulating star clusters forming from low mass clouds $\leq 10^5~M_{\odot}$. This leaves a sizable gap compared to the observed mass range of GMCs. While clusters with mass $<10^5~M_{\odot}$ are comparable to local group observations, YMC and GC formation is out of their reach. Furthermore, most star formation takes place in GMCs of mass $\geq 10^5~M_{\odot}$ (McKee & Williams 1997; Murray & Rahman 2009).

The goal of this work is to model the formation of massive clusters from their initial GMCs while following the formation of individual stars and their feedback. We aim to answer how, and in what conditions, YMCs can form while remaining bound throughout the onset of gas expulsion. How efficient is the process of star formation in a cluster, and what is the timescale over which star formation occurs? Do the clusters formed from these massive clouds survive and remain bound, or quickly disperse? How do our results compare to those that use a sub-cluster formation and feedback model?

To do this, we use the TORCH framework (Wall et al. 2019, 2020). TORCH employs the Astrophysical Multipurpose Software Environment (AMUSE) framework to couple separate physics codes that handle MHD, radiative transfer, stellar evolution, and N-body dynamics. TORCH uses the MHD code FLASH (Fryxell et al. 2000; Dubey et al. 2014), which accounts for the evolution of the gas and the formation of sink particles and stars. Stellar feedback in the form of winds and SNe is included, and the effect of ionizing and non-ionizing radiation is followed using a raytracing approach (Baczynski et al. 2015). The star formation model samples the Kroupa (2002) initial mass function (IMF) to form stars from sink mass reservoirs (Wall et al. 2019). SEBA (Portegies Zwart & Verbunt 1996) tracks stellar evolution from the zero-age main sequence, and, in the original version of TORCH, PH4 (McMillan et al. 2012) + MULTIPLES (Portegies Zwart & McMillan 2018) handled the stellar dynamics.

TORCH in that version (Wall et al. 2019, 2020) could not handle the hundreds of thousands of stars that form in massive GMCs $>10^5~M_{\odot}$. In this work, we solve this problem by making three updates: 1) we replace the combination of the N-body code PH4 and the higher-order interactions MULTIPLES with the code PETAR (Wang et al. 2020a); 2) we agglomerate stars with masses $<4~M_{\odot}$ into summed-mass dynamic star particles with masses of $\geq4~M_{\odot}$; and 3) we mass load the stellar winds to reduce the peak temperatures beyond their termination shocks. With these modifications, TORCH can now model clouds with an initial mass of up to $10^6~M_{\odot}$ that form hundreds of thousands of individual stars.

We present simulations of star clusters forming from turbulent spherical clouds with masses of 10^4 , 10^5 , and $10^6 M_{\odot}$. Each of these clouds is almost identical in terms of their initial properties, with only mass and density scaled between them. Our study investigates whether the formation of YMCs parallels that of low-mass clusters, or if it varies significantly with initial cloud mass and density.

This paper is the first in a series exploring the results of these simulations. In this paper, we describe the TORCH code, the new features integrated into TORCH for handling massive GMCs, and the initial conditions of our three clouds in Sect. 2. We analyze the time evolution of global gas and stellar properties in Sect. 3. In Sect. 4, we discuss the results of our analysis, and in Sect. 5 we conclude with a summary of the most important results.

2. Methods

2.1. Standard TORCH

TORCH¹ is built upon the AMUSE framework, which couples multiple autonomous astrophysical codes. We choose codes that allow efficient calculation of the disparate physical processes at work in star cluster formation.

The TORCH framework incorporates the adaptive mesh refinement MHD code FLASH v4.6.2 (Fryxell et al. 2000; Dubey et al. 2014), with a number of enhancements to the base version of FLASH. Base FLASH handles the MHD, and sink particle formation and evolution. The modifications to FLASH presented in Wall et al. (2019, 2020) include: heating and cooling, ionization, radiation transfer (using raytracing; see Baczynski et al. 2015), and feedback injection from stars. Stellar feedback is implemented in FLASH in the form of ionizing extreme ultraviolet (EUV) and nonionizing far ultraviolet (FUV) radiation in the form of radiative heating and radiation pressure, as well as mechanical feedback from stellar winds and SNe. We use the HLLD Riemann solver (Miyoshi & Kusano 2005) in FLASH paired with third-order piecewise parabolic method reconstruction (Colella & Woodward 1984).

To avoid artificial fragmentation, the Jeans length

$$\lambda_{\rm J} = \sqrt{\pi c_s^2 / (G\rho)} \tag{1}$$

must be resolved by at least four cells (Truelove et al. 1997). As density increases during collapse, the Jeans length will decrease until this criterion is no longer met at the highest level of AMR refinement. Sink particles are needed to collect the gas that surpasses this density. This constraint sets the sink radius to $R_{\rm sink} = 2.5\Delta x_{\rm min}$ and gives the sink threshold density as

$$\rho_{\rm sink} = \frac{\pi c_s^2}{G\lambda_{\rm J}^2} = \frac{\pi c_s^2}{G(5\Delta \mathbf{x}_{\rm min})^2} \,, \tag{2}$$

where $c_{\rm s}$ is evaluated using the initial temperature of the gas.

On each timestep, the mass of gas within a distance $R_{\rm sink}$ of a sink particle that satisfies the criteria outlined in Federrath et al. (2010) is added to that sink's mass reservoir for creating stars. When a sink forms, it randomly samples the Kroupa IMF (Kroupa 2002) and stores a list of star masses to form (see also Sormani et al. 2017). Each timestep, the sink forms stars from this mass list until its mass reservoir is depleted. It again forms one or more stars the next time it has accreted enough mass for at least the next star on the list. This is the standard stellar mass sampling method used in TORCH (Wall et al. 2019). Star positions are randomly sampled from a uniform spherical distribution within the sink's accretion radius. Star velocities are set by the sink velocity and an additional isotropic velocity dispersion with a Gaussian distribution with standard deviation of the local sound speed.

Star particles are initially formed as zero-age main sequence stars, neglecting pre-main sequence evolution. Subsequent stellar evolution is tracked with SEBA (Portegies



Fig. 1. Wall-clock time for an evolution step for PETAR and PH4+MULTIPLES given the number of stars.

Zwart & Verbunt 1996), which passes the evolutionary properties informing stellar feedback to FLASH. The N-body dynamics of the stars are calculated using PETAR (Wang et al. 2020a), which is discussed further in the next section. Stars dynamically interact with the gas in FLASH through the AMUSE hierarchical coupling (Portegies Zwart et al. 2009) based on the gravity-bridge algorithm of Fujii et al. (2007).

2.2. PETAR N-body

TORCH was first designed to use the N-body code PH4 (McMillan et al. 2012) to handle direct stellar dynamics, paired with MULTIPLES (Portegies Zwart & McMillan 2018) to track binary orbital evolution and higher order perturbations. For TORCH runs using an initial cloud of $10^4 M_{\odot}$ and producing only a few thousand stars, this works well. However, the computational cost becomes unfeasible when pushing to higher initial cloud masses that produce far more than 10^4 stars with over a few hundred binary systems. This is because MULTIPLES is a serial Python code, so with many interacting stars computation times become impractical. To speed up TORCH, we replaced PH4 and MULTIPLES with PETAR (Wang et al. 2020a).

PETAR is a state-of-the-art gravitational dynamics code optimized for solving the stellar dynamics of systems with millions of stars. It accomplishes this by dividing gravitational interactions into three regimes: distant interactions calculated with a Barnes & Hut (1986) tree and handled by the framework for developing parallel particle simulation codes (FDPS; Iwasawa et al. 2016, 2020), nearby interactions solved with a fourth-order Hermite direct N-body integrator (Makino & Aarseth 1992), and close interactions (binaries and higher order systems/perturbations) solved using the Slow Down Algorithmic Regularization (SDAR) technique (Wang et al. 2020b). For each particle, the force from neighboring particles will be solved depending on what distance regime they are in, with a mass-dependent factor to increase the distance over which massive particles are considered close neighbors. The SDAR feature for handling higher-order dynamics is the novel component of PETAR,

¹ Available at https://bitbucket.org/Torch-sf/Torch/src/main/

enabling it to handle large numbers of binaries and higher order systems in parallel.

In Figure 1, we plot the wall-clock time per evolution step for each of the N-body codes considered. For reproducibility, this test was done with the parameters $r_{out} = 0.001 \,\mathrm{pc}$, $r_{bin} = 100 \,\mathrm{AU}$ for PETAR and the stellar interaction radius $r_{\rm int} = 15 \,\mathrm{R}_{\odot}$ for MULTIPLES. PETAR is significantly faster and consistently performs well as the number of stars increases. The variability in the performance of PH4 and MULTIPLES is due to MULTIPLES taking longer if there are many third-body perturbations in a given step. Note that this test was done with single stars only; the scaling for a run with primordial binaries will be different.

When running PETAR in TORCH, the timestep of the long distance particle tree must be set (dt_soft), as well as the changeover radius between direct N-body and tree method for force calculations (r_out). If the user sets these two parameters, all other parameters are set automatically. In TORCH, the MHD code sets the global timestep for all worker codes based on the Courant condition. The tree timestep is set as the nearest power of two in code units below the requested timestep, as a power of two is required by PETAR (like most N-body codes). This sets dt_soft. We set the outer radius r_out to 0.001 pc, the standard value used in PETAR simulations. We use a softening length of $\ell = 15 \, \mathrm{R}_{\odot}$.

The code handling stellar mergers within PETAR is not active within the AMUSE framework, which can result in star particles that approach within each others' softening radius and should merge instead ending up with identical positions, leading to a halt in code execution. We have implemented code to check for particles in this state, and merge them. We use SEBA to model the merger and calculate the stellar properties of the remaining star, and then we remove the swallowed companion star.

2.3. Stellar Modifications

We have made three alterations to the star formation and evolution procedures that vary from standard TORCH to accommodate the several orders of magnitude increase in number of stars formed when increasing the initial cloud mass from $10^4\,M_\odot$ to $10^6\,M_\odot$.

- I. We agglomerate low-mass star particles below $M_{\text{agg}} = 4 \,\mathrm{M}_{\odot}$ as they form until their summed mass is $\geq M_{\text{agg}}$. Then, a star particle is formed with a mass equal to the sum of the low-mass stars. This reduces the strain on the N-body calculations by reducing the number of dynamical star particles by 90%.
- II. We mass-load stellar winds to raise the Courant timestep by limiting the temperature of wind-blown bubbles to $T_w = 3 \times 10^5$ K, which significantly speeds up the simulations. This results in smaller, cooler, momentum-conserving bubbles instead of hot energy-conserving bubbles. The primary effect of wind feedback in cluster formation is to clear out extremely dense gas to allow ionizing radiation to form expanding H 2 regions. In this dense gas even hot stellar wind bubbles cool quickly, so there is little change in behavior in this regime.
- III. We only inject feedback from stars above $20 \, M_{\odot}$ to reduce the cost of ray-tracing. Massive stars output most of the ionizing radiation and mechanical wind

Article number, page 4 of 18

energy in clusters: by setting this limit we lose less than 20% of the total feedback energy. Stars below the feedback cutoff mass will not go SN within the time frame of our simulations (≤ 10 Myr).

We further explain and examine the effects of these modifications in Appendix B, including providing a quantitative analysis of the amount of total energy lost by excluding feedback for stars $< 20\,M_{\odot}$ in the M6 model.

2.4. Initial Conditions

The initial properties of our three clouds are listed in Tables 1 and 2. We chose to keep the radius of all three clouds constant at $R_{\rm cl} = 11.7$ pc. The radius was kept the same to have the same spatial distribution of star formation for each run. Constant radius allows the cell resolution and size of sink particles to be the same between the three simulations, and it facilitates directly comparing the morphology and dynamics of the forming clusters.

Consequently, the average initial densities of the clouds are 1.5, 15, and 150 $M_{\odot} \text{ pc}^{-3}$, or 10^{-22} , 10^{-21} , and $10^{-20} \text{ g cm}^{-3}$. The column densities of these clouds are $2.325 \times 10^{1,2,3} \text{ M}_{\odot} \text{ pc}^{-2}$, respectively. Assuming a 9:1 number ratio of H:He, resulting in a mean molecular weight of $\mu = 1.3$, this gives total particle number densities of n = 46, 460, and 4600 cm⁻³. Each cloud has a column density consistent with observations. Observations show a strong positive correlation between the mass and density of GMCs in PHANGS galaxies (Sun et al. 2022), suggesting mass and density should be varied together.

The initial clouds must be in pressure equilibrium with their surroundings to avoid unphysical shocks from pressure imbalances at their surfaces. The M4 and M5 clouds are in the pressure regime where there is a stable two-phase medium at solar metallicity and Milky Way background UV field (Field et al. 1969; Wolfire et al. 2003), meaning there is a set of temperatures for the cloud and background for a given cloud density where the cold dense cloud and the warm ambient medium are both in thermal equilibrium at equal pressure. The cloud temperatures for the M4 and M5 clouds are $T_{\rm cl}=103$ K and 28 K, respectively, and the corresponding background temperatures and number densities are $T_{\text{amb}} = 9,000$ and 4,000 K, and $n_{\text{amb}} = 3$ and 1 cm^{-3} . The M6 cloud, however, is at a high enough pressure that a two-phase medium no longer exists. Only the cold phase can be in thermal equilibrium at the M6 cloud's pressure. This means that the low-density envelope of the M6 cloud is inherently not in thermal equilibrium. To minimize the pressure imbalance with the core, we therefore raise the background density to $n_{\rm amb} = 100 \text{ cm}^{-3}$. Both the cloud and background medium for M6 are at a temperature of $T_{\rm cl} = T_{\rm amb} = 50 \text{ K}.$

The initial conditions described in the rest of this section and summarized in Table 2 apply to all three clouds. The clouds have a Gaussian density profile (Bate et al. 1995; Goodwin et al. 2004) with the standard deviation set such that the ratio of the cloud's central to edge density is 3:1. The simulation domain is a cube of half-width $R_{\rm box} = 20$ pc with outflow boundary conditions. The outflow boundaries do allow gas flow onto the grid from ghost zones if the velocity at the edge of the grid is directed inward. We use three refinement levels, yielding cell sizes that range from $\Delta x_{\rm min} = 0.3125$ pc to $\Delta x_{\rm max} = 1.25$ pc. Refinement and



Fig. 2. Slice plots of the three simulations in the x-y plane over time. The plane of the slices for a given cloud is the center of stellar mass in the final snapshot. Stellar positions are shown by white dots. The free-fall times $t_{\rm ff}$ are given in Table 1. The number of stars shows the amount of star particles in the domain, not the number sampled from the IMF. Due to our agglomeration of low-mass star particles, the number of stars sampled from the IMF is ~ 10× greater. Article number, page 5 of 18

 Table 1. Model Parameters

Parameter	M4	M5	M6
$M_{\rm cl} [{\rm M}_{\odot}]$	10^{4}	10^{5}	10^{6}
$\rho_c, \bar{\rho} [\mathrm{M}_\odot \mathrm{pc}^{-3}]$	2.8, 1.5	28, 15	280, 150
$\Sigma [{ m M}_{\odot}{ m pc}^{-2}]$	23.25	232.5	2325
$\lambda_{\rm J}$ [pc]	10.0	3.2	1.0
$t_{\rm ff}$ [Myr]	6.7	2.1	0.67

Notes. Rows: cloud mass, cloud central and average volume density, initial column density, Jeans length (Eq. 1) at initial temperature, free-fall time.

 Table 2. Control Parameters

Parameter	Value	Units
$R_{\rm cloud}$	11.7	pc
$R_{ m box}$	20.0	\mathbf{pc}
$lpha_{ m v}$	0.15	-
Δx_{min}	0.3125	\mathbf{pc}
Δx_{max}	1.25	\mathbf{pc}
$R_{\rm sink}$	0.78125	\mathbf{pc}
$ ho_{ m sink}$	8×10^{-21}	${ m g~cm^{-3}}$
$M_{\rm sink}$	246	M_{\odot}
$M_{\rm feedback}$	20	M_{\odot}
$M_{\rm n-body}$	4	M_{\odot}
$M_{ m IMF}$	0.08 - 100	M_{\odot}

Notes. Rows: Radius of cloud, half-width of box, virial parameter, minimum cell width, maximum cell width, sink radius, sink threshold density, initial sink mass, minimum feedback star mass, agglomeration mass of low-mass stars, mass sampling range of IMF.

de-refinement of the grid is determined by the Jeans criterion described in Sect. 2.1 and on temperature and pressure gradients.

We initially impose a Kolmogorov (1941) turbulent velocity spectrum on all the gas in the domain. The same random seed is used to generate the turbulent velocity spectrum for all three clouds. This ensures the same turbulent collapse patterns, minimizing differences in the formation, location, and morphology of dense cores. From the edge of the cloud to the domain boundary, we linearly taper the magnitude of the turbulent velocities from 100% to 25%. This tapering does not affect the low-density ambient background of the M4 and M5 cloud, but helps with stability in the M6 cloud by mixing the border of the cloud, where there is a small pressure jump.

The sink accretion radius and threshold density, derived in Sect. 2.1, are $r_{\rm sink} = 2.5\Delta x_{\rm min} = 0.78$ pc and $\rho_{\rm sink} = 8.35 \times 10^{-21}$ g cm⁻³. This gives an initial sink mass resolution of $m_{\rm sink} = 245 M_{\odot}$, meaning that when a sink initially forms it will accrete and form approximately $m_{\rm sink}$ worth of stellar mass, given the sink's threshold density and accretion radius. The IMF sampling mass range is 0.08–100 M_{\odot}. The lower end is the hydrogen-burning limit, while the upper end is the most massive star thought to form in a star cluster with stellar mass $\approx 10^4 \, {\rm M}_{\odot}$ (Weidner et al. 2009).

The critical virial ratio for stability is $\alpha_v = E_{\rm kin}/|E_{\rm pot}| = 0.5$, below which collapse occurs. Massive clouds tend to be sub-virial, with clouds of $10^6 \,{\rm M_{\odot}}$ observed to have virial parameters of $\alpha_v \approx 0.05 - 0.35$ (Kauffmann et al. 2013), though some surveys see super-virial

massive clouds (see Fig. 2 of Chevance et al. 2023). Note that these values have been converted from the different virial parameter definition in Kauffmann et al. (2013). Therefore, we choose a fiducial virial parameter value of $\alpha_v = 0.15$ for our models to promote rapid onset of collapse.

Magnetic fields are prevalent in the interstellar medium (Crutcher et al. 2003) and affect the collapse of GMCs and subsequent star formation. Although they are not the dominant factor in determining how star formation proceeds within a cloud, their presence has been shown to alter the fragmentation of cores (Price & Bate 2008; Peters et al. 2011) and slow down the global evolution of the cloud (Heitsch et al. 2001). With a strong enough field, clouds can be supported against gravitational collapse (Heiles 1976), although generally observed magnetic fields are not strong enough to inhibit collapse (Klessen & Glover 2016). The critical value of the mass-to-flux ratio for a cloud to be supported by magnetic fields against gravitational collapse is given by (Mouschovias & Spitzer 1976; Mouschovias 1991)

$$\left[\frac{M}{\Phi}\right]_c = \frac{\zeta}{3\pi} \sqrt{\frac{5}{G}} = 490 \ \frac{g}{Gauss \ cm^2} , \qquad (3)$$

where G is the gravitational constant, and a correction factor $\zeta = 0.53$ for a uniform sphere is used (Strittmatter 1966).

In our simulations, each cloud's initial magnetic field $\boldsymbol{B} = B_z \hat{z}$ is uniform in z and decreases radially in the x-y plane, following the mid-plane density $\rho(x, y, z = 0)$, as:

$$B_z(x,y) = B_0 \exp\left[-(x^2 + y^2)\ln(3)/R_{\rm cl}^2\right]$$
(4)

with $B_0 = 0.185, 1.85, 18.5 \,\mu\text{G}$ for the M4, M5, M6 clouds respectively. These values match observations for M5 and M6, while the field is a factor 10 weaker for M4 (Crutcher et al. 2010). The integrated magnetic flux $\Phi = 2\pi B_0 R_{\rm cl}^2/(3 \ln(3))$, so all clouds have initial mass-toflux ratio $M_{\rm cl}/\Phi = 4.5 \times 10^4 \,\text{g Gauss}^{-1} \,\text{cm}^{-2}$ much larger than Equation (3). The initial magnetic fields are thus weak and do not inhibit collapse in any of our simulations.

3. Results

3.1. Cluster Formation Overview

At the onset of the simulation, each cloud begins to gravitationally collapse. Turbulent velocities fragment the cloud and create overdense hubs and filaments. Because the same random seed is used in all three clouds to generate the initial turbulent velocity spectrum, the web of dense gas is the same for each cloud. This means that the spatial distribution of star formation is similar for all three clouds. This can be seen in the time evolution of the three clouds in Figure 2. The first stars all form in the largest over-density in the middle of the bottom of the cloud. Then, more stars form along the filaments of the dense cloud forming a V shape. The M5 and M6 clusters in particular look very similar in terms of sub-clustering and merging. The M4 cluster forms significantly fewer stars and therefore fewer sub-clusters.

By a free-fall time $t_{\rm ff}$, the sub-clusters in M4 and M5 have mostly merged, forming a single central spherical cluster. The M6 model is still forming stars in various subclusters and has not assembled its main central cluster yet. By looking at the spatial distribution of sub-clusters and the density of the gas, one can see that stellar feedback becomes most efficient once the sub-clusters have merged into a single cluster. Whether feedback is only strong enough to disperse gas when clustered or this is coincidental with the timing of feedback needs further examination, but this is outside of the scope of this introductory paper. Low density bubbles begin to occupy a significant fraction of the cloud volume once the central star cluster has been assembled.

Once most of the stars have formed, the efficiency of stellar feedback for dispersing the natal gas varies greatly for the three cloud masses. The final row in Figure 2 shows the M4, M5, and M6 systems at $1.2t_{\rm ff}$. At this point, only the M4 and M5 clusters have blown large bubbles. The feedback from the M6 cluster has hardly slowed the collapse of the densest gas, and rapid star formation continues. The M4 cloud has dispersed nearly all of the remaining gas, and star formation has halted completely.

3.2. Visualizing Cluster Morphology

The complex 3D structure of star clusters is hard to visualize using 2D plots. Figure 3 shows a still of an interactive plot of the M4, M5, and M6 simulations after one free-fall time generated with Plotly (Plotly Technologies Inc. 2015). After downloading the HTML file available in the online version of this paper, readers can zoom, pan, and rotate for a complete look at the morphology of each cluster. The color is an isosurface of the gas density in log scale, and the points are stars with sizes scaled to their stellar radius. This tool makes it clear just how non-spherical these clusters are. Comparing the still of the interactive plot to the slice plots in Figure 2, one can already extract much more information on the system's morphology.

Zooming into the core of each cluster shows the immense stellar density of the M6 cluster, whereas in the M4 cluster one can easily distinguish individual stars. The gas is also far less dense in the M4 system compared to M5 and M6.

The shape of the M6 cluster is highly irregular. Stemming from the largest cluster, one can see a row of subclusters forming along a filament. Branching perpendicularly off this filamentary cluster are two more star forming filaments in a configuration resembling the letter "F". The M5 cluster has a shape congruous to the shape of M6, but with fewer stars bridging the gaps between clusters in the main filament. The M5 cluster also has only one finger perpendicular to the main filament, which contains many fewer stars than the fingers of the M6 cluster. The M4 cluster is much less dense, with its few stars outlining the same core filament cluster seen in M5 and M6. However, in M4 subclusters can no longer be distinguished. The M4 sub-clusters already merged into a singular central cluster.

3.3. Star Formation History

The global properties of the star clusters that form from the M4, M5 and M6 clouds over the period of star formation are shown in Figure 4 as a function of time in units of the global free-fall time of the cloud $t_{\rm ff}$ (see Table 1). For reference, this same figure is shown as a function of physical time in Appendix A. We analyze these properties and discuss how they compare across the three clouds to assess the effect of the initial cloud mass and density on the resultant cluster

properties. Table 3 highlights the final properties of the clusters formed in the three models.

Our results suggest that star formation in a GMC is a fast and efficient process regardless of initial cloud mass and density, with all three clouds converting at least 30% of their initial gas into stars within an initial free-fall time. Star formation becomes faster and more efficient as the mass and density of the GMC increases.

Stars begin forming in the M5 and M6 clouds at $t = 0.3 t_{\rm ff}$, while in the M4 run it is delayed until $t = 0.5 t_{\rm ff}$. Because of the turbulent field, regions of the clouds have locally shorter free-fall times leading to star formation earlier than the global free-fall time. The duration of star formation is the shortest for the M4 cloud, lasting $t_{\rm sf} = 0.7 t_{\rm ff}$. The M5 cloud forms stars for a longer period in terms of its initial free-fall time, $t_{\rm sf} = 1.3 t_{\rm ff}$. M6 is still forming stars 1.3 $t_{\rm ff}$ after the onset of SF.

The SFR (Fig. 4(a)) increases with cloud mass. The peak SFR for the M4, M5 and M6 clouds are SFR_{peak} = 0.4, 5.5, and 390 M_{\odot} yr⁻¹ respectively. The average SFRs also increase with mass, with values of SFR_{ave} = 0.01, 0.06 and 2.2 M_{\odot} yr⁻¹.

The total star formation efficiency (SFE; Fig. 4(b)) we discuss here is simply the ratio of stellar mass formed to the initial gas mass of the cloud $\epsilon_{\star} = M_{\star}/M_c$. Typical SFE values of nearby clusters in the Milky Way lie between 0.1–0.3 (Lada & Lada 2003). Note that observed SFEs are the instantaneous ($\epsilon_{inst} = M_{\star}/(M_{\star} + M_{gas})$) rather than integrated SFEs we derive from our results. The M4 cloud is representative of typical GMCs at the solar circle, and its SFE lies at the top of this range at 0.3. In the higher mass clouds of M5 and M6, the SFE is much higher. The M5 cloud converted 0.65 of its gas into stars, and the M6 cloud converted a surprising 0.85 of gas into stars. This suggests that the SFE in high-mass, high-density environments can be much higher than seen in low-mass local clusters.

The free-fall time becomes so short in these high mass clouds that the stellar feedback simply does not act quickly enough to stop star formation before most of the gas has formed into stars. Free-fall times of dense environments that are shorter than the development times for winds and SNe have indeed been shown to result in high SFE (Dekel et al. 2023).

Figure 4(c) shows the mass of the most massive star that has formed from random draws from the IMF. By a free-fall time, each cloud has already formed the most massive star in its cluster. We find that the mass of the most massive star increases with cluster mass. For the M5 and M6 clouds, the most massive star is at the maximum sampling mass of $100 \,\mathrm{M}_{\odot}$, while the M4 cloud's most massive star is around $80 \,\mathrm{M}_{\odot}$. This is a stochastic effect; as more stars are sampled from the IMF, you will eventually sample the most massive star in the distribution. This ends up reproducing the effect suggested by Weidner et al. (2009) and Yan et al. (2023) that the cluster mass limits the most massive star mass. In each cloud, it is interesting to note that each instance of the formation of a very massive star, i.e. above $40 \,\mathrm{M}_{\odot}$, correlates with a slowing of star formation indicated by a reduction in the SFE slope.

The M6 cloud forms from IMF sampling $\sim 10^6$ stars, M5 forms $\sim 10^5$ stars, and M4 forms $\sim 10^4$ stars. With agglomeration, the number of stars in the simulation are about 10% of these numbers, so the improved version of



Fig. 3. Still of the interactive plot of the embedded M4, M5, and M6 clusters (left to right) at $1 t_{\rm ff}$. The interactive plot file is available for downloading in the online publication page.

TORCH with PETAR can simulate clusters of $> 10^5$ individual stars.

3.4. Cluster Evolution

The evolution of the global properties of the formed star clusters occurs quite similarly for all three clouds, but the magnitude of their values depends greatly on the cloud's initial mass.

The stellar velocity dispersion (Fig. 4(e)) also increases roughly linearly with initial cloud mass. The velocities of stars increase at a slow pace before leveling out after $1 t_{\rm ff}$. For the M4 cluster, the velocity dispersion levels out at just 0.9 km s⁻¹. The M5 cluster reaches a velocity dispersion of 5 km s⁻¹ and the M6 cluster has a velocity dispersion of 20 km s⁻¹. The deeper potential wells of the higher mass clusters, going as the square root of the mass for these similar sized objects, drive the higher velocity dispersions. In the case of M6, the potential well depth exceeds the sound speed of ionized gas, preventing gas from escaping even after ionization.

The evolution of the half-mass radius $R_{1/2}$ of all the stars in the cluster (Fig. 4(f)) seems to be split into a high and low-mass regime. The M5 and M6 clusters follow the same track closely. From $0.3 t_{\rm ff}$ to $0.5 t_{\rm ff}$, $R_{1/2}$ is steady at 2 pc. Then, from $0.5 t_{\rm ff}$ to $1.0 t_{\rm ff}$ it increases to a peak $R_{1/2}$ of 4 pc at $\sim 0.6 t_{\rm ff}$ then goes back down to 2 pc. From there the two clusters begin to diverge, with the size of the M5 cluster steadily increasing. The similarities in the evolution of the two clusters are most likely due to the fact that both clouds at early times form enough stars for distinct subclusters to form and merge. The sub-clusters have formed in the same places so both clusters peak at roughly the same $R_{1/2}$. Though the clusters are expanding, all of the stars remain bound, suggesting that the clusters are not dissolving but rather relaxing into gravitational equilibrium. Longer runs following just the stars after gas dispersal will ultimately be needed to demonstrate this.

The M4 cluster grows differently. It also hovers at roughly 2 pc until 1 $t_{\rm ff}$, after which it linearly increases to 6 pc by 1.5 $t_{\rm ff}$. The M4 cluster is clearly expanding, but 60% of its stars remain bound, so complete dissolution has not yet occurred (Fig. 4(i)).

The onset and duration of gas dispersal from the star clusters depends strongly on the initial mass and density of the cloud. Figure 4(q-i) shows the time evolution of the mass, virial parameter, and bound mass fraction of the gas and stars. With these three plots we can track the degree of gas dispersal. From the mass plot we see that by the end of star formation, only the M4 cluster has expelled a significant fraction of the cloud. By $1.25 t_{\rm ff}$, none of the gas is bound. Although the M5 cluster has only expelled 10%of its gas by the end of the simulation, it is well on its way to full gas expulsion, as only 10% of the gas still on the grid is bound. The gas in the M4 and M5 systems become super-virial by a free-fall time. The gas in M5 takes $\sim 10\%$ longer to become unbound, but progresses identically to the M4 gas. The gas in M6 differs significantly as it remains sub-virial even beyond one free-fall time. The potential well created by the massive cluster is enough to keep the remaining gas infalling.

The only star cluster that is starting to disperse is the M4 cluster (Fig. 4(i)). The other two star clusters remain fully bound, although their virial parameter follows the same trajectory of the M4 cluster. The stars in all three systems are slightly super-virial by a free-fall time (Fig. 4(h)). The dispersal time of the gas and stars increases with initial cloud mass as there is more gravity for the stellar feedback to counteract. Although massive clusters have more stars injecting feedback, the increasing gravity overpowers the feedback. At high densities, where the potential well depth exceeds the sound speed of ionized gas, ionization feedback cannot disperse gas, while the short free-fall time assembles dense gas more quickly than feedback can work against gravity.

Simulation data and plotting scripts will be available for download through the online publication of the paper.



Fig. 4. Global properties of the clusters and gas over time for models M4 (orange), M5 (maroon), and M6 (blue-violet) in units of free-fall time $t_{\rm ff}$ of the initial cloud (given in Table 1). From top left to bottom right: (a) SFR, where the transparent lines show the SFR at each star formation event, and the solid lines give the SFR smoothed using a Gaussian filter with $\sigma = 0.005t_{\rm ff}$. (b) SFE of the cloud, the ratio of total stellar mass formed to the initial gas mass. (c) Most massive star formed. (d) Number of formed stars. Dashed line: actual number of stars that would form from sampling the IMF given the amount of gas mass collected for star formation by sink particles. Solid line: number of stars followed in TORCH after the sampled stellar population below $4 M_{\odot}$ has been agglomerated. Dotted line: number of stars above $20 M_{\odot}$ on the grid that are generating feedback. The number of stars can drop due to SNe, mass loss, or exiting the grid. (e) Three-dimensional stellar velocity dispersion. (f) Half-mass radius of the entire star cluster. (g) Total mass of stars (dashed line) and gas (solid line) on the grid. (h) Virial parameter of stars (dashed line) and gas (solid line), where $\alpha_{\rm vir} = 0.5$ is the equilibrium value. (i) Fraction of mass bound for stars (dotted line) and gas (solid line).

Run	M _*	N_{\star}	ϵ_{\star}	$\langle SFR \rangle$	SFR _{pk}
M4	3,180	5,332	0.32	0.012	0.413
M5	64,733	$112,\!661$	0.65	0.063	5.535
M6	845,815	$1,\!468,\!969$	0.85	1.846	392.0

Notes. Column values and units: stellar mass $[M_{\odot}]$, number of stars formed from IMF sampling (number of stars in the simulation after agglomeration of stars $< 4 \, M_{\odot}$ is $\sim 10\%$ of this value), SFE $[M_{\star}/M_c]$, average SFR $[M_{\odot} \, \mathrm{yr}^{-1}]$, peak SFR $[M_{\odot} \, \mathrm{yr}^{-1}]$.

4. Discussion

4.1. Observations

Galactic surveys of embedded clusters in the Milky Way typically find the SFE to be $\leq 30\%$ (Lada & Lada 2003), with some studies finding lower values of $\leq 8\%$ (Evans et al. 2009; Peters et al. 2011). The M4 cloud, which is a good representative of galactic GMCs², agrees well with this SFE albeit at the high end of observed values. This is most likely due to the low virial parameter used, which is appropriate for M6 clouds but lower than the average value of $\alpha_v = 1$ in M4 clouds.

 $^{^{2}}$ See Rice et al. (2016) for a catalogue of Milky-Way molecular cloud properties.

The higher mass M5 and M6 clouds, however, have SFEs well above 30%. While there are no Milky-Way analogues to the M6 cloud, there are a few for M5. There is the W43 GMC with $1.32 \times 10^5 \,\mathrm{M_{\odot}}$ of gas within $R \sim 10 \,\mathrm{pc}$ (Lin et al. 2016), similar to the M5 cloud with $R = 11.7 \,\mathrm{pc}$. The W49 star forming region has a central YMC with stellar mass $\gtrsim 5 \times 10^4 \,\mathrm{M_{\odot}}$ and gas mass $\sim 2 \times 10^5 \,\mathrm{M_{\odot}}$ and $\sim 1.1 \times 10^6 \,\mathrm{M_{\odot}}$ within 6 and 60 pc respectively (Galván-Madrid et al. 2013). This gives a current SFE of 20% in the inner region. With its ongoing star formation, W49 may reach SFEs comparable to the 65% reached in the M5 cloud.

Though conditions required to form the M6 cloud are not observed in the Milky Way, they are present in other galaxies. For example, the disks of gas-rich high redshift galaxies can be violently unstable and are thought to form such clouds (see Tacconi et al. 2020). We can now directly observe the high redshift environment of forming GCs with JWST. Recent JWST observations uncovered "younger" populations of GCs in galaxies at redshift z = 0.38 (Harris & Reina-Campos 2023), and more are expected to observe GCs up to z = 1 without lensing (Reina-Campos & Harris 2023). With lensing, clumps that are likely proto-GCs can be observed at redshift z > 1 (Adamo et al. 2023; Claeyssens et al. 2023).

Another such situation that can form M6-like GMCs is major galaxy mergers with small mass ratios. Tidal interactions of major galaxies are linked to bursts in star formation (Larson & Tinsley 1978; Lonsdale et al. 1984; Barton et al. 2000; Ellison et al. 2008; Renaud et al. 2019). Since most massive galaxies are believed to undergo at least one merger in their lifetime, this is not a rare occurrence. Galaxy mergers have been suggested as the progenitors of YMCs and younger GCs (Ashman & Zepf 1992; van den Bergh 2001). Note that this only applies to major galaxy interactions: minor galaxy interactions with large mass ratios produce little to no enhancement of the overall SFR (Cox et al. 2008; Tress et al. 2020).

In the interacting Antennae galaxies, a cloud named the "Firecracker" cloud resembling M6 was observed by Whitmore et al. (2014). Finn et al. (2019) constrained its mass and characteristic radius to $(1-9) \times 10^6 M_{\odot}$ and 22 pc. The Firecracker cloud is in the very early stages of star formation, as it is estimated to have only formed $M_{\star} \leq 10^4 M_{\odot}$. This is less than 10% of the expected stellar mass of the final star cluster (Johnson et al. 2015). These observations show that progenitor clouds similar to M6 can form before any significant amount of star formation occurs.

A survey of the molecular clouds in the Antennae galaxies done by Wei et al. (2012) revealed two populations of MCs, with a distinct break in the differential mass function at $\log(M_{\rm cl}/{\rm M}_{\odot}) = 6.5$. Clouds above this mass were found in the regions of intense star formation, while the lower mass clouds were in more dormant regions. The large velocities seen in the high SF regions suggest compression by shocks, supporting the idea that galaxy mergers lead to high mass GMCs that become sites of extreme star formation.

Finn et al. (2019) measured the velocity dispersion in the Firecracker cloud and found it to be neither in virial equilibrium nor free-fall. They concluded that there must be a high pressure background to contain the gas at such high densities in equilibrium.

We compare this velocity dispersion to those in our clouds over time to test whether we reach such high ve-



Fig. 5. The size-line width coefficient vs. surface density for the Firecracker cloud (Finn et al. 2019) and the M4, M5, and M6 clouds. The four red diamonds are observations, done using apertures (1-4) with radii R = 6.4, 15, 26, and 37 pc. The other points are our simulations, with connected points corresponding to apertures (1-4) of radii R = 5, 10, 15 and 20 pc. The colors indicate how much stellar mass has been produced (with empty points indicating no star formation). The lines correspond to virial equilibrium and free-fall as labeled.

locities through the addition of stellar feedback to free-fall collapse alone or whether a high pressure background is indeed needed. The results of this comparison are shown in Figure 5. We plot the size-line width coefficient σ_v^2/R against surface density $\Sigma = M/\pi R^2$ for each of our clouds to compare with the observations from Finn et al. (2019). The observations are shown by the red points in Figure 5 corresponding to 4 aperture sizes they used for R: 6.4, 15, 26, and 37 pc. We use four smaller aperture sizes: 5, 10, 15, and 20 pc, as our cloud is half the radius of the Firecracker. We plot four times for each of our simulations corresponding to $t = 0.0, 0.5, 1.0, 1.25 t_{\rm ff}$, with a line connecting the points showing the set of apertures. The apertures for each set increase from right to left with decreasing surface density. The colors of the points indicate the amount of stellar mass formed, colored white when no SF has occurred vet. The sizes indicate the initial gas mass of the cloud with the smallest being the M4 simulations and the largest being M6.

Though the Firecracker cloud is as massive as M6, its surface density is more comparable to M5. This is why we see an overlap between M5 and the Firecracker cloud when M5 has formed ~ $10^4 M_{\odot}$ of stars. This is also the amount of stellar mass estimated to have already formed in the Firecracker cloud. This supports the possibility that the Firecracker gas velocities could be caused by the contribution of stellar feedback to the velocity dispersion in addition to free-fall collapse. The fast dynamical evolution of our models suggests that these are not equilibrium objects, making it unnecessary to invoke a high-pressure background to keep the cloud from expanding. This suggests objects like the Firecracker cloud can form from collapse with observed velocity dispersions without invoking a high pressure background medium.

4.2. Other Simulations

For massive star clusters to form, they must survive the epoch of gas dispersal and remain bound. Numerical studies done by Geyer & Burkert (2001) found that if the stars are initially in virial equilibrium with the remaining gas, only clusters with SFE \geq 50% remain bound against the outflow of the gas. Li et al. (2018) found in their cosmological galaxy formation models that though galactic properties are unaffected by varying ϵ_{\star} , the properties of star clusters are. Particularly, they found that the initial bound fraction of stars increases with ϵ_{\star} and cloud mass. Farias et al. (2023) ran cluster formation models from low-mass $(2\times 10^4\,{\rm M_{\odot}})$ clouds and found SFE and gas expulsion time correlate with global bound fraction, with all SFEs $\leq 20\%$ and all bound fractions $\leq 40\%$. Although it is still possible to form bound clusters with low SFE, these studies imply massive bound star clusters were most likely formed with high SFEs.

Menon et al. (2023) also found high SFEs of ~ 80% for $10^6 M_{\odot}$ clouds with feedback in the form of radiation pressure solved using a variable Eddington tensor approach as opposed to our ray-tracing method. In this density regime, radiation pressure is the dominant feedback mechanism³. They concluded that radiation pressure simply cannot regulate star formation for clouds with surface densities $\Sigma \gtrsim 10^3 M_{\odot} \ pc^{-2}$.

Our results are more constraining: we include more feedback physics and we still achieve SFEs of $\epsilon_{\star} > 80\%$. Our M6 cloud is above this surface density with $\Sigma = 2.3 \times 10^3 \,\mathrm{M_{\odot} \ pc^{-2}}$. They also tested a larger $10^6 \,\mathrm{M_{\odot}}$ cloud with roughly the same surface density as our M5 cloud, and find an SFE of $\epsilon_{\star} \approx 60\%$ comparable to the SFE of our M5 cluster of $\epsilon_{\star} = 65\%$.

Other simulations of massive star cluster formation with initial cloud mass of $10^6 \,\mathrm{M_{\odot}}$ find high SFEs of ~ 65% (Grudić et al. 2018) and 38% (Kim et al. 2018) for surface densities of $\Sigma = 1.27 \times 10^4 \,\mathrm{M_{\odot}} \,\mathrm{pc^{-2}}$ and $\Sigma = 500 \,\mathrm{M_{\odot}} \,\mathrm{pc^{-2}}$, respectively. Kim et al. (2018) found an even higher SFE of 51% for a $10^5 \,\mathrm{M_{\odot}}$ cloud but with a surface density of $\Sigma = 1.27 \times 10^3 \,\mathrm{M_{\odot}} \,\mathrm{pc^{-2}}$. The SFE in both studies increases strongly with surface density and slightly with initial cloud mass. These results combined with ours suggest that the formation of bound super-massive star clusters requires both high cloud mass and high surface density.

Protostellar jets are not implemented in our feedback model, although they contribute to the dispersal of gas around low-mass stars at small scales. Due to the quantity of low-mass stars and the collimated shape of the outflow, jets are drivers of turbulence at large scales in GMCs (e.g. Nakamura & Li 2007; Federrath 2015). Despite their ubiquity, they do not prevent gas in high-density GMCs from forming stars eventually nor contain the power needed to disperse GMCs (see Chevance et al. 2023). On the other hand, studies also show that jets are an important factor in slowing the growth rate of the integrated SFE, though the final SFE is not known due to the duration of the simulations (Federrath 2015; Appel et al. 2022).

Guszejnov et al. (2021) performed simulations of starby-star cluster formation from low mass clouds $(2 \times 10^4 M_{\odot})$ with stellar feedback, including protostellar jets as well as radiation, winds, and SNe. Simulations were repeated that



Fig. 6. WARPFIELD evolution of shell radius vs time with different initial SFE, with the same parameters described in M6. In all cases, stellar feedback is inefficient in dispersing the surrounding dense cloud, and the shell eventually undergoes re-collapse.

isolated each form of feedback. They found jets to be important in regulating the growth of low-mass stars and constraining the IMF. Radiation and jets were the primary form of feedback that slowed star formation and dispersed the cloud. However, the simulations were not run until the end of star formation, so the degree to which each affect the final SFE is uncertain. This suggests the SFE of M4 may be overestimated due the exclusion of jets. The effect that jets would have on more massive clouds is unclear. Analytic work by Matzner (2002) suggests that the density of more massive clouds would be resistant to dispersal by jets, consistent with simulations by Guszejnov et al. (2022).

After the M6 simulation resulted in such a high SFE, we wanted to verify that this many stars would not just immediately blow away the gas. To do this, we have directly compared our 3D results to a followup calculation using the 1D code Winds And Radiation Pressure: Feedback Induced Expansion. colLapse and Dissolution (WARPFIELD: Rahner et al. 2019). This code models the effect of stellar feedback from young clusters on their natal gas cloud. WARPFIELD is designed to solve for the self-consistent motion of a 1D spherical gas shell evolving under the influence of feedback mechanisms including stellar winds, SNe, and radiation pressure, with consideration of gravity. We run WARPFIELD using the same physical conditions as chosen for the M6 run (i.e., mass, density, temperature), with the addition that we vary the SFE from $\epsilon_{\star} = 0.1$ –0.9 in bins of 0.1, as shown in Figure 6.

For all SFE values, the shocked gas eventually recollapses. At this high density, the included feedback is not strong enough to completely disperse the cloud. In the 1D model, all feedback occurs at a single point, so it is more effective than in our star-by-star three-dimensional model, as in multiple dimensions channels that vent thermal energy can exist. Nevertheless, the gas still re-collapses, promoting further star formation. The expanding gas is not accelerated fast enough to escape the deep potential well of the massive cloud and the cluster that forms from it.

Our M6 cluster mostly closely resembles the WARP-FIELD runs with $\epsilon_{\star} = 80\%\&90\%$ in Figure 6, which reach

 $[\]overline{}^{3}$ See extended data Figure 5 of Howard et al. (2018) and Figure 12 of Krumholz et al. (2019).

a maximum radius of $R \sim 55$ pc& at $t \sim 6.5$ Myr, and collapse back to R = 0 pc by $t \sim 11.25$ Myr.

However, our results do suggest that more dispersed star formation leading to increased energy dissipation by radiative cooling may not even allow that much expansion. To resolve SFE well, feedback must be modelled for individual stars instead of for entire clusters. Approximating feedback as a sum for an entire cluster underestimates the SFE.

5. Conclusions

We performed numerical simulations of star cluster formation from gas clouds that run until star formation ceases or slows significantly due to stellar feedback dispersing any remaining gas. We tested initial cloud masses of 10^4 , 10^5 , and $10^6 M_{\odot}$ with radius $R = 11.7 \,\mathrm{pc}$, holding all other characteristics of the initial cloud and simulation parameters the same. We analyzed the star formation histories and followed the evolution of the gas and forming star clusters. From this study, we conclude the following:

- GMCs with surface density $\Sigma \ge 10^2 \,\mathrm{M_{\odot} \ pc^{-2}}$ and mass $M_c \ge 10^5 \,\mathrm{M_{\odot}}$ can form fully bound star clusters with stellar mass $M_{\star} \ge 10^4 \,\mathrm{M_{\odot}}$ with high SFE $\epsilon_{\star} \ge 65\%$ over a short time $t_{\rm sf} \approx 1t_{\rm ff}$, as seen by M5 and M6. The lower mass and density M4 cloud forms a cluster with a lower bound mass fraction of 60%.
- The Firecracker cloud in the Antennae galaxies, with a mass of $1-9\times10^6$ M_☉ and a radius of 22 pc (Finn et al. 2019) is a close analog to our M6 cloud, though with a surface density more closely matching our M5 cloud. From our results we can estimate that the Firecracker cloud will convert 65–85% of its mass into stars within a free-fall time, and that it will form a YMC.
- It has been suggested that the Firecracker cloud must be surrounded by a high pressure medium to contain it because of its high surface density and size-line width coefficient σ_v^2/R (Johnson et al. 2015; Finn et al. 2019). However, the M5 cluster reaches the same values by the time it forms $M_{\star} \approx 10^4 \,\mathrm{M}_{\odot}$ worth of stars, the same amount of stellar mass estimated to have formed in the Firecracker cloud. This suggests another possibility: rather than being an equilibrium object confined in a high pressure environment, the Firecracker cloud is actually dynamically collapsing and forming stars, and the high velocity dispersion of the gas is from the combination of free-fall collapse and stellar feedback.
- Star formation from GMCs is capable of achieving up to 85% efficiency at high densities. Our M6 cloud is the most efficient of our models, converting $\epsilon_{\star} = 85\%$ of its gas into stars. Even with hundreds of massive stars producing feedback, the short timescale of gravitational collapse for dense massive clouds renders the stellar feedback inefficient at slowing early star formation. However, even at much lower densities and masses, the M5 and M4 cloud achieved high SFEs of $\epsilon_{\star} = 65\%$ and 30% respectively.
- The M4 cloud is a typical mass and size of Milky-Way GMCs. The SFE of M4 matches the maximum observed SFE values. This high SFE could be because of the low initial virial parameter of the cloud. Alternatively, the exclusion of the protostellar jet feedback mechanism may be important for M4-like clouds, as suggested, for example, by Chevance et al. (2023). Further studies

Article number, page 12 of 18

must be done to constrain the effect of varying the virial parameter and including protostellar jets on integrated SFE.

- Star formation is fast in our models of clouds with low α_v . Regardless of initial mass or density, the majority of star formation occurs within the first global free-fall time of the collapsing GMC. Collapse occurs and produces stars so rapidly that by the time stellar feedback is prevalent and strong enough to clear dense gas, most of the star formation has already transpired. The speed of star formation may also depend strongly on initial virial parameter and inclusion of jets.
- A one-dimensional stellar feedback model WARPFIELD was run using the same mass and density as the M6 simulation. In it the gas re-collapses even for SFEs up to 90%. Even centralized feedback cannot expel the gas from the potential well of the massive cluster that forms. The WARPFIELD results indicate that the expanding gas shell for $\epsilon_{\star} = 85\%$ collapses back to R = 0 by 11 Myr.
- Including feedback for individual stars rather than adding the total energy for the cluster at a single point is important for correctly constraining star formation histories. Modelling individual stellar feedback spreads the feedback energy enough to greatly reduce its effectiveness at clearing the natal gas because of the resulting enhanced radiative cooling. Models that add stellar feedback for the entire star cluster at a single point appear to overestimate the effect of the feedback on the gas and the star formation timescale and to underestimate the final SFE.

In conclusion, bound massive star clusters like YMCs and GCs readily form from high mass, dense GMCs. These GMCs can become this dense and massive naturally, even in the present day, as shown, for example, by the Firecracker cloud in the Antennae galaxies. In the early Universe where galaxies were much more gravitationally unstable these conditions would be much more common. The subsequent star formation from these dense high mass clouds is highly efficient, converting $\geq 40\%$ of the gas mass into stars. The short timescales of star formation render stellar feedback unable to significantly slow star formation, leading to integrated efficiencies as high as 85% for more massive clouds. After their formation, the clusters born in these environments remain bound after 90% of the gas is expelled. These results indicate that this mode of star formation is a probable path for the formation YMCs and GCs in the present and early Universe, respectively.

Acknowledgements. B.P. was partly supported by a fellowship from the International Max Planck Research School for Astronomy and Cosmic Physics at the University of Heidelberg (IMPRS-HD). B.P. is grateful to Steven Rieder for his support with using PETAR in AMUSE. B.P. would also like to thank Gastón Escobar and Michela Mapelli for the helpful discussions that led to solving a persistent bug allowing continuation of the simulations. M.-M.M.L., B.P., and A.T. were partly supported by NSF grants AST18-15461 and AST23-07950. C.C.C. is supported by a Canada Graduate Scholarship - Doctoral (CGS D) from the Natural Sciences and Engineering Research Council of Canada (NSERC). This work used Stampede 2 at TACC through allocation PHY220160 from the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) program, which is supported by National Science Foundation grants 21-38259, 21-38286, 21-38307, 21-37603, and 21-38296. The code development that facilitated this study was done on Snellius through the Dutch National Supercomputing Center SURF grants 15220 and 2023/ENW/01498863. S.A. acknowledges the support of NSF grant AST-2009679. M.W. acknowledges the support of NOVA project 10.2.5.12. R.S.K. and

S.C.O.G. acknowledge financial support from the European Research Council via the ERC Synergy Grant "ECOGAL" (project ID 855130), from the Heidelberg Cluster of Excellence (EXC 2181 - 390900948) "STRUCTURES", funded by the German Excellence Strategy, and from the German Ministry for Economic Affairs and Climate Action in project "MAINN" (funding ID 50002206). The team in Heidelberg also thanks The Länd and the German Science Foundation (DFG) for computing resources provided in bwHPC supported by grant INST 35/1134-1 FUGG and for data storage at SDS@hd supported by grant INST 35/1314-1 FUGG. L.W. thanks the National Natural Science Foundation of China for support through grants 21BAA00619, 12073090 and 12233013, and the one-hundred-talent project of Sun Yat-sen University, the Fundamental Research Funds for the Central Universities, Sun Yat-sen University (22hytd09).

References

- Adamo, A., Usher, C., Pfeffer, J., & Claeyssens, A. 2023, MNRAS, 525, L6
- Adamo, A., Zeidler, P., Kruijssen, J. M. D., et al. 2020, Space Sci. Rev., 216, 69
- Appel, S. M., Burkhart, B., Semenov, V. A., Federrath, C., & Rosen, A. L. 2022, ApJ, 927, 75
- Arthur, S. J. 2007, in Astrophysics and Space Science Proceedings, Vol. 1, Diffuse Matter from Star Forming Regions to Active Galaxies - A Volume Honouring John Dyson, 183
- Arthur, S. J., Dyson, J. E., & Hartquist, T. W. 1993, MNRAS, 261, 425
- Arthur, S. J., Henney, W. J., & Dyson, J. E. 1996, A&A, 313, 897
- Ashman, K. M. & Zepf, S. E. 1992, ApJ, 384, 50
- Baczynski, C., Glover, S. C. O., & Klessen, R. S. 2015, Monthly Notices of the Royal Astronomical Society, 454, 380
- Barnes, J. & Hut, P. 1986, Nature, 324, 446
- Barton, E. J., Geller, M. J., & Kenyon, S. J. 2000, ApJ, 530, 660
- Bate, M. R., Bonnell, I. A., & Price, N. M. 1995, MNRAS, 277, 362
- Bochkarev, N. G. 1988, Nature, 332, 518
- Brodie, J. P. & Strader, J. 2006, ARA&A, 44, 193
- Chevance, M., Krumholz, M. R., McLeod, A. F., et al. 2023, in Astronomical Society of the Pacific Conference Series, Vol. 534, Protostars and Planets VII, ed. S. Inutsuka, Y. Aikawa, T. Muto, K. Tomida, & M. Tamura, 1
- Chu, Y.-H., Guerrero, M. A., Gruendl, R. A., García-Segura, G., & Wendker, H. J. 2003, ApJ, 599, 1189
- Claeyssens, A., Adamo, A., Richard, J., et al. 2023, MNRAS, 520, 2180
- Colella, P. & Woodward, P. R. 1984, Journal of Computational Physics, 54, 174
- Colín, P., Vázquez-Semadeni, E., & Gómez, G. C. 2013, Monthly Notices of the Royal Astronomical Society, 435, 1701
- Cournoyer-Cloutier, C., Sills, A., Harris, W. E., et al. 2023, MNRAS, 521, 1338
- Cox, T. J., Jonsson, P., Somerville, R. S., Primack, J. R., & Dekel, A. 2008, MNRAS, 384, 386
- Crutcher, R., Heiles, C., & Troland, T. 2003, in Lecture Notes in Physics, Vol. 614, Turbulence and Magnetic Fields in Astrophysics, ed. E. Falgarone & T. Passot (Berlin, Heidelberg: Springer), 155-181
- Crutcher, R. M., Wandelt, B., Heiles, C., Falgarone, E., & Troland, T. H. 2010, ApJ, 725, 466
- Dekel, A., Sarkar, K. C., Birnboim, Y., Mandelker, N., & Li, Z. 2023, MNRAS, 523, 3201
- Dubey, A., Antypas, K., Calder, A. C., et al. 2014, The International Journal of High Performance Computing Applications, 28, 225
- Dyson, J. E. & Hartquist, T. W. 1992, Astrophysical Letters and Communications, 28, 301
- Ellison, S. L., Patton, D. R., Simard, L., & McConnachie, A. W. 2008, AJ, 135, 1877
- Evans, Neal J., I., Dunham, M. M., Jørgensen, J. K., et al. 2009, ApJS, 181, 321
- Fall, S. M. 2006, ApJ, 652, 1129
- Fall, S. M., Chandar, R., & Whitmore, B. C. 2005, ApJ, 631, L133
- Fall, S. M. & Zhang, Q. 2001, ApJ, 561, 751
- Farias, J. P., Offner, S. S. R., Grudić, M. Y., et al. 2023, MNRAS
- Federrath, C. 2015, MNRAS, 450, 4035
- Federrath, C., Banerjee, R., Clark, P. C., & Klessen, R. S. 2010, ApJ, 713.269

- Federrath, C., Banerjee, R., Clark, P. C., & Klessen, R. S. 2010, The Astrophysical Journal, 713, 269
- Field, G. B., Goldsmith, D. W., & Habing, H. J. 1969, ApJ, 155, L149 Finn, M. K., Johnson, K. E., Brogan, C. L., et al. 2019, ApJ, 874, 120
- Fryxell, B., Olson, K., Ricker, P., et al. 2000, ApJs, 131, 273
- Fujii, M., Iwasawa, M., Funato, Y., & Makino, J. 2007, PASJ, 59,
- 1095
- Fujii, M. S. & Portegies Zwart, S. 2011, Science, 334, 1380
- Galván-Madrid, R., Liu, H. B., Zhang, Z. Y., et al. 2013, ApJ, 779, 121
- Geyer, M. P. & Burkert, A. 2001, MNRAS, 323, 988
- Girichidis, P., Offner, S. S. R., Kritsuk, A. G., et al. 2020, Space Sci. Rev., 216, 68
- Goodwin, S. P., Whitworth, A. P., & Ward-Thompson, D. 2004, A&A, 414, 633
- Grudić, M. Y., Guszejnov, D., Hopkins, P. F., Offner, S. S. R., & Faucher-Giguère, C.-A. 2021, MNRAS, 506, 2199
- Grudić, M. Y., Hopkins, P. F., Faucher-Giguère, C.-A., et al. 2018, MNRAS, 475, 3511
- Grudić, M. Y., Hopkins, P. F., Faucher-Giguère, C.-A., et al. 2018, Monthly Notices of the Royal Astronomical Society, 475, 3511
- Guszejnov, D., Grudić, M. Y., Hopkins, P. F., Offner, S. S. R., & Faucher-Giguère, C.-A. 2021, MNRAS, 502, 3646 Guszejnov, D., Grudić, M. Y., Offner, S. S. R., et al. 2022, MNRAS,
- 515, 4929
- Harris, W. E. & Reina-Campos, M. 2023, MNRAS, 526, 2696
- Hartquist, T. W. & Dyson, J. E. 1996, Ap&SS, 245, 263
- Hartquist, T. W., Dyson, J. E., Pettini, M., & Smith, L. J. 1986, MNRAS, 221, 715
- Heiles, C. 1976, Annual Review of Astronomy and Astrophysics, 14, 1
- Heitsch, F., Mac Low, M.-M., & Klessen, R. S. 2001, ApJ, 547, 280 Howard, C. S., Pudritz, R. E., & Harris, W. E. 2017, Monthly Notices
- of the Royal Astronomical Society, 470, 3346
- Howard, C. S., Pudritz, R. E., & Harris, W. E. 2018, Nature Astronomy, 2, 725
- Iwasawa, M., Namekata, D., Nitadori, K., et al. 2020, PASJ, 72, 13 Iwasawa, M., Tanikawa, A., Hosono, N., et al. 2016, PASJ, 68, 54
- Johnson, K. E., Leroy, A. K., Indebetouw, R., et al. 2015, ApJ, 806, 35

- Kauffmann, J., Pillai, T., & Goldsmith, P. F. 2013, ApJ, 779, 185 Kim, J.-G., Kim, W.-T., & Ostriker, E. C. 2018, ApJ, 859, 68 Kim, J.-G., Kim, W.-T., Ostriker, E. C., & Skinner, M. A. 2017, The Astrophysical Journal, 851, 93
- Klessen, R. S. & Glover, S. C. O. 2016, in Saas-Fee Advanced Course, Vol. 43, Saas-Fee Advanced Course, ed. Y. Revaz, P. Jablonka, R. Teyssier, & L. Mayer, 85
- Kolmogorov, A. 1941, Akademiia Nauk SSSR Doklady, 30, 301 Krause, M. G. H., Offner, S. S. R., Charbonnel, C., et al. 2020, Space Sci. Rev., 216, 64
- Kroupa, P. 2002, Science, 295, 82
- Krumholz, M. R., McKee, C. F., & Bland-Hawthorn, J. 2019, ARA&A, 57, 227
- Kudritzki, R.-P. & Puls, J. 2000, ARA&A, 38, 613 Lada, C. J. & Lada, E. A. 2003, ARA&A, 41, 57
- Lahén, N., Naab, T., Johansson, P. H., et al. 2019, ApJ, 879, L18
- Lancaster, L., Ostriker, E. C., Kim, J.-G., & Kim, C.-G. 2021, ApJ, 914, 89
- Larson, R. B. 1981, MNRAS, 194, 809
- Larson, R. B. & Tinsley, B. M. 1978, ApJ, 219, 46
- Lewis, S. C., McMillan, S. L. W., Low, M.-M. M., et al. 2023, ApJ,
- 944, 211
- Li, H., Gnedin, O. Y., & Gnedin, N. Y. 2018, ApJ, 861, 107
- Li, H., Vogelsberger, M., Marinacci, F., & Gnedin, O. Y. 2019, Monthly Notices of the Royal Astronomical Society, 487, 364
- Lin, Y., Liu, H. B., Li, D., et al. 2016, ApJ, 828, 32
- Lonsdale, C. J., Persson, S. E., & Matthews, K. 1984, ApJ, 287, 95
- Mac Low, M.-M. & Klessen, R. S. 2004, Rev. Mod. Phys., 76, 125
- Makino, J. & Aarseth, S. J. 1992, PASJ, 44, 141
- Matzner, C. D. 2002, ApJ, 566, 302
- McKee, C. F. & Ostriker, E. C. 2007, Annual Review of Astronomy and Astrophysics, 45, 565
- McKee, C. F. & Williams, J. P. 1997, The Astrophysical Journal, 476, 144
- McMillan, S., Portegies Zwart, S., van Elteren, A., & Whitehead, A. 2012, in Astronomical Society of the Pacific Conference Series, Vol. 453, Advances in Computational Astrophysics: Methods, Tools, and Outcome, ed. R. Capuzzo-Dolcetta, M. Limongi, & A. Tornambè, 129

Article number, page 13 of 18

- Menon, S. H., Federrath, C., & Krumholz, M. R. 2023, MNRAS, 521, 5160
- Miyoshi, T. & Kusano, K. 2005, Journal of Computational Physics, 208, 315
- Mouschovias, T. C. 1991, Cosmic Magnetism and the Basic Physics of the Early Stages of Star Formation (Dordrecht: Springer Netherlands), 61–122
- Mouschovias, T. C. & Spitzer, L., J. 1976, ApJ, 210, 326
- Murray, N. & Rahman, M. 2009, The Astrophysical Journal, 709, 424 Nakamura, F. & Li, Z.-Y. 2007, ApJ, 662, 395
- Peters, T., Banerjee, R., Klessen, R. S., & Mac Low, M.-M. 2011, ApJ, 729, 72
- Pittard, J. M., Hartquist, T. W., & Dyson, J. E. 2001, A&A, 373, 1043
- Plotly Technologies Inc. 2015, Collaborative data science
- Portegies Zwart, S. & McMillan, S. 2018, Astrophysical Recipes, 2514-3433 (IOP Publishing)
- Portegies Zwart, S., McMillan, S., Harfst, S., et al. 2009, New Astronomy, 14, 369
- Portegies Zwart, S. F., McMillan, S. L. W., & Gieles, M. 2010, ARA&A, 48, 431
- Portegies Zwart, S. F. & Verbunt, F. 1996, A&A, 309, 179
- Price, D. J. & Bate, M. R. 2008, MNRAS, 385, 1820
- Rahner, D., Pellegrini, E. W., Glover, S. C. O., & Klessen, R. S. 2019, MNRAS, 483, 2547
- Reina-Campos, M. & Harris, W. E. 2023, arXiv e-prints, arXiv:2310.02307
- Renaud, F. 2020, in Star Clusters: From the Milky Way to the Early Universe, ed. A. Bragaglia, M. Davies, A. Sills, & E. Vesperini, Vol. 351, 40–46
- Renaud, F., Agertz, O., & Gieles, M. 2017, MNRAS, 465, 3622
- Renaud, F., Bournaud, F., Agertz, O., et al. 2019, A&A, 625, A65
- Rice, T. S., Goodman, A. A., Bergin, E. A., Beaumont, C., & Dame, T. M. 2016, ApJ, 822, 52
- Simpson, C. M., Bryan, G. L., Hummels, C., & Ostriker, J. P. 2015, ApJ, 809, 69
- Smith, L. J., Pettini, M., Dyson, J. E., & Hartquist, T. W. 1984, MNRAS, 211, 679
- Sormani, M. C., Treß, R. G., Klessen, R. S., & Glover, S. C. O. 2017, MNRAS, 466, 407
- Strittmatter, P. A. 1966, MNRAS, 132, 359
- Su, K.-Y., Hopkins, P. F., Hayward, C. C., et al. 2018, Monthly Notices of the Royal Astronomical Society, 480, 1666
- Sun, J., Leroy, A. K., Rosolowsky, E., et al. 2022, AJ, 164, 43
- Tacconi, L. J., Genzel, R., & Sternberg, A. 2020, ARA&A, 58, 157
- Tress, R. G., Smith, R. J., Sormani, M. C., et al. 2020, MNRAS, 492, 2973
- Truelove, J. K., Klein, R. I., McKee, C. F., et al. 1997, ApJ, 489, L179 van den Bergh, S. 2001, ApJ, 559, L113
- Vink, J. S., de Koter, A., & Lamers, H. J. G. L. M. 2000, A&A, 362, 295
- Wall, J. E., Mac Low, M.-M., McMillan, S. L. W., et al. 2020, ApJ, 904, 192
- Wall, J. E., McMillan, S. L. W., Mac Low, M.-M., Klessen, R. S., & Portegies Zwart, S. 2019, ApJ, 887, 62
- Wang, L., Iwasawa, M., Nitadori, K., & Makino, J. 2020a, MNRAS, 497, 536
- Wang, L., Nitadori, K., & Makino, J. 2020b, MNRAS, 493, 3398
- Weaver, R., McCray, R., Castor, J., Shapiro, P., & Moore, R. 1977, ApJ, 218, 377
- Wei, L. H., Keto, E., & Ho, L. C. 2012, ApJ, 750, 136
- Weidner, C., Kroupa, P., & Bonnell, I. A. D. 2009, Monthly Notices of the Royal Astronomical Society, 401, 275
- Whitmore, B. C., Brogan, C., Chandar, R., et al. 2014, ApJ, 795, 156
 Whitmore, B. C., Chandar, R., Rodríguez, M. J., et al. 2023, ApJ, 944, L14
- Wolfire, M. G., McKee, C. F., Hollenbach, D., & Tielens, A. G. G. M. 2003, ApJ, 587, 278
- Wrigge, M. 1999, A&A, 343, 599
- Wrigge, M., Wendker, H. J., & Wisotzki, L. 1994, A&A, 286, 219
- Yan, Z., Jerabkova, T., & Kroupa, P. 2023, A&A, 670, A151

Appendix A: Stellar Properties: Physical times

Figure A.1 reproduces Figure 4 using physical time rather than free-fall times to show global stellar properties over time. This demonstrates how much the duration and intensity of star formation are increased as the cloud mass increases.

Appendix B: Stellar Modifications

B.1. Low-mass Star Agglomeration

Upwards of 10^6 stars can be expected to form from a $10^6 \,\mathrm{M_{\odot}}$ cloud with a peak number density of $n \approx 1000 \,\mathrm{cm^{-3}}$. Even with the best modern N-body codes, evolving this many single stars and higher order stellar systems in such a dense stellar environment with a gravity bridge from each star to the gas in a separate code is immensely computationally taxing. To reduce the strain on the N-body portion of the calculations, we choose to agglomerate all stars under a given mass into gravitational super-star particles of equivalent mass to their sum. We refer to this mass cutoff as the agglomerate mass.

When a sink progresses through the list of stellar masses it will form, stars with masses under $M_{\rm agg}$ are put as ide until the sum of their masses is above M_{agg} . Then a star particle is formed with the summed mass. Figure B.1 shows the reduction in number of stars formed in a cloud for a given agglomerate mass. For our choice of 4 M_{\odot} , we only have 10% of the stars undergoing gravitational interactions compared to the case with no agglomeration. This reduces our N-body execution time by a factor of somewhere between the $10 \log 10$ expected for the tree and 10^3 expected for the direct N portion of the PETAR algorithm. Note that the feedback from these low-mass stars is shown in Appendix B.2 to be negligible compared to that of the higher-mass stars, and in any case TORCH currently only models the feedback from stars $\geq 8 \,\mathrm{M}_{\odot}$ as we neglect jets and the ionizing radiation from such low mass stars is negligible. The primary missing contribution from low-mass stars physically is their mutual gravitational interactions, which could potentially lead to the ejection of some fraction of them. However, the dynamics driven by those lowmass stars is also expected to be negligible in comparison to the effect of gas and more massive stars in the cluster. TORCH simulations with no mass agglomeration were done by Cournoyer-Cloutier et al. (2023), and in analyzing the morphology of clusters they found that the gravitational effects of the gas dominate over any stellar dynamics effect for the overall evolution of the cluster while it remains embedded.

B.2. Feedback Mass Limit

We limit all forms of stellar feedback—winds, radiation, and SNe—to stars above 20 M_{\odot} instead of the value of 8 M_{\odot} (lower bound for SN explosions) usually adopted in TORCH. This is necessary to significantly reduce the number of rays on the grid, which greatly decreases the calculation time and memory overhead for the ray-tracing algorithm. We quantify the effects of excluding radiation and winds from stars with masses below 20 M_{\odot} by comparing the power output in the form of winds and radiation from all stars above 8 M_{\odot} and above 20 M_{\odot} .

The power as a function of mass in the form of EUV radiation, non-ionizing FUV radiation, and stellar wind is shown in Figure B.2. We calculate these powers by taking stars from 8 to 100 M_{\odot} in 1 M_{\odot} increments, evolving them in SEBA for 1 Myr, and summing the energy output of each feedback channel. From this figure we can see that the power output of stellar winds and UV radiation is several orders of magnitude higher for stars above 20 M_{\odot} than for stars closer to $8 M_{\odot}$. Although stars in the 8–20 M_{\odot} mass range still output a considerable amount of FUV radiation, stars above 20 M_{\odot} account for over 80% of the total radiation power.

Although the feedback power is much stronger for stars above $20 \,\mathrm{M}_{\odot}$, stars with masses $8-20 \,\mathrm{M}_{\odot}$ greatly outnumber them. To find the ratio of feedback power for stars below and above 20 M_{\odot} , we convolve the number of stars of each mass with the power output for each stellar mass (Fig. B.3). In the top left histogram, we show the ratio of stars with mass 8–20 $\,\rm M_{\odot}$ to stars with mass 20–100 $\,\rm M_{\odot}$ in all three simulations, sampled at their respective initial free-fall times. All three runs have more stars in the lowermass bin. Note that the number of stars in the two bins is almost equal in the M4 simulation: this is a statistical artifact as there are only 11 stars above 8 M_{\odot} at this point in this simulation. In the top right plot, we show the ratio of total stellar feedback power $P_{\rm FB}$ (excluding SNe) for the stars in the two mass bins considered. We can see that although the lower-mass stars outnumber the higher-mass stars, the higher-mass stars still account for > 80% of the total stellar feedback energy. This shows that only including feedback from stars above 20 $\,\rm M_\odot$ still retains almost all of the feedback energy produced after the formation of all three star clusters.

The bottom panel in Figure B.3 shows the feedback power per mass bin for each separate feedback process. For the EUV radiation and wind feedback, the low mass stars contribute practically nothing to the feedback energy in comparison to the high mass stars. The FUV feedback of low mass stars is not negligible, but is still well below 20% of the total FUV feedback energy from all stars.

The exclusion of SNe for the low mass stars is inconsequential, as our simulations run for $\leq 10\,\rm Myr$ which is roughly the main-sequence lifetime of a $20\,\rm M_{\odot}$ star. Less massive stars would not go SNe in the time-frame of our simulations, so whether we include them or not makes no practical difference.

B.3. Mass-loading Stellar Winds

In TORCH, the stellar wind feedback implementation is inspired by Simpson et al. (2015), using a method of momentum injection, the details of which can be found in Wall et al. (2020). The energy of the cells within the wind injection radius of the star is increased based on the mechanical luminosity of the wind $L_{\rm w} = (1/2)\dot{M}v_{\rm w}^2$, where \dot{M} is the stellar mass loss rate and $v_{\rm w}$ is the terminal wind velocity (Kudritzki & Puls 2000; Vink et al. 2000). The wind injection radius is set by comparing the cell width Δx to the wind termination shock radius (Weaver et al. 1977)

$$R_{\rm w} = 0.74 \left(\frac{\dot{M}}{\rho_0}\right)^{3/10} v_{\rm w}^{1/10} t_{\rm w}^{2/5},\tag{B.1}$$

Article number, page 15 of 18

A&A proofs: manuscript no. main



Fig. A.1. Global properties of the clusters and gas over time for models M4 (orange), M5 (maroon), and M6 (blue-violet) for comparison to Figure 4 where units of free-fall time (see Table 1) are used. From top left to bottom right: (a) SFR, where the transparent lines show the SFR at each star formation event, and the solid lines give the SFR smoothed using a Gaussian filter with $\sigma = 0.005t_{\rm ff}$. (b) SFE of the cloud, the ratio of total stellar mass formed to the initial gas mass. (c) Most massive star formed. (d) Number of formed stars. Dashed line: actual number of stars that would form from sampling the IMF given the amount of gas mass collected for star formation by sink particles. Solid line: number of stars followed in TORCH after the sampled stellar population below $4 M_{\odot}$ has been agglomerated. Dotted line: number of stars above $20 M_{\odot}$ on the grid that are generating feedback. The number of stars can drop due to SNe, mass loss, or exiting the grid. (e) Three-dimensional stellar velocity dispersion. (f) Half-mass radius of the entire star cluster. (g) Total mass of stars (dashed line) and gas (solid line) on the grid. (h) Virial parameter of stars (dashed line) and gas (solid line), where $\alpha_{\rm vir} = 0.5$ is the equilibrium value. (i) Fraction of mass bound for stars (dotted line) and gas (solid line).

where ρ_0 is the background density and t_w is the wind lifetime. If $R_w < \Delta x$ the injection radius is set to Δx , otherwise it is set to a maximum value of $6\sqrt{3}\Delta x$, at which we have found that spherical winds are well resolved. Momentum and energy are conserved when injecting stellar winds.

Within a stellar wind bubble, in dense clumpy regions of star formation such as the ones in our simulations, material will be swept up into the flow of the hot bubble by mass loading processes such as photoevaporation and hydrodynamic ablation (Dyson & Hartquist 1992; Hartquist & Dyson 1996; Pittard et al. 2001; Lancaster et al. 2021). With enough mass loading, the density increase will result in much more efficient cooling and create momentum-driven rather than energy-driven bubbles. The amount of massloading in the case of hydrodynamic ablation depends on the prevalence of dense clumps within the wind region as well as the Mach number M of the flow around the clump. With a supersonic flow, the mass-loading rate saturates. With a subsonic flow, the mass-loading rate is proportional to $M^{4/3}$ (Smith et al. 1984; Hartquist et al. 1986). Accounting for mass loading in stellar wind models has been shown to successfully reproduce the kinematic properties of the observed stellar wind bubble of the Wolf-Rayet star RCW 58 (Arthur et al. 1993, 1996; Arthur 2007).

Simply injecting winds at $v_{\rm w}$ does not account for these mass-loading processes and results in unphysically hot bubbles. Therefore we choose a lower temperature target for our bubbles and lower the wind velocity $v_{\rm w}$ such that the final temperature of the wind bubble is the correct one. We conserve momentum and energy when injecting stellar



Fig. B.1. Fraction of the number of stars formed with agglomeration of stars below the mass on the x-axis over the total number of stars sampled by the IMF. We use $M_{\rm agg} = 4 \, {\rm M}_{\odot}$, which means the number of stars on the grid is 10% of the number that are formed by our IMF.



Fig. B.2. Power of stellar feedback in the form of winds, and FUV and EUV radiation for different stellar masses. Left of the vertical line shows the amount of feedback power lost per star by only including feedback from stars above 20 M_{\odot} .

winds, so while lowering the wind velocity, we also infuse correspondingly more mass into the bubble than the stellar mass loss calculated. This mass is not taken off the grid elsewhere, meaning mass is not entirely conserved.

Observed circumstellar bubbles cooled by suspected mass loading have been seen with temperatures as low as $T_{\rm b} \approx 1.1 \times 10^6$ K in the S308 bubble (Chu et al. 2003). The spectra of the NGC 6888 bubble indicates a dominant component almost as cool, with $T_{\rm b} = 1.5 \times 10^6$ K (Bochkarev 1988; Wrigge et al. 1994; Wrigge 1999).

In the simulations presented here, we heavily mass load the stellar winds to achieve a lower than observed bubble temperature of $T_{\rm b} = 3 \times 10^5$ K. This temperature is at the peak of the cooling curve, so the shocked wind rapidly cools, resulting in smaller, cooler, momentum-driven bubbles instead of hot bubbles filled with 10^6 K gas. We choose to do so because the high sound speeds in hot wind bubbles lower the Courant timestep significantly, making the computation impractical. Since we do not follow X-rays through ray-tracing, having cooler bubbles is adequate. Bubbles at this temperature also do not affect the ionization of the surrounding gas. The primary action of wind feedback during cluster formation is to clear out dense regions of gas so that radiatively ionized H II regions can expand. The only hot gas ($\geq 10^6$ K) on the grid comes from SNe. Capping the temperature of gas on the grid at 3×10^5 K until SNe occur significantly speeds up the simulations.

B.4. Effect on SFE

Limiting the temperature of stellar winds and only modelling feedback for stars above $20 \,\mathrm{M}_{\odot}$ could potentially lead to un-physical runaway star formation. To test this, we reran the M6 model at early times to see if these two approximations are the cause for the extremely high SFE of 85%. For the first new M6 run we raised the wind temperature from $3 \times 10^5 \,\mathrm{K}$ to $5 \times 10^6 \,\mathrm{K}$. For the second test, we both raised the wind temperature and modelled feedback for all stars above $8 \,\mathrm{M}_{\odot}$. The SFE over time for the fiducial M6 run with our standard approximations and the new M6 models are shown in Figure B.4.

The two runs without the approximations that reduce the strength of the stellar feedback have similar SFEs as the M6 model with the aforementioned approximations. This validates our approximations and supports our argument that the high SFE in model M6 is not an artifact of underestimating the strength of stellar feedback.

A&A proofs: manuscript no. main



Fig. B.3. (Top Left) Histogram showing the fractional stellar population of the three runs at one free-fall time, split into the mass regimes of 8–20 and 20–100 M_{\odot} . (Top Right) Fraction of feedback power in each mass regime. (Bottom) Histograms showing the fraction of feedback power for FUV, EUV, and winds in each mass regime. Although there are more lower-mass stars, the feedback produced by them is less than 20% of the total feedback energy for all stars.



Fig. B.4. Star formation efficiency over time for the fiducial M6 cloud with $T_{\rm w} = 300,000$ K and $M_{\rm feedback} \ge 20 \, {\rm M}_{\odot}$ (*cool winds*), the M6 cloud with $T_{\rm w} = 5,000,000$ K and $M_{\rm feedback} \ge 20 \, {\rm M}_{\odot}$ (*hot winds*), and the M6 cloud with $T_{\rm w} = 5,000,000$ K and $M_{\rm feedback} \ge 8 \, {\rm M}_{\odot}$ (*hot winds*, $M_{\rm fb} = 8 \, {\rm M}_{\odot}$).

Article number, page 18 of 18