

Convergence of the Chambolle–Pock Algorithm in the Absence of Monotonicity

Brecht Evens

Puya Latafat

Panagiotis Patrinos

Abstract

The Chambolle–Pock algorithm (CPA), also known as the primal-dual hybrid gradient method, has gained popularity over the last decade due to its success in solving large-scale convex structured problems. This work extends its convergence analysis for problems with varying degrees of (non)monotonicity, quantified through a so-called oblique weak Minty condition on the associated primal-dual operator. Our results reveal novel stepsize and relaxation parameter ranges which do not only depend on the norm of the linear mapping, but also on its other singular values. In particular, in nonmonotone settings, in addition to the classical stepsize conditions, extra bounds on the stepsizes and relaxation parameters are required. On the other hand, in the strongly monotone setting, the relaxation parameter is allowed to exceed the classical upper bound of two. Moreover, we build upon the recently introduced class of semimonotone operators, providing sufficient convergence conditions for CPA when the individual operators are semimonotone. Since this class of operators encompasses traditional operator classes including (hypo)- and co(hypo)-monotone operators, this analysis recovers and extends existing results for CPA. Tightness of the proposed stepsize ranges is demonstrated through several examples.

Keywords. convex/nonconvex optimization · monotone/nonmonotone inclusion problems · Chambolle–Pock algorithm · primal-dual hybrid gradient · semimonotone operators

AMS subject classifications. 47H04 · 49J52 · 49J53 · 65K15 · 90C26.

1 Introduction

This paper considers composite inclusion problems of the form

$$\text{find } x \in \mathbb{R}^n \quad \text{such that} \quad 0 \in T_{\text{p}}x := Ax + L^{\top}BLx, \quad (\text{P-I})$$

where $A : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$, $B : \mathbb{R}^m \rightrightarrows \mathbb{R}^m$ are two (possibly nonmonotone) operators and $L \in \mathbb{R}^{m \times n}$ is a nonzero matrix. Problems of this form emerge naturally in a wide variety of applications in optimization and variational analysis. For instance, in the framework of convex optimization, the first-order optimality condition for minimizing $g(x) + h(Lx)$ is of the form (P-I), with $A = \partial g$ and $B = \partial h$ representing the subdifferentials of proper lsc convex functions g and h .

One of the central algorithms for solving (P-I) is the Chambolle–Pock algorithm (CPA) [12] (also known as the primal-dual hybrid gradient (PDHG) method [54, 18, 24]). Given strictly positive stepsizes $\gamma, \tau > 0$, a sequence of strictly positive relaxation parameters $(\lambda_k)_{k \in \mathbb{N}}$ and an initial guess $(x^0, y^0) \in \mathbb{R}^{n+m}$, this algorithm consists of the following iterates.

$$\begin{cases} \bar{x}^k \in J_{\gamma A}(x^k - \gamma L^{\top}y^k) \\ \bar{y}^k \in J_{\tau B^{-1}}(y^k + \tau L(2\bar{x}^k - x^k)) \\ x^{k+1} = x^k + \lambda_k(\bar{x}^k - x^k) \\ y^{k+1} = y^k + \lambda_k(\bar{y}^k - y^k) \end{cases} \quad (\text{CPA})$$

The convergence analysis of CPA in literature largely relies upon an underlying monotonicity assumption. In this work, we identify classes of nonmonotone problems along with corresponding stepsize and relaxation

Department of Electrical Engineering (ESAT-STADIUS) – KU Leuven, Kasteelpark Arenberg 10, 3001 Leuven, Belgium.
E-mail: {brecht.evens, puya.latafat, panos.patrinos}@kuleuven.be.

This work was supported by: the Research Foundation Flanders (FWO) PhD grant No. 1183822N; postdoctoral grant No. 12Y7622N and research projects G081222N, G033822N, G0A0920N; European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 953348.

parameter conditions for which CPA remains convergent. To this end, we rely on casting CPA as an instance of the preconditioned proximal point algorithm (PPPA). This connection was previously exploited in [18, 24, 14, 29, 11] in the monotone setting. Many other widely used numerical methods can also be interpreted as special cases of PPPA, see e.g. [16, 43, 46, 17, 14]. In particular, consider the inclusion problem of finding a zero of a set-valued operator $T : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$, i.e.,

$$\text{find } z \in \mathbb{R}^n \text{ such that } 0 \in Tz. \quad (\text{G-I})$$

Then, given a symmetric positive *semidefinite* preconditioning matrix $P \in \mathbb{R}^{n \times n}$ and a sequence of strictly positive relaxation parameters $(\lambda_k)_{k \in \mathbb{N}}$, the (relaxed) preconditioned proximal point algorithm applied to (G-I) consists of the fixed point iterations

$$\begin{cases} \bar{z}^k \in (P + T)^{-1} P z^k \\ z^{k+1} = z^k + \lambda_k (\bar{z}^k - z^k). \end{cases} \quad (\text{PPPA})$$

The Chambolle–Pock algorithm can be retrieved for certain choice of the preconditioner P and the operator T . Specifically, consider the so-called *primal-dual inclusion*

$$\text{find } z = (x, y) \in \mathbb{R}^{n+m} \text{ such that } 0 \in T_{\text{PD}}(z) := \begin{bmatrix} Ax \\ B^{-1}y \end{bmatrix} + \begin{bmatrix} L^\top y \\ -Lx \end{bmatrix}. \quad (\text{PD-I})$$

Then, letting $z^k = (x^k, y^k)$ and $\bar{z}^k = (\bar{x}^k, \bar{y}^k)$, CPA is equivalent to applying PPPA to (PD-I), with preconditioner

$$P = \begin{bmatrix} \frac{1}{\gamma} I_n & -L^\top \\ -L & \frac{1}{\tau} I_m \end{bmatrix}. \quad (1.1)$$

As a result of this equivalence, the convergence properties of CPA can be inferred from those of PPPA. In the monotone setting, convergence of PPPA is well understood, not only for positive definite preconditioners [32, 43, 44, 45] but also for positive semidefinite ones [29, Thm. 3.4], [11, §2.1]. Analogously, the convergence of CPA for monotone inclusions is relatively well-understood, provided that the stepsizes γ and τ satisfy a certain stepsize condition. The standard assumption in the first works on CPA such as [12, 18, 24] was that the stepsizes γ and τ satisfy $\gamma\tau\|L\|^2 < 1$. This assumption was later relaxed to $\gamma\tau\|L\|^2 \leq 1$ in [14, 29, 37], broadening the scope of the analysis to Douglas-Rachford splitting (DRS), for which $\tau = 1/\gamma$ and $L = I$. Interestingly, when interpreting CPA as a particular instance of PPPA, the stepsize condition discussed in these works is directly linked to the positive definiteness of the preconditioning matrix P in PPPA. This connection becomes evident by observing that, owing to the Schur complement lemma, P is positive definite under the traditional stepsize condition $\gamma\tau\|L\|^2 < 1$ and positive semidefinite under the relaxed stepsize condition $\gamma\tau\|L\|^2 \leq 1$.

Recently, convergence of PPPA in the nonmonotone setting has been considered in [19] under the assumption that T admits a set of oblique weak Minty solutions, defined as follows.

Definition 1.1 (*V-oblique weak Minty solutions* [19]). *An operator $T : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is said to have V-oblique weak Minty solutions at (a nonempty set) $S^* \subseteq \text{zer } T$ for some symmetric matrix $V \in \mathbb{R}^{n \times n}$ if*

$$\langle v, z - z^* \rangle \geq q_V(v), \quad \text{for all } z^* \in S^*, (z, v) \in \text{gph } T, \quad (1.2)$$

where the quadratic form $q_V(v) := \langle v, Vv \rangle$. Whenever $V = \rho I$ for some $\rho \in \mathbb{R}$, we refer to them as ρ -weak Minty solutions.

One key aspect of this assumption is its generality, as V is allowed to be any (possibly indefinite) symmetric matrix. For instance, if V is equal to the zero matrix, (1.2) reduces to the classic Minty variational inequality (MVI) [34, 20], while if $V = \rho I$ the so-called weak MVI is retrieved. In literature, weak MVI and the closely related notion of cohypomonotonicity have been employed in the context of the extragradient and the forward-backward-forward method [15, 42, 41, 9, 22], as well as the classic PPA method [39, 26, 13, 22].

Leveraging the results from [19] and the primal-dual connection between CPA and PPPA, the first part of this work will focus on establishing convergence of CPA under the assumption that the primal-dual operator T_{PD} admits a set of V -oblique weak Minty solutions. To account for the inherent structure present within T_{PD} , we impose a specific block diagonal form for $V = \text{blkdiag}(V_P, V_D)$, which depends on the fundamental subspaces of L (see (3.3) and the discussion thereafter). Furthermore, we demonstrate that by restricting our

obtained results to the case where $L = I$ and $\tau = 1/\gamma$, the convergence results for nonmonotone DRS from [19, Sec. 3] are retrieved.

In contrast to the DRS setting, where the convergence results follow in a straightforward manner from those of PPPA (see proof of [19, Thm. 3.3]), convergence results for CPA are more challenging to obtain, not only due to additional stepsize parameter τ , but mainly due to the additional complexity in the algorithm introduced by the matrix L . This difficulty is overcome by considering the singular value decomposition of L and using the corresponding orthonormal basis to carefully decompose the preconditioner P and the oblique weak Minty matrix V (see proof of Theorem 3.4).

In practice, it might be difficult to determine whether the associated primal-dual operator of a given inclusion problem admits V -oblique weak Minty solutions. This issue will be addressed in the second part of this work, where we introduce the class of (M, R) -semimonotone operators and provide several calculus rules for this class, allowing to verify the existence of V -oblique weak Minty solutions based on the semimonotonicity properties of the underlying operators A and B . The class of semimonotone operators is defined as follows.

Definition 1.2 (semimonotonicity). *Let $M, R \in \mathbb{R}^{n \times n}$ be symmetric (possibly indefinite) matrices. An operator $A : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is said to be (M, R) -semimonotone at $(x', y') \in \text{gph } A$ if*

$$\langle x - x', y - y' \rangle \geq q_M(x - x') + q_R(y - y'), \quad \text{for all } (x, y) \in \text{gph } A, \quad (1.3)$$

where $q_X(\cdot) := \langle \cdot, \cdot \rangle_X$ for any symmetric matrix $X \in \mathbb{R}^{n \times n}$. An operator A is said to be (M, R) -semimonotone if it is (M, R) -semimonotone at all $(x', y') \in \text{gph } A$. It is said to be maximally (M, R) -semimonotone if its graph is not strictly contained in the graph of another (M, R) -semimonotone operator.

Throughout, whenever $M = \mu I_n$ and $R = \rho I_n$ where $\mu, \rho \in \mathbb{R}$, the prefix (M, R) is replaced by (μ, ρ) and condition (1.3) reduces to

$$\langle x - x', y - y' \rangle \geq \mu \|x - x'\|^2 + \rho \|y - y'\|^2, \quad \text{for all } (x, y) \in \text{gph } A. \quad (1.4)$$

The class of (μ, ρ) -semimonotone operators was introduced in [19, Sec. 4] and enjoys a lot of additional freedom compared to more traditional operators classes. For instance, it encompasses the classes of (hypo)monotone, co(hypomonotone), ρ -semimonotone [38, Def. 2], averaged and firmly nonexpansive operators (see [19, Rem. 4.2 & Fig. 4]).

In this work, this notion is generalized by characterizing the operator class with matrices (M, R) instead of scalars (μ, ρ) . This generalization is crucial to capture and exploit the specific structure emerging in CPA. To illustrate this, the next theorem provides a simplified version of our main result (see Corollary 5.4 for the full statement). For instance, if μ_A is positive, $(\mu_A L^\top L, \rho_A I_n)$ -semimonotonicity of A in Theorem 1.3 could be replaced by $(\mu_A \|L\|^2, \rho_A)$ -semimonotonicity, which is in general a much more restrictive assumption.

Theorem 1.3 (convergence of CPA under semimonotonicity (simplified)). *Let operators $A : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ and $B : \mathbb{R}^m \rightrightarrows \mathbb{R}^m$ be outer semicontinuous. Suppose that there exists $(x^*, y^*) \in \text{zer } T_{\text{PD}}$ such that A is $(\mu_A L^\top L, \rho_A I_n)$ -semimonotone at $(x^*, -L^\top y^*) \in \text{gph } A$, B is $(\mu_B I_m, \rho_B L L^\top)$ -semimonotone at $(L x^*, y^*) \in \text{gph } B$ and the semimonotonicity moduli $(\mu_A, \mu_B, \rho_A, \rho_B) \in \mathbb{R}^4$ satisfy one of the following conditions.*

- (i) (either) $\mu_A = \mu_B = 0$ and $\rho_A = \rho_B = 0$.
- (ii) (or) $\mu_A + \mu_B > 0$ and $\rho_A = \rho_B = 0$.
- (iii) (or) $\rho_A + \rho_B > 0$ and $\mu_A = \mu_B = 0$.
- (iv) (or) $\mu_A + \mu_B > 0$, $\rho_A + \rho_B > 0$ and $\min\{0, \frac{\mu_A \mu_B}{\mu_A + \mu_B}\} \min\{0, \frac{\rho_A \rho_B}{\rho_A + \rho_B}\} < \frac{1}{4\|L\|^2}$.

Then, there exist positive stepsizes γ, τ and relaxation sequences $(\lambda_k)_{k \in \mathbb{N}}$ such that if the resolvents $J_{\gamma A}, J_{\tau B^{-1}}$ have full domain¹, any sequence $(\bar{z}^k)_{k \in \mathbb{N}} = (\bar{x}^k, \bar{y}^k)_{k \in \mathbb{N}}$ generated by CPA starting from $z^0 \in \mathbb{R}^n$ either reaches a point $\bar{z}^k \in \text{zer } T_{\text{PD}}$ in a finite number of iterations or every limit point of $(\bar{z}^k)_{k \in \mathbb{N}}$ belongs to $\text{zer } T_{\text{PD}}$.

This convergence result possesses two primary attributes that deserve attention. First of all, it only requires semimonotonicity of the involved operators at a single point, as opposed to the traditional global assumptions of (hypo)monotonicity and co(hypo)monotonicity. Secondly, by considering the more general class of semimonotone operators, we obtain fundamentally new convergence results (see case 1.3(iv)), not covered by any

¹The full domain assumption is imposed to ensure that the iterates of CPA are well-defined.

existing theory for CPA. Most notably, this includes examples where $\mu_A\mu_B < 0$ and $\rho_A\rho_B < 0$, for which neither the primal nor the dual nor the primal-dual inclusion are monotone (see e.g. Example 5.7).

As first observed in [48], CPA can be viewed as a particular instance of proximal ADMM. Exploiting this connection it is possible to obtain convergence results for CPA based on those for nonconvex proximal ADMM, see [30, 8]. This approach leads to requirements for L such as full row rank assumption, and restrictions on its condition number (see [8, Ass. 1, Rem. 2(c)]). Our convergence results for CPA also do not depend on any explicit rank conditions on L , allowing to cover rank-deficient cases without introducing a switching mechanism.

1.1 Contributions

The main contribution of the paper is to establish convergence of CPA under the assumption that the primal-dual operator T_{PD} admits a set of V -oblique weak Minty solutions, which leads to novel stepsize and relaxation parameter ranges in both strongly monotone and nonmonotone settings (see Theorem 3.4 and the preceding discussion). Interestingly, in contrast to the classical stepsize condition $\gamma\tau\|L\|^2 \leq 1$ in the monotone setting, the conditions obtained through our analysis not only depend on the norm of L but also on its other singular values. The tightness of our main convergence theorem is demonstrated through Examples 3.6 and 3.7.

As our second main contribution, convergence results are provided for the class of semimonotone operators [19, Sec. 4], which can be viewed as a natural extension of the (hypo)- and co(hypo)monotone operators. We show that the stepsize requirements reduce to a look-up table depending on the level of (hypo)- and co(hypo)monotonicity (see Corollary 5.4). These results are made possible by establishing a link between the oblique weak Minty assumption for the primal-dual operator and semimonotonicity of the underlying operators A and B , relying on the extended calculus rules developed in Section 4 (see also Theorem 5.1).

1.2 Organization

The paper is structured in the following manner. In Section 1.3, some notation and standard definitions are provided. Section 2 recalls the main convergence results from [19] for PPPA in the nonmonotone setting. In Section 3, the primal-dual equivalence between CPA and PPPA is established, which lead to convergence of CPA under an oblique weak Minty assumption on the associated primal-dual operator. Section 4 discusses and introduces various calculus rules for the class of (M, R) -semimonotone operators. Leveraging these calculus rules, Section 5 presents a set of sufficient conditions for the convergence of CPA, based on the semimonotonicity of the underlying operators, along with several examples. Finally, Section 6 concludes the paper. Several proofs and auxiliary results are deferred to the Appendix.

1.3 Notation

The set of natural numbers including zero is denoted by $\mathbb{N} := \{0, 1, \dots\}$. The set of real and extended-real numbers are denoted by $\mathbb{R} := (-\infty, \infty)$ and $\overline{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$, while the positive and strictly positive reals are $\mathbb{R}_+ := [0, \infty)$ and $\mathbb{R}_{++} := (0, \infty)$. We use the notation $(w^k)_{k \in I}$ to denote a sequence with indices in the set $I \subseteq \mathbb{N}$. When dealing with scalar sequences we use the subscript notation $(\gamma_k)_{k \in I}$. We denote the positive part of a real number by $[\cdot]_+ := \max\{0, \cdot\}$ and the negative part by $[\cdot]_- := \min\{0, \cdot\}$. With id we indicate the identity function $x \mapsto x$ defined on a suitable space. The identity matrix is denoted by $I_n \in \mathbb{R}^{n \times n}$ and the zero matrix by $0_{m \times n} \in \mathbb{R}^{m \times n}$; we write respectively I and 0 when no ambiguity occurs. Adopting the notation from [7], we say a matrix $P \in \mathbb{R}^{m \times n}$ is empty if $\min(m, n) = 0$ and use the conventions $P0_{n \times 0} = 0_{m \times 0}$, $0_{0 \times m}P = 0_{0 \times n}$ and $0_{m \times 0}0_{0 \times n} = 0_{m \times n}$. Moreover, we use the convention that $\frac{1}{0} = \infty$. Given a matrix $P \in \mathbb{R}^{m \times n}$, we denote the range of P by $\mathcal{R}(P)$ and the kernel of P by $\mathcal{N}(P)$. The trace of a square matrix $P \in \mathbb{R}^{n \times n}$ is denoted by $\text{tr } P$.

We denote by \mathbb{R}^n the standard n -dimensional Euclidean space with inner product $\langle \cdot, \cdot \rangle$ and induced norm $\|\cdot\|$. The set of symmetric n -by- n matrices is denoted by \mathbb{S}^n . Given a symmetric matrix $P \in \mathbb{S}^n$, we write $P \geq 0$ and $P > 0$ to denote that P is positive semidefinite and positive definite, respectively. Furthermore, for any $P \in \mathbb{S}^n$ we define the quadratic form $q_P(x) := \langle x, Px \rangle$. Let $\text{diag}(\cdot)$ denote the diagonal matrix whose arguments constitute its diagonal elements. For arbitrary matrices A and B , we define the direct sum $A \oplus B = \text{blkdiag}(A, B)$, where $\text{blkdiag}(\cdot)$ denotes the block diagonal matrix whose arguments constitute its diagonal blocks. We denote the Kronecker product between two matrices of arbitrary size by \otimes .

Two vectors $u, v \in \mathbb{R}^n$ are said to be orthogonal if $\langle u, v \rangle = 0$, and orthonormal if they are orthogonal and $\|u\| = \|v\| = 1$. Two linear subspaces $\mathbf{U} \subseteq \mathbb{R}^n$ and $\mathbf{V} \subseteq \mathbb{R}^n$ are said to be orthogonal if any $u \in \mathbf{U}$ and any $v \in \mathbf{V}$ are orthogonal. We say that $U \in \mathbb{R}^{n \times m}$ is an orthonormal basis for a linear subspace $\mathbf{U} \subseteq \mathbb{R}^n$ if U has orthonormal columns and $\mathcal{R}(U) = \mathbf{U}$.

The effective domain of an extended-real-valued function $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ is given by the set $\text{dom } f := \{x \in \mathbb{R}^n \mid f(x) < \infty\}$. We say that f is proper if $\text{dom } f \neq \emptyset$ and that f is lower semicontinuous (lsc) if the epigraph $\text{epi } f := \{(x, \alpha) \in \mathbb{R}^n \times \mathbb{R} \mid f(x) \leq \alpha\}$ is a closed subset of \mathbb{R}^{n+1} . We denote the limiting subdifferential of f by ∂f . We denote the normal cone of a set $E \subseteq \mathbb{R}^n$ by N_E and the projection onto E is denoted by $\Pi_E(x) := \arg \min_{z \in E} \|z - x\|$. An operator or set-valued mapping $A : \mathbb{R}^n \rightrightarrows \mathbb{R}^d$ maps each point $x \in \mathbb{R}^n$ to a subset $A(x)$ of \mathbb{R}^d . We will use the notation $A(x)$ and Ax interchangeably. We denote the domain of A by $\text{dom } A := \{x \in \mathbb{R}^n \mid Ax \neq \emptyset\}$, its graph by $\text{gph } A := \{(x, y) \in \mathbb{R}^n \times \mathbb{R}^d \mid y \in Ax\}$, and the set of its zeros by $\text{zer } A := \{x \in \mathbb{R}^n \mid 0 \in Ax\}$. The inverse of A is defined through its graph: $\text{gph } A^{-1} := \{(y, x) \mid (x, y) \in \text{gph } A\}$. The *resolvent* of A is defined by $J_A := (\text{id} + A)^{-1}$. We say that A is *outer semicontinuous (osc)* at $x' \in \text{dom } A$ if

$$\limsup_{x \rightarrow x'} Ax := \{y \mid \exists x^k \rightarrow x', \exists y^k \rightarrow y \text{ with } y^k \in Ax^k\} \subseteq Ax'.$$

Outer semicontinuity of A everywhere is equivalent to its graph being a closed subset of $\mathbb{R}^n \times \mathbb{R}^d$.

Definition 1.4 ((co)monotonicity). *An operator $A : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is said to be μ -monotone for some $\mu \in \mathbb{R}$ if*

$$\langle x - x', y - y' \rangle \geq \mu \|x - x'\|^2, \quad \text{for all } (x, y), (x', y') \in \text{gph } A,$$

and it is said to be ρ -comonotone for some $\rho \in \mathbb{R}$ if

$$\langle x - x', y - y' \rangle \geq \rho \|y - y'\|^2, \quad \text{for all } (x, y), (x', y') \in \text{gph } A.$$

A is said to be maximally (co-)monotone if its graph is not strictly contained in the graph of another (co-)monotone operator. We say that A is monotone if it is 0-monotone.

2 Preliminaries on the preconditioned proximal point method

Departing from the classical monotone setting, convergence of relaxed PPPA was established in [19] for a class of nonmonotone operators that admit a set of *oblique weak Minty solutions* (see Definition 1.1). This result will serve as our primary tool for establishing convergence of CPA in the nonmonotone setting, which is why we will reiterate it here. In particular, its analysis involves the following assumptions.

Assumption I. *The operator T in (G-I) and the symmetric positive semidefinite preconditioner P in (PPPA) satisfy the following properties.*

A1 $T : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is outer semicontinuous.

A2 The preconditioned resolvent $(P + T)^{-1} \circ P$ has full domain.

A3 There exists a nonempty set $\mathcal{S}^* \subseteq \text{zer } T$ and a symmetric, possibly indefinite matrix $V \in \mathbb{S}^n$ such that T has V -oblique weak Minty solutions at \mathcal{S}^* for V .

A4 $P \in \mathbb{S}^n$ is a symmetric positive semidefinite matrix such that

$$1 + \bar{\eta} > 0 \quad \text{with} \quad \bar{\eta} := \lambda_{\min}(U^T V P U),^2 \quad (2.1)$$

where U is any orthonormal basis for the range of P .

In contrast to the convergence analysis techniques relying on firm nonexpansiveness of the resolvent mapping, the analysis of [19] relies on a projective interpretation of the preconditioned proximal point algorithm, which dates back to [50, 49, 28]. Most notably, it was demonstrated in [19, Lem. 2.2] that the update rule for the (shadow) sequence generated by PPPA can be interpreted as a relaxed projection onto a certain halfspace, and that if any iterate belongs to this halfspace, which contains the set of projected oblique weak Minty solutions $\Pi_{\mathcal{R}(P)} \mathcal{S}^*$, this implies its optimality. Based on this insight, the following convergence result for PPPA was established.

²As $U^T V P U$ is similar to a symmetric matrix, its eigenvalues are real [19, Rem. 2.2].

Theorem 2.1 ([19, Thm. 2.3] convergence of PPPA). *Suppose that Assumption I holds, and consider a sequence $(z^k, \bar{z}^k)_{k \in \mathbb{N}}$ generated by PPPA starting from $z^0 \in \mathbb{R}^n$ with relaxation parameters $\lambda_k \in (0, 2(1 + \bar{\eta}))$ such that $\liminf_{k \rightarrow \infty} \lambda_k(2(1 + \bar{\eta}) - \lambda_k) > 0$, where $\bar{\eta}$ is defined as in (2.1). Then, either a point $\bar{z}^k \in \text{zer } T$ is reached in a finite number of iterations or the following hold for the sequence $(z^k, \bar{z}^k)_{k \in \mathbb{N}}$.*

- (i) $\bar{v}^k := P(z^k - \bar{z}^k) \in T\bar{z}^k$ for all k and $(\bar{v}^k)_{k \in \mathbb{N}}$ converges to zero.
- (ii) Every limit point (if any) of $(\bar{z}^k)_{k \in \mathbb{N}}$ belongs to $\text{zer } T$.
- (iii) The sequences $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$, $(\Pi_{\mathcal{R}(P)} \bar{z}^k)_{k \in \mathbb{N}}$ are bounded and their limit points belong to $\Pi_{\mathcal{R}(P)} \text{zer } T$.
- (iv) If in Assumption I.A3 the set $\Pi_{\mathcal{R}(P)} \mathcal{S}^*$ is equal to $\Pi_{\mathcal{R}(P)} \text{zer } T$, then $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ converges to some element of $\Pi_{\mathcal{R}(P)} \text{zer } T$. If additionally $(P + T)^{-1} \circ P$ is (single-valued) continuous, then $(\bar{z}^k)_{k \in \mathbb{N}}$ converges to some $z^* \in \text{zer } T$. Finally, if λ_k is additionally uniformly bounded in the interval $(0, 2)$, then $(z^k)_{k \in \mathbb{N}}$ also converges to $z^* \in \text{zer } T$.

As mentioned in [19], Assumption I.A3 can be further relaxed by only requiring (1.2) to hold on $(z, v) \in \text{gph } T \cap (\mathcal{R}((P + T)^{-1}P) \times \mathcal{R}(P))$ instead. Under this relaxed assumption, all results from Theorem 2.1 remain valid, as the proof of Theorem 2.1 only involves invoking (1.2) at points in this restricted set. This relaxation will prove to be relevant in Example 3.6.

3 Chambolle–Pock under oblique weak Minty

In the monotone setting, it is well-known that CPA can be interpreted as applying PPPA to the primal-dual operator T_{PD} [18, 24, 14, 29, 11]. Relying upon the abstract duality framework from [1], [2, Sec. 6.9], this equivalence can be extended to the nonmonotone setting. Within this framework, inclusion problems (P-I) and (PD-I) are labelled as the *primal* and the *primal-dual inclusion*, respectively. Related to these two inclusions is the *dual inclusion*, given by

$$\text{find } y \in \mathbb{R}^n \quad \text{such that} \quad 0 \in T_{\text{D}}y := (-L)A^{-1}(-L^\top)(y) + B^{-1}(y), \quad (\text{D-I})$$

A fundamental equivalence property for these inclusions is summarized below.

Proposition 3.1 ([2, Prop. 6.9.2]). *Let $(x, y) \in \mathbb{R}^n \times \mathbb{R}^m$. The following statements are equivalent.*

- (i) $(x, y) \in \text{zer } T_{\text{PD}}$.
- (ii) $x \in \text{zer } T_{\text{P}}$ and $y \in \text{zer } T_{\text{D}}$.
- (iii) $(x, -L^\top y) \in \text{gph } A$ and $(Lx, y) \in \text{gph } B$.

Furthermore, it holds that $\text{zer } T_{\text{P}} = \{x \mid \exists y : (x, y) \in \text{zer } T_{\text{PD}}\}$ and $\text{zer } T_{\text{D}} = \{y \mid \exists x : (x, y) \in \text{zer } T_{\text{PD}}\}$.

A solution of the primal inclusion (P-I) (and of the dual inclusion (D-I)) can thus be obtained by finding a solution of the associated primal-dual inclusion. Now, consider applying PPPA to the primal-dual inclusion (PD-I), with the preconditioner P given by (1.1). Then, each iteration corresponds to first finding \bar{x}^k and \bar{y}^k satisfying the inclusions

$$\frac{1}{\gamma}x^k - L^\top y^k \in (\frac{1}{\gamma}\text{id} + A)\bar{x}^k \quad \text{and} \quad \frac{1}{\tau}y^k - Lx^k \in -2L\bar{x}^k + (\frac{1}{\tau}\text{id} + B^{-1})\bar{y}^k$$

and then performing a relaxation step $z^{k+1} = z^k + \lambda_k(\bar{z}^k - z^k)$. Multiplying the two relations by γ and τ , respectively, and reordering the terms, the update rule for CPA is retrieved. This result is summarized in the following lemma.

Lemma 3.2 (equivalence of CPA and PPPA). *Let $z^0 = (x^0, y^0) \in \mathbb{R}^{n+m}$ be the initial guess for CPA and for PPPA applied to the primal-dual inclusion (PD-I), with the preconditioner P given by (1.1). Then, the sequences $(z^k)_{k \in \mathbb{N}} = (x^k, y^k)_{k \in \mathbb{N}}$, $(\bar{z}^k)_{k \in \mathbb{N}} = (\bar{x}^k, \bar{y}^k)_{k \in \mathbb{N}}$ generated by CPA satisfy update rule PPPA.*

Leveraging this connection, we will establish the convergence of CPA based on [Theorem 2.1](#) for PPPA. In contrast to the classical stepsize condition $\gamma\tau\|L\|^2 \leq 1$ in the monotone setting, our upcoming analysis will demonstrate that the stepsize condition on γ and τ for CPA in general does not only depend on $\|L\|$, i.e., the largest singular value of L , but also its other singular values. Therefore, let r denote the rank of L , and without loss of generality, let $\sigma_1, \dots, \sigma_d$ denote its distinct strictly positive singular values in descending order with respective multiplicities m_1, \dots, m_d . Then, it holds that $r = \sum_{i=1}^d m_i$. Define $\Sigma = \sigma_1 I_{m_1} \oplus \dots \oplus \sigma_d I_{m_d} \in \mathbb{R}^{r \times r}$ and consider the singular value decomposition

$$L = \begin{bmatrix} Y & Y' \end{bmatrix} \begin{bmatrix} \Sigma & \\ & 0 \end{bmatrix} \begin{bmatrix} X^\top \\ X'^\top \end{bmatrix}, \quad Y = \begin{bmatrix} Y_1 & \dots & Y_d \end{bmatrix}, \quad X = \begin{bmatrix} X_1 & \dots & X_d \end{bmatrix}, \quad (3.1)$$

where the zero matrix is in $\mathbb{R}^{(m-r) \times (n-r)}$, $Y_i \in \mathbb{R}^{m \times m_i}$ and $X_i \in \mathbb{R}^{n \times m_i}$, $i \in [d]$, have orthonormal columns that span the eigenspace corresponding to eigenvalue σ_i^2 of LL^\top and $L^\top L$, respectively, and $Y' \in \mathbb{R}^{m \times (m-r)}$ and $X' \in \mathbb{R}^{n \times (n-r)}$ have orthonormal columns which span the null space of L^\top and L , respectively. The projection onto the range and the kernel of L and L^\top can be expressed as [[21](#), Sec. 2.5.2]

$$\Pi_{\mathcal{R}(L)} = YY^\top, \quad \Pi_{\mathcal{R}(L^\top)} = XX^\top, \quad \Pi_{\mathcal{N}(L)} = I_n - XX^\top = X'X'^\top \quad \text{and} \quad \Pi_{\mathcal{N}(L^\top)} = I_m - YY^\top = Y'Y'^\top. \quad (3.2)$$

These projections will play a central role in our upcoming analysis. We will work under the following assumptions on the individual operators A and B and the (nonzero) matrix L .

Assumption II. *In problem (P-I), the following hold.*

A1 *Operators A and B are outer semicontinuous.*

A2 *For the selected positive stepsizes the corresponding resolvents have full domain, i.e., $\text{dom } J_{\gamma A} = \mathbb{R}^n$ and $\text{dom } J_{\tau B^{-1}} = \mathbb{R}^m$.*

A3 *The set $\text{zer } T_{\text{PD}}$ is nonempty and there exist parameters $\beta_P, \beta'_P, \beta_D, \beta'_D \in \mathbb{R}$ and a nonempty set $\mathcal{S}^* \subseteq \text{zer } T_{\text{PD}}$ such that the primal-dual operator T_{PD} has V -oblique weak Minty solutions at \mathcal{S}^* , where*

$$V := V_P \oplus V_D = (\beta_P \Pi_{\mathcal{R}(L^\top)} + \beta'_P \Pi_{\mathcal{N}(L)}) \oplus (\beta_D \Pi_{\mathcal{R}(L)} + \beta'_D \Pi_{\mathcal{N}(L^\top)}) \in \mathbb{S}^{n+m}. \quad (3.3)$$

Moreover, defining

$$\delta := 1 + [\beta_P \beta_D]_-(\|L\|^2 - \sigma_d^2) = \begin{cases} 1, & \text{if } \beta_P \beta_D \geq 0, \\ 1 + \beta_P \beta_D (\|L\|^2 - \sigma_d^2), & \text{if } \beta_P \beta_D < 0, \end{cases} \quad (3.4)$$

the following conditions on $\beta_P, \beta'_P, \beta_D$ and β'_D hold.

$$[\beta_P]_- [\beta_D]_- < \frac{1}{4\|L\|^2}, \quad \beta'_P \geq \begin{cases} 0, & \text{if } \beta_P \geq 0, \\ \frac{\beta_P}{\delta - \beta_P \beta_D \|L\|^2}, & \text{if } \beta_P < 0, \end{cases} \quad \text{and} \quad \beta'_D \geq \begin{cases} 0, & \text{if } \beta_D \geq 0, \\ \frac{\beta_D}{\delta - \beta_P \beta_D \|L\|^2}, & \text{if } \beta_D < 0. \end{cases} \quad (3.5)$$

In [Assumption II.A3](#), the matrix V is defined as in (3.3) with the understanding that in our convergence analysis we shall set $\beta'_P = 0$ when L is full column rank and $\beta'_D = 0$ when L is full row rank, as in these cases V is not affected by their choice. Note also that the conditions on β'_P and β'_D specified in (3.5) can be slightly relaxed (see [Theorem A.1](#)). This relaxation, however, results in more complicated stepsize rules for γ and τ , and for this reason is not pursued further.

The imposed block structure on V in (3.3) is not simply an arbitrary choice, but it aligns perfectly with the inherent structure present within the primal-dual operator itself. To illustrate this, consider the following lemma, which shows that the blocks V_P and V_D from (3.3) can be interpreted as the primal and the dual blocks of V , respectively. This lemma extends [[19](#), Lem. 3.2], which considers the case $L = I$. The proof is deferred to [Appendix C](#).

Lemma 3.3 (oblique weak Minty for primal and dual operator). *Let $V_P \in \mathbb{S}^n$ and $V_D \in \mathbb{S}^m$. Suppose that there exists a nonempty set $\mathcal{S}^* \subseteq \text{zer } T_{\text{PD}}$ such that the primal-dual operator T_{PD} has $(V_P \oplus V_D)$ -oblique weak Minty solutions at \mathcal{S}^* and let*

$$\mathcal{S}_P^* := \{x^* \mid \exists y^* : (x^*, y^*) \in \mathcal{S}^*\} \subseteq \text{zer } T_P \quad \text{and} \quad \mathcal{S}_D^* := \{y^* \mid \exists x^* : (x^*, y^*) \in \mathcal{S}^*\} \subseteq \text{zer } T_D.$$

Then, the primal operator T_P has V_P -oblique weak Minty solutions at \mathcal{S}_P^ and the dual operator T_D has V_D -oblique weak Minty solutions at \mathcal{S}_D^* .*

Table 1: Range of the stepsizes γ and τ for CPA.

| | $\beta_D \geq 0$ | $\beta_D < 0$ |
|------------------|---|---|
| $\beta_P \geq 0$ | $\gamma \in (0, +\infty)$ $\tau \in \left(0, \frac{1}{\gamma\ L\ ^2}\right]$ | $\gamma \in (0, \gamma_{\max})$ $\tau \in \left(\tau_{\min}(\gamma), \frac{1}{\gamma\ L\ ^2}\right]$ |
| $\beta_P < 0$ | $\gamma \in (\gamma_{\min}, +\infty)$ $\tau \in \left(\tau_{\min}(\gamma), \frac{1}{\gamma\ L\ ^2}\right]$ | $\gamma \in (\gamma_{\min}, \gamma_{\max})$ $\tau \in \left(\tau_{\min}(\gamma), \frac{1}{\gamma\ L\ ^2}\right]$ |

Table 2: Definition of η' in Relaxation parameter rule I.

| | rank $L = n$ | rank $L < n$ |
|--------------|--------------------------|---|
| rank $L = m$ | $+\infty$ | $\frac{1}{\gamma}\beta'_P$ |
| rank $L < m$ | $\frac{1}{\tau}\beta'_D$ | $\min\left\{\frac{1}{\gamma}\beta'_P, \frac{1}{\tau}\beta'_D\right\}$ |

As shown in the proof of [Lemma 3.3](#), the quadratic terms $q_{V_P}(y_A + L^\top y_B)$ and $q_{V_D}(x_B - Lx_A)$ emerging in oblique weak Minty inequality correspond to the primal and the dual problems, respectively. By selecting V_P and V_D as in (3.3), these terms can be written as

$$\begin{aligned} q_{V_P}(y_A + L^\top y_B) &= \beta_P \|\Pi_{\mathcal{R}(L^\top)} y_A + L^\top y_B\|^2 + \beta'_P \|\Pi_{\mathcal{N}(L)} y_A\|^2, \\ q_{V_D}(x_B - Lx_A) &= \beta_D \|\Pi_{\mathcal{R}(L)} x_B - Lx_A\|^2 + \beta'_D \|\Pi_{\mathcal{N}(L^\top)} x_B\|^2, \end{aligned}$$

reducing to the norm of the scaled sum of a vector belonging to the range of L^\top and another to its nullspace (resp., range of L and nullspace of L^\top). This decomposition proves essential in the proof of [Theorem 3.4](#), as it enables to split condition (2.1) into two terms, one depending only on β_P and β_D and the other only depending on β'_P and β'_D (see (3.12)).

One of the main aspects of the upcoming convergence proof for CPA is showing that [Assumption I](#) holds for the operator T_{PD} and preconditioner P from (1.1). To this end, the stepsizes γ and τ and the relaxation parameters λ_k need to be selected as follows.

Stepsize rule I. The stepsizes γ and τ satisfy the bounds provided in [Table 1](#), where δ is defined as in (3.4) and

$$\gamma_{\min} := \frac{-2\beta_P}{\delta + \sqrt{\delta^2 - 4\beta_P\beta_D\|L\|^2}}, \quad \gamma_{\max} := \frac{\delta + \sqrt{\delta^2 - 4\beta_P\beta_D\|L\|^2}}{-2\beta_D\|L\|^2} \quad \text{and} \quad \tau_{\min}(\gamma) := \frac{[-\beta_D(\gamma + \beta_P)]_+}{\gamma(\delta - \beta_P\beta_D\|L\|^2) + \beta_P}. \quad (3.6)$$

Relaxation parameter rule I. Define

$$\theta_{\gamma\tau}(\sigma) := \sqrt{\left(\frac{1}{2\gamma}\beta_P - \frac{1}{2\tau}\beta_D\right)^2 + \beta_P\beta_D\sigma^2}. \quad (3.7)$$

Let $\bar{\eta} := \min\{\eta, \eta'\}$, where η' is defined as in [Table 2](#) and

$$\eta := \begin{cases} \left. \begin{aligned} &\left\{ \begin{aligned} &\frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D - \theta_{\gamma\tau}(\sigma_d), & \text{if } \beta_P\beta_D < 0 \\ &\frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D - \theta_{\gamma\tau}(\|L\|), & \text{if } \beta_P\beta_D \geq 0 \end{aligned} \right\} & \text{if } \gamma\tau < \frac{1}{\|L\|^2}, \\ &\frac{1}{\gamma}\beta_P + \frac{1}{\tau}\beta_D, & & \text{if } \gamma\tau = \frac{1}{\|L\|^2} \text{ and } d = 1, \\ &\left\{ \begin{aligned} &\frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D - \theta_{\gamma\tau}(\sigma_d), & \text{if } \beta_P\beta_D < 0 \\ &\frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D - \theta_{\gamma\tau}(\sigma_2), & \text{if } \min\{\beta_P, \beta_D\} \geq 0 \end{aligned} \right\} & \text{if } \gamma\tau = \frac{1}{\|L\|^2} \text{ and } d > 1. \\ &\frac{1}{\gamma}\beta_P + \frac{1}{\tau}\beta_D, & \text{otherwise} & \end{aligned} \right\} \end{cases} \quad (3.8)$$

The sequence $(\lambda_k)_{k \in \mathbb{N}}$ satisfies $\lambda_k \in (0, 2(1 + \bar{\eta}))$ and $\liminf_{k \rightarrow \infty} \lambda_k(2(1 + \bar{\eta}) - \lambda_k) > 0$.

When $L = I$ and $\tau = 1/\gamma$, [Stepsize rule I](#) matches the stepsize range for Douglas–Rachford splitting from [19, Thm. 3.3]. In [Relaxation parameter rule I](#) there is an interplay between the stepsizes γ and τ and the range

of admissible relaxation parameters λ_k . In the monotone setting where $\beta_p = \beta'_p = \beta_D = \beta'_D = 0$, this interplay vanishes as the relaxation rule reduces to the classical condition $\lambda_k \in (0, 2)$. In strongly monotone settings, this interplay allows us to select relaxation parameters beyond the classical upper bound of two. For instance, when $\beta_p > 0$ and $\beta'_p > 0$, then for small enough stepsizes γ the upper bound on λ_k will be larger than two (see [19, Ex. 6.2] for an example in the DRS setting where $L = I$ and $\tau = 1/\gamma$). Conversely, when $\beta_D > 0$ and $\beta'_D > 0$, this phenomenon will occur for small enough τ . Finally, when all β parameters are strictly positive then the upper bound for λ_k is larger than two for all valid stepsizes γ and τ (see e.g. [Example 3.7](#)).

Having discussed our underlying assumptions and stepsize/relaxation parameter rules, we will now present our main convergence theorem for CPA. The proof relies on carefully decomposing both the preconditioner P and the oblique weak Minty matrix V into two separate, orthogonal matrices. Exploiting the inherent structure present in these orthogonal matrices, the conditions from [Assumption I](#) are reduced to a set of eigenvalue problems of two-by-two matrices (see equation (3.13)).

Theorem 3.4. *Suppose that [Assumption II](#) holds, that γ and τ are selected according to [Stepsize rule I](#) and that the relaxation sequence $(\lambda_k)_{k \in \mathbb{N}}$ is selected according to [Relaxation parameter rule I](#). Consider the sequences $(z^k)_{k \in \mathbb{N}} = (x^k, y^k)_{k \in \mathbb{N}}$ and $(\bar{z}^k)_{k \in \mathbb{N}} = (\bar{x}^k, \bar{y}^k)_{k \in \mathbb{N}}$ generated by CPA starting from $z^0 \in \mathbb{R}^{n+m}$. Then, either a point $\bar{z}^k \in \text{zer } T_{\text{PD}}$ is reached in a finite number of iterations or the following hold for the sequences $(z^k)_{k \in \mathbb{N}}$ and $(\bar{z}^k)_{k \in \mathbb{N}}$.*

- (i) $\bar{v}^k := P(z^k - \bar{z}^k) \in T_{\text{PD}}\bar{z}^k$ for all k and $(\bar{v}^k)_{k \in \mathbb{N}}$ converges to zero.
- (ii) Every limit point (if any) of $(\bar{z}^k)_{k \in \mathbb{N}}$ belongs to $\text{zer } T_{\text{PD}}$.
- (iii) The sequences $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$, $(\Pi_{\mathcal{R}(P)} \bar{z}^k)_{k \in \mathbb{N}}$ are bounded and their limit points are in $\Pi_{\mathcal{R}(P)} \text{zer } T_{\text{PD}}$.
- (iv) If in [Assumption II.A3](#) the set $\Pi_{\mathcal{R}(P)} \mathcal{S}^*$ is equal to $\Pi_{\mathcal{R}(P)} \text{zer } T$, then $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ converges to some element of $\Pi_{\mathcal{R}(P)} \text{zer } T_{\text{PD}}$. If additionally $J_{\gamma A}$ and $J_{\tau B^{-1}}$ are (single-valued) continuous, then $(\bar{z}^k)_{k \in \mathbb{N}}$ converges to some $z^* \in \text{zer } T_{\text{PD}}$. Finally, if λ_k is additionally uniformly bounded in the interval $(0, 2)$, then $(z^k)_{k \in \mathbb{N}}$ also converges to $z^* \in \text{zer } T_{\text{PD}}$.

Proof. [Assumption I.A3](#) is immediate. Outer semicontinuity of T_{PD} follows from that of A and B [47, Theorem 5.7(a)], showing that [Assumption I.A1](#) holds. Since $J_{\gamma A}$ and $J_{\tau B^{-1}}$ have full domain the preconditioned resolvent $(P + T_{\text{PD}})^{-1}P$ has full domain owing to [47, Lemma 12.14], establishing [Assumption I.A2](#). It only remains to show that [Assumption I.A4](#) holds. Before that, we first show that $\bar{\eta}$ as defined in (2.1) is characterized by the expression given in [Relaxation parameter rule I](#).

Let $Z_i := X_i \oplus Y_i \in \mathbb{R}^{(m+n) \times (2m_i)}$ and $Z' := X' \oplus Y' \in \mathbb{R}^{(m+n) \times (m+n-2r)}$ where Y_i, X_i, Y' and X' are given by (3.1). Let

$$Z := \begin{bmatrix} Z_1 & \cdots & Z_d \end{bmatrix} \in \mathbb{R}^{(m+n) \times (2r)},$$

which by construction has orthonormal columns. The preconditioner P in (1.1) can be decomposed as

$$P = \begin{bmatrix} \frac{1}{\gamma} I_n & -L^\top \\ -L & \frac{1}{\tau} I_m \end{bmatrix} \stackrel{(3.1)}{=} \begin{bmatrix} \frac{1}{\gamma} I_n & -X \Sigma Y^\top \\ -Y \Sigma X^\top & \frac{1}{\tau} I_m \end{bmatrix} \stackrel{(3.2)}{=} \underbrace{\begin{bmatrix} \frac{1}{\gamma} X X^\top & -X \Sigma Y^\top \\ -Y \Sigma X^\top & \frac{1}{\tau} Y Y^\top \end{bmatrix}}_{=: Z \hat{P} Z^\top} + \underbrace{\begin{bmatrix} \frac{1}{\gamma} X' X'^\top \\ \frac{1}{\tau} Y' Y'^\top \end{bmatrix}}_{=: Z' P' Z'^\top}, \quad (3.9)$$

where

$$\hat{P}_i = \begin{bmatrix} \frac{1}{\gamma} & -\sigma_i \\ -\sigma_i & \frac{1}{\tau} \end{bmatrix}, \quad \hat{P} = \left((\hat{P}_1 \otimes I_{m_1}) \oplus \cdots \oplus (\hat{P}_d \otimes I_{m_d}) \right) \quad \text{and} \quad P' = \frac{1}{\gamma} I_{n-r} \oplus \frac{1}{\tau} I_{m-r}.$$

Using this decomposition, an orthonormal basis for $\mathcal{R}(P)$ can be constructed. Let $\hat{U} = (\hat{U}_1 \otimes I_{m_1}) \oplus \cdots \oplus (\hat{U}_d \otimes I_{m_d})$, where \hat{U}_i is an orthonormal basis for $\mathcal{R}(\hat{P}_i)$, for $i \in [d]$. Since $\hat{P}_i > 0$ for $i \in [2, d]$, select $\hat{U}_i = I_2$ for $i \in [2, d]$, so that $\hat{U} = (\hat{U}_1 \otimes I_{m_1}) \oplus I_{(2r-2m_1)}$. As \hat{P} conforms to the same block-diagonal structure, \hat{U} is an orthonormal basis for $\mathcal{R}(\hat{P})$. For any $A \in \mathbb{R}^{k \times l}$ and any full row rank $B \in \mathbb{R}^{l \times p}$, it holds that $\mathcal{R}(AB) = \mathcal{R}(A)$ [7, Prop. 2.6.3]. Since Z and Z' are full column rank and P' is full rank, this implies that $\mathcal{R}(Z \hat{P} Z^\top) = \mathcal{R}(Z \hat{U})$ and $\mathcal{R}(Z' P' Z'^\top) = \mathcal{R}(Z')$. Finally, since $Z \hat{U}$ has orthonormal columns (as $(Z \hat{U})^\top Z \hat{U} = I$), an orthonormal basis for $\mathcal{R}(P)$ is given by

$$U = \begin{bmatrix} Z \hat{U} & Z' \end{bmatrix} = \begin{bmatrix} Z_1 (\hat{U}_1 \otimes I_{m_1}) & Z_2 & \cdots & Z_d & Z' \end{bmatrix}. \quad (3.10)$$

Analogously, the V -oblique weak Minty matrix as defined in (3.3) can be decomposed as

$$V \stackrel{(3.2)}{=} \underbrace{(\beta_P X X^\top) \oplus (\beta_D Y Y^\top)}_{=: Z \hat{V} Z^\top} + \underbrace{(\beta'_P X' X'^\top) \oplus (\beta'_D Y' Y'^\top)}_{=: Z' V' Z'^\top}, \quad (3.11)$$

where

$$\hat{V}_i = \text{diag}(\beta_P, \beta_D), \quad \hat{V} = (\hat{V}_1 \otimes \mathbf{I}_{m_1}) \oplus \cdots \oplus (\hat{V}_d \otimes \mathbf{I}_{m_d}) \quad \text{and} \quad V' = \beta'_P \mathbf{I}_{n-r} \oplus \beta'_D \mathbf{I}_{m-r}.$$

Since Z and Z' both have orthonormal columns, i.e., $Z^\top Z = \mathbf{I}_{2r}$ and $Z'^\top Z' = \mathbf{I}_{m+n-2r}$, and since $\mathcal{R}(Z)$ and $\mathcal{R}(Z')$ are orthogonal, it follows from (3.9)-(3.11) that

$$U^\top V P U = \begin{bmatrix} \hat{U}^\top Z^\top \\ Z'^\top \end{bmatrix} (Z \hat{V} Z^\top + Z' V' Z'^\top) (Z \hat{P} Z^\top + Z' P' Z'^\top) \begin{bmatrix} Z \hat{U} & Z' \end{bmatrix} = \hat{U}^\top \hat{V} \hat{P} \hat{U} \oplus Z'^\top V' P' Z'.$$

Therefore, condition (2.1) becomes

$$1 + \bar{\eta} = 1 + \lambda_{\min}(U^\top V P U) = 1 + \min\left(\overbrace{\lambda_{\min}(\hat{U}^\top \hat{V} \hat{P} \hat{U})}^{=: \eta}, \overbrace{\lambda_{\min}(Z'^\top V' P' Z')}^{=: \eta'}\right) > 0. \quad (3.12)$$

Due to the block diagonal structure of $\hat{U}^\top \hat{V} \hat{P} \hat{U}$, it follows that

$$\begin{aligned} \eta &= \min \left\{ \lambda_{\min}(\hat{U}_i^\top \hat{V}_i \hat{P}_i \hat{U}_i) \right\}_{i=1}^d = \min \left\{ \lambda_{\min}(\hat{U}_1^\top \hat{V}_1 \hat{P}_1 \hat{U}_1), \left\{ \lambda_{\min}(\hat{V}_i \hat{P}_i) \right\}_{i=2}^d \right\} \\ &= \min \left\{ \lambda_{\min}(\hat{U}_1^\top \begin{bmatrix} \frac{1}{\gamma} \beta_P & -\beta_P \sigma_1 \\ -\beta_D \sigma_1 & \frac{1}{\tau} \beta_D \end{bmatrix} \hat{U}_1), \left\{ \frac{1}{2\gamma} \beta_P + \frac{1}{2\tau} \beta_D - \theta_{\gamma\tau}(\sigma_i) \right\}_{i=2}^d \right\}, \end{aligned} \quad (3.13)$$

where $\theta_{\gamma\tau}(\cdot)$ is defined as in (3.7). Consider the following cases.

♣ $\gamma\tau\|L\|^2 < 1$: Then, $\hat{P}_1 > 0$ so that $\hat{U}_1 = \mathbf{I}_2$ and

$$\eta = \min \left\{ \lambda_{\min}(\hat{V}_i \hat{P}_i) \right\}_{i=1}^d = \min \left\{ \frac{1}{2\gamma} \beta_P + \frac{1}{2\tau} \beta_D - \theta_{\gamma\tau}(\sigma_i) \right\}_{i=1}^d,$$

which matches the definition of η provided in (3.8).

♣ $\gamma\tau\|L\|^2 = 1$: Then, the matrix \hat{P}_1 has $\frac{1}{\gamma} + \frac{1}{\tau}$ and zero as eigenvalues, so that

$$\hat{U}_1 = \sqrt{\frac{\tau}{\gamma+\tau}} \begin{bmatrix} 1 & \\ & -\frac{1}{\sqrt{\gamma\tau}} \end{bmatrix} \quad \text{and} \quad \hat{P}_1 = \left(\frac{1}{\gamma} + \frac{1}{\tau}\right) \hat{U}_1 \hat{U}_1^\top. \quad (3.14)$$

Therefore,

$$\begin{aligned} \eta &= \min \left\{ \lambda_{\min}\left(\frac{\tau}{\gamma+\tau} \begin{bmatrix} 1 & \\ & -\sqrt{\gamma\tau} \end{bmatrix} \begin{bmatrix} \frac{1}{\gamma} \beta_P & -\beta_P \sigma_1 \\ -\beta_D \sigma_1 & \frac{1}{\tau} \beta_D \end{bmatrix} \begin{bmatrix} 1 \\ -\sqrt{\gamma\tau} \end{bmatrix}\right), \left\{ \frac{1}{2\gamma} \beta_P + \frac{1}{2\tau} \beta_D - \theta_{\gamma\tau}(\sigma_i) \right\}_{i=2}^d \right\} \\ &= \min \left\{ \frac{1}{\gamma} \beta_P + \frac{1}{\tau} \beta_D, \left\{ \frac{1}{2\gamma} \beta_P + \frac{1}{2\tau} \beta_D - \theta_{\gamma\tau}(\sigma_i) \right\}_{i=2}^d \right\}, \end{aligned}$$

where we used that $\sqrt{\gamma\tau}\sigma_1 = 1$. This matches the definition of η provided in (3.8).

By definition of Z' , V' and P' , it holds that

$$\eta' = \lambda_{\min}\left(\left(\frac{1}{\gamma} \beta'_P X'^\top X'\right) \oplus \left(\frac{1}{\tau} \beta'_D Y'^\top Y'\right)\right) = \lambda_{\min}\left(\left(\frac{1}{\gamma} \beta'_P \mathbf{I}_{n-r}\right) \oplus \left(\frac{1}{\tau} \beta'_D \mathbf{I}_{m-r}\right)\right), \quad (3.15)$$

which matches the definition of η' provided in Table 2. Having shown that the values of η and η' match the ones provided in Relaxation parameter rule I, it remains to show that under the prescribed Stepsize rule I the preconditioner is positive semidefinite (equivalently $\gamma\tau \in (0, 1/\|L\|^2]$) and $1 + \bar{\eta} := 1 + \min\{\eta, \eta'\} > 0$ as required by Assumption I.A4. We proceed with two intermediate claims.

Claim 1: The pair $(\gamma, \tau) \in \mathbb{R}_{++}^2$ satisfies $\gamma\tau \in (0, 1/\|L\|^2]$ and $1 + \eta > 0$ if and only if γ and τ comply with Stepsize rule I.

Consider the six different cases for η from (3.8). For the first and fourth case this follows from Lemma B.2(ii), for the second case from Lemmas B.2(i) and B.2(iii) since $[\beta_P]_- - [\beta_D]_- < \frac{1}{4\|L\|^2}$ by (3.5) and for the fifth case from Lemma B.2(i). Finally, for the third and the sixth case use Fact B.1, by plugging in $\tau = \frac{1}{\gamma\|L\|^2}$ into $1 + \frac{1}{\gamma} \beta_P + \frac{1}{\tau} \beta_D > 0$ and observing that $\delta = 1$.

Claim 2: If γ and τ comply with [Stepsize rule I](#) then $1 + \eta' > 0$.

Recall that as noted in the discussion after [Assumption II](#), the value of β'_P (resp. β'_D) is irrelevant when L is full column rank (resp. full row rank), and thus can be set equal to zero. In view of (3.15), γ and τ satisfy $1 + \eta' > 0$ if and only if

$$\gamma > -\beta'_P \quad \text{and} \quad \tau > -\beta'_D. \quad (3.16)$$

If $\min\{\beta_P, \beta_D\} \geq 0$, then owing to (3.5) it holds that $\beta'_P \geq 0$ and $\beta'_D \geq 0$, so that (3.16) is implied trivially by [Stepsize rule I](#). Consider the case where $\min\{\beta_P, \beta_D\} < 0$. By definition of δ from (3.4) and using $\beta_P\beta_D = [\beta_P\beta_D]_- + [\beta_P\beta_D]_+$ it holds that for any parameter $\kappa \geq 1$

$$\delta - \kappa\beta_P\beta_D\|L\|^2 = 1 - [\beta_P\beta_D]_-(\sigma_d^2 + (\kappa - 1)\|L\|^2) - \kappa[\beta_P\beta_D]_+\|L\|^2 \geq 1 - \frac{\kappa}{4}, \quad (3.17)$$

where the inequality follows from the fact that either both β_P and β_D are negative in which (3.5) is used, or only one is negative, in which case the quantity is larger than 1. Moreover,

$$\begin{aligned} 0 \leq \delta^2 - 4\beta_P\beta_D\|L\|^2 &= (\delta - 2\beta_P\beta_D\|L\|^2)^2 + 4\beta_P\beta_D\|L\|^2(\delta - 1 - \beta_P\beta_D\|L\|^2) \\ &= (\delta - 2\beta_P\beta_D\|L\|^2)^2 - 4\beta_P\beta_D\|L\|^2([\beta_P\beta_D]_+\|L\|^2 + [\beta_P\beta_D]_-\sigma_d^2) \\ &\leq (\delta - 2\beta_P\beta_D\|L\|^2)^2, \end{aligned}$$

where the last inequality follows from the trivial observation that for any $a \in \mathbb{R}$, both $a[a]_- \geq 0$ and $a[a]_+ \geq 0$. Therefore, $\delta + \sqrt{\delta^2 - 4\beta_P\beta_D\|L\|^2} \leq 2(\delta - \beta_P\beta_D\|L\|^2)$ and

$$\beta_P < 0 \implies -\beta'_P \stackrel{(3.5)}{\leq} \frac{-\beta_P}{\delta - \beta_P\beta_D\|L\|^2} \leq \gamma_{\min}. \quad (3.18)$$

Observe that regardless of the sign of β_P it holds that $\gamma(\delta - \beta_P\beta_D\|L\|^2) + \beta_P > 0$; if $\beta_P \geq 0$, then this holds owing to (3.17) since $\gamma > 0$, if $\beta_P < 0$, then this inequality remains true since in that case $\gamma > \gamma_{\min}$ by [Stepsize rule I](#) and using (3.18). We proceed by first showing that

$$\tau_{\min}(\gamma) = \frac{[-\beta_D(\gamma + \beta_P)]_+}{\gamma(\delta - \beta_P\beta_D\|L\|^2) + \beta_P} \geq \frac{[-\beta_D]_+}{\delta - \beta_P\beta_D\|L\|^2}. \quad (3.19)$$

If $\beta_D \geq 0$, then (3.19) holds trivially. If $\beta_D < 0$ and $\beta_P \geq 0$, then $\gamma + \beta_P > 0$ since $\gamma > 0$. If $\beta_D < 0$ and $\beta_P < 0$, then $\gamma + \beta_P > 0$ since in that case $\gamma > \gamma_{\min} \geq \frac{-\beta_P}{1 - \beta_P\beta_D\|L\|^2} > -\beta_P$ by [Stepsize rule I](#) and using (3.18). In either case, if $\beta_D < 0$ then $\gamma + \beta_P > 0$ and using the fact that the denominators in (3.19) are strictly positive by (3.17) it reduces to

$$-\beta_D(\gamma + \beta_P)(\delta - \beta_P\beta_D\|L\|^2) \geq -\beta_D(\gamma(\delta - \beta_P\beta_D\|L\|^2) + \beta_P).$$

Substituting δ from (3.4) and using $\beta_P\beta_D = [\beta_P\beta_D]_- + [\beta_P\beta_D]_+$, rearrange it to obtain $[\beta_P\beta_D]_-^2\sigma_d^2 + [\beta_P\beta_D]_+^2\|L\|^2 \geq 0$, which holds trivially. Combining (3.18), (3.19) and (3.5) shows that (3.16) holds and the claim is established.

Combining [Claims 1](#) and [2](#), it follows that [Stepsize rule I](#) is equivalent to the stepsizes being strictly positive, $P \geq 0$ and $1 + \bar{\eta} = 1 + \min\{\eta, \eta'\} > 0$, establishing that [Assumption I.A4](#) holds. Finally, $(P + T_{PD})^{-1} \circ P$ is (single-valued) continuous if $J_{\gamma A}$ and $J_{\tau B^{-1}}$ are (single-valued) continuous, since $\bar{z}^k \in (P + T_{PD})^{-1}Pz^k$ if and only if $\bar{x}^k \in J_{\gamma A}(x^k - \gamma L^\top y^k)$ and $\bar{y}^k \in J_{\tau B^{-1}}(y^k + \tau L(2\bar{x}^k - x^k))$. All claims for [CPA](#) follow directly from [Theorem 2.1](#), using the equivalence from [Lemma 3.2](#). \square

Note that when λ_k is uniformly bounded in the interval $(0, 2(1 + \bar{\eta}))$, a rate of $O(\frac{1}{N})$ can be obtained for $\min_{k=0,1,\dots,N} \|\bar{v}^k\|^2$ by telescoping (see [[19](#), thm. 2.3(iv)]).

Observe that [Theorem 3.4](#) discusses not only the convergence of $(z^k)_{k \in \mathbb{N}}$, but also of its projection onto the range of the preconditioner $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$. Notably, convergence of $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ is established under weaker assumptions than for $(z^k)_{k \in \mathbb{N}}$. When P is positive definite, meaning that $\gamma\tau < 1/\|L\|^2$, this is irrelevant because in this case the range of P is full. However, in the positive semidefinite case, when $\gamma\tau = 1/\|L\|^2$, these sequences are no longer identical. This observation is not surprising, as it is a natural extension of the convergence results for DRS, i.e., when $L = I$ and $\gamma = 1/\tau$. In particular, in the DRS setting it was shown that the convergence of $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ to $\Pi_{\mathcal{R}(P)} \text{zer } T_{PD}$ is equivalent to the convergence of the *shadow sequence* $(s^k)_{k \in \mathbb{N}} := (x^k - \gamma y^k)_{k \in \mathbb{N}}$ to the set $\{x^* - \gamma y^* \mid (x^*, y^*) \in \text{zer } T_{PD}\}$ [[19](#), Thm. 3.3]. In the following proposition, this shadow sequence interpretation is generalized to arbitrary L matrices.

Proposition 3.5 (convergent sequences). *Using the SVD of L from (3.1), define the function $\psi : \mathbb{R}^{m+n} \rightarrow \mathbb{R}^{m+n-m_1}$ as*

$$\psi(x, y) := \left(X_1^\top x - \sqrt{\frac{\tau}{\gamma}} Y_1^\top y, X_2^\top x, Y_2^\top y \right), \quad (3.20)$$

where $X_2 := [X_2 \ \cdots \ X_d \ X']$ and $Y_2 := [Y_2 \ \cdots \ Y_d \ Y']$ ³. Consider a sequence $(z^k)_{k \in \mathbb{N}} = (x^k, y^k)_{k \in \mathbb{N}}$ generated by CPA starting from $z^0 \in \mathbb{R}^{m+n}$, where $\gamma\tau = \frac{1}{\|L\|^2}$ and define

$$s^k := \psi(x^k, y^k) \quad \text{and} \quad \mathcal{T} := \left\{ \psi(x^*, y^*) \mid (x^*, y^*) \in \text{zer } T_{\text{PD}} \right\}. \quad (3.21)$$

Then, the following statements hold.

- (i) The limit points of $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ are in $\Pi_{\mathcal{R}(P)} \text{zer } T_{\text{PD}}$ if and only if the limit points of $(s^k)_{k \in \mathbb{N}}$ are in \mathcal{T} .
- (ii) The sequence $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ converges to $\Pi_{\mathcal{R}(P)} \text{zer } T_{\text{PD}}$ if and only if $(s^k)_{k \in \mathbb{N}}$ converges to \mathcal{T} .

Proof. First, by plugging in (3.14) into (3.10), observe that

$$U = \left[\sqrt{\frac{\tau}{\gamma+\tau}} Z_1 \begin{bmatrix} I_{m_1} \\ -\sqrt{\gamma/\tau} I_{m_1} \end{bmatrix} \quad Z_2 \quad \cdots \quad Z_d \quad Z' \right] = \begin{bmatrix} \sqrt{\frac{\tau}{\gamma+\tau}} X_1 & X_2 & 0 & \cdots & X_d & 0 & X' & 0 \\ -\sqrt{\frac{\gamma}{\gamma+\tau}} Y_1 & 0 & Y_2 & \cdots & 0 & Y_d & 0 & Y' \end{bmatrix}. \quad (3.22)$$

Therefore, ψ corresponds to the linear mapping

$$\psi(x, y) = \left(\sqrt{\frac{\gamma+\tau}{\tau}} I_{m_1} \oplus I_{m+n-2m_1} \right) U^\top \begin{bmatrix} x \\ y \end{bmatrix}.$$

As a result, the claims follow from (3.21), using that $\Pi_{\mathcal{R}(P)} = UU^\top$ and that U has orthonormal columns. \square

Notably, Proposition 3.5 along with Theorem 3.4 establishes the convergence of an $(m+n-m_1)$ -dimensional sequence $(s^k)_{k \in \mathbb{N}}$ of CPA when $\gamma\tau = 1/\|L\|^2$. Since $s^k = x^k - \gamma y^k$ when $L = I$ and $\tau = 1/\gamma$, it follows immediately that Theorem 3.4 matches the convergence results for DRS obtained in [19, Thm. 3.3]. A simple example where $(s^k)_{k \in \mathbb{N}}$ converges while $(z^k)_{k \in \mathbb{N}}$ diverges is provided in Example 3.7.

In the remainder of this section, two theoretical examples are presented to demonstrate some of the main attributes of the convergence results in Theorem 3.4. Supplementary Python code verifying these results can be found on GitHub⁴ and the proofs are deferred to Appendix C.

In the first example, the tightness of the bounds on λ_k from Theorem 3.4 is demonstrated through a simple system of linear equations. In this setting, the iterations of CPA can be expressed as a linear dynamical system, so that tight bounds on λ_k can be obtained by ensuring stability. In this example, an artificial parameter c is introduced when splitting the problem into the form $A + L^\top BL$. While this parameter may appear inconsequential at first sight, it controls the value of β'_D within the problem (see equation (C.5)).

Example 3.6 (saddle point problem). Consider the problem of finding a zero of the following structured linear inclusion

$$0 \in T_{\text{P}}x = \begin{bmatrix} b\ell^2 & a \\ -a & b\ell^2 \end{bmatrix} x = \overbrace{\begin{bmatrix} 0 & a \\ -a & 0 \end{bmatrix}}^A x + \overbrace{\begin{bmatrix} \ell & 0 & 0 \\ 0 & \ell & 0 \end{bmatrix}}^{L^\top} \overbrace{\begin{bmatrix} b & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}}^B \overbrace{\begin{bmatrix} \ell & 0 \\ 0 & \ell \\ 0 & 0 \end{bmatrix}}^L x, \quad (3.23)$$

where $a, b, \ell \in \mathbb{R} \setminus \{0\}$ and $c \in \mathbb{R}$. Any solution to the inclusion problem $0 \in T_{\text{P}}x$ is a minimax solution of $f(x_1, x_2) := ax_1x_2 + \frac{b\ell^2}{2}(x_1^2 - x_2^2)$ when $b > 0$ and a maximin solution when $b < 0$. Consider the sequence $(z^k)_{k \in \mathbb{N}} = (x^k, y^k)_{k \in \mathbb{N}}$ generated by applying CPA to (3.23) with $\tau = \frac{1}{\gamma\|L\|^2}$ and fixed relaxation parameter λ . Then, the following assertions hold.

³When L has orthogonal rows (resp. columns) with identical norm, then X_2 (resp. Y_2) are empty and the terms $X_2^\top x$ (resp. $Y_2^\top y$) vanish.

⁴<https://github.com/brechtveens/Minty-CP-examples>.




Figure 1: Convergence of the sequence $(s^k)_{k \in \mathbb{N}} = (X_1^T x^k - Y_1^T y^k, X_2^T x^k, Y_2^T y^k)_{k \in \mathbb{N}}$ from Example 3.7 for $n = 3$, $\ell_2 = 1/2$, $\ell_3 = 1/5$ and $\lambda = 2.1$. (a) Norm of the sequence $(z^k)_{k \in \mathbb{N}} = (x^k, y^k)_{k \in \mathbb{N}}$. This sequence does not converge, since λ has been selected larger than two (see Theorem 3.4(iv)). (b) Norm of the sequence $(X_1^T x^k - Y_1^T y^k)_{k \in \mathbb{N}}$, which converges to zero. (c) Visualization of the primal sequences $(x^k)_{k \in \mathbb{N}}$ and $(X_2^T x^k)_{k \in \mathbb{N}}$. It can be seen that although $(x^k)_{k \in \mathbb{N}}$ does not converge (its first coordinate diverges), its projection onto the 2-dimensional space spanned by the columns of X_2 does converge to zero (marked by a red dot). (d) Visualization of the dual sequences $(y^k)_{k \in \mathbb{N}}$ and $(Y_2^T y^k)_{k \in \mathbb{N}}$. Analogous to the primal setting, $(y^k)_{k \in \mathbb{N}}$ diverges while $(Y_2^T y^k)_{k \in \mathbb{N}}$ converges to zero.

(i) Examining the spectral radius of the algorithmic operator reveals that $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ converges if and only if $\lambda \in (0, \bar{\lambda})$ and that $(z^k)_{k \in \mathbb{N}}$ converges if and only if $\lambda \in (0, \min\{2, \bar{\lambda}\})$, where

$$\bar{\lambda} := \min \left\{ 2 \left(1 + \frac{b\ell^2}{\gamma(a^2 + b^2\ell^4)} + \frac{ba^2\ell^2\gamma}{a^2 + b^2\ell^4} \right), 2(1 + \gamma c\ell^2) \right\}. \quad (3.24)$$

This upper bound is strictly positive if and only if $\gamma < -\frac{1}{|c|\ell^2}$ and either $b > 0$ or $b < 0$, $a^2 \neq b^2\ell^4$ and $\gamma \in \left(\min\{-\frac{1}{b\ell^2}, -\frac{b\ell^2}{a^2}\}, \max\{-\frac{b\ell^2}{a^2}, -\frac{b\ell^2}{a^2}\} \right)$.

(ii) Theorem 3.4(iv) is tight in the sense that it matches the bounds on λ from (3.24), when in (1.2) the vector v is restricted to $\mathcal{R}(P)$ (see the remark below Theorem 2.1).

(iii) The range of parameters a, b, c and ℓ for which CPA converges includes cases where neither the primal, nor the dual, nor the primal-dual problem are monotone. For instance, when $a = 10$, $b = c = -\frac{1}{4}$ and $\ell = 2$, by Example 3.6(i), $(z^k)_{k \in \mathbb{N}}$ then converges if and only if $\gamma \in (\frac{1}{100}, 1)$ and $\lambda \in (0, 2 - \frac{2}{101\gamma} - \frac{200\gamma}{101})$. \square

The second example focusses on a particular instance of Theorem 3.4 where β_P and β_D are both strictly positive, the number of distinct singular values of L is strictly larger than 1 and $\gamma\tau\|L\|^2 = 1$. Then, the admissible range for λ depends on the second largest singular value of L (see Relaxation parameter rule I). Although the following simple example involves n separable inclusions, it proves sufficient to demonstrate that this result is not merely a consequence of our analysis, but that this is also observed in practice.

Example 3.7 (influence of singular values). Let $n \in \{2, 3, \dots\}$ and $L = \text{diag}(1, \ell_2 \dots \ell_n)$, where $|\ell_k| < 1$, $\forall k \in \{2, \dots, n\}$. Let

$$A = \text{diag}\left(1, 1 + \sqrt{1 - \ell_2^2}, \dots, 1 + \sqrt{1 - \ell_n^2}\right) \quad \text{and} \quad B = \text{diag}\left(1, \frac{1}{1 + \sqrt{1 - \ell_2^2}}, \dots, \frac{1}{1 + \sqrt{1 - \ell_n^2}}\right).$$

Consider the sequences $(z^k)_{k \in \mathbb{N}} = (x^k, y^k)_{k \in \mathbb{N}}$ and $(s^k)_{k \in \mathbb{N}}$ generated by applying CPA to $0 \in Ax + L^T B L x$ with $\gamma = \tau = 1$ and constant λ , where s^k is defined as in (3.21). Then, the following assertions hold.

- (i) The associated primal-dual operator T_{PD} has a $(\frac{1}{2}\mathbf{I}_n \oplus \frac{1}{2}\mathbf{I}_n)$ -oblique weak Minty solution at $(0, 0) = \text{zer } T_{\text{PD}}$.
- (ii) By [Theorem 3.4\(iv\)](#) and [Proposition 3.5](#), both $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ and $(s^k)_{k \in \mathbb{N}}$ converge to zero if λ is selected according to [Relaxation parameter rule I](#), i.e., if $\lambda \in (0, \bar{\lambda})$, where

$$\begin{aligned}\bar{\lambda} &= 2\left(1 + \frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D - \theta_{\gamma\tau}(\max\{|\ell_2|, \dots, |\ell_n|\})\right) \\ &= 3 - \max\{|\ell_2|, \dots, |\ell_n|\}.\end{aligned}$$

- (iii) Let $n = 3$, $\ell_2 \in (0, 1)$ and $\ell_3 = 1/5$. Then, examining the spectral radius of the algorithmic operator reveals that the set of relaxation parameters for which the sequences $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ and $(s^k)_{k \in \mathbb{N}}$ converge is almost entirely covered by [Theorem 3.4\(iv\)](#) (see [Figure 2](#)).




Figure 2: The upper bounds $\bar{\lambda}$ and $\bar{\lambda}_{\text{spectral}}$ for [Example 3.7](#), where $\bar{\lambda}_{\text{spectral}}$ is obtained by examining the spectral radius of the algorithmic operator.

□

4 Semimonotone operators

In this section, we provide calculus rules for the class of (M, R) -semimonotone operators defined in [Definition 1.2](#), generalizing the class of (μ, ρ) -semimonotone operators introduced in [\[19, Sec. 4\]](#). Sufficient conditions for the convergence of [CPA](#) applied to [\(P-I\)](#) for (M, R) -semimonotone operators A and B will be provided in [Section 5](#). The proofs of the calculus rules in this section are deferred to [Appendix C](#).

For some choices of M and R , it follows from the Fenchel-Young inequality that all operators satisfy the definition of (M, R) -semimonotonicity, as stated below.

Proposition 4.1. *Let $M, R \in \mathbb{S}^n$. If $M < 0$, $R < 0$ and $M \leq \frac{1}{4}R^{-1}$, then all operators $A : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ satisfy the definition of (M, R) -semimonotonicity.*

In what follows, various basic properties of (M, R) -semimonotone operators will be provided. For instance, by definition, their inverses belong to the same class of operators, with the roles of M and R reversed. Additionally, the following proposition analyzes scaling and shifting of semimonotone operators, as well as the cartesian product of two semimonotone operators.

Proposition 4.2 (inverting, shifting, scaling and cartesian product). *Let operator $A : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ be (maximally) (M_A, R_A) -semimonotone [at $(x'_A, y'_A) \in \text{gph } A$] and operator $B : \mathbb{R}^m \rightrightarrows \mathbb{R}^m$ be (maximally) (M_B, R_B) -semimonotone [at $(x'_B, y'_B) \in \text{gph } B$]. Let $\alpha \in \mathbb{R}_{++}$. Then, the following hold.*

- (i) *The inverse operator A^{-1} is (maximally) (R_A, M_A) -semimonotone [at $(y'_A, x'_A) \in \text{gph } A^{-1}$].*
- (ii) *For all $u, w \in \mathbb{R}^n$, operator $T(x) := w + \alpha A(x + u)$ is (maximally) $(\alpha M_A, \alpha^{-1} R_A)$ -semimonotone [at $(x'_A - u, w + \alpha y'_A)$].*
- (iii) *Operator $T := A \times B$ is (maximally) $(M_A \oplus M_B, R_A \oplus R_B)$ -semimonotone [at $(x', y') \in \text{gph } T$ where $x' = (x'_A, x'_B)$ and $y' = (y'_A, y'_B)$].*

In [Definition 1.2](#), there is some freedom in selecting the matrices M and R , which might lead to a tradeoff between both. One particular class of operators for which this is true is the class of linear operators. This is summarized in the following proposition, which generalizes [\[6, Prop. 5.1\]](#) for μ -monotone and ρ -comonotone operators and [\[19, Prop. 4.5\]](#) for (μ, ρ) -semimonotone operators.

Proposition 4.3 (linear operator). *Let $D \in \mathbb{R}^{n \times n}$ and let $M, R \in \mathbb{S}^n$. Then, D is (M, R) -semimonotone if and only if $\frac{1}{2}(D + D^\top) - M - D^\top R D \geq 0$.*

Given a certain matrix D and a desired semimonotonicity modulus M , it might be difficult to determine whether there exists an R satisfying $D^\top R D \leq \frac{1}{2}(D + D^\top) - M$, as this corresponds to solving a linear matrix inequality (LMI). The study of LMIs in general form has been extensively explored within the control and

systems theory communities, leading to well-known results such as the Kalman–Yakubovich–Popov lemma, Finsler’s lemma and the (nonstrict) projection lemma [10, 25, 4, 33]. In this work, we rely on a particular result for LMIs of the form $D^\top X D \leq Y$, which is due to [51, 52] and relies upon the classical results from [40, 27, 3] for the linear matrix equality $D^\top X D = Y$.

Proposition 4.4 (symmetric solution of $D^\top X D \leq Y$). *Let $D \in \mathbb{R}^{m \times n}$ and $Y \in \mathbb{S}^n$. Then, the following hold.*

(i) *The set of solutions $C := \{X \in \mathbb{S}^m \mid D^\top X D \leq Y\}$ is nonempty if and only if*

$$\Pi_{\mathcal{N}(D)} Y \Pi_{\mathcal{N}(D)} \geq 0 \quad \text{and} \quad \text{rank}(\Pi_{\mathcal{N}(D)} Y \Pi_{\mathcal{N}(D)}) = \text{rank}(\Pi_{\mathcal{N}(D)} Y). \quad (4.1)$$

(ii) *If (4.1) holds, then $X^* \in C$, where*

$$X^* = \begin{bmatrix} 0 & \mathbf{I} \end{bmatrix} \begin{bmatrix} -Y & D^\top \\ D & 0 \end{bmatrix}^\dagger \begin{bmatrix} 0 \\ \mathbf{I} \end{bmatrix} = (D^\dagger)^\top (Y - Y \Pi_{\mathcal{N}(D)} (\Pi_{\mathcal{N}(D)} Y \Pi_{\mathcal{N}(D)})^\dagger \Pi_{\mathcal{N}(D)} Y) D^\dagger. \quad (4.2)$$

Moreover, $D^\top X D \leq D^\top X^ D \leq Y$ for all $X \in C$.*

(iii) *If the matrix equation $D^\top X D = Y$ is consistent, i.e. if $\mathcal{R}(Y) \subseteq \mathcal{R}(D^\top)$, then $X^* = (D^\dagger)^\top Y D^\dagger$ is the solution of $D^\top X D = Y$ with minimal trace $\text{tr } X^2$.*

Applying this result to [Proposition 4.3](#), the following corollary for linear operators is obtained.

Corollary 4.5 (linear operator). *Let $D \in \mathbb{R}^{n \times n}$ and $M \in \mathbb{S}^n$. Then, the following hold.*

(i) *There exists $R \in \mathbb{S}^n$ such that D is (M, R) -semimonotone if and only if*

$$\Pi_{\mathcal{N}(D)} M \Pi_{\mathcal{N}(D)} \leq 0 \quad \text{and} \quad \text{rank}(\Pi_{\mathcal{N}(D)} M \Pi_{\mathcal{N}(D)}) = \text{rank}(\Pi_{\mathcal{N}(D)} (\frac{1}{2}D - M)). \quad (4.3)$$

(ii) *If (4.3) holds, then D is (M, R^*) -semimonotone, where*

$$R^* = \begin{bmatrix} 0 & \mathbf{I} \end{bmatrix} \begin{bmatrix} M - \frac{1}{2}(D + D^\top) & D^\top \\ D & 0 \end{bmatrix}^\dagger \begin{bmatrix} 0 \\ \mathbf{I} \end{bmatrix}.$$

In particular, when D is either symmetric or skew-symmetric, it holds that

$$R^* = 2(D + D^\top)^\dagger - D^{\dagger\top} M D^\dagger + (D^\dagger)^\top M \Pi_{\mathcal{N}(D)} (\Pi_{\mathcal{N}(D)} M \Pi_{\mathcal{N}(D)})^\dagger \Pi_{\mathcal{N}(D)} M D^\dagger. \quad (4.4)$$

Note that R^* can be seen as an optimal choice for R , as it solves the LMI from [Proposition 4.3](#) as tightly as possible. A second consequence of [Proposition 4.4](#) is the following result on operators of the form DTD^\top .

Corollary 4.6 (semimonotonicity of DTD^\top). *Let $D \in \mathbb{R}^{n \times m}$, $M, Y \in \mathbb{S}^m$ and let operator $T : \mathbb{R}^m \rightrightarrows \mathbb{R}^n$ be (M, Y) -semimonotone [at $(D^\top x', y') \in \text{gph } T$]. If (4.1) holds for D and T , then DTD^\top is (DMD^\top, X^*) -semimonotone [at (x', Dy')], where X^* is given by (4.2).*

Leveraging the previous result for the semimonotonicity of DTD^\top , the semimonotonicity of the sum and parallel sum of two semimonotone operators is investigated next. Recall that the parallel sum of two operators $A, B : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is defined as $A \square B := (A^{-1} + B^{-1})^{-1}$. We shall also use the parallel sum between symmetric matrices $X, Y \in \mathbb{S}^n$ as defined below.

Definition 4.7 (parallel sum between symmetric matrices). *Let $X, Y \in \mathbb{S}^n$. We say that X and Y are parallel summable if $\mathcal{R}(X) \subseteq \mathcal{R}(X + Y)$ or equivalently $\mathcal{R}(Y) \subseteq \mathcal{R}(X + Y)$. For parallel summable matrices X and Y , their parallel sum is defined as [35, Cor. 9.2.5]*

$$X \square Y := X(X + Y)^\dagger Y = Y(X + Y)^\dagger X = X - X(X + Y)^\dagger X = Y - Y(X + Y)^\dagger Y.$$

For scalars $\alpha, \beta \in \mathbb{R}$, we say that α and β are parallel summable if either $\alpha = \beta = 0$ or $\alpha + \beta \neq 0$ and their parallel sum is defined as

$$\alpha \square \beta := \alpha(\alpha + \beta)^\dagger \beta = \begin{cases} 0, & \text{if } \alpha = \beta = 0, \\ \frac{\alpha\beta}{\alpha + \beta}, & \text{otherwise.} \end{cases}$$

In addition, consider the following set, which will be referred to as the effective domain of the parallel sum between two symmetric matrices.

Definition 4.8 (effective domain of parallel sum). *The set*

$$\text{dom}_{\square} := \{(X, Y) \in \mathbb{S}^n \times \mathbb{S}^n \mid X + Y \geq 0, X \text{ and } Y \text{ are parallel summable}\} \quad (4.5)$$

is the effective domain of the parallel sum between two symmetric (possibly indefinite) matrices. Let $X = \alpha \mathbf{I}_n$ and $Y = \beta \mathbf{I}_n$ where $\alpha, \beta \in \mathbb{R}$. Then, $(X, Y) \in \text{dom}_{\square}$ reduces to

$$(\alpha, \beta) \in \text{dom}_{\square} = \{(\alpha, \beta) \mid \alpha + \beta > 0 \text{ or } \alpha = \beta = 0\}.$$

In the upcoming two propositions, it is shown that the sum and parallel sum of two semimonotone operators are also semimonotone operators, if the involved semimonotonicity matrices belong to the effective domain of the parallel sum. The first result generalizes [19, Prop. 4.7] for the sum of two (μ, ρ) -semimonotone operators.

Proposition 4.9 (sum and parallel sum). *Let operator $A : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ be (M_A, R_A) -semimonotone [at $(x'_A, y'_A) \in \text{gph } A$] and operator $B : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ be (M_B, R_B) -semimonotone [at $(x'_B, y'_B) \in \text{gph } B$].*

- (i) *If $(R_A, R_B) \in \text{dom}_{\square}$ [and $x'_A = x'_B =: x'$], then $A + B$ is $(M_A + M_B, R_A \square R_B)$ -semimonotone [at $(x', y'_A + y'_B)$].*
- (ii) *If $(M_A, M_B) \in \text{dom}_{\square}$ [and $y'_A = y'_B =: y'$], then $A \square B$ is $(M_A \square M_B, R_A + R_B)$ -semimonotone [at $(x'_A + x'_B, y')$].*

When one of the two involved operators is linear, more precise statements for the resulting semimonotonicity matrices can be derived. For instance, consider the following lemma for the sum of a semimonotone operator and a skew-symmetric matrix. This result will be used later in [Theorem 5.1](#) for analyzing the primal-dual operator T_{PD} .

Lemma 4.10 (sum with skew-symmetric matrix). *Let $D \in \mathbb{R}^{n \times n}$ be a skew-symmetric matrix and operator $T : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ be $(D^{\top} M D, R + R')$ -semimonotone [at $(x', y') \in \text{gph } T$], where $\mathcal{R}(R') \subseteq \mathcal{N}(D)$ and $(M, R) \in \text{dom}_{\square}$. Then, $T + D$ is $(0, R' + M \square R)$ -semimonotone [at $((x'_A, y'_B), (y'_A + L^{\top} y'_B, x'_B - L x'_A))$].*

5 Chambolle–Pock for semimonotone operators

In [Section 3](#), convergence of [CPA](#) was established under an oblique weak Minty assumption on the underlying primal-dual operator. This section provides a set of sufficient conditions for the convergence of [CPA](#) for composite inclusion problems involving semimonotone operators.

5.1 Existence of oblique weak Minty solutions

The main tool for establishing simplified conditions for [CPA](#) for semimonotone operators is the following calculus rule, which connects the semimonotonicity of the individual operators A and B to the existence of V -oblique weak Minty solutions of the primal-dual operator T_{PD} .

Theorem 5.1 (primal-dual operator). *In the primal-dual inclusion (PD-I), suppose that there exists a nonempty set $\mathcal{S}^* \subseteq \text{zer } T_{\text{PD}}$ and matrices $R_A, R'_A, R_B \in \mathbb{S}^n$ and $M_A, M_B, M'_B \in \mathbb{S}^m$ such that for every $z^* = (x^*, y^*) \in \mathcal{S}^*$ the following hold.*

- (i) $\mathcal{R}(R_A) \subseteq \mathcal{R}(L^{\top})$, $\mathcal{R}(R'_A) \subseteq \mathcal{N}(L)$, $\mathcal{R}(M_B) \subseteq \mathcal{R}(L)$ and $\mathcal{R}(M'_B) \subseteq \mathcal{N}(L^{\top})$.
- (ii) $(M_A, M_B) \in \text{dom}_{\square}$ and $(R_A, R_B) \in \text{dom}_{\square}$.
- (iii) Operator A is $(L^{\top} M_A L, R_A + R'_A)$ -semimonotone at $(x^*, -L^{\top} y^*) \in \text{gph } A$.
- (iv) Operator B is $(M_B + M'_B, L R_B L^{\top})$ -semimonotone at $(L x^*, y^*) \in \text{gph } B$.

Then, T_{PD} has $((R_A \square R_B + R'_A) \oplus (M_A \square M_B + M'_B))$ -oblique weak Minty solutions at \mathcal{S}^* .

Proof. Let $(x^*, y^*) \in \mathcal{S}^*$ and decompose the primal-dual operator as $T_{\text{PD}} = T + D$, where $T := A \times B^{-1}$ and $D(x, y) := (L^\top y, -Lx)$. By [Proposition 4.2\(i\)](#), B^{-1} is $(LR_B L^\top, M_B + M'_B)$ -semimonotone at $(y^*, Lx^*) \in \text{gph } B^{-1}$, so that T is $(D^\top(M_A \oplus R_B)D, (R_A + R'_A) \oplus (M_B + M'_B))$ -semimonotone at $((x^*, y^*), (-L^\top y^*, Lx^*)) \in \text{gph } T$ due to [Proposition 4.2\(iii\)](#). Consequently, by [Lemma 4.10](#) it follows that T_{PD} is $(0, (R_A \square R_B + R'_A) \oplus (M_A \square M_B + M'_B))$ -semimonotone at $((x^*, y^*), 0)$. The claim then follows by [Definition 1.1](#). \square

Suppose that the underlying assumptions from [Theorem 5.1](#) hold, in which case $\mathcal{R}(R_A \square R_B) \subseteq \mathcal{R}(R_A) \subseteq \mathcal{R}(L^\top)$ and $\mathcal{R}(M_A \square M_B) \subseteq \mathcal{R}(M_B) \subseteq \mathcal{R}(L)$. Then, by virtue of [\(3.2\)](#) and the particular form of $V = ((R_A \square R_B + R'_A) \oplus (M_A \square M_B + M'_B))$ from [Theorem 5.1](#), the primal-dual operator T_{PD} has V -oblique weak Minty solutions at \mathcal{S}^* , with V given by [\(3.3\)](#) and

$$\beta_P = \lambda_{\min}(X^\top(R_A \square R_B)X), \quad \beta_D = \lambda_{\min}(Y^\top(M_A \square M_B)Y), \quad \beta'_P = \lambda_{\min}(X'^\top R'_A X'), \quad \beta'_D = \lambda_{\min}(Y'^\top M'_B Y'), \quad (5.1)$$

where X, X', Y, Y' are defined as in [\(3.1\)](#). Hence, [Assumption II.A3](#) holds if the parameters from [\(5.1\)](#) satisfy condition [\(3.5\)](#).

As an implication of [Theorem 5.1](#), one can easily verify the underlying assumptions of [Theorem 3.4](#). For instance, consider the following numerical example, where [Theorem 3.4](#) is applied to a nonconvex QP with an indefinite Q matrix, where L is full row rank. An example where L is rank-deficient is provided later in [Example 5.6](#).

Example 5.2. Consider the quadratic program

$$\underset{x \in \mathbb{R}^n}{\text{minimize}} \quad \frac{1}{2} x^\top Q x + q^\top x \quad \text{subject to} \quad Lx \in C, \quad (5.2)$$

where $Q = \text{diag}(1, -1, 2)$, $q = [-1 \ 1 \ -1]^\top$, $L = \begin{bmatrix} 1 & 1/4 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ and $C := \{x \in \mathbb{R}^2 \mid 2 \leq x_i \leq 4, i = 1, 2\}$. Then, the associated first-order optimality conditions are given by $0 \in Ax + L^\top B Lx$, where $A : x \mapsto Qx + q$ and $B := N_C$, and the global minimizer of [\(5.2\)](#) is given by $x^* = [1 \ 4 \ 1/2]^\top$. The following assertions hold.

(i) Operator A is $(\text{diag}(1, -1, 0), \text{diag}(0, 0, 1/2))$ -semimonotone.

(ii) Operator B is $(\text{diag}(0, 3/2), 0)$ -semimonotone at $(Lx^*, -L^\top A x^*) = \left(\begin{bmatrix} 2 \\ 4 \end{bmatrix}, \begin{bmatrix} 0 \\ 3 \end{bmatrix}\right) \in \text{gph } B$.

(iii) The primal-dual pair $(x^*, -L^\top A x^*) \in \text{zer } T_{\text{PD}}$ is a V -oblique weak Minty solution of T_{PD} with V given by [\(3.3\)](#), where $\beta_P = 0, \beta_D = -3, \beta'_P = 1/2$ and $\beta'_D = 0$ as in [\(5.1\)](#).

(iv) The sequence $(z^k)_{k \in \mathbb{N}} = (x^k, y^k)_{k \in \mathbb{N}}$ generated by [CPA](#) with fixed relaxation parameter λ converges for

$$\gamma \in \left(0, \frac{-1}{\beta_D \|L\|^2}\right) \approx (0, 0.26), \quad \tau \in \left(-\beta_D, \frac{1}{\gamma \|L\|^2}\right) \approx \left(3, \frac{0.779}{\gamma}\right], \quad \lambda \in (0, 2 + \frac{2}{\tau} \beta_D) = \left(0, 2 - \frac{6}{\tau}\right),$$

where we used that $\|L\|^2 = \frac{33 + \sqrt{65}}{32} \approx 1.28$. \square

Proof. [5.2\(i\)](#) follows from [Propositions 4.2\(ii\)](#) and [4.3](#); [5.2\(ii\)](#) follows from [Proposition B.5](#) since $L^\dagger = \begin{bmatrix} 1 & 0 & 0 \\ -1/4 & 1 & 0 \end{bmatrix}^\top$; [5.2\(iii\)](#) holds by [Theorem 5.1](#) with $M_A = \frac{1}{16} \begin{bmatrix} 16 & -4 \\ -4 & -15 \end{bmatrix}$, $M_B = \text{diag}(0, 3/2)$, $M'_B = 0_{2 \times 2}$, $R'_A = \text{diag}(0, 0, 1/2)$ and $R_A = R_B = 0_{3 \times 3}$, using that $\Pi_{\mathcal{R}(L^\top)} = \text{diag}(1, 1, 0)$, $L^\top M_A L = \text{diag}(1, -1, 0)$ and $M_A \square M_B = \text{diag}(0, -3)$; [5.2\(iv\)](#) follows directly from [Theorem 3.4](#) by continuity of $J_{\gamma A}$ and $J_{\tau B^{-1}}$. \square

5.2 Sufficient conditions for convergence of CPA

[Theorem 5.1\(i\)](#) imposes range conditions on the semimonotonicity matrices of A and B . In this subsection, it is shown that this can be achieved by imposing a certain structure on these matrices. In particular, consider the following set of assumptions.

Assumption III. In problem [\(P-I\)](#), suppose that $\text{zer } T_{\text{PD}}$ is nonempty and that there exists a nonempty set $\mathcal{S}^* \subseteq \text{zer } T_{\text{PD}}$ such that for every $z^* = (x^*, y^*) \in \mathcal{S}^*$ it holds that A is $(\mu_A L^\top L, \rho_A I_n)$ -semimonotone at $(x^*, -L^\top y^*) \in \text{gph } A$, B is $(\mu_B I_m, \rho_B L L^\top)$ -semimonotone at $(Lx^*, y^*) \in \text{gph } B$ and the semimonotonicity moduli $(\mu_A, \mu_B, \rho_A, \rho_B) \in \mathbb{R}^4$ satisfy either one of the following conditions.

(i) (either) $\mu_A = \mu_B = 0$ and $\rho_A = \rho_B = 0$.

Table 3: Range of the stepsizes γ and τ for CPA involving semimonotone operators.

| | | $\mu_A = \mu_B = 0$ | $\mu_A + \mu_B > 0$ | |
|-----------------------|------------------------|--|---|---|
| | | | $\mu_A \mu_B \geq 0$ | $\mu_A \mu_B < 0$ |
| $\rho_A = \rho_B = 0$ | | $\gamma \in (0, +\infty)$ $\tau \in (0, \frac{1}{\gamma \ L\ ^2}]$ | $\gamma \in (0, -\frac{1}{(\mu_A \square \mu_B) \ L\ ^2})$ $\tau \in (-\mu_A \square \mu_B, \frac{1}{\gamma \ L\ ^2}]$ | |
| $\rho_A + \rho_B > 0$ | $\rho_A \rho_B \geq 0$ | | $\gamma \in (0, \gamma_{\max})$ $\tau \in (\tau_{\min}(\gamma), \frac{1}{\gamma \ L\ ^2}]$ | |
| | $\rho_A \rho_B < 0$ | $\gamma \in (-\rho_A \square \rho_B, +\infty)$ $\tau \in (0, \frac{1}{\gamma \ L\ ^2}]$ | $\gamma \in (\gamma_{\min}, +\infty)$ $\tau \in (\tau_{\min}(\gamma), \frac{1}{\gamma \ L\ ^2}]$ | $\gamma \in (\gamma_{\min}, \gamma_{\max})$ $\tau \in (\tau_{\min}(\gamma), \frac{1}{\gamma \ L\ ^2}]$ |

(ii) (or) $\mu_A + \mu_B > 0$ and $\rho_A = \rho_B = 0$.

(iii) (or) $\rho_A + \rho_B > 0$ and $\mu_A = \mu_B = 0$.

(iv) (or) $\mu_A + \mu_B > 0$, $\rho_A + \rho_B > 0$ and $[\mu_A \square \mu_B]_- [\rho_A \square \rho_B]_- < \frac{1}{4\|L\|^2}$.

Assumption III provides a set of sufficient conditions on the operators A and B so that the primal-dual operator admits oblique weak Minty solutions. This key result is stated in the following corollary.

Corollary 5.3. Suppose that **Assumption III** holds. Then, the primal-dual operator T_{PD} has V -oblique weak Minty solutions at \mathcal{S}^* , where V is given by (3.3) and

$$\beta_{\text{P}} = \rho_A \square \rho_B, \quad \beta_{\text{D}} = \mu_A \square \mu_B, \quad \beta'_{\text{P}} = \begin{cases} 0, & \text{if rank } L = n, \\ \rho_A, & \text{if rank } L < n, \end{cases} \quad \beta'_{\text{D}} = \begin{cases} 0, & \text{if rank } L = m, \\ \mu_B, & \text{if rank } L < m. \end{cases} \quad (5.3)$$

Proof. Note that $(\mu_A, \mu_B) \in \text{dom}_{\square}$ and $(\rho_A, \rho_B) \in \text{dom}_{\square}$ by **Assumption III** and that $\rho_A \mathbf{I}_n = \rho_A \Pi_{\mathcal{R}(L^{\top})} + \beta'_{\text{P}} \Pi_{\mathcal{N}(L)}$ and $\mu_B \mathbf{I}_m = \mu_B \Pi_{\mathcal{R}(L)} + \beta'_{\text{D}} \Pi_{\mathcal{N}(L^{\top})}$. Therefore, the claim follows directly from **Theorem 5.1**, with $M_A = \mu_A \Pi_{\mathcal{R}(L)}$, $R_A = \rho_A \Pi_{\mathcal{R}(L^{\top})}$, $R'_A = \beta'_{\text{P}} \Pi_{\mathcal{N}(L)}$, $M_B = \mu_B \Pi_{\mathcal{R}(L)}$, $M'_B = \beta'_{\text{D}} \Pi_{\mathcal{N}(L^{\top})}$ and $R_B = \rho_B \Pi_{\mathcal{R}(L^{\top})}$. \square

Leveraging this key result, the convergence of CPA for semimonotone operators is analyzed in **Corollary 5.4**, for which the stepsize rule corresponds to a simple look-up table. The proof is deferred to **Appendix C**.

Stepsize rule II. The stepsizes γ and τ satisfy the bounds provided in **Table 3**, where γ_{\min} , γ_{\max} and $\tau_{\min}(\gamma)$ are defined as in (3.6) with $\beta_{\text{D}} = \mu_A \square \mu_B$ and $\beta_{\text{P}} = \rho_A \square \rho_B$.

Relaxation parameter rule II. Let η' be defined as in **Table 4** and define

$$\Delta_{\gamma, \tau} := \frac{1}{2\gamma}(\rho_A \square \rho_B) + \frac{1}{2\tau}(\mu_A \square \mu_B) \quad \text{and} \quad \theta_{\gamma, \tau}(\sigma) := \sqrt{\Delta_{\gamma, \tau}^2 + (\mu_A \square \mu_B)(\rho_A \square \rho_B)\sigma^2}.$$

The sequence $(\lambda_k)_{k \in \mathbb{N}}$ satisfies $\lambda_k \in (0, 2(1 + \bar{\eta}))$ and $\liminf_{k \rightarrow \infty} \lambda_k(2(1 + \bar{\eta}) - \lambda_k) > 0$, where

$$\bar{\eta} := \begin{cases} \left\{ \begin{array}{ll} \left(\Delta_{\gamma, \tau} - \theta_{\gamma, \tau}(\|L\|), & \text{if } \mu_A \mu_B \rho_A \rho_B \geq 0 \\ \min\{\Delta_{\gamma, \tau} - \theta_{\gamma, \tau}(\sigma_d), \eta'\}, & \text{otherwise} \end{array} \right\} & \text{if } \gamma\tau < \frac{1}{\|L\|^2}, \\ \left\{ \begin{array}{ll} 2\Delta_{\gamma, \tau}, & \text{if } \max\{\mu_A \mu_B, \rho_A \rho_B\} \leq 0 \\ \min\{2\Delta_{\gamma, \tau}, \eta'\}, & \text{otherwise} \end{array} \right\} & \text{if } \gamma\tau = \frac{1}{\|L\|^2} \text{ and } d = 1, \\ \left\{ \begin{array}{ll} 2\Delta_{\gamma, \tau}, & \text{if } \max\{\mu_A \mu_B, \rho_A \rho_B\} \leq 0 \\ \Delta_{\gamma, \tau} - \theta_{\gamma, \tau}(\sigma_2), & \text{if } \min\{\mu_A \mu_B, \rho_A \rho_B\} > 0 \\ \min\{\Delta_{\gamma, \tau} - \theta_{\gamma, \tau}(\sigma_d), \eta'\}, & \text{otherwise} \end{array} \right\} & \text{if } \gamma\tau = \frac{1}{\|L\|^2} \text{ and } d > 1. \end{cases}$$

Corollary 5.4 (convergence of CPA under semimonotonicity). Suppose that **Assumption II.A1**, **Assumption II.A2** and **Assumption III** hold, that γ and τ are selected according to **Stepsize rule II** and that the relaxation sequence $(\lambda_k)_{k \in \mathbb{N}}$ is selected according to **Relaxation parameter rule II**. Then, all the claims of **Theorem 3.4** hold.

Table 4: Definition of η' in *Relaxation parameter rule II*.

| | rank $L = n$ | rank $L < n$ |
|--------------|-----------------------|---|
| rank $L = m$ | $+\infty$ | $\frac{1}{\gamma}\rho_A$ |
| rank $L < m$ | $\frac{1}{\tau}\mu_B$ | $\min\{\frac{1}{\gamma}\rho_A, \frac{1}{\tau}\mu_B\}$ |

Corollary 5.4 serves as a universal framework for analyzing the convergence of **CPA**, both in monotone and nonmonotone settings. Notably, it encompasses and extends many of the existing results in literature. Several examples are provided below.

Remark 5.5 (connection to existing theory). Case (i) of **Assumption III** can be interpreted as a relaxation of the classical monotonicity assumption for **CPA** [12], as it does not impose monotonicity between any two pairs in the graph of A and B . In case (ii) of **Assumption III**, a monotone problem is split in a nonmonotone fashion. In the optimization setting, this setting was studied in [36]. To see this, let g be a proper lsc μ_g -convex function with $\mu_g > 0$ and h be a proper lsc μ_h -convex function. Then, $A = \partial g$ is $(\mu_g/\|L\|^2 L^\top L, 0)$ -semimonotone and $B = \partial h$ is $(\mu_h, 0)$ -semimonotone. Then, **Corollary 5.4** requires that $\mu_g/\|L\|^2 + \mu_h > 0$, which matches [36, Thm. 2.8]. Note that case (iii) of **Assumption III** can be interpreted as the dual counterpart of case (ii), as the assumptions of the latter hold for A and B in the primal inclusion problem if and only if the assumptions of the former hold for B^{-1} and A^{-1} in the dual one (see **(D-I)**). Up to the knowledge of the authors, no particular instances of case (iv) of **Assumption III** have been covered in literature, even in the minimization setting. \square

In what follows, **Corollary 5.4** will be applied to several examples previously discussed in this paper. First of all, consider the following constrained QP, where Q is an indefinite matrix and L is rank-deficient. Previously, it was shown for the nonconvex QP from **Example 5.2** that convergence of **CPA** can be established using **Theorem 5.1** and **Theorem 3.4**. This example demonstrates that if the monotonicity of A can be expressed in the form $\mu_A L^\top L$, then this result can be obtained directly using **Corollary 5.4**.

Example 5.6. Consider a QP of the same form as in **Example 5.2**, where now $Q = \text{diag}(-3, -2, 1)$, $q = [0 \ 1 \ 0]^\top$, $L = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & -1 & 0 \end{bmatrix}$ and $C = \{x \in \mathbb{R}^2 \mid 1/2 \leq x_i \leq 1, i = 1, 2, 3\}$. Then, the global minimizer is given by $x^* = [1 \ 0 \ 0]^\top$ and the following hold.

(i) Operator A is $(\mu_A L^\top L, \text{diag}(0, 0, 1))$ -semimonotone where $\mu_A = -1$.

(ii) Operator B is $(\mu_B I, 0)$ -semimonotone at $(Lx^*, -L^\top A x^*) = ([1 \ 1 \ 1]^\top, [1 \ 1 \ 1]^\top) \in \text{gph } B$ where $\mu_B = 2$.

(iii) The sequence $(z^k)_{k \in \mathbb{N}} = (x^k, y^k)_{k \in \mathbb{N}}$ generated by **CPA** with fixed relaxation parameter λ converges for

$$\gamma \in (0, \frac{-1}{(\mu_A \square \mu_B) \|L\|^2}) = (0, \frac{1}{6}), \quad \tau \in (-(\mu_A \square \mu_B), \frac{1}{\gamma \|L\|^2}) = (2, \frac{1}{3\gamma}], \quad \lambda \in (0, 2 + \frac{2}{\tau}(\mu_A \square \mu_B)) = (0, 2 - \frac{4}{\tau}). \quad \square$$

Proof. **5.6(i)** follows from **Propositions 4.2(ii)** and **4.3**; **5.6(ii)** holds by **Proposition B.5** since $L^\dagger = \frac{1}{6} \begin{bmatrix} 2 & 2 & 2 \\ 0 & 3 & -3 \\ 0 & 0 & 0 \end{bmatrix}$; **5.6(iii)** follows from **Corollary 5.4**, using that A is $(\mu_A L^\top L, 0)$ -semimonotone and continuity of $J_{\gamma A}$ and $J_{\tau B^{-1}}$. \square

Next, we revisit the two theoretical examples of **Section 3**, this time under the lens of semimonotonicity. First of all, consider the linear inclusion problem from **Example 3.6**, where the parameters a, b, c, l are selected as in **Example 3.6(iii)**.

Example 5.7 (saddle point problem (revisited)). Consider inclusion problem (3.23) with $a = 10$, $b = c = -\frac{1}{4}$ and $\ell = 2$. Using **Proposition 4.3**, it follows that A is $(L^\top L, -\frac{1}{25} I_n)$ -semimonotone and B is $(-\frac{3}{10} I_m, \frac{1}{5} L^\top L)$ -semimonotone. By **Corollary 5.4**, the sequence $(z^k)_{k \in \mathbb{N}}$ generated by applying **CPA** to (3.23) with $\tau = \frac{1}{\gamma \|L\|^2}$ and fixed relaxation parameter λ converges for

$$\gamma \in (\gamma_{\min}, \gamma_{\max}) \approx (0.055, 0.528) \quad \text{and} \quad \lambda \in (0, 2 - \frac{1}{10\gamma} - \frac{24\gamma}{7}). \quad \square$$

The obtained range of stepsize parameters is only a subset of the tight range obtained in [Example 3.6\(iii\)](#). However, this should not come as a surprise, since part of the information about operators A and B is lost by analyzing them under the lens of semimonotonicity. This is also observed in the second example.

Example 5.8 (influence of singular values (revisited)). Consider the composite inclusion problem $0 \in Ax + L^\top BLx$ from [Example 3.7](#). It follows from [Proposition 4.3](#) that A is $(\frac{1}{2}L^\top L, \frac{1}{2}I_n)$ -semimonotone and B is $(\frac{1}{2}I_n, \frac{1}{2}LL^\top)$ -semimonotone. By [Corollary 5.4](#) and [Proposition 3.5](#), the sequences $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ and $(s^k)_{k \in \mathbb{N}}$ generated by applying CPA with $\gamma = \tau = 1$ and fixed relaxation parameter λ , converge to zero if λ is selected according to [Relaxation parameter rule II](#), which reduces to

$$\lambda \in \left(0, 2\left(1 + \frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D - \theta_{\gamma\tau}(\max\{\ell_2, \dots, \ell_n\})\right)\right) = \left(0, 5/2 - 1/2 \max\{\ell_2, \dots, \ell_n\}\right). \quad \square$$

In [Example 3.7\(i\)](#), it was shown that T_{PD} has $(\frac{1}{2}I_n \oplus \frac{1}{2}I_n)$ -oblique weak Minty solutions at $\text{zer } T_{PD}$. On the other hand, here it is shown that A is $(\frac{1}{2}L^\top L, \frac{1}{2}I_n)$ -semimonotone and B is $(\frac{1}{2}I_n, \frac{1}{2}LL^\top)$ -semimonotone. Applying [Corollary 5.3](#), this implies that T_{PD} only has a $(\frac{1}{4}I_n \oplus \frac{1}{4}I_n)$ -oblique weak Minty solution at $(0_n, 0_n) = \text{zer } T_{PD}$. By analyzing A and B under the lens of semimonotonicity, some additional looseness is inevitably introduced. More specifically, the information that A and B are linear and symmetric, that $A = B^{-1}$ and that $A = 1/2L^\top L + 1/2A^\top A$ are lost in this process.

6 Conclusion

In this work, convergence of the Chambolle–Pock algorithm (CPA) was established for a class of nonmonotone problems, characterized by an oblique weak Minty assumption on the associated primal-dual operator. To facilitate the verification of this underlying assumption, a generalization of the class of semimonotone operators (see [\[19\]](#)) was introduced, and sufficient conditions for the convergence of CPA were provided for inclusion problems involving operators belonging to this class. When restricting to minimization problems, our results reveal that for certain problem classes no explicit rank or condition number restriction on the linear mapping is required.

It would be interesting to explore if stronger results can be obtained when the operators are known to be subdifferentials. Other future research directions include extensions to the setting where the preconditioning is indefinite, allowing to cover the extended Chambolle–Pock stepsize range $\gamma\tau\|L\|^2 \leq 4/3$ from [\[31, 5\]](#), as well as analyzing other splitting methods in nonmonotone settings.

A Supplementary results

Theorem A.1. *Suppose that [Assumption II](#) holds, with condition [\(3.5\)](#) relaxed to*

$$[\beta_P]_-, [\beta_D]_- < \frac{1}{4\|L\|^2}, \quad [\beta'_P]_-, [\beta'_D]_- < \frac{1}{\|L\|^2}, \quad [\beta'_P]_- > \frac{\delta + \sqrt{\delta^2 - 4\beta_P\beta_D\|L\|^2}}{2[\beta_D]_- \|L\|^2} \quad \text{and} \quad [\beta'_D]_- > \frac{\delta + \sqrt{\delta^2 - 4\beta_P\beta_D\|L\|^2}}{2[\beta_P]_- \|L\|^2}. \quad (\text{A.1})$$

Suppose that γ and τ are selected within the intersection of the bounds provided in [Stepsize rule I](#) and the set

$$\left\{(\gamma, \tau) \in \mathbb{R}^2 \mid \gamma \in \left(-[\beta'_P]_-, \frac{1}{-[\beta'_D]_- \|L\|^2}\right), \tau \in \left(-[\beta'_D]_-, \frac{1}{\gamma \|L\|^2}\right)\right\}. \quad (\text{A.2})$$

Moreover, suppose that the relaxation sequence $(\lambda_k)_{k \in \mathbb{N}}$ is selected according to [Relaxation parameter rule I](#). Then, all the claims from [Theorem 3.4](#) hold.

Proof. The proof is identical to the proof of [Theorem 3.4](#), except that in the item discussing $1 + \eta > 0$ the occurrence of [\(3.5\)](#) is replaced by [\(A.1\)](#) and that the item discussing $1 + \bar{\eta} > 0$ is replaced by the following.

♣ *The set of pairs $(\gamma, \tau) \in \mathbb{R}_{++}^2$ satisfying $\gamma\tau \in (0, 1/\|L\|^2]$ and $1 + \bar{\eta} > 0$ is given by the intersection of the bounds provided in [Stepsize rule I](#) and the set [\(A.2\)](#), which is nonempty.*

With the understanding that $\beta'_P = 0$ when L is full column rank and $\beta'_D = 0$ when L is full row rank, as discussed below [Assumption II](#), the set of pairs $(\gamma, \tau) \in \mathbb{R}_{++}^2$ satisfying $\gamma\tau \in (0, 1/\|L\|^2]$ and $1 + \eta' > 0$ is given by [\(A.2\)](#). Hence, it only remains to verify that the intersection of the bounds provided in [Stepsize rule I](#) and the set [\(A.2\)](#) is nonempty, which is ensured by the conditions on β'_P and β'_D provided in [\(A.1\)](#). \square

B Auxiliary results

Fact B.1 (solution of quadratic inequality). *Let $\beta_P, \beta_D \in \mathbb{R}$, $\|L\| > 0$, $\sigma_d \in (0, \|L\|]$, let δ be defined as in (3.4) and let γ_{\min} and γ_{\max} be defined as in [Stepsize rule 1](#). Then, the following hold.*

- (i) *There exists a $\gamma > 0$ satisfying $\beta_D \|L\|^2 \gamma^2 + \delta \gamma + \beta_P > 0$ if and only if $[\beta_P]_- [\beta_D]_- < \frac{1}{4\|L\|^2}$.*
- (ii) *If $[\beta_P]_- [\beta_D]_- < \frac{1}{4\|L\|^2}$, then $\gamma > 0$ satisfies $\beta_D \|L\|^2 \gamma^2 + \delta \gamma + \beta_P > 0$ if and only if γ lies within the (nonempty) stepsize interval provided in [Table 1](#).*

Lemma B.2. *Let $\beta_P, \beta_D \in \mathbb{R}$, $\|L\| > 0$, $\sigma_d \in (0, \|L\|]$ and for any $\sigma \in [\sigma_d, \|L\|]$ define the set*

$$\Gamma(\sigma) := \left\{ (\gamma, \tau) \in \mathbb{R}_{++}^2 \mid \gamma\tau \in (0, 1/\|L\|^2] \text{ and } 1 + \frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D > \theta_{\gamma\tau}(\sigma) \right\},$$

where $\theta_{\gamma\tau}(\cdot)$ is defined as in (3.7). Then, the following hold.

- (i) *If $\min\{\beta_P, \beta_D\} \geq 0$, then, for any $\sigma \in [\sigma_d, \|L\|]$, all stepsizes within the nonempty set $\Gamma(\sigma)$ are characterized by [Table 1](#).*
- (ii) *If $\min\{\beta_P, \beta_D\} < 0$ and $\max\{\beta_P, \beta_D\} \geq 0$, then all stepsizes within the nonempty set $\Gamma(\sigma_d)$ are characterized by [Table 1](#).*
- (iii) *If $\max\{\beta_P, \beta_D\} < 0$, then the set $\Gamma(\|L\|)$ is nonempty if and only if $[\beta_P]_- [\beta_D]_- < \frac{1}{4\|L\|^2}$, in which case all stepsizes within $\Gamma(\|L\|)$ are characterized by [Table 1](#).*

Proof. Let $\sigma \in [\sigma_d, \|L\|]$. Solving the square root inequality $1 + \frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D > \theta_{\gamma\tau}(\sigma)$, it follows that

$$\Gamma(\sigma) = \left\{ (\gamma, \tau) \in \mathbb{R}_{++}^2 \mid \gamma\tau \in (0, 1/\|L\|^2], \overbrace{1 + \frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D}^{\Gamma_1 :=} > 0 \text{ and } \overbrace{1 + \frac{1}{\gamma}\beta_P + \frac{1}{\tau}\beta_D + \frac{1}{\gamma\tau}\beta_P\beta_D(1 - \gamma\tau\sigma^2)}^{\Gamma_2(\sigma) :=} > 0 \right\}.$$

Define $c_1(\sigma, \gamma) := \gamma(1 - \beta_P\beta_D\sigma^2) + \beta_P$ and $c_2(\gamma) := \beta_D(\gamma + \beta_P)$, so that $\gamma\tau\Gamma_2(\sigma) = c_1(\sigma, \gamma)\tau + c_2(\gamma)$.

♣ **B.2(i):** If $\min\{\beta_P, \beta_D\} \geq 0$, then $\Gamma_1 > 0$ and $\Gamma_2(\sigma) > 0$ since $1 - \gamma\tau\sigma^2 \geq 0$. Therefore, it follows by algebraic manipulation that $\Gamma(\sigma)$ is nonempty and equal to

$$\Gamma(\sigma) = \left\{ (\gamma, \tau) \in \mathbb{R}^2 \mid \gamma \in (0, +\infty) \text{ and } \tau \in (0, 1/\gamma\|L\|^2] \right\}.$$

♣ **B.2(ii):** If $\min\{\beta_P, \beta_D\} < 0$ and $\max\{\beta_P, \beta_D\} \geq 0$, then either $\frac{1}{\gamma}\beta_P + \frac{1}{\tau}\beta_D \geq 0$, in which case by definition $\Gamma_1 > 0$, or $\frac{1}{\gamma}\beta_P + \frac{1}{\tau}\beta_D < 0$, in which case $\Gamma_1 > \Gamma_2(\sigma_d)$ since $1 - \gamma\tau\sigma_d^2 \geq 0$. Therefore, regardless of the sign of $\frac{1}{\gamma}\beta_P + \frac{1}{\tau}\beta_D$ it holds that $\Gamma(\sigma_d)$ reduces to $\left\{ (\gamma, \tau) \in \mathbb{R}_{++}^2 \mid \gamma\tau \in (0, 1/\|L\|^2] \text{ and } \Gamma_2(\sigma_d) > 0 \right\}$.

◇ $\beta_P \geq 0, \beta_D < 0$: Then, by construction $c_1(\sigma_d, \gamma) > 0$ and $c_2(\gamma) < 0$ for all $\gamma \in \mathbb{R}_+$. Consequently, $\Gamma_2(\sigma_d) > 0 \iff c_1(\sigma_d, \gamma)\tau + c_2(\gamma) > 0 \iff \tau > -c_2(\gamma)/c_1(\sigma_d, \gamma) = \tau_{\min}(\gamma)$ and the set $\Gamma(\sigma_d)$ is nonempty if for some $\gamma > 0$ it holds that

$$\tau_{\min}(\gamma) < 1/\gamma\|L\|^2 \iff c_2(\gamma)\gamma\|L\|^2 + c_1(\sigma_d, \gamma) = \beta_D\|L\|^2\gamma^2 + (1 + \beta_P\beta_D(\|L\|^2 - \sigma_d^2))\gamma + \beta_P > 0, \quad (\text{B.1})$$

which is guaranteed by [Fact B.1\(i\)](#). Therefore, it follows from [Fact B.1\(ii\)](#) that

$$\Gamma(\sigma_d) = \left\{ (\gamma, \tau) \in \mathbb{R}^2 \mid \gamma \in (0, \gamma_{\max}) \text{ and } \tau \in (\tau_{\min}(\gamma), 1/\gamma\|L\|^2] \right\}.$$

◇ $\beta_P < 0, \beta_D \geq 0$: Observe that for all $\gamma \leq -\beta_P/(1 - \beta_P\beta_D\sigma_d^2) < -\beta_P$ it holds that $c_1(\sigma_d, \gamma) \leq 0$ and $c_2(\gamma) < 0$. Therefore, if $\Gamma_2(\sigma_d) > 0$ then necessarily $\gamma > -\beta_P/(1 - \beta_P\beta_D\sigma_d^2)$, implying $c_1(\sigma_d, \gamma) > 0$, so that $\Gamma(\sigma_d)$ reduces to $\left\{ (\gamma, \tau) \in \mathbb{R}_{++}^2 \mid \gamma\tau \in (0, 1/\|L\|^2], \gamma > -\beta_P/(1 - \beta_P\beta_D\sigma_d^2) \text{ and } \tau > \tau_{\min}(\gamma) \right\}$. Ensuring that $\tau_{\min}(\gamma) < 1/\gamma\|L\|^2$, i.e., solving (B.1) as before, yields $\gamma \in (\gamma_{\min}, +\infty)$. Finally, by observing that $\gamma_{\min} > -\beta_P > -\beta_P/(1 - \beta_P\beta_D\sigma_d^2)$, it follows that

$$\Gamma(\sigma_d) = \left\{ (\gamma, \tau) \in \mathbb{R}^2 \mid \gamma \in (\gamma_{\min}, +\infty) \text{ and } \tau \in (\tau_{\min}(\gamma), 1/\gamma\|L\|^2] \right\}.$$

◆ **B.2(iii)**: If $\max\{\beta_P, \beta_D\} < 0$, then $\Gamma_1 > 0$ for $(\gamma, \tau) \in \mathbb{R}_{++}^2$ if and only if $\gamma > -\beta_P/2$ and $\tau > -\gamma\beta_D/(2\gamma+\beta_P)$. As a result, the set $\Gamma(\|L\|)$ is empty when $\beta_P\beta_D \geq 1/\|L\|^2$, as in this case the inequality $-\gamma\beta_D/(2\gamma+\beta_P) < 1/\gamma\|L\|^2$ does not have a positive solution for $\gamma > -\beta_P/2$. Consider the following cases, assuming that $\beta_P\beta_D < 1/\|L\|^2$.

◇ $\gamma \in (-\beta_P/2, -\beta_P/(1-\beta_P\beta_D\|L\|^2))$: Then, $c_1(\|L\|, \gamma) < 0$ and thus $\Gamma_2(\|L\|) > 0$ if and only if $\tau < \tau_{\min}(\gamma)$. Since it is easy to verify that in this case $-\gamma\beta_D/(2\gamma+\beta_P) > \tau_{\min}(\gamma)$, no such γ belong to the set $\Gamma(\|L\|)$.

◇ $\gamma = -\beta_P/(1-\beta_P\beta_D\|L\|^2)$: Then, $c_1(\|L\|, \gamma) = 0$ and $c_2(\gamma) < 0$, so that $\Gamma_2(\|L\|) < 0$ for all $\tau \in \mathbb{R}$.

◇ $\gamma > -\beta_P/(1-\beta_P\beta_D\|L\|^2)$: Then, $c_1(\|L\|, \gamma) > 0$ and thus $\Gamma_2(\|L\|) > 0$ if and only if $\tau > \tau_{\min}(\gamma)$, which is by construction larger than $-\gamma\beta_D/(2\gamma+\beta_P)$. The set $\Gamma(\sigma_d)$ is nonempty if for some $\gamma > 0$ it holds that $\tau_{\min}(\gamma) < 1/\gamma\|L\|^2$, which holds by **Fact B.1(i)** if and only if $[\beta_P]_-[\beta_D]_- < 1/4\|L\|^2$. Finally, noting that $\gamma_{\min} > -\beta_P/2$, it follows from **Fact B.1(ii)** that

$$\Gamma(\|L\|) = \{(\gamma, \tau) \in \mathbb{R}^2 \mid \gamma \in (\gamma_{\min}, \gamma_{\max}) \text{ and } \tau \in (\tau_{\min}(\gamma), 1/\gamma\|L\|^2)\}.$$

Since these sets corresponds to the stepsize ranges provided by **Table 1**, the proof is completed. \square

Lemma B.3. Let $D := [I_n \ I_n]$, $(Y_1, Y_2) \in \text{dom}_{\square}$ and define $Y = Y_1 \oplus Y_2$. Then, (4.1) holds and X^* as defined in (4.2) is equal to $Y_1 \square Y_2$.

Proof. Let $E = \frac{1}{2} \begin{bmatrix} 1 & \\ & -1 \end{bmatrix}$. Observe that $D^\dagger = \frac{1}{2} \begin{bmatrix} I_n & \\ & I_n \end{bmatrix}$ and $\Pi_{\mathcal{N}(D)} = \frac{1}{2} \begin{bmatrix} I_n & -I_n \\ & -I_n & I_n \end{bmatrix}$, so that

$$\begin{aligned} \Pi_{\mathcal{N}(D)} Y &= (E \otimes I_n) \begin{bmatrix} Y_1 & -Y_2 \end{bmatrix}, & \Pi_{\mathcal{N}(D)} Y \Pi_{\mathcal{N}(D)} &= (EE^\top) \otimes (Y_1 + Y_2), \\ \text{rank}(\Pi_{\mathcal{N}(D)} Y) &= \text{rank} \begin{bmatrix} Y_1 & Y_2 \end{bmatrix}, & \text{rank}(\Pi_{\mathcal{N}(D)} Y \Pi_{\mathcal{N}(D)}) &= \text{rank}(Y_1 + Y_2). \end{aligned}$$

Consequently, (4.1) holds owing to [35, Thm. 9.2.4], since $Y_1 + Y_2 \geq 0$ and Y_1 and Y_2 are parallel summable. The claim for X^* follows from **Proposition 4.4(ii)**, since

$$\begin{aligned} X^* &= \frac{1}{4}(Y_1 + Y_2) - \frac{1}{4}(Y_1 - Y_2)(E \otimes I_n)^\top \left((EE^\top) \otimes (Y_1 + Y_2) \right)^\dagger (E \otimes I_n)(Y_1 - Y_2) \\ &= \frac{1}{4}(Y_1 + Y_2) - \frac{1}{4}(Y_1 - Y_2)(Y_1 + Y_2)^\dagger (Y_1 - Y_2) \\ &= \frac{1}{4}(Y_1 + Y_2) - \frac{1}{4}(Y_1 + Y_2 - 2Y_2)(Y_1 + Y_2)^\dagger (Y_1 + Y_2 - 2Y_2) \\ &= \frac{1}{2}Y_2(Y_1 + Y_2)^\dagger (Y_1 + Y_2) + \frac{1}{2}(Y_1 + Y_2)(Y_1 + Y_2)^\dagger Y_2 - Y_2(Y_1 + Y_2)^\dagger Y_2 = Y_1 \square Y_2, \end{aligned}$$

where the second equality holds since for arbitrary matrices Z_1, Z_2 it holds that $(Z_1 \otimes Z_2)^\dagger = Z_1^\dagger \otimes Z_2^\dagger$ and the final equality holds by definition of the parallel sum and parallel summability. \square

Lemma B.4. Suppose that **Assumption III** holds and that the sets $\text{gph}(\Pi_{\mathcal{R}(L^\top)} A^{-1})$ and $\text{gph}(\Pi_{\mathcal{R}(L)} B)$ are not singletons. Then, it holds that $[\mu_A]_+[\rho_A]_+ \leq 1/4\sigma_d^2$ and $[\mu_B]_+[\rho_B]_+ \leq 1/4\sigma_d^2$.

Proof. Suppose that $\mu_A, \rho_A > 0$ and $\mu_B, \rho_B > 0$, for otherwise the two claims hold trivially. Consider $y_D \in \text{dom}(A^{-1} \circ (-L^\top)) \cap \text{dom}(B^{-1}) = \text{dom } T_D \neq \emptyset$ and let $y = -L^\top y_D$. By semimonotonicity of A at $(x^*, -L^\top y^*)$ it holds for all $(x^*, y^*) \in \mathcal{S}^*$, $x \in A^{-1}(y) = A^{-1} \circ (-L^\top)(y_D)$ that

$$\langle x - x^*, y + L^\top y^* \rangle \geq q_{\mu_A L^\top L}(x - x^*) + \rho_A \|y + L^\top y^*\|^2 \geq \mu_A \sigma_d^2 \|\Pi_{\mathcal{R}(L^\top)}(x - x^*)\|^2 + \rho_A \|y + L^\top y^*\|^2. \quad (\text{B.2})$$

Noting that by definition $(y, \Pi_{\mathcal{R}(L^\top)} x) \in \text{gph}(\Pi_{\mathcal{R}(L^\top)} A^{-1})$ and since $\text{gph}(\Pi_{\mathcal{R}(L^\top)} A^{-1})$ is not equal to the singleton $\{(-L^\top y^*, \Pi_{\mathcal{R}(L^\top)} x^*)\}$, both of the above involved norms are nonzero.

On the other hand, since $y = -L^\top y_D \in \mathcal{R}(L^\top)$, it holds by the Fenchel-Young inequality with modulus $2\mu_A \sigma_d^2 > 0$ that

$$\langle x - x^*, y + L^\top y^* \rangle = \langle \Pi_{\mathcal{R}(L^\top)}(x - x^*), y + L^\top y^* \rangle \leq \mu_A \sigma_d^2 \|\Pi_{\mathcal{R}(L^\top)}(x - x^*)\|^2 + \frac{1}{4\mu_A \sigma_d^2} \|y + L^\top y^*\|^2. \quad (\text{B.3})$$

Combining (B.2) and (B.3), it follows that $\rho_A \leq \frac{1}{4\mu_A \sigma_d^2}$.

Analogously, consider $x_P \in \text{dom}(A) \cap \text{dom}(B \circ L) = \text{dom } T_P \neq \emptyset$ and let $x = Lx_P$. Then, it holds for all $(x^*, y^*) \in \mathcal{S}^*$, $y \in B(x) = B(Lx_P)$ by the semimonotonicity assumption of B at (Lx^*, y^*) that

$$\langle x - Lx^*, y - y^* \rangle \geq \mu_B \|x - Lx^*\|^2 + q_{\rho_B LL^\top}(y - y^*) \geq \mu_B \|x - Lx^*\|^2 + \rho_B \sigma_d^2 \|\Pi_{\mathcal{R}(L)}(y - y^*)\|^2, \quad (\text{B.4})$$

where the involved norms are nonzero since $(x, \Pi_{\mathcal{R}(L)} y) \in \text{gph}(\Pi_{\mathcal{R}(L)} B)$ and $\text{gph}(\Pi_{\mathcal{R}(L)} B)$ is not equal to the singleton $\{Lx^*, \Pi_{\mathcal{R}(L)} y^*\}$. On the other hand, since $x = Lx_P \in \mathcal{R}(L)$, it holds by the Fenchel-Young inequality with modulus $2\mu_B > 0$ that

$$\langle x - Lx^*, y - y^* \rangle = \langle x - Lx^*, \Pi_{\mathcal{R}(L)}(y - y^*) \rangle \leq \mu_B \|x - Lx^*\|^2 + \frac{1}{4\mu_B} \|\Pi_{\mathcal{R}(L)}(y - y^*)\|^2. \quad (\text{B.5})$$

Finally, combining (B.4) and (B.5), it follows that $\rho_B \sigma_d^2 \leq \frac{1}{4\mu_B}$, establishing the claim. \square

Proposition B.5 (normal cone of a box). *The normal cone operator $N_C : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ of the n -dimensional box $C := \{x \in \mathbb{R}^n \mid l_i \leq x_i \leq u_i, i = 1, \dots, n\}$ is $(\text{diag}(\frac{|\tilde{v}_1|}{u_1 - l_1}, \dots, \frac{|\tilde{v}_n|}{u_n - l_n}), 0)$ -semimonotone at $(\tilde{x}, \tilde{v}) \in \text{gph } N_C$.*

Proof. By Proposition 4.2(iii) it suffices to show that N_{C_i} is $(\frac{|\tilde{v}_i|}{u_i - l_i}, 0)$ -semimonotone at $(\tilde{x}_i, \tilde{v}_i) \in \text{gph } N_{C_i}$. Using the fact that $|\tilde{x}_i - x_i| \leq u_i - l_i$ and monotonicity of N_{C_i} , we have for all $x_i \in C_i$ that $\frac{|\tilde{v}_i|}{u_i - l_i} |\tilde{x}_i - x_i|^2 \leq |\tilde{v}_i| |\tilde{x}_i - x_i| = \langle \tilde{v}_i, \tilde{x}_i - x_i \rangle$. \square

C Omitted proofs

Proof of Lemma 3.3 (oblique weak Minty for primal and dual operator). Note that $\text{gph } T_{\text{PD}}$ is equal to the set $\{(x_A, y_B), (y_A + L^\top y_B, x_B - Lx_A) \mid (x_A, y_A) \in \text{gph } A, (x_B, y_B) \in \text{gph } B\}$. Consequently, it holds by assumption for all $(x^*, y^*) \in \mathcal{S}^*$, $(x_A, y_A) \in \text{gph } A$ and $(x_B, y_B) \in \text{gph } B$ that

$$\langle y_A + L^\top y_B, x_A - x^* \rangle + \langle x_B - Lx_A, y_B - y^* \rangle \geq q_{V_P}(y_A + L^\top y_B) + q_{V_D}(x_B - Lx_A), \quad (\text{C.1})$$

◆ In (C.1), consider $x_A \in \text{dom}(A) \cap \text{dom}(B \circ L) = \text{dom } T_P \neq \emptyset$ and let $x_B = Lx_A$. Then, it holds for all $x^* \in \mathcal{S}_P^*$, $y_A \in A(x_A)$ and $y_B \in B(Lx_A)$ that

$$\langle y_A + L^\top y_B, x_A - x^* \rangle \geq q_{V_P}(y_A + L^\top y_B). \quad (\text{C.2})$$

Since $(x_A, y_A + L^\top y_B) \in \text{gph } T_P$ by construction and $\mathcal{S}_P^* \subseteq \text{zer } T_P$ by Proposition 3.1, it follows by definition that T_P has V_P -oblique weak Minty solutions at \mathcal{S}_P^* .

◆ Analogously, consider $y_B \in \text{dom}(A^{-1} \circ (-L^\top)) \cap \text{dom}(B^{-1}) = \text{dom } T_D \neq \emptyset$ and let $y_A = -L^\top y_B$ in (C.1). Then, it holds for all $y^* \in \mathcal{S}_D^*$, $x_A \in A^{-1} \circ (-L^\top)(y_B)$ and $x_B \in B^{-1}(y_B)$ that

$$\langle x_B - Lx_A, y_B - y^* \rangle \geq q_{V_D}(x_B - Lx_A). \quad (\text{C.3})$$

Since $(y_B, x_B - Lx_A) \in \text{gph } T_D$ by construction and $\mathcal{S}_D^* \subseteq \text{zer } T_D$ by Proposition 3.1, it follows by definition that T_D has V_D -oblique weak Minty solutions at \mathcal{S}_D^* , completing the proof \square

Proof of Example 3.6 (saddle point problem).

◆ 3.6(i): By defining $H := I + \lambda((P + T_{\text{PD}})^{-1}P - I)$ and substituting $\tau = \frac{1}{\gamma \ell^2}$, the update rule for z^k corresponds to the linear dynamical system $z^{k+1} = Hz^k$. Global asymptotic stability of this system is achieved if and only if the spectral radius of the matrix H is strictly less than one, which holds iff $\lambda \in (0, \min\{2, \bar{\lambda}\})$. Analogously, the convergence result for $(\Pi_{\mathcal{R}(P)} z^k)_{k \in \mathbb{N}}$ can be obtained by analyzing the spectral radius of $\Pi_{\mathcal{R}(P)} H$.

◆ 3.6(ii): The primal-dual operator and its inverse are given by

$$T_{\text{PD}} = \begin{bmatrix} A & L^\top \\ -L & B^{-1} \end{bmatrix} \quad \text{and} \quad T_{\text{PD}}^{-1} = \begin{bmatrix} (A + L^\top BL)^{-1} & -(A + L^\top BL)^{-1} L^\top B \\ BL(A + L^\top BL)^{-1} & B - BL(A + L^\top BL)^{-1} L^\top B \end{bmatrix}$$

owing to the Schur complement lemma. Therefore, when the vector v in (1.2) is restricted to $\mathcal{R}(P)$, Assumption II.A3 is equivalent to

$$z^\top \left(\frac{T_{\text{PD}} + T_{\text{PD}}^\top}{2} - T_{\text{PD}}^\top V T_{\text{PD}} \right) z \geq 0, \quad \text{for all } z \in \mathbb{R}^n : z \in T_{\text{PD}}^{-1} \mathcal{R}(P), \quad (\text{C.4})$$

where V is given by (3.3). Using that $L = Y\Sigma X^\top$, where $Y = I_3$, $\Sigma = \begin{bmatrix} \ell I_2 \\ 0 \end{bmatrix}$ and $X = \text{sgn}(\ell)I_2$, it follows from (3.22) that

$$U = \begin{bmatrix} \frac{1}{\sqrt{1+\gamma^2\ell^2}}I_2 & 0 \\ -\frac{\gamma\ell}{\sqrt{1+\gamma^2\ell^2}}I_2 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad V = \beta_P I_2 \oplus \beta_D I_2 \oplus \beta'_D I_1,$$

where U is an orthonormal basis for $\mathcal{R}(P)$. As a result, (C.4) is satisfied if and only if

$$\begin{aligned} (T_{\text{PD}}^{-1}U)^\top \left(\frac{T_{\text{PD}} + T_{\text{PD}}^\top}{2} - T_{\text{PD}}^\top V T_{\text{PD}} \right) T_{\text{PD}}^{-1}U \geq 0 &\iff \frac{b\ell^2(1+a^2\gamma^2) - (\beta_P + \gamma^2\ell^2\beta_D)(a^2+b^2\ell^4)}{(a^2+b^2\ell^4)(1+\gamma^2\ell^2)} I_2 \oplus (c - \beta'_D)I_1 \geq 0 \\ &\iff \frac{1}{\gamma}\beta_P + \gamma\ell^2\beta_D \leq \frac{b\ell^2(1+a^2\gamma)}{\gamma(a^2+b^2\ell^4)} \quad \text{and} \quad \beta'_D \leq c. \end{aligned}$$

Therefore, the upper bound on λ implied by Theorem 3.4 is given by

$$\lambda_{\max} := 2(1 + \min\{\eta, \eta'\}) = 2\left(1 + \min\left\{\frac{1}{\gamma}\beta_P + \gamma\ell^2\beta_D, \gamma\ell^2\beta'_D\right\}\right) \leq \min\left\{2 + 2\frac{b\ell^2(1+a^2\gamma)}{\gamma(a^2+b^2\ell^4)}, 2(1 + \gamma c\ell^2)\right\} = \bar{\lambda}. \quad (\text{C.5})$$

♣ 3.6(iii): For this particular instance $\text{tr } T_P = -2$ and $\text{tr } T_D = \text{tr } T_{\text{PD}} = -12$. Since the trace of a matrix equals the sum of its eigenvalues, the proof is completed. \square

Proof of Example 3.7 (influence of singular values).

♣ 3.7(i): By [6, Prop. 5.1(ii)] and using that A , B and L are symmetric, T_{PD} is $\frac{1}{2}$ -comonotone if and only if

$$\frac{T_{\text{PD}} + T_{\text{PD}}^\top}{2} - \frac{1}{2}T_{\text{PD}}^\top T_{\text{PD}} = \begin{bmatrix} A - \frac{1}{2}(A^\top A + L^\top L) & -\frac{1}{2}(AL - LB^{-1}) \\ -\frac{1}{2}(LA - B^{-1}L) & B^{-1} - \frac{1}{2}(B^{-1\top}B^{-1} + LL^\top) \end{bmatrix} \geq 0. \quad (\text{C.6})$$

Using that $B^{-1} = A$, that L is symmetric and that A and L commute, i.e., $AL = LA$, this condition reduces to $A - \frac{1}{2}(A^\top A + L^\top L) \geq 0$, which holds by definition of A and L . Noting that $\text{zer } T_{\text{PD}} = (0_n, 0_n)$, the claim is established.

♣ 3.7(ii): Follows from Theorem 3.4 and Proposition 3.5, using that $\|L\| = 1$ and $\gamma\tau = \frac{1}{\|L\|^2}$.

♣ 3.7(iii): Analogous to the setting of Example 3.6(i), the update rule for $\Pi_{\mathcal{R}(P)} z^k$ can be expressed as the linear dynamical system $\Pi_{\mathcal{R}(P)} z^{k+1} = \Pi_{\mathcal{R}(P)} H \Pi_{\mathcal{R}(P)} z^k$, where $H := I + \lambda((P + T_{\text{PD}})^{-1}P - I)$ and P is defined as in (1.1). This system is globally asymptotically stable if and only if the spectral radius of $\Pi_{\mathcal{R}(P)} H$ is strictly less than one, i.e., if and only if $\lambda \in (0, \bar{\lambda}_{\text{spectral}})$, where

$$\bar{\lambda}_{\text{spectral}} \in \arg \max_{\lambda} \lambda \quad \text{subject to} \quad \|\Pi_{\mathcal{R}(P)} H(\lambda)\|_2 < 1.$$

The values for $\bar{\lambda}_{\text{spectral}}$ reported in Figure 2 are obtained by solving this problem using SymPy. \square

Proof of Proposition 4.1. By the Fenchel-Young inequality, it holds for any $R > 0$ that

$$\langle x - x', y - y' \rangle \geq \frac{1}{4} q_{R^{-1}}(x - x') - q_R(y - y'), \quad \text{for all } (x, y), (x', y') \in \text{gph } A.$$

Therefore, (1.3) is satisfied for all $M \leq \frac{1}{4}R^{-1}$. \square

Proof of Proposition 4.2 (inverting, shifting, scaling and cartesian product).

♣ 4.2(ii): First, consider the assertion where A is semimonotone only at (x'_A, y'_A) . Define $s' = x'_A - u$ and $t' = y + \alpha y'_A$, such that $(s', t') \in \text{gph } T$. Then, it holds for all $(s, t) \in \text{gph } T$ that

$$\begin{aligned} \langle s - s', t - t' \rangle &= \alpha \langle (s + u) - (s' + u), \alpha^{-1}(t - y) - \alpha^{-1}(t' - y) \rangle \\ (\text{semimonotonicity of } A \text{ at } (x'_A, y'_A)) &\geq \alpha q_M((s + u) - (s' + u)) + \alpha q_R(\alpha^{-1}(t - y) - \alpha^{-1}(t' - y)) \\ &= q_{\alpha M}(s - s') + q_{\alpha^{-1}R}(t - t'), \end{aligned}$$

where we used that $(s' + u, \alpha^{-1}(t' - y)) = (x'_A, y'_A)$ and $(s + u, \alpha^{-1}(t - y)) \in \text{gph } A$. Hence, it follows that T is $(\alpha M_A, \alpha^{-1} R_A)$ -semimonotone at (s', t') .

If A is (M_A, R_A) -semimonotone at all $(x'_A, y'_A) \in \text{gph } A$, we then know that T is $(\alpha M_A, \alpha^{-1} R_A)$ -semimonotone at all points in the set $\{(x'_A - u, y + \alpha y'_A) \mid (x'_A, y'_A) \in \text{gph } A\}$. Since this set is equal to $\text{gph } T$, it follows that T is $(\alpha M_A, \alpha^{-1} R_A)$ -semimonotone (everywhere).

♣ **4.2(iii)**: Let A and B be semimonotone at respectively (x'_A, y'_A) and (x'_B, y'_B) . Since $\text{gph } T$ is equal to the set $\{(x_A, x_B), (y_A, y_B) \mid y_A \in Ax_A, y_B \in Bx_B\}$, it holds for all $(x, y) \in \text{gph } T$ that

$$\begin{aligned} \langle x - x', y - y' \rangle &= \langle x_A - x'_A, y_A - y'_A \rangle + \langle x_B - x'_B, y_B - y'_B \rangle \\ (\text{semimonotonicity of } A \text{ and } B) &\geq \mathbf{q}_{M_A}(x_A - x'_A) + \mathbf{q}_{R_A}(y_A - y'_A) + \mathbf{q}_{M_B}(x_B - x'_B) + \mathbf{q}_{R_B}(y_B - y'_B) \\ &= \mathbf{q}_{M_A \oplus M_B}(x - x') + \mathbf{q}_{R_A \oplus R_B}(y - y'), \end{aligned}$$

and thus T is $(M_A \oplus M_B, R_A \oplus R_B)$ -semimonotone at $(x', y') \in \text{gph } T$.

If A and B are semimonotone at all points in their graph, then T is $(M_A \oplus M_B, R_A \oplus R_B)$ -semimonotone at all points in $\text{gph } T$, which completes the proof. \square

Proof of Proposition 4.3 (linear operator). Owing to the linearity of D , (M, R) -semimonotonicity corresponds to having $\langle x, Dx \rangle \geq \mathbf{q}_M(x) + \mathbf{q}_R(Dx)$ for all $x \in \mathbb{R}^n$, which is equivalent to the claimed LMI. \square

Proof of Proposition 4.4 (symmetric solution of $D^\top X D \leq Y$).

♣ **4.4(i)**: First, note that the problem of finding an $X \in \mathbb{S}^n$ such that $D^\top X D \leq Y$ is equivalent to the problem of finding a pair $(X, Z) \in \mathbb{S}^n \times \mathbb{S}^m$ such that

$$D^\top X D = Y - Z \tag{C.7}$$

and $Z \geq 0$. Second, observe that by [40, Thm. 2], [23, Prop. 1] the involved linear matrix equality is solvable for $X \in \mathbb{S}^n$ if and only if $\mathcal{R}(Y - Z) \subseteq \mathcal{R}(D^\top)$, i.e., $\Pi_{\mathcal{N}(D)} Z = \Pi_{\mathcal{N}(D)} Y$. By [27, Thm. 2.2], a matrix $Z \geq 0$ satisfying this condition exists if and only if (4.1) holds, and the general solution is given by

$$Z = Y \Pi_{\mathcal{N}(D)} (\Pi_{\mathcal{N}(D)} Y \Pi_{\mathcal{N}(D)})^\dagger \Pi_{\mathcal{N}(D)} Y + D^\top (D^\dagger)^\top G D^\top (D^\dagger)^\top, \tag{C.8}$$

where $G \in \mathbb{S}^n$ is an arbitrary symmetric positive semidefinite matrix.

♣ **4.4(ii)**: Substituting (C.8) into (C.7) yields

$$D^\top X D = Y - Y \Pi_{\mathcal{N}(D)} (\Pi_{\mathcal{N}(D)} Y \Pi_{\mathcal{N}(D)})^\dagger \Pi_{\mathcal{N}(D)} Y - D^\top (D^\dagger)^\top G D^\top (D^\dagger)^\top,$$

of which the general solution is given by [23, Prop. 1]

$$X = \underbrace{(D^\dagger)^\top (Y - Y \Pi_{\mathcal{N}(D)} (\Pi_{\mathcal{N}(D)} Y \Pi_{\mathcal{N}(D)})^\dagger \Pi_{\mathcal{N}(D)} Y - G) D^\dagger}_{= X^* - (D^\dagger)^\top G D^\dagger} + H - \Pi_{\mathcal{R}(D)} H \Pi_{\mathcal{R}(D)}, \tag{C.9}$$

where $H \in \mathbb{S}^n$ is an arbitrary matrix. Substituting (C.9) into $D^\top X D \leq Y$ shows that

$$Y - D^\top X D = Y - D^\top X^* D + \Pi_{\mathcal{R}(D^\top)} G \Pi_{\mathcal{R}(D^\top)} \geq Y - D^\top X^* D \geq 0.$$

Finally, the alternative expression for X^* given in (4.2) follows directly from [53, Lem. 3].

♣ **4.4(iii)**: [7, Fact 6.4.38] \square

Proof of Corollary 4.6 (semimonotonicity of DTD^\top). If (4.1) holds, then it follows from Proposition 4.4 that $D^\top X^* D \leq Y$. Therefore, it only remains to be shown that this implies (DMD^\top, X^*) -semimonotonicity of DTD^\top [at (x', Dy')].

First, consider the case where T is semimonotone only at a single point $(D^\top x', y')$. Let $(D^\top x, y) \in \text{gph } T$ and denote $u = Dy$ and $u' = Dy'$. Then, $(x, u), (x', u') \in \text{gph } DTD^\top$ and it holds that

$$\langle x - x', u - u' \rangle = \langle D^\top (x - x'), y - y' \rangle$$

$$\begin{aligned}
(\text{semi. of } T \text{ at } (D^\top x', y')) &\geq \mathfrak{q}_M(D^\top(x - x')) + \mathfrak{q}_Y(y - y') \\
&\geq \mathfrak{q}_{DMD^\top}(x - x') + \mathfrak{q}_{X^*}(D(y - y')) = \mathfrak{q}_{DMD^\top}(x - x') + \mathfrak{q}_{X^*}(u - u'),
\end{aligned}$$

where $D^\top X^* D \leq Y$ was used in the second inequality, showing that DTD^\top is (DMD^\top, X^*) -semimonotone at (x', Dy') . Finally, if T is (M, Y) -semimonotone at all $(D^\top x', y') \in \text{gph } T$, then DTD^\top is (DMD^\top, X^*) -semimonotone at all points in $\{(x', Dy') \mid (D^\top x', y') \in \text{gph } T\}$, which equals $\text{gph } DTD^\top$. \square

Proof of Proposition 4.9 (sum and parallel sum). Let $D = [I_n \ 0]$. Then, $A + B$ is equal to DTD^\top , where $T := A \times B$. By Proposition 4.2(iii), operator T is $(M, R) = (M_A \oplus M_B, R_A \oplus R_B)$ -semimonotone [at $((x', x'), (y'_A, y'_B)) \in \text{gph } T$]. Consequently, it follows from Corollary 4.6 and Lemma B.3 that $DTD^\top = A + B$ is $(M_A + M_B, R_A \square R_B)$ -semimonotone [at $(x', y'_A + y'_B)$]. Finally, the claim for the parallel sum follows directly from those for the sum and Proposition 4.2(i), since $A \square B := (A^{-1} + B^{-1})^{-1}$. \square

Proof of Lemma 4.10 (sum with skew-symmetric matrix). First, consider the assertion where T is semimonotone at (x', y') . Let $(x, y) \in \text{gph } T$. Then, $(x, y + Dx), (x', y' + Dx') \in \text{gph } (T + D)$ and

$$\langle x - x', y - y' + D(x - x') \rangle \geq \mathfrak{q}_{D^\top MD}(x - x') + \mathfrak{q}_{R+R'}(y - y') + \langle x - x', D(x - x') \rangle.$$

By Corollary 4.5(ii) and skew-symmetry of D , it follows that D is $(-D^\top MD, \Pi_{\mathcal{R}(D)} M \Pi_{\mathcal{R}(D)})$ -semimonotone. Consequently,

$$\begin{aligned}
\langle x - x', y - y' + D(x - x') \rangle &\geq \mathfrak{q}_{R+R'}(y - y') + \mathfrak{q}_{\Pi_{\mathcal{R}(D)} M \Pi_{\mathcal{R}(D)}}(Dx - Dx') \\
&= \mathfrak{q}_{R'}(y - y' + D(x - x')) + \mathfrak{q}_R(y - y') + \mathfrak{q}_M(D(x - x')) \\
&\geq \mathfrak{q}_{R'+M \square R}(y - y' + D(x - x'))
\end{aligned}$$

where the fact that $\mathcal{R}(R') \subseteq \mathcal{N}(D)$ was used in the equality, and the final inequality follows from Proposition 4.4. Finally, when T is semimonotone, the claim follows analogously by considering all $(x', y') \in \text{gph } T$, completing the proof. \square

Proof of Corollary 5.4 (convergence of CPA under semimonotonicity). We begin by showing two intermediate results.

Claim 1: It holds that $\beta_P = \rho_A \square \rho_B \leq \rho_A$ and $\beta_D = \mu_A \square \mu_B \leq \mu_B$.

This claim follows by Assumption III and the definition of the parallel sum.

Claim 2: The graphs $\text{gph}(\Pi_{\mathcal{R}(L^\top)} \circ A^{-1})$ and $\text{gph}(\Pi_{\mathcal{R}(L)} \circ B)$ are not singletons.

By Assumption II.A2 the resolvents $J_{\gamma A}$ and $J_{\tau B^{-1}}$ have full domain. Consequently, $\gamma \text{id} + A^{-1}$ and $\tau \text{id} + B$ have full range [6, Prop. 2.11(ii)], which implies that $\mathcal{R}(\Pi_{\mathcal{R}(L^\top)} \circ (\gamma \text{id} + A^{-1})) = \mathcal{R}(L^\top)$ and $\mathcal{R}(\Pi_{\mathcal{R}(L)} \circ (\tau \text{id} + B)) = \mathcal{R}(L)$. Since $\mathcal{R}(\Pi_{\mathcal{R}(L^\top)} \circ (\gamma \text{id} + A^{-1}))$ is a singleton whenever $\text{gph}(\Pi_{\mathcal{R}(L^\top)} \circ A^{-1})$ is a singleton and $\mathcal{R}(\Pi_{\mathcal{R}(L)} \circ (\tau \text{id} + B))$ is a singleton whenever $\text{gph}(\Pi_{\mathcal{R}(L)} \circ B)$ is a singleton, the claim is established.

Next, we show that the parameters $\beta_P, \beta'_P, \beta_D$ and β'_D as defined in (5.3) satisfy condition (3.5) and thus ensure that Assumption II.A3 holds, see Corollary 5.3.

For the first inequality of (3.5), this follows directly from Assumption III(iv), since $[\beta_P]_- [\beta_D]_- = [\rho_A \square \rho_B]_- [\mu_A \square \mu_B]_- \leq 1/4\|L\|^2$. When $\beta'_P \geq 0$, the second inequality of (3.5) holds by construction. Let $\beta'_P = \rho_A < 0$. Then $\rho_B > -\rho_A > 0$ owing to Assumption III. If $\mu_A = \mu_B = 0 = \beta_D$, this inequality reduces to $\rho_A \geq \beta_P$, which holds by Claim 1. Otherwise, $\mu_A + \mu_B > 0$ and it reduces to

$$\rho_A - \rho_A(\rho_A \square \rho_B) \left([\mu_A \square \mu_B]_+ \sigma_d^2 + [\mu_A \square \mu_B]_- \|L\|^2 \right) \geq \rho_A \square \rho_B. \quad (\text{C.10})$$

Multiplying both sides of the inequality by $(\mu_A + \mu_B)(\rho_A + \rho_B)^{1/\rho_A^2} > 0$ and reordering, this condition reduces to $\mu_A + \mu_B - [\mu_A \mu_B]_+ \rho_B \sigma_d^2 - [\mu_A \mu_B]_- \rho_B \|L\|^2 \geq 0$, which is vacuously satisfied when $[\mu_A \mu_B]_+ = 0$. Otherwise, it holds that $\mu_A > 0$ and $\mu_B > 0$, so that it becomes $\mu_B + \mu_A(1 - \mu_B \rho_B \sigma_d^2) \geq 0$, which is satisfied owing to Lemma B.4 and Claim 2. The argument for the final inequality of (3.5) is analogous.

It is easy to verify that the stepsize intervals provided in Table 3 match those from Step size rule I when β_P and β_D are given by (5.3). Therefore, it only remains to verify that that Relaxation parameter rule II is equivalent to Relaxation parameter rule I. First, observe that the definition of η' from Table 4 is obtained by plugging in β'_P and β'_D from (5.3) into Table 2. Moreover, as a consequence of Claim 1, $\min\{\frac{1}{\gamma}\beta_P, \frac{1}{\tau}\beta_D\} \leq \min\{\frac{1}{\gamma}\rho_A, \frac{1}{\tau}\mu_B\} \leq \eta'$ and the following assertions hold.

(i) If $\max\{\beta_P, \beta_D\} \leq 0$, then $\frac{1}{\gamma}\beta_P + \frac{1}{\tau}\beta_D \leq \eta'$.

(ii) if $\beta_P\beta_D \geq 0$, then $\frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D - \theta_{\gamma\tau}(\sigma) \leq \eta'$ for any $\sigma \in (0, \|L\|]$, since

$$\theta_{\gamma\tau}(\sigma) = \sqrt{\left(\frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D\right)^2 - \frac{1}{\gamma\tau}\beta_P\beta_D(1 - \gamma\tau\sigma^2)} \geq \left|\frac{1}{2\gamma}\beta_P + \frac{1}{2\tau}\beta_D\right|.$$

The claimed equivalence follows immediately from these two assertions, completing the proof. \square

References

- [1] H. ATTOUCH AND M. THÉRA, *A general duality principle for the sum of two operators*, Journal of Convex Analysis, 3 (1996), pp. 1–24.
- [2] A. AUSLENDER AND M. TEBoulLE, *Asymptotic cones and functions in optimization and variational inequalities*, Springer Science & Business Media, Berlin, 2006.
- [3] J. K. BAKSALARY, F. PUKELSHEIM, AND G. P. STYAN, *Some properties of matrix partial orderings*, Linear Algebra and its Applications, 119 (1989), pp. 57–85.
- [4] V. BALAKRISHNAN AND L. VANDENBERGHE, *Semidefinite programming duality and linear time-invariant systems*, IEEE Transactions on Automatic Control, 48 (2003), pp. 30–41.
- [5] S. BANERT, M. UPADHYAYA, AND P. GISELSSON, *The Chambolle–Pock method converges weakly with $\theta > 1/2$ and $\tau\sigma\|l\|^2 < 4/(1 + 2\theta)$* , 2023, arXiv preprint arXiv:2309.03998.
- [6] H. H. BAUSCHKE, W. M. MOURSI, AND X. WANG, *Generalized monotone operators and their averaged resolvents*, Mathematical Programming, 189 (2021), pp. 55–74.
- [7] D. S. BERNSTEIN, *Matrix Mathematics*, Princeton University Press, Princeton, 2009.
- [8] R. I. BOŦ AND D.-K. NGUYEN, *The proximal alternating direction method of multipliers in the nonconvex setting: Convergence analysis and rates*, Mathematics of Operations Research, 45 (2020), pp. 682–712.
- [9] A. BÖHM, *Solving nonconvex-nonconcave min-max problems exhibiting weak Minty solutions*, 2022, arXiv preprint arXiv:2201.12247.
- [10] S. BOYD, L. EL GHAOU, E. FERON, AND V. BALAKRISHNAN, *Linear matrix inequalities in system and control theory*, SIAM, 1994.
- [11] K. BREDIES, E. CHENCHENE, D. A. LORENZ, AND E. NALDI, *Degenerate preconditioned proximal point algorithms*, SIAM Journal on Optimization, 32 (2022), pp. 2376–2401.
- [12] A. CHAMBOLLE AND T. POCK, *A first-order primal-dual algorithm for convex problems with applications to imaging*, Journal of Mathematical Imaging and Vision, 40 (2011), pp. 120–145.
- [13] P. L. COMBETTES AND T. PENNANEN, *Proximal methods for cohypomonotone operators*, SIAM Journal on Control and Optimization, 43 (2004), pp. 731–742.
- [14] L. CONDAT, *A primal–dual splitting method for convex optimization involving Lipschitzian, proximable and linear composite terms*, Journal of Optimization Theory and Applications, 158 (2013), pp. 460–479.
- [15] J. DIAKONIKOLAS, C. DASKALAKIS, AND M. I. JORDAN, *Efficient methods for structured nonconvex-nonconcave min-max optimization*, in International Conference on Artificial Intelligence and Statistics, 2021, pp. 2746–2754.
- [16] J. ECKSTEIN, *The Lions–Mercier splitting algorithm and the alternating direction method are instances of the proximal point algorithm*, tech. report, LIDS-P-1769, 1988.

- [17] J. ECKSTEIN AND D. P. BERTSEKAS, *On the Douglas–Rachford splitting method and the proximal point algorithm for maximal monotone operators*, *Mathematical Programming*, 55 (1992), pp. 293–318.
- [18] E. ESSER, X. ZHANG, AND T. F. CHAN, *A general framework for a class of first order primal-dual algorithms for convex optimization in imaging science*, *SIAM Journal on Imaging Sciences*, 3 (2010), pp. 1015–1046.
- [19] B. EVENS, P. PAS, P. LATAFAT, AND P. PATRINOS, *Convergence of the preconditioned proximal point method and Douglas-Rachford splitting in the absence of monotonicity*, 2023, arXiv preprint arXiv:2305.03605.
- [20] F. GIANNESI, *On Minty variational principle*, in *New Trends in Mathematical Programming: Homage to Steven Vajda*, Springer US, Boston, MA, 1998, pp. 93–99.
- [21] G. H. GOLUB AND C. F. VAN LOAN, *Matrix Computations*, JHU press, 2013.
- [22] E. GORBUNOV, A. TAYLOR, S. HORVÁTH, AND G. GIDEL, *Convergence of proximal point and extragradient-based methods beyond monotonicity: the case of negative comonotonicity*, in *International Conference on Machine Learning*, PMLR, 2023, pp. 11614–11641.
- [23] J. GROSZ, *A note on the general hermitian solution to $AXA^* = B$* , *Bulletin of the Malaysian Mathematical Sciences Society*, 21 (1998).
- [24] B. HE AND X. YUAN, *Convergence analysis of primal-dual algorithms for a saddle-point problem: From contraction perspective*, *SIAM Journal on Imaging Sciences*, 5 (2012), pp. 119–149.
- [25] A. HELMERSSON, *Methods for robust gain scheduling*, PhD thesis, Linköping University Electronic Press, 1995.
- [26] A. N. IUSEM, T. PENNANEN, AND B. F. SVAITER, *Inexact variants of the proximal point algorithm without monotonicity*, *SIAM Journal on Optimization*, 13 (2003), pp. 1080–1097.
- [27] C. G. KHATRI AND S. K. MITRA, *Hermitian and nonnegative definite solutions of linear matrix equations*, *SIAM Journal on Applied Mathematics*, 31 (1976), pp. 579–585.
- [28] I. V. KONNOV, *A class of combined iterative methods for solving variational inequalities*, *Journal of Optimization Theory and Applications*, 94 (1997), pp. 677–693.
- [29] P. LATAFAT AND P. PATRINOS, *Asymmetric forward–backward–adjoint splitting for solving monotone inclusions involving three operators*, *Computational Optimization and Applications*, 68 (2017), pp. 57–93.
- [30] G. LI AND T. PONG, *Global convergence of splitting methods for nonconvex composite optimization*, *SIAM Journal on Optimization*, 25 (2015), pp. 2434–2460.
- [31] Y. LI AND M. YAN, *On the improved conditions for some primal-dual algorithms*, 2022, arXiv preprint arXiv:2201.00139.
- [32] B. MARTINET, *Régularisation d’inéquations variationnelles par approximations successives*, *Revue française d’informatique et de recherche opérationnelle. Série rouge*, 4 (1970), pp. 154–158.
- [33] T. J. MEIJER, T. HOLICKI, S. J. A. M. VAN DEN EIJNDEN, C. W. SCHERER, AND W. P. M. H. HEEMELS, *The non-strict projection lemma*, 2023, arXiv preprint arXiv:2305.08735.
- [34] G. J. MINTY, *Monotone (nonlinear) operators in Hilbert space*, *Duke Mathematical Journal*, 29 (1962), pp. 341 – 346.
- [35] S. K. MITRA, P. BHIMASANKARAM, AND S. B. MALIK, *Matrix partial orders, shorted operators and applications*, vol. 10, World Scientific, 2010.
- [36] T. MÖLLENHOFF, E. STREKALOVSKIY, M. MOELLER, AND D. CREMERS, *The primal-dual hybrid gradient method for semiconvex splittings*, *SIAM Journal on Imaging Sciences*, 8 (2015), pp. 827–857.

- [37] D. O’CONNOR AND L. VANDENBERGHE, *On the equivalence of the primal-dual hybrid gradient method and Douglas–Rachford splitting*, *Mathematical Programming*, 179 (2020), pp. 85–108.
- [38] R. G. OTERO AND A. IUSEM, *Regularity results for semimonotone operators*, *Computational & Applied Mathematics*, 30 (2011), p. 13.
- [39] T. PENNANEN, *Local convergence of the proximal point algorithm and multiplier methods without monotonicity*, *Mathematics of Operations Research*, 27 (2002), pp. 170–191.
- [40] R. PENROSE, *A generalized inverse for matrices*, in *Mathematical proceedings of the Cambridge philosophical society*, vol. 51, Cambridge University Press, 1955, pp. 406–413.
- [41] T. PETHICK, O. FERCOQ, P. LATAFAT, P. PATRINOS, AND V. CEVHER, *Solving stochastic weak Minty variational inequalities without increasing batch size*, 2023, arXiv preprint arXiv:2302.09029.
- [42] T. PETHICK, P. LATAFAT, P. PATRINOS, O. FERCOQ, AND V. CEVHER, *Escaping limit cycles: Global convergence for constrained nonconvex-nonconcave minimax problems*, in *International Conference on Learning Representations*, 2021.
- [43] R. T. ROCKAFELLAR, *Augmented Lagrangians and applications of the proximal point algorithm in convex programming*, *Mathematics of Operations Research*, 1 (1976), pp. 97–116.
- [44] R. T. ROCKAFELLAR, *Monotone operators and the proximal point algorithm*, *SIAM Journal on Control and Optimization*, 14 (1976), pp. 877–898.
- [45] R. T. ROCKAFELLAR, *Progressive decoupling of linkages in optimization and variational inequalities with elicitable convexity or monotonicity*, *Set-Valued and Variational Analysis*, 27 (2019), pp. 863–893.
- [46] R. T. ROCKAFELLAR AND J. SUN, *Solving monotone stochastic variational inequalities and complementarity problems by progressive hedging*, *Mathematical Programming*, 174 (2019), pp. 453–471.
- [47] R. T. ROCKAFELLAR AND R. J.-B. WETS, *Variational Analysis*, Springer Science & Business Media, Berlin, 2009.
- [48] R. SHEFI AND M. TEBoulLE, *Rate of convergence analysis of decomposition methods based on the proximal method of multipliers for convex minimization*, *SIAM Journal on Optimization*, 24 (2014), pp. 269–297.
- [49] M. V. SOLODOV AND B. F. SVAITER, *A hybrid projection-proximal point algorithm.*, *Journal of Convex Analysis*, 6 (1999), pp. 59–70.
- [50] M. V. SOLODOV AND P. TSENG, *Modified projection-type methods for monotone variational inequalities*, *SIAM Journal on Control and Optimization*, 34 (1996), pp. 1814–1830.
- [51] Y. TIAN, *Equalities and inequalities for inertias of Hermitian matrices with applications*, *Linear algebra and its applications*, 433 (2010), pp. 263–296.
- [52] Y. TIAN, *Analytical solutions to some optimization problems on ranks and inertias of matrix-valued functions subject to linear matrix inequalities*, 2013, arXiv preprint arXiv:1301.0986.
- [53] C. YONGLIN AND Z. BINGJUN, *On g -inverses and the nonsingularity of a bordered matrix*, *Linear Algebra and its Applications*, 133 (1990), pp. 133–151.
- [54] M. ZHU AND T. CHAN, *An efficient primal-dual hybrid gradient algorithm for total variation image restoration*, *Ucla Cam Report*, 34 (2008), pp. 8–34.