

# ON THE INJECTIVITY OF EULER INTEGRAL TRANSFORMS WITH HYPERPLANES AND QUADRIC HYPERSURFACES

MATTIE JI

**ABSTRACT.** The Euler characteristic transform (ECT) is an integral transform used widely in topological data analysis. Previous efforts by Curry et al. and Ghrist et al. have independently shown that the ECT is injective on all compact definable sets. In this work, we first study the injectivity of the ECT on definable sets that are not necessarily compact and prove a complete classification of constructible functions that the Euler characteristic transform is not injective on. We then introduce the quadric Euler characteristic transform (QECT) as a natural generalization of the ECT by detecting definable shapes with quadric hypersurfaces rather than hyperplanes. We also discuss some criteria for the injectivity of QECT.

## 1. INTRODUCTION

The Euler characteristic transform (ECT) is an integral transform in topological data analysis (TDA) introduced in Turner et al. (2014). Since then, the ECT itself and its variants have been widely used in the applied science (Amézquita et al., 2021; Crawford et al., 2020; Wang et al., 2021; Nadimpalli et al., 2023; Hacquard and Lebovici, 2023; Roell and Rieck, 2023). On a high level, the ECT takes in a shape  $S$  in  $\mathbb{R}^n$ , “scans”  $S$  through each direction  $v \in \mathbb{S}^{n-1}$ , and keeps track of the Euler characteristics of the sublevel sets of  $S$ . Concretely, the Euler characteristic transform of  $S$  may be formalized into a function  $\text{ECT}(S) : \mathbb{S}^{n-1} \times \mathbb{R} \rightarrow \mathbb{Z}$  defined by

$$(\nu, t) \mapsto \text{ECT}(S)(\nu, t) = \chi(\{x \in S \mid x \cdot \nu \leq t\}),$$

where  $\chi(\bullet)$  denotes the combinatorial Euler characteristic (see Definition 2.6). Note that  $x \cdot \nu = t$  defines the equation of a hyperplane in  $\mathbb{R}^n$ .

In the conclusion of Curry et al. (2022), the authors posed the question of how the Euler characteristic transform (ECT) would behave on shapes cut out by quadratic equations rather than linear equations. Inspired by this question, we consider a “converse” of this question in this work - *what if we replace hyperplanes in the ECT with quadric hypersurfaces?*

The equation of a quadric hypersurface may be written as  $x^T A x + \nu \cdot x = t$ , where  $A$  is a symmetric  $n \times n$  real matrix and  $\nu$  is a vector in  $\mathbb{R}^n$ . Based on this notion, we can define the quadric Euler characteristic transform (QECT) of the shape  $S$  as a function given by  $(A, \nu, t) \mapsto \text{QECT}(S)(A, \nu, t) := \chi(\{x \in S \mid x^T A x + x \cdot \nu \leq t\})$ . This definition will be made more precise in Section 4. By extending the class of hyperplanes to quadric hypersurfaces, the hope is that the QECT would add an extra variable that takes into account of curvatures.

In this work, a central question we are interested in is the injectivity of the ECT and the QECT. Previous efforts by Curry et al. (2022) and Ghrist et al. (2018) have independently shown that the ECT is injective on a “reasonable” class of compact shapes. Furthermore, the work in Ghrist et al. (2018) showed that this injectivity result extends to finite sums of indicator functions on a collection of shapes (known as *constructible functions*) that are compactly supported.

We will first investigate the injectivity of the ECT on constructible functions that are not compactly supported. As we will see in Example 3.2 and Theorem 3.1 in Section 3, there are many pairs of constructible functions that the ECT is not injective on. We then extend the ECT to the QECT and discuss its injectivity in Theorem 4.7 and Theorem 4.8. Specifically, we will prove the following main results in this paper.

- (1) We completely classify all the pairs of constructible functions that the Euler characteristic transform is not injective on in Theorem 3.3.
- (2) Suppose  $v = 0$  is fixed, we show that the function  $S \mapsto \{(A, t) \mapsto \text{QECT}(S)(A, 0, t)\}$  is injective up to sign in Theorem 4.7.
- (3) Suppose the classes of “reasonable” shapes (see Definition 2.1) we are considering are all contained in  $B_R(0) := \{x \in \mathbb{R}^n \mid |x| \leq R\}$  for some  $R \geq 0$ . For a fixed  $A$  such that  $\|A\|_{op} < \frac{1}{1+2R^2}$ , we show that the function  $S \mapsto \{(v, t) \mapsto \text{QECT}(S)(A, v, t)\}$  is injective in Theorem 4.8. In particular, this serves as an interpolation between the injectivity of the ECT and Theorem 4.7 (see Remark 4.9).

These statements will be made more precise in their respective theorems.

**1.1. Outline.** The paper is organized as follows. In Section 2, we introduce the relevant backgrounds in o-minimal structures, Euler calculus, and the ECT. In Section 3, we discuss the injectivity of the ECT to all constructible functions, leading to a complete characterization of injectivity in Theorem 3.1. In Section 4, we extend the ECT to the QECT by considering quadric surfaces rather than hyperplanes in the sublevel sets of the integral transform and discuss several results on the injectivity of the QECT, leading to Theorem 4.7 and Theorem 4.8 in the end.

**Acknowledgements.** M.J. would like to thank Professor Kun Meng and Professor Richard Schwartz for their helpful comments and discussions. M.J. would also like to thank Cheng Chen for helpful conversations on functional analysis. M.J. would also like to thank Nir Elber and Riley Guyett for proofreading the paper and providing feedback and suggestions.

## 2. BACKGROUND

In this section, we will cover the necessary backgrounds in o-minimal structures, Euler calculus, and the ECT. We refer the reader to van den Dries (1998) for a comprehensive introduction to o-minimal structures, Curry et al. (2012) and Gusein-Zade (2010) for more details in Euler calculus, and Munch (2023) for a general review of the Euler characteristic transform.

**2.1. O-minimal structures.** O-minimal structures are widely used as the mathematical representation of a shape (Ghrist, 2014; Curry et al., 2022; Kirveslahti and Mukherjee, 2023; Meng et al., 2023) in applied topology and topological data analysis. Often in integral geometry, we want to consider shapes that have some level of “tameness” to avoid pathological examples, and o-minimal structures offer one way to capture the idea of “tameness”.

**Definition 2.1.** Let  $\mathcal{O}_n$  be a collection of subsets of  $\mathbb{R}^n$  and  $\mathcal{O} = \{\mathcal{O}_n\}_{n \geq 1}$ , we say that  $\mathcal{O}$  is an *o-minimal structure* if it satisfies the following seven axioms.

- (1)  $\mathcal{O}_n$  is a Boolean algebra.
- (2) If  $A \in \mathcal{O}_n$ , then  $A \times \mathbb{R} \in \mathcal{O}_{n+1}$  and  $\mathbb{R} \times A \in \mathcal{O}_{n+1}$ .
- (3)  $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i = x_j\} \in \mathcal{O}_n$  for  $1 \leq i < j \leq n$ .
- (4)  $\mathcal{O}$  is closed under axis-aligned projections.
- (5)  $\{r\} \in \mathcal{O}_1$  for all  $r \in \mathbb{R}$  and  $\{(x, y) \in \mathbb{R}^2 \mid x < y\} \in \mathcal{O}_2$ .
- (6)  $\mathcal{O}_1$  is exactly the finite unions of points and open intervals.
- (7)  $\mathcal{O}$  contains all real algebraic sets.

An element of  $\mathcal{O}_n$  is called a *definable set*.

In this paper, we will fix an arbitrary o-minimal structure  $\mathcal{O}_n$ . In particular, Definition 2.1 implies that any o-minimal structure has to contain all semialgebraic sets (see Remark 2.2 of Curry et al. (2022)). We also want a notion of “definability” for functions between definable sets.

**Definition 2.2.** Let  $f : X \rightarrow Y$  be a function between definable spaces.

- (1)  $f$  is called *definable* if its graph is a definable set.
- (2) If  $f$  is continuous definable with continuous definable inverse, then  $f$  is called a *definable homeomorphism*, and  $X$  and  $Y$  are said to be *definably homeomorphic*.
- (3) If  $f$  is an integer valued function, then  $f : X \rightarrow \mathbb{Z}$  is called a *constructible function*. Let  $\text{CF}(X)$  denote the space of constructible functions on  $X$ . Note that Definition 2.1(4) implies that the image of  $f$  is a discrete definable subset of  $\mathcal{O}_1$  and is thus finite.

When we define the quadric Euler characteristic transform later, we want to consider a suitable norm on the space of symmetric  $n \times n$  matrices. There are many choices of norms for matrices that are popular in machine learning, such as the Schatten norm, cut norms, and  $L_{p,q}$  norms, or the operator norm (see Fan et al. (2020)), so we need to consider norms that are compatible with our  $\mathcal{o}$ -minimal structure.

**Definition 2.3.** Let  $(V, \|\bullet\|)$  be a finite-dimensional normed real vector space.

- (1)  $\|\bullet\|$  is called a *definable norm* if the norm function  $\|\bullet\| : V \rightarrow \mathbb{R}_{\geq 0}$  is definable.
- (2)  $S^V := \{x \in V \mid \|x\| = 1\}$  is called the *unit sphere* with respect to  $\|\bullet\|$ . Note that  $S^V$  is a compact definable set when  $\|\bullet\|$  is a definable norm.
- (3) In particular, we will use  $|\bullet|$  to denote the usual  $\ell_2$  norm on  $\mathbb{R}^n$ , and  $\mathbb{S}^{n-1}$  to denote the usual unit sphere in  $\mathbb{R}^n$  with respect to the  $\ell_2$  norm.

**Example 2.4.** Here are some examples of definable norms that will be relevant to our discussions in Section 4.

- (1) Let  $|\bullet|$  be the  $\ell_2$  norm on  $\mathbb{R}^n$ , then the set

$$\{(x_1, \dots, x_n, y) \in \mathbb{R}^n \times \mathbb{R} \mid x_1^2 + \dots + x_n^2 = y^2 \text{ and } y \geq 0\}$$

is a semialgebraic set and is hence definable. This is the graph of  $|\bullet| : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ .

- (2) Let  $V$  be the vector space of  $n \times n$  symmetric real matrices and  $\|\bullet\|_{\text{op}}$  be the operator norm on  $V$ . The graph of  $\|\bullet\|_{\text{op}} : V \rightarrow \mathbb{R}_{\geq 0}$  may be realized as an axis-aligned projection of the following definable set

$$\{(A, \lambda_1, \dots, \lambda_n) \in V \times \mathbb{R}^n \mid \det(A - \lambda_i I) = 0, e_i(\lambda_1, \dots, \lambda_n) = \frac{a_i(A)}{a_n(A)} \text{ for all } i = 1, \dots, n \text{ and } \lambda_1^2 \leq \dots \leq \lambda_n^2\},$$

where  $e_i$  denotes the  $i$ -th elementary symmetric polynomial,  $a_i(A)$  denotes the  $i$ -th coefficient of the characteristic polynomial  $\det(A - xI)$ . Note that  $a_i$  is a polynomial function on the components of the matrix  $A$ . Thus, the operator norm is definable.

We also state the following technical lemma on  $\mathcal{o}$ -minimal structures that will be used later in the paper.

**Lemma 2.5** (Rephrased from Proposition 2.10 of Chapter 4 of van den Dries (1998)). *Let  $S \subseteq \mathbb{R}^{m+n}$  be a definable set. For any  $a \in \mathbb{R}^m$ , define  $S_a := \{x \in \mathbb{R}^n \mid (a, x) \in S\}$ . Then  $\chi(S_a)$  takes only finitely many values as  $a$  runs through  $\mathbb{R}^m$ , and for each integer  $e$  the set  $\{a \in \mathbb{R}^m \mid \chi(S_a) = e\}$  is definable.*

**2.2. Euler Calculus and the Euler Characteristic Transform.** Let  $S \subseteq \mathbb{R}^n$  be a definable set, the cell decomposition theorem (van den Dries, 1998, Chapter 3, Theorem 2.11) asserts that there is a disjoint partition of  $S$  into open-cells  $C_1, \dots, C_N$  such that each  $C_i$  is definably homeomorphic to  $\mathbb{R}^{a_i}$  for some  $a_i$ .

**Definition 2.6.** Choose  $S$  as above, the *Euler characteristic* of  $S$  is  $\chi(S) := \sum_{i=1}^N (-1)^{a_i}$ . This quantity is independent of the cell partition and is preserved under definable homeomorphisms (see Chapter 4 of van den Dries (1998)).

Euler calculus is an integral calculus based on the observation that the *Euler characteristics*  $\chi(\bullet)$  exhibits a finitely additive property similar to a signed measure:

$$\chi(A \cup B) = \chi(A) + \chi(B) - \chi(A \cap B).$$

The field seeks to develop a theory of integration for constructible functions, similar to how regular calculus developed a theory of integration for measurable functions.

**Definition 2.7.** Let  $X$  be a definable function and  $f : X \rightarrow \mathbb{Z}$  be a constructible function. The *Euler integral* of  $f$  is

$$\int_X f(x) d\chi(x) := \sum_{n=-\infty}^{\infty} n\chi(\{x \in X \mid f(x) = n\}).$$

Note that this quantity is well-defined by the discussions in Definition 2.2(3). The *Euler characteristic transform* of  $f$  is defined as

$$\text{ECT}(f) : \mathbb{S}^{n-1} \times \mathbb{R} \rightarrow \mathbb{Z}, (\nu, t) \mapsto \text{ECT}(f)(\nu, t) = \int_X f(x) \mathbb{1}_{X_t^\nu}(x) d\chi(x),$$

where  $X_t^\nu$  denotes the set  $\{x \in X \mid \nu \cdot x \leq t\}$ . For a definable subset  $S \subseteq X$ , We use  $\text{ECT}(S)$  to indicate the Euler characteristic transform of the indicator function on  $S$ .

Here is an example of computation with the Euler characteristic transform.

**Example 2.8.** Take  $B_1(0) = \{x \in \mathbb{R}^n \mid |x| \leq 1\}$  to be the closed unit ball with respect to the  $\ell_2$  norm. For any  $\nu \in \mathbb{S}^{n-1}$ , we have that  $\text{ECT}(B_1(0))(\nu, t) = 1$  if  $-1 \leq t$  and  $\text{ECT}(B_1(0))(\nu, t) = 0$  if  $t < -1$ .

Euler calculus also enjoys its version of Fubini's Theorem.

**Theorem 2.9** (Fubini's Theorem for Euler integrals). *Let  $f : X \rightarrow Y$  be a definable function between definable sets and  $h : X \rightarrow \mathbb{Z}$  be a constructible function, then*

$$\int_X h(x) d\chi(x) = \int_Y \left( \int_{f^{-1}(y)} h(x) d\chi(x) \right) d\chi(y).$$

A proof of Theorem 2.9 may be found in Theorem 4.5 of Curry et al. (2012). Note that while the authors assumed  $h$  to be compactly supported, the condition is not strictly required in the proof (see Page 5 of Curry et al. (2012)). Theorem 1 of Gusein-Zade (2010) presents an explicit proof of Theorem 2.9 for the case of semialgebraic sets without the assuming  $h$  to be compactly supported, and the case for a general  $o$ -minimal structure follows similarly.

For convenience, we will also briefly explain what a Radon transform is and how it relates to the Euler characteristic transform.

**Definition 2.10.** Let  $(X, Y)$  be a pair of definable sets and  $K \in \text{CF}(X \times Y)$  (known as a *kernel function*), then the *Radon transform* is a function  $R_K : \text{CF}(X) \rightarrow \text{CF}(Y)$  defined by

$$(R_K h)(y) = \int_X h(x) K(x, y) d\chi(x), h(x) \in \text{CF}(X), \text{ for all } y \in Y.$$

In particular, when  $Y = \mathbb{S}^{n-1} \times \mathbb{R}$  and  $K$  is the indicator function on  $\{(x, \nu, t) \in X \times Y \mid \nu \cdot x \leq t\}$ , then  $R_K$  is the ECT.

### 3. EULER CHARACTERISTIC TRANSFORM WITH HYPERPLANES

In Ghrist et al. (2018), the authors proved the following result on the injectivity of the Euler characteristic transform based on the Schapira inversion formula in Schapira (1995).

**Theorem 3.1** (Theorem 1 of Ghrist et al. (2018), Modified). *Let  $X = \mathbb{R}^n$ ,  $Y = \mathbb{S}^{n-1} \times \mathbb{R}$ ,  $K \in \text{CF}(X \times Y)$  be the indicator function on  $\{(x, \nu, t) \in X \times Y \mid \nu \cdot x \leq t\}$  and  $K' \in \text{CF}(Y \times X)$  be the indicator function on  $\{(\nu, t, x) \in Y \times X \mid \nu \cdot x \geq t\}$ , then for any  $h \in \text{CF}(X)$ , the following formula holds*

$$(R_{K'} \circ R_K)h = (\mu - \lambda)h + \lambda \left( \int_X h d\chi \right) \mathbb{1}_X, \quad (1)$$

where  $\mu = \chi(\mathbb{S}^{n-1})$  and  $\lambda = 1$ . Moreover, when restricted to the class of compactly supported functions on  $X$ ,  $R_K = \text{ECT}$  is injective.

In the original proof by the authors of Ghrist et al. (2018), this formula is only stated in the case where  $h$  is compactly supported. However, the formula still holds when  $h$  is not compactly supported. Please see the Appendix (Section 5) for a proof of Equation 1 without assuming that  $h$  is compactly supported. While the ECT is injective on compactly supported constructible functions, it is not injective on  $\text{CF}(\mathbb{R}^n)$ . We illustrate this with the following counter-example.

**Example 3.2.** Let  $X = \mathbb{R}^n$ ,  $S_1 = \mathbb{R}^n$ , and  $S_2 = \emptyset$ , then  $\text{ECT}(S_1)(\nu, t) = \text{ECT}(S_2)(\nu, t)$  for all  $(\nu, t) \in \mathbb{S}^n \times \mathbb{R}$ . Indeed, the set  $\{x \in \mathbb{R}^n \mid x \cdot \nu = t\}$  is definably homeomorphic to  $\mathbb{R}^{n-1}$ , and the set  $\{x \in \mathbb{R}^n \mid x \cdot \nu > t\}$  is definably homeomorphic to  $\mathbb{R}^n$ . Hence, the additivity of Euler characteristic implies that

$$\text{ECT}(S_1)(\nu, t) = \chi(\{x \in \mathbb{R}^n \mid x \cdot \nu \leq t\}) = \chi(\mathbb{R}^{n-1}) + \chi(\mathbb{R}^n) = 0.$$

Hence  $\text{ECT}(S_1)$  is the zero function. On the other hand, the Euler characteristic of the empty set is always zero, so  $\text{ECT}(S_2)$  is also the zero function.

Fortunately, we can classify how non-injective is the ECT with the following theorem. From the theorem, we will also obtain a corollary that shows Example 3.2 is the only such counter-example for the case of definable sets.

**Theorem 3.3.** *Let  $f, g : \mathbb{R}^n \rightarrow \mathbb{Z}$  be constructible functions, then  $\text{ECT}(f) = \text{ECT}(g)$  if and only if there exists some  $c \in \mathbb{Z}$  such that*

$$f(x) = \sum_{n=-\infty}^{+\infty} n \mathbb{1}_{\{f^{-1}(n)\}}(x) \text{ and } g(x) = \sum_{n=-\infty}^{+\infty} (n + c) \mathbb{1}_{\{f^{-1}(n)\}}(x).$$

*In particular, suppose  $f(x) = \mathbb{1}_{S_1}(x)$  and  $g(x) = \mathbb{1}_{S_2}(x)$  for distinct definable sets  $S_1, S_2 \subseteq \mathbb{R}^n$  and  $\text{ECT}(f) = \text{ECT}(g)$ , then  $S_1 = \mathbb{R}^n$  and  $S_2 = \emptyset$  up to renaming of variables.*

*Proof.* Suppose  $\text{ECT}(f) = \text{ECT}(g)$ , then Equation 1 implies that there exists integers  $\mu \neq \lambda$  such that

$$(\mu - \lambda)f(x) + \lambda \int_{\mathbb{R}^n} f(x) d\chi = (\mu - \lambda)g(x) + \lambda \int_{\mathbb{R}^n} g(x) d\chi,$$

for all  $x \in \mathbb{R}^n$ . Hence, the difference  $f(x) - g(x)$  is a constant integer, say  $c$ , and may be expressed as

$$g(x) - f(x) = c := \frac{\lambda}{\mu - \lambda} \left( \int_{\mathbb{R}^n} f(x) d\chi - \int_{\mathbb{R}^n} g(x) d\chi \right). \quad (2)$$

Since the images of constructible functions are finite, we can write  $f(x) = \sum_{i=1}^n a_i \mathbb{1}_{A_i}(x)$  such that  $A_i = f^{-1}(a_i)$  and  $a_i$  ranges through the image of  $f(x)$ . Similarly, we can write  $g(x) = \sum_{j=1}^m b_j \mathbb{1}_{B_j}(x)$  such that  $B_j = g^{-1}(b_j)$  and  $b_j$  ranges through the image of  $g(x)$ .

Let  $x \in A_i$ , then  $f(x) - g(x) = c$  by Equation 2. On the other hand  $f(x) = a_i$ , so  $g(x) = b_j = a_i + c$  for some  $b_j$  in the image of  $g(x)$ . Thus, the set function  $\{a_1, \dots, a_n\} \mapsto \{b_1, \dots, b_m\}$  by  $a_i \mapsto a_i + c$  is a well-defined

injective set function. Similarly, the set function  $\{b_1, \dots, b_m\} \mapsto \{a_1, \dots, a_n\}$  is also a well-defined inverse of the previous set function. Thus, we conclude that  $n = m$  and  $b_i = a_i + c$  for  $i = 1, \dots, n$  up to reordering.

Now for any  $x \in A_i$ ,  $g(x) = c + f(x) = c + a_i = b_i$ , so  $x \in B_i$ . Similarly for any  $x \in B_i$ ,  $f(x) = g(x) - c = a_i$ , thus  $x \in A_i$ . Hence  $A_i = B_i$ . Thus, we conclude that

$$f(x) = \sum_{i=1}^n a_i \mathbb{1}_{A_i}(x) \text{ and } g(x) = \sum_{i=1}^n (a_i + c) \mathbb{1}_{A_i}(x).$$

This concludes the proof of the “only if” direction.

Conversely, suppose  $f(x) = \sum_{n=-\infty}^{+\infty} n \mathbb{1}_{\{f^{-1}(n)\}}(x)$  and  $g(x) = \sum_{n=-\infty}^{+\infty} (n + c) \mathbb{1}_{\{f^{-1}(n)\}}(x)$ , then for any  $(\nu, t) \in \mathbb{S}^{n-1} \times \mathbb{R}$ , we will compute the difference of their respective Euler characteristic transforms.

$$\begin{aligned} \text{ECT}(g)(\nu, t) - \text{ECT}(f)(\nu, t) &= \sum_{n=-\infty}^{+\infty} \int_{\mathbb{R}^n} (n + c) \mathbb{1}_{\{f^{-1}(n)\} \cap \{x \cdot \nu \leq t\}}(x) d\chi - \int_{\mathbb{R}^n} n \mathbb{1}_{\{f^{-1}(n)\} \cap \{x \cdot \nu \leq t\}}(x) d\chi \\ &= \sum_{n=-\infty}^{+\infty} \int_{\mathbb{R}^n} (n + c - n) \mathbb{1}_{\{f^{-1}(n)\} \cap \{x \cdot \nu \leq t\}}(x) d\chi \\ &= c \sum_{n=-\infty}^{+\infty} \int_{\mathbb{R}^n} \mathbb{1}_{\{f^{-1}(n)\} \cap \{x \cdot \nu \leq t\}}(x) d\chi \\ &= c \int_{\mathbb{R}^n} \mathbb{1}_{\{x \cdot \nu \leq t\}}(x) d\chi \\ &= c \text{ECT}(\mathbb{R}^n)(\nu, t) \\ &= 0, \end{aligned}$$

where the fourth line follows from the fact that the sets  $\{f^{-1}(n)\}_{n \in \mathbb{Z}}$  form a finite (disregarding empty sets) partition of  $\mathbb{R}^n$ , and the sixth line follows from Example 3.2.

Finally, we will focus on the specific case that  $f(x) = \mathbb{1}_{S_1}(x)$  and  $g(x) = \mathbb{1}_{S_2}(x)$ . Without loss of generality, we will assume that  $S_1$  is non-empty. Since  $\text{ECT}(f) = \text{ECT}(g)$ , there exists some  $c \in \mathbb{Z}$  such that  $g(x) = 1 + c$  for all  $x \in S_1$  and  $g(x) = 0 + c$  for all  $x \notin S_1$ .

Since  $S_1$  is not empty, then let  $y$  be any point in  $S_1$ .  $f(y) = 1$  implies that  $g(y) = 1 + c$ . If  $g(y) = 1$ , then  $c = 0$  and  $g(x)$  becomes the indicator function on  $S_1$ , which is a contradiction to the assumption that  $f(x) \neq g(x)$ . If  $g(y) = 1 + c = 0$ , then it follows that  $c = -1$  and  $g(x) = -1$  for all  $x \notin S_1$ . Since  $g(x)$  takes values only between 0 and 1, this can occur only when  $S_1 = \mathbb{R}^n$ .

Thus,  $S_1 = \mathbb{R}^n$  and  $f(x)$  is the constant function with value 1. Since  $c = -1$ ,  $g(x)$  is the constant function with value 0, which implies that  $S_2$  is the empty set.  $\square$

#### 4. EULER CHARACTERISTIC TRANSFORM WITH QUADRIC HYPERSURFACES

In this section, we will define the quadric Euler characteristic transform and prove several injectivity results on this transform. Before going into the QECT specifically, we will first discuss some results on Radon transforms in Section 4.1 that will be useful in Section 4.2 (and the Appendix).

**4.1. Generalized Kernel Spaces.** Here is the general setup we will consider.

**Definition 4.1.** Let  $X \subseteq \mathbb{R}^n$  be a definable set,  $P \subseteq \mathbb{R}^k$  be a compact definable set (called the “parameter space”), and  $f : X \times P \rightarrow \mathbb{R}$  a definable function.

- (1) We define  $K_f(x, (\xi, t)) \in \text{CF}(X \times P \times \mathbb{R})$  as the indicator function on  $\{(x, (\xi, t)) \in X \times P \times \mathbb{R} \mid f(x, \xi) \leq t\}$  (the kernel function).
- (2) We define  $K'_f((\xi, t), x) \in \text{CF}(P \times \mathbb{R} \times X)$  as the indicator function on  $\{(\xi, t), x \in P \times \mathbb{R} \times X \mid f(x, \xi) \geq t\}$  (the dual kernel function).

- (3) We also define the fiber  $K_{x,f} = \{(\xi, t) \in P \times \mathbb{R} \mid f(x, \xi) \leq t\}$  and the dual fiber  $K'_{x',f} = \{(\xi, t) \in P \times \mathbb{R} \mid f(x', \xi) \geq t\}$ .

Given a constructible function  $h : X \rightarrow \mathbb{Z}$ , we are interested in what the function  $(R_{K'_f} \circ R_{K_f})h$  is to be able to prove injectivity results similar to that of Theorem 3.1. We first prove a technical lemma.

**Lemma 4.2.** *Let  $x, x' \in X$ .*

- (1)  $\chi(K_{x,f} \cap K'_{x',f}) = \chi(\{\xi \in P \mid f(x', \xi) - f(x, \xi) \geq 0\})$ .  
(2) *If  $x = x'$ , then  $\chi(K_{x,f} \cap K'_{x',f}) = \chi(P)$ .*  
(3) *As  $(x, x')$  ranges through  $X \times X$ , the function  $(x, x') \mapsto \chi(K_{x,f} \cap K'_{x',f})$  can only take on finitely many values  $c_1, \dots, c_n$ . Furthermore, the preimage  $S_i$  of each  $c_i$  is a definable subset of  $X \times X$ .*

*Proof.* For Lemma 4.2(1), we first rewrite the set  $K_{x,f} \cap K'_{x',f}$  as follows.

$$\begin{aligned} K_{x,f} \cap K'_{x',f} &= \{(\xi, t) \in P \times \mathbb{R} \mid t \geq f(x, \xi) \text{ and } f(x', \xi) \geq t\} \\ &= \{(\xi, t) \in P \times \mathbb{R} \mid f(x, \xi) \leq t \leq f(x', \xi)\} \\ &= \{(\xi, t) \in P \times \mathbb{R} \mid t \in [f(x, \xi), f(x', \xi)]\}. \end{aligned}$$

By considering the definable homeomorphism  $\varphi : P \times \mathbb{R} \rightarrow P \times \mathbb{R}$  by  $\varphi(\xi, t) = (\xi, t - f(x, \xi))$ , we have that

$$\begin{aligned} \chi(K_{x,f} \cap K'_{x',f}) &= \chi(\varphi(K_{x,f} \cap K'_{x',f})) \\ &= \chi(\{(\xi, t) \in P \times \mathbb{R} \mid t \in [0, f(x', \xi) - f(x, \xi)]\}). \end{aligned}$$

Since  $P$  is compact and definable, the set  $A := \{(\xi, t) \in P \times \mathbb{R} \mid t \in [0, f(x', \xi) - f(x, \xi)]\}$  is compact and definable. Define the straight-line homotopy  $H : A \times [0, 1] \rightarrow A$  as  $H((\xi, t), s) = (\xi, (1-s)t)$  for all  $((\xi, t), s) \in A \times [0, 1]$ , this produces a deformation retract of  $A$  onto the set  $\{\xi \in P \mid f(x', \xi) - f(x, \xi) \geq 0\}$ , which preserves the Euler characteristic because both sets are compact and definable. The proof of Lemma 4.2(1) is thus completed.

For Lemma 4.2(2),  $x = x'$  implies that  $f(x', p) - f(x, p) = 0$  for any  $p \in P$ . It then follows from Lemma 4.2(1) that  $\chi(K_{x,f} \cap K'_{x',f}) = \chi\{p \in P \mid 0 = 0\} = \chi(P)$ .

For Lemma 4.2(3), we implement Lemma 2.5 as follows. We define  $S$  as the definable set

$$S := \{(x, x', \xi) \in X \times X \times P \subseteq \mathbb{R}^{2n+k} \mid f(x', \xi) - f(x, \xi) \geq 0\}.$$

In this case,  $S_{(x,x')} = \{\xi \in \mathbb{R}^k \mid f(x', \xi) - f(x, \xi) \geq 0\}$ . Thus, Lemma 4.2(3) follows directly from Lemma 2.5.  $\square$

It is not generally true that for  $x \neq x' \in P$ , the value of  $\chi(K_{x,f} \cap K'_{x',f})$  remains constant, as will be shown in the proof of Theorem 4.7. However, we can still compute the function  $(R_{K'_f} \circ R_{K_f})h$ .

**Lemma 4.3.** *Following the context of Definition 4.1 and Lemma 4.2(3) and fix  $c_1 = \chi(P)$ , then for any  $x' \in X$ ,*

$$(R_{K'_f} \circ R_{K_f})h(x') = \chi(P) \int_X h(x) \mathbb{1}_{S_1}(x, x') d\chi(x) + \sum_{i=2}^n c_i \int_X h(x) \mathbb{1}_{S_i}(x, x') d\chi(x).$$

*In particular, if  $S_1 = \Delta$  is the diagonal of  $X \times X$ , then*

$$(R_{K'} \circ R_K)h(x') = \chi(P)h(x') + \sum_{i=2}^n c_i \int_X h(x) \mathbb{1}_{S_i}(x, x') d\chi(x).$$

*Proof.* By Lemma 4.2(3), we may write  $\chi(K_{x,f} \cap K'_{x',f}) = \sum_{i=1}^n c_i \mathbb{1}_{S_i}(x, x')$  as a function of  $x$  and  $x'$ . By Lemma 4.2, the diagonal  $\Delta \subseteq X \times X$  is contained in exactly one of the  $S_i$ , say  $S_1$ . Then

$$\begin{aligned}
(R_{K'_f} \circ R_{K_f})h(x') &= \int_{P \times \mathbb{R}} K'(y, x') \left[ \int_X h(x) K(x, y) d\chi(x) \right] d\chi(y) \\
&= \int_X h(x) \left[ \int_{P \times \mathbb{R}} K'_f(y, x') K_f(x, y) d\chi(y) \right] d\chi(x) \\
&= \int_X h(x) \chi(K_{x,f} \cap K'_{x',f}) \\
&= \int_X h(x) \sum_{i=1}^n c_i \mathbb{1}_{S_i}(x, x') d\chi(x) \\
&= c_1 \int_X h(x) \mathbb{1}_{S_1}(x, x') d\chi(x) + \sum_{i=2}^n c_i \int_X h(x) \mathbb{1}_{S_i}(x, x') d\chi(x) \\
&= \chi(P) \int_X h(x) \mathbb{1}_{S_1}(x, x') d\chi(x) + \sum_{i=2}^n c_i \int_X h(x) \mathbb{1}_{S_i}(x, x') d\chi(x),
\end{aligned}$$

where the second line follows from Theorem 2.9.  $\square$

**4.2. Quadric Euler Characteristic Transform.** Let  $V$  be the space of real  $n \times n$  symmetric matrices equipped with a definable norm  $\|\bullet\|_V$  (recall the notations in Definition 2.3). Recall that a general quadric surface is given by

$$x^T A x + v \cdot x = t$$

where  $A \in V$ ,  $v \in \mathbb{R}^n$ , and  $t \in \mathbb{R}$ . Then it would seem that a natural definition of QECT on a constructible function  $f : \mathbb{R}^n \rightarrow \mathbb{Z}$  would be

$$\text{QECT}(f) : S^V \times \mathbb{S}^{n-1} \times \mathbb{R} \rightarrow \mathbb{Z}, \text{QECT}(f)(A, v, t) = \int_X f(x) \mathbb{1}_{X_t^{A,v}}(x) d\chi(x),$$

where  $X_t^{A,v}$  denotes the set  $\{x^T A x + v \cdot x \leq t\}$ .

There is a question of whether our domain of choice is the best choice of domain. On one hand, QECT seems like a natural thematic generalization of the ECT. However, there are no choices of  $A \in S^V$  such that  $\|A\|_V = 0$ , so we cannot recover the ECT from this definition of the QECT.

The domain of  $\text{ECT}(f)$  is  $\mathbb{S}^{n-1} \times \mathbb{R}$  avoids the degenerate case of the zero vector. Thus, what our more general QECT wants is to consider the case where  $A$  and  $v$  are both not identically zero. This suggests that we should think of the norm of  $(A, v)$  as an element of  $V \times \mathbb{R}^n$ , which we will refer to as the space  $W$ . We define the norm  $\|(A, v)\|_W = \|A\|_V + |v|$  for all  $(A, v) \in W = V \times \mathbb{R}^n$ .

Thus, we adjust our definition to the following.

**Definition 4.4.** Let  $X \subseteq \mathbb{R}^n$  be a definable set and  $f : X \rightarrow \mathbb{Z}$  be a constructible function, the quadric Euler characteristic transform of  $f$  is the function  $\text{QECT}(f) : S^W \times \mathbb{R} \rightarrow \mathbb{Z}$  defined by

$$\text{QECT}(f)(A, v, t) = \int_X \left( f(x) \mathbb{1}_{X_t^{A,v}}(x) \right) d\chi(x).$$

**Example 4.5.** Let  $S_1 = \mathbb{R}^n$ ,  $S_2 = \emptyset$ , and  $I$  be the  $n \times n$  identity matrix, then  $\text{QECT}(S_1)(I, 0, t) \neq \text{QECT}(S_2)(I, 0, t)$ . Hence, the QECT can tell the difference between  $S_1$  and  $S_2$  compared to Example 3.2.

Now we will analyze a few properties of the QECT. First of all, when  $v = 0$  is fixed to be the zero vector, we note that the function  $f \mapsto \{(A, t) \mapsto \text{QECT}(f)(A, 0, t)\}$ , which we will refer to as  $\text{QECT}(-, 0, -) : \text{CF}(\mathbb{R}^n) \rightarrow \text{CF}(S^V \times \mathbb{R})$ , is not injective.

**Example 4.6.** Let  $p \in \mathbb{R}^n$  be the vector whose components are all unity,  $f(x) = \mathbb{1}_p(x)$ , and  $g(x) = \mathbb{1}_{-p}(x)$ , then  $\text{QECT}(f)(A, 0, t) = \text{QECT}(g)(A, 0, t)$  for all  $(A, t) \in S^V \times \mathbb{R}$ .

However, we can see that the failure to detect signs in Example 4.6 is the only such locus of non-injectivity with the following theorem.

**Theorem 4.7.** *Let  $v = 0$  be fixed, the function  $\text{QECT}(-, 0, -) : \text{CF}(\mathbb{R}^n) \rightarrow \text{CF}(S^V \times \mathbb{R})$  is “injective up to sign”. More precisely, let  $h : X \rightarrow \mathbb{R}^n$ , we obtain an inversion formula reminiscent of the Schapira inversion formula,*

$$(R_{K'_f} \circ R_{K_f} h)(x') = (\mu - \lambda) \sum_{z \in \{+x', -x'\}} h(z) + \lambda \left( \int_X h d\chi \right) \mathbb{1}_X,$$

where  $\mu$  and  $\lambda$  are distinct integers.

*Proof.* Since  $v = 0$ ,  $\|A\|_V = 1 - |v| = 1$ . Thus, the function  $(A, t) \mapsto \text{QECT}(f)(A, 0, t)$  has domain  $S^V \times \mathbb{R}$ . Following the setup of Definition 4.1, we choose  $X = \mathbb{R}^n$ ,  $P = S^V$ , and  $f : X \times P \rightarrow \mathbb{R}$  to be the function  $(x, A) \mapsto x^T A x$ . By Lemma 4.2 and the property that  $A$  is symmetric,  $\chi(K_{x,f} \cap K'_{x',f}) = \chi(\{A \in P \mid (x')^T A(x') - (x)^T A(x) \geq 0\}) = \chi(\{A \in P \mid (x' + x)^T A(x' - x) \geq 0\})$ .

If  $x = x'$  or  $x = -x'$ , then  $(x' + x)^T A(x' - x) = 0$  so  $\chi(K_{x,f} \cap K'_{x',f}) = \chi(S^V)$ .

Otherwise, suppose  $x \notin \{\pm x'\}$ , then consider the function  $\varphi : V \rightarrow \mathbb{R}$  given by  $\varphi(A) = (x' + x)^T A(x' - x)$ . Since  $(x + x')$  and  $(x' - x)$  are both non-zero vectors,  $\varphi$  is a surjective linear transformation, and hence  $\ker(\varphi)$  is a codimension 1 linear subspace of  $V$ .  $\ker(\varphi)$  inherits a natural norm from  $V$  and hence  $S^V \cap \ker(\varphi) = S^{\ker(\varphi)}$ .

Since all norms on a finite-dimensional real vector space are equivalent, the map  $\psi : S^V \rightarrow \mathbb{S}^{\dim V - 1}$  by  $\psi(x) = \frac{x}{|x|}$  is a definable homeomorphism whose restriction to  $S^{\ker(\varphi)}$  gives a homeomorphism between  $S^{\ker(\varphi)}$  and the  $\ell_2$  unit sphere of  $\ker(\varphi)$ .

By Alexander duality,  $\tilde{H}^0(S^V \setminus S^{\ker(\varphi)}) \cong \tilde{H}^{\dim V - 1 - 1}(S^{\ker(\varphi)}) \cong \mathbb{Z}$ , so  $S^V \setminus S^{\ker(\varphi)}$  has two connected components. Since  $\varphi$  is an odd function, we will denote the two connected components as  $S^V_+$  and  $S^V_-$  corresponding to the locus where  $\varphi$  is positive and negative respectively. Hence, the map  $\psi : S^V \rightarrow \mathbb{S}^{\dim V - 1}$  brings  $\ker(f) \cup S^V_+$  homeomorphically to a closed hemisphere of  $\mathbb{S}^{\dim V - 1}$ . Thus, the set  $\ker(f) \cup S^V_+$  is compact and contractible, so we conclude that  $\chi(K_{x,f} \cap K'_{x',f}) = 1$ .

Let  $\mu = \chi(P)$  and  $\lambda = 1$ , by Lemma 4.3, we can write

$$(R_{K'_f} \circ R_{K_f} h)(x') = \mu \int_{\mathbb{R}^n} h(x) \mathbb{1}_{\pm\Delta}(x, x') d\chi(x) + \lambda \int_{\mathbb{R}^n} h(x) \mathbb{1}_{\{X \times X - \pm\Delta\}}(x, x') d\chi(x),$$

where  $\mathbb{1}_{\pm\Delta}(x, x') = 1$  if  $x \in \{\pm x'\}$  and is 0 otherwise. Finally, we can furthermore simplify the expression as

$$\begin{aligned} (R_{K'_f} \circ R_{K_f} h)(x') &= \mu \int_{\{\pm x'\}} h(x) d\chi(x) + \lambda \int_{\mathbb{R}^n - \{\pm x'\}} h(x) d\chi(x) \\ &= (\mu - \lambda) \int_{\{\pm x'\}} h(x) d\chi(x) + \lambda \int_{\mathbb{R}^n} h(x) d\chi(x) \\ &= (\mu - \lambda) \sum_{z \in \{\pm x'\}} h(z) + \lambda \int_{\mathbb{R}^n} h(x) d\chi(x). \end{aligned}$$

This concludes the proof of Theorem 4.7. □

Theorem 4.7 examined what happens to the QECT when its vector component is fixed. We are also interested in what happens to the QECT when its matrix component is fixed.

**Theorem 4.8.** *Let  $B_R(0)$  denote the closed ball of radius  $R$  for some  $R \geq 0$ . Fix the norm on  $V$  to be the operator norm  $\|\bullet\|_{op}$ , and let  $A \in V$  be fixed such that  $\|A\|_{op} < \frac{1}{1+2R^2}$ . Define the function  $\text{QECT}(A, -, -) : \text{CF}(B_R(0)) \rightarrow \text{CF}(P \times \mathbb{R})$  by  $f \mapsto \{(v, t) \mapsto \text{QECT}(f)(A, v, t)\}$ , where  $P$  is the sphere of radius  $1 - \|A\|_{op}$  in  $\mathbb{R}^n$ . Then,  $\text{QECT}(A, -, -)$  is injective.*

**Remark 4.9.** Theorem 4.8 interpolates between Theorem 4.7 and Theorem 3.1 in the following sense.

- (1) When  $\|A\|_{op} = 1$ , the inequality  $1 < \frac{1}{1+2R^2}$  does not hold for any value of  $R$ . This is reflective of the fact that  $(A, t) \mapsto \text{QECT}(A, 0, t)$  is only injective up to signs in Theorem 4.7 and Example 4.6.
- (2) When  $\|A\|_{op} = 0$ ,  $A$  is the zero matrix and the QECT becomes the usual ECT, which is injective by Theorem 3.1 no matter what  $R$  is. This reflects the fact that the inequality  $0 < \frac{1}{1+2R^2}$  is satisfied for any value of  $R$ .

The requirement for the norm on  $V$  to be the operator norm is not strictly necessary. The same statement holds for definable norm  $\|\bullet\|_V$  on  $V$  that satisfies the property  $\|x^T Ax\|_V \leq \|A\|_V |x|^2$ . Common examples include the Frobenius norm and the nuclear norm. Furthermore, by adjusting the constant  $\frac{1}{1+2R^2}$  appropriately, similar statements for any definable norm on  $V$  will hold.

Now we will prove Theorem 4.8.

*Proof of Theorem 4.8.* Following the setup of Definition 4.1, we choose  $X = B_R(0)$ ,  $P = \{v \in \mathbb{R}^n \mid |v| = 1 - \|A\|_{op}\}$ , and  $f : X \times P \rightarrow \mathbb{R}$  to be the function  $(x, v) \mapsto x^T Ax + v \cdot x$ . By Lemma 4.2 and the property that  $A$  is symmetric,  $\chi(K_{x,f} \cap K'_{x',f}) = \chi(\{v \in P \mid (x')^T A(x') + v \cdot (x') - (x)^T A(x) - v \cdot x \geq 0\}) = \chi(\{v \in P \mid (x' + x)^T A(x' - x) + v \cdot (x' - x) \geq 0\})$ .

If  $x = x'$ , then Lemma 4.2 tells us that  $\chi(K_{x,f} \cap K'_{x',f}) = \chi(P)$ . Otherwise, if  $x \neq x'$ , we can consider the function  $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$  by  $\varphi(v) = (x' + x)^T A(x' - x) + v \cdot (x' - x)$ . Since  $x' - x$  is not the zero vector,  $\varphi$  is a surjective affine map and  $\ker(\varphi)$  is a hyperplane in  $\mathbb{R}^n$ . Furthermore, the vector  $\frac{(x' - x)}{|x - x'|}$  is a unit normal vector to  $\ker(\varphi)$ , and  $\ker(\varphi)$  may be written as the sum

$$\ker\{v \mapsto v \cdot (x' - x)\} - \frac{(x' + x)^T A(x' - x)}{|x - x'|}(x' - x).$$

This is because for any  $v - \frac{(x' + x)^T A(x' - x)}{|x - x'|}(x - x')$  in the sum above,

$$\begin{aligned} \varphi\left(v - \frac{(x' + x)^T A(x' - x)}{|x - x'|}(x' - x)\right) &= (x' + x)^T A(x' - x) + \left[v - \frac{(x' + x)^T A(x' - x)}{|x - x'|}(x - x')\right] \cdot (x' - x) \\ &= (x' + x)^T A(x' - x) + 0 - \frac{(x' + x)^T A(x' - x)}{|x - x'|}|x' - x| \\ &= (x' + x)^T A(x' - x) - (x' + x)^T A(x' - x) \\ &= 0. \end{aligned}$$

Now we observe that

$$\begin{aligned} |((x' + x)^T A(x' - x))| &= |(x')^T A(x') - (x)^T A(x)| \\ &\leq |x^T Ax| + |(x')^T A(x')| \\ &\leq \|A\|_{op}|x|^2 + \|A\|_{op}|x'|^2 \\ &\leq \|A\|_{op}(2R^2) \\ &< \frac{2R^2}{1 + 2R^2} \\ &= 1 - \frac{1}{1 + 2R^2} \\ &< 1 - \|A\|_{op}, \end{aligned}$$

where the fifth and last line both follow from the assumption  $\|A\|_{op} < \frac{1}{1+2R^2}$ . The radius of  $P$  is  $1 - \|A\|_{op}$ , so  $|(x + x')^T A(x - x')| < 1 - \|A\|_{op}$  implies that  $\{v \in P \mid \varphi(v) \geq 0\}$  is a compact and contractible subset of  $P$ . Thus,  $\chi(K_{x,f} \cap K'_{x',f}) = \chi(\{v \in P \mid \varphi(v) \geq 0\}) = 1$ .

Thus, by Lemma 4.3, for all  $x' \in \mathbb{R}^n$ ,

$$\begin{aligned}
R_{K'_f} \circ R_{K_f} h &= [1 + (-1)^{n-1}]h(x') + \int_{\mathbb{R}^n} h(x) \mathbb{1}_{\{(x,x') \notin \Delta\}} d\chi(x) \\
&= [1 + (-1)^{n-1}]h(x') + \int_{\mathbb{R}^n - \{x'\}} h(x) d\chi(x) \\
&= [1 + (-1)^{n-1}]h(x') + \int_{\mathbb{R}^n} h(x) d\chi(x) - \int_{\{x'\}} h(x) d\chi(x) \\
&= [(-1)^{n-1}]h(x') + \int_{\mathbb{R}^n} h(x) d\chi(x).
\end{aligned}$$

Since  $h(x) \in \text{CF}(B_R(0))$  has compact support, QECT would determine the value of  $\int_{\mathbb{R}^n} h(x) d\chi(x)$ . Thus, the function  $\text{QECT}(A, -, -) : \text{CF}(B_R(0)) \rightarrow \text{CF}(\mathbb{S}^{n-1} \times \mathbb{R})$  is injective.  $\square$

Finally, we will also discuss some auxiliary properties of the QECT.

**Proposition 4.10.** *Let  $f : \mathbb{R}^n \rightarrow \mathbb{Z}$  be a constructible function, then*

- (1) *QECT( $f$ ) takes only finitely many values as  $(A, \nu, t)$  runs through  $S^W \times \mathbb{R}$ .*
- (2) *For a fixed  $(A, \nu) \in S^W$ , the function  $t \mapsto \text{QECT}(f)(A, \nu, t)$  is right continuous.*

*Proof.* Proposition 4.10(1) is a direct application of Lemma 2.5 whose proof is similar to that of Lemma 4.2(3). Thus, the proof is omitted here. Proposition 4.10(2) is a direct application of Theorem 3.1 of Ji et al. (2023).  $\square$

## 5. APPENDIX

**5.1. Inversion Formula of ECT without Compact Support.** Here we reprove the inversion formula of ECT in Ghrist et al. (2018) without the assumption that  $h : \mathbb{R}^n \rightarrow \mathbb{Z}$  is compactly supported. The proof will be similar to that of Ghrist et al. (2018).

*Proof of Equation 1.* Following the setup of Definition 4.1, we choose  $X = \mathbb{R}^n$ ,  $P = \mathbb{S}^{n-1}$ , and  $f : X \times P \rightarrow \mathbb{R}$  to be the function  $(x, \nu) \mapsto x \cdot \nu$ . By Lemma 4.2, we can compute that

$$\begin{aligned}
\chi(K_{x,f} \cap K'_{x'}, f) &= \chi(\{\nu \in \mathbb{S}^{n-1} \mid f(x', \nu) - f(x, \nu) \geq 0\}) \\
&= \chi(\{\nu \in \mathbb{S}^{n-1} \mid (x' - x) \cdot \nu \geq 0\}) \\
&= \begin{cases} \chi(\mathbb{S}^{n-1}), & x = x' \\ \chi(\mathbb{S}_+^{n-1}), & x \neq x' \end{cases} \\
&= \begin{cases} 1 + (-1)^{n-1}, & x = x' \\ 1, & x \neq x' \end{cases},
\end{aligned}$$

where  $\mathbb{S}_+^{n-1}$  denotes the closed upper hemisphere of  $\mathbb{S}^{n-1}$ . By Lemma 4.3, for all  $x' \in \mathbb{R}^n$ ,

$$\begin{aligned}
(R_{K'} \circ R_K)h &= [1 + (-1)^{n-1}]h(x') + \int_{\mathbb{R}^n} h(x) \mathbb{1}_{\{(x,x') \notin \Delta\}} d\chi(x) \\
&= [1 + (-1)^{n-1}]h(x') + \int_{\mathbb{R}^n} h(x) d\chi(x) - \int_{\{x'\}} h(x) d\chi(x) \\
&= [(-1)^{n-1}]h(x') + \int_{\mathbb{R}^n} h(x) d\chi(x).
\end{aligned}$$

The proof of Equation 1 is completed.  $\square$

## REFERENCES

- E. J. Amézquita, M. Y. Quigley, T. Ophelders, J. B. Landis, D. Koenig, E. Munch, and D. H. Chitwood. Measuring hidden phenotype: quantifying the shape of barley seeds using the Euler characteristic transform. *in silico Plants*, 4(1):diab033, 12 2021. ISSN 2517-5025.
- L. Crawford, A. Monod, A. X. Chen, S. Mukherjee, and R. Rabadán. Predicting Clinical Outcomes in Glioblastoma: An Application of Topological and Functional Data Analysis. *Journal of the American Statistical Association*, 115(531):1139–1150, 2020.
- J. Curry, R. Ghrist, and M. Robinson. Euler calculus with applications to signals and sensing, 2012.
- J. Curry, S. Mukherjee, and K. Turner. How Many Directions Determine a Shape and Other Sufficiency Results for Two Topological Transforms. *Transactions of the American Mathematical Society, Series B*, 9(32):1006–1043, 2022.
- J. Fan, R. Li, C.-H. Zhang, and H. Zou. *Statistical Foundation of Data Science*. CRC Press, Taylor et Francis Group, 2020.
- R. Ghrist, R. Levanger, and H. Mai. Persistent Homology and Euler Integral Transforms. *Journal of Applied and Computational Topology*, 2:55–60, 2018.
- R. W. Ghrist. *Elementary Applied Topology*, volume 1. Createspace Seattle, 2014.
- S. M. Gusein-Zade. Integration with respect to the euler characteristic and its applications. *Russian Mathematical Surveys*, 65(3):399, 2010.
- O. Hacquard and V. Lebovici. Euler Characteristic Tools For Topological Data Analysis, 2023.
- M. Ji, K. Meng, and K. Ding. Euler characteristics and homotopy types of definable sublevel sets, with applications to topological data analysis, 2023.
- H. Kirveslahti and S. Mukherjee. Representing Fields without Correspondences: the Lifted Euler Characteristic Transform. *Journal of Applied and Computational Topology*, pages 1–34, 2023.
- K. Meng, M. Ji, J. Wang, K. Ding, H. Kirveslahti, A. Eloyan, and L. Crawford. Statistical Inference on Grayscale Images via the Euler-Radon Transform. *arXiv preprint arXiv:2308.14249v1*, 2023.
- E. Munch. An Invitation to the Euler Characteristic Transform. *arXiv preprint arXiv:2310.10395*, 2023.
- K. V. Nadimpalli, A. Chattopadhyay, and B. Rieck. Euler characteristic transform based topological loss for reconstructing 3d images from single 2d slices. In *2023 IEEE/CVF Conference on Computer Vision and Pattern Recognition Workshops (CVPRW)*, pages 571–579, Los Alamitos, CA, USA, jun 2023. IEEE Computer Society.
- E. Roell and B. Rieck. Differentiable euler characteristic transforms for shape classification, 2023.
- P. Schapira. Tomography of Constructible Functions. In *International Symposium on Applied Algebra, Algebraic Algorithms, and Error-Correcting Codes*, pages 427–435. Springer, 1995.
- K. Turner, Y. Mileyko, S. Mukherjee, and J. Harer. Fréchet Means for Distributions of Persistence Diagrams. *Discrete & Computational Geometry*, 52(1):44–70, 2014.
- L. van den Dries. *Tame Topology and O-minimal Structures*, volume 248. Cambridge university press, 1998.
- B. Wang, T. Sudijono, H. Kirveslahti, T. Gao, D. M. Boyer, S. Mukherjee, and L. Crawford. A Statistical Pipeline for Identifying Physical Features That Differentiate Classes of 3d Shapes. *The Annals of Applied Statistics*, 15(2):638–661, 2021.

BROWN UNIVERSITY, DEPARTMENT OF MATHEMATICS, BOX 1917, 151 THAYER STREET, PROVIDENCE, RI 02912, USA

Email address: mattie\_ji@brown.edu