Deep Neural Networks and Finite Elements of Any Order on Arbitrary Dimensions

Juncai He[∗] Jinchao Xu∗†

Abstract

In this study, we establish that deep neural networks employing ReLU and ReLU² activation functions can effectively represent Lagrange finite element functions of any order on various simplicial meshes in arbitrary dimensions. We introduce two novel formulations for globally expressing the basis functions of Lagrange elements, tailored for both specific and arbitrary meshes. These formulations are based on a geometric decomposition of the elements, incorporating several insightful and essential properties of high-dimensional simplicial meshes, barycentric coordinate functions, and global basis functions of linear elements. This representation theory facilitates a natural approximation result for such deep neural networks. Our findings present the first demonstration of how deep neural networks can systematically generate general continuous piecewise polynomial functions on both specific or arbitrary simplicial meshes.

1 Introduction

Deep neural networks (DNNs) with specific activation functions have attracted significant attention in scientific computing, particularly in the context of solving partial differential equations (PDEs) [\[21](#page-26-0), [13,](#page-26-1) [35,](#page-28-0) [41](#page-28-1), [39](#page-28-2)]. To capitalize on the comprehensive mathematical findings and insights from traditional numerical analysis methods, exploring the expressivity properties of deep neural networks for classical methods is valuable. Specifically, we aim to study the relationships between DNNs and finite

[∗]Computer, Electrical and Mathematical Science and Engineering Division, King Abdullah University of Science and Technology, Thuwal 23955, Saudi Arabia.

[†]Department of Mathematics, The Pennsylvania State University, University Park, PA 16802, USA.

element functions [\[11\]](#page-25-0). For instance, it is recognized that DNNs with the rectified lin-ear unit (ReLU) activation function [\[32\]](#page-27-0), defined as $ReLU(t) := max\{0, t\}$, represent a class of continuous piecewise linear (CPwL) functions. It has been shown that any CPwL function on \mathbb{R}^d can be represented by a ReLU DNN with at most $\lceil \log_2(d+1) \rceil$ hidden layers [\[5\]](#page-25-1). However, as indicated in [\[17\]](#page-26-2), such representations may require an extraordinarily large parameter count. By leveraging properties of linear finite element functions, we developed a more succinct and efficient representation for linear finite element functions (CPwL functions with simplex meshes) in [\[17\]](#page-26-2). Beyond representing general CPwL functions, several studies [\[31](#page-27-1), [40,](#page-28-3) [29](#page-27-2), [26,](#page-27-3) [28](#page-27-4)] have illustrated that ReLU networks with adequate depth and $\mathcal{O}(N)$ parameters can generate specific CPwL functions with segment counts exceeding any polynomial of N . These discoveries are pivotal in a body of research focusing on the approximation properties of very deep ReLU DNNs [\[44](#page-28-4), [33,](#page-27-5) [30](#page-27-6), [25](#page-27-7), [27\]](#page-27-8). A hierarchical basis viewpoint, originating from finite elements, was suggested in [\[16\]](#page-26-3) to enhance the understanding of these approximations. Nonetheless, as shown in [\[12,](#page-25-2) [15\]](#page-26-4), such super-expressivity is confined to particular types of CPwL functions. More explicitly, whether ReLU DNNs with $\mathcal{O}(N)$ parameters of various widths and depths can accurately replicate any linear finite element function with N degrees of freedom remains unanswered. Currently, positive evidence exists only for the interval $[0, 1]$, as also demonstrated in [\[12](#page-25-2), [15](#page-26-4)].

On the other hand, the expressive capabilities of neural networks employing ReLU^k activation functions for $k \geq 2$ have also been the subject of research in works such as $[22, 41, 10, 34]$ $[22, 41, 10, 34]$ $[22, 41, 10, 34]$ $[22, 41, 10, 34]$ $[22, 41, 10, 34]$ $[22, 41, 10, 34]$ $[22, 41, 10, 34]$. These inquiries focus mainly on the representation of high-order splines within the interval [0, 1] or global polynomials on \mathbb{R}^d . In particular, [\[18](#page-26-6)] provided a constructive proof with explicit bounds for weight values on how to uniformly express polynomials of arbitrary orders using ReLU^k activation functions for any $k > 2$.

In summary of the preceding studies on the expressivity of ReLU and $ReLU^k$ $(k \geq 2)$ DNNs, we can draw the following two intuitive conclusions:

- 1. ReLU DNNs can represent linear finite element functions with any mesh.
- 2. ReLU^k DNNs for $k \geq 2$ can represent global polynomials of any order.

Consequently, a natural and fundamental question emerges:

Question 1.1 Is it possible to represent any continuous piecewise polynomials, particularly Lagrange finite element functions of any dimension and order, using a combination of ReLU and $ReLU^k$ in DNNs?

A potential strategy for combining ReLU and $ReLU^k$ in DNNs involves the socalled DNNs with learnable or arbitrary activation functions, which have been examined in [\[37](#page-28-5), [23](#page-26-7), [46,](#page-28-6) [10,](#page-25-3) [24,](#page-27-10) [43,](#page-28-7) [42\]](#page-28-8). Notably, the authors in [\[10\]](#page-25-3) suggest learning the activation function for each layer as a composition of $ReLU^k$ for $k = 0$ to some positive integer $n \geq 2$ to improve the efficiency of neural networks, where ReLU⁰ $(t) = t$. The work in $[24]$ then demonstrated the use of ReLU, ReLU^k, and Heaviside activation functions to reproduce the lowest $H(\text{div})$ and $H(\text{curl})$ finite element functions in two or three dimensions. This study also showed how to represent discontinuous piecewise polynomials by using $ReLU^k$ to recover polynomials and Heaviside functions to represent the index functions for simplicial elements. More recently, [\[43](#page-28-7), [42\]](#page-28-8) explored the approximation properties of DNNs with arbitrary ReLU or ReLU^2 activation functions. Nevertheless, most of these studies concentrate on the approximation capabilities of such DNNs or how this innovative architecture, in terms of activation functions, can enhance the performance of neural networks in practical applications. In particular, the expressiveness of DNNs with learnable or arbitrary activation functions has not been thoroughly explored, particularly in terms of their connection to continuous piecewise polynomials (Lagrange finite element functions) of any dimension and order.

To better present the main study of this paper, we introduce the following fully connected DNN architecture with learnable or arbitrary activation functions:

$$
\begin{cases}\n u^{0}(\boldsymbol{x}) = \boldsymbol{x}, \\
 [u^{j}(\boldsymbol{x})]_{i} = \sigma_{i}^{j} ([W^{j}u^{j-1}(\boldsymbol{x}) + b^{j}]_{i}), \quad i = 1, 2, \cdots, n_{j}, \ j = 1, 2, \cdots, L,\n\end{cases} (1.1)
$$
\n
$$
u(\boldsymbol{x}) = W^{L+1}u^{L}(\boldsymbol{x}) + b^{L+1},
$$

where σ_i^j $i : \mathbb{R} \to \mathbb{R}$ denotes the activation function of the *i*-th neuron in the *j*-th hidden layer for all $i = i = 1, 2, \cdots, n_j$ and $j = 1, 2, \cdots, L$.

Notation 1.1 Specifically, we introduce the following two types of DNNs:

• Standard DNNs with fixed ReLU^k activation function:

$$
\Sigma_{n_{1:L}}^k := \{DNNs \text{ defined by (1.1) with } \sigma_i^j = \text{ReLU}^k, \ \forall \ i, j \}
$$
 (1.2)

for any $k \in \mathbb{N}^+$. In particular, $\Sigma^1_{n_{1:L}}$ and $\Sigma^2_{n_{1:L}}$ denote the standard DNNs with $ReLU$ and $ReLU²$ activation functions, respectively.

• DNNs with a special learnable activation function σ_i^{ℓ} :

$$
\Sigma_{n_{1:L}}^{1,2} := \begin{cases}\nDNNs \text{ defined by (1.1) with} \\
\sigma_i^j(t) = a_i^j \text{ReLU}(t) + b_i^j \text{ReLU}^2(t),\n\end{cases}
$$
\n(1.3)

where $t \in \mathbb{R}$ and a_i^j $\mathbf{z}_i^j, \mathbf{b}_i^j \in \mathbb{R}$ are additional trainable parameters. Generally, this indicates that the activation function for each neuron is a combination of $ReLU$ and $ReLU²$.

Consequently, we have

$$
\Sigma_{n_{1:L}}^k \subseteq \Sigma_{n_{1:L}}^{1,2},\tag{1.4}
$$

for $k = 1$ and 2. Thus, a more specific question arises: Can $\sum_{n=1}^{1,2}$ accurately represent any Lagrange finite element functions?

In this paper, we present a positive answer to the expressive capabilities of $\Sigma_{n_{1:L}}^{1,2}$ for representing Lagrange finite element functions of any dimension and order. This result is derived from two novel formulations of the global basis functions for Lagrange finite elements on specific or arbitrary simplicial meshes, incorporating a geometric decomposition of these elements. We initially examine the representation for simplicial meshes with a convex constraint for the support set (patch) of basis functions. Central to this development are two foundational lemmas concerning high-dimensional simplicial meshes and the properties of barycentric functions. For arbitrary simplicial meshes, we propose another uniform formulation using the global basis functions of the linear finite element space on the same mesh. The key of this construction lies in an intrinsic property in the multiplication structure of linear basis functions on a certain sub-simplex, seen as a generalization of the previous result for high-dimensional simplicial meshes with a convex constraint. By integrating these new formulations with the properties of $\Sigma_{n_{1:L}}^{1,2}$, we initially achieve two representations of the basis functions. Advancing this approach with a modified structure of $\Sigma_{n_1,L}^{1,2}$, we also capture representations of any Lagrange finite element functions with parameter levels same to their degrees of freedom. We will demonstrate that the representation of Lagrange finite element functions on simplicial meshes with a convex constraint is more concise and efficient. Furthermore, by examining a uniform simplex mesh over $[0, 1]^d$ that satisfies the convex constraint and exploring the approximation characteristics of these functions, we obtain an additional approximation result for $\Sigma_{n_{1:L}}^{1,2}$. This offers a more intuitive and direct method for obtaining $W^{s,p}$ error estimates for $s = 0, 1$ and $p \in [2, \infty]$.

The structure of this paper is organized as follows. In Section [2,](#page-4-0) we present some preliminaries about high-dimensional simplicial meshes and the geometric decomposition of Lagrange finite elements. In Section [3,](#page-8-0) we introduce these two new formulas to explicitly and globally define the basis functions of Lagrange finite elements for both specific and arbitrary simplicial meshes. In Section [4,](#page-16-0) we show how DNNs can be employed to represent Lagrange finite element functions in any dimension and of any order. In Section [5,](#page-22-0) we show an approximation result that is a corollary of our representation insights. We draw some concluding remarks in Section [6,](#page-24-0) reflecting on the implications of our findings.

2 Preliminaries

In this section, we introduce some preliminary results about simplicial meshes, Lagrange finite elements, and the geometric decomposition of these spaces [\[4](#page-25-4), [8](#page-25-5)].

Simplex, sub-simplex, simplicial mesh, and their properties in \mathbb{R}^d . For any finite set $\{\boldsymbol{v}_1, \boldsymbol{v}_2, \cdots, \boldsymbol{v}_m\} \subset \mathbb{R}^d$, we denote the convex combination as

Conv
$$
(\mathbf{v}_1, \mathbf{v}_2, \cdots, \mathbf{v}_m) := \left\{ \mathbf{x} = \sum_{i=1}^m a_i \mathbf{v}_i : \sum_{i=1}^m a_i = 1, a_i \ge 0, \forall i \right\}.
$$
 (2.1)

Then, for any set $S \subseteq \mathbb{R}^d$, the convex hull of S Conv (S) is defined as the set of all convex combinations of points in S. A set $T \subset \mathbb{R}^d$ is called a simplex if it is the closed convex hull of $d+1$ affinely independent points $\{v_0, v_1, \dots, v_d\}$, i.e.,

$$
T = \text{Conv}(\boldsymbol{v}_0, \boldsymbol{v}_1, \cdots, \boldsymbol{v}_d) = \left\{ \boldsymbol{x} \; : \; \boldsymbol{x} = \sum_{i=0}^d \lambda_i(\boldsymbol{x}) \boldsymbol{v}_i \right\}.
$$
 (2.2)

These points are the vertices of T, and these affine functions $\lambda_i(\boldsymbol{x})$ are the barycentric coordinate functions. For any simplex $T = \text{Conv}(\boldsymbol{v}_0, \boldsymbol{v}_1, \cdots, \boldsymbol{v}_d) \subseteq \mathbb{R}^d$, we call $f =$ Conv $\left(v_0^f\right)$ $_0^f, \bm v_1^f$ $_{1}^{f},\cdots, {\boldsymbol v}_{\ell}^{f}$ ℓ) as a ℓ -dimensional sub-simplex of T for any $\big\{\boldsymbol{v}_0^f\big\}$ $_0^f, \bm{v}_1^f$ $_{1}^{f},\cdots, {\bm v}_{\ell}^{f}$ ℓ o ⊆ $\{v_0, v_1, \dots, v_d\}$ with some $0 \leq \ell \leq d$. Following [\[4](#page-25-4)], let $\Delta(T)$ denote all the subsimplices of T, while $\Delta_{\ell}(T)$ denotes the set of sub-simplices of dimension ℓ for $\ell=0,1,\ldots,d.$

From then on, we denote $\Omega \subset \mathbb{R}^d$ as a bounded polytope domain. We say that \mathcal{T} is a simplicial mesh of Ω if it consists of d-dimensional simplices, that is, $\Omega = \bigcup_{T \in \mathcal{T}} T$, and the intersection of any two simplices is a face (a sub-simplex) common to these two simplices if the intersection is not empty. Correspondingly, for the simplicial mesh \mathcal{T} , we define

$$
\Delta_{\ell}(\mathcal{T}) := \bigcup_{T \in \mathcal{T}} \Delta_{\ell}(T). \tag{2.3}
$$

That is, $\Delta_{\ell}(\mathcal{T})$ denotes the collection of all ℓ -dimensional sub-simplices in \mathcal{T} . For example, $\Delta_0(\mathcal{T})$ represents the set of all vertices in $\mathcal{T}, \Delta_1(\mathcal{T})$ corresponds to the set of all edges in \mathcal{T} , and so on, up to $\Delta_d(\mathcal{T}) = \mathcal{T}$.

Lagrange finite elements and global basis functions in classical definition. As introduced in [\[4\]](#page-25-4), we first define the simplicial lattice, which will be used later in establishing the local basis functions for Lagrange finite elements with k -th order $(P_k$ element). For any $k \geq 1$, the simplicial lattice of degree k is given by

$$
\mathbb{T}_k^d := \left\{ \alpha \in \mathbb{N}^{d+1} \ : \ |\alpha| = k \right\},\tag{2.4}
$$

where $\boldsymbol{\alpha} = (\alpha_0, \alpha_1, \dots, \alpha_d)$ and $|\boldsymbol{\alpha}| = \sum_{i=0}^d \alpha_i$. For any simplex $T = \text{Conv}(\boldsymbol{v}_0, \boldsymbol{v}_1, \cdots, \boldsymbol{v}_d)$, we have the following interpolation points of P_k elements defined by

$$
\mathcal{X}_k(T) = \left\{ \boldsymbol{x}_{\alpha} = \frac{1}{k} \sum_{i=0}^d \alpha_i \boldsymbol{v}_i \ : \ \boldsymbol{\alpha} \in \mathbb{T}_k^d \right\}.
$$
 (2.5)

 $\mathcal{X}_k(T)$ can be interpreted as the geometric embedding of the algebraic set \mathbb{T}_k^d . Geometrically, $\mathcal{X}_k(T)$ represents the intersection points within T of the hyperplanes defined by $\{\boldsymbol{x}: k\lambda_i(\boldsymbol{x}) - j = 0\}$ for all $i = 0, 1, \dots, d$ and $j = 0, 1, \dots, k$. The cardinality of $\mathcal{X}_k(T)$ is $\binom{d+k}{k}$ $\binom{+k}{k}$, which is also the cardinality of \mathbb{T}_k^d and the dimension of $\mathbb{P}_k(T)$ the polynomial space on $T \subset \mathbb{R}^d$ of degree at most k.

For any $p \in \mathcal{X}_k(T)$, we denote $\varphi_p(x) \in \mathbb{P}_k(T)$ as the interpolation basis function corresponding to p such that

$$
\varphi_{\mathbf{p}}(\mathbf{p}') = \begin{cases} 1, & \text{if } \mathbf{p} = \mathbf{p}', \\ 0, & \text{if } \mathbf{p} \neq \mathbf{p}', \end{cases}
$$
 (2.6)

for all $p, p' \in \mathcal{X}_k(T)$. The following theorem provides a uniform representation of the local basis function $\varphi_p(x)$ on T for any $p \in \mathcal{X}_k(T)$.

Theorem 2.1 ([\[7](#page-25-6), [8](#page-25-5)]) For any $p = x_\alpha \in \mathcal{X}_k(T)$ with $\alpha \in \mathbb{T}_k^d$ and $x \in T$, the basis function is given by

$$
\varphi_{\mathbf{p}}(\mathbf{x}) = \frac{1}{\alpha!} \prod_{i=0}^{d} \prod_{j=0}^{\alpha_i - 1} (k \lambda_i(\mathbf{x}) - j), \qquad (2.7)
$$

where $\boldsymbol{\alpha}! = \prod_{i=0}^{d} (\alpha_i!).$

Definition 2.1 For any simplicial mesh \mathcal{T} of Ω with N total elements (simplices) on Ω , let us assume the simplices in $\mathcal T$ are ordered as T_i for $i = 1, \cdots, N$ and denote

$$
I_S := \{ i : S \subseteq T_i \} \quad and \quad P_S := \bigcup_{S \subseteq T \in \mathcal{T}} T = \bigcup_{i=I_S} T_i \tag{2.8}
$$

as the index set and patch set for any $S \subseteq \Omega$. With a slight abuse of notation, we denote I_x and P_x if $S = \{x\}$ is a set of a single point.

By aggregating \mathcal{X}_T for all $T \in \mathcal{T}$, we obtain the set

$$
\mathcal{X}_k(\mathcal{T}) := \bigcup_{T \in \mathcal{T}} \mathcal{X}_k(T). \tag{2.9}
$$

Therefore, the global basis function $\varphi_p(x)$ for any $p \in \mathcal{X}_k(\mathcal{T})$ constructed by gluing $\varphi_p(x)|_{T_i}$ together for all $i \in I_p$. More precisely, $\varphi_p(x) = 0$ if $x \notin P_p$ and for any $s \in I_p$ with $p = x_\alpha \in \mathcal{X}_k(T_s)$, we have

$$
\varphi_{\mathbf{p}}(\mathbf{x}) = \frac{1}{\alpha!} \prod_{i=0}^{d} \prod_{j=0}^{\alpha_i - 1} (k \lambda_{s,i}(\mathbf{x}) - j), \quad \forall \mathbf{x} \in T_s,
$$
\n(2.10)

where $\lambda_{s,i}(\boldsymbol{x})$ are the barycentric coordinate functions corresponding to the *i*-th vertex in T_s for $i = 0, 1, \dots, d$. Obviously, we have $\text{supp}(\varphi_p) = P_p$ for any $p \in \mathcal{X}_k(\mathcal{T})$, where $\text{supp}(f(\boldsymbol{x})) := \{ \boldsymbol{x} : f(\boldsymbol{x}) \neq 0 \}$ denotes the support set of $f(\boldsymbol{x})$.

Geometric Decomposition of $\mathcal{X}_k(\mathcal{T})$. As seen from the definition of the global basis function $\varphi_p(x)$, it is necessary to determine the corresponding index α such that $p = x_\alpha \in \mathcal{X}_k(T_s)$ for different T_s with $s \in I_p$. The key observation is that some invariant properties of α for p across different T_s can be identified using a geometric decomposition of $\mathcal{X}_k(\mathcal{T})$. To better formulate this decomposition, for any two sets A and B, we denote $A \oplus B = A \cup B$ if $A \cap B = \emptyset$.

First, we define the local decomposition of $\mathcal{X}_k(T)$ for any $T \in \mathcal{T}$. Here, $\mathcal{X}_k^{\ell}(T)$ is defined by

$$
\mathcal{X}_k^{\ell}(T) = \bigoplus_{f \in \Delta_{\ell}(T)} \mathcal{X}_k(\mathring{f}),\tag{2.11}
$$

where $\mathcal{X}_k(\mathring{f}) = \mathcal{X}_k(T) \cap \mathring{f}$. Here, \mathring{f} denotes the interior of the ℓ -dimensional subsimplex f, and $\mathring{f} = f$ if $\ell = 0$. That is, $\mathcal{X}_k^{\ell}(T)$ represents those interpolation points in $\mathcal{X}_k(T)$ that are part of some ℓ -dimensional sub-simplices in $\Delta_{\ell}(T)$ but do not belong to any sub-simplex in $\Delta_{\ell-1}(T)$. Then, the decomposition of $\mathcal{X}_k(T)$ is defined as

$$
\mathcal{X}_k(T) = \bigoplus_{\ell=0}^d \mathcal{X}_k^{\ell}(T). \tag{2.12}
$$

Figure [2.1](#page-7-0) illustrates an example of the decomposition of $\mathcal{X}_4(T)$ on a 2D simplex T.

Figure 2.1: An example of the decomposition of $\mathcal{X}_4(T)$ on a 2D simplex T is illustrated. Here, $\mathcal{X}_4^0(T)$ denotes all the vertices of T (marked in blue), $\mathcal{X}_4^1(T)$ represents all nodes on the interiors of edges (marked in green), and $\mathcal{X}_4^2(T)$ includes all nodes in the interior of the simplex T (marked in yellow).

Finally, by defining $\mathcal{X}_{k}^{\ell}(\mathcal{T}) = \bigcup_{T \in \mathcal{T}} \mathcal{X}_{k}^{\ell}(T) = \bigoplus_{f \in \Delta_{\ell}(\mathcal{T})} \mathcal{X}_{k}(\mathring{f}),$ we have

$$
\mathcal{X}_k(\mathcal{T}) = \bigoplus_{\ell=0}^d \mathcal{X}_k^{\ell}(\mathcal{T}) = \bigoplus_{\ell=0}^d \bigoplus_{f \in \Delta_{\ell}(\mathcal{T})} \mathcal{X}_k(\mathring{f}). \tag{2.13}
$$

One of the first properties given by the above decomposition is that

$$
I_p = I_f \quad \text{and} \quad P_p = P_f \tag{2.14}
$$

if $p \in \mathcal{X}_{k}^{\ell}(\mathcal{T})$ for some $f \in \Delta_{\ell}(\mathcal{T})$ and $p \in \mathring{f}$, or $p \in \mathcal{X}_{k}(\mathring{f})$ for some $f \in \Delta_{\ell}(\mathcal{T})$.

The geometric decomposition for Lagrange elements has been widely studied in [\[3,](#page-25-7) [4,](#page-25-4) [9,](#page-25-8) [8\]](#page-25-5). However, these studies mainly focus on the local decomposition of polynomial spaces $\mathbb{P}_k(T)$. In the following, we will demonstrate how to use this decomposition and global basis for linear elements to derive two new formulations for global basis functions of Lagrange finite elements of any order on both specific and arbitrary simplicial meshes.

3 New formulations of global Lagrange element basis functions of any order on any dimensions

In this section, we introduce two innovative formulations of the global basis functions for Lagrange finite elements. The first formulation emerges from a geometric constraint for the mesh \mathcal{T} , coupled with two foundational insights pertaining to high-dimensional simplicial meshes and barycentric functions. The second formulation generalizes the result by removing the geometric constraint for the mesh \mathcal{T} .

Given the decomposition of $\mathcal{X}_k(\mathcal{T})$ mentioned above, for any $p \in \mathcal{X}_k^{\ell}(\mathcal{T})$ and k $\boldsymbol{p} \in f = {\rm Conv}(\boldsymbol{v}_0^f$ $_0^f, \bm v_1^f$ $\frac{f}{1},\ldots,\bm{v}_\ell^f$ ℓ), we first observe that there uniquely exist $\alpha_i \in \mathbb{N}^+$ for $i = 0, 1, \ldots, \ell$ with $\sum_{i=0}^{\ell} \alpha_i = k$ such that

$$
\boldsymbol{p} = \frac{1}{k} \sum_{i=0}^{\ell} \alpha_i \boldsymbol{v}_i^f. \tag{3.1}
$$

To simplify the expression of our main theorem, we introduce the following definition.

Definition 3.1 For any $0 \leq \ell \leq d$, $p \in \mathcal{X}_k^{\ell}(\mathcal{T})$, and $p \in \mathring{f}$ with $f \in \Delta_{\ell}(\mathcal{T})$, we always define the $\ell + 1$ vertices in f as the first $\ell + 1$ vertices in T_s for each $s \in I_p$.

Consequently, for any $i \in I_p$, we have $p \in \mathcal{X}_k^{\ell}(T_i)$ and $p = x_\alpha$ where

$$
\boldsymbol{\alpha} = (\alpha_0, \cdots, \alpha_\ell, 0, \dots, 0) \in \mathbb{T}_k^d.
$$
\n(3.2)

That is, the index α of $p \in \mathcal{X}_k^{\ell}(\mathcal{T})$ depends only on the f where $p \in \mathring{f}$ and is an invariant on different T_s for $s \in I_p$ by applying Definition [3.1.](#page-8-1)

A novel formulation for global basis functions on vertex-convex meshes.

Definition 3.2 For any $0 \leq \ell \leq d$, we call a simplicial mesh \mathcal{T} of Ω is " ℓ -convex" if $P_f = \bigcup_{f \subseteq T \in \mathcal{T}} T$ is convex for any ℓ -dimensional sub-simplex $f \in \Delta_{\ell}(\mathcal{T})$. In particular, we call $\mathcal T$ "vertex-convex" if it is 0-convex. Obviously, $\mathcal T$ is d-convex.

Lemma 3.1 For any simplicial mesh \mathcal{T} on Ω and $1 \leq \ell \leq d$, \mathcal{T} is ℓ -convex if it is $(\ell-1)$ -convex.

Proof For any $f = \text{Conv}(\boldsymbol{v}_0^f)$ $_0^f, \bm v_1^f$ $_{1}^{f}\ldots, \cdots$ $\boldsymbol{v}_{\ell}^{f}$ $(\mathcal{L}_{\ell}) \in \Delta_{\ell}(\mathcal{T})$ with some $0 < \ell < d$, let us define $\widetilde{f}_i = \text{Conv}(\boldsymbol{v}_0^f)$ $_{0}^{f},\cdots,$ \boldsymbol{v}_{i}^{f} $_{i-1}^f, \bm{v}^f_{i+1}, \cdots \bm{v}^f_{\ell}$ $\mathcal{L}_{\ell}^{J}(\mathcal{T})\in \Delta_{\ell-1}(\mathcal{T}).$

Thus, $P_{\tilde{f}_i}$ are convex for all $i = 0, 1, \ldots, \ell$. To complete the proof, we show that

$$
P_f = \bigcap_{i=0}^{\ell} P_{\tilde{f}_i},\tag{3.3}
$$

where P_f and $P_{\tilde{f}_i}$ are defined as in Definition [2.1.](#page-5-0) This will be sufficient to demonstrate that P_f is convex, given the convexity of $P_{\tilde{f}_i}$.

First, for any $T \in \mathcal{T}$ such that $f \subseteq T$, we have $f_i \subset T$ for all $i = 0, 1, \ldots, \ell$. That is, $T \subseteq P_{\tilde{f}_i}$ for all $i = 0, 1, \ldots, \ell$, i.e.,

$$
P_f \subseteq \bigcap_{i=0}^{\ell} P_{\tilde{f}_i}.\tag{3.4}
$$

Furthermore, by using the associative property of set algebra, we have

$$
\bigcap_{i=0}^{\ell} P_{\widetilde{f}_i} = \bigcap_{i=0}^{\ell} \bigcup_{j_i \in I_{\widetilde{f}_i}} T_{j_i} = \bigcup_{\substack{j_0 \in I_{\widetilde{f}_0} \\ j_\ell \in I_{\widetilde{f}_0}}} \bigcap_{i=0}^{\ell} T_{j_i}.
$$

Given the property of simplicial mesh, $\bigcap_{i=0}^{\ell} T_{j_i}$ can only be a sub-simplex or a simplex. Consequently, we have

$$
\bigcap_{i=0}^{\ell} P_{\tilde{f}_i} = \bigcup_{\ell=0}^{d} \mathcal{S}_{\ell} = \mathcal{S}_d \bigcup \mathcal{S}',\tag{3.5}
$$

where $\mathcal{S}_{\ell} \subset \Delta_{\ell}(\mathcal{T})$ denote the ℓ -dimensional sub-simplices for all $\ell = 0, 1, \ldots, d$ and $\mathcal{S}' = \bigcup_{\ell=0}^{d-1} \mathcal{S}_{\ell}$. We first claim that $\mathcal{S}_d = P_f$. For any $T \in \mathcal{T}$ and $T \subseteq \mathcal{S}_d \subseteq \bigcap_{i=0}^{\ell} P_{\tilde{f}_i}$, we have $\widetilde{f}_i \subseteq T$. Thus, $f \subset T$ and $\mathcal{S}_d = P_f$.

Now, we prove that $\mathcal{S}' = \emptyset$. If not, there exists $x \in \mathcal{S}'$ and $x \notin \mathcal{S}_d$. Hence, we have

$$
\text{dist}(\boldsymbol{x},\mathcal{S}_d) := \inf_{\boldsymbol{x}'\in\mathcal{S}_d} \|\boldsymbol{x}-\boldsymbol{x}'\| = \delta > 0
$$

since $S_d = P_f$ is a closed set. Then, for any $T \in \mathcal{T}$ and $T \subseteq S_d$, we have that

$$
B_{\boldsymbol{x}} := B(\boldsymbol{x},\delta/2) \bigcap \text{Conv} \left(\{\boldsymbol{x}\} \cup \mathcal{S}_d \right)
$$

has nonzero d-dimensional volume and $B_x \cap S_d = \emptyset$ where $B(x, \delta/2)$ denotes the d-dimensional ball centered at x with radius $\delta/2$. However, we have

$$
B_{\boldsymbol{x}} \subseteq \text{Conv}\left(\mathcal{S}' \bigcup \mathcal{S}_d\right) = \text{Conv}\left(\bigcap_{i=0}^{\ell} P_{\tilde{f}_i}\right) = \bigcap_{i=0}^{\ell} P_{\tilde{f}_i} = \mathcal{S}' \bigcup \mathcal{S}_d \tag{3.6}
$$

since $P_{\tilde{f}_i}$ is convex for each $i = 0, 1, \ldots, \ell$. Thus, we have $B_{\boldsymbol{x}} \subset \mathcal{S}'$ since $B_{\boldsymbol{x}} \cap \mathcal{S}_d = \emptyset$. This is impossible since \mathcal{S}' consists of low-dimensional sub-simplices which have only zero d-dimensional volume. This finishes the proof. \Box

Theorem 3.1 For any vertex-convex simplicial mesh \mathcal{T} of Ω for the P_k element and $0 \leq \ell \leq d$, by taking Definition [3.1,](#page-8-1) the basis function φ_p corresponding to the interpolation point $p \in \mathcal{X}_k^{\ell}(\mathcal{T})$ can be written as

$$
\varphi_{\mathbf{p}}(\mathbf{x}) = \frac{1}{\alpha!} \min_{i' \in I_{\mathbf{p}}} \left\{ \prod_{i=0}^{\ell} k \text{ReLU}(\lambda_{i',i}(\mathbf{x})) \right\} \times \prod_{i=0}^{\ell} \prod_{j=1}^{\alpha_{i}-1} \left(k \min_{i' \in I_{\mathbf{p}}} \left\{ \lambda_{i',i}(\mathbf{x}) \right\} - j \right), \quad (3.7)
$$

where $\lambda_{i',i}(\boldsymbol{x})$ denote the barycentric coordinates functions corresponding to the *i*-th vertex in the element $T_{i'}$ for all $i' \in I_p$ and $i = 0, 1, \ldots, \ell$. With a slight abuse of notation, we assume $\lambda_{i',i}(\boldsymbol{x})$ are defined on \mathbb{R}^d .

Lemma 3.2 With the same assumptions as in Theorem [3.7,](#page-10-0) we have

$$
supp(\varphi_{\mathbf{p}}) = P_{\mathbf{p}} = P_f = \bigcap_{i' \in I_{\mathbf{p}}} \bigcap_{i=0}^{\ell} \left\{ \mathbf{x} \in \Omega : \lambda_{i',i}(\mathbf{x}) \ge 0 \right\},\tag{3.8}
$$

where $p \in \mathring{f}$ for some $f \in \Delta_{\ell}(\mathcal{T})$.

Proof Given the definition of $\varphi_n(x)$ in [\(2.10\)](#page-6-0) and properties in [\(2.14\)](#page-7-1), the first two identities hold. For the last identity, we first observe the convexity of supp $(\varphi_p) = P_f$ since T is vertex-convex. Moreover, the boundary set $\partial(\text{supp}(\varphi_p))$ consists of faces

$$
F_{i',i} := \{ \mathbf{x} \in T_{i'} : \lambda_{i',i}(\mathbf{x}) = 0 \},\tag{3.9}
$$

for all $i' \in I_p$ and $i = 0, 1, \ldots, \ell$. Additionally, we have $\{x \in \Delta_i : \lambda_{i,j}(x) \geq 0\} \subseteq$ supp(φ_n). Thus, the d-dimensional polytope P_f can be expressed as the intersection of half-spaces as in (3.8) .

Then, to prove the main theorem, let us establish the following lemmas regarding $\text{important properties of } \min_{i' \in I_p} \{ \lambda_{i',i}(\boldsymbol{x}) \} \text{ in } \varphi_{\boldsymbol{p}}(\boldsymbol{x}) \text{ and } \min_{i' \in I_p} \left\{ \prod_{i=0}^{\ell} \text{ReLU}(\lambda_{i',i}(\boldsymbol{x})) \right\},$ under the assumption that $\mathcal T$ is a vertex-convex mesh.

Lemma 3.3 For any vertex-convex simplicial mesh \mathcal{T} of Ω for the P_k element and $\boldsymbol{p} \in \mathcal{X}_{k}^{\ell}(\mathcal{T})$ with some $0 \leq \ell \leq d$, by taking Definition [3.1,](#page-8-1) we have

$$
\min_{i' \in I_p} \left\{ \lambda_{i',i}(\boldsymbol{x}) \right\} = \lambda_{s,i}(\boldsymbol{x}) \tag{3.10}
$$

for any $0 \leq i \leq \ell$ and $\boldsymbol{x} \in T_s$ with $s \in I_p$.

Proof Assuming that $p \in \mathring{f}$ with some $f = \text{Conv}\left(\boldsymbol{v}_0^f\right)$ $_0^f, \bm v_1^f$ $i_1^f,\ldots,\bm{v}_\ell^f$ ℓ $\Big) \in \Delta_{\ell}(\mathcal{T}), \text{ for }$ any $i = 0, 1, \ldots, \ell$, we have

$$
\lambda_{i',i}(\mathbf{v}_i^f) = 1, \quad \forall i' \in I_p. \tag{3.11}
$$

For any $x \in T_s$ with some $s \in I_p$ and $0 \leq i \leq \ell$, given Lemma [3.2,](#page-10-2) there is a unique point $y_x \in F_{s,i} = \{x \in T_s : \lambda_{s,i}(x) = 0\}$ such that there exists $t_x \in [0,1]$ and

$$
\boldsymbol{x} = t_{\boldsymbol{x}} \boldsymbol{v}_i^f + (1 - t_{\boldsymbol{x}}) \boldsymbol{y}_{\boldsymbol{x}}.
$$
\n(3.12)

Since $\text{supp}(\varphi_p) = P_p$ is convex as shown in Lemma [3.1,](#page-8-2) we have

$$
\lambda_{i',i}(\mathbf{y_x}) \ge 0, \quad \forall i' \in I_p,\tag{3.13}
$$

and in particular,

$$
\lambda_{s,i}(\mathbf{y_x}) = 0. \tag{3.14}
$$

Since $\lambda_{i,j}(\boldsymbol{x})$ are all affine maps, it follows that

$$
\lambda_{i',i}(\boldsymbol{x}) = \lambda_{i',i} \left(t_{\boldsymbol{x}} \boldsymbol{v}_i^f + (1 - t_{\boldsymbol{x}}) \boldsymbol{y}_{\boldsymbol{x}} \right)
$$

\n
$$
= t_{\boldsymbol{x}} \lambda_{i',i} (\boldsymbol{v}_i^f) + (1 - t_{\boldsymbol{x}}) \lambda_{i',i} (\boldsymbol{y}_{\boldsymbol{x}})
$$

\n
$$
\geq t_{\boldsymbol{x}}
$$

\n
$$
= t_{\boldsymbol{x}} \lambda_{s,i} (\boldsymbol{v}_i^f) + (1 - t_{\boldsymbol{x}}) \lambda_{s,i} (\boldsymbol{y}_{\boldsymbol{x}})
$$

\n
$$
= \lambda_{s,i}(\boldsymbol{x})
$$
\n(3.15)

for any $i \in I_p$. Thus, the proof is complete.

Remark 1 Here, we note that a core identity used in [\[17\]](#page-26-2) and [\[45\]](#page-28-9) to represent linear finite element functions using ReLU DNNs is essentially a special case of $\ell = 0$ in the above lemma. An illustration of the proof is provided in Fig [3.1](#page-12-0) for $p \in \mathcal{X}_2^1(\mathcal{T})$ within a 2D simplicial mesh.

Figure 3.1: An example of $p \in \mathcal{X}_2^1(\mathcal{T})$ on 2D, where $I_p = \{s, s'\}$ and $p \in f =$ $\mathrm{Conv}(\bm{v}_0^f$ $_0^f, \bm v_1^f$ ^{*I*}</sup>). We have min $\{\lambda_{s,i}(\boldsymbol{x}), \lambda_{s',i}(\boldsymbol{x})\} = \lambda_{s,i}(\boldsymbol{x})$ for any $\boldsymbol{x} \in T_s$ and $i = 0, 1$.

Lemma 3.4 For any vertex-convex simplicial mesh \mathcal{T} of Ω for the P_k element and $\boldsymbol{p} \in \mathcal{X}_{k}^{\ell}(\mathcal{T})$ with some $0 \leq \ell \leq d$, by taking Definition [3.1,](#page-8-1) we have

$$
\min_{i' \in I_p} \left\{ \prod_{i=0}^{\ell} \text{ReLU} \left(\lambda_{i',i}(\boldsymbol{x}) \right) \right\} = \begin{cases} \prod_{i=0}^{\ell} \lambda_{s,i}(\boldsymbol{x}) & \text{if } \boldsymbol{x} \in T_s \text{ and } s \in I_p \\ 0 & \text{if } \boldsymbol{x} \notin P_p. \end{cases} \tag{3.16}
$$

Proof Given Lemma [3.2,](#page-10-2) we observe that $\prod_{i=0}^{\ell}$ ReLU $(\lambda_{i',i}(\boldsymbol{x})) \geq 0$ for all $i' \in I_p$. Then, if $x \notin P_p$, there exists a $\lambda_{i,i}$ for some $i' \in I_p$ and $0 \leq i \leq \ell$ such that $\lambda_{i',i}(\boldsymbol{x}) < 0.$ Therefore, $\min_{i' \in I_{\boldsymbol{p}}} \left\{ \prod_{i=0}^{\ell} \text{ReLU}(\lambda_{i',i}(\boldsymbol{x})) \right\} = 0$ if $\boldsymbol{x} \notin P_{\boldsymbol{p}}$.

For any $x \in \text{supp}(\varphi_p) = P_p$, there exists $s \in I_p$ such that $x \in T_s$. Moreover, we have $\lambda_{i',i}(\boldsymbol{x}) \geq 0$ for all $i' \in I_p$ and $i = 0, 1, \ldots, \ell$ due to the connection in [\(3.8\)](#page-10-1). Combined with Lemma [3.3,](#page-11-0) it follows that

$$
\min_{i' \in I_p} \left\{ \prod_{i=0}^{\ell} \text{ReLU} \left(\lambda_{i',i}(\boldsymbol{x}) \right) \right\} = \prod_{i=0}^{\ell} \lambda_{s,i}(\boldsymbol{x}) \tag{3.17}
$$

when $x \in T_s$ for some $s \in I_p$. This completes the proof.

Proof of Theorem [3.1](#page-10-3) Proof Given Lemma [3.3](#page-11-0) and Lemma [3.4,](#page-12-1) for any $x \in T_s$

with some $s \in I_p$, we have

$$
\frac{1}{\alpha!} \min_{i' \in I_p} \left\{ \prod_{i=0}^{\ell} \text{ReLU} \left(\lambda_{i',i}(\boldsymbol{x}) \right) \right\} \times \prod_{i=0}^{\ell} \prod_{j=1}^{\alpha_i - 1} \left(k \min_{i' \in I_p} \left\{ \lambda_{i',i}(\boldsymbol{x}) \right\} - j \right)
$$
\n
$$
= \frac{1}{\alpha!} \left\{ \prod_{i=0}^{\ell} \text{ReLU} \left(\lambda_{s,i}(\boldsymbol{x}) \right) \right\} \times \prod_{i=0}^{\ell} \prod_{j=1}^{\alpha_i - 1} \left(k \lambda_{s,i}(\boldsymbol{x}) - j \right)
$$
\n
$$
= \frac{1}{\alpha!} \prod_{i=0}^{\ell} \prod_{j=0}^{\alpha_i - 1} \left(k \left\{ \lambda_{s,i}(\boldsymbol{x}) \right\} - j \right) = \frac{1}{\alpha!} \prod_{i=0}^{d} \prod_{j=0}^{\alpha_i - 1} \left(k \left\{ \lambda_{s,i}(\boldsymbol{x}) \right\} - j \right).
$$
\n(3.18)

A novel formulation for global basis functions on arbitrary simplicial meshes. In this section, we propose another new formulation for the global basis functions of Lagrange elements on arbitrary simplicial meshes. We construct the piecewise polynomial basis functions by using the multiplication of piecewise linear functions, which are the basis functions of linear finite elements on the same simplicial mesh. The most crucial step in this construction is based on a generalization of Lemma [3.1.](#page-8-2)

We start our construction with the following observation. For $\ell = 0$, i.e., $p \in$ $\mathcal{X}_{k}^{0}(\mathcal{T}) = \Delta_{0}(\mathcal{T})$, we have

$$
\varphi_{\mathbf{p}}(\mathbf{x}) = \frac{1}{\alpha_0!} \prod_{j=0}^{\alpha_0-1} (k \lambda_{s,0}(\mathbf{x}) - j)
$$

for any $\mathbf{x} \in T_s$ for $s \in I_p$.

Definition 3.3 For any simplicial mesh \mathcal{T} of $\Omega \subset \mathbb{R}^d$, let ϕ_v denote the piecewise linear basis function at $v \in \mathcal{X}_{k}^{0}(\mathcal{T}) = \Delta_{0}(\mathcal{T})$, i.e.,

$$
\phi_{\mathbf{v}}(\mathbf{x}) = \begin{cases} \lambda_{s,j} & \text{if } \mathbf{x} \in T_s, \text{ for some } s \in I_{\mathbf{v}} \text{ and } \mathbf{v} \text{ is the } j\text{-th vertex in } T_i \\ 0 & \text{if } x \notin \bigcup_{s \in I_{\mathbf{v}}} T_s = P_{\mathbf{v}}.\end{cases}
$$

Consequently, the aforementioned global basis function $\varphi_p(x)$ for any $p \in \mathcal{X}_{k}^0(\mathcal{T})$ can be represented as:

$$
\varphi_{\mathbf{p}}(\mathbf{x}) = \frac{1}{\alpha_0!} \prod_{j=0}^{\alpha_0 - 1} \left(k \phi_{\mathbf{p}}(\mathbf{x}) - j \right), \quad \forall \mathbf{x} \in \Omega.
$$
 (3.19)

Here, we notice that $\varphi_p(\boldsymbol{x}) = 0$ for any $\boldsymbol{x} \notin P_{\boldsymbol{v}}$ since $\prod_{j=0}^{\alpha_0-1} (k\phi_p(\boldsymbol{x}) - j) = k\phi_p(\boldsymbol{x}) \times$ $\prod_{j=1}^{\alpha_0-1} (k\phi_p(\boldsymbol{x})-j).$

A natural question arises as to whether we can generalize (3.19) to arbitrary $\ell > 0$. A main observation of this effort is summarized as the following lemma, which can be understood as a generalization of Lemma [3.1.](#page-8-2)

Lemma 3.5 For any simplicial mesh \mathcal{T} and $f \in \Delta_{\ell}(\mathcal{T})$ with $0 \leq \ell \leq d$ and $f =$ $\mathrm{Conv}(\bm{v}_0^f$ $_0^f, \bm v_1^f$ $_{1}^{f}\ldots, \bm{v}_{\ell}^{f}$ $\binom{J}{\ell}$, we have

$$
\prod_{i=0}^{\ell} \phi_{\mathbf{v}_i^f}(\mathbf{x}) = 0, \quad \text{if} \quad \mathbf{x} \notin \bigcup_{f \subseteq T \in \mathcal{T}} T = P_f.
$$

Proof This proof mainly follows the proof of Lemma [3.1.](#page-8-2) First, we have

$$
\prod_{i=0}^{\ell} \phi_{\mathbf{v}_i^f}(\mathbf{x}) = 0 \quad \text{if} \quad \mathbf{x} \notin \bigcap_{i=0}^{\ell} P_{v_i^f}.
$$

Then, similar to the proof of Lemma [3.1,](#page-8-2) we have the following decomposition

$$
\bigcap_{i=0}^{\ell} P_{\boldsymbol{v}_i^f} = \bigcup_{\ell=0}^d \mathcal{S}_{\ell} = \mathcal{S}_d \bigcup \mathcal{S}'.
$$

If \mathcal{S}' is nonempty, a key observation here is that

$$
f \not\subset f'
$$
 for any $f' \in \Delta_{\ell'}(\mathcal{T})$ and $f' \subseteq \mathcal{S}'$. (3.20)

Thus, we can complete the proof by showing that $\prod_{i=0}^{\ell} \phi_{\mathbf{v}_i^f}(\mathbf{x}) = 0$ for all $\mathbf{x} \in \mathbb{R}$ $f' \subseteq \mathcal{S}'$ for any $f' \in \Delta_{\ell'}(\mathcal{T})$ and $f' \subset \mathcal{S}'$. For this purpose, we claim

$$
f' \subset \bigcup_{i=0}^{\ell} \left(\partial P_{v_i^f} \right) \tag{3.21}
$$

for any $f' \subseteq S'$ with $f' \in \Delta_{\ell'}(\mathcal{T})$. If not, there exists $x \in f' \subseteq S'$ for some $f' \in \Delta_{\ell'}(\mathcal{T})$ with $\ell' < d$ such that $\boldsymbol{x} \in \bigcap_{i=0}^{\ell} \mathring{P}_{\boldsymbol{v}_i^f}$. That is, there exists a d-dimensional ball $B(\mathbf{x}, r)$ centered at \mathbf{x} with radius $r > 0$ such that $B(\mathbf{x}, r) \subset \mathring{P}_{\{v_i^f\}}$ for all $i = 0, 1, \ldots, \ell$. Hence, we have $f' \cap B(x, r) \subset \mathring{P}_{\{v_i^f\}}$. This implies that $f' \subset P_{v_i^f}$ since $\mathring{f}' \cap \mathring{P}_{v_i^f} \neq \emptyset$ for all $i = 0, 1, \ldots, \ell$. Therefore, there exists $T_{x,i} \in \mathcal{T}$ with $T_{x,i} \subset P_{\{v_i^f\}}$ such that $\boldsymbol{x} \in f' \subset \partial T_{\boldsymbol{x},i}$ for all $i = 0, 1, \ldots, \ell$. Furthermore, noting that $F_{\boldsymbol{x},i} \subset \partial P_{\boldsymbol{v}'_i}$

where $F_{\boldsymbol{x},i}$ denotes the $(d-1)$ -hyperplane face corresponding to vertex \boldsymbol{v}_i^f $_i^J$ in $T_{\boldsymbol{x},i}$, we have $x \notin F_{x,i} \in \Delta_{d-1}(T_{x,i})$. Thus, any sub-simplex that contains x must also contain vertex \boldsymbol{v}_i^f f. That is, $v_i^f \in f'$ for all $i = 0, 1, \ldots, \ell$ since $x \in f'$ is a ℓ' -dimensional subsimplex. Therefore, $f = \text{Conv}(\boldsymbol{v}_0^f)$ $_0^f, \bm v_1^f$ $_{1}^{f}\ldots, \bm{v}_{\ell}^{f}$ \mathcal{L}_{ℓ}^{f}) \subseteq f'. This contradicts the observation [\(3.20\)](#page-14-0). Thus, the claim [\(3.21\)](#page-14-1) is correct. Consequently, $\prod_{i=0}^{\ell} \phi_{v_i^f}(\boldsymbol{x}) = 0$ for all $\mathbf{x} \in \mathcal{S}'$ since $\phi_{\mathbf{v}_i^f}(\mathbf{x}) = 0$ for all $\mathbf{x} \in \partial P_{\mathbf{v}_i^f}$ for any $i = 0, 1, \dots, \ell$. This finishes the proof. \Box

Remark 2 Here, we note that S' can be nonempty if \mathcal{T} is not a vertex-convex mesh. Fig. [3.2](#page-15-0) demonstrates an example on $[0,1]^2$.

Figure 3.2: An example on 2D showing that S' is nonempty. In this example, $P_{v_0^f}$ and $P_{\mathbf{v}_1^f}$ are not convex, and $S' = f' \in \Delta_1(\mathcal{T})$ is a 1-dimensional sub-simplex marked as the red segment in the diagram. Here, we have $f' \subseteq \partial P_{\mathbf{v}_0^f} \cap \partial P_{\mathbf{v}_1^f}$.

Remark 3 For any $T \in \mathcal{T}$ and $T \subseteq P_f$, we note that $b(\boldsymbol{x}) := \left. \prod_{i=0}^{\ell} \phi_{\boldsymbol{v}_i^f}(\boldsymbol{x}) \right|_T =$ $\lambda_0(\bm{x})\lambda_1(\bm{x})\cdots\lambda_\ell(\bm{x})$ (by using Definition [3.1\)](#page-8-1) is the local bubble function as studied in the geometric decomposition of Lagrange finite elements in $[4, 8]$ $[4, 8]$. The main property of this local bubble function is $b(x) = 0$ for any $x \in \partial f$. By gluing these local barycentric coordinate functions $\lambda_i(\boldsymbol{x})$ into linear basis functions $\phi_{\boldsymbol{v}_i^f}(\boldsymbol{x})$, the above lemma presents an important property similar to the local bubble function.

Now, we have the following main theorem about the new global formula for Lagrange basis functions on an arbitrary simplicial mesh.

Theorem 3.2 For any simplicial mesh \mathcal{T} of Ω for the P_k element and interpolation point $p \in \mathcal{X}_k^{\ell}(\mathcal{T})$ for any $\ell = 0, 1, ..., d$, by taking Definition [3.1,](#page-8-1) the basis function $\varphi_{\bm{p}}$ can be written as

$$
\varphi_{\mathbf{p}}(\mathbf{x}) = \frac{1}{\alpha!} \prod_{i=0}^{\ell} \prod_{j=0}^{\alpha_i - 1} \left(k \phi_{\mathbf{v}_i^f}(\mathbf{x}) - j \right), \quad \forall \mathbf{x} \in \Omega,
$$
\n(3.22)

where $f = \text{Conv}(\boldsymbol{v}_0^f)$ $_0^f, \bm v_1^f$ $_{1}^{f},\ldots,\boldsymbol{v}_{\ell}^{f}$ \mathbf{y}_{ℓ}^f) with $\mathbf{p} \in \mathring{f}$ and $\phi_{\mathbf{v}_i^f}$ is the piecewise linear basis function at $\boldsymbol{v}_i^f\in\mathcal{X}_k^0(\mathcal{T})$ regarding the simplicial mesh $\mathcal{T}.$

Proof of Theorem [3.2](#page-16-1) *Proof* First, we can rewrite (3.22) to

$$
\varphi_{p}(\boldsymbol{x}) = \frac{1}{\alpha!} \prod_{i=0}^{\ell} k \phi_{\boldsymbol{v}_{i}^{f}}(\boldsymbol{x}) \times \prod_{i=0}^{\ell} \prod_{j=1}^{\alpha_{i}-1} \left(k \phi_{\boldsymbol{v}_{i}^{f}}(\boldsymbol{x}) - j \right), \quad \forall \boldsymbol{x} \in \Omega \tag{3.23}
$$

since $\alpha_i \geq 1$ for all $i = 0, 1, \ldots, \ell$. Given Lemma [3.5,](#page-14-2) we have $\varphi_p(x) = 0$ for any $\boldsymbol{x} \notin \bigcup_{f \subseteq T \in \mathcal{T}} T = P_f = P_{\boldsymbol{p}}$ since $\boldsymbol{p} \in \mathring{f}$. Furthermore, for any $s \in T_s$ and $\boldsymbol{x} \in T_s$, we have $\phi_{\mathbf{v}_i^f}(\mathbf{x}) = \lambda_{s,i}(\mathbf{x})$ by the definition of $\phi_{\mathbf{v}_i^f}(\mathbf{x})$. This completes the proof. \square

Remark 4 Compared to formula [\(3.7\)](#page-10-0) for vertex-convex meshes, the formulation [\(3.22\)](#page-16-2) for arbitrary simplicial meshes appears more general and unified from the finite element perspective. However, regarding the representation by DNNs, equation [\(3.7\)](#page-10-0) offers a more straightforward construction. Furthermore, it provides a more efficient implementation for Lagrange finite element functions with vertex-convex meshes, which will be discussed in the following section.

4 Representations of Lagrange elements using DNNs with ReLU and ReLU² activation functions

In this section, we demonstrate the use of $\Sigma_{n_{1:L}}^{1,2}$ to represent Lagrange elements on both vertex-convex and arbitrary simplicial meshes by employing the global representation formulas presented in Theorem [3.1](#page-10-3) and Theorem [3.2,](#page-16-1) leveraging two fundamental properties of $\Sigma_{n_{1:L}}^{1,2}$.

Initially, we introduce the following crucial properties of ReLU and $ReLU^2$.

Properties 4.1 For any $(x, y) \in \mathbb{R}^2$, we have

$$
\min\{x, y\} = a_1 \text{ReLU}\left(W\begin{pmatrix} x \\ y \end{pmatrix}\right) \in \Sigma^1_4 \subset \Sigma^{1, 2}_4,
$$

and

$$
xy = a_2 \text{ReLU}^2 \left(W \begin{pmatrix} x \\ y \end{pmatrix} \right) \in \Sigma_4^2 \subset \Sigma_4^{1,2},
$$

where

$$
W = \begin{pmatrix} 1 & 1 \\ -1 & -1 \\ 1 & -1 \\ -1 & 1 \end{pmatrix}
$$

and

$$
a_1 = \frac{1}{2} (1 \ -1 \ -1 \ -1), \quad a_2 = \frac{1}{4} (1 \ 1 \ -1 \ -1).
$$

Proof The first identity holds since

$$
\min\{x, y\} = \frac{x+y}{2} - \frac{|x-y|}{2}
$$

and

$$
x = \text{ReLU}(x) - \text{ReLU}(-x)
$$
 and $|x| = \text{ReLU}(x) + \text{ReLU}(-x)$.

The second identity holds since

$$
xy = \frac{(x+y)^2}{4} - \frac{(x-y)^2}{4}
$$

and

$$
x^2 = \text{ReLU}^2(x) + \text{ReLU}^2(-x).
$$

 \Box

To simplify our presentation, for any $p \in \mathcal{X}_k^{\ell}(\mathcal{T})$ with $0 \leq \ell \leq d$, let us define

$$
\tau(\boldsymbol{p}) = \#I_{\boldsymbol{p}},
$$

as the number of simplices in the mesh $\mathcal T$ that contain the interpolation point $\boldsymbol p$.

Representation of basis functions on vertex-convex meshes. By combining our main result in Theorem [3.1](#page-10-3) and Properties [4.1,](#page-17-0) we have the following lemma for representing basis functions $\varphi_p(x)$ for Lagrange elements on vertex-convex meshes.

Lemma 4.1 For any vertex-convex simplicial mesh \mathcal{T} of Ω for the P_k element and $\boldsymbol{p} \in \mathcal{X}_k^{\ell}(\mathcal{T})$ for some $0 \leq \ell \leq d,$ there exists

$$
\widetilde{\varphi}_{\mathbf{p}}(\mathbf{x}) \in \Sigma_{n_{1:L}}^{1,2},\tag{4.1}
$$

where

$$
L \leq \lceil \log_2(\tau(\mathbf{p})) \rceil + \max \{ \lceil \log_2(\ell+1) \rceil, \lceil \log_2(k-\ell-1) \rceil \} + 3 \tag{4.2}
$$

and

$$
n_j \le 2^{3-j} \max\{\ell+1, k-\ell-1\}\tau(\pmb{p}).\tag{4.3}
$$

for all $j = 1, \ldots, L$ such that $\widetilde{\varphi}_{p}(x) = \varphi_{p}(x)$ for all $x \in \Omega$. Furthermore, $\widetilde{\varphi}_{p}(x) =$ $\varphi_{\bf p}({\bf x})$ for any ${\bf x} \in \mathbb{R}^d$ if ${\bf p} \in \mathring{\Omega}$.

Proof As demonstrated in [\(3.7\)](#page-10-0) from Theorem [3.1,](#page-10-3) we have

$$
\varphi_{\mathbf{p}}(\mathbf{x}) = \frac{1}{\alpha!} \min_{i' \in I_{\mathbf{p}}} \left\{ \prod_{i=0}^{\ell} k \text{ReLU}(\lambda_{i',i}(\mathbf{x})) \right\} \times \prod_{i=0}^{\ell} \prod_{j=1}^{\alpha_{i}-1} \left(k \min_{i' \in I_{\mathbf{p}}} \left\{ \lambda_{i',i}(\mathbf{x}) \right\} - j \right). \tag{4.4}
$$

This formulation allows us to construct a DNN where $\lambda_{i',i}(\boldsymbol{x})$ is the input of the first layer with $\tau(\bm{p}) \times (\ell+1)$ neurons. Subsequent operations such as $\prod_{i=0}^{\ell} k\text{ReLU} \left(\lambda_{i',i}(\bm{x})\right)$, $\min_{i' \in I_p} {\lambda_{i',i}(\boldsymbol{x})}$, $\min_{i' \in I_p} {\{\cdot\}}$, and $\prod_{i=0}^{\ell} \prod_{j=1}^{\alpha_i-1} (\cdot)$ can be implemented utilizing Property [4.1.](#page-17-0) Specifically, we have

$$
\min_{i' \in I_p} \left\{ \prod_{i=0}^{\ell} k \text{ReLU} \left(\lambda_{i',i}(\boldsymbol{x}) \right) \right\} \in \Sigma_{n_{1:L_1}}^{1,2},\tag{4.5}
$$

where $L_1 \leq \lceil \log_2(\tau(\mathbf{p})) \rceil + \lceil \log_2(\ell+1) \rceil + 1$ and $n_j \leq 2^{3-j}(k-\ell-1)\tau(\mathbf{p})$, and

$$
\prod_{i=0}^{\ell} \prod_{j=1}^{\alpha_i - 1} \min_{i' \in I_p} \{ k \lambda_{i',i}(\boldsymbol{x}) - j \} \in \Sigma_{n_1, L_2}^{1, 2},
$$
\n(4.6)

where $L_2 \leq \lceil \log_2(\tau(\mathbf{p})) \rceil + \lceil \log_2(k - \ell - 1) \rceil + 1$ and $n_j \leq 2^{3-j}(\ell+1)\tau(\mathbf{p})$. By incorporating an additional hidden layer to combine (multiply) these two components, we establish that

$$
\widetilde{\varphi}_{\mathbf{p}}(\mathbf{x}) \in \Sigma_{n_{1:L}}^{1,2},\tag{4.7}
$$

where $L \leq \lceil \log_2(\tau(\mathbf{p})) \rceil + \max \{ \lceil \log_2(\ell+1) \rceil, \lceil \log_2(k-\ell-1) \rceil \} + 3$ and $n_j \leq 2^{3-j} \max\{\ell+1\}$ $1, k - \ell - 1$ } $\tau(\mathbf{p}).$

Representation of basis functions on arbitrary simplicial meshes. For general simplicial mesh $\mathcal T$ that may not be vertex-convex, we first introduce how to use $\Sigma_{n_{1:L}}^{1,2}$ to represent the basis functions for linear finite elements on \mathcal{T} .

Theorem 4.1 ([\[24\]](#page-27-10), Theorem 4.3) For any simplicial mesh \mathcal{T} of Ω , $\boldsymbol{v} \in \mathcal{X}_{k}^{0}(\mathcal{T}) =$ $\Delta_0(\mathcal{T})$, and ϕ_v the basis function of linear finite element at v, there exist

$$
\widetilde{\phi}_{\boldsymbol{v}}(\boldsymbol{x}) \in \Sigma_{n_{1:L}}^1 \subset \Sigma_{n_{1:L}}^{1,2} \tag{4.8}
$$

where $L \leq \lceil \log_2(\tau(\boldsymbol{v})) \rceil + \lceil \log_2(d+1) \rceil + 7$ and $n_j \leq 2^{3-j}(d+1)\tau(\boldsymbol{v})$ for all $j =$ $1,\ldots,L$ such that $\phi_{\bm{v}}(\bm{x}) = \phi_{\bm{v}}(\bm{x})$ for all $\bm{x} \in \Omega$. Furthermore, $\phi_{\bm{v}}(\bm{x}) = \phi_{\bm{v}}(\bm{x})$ for any $\boldsymbol{x} \in \mathbb{R}^d$ if $\boldsymbol{v} \in \mathring{\Omega}$.

Remark 5 Theorem [4.1](#page-19-0) is based on a main theorem in [\[17\]](#page-26-2) about the representation of the linear basis function $\phi_{\boldsymbol{v}}(\boldsymbol{x})$ by ReLU DNNs when $P_{\boldsymbol{v}}$ is convex for $\boldsymbol{v} \in \mathcal{X}_k^0(\mathcal{T})$. The key step is the overlapping decomposition $P_v = \bigcup_{i \in I_v} P_{v,i}$, where $v \in \mathring{P}_{v,i}$ and $P_{v,i}$ is a simplex consisting of $d+1$ simplices. That is, we can understand $P_{v,i}$ as local vertex-convex meshes of **v** for all $i \in I_v$. Then, by using the result in [\[17](#page-26-2)], we can construct ReLU DNNs $\phi_{v,i}(x)$ for all $i \in I_v$ such that $\phi_{v,i}(x)$ is the linear basis function in terms of the local vertex-convex meshes $P_{v,i}$. Finally, one can have $\phi_{\boldsymbol{v}}(\boldsymbol{x}) = \max_{i \in I_{\boldsymbol{v}}} {\{\phi_{\boldsymbol{v},i}(\boldsymbol{x})\}}$. More details can be found in [\[24\]](#page-27-10).

By combining our main result in Theorem [3.2](#page-16-1) with Theorem [4.1](#page-19-0) and Proper-ties [4.1,](#page-17-0) we have the following lemma for representing basis functions $\varphi_p(x)$ for Lagrange elements on arbitrary simplicial meshes.

Lemma 4.2 For any simplicial mesh \mathcal{T} of Ω for the P_k element and $p \in \mathcal{X}_k^{\ell}(\mathcal{T})$ for some $0 \leq \ell \leq d$, let us assume $f = \text{Conv}(\boldsymbol{v}_0^f)$ $_0^f, \bm v_1^f$ $_{1}^{f},\ldots,\boldsymbol{v}_{\ell}^{f}$ ℓ) and $p \in \mathring{f}$ and $\phi_{\mathbf{v}_{i}^{f}}$. Then, there exists

$$
\widetilde{\varphi}_{\mathbf{p}}(\mathbf{x}) \in \Sigma^{1,2}_{n_{1:L}},\tag{4.9}
$$

where

$$
L \le \max_{i=0,1,\dots,\ell} \left[\log_2(\tau(\boldsymbol{v}_i^f)) \right] + \left[\log_2(d+1) \right] + \left[\log_2(k) + 7 \right] \tag{4.10}
$$

and

$$
n_j \le 2^{3-j}(d+1) \sum_{i=0,1,\dots,\ell} \tau(\mathbf{v}_i^f), \tag{4.11}
$$

for all $j = 1, ..., L$ such that $\widetilde{\varphi}_{p}(x) = \varphi_{p}(x)$ for all $x \in \Omega$. Furthermore, $\widetilde{\varphi}_{p}(x) =$ $\varphi_{\bf p}({\bf x})$ for any ${\bf x} \in \mathbb{R}^d$ if ${\bf p} \in \mathring{\Omega}$.

Proof As demonstrated in [\(3.22\)](#page-16-2) from Theorem [3.2,](#page-16-1) we have

$$
\varphi_{\mathbf{p}}(\mathbf{x}) = \frac{1}{\alpha!} \prod_{i=0}^{\ell} \prod_{j=0}^{\alpha_i - 1} \left(k \phi_{\mathbf{v}_i^f}(\mathbf{x}) - j \right) \quad \forall \mathbf{x} \in \Omega.
$$

Given that $\phi_{\mathbf{v}_i^f}$ can be represented by $\Sigma^1_{n_{1:L}} \subset \Sigma^{1,2}_{n_{1:L}}$ and $\Sigma^{1,2}_{n_{1:L}}$ can handle the multiplication operation, the above formulation allows us to construct a DNN function in $\Sigma^{1,2}_{n_{1:L}}$ to recover $\varphi_p(x)$ by first representing $\phi_{v_i^f}(x)$. According to Theorem [4.1,](#page-19-0) $\phi_{\mathbf{v}_i^f}(\mathbf{x})$ can be reproduced with $\lceil \log_2(\tau(\mathbf{v})) \rceil + \lceil \log_2(d+1) \rceil + 7$ hidden layers and $2^{3-j}(d+1)\tau(v)$ neurons in the j-th hidden layer. Noting that there are extra $\sum_{j=0}^{\ell} \alpha_i = k$ multiplications from $\phi_{\mathbf{v}_i^f}(\mathbf{x})$ to $\varphi_{\mathbf{p}}(\mathbf{x})$, we require only an additional $\lceil \log_2(k) \rceil$ hidden layers to recover these multiplication operations and to parallelly concatenate hidden layers of different $\phi_{\mathbf{v}_i^f}(\mathbf{x})$. Thus, we can establish bounds for L and n_j . . The contract of \Box

Representation of Lagrange finite element functions using $\Sigma_{n_{1:L}}^{1,2}$. With the representation results for basis functions in place, we can state our main theorem as follows.

Theorem 4.2 For any P_k Lagrange finite element function $u_h(x)$ on a simplicial mesh $\mathcal T$ of Ω , there is

$$
\widetilde{u}_h(\boldsymbol{x}) \in \Sigma_{n_{1:L}}^{1,2},\tag{4.12}
$$

such that $\widetilde{u}_h(\boldsymbol{x}) = u_h(\boldsymbol{x})$ for any $\boldsymbol{x} \in \Omega$. More precisely, we have the following bounds of $\Sigma^{1,2}_{n_{1:L}}$:

If $\mathcal T$ is vertex-convex,

$$
L \le \sum_{\ell=0}^{d} \eta_{\ell} \left(\lceil \log_2(\tau_{\ell}) \rceil + \max \left\{ \lceil \log_2(\ell+1) \rceil, \lceil \log_2(k-\ell-1) \rceil \right\} + 3 \right) \tag{4.13}
$$

and

$$
\max_{j} n_j \le 4 \max_{\ell} \{ \max\{\ell+1, k-\ell-1\} \tau_{\ell} \} + 2d + 1; \tag{4.14}
$$

If $\mathcal T$ is arbitrary,

$$
L \leq (\lceil \log_2(\tau_0) \rceil + \lceil \log_2(d+1) \rceil + \lceil \log_2(k) \rceil + 7) \sum_{\ell=0}^d \eta_\ell \tag{4.15}
$$

and

$$
\max_{j} n_j \le 4(d+1)\min\{d, k+1\}\tau_0 + 2d + 1,\tag{4.16}
$$

where $\tau_{\ell} := \max_{\mathbf{p} \in \mathcal{X}_{k}^{\ell}(\mathcal{T})} \tau(\mathbf{p})$ and $\eta_{\ell} = \# \mathcal{X}_{k}^{\ell}(\mathcal{T})$ for all $\ell = 0, 1, ..., d$.

To simplify the proof, we introduce the following revised neural network architecture:

$$
\begin{cases}\n\widetilde{u}^{0}(\boldsymbol{x}) = \boldsymbol{x}, \\
[\widetilde{u}^{j}(\boldsymbol{x})]_{i} = \sigma_{i}^{j} \left(\left[\widetilde{W}^{\ell} \left(\widetilde{u}^{j-1}(\boldsymbol{x}), \boldsymbol{x} \right) + \widetilde{b}^{j} \right]_{i} \right), \quad i = 1, \cdots, n_{j}, \ j = 1, \cdots, L,\n\tag{4.17} \\
\widetilde{u}(\boldsymbol{x}) = \widetilde{W}^{L+1} \left(\widetilde{u}^{0}, \widetilde{u}^{1}, \cdots, \widetilde{u}^{L} \right) + \widetilde{b}^{L+1},\n\end{cases}
$$

and denote it as $\Sigma_{n_1,L}^{1,2}$ when σ_i^{ℓ} shares the same definition as in [\(1.3\)](#page-2-1). This type of network architecture is widely studied in the literature, such as in [\[12](#page-25-2), [16](#page-26-3)], where $\sigma_i^j =$ ReLU. Here, we generalize this structure to the learnable ReLU-ReLU^2 activation function while preserving the following important properties.

Properties 4.2 For $\tilde{\Sigma}_{n_{1:L}}^{1,2}$ on \mathbb{R}^{d} , we have:

$$
\widetilde{\Sigma}_{n_{1:L_1}}^{1,2} + \widetilde{\Sigma}_{m_{1:L_2}}^{1,2} \subseteq \widetilde{\Sigma}_{(n_{1:L_1}, m_{1:L_2})}^{1,2}, \tag{4.18}
$$

where $(n_{1:L_1}, m_{1:L_2})$ equals $(n_1, \dots, n_{L_1}, m_1, \dots, m_{L_2});$

2.

1.

$$
\Sigma_{\tilde{n}_{1:L}}^{1,2} \subseteq \tilde{\Sigma}_{\tilde{n}_{1:L}}^{1,2} \subseteq \Sigma_{n_{1:L}}^{1,2},\tag{4.19}
$$

provided $n_j \geq \tilde{n}_j + 2d + 1$ for all j.

Proof The first property is a direct consequence of the definition of $\Sigma_{n_{1:L}}^{1,2}$. The proof follows the same lines as in [\[16](#page-26-3)]. The second property is valid because σ_i^j $\frac{j}{i}$ can replicate the identity function, which can be demonstrated using a similar argument to that in [\[16](#page-26-3)].

Proof of Themorem [4.2](#page-20-0) *Proof* For any $u_h(x)$, it can be written as

$$
u_h(\boldsymbol{x}) = \sum_{\boldsymbol{p} \in \mathcal{X}_k(\mathcal{T})} u_h(\boldsymbol{p}) \varphi_{\boldsymbol{p}}(\boldsymbol{x}) = \sum_{\ell=0}^d \sum_{\boldsymbol{p} \in \mathcal{X}_k(\mathcal{T})} u_h(\boldsymbol{p}) \varphi_{\boldsymbol{p}}(\boldsymbol{x}). \tag{4.20}
$$

Given Properties [4.2](#page-21-0) and the estimates in Lemma [4.1](#page-18-0) and Lemma [4.2,](#page-19-1) we have

$$
\widetilde{u}_h(\boldsymbol{x}) = \sum_{\ell=0}^d \sum_{\boldsymbol{p} \in \mathcal{X}_k^{\ell}(\mathcal{T})} u_h(\boldsymbol{p}) \widetilde{\varphi}_{\boldsymbol{p}}(\boldsymbol{x}) \in \widetilde{\Sigma}_{\tilde{n}_{1:L}}^{1,2} \subset \Sigma_{n_{1:L}}^{1,2} \tag{4.21}
$$

where L and n_i are bounded as above.

Remark 6 For simplicial meshes, we observe that $\tau_{\ell} \geq \tau_{\ell'}$ for any $0 \leq \ell \leq \ell' \leq$ d. In particular, τ_0 is typically the largest and often scales as the exponent (or even factorial) of d. Thus, we note that the depths and widths required for vertexconvex meshes are significantly smaller than those for arbitrary simplicial meshes, as demonstrated in Theorem 4.2 .

5 An application of the representation result for approximation properties of $\Sigma^{1,2}_{n_1}$ $n_{1:L}$

In this section, we present an approximation result for $\Sigma_{n_{1:L}}^{1,2}$ utilizing representation theory on a special vertex-convex mesh. We assume that $u \in W^{q+1,p}(\Omega)$ for some $k \geq 1$ and $p \in [2,\infty]$ with $\Omega = [0,1]^d$. The space $W^{k+1,p}(\Omega)$ denotes the standard Sobolev space [\[1\]](#page-25-9). We establish the following approximation property of $\Sigma_{n_1,L}^{1,2}$.

Theorem 5.1 For any function $u \in W^{k+1,p}(\Omega)$ with $p \in [2,\infty]$ and $N > 0$, we obtain the approximation rate

$$
\inf_{\widetilde{u}\in\Sigma_{n_{1:L}}^{1,2}}\|u-\widetilde{u}\|_{W^{s,p}(\Omega)}=\|u\|_{W^{k+1,p}(\Omega)}\mathcal{O}\left(N^{-\frac{k+1-s}{d}}\right),\tag{5.1}
$$

for any $s = 0, 1$. Here, $L \leq C_1 \log(k) k^d N$ and $n_j \leq C_2 k$, where C_1 and C_2 depend only on d.

Uniform simplex mesh on $[0, 1]^d$ with size $h = \frac{1}{m}$ To prove Theorem [5.1,](#page-22-1) we introduce some special simplices with size $h = \frac{1}{m}$ $\frac{1}{m}$ for some $m \in \mathbb{N}^+$ in $[0, 1]^d$ as

$$
\Delta_{\boldsymbol{v},\rho}^h := \left\{ \boldsymbol{x} \in [0,1]^d : 0 \le x_{\rho(1)} - v_{\rho(1)} \le \cdots \le x_{\rho(d)} - v_{\rho(d)} \le h \right\} \tag{5.2}
$$

for any $\boldsymbol{v} \in \mathbf{Z}_h^d := \{0, h, 2h, \cdots, (m-1)h, 1\}^d$ and any permutation ρ of $\{1, 2, \cdots, d\}$. The corresponding uniform simplicial mesh \mathcal{T}_h^d with size h on $[0,1]^d$ is

$$
\mathcal{T}_h^d := \left\{ \Delta_{\boldsymbol{v},\rho}^h : \boldsymbol{v} \in \mathbf{Z}_h^d, \rho \text{ is a permutation of } \{1,2,\cdots,d\} \right\} = [0,1]^d. \tag{5.3}
$$

This particular mesh, \mathcal{T}_h^d , is known as the Freudenthal triangulation [\[14](#page-26-8)] and is also referred to as the Kuhn partition, as described in the paper [\[20\]](#page-26-9). This triangulation can also result from decomposing $[0, 1]^d$ with hyperplanes defined by $x_i - x_j = kh$ and $x_i = sh$ for any $1 \leq i, j \leq d$ and $k, s = 0, \dots, m$. Hence, the set of all vertices of \mathcal{T}_h^d is \mathbf{Z}_h^d . As demonstrated in [\[45\]](#page-28-9), supp (φ_v) is convex for any vertex $v \in \mathbf{Z}_h^d = \mathcal{X}_k^0(\mathcal{T}_h^d)$. That is, \mathcal{T}_h^d is vertex-convex.

Proof of Theorem [5.1](#page-22-1) Proof Let V_h^k denote the space of k-th order Lagrange elements on the uniform mesh \mathcal{T}_h^d of $[0,1]^d$ with size $h=\frac{1}{m}$ $\frac{1}{m}$. Given the estimate for the vertex-convex mesh in Theorem [4.2](#page-20-0) and the self-similarity of \mathcal{T}_h^d , we observe that τ_{ℓ} for all $\ell = 0, 1, \ldots, d$ are determined only by the dimension d and is independent of the mesh size h . Hence, we have

$$
\max_{j} n_j \le C_W k,\tag{5.4}
$$

where C_W is a constant depending only on d. Additionally, we find that

$$
\lceil \log_2(\tau_\ell) \rceil + \max_{1 \le \ell \le \min\{k, d+1\}} \left\{ \lceil \log_2(\ell) \rceil, \lceil \log_2(k - \ell - 1) \rceil \right\} + 3
$$

$$
\le \lceil \log_2(\tau_\ell) \rceil + \max \left\{ \lceil \log_2(d+1) \rceil, \lceil \log_2(k) \rceil \right\} \le C_L \log(k) + 3,
$$
 (5.5)

where C_L is a constant dependent only on d. Consequently, we obtain

$$
L \le C_L \log(k) \sum_{\ell=0}^d \eta_\ell,\tag{5.6}
$$

noting that $\sum_{\ell=0}^d \eta_\ell$ is the total number of degrees of freedom for V_h^k , which can be bounded by

$$
\sum_{\ell=0}^{d} \eta_{\ell} = \dim(V_h^k) \le C_T k^d m^d,\tag{5.7}
$$

where C_T depends only on d. As a result, we have

$$
V_h^k \subseteq \Sigma_{n_{1:L}}^{1,2},\tag{5.8}
$$

with $L \leq C_1 \log(k) k^d m^d$ and $n_j \leq C_2 k$, where C_1 and C_2 are constants depending only on d.

Utilizing the classical approximation results of V_h^k for functions in $W^{k+1,p}(\Omega)$, as shown in $[11, 38]$ $[11, 38]$, we have

$$
\inf_{\widetilde{u}\in\Sigma_{n_{1:L}}^{1,2}}\|u-\widetilde{u}\|_{W^{k,p}(\Omega)}\leq \inf_{\widetilde{u}\in V_h^k}\|u-\widetilde{u}\|_{W^{s,p}(\Omega)}=\mathcal{O}(\|u\|_{W^{k+1,p}}h^{k+1-s}),\tag{5.9}
$$

for any $s = 0, 1$ and $p \in [2, \infty]$, where $L \leq C_1 \log(k) k^d m^d$ and $n_j \leq C_2 k$, and C_1 and C_2 depend only on d. The proof is completed by noting that $h = \frac{1}{n}$ $\frac{1}{m}$ and substituting m^d with N. **Remark 7** In the approximation estimate given by (5.1) , the case for $s = 0$ and $k = 1$, that is, the L^p estimate for linear finite element interpolation, has already been established in $[17, 45]$ $[17, 45]$. By incorporating bit-extraction techniques, the L^p approximation rate for ReLU DNNs (i.e., $k = 1$) can be enhanced to $N^{-\frac{2}{d}}$ for functions in $C(\Omega)$, as shown in [\[45,](#page-28-9) [36](#page-28-11)], and to $N^{-\frac{2k}{d}}$ for functions in $C^k(\Omega)$, as demonstrated in [\[25\]](#page-27-7). Notably, the $W^{s,p}$ estimate for $s=1$ is first introduced in this work.

6 Concluding remarks

In conclusion, this paper validates the capability of $\Sigma_{n_1,\mathcal{L}}^{1,2}$ to represent Lagrange finite element functions across dimensions and orders, leveraging two novel formulations of global basis functions on both vertex-convex and arbitrary simplicial meshes, geometric decomposition, and insights from high-dimensional simplicial meshes, barycentric functions, and multiplication of linear basis functions. This framework not only clarifies the basis function representations but also encompasses the full scope of Lagrange finite element functions with efficient parameterizations. Moreover, our investigation into the uniform simplex mesh and the associated approximation properties has yielded a straightforward proof for the approximation result for $\Sigma_{n_1,\iota}^{1,2}$.

Additionally, this work opens the door to several interesting mathematical inquiries in the domains of deep neural networks and finite element methods. A pertinent question is whether $\Sigma_{n_1,L}^{1,2}$ with $\mathcal{O}(N)$ parameters can effectively represent Lagrange finite element functions with N degrees of freedom. Efforts to improve the approximation rate could also involve combining the representation results with bit-extraction techniques, as shown for linear finite elements in [\[45](#page-28-9)]. Furthermore, the expressiveness of $\Sigma_{n_1,L}^{1,2}$ could potentially be generalized to encompass a wider array of finite elements, such as those from finite element exterior calculus [\[3](#page-25-7), [2](#page-25-10), [24\]](#page-27-10), elements with higher smoothness $[19, 9]$ $[19, 9]$ $[19, 9]$, *h-p* adaptive elements $[6]$, and even nonconforming elements [\[38\]](#page-28-10). On the other hand, in contrast to traditional methods that rely on domain partitioning and function expression via bases and degrees of freedom, our approach employs deep neural networks for finite element function representation within a cohesive framework. This innovation could revolutionize finite element implementation and inspire new "adaptive" methods for solving PDEs, as shown in the study of linear finite elements in one dimension in [\[17](#page-26-2)].

References

- [1] Robert A Adams and John JF Fournier. Sobolev spaces. Elsevier, 2003.
- [2] Douglas Arnold, Richard Falk, and Ragnar Winther. Finite element exterior calculus: from hodge theory to numerical stability. Bulletin of the American mathematical society, 47(2):281–354, 2010.
- [3] Douglas N Arnold, Richard S Falk, and Ragnar Winther. Finite element exterior calculus, homological techniques, and applications. Acta numerica, 15:1–155, 2006.
- [4] Douglas N Arnold, Richard S Falk, and Ragnar Winther. Geometric decompositions and local bases for spaces of finite element differential forms. Computer Methods in Applied Mechanics and Engineering, 198(21-26):1660–1672, 2009.
- [5] Raman Arora, Amitabh Basu, Poorya Mianjy, and Anirbit Mukherjee. Understanding deep neural networks with rectified linear units. In International Conference on Learning Representations, 2018.
- [6] Ivo Babuˇska and Manil Suri. The p and h-p versions of the finite element method, basic principles and properties. SIAM review, 36(4):578–632, 1994.
- [7] John Burkardt. The finite element basis for simplices in arbitrary dimensions. Technical report, Technical Report, Florida State University, Department of Scientific Computing, 2013.
- [8] Chunyu Chen, Long Chen, Xuehai Huang, and Huayi Wei. Geometric decomposition and efficient implementation of high order face and edge elements. $arXiv$ preprint arXiv:2309.13843, 2023.
- [9] Long Chen and Xuehai Huang. Geometric decompositions of the simplicial lattice and smooth finite elements in arbitrary dimension. arXiv preprint arXiv:2111.10712, 2021.
- [10] Qipin Chen, Wenrui Hao, and Juncai He. Power series expansion neural network. Journal of Computational Science, 59:101552, 2022.
- [11] Philippe G Ciarlet. The finite element method for elliptic problems. SIAM, 2002.
- [12] Ingrid Daubechies, Ronald DeVore, Simon Foucart, Boris Hanin, and Guergana Petrova. Nonlinear approximation and (deep) relu networks. Constructive Approximation, 55(1):127–172, 2022.
- [13] Weinan E and Bin Yu. The deep ritz method: a deep learning-based numerical algorithm for solving variational problems. Communications in Mathematics and Statistics, 6(1):1–12, 2018.
- [14] Hans Freudenthal. Simplizialzerlegungen von beschrankter flachheit. Annals of Mathematics, pages 580–582, 1942.
- [15] Juncai He. On the optimal expressive power of relu dnns and its application in approximation with kolmogorov superposition theorem. *arXiv preprint* arXiv:2308.05509, 2023.
- [16] Juncai He, Lin Li, and Jinchao Xu. Relu deep neural networks from the hierarchical basis perspective. Computers $\mathcal{C}_{\mathcal{A}}$ Mathematics with Applications, 120:105– 114, 2022.
- [17] Juncai He, Lin Li, Jinchao Xu, and Chunyue Zheng. Relu deep neural networks and linear finite elements. Journal of Computational Mathematics, 38(3):502– 527, 2020.
- [18] Juncai He, Tong Mao, and Jinchao Xu. Expressivity and approximation properties of deep neural networks with reluk activation. $arXiv$ preprint arXiv:2312.16483, 2023.
- [19] Jun Hu, Ting Lin, and Qingyu Wu. A construction of c r conforming finite element spaces in any dimension. Foundations of Computational Mathematics, pages 1–37, 2023.
- [20] Harold W Kuhn. Some combinatorial lemmas in topology. IBM Journal of research and development, 4(5):518–524, 1960.
- [21] Isaac E Lagaris, Aristidis Likas, and Dimitrios I Fotiadis. Artificial neural networks for solving ordinary and partial differential equations. IEEE transactions on neural networks, 9(5):987–1000, 1998.
- [22] Bo Li, Shanshan Tang, and Haijun Yu. Better approximations of high dimensional smooth functions by deep neural networks with rectified power units. Communications in Computational Physics, 27(2):379–411, 2019.
- [23] Senwei Liang, Liyao Lyu, Chunmei Wang, and Haizhao Yang. Reproducing activation function for deep learning. arXiv preprint arXiv:2101.04844, 2021.
- [24] Marcello Longo, Joost AA Opschoor, Nico Disch, Christoph Schwab, and Jakob Zech. De rham compatible deep neural network fem. Neural Networks, 2023.
- [25] Jianfeng Lu, Zuowei Shen, Haizhao Yang, and Shijun Zhang. Deep network approximation for smooth functions. SIAM Journal on Mathematical Analysis, 53(5):5465–5506, 2021.
- [26] Zhou Lu, Hongming Pu, Feicheng Wang, Zhiqiang Hu, and Liwei Wang. The expressive power of neural networks: A view from the width. In Advances in Neural Information Processing Systems, pages 6231–6239, 2017.
- [27] Carlo Marcati, Joost AA Opschoor, Philipp C Petersen, and Christoph Schwab. Exponential relu neural network approximation rates for point and edge singularities. Foundations of Computational Mathematics, 23(3):1043–1127, 2023.
- [28] Mohammad Mehrabi, Aslan Tchamkerten, and Mansoor Yousefi. Bounds on the approximation power of feedforward neural networks. In International Conference on Machine Learning, pages 3453–3461. PMLR, 2018.
- [29] Hrushikesh N Mhaskar and Tomaso Poggio. Deep vs. shallow networks: An approximation theory perspective. Analysis and Applications, 14(06):829–848, 2016.
- [30] Hadrien Montanelli, Haizhao Yang, and Qiang Du. Deep relu networks overcome the curse of dimensionality for generalized bandlimited functions. Journal of Computational Mathematics, 39(6), 2021.
- [31] Guido F Montufar, Razvan Pascanu, Kyunghyun Cho, and Yoshua Bengio. On the number of linear regions of deep neural networks. In Advances in Neural Information Processing Systems, pages 2924–2932, 2014.
- [32] Vinod Nair and Geoffrey E Hinton. Rectified linear units improve restricted boltzmann machines. In Proceedings of the 27th international conference on machine learning (ICML-10), pages 807–814, 2010.
- [33] Joost AA Opschoor, Philipp C Petersen, and Christoph Schwab. Deep relu networks and high-order finite element methods. Analysis and Applications, pages 1–56, 2020.
- [34] Joost AA Opschoor, Ch Schwab, and Jakob Zech. Exponential relu dnn expression of holomorphic maps in high dimension. Constructive Approximation, 55(1):537–582, 2022.
- [35] Maziar Raissi, Paris Perdikaris, and George E Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. Journal of Computational physics, 378:686–707, 2019.
- [36] Zuowei Shen, Haizhao Yang, and Shijun Zhang. Deep network approximation characterized by number of neurons. Communications in Computational Physics, 28(5):1768–1811, 2020.
- [37] Zuowei Shen, Haizhao Yang, and Shijun Zhang. Neural network approximation: Three hidden layers are enough. Neural Networks, 141:160–173, 2021.
- [38] Zhongci Shi and Ming Wang. Finite Element Methods. Science Press, Beijing, China, 2013.
- [39] Jonathan W Siegel, Qingguo Hong, Xianlin Jin, Wenrui Hao, and Jinchao Xu. Greedy training algorithms for neural networks and applications to pdes. Journal of Computational Physics, 484:112084, 2023.
- $[40]$ Matus Telgarsky. Representation benefits of deep feedforward networks. $arXiv$ preprint arXiv:1509.08101, 2015.
- [41] Jinchao Xu. Finite neuron method and convergence analysis. Communications in Computational Physics, 28(5), 2020.
- [42] Yahong Yang, Yue Wu, Haizhao Yang, and Yang Xiang. Nearly optimal approximation rates for deep super relu networks on sobolev spaces. *arXiv preprint* arXiv:2310.10766, 2023.
- [43] Yahong Yang, Haizhao Yang, and Yang Xiang. Nearly optimal vc-dimension and pseudo-dimension bounds for deep neural network derivatives. arXiv preprint arXiv:2305.08466, 2023.
- [44] Dmitry Yarotsky. Error bounds for approximations with deep relu networks. Neural Networks, 94:103–114, 2017.
- [45] Dmitry Yarotsky. Optimal approximation of continuous functions by very deep relu networks. In Conference on learning theory, pages 639–649. PMLR, 2018.
- [46] Dmitry Yarotsky. Elementary superexpressive activations. In International Conference on Machine Learning, pages 11932–11940. PMLR, 2021.