

COMPLETE SPACE-LIKE SELF-EXPANDERS IN THE MINKOVSKI SPACE

ZHI LI AND GUOXIN WEI

ABSTRACT. It is our purpose to study complete space-like self-expanders in the Minkovski space. By use of maximum principle of Omori-Yau type, we can obtain the rigidity theorems on n -dimensional complete space-like self-expanders in the Minkovski space \mathbb{R}_1^{n+1} . For complete space-like self-expanders of dimension 2, we give a classification of them under assumption of constant squared norm of the second fundamental form.

1. INTRODUCTION

An n -dimensional smooth immersed hypersurface $x : M^n \rightarrow \mathbb{R}_1^{n+1}$ is called a self-expanders of mean curvature flow if its mean curvature vector \vec{H} satisfies the equation

$$(1.1) \quad \vec{H} = x^\perp,$$

where $\vec{H} = He_{n+1}$, x and e_{n+1} denote the position vector and the inward unit normal field, respectively. If we denote by $\langle \cdot, \cdot \rangle$ the standard inner product on \mathbb{R}_1^{n+1} , then the equation (1.1) is equivalent to

$$(1.2) \quad H = -\langle x, e_{n+1} \rangle, \quad \langle e_{n+1}, e_{n+1} \rangle = -1.$$

Self-expanders are self similar solutions of mean curvature flow, that is, the family of hypersurfaces $x_t = \sqrt{2t}x$, $t > 0$ satisfying the equation of mean curvature flow. Self-expanders are important as they model the behavior of a mean curvature flow coming out of a conical singularity ([3]), and also model the long time behaviors of the flows starting from entire graphs ([11]). There are many classification results about the self-expanders. Ishimura [13] and Halldorsson [12] have classified self-expander curves in \mathbb{R}^2 . In [12], Halldorsson states that each of the complete self-expander curves immersed in \mathbb{R}^2 is convex, properly embedded and asymptotic to the boundary of a cone with vertex at the origin. In 2018, Cheng and Zhou [7] studied some properties of complete properly immersed self-expanders. Recently, Ancari and Cheng [2] studied immersed self-expander hypersurfaces whose mean curvatures have some linear growth controls and obtained some rigidity property of hyperplanes as self-expanders in Euclidean space. More other results about the self-expanders have been done (cf. [4], [8], [9], [10], [14], [15], [16]).

In this paper, by use of maximum principle of Omori-Yau type, we obtain the classification of complete space-like self-expanders in the Minkowski space \mathbb{R}_1^{n+1} .

2020 *Mathematics Subject Classification*: 53E10, 53C40.

Key words and phrases: mean curvature flow, self-expander, maximum principle.

Theorem 1.1. *Let $x : M^n \rightarrow \mathbb{R}_1^{n+1}$ be a complete space-like self-expander with the constant squared norm S of the second fundamental form. If $\inf H^2 > 0$, then $S = 1$ and $x(M^n)$ is the hyperbolic space $\mathbb{H}^n(\sqrt{n})$ or the hyperbolic cylinder $\mathbb{H}^k(\sqrt{k}) \times \mathbb{R}^{n-k}$, $1 \leq k \leq n-1$, where h_{ij} are components of the second fundamental form, $S = \sum_{i,j} (h_{ij})^2$.*

When $n = 2$, we completely classify complete self-expanders with constant squared norm of the second fundamental form in the Minkovski space \mathbb{R}_1^3 . The study is as follows:

Theorem 1.2. *Let $x : M^2 \rightarrow \mathbb{R}_1^3$ be a 2-dimensional complete space-like self-expander in \mathbb{R}_1^3 . If the squared norm S of the second fundamental form is constant, then $x : M^2 \rightarrow \mathbb{R}_1^3$ is one of the space-like affine plane \mathbb{R}^2 through the origin, the hyperbolic cylinder $\mathbb{H}^1(1) \times \mathbb{R}^1$ or the hyperbolic space $\mathbb{H}^2(\sqrt{2})$, where h_{ij} are components of the second fundamental form, $S = \sum_{i,j} (h_{ij})^2$.*

Remark 1.1. *In fact, using the proof method of Theorem 1.1, we can also give the classification of complete self-expanders in the Euclidean space \mathbb{R}^3 under the condition of the constant squared norm of the second fundamental form, and the classification result is only plane \mathbb{R}^2 through the origin, which indicates that the conclusion of Ancari and Cheng [2] still holds without assumption on the scalar curvature.*

2. PRELIMINARIES

Let $x : M^n \rightarrow \mathbb{R}_1^{n+1}$ be an n -dimensional space-like hypersurface of $(n+1)$ -dimensional Minkovski space \mathbb{R}_1^{n+1} . We choose a local orthonormal frame field $\{e_A\}_{A=1}^{n+1}$ in \mathbb{R}_1^{n+1} with dual coframe field $\{\omega_A\}_{A=1}^{n+1}$, such that, restricted to M^n , e_1, \dots, e_n are tangent to M^n . In this paper, we shall make use of the following conventions on the ranges of indices,

$$1 \leq i, j, k, l \leq n.$$

Then we have the structure equations

$$dx = \sum_i \omega_i e_i, \quad de_i = \sum_j \omega_{ij} e_j + \omega_{in+1} e_{n+1},$$

$$de_{n+1} = \omega_{n+1i} e_i, \quad \omega_{n+1i} = \omega_{in+1}$$

where ω_{ij} is the Levi-Civita connection of the hypersurface. By restricting these forms to M , we get

$$(2.1) \quad \omega_{n+1} = 0.$$

Taking exterior derivatives of (2.1), we obtain

$$(2.2) \quad \omega_{in+1} = \sum_j h_{ij} \omega_j, \quad h_{ij} = h_{ji}.$$

$$h = \sum_{i,j} h_{ij} \omega_i \otimes \omega_j, \quad H = \sum_i h_{ii}$$

are called the second fundamental form and the mean curvature of $x : M \rightarrow \mathbb{R}_1^{n+1}$, respectively. Let $S = \sum_{i,j} (h_{ij})^2$ be the squared norm of the second fundamental form

of $x : M \rightarrow \mathbb{R}_1^{n+1}$. The Gauss equations are given by

$$(2.3) \quad R_{ijkl} = -(h_{ik}h_{jl} - h_{il}h_{jk}).$$

Defining the covariant derivative of h_{ij} by

$$(2.4) \quad \sum_k h_{ijk} \omega_k = dh_{ij} + \sum_k h_{ik} \omega_{kj} + \sum_k h_{kj} \omega_{ki},$$

we obtain the Codazzi equations

$$(2.5) \quad h_{ijk} = h_{ikj}.$$

By taking exterior differentiation of (2.4), and defining

$$(2.6) \quad \sum_l h_{ijkl} \omega_l = dh_{ijk} + \sum_l h_{ljk} \omega_{li} + \sum_l h_{ilk} \omega_{lj} + \sum_l h_{ijl} \omega_{lk},$$

we have the following Ricci identities

$$(2.7) \quad h_{ijkl} - h_{ijlk} = \sum_m h_{mj} R_{mikl} + \sum_m h_{im} R_{mjkl}.$$

Defining

$$(2.8) \quad \begin{aligned} \sum_m h_{ijklm} \omega_m &= dh_{ijkl} + \sum_m h_{mjkl} \omega_{mi} + \sum_m h_{imkl} \omega_{mj} + \sum_m h_{ijml} \omega_{mk} \\ &+ \sum_m h_{ijkml} \omega_{ml} \end{aligned}$$

and taking exterior differentiation of (2.6), we get

$$(2.9) \quad h_{ijklq} - h_{ijkql} = \sum_m h_{mjk} R_{milq} + \sum_m h_{imk} R_{mjql} + \sum_m h_{ijm} R_{mklq}.$$

Let V be a tangent C^1 -vector field on M^n and denote by $Ric_V := Ric - \frac{1}{2}L_V g$ the Bakry-Emery Ricci tensor with L_V to be the Lie derivative along the vector field V . Define a differential operator

$$\mathcal{L}f = \Delta f + \langle V, \nabla f \rangle,$$

where Δ and ∇ denote the Laplacian and the gradient operator, respectively. The following maximum principle of Omori-Yau type concerning the operator \mathcal{L} will be used in this paper, which was proved by Chen and Qiu [5].

Lemma 2.1. *Let (M^n, g) be a complete Riemannian manifold, and V is a C^1 vector field on M^n . If the Bakry-Emery Ricci tensor Ric_V is bounded from below, then for any $f \in C^2(M^n)$ bounded from above, there exists a sequence $\{p_t\} \subset M^n$, such that*

$$\lim_{m \rightarrow \infty} f(p_t) = \sup f, \quad \lim_{m \rightarrow \infty} |\nabla f|(p_t) = 0, \quad \lim_{m \rightarrow \infty} \mathcal{L}f(p_t) \leq 0.$$

We next suppose that $x : M \rightarrow \mathbb{R}_1^{n+1}$ is a self-expander, that is, $H = -\langle x, e_{n+1} \rangle$. By a simple calculation, we have the following basic formulas.

$$(2.10) \quad \nabla_i H = -\sum_k h_{ik} \langle x, e_k \rangle, \quad \nabla_j \nabla_i H = -\sum_k h_{ijk} \langle x, e_k \rangle - h_{ij} + H \sum_k h_{ik} h_{kj}.$$

In this paper, we shall always take $V = x^\top$ for any formulas. Using (2.10) and the Ricci identities, we can get the following lemmas.

Lemma 2.2. *Let $x : M^n \rightarrow \mathbb{R}_1^{n+1}$ be an n -dimensional complete space-like self-expander in \mathbb{R}_1^{n+1} . We have*

$$(2.11) \quad \mathcal{L}H = H(S - 1),$$

$$(2.12) \quad \frac{1}{2} \mathcal{L}H^2 = |\nabla H|^2 + H^2(S - 1),$$

$$(2.13) \quad \frac{1}{2} \mathcal{L}S = \sum_{i,j,k} h_{ijk}^2 + S(S - 1).$$

Lemma 2.3. *Let $x : M^n \rightarrow \mathbb{R}_1^{n+1}$ be an n -dimensional complete space-like self-expander in \mathbb{R}_1^{n+1} . If S is constant, we have*

$$(2.14) \quad \begin{aligned} \sum_{i,j,k,l} (h_{ijkl})^2 &= (2 - S) \sum_{i,j,k} (h_{ijk})^2 + 6 \sum_{i,j,k,l,p} h_{ijk} h_{il} h_{jp} h_{klp} \\ &\quad - 3 \sum_{i,j,k,l,p} h_{ijk} h_{ijl} h_{kp} h_{lp}. \end{aligned}$$

In order to use the maximum principle of Omori-Yau type, we need the following lemma.

Lemma 2.4. *Let $x : M^n \rightarrow \mathbb{R}_1^{n+1}$ be an n -dimensional complete space-like self-expander in \mathbb{R}_1^{n+1} . If the squared norm S of the second fundamental form is constant, the Bakry-Emery Ricci tensor Ric_V is bounded from below, where $V = x^\top$.*

Proof. For any unit vector $e \in TM^n$, we can choose a local tangent orthonormal frame field $\{e_i\}_{i=1}^n$ such that $e = e_i$. Then by the definition of self-expander, direct computation gives

$$\begin{aligned} \frac{1}{2} L_{x^\top} g(e_i, e_i) &= \frac{1}{2} x^\top (g(e_i, e_i)) - g([x^\top, e_i], e_i) \\ &= g(\nabla_{e_i} (x - x^\perp), e_i) \\ &= 1 - H g(\nabla_{e_i} e_{n+1}, e_i) \\ &= 1 - H h_{ii}. \end{aligned}$$

Then (2.3) yields

$$\begin{aligned} Ric_{x^\top}(e_i, e_i) &= Ric(e_i, e_i) - \frac{1}{2}L_{x^\top}g(e_i, e_i) \\ &= -(Hh_{ii} - \sum_j h_{ij}^2) - (1 - Hh_{ii}) \\ &= \sum_j h_{ij}^2 - 1 \geq -1. \end{aligned}$$

The proof of the Lemma 2.4 is finished. \square

3. PROOF OF MAIN THEOREMS

For an arbitrary fixed point p in M^n , we always choose a local frame field $\{e_i\}_{i=1}^n$ such that

$$h_{ij} = \lambda_i \delta_{ij}, \quad S = \sum_i \lambda_i^2, \quad H^2 \leq nS,$$

and the equality of the above inequality holds if and only if $\lambda_1 = \lambda_2 = \dots = \lambda_n$. Taking exterior derivative of $S = \sum_{i,j} (h_{ij})^2$, we know that

$$(3.1) \quad \sum_{i,j} h_{ij} h_{ijk} = 0, \quad \sum_{i,j} h_{ij} h_{ijkl} + \sum_{i,j} h_{ijk} h_{ijl} = 0, \quad k, l = 1, \dots, n.$$

Besides, from Ricci identities (2.7), we obtain

$$h_{ijkl} - h_{ijlk} = -(\lambda_i \lambda_j \lambda_k \delta_{il} \delta_{jk} - \lambda_i \lambda_j \lambda_l \delta_{ik} \delta_{jl} + \lambda_i \lambda_j \lambda_k \delta_{ik} \delta_{jl} - \lambda_i \lambda_j \lambda_l \delta_{il} \delta_{jk}).$$

That is,

$$(3.2) \quad h_{iijj} - h_{jjii} = -\lambda_i \lambda_j (\lambda_i - \lambda_j), \quad h_{iikl} - h_{iilk} = 0, \quad k \neq l.$$

In 2015, Cheng and Peng [6] obtain the rigidity theorems on complete self-shrinkers without the assumption on polynomial volume growth. Motivated by the above work, we obtain the following result on self-expanders.

Theorem 3.1. *(Theorem 1.1) Let $x : M^n \rightarrow \mathbb{R}_1^{n+1}$ be a complete space-like self-expander with the constant squared norm S of the second fundamental form. If $\inf H^2 > 0$, then $S = 1$ and $x(M^n)$ is the hyperbolic space $\mathbb{H}^n(\sqrt{n})$ or the hyperbolic cylinder $\mathbb{H}^k(\sqrt{k}) \times \mathbb{R}^{n-k}$, $1 \leq k \leq n-1$.*

Proof. Since $\inf H^2 > 0$, without loss of generality, we can assume that $\inf H > 0$. For the constant S , from (2.13) and the Lemma 2.4, we infer that

$$(3.3) \quad \sum_{i,j,k} h_{ijk}^2 + S(S-1) = 0.$$

and the Bakry-Emery Ricci curvature of $x : M^n \rightarrow \mathbb{R}_1^{n+1}$ is bounded from below. By the maximum principle of Omori-Yau type concerning the operator \mathcal{L} to the function $-H$, there exists a sequence $\{p_t\} \in M^2$ such that

$$0 \leq \lim_{t \rightarrow \infty} \mathcal{L}H(p_t) = \inf H(S-1).$$

It is obvious that $S \geq 1$, and then $S = 1$ and $\sum_{i,j,k} h_{ijk}^2 = 0$ from (3.3). Namely, the second fundamental form of $x(M^n)$ is parallel and H is constant. It follows from (2.10) that $\lambda_i(H\lambda_i - 1) = 0$, $1 \leq i \leq n$. Then we infer that $\lambda_i = 0$ or $\frac{1}{H}$ for any $1 \leq i \leq n$, which indicates that the number of distinct principal curvatures is at most two. If $x(M^n)$ has only one principal curvature, that is, $x(M^n)$ is totally umbilic, then $x(M^n)$ is the hyperbolic space $\mathbb{H}^n(\sqrt{n})$. If $x(M^n)$ has just two distinct constant principal curvatures, by the congruence theorem of Abe, Koike and Yamaguchi [1], $x(M^n)$ is the hyperbolic cylinder $\mathbb{H}^k(\sqrt{k}) \times \mathbb{R}^{n-k}$, $1 \leq k \leq n-1$. \square

In order to prove the main theorem of this paper, we need the following theorem.

Theorem 3.2. *Let $x : M^2 \rightarrow \mathbb{R}_1^3$ be a complete space-like self-expander with the non-zero constant squared norm S of the second fundamental form, then $\inf H^2 > 0$.*

Proof. We will use the proof by contradiction. Suppose $\inf H^2 = 0$, there exists a sequence $\{p_t\}$ in M^2 such that

$$\lim_{t \rightarrow \infty} H^2(p_t) = \inf H^2 = \bar{H}^2 = 0.$$

By (2.13), (2.14) and S being constant, we know that $\{h_{ij}(p_t)\}$, $\{h_{ijk}(p_t)\}$ and $\{h_{ijkl}(p_t)\}$ are bounded sequences, one can assume

$$\lim_{t \rightarrow \infty} h_{ij}(p_t) = \bar{h}_{ij} = \bar{\lambda}_i \delta_{ij}, \quad \lim_{t \rightarrow \infty} h_{ijk}(p_t) = \bar{h}_{ijk}, \quad \lim_{t \rightarrow \infty} h_{ijkl}(p_t) = \bar{h}_{ijkl}, \quad i, j, k, l = 1, 2.$$

It follows from (2.10) that

$$(3.4) \quad \bar{h}_{11i} + \bar{h}_{22i} = -\bar{\lambda}_i \lim_{t \rightarrow \infty} \langle X, e_i \rangle(p_t), \quad i = 1, 2$$

and

$$(3.5) \quad \bar{h}_{11ij} + \bar{h}_{22ij} = -\sum_k \bar{h}_{ijk} \lim_{t \rightarrow \infty} \langle X, e_k \rangle(p_t) - \bar{\lambda}_i \delta_{ij}, \quad i, j = 1, 2.$$

By use of (3.1), we obtain

$$(3.6) \quad \bar{\lambda}_1 \bar{h}_{11k} + \bar{\lambda}_2 \bar{h}_{22k} = 0$$

and

$$(3.7) \quad \bar{\lambda}_1 \bar{h}_{11kl} + \bar{\lambda}_2 \bar{h}_{22kl} = -\sum_{ij} \bar{h}_{ijk} \bar{h}_{ijl}, \quad k, l = 1, 2.$$

(3.2) yields

$$(3.8) \quad \bar{h}_{1112} - \bar{h}_{1121} = 0, \quad \bar{h}_{2221} - \bar{h}_{2212} = 0.$$

Since $\bar{H}^2 = 0$ and S is non-zero constant, we draw that

$$\bar{\lambda}_1 = -\bar{\lambda}_2 \neq 0, \quad S = 2\bar{\lambda}_1^2.$$

Then (3.4) and (3.6) imply

$$(3.9) \quad \bar{h}_{11k} = \bar{h}_{22k}, \quad \lim_{t \rightarrow \infty} \langle X, e_k \rangle(p_t) = -\frac{2\bar{h}_{11k}}{\bar{\lambda}_k}, \quad k = 1, 2.$$

It follows from (3.5), (3.7) and (3.9) that

$$\begin{aligned}\bar{h}_{1111} + \bar{h}_{2211} &= \frac{2(\bar{h}_{111}^2 - \bar{h}_{112}^2)}{\bar{\lambda}_1} - \bar{\lambda}_1, & \bar{h}_{1122} + \bar{h}_{2222} &= \frac{2(\bar{h}_{111}^2 - \bar{h}_{112}^2)}{\bar{\lambda}_1} + \bar{\lambda}_1, \\ \bar{h}_{1112} + \bar{h}_{2212} &= 0\end{aligned}$$

and

$$\begin{aligned}\bar{\lambda}_1(\bar{h}_{1111} - \bar{h}_{2211}) &= -2\bar{h}_{111}^2 - 2\bar{h}_{112}^2, & \bar{\lambda}_1(\bar{h}_{1122} - \bar{h}_{2222}) &= -2\bar{h}_{111}^2 - 2\bar{h}_{112}^2, \\ \bar{\lambda}_1(\bar{h}_{1112} - \bar{h}_{2212}) &= -4\bar{h}_{111}\bar{h}_{112}.\end{aligned}$$

Thus,

$$(3.10) \quad \begin{cases} \bar{h}_{1111} = -\frac{2\bar{h}_{112}^2}{\bar{\lambda}_1} - \frac{\bar{\lambda}_1}{2}, & \bar{h}_{2211} = \frac{2\bar{h}_{111}^2}{\bar{\lambda}_1} - \frac{\bar{\lambda}_1}{2}, & \bar{h}_{1122} = -\frac{2\bar{h}_{112}^2}{\bar{\lambda}_1} + \frac{\bar{\lambda}_1}{2}, \\ \bar{h}_{2222} = \frac{2\bar{h}_{111}^2}{\bar{\lambda}_1} + \frac{\bar{\lambda}_1}{2}, & \bar{h}_{1112} = -\frac{2\bar{h}_{111}\bar{h}_{112}}{\bar{\lambda}_1}, & \bar{h}_{2212} = \frac{2\bar{h}_{111}\bar{h}_{112}}{\bar{\lambda}_1}. \end{cases}$$

It follows from (2.13) that

$$(3.11) \quad \sum_{i,j,k} h_{ijk}^2 = S(1-S), \quad \sum_{i,j,k} \bar{h}_{ijk}\bar{h}_{ijkl} = 0, \quad l = 1, 2.$$

Specifically,

$$\begin{aligned}\bar{h}_{111}\bar{h}_{1111} + \bar{h}_{222}\bar{h}_{2221} + 3\bar{h}_{112}\bar{h}_{1121} + 3\bar{h}_{221}\bar{h}_{2211} &= 0, \\ \bar{h}_{111}\bar{h}_{1112} + \bar{h}_{222}\bar{h}_{2222} + 3\bar{h}_{112}\bar{h}_{1122} + 3\bar{h}_{221}\bar{h}_{2212} &= 0.\end{aligned}$$

Then by (3.8) and (3.9), we conclude that

$$(3.12) \quad \begin{aligned}\bar{h}_{111}\bar{h}_{1111} + \bar{h}_{112}\bar{h}_{2212} + 3\bar{h}_{112}\bar{h}_{1112} + 3\bar{h}_{111}\bar{h}_{2211} &= 0, \\ \bar{h}_{111}\bar{h}_{1112} + \bar{h}_{112}\bar{h}_{2222} + 3\bar{h}_{112}\bar{h}_{1122} + 3\bar{h}_{111}\bar{h}_{2212} &= 0.\end{aligned}$$

Combining (3.10) with (3.12), we infer

$$(3.13) \quad \bar{h}_{111}(3\bar{h}_{111}^2 - 3\bar{h}_{112}^2 - \bar{\lambda}_1^2) = 0, \quad \bar{h}_{112}(3\bar{h}_{111}^2 - 3\bar{h}_{112}^2 + \bar{\lambda}_1^2) = 0.$$

If $\bar{h}_{111}\bar{h}_{112} \neq 0$, (3.13) implies that $\bar{\lambda}_1^2 = S = 0$. It is a contradiction. If $\bar{h}_{111}\bar{h}_{112} = 0$, we can also obtain contradictions. In fact, we can declare that $\sum_{i,j,k} \bar{h}_{ijk}^2 \neq 0$. Otherwise, we know that $\bar{h}_{ijkl} = 0$ for $i, j, k, l = 1, 2$ from (2.14). Then by (3.10), we infer $\bar{\lambda}_1 = 0$. It is a contradiction. Assuming $\bar{h}_{111} \neq 0$, $\bar{h}_{112} = 0$. From the first equation of (3.11) and (3.13), we draw

$$\bar{h}_{111}^2 = \frac{1}{4}S(1-S), \quad \bar{h}_{111}^2 = \frac{1}{3}\bar{\lambda}_1^2 = \frac{1}{6}S.$$

Then

$$(3.14) \quad S = \frac{1}{3}, \quad \bar{\lambda}_1^2 = \frac{1}{6}, \quad \bar{h}_{111}^2 = \frac{1}{18}.$$

Consequently the following relationship

$$(3.15) \quad \bar{h}_{1111} = -\frac{\bar{\lambda}_1}{2}, \quad \bar{h}_{2211} = \frac{\bar{\lambda}_1}{6}, \quad \bar{h}_{1122} = \frac{\bar{\lambda}_1}{2}, \quad \bar{h}_{2222} = \frac{7\bar{\lambda}_1}{6}, \quad \bar{h}_{1112} = \bar{h}_{2212} = 0$$

can be derived by the simple calculation from (3.10).

It follows from (2.14), (3.14) and (3.15) that

$$\sum_{i,j,k,l} (\bar{h}_{ijkl})^2 = \bar{h}_{1111}^2 + \bar{h}_{2222}^2 + 3\bar{h}_{1122}^2 + 3\bar{h}_{2211}^2 + 4\bar{h}_{1112}^2 + 4\bar{h}_{2212}^2 = \frac{11}{27}$$

and

$$\begin{aligned} \sum_{i,j,k,l} (\bar{h}_{ijkl})^2 &= (2-S) \sum_{i,j,k} (\bar{h}_{ijk})^2 + 6 \sum_{i,j,k} \bar{\lambda}_i \bar{\lambda}_j \bar{h}_{ijk}^2 - 3 \sum_{i,j,k} \bar{\lambda}_k^2 \bar{h}_{ijk}^2 \\ &= 4(2-S)\bar{h}_{111}^2 - 12\bar{\lambda}_1^2 \bar{h}_{111}^2 = \frac{7}{27}. \end{aligned}$$

This is impossible. Assuming $\bar{h}_{111} = 0$, $\bar{h}_{112} \neq 0$. Using the similar methods to also obtain contradictions. The proof of the Theorem 3.2 is thus finished. \square

Proof of Theorem 1.2. If $S = 0$, we know that $x : M^2 \rightarrow \mathbb{R}_1^3$ is a space-like affine plane \mathbb{R}^2 . If $S \neq 0$, it is obvious that either $x(M^2)$ is $\mathbb{H}^1(1) \times \mathbb{R}^1$ or $\mathbb{H}^2(\sqrt{2})$ from the Theorem 3.1 and the Theorem 3.2. \square

Acknowledgements The first author was partially supported by the China Post-doctoral Science Foundation Grant No.2022M711074. The second author was partly supported by grant No.12171164 of NSFC, GDUPS (2018), Guangdong Natural Science Foundation Grant No.2023A1515010510.

REFERENCES

- [1] N. Abe, N. Koike and S. Yamaguchi, Congruence theorems for proper semi-Riemannian hypersurfaces in a real space form. *Yokohama Math. J.*, **35**(1987), no. 1-2, 123-136.
- [2] S. Ancari and X. Cheng, Volume properties and rigidity on self-expanders of mean curvature flow, *Geom. Dedicata*, **216**,(2022), no. 2, Paper No. 24, 25 pp.
- [3] S. Angenent, T. Ilmanen and D. L. Chopp, A computed example of nonuniqueness of mean curvature flow in \mathbb{R}^3 , *Comm. Partial Differential Equations*, **20** (1995), no. 11-12, 1937-1958.
- [4] J. Bernstein and L. Wang, Smooth compactness for spaces of asymptotically conical self-expanders of mean curvature flow, *Int. Math. Res. Not.*, **2021**(2021), no. 12, 9016-9044.
- [5] Q. Chen and H. Qiu, Rigidity of self-shrinkers and translating solitons of mean curvature flows, *Adv. Math.*, **294**(2016), 517-531.
- [6] Q. -M. Cheng and Y. J. Peng, Complete self-shrinkers of the mean curvature flow, *Calc. Var. Partial Differential Equations*, **52** (2015), no. 3-4, 497-506.
- [7] X. Cheng and D. Zhou, Spectral properties and rigidity for self-expanding solutions of the mean curvature flows, *Math. Ann.*, **371**(2018), no. 1-2, 371-389.
- [8] K. Ecker and G.Huisken, Mean curvature evolution of entire graphs, *Ann. Math.*, **130**(1989), no.3, 453-471.
- [9] Q. Ding, Minimal cones and self-expanding solutions for mean curvature flows, *Math. Ann.*, **376**(2020), no. 1-2, 359-405.
- [10] F. T. -H. Fong and P. McGrath, Rotational symmetry of asymptotically conical mean curvature flow self-expanders, *Comm. Anal. Geom.*, **27**(2019), no. 3, 599-618.
- [11] S. -H. Guo, Asymptotic behavior and stability of mean curvature flow with a conical end, *Adv. Math.*, **375** (2020), 107408, 63 pp.
- [12] P. H. Halldorsson, Self-similar solutions to the curve shortening flow, *Trans. Amer. Math. Soc.*, **364** (2012), 5285-5309.
- [13] N. Ishimura, Curvature evolution of plane curves with prescribed opening angle, *Bull Austral. Math. Soc.*, **52** (1995), 287-296.
- [14] K. Smoczyk, Self-expanders of the mean curvature flow, *Vietnam J. Math.*, **49**(2021), 433-445.

- [15] N. Stavrou, Selfsimilar solutions to the mean curvature flow, *J. Reine Angew. Math.*, **499**(1998), 189-198.
- [16] J. M. Xie and J. T. Yu, Convexity of 2-convex translating and expanding solitons to the mean curvature flow in \mathbb{R}^{n+1} , *J. Geom. Anal.*, **33**(2023), no, 8, 252.

ZHI LI,
COLLEGE OF MATHEMATICS AND INFORMATION SCIENCE, HENAN NORMAL UNIVERSITY,
453007, XINXIANG, HENAN, CHINA.
Email address: lizhihnsd@126.com

GUOXIN WEI,
SCHOOL OF MATHEMATICAL SCIENCES, SOUTH CHINA NORMAL UNIVERSITY,
510631, GUANGZHOU, CHINA.
Email address: weiguoxin@tsinghua.org.cn