

# Cluster algebras and monotone Lagrangian tori

Yunhyung Cho, Myungho Kim, Yoosik Kim, and Euiyong Park

**Abstract.** Motivated by recent developments in the construction of Newton–Okounkov bodies and toric degenerations via cluster algebras in [GHKK18, FO20], we consider a family of Newton–Okounkov polytopes of a complex smooth Fano variety  $X$  related by a composition of tropicalized cluster mutations. According to the work of [HK15], the toric degeneration associated with each Newton–Okounkov polytope  $\Delta$  in the family produces a Lagrangian torus fibration of  $X$  over  $\Delta$ . We investigate circumstances in which each Lagrangian torus fibration possesses a monotone Lagrangian torus fiber. We provide a sufficient condition, based on the data of tropical integer points and exchange matrices, for the family of constructed monotone Lagrangian tori to contain infinitely many monotone Lagrangian tori, no two of which are related by any symplectomorphisms. By employing this criterion and exploiting the correspondence between the tropical integer points and the dual canonical basis elements, we generate infinitely many distinct monotone Lagrangian tori on flag manifolds of arbitrary type except in a few cases.

## CONTENTS

1. Introduction	1
2. Newton–Okounkov bodies, toric degenerations, and Lagrangian tori	6
3. $\mathbb{Q}$ -Gorenstein Fano polytopes and Monotone Lagrangian tori	9
4. Infinitely many distinct monotone Lagrangian tori via cluster mutations	17
5. Cluster polytopes for flag manifolds	26
6. Exchange matrix with a large entry between a frozen and unfrozen	34
References	40

## 1. Introduction

### 1.1. Backgrounds

Cluster algebras, introduced by Fomin–Zelevinsky [FZ02], are a class of  $\mathbb{Z}$ -subalgebras of a rational function field with a special combinatorial structure. This algebra possesses a generating set divided into subsets called *clusters*, and these clusters are connected by special combination procedures called *mutations*. The generating set can be recursively constructed from a *seed* consisting of a cluster together and an *exchange matrix*, a skew-symmetrizable matrix that determines mutations. Cluster algebras have naturally appeared in many research areas

---

The research of Y. Cho was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government (MSIT) (NRF-2020R1C1C1A01010972) and (NRF-2020R1A5A1016126).

The research of M. Kim was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government (MSIT) (NRF-2022R1F1A1076214 and NRF-2020R1A5A1016126).

The research of Y. Kim was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government (MSIT) (NRF-2021R1F1A1057739 and NRF-2020R1A5A1016126).

The research of E. Park was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government (MSIT) (RS-2023-00273425 and NRF-2020R1A5A1016126).

including geometry, representation theory, and combinatorics, numerous applications have been studied, and now become an active area of mathematical research.

The algebro-geometric counterpart of a cluster algebra is a *cluster variety*. By gluing split algebraic tori via two different types of mutations, one produces a pair of cluster varieties; an  $\mathcal{A}$ -cluster variety and an  $\mathcal{X}$ -cluster variety. The full Fock–Goncharov (FG) conjecture [FG06] asserts that the ring of regular functions of one cluster variety admits a basis whose elements are parametrized by tropical integer points of the other cluster variety. A foundational work of Gross–Hacking–Keel–Kontsevich (GHKK) [GHKK18] reformulates and resolves the conjecture by constructing a *theta basis*, whose elements correspond to a broken line in a scattering diagram associated with the cluster structure. The pair of cluster varieties is conceived as a mirror pair, see [GHK15b] for a precise formulation. Additionally, GHKK expected that the mirror of a compactified  $\mathcal{A}$ -cluster variety is given by the *disk potential function* on the dual cluster  $\mathcal{X}$ -variety, a generating function for invariants counting holomorphic disks in [CO06, Aur07, FOOO10].

This paper focuses on a situation where the FG conjecture holds and explores an application of the theory of cluster algebras to symplectic topology. To put our work in the above context, we formulate the conjecture on mirror symmetry of the dual pair of cluster varieties more concretely. A recent work of Fujita–Oya [FO20] revealed that a refinement of the Qin’s dominance order in [Qin17] on a lattice for an  $\mathcal{A}$ -cluster variety produces a Newton–Okounkov body  $\Delta$ . It is a rational polytope and generates GHKK’s toric degeneration on a compactification  $X$  of the  $\mathcal{A}$ -cluster variety. The tropical integer points of  $\Delta$  parametrize a theta basis respecting the  $\mathcal{X}$ -cluster structure. Moreover, the polytope  $\Delta$  produces a Lagrangian torus fibration (more precisely, completely integrable system) on  $X$  over  $\Delta$  by Harada–Kaveh [HK15]. In particular, we obtain a Lagrangian torus fibration on (an open dense part of)  $X$  associated with each seed. Conjecturally, by gluing the (Floer theoretical) SYZ mirrors together with disk potential functions of those Lagrangian torus fibrations, one should (partially) recover the dual  $\mathcal{X}$ -cluster variety and the superpotential given by the sum of theta basis elements corresponding to an irreducible component of the anticanonical divisor for  $X$ .

In this paper, we are concerned with Lagrangian torus fibrations on the compactified  $\mathcal{A}$ -cluster variety  $X$ , constructed from a cluster algebra as described above. Specifically, our focus lies on *monotone* Lagrangian tori, refer to Definition 3.1 for a precise definition. However, *not* every toric degeneration can yield a monotone Lagrangian torus. To describe situations where a monotone Lagrangian torus is constructed, we recall some notions. A full-dimensional rational polytope  $\Delta \subset \mathbb{R}^m$  is called  *$\mathbb{Q}$ -Gorenstein Fano* if there exists a vector  $\mathbf{u}_0 \in \mathbb{R}^m$  such that the translated polytope  $\Delta - \mathbf{u}_0$  is a constant multiple of the polar dual of some Fano polytope. Such a point  $\mathbf{u}_0$  uniquely exists and is called the *center* of  $\Delta$ . The following proposition specifies a circumstance in which the toric degeneration produces a monotone Lagrangian torus, see Proposition 3.13 for a more precise statement.

**Proposition A** (Proposition 3.13). *Let  $X$  be a smooth Fano variety equipped with a Kähler form obtained from a very ample line bundle  $\mathcal{L}$ , a positive power of the anticanonical bundle of  $X$ . Suppose that  $X$  has a toric degeneration constructed from a  $\mathbb{Q}$ -Gorenstein Fano Newton–Okounkov polytope  $\Delta$  of  $\mathcal{L}$  such that the central toric variety is normal. Let  $\Phi: X \rightarrow \Delta$  be a Lagrangian torus fibration constructed from this toric degeneration. Then the Lagrangian torus fiber of  $\Phi$  located at the center of  $\Delta$  is a monotone Lagrangian torus.*

**Remark.** Galkin–Mikhalkin in [GM22] proved an analog statement, see Remark 3.14.

The monotonicity condition ensures that disks bounded by a monotone Lagrangian torus  $L$  do *not* pose any serious obstruction to produce mirror spaces via Lagrangian Floer theory of  $L$ . Moreover, each monotone Lagrangian torus  $L$  gives rise to a complex algebraic torus as mirror spaces, which fits into our interest in cluster varieties.

Beyond the aspects of cluster duality and Floer theoretical SYZ mirror symmetry, there are additional motivations for considering monotone Lagrangian tori. One of the versions of mirror symmetry conjecture for Fano varieties asserts that for each  $\mathbb{Q}$ -factorial Fano variety  $X$  having at worst terminal singularities, there exists a Laurent polynomial  $W: (\mathbb{C}^*)^n \rightarrow \mathbb{C}$  called a *weak Landau–Ginzburg mirror* such that its period integral

$$\pi_W(t) := \int_{|x_1|=\dots=|x_n|=1} \frac{1}{1-tW} \Omega, \quad \Omega = \frac{dx_1}{x_1} \wedge \dots \wedge \frac{dx_n}{x_n}$$

is equal to a *regularized quantum period*, i.e., a generating function of gravitational descendant one-pointed Gromov–Witten invariants of  $X$ , see [Prz07, KP11] for instance. In [Ton18], Tonkonog proved that  $W$  can be obtained as the disk potential function of a monotone Lagrangian torus in  $X$ . Thus the mirror symmetry conjecture will be proved if one can succeed in finding a monotone Lagrangian torus in a given Fano manifold. Proposition A extends this result to an arbitrary smooth Fano variety admitting a  $\mathbb{Q}$ -Gorenstein Fano and *normal toric degeneration*.

The notion of monotonicity was originally introduced and used to construct Lagrangian Floer homology by Oh [Oh93] and Biran–Cornea [BC09]. In symplectic topology, it is an interesting problem to construct a new monotone Lagrangian submanifold not related to any pre-existing monotone Lagrangian submanifold through any symplectomorphisms. In [Che96], Chekanov first constructed a monotone Lagrangian torus (called a *Chekanov torus*) that is not Hamiltonian isotopic to any standard product torus in  $\mathbb{R}^{2n}$ . By suitably embedding a Chekanov torus into  $\mathbb{C}P^2$ , one obtains a monotone Lagrangian torus distinct from the Clifford torus, the most standard monotone torus in  $\mathbb{C}P^2$ . Later, in [Via14, Via16], Vianna constructed an infinite family of new monotone tori in  $\mathbb{C}P^2$ , no two of which are related by any symplectomorphisms. Also, in [Aur15], Auroux constructed infinitely many distinct monotone tori with the same monotonicity constant in the Euclidean space  $\mathbb{R}^6$ . The following list attempts to include recent developments of construction exotic tori in different directions, see [Via17, Cas23, GM22, CHW23, Bre23].

One of our motivations is to produce a general framework to produce infinitely many Lagrangian tori in a complex smooth projective variety with a monotone Kähler form obtained from different limits in the deformation space of  $X$  and to find a novel class of Fano varieties admitting infinitely many distinct monotone Lagrangian tori.

## 1.2. Main results

The first half of this paper discusses a sufficient condition for a family of monotone Lagrangian tori constructed from a cluster structure to have *infinitely* many monotone Lagrangian tori, no two of which are related by any symplectomorphisms.

Throughout this paper, all varieties are defined over the field  $\mathbb{C}$  of complex numbers. Let  $X$  be a smooth projective variety of complex dimension  $m$ . Consider an oriented rooted regular tree  $\mathbb{T}$  whose vertices parametrize the seeds of a cluster algebra with outgoing edges indicating the direction of mutation from each vertex. Let  $t_0$  be the root corresponding to the initial seed. Suppose that for each vertex  $t \in \mathbb{T}$ , there is a valuation  $v_t: \mathbb{C}(X) \setminus \{0\} \rightarrow \mathbb{Z}^m$  with one-dimensional leaves associated with  $t$ . We denote by  $S_t$  (resp.  $\Delta_t$ ) the semigroup (resp. the Newton–Okounkov polytope) constructed from  $v_t$ .

The family  $\{\Delta_t \mid t \in \mathbb{T}\}$  is said to have a *tropical cluster structure* if, whenever two vertices  $t$  and  $t'$  are connected by an oriented edge, the Newton–Okounkov polytopes  $\Delta_t$  and  $\Delta_{t'}$  are related by a tropicalized cluster mutation in the direction corresponding to the oriented edge. Similarly, the family  $\{S_t \mid t \in \mathbb{T}\}$  is said to have a *tropical cluster structure* if, for each pair  $(t, t')$  of vertices connected by an oriented edge, the semigroups  $S_t$  and  $S_{t'}$  are related by a level-wise tropicalized cluster mutation in the direction corresponding to the connecting oriented edge. Refer to Definition 4.3 for the precise definition. In the case where each polytope is given by the convex hull of the tropical integer points, the Newton–Okounkov polytopes have a tropical cluster structure if the FG conjecture holds.

Assume in addition that the Newton–Okounkov polytope  $\Delta_{t_0}$  at the initial seed is  $\mathbb{Q}$ -Gorenstein Fano and normal. The following proposition claims that the other Newton–Okounkov polytopes are  $\mathbb{Q}$ -Gorenstein Fano and normal.

**Proposition B** (Proposition 4.10). *For each  $t \in \mathbb{T}$ , let  $\Delta_t$  be the Newton–Okounkov polytope of  $X$  corresponding to a semigroup  $S_t$ . Suppose that the family  $\{S_t \mid t \in \mathbb{T}\}$  of semigroups has a tropical cluster structure. If*

- (1) *the center of the polytope  $\Delta_{t_0}$  is fixed under any tropicalized cluster mutations,*
- (2) *the Newton–Okounkov polytope  $\Delta_{t_0}$  at the initial seed is  $\mathbb{Q}$ -Gorenstein Fano, and*
- (3) *the associated toric degeneration is normal,*

*then the other Newton–Okounkov polytopes in the family are also  $\mathbb{Q}$ -Gorenstein Fano and the corresponding toric degenerations are normal.*

Suppose that the Kähler form on  $X$  inherited from the ambient space of the toric degeneration is monotone. By Proposition A and B, there are as many monotone Lagrangian tori in  $X$  as seeds in the cluster algebra once the initial Newton–Okounkov body is  $\mathbb{Q}$ -Gorenstein Fano and the associated toric degeneration is normal. If the cluster algebra is of infinite type, we obtain infinitely many Lagrangian tori, some of which could be related by a symplectomorphism.

The next crucial question, therefore, is how to distinguish the constructed monotone tori. The Newton polytope of invariants counting holomorphic disks or displacement energy of neighboring Lagrangians has been employed to compare the constructed Lagrangians, see [EP97, Che96] for instance. To compute (or estimate) them, a description of the facets of Newton–Okounkov polytope is quite crucial. In our case, *however*, each Newton–Okounkov polytope is given by the convex hull of a discrete set of tropical integer points so that we do *not* have a preferred description at hand. In fact, finding such an explicit “polyhedral” description of a family of polytopes is a challenging problem, see [BZ01, RW19] for results on certain classes of examples in this problem. This difficulty motivates us to devise a *practical* criterion that relies solely on the data of tropical integer points along with the exchange matrices (*not* on the data of facets).

**Theorem C** (Theorem 4.11). *Let  $X$  be a smooth Fano variety equipped with a Kähler form obtained from a very ample line bundle  $\mathcal{L}$  given by a positive power of the anticanonical bundle of  $X$ . Suppose that  $X$  has a family  $\{\Delta_t \mid t \in \mathbb{T}\}$  of Newton–Okounkov polytopes of  $\mathcal{L}$  having a tropical cluster structure and satisfying all conditions in Proposition B. Let  $L_t$  be a monotone Lagrangian torus constructed from  $\Delta_t$  by Proposition A. Assume that the Newton–Okounkov polytope  $\Delta_{t_0}$  at the initial seed contains the origin.*

*If there exists a sequence  $(t_\ell)_{\ell \in \mathbb{N}}$  of seeds and a sequence  $(n_\ell)_{\ell \in \mathbb{N}}$  of integers such that*

- (1) *the integer  $n_\ell$  is an entry  $\varepsilon_{r_\ell, s_\ell}$  of the exchange matrix  $\varepsilon_{t_\ell}$  associated with  $t_\ell$  such that  $r_\ell$  indices a unfrozen variable and the sequence  $(n_\ell)_{\ell \in \mathbb{N}}$  diverges to  $-\infty$  as  $\ell \rightarrow \infty$ ,*

- (2) both the Newton–Okounkov polytope  $\Delta_{t_\ell}$  and the image of  $\Delta_{t_\ell}$  under the tropicalized cluster mutation in the  $r_\ell$ -direction are contained in the half-space

$$\{\mathbf{u} \in \mathbb{R}^m \mid u_{s_\ell} \geq 0\},$$

then the family  $\{L_{t_\ell} \mid \ell \in \mathbb{N}\}$  contains infinitely many monotone Lagrangian tori, no two of which are related by any symplectomorphisms.

To derive this criterion, we explore relations between various polytopes obtained from Newton–Okounkov polytope  $\Delta_{t_\ell}$ . Suppose that the Newton–Okounkov polytope  $\Delta_{t_0}$  is  $\mathbb{Q}$ -Gorenstein Fano. Assume that the center  $\mathbf{u}_0$  is preserved under the tropicalized mutation in each direction as in Proposition B. It implies that every polytope  $\Delta_{t_\ell}$  is a  $\mathbb{Q}$ -Gorenstein Fano polytope with the same center  $\mathbf{u}_0$ . Under this circumstance, we study a relation between the entry  $-\varepsilon_{r_\ell, s_\ell}$  exchange matrix and the number of lattice points of the polar dual of  $\Delta_{t_\ell} - \mathbf{u}_0$  in the lattice  $\frac{1}{q}\mathbb{Z}$ . Here the integer  $q$  is a certain natural number completely determined by the center  $\mathbf{u}_0$ .

With the relation at hand, to extract a geometric consequence from the polar dual, we introduce a *refined disk potential* of  $L_{t_\ell}$ . As every counting invariant bounded by  $L_{t_\ell}$  is *not* known, we only have partial information on counting invariants in general. The refined disk potential is designed to avoid undesired cancellations and the Newton polytope of the refined disk potential is also invariant under symplectomorphisms module unimodular equivalences. As a consequence, (a multiple of) the polar dual  $(\Delta_{t_\ell} - \mathbf{u}_0)^\circ$  is contained in the Newton polytope of its refined disk potential of  $L_{t_\ell}$ , see Proposition 3.25. Therefore, to show that there are infinitely many *distinct* monotone Lagrangian tori, it suffices to show that there is a polar dual with an arbitrarily large number of lattice points. The problem turns into searching an exchange matrix with an arbitrarily large entry in the same mutation equivalence class because of the derived relation between the lattice points and the entry of the exchange matrix.

In the second part of the paper, we apply the above criterion to the flag manifolds of *arbitrary* type to show that they have infinitely many distinct monotone Lagrangian tori.

**Theorem D** (Theorem 5.1). *Let  $G$  be a simply connected and simple complex Lie group and  $B$  a Borel subgroup. Let  $2\rho$  be the anticanonical regular dominant weight. Then every flag manifold  $X = G/B$  equipped with the Kähler form  $\omega_{2\rho}$  other than  $A_1, A_2, A_3, A_4$ , and  $B_2 = C_2$  type, that is,*

$$(1.1) \quad G \neq \mathrm{SL}_2(\mathbb{C}), \mathrm{SL}_3(\mathbb{C}), \mathrm{SL}_4(\mathbb{C}), \mathrm{SL}_5(\mathbb{C}), \mathrm{Spin}_5(\mathbb{C}) = \mathrm{Sp}_4(\mathbb{C}).$$

*has infinitely many monotone Lagrangian tori, no two of which are related by any symplectomorphisms.*

To construct *infinitely* many monotone Lagrangian tori in flag manifolds, we exploit a recent construction of Newton–Okounkov bodies of Schubert varieties via the theory of cluster algebras by Fujita–Oya [FO20], which will be briefly recalled.

Let  $G$  be a simply connected and simple complex Lie group. The coordinate ring of a unipotent cell  $U_w^-$  is isomorphic to an upper cluster algebra generated by unipotent minors, constructed by [BFZ05, Wil13, GLS11]. In [FO20], Fujita–Oya showed that each seed gives rise to a valuation on the function field  $\mathbb{C}(U_w^-)$  and produces a Newton–Okounkov polytope of  $X_w$  in  $G/B$ . We call this polytope a *cluster polytope*. Moreover, they proved that the lattice points of each cluster polytope parametrize the elements of a certain basis on  $\mathbb{C}[U_w^-]$ . This basis arises from the dual canonical basis of Lusztig in [Lus90] or the upper global basis of Kashiwara in [Kas90] of the quantum group of  $G$ . By specializing the basis at  $q = 1$ , it gives rise to a basis on the coordinate ring of the unipotent radical and induces a basis

on  $\mathbb{C}[U_w^-]$ . The dual canonical basis has remarkable properties, satisfying the axioms of a triangular basis in Qin [Qin17, Qin20]. For instance, for each choice of seed, we obtain a parametrization of the induced basis given by the extended  $g$ -vectors. In particular, two parametrizations are related by a finite sequence of tropicalized cluster mutation. It in turn implies that each pair of cluster polytopes are related by a finite sequence of tropicalized cluster mutations. Therefore, the family of cluster polytopes has a tropical cluster structure.

This construction of cluster polytopes is crucial for Theorem D as the cluster polytopes enable us to construct *infinitely* many toric degenerations if the cluster algebra is of infinite type, which happens in (1.1). Recall that the previous known toric degenerations of  $G/B$  arise from the Gelfand–Zeitlin polytope or string polytopes, see [GL96, KM05, Cal02] for instance. For a fixed group  $G$  and a regular integral dominant weight  $\lambda$ , there are only *finitely* many string polytopes and hence there are only finitely many toric degenerations of  $G/B$  arising from a string polytope. Cluster polytopes are a generalization of string polytopes. The polyhedral description of a “standard” string polytope  $\Delta$  was known by Littelmann [Lit98]. Using this description, we prove that the polytope  $\Delta$  is  $\mathbb{Q}$ -Gorenstein Fano. Moreover, every cluster polytope is  $\mathbb{Q}$ -Gorenstein Fano by Proposition B and hence we have infinitely many monotone Lagrangian tori.

To apply Theorem C, we need to check the two conditions. One of the key ingredients for the first condition is a classification of mutation finite skew-symmetrizable matrices with frozen indices, recently established in [FT21]. Assume that a skew-symmetrizable matrix with one frozen index  $s$  is mutation infinite but the submatrix of unfrozen part is mutation-finite. Then its mutation equivalence class contains a sequence matrices  $(\varepsilon^\ell)_{\ell \in \mathbb{N}}$  such that  $\varepsilon_{r_\ell, s}^\ell \rightarrow -\infty$  as  $\ell \rightarrow \infty$  for a sequence of unfrozen indices  $(r_\ell)_{\ell \in \mathbb{N}}$ . Since the mutation of matrices is compatible with the restriction, the problem is reduced to finding an exchange matrix that contains a submatrix with the above property. Using the compatibility of Dynkin diagram embedding with the seed arising from a reduced expression of the longest element in the Weyl group, we reduce the problem into the five cases  $\mathfrak{g} = A_5, B_3, C_3, D_4$  and  $G_2$ . Finally, we provide a case-by-case analysis for those cases.


The second condition is extracted from the correspondence between the lattice points of Newton–Okounkov bodies of a flag manifold  $G/B$  are parametrized by the dual canonical basis. In the case of  $G/B$ , the frozen components of the extended  $g$ -vectors of dual canonical basis elements are always *non-negative*. It in turn yields that a cluster polytope and its mutations in any direction are simultaneously supported by the half-space associated with each frozen variable. This observation determines a lower bound for the lattice points the polar dual of the cluster polytope.

### Acknowledgement

The author would like to thank Cheol-Hyun Cho and Jae-Hoon Kwon for their helpful discussions, comments, and encouragement. The third named author would like to thank Hyun Kyu Kim and Grigory Mikhalkin for their helpful discussions.

## 2. Newton–Okounkov bodies, toric degenerations, and Lagrangian tori

In this section, we briefly review a way of constructing a Lagrangian torus fibration on a given smooth projective variety via the theory of Newton–Okounkov bodies, following [HK15, And13]. The outline of the construction is depicted below.



### 2.1. Lagrangian tori from toric degenerations

Recall that a *toric degeneration* of a smooth projective variety  $X$  is a flat family  $\mathfrak{X} \subset \mathbb{P}^N \times \mathbb{C}$  of projective varieties in an ambient space  $\mathbb{C}P^N$  with the following commutative diagram

$$(2.1) \quad \begin{array}{ccc} \mathfrak{X} & \xrightarrow{i} & \mathbb{C}P^N \times \mathbb{C} \\ & \searrow \pi & \downarrow \text{pr} \\ & & \mathbb{C} \end{array}$$

such that

- (1) the family is trivial over  $\mathbb{C}^*$ ,  $\pi^{-1}(\mathbb{C}^*) \cong X \times \mathbb{C}^*$ , and
- (2)  $\pi^{-1}(0)$  is a (not necessarily normal) projective toric variety.

We denote by  $X_t$  the fiber  $\pi^{-1}(t)$  over  $t \in \mathbb{C}$ .

**Example 2.1.** Let  $C = \{[x : y : z] \in \mathbb{C}P^2 \mid y^3 = x^3 + z^3\}$  be an elliptic curve in  $\mathbb{C}P^2$ . For example, a toric degeneration of  $C$  is given by

$$\mathfrak{X} = \{([x : y : z], t) \in \mathbb{C}P^2 \times \mathbb{C} \mid y^2 z = x^3 + tz^3\},$$

a flat family of curves such that  $X_t \cong C$  for  $t \in \mathbb{C}^*$ . The variety  $X_0$  over the origin is the cuspidal cubic. It is a (non-normal) toric variety because it admits the obvious  $\mathbb{C}^*$ -action on  $X_0$  induced from the  $\mathbb{C}^*$ -action on  $\mathbb{P}^2$  given by

$$t \cdot [x : y : z] = [x : ty : t^{-2}z] \quad \text{for } t \in \mathbb{C}^*$$

with two fixed points  $[0 : 1 : 0]$  and  $[0 : 0 : 1]$ .

Let  $X$  equip with a Kähler form  $\omega$  induced from the Fubini–Study form on  $\mathbb{C}P^N$ . Consider a Kähler form  $\Omega$  on the smooth part  $\mathring{\mathfrak{X}}$  of  $\mathfrak{X}$  induced from the standard Kähler form  $\omega_{\text{FS}} \oplus \omega_{\text{std}}$  on  $\mathbb{C}P^N \times \mathbb{C}$  where  $\omega_{\text{FS}}$  and  $\omega_{\text{std}}$  are the Fubini–Study form on  $\mathbb{C}P^N$  and the standard symplectic form on  $\mathbb{C}$ , respectively. Then each fiber  $X_t$  inherits a Kähler form  $\omega_t$  from  $(\mathring{\mathfrak{X}}, \Omega)$ , i.e.,  $\omega_t = \Omega|_{X_t}$ . We call  $\pi$  a *toric degeneration of  $(X, \omega)$*  if the equipped Kähler form  $\omega$  agrees with the restricted form  $\omega_1$ .

The following theorem produces Lagrangian tori in  $X$  from a toric degeneration of  $(X, \omega)$ .

**Theorem 2.2** ([HK15]). *Suppose that  $\pi: \mathfrak{X} \rightarrow \mathbb{C}$  is a toric degeneration of  $(X, \omega)$ . Then there is a continuous map  $\phi: X = X_1 \rightarrow X_0$  satisfying the following:*

- (1) *Let  $U_0$  be the smooth locus of  $X_0$  and  $U := \phi^{-1}(U_0) \subset X$ . Then the map  $\phi$  restricted to  $U$  is a symplectomorphism onto  $U_0$ .*
- (2) *Let  $\Phi_0: X_0 \rightarrow \Delta$  be the moment map for  $(X_0, \omega_0)$ . Then the composition*

$$(2.2) \quad \Phi_1 := \Phi_0 \circ \phi: X \longrightarrow \Delta$$

*is a Lagrangian torus fibration on  $U$ , an open dense subset of  $X$ .*

Note that the toric variety  $X_0$  carries a Lagrangian torus fibration  $\Phi_0$  on  $U_0$  generated by the real torus action induced from one in  $\mathbb{C}P^N$ . By pulling this fibration on  $U_0$  back to  $U$  via  $\phi$ , we produce the Lagrangian torus fibration  $\Phi_1$  in (2.2).

**Remark 2.3.** If the toric variety  $X_0$  in a toric degeneration  $\pi$  is normal as an algebraic variety,  $U_0$  contains  $\Phi_0^{-1}(\Delta \setminus \Delta_2)$  where  $\Delta_2$  is the union of codimension two faces of  $\Delta$ . Consequently, the Lagrangian fibration  $\Phi_1: X \rightarrow \Delta$  can be identified with the moment map  $\Phi_0: X_0 \rightarrow \Delta$  on the open dense subset  $\Phi_1^{-1}(\Delta \setminus \Delta_2) \subset X$  under the symplectomorphism  $\phi$ .

## 2.2. Toric degenerations via Newton–Okounkov bodies

Let  $X$  be a complex  $m$ -dimensional smooth projective variety with a very ample line bundle  $\mathcal{L}$  on  $X$ . Fix a total order  $\geq$  on  $\mathbb{Z}^m$  respecting the addition. To construct a Newton–Okounkov body, in addition to  $\mathcal{L}$ , we need to choose two data; a valuation  $v$  and a reference section  $h$ .

Suppose that  $v$  is a valuation on the function field, that is, a function  $v: \mathbb{C}(X) \setminus \{0\} \rightarrow \mathbb{Z}^m$  such that for all rational functions  $f, g \in \mathbb{C}(X) \setminus \{0\}$  and  $c \in \mathbb{C} \setminus \{0\}$ ,

- (1)  $v(fg) = v(f) + v(g)$ ,
- (2)  $v(f + g) \geq \min(v(f), v(g))$ , and
- (3)  $v(cf) = v(f)$ .

We additionally assume that the valuation  $v$  has *one-dimensional leaves*, that is,  $v$  satisfies that for all rational functions  $f, g \in \mathbb{C}(X) \setminus \{0\}$  with  $v(f) = v(g)$ , there is  $c \in \mathbb{C} \setminus \{0\}$  such that  $v(g - cf) > v(g)$  or  $g - cf = 0$ . Equivalently, setting  $\mathbb{C}(x)_\alpha = \{f \in \mathbb{C}(x) \mid v(f) \geq \alpha \text{ or } f = 0\}$ ,

$$(2.3) \quad \dim_{\mathbb{C}} \left( \mathbb{C}(x)_\alpha / \sum_{\beta; \beta > \alpha} \mathbb{C}(x)_\beta \right) \leq 1, \quad \text{for each } \alpha \in \mathbb{Z}^m.$$

Note that every divisorial valuation has one-dimensional leaves. For instance, a valuation obtained from a flag  $X_\bullet = X = X_m \supset X_{m-1} \supset \cdots \supset X_1 \supset X_0$  of subvarieties in  $X$  has one-dimensional leaves.

Let  $\mathcal{L} := H^0(X, \mathcal{L})$  be the space of global sections. Then

$$R = \bigoplus_{k \geq 0} R_k, \quad R_k = \mathcal{L}^{\otimes k}, \quad R_0 := \mathbb{C}$$

is the homogeneous coordinatizing ring of  $X$ . Choose a non-zero reference section  $h \in \mathcal{L}$  and define a semigroup

$$(2.4) \quad S(\mathcal{L}, v, h) = \bigcup_{k \geq 1} \{(k, v(f/h^k)) \mid f \in \mathcal{L}^k \setminus \{0\}\} \subset \mathbb{Z}_{\geq 0} \times \mathbb{Z}^m \subset \mathbb{R}_{\geq 0} \times \mathbb{R}^m.$$

The *Newton–Okounkov body* associated with the triple  $(\mathcal{L}, v, h)$  is defined by

$$(2.5) \quad \Delta(\mathcal{L}, v, h) := \overline{\text{conv} \left( \bigcup_{k \in \mathbb{N}} \{x/k \mid (k, x) \in S(\mathcal{L}, v, h)\} \right)} \subset \mathbb{R}^m.$$

We call  $\Delta$  a *Newton–Okounkov body of  $\mathcal{L}$*  if  $\Delta = \Delta(\mathcal{L}, v, h)$  for some pair  $(v, h)$  of a valuation and a reference section.

One can systematically produce a toric degeneration of  $X$  from a Newton–Okounkov body.

**Theorem 2.4** ([And13]). *If the semigroup  $S(\mathcal{L}, v, h)$  in (2.4) is finitely generated, then the Newton–Okounkov body  $\Delta(\mathcal{L}, v, \tau)$  is a rational polytope. Furthermore, there is a toric degeneration  $\pi: \mathfrak{X} \rightarrow \mathbb{C}$  of  $X$  where the central fiber  $X_0$  is a projective toric variety whose normalization is the toric variety associated with  $\Delta(\mathcal{L}, v, \tau)$ .*



### 3. $\mathbb{Q}$ -Gorenstein Fano polytopes and Monotone Lagrangian tori

The aim of this section is to construct a monotone Lagrangian torus in a smooth projective variety  $X$  equipped with a monotone Kähler form when  $X$  admits a *normal* toric degeneration arising from a  *$\mathbb{Q}$ -Gorenstein Fano* Newton–Okounkov polytope. We also define a *refined* version of Newton polytopes from counting invariants of a monotone Lagrangian torus, which will be employed to distinguish the constructed Lagrangian tori later on.

#### 3.1. Monotone Lagrangians and gradient holomorphic disks

Let  $X$  be a symplectic manifold with a symplectic form  $\omega$ . The symplectic form  $\omega$  is said to be *monotone* if  $c_1(TX) = \lambda \cdot [\omega] \in H^2(X; \mathbb{Z})$  for some real number  $\lambda > 0$ . By scaling the symplectic form  $\omega$  if necessary, we may assume that  $\omega$  is *normalized*, that is,

$$(3.1) \quad c_1(TX) = [\omega] \quad \text{in } H^2(X; \mathbb{Z}).$$

The cohomology classes  $[\omega]$  and  $c_1(TX)$  can be regarded as a homomorphism assigning the symplectic area and the Chern number of each spherical homotopy class in  $\pi_2(X)$ , respectively, that is,  $[\omega]$  and  $c_1(TX) \in \text{Hom}(\pi_2(X), \mathbb{Z})$ . Note that the symplectic form  $\omega$  is monotone if  $\omega(\alpha) = c_1(TX)(\alpha)$  for all  $\alpha \in \pi_2(X)$ .

Recall that a submanifold  $L$  of  $X$  is *Lagrangian* if  $\dim L = (\dim X)/2$  and  $\omega|_L \equiv 0$ . Let  $\mathbb{D} = \{z \in \mathbb{C} \mid |z| \leq 1\}$  be the unit disk. To each continuous map  $\varphi: \mathbb{D} \rightarrow X$  with the Lagrangian boundary condition  $\varphi(\partial\mathbb{D}) \subset L$ , one can assign an integer  $\mu_L([\varphi])$ , called the *Maslov index*. It can be thought as a Chern number of the disk  $\varphi$  in the sense that if the image of a continuous map sphere  $\varphi: S^2 \rightarrow X$  intersects  $L$  so that  $\varphi(S^2)$  divides into two disks  $\varphi^+: \mathbb{D}^2 \rightarrow X$  and  $\varphi^-: \mathbb{D}^2 \rightarrow X$ , then the following holds:

$$\mu_L([\varphi^+]) + \mu_L([\varphi^-]) = 2 \cdot c_1(TX)([\varphi(S^2)]).$$

The Maslov index is well-defined up to homotopy and we may think of the Maslov index as a homomorphism  $\mu_L: \pi_2(X, L) \rightarrow \mathbb{Z} \subset \mathbb{R}$ . Because of the Lagrangian boundary condition, we have a well-defined symplectic area homomorphism  $\omega: \pi_2(X, L) \rightarrow \mathbb{R}$  defined by  $\omega([\varphi]) := \int_{\mathbb{D}} \varphi^* \omega$ .

**Definition 3.1.** A Lagrangian submanifold  $L$  of  $(X, \omega)$  is called *monotone* if there exists  $\delta > 0$  such that

$$(3.2) \quad \mu_L(\beta) = \delta \cdot \omega(\beta) \quad \text{for all } \beta \in \pi_2(X, L).$$

Assume that  $(X, \omega)$  is not symplectically aspherical, that is, there exists  $\alpha \in \pi_2(X)$  such that  $\omega(\alpha) \neq 0$ . It is known that a monotone Lagrangian can exist only when  $\omega$  is monotone. Moreover, if  $c_1(TX) = \lambda \cdot [\omega]$  and  $X$  admits a monotone Lagrangian submanifold  $L$  satisfying (3.2), then  $\delta = 2\lambda$ , see [Oh93, Remark 2.3]. Thus if  $\omega$  is normalized as in (3.1), we then have

$$\mu_L(\beta) = 2 \cdot \omega(\beta) \quad \text{for all } \beta \in \pi_2(X, L).$$

**Example 3.2.** Let  $X$  be the complex projective line  $\mathbb{C}P^1 \simeq S^2$  with the Fubini–Study form  $\omega_{\text{FS}}$ . Then every closed Lagrangian submanifold  $L$  of  $(X, \omega_{\text{FS}})$  is diffeomorphic to a circle. As  $L$  divides  $X$  into two pieces  $\mathbb{D}^+$  and  $\mathbb{D}^-$  of disks, we have two holomorphic disks  $\varphi^+: \mathbb{D} \rightarrow \mathbb{D}^+$  and  $\varphi^-: \mathbb{D} \rightarrow \mathbb{D}^-$ , each of which are both Maslov index two. Therefore  $L$  is monotone if and only if the symplectic areas of  $\mathbb{D}^+$  and  $\mathbb{D}^-$  are equal.

In [CK21], the first and the third named authors developed a method of computing the Maslov index of a *gradient holomorphic disks*, which will be briefly recalled. Let  $(X, \omega)$  be a symplectic manifold with a Hamiltonian  $S^1$ -action and an  $S^1$ -invariant  $\omega$ -compatible almost

complex structure  $J$  on  $X$ . Let  $p \in X$  and  $C$  be the  $S^1$ -orbit containing  $p$ . If  $C$  flows along the gradient vector field of a moment map with respect to  $\omega$  and  $J$ , then the trajectory of  $C$  defines an embedded  $J$ -holomorphic half-cylinder  $Y$  diffeomorphic to  $S^1 \times \mathbb{R}_{\geq 0}$ . In the case where  $C$  converges to some point, a fixed point of the action indeed, then it gives rise to a  $J$ -holomorphic disk

$$\varphi: \mathbb{D} \rightarrow X \quad \text{satisfying } \varphi(\mathbb{D} \setminus \{0\}) = Y.$$

Such a disk  $\varphi$  is called a *gradient holomorphic disk*. Let us recall the following.

**Theorem 3.3** (Theorem A in [CK21]). *Let  $(X, \omega)$  be a symplectic manifold with a Hamiltonian  $S^1$ -action. Let  $H: X \rightarrow \mathbb{R}$  be a moment map of the  $S^1$ -action. Suppose that  $\varphi: \mathbb{D} \rightarrow X$  is a gradient holomorphic disk such that*

- (1)  $\varphi(0)$  is a fixed point of the  $S^1$ -action,
- (2)  $L$  is a Lagrangian submanifold of  $(X, \omega)$  lying on some level set  $H^{-1}(c)$ , and
- (3)  $\varphi: \mathbb{D} \rightarrow X$  satisfies the Lagrangian boundary condition, that is,  $\varphi(\partial\mathbb{D}) \subset L$ .

*Then the Maslov index of  $[\varphi]$  is equal to  $-2n_0$ , where  $n_0$  is the sum of negative weights of the tangential  $S^1$ -representation at  $\varphi(0)$ .*

Gradient holomorphic disks are useful to test whether a given torus invariant Lagrangian submanifold (Lagrangian toric fibers, for instance) is monotone or not. Here is an example.

**Example 3.4** (Fano toric varieties). Let  $X$  be an  $m$ -dimensional smooth Fano toric variety with a moment map  $\Phi: X \rightarrow \Delta \subset \mathfrak{t}^* \cong \mathbb{R}^m$  where  $\Delta$  is a smooth reflexive polytope and contains a unique interior lattice point  $\mathbf{u}_0 \in \Delta$ . By the reflexivity of  $\Delta$ , the variety  $X$  carries a Kähler form  $\omega$  invariant under the torus action and satisfying  $c_1(TX) = [\omega]$ . Let  $F_1, \dots, F_\kappa$  be the facets of  $\Delta$ . For an interior point  $\mathbf{u}$  of  $\Delta$ , Cho and Oh proved that a Lagrangian fiber  $L_{\mathbf{u}} := \Phi^{-1}(\mathbf{u})$  is monotone if and only if  $\mathbf{u} = \mathbf{u}_0$  in [CO06]. They proved that for each facet  $F_i$  and a generic point  $p \in L_{\mathbf{u}}$ , there exists a unique holomorphic disk  $\varphi'_i: \mathbb{D} \rightarrow X$  of Maslov index two passing through  $p$  such that  $\varphi'_i(\partial\mathbb{D}) \subset L_{\mathbf{u}}$  and the area of  $\varphi'_i(\mathbb{D})$  is precisely the affine distance from  $\mathbf{u}$  to  $F_i$ . Therefore,  $L_{\mathbf{u}}$  is monotone if and only if the affine distance from  $\mathbf{u}$  to all facets are all equal, that is,  $\mathbf{u} = \mathbf{u}_0$ .

In fact,  $\varphi'_i$  can be described as a gradient holomorphic disk obtained as follows. We first take the point  $p \in L_{\mathbf{u}}$  in the previous paragraph and  $S^1$  denotes the circle subgroup of  $T$  whose Lie algebra is the one-dimensional subspace of  $\mathfrak{t}^* \cong \mathbb{R}^n$  perpendicular to the facet  $F_i$ . We take the standard complex structure, which is  $S^1$ -invariant. Let  $C$  be the  $S^1$ -orbit containing  $p$ . Then the orbit  $C$  flows along the gradient vector field of a moment map for the  $S^1$ -action and it converges to some fixed point whose moment map image is in the relative interior of  $F_i$ . This trajectory gives a gradient holomorphic disk which we denote by  $\varphi_i: \mathbb{D} \rightarrow X$ . As the fixed component of the  $S^1$ -action corresponding to  $F_i$  is of codimension two, the  $S^1$ -action near  $\pi^{-1}(F_i)$  is semifree and hence  $\varphi_i$  is of Maslov index two. By the classification of holomorphic disks of Maslov index two in [CO06], we conclude that  $\varphi'_i = \varphi_i$ .

### 3.2. $\mathbb{Q}$ -Gorenstein Fano and normal toric degenerations

Let  $X$  be a smooth projective variety of complex dimension  $m$ . Let  $v$  be a valuation on the function field  $\mathbb{C}(X)$  with one-dimensional leaves (see (2.3)),  $\mathcal{L}$  a very ample line bundle on  $X$ , and  $h$  a non-zero reference section in  $\mathcal{L} = H^0(X, \mathcal{L})$ . From the choice  $(\mathcal{L}, v, h)$ , we produce a triple; a semigroup  $S(\mathcal{L}, v, h)$ , a Newton–Okounkov body  $\Delta(\mathcal{L}, v, h)$ , and a toric degeneration  $\pi: \mathfrak{X} \rightarrow \mathbb{C}$  as in Section 2.2. To produce a *monotone* Lagrangian torus from the toric degeneration  $\pi$  of  $X$ , we require the toric degeneration  $\pi$  to be *normal* and the Newton–Okounkov body  $\Delta(\mathcal{L}, v, h)$  to be  *$\mathbb{Q}$ -Gorenstein Fano*, which will be defined in this section.

To begin with, we review some notions of polyhedral geometry. Let  $N$  be a lattice of rank  $m$  and  $M := \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$  the dual lattice of  $N$ . Let us fix a basis  $\mathbf{s}$  for the free abelian group  $N$ . Using the fixed basis  $\mathbf{s}$ ,  $N$  can be identified with  $\mathbb{Z}^m$ . The dual lattice  $M$  can be identified with  $\mathbb{Z}^m$  via the dual basis of  $\mathbf{s}$ . We also have the identifications  $N_{\mathbb{R}} := N \otimes \mathbb{R} \simeq \mathbb{R}^m$  and  $M_{\mathbb{R}} := M \otimes \mathbb{R} \simeq \mathbb{R}^m$ . We take the Cartesian coordinate system  $\mathbf{v} := (v_1, v_2, \dots, v_m)$  (resp.  $\mathbf{u} := (u_1, u_2, \dots, u_m)$ ) for  $N_{\mathbb{R}} \simeq \mathbb{R}^m$  (resp.  $M_{\mathbb{R}} \simeq \mathbb{R}^m$ ). Let  $\langle \cdot, \cdot \rangle: M_{\mathbb{R}} \times N_{\mathbb{R}} \rightarrow \mathbb{R}$  be the canonical pairing. For a non-zero vector  $\mathbf{v} \in N_{\mathbb{R}}$  and a real number  $\alpha \in \mathbb{R}$ ,

- the *hyperplane*  $H_{\mathbf{v}, \alpha}$  is defined by  $H_{\mathbf{v}, \alpha} := \{\mathbf{u} \in M_{\mathbb{R}} \mid \langle \mathbf{u}, \mathbf{v} \rangle + \alpha = 0\}$  and
- the (*closed*) *half-space*  $H_{\mathbf{v}, \alpha}^+$  is defined by  $H_{\mathbf{v}, \alpha}^+ := \{\mathbf{u} \in M_{\mathbb{R}} \mid \langle \mathbf{u}, \mathbf{v} \rangle + \alpha \geq 0\}$ .

Suppose that  $\Delta \subset M_{\mathbb{R}}$  is a full-dimensional polytope that contains the origin in its interior, that is,  $\mathbf{0} \in \text{Int}(\Delta)$ . Then the polytope  $\Delta$  in  $M_{\mathbb{R}}$  can be uniquely expressed as an intersection of half-spaces

$$(3.3) \quad \Delta = \bigcap_{j=1}^{\kappa} H_{\mathbf{v}_j, 1}^+$$

satisfying the following conditions;

- (1) each vector  $\mathbf{v}_j \in N_{\mathbb{R}}$ ,
- (2) each hyperplane  $H_{\mathbf{v}_j, \alpha}$  contains a facet of  $\Delta$ , and
- (3)  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{\kappa}\}$  is pairwise distinct.

In the presentation of  $\Delta$  in (3.3), note that each vector  $\mathbf{v}_j$  is an *inward* normal vector to a facet of  $\Delta$  and the number of facets is equal to  $\kappa$ .

Let  $C$  be a subset of  $M_{\mathbb{R}}$ . For a non-zero vector  $\mathbf{v} \in N_{\mathbb{R}}$  and a real number  $\alpha \in \mathbb{R}$ , the half-space  $H_{\mathbf{v}, \alpha}^+$  is called a *supporting half-space* of the subset  $C$  if  $C \subset H_{\mathbf{v}, \alpha}^+$  and  $C \cap H_{\mathbf{v}, \alpha} \neq \emptyset$ . In this case, we call  $H_{\mathbf{v}, \alpha}$  a *supporting hyperplane* of  $C$ . Note that if  $C$  is a polytope  $\Delta$ , then a hyperplane containing a lower dimensional stratum can be a supporting hyperplane. We emphasize that a supporting hyperplane need *not* contain a facet of  $\Delta$ .

We recall the “polar dual” of a polytope  $\Delta$ .

**Definition 3.5.** Let  $\Delta$  be a full-dimensional polytope in  $M_{\mathbb{R}}$  containing the origin in its interior. The *polar dual* of  $\Delta$  is defined by

$$\Delta^{\circ} := \{\mathbf{v} \in N_{\mathbb{R}} \mid \langle \mathbf{u}, \mathbf{v} \rangle + 1 \geq 0 \text{ for all } \mathbf{u} \in \Delta\}.$$

Indeed,  $\Delta^{\circ}$  is also a polytope containing the origin and it contains the origin in its interior. Moreover, it satisfies  $(\Delta^{\circ})^{\circ} = \Delta$ , see [Brø83, Theorem 6.2] for the proof. The following proposition provides an alternative way of presenting the polar dual of  $\Delta$ .

**Proposition 3.6.** *Suppose that  $\Delta \subset M_{\mathbb{R}} \simeq \mathbb{R}^m$  is a full-dimensional polytope containing the origin  $\mathbf{0}$  in the interior of  $\Delta$ . Then the polar dual  $\Delta^{\circ}$  is equal to the convex hull of  $\{\mathbf{v}_j \mid j = 1, 2, \dots, \kappa\}$  in  $N_{\mathbb{R}}$  where  $\mathbf{v}_j$ 's are from the unique presentation (3.3) of  $\Delta$ .*

*Proof.* Let  $\Delta'$  be the convex hull of  $\{\mathbf{v}_j \mid j = 1, 2, \dots, \kappa\}$ . Since  $\langle \mathbf{u}, \mathbf{v}_j \rangle \geq -1$  for all  $\mathbf{u} \in \Delta$ , the polar dual  $\Delta^{\circ}$  contains all  $\mathbf{v}_j$ 's. Since  $\Delta^{\circ}$  is convex, we have  $\Delta' \subset \Delta^{\circ}$ . On the other hand, if  $\mathbf{u} \in (\Delta')^{\circ}$ , then  $\langle \mathbf{u}, \mathbf{v} \rangle + 1 \geq 0$  for all  $\mathbf{v} \in \Delta'$ . In particular,  $\langle \mathbf{u}, \mathbf{v}_j \rangle + 1 \geq 0$  for all  $j = 1, 2, \dots, \kappa$ . Hence  $(\Delta')^{\circ} \subset \bigcap_{j=1}^{\kappa} H_{\mathbf{v}_j, 1}^+ = \Delta$  so that we have

$$\Delta^{\circ} \subset ((\Delta')^{\circ})^{\circ}.$$

Since  $\Delta'$  is a closed and convex set containing  $\mathbf{0}$ ,  $\Delta^{\circ} \subset \Delta'$  follows from  $((\Delta')^{\circ})^{\circ} = \Delta'$ .  $\square$

As we are interested in Newton–Okounkov bodies constructed from Theorem 2.4, we are only concerned with rational polytopes. From now on, every polytope  $\Delta$  is assumed to be *rational*, which means that all vertices of  $\Delta$  have rational coordinates.

**Definition 3.7.** A full-dimensional rational polytope  $\Delta \subset M_{\mathbb{R}} \simeq \mathbb{R}^m$  is called  *$\mathbb{Q}$ -Gorenstein Fano* if there exists a vector  $\mathbf{u}_0 \in M_{\mathbb{R}}$  and a number  $\lambda \in \mathbb{Q}_{>0}$  such that  $\lambda \cdot (\Delta - \mathbf{u}_0)^\circ$  is a *Fano* polytope, a lattice polytope each of which vertex is a *primitive lattice* vector, see [KN13]. Equivalently, the translated polytope  $\Delta - \mathbf{u}_0$  has a presentation of the form (3.3) satisfying that the  $\lambda$ -multiple of each inward facet normal vector  $\mathbf{v}_j$  is primitive. In this circumstance, we call that point  $\mathbf{u}_0$  the *center* of  $\Delta$ , which maps to the origin  $\mathbf{0}$  under the translation  $\mathbf{u} \mapsto \mathbf{u} - \mathbf{u}_0$ . A  $\mathbb{Q}$ -Gorenstein Fano polytope  $\Delta \subset M_{\mathbb{R}} \simeq \mathbb{R}^m$  is said to be *normalized* if  $\lambda = 1$ .

**Definition 3.8.** For a  $\mathbb{Q}$ -Gorenstein Fano polytope  $\Delta$ , the Fano polytope  $\lambda \cdot (\Delta - \mathbf{u}_0)^\circ \subset N_{\mathbb{R}}$  in Definition 3.7 is called the *combinatorial dual* of  $\Delta$  and denoted by  $\Delta^\vee$ . In other words,  $\Delta^\vee$  is the convex hull of all 1-cones of the normal fan of  $\Delta$ .

**Remark 3.9.** The name “ $\mathbb{Q}$ -Gorenstein Fano” is originated from the following geometric fact. The toric variety associated with the normal fan of a  $\mathbb{Q}$ -Gorenstein Fano polytope is  $\mathbb{Q}$ -Gorenstein Fano as an algebraic variety. Note that the  $\mathbb{Q}$ -Gorenstein Fano condition is more general than the “reflexive” condition. Namely, every reflexive polytope is normalized  $\mathbb{Q}$ -Gorenstein Fano. In particular, a  $\mathbb{Q}$ -Gorenstein Fano polytope need *not* be a lattice polytope.

**Remark 3.10.** When  $\Delta$  is a normalized  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{u}_0$ , we can compute the polar dual  $(\Delta - \mathbf{u}_0)^\circ$  using the facet normal vectors of  $\Delta$  by Proposition 3.6 as follows. Let  $\{F_1, F_2, \dots, F_\kappa\}$  be the set of facets of  $\Delta$  and  $\mathbf{v}_{F_j}$  the primitive inward normal vector to the facet  $F_j$ . Then  $\Delta^\vee = (\Delta - \mathbf{u}_0)^\circ$  is expressed as

$$(\Delta - \mathbf{u}_0)^\circ = \text{Convex hull of } \{\mathbf{v}_{F_j} \mid j = 1, 2, \dots, \kappa\}.$$

If  $\Delta$  is a (not necessarily normalized)  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{u}_0$ , by Proposition 3.6, the combinatorial dual  $\Delta^\vee$  of  $\Delta$  is equal to

$$\text{Convex hull of } \{\lambda \cdot \mathbf{v}_j \mid j = 1, 2, \dots, \kappa\} = \text{Convex hull of } \{\mathbf{v}_{F_j} \mid j = 1, 2, \dots, \kappa\}.$$

Let  $X$  be a smooth projective variety polarized by a very ample line bundle  $\mathcal{L}$  with  $N := \dim_{\mathbb{C}} H^0(X, \mathcal{L}) - 1$ . Via the Kodaira embedding  $\varphi_{\mathcal{L}}: X \rightarrow \mathbb{C}P^N$ , the variety  $X$  can be regarded as a subvariety of the projective space  $\mathbb{C}P^N$ . Recall that we have chosen a valuation  $v: \mathbb{C}(X) \setminus \{0\} \rightarrow \mathbb{C}$  with one-dimensional leaves and a nonzero section  $h \in H^0(X, \mathcal{L})$ . We then have a semigroup  $S(\mathcal{L}, v, h)$  associated with the triple  $(\mathcal{L}, v, h)$ .

**Assumption 3.11.** Throughout this section, the associated semigroup  $S(\mathcal{L}, v, h)$  is always assumed to be finitely generated.

By Theorem 2.4, there exists a toric degeneration  $\pi: \mathfrak{X} \rightarrow \mathbb{C}$  of  $X$  associated with the Newton–Okounkov polytope  $\Delta(\mathcal{L}, v, h)$  such that the diagram (2.1) commutes. Let us choose a Kähler form  $\omega$  on  $X$  induced from the Fubini–Study form on the ambient projective space, that is,  $\omega = \varphi_{\mathcal{L}}^* \omega_{\text{FS}}$ . Such a form  $\omega$  is said to be *inherited from* the very ample line bundle  $\mathcal{L}$ . When considering a Newton–Okounkov polytope  $\Delta(\mathcal{L}, v, h)$  of  $X$ , we equip  $X$  with the Kähler form inherited from  $\mathcal{L}$ .

**Definition 3.12.** Let  $X$  be a smooth projective variety with a toric degeneration  $\pi: \mathfrak{X} \rightarrow \mathbb{C}$  arising from a Newton–Okounkov polytope  $\Delta(\mathcal{L}, v, h)$ .

- We say that  $\pi$  is  $\mathbb{Q}$ -Gorenstein Fano if the central fiber is  $\mathbb{Q}$ -Gorenstein Fano as an algebraic variety. Equivalently, the Newton–Okounkov polytope  $\Delta(\mathcal{L}, v, h)$  is  $\mathbb{Q}$ -Gorenstein Fano in the sense of Definition 3.7.
- We say that  $\pi$  is *normal* if the central fiber of  $\pi$  is normal as an algebraic variety. Equivalently,  $S(\mathcal{L}, v, h)$  is saturated.

We now construct a monotone Lagrangian torus of a smooth Fano variety equipped with a Kähler form induced from a very ample line bundle  $\mathcal{L}$ , which is defined as a positive power of the anticanonical bundle of  $X$ .

**Proposition 3.13** (Proposition A). *Let  $X$  be a smooth Fano variety equipped with a Kähler form  $\omega$  inherited from a very ample line bundle  $\mathcal{L}$ , a positive power of the anticanonical bundle of  $X$ . Suppose that the Newton–Okounkov polytope  $\Delta = \Delta(\mathcal{L}, v, h)$  of  $\mathcal{L}$  with a choice of  $(v, h)$  in Section 2.2 is  $\mathbb{Q}$ -Gorenstein Fano and the associated toric degeneration is normal. Let  $\Phi: X \rightarrow \Delta$  be a Lagrangian torus fibration constructed from Theorem 2.2. Then the fiber  $L := \Phi^{-1}(\mathbf{u}_0)$  of  $\Phi$  located at the center  $\mathbf{u}_0$  of  $\Delta$  is a monotone Lagrangian torus.*

*Proof.* Observe that the Kähler form inherited from  $\mathcal{L}$  is monotone. The very line bundle  $\mathcal{L}$  is a positive power of the anticanonical bundle  $K_X^{-1}$  of  $X$ , i.e.,  $\mathcal{L} = (K_X^{-1})^{\otimes \lambda}$  for some  $\lambda \in \mathbb{N}$ . We have

$$(3.4) \quad [\omega] = c_1(\mathcal{L}) = \lambda \cdot c_1(TX)$$

as desired. Since every smooth Fano variety is simply connected, by the long exact sequence of the pair  $(X, L)$ , we have

$$(3.5) \quad \pi_2(X, L) \simeq \pi_2(X) \oplus \pi_1(L).$$

Regarding  $\pi_2(X)$  as  $H_2(X)$  via the Hurewicz map, (3.4) yields

$$(3.6) \quad \lambda \cdot \mu_L(\alpha) = 2 \cdot \omega(\alpha) \quad \text{for every spherical class } \alpha \in \pi_2(X).$$

Note that  $\lambda$  is exactly the scaling factor associated with  $\Delta$  in Definition 3.7.

Consider the Lagrangian toric fiber  $L_0 := \Phi_0^{-1}(\mathbf{u}_0)$  in the toric variety  $X_0$ . Since  $X_0$  is normal, the singular locus of  $X_0$  has complex codimension  $\geq 2$ . In particular, the smooth locus  $U_0$  of  $X_0$  contains the inverse image  $\Phi_0^{-1}(\text{int}(F))$  of each facet  $F$  of  $\Delta$ . Let  $\mathbf{v}_F \in N$  be a primitive integral vector normal to  $F$  and  $S_F^1$  the circle subgroup generated by  $\mathbf{v}_F$ . As in Example 3.4, there is a gradient holomorphic disk of Maslov index two

$$u_F: \mathbb{D} \rightarrow X$$

such that  $u_F(\partial\mathbb{D}) \subset L_0$  and  $u_F(\mathbb{D}) \subset U_0$ .

Let  $U_1 := \phi^{-1}(U_0) \subset X_1 = X$  be the inverse image of the smooth locus  $U_0$  of  $X_0$  via the map  $\phi$  in Theorem 2.2. The map restricted to  $U_1$  is a symplectomorphism from  $U_1$  to  $U_0$ . Moreover,  $L$  maps to  $L_0$  via  $\phi$ . Since the polytope  $\Delta$  in  $M_{\mathbb{R}} \simeq \mathbb{R}^m$  is full dimensional and  $\mathbb{Q}$ -Gorenstein Fano, we may choose  $m$  facets  $\{F_1, F_2, \dots, F_m\}$  of  $\Delta$  such that the normal vectors  $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\} \subset N$  in the expression (3.3) are primitive integral and they form a basis for  $N_{\mathbb{Q}} \simeq \mathbb{Q}^m$ . For each  $j = 1, 2, \dots, m$ , we denote by  $u_{F_j}: \mathbb{D} \rightarrow X_0$  a gradient holomorphic disk of Maslov index two bounded by  $L_0$  and let  $\varphi_j := u_{F_j} \circ \phi$  be the pull-back of  $u_{F_j}$  along  $\phi: U_1 \rightarrow U_0$ . Note that  $\{[\varphi_j] \mid j = 1, 2, \dots, m\}$  is a  $\mathbb{Q}$ -basis of  $\mathbb{Q} \otimes_{\mathbb{Z}} \pi_1(L)$ .

Since  $\phi$  is a symplectomorphism,  $\varphi_j$  is of Maslov index two and its symplectic area is the same as that of  $u_j$ . Recall that the area of  $u_j$  is equal to the affine distance  $\lambda$  between the center and the facet  $F_j$  due to Archimedes. Therefore, we have

$$(3.7) \quad \lambda \cdot \mu_L([\varphi_j]) = 2 \cdot \omega([\varphi_j]).$$

In conclusion, combining (3.5), (3.6), and (3.7) together with the fact that  $\mu_L$  and  $\omega$  are homomorphisms, we have shown that

$$\lambda \cdot \mu_L(\beta) = 2 \cdot \omega(\beta)$$

for all  $\beta \in \pi_2(X, L)$ . Hence,  $L$  is monotone.  $\square$

**Remark 3.14.** In [GM22], Galkin–Mikhalkin proved an analog statement. They showed the monotonicity of a Lagrangian torus obtained from a symplectic parallel transport of the toric fiber at the center  $\mathbf{u}_0$  to a generic fiber.

### 3.3. Newton polytopes from disk counting

The goal of this subsection is to introduce two dual polytopes obtained by counting disks bounded by a monotone Lagrangian torus. One has been widely used to distinguish monotone Lagrangian tori in the previous literature and the other is its slight modification.

To each monotone Lagrangian torus  $L$  in a  $2m$ -dimensional monotone symplectic manifold  $(X, \omega)$ , one can assign a *disk potential*

$$W_L \in \mathbb{C}[z_1^\pm, \dots, z_m^\pm],$$

a Laurent polynomial which encodes counting invariants (a.k.a open Gromov-Witten invariants) of  $L$ . We briefly review a construction of the disk potential  $W_L$  of  $L$ .

Choose a  $\omega$ -compatible almost complex structure  $J$ . For each homotopy class  $\beta \in \pi_2(X, L)$  of Maslov index two, we denote by  $\mathcal{M}_1(L, J; \beta)$  the moduli space of  $J$ -holomorphic disks with one boundary marked point  $(\varphi, z_0)$  in the class  $\beta$  (modulo the automorphisms), that is,

$$\mathcal{M}_1(L, J; \beta) = \{(\varphi, z_0) \mid \varphi: (\mathbb{D}, \partial\mathbb{D}) \longrightarrow (X, L), z_0 \in \partial\mathbb{D}, J \circ d\varphi = d\varphi \circ j \text{ and } [\varphi] = \beta\} / \sim$$

where  $j$  is the standard complex structure on  $\mathbb{D}$ . The expected dimension of  $\mathcal{M}_1(L, J; \beta)$  is  $\dim_{\mathbb{R}} L + \#\{\text{marked points}\} + \mu(\beta) - \dim_{\mathbb{R}} \text{PSL}(2, \mathbb{R})$ . In this monotone case, by choosing an almost complex structure  $J$  generically, the moduli space becomes transversal when  $\mu(\beta) = 2$ . Moreover, the monotonicity condition ensures that the moduli space  $\mathcal{M}_1(L, J; \beta)$  with  $\mu(\beta) = 2$  is a compact smooth manifold *without* boundary. Taking an orientation and relative spin structure on the Lagrangian  $L$ , the degree of the evaluation map

$$\text{ev}: \mathcal{M}_1(L, J; \beta) \longrightarrow L, \quad (\varphi, z_0) \mapsto \varphi(z_0)$$

is defined and denoted by  $n(\beta, L, J)$ . In other words, the number  $n(\beta, L, J)$  is the (signed) counting of  $J$ -holomorphic disks passing through a generic point of  $L$ .

In general, for a general Lagrangian submanifold  $L$  (with a fixed choice of an orientation and a relative spin structure), the number  $n(\beta, L, J)$  *does* depend on a choice of  $J$ . When  $L$  is monotone, one can construct an oriented cobordism between two moduli spaces (with different choices of a generic almost complex structure) and hence the number  $n(\beta, L, J)$  becomes a deformation invariant as stated below.

**Proposition 3.15** ([EP97]). *Suppose that  $L$  is a monotone Lagrangian submanifold of  $(X, \omega)$ . For any class  $\beta \in \pi_2(X, L)$  of Maslov index two and generic compatible almost complex structures  $J$  and  $J'$ , the counting invariants  $n(\beta; L, J)$  and  $n(\beta; L, J')$  are equal.*

From now on, for simplicity,  $n(\beta; L, J)$  is denoted by  $n_\beta$ . To define a disk potential of  $L$ , we make a choice on a tuple  $\mathcal{B} = (\vartheta_1, \vartheta_2, \dots, \vartheta_m)$  of oriented loops that form a  $\mathbb{Z}$ -basis for  $\pi_1(L)$ . The choice leads to the identification  $\pi_1(L)$  with  $N := \mathbb{Z}^m$ . Namely, the basis elements in  $\mathcal{B}$  then correspond to the standard lattice vectors in  $N$ . Let  $\mathcal{L}$  be a trivial complex line bundle over  $L$  and consider a flat  $\mathbb{C}^*$ -connection  $\nabla$  on  $\mathcal{L}$ . We take a monomial

$z_j := \text{hol}_\nabla(\vartheta_j) \in (\mathbb{C}^*)^m$  for  $j = 1, 2, \dots, m$ . The set of those monomials forms a coordinate system for the moduli space of flat  $\mathbb{C}^*$ -connections on  $\mathcal{L}$  (modulo gauge equivalences).

For a lattice vector  $\mathbf{v} \in N$ , we set  $\mathbf{z}^{\mathbf{v}} = \mathbf{z}^{(v_1, \dots, v_m)}$  to be the monomial  $z_1^{v_1} \cdots z_m^{v_m}$ . The *disk potential* of  $L$  in [CO06, FOOO10] is defined by

$$(3.8) \quad W_{L, \mathcal{B}}: (\mathbb{C}^*)^m \longrightarrow \mathbb{C}, \quad W_{L, \mathcal{B}}(\mathbf{z}) = \sum_{\beta \in \pi_2(X, L)} n_\beta \cdot \mathbf{z}^{\partial\beta}$$

where the sum is taken over all homotopy classes of Maslov index two. We define the first combinatorial object associated to the disk potential  $W_{L, \mathcal{B}}$ .

**Definition 3.16.** We denote by  $\Delta_{L, \mathcal{B}}^{\text{disk}}$  the Newton polytope of the disk potential  $W_{L, \mathcal{B}}$  of a monotone Lagrangian torus  $L$ , that is,

$$\Delta_{L, \mathcal{B}}^{\text{disk}} := \text{the convex hull of } \left\{ \mathbf{v} \mid \sum_{\beta: \partial\beta = \mathbf{v}} n_\beta \neq 0 \right\} \text{ in } N_{\mathbb{R}}.$$

**Remark 3.17.** By the Gromov's compactness theorem, the number of  $J$ -holomorphic disks of Maslov index two is finite. Hence, the convex hull  $\Delta_{L, \mathcal{B}}^{\text{disk}}$  is compact (and hence a polytope).

**Remark 3.18.** In general, it is hard to find an explicit expression of  $W_{\text{disk}}$ . It is known that the disk potential can be immediately written from the facets of  $\Delta$  if the singular loci of  $X_0$  are of particular type, see [NNU10, BGM22].

Suppose that  $\mathcal{B}$  and  $\mathcal{B}'$  are two  $\mathbb{Z}$ -basis for  $\pi_1(L) \simeq N$ . Then there exists an invertible matrix  $A \in \text{GL}(m, \mathbb{Z})$  such that  $\mathcal{B}$  maps to  $\mathcal{B}'$  under the linear transformation  $\mathbf{v} \mapsto A\mathbf{v}$ . Two disk potentials  $W_{L, \mathcal{B}}$  and  $W_{L, \mathcal{B}'}$  are simply related by the monomial coordinate change defined by  $\mathbf{z}^{\mathbf{v}} \mapsto \mathbf{z}^{A\mathbf{v}}$ . In particular, the Newton polytopes  $\Delta_{L, \mathcal{B}}^{\text{disk}}$  and  $\Delta_{L, \mathcal{B}'}^{\text{disk}}$  are unimodularly equivalent.

**Definition 3.19.** Two polytopes  $\Delta$  and  $\Delta'$  are called *unimodularly equivalent* if  $\Delta$  maps to  $\Delta'$  under a unimodular equivalence, that is, an affine transformation  $\mathbf{v} \mapsto A\mathbf{v} + \mathbf{v}_0$  for some  $A \in \text{GL}(m, \mathbb{Z})$  and  $\mathbf{v}_0 \in \mathbb{R}^m$ .

From now on, two Newton polytopes are regarded as the same if they are unimodularly equivalent. We simply suppress  $\mathcal{B}$  in the notation  $\Delta_{L, \mathcal{B}}^{\text{disk}}$ , that is,  $\Delta_L^{\text{disk}} := \Delta_{L, \mathcal{B}}^{\text{disk}}$  for simplicity.

Suppose that  $\phi: X \rightarrow X$  is a symplectomorphism such that  $\phi(L_1) = L_2$ . We take a  $\mathbb{Z}$ -basis  $\phi_*\mathcal{B} = (\phi \circ \vartheta_1, \phi \circ \vartheta_2, \dots, \phi \circ \vartheta_m)$  consisting of oriented loops for  $\pi_1(L_2)$  and set  $z'_j := \text{hol}_\nabla(\phi \circ \vartheta_j)$ . By Proposition 3.15, two counting invariants  $n(\beta_1; L_1, J_1)$  and  $n(\phi_*\beta_1; L_2, \phi_*J_1)$  are equal and hence  $W_{L_1, \mathcal{B}}$  agrees with  $W_{L_2, \phi_*\mathcal{B}}$  under the transformation  $z_j \mapsto z'_j$  for  $j = 1, 2, \dots, m$ . We then have the following corollary.

**Corollary 3.20.** *Consider two monotone Lagrangian tori  $L_1$  and  $L_2$  in a symplectic manifold  $(X, \omega)$ . If there is a symplectomorphism  $\phi: X \rightarrow X$  such that  $\phi(L_1) = L_2$ , then the Newton polytopes  $\Delta_{L_1}^{\text{disk}}$  and  $\Delta_{L_2}^{\text{disk}}$  of disk potentials are unimodularly equivalent.*

**Remark 3.21.** This corollary has been employed to distinguish the constructed monotone Lagrangian tori, see [Aur15, Via16, Via17] for instance.

We introduce the Newton polytope  $\Delta_L^{\text{ref}}$  of the ‘‘refined’’ version of the disk potential  $W_L$  in the sense of effective disk classes. For motivation, we now discuss some drawbacks of this disk potential  $W_{\text{disk}}$  in (3.8) to distinguish monotone Lagrangian tori. Originally, the disk potential function  $W_{\text{disk}}$  in [FOOO09] was defined on the Maurer–Cartan space of  $L$ . To make the expression malleable, it is usually written in the form of a Laurent polynomial or series by restricting the Maurer–Cartan space to  $H^1(L)$  (provided  $H^1(L)$  is a subset of the

Maurer–Cartan space) and taking a basis  $\mathcal{B} = \{\vartheta_1, \vartheta_2, \dots, \vartheta_m\}$  for  $\pi_1(L) = \text{Hom}(H^1(L); \mathbb{Z})$ . However, throughout this process, we lose some information. Suppose that there are two distinct classes  $\beta$  and  $\beta'$  that can be realized as a holomorphic disk satisfying

- (1)  $\partial(\beta) = \partial(\beta')$  and
- (2)  $\mu(\beta) = \mu(\beta') = 2$ .

If their counting invariants are canceled, that is,  $n_\beta + n_{\beta'} = 0$ , then the disk potential  $W_{\text{disk}}$  cannot capture the existence of such disks. To capture this piece of information on the existence, we shall modify this disk potential which is more relevant to distinguish the constructed monotone Lagrangian tori in our situation.

**Definition 3.22.** Identify  $N \simeq \mathbb{Z}^m$  with the fundamental group  $\pi_1(L)$ . For a lattice vector  $\mathbf{v} \in N$ , we define

$$c_{\mathbf{v}} := \begin{cases} 1 & \text{if there exists } \beta \in \pi_2(X, L) \text{ such that } \partial\beta = \mathbf{v} \text{ and } n_\beta \neq 0, \\ 0 & \text{otherwise.} \end{cases}$$

The *refined disk potential* of  $L$  is defined by

$$(3.9) \quad W_L^{\text{ref}}(\mathbf{z}) := \sum_{\mathbf{v} \in N} c_{\mathbf{v}} \cdot z^{\mathbf{v}}.$$

**Definition 3.23.** We denote by  $\Delta_L^{\text{ref}}$  the Newton polytope of the refined disk potential  $W_L^{\text{ref}}$  of a monotone Lagrangian torus  $L$ , that is,

$$\Delta_L^{\text{ref}} := \text{the convex hull of } \{\mathbf{v} \mid \exists \beta \text{ such that } \mathbf{v} = \partial\beta \text{ and } n_\beta \neq 0\} \text{ in } N_{\mathbb{R}}.$$

We also have the invariance of the Newton polytope of a refined disk potential by Proposition 3.15.

**Corollary 3.24.** *Suppose that there are two monotone Lagrangian tori  $L_1$  and  $L_2$  in a symplectic manifold  $(X, \omega)$ . If there is a symplectomorphism  $\phi: X \rightarrow X$  such that  $\phi(L_1) = L_2$ , then the Newton polytopes  $\Delta_{L_1}^{\text{ref}}$  and  $\Delta_{L_2}^{\text{ref}}$  of refined disk potentials are unimodularly equivalent.*

In the remaining part of this section, we explore the relationship between  $\Delta^\vee$  in Definition 3.8 and  $\Delta_{L, \mathcal{B}}^{\text{ref}}$  in Definition 3.23. Recall that the Lagrangian torus  $L$  is constructed from the Newton–Okounkov polytope  $\Delta \subset M_{\mathbb{R}} \simeq \mathbb{R}^m$ . The set  $\mathcal{B}_0$  is taken as the oriented loops  $\{\vartheta_1, \vartheta_2, \dots, \vartheta_m\}$  in  $L$  respectively corresponding to the dual basis elements of the standard basis for  $M$  via  $N \simeq \pi_1(L)$ . From now on, we set  $\Delta_L^{\text{ref}} := \Delta_{L, \mathcal{B}_0}^{\text{ref}}$  for notational simplicity.

**Proposition 3.25.** *The combinatorial dual  $\Delta^\vee$  of a  $\mathbb{Q}$ -Gorenstein Fano polytope  $\Delta$  is contained in the Newton polytope of the refined disk potential  $W_L^{\text{ref}}$ , that is,  $\Delta^\vee \subset \Delta_L^{\text{ref}}$ .*

**Remark 3.26.** Recall that  $\Delta_{L, \mathcal{B}_0}^{\text{disk}}$  denotes the Newton polytope of the disk potential  $W_{L, \mathcal{B}_0}$  of  $L$ . The authors do not know whether  $\Delta^\vee \subset \Delta_{L, \mathcal{B}_0}^{\text{disk}}$  holds or not in a general setting. Recently, Galkin–Mikhalkin in [GM22] discovered an algebro-geometric condition on  $\Delta^\vee = \Delta_{L, \mathcal{B}_0}^{\text{disk}}$ . To apply their criterion, one needs to verify that both the toric variety  $(X_0, \text{Sing}(X_0))$  and the total space  $(\mathfrak{X}, \text{Sing}(\mathfrak{X}) = \text{Sing}(X_0))$  are  $\mathbb{Q}\Gamma$ -spaces, see the definition therein. In our criterion, we do *not* need to check that  $(\mathfrak{X}, \text{Sing}(\mathfrak{X}))$  is a  $\mathbb{Q}\Gamma$ -space because we introduce the refined disk potential and make use of the map  $\phi$  in Theorem 2.2.

We denoted by  $F_j$  the facet of  $\Delta$  contained in the hyperplane  $H_{\mathbf{v}_j, 1}$  in (3.3). Let  $\beta_j \in \pi_2(X, L)$  be a homotopy class represented by a gradient holomorphic disk intersecting the inverse image of  $F_j$ . Such a class is called *basic*. To orient the moduli space  $\mathcal{M}(L; \beta_j)$ , we take a spin structure and an orientation of  $L$  given by the torus action.



**Lemma 3.27.** *For each  $j = 1, 2, \dots, \kappa$ , the counting invariant of  $\beta_j$  is one, that is,  $n_{\beta_j} = 1$ . In particular, the refined disk potential of  $L$  is of the form*

$$W_L^{\text{ref}}(\mathbf{z}) = \sum_{j=1}^{\kappa} \mathbf{z}^{\mathbf{v}_j} + \widetilde{W}_L^{\text{ref}}(\mathbf{z})$$

where  $\widetilde{W}_L^{\text{ref}}(\mathbf{z})$  is a subtraction-free Laurent polynomial.

*Proof.* Let  $\beta$  be a basic class of  $(X, L)$  corresponding to a facet  $F$ . To compute the counting invariant  $n_{\beta}$ , we exploit the toric degeneration  $\mathfrak{X}$  of  $X$ . Set  $L_{\tau} := \phi_{1,\tau}(L)$ . By using the toric structure on  $X_0$ , we generate a gradient holomorphic disk  $\varphi: (\mathbb{D}, \partial\mathbb{D}) \rightarrow (X_0, L_0)$  that intersects with the inverse image of the relative interior of  $F$ . The image of  $\varphi$  is contained in the smooth locus  $U_0 := X_0 \setminus \text{Sing}(X_0)$ .

Choose a compatible almost complex structure  $J_1$  on  $X_1$  as an extension of the pull-back of the toric complex structure on an open subset of  $\phi^{-1}(U_0)$  by the commutativity of (2.2). By taking the interpolation part sufficiently small enough for the extension, we may assume the composition  $\phi^{-1} \circ \varphi$  becomes a  $J_1$ -holomorphic disk. In particular,  $\phi_*(\beta) = [\varphi]$ .

Let  $\widetilde{X}_0$  be a simplicialization of the central toric variety  $X_0$ . It comes with a toric morphism  $\Pi: \widetilde{X}_0 \rightarrow X_0$ . The inverse image  $\widetilde{L}_0 := \Pi^{-1}(L_0)$  is a Lagrangian toric fiber of  $\widetilde{X}_0$  because the algebraic torus of  $\widetilde{X}_0$  maps into the algebraic torus of  $X_0$ . We denote by  $\widetilde{\beta}$  the class represented by a strict transformation  $\widetilde{\varphi}$  of  $\varphi$ . On the other hand, since the toric morphism  $\Pi$  is holomorphic, the map  $\Pi$  induces

$$\Pi_*: \mathcal{M}(\widetilde{L}_0, \widetilde{J}_0; \widetilde{\beta}) \longrightarrow \mathcal{M}(L_0, J_0; \phi_*(\beta)) \quad \varphi' \mapsto \Pi \circ \varphi'$$

where  $J_0$  and  $\widetilde{J}_0$  are the standard complex structure.

We claim that the map  $\Pi_*$  is an orientation preserving diffeomorphism. To see it, we first note that  $\varphi$  is equal to  $\Pi \circ \widetilde{\varphi}$  in  $\mathcal{M}(L_0, J_0; \phi_*(\beta))$  by the construction of strict transformations. Also,  $\varphi'$  is equal to the strict transformation of  $\Pi \circ \varphi'$  in  $\mathcal{M}(\widetilde{L}_0, \widetilde{J}_0; \widetilde{\beta})$ . It is because the class  $\widetilde{\beta}$  of Maslov index two is regular and hence spheres cannot bubble off. The classification of holomorphic disks in [CP14] yields that two holomorphic disks in the basic class  $\widetilde{\beta}$  having the same boundary are equal. Thus, the map  $\Pi_*$  is a diffeomorphism. By choosing a spin structure and orientation of  $L$  induced from the torus action, the counting invariant is one, that is,  $n_{\widetilde{\beta}} = 1$  again by the classification of holomorphic disks in [CP14]. Then the map  $\Pi_*$  is orientation preserving.

Combining the above, we obtain the following commutative diagram

$$\begin{array}{ccccc} \mathcal{M}_1(L_1, J_1; \beta) & \xrightarrow{\phi_*} & \mathcal{M}_1(L_0, J_0; \phi_*(\beta)) & \xleftarrow{\Pi_*} & \mathcal{M}_1(\widetilde{L}_0, \widetilde{J}_0; \widetilde{\beta}) \\ \text{ev} \downarrow & & \downarrow \text{ev} & & \text{ev} \downarrow \\ L_1 & \xrightarrow{\phi} & L_0 & \xleftarrow{\Pi} & \widetilde{L}_0. \end{array}$$

such that every arrow is an orientation preserving diffeomorphism. In particular,  $n_{\beta} = 1$ .  $\square$

*Proof of Proposition 3.25.* By Lemma 3.27, the inclusion is derived.  $\square$

#### 4. Infinitely many distinct monotone Lagrangian tori via cluster mutations

Let  $X$  be a smooth Fano variety of complex dimension  $m$  polarized by a very ample line bundle  $\mathcal{L}$ , a positive power of the anticanonical bundle of  $X$ . Thanks to Proposition 3.13,  $X$

possesses a monotone Lagrangian torus if  $X$  admits the normal toric degeneration associated with a  $\mathbb{Q}$ -Gorenstein Fano Newton–Okounkov polytope. Suppose that  $X$  has an *infinite* family of such  $\mathbb{Q}$ -Gorenstein Fano and normal toric degenerations. Then we produce an infinite family of monotone Lagrangian tori of  $X$ . The goal of this section is to devise a criterion for the family to contain infinitely many *distinct* monotone Lagrangian tori when the family of Newton–Okounkov polytopes has a *tropical cluster structure*, see Definition 4.3 for the precise meaning.

#### 4.1. Cluster varieties and cluster algebras

We begin by reviewing some notions on skew-symmetrizable cluster algebras and varieties, following Fock–Goncharov [FG09] and Gross–Hacking–Keel [GHK15a].

We fix the following data  $\Gamma$  (called a *fixed data*) consisting of

- a lattice  $N$  of finite rank with a skew-symmetric bilinear form

$$\{\cdot, \cdot\}: N \times N \longrightarrow \mathbb{Q},$$

- an *unfrozen* sublattice  $N_{\text{uf}} \subset N$ , a saturated sublattice of  $N$ ,
- an index set  $J$  such that  $|J| = \text{rank } N$ ,
- a subset  $J_{\text{uf}}$  of  $J$  such that  $|J_{\text{uf}}| = \text{rank } N_{\text{uf}}$ ,
- a sublattice  $N^\circ \subset N$  of finite index such that

$$\{N_{\text{uf}}, N^\circ\} \subset \mathbb{Z} \text{ and } \{N, N_{\text{uf}} \cap N^\circ\} \subset \mathbb{Z},$$

- positive integers  $d_j$  for  $j \in J$  of which greatest common divisor is one.

For the fixed data  $\Gamma$ , a *seed* is a labeled collection of elements of  $N$

$$(4.1) \quad \mathbf{s} := (\mathbf{e}_j \mid j \in J)$$

such that  $\{\mathbf{e}_j \mid j \in J\}$  is  $\mathbb{Z}$ -basis for  $N$ ,  $\{\mathbf{e}_j \mid j \in J_{\text{uf}}\}$  is a  $\mathbb{Z}$ -basis of  $N_{\text{uf}}$ , and  $\{d_j \mathbf{e}_j \mid j \in J\}$  is a  $\mathbb{Z}$ -basis for  $N^\circ$ .

We set  $\varepsilon_{r,s} := \{\mathbf{e}_r, \mathbf{e}_s\} d_s$ . Note that  $\varepsilon_{r,s} \in \mathbb{Z}$  if at least one of  $r$  and  $s$  is in  $J_{\text{uf}}$ . We then obtain a (not necessarily skew-symmetric) matrix  $\varepsilon = (\varepsilon_{r,s})_{r,s \in J} \in M_{J \times J}(\mathbb{Q})$ . The matrix is called an *exchange matrix* of  $\Gamma$ . Note that

$$(4.2) \quad \varepsilon_{r,s} d_r + \varepsilon_{s,r} d_s = 0.$$

Let  $M := \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$  be the dual lattice of  $N$ . We denote by  $\{\mathbf{e}_j^* \mid j \in J\}$  the dual basis of  $\{\mathbf{e}_j \mid j \in J\}$  for  $M$ . Let  $M^\circ := \text{Hom}_{\mathbb{Z}}(N^\circ, \mathbb{Z})$  be the dual lattice of  $N^\circ$ . We also have a  $\mathbb{Z}$ -basis  $\{\mathbf{f}_j := d_j^{-1} \mathbf{e}_j^* \mid j \in J\}$  for  $M^\circ$ . To each seed  $\mathbf{s}$ , we associate the dual pair of an  $\mathcal{A}$ -torus  $\mathcal{A}_{\mathbf{s}}$  and an  $\mathcal{X}$ -torus  $\mathcal{A}_{\mathbf{s}}^\vee$  defined by

$$(4.3) \quad \mathcal{A}_{\mathbf{s}} := T_{N^\circ} := \text{Spec}(\mathbb{C}[M^\circ]) \text{ and } \mathcal{A}_{\mathbf{s}}^\vee := T_M := \text{Spec}(\mathbb{C}[N]).$$

For a given fixed data  $\Gamma$  and a chosen seed  $\mathbf{s}_0$ , we can produce seeds and exchange matrices by mutating the seed  $\mathbf{s}_0$  into all possible directions in  $J_{\text{uf}}$  inductively. The chosen seed  $\mathbf{s}_0$  at the beginning is called an *initial seed*. For each index  $k \in J_{\text{uf}}$ , the *mutation in the  $k$ -direction* of various data is defined below.

**Definition 4.1.** Let  $[a]_+ := \max\{a, 0\}$  and  $\text{sgn}(a) :=$  the sign of  $a$ . For an index  $k \in J_{\text{uf}}$ , the *mutation of a seed  $\mathbf{s}$  in the  $k^{\text{th}}$ -direction* is a seed  $\mu_k(\mathbf{s}) := \mathbf{s}'$  defined by

$$(4.4) \quad \mathbf{s}' := (\mathbf{e}'_j \mid j \in J), \quad \mathbf{e}'_j := \begin{cases} -\mathbf{e}_j & \text{if } j = k, \\ \mathbf{e}_j + [\varepsilon_{j,k}]_+ \mathbf{e}_k & \text{if } j \neq k. \end{cases}$$

Here, the exchange matrix  $(\varepsilon_{j,k})$  is associated with the seed  $\mathbf{s}$ . The basis element  $\mathbf{e}'_j$  is also denoted by  $\mu_k(\mathbf{e}_j)$ .

Then the set  $\{\mathbf{e}'_j \mid j \in J\}$  is still a  $\mathbb{Z}$ -basis for  $N$ . The subset  $\{\mathbf{e}'_j \mid j \in J_{\text{uf}}\}$  is a  $\mathbb{Z}$ -basis for  $N_{\text{uf}}$  and  $\{d_j \mathbf{e}'_j \mid j \in J\}$  also forms a  $\mathbb{Z}$ -basis for  $N^\circ$ . Note that the basis  $\{\mathbf{f}'_j \mid j \in J\}$  is obtained from  $\{\mathbf{f}_j \mid j \in J\}$  via the formula

$$(4.5) \quad \mathbf{f}'_j := \begin{cases} -\mathbf{f}_k + \sum_i [\varepsilon_{k,i}]_+ \mathbf{f}_i & \text{if } j = k, \\ \mathbf{f}_j & \text{if } j \neq k \end{cases}$$

because  $\varepsilon_{r,s} d_r + \varepsilon_{s,r} d_s = 0$ . Also, under the mutation  $\mu_k$  in (4.4), the exchange matrix is also mutated via the formula  $\mu_k(\varepsilon) := \varepsilon'$  given by

$$(4.6) \quad \varepsilon' := (\varepsilon'_{r,s})_{r,s \in J}, \quad \varepsilon'_{r,s} := \{\mathbf{e}'_r, \mathbf{e}'_s\} d_s = \begin{cases} -\varepsilon_{r,s} & \text{if } r = k \text{ or } s = k, \\ \varepsilon_{r,s} + \text{sgn}(\varepsilon_{k,s}) [\varepsilon_{r,k} \varepsilon_{k,s}]_+ & \text{otherwise.} \end{cases}$$

As in (4.3), each seed assigns an  $\mathcal{A}$ -torus and an  $\mathcal{X}$ -torus. We define a birational transformation between  $\mathcal{A}$ -tori (resp. between  $\mathcal{X}$ -tori). For each seed  $\mathbf{s} = (\mathbf{e}_j \mid j \in J)$ , we associate the coordinate functions  $(X_j := z^{\mathbf{e}_j} \mid j \in J)$  on the  $\mathcal{X}$ -torus  $\mathcal{A}_{\mathbf{s}}^\vee$ . Dually, we associate the coordinate functions  $(A_j := z^{\mathbf{f}_j} \mid j \in J)$  on the  $\mathcal{A}$ -torus  $\mathcal{A}_{\mathbf{s}}$ . For  $\mathbf{s}' = \mu_k(\mathbf{s})$ , we denote by  $A'_j$  the coordinate function  $z^{\mathbf{f}'_j}$  on  $\mathcal{A}_{\mathbf{s}'}$  corresponding to  $\mathbf{f}'_j$  and have

$$(4.7) \quad \mu_k: \mathcal{A}_{\mathbf{s}} \dashrightarrow \mathcal{A}_{\mathbf{s}'}, \quad \mu_k^*(A'_j) = \begin{cases} A_k^{-1} \left( \prod_{i \in J} A_i^{[\varepsilon_{k,i}]_+} + \prod_{i \in J} A_i^{[-\varepsilon_{k,i}]_+} \right) & \text{if } j = k, \\ A_j & \text{if } j \neq k. \end{cases}$$

For  $\mathbf{s}' = \mu_k(\mathbf{s})$ , we denote by  $X'_j$  the coordinate function  $z^{\mathbf{e}'_j}$  on  $\mathcal{A}_{\mathbf{s}'}^\vee$  and have

$$(4.8) \quad \mu_k: \mathcal{A}_{\mathbf{s}}^\vee \dashrightarrow \mathcal{A}_{\mathbf{s}'}^\vee, \quad \mu_k^*(X'_j) = \begin{cases} X_j^{-1} & \text{if } j = k, \\ X_j \left( 1 + X_k^{-\text{sgn}(\varepsilon_{j,k})} \right)^{-\varepsilon_{j,k}} & \text{if } j \neq k. \end{cases}$$

To parametrize the seeds, let  $\mathbb{T}$  be the oriented rooted tree with  $|J_{\text{uf}}|$  outgoing edges from each vertex labeled by the elements of  $J_{\text{uf}}$ . The root  $t_0$  corresponds to the initial seed  $\mathbf{s}_0$ . If two vertices  $t$  and  $t'$  are connected by a directed edge from  $t$  to  $t'$ , then we denote by

$$(4.9) \quad \mu_k(t) = t'.$$

Also, the seed and the exchange matrix corresponding to  $t$  are denoted by  $\mathbf{s}_t$  and  $\varepsilon_t$ , respectively. By gluing the  $\mathcal{A}$ -tori via the birational transformations (4.7), we obtain a scheme

$$(4.10) \quad \mathcal{A}_{\Gamma, \mathbf{s}_0} = \bigcup_{t \in \mathbb{T}} \mathcal{A}_{\mathbf{s}_t},$$

which is called an  $\mathcal{A}$ -cluster variety of the chosen data  $(\Gamma, \mathbf{s}_0)$ . Dually, an  $\mathcal{X}$ -cluster variety of the data  $(\Gamma, \mathbf{s}_0)$  is defined by gluing the  $\mathcal{X}$ -tori via the birational transformations (4.8)

$$(4.11) \quad \mathcal{A}_{\Gamma, \mathbf{s}_0}^\vee = \bigcup_{t \in \mathbb{T}} \mathcal{A}_{\mathbf{s}_t}^\vee.$$

The  $\mathcal{X}$ -cluster variety  $\mathcal{A}_{\Gamma, \mathbf{s}_0}^\vee$  is also called the (Fock–Goncharov) dual of  $\mathcal{A}_{\Gamma, \mathbf{s}_0}$ .

We also deal with cluster algebras and upper cluster algebras later on and we briefly recall them. From the fixed data  $\Gamma$  and an initial seed  $\mathbf{s}_0$ , consider the function field  $\mathbb{C}(\mathcal{A}_{\mathbf{s}_0})$  of the  $\mathcal{A}$ -torus  $\mathcal{A}_{\mathbf{s}_0}$ . Suppose that  $\mathbf{s}_t$  is obtained by applying a finite sequence of mutations in (4.4) to  $\mathbf{s}$ , that is,  $\mathbf{s}_t = \mu_{t_0, t}(\mathbf{s}_{t_0})$  for some  $\mu_{t_0, t} := \mu_{k_\ell} \circ \cdots \circ \mu_{k_2} \circ \mu_{k_1}$ . For  $j \in J$ , we denote by  $A_{j, \mathbf{s}_t}$  the coordinate function corresponding to  $\mathbf{f}_{j, \mathbf{s}_t}$  on the torus  $\mathcal{A}_{\mathbf{s}_t}$  associated with  $t$ . By the Laurent phenomenon in [FZ02], every coordinate function  $A_{j, \mathbf{s}_t}$  for  $\mathcal{A}_{\mathbf{s}_t}$  can be expressed as a Laurent polynomial of the variables  $\{A_{j, t_0} \mid j \in J\}$  by pulling  $A_{j, t}$  via the sequence of mutations corresponding to  $\mu_{t_0, t}$  for  $\mathcal{A}$ -cluster charts in (4.7). By abuse of notation, we

denote by  $\mu_{t_0,t}$  the corresponding sequence of mutations in (4.7) and denote the Laurent polynomial

$$(4.12) \quad A_{j,t} := \mu_{t,t_0}^*(A_{j,s_t}) \in \mathbb{C}(\mathcal{A}_{s_0}).$$

We define upper cluster algebras and ordinary cluster algebras associated with  $(\Gamma, s_0)$ .

**Definition 4.2.** An *upper cluster algebra*  $\text{up}(\mathcal{A})$  is defined by

$$\text{up}(\mathcal{A}) = \bigcap_{t \in \mathbb{T}} \mathbb{C}[A_{j,t}^{\pm 1} \mid j \in J] \subset \mathbb{C}(\mathcal{A}_{s_0}).$$

An (*ordinary*) *cluster algebra*  $\text{ord}(\mathcal{A})$  is defined by the  $\mathbb{C}$ -subalgebra of  $\mathbb{C}(\mathcal{A}_{s_0})$  generated by

$$\{A_{j,t} \mid t \in \mathbb{T}, j \in J_{\text{uf}}\} \cup \left\{ A_{j,t}^{\pm 1} \mid t \in \mathbb{T}, j \in J \setminus J_{\text{uf}} \right\}.$$

Each element of the tuple  $(A_{j,t})_{j \in J}$  is called a *cluster variable*.

Indeed, the upper/ordinary cluster algebras in Definition 4.2 can be defined from smaller pieces of information than a fixed data  $\Gamma$  and a seed  $s_0$  in (4.1). For the later usage of cluster algebras in Chapter 5 and 6, we recall Fomin–Zelevinsky’s way of constructing cluster algebras in [FZ02]. Let  $\mathbb{C}(z_j \mid j \in J)$  be the field of rational functions. A *Fomin–Zelevinsky’s seed*  $s$  consists of

- a  $J$ -tuple  $(A_j)_{j \in J}$  of elements that is a free generating set of  $\mathbb{C}(z_j \mid j \in J)$
- a matrix  $\varepsilon = (\varepsilon_{r,s})_{r \in J_{\text{uf}}, s \in J} \in M_{J_{\text{uf}} \times J}(\mathbb{Z})$  such that (4.2) holds for all  $r, s \in J_{\text{uf}}$ .

The matrix  $\varepsilon$  is called an *extended exchange matrix*. Out of a seed  $((A_j)_{j \in J}, \varepsilon)$ , we can produce Fomin–Zelevinsky’s seeds by mutating the seed  $s_0$  into all possible directions in  $J_{\text{uf}}$  inductively. The mutation formulas for Fomin–Zelevinsky’s seeds are given in (4.7) and (4.6). We then can produce  $\text{up}(\mathcal{A})$  and  $\text{ord}(\mathcal{A})$  in Definition 4.2.

In this case, the mutations are involutive and hence it suffices to employ an unoriented  $|J_{\text{uf}}|$ -regular tree whose edges are labeled by  $J_{\text{uf}}$  to parametrize the Fomin–Zelevinsky’s seeds. By abuse of notation, we also denote by  $\mathbb{T}$  this tree and call it the *exchange graph*.

## 4.2. Tropicalized cluster mutations and $\mathbb{Q}$ -Gorenstein Fano polytopes

Let  $X$  be a smooth projective variety polarized by a very ample line bundle  $\mathcal{L}$ . Let us take a reference section  $h \in \mathcal{L} := H^0(X, \mathcal{L})$  as in (2.4). Suppose that there exist a fixed data  $\Gamma$  and an initial seed  $s_0 = s_{t_0}$ . We inductively produce the set  $\{s_t \mid t \in \mathbb{T}\}$  of seeds generated by  $s_{t_0}$  as in Section 4.1 and parametrized by the vertices of an oriented rooted tree  $\mathbb{T}$  with  $|J_{\text{uf}}|$  outgoing edges labeled by the elements of  $J_{\text{uf}}$  from each vertex.

Suppose that we have a family of valuations with one-dimensional leaves and parametrized by the vertices of  $\mathbb{T}$ , say  $\{v_t \mid t \in \mathbb{T}\}$ . Assume that the semigroup  $S_t := S(\mathcal{L}, v_t, h)$  constructed from  $v_t$  is finitely generated. (Recall that every semigroup obtained from a valuation generated by a cluster algebra is always finitely generated, see [GHKK18, Section 8.3].) Let  $\Delta_t := \Delta(\mathcal{L}, v_t, h)$  be the Newton–Okounkov polytope of  $X$  corresponding to  $S_t$  for each  $t \in \mathbb{T}$  in (2.5). Using the seed  $s_t$ , each Newton–Okounkov polytope  $\Delta_t$  (resp. semigroup  $S_t$ ) in the family can be regarded as a subset of  $M_{\mathbb{R}} = \mathbb{R}\langle \mathbf{e}_j^* \mid j \in J \rangle = \left\{ \sum_j u_j \mathbf{e}_j^* \mid u_j \in \mathbb{R} \text{ for all } j \in J \right\} \simeq \mathbb{R}^J$  (resp.  $\mathbb{R}_{\geq 0} \times M_{\mathbb{R}} \simeq \mathbb{R}_{\geq 0} \times \mathbb{R}^J$ ).

We now describe relation between semigroups  $S_t$  and  $S_{t'}$  and between Newton–Okounkov polytopes  $\Delta_t$  and  $\Delta_{t'}$ .

**Definition 4.3.** Consider a family  $\{S_t \subset \mathbb{R}_{\geq 0} \times M_{\mathbb{R}} \mid t \in \mathbb{T}\}$  of semigroups parametrized by  $\mathbb{T}$  and the family  $\{\Delta_t \subset M_{\mathbb{R}} \mid t \in \mathbb{T}\}$  of associated Newton–Okounkov bodies, which are also parametrized by  $\mathbb{T}$ .

- (1) For an unfrozen index  $k \in J_{\text{uf}}$  and a pair  $(t, t')$  of vertices with  $\mu_k(t) = t'$  in  $\mathbb{T}$ , a *tropicalized cluster mutation in the  $k^{\text{th}}$ -direction*

$$\mu_k^T : M_{\mathbb{R}} \simeq_{s_t} \mathbb{R}^J \longrightarrow M_{\mathbb{R}} \simeq_{s_{t'}} \mathbb{R}^J$$

is defined to be a piecewise-linear transformation defined by

$$(4.13) \quad \mathbf{u} := (u_j) \mapsto \mathbf{u}' := (u'_j), \quad u'_j = \begin{cases} -u_j & \text{if } j = k \\ u_j + [\varepsilon_{k,j}]_+ u_k & \text{if } j \neq k \text{ and } u_k \geq 0 \\ u_j + [-\varepsilon_{k,j}]_+ u_k & \text{if } j \neq k \text{ and } u_k \leq 0 \end{cases}$$

where  $\varepsilon_t = (\varepsilon_{i,j})_{i,j \in J}$  is the exchange matrix associated with  $t$ .

- (2) We also define

$$(4.14) \quad \widehat{\mu}_k^T : \mathbb{R}_{\geq 0} \times M_{\mathbb{R}} \simeq_{s_t} \mathbb{R}_{\geq 0} \times \mathbb{R}^J \longrightarrow \mathbb{R}_{\geq 0} \times M_{\mathbb{R}} \simeq_{s_{t'}} \mathbb{R}_{\geq 0} \times \mathbb{R}^J$$

by  $\widehat{\mu}_k^T(r, \mathbf{u}) := (r, \mu_k^T(\mathbf{u}))$ . This map  $\widehat{\mu}_k^T$  is also called a *tropicalized cluster mutation in the  $k^{\text{th}}$ -direction*.

- (3) The family of semigroups parametrized by  $\mathbb{T}$  is said to have a *tropical cluster structure* if for every pair  $(t, t')$  of vertices with  $t' = \mu_k(t)$ , the associated semigroups  $S_t$  and  $S_{t'}$  are related by the tropicalized cluster mutation  $\widehat{\mu}_k^T$  in the  $k^{\text{th}}$ -direction, that is,  $S_{t'} = \widehat{\mu}_k^T(S_t)$ .
- (4) The family of Newton–Okounkov bodies of  $X$  parametrized by  $\mathbb{T}$  is said to have a *tropical cluster structure* if for every pair  $(t, t')$  of vertices with  $t' = \mu_k(t)$ , the associated Newton–Okounkov bodies  $\Delta_t$  and  $\Delta_{t'}$  are related by the tropicalized cluster mutation  $\mu_k^T$  in the  $k^{\text{th}}$ -direction, that is,  $\Delta_{t'} = \mu_k^T(\Delta_t)$ .

**Remark 4.4.** We outline a scenario where a smooth projective variety  $X$  has a tropical cluster structure. The reader is referred to [FO20, Section 4] for a detailed explanation. Consider a cluster ensemble  $(\mathcal{A}_{\Gamma, s_0}, \mathcal{A}_{\Gamma, s_0}^{\vee})$  defined in (4.10) and (4.11). Assume that  $X$  is a compactification of the  $\mathcal{A}$ -cluster variety  $\mathcal{A}_{\Gamma, s_0}$  and let  $X^{\vee}$  be a (partial compactification) of its dual  $\mathcal{X}$ -cluster variety  $\mathcal{A}_{\Gamma, s_0}^{\vee}$ . Suppose that the compactification  $X$  is a smooth Fano projective variety and a Landau–Ginzburg mirror of  $X$  is given by a regular function  $W$  on the  $\mathcal{X}$ -cluster variety  $X^{\vee}$ . In [GHKK18], they constructed a theta basis for a certain class of cluster algebras from a scattering diagram. When a seed  $\mathbf{s}$  is fixed, the  $g$ -vectors of this theta basis are parametrized by the tropical integer points of the Newton–Okounkov polytope, which is constructed from a valuation from the exchange matrix associated with  $\mathbf{s}$ . In this case, every set of  $g$ -vectors from a seed and the set of  $g$ -vectors from the initial seed are related by a finite sequence of tropicalized ( $\mathcal{X}$ -)cluster mutations and so are the corresponding Newton–Okounkov polytopes. Also, it is expected that the Newton–Okounkov polytopes are given by the tropicalization of the LG mirror  $(X^{\vee}, W)$  restricted to a cluster chart.

Suppose that the family  $\{S_t \mid t \in \mathbb{T}\}$  of finitely generated semigroups has a tropical cluster structure and hence the family  $\{\Delta_t \mid t \in \mathbb{T}\}$  of corresponding Newton–Okounkov bodies of  $X$  also has a tropical cluster structure. By Theorem 2.4, each  $\Delta_t$  is a rational polytope. The following lemmas discuss that if the toric degeneration at the initial seed is  $\mathbb{Q}$ -Gorenstein Fano and normal, then so are all the other toric degenerations.

**Lemma 4.5.** *Suppose that the semigroup  $S_{t_0}$  at the initial seed is saturated. Then, for each  $t \in \mathbb{T}$ , the semigroup  $S_t$  is also saturated.*

*Proof.* By induction, it suffices to prove that if  $\mu_k(t) = t'$  for  $k \in J_{\text{uf}}$  and  $S_t$  is saturated, then so is  $S_{t'}$ . We need to check that if  $n \cdot (r, \mathbf{u}') \in S_{t'}$  for  $(r, \mathbf{u}') \in \mathbb{N} \times \mathbb{Z}^J$  and  $n \in \mathbb{N}$ ,

then  $(r, \mathbf{u}') \in S_{t'}$ . Since  $\mu_k^T$  is a bijection on the lattice  $\mathbb{Z}^J$  and  $\widehat{\mu}_k^T$  is a bijection from  $S_t$  to  $S_{t'}$ , there exists  $(r, \mathbf{u}) \in \mathbb{N} \times \mathbb{Z}^J$  such that  $\widehat{\mu}_k^T(r, \mathbf{u}) = (r, \mathbf{u}')$  and  $n \cdot (r, \mathbf{u}) \in S_t$ . Since  $S_t$  is saturated,  $(r, \mathbf{u}) \in S_t$  and hence  $(r, \mathbf{u}') \in S_{t'}$ .  $\square$

**Lemma 4.6.** *Suppose that the Newton–Okounkov polytope  $\Delta_{t_0}$  at the initial seed is (resp. normalized)  $\mathbb{Q}$ -Gorenstein Fano with the center  $\mathbf{0}$ . Then, for every  $t \in \mathbb{T}$ , the corresponding Newton–Okounkov polytope  $\Delta_t$  is also (resp. normalized)  $\mathbb{Q}$ -Gorenstein Fano with the center  $\mathbf{0}$ .*

*Proof.* By induction, it suffices to prove that if  $\mu_k(t) = t'$  for some  $k \in J_{\text{uf}}$  and  $\Delta_t$  is  $\mathbb{Q}$ -Gorenstein Fano with the center at the origin  $\mathbf{0}$ , then so is  $\Delta_{t'}$ . Since  $\Delta_t$  is  $\mathbb{Q}$ -Gorenstein Fano, there are  $c \in \mathbb{Q}_{>0}$  and *primitive* lattice vectors  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_\kappa$  such that  $\Delta_t$  is expressed as

$$\Delta_t = \bigcap_{j=1}^{\kappa} H_{c\mathbf{v}_j, 1}^+$$

satisfying the conditions for (3.3). Since  $\Delta_{t'}$  also contains the origin in the interior,  $\Delta_{t'}$  can be expressed as

$$(4.15) \quad \Delta_{t'} = \bigcap_{j=1}^{\kappa'} H_{c\mathbf{v}'_j, 1}^+ \quad \text{where } \mathbf{v}'_j \in N_{\mathbb{R}} \text{ for } j = 1, 2, \dots, \kappa'$$

satisfying the conditions for (3.3).

We claim that every  $\mathbf{v}'_j$  in (4.15) is a primitive lattice vector. From the explicit expression of the tropicalized cluster mutation  $\mu_k^T$  in (4.13), depending on the sign of  $u_k$ , the image  $\mu_k^T(\mathbf{u})$  is determined by two linear transformations. Let us denote  $\mu_k^T(\mathbf{u})$  by  $A\mathbf{u}$  or  $B\mathbf{u}$  for some  $A, B \in \text{GL}(J, \mathbb{Z})$ . Then the half space  $H_{c\mathbf{v}_j, 1}^+$  corresponds to  $H_{c(A^T)^{-1}\mathbf{v}_j, 1}^+$ ,  $H_{c(B^T)^{-1}\mathbf{v}_j, 1}^+$ , or both. Consequently,

$$\{\mathbf{v}'_1, \mathbf{v}'_2, \dots, \mathbf{v}'_{\kappa'}\} \subset \{(A^T)^{-1}\mathbf{v}_1, \dots, (A^T)^{-1}\mathbf{v}_\kappa\} \cup \{(B^T)^{-1}\mathbf{v}_1, \dots, (B^T)^{-1}\mathbf{v}_\kappa\}.$$

If  $A \in \text{GL}(J, \mathbb{Z})$  and  $\mathbf{v}$  is a primitive lattice vector, then  $A\mathbf{v}$  is also primitive. Thus, every vector  $\mathbf{v}'_j$  is a primitive lattice vector and  $\Delta_{t'}$  is also  $\mathbb{Q}$ -Gorenstein Fano with the center  $\mathbf{0}$ .  $\square$

In general, the origin  $\mathbf{0} \in M_{\mathbb{R}}$  may *not* be the center of a  $\mathbb{Q}$ -Gorenstein Fano Newton–Okounkov polytope  $\Delta$  according to our definition of  $\mathbb{Q}$ -Gorenstein Fano polytope in Definition 3.7. To apply Lemma 4.6, we need to translate  $\Delta$  to position the center at the origin  $\mathbf{0}$ . In general, however, the translations and the tropicalized cluster mutations may *not* commute. We explore a condition that ensures commutativity.

Recall that for each  $t \in \mathbb{T}$ , a point  $\mathbf{u}$  of  $M_{\mathbb{R}}$  can be regarded as a point in  $\mathbb{R}^J \simeq_{s_t} M_{\mathbb{R}}$ . Let  $(t, t')$  be a pair of vertices with  $\mu_k(t) = t'$  in  $\mathbb{T}$  for an index  $k \in J_{\text{uf}}$ . Fix a point  $\mathbf{u}_0 \in M_{\mathbb{R}}$ . For each  $t \in \mathbb{T}$ , let  $\mathbf{u}_{t,0}$  be the image of  $\mathbf{u}_0$  under the identification given by  $s_t$ :

$$(4.16) \quad M_{\mathbb{R}} \simeq_{s_t} \mathbb{R}^J \quad (\mathbf{u}_0 \mapsto \mathbf{u}_{t,0}).$$

We denote by  $\tau_t: \mathbb{R}^J \rightarrow \mathbb{R}^J$  the translation defined by  $\mathbf{u} \mapsto \mathbf{u} - \mathbf{u}_{t,0}$ .

**Lemma 4.7.** *Let  $(t, t')$  be a pair of vertices with  $\mu_k(t) = t'$  in  $\mathbb{T}$  for an index  $k \in J_{\text{uf}}$ . Fix a point  $\mathbf{u}_0 \in M_{\mathbb{R}}$ . Let  $\mathbf{u}_{t,0}$  be the image of the point  $\mathbf{u}_0$  under (4.16). Then the following are equivalent.*

- (1) *The point  $\mathbf{u}_{t,0}$  is fixed under the tropicalized cluster mutation  $\mu_k^T$ , that is,*

$$(4.17) \quad \mu_k^T(\mathbf{u}_{t,0}) = \mathbf{u}_{t,0}.$$

(2) The translations and the tropicalized cluster mutation are commutative, that is,

$$\tau_{t'} \circ \mu_k^T = \mu_k^T \circ \tau_t \text{ on } \mathbb{R}^J.$$

(3) The  $k$ -th component of  $\mathbf{u}_{t,0}$  is equal to zero, that is,  $u_{t,0,k} = 0$  where  $\mathbf{u}_{t,0} = (u_{t,0,j})_{j \in J}$ .

*Proof.* The statement (1)  $\Rightarrow$  (2) follows from

$$(\mu_k^T \circ \tau_t)(\mathbf{u}) = \mu_k^T(\mathbf{u} - \mathbf{u}_{t,0}) = \mu_k^T(\mathbf{u}) - \mathbf{u}_{t',0} = (\tau_{t'} \circ \mu_k^T)(\mathbf{u}).$$

For (2)  $\Rightarrow$  (3), we observe

$$(4.18) \quad -\mathbf{u}_{t',0} = \tau_{t'}(\mathbf{0}) = (\tau_{t'} \circ \mu_k^T)(\mathbf{0}) = (\mu_k^T \circ \tau_t)(\mathbf{0}) = \mu_k^T(-\mathbf{u}_{t,0}) = -\mu_k^T(\mathbf{u}_{t,0}).$$

Restricting to the  $k$ -th component of (4.18), we obtain

$$u_{t',0,k} = \mu_k^T(\mathbf{u}_{t,0})_k = -u_{t,0,k},$$

It in turn implies that  $u_{t,0,k} = 0$  by the (piecewise)-linearity of  $\mu_k^T$ . Finally, suppose that (3) holds. Then (4.17) is obtained by the expression (4.13) for  $\mu_k^T$ .  $\square$

**Corollary 4.8.** *Let  $\mathbf{u}_0 \in M_{\mathbb{R}}$  and  $\mathbf{u}_{t,0}$  be the image of  $\mathbf{u}_0$  under (4.16). Let  $M_{\text{fr},\mathbb{R}}$  be the  $\mathbb{R}$ -vector space generated by  $\{\mathbf{e}_j^* \mid j \in J \setminus J_{\text{uf}}\}$ . The following are equivalent.*

- (1) For each index  $k \in J_{\text{uf}}$ , the point  $\mathbf{u}_{t,0}$  is fixed under the tropicalized cluster mutation  $\mu_k^T$ , that is,  $\mu_k^T(\mathbf{u}_{t,0}) = \mathbf{u}_{t',0}$  for  $\mu_k(t) = t'$ .
- (2) For each index  $k \in J_{\text{uf}}$ , the translations and the tropicalized cluster mutations are commutative, that is,  $\tau_{t'} \circ \mu_k^T = \mu_k^T \circ \tau_t$  on  $\mathbb{R}^J$  for  $\mu_k(t) = t'$ .
- (3) The point  $\mathbf{u}_{t,0}$  is contained in the space  $M_{\text{fr},\mathbb{R}}$ .

Let  $\mathbf{u}_{t_0,0}$  be the center of the  $\mathbb{Q}$ -Gorenstein Fano Newton–Okounkov polytope  $\Delta_{t_0}$  at the initial seed. We define a point  $\mathbf{u}_{t,0}$  of  $\Delta_t$  by the image of  $\mathbf{u}_{t_0,0}$  under the composition of the identifications

$$(4.19) \quad \mathbb{R}^J \simeq_{s_{t_0}} M_{\mathbb{R}} \simeq_{s_t} \mathbb{R}^J \quad (\mathbf{u}_{t_0,0} \mapsto \mathbf{u}_0 \mapsto \mathbf{u}_{t,0}).$$

**Lemma 4.9.** *For  $t \in \mathbb{T}$ , let  $\Delta_t$  be a  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{u}_{t,0}$ . Suppose that for a pair  $(t, t')$  of vertices with  $\mu_k(t) = t'$  in  $\mathbb{T}$  and an index  $k \in J_{\text{uf}}$ , (4.17) holds. Then  $\mu_k^T(\Delta_t) = \Delta_{t'}$  is also a  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{u}_{t',0}$ .*

*Proof.* Consider the translated polytope  $\Delta_t - \mathbf{u}_{t,0} = \tau_t(\Delta_t)$ , which is a  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{0}$ . By Lemma 4.6 and 4.7,  $(\tau_{t'} \circ \mu_k)(\Delta_t) = (\mu_k \circ \tau_t)(\Delta_t)$  is also  $\mathbb{Q}$ -Gorenstein Fano with the center  $\mathbf{0}$ . Therefore,  $\Delta_{t'}$  is a  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{u}_{t',0}$ .  $\square$

In summary, by combining Lemma 4.5 and 4.9, we deduce the following proposition.

**Proposition 4.10.** *Suppose that the family  $\{S_t \mid t \in \mathbb{T}\}$  of finitely generated semigroups has a tropical cluster structure and hence the family  $\{\Delta_t \mid t \in \mathbb{T}\}$  of corresponding Newton–Okounkov bodies of  $X$  also has a tropical cluster structure. If*

- (1) the semigroup  $S_{t_0}$  is saturated,
- (2) the Newton–Okounkov polytope  $\Delta_{t_0}$  is  $\mathbb{Q}$ -Gorenstein Fano with the center  $\mathbf{u}_0$ , and
- (3) the center  $\mathbf{u}_0$  is fixed under the tropicalized cluster mutation of each direction in the sense of (4.17),

*then every polytope  $\Delta_t$  in the family is  $\mathbb{Q}$ -Gorenstein Fano with the center  $\mathbf{u}_{t,0}$  and the toric degeneration associated with  $\Delta_t$  is normal.*

### 4.3. Infinitely many tori from tropically related Newton–Okounkov polytopes

Let  $X$  be a smooth Fano variety of complex dimension  $m$  equipped with the Kähler form of a very ample line bundle  $\mathcal{L}$ , a positive power of the anticanonical bundle of  $X$ . Suppose that we have a family  $\{S_t \mid t \in \mathbb{T}\}$  of finitely generated semigroups, each of which gives rise to a  $\mathbb{Q}$ -Gorenstein Fano Newton–Okounkov polytope  $\Delta_t$  and a normal toric degeneration of a smooth projective variety  $X$ . If the family has a tropical cluster structure, then have the family of monotone Lagrangian tori of  $X$  by Proposition 3.13. The following theorem provides a criterion for the existence of infinitely many *distinct* monotone Lagrangian tori.

**Theorem 4.11** (Theorem C). *Let  $X$  be a smooth Fano variety equipped with a Kähler form inherited from a very ample line bundle  $\mathcal{L}$  given by a positive power of the anticanonical bundle of  $X$ . Suppose that  $X$  admits the family of normal toric degenerations constructed from a family  $\{\Delta_t \mid t \in \mathbb{T}\}$  of  $\mathbb{Q}$ -Gorenstein Fano Newton–Okounkov polytopes of  $\mathcal{L}$  having a tropical cluster structure. Let  $L_t$  be a monotone Lagrangian torus constructed from  $\Delta_t$  by Proposition 3.13. Assume that the Newton–Okounkov polytope  $\Delta_{t_0}$  at the initial seed contains the origin  $\mathbf{0}$  and the center  $\mathbf{u}_0$  of  $\Delta_{t_0}$  is fixed under the tropicalized cluster mutation of each direction. If there exists a sequence  $(t_\ell)_{\ell \in \mathbb{N}}$  of seeds and a sequence  $(r_\ell, s_\ell)_{\ell \in \mathbb{N}}$  of indices in  $J_{\text{uf}} \times J$  such that*

- (1) *the sequence  $(\varepsilon_{r_\ell, s_\ell})_{\ell \in \mathbb{N}}$  consisting of the  $(r_\ell, s_\ell)$ -entry in the exchange matrix  $\varepsilon_{t_\ell}$  associated with  $t_\ell$  diverges to  $-\infty$  as  $\ell \rightarrow \infty$ ,*
- (2) *the Newton–Okounkov polytope  $\Delta_{t_\ell}$  is contained in the half-space  $H_{\mathbf{e}_{s_\ell}, 0}^+$  where  $\mathbf{s}_{t_\ell} = (\mathbf{e}_j \mid j \in J)$ , and*
- (3) *the image of the polytope  $\Delta_{t_\ell}$  under the tropicalized cluster mutation  $\mu_{r_\ell}^T$  in the  $r_\ell$ -direction is contained in the half-space  $H_{\mathbf{e}'_{s_\ell}, 0}^+$  where  $\mathbf{s}_{\mu_{r_\ell}^T(t_\ell)} = (\mathbf{e}'_j \mid j \in J)$ ,*

*then the family  $\{L_\ell \mid \ell \in \mathbb{N}\}$  contains infinitely many monotone Lagrangian tori, no two of which are related by any symplectomorphisms.*

Here are some remarks on the conditions of Theorem 4.11.

**Remark 4.12.** We emphasize that the corresponding hyperplane  $H_{\mathbf{e}_s, 0}$  (resp.  $H_{\mathbf{e}'_s, 0}$ ) need *not* contain a facet of  $\Delta_{t_\ell}$  (resp.  $\mu_{r_\ell}^T(\Delta_{t_\ell})$ ) in the condition (2) (resp. (3)). It would be harder to apply Theorem 4.11 if one has to verify that the hyperplane contains a facet. Indeed, the conditions (1), (2), and (3) can be checked by information from the lattice points contained in  $\Delta$  *without* figuring out facets of  $\Delta_{t_\ell}$  and  $\mu_{r_\ell}^T(\Delta_{t_\ell})$ .

Note that if a polytope  $\Delta'$  is obtained from another polytope  $\Delta$  by applying the tropicalized cluster mutation  $\mu_r^T$  in the  $r^{\text{th}}$ -direction and  $\Delta$  contains the origin  $\mathbf{0} \in M$ , then  $\Delta'$  also contains the origin  $\mathbf{0} \in M$ . Hence, if the polytope  $\Delta_{t_0}$  at the initial seed contains the origin  $\mathbf{0}$ , then so do the others.

**Remark 4.13.** Suppose that the positive numbers  $d_j$ 's in the fixed data  $\Gamma$  are *all one* and hence every exchange matrix is skew-symmetric. In this case, the exchange matrix can be represented as a quiver. The condition (1) says that there is a sequence of quivers in the mutation class of the initial quiver  $q_{t_0}$  such that the multiplicity of arrows from  $r$  to  $s$  goes to infinity. From a polyhedron-geometric perspective, the condition (1) means that the Newton–Okounkov polytopes get thinner and thinner through the sequence of mutations in a specific direction (as the volume is equal to the degree of the projective embedding of  $(X, \mathcal{L})$  and hence invariant under the mutations).

For the existence of such half-spaces in the (2) and (3), we consider the scattering diagram for the cluster variety  $\mathcal{A}_{\Gamma, s_0}$  in Gross–Siebert's program in [GS11]. The cluster variety  $\mathcal{A}_{\Gamma, s_0}$



is contained in the log Calabi–Yau manifold  $X \setminus D$  where  $D$  is a normal crossing anticanonical divisor. To each irreducible component of  $D$ , there is the corresponding initial ray getting into a scattering diagram. Although the initial ray can be scattered into multiple broken lines, there should be one single ray parallel to the initial ray. It corresponds to a Laurent monomial in the superpotential  $W$  on  $X^\vee$ , which corresponds to a hyperplane. It means that no matter what you take a chamber in the scattering diagram (resp. a Newton–Okounkov polytope in the family) there exists the specific initial ray (resp. a facet corresponding to the specific component of  $D$ .)

A key part of the proof of Theorem 4.11 is to extract pieces of data from the iterative process (4.13) that enable us to distinguish Lagrangian tori *without* probing *individual* toric degeneration and Newton–Okounkov polytope. We begin by collecting some lemmas.

**Lemma 4.14** (Theorem 6.4 in [Brø83]). *Suppose that a polytope  $\Delta$  contains the origin  $\mathbf{0}$  in its interior. Let  $\Delta^\circ$  be the polar of  $\Delta$ . Then the following are equivalent.*

- (1) *The hyperplane  $H_{\mathbf{v},1}$  is a supporting hyperplane of the polytope  $\Delta$ .*
- (2) *The vector  $\mathbf{v}$  is contained in the boundary of  $\Delta^\circ$ , that is,  $\mathbf{v} \in \Delta^\circ \setminus \text{Int}(\Delta^\circ)$ .*

**Lemma 4.15.** *Suppose that a polytope  $\Delta'$  is the image of a polytope  $\Delta$  under the tropicalized cluster mutation  $\mu_r^T$  in (4.13), that is,  $\Delta' = \mu_r^T(\Delta)$ . Assume that the polytope  $\Delta$  contains the origin  $\mathbf{0}$ . If  $\Delta$  is contained in the half-space  $H_{\mathbf{e}_s,0}^+$ ,  $\Delta'$  is contained in the half-space  $H_{\mathbf{e}'_s,0}^+$ , and  $\varepsilon_{r,s} \leq 0$ , then the half-space  $H_{\mathbf{e}_s - \varepsilon_{r,s}\mathbf{e}_r,0}^+$  is a supporting half-space of  $\Delta$ .*

*Proof.* To show that  $H_{\mathbf{e}_s - \varepsilon_{r,s}\mathbf{e}_r,0}^+$  is a supporting half-space of  $\Delta$ , consider the mutated polytope  $\Delta'$ . Since  $\Delta'$  is contained in  $H_{\mathbf{e}'_s,0}^+$ , we have

$$(\mu_r^T(\mathbf{u}))_s \geq 0.$$

By the relation (4.13), each point  $\mathbf{u}$  in the Newton–Okounkov body  $\Delta$  satisfies

$$(\mu_r^T(\mathbf{u}))_s = \begin{cases} u_s & \text{if } u_r \geq 0 \\ u_s - \varepsilon_{r,s}u_r & \text{if } u_r \leq 0. \end{cases}$$

In both cases, we claim that  $u_s - \varepsilon_{r,s}u_r \geq 0$ .

- (1) If  $u_r \leq 0$ , then  $u_s - \varepsilon_{r,s}u_r = (\mu_r^T(\mathbf{u}))_s \geq 0$ .
- (2) If  $u_r \geq 0$ , then  $u_s - \varepsilon_{r,s}u_r \geq u_s = (\mu_r^T(\mathbf{u}))_s \geq 0$  because  $\varepsilon_{r,s} \leq 0$ .

Moreover, the origin  $\mathbf{0}$  is contained in the intersection  $H_{\mathbf{e}_s - \varepsilon_{r,s}\mathbf{e}_r,0}^+ \cap \Delta$  and hence  $H_{\mathbf{e}_s - \varepsilon_{r,s}\mathbf{e}_r,0}^+$  is a supporting half-space of  $\Delta$ .  $\square$

We are ready to prove Theorem 4.11. Before presenting its proof, we address an issue that one should be cautious about. Suppose that  $\Delta$  is a normalized  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{u}_0$ . Let  $H_{\mathbf{v},a}^+$  be a supporting half-space of the polytope  $\Delta$  where  $\mathbf{v}$  is a *primitive* lattice vector. Let  $\tau: \mathbb{R}^J \rightarrow \mathbb{R}^J$  be the translation  $\mathbf{u} \mapsto \mathbf{u} - \mathbf{u}_0$ . If the hyperplane  $H_{\mathbf{v},a}$  contains a *facet*, then  $\tau(\Delta) = \Delta - \mathbf{u}_0$  also has a supporting half-space  $H_{\mathbf{v},1}^+$  by Remark 3.10. However, if  $H_{\mathbf{v},a}$  does *not* contain a facet, depending on  $\mathbf{u}_0$ ,  $H_{\mathbf{v},1}^+$  may *not* be a supporting half-space because

$$\tau(H_{\mathbf{v},a}^+) = H_{\frac{\mathbf{v}}{\langle \mathbf{u}_0, \mathbf{v} \rangle + a}, 1}^+.$$

More generally, suppose that  $\Delta$  is a (not necessarily normalized)  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{u}_0$ . There exists  $c \in \mathbb{Q}$  such that  $c \cdot (\Delta - \mathbf{u}_0)$  is normalized. Set  $\lambda = c^{-1}$ . If  $\Delta$

has a supporting half-space  $H_{\mathbf{v},a}^+$ , then the corresponding half-space supporting  $c \cdot (\Delta - \mathbf{u}_0)$  is

$$(4.20) \quad H_{\frac{\lambda \mathbf{v}}{\langle \mathbf{u}_0, \mathbf{v} \rangle + a}, 1}^+.$$

To obtain an easy-to-use criterion, we do *not* require the half-spaces to contain a facet of  $\Delta$ . Extra care is necessary for this reason. With this issue in mind, we prove Theorem 4.11.

*Proof of Theorem 4.11.* By taking a subsequence if necessary, without loss of generality, we may assume that  $s_\ell$  are all same in  $((r_\ell, s_\ell) \in J_{\text{uf}} \times J \setminus J_{\text{uf}})_{\ell \in \mathbb{N}}$ . Let  $s = s_\ell$  for all  $\ell \in \mathbb{N}$ . For notational simplicity, let  $\Delta_\ell := \Delta_{t_\ell}$  and  $\mathbf{u}_{\ell,0} := \mathbf{u}_{t_\ell,0}$ . By the proof of Lemma 4.6, for each  $t \in \mathbb{T}$ ,  $\lambda \cdot (\Delta_t - \mathbf{u}_{t,0})^\circ$  is also Fano if  $\lambda \cdot (\Delta_{t_0} - \mathbf{u}_0)^\circ$  is Fano.

Let  $\Delta_\ell^\vee = \lambda \cdot (\Delta_\ell - \mathbf{u}_{\ell,0})^\circ$  be the combinatorial dual polytope of  $\Delta_\ell$ . Since  $\mathbf{u}_{\ell,0}$  is the same element in  $M_{\mathbb{R}}$  by the choice of  $\mathbf{u}_{t,0}$  in (4.19), we have  $\langle \mathbf{u}_0, \mathbf{v} \rangle = \langle \mathbf{u}_{t,0}, \mathbf{v} \rangle$  for all  $t \in \mathbb{T}$ . According to Corollary 4.8,  $\mathbf{u}_{t,0}$  can be expressed as

$$\mathbf{u}_{t,0} = \sum_{j \in J \setminus J_{\text{uf}}} a_j \mathbf{e}_j^*.$$

For  $\mathbf{v} = \mathbf{e}_s$  or  $\mathbf{e}_s - \varepsilon_{r_\ell, s} \mathbf{e}_r$ , we have

$$\langle \mathbf{u}_{\ell,0}, \mathbf{v} \rangle = \langle \mathbf{u}_0, \mathbf{v} \rangle = a_s \in \mathbb{Q} \quad \text{for all } \ell.$$

By Lemma 4.15 together with (4.20), we have the following supporting half-spaces of  $\Delta_\ell$

$$(4.21) \quad H_{\frac{\lambda}{a_s} \mathbf{v}, 1}^+ \quad \text{for } \mathbf{v} = \mathbf{e}_s \text{ or } \mathbf{e}_s - \varepsilon_{r_\ell, s} \mathbf{e}_r.$$

Thus, the polytope  $\Delta_\ell^\vee$  contains the line segment joining the points  $\frac{\lambda}{a_s} \mathbf{e}_s$  and  $\frac{\lambda}{a_s} (\mathbf{e}_s - \varepsilon_{r_\ell, s} \mathbf{e}_r)$ . Let  $\frac{\lambda}{a_s} = \frac{p}{q}$  for  $p \in \mathbb{Z}, q \in \mathbb{N}$  with  $\gcd(p, q) = 1$ . Consider the lattice  $\frac{1}{q}M \supset M$ . Since the condition  $-\varepsilon_{r_\ell, s} \rightarrow \infty$  as  $\ell \rightarrow \infty$ , the number of lattice points of  $\Delta_\ell^\vee$  in  $\frac{1}{q}M$  diverges to infinity because of (4.21).

Let  $L_\ell := L_{t_\ell}$  be a monotone Lagrangian torus constructed from  $\Delta_\ell$  in Proposition 3.13. We claim that the family  $\{L_\ell \mid \ell \in \mathbb{N}\}$  contains *distinct* infinitely many monotone Lagrangian tori. Because of Proposition 3.25, we have

$$\Delta_\ell^\vee \subset \Delta_{L_\ell}^{\text{ref}}$$

and hence the number of lattice points of  $\Delta_{L_\ell}^{\text{ref}}$  in  $\frac{1}{q}M$  also diverges to infinity as  $\ell \rightarrow \infty$ . It means that the family contains infinitely many polytopes, no two of which are related by any unimodular equivalences because the number of lattice points in  $\frac{1}{q}M$  is invariant under unimodular equivalences. Hence Corollary 3.24 concludes that there are infinitely many distinct monotone Lagrangian tori in  $\{L_\ell \mid \ell \in \mathbb{N}\}$  as desired.  $\square$

## 5. Cluster polytopes for flag manifolds

In the remaining sections, we shall apply Theorem 4.11 to a flag manifold to show that it carries infinitely many Lagrangian tori, no two of which are related by any symplectomorphisms.

We set up some notations which will be used in the remaining sections. Let  $G$  be a simply connected and semisimple algebraic group over  $\mathbb{C}$  and  $\mathfrak{g}$  the corresponding Lie algebra. Let  $A = (a_{i,j})_{i,j \in I}$  be the Cartan matrix of  $\mathfrak{g}$ . Let  $B$  be a Borel subgroup of  $G$ ,  $H$  a maximal torus in  $B$ ,  $B^-$  the opposite Borel subgroup, and  $U^-$  the unipotent radical of  $B^-$ . We then have the complex flag manifold  $G/B$ . We denote by  $W = \langle s_i \mid i \in I \rangle$  the Weyl group

associated with  $\mathfrak{g}$ , where  $s_i$  is the  $i^{\text{th}}$  simple reflection. The discussion of this section will be centered around the following objects. For each element  $w \in W$ ,

- the *Schubert variety*  $X_w$  associated to  $w$  by the Zariski closure of  $BwB/B$  in  $G/B$
- the *unipotent cell*  $U_w^-$  associated to  $w$  by  $U^- \cap BwB$  in  $G$  where we identify the Weyl group  $W$  with the normalizer of the maximal torus  $H$  in the group  $G$ .

The goal of this section is to present the main result of this paper, which claims that there are infinitely many distinct monotone Lagrangian tori in every flag manifold  $G/B$  of arbitrary type except a few low-dimensional cases. To construct such a family of Lagrangian tori, we employ a family of Newton–Okounkov polytopes of  $G/B$  constructed by cluster algebra. We discuss the cluster structure on the function field of  $G/B$  and the relationship between the dual canonical basis of the quantum group and the Newton–Okounkov polytopes. The connection leads to the proof of some conditions to apply the distinguishing criterion (Theorem 4.11) to the family.

### 5.1. Infinitely many Lagrangian tori in flag manifolds

In the early stages, the construction problem for a toric degeneration of an algebraic variety was tackled by the theory of Gröbner basis or SAGBI basis. By constructing such a basis, a toric degeneration of a flag variety  $SL_n/B$  of type  $A$  (and Schubert varieties therein) was constructed by Gonciulea–Lakshmibai [GL96] and Kogan–Miller [KM05] for instance. Indeed, the toric variety at the central fiber in this toric degeneration corresponds to the toric variety associated with the Gelfand–Zeitlin polytope.

In [Cal02], Caldero brought a new approach to the degeneration problem on Schubert varieties based on the string parametrizations of the dual canonical basis or the upper global basis. The string parametrizations of the dual canonical basis in the irreducible representation of  $G$  with the highest weight  $\lambda$  were introduced by Littelmann [Lit98] and Berenstein–Zelevinsky [BZ01]. A *string polytope* is defined by the convex hull of the string parametrizations in the irreducible representation of  $G$ . Caldero constructed a family of toric degenerations of a flag variety (and its Schubert varieties) of arbitrary type corresponding to a string polytope. Indeed, string polytopes are important examples of Newton–Okounkov bodies developed by [Oko96, LM09, KK12]. Later on, Kaveh [Kav15] proved how string polytopes can be viewed as a Newton–Okounkov polytope by finding a suitable valuation on the functions field of the flag manifold, see Fujita [Fuj18] for a generalization to Schubert varieties.

To construct a string polytope, we need to choose two data. One is a dominant integral weight  $\lambda$ , a non-negative linear combination of fundamental weights, and the other is a reduced expression  $\underline{w}$  of a Weyl group element  $w \in W$ . From the geometric perspective, the choice of  $\lambda$  determines an ambient partial flag variety and its adorned Kähler form  $\omega_\lambda$ . Next, the choice of a Weyl group element  $w$  determines a Schubert variety  $X_w$ . In particular,  $X_{w_0}$  is the ambient flag manifold if  $w_0$  is the longest element of  $W$ . Finally, the choice of a reduced expression  $\underline{w}$  of the element  $w$  gives rise to a valuation on  $\mathbb{C}(X)$  arising from the sequence of (resolutions of) Schubert subvarieties of  $X_w$  given by the truncations of  $\underline{w}$ . In sum, a string polytope associated with the choice  $(\lambda, \underline{w})$  can be used to understand a toric degeneration of  $X_w$ . One important point is that we obtain possibly different Newton–Okounkov bodies if we make a different choice of a reduced expression  $\underline{w}$  leaving the other choices  $\lambda$  and  $w$  fixed. In representation terminology, the reduced expression selects the order of types of the Kashiwara operators on the crystal graph so that they give rise to different string parametrizations (and they produce different convex hulls.)

Yet, there are *only* finitely many reduced expressions of the longest element of  $W$ . Thus, we can have only finitely many distinct monotone Lagrangian tori at best. We need to find other sources for toric degenerations to produce infinitely distinct objects.

Gross–Hacking–Keel–Kontsevich [GHKK18] laid down a general framework for constructing toric degenerations via cluster algebra. This framework can be very useful to construct meaningful Lagrangian tori of a smooth projective variety thanks to the work of Harada–Kaveh [HK15]. In [FO20], Fujita–Oya showed that toric degenerations of a Schubert variety  $X_w$  in a flag manifold arising from the cluster structure on the unipotent cell  $U_w$  in Berenstein–Fomin–Zelevinsky [BFZ05] are typical examples for the GHKK construction. In particular, they demonstrated how the toric degenerations can be realized as a Newton–Okounkov body on  $X_w$ , relying on Anderson’s construction [And13].

An upper cluster algebra structure on the unipotent coordinate ring  $\mathbb{C}[U_w^-]$  was discovered by Berenstein–Fomin–Zelevinsky [BFZ05]. It yields that  $U_w^-$  is birational to the  $\mathcal{A}$ -cluster variety. By using the  $\mathcal{A}$ -cluster structure, for each seed  $\mathfrak{s}$ , Fujita–Oya [FO20] constructed a valuation  $v_{\mathfrak{s}}$  on the function field of  $\mathcal{A}$ . Since it is birational to the Schubert variety  $X_w$ , the valuation on  $\mathbb{C}(\mathcal{A})$  defines the valuation on  $\mathbb{C}(X_w)$ . Therefore each seed gives rise to a Newton–Okounkov body of the Schubert variety  $X_w$ .

Moreover, the constructed Newton–Okounkov bodies can be described via the Fock–Goncharov dual  $\mathcal{A}^\vee$ . By the recent work of Qin [Qin20] it was shown that the dual canonical basis of  $\mathbb{C}[U_w^-]$  is *pointed* and the *extended  $g$ -vector* agrees with the valuation  $v_{\mathfrak{s}}$  of a seed  $\mathfrak{s}$ . Through the relation, we can extract data of lattice points of Newton–Okounkov bodies via properties of extended  $g$ -vectors. In fact, for a special choice of seeds, the corresponding Newton–Okounkov polytope is unimodularly equivalent to a string polytope, and hence the family of polytopes can be thought of as a generalization of string polytopes.

With this background in mind, we choose an anticanonical regular dominant weight  $\lambda = 2\rho$  where  $\rho$  is the sum of fundamental weights and the longest element  $w = w_0$  of the Weyl group  $W$ . Hence the Schubert variety  $X_w$  becomes a flag manifold equipped with a monotone Kähler form  $\omega_{2\rho}$ . The toric degeneration arising from each Newton–Okounkov body can be shown to be  $\mathbb{Q}$ -Gorenstein Fano and normal so that for each seed  $\mathfrak{s}$ , we obtain a monotone Lagrangian torus  $L_{\mathfrak{s}}$  by Proposition 3.13 and 4.10. By applying Theorem 4.11, we shall verify that the family contains infinitely many distinct monotone Lagrangian tori, as stated below.

**Theorem 5.1.** *Suppose that  $G$  is a simply connected and simple complex Lie group other than  $A_1, A_2, A_3, A_4$ , and  $B_2 = C_2$  type, that is,*

$$G \neq \mathrm{SL}_2(\mathbb{C}), \mathrm{SL}_3(\mathbb{C}), \mathrm{SL}_4(\mathbb{C}), \mathrm{SL}_5(\mathbb{C}), \mathrm{Spin}_5(\mathbb{C}) = \mathrm{Sp}_4(\mathbb{C}).$$

*Let  $\rho$  be the sum of fundamental weights and consider the flag manifold  $X := G/B$  equipped with a monotone Kähler form  $\omega_{2\rho}$ . Let  $\{L_{\mathfrak{s}}\}$  be the family of monotone Lagrangian tori of the flag manifold  $X$  constructed by the toric degenerations from the upper cluster structure on the coordinate ring of the unipotent cell  $U_{w_0}^-$ . Then the family  $\{L_{\mathfrak{s}}\}$  contains infinitely many monotone Lagrangian tori, no two of which are related by any symplectomorphisms.*

**Remark 5.2.** If  $G = A_1$ , then the flag manifold  $G/B$  is the projective space  $\mathbb{C}P^1 \simeq S^2$ . In this case, every simply closed curve dividing the sphere into two pieces having the same area is Hamiltonian isotopic to a great circle. In other words, the sphere has a unique monotone Lagrangian circle. It would be interesting to see whether  $G/B$  carries infinitely many distinct monotone Lagrangian tori or not when  $G$  is of type  $A_2, A_3, A_4$ , and  $B_2 = C_2$ .

The proof of Theorem 5.1 will be given in Section 6 and here is the outline of the proof of Theorem 5.1.

- (1) We briefly recall a construction of Newton–Okounkov polytopes of a flag manifold  $G/B$  via the upper cluster structure on the coordinate ring of the unipotent cell  $U_{w_0}^-$ . A constructed Newton–Okounkov polytope is called a *cluster polytope*. We discuss the tropical cluster structure on the family of cluster polytopes of flag manifolds, see Section 5.2.
- (2) As an initial step, we first prove that the cluster polytope  $\Delta_{t_0}$  at the initial seed is  $\mathbb{Q}$ -Gorenstein Fano, see Proposition 5.11. Inductively, Lemma 4.6 shows that every cluster polytope is also  $\mathbb{Q}$ -Gorenstein Fano. As a consequence of Proposition 4.10, we have a family of infinitely many monotone Lagrangian tori in  $G/B$ , see Corollary 5.12.
- (3) By exploiting the correspondence between the tropical integer points of a cluster polytope and the dual canonical (or upper global) basis for the negative half  $U_q^-(\mathfrak{g})$  of the quantum group  $U_q(\mathfrak{g})$ , we shall check the conditions (2) and (3) for the criterion in Theorem 4.11, see Section 5.4.
- (4) Section 6 confirms the remaining condition (1) for Theorem 4.11. We shall find a sequence of exchange matrices in the same mutation class with an arbitrarily large entry between an unfrozen variable and frozen variables. Consequently, Theorem 4.11 shows that the constructed family has infinitely many distinct monotone Lagrangian tori in  $G/B$ .

## 5.2. Cluster polytopes and the dual canonical basis

In this subsection, we review the cluster algebra structure on the coordinate ring  $\mathbb{C}[U_w^-]$  of the unipotent cell  $U_w^-$  in [BFZ05, Wil13, GLS11] to construct a family of Newton–Okounkov bodies of the Schubert variety  $X_w$ . We also recall some results on these Newton–Okounkov bodies in Fujita–Oya [FO20], which will be key ingredients for the proof of Theorem 5.1.

Let  $G$  be a simply connected and semisimple algebraic group over  $\mathbb{C}$ . For a Weyl group element  $w = s_{i_1} s_{i_2} \cdots s_{i_m} \in W$ , we set  $\text{supp}(w) := \{i_1, i_2, \dots, i_m\} \subset I$  where  $I$  is the set of indices for the simple roots. Note that  $\text{supp}(w)$  does *not* depend on the choice of a reduced expression of  $w$ . Assume that  $\text{supp}(w) = I$  for simplicity. For a dominant integral weight  $\lambda$  and  $u, v \in W$ , we denote by  $\Delta_{u\lambda, v\lambda}$  the *generalized minor* associated with  $u, v$  and  $\lambda$  and set

$$D_{u\lambda, v\lambda} := \Delta_{u\lambda, v\lambda}|_{U_w^-},$$

which is called a *unipotent minor*.

For a reduced expression  $\underline{w} = s_{i_1} s_{i_2} \cdots s_{i_m}$  of  $w$  and  $1 \leq k \leq m$ , we set

$$\begin{aligned} w_{\leq k} &:= s_{i_1} s_{i_2} \cdots s_{i_k}, \\ k^+ &:= \min(\{m+1\} \cup \{k+1 \leq j \leq m \mid i_j = i_k\}), \\ k^- &:= \max(\{0\} \cup \{1 \leq j \leq k-1 \mid i_j = i_k\}). \end{aligned}$$

Let  $J := \{1, 2, \dots, m\}$ ,  $J_{\text{fr}} := \{j \in J \mid j^+ = m+1\}$  and  $J_{\text{uf}} := J \setminus J_{\text{fr}}$ . We set

$$D_j := D_{w_{\leq j} \varpi_{i_j}, \varpi_{i_j}} \quad \text{for } 1 \leq j \leq m$$

and define the extended exchange matrix  $\varepsilon_0 = (\varepsilon_{r,s})_{r \in J_{\text{uf}}, s \in J}$  by

$$(5.1) \quad \varepsilon_{r,s} = \begin{cases} -1 & \text{if } r = s^+, \\ -a_{i_s, i_r} & \text{if } s < r < s^+ < r^+, \\ 1 & \text{if } r^+ = s, \\ a_{i_s, i_r} & \text{if } r < s < r^+ < s^+, \\ 0 & \text{otherwise} \end{cases}$$

where  $\mathbf{A} = (a_{i,j})_{i,j \in J}$  is the Cartan matrix of  $G$ .

It turns out that the coordinate ring  $\mathbb{C}[U_w^-]$  has a cluster algebra structure. The set  $D_w := \{D_j \mid j = 1, 2, \dots, m\}$  together with the extended exchange matrix  $\varepsilon_0$  forms a (Fomin–Zelevinsky’s) seed. Then  $\mathbb{C}[U_w^-]$  is isomorphic to the upper cluster algebra generated by the initial seed  $\mathbf{s}_0 := (D_w, \varepsilon_0)$ , see Definition 4.2. Let  $\mathbb{T}$  be the exchange graph associated with the cluster algebra  $\mathbb{C}[U_w^-]$ . Let  $(A_{j,t})_{j \in J}$  be the cluster variables associated with  $t \in \mathbb{T}$ , defined in (4.12). Note that  $A_{j,t_0} = D_j$ .

By utilizing the above cluster algebra structure, for a fixed  $t \in \mathbb{T}$ , we define a valuation  $v_t$  on the function field of  $U_w^-$  that is isomorphic to the function field of  $X_w$ . Let  $\mathbf{s}_t = ((A_{j,t})_{j \in J}, \varepsilon)$  be the seed of  $\mathbb{C}[U_w^-]$  associated with  $t$ . For  $\mathbf{a}, \mathbf{b} \in \mathbb{Z}^J$ , we write

$$\mathbf{a} \preceq_\varepsilon \mathbf{b} \iff \mathbf{a} = \mathbf{b} + \mathbf{v}\varepsilon \text{ for some } \mathbf{v} \in \mathbb{Z}_{\geq 0}^{J_{\text{uf}}}.$$

The order  $\preceq_\varepsilon$  on  $\mathbb{Z}^J$  is called the *dominance order* with respect to  $\varepsilon$  in [Qin17]. We consider the Laurent polynomial ring  $\mathcal{F} := \mathbb{C}[A_{j,t}^{\pm 1} \mid j \in J]$ . By identifying a Laurent monomial  $\prod_{j \in J} A_{j,t}^{a_j}$  with  $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{Z}^J$ , we obtain the induced order  $\preceq_\varepsilon$  on the set of Laurent monomials in  $\mathcal{F}$ . We denote by  $v_{\mathbf{s}_t}$  the highest term valuation on  $\mathcal{F}$  with respect to a total order  $<_t$  refined from  $\preceq_\varepsilon$ . We sometimes write  $v_t$  for  $v_{\mathbf{s}_t}$  if no confusion arises.

For a certain class of elements in  $\mathcal{F}$ , the valuation can be calculated by the extended  $g$ -vector, which we are about to recall. Following [FZ07, FG09], we set

$$X_{i,t} := \prod_{j \in J} A_{j,t}^{\varepsilon_{i,j}}.$$

An element  $f \in \mathcal{F}$  is said to be *weakly pointed* at  $(g_j)_{j \in J} \in \mathbb{Z}^J$  if  $f$  can be expressed as

$$(5.2) \quad f = \left( \prod_{j \in J} A_{j,t}^{g_j} \right) \left( \sum_{\mathbf{a}=(a_j) \in \mathbb{Z}_{\geq 0}^{J_{\text{uf}}}} c_{\mathbf{a}} \prod_{j \in J_{\text{uf}}} X_{j,t}^{a_j} \right)$$

for some nonzero  $c_{\mathbf{a}} \in \mathbb{C}$  with  $c_{\mathbf{0}} \neq 0$ . In this case, the  $g_{\mathbf{s}_t}(f) := (g_j)_{j \in J}$  is called the *extended  $g$ -vector* of  $f$ . If  $c_{\mathbf{0}} = 1$  in addition, then the element  $f$  is called *pointed*. We sometimes write  $g_t$  for  $g_{\mathbf{s}_t}$  if no confusion arises. By [FO20, Proposition 3.9], for every weakly pointed element  $f \in \mathcal{F}$ , we have

$$(5.3) \quad v_t(f) = g_t(f).$$

**Remark 5.3.** Note that the extended  $g$ -vector  $g_t$  corresponds to  $\mathbf{g}_t^{\text{I}}$  defined in [KK19] under the categorification using quiver Hecke algebras (see [FO20, Remark C.4]).

For a dominant integral weight  $\lambda \in P^+$ , we define a line bundle  $\mathcal{L}_\lambda := (G \times \mathbb{C})/B$  over the flag manifold  $G/B$ , where  $B$  acts on  $G \times \mathbb{C}$  from the right as  $(g, c) \cdot b := (gb, \lambda(b)c)$  for  $g \in G, c \in \mathbb{C}$ , and  $b \in B$ . Restricting to  $X_w$ , we obtain a line bundle on  $X_w$  which is also denoted by  $\mathcal{L}_\lambda$ . If  $\lambda$  is regular in addition, that is,  $\langle h_i, \lambda \rangle > 0$  for every  $i \in I$ , then the line bundle  $\mathcal{L}_\lambda$  is very ample. We fix a lowest weight vector  $\tau_\lambda$  in  $H^0(G/B, \mathcal{L}_\lambda)$  and restrict it to  $X_w$ . Following Section 2.2 and using the valuation  $v_t$  on the function field  $\mathcal{F} \simeq \mathbb{C}(X_w)$ , one can produce the semigroup  $S(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)$  defined in (2.4) and the Newton–Okounkov body  $\Delta(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)$  defined by (2.5). Let  $C_{\mathbf{s}_t}(w)$  be the smallest real closed cone containing  $v_t(\mathbb{C}[U^- \cap X_w] \setminus \{0\})$  in  $\mathbb{R}^J$ , which is called the *cluster cone* of  $X_w$  associated with the seed  $\mathbf{s}_t$  of  $t \in \mathbb{T}$ .

**Theorem 5.4** (Theorem 6.8, Corollary 6.9, Corollary 6.10 in [FO20]). *For each  $t$  in  $\mathbb{T}$ , the following hold.*

- (1) If a dominant integral weight  $\lambda$  is regular, then  $\mathcal{L}_\lambda$  is very ample.
- (2)  $S(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)$  is finitely generated and saturated (and hence  $\Delta(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)$  is a rational polytope and there exists a normal toric degeneration of  $X_w$  corresponding to  $\Delta(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)$  by Theorem 2.4.)
- (3)  $C_{s_t}(w) \cap \mathbb{Z}^J = v_t(\mathbb{C}[U^- \cap X_w] \setminus \{0\})$ .
- (4)  $C_{s_t}(w) = \bigcup_{\lambda \in P^+} \Delta(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)$ .

**Definition 5.5.** We simply call  $\Delta(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)$  a *cluster polytope*. If no confusion arises, we simply write  $\Delta_{s_t}(w, \lambda)$  for  $\Delta(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)$ .

To describe the relation between the Newton–Okounkov polytopes from two different choices of seeds, we consider a “nice” basis on  $\mathbb{C}[U_w^-]$ . More precisely, the unipotent coordinate ring  $\mathbb{C}[U_w^-]$  admits a  $\mathbb{C}$ -basis  $\mathcal{B}_w$  satisfying the following properties.

- (1) Every basis element in  $\mathcal{B}_w$  is weakly pointed for all  $t \in \mathbb{T}$ .
- (2) For each  $t \in \mathbb{T}$ , the map  $\mathcal{B}_w \rightarrow \mathbb{Z}^J$  given by  $b \mapsto g_t(b)$  is injective.
- (3) If  $t' = \mu_k(t)$  for some  $k \in J_{\text{uf}}$  and  $b \in \mathcal{B}_w$ , then the extended  $g$ -vector  $g_t(b) = (g_j)_{j \in J}$  at  $t$  and the extended  $g$ -vector  $g_{t'}(b) = (g'_j)_{j \in J}$  at  $t'$  are related by the tropicalized cluster mutation  $\mu_k^T$  in (4.13).
- (4) For each dominant integral weight  $\lambda$ , there exists a subset  $\mathcal{B}_w(\lambda)$  of  $\mathcal{B}_w$  such that  $\mathcal{B}_w(\lambda)$  is a  $\mathbb{C}$ -basis for the space  $\{\sigma/\tau_\lambda \mid \sigma \in H^0(X_w, \mathcal{L}_\lambda)\}$ .

Indeed, the dual canonical basis/upper global basis on the negative half  $U_q^-(\mathfrak{g})$  of the quantized enveloping algebra induces such a basis of  $\mathbb{C}[U_w^-]$  via the process of specialization and localization. The induced basis on  $\mathbb{C}[U_w^-]$  carries nice properties inherited from the dual canonical basis, the above properties (1) – (4), see Lusztig [Lus90, Lus91], Kashiwara [Kas90, Kas91, Kas93], and see also [KK19, Qin20], [FO20, Appendix C].

By (1), every element  $f$  of the dual canonical basis is weakly pointed. Because of (5.3) and (4), the valuation  $v_t$  of the basis element is equal to its extended  $g$ -vector  $g_t$ . From the relation (3) on the extended  $g$ -vectors from two different choices  $t$  and  $t'$ , it follows the relation between two sets of integral points realized by  $v_t$  and  $v_{t'}$ . It in turn yields that the semigroups and cluster polytopes are related by a sequence of tropicalized cluster mutations in (4.14) and (4.13), respectively. Here are more precise statements.

**Theorem 5.6** (Corollary 5.7 and 5.8 in [FO20]). *If  $t' = \mu_k(t)$  for some  $k \in J_{\text{uf}}$ , then*

- (1) *the associated semigroups  $S(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)$  and  $S(X_w, \mathcal{L}_\lambda, v_{t'}, \tau_\lambda)$  are related by the tropicalized cluster mutation  $\widehat{\mu}_k^T$  in the  $k^{\text{th}}$ -direction, that is,*

$$S(X_w, \mathcal{L}_\lambda, v_{t'}, \tau_\lambda) = \widehat{\mu}_k^T(S(X_w, \mathcal{L}_\lambda, v_t, \tau_\lambda)).$$

*Therefore, the family of semigroups has a tropical cluster structure.*

- (2) *the associated cluster polytopes  $\Delta_{s_t}(w, \lambda)$  and  $\Delta_{s_{t'}}(w, \lambda)$  are related by the tropicalized cluster mutation  $\mu_k^T$  in the  $k^{\text{th}}$ -direction, that is,*

$$\Delta_{s_{t'}}(w, \lambda) = \mu_k^T(\Delta_{s_t}(w, \lambda)).$$

*Therefore, the family of cluster polytopes has a tropical cluster structure.*

### 5.3. Cluster polytopes are $\mathbb{Q}$ -Gorenstein Fano.

The cluster polytopes can be thought of as a generalization of string polytopes in the following sense. Fix a dominant integral weight  $\lambda$ . Let  $\underline{w} = s_{i_1} s_{i_2} \cdots s_{i_m}$  and  $\mathfrak{s}_{t_0} := (D_{\underline{w}}, \varepsilon_0)$  be the corresponding seed. On one hand, the seed gives rise to a cluster polytope  $\Delta_{\mathfrak{s}_{t_0}}(w, \lambda)$  in Definition 5.5. On the other hand, setting  $\mathbf{i} := (i_1, i_2, \dots, i_m) \in I^m$ , and there is a *string*

polytope  $\Delta_{\mathbf{i}}(\lambda)$  associated with  $\mathbf{i}$ , see [BZ97, Lit98] for the precise definition. Then they are unimodularly equivalent.

**Theorem 5.7** (Corollary 6.6, Corollary 6.28 in [FO20]). *Let  $\lambda$  be a dominant integral weight. Then the Newton–Okounkov body  $\Delta_{\mathbf{s}_{t_0}}(w, \lambda)$  is unimodularly equivalent to the string polytope  $\Delta_{\mathbf{i}}(\lambda)$ .*

**Remark 5.8.** The string polytope of type  $A$  can be described by the combinatorics of wiring diagrams in [GP00]. One can associate a quiver to each wiring diagram and the braid 3-move is compatible with a quiver mutation. Indeed, the cluster polytopes arise from the quiver mutations of an initial quiver.

Throughout this subsection, we assume that  $w$  is the longest element  $w_0$  and hence  $X_{w_0} = G/B$ . The main goal is to prove that every cluster polytope  $\Delta_{\mathbf{s}_t}(w_0, \lambda)$  is  $\mathbb{Q}$ -Gorenstein Fano if we take the line bundle  $\mathcal{L}_\lambda$  with the anticanonical weight  $\lambda = 2\rho$  where  $\rho$  is the sum of fundamental weights. We set  $\Delta_{\mathbf{s}_t}(\lambda) := \Delta_{\mathbf{s}_t}(w_0, \lambda)$  for a seed  $\mathbf{s}_t$  ( $t \in \mathbb{T}$ ). As the base step for the induction, we prove that  $\Delta_{\mathbf{i}_{\text{std}}}(2\rho)$  is  $\mathbb{Q}$ -Gorenstein Fano and hence so is  $\Delta_{\mathbf{s}_{t_0}}(\lambda)$  because the  $\mathbb{Q}$ -Gorenstein Fano condition is preserved under the unimodular equivalence.

**Lemma 5.9.** *The string polytope  $\Delta_{\mathbf{i}_{\text{std}}}(2\rho)$  is a normalized  $\mathbb{Q}$ -Gorenstein Fano polytope.*

*Proof.* In [Lit98], Littelmann provided two sets of lattice vectors  $\{\mathbf{z}_p\}_{1 \leq p \leq m}, \{\mathbf{w}_q\}_{1 \leq q \leq \ell}$  in  $N$  such that

$$\Delta_{\mathbf{i}_{\text{std}}}(2\rho) = \left( \bigcap_{p=1}^m H_{\mathbf{z}_p, 0}^+ \right) \cap \left( \bigcap_{q=1}^{\ell} H_{\mathbf{w}_q, 2}^+ \right).$$

Here the vectors  $\mathbf{z}_p$  are the coefficients of the inequalities defining the string cone associated with  $\mathbf{i}_{\text{std}}$ , and the vectors  $\mathbf{w}_q$  are the coefficients of the additional inequalities to define the string polytope associated with the highest weight  $2\rho$ . One may assume that each vector  $\mathbf{z}_p$  is primitive, since  $H_{\mathbf{z}_p, 0}^+ = H_{a\mathbf{z}_p, 0}^+$  for any  $a \in \mathbb{R}_{>0}$ . Each vector  $\mathbf{w}_q$  contains a component  $-1$  (see the Definition [Lit98, Page 149]) so that  $\mathbf{w}_q$  is primitive.

By applying the main theorem of Steinert in [Ste22] to  $\Delta_{\mathbf{i}_{\text{std}}}(2\rho)$ , we conclude that it has a unique interior lattice point  $\mathbf{a}$  and the polar  $(\Delta_{\mathbf{i}_{\text{std}}}(2\rho) - \mathbf{a})^\circ$  is a lattice polytope. Since

$$\Delta_{\mathbf{i}_{\text{std}}}(2\rho) - \mathbf{a} = \left( \bigcap_{p=1}^m H_{\frac{\mathbf{z}_p}{\langle \mathbf{z}_p, \mathbf{a} \rangle}, 1}^+ \right) \cap \left( \bigcap_{q=1}^{\ell} H_{\frac{\mathbf{w}_q}{\langle \mathbf{w}_q, \mathbf{a} \rangle + 2}, 1}^+ \right),$$

if  $H_{\mathbf{z}_p, 0}$  (resp.  $H_{\mathbf{w}_q, 2}$ ) contains a facet of  $\Delta_{\mathbf{i}_{\text{std}}}(2\rho)$ , then  $\frac{\mathbf{z}_p}{\langle \mathbf{z}_p, \mathbf{a} \rangle}$  (resp.  $\frac{\mathbf{w}_q}{\langle \mathbf{w}_q, \mathbf{a} \rangle + 2}$ ) is a lattice vector. Since  $\mathbf{z}_p$  (respectively  $\mathbf{w}_q$ ) is primitive, we have  $\langle \mathbf{z}_p, \mathbf{a} \rangle = \pm 1$  (respectively,  $\langle \mathbf{w}_q, \mathbf{a} \rangle + 2 = \pm 1$ ) and  $\frac{\mathbf{z}_p}{\langle \mathbf{z}_p, \mathbf{a} \rangle}$  (respectively,  $\frac{\mathbf{w}_q}{\langle \mathbf{w}_q, \mathbf{a} \rangle + 2}$ ) is a primitive vector, as desired.  $\square$

The following lemma describes the center point of the cluster polytopes:

**Lemma 5.10** (Corollary 4.18, Proposition 4.20 in [FH21]). *Let*

$$(5.4) \quad \mathbf{u}_0 := \sum_{j \in J \setminus J_{\text{uf}}} \mathbf{e}_j^*.$$

*For each  $t \in \mathbb{T}$ , let  $\mathbf{u}_{t,0}$  be the point in (4.16). Then the cluster polytope  $\Delta_{\mathbf{s}_t}(2\rho)$  satisfies*

- (1) *the point  $\mathbf{u}_{t,0}$  in (5.4) is the unique interior point of  $\Delta_{\mathbf{s}_t}(2\rho)$ ,*
- (2) *the polar dual of  $\Delta_{\mathbf{s}_t}(2\rho) - \mathbf{u}_{t,0}$  is a lattice polytope, and*
- (3)  *$\mathbf{u}_{t,0}$  is fixed under the tropicalized cluster mutation of each direction.*



The following proposition further claims that the polar dual of  $\Delta_{\mathfrak{s}_t}(2\rho) - \mathbf{u}_{t,0}$  is the convex hull of primitive lattice vectors. It in turn yields that  $\Delta_{\mathfrak{s}_t}(2\rho)$  is a normalized  $\mathbb{Q}$ -Gorenstein Fano polytope.

**Proposition 5.11.** *For each  $t \in \mathbb{T}$ , the cluster polytope  $\Delta_{\mathfrak{s}_t}(2\rho)$  is a normalized  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{u}_{t,0}$  and the associated toric degeneration is normal.*

*Proof.* For each seed  $\mathfrak{s}_t$  ( $t \in \mathbb{T}$ ), let  $S_t := S(X_{w_0}, \mathcal{L}_{2\rho}, v_t, \tau_{2\rho})$  be the semigroup associated to  $v_t$ . By Theorem 5.4, each semigroup  $S_t$  is finitely generated and saturated. Hence, the toric degeneration associated with the corresponding Newton–Okounkov polytope  $\Delta_{\mathfrak{s}_t}(2\rho)$  is normal by [And13].

By Theorem 5.7,  $\Delta_{\mathfrak{s}_{t_0}}(2\rho)$  is a normalized  $\mathbb{Q}$ -Gorenstein Fano polytope with the center  $\mathbf{u}_{t_0,0}$ . By Lemma 4.6 and Corollary 4.8, so are all the other string polytopes.  $\square$

**Corollary 5.12.** *For each  $t \in \mathbb{T}$ , consider a Lagrangian torus fibration  $\Phi_t: G/B \rightarrow \Delta_{\mathfrak{s}_t}(2\rho)$  constructed from Theorem 2.2. Then the fiber  $\Phi_t^{-1}(\mathbf{u}_{t,0})$  at the center is a monotone Lagrangian torus.*

*Proof.* The line bundle  $\mathcal{L}_{2\rho}$  is very ample and anticanonical and hence the Kähler form inherited from  $\mathcal{L}_{2\rho}$  is monotone. The statement follows from Proposition 5.11 and Theorem 3.13.  $\square$

#### 5.4. Nonnegativity of the exponents of the frozen variables

The goal of this subsection is to verify the conditions (2) and (3) in Theorem 4.11 in the case of cluster polytopes. Throughout this subsection, we assume that  $w$  is the longest element  $w_0$  and hence  $X_{w_0} = G/B$ . Set  $\Delta_{\mathfrak{s}_t}(\lambda) := \Delta_{\mathfrak{s}_t}(w_0, \lambda)$  and  $C_{\mathfrak{s}_t} := C_{\mathfrak{s}_t}(w_0)$  for a seed  $\mathfrak{s}_t$  ( $t \in \mathbb{T}$ ).

Let  $B(\infty)$  be the infinite crystal of the negative half  $U_q^-(\mathfrak{g})$  of the quantum group  $U_q(\mathfrak{g})$ . We consider the *dual canonical basis* (or *upper global basis*) of  $U_q^-(\mathfrak{g})$

$$\mathbf{G}^{\text{up}}(\infty) := \{\mathbf{G}^{\text{up}}(b) \mid b \in B(\infty)\},$$

where  $\mathbf{G}^{\text{up}}(b)$  is the element corresponding to  $b$  in  $B(\infty)$ . Then the specialization  $\mathbf{G}_{q=1}^{\text{up}}(\infty) := \{\mathbf{G}_{q=1}^{\text{up}}(b) \mid b \in B(\infty)\}$  of  $\mathbf{G}^{\text{up}}(\infty)$  at  $q = 1$  forms a basis of the coordinate ring  $\mathbb{C}[U^-]$ . Since the dual canonical basis  $\mathbf{G}^{\text{up}}(\infty)$  is pointed ([Qin20]), (5.3) yields

$$v_{\mathfrak{s}_t}(b) = g_{\mathfrak{s}_t}(b) \quad \text{for any } b \in \mathbf{G}_{q=1}^{\text{up}}(\infty).$$

Thanks to Theorem 5.4, we have

$$(5.5) \quad C_{\mathfrak{s}_t} \cap \mathbb{Z}^J = \{g_{\mathfrak{s}_t}(b) \mid b \in \mathbf{G}_{q=1}^{\text{up}}(\infty)\}.$$

The unipotent coordinate ring  $\mathbb{C}[U^-]$  is a polynomial ring, and it is equipped with a cluster algebra structure on which the frozen variables are *not* invertible, see [GLS11, Theorem 3.3] and [GY17, GY20]. Indeed, for each reduced expression  $\underline{w}_0 = s_{i_1} s_{i_2} \cdots s_{i_\ell}$ , there exists a seed of  $\mathbb{C}[U^-]$  consisting of the cluster variables

$$(5.6) \quad \left\{ \Delta_{w_{\leq k} \varpi_{i_k}, \varpi_{i_k}} |_{U^-} \mid 1 \leq k \leq \ell \right\}$$

with the same set of indices of frozen variables and the same exchange matrix (5.1) given in the previous section.

On the other hand, we have

$$U_{w_0}^- = U^- \cap \mathbf{O}_{w_0},$$

where

$$\mathbf{O}_{w_0} = \{g \in U^- \mid \Delta_{w_0 \varpi_i, \varpi_i}(g) \neq 0 \text{ for } i \in I\},$$

see [BZ97], [GLS11, Propositions 8.4 and 8.5] and see also [KO21, Proposition 2.20]. Hence, the coordinate ring  $\mathbb{C}[U_{w_0}^-]$  is the localization of  $\mathbb{C}[U^-]$  at  $\{\Delta_{w_0\varpi_i, \varpi_i}|_{U^-} \mid i \in I\}$ , which is the set of frozen variables of the cluster algebra  $\mathbb{C}[U^-]$ . It follows that the cluster  $\{A_j \mid j \in J\}$  of a seed  $\mathfrak{s}_t$  of  $\mathbb{C}[U_{w_0}^-]$  is also a cluster of  $\mathbb{C}[U^-]$ . Hence any element in  $\mathbb{C}[U^-] \subset \mathbb{C}[U_{w_0}^-]$  can be written as a Laurent polynomial in the cluster variables in  $\{A_j \mid j \in J\}$ , in which the exponents of the frozen variables in any monomials are nonnegative by the Laurent phenomenon ([FZ02]). Since  $\mathbf{G}_{q=1}^{\text{up}}(\infty) \subset \mathbb{C}[U^-]$ , we deduce the following proposition.

**Proposition 5.13.** *For every  $b \in \mathbf{G}_{q=1}^{\text{up}}(\infty)$  and for each frozen index  $k \in J_{\text{fr}}$ , the  $k^{\text{th}}$ -component of the extended  $g$ -vector of  $b$  is nonnegative, that is,*

$$(5.7) \quad (g_{\mathfrak{s}_t}(b))_k \geq 0.$$

For a dominant integral weight  $\lambda$ , let  $B(\lambda)$  be the crystal of the irreducible highest weight  $U_q(\mathfrak{g})$ -module  $V(\lambda)$  and let  $\mathbf{G}^{\text{up}}(\lambda)$  be the *dual canonical basis* (or *upper global basis*) of  $V(\lambda)$ . We regard  $\mathbf{G}^{\text{up}}(\lambda)$  as a subset of  $\mathbf{G}^{\text{up}}(\infty)$  via the dual map of the natural projection  $\pi: U_q^-(\mathfrak{g}) \rightarrow V(\lambda)$  sending 1 to the highest weight vector  $v_\lambda$  of  $V(\lambda)$ . By [FO20, Theorem 6.8 (2), Corollary 6.10], we have

$$(5.8) \quad \Delta_{\mathfrak{s}_t}(\lambda) \cap \mathbb{Z}^J = \{g_{\mathfrak{s}_t}(b) \mid b \in \mathbf{G}_{q=1}^{\text{up}}(\lambda)\}.$$

**Proposition 5.14.** *Let  $\mathfrak{s}_t$  be a seed for  $t \in \mathbb{T}$ , and let  $k \in J_{\text{fr}}$ . We consider*

$$H_{\mathbf{e}_k, 0} = \{\mathbf{u} \in \mathbb{R}^J \mid \langle \mathbf{u}, \mathbf{e}_k \rangle = 0\} \text{ and } H_{\mathbf{e}_k, 0}^+ = \{\mathbf{u} \in \mathbb{R}^J \mid \langle \mathbf{u}, \mathbf{e}_k \rangle \geq 0\},$$

where  $\mathbf{e}_j := (\delta_{i,j})_{i \in J}$  is the standard unit vector for  $j \in J$ . Then the following hold.

- (1) *The half-space  $H_{\mathbf{e}_k, 0}^+$  contains the cluster cone  $C_{\mathfrak{s}_t}$  and the hyperplane  $H_{\mathbf{e}_k, 0}$  is a supporting hyperplane of  $C_{\mathfrak{s}_t}$ . Moreover,  $H_{\mathbf{e}_k, 0}$  contains a facet of  $C_{\mathfrak{s}_t}$ .*
- (2) *For any dominant integral weight  $\lambda$ , the half-space  $H_{\mathbf{e}_k, 0}^+$  contains the cluster polytope  $\Delta_{\mathfrak{s}_t}(\lambda)$  and  $H_{\mathbf{e}_k, 0}$  is a supporting hyperplane of  $\Delta_{\mathfrak{s}_t}(\lambda)$ .*

*Proof.* (1) Let  $\mathfrak{s}_t = (\{A_j\}_{j \in J}, \varepsilon)$ . Since the cluster variable  $A_j$  ( $j \in J$ ) is contained in  $\mathbf{G}_{q=1}^{\text{up}}(\infty)$  (see [McN21, Qin20]), by the definition of the extended  $g$ -vector  $g_{\mathfrak{s}_t}$  and (5.5), we have

$$g_{\mathfrak{s}_t}(A_j) = \mathbf{e}_j \in C_{\mathfrak{s}_t} \quad \text{for any } j \in J.$$

Since  $C_{\mathfrak{s}_t}$  is a convex polyhedral cone, we have

$$(5.9) \quad F_k := \text{Span}_{\mathbb{R}_{\geq 0}}(B \setminus \{\mathbf{e}_k\}) \subset C_{\mathfrak{s}_t} \cap H_{\mathbf{e}_k, 0} \quad \text{for any frozen index } k \in J_{\text{fr}},$$

where we set  $B := \{\mathbf{e}_j \mid j \in J\}$ . Note that  $F_k$  has codimension one. It follows from (5.7) and (5.9) that

$$C_{\mathfrak{s}_t} \subset H_{\mathbf{e}_k, 0}^+,$$

and  $H_{\mathbf{e}_k, 0}$  contains a facet  $F_k$  of  $C_{\mathfrak{s}_t}$ .

- (2) It follows from  $0 \in \Delta_{\mathfrak{s}_t}(\lambda) \cap H_{\mathbf{e}_k, 0}$  and  $\Delta_{\mathfrak{s}_t}(\lambda) \subset C_{\mathfrak{s}_t} \subset H_{\mathbf{e}_k, 0}^+$ . □

Note that Proposition 5.14 confirms that the conditions (2) and (3) for the criterion in Theorem 4.11 in the case of cluster polytopes.

## 6. Exchange matrix with a large entry between a frozen and unfrozen

The aim of this section is to prove the existence of exchange matrices with an arbitrarily large entry between an unfrozen variable and frozen variables and complete the proof of Theorem 5.1.

**Proposition 6.1.** *Let  $\mathfrak{g}$  be a complex simple Lie algebra of type other than  $A_1, A_2, A_3, A_4$  or  $B_2 = C_2$ . Then there is  $s \in J \setminus J_{\text{uf}}$  such that the extended exchange matrix  $\varepsilon_0 = (\varepsilon_{i,j})_{i \in J, j \in J_{\text{uf}}}$  in (5.1) of the cluster structure of  $\mathbb{C}[U_{w_0}^-]$  satisfies the following property*

$$(6.1) \quad \text{for any } \ell \geq 0, \text{ there exists a matrix } \varepsilon^\ell = (\varepsilon_{i,j}^\ell)_{i \in J, j \in J_{\text{uf}}} \text{ which is mutation equivalent to } \varepsilon_0 = (\varepsilon_{i,j})_{i \in J, j \in J_{\text{uf}}} \text{ such that } -\varepsilon_{r_\ell, s}^\ell \geq \ell \text{ for some } r_\ell \in J_{\text{uf}}.$$

Now, assuming Proposition 6.1, we wrap up the proof of Theorem 5.1.

*Proof of Theorem 5.1.* By Corollary 5.12, we have the family of infinitely many monotone Lagrangian tori on  $G/B$ . We can apply Theorem 4.11 to this family because of Theorem 5.6, Proposition 5.14, and Proposition 6.1. Therefore, the family contains infinitely many monotone Lagrangian tori, no two of which are related by any symplectomorphisms.  $\square$

Let  $J = J_{\text{uf}} \sqcup J_{\text{fr}}$  be a finite set and  $\varepsilon := (\varepsilon_{i,j})_{i \in J, j \in J_{\text{uf}}}$  be an extended skew-symmetrizable matrix. We say that  $\varepsilon$  is *mutation finite* if the mutation equivalence class  $[\varepsilon]$  of  $\varepsilon$  is finite. If  $[\varepsilon]$  is infinite, then we say that  $\varepsilon$  is *mutation infinite*.

For a subset  $J' \subset J$ , the restriction  $\varepsilon|_{J'}$  of  $\varepsilon$  is the matrix obtained from  $\varepsilon$  by restricting to the column set  $J'$  and to the row set  $J' \cap J_{\text{uf}}$ . Note that if  $k \in J'$  is mutable, then we have  $\mu_k(\varepsilon|_{J'}) = (\mu_k(\varepsilon))|_{J'}$ . That is, the restriction commutes with the mutation.

**Lemma 6.2.** *Let  $J = J_{\text{uf}} \sqcup J_{\text{fr}}$  be a finite set and  $\varepsilon := (\varepsilon_{i,j})_{i \in J, j \in J_{\text{uf}}}$  be an extended skew-symmetrizable matrix. Assume that  $\varepsilon$  is mutation infinite and  $\varepsilon|_{J_{\text{uf}}}$  is mutation finite. Then  $\varepsilon$  satisfies (6.2).*

$$(6.2) \quad \text{for any } \ell \geq 0, \text{ there exists a matrix } \varepsilon^\ell = (\varepsilon_{i,j}^\ell)_{i \in J, j \in J_{\text{uf}}} \text{ which is mutation equivalent to } (\varepsilon_{i,j})_{i \in J, j \in J_{\text{uf}}} \text{ such that } -\varepsilon_{r,s}^\ell \geq \ell \text{ for some } r \in J_{\text{uf}} \text{ and } s \in J_{\text{fr}}.$$

Note that if a matrix satisfies (6.2), then so does any matrix mutation equivalent to it.

*Proof.* Let  $\varepsilon = \varepsilon^0, \varepsilon^1, \varepsilon^2, \varepsilon^3, \dots$  be an infinite sequence of pairwise different matrices in the mutation equivalence class of  $\varepsilon$ . Since  $\varepsilon|_{J_{\text{uf}}}$  is mutation finite, the set  $\{\varepsilon^n|_{J_{\text{uf}}} \mid n \geq 0\}$  is finite. It follows that there exists  $m$  such that  $\varepsilon^m|_{J_{\text{uf}}}$  appears infinitely many times in the sequence  $(\varepsilon^n|_{J_{\text{uf}}} \mid n \geq 0)$ . Hence there is an infinite sequence  $m = p_0 < p_1 < p_2 < \dots$  such that  $\varepsilon^{p_k}|_{J_{\text{uf}}} = \varepsilon^m|_{J_{\text{uf}}}$  for all  $k \geq 0$ . Set  $M_k := \max\{|\varepsilon_{r,s}^{p_k}| \mid r \in J_{\text{uf}}, s \in J_{\text{fr}}\}$ . Then  $M_k$  has no upper bound since  $\varepsilon^m = \varepsilon^{p_0}, \varepsilon^{p_1}, \varepsilon^{p_2}, \varepsilon^{p_3}, \dots$  is an infinite sequence of pairwise distinct matrices. We may assume that there are infinitely many  $k$ 's such that  $-\varepsilon_{r,s}^{p_k} = M_k$  for some  $r \in J_{\text{uf}}, s \in J_{\text{fr}}$ . Otherwise, we can change the sign of  $\varepsilon_{r,s}^{p_k}$  by mutating the exchange matrix  $\varepsilon^{p_k}$  in the  $r^{\text{th}}$ -direction. Hence  $\varepsilon^m$  satisfies (6.2).  $\square$

Recall that if  $\varepsilon$  is an extended skew-symmetric matrix, then one can associate a quiver  $Q = Q(\varepsilon)$  with vertices  $J = J_{\text{uf}} \sqcup J_{\text{fr}}$  given by

$$\text{there are } \varepsilon_{j,i}\text{-many arrows from } i \text{ to } j \text{ whenever } \varepsilon_{j,i} \geq 0.$$

To prove Proposition 6.1, we make use of recent classification results of Felikson–Tumarkin on mutation finite quivers. The following is a half of [FT21, Theorem 1.11] when  $\varepsilon$  is an extended skew-symmetric matrix.

**Lemma 6.3.** *Let  $Q$  be a quiver associated with an extended skew-symmetric matrix. Assume that  $J_{\text{fr}} = \{f\}$  and  $Q|_{J_{\text{uf}}}$  is mutation finite. If there exists  $Q'$  mutation equivalent to  $Q$  such that*

- (1)  $Q'$  contains a double arrow  $v_1 \rightrightarrows v_2$ ,
- (2) either  $b_1 \neq -b_2$  or  $b_2 < 0$ , where  $b_i$  is the number of arrows from  $v_i$  to  $f$ ,

then  $Q$  is mutation infinite.


**Remark 6.4.** Indeed, there is an analog version of Lemma 6.3 when  $\varepsilon$  is extended skew-symmetrizable matrix and  $Q$  is the diagram associated with  $\varepsilon$ , see [FT21, Section 10].

**Lemma 6.5** (cf. Lemma 4.1 in [FT21]). *Let  $Q$  be of type  $\tilde{A}_{p,q}$ , that is, the underlying graph is  $(p+q)$ -gon, there are  $p$ -many clockwise arrows and  $q$ -many counterclockwise arrows, and all the vertices of  $Q$  are mutable. Assume that  $p > 0$  and  $q > 0$ . Then the following hold.*

- (1) *For any vertex  $a$  of  $Q$ , there exists a sequence of mutations at vertices different from  $a$  such that the resulting quiver contains a double arrow  $a \Rightarrow v$  for some vertex  $v$ .*
- (2) *Let  $Q'$  be the quiver obtained from  $Q$  by adding a frozen vertex  $f$  with arrows connecting  $f$  to a single vertex  $a$  in  $Q$ . Then  $Q'$  is mutation-infinite.*
- (3) *For any  $\ell \geq 0$ , there is a quiver  $Q^\ell$  mutation equivalent to  $Q'$  such that there exists a mutable vertex  $r_\ell$  with arrows more than  $\ell$  connecting to  $f$ .*

*Proof.* It is well-known that  $Q$  is mutation finite.

(1) Let  $J$  be the set of vertices of  $Q$ . Then there is a sequence of mutations of  $Q$  at vertices  $J \setminus \{a\}$ , all of which are mutations at a sink or a source to obtain the following quiver:



That is,  $a$  is a sink,  $v$  is a source, and there are  $q$ -many counterclockwise arrows (respectively  $p$ -many clockwise arrows) between  $a$  and  $v$ . Then the sequence of mutations

$$\left(\mu_{c_{p-1}} \circ \mu_{c_{p-2}} \circ \cdots \mu_{c_2} \circ \mu_{c_1}\right) \circ \left(\mu_{b_{q-1}} \circ \mu_{b_{q-2}} \circ \cdots \mu_{b_2} \circ \mu_{b_1}\right)$$

yields the desired quiver.

(2) Let  $\mu$  be a mutation sequence in (1). Then  $\mu(Q')$  contains a double arrow  $a \Rightarrow v$  and the vertex  $a$  is the only vertex connected to  $f$ . Hence we have  $b_a > 0$  and  $b_v = 0$  so that  $b_a \neq -b_v$ . Hence  $Q'$  is mutation infinite by the above lemma.

(3) It follows from Lemma 6.2 that  $Q'$  has the desired property since  $Q'$  has only one frozen vertex.  $\square$

**Corollary 6.6.** *Let  $Q$  be a quiver associated with an extended skew-symmetric matrix  $\varepsilon$ . Assume that there is a full subquiver  $Q'$  of  $Q$  such that*

- (1) *the mutable part of  $Q'$  is of type  $\tilde{A}_{p,q}$  with  $p, q > 0$ , and*

- (2)  $Q'$  contains only one frozen vertex  $s$  and there is only one unfrozen vertex connected to  $s$ .

Then we have the property (6.1)

*Proof.* As the restriction commutes with the mutation, (6.1) follows from Lemma 6.5 (3).  $\square$

We are ready to prove Proposition 6.1.

*Proof of Proposition 6.1.* If the Dynkin diagram  $D$  of  $\mathfrak{g}$  contains another Dynkin diagram  $D'$  as a full subgraph, then the longest element  $w_0$  of the Weyl group of  $\mathfrak{g}$  can be decomposed into

$$w_0 = vw' \quad \text{where } w' \text{ is the longest element of the parabolic subgroup of type } D'.$$


Let  $\underline{w}'$  be a reduced expression of  $w'$  and let  $\underline{w}_0$  be a reduced expression of  $w_0$  extending  $\underline{w}'$  with respect to the above decomposition. Then by (5.1), the matrix attached to  $\underline{w}'$  is a restriction of the one attached to  $\underline{w}_0$ . Moreover, the frozen indices are preserved in the restriction. Hence one may assume that  $\mathfrak{g}$  is of type  $A_5$ ,  $D_4$ ,  $C_3$ ,  $B_3$ , or  $G_2$ . We deal with the five cases in Lemma 6.7, 6.8, 6.9, 6.10, and 6.11.  $\square$

**Lemma 6.7.** *If  $\mathfrak{g}$  is of type  $A_5$ , then (6.1) holds.*


*Proof.* For type  $A_5$ , take

$$\underline{w}_0 = s_1 s_2 s_1 s_3 s_2 s_1 s_4 s_3 s_2 s_1 s_5 s_4 s_3 s_2 s_1$$

where the Dynkin diagram is



The quiver attached to  $\underline{w}_0$  is




and the full subquiver with the vertices 14, 9, 8, 4, 2, 3 and 6 satisfies the condition in Corollary 6.6. Here the boxed entries are frozen vertices.  $\square$

**Lemma 6.8.** *If  $\mathfrak{g}$  is of type  $D_4$ , then (6.1) holds.*


*Proof.* For the case  $\mathfrak{g} = D_4$ , take

$$\underline{w}_0 = s_2 s_4 s_1 s_2 s_4 s_3 s_2 s_4 s_1 s_2 s_3 s_4$$

where the Dynkin diagram is



The quiver attached to  $\underline{w}_0$  is



and the full subquiver with the vertices 10, 7, 3, 1, 2 and 5 satisfies the condition in Corollary 6.6.  $\square$

**Lemma 6.9.** *If  $\mathfrak{g}$  is of type  $C_3$ , then (6.1) holds.*

*Proof.* In the case  $\mathfrak{g} = C_3$ , the Dynkin diagram and Cartan matrix are given by

$$C_3 \quad \begin{array}{ccc} \circ & \circ & \circ \\ 1 & 2 & 3 \end{array} \quad \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -2 \\ 0 & -1 & 2 \end{pmatrix}.$$

Take

$$\underline{w}_0 = s_3 s_2 s_3 s_2 s_1 s_2 s_3 s_2 s_1$$

Then

$$\varepsilon = \begin{pmatrix} 0 & -2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 2 & 0 & 0 & 0 & -2 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 & 0 & 0 & -1 & 1 & 0 \end{pmatrix}$$

Let

$$J' = \{1, 2, 3, 6, 8\}$$

Then we have

$$(6.3) \quad \varepsilon|_{J'} = \begin{pmatrix} 0 & -2 & 1 & 0 & 0 \\ 1 & 0 & -1 & 0 & 0 \\ -1 & 2 & 0 & -2 & 0 \\ 0 & 0 & 1 & 0 & 1 \end{pmatrix}.$$

Then by taking a sequence of mutation we get

$$\mu_3 \circ \mu_2 \circ \mu_6(\varepsilon|_{J'}) = \begin{pmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 2 & 0 \\ -1 & 2 & 0 & -2 & 0 \\ 0 & -2 & 1 & 0 & -2 \end{pmatrix}$$

Note that the corresponding *diagram* (see [FZ03], [FT21, Section 9.1] for the definition of diagram associated with an extended skew-symmetrizable matrix with one frozen index) is

$$(6.4) \quad \begin{array}{ccccccc} & & 2 & & & & \\ & & \downarrow & \swarrow & & & \\ & & 2 & & & & \\ 1 & \longleftarrow & 3 & \xrightarrow{2} & 6 & \longrightarrow & 8 \end{array}$$

The mutable part of the diagram is of type  $\tilde{C}_3$  (see [FT21, Figure 9.2]), which is mutation finite. By applying [FT21, Theorem 9.4] (which is an analog of Lemma 6.3 for skew-symmetrizable cases), the matrix  $\varepsilon|_{J'}$  is mutation infinite and hence the matrix  $\varepsilon$  has the desired property by Lemma 6.2.  $\square$

**Lemma 6.10.** *If  $\mathfrak{g}$  is of type  $B_3$ , then (6.1) holds.*

*Proof.* In the case  $\mathfrak{g} = B_3$ , the Dynkin diagram and Cartan matrix are given by

$$B_3 \quad \begin{array}{ccc} \circ & \text{---} & \circ \rightleftarrows \circ \\ 1 & 2 & 3 \end{array} \quad \begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -2 & 2 \end{pmatrix}$$

Take

$$\underline{w}_0 = s_3 s_2 s_3 s_2 s_1 s_2 s_3 s_2 s_1.$$

Then we have

$$\varepsilon = \begin{pmatrix} 0 & -1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 2 & 0 & -2 & 1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 1 & 0 & 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & -1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 2 & -1 & 0 & 0 & -2 & 1 & 0 \end{pmatrix}$$

Let

$$J' = \{1, 2, 3, 6, 8\}$$

Then we have

$$\varepsilon|_{J'} = \begin{pmatrix} 0 & -1 & 1 & 0 & 0 \\ 2 & 0 & -2 & 0 & 0 \\ -1 & 1 & 0 & -1 & 0 \\ 0 & 0 & 2 & 0 & 1 \end{pmatrix}$$

and we have

$$\mu_3(\varepsilon|_{J'}) = \begin{pmatrix} 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 2 & -2 & 0 \\ 1 & -1 & 0 & 1 & 0 \\ 0 & 2 & -2 & 0 & 1 \end{pmatrix}$$

Note that the corresponding diagram is again (6.4) so that we have the desired property.  $\square$

**Lemma 6.11.** *If  $\mathfrak{g}$  is of type  $G_2$ , then (6.1) holds.*

*Proof.* In the case of  $G_2$  the Dynkin diagram and Cartan matrix are

$$G_2 \quad \begin{array}{c} \circ \rightleftarrows \circ \\ 1 \quad 2 \end{array} \quad \begin{pmatrix} 2 & -3 \\ -1 & 2 \end{pmatrix}$$


We take

$$\underline{w}_0 = s_1 s_2 s_1 s_2 s_1 s_2.$$


to obtain the exchange matrix

$$\varepsilon = \begin{pmatrix} 0 & -1 & 1 & 0 & 0 & 0 \\ 3 & 0 & -3 & 1 & 0 & 0 \\ -1 & 1 & 0 & -1 & 1 & 0 \\ 0 & -1 & 3 & 0 & -3 & 1 \end{pmatrix}$$

Taking the restriction to  $J' = \{1, 2, 3, 5\}$ , we obtain the diagram



which is mutation equivalent to



by taking the composition of mutations  $\mu_2 \circ \mu_1 \circ \mu_3 \circ \mu_2 \circ \mu_1$ . Since the mutable part of the above diagram is of type  $\tilde{G}_2$  (see [FT21, Figure 9.2]), which is mutation-finite, by Lemma 6.3 we conclude that it is mutation infinite. Thus we obtain the desired property by Lemma 6.2.  $\square$

**Remark 6.12.** If  $\mathfrak{g}$  is of type  $A_1, A_2, A_3, A_4$  or  $B_2$ , then the cluster algebra  $\mathbb{C}[U_{w_0}^-]$  is finite type. Hence the exchange matrices of  $\mathbb{C}[U_{w_0}^-]$  are mutation finite so that they do *not* satisfy (6.1)

## References

- [And13] Dave Anderson, *Okounkov bodies and toric degenerations*, Math. Ann. **356** (2013), no. 3, 1183–1202.
- [Aur07] Denis Auroux, *Mirror symmetry and T-duality in the complement of an anticanonical divisor.*, J. Gökova Geom. Topol. GGT **1** (2007), 51–91.
- [Aur15] Denis Auroux, *Infinitely many monotone Lagrangian tori in  $\mathbb{R}^6$* , Invent. Math. **201** (2015), no. 3, 909–924.
- [BC09] Paul Biran and Octav Cornea, *A Lagrangian quantum homology*, New perspectives and challenges in symplectic field theory, CRM Proc. Lecture Notes, vol. 49, Amer. Math. Soc., Providence, RI, 2009, pp. 1–44.
- [BFZ05] Arkady Berenstein, Sergey Fomin, and Andrei Zelevinsky, *Cluster algebras. III. Upper bounds and double Bruhat cells*, Duke Math. J. **126** (2005), no. 1, 1–52.
- [BGM22] Pieter Belmans, Sergey Galkin, and Swarnava Mukhopadhyay, *Graph potentials and symplectic geometry of moduli spaces of vector bundles*, preprint (2022), arXiv:2206.11584.
- [Bre23] Joé Brendel, *Local exotic tori*, preprint (2023), arXiv:2310.11359.
- [Brø83] Arne Brøndsted, *An introduction to convex polytopes*, Graduate Texts in Mathematics, vol. 90, Springer-Verlag, New York-Berlin, 1983.
- [BZ97] Arkady Berenstein and Andrei Zelevinsky, *Total positivity in Schubert varieties*, Comment. Math. Helv. **72** (1997), no. 1, 128–166.



- [BZ01] Arkady Berenstein and Andrei Zelevinsky, *Tensor product multiplicities, canonical bases and totally positive varieties*, Invent. Math. **143** (2001), no. 1, 77–128.
- [Cal02] Philippe Caldero, *Toric degenerations of Schubert varieties*, Transform. Groups **7** (2002), no. 1, 51–60.
- [Cas23] Marco Castronovo, *Exotic Lagrangian tori in Grassmannians*, Quantum Topol. **14** (2023), no. 1, 65–99.
- [Che96] Yu. V. Chekanov, *Lagrangian tori in a symplectic vector space and global symplectomorphisms*, Math. Z. **223** (1996), no. 4, 547–559.
- [CHW23] Soham Chanda, Amanda Hirschi, and Luya Wang, *Infinitely many monotone Lagrangian tori in higher projective spaces*, preprint (2023), arXiv:2307.06934.
- [CK21] Yunhyung Cho and Yoosik Kim, *Monotone Lagrangians in flag varieties*, Int. Math. Res. Not. IMRN (2021), no. 18, 13892–13945.
- [CO06] Cheol-Hyun Cho and Yong-Geun Oh, *Floer cohomology and disc instantons of Lagrangian torus fibers in Fano toric manifolds*, Asian J. Math. **10** (2006), no. 4, 773–814.
- [CP14] Cheol-Hyun Cho and Mainak Poddar, *Holomorphic orbi-discs and Lagrangian Floer cohomology of symplectic toric orbifolds*, J. Differential Geom. **98** (2014), no. 1, 21–116.
- [EP97] Yakov Eliashberg and Leonid Polterovich, *The problem of Lagrangian knots in four-manifolds*, Geometric topology (Athens, GA, 1993), AMS/IP Stud. Adv. Math., vol. 2, Amer. Math. Soc., Providence, RI, 1997, pp. 313–327.
- [FG06] Vladimir Fock and Alexander Goncharov, *Moduli spaces of local systems and higher Teichmüller theory*, Publ. Math. Inst. Hautes Études Sci. (2006), no. 103, 1–211.
- [FG09] Vladimir V. Fock and Alexander B. Goncharov, *Cluster ensembles, quantization and the dilogarithm*, Ann. Sci. Éc. Norm. Supér. (4) **42** (2009), no. 6, 865–930.
- [FH21] Naoki Fujita and Akihiro Higashitani, *Newton–Okounkov bodies of flag varieties and combinatorial mutations*, Int. Math. Res. Not. IMRN (2021), no. 12, 9567–9607.
- [FO17] Naoki Fujita and Hironori Oya, *A comparison of Newton–Okounkov polytopes of Schubert varieties*, J. Lond. Math. Soc. (2) **96** (2017), no. 1, 201–227.
- [FO20] Naoki Fujita and Hironori Oya, *Newton–Okounkov polytopes of Schubert varieties arising from cluster structures*, preprint (2020), arXiv:2002.09912.
- [FOOO09] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono, *Lagrangian intersection Floer theory: anomaly and obstruction. Part I*, AMS/IP Studies in Advanced Mathematics, vol. 46.1, American Mathematical Society, Providence, RI; International Press, Somerville, MA, 2009.
- [FOOO10] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono, *Lagrangian Floer theory on compact toric manifolds. I*, Duke Math. J. **151** (2010), no. 1, 23–174.
- [FT21] Anna Felikson and Pavel Tumarkin, *Cluster algebras of finite mutation type with coefficients*, preprint (2021), arXiv:2110.12917.
- [Fuj18] Naoki Fujita, *Newton–Okounkov bodies for Bott–Samelson varieties and string polytopes for generalized Demazure modules*, J. Algebra **515** (2018), 408–447.
- [FZ02] Sergey Fomin and Andrei Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. **15** (2002), no. 2, 497–529.
- [FZ03] Sergey Fomin and Andrei Zelevinsky, *Cluster algebras. II. Finite type classification*, Invent. Math. **154** (2003), no. 1, 63–121.
- [FZ07] Sergey Fomin and Andrei Zelevinsky, *Cluster algebras. IV. Coefficients*, Compos. Math. **143** (2007), no. 1, 112–164.
- [GHK15a] Mark Gross, Paul Hacking, and Sean Keel, *Birational geometry of cluster algebras*, Algebr. Geom. **2** (2015), no. 2, 137–175.
- [GHK15b] Mark Gross, Paul Hacking, and Sean Keel, *Mirror symmetry for log Calabi–Yau surfaces I*, Publ. Math. Inst. Hautes Études Sci. **122** (2015), 65–168.
- [GHKK18] Mark Gross, Paul Hacking, Sean Keel, and Maxim Kontsevich, *Canonical bases for cluster algebras*, J. Amer. Math. Soc. **31** (2018), no. 2, 497–608.
- [GL96] N. Gonciulea and V. Lakshmibai, *Degenerations of flag and Schubert varieties to toric varieties*, Transform. Groups **1** (1996), no. 3, 215–248.
- [GLS11] Christof Geiß, Bernard Leclerc, and Jan Schröer, *Kac–Moody groups and cluster algebras*, Adv. Math. **228** (2011), no. 1, 329–433.
- [GM22] Sergey Galkin and Grigory Mikhalkin, *Singular symplectic spaces and holomorphic membranes*, Eur. J. Math. **8** (2022), no. 3, 932–951.

- [GP00] Oleg Gleizer and Alexander Postnikov, *Littlewood-Richardson coefficients via Yang-Baxter equation*, Internat. Math. Res. Notices (2000), no. 14, 741–774.
- [GS11] Mark Gross and Bernd Siebert, *From real affine geometry to complex geometry*, Ann. of Math. (2) **174** (2011), no. 3, 1301–1428.
- [GY17] K. R. Goodearl and M. T. Yakimov, *Quantum cluster algebra structures on quantum nilpotent algebras*, Mem. Amer. Math. Soc. **247** (2017), no. 1169, vii+119.
- [GY20] K. R. Goodearl and M. T. Yakimov, *Integral quantum cluster structures*, preprint (2020), arXiv:2003.04434.
- [HK15] Megumi Harada and Kiumars Kaveh, *Integrable systems, toric degenerations and Okounkov bodies*, Invent. Math. **202** (2015), no. 3, 927–985.
- [Kas90] Masaki Kashiwara, *Crystalizing the  $q$ -analogue of universal enveloping algebras*, Comm. Math. Phys. **133** (1990), no. 2, 249–260.
- [Kas91] Masaki Kashiwara, *On crystal bases of the  $Q$ -analogue of universal enveloping algebras*, Comm. Math. Phys. **133** (1990), no. 2, 249–260.
- [Kas93] Masaki Kashiwara, *Global crystal bases of quantum groups*, Duke Math. J. **69** (1993), no. 2, 455–485.
- [Kav15] Kiumars Kaveh, *Crystal bases and Newton-Okounkov bodies*, Duke Math. J. **164** (2015), no. 13, 2461–2506.
- [KK12] Kiumars Kaveh and A. G. Khovanskii, *Newton-Okounkov bodies, semigroups of integral points, graded algebras and intersection theory*, Ann. of Math. (2) **176** (2012), no. 2, 925–978.
- [KK19] Masaki Kashiwara and Myungho Kim, *Laurent phenomenon and simple modules of quiver Hecke algebras*, Compos. Math. **155** (2019), no. 12, 2263–2295.
- [KM05] Mikhail Kogan and Ezra Miller, *Toric degeneration of Schubert varieties and Gelfand-Tsetlin polytopes*, Adv. Math. **193** (2005), no. 1, 1–17.
- [KN13] Alexander M. Kasprzyk and Benjamin Nill, *Fano polytopes*, Strings, gauge fields, and the geometry behind, World Sci. Publ., Hackensack, NJ, 2013, pp. 349–364.
- [KO21] Yoshiyuki Kimura and Hironori Oya, *Twist automorphisms on quantum unipotent cells and dual canonical bases*, Int. Math. Res. Not. IMRN (2021), no. 9, 6772–6847.
- [KP11] Ludmil Katzarkov and Victor Przyjalkowski, *Landau-Ginzburg models—old and new*, Proceedings of the Gökova Geometry-Topology Conference 2011, (2012), 97–124.
- [Lit98] P. Littelmann, *Cones, crystals, and patterns*, Transform. Groups **3** (1998), no. 2, 145–179.
- [LM09] Robert Lazarsfeld and Mircea Mustață, *Convex bodies associated to linear series*, Ann. Sci. Éc. Norm. Supér. (4) **42** (2009), no. 5, 783–835.
- [Lus90] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), no. 2, 447–498.
- [Lus91] G. Lusztig, *Quivers, perverse sheaves, and quantized enveloping algebras*, J. Amer. Math. Soc. **4** (1991), no. 2, 365–421.
- [McN21] Peter J. McNamara, *Cluster Monomials are Dual Canonical*, preprint (2021), arXiv:2112.04109.
- [NNU10] Takeo Nishinou, Yuichi Nohara, and Kazushi Ueda, *Toric degenerations of Gelfand-Cetlin systems and potential functions*, Adv. Math. **224** (2010), no. 2, 648–706.
- [Oh93] Yong-Geun Oh, *Floer cohomology of Lagrangian intersections and pseudo-holomorphic disks. I*, Comm. Pure Appl. Math. **46** (1993), no. 7, 949–993.
- [Oko96] Andrei Okounkov, *Brunn-Minkowski inequality for multiplicities*, Invent. Math. **125** (1996), no. 3, 405–411.
- [Prz07] Victor Przyjalkowski, *On Landau-Ginzburg models for Fano varieties*, Commun. Number Theory Phys. **1** (2007), no. 4, 713–728.
- [Qin17] Fan Qin, *Triangular bases in quantum cluster algebras and monoidal categorification conjectures*, Duke Math. J. **166** (2017), no. 12, 2337–2442.
- [Qin20] Fan Qin, *Dual canonical bases and quantum cluster algebras*, preprint (2020), arXiv:2003.13674.
- [RW19] K. Rietsch and L. Williams, *Newton-Okounkov bodies, cluster duality, and mirror symmetry for Grassmannians*, Duke Math. J. **168** (2019), no. 18, 3437–3527.
- [Ste22] Christian Steinert, *Reflexivity of Newton-Okounkov bodies of partial flag varieties* Representation Theory of the American Mathematical Society **26** (2022) 859–873.
- [Ton18] Dmitry Tonkonog, *String topology with gravitational descendants, and periods of Landau-Ginzburg potentials*, preprint (2018), arXiv:1801.06921.
- [Via14] Renato Vianna, *On exotic Lagrangian tori in  $\mathbb{C}P^2$* , Geom. Topol. **18** (2014), no. 4, 2419–2476.

- [Via16] Renato Ferreira de Velloso Vianna, *Infinitely many exotic monotone Lagrangian tori in  $\mathbb{C}\mathbb{P}^2$* , J. Topol. **9** (2016), no. 2, 535–551.
- [Via17] Renato Vianna, *Infinitely many monotone Lagrangian tori in del Pezzo surfaces.*, Selecta Math. (N.S.) **23** (2017), no. 3, 1955–1996.
- [Wil13] Harold Williams, *Cluster ensembles and Kac-Moody groups*, Adv. Math. **247** (2013), 1–40.

DEPARTMENT OF MATHEMATICS EDUCATION, SUNGKYUNKWAN UNIVERSITY, SEOUL, REPUBLIC OF KOREA  
*Email address:* `yunhyung@skku.edu`

DEPARTMENT OF MATHEMATICS, KYUNG HEE UNIVERSITY, SEOUL, REPUBLIC OF KOREA  
*Email address:* `mkim@khu.ac.kr`

DEPARTMENT OF MATHEMATICS AND INSTITUTE OF MATHEMATICAL SCIENCE, PUSAN NATIONAL UNIVERSITY, BUSAN, REPUBLIC OF KOREA  
*Email address:* `yosik@pusan.ac.kr`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SEOUL, SEOUL, REPUBLIC OF KOREA  
*Email address:* `epark@uos.ac.kr`