

A PRIORI ENERGY ESTIMATE WITH DECAY IN WEIGHTED NORMS FOR THE WATER-WAVES PROBLEM WITH CONTACT POINTS

MEI MING

ABSTRACT. We prove a weighted a priori energy estimate for the two dimensional water-waves problem with contact points in the absence of gravity and surface tension. When the surface graph function and its time derivative have some decay near the contact points, we show that there is corresponding decay for the velocity, the pressure and other quantities in a short time interval. As a result, we have fixed contact points and contact angles. To prove the energy estimate, a conformal mapping is used to transform the equation for the mean curvature into an equivalent equation in a flat strip with some weights. Moreover, the weighted limits at contact points for the velocity, the pressure etc. are tracked and discussed. Our formulation can be adapted to deal with more general cases.

CONTENTS

1. Introduction	1
1.1. Main result of the paper	4
1.2. Organization of the paper	6
1.3. Notations	6
2. Preliminaries	7
3. Weighted elliptic estimates related to D-N operator	12
3.1. Weighted elliptic estimates on the strip domain	12
3.2. Weighted estimates for D-N operator	17
4. Reformulation of the problem and preparations	19
4.1. Geometric formulation in the strip domain	19
4.2. The proper weights	22
4.3. Preparations for the energy estimate	23
4.4. Weighted estimates for the conformal mapping \mathcal{T}	24
4.5. Weighted estimates for the velocity and other quantities	35
4.6. Modification of the equation	42
5. A priori estimates	43
Acknowledgements	50
References	50

1. INTRODUCTION

We consider a two-dimensional domain Ω with two fixed contact points p_l, p_r in the absence of gravity and surface tension. In fact, the domain

$$\Omega = \{(x, z) \mid b(x) \leq z \leq \zeta(t, x), x_l \leq x \leq x_r\}$$

is changing with time, where

$$\Gamma_t = \{(x, z) \mid z = \zeta(t, x), x_l \leq x \leq x_r\}$$

is the free surface curve denoted by a graph function ζ and x_l, x_r are two constants.

The bottom is denoted by

$$\Gamma_b = \{(x, z) \mid z = b(x), x_l \leq x \leq x_r\},$$

where we assume that $b(x)$ can be smooth enough, and Γ_b becomes some line segment near the corners for the sake of simplicity.

The points of intersection p_l, p_r between Γ_t and Γ_b take place only at $x = x_l, x_r$ (the left and the right), which means

$$b(x) < \zeta(t, x) \quad \text{when} \quad x_l < x < x_r, \quad t \in [0, T]$$

and $b(x_i) = \zeta(t, x_i)$ for $i = l, r$ and some $T > 0$.




Figure 1. The physical domain

Meanwhile, we have some more assumptions on the free surface Γ_t :

Assumption 1. Γ_t are assumed to intersect with Γ_b at two fixed contact points when $t \in [0, T]$ for some $T > 0$: There are the left contact point p_l with fixed contact angle ω and the right contact point p_r with fixed angle ω_r , and both $\omega, \omega_r \in (0, \pi/2)$. Moreover, we take

$$\partial_x \zeta|_{x_i} = 0, \quad i = l, r, \quad t \in [0, T]$$

which means the tangential lines at the contact points on Γ_t are both horizontal.

We plan to investigate possible behaviors of ideal incompressible irrotational water waves in the case of fixed contact angles (some decay of the free surface indeed) and zero gravity. To present the water-waves system, we start with Euler's equation:

$$(1) \quad \partial_t v + v \cdot \nabla v = -\nabla P \quad \text{in } \Omega, \quad t \geq 0$$

where $v = (v_1, v_2)^T$ is the velocity field and P is the pressure to be determined. We use the notation of material derivative frequently: $D_t = \partial_t + v \cdot \nabla$.

The incompressibility and the irrotationality of the fluid are expressed by

$$(2) \quad \nabla \cdot v = 0, \quad \nabla \times v = 0 \quad \text{in } \Omega, \quad t \geq 0.$$

At the bottom, there is no fluid particles transported, which means

$$(3) \quad v \cdot n_b|_{\Gamma_b} = 0 \quad \text{for } t \geq 0,$$

where n_b is the unit outward normal vector on Γ_b .

The kinematic condition holds at the free surface, i.e.

$$(4) \quad \partial_t \zeta = (1 + |\partial_x \zeta|^2)^{\frac{1}{2}} v \cdot n_t|_{\Gamma_t: z=\zeta} \quad \text{for } t \geq 0,$$

where n_t is the unit outward normal vector on Γ_t .

Moreover, ignoring the effect of surface tension, we have

$$(5) \quad P|_{\Gamma_t} = 0 \quad \text{for } t \geq 0$$

without loss of generality.

Before introducing our result, we recall briefly earlier works on the well-posedness of the water-waves problem with smooth boundaries, where one has smooth surface Γ_t satisfying

$$\Gamma_t \cap \Gamma_b = \emptyset.$$

There is a very rich literature on various local well-posedness results for this problem, see [51, 69, 70, 19, 65, 66, 40, 32, 29] and [3, 14, 45, 17, 57, 58, 55, 9, 52, 7, 71, 6, 41, 64] etc.. Starting around 2009, there are quite a few results focusing on the global well-posedness with small initial data and some long-time behaviors, see [68, 24, 5, 33, 21, 28, 63, 61] etc.. Moreover, there are also some works concerning “splash” or “splat” singularities on the free surfaces, see [12, 13, 18] etc..

Compared to the various results for smooth-boundary water waves, there are only very limited results on the mathematical theory of well-posedness when the boundary is Lipschitz, i.e. water waves with non-smooth boundaries (non-smooth-boundary water waves). In fact, the mathematical research for this problem just started these years, and there are in general two kinds of problems depending on the location of the corners.

The first kind of problem is the contact-line (contact-point in two dimensional case) problem, where there are intersections between the free surface and the fixed bottom, i.e.

$$\Gamma_t \cap \Gamma_b \neq \emptyset.$$

As a pioneer work for this problem, Alazard, Burq and Zuily mention in [4] a special case with zero surface tension, where the contact angle with respect to a vertical wall is equal to $\pi/2$ (“the right angle” to keep the Taylor’s sign condition, i.e. $\mathbf{a} = -\nabla_{n_t} P|_{\Gamma_t} > 0$). They use a periodic extension to turn this problem into a classical smooth periodic case. de Poyferré [20] proves an a priori energy estimate in Sobolev spaces in general n -dimensional bounded domains without surface tension. The contact angles there are assumed to be small to make sure that the solution has sufficient regularity near the corner. Under a similar assumption of small contact angles, Ming and Wang obtain the a priori estimate as well as the local well-posedness in two-dimensional corner domains in the presence of surface tension, see [47, 48]. Meanwhile, notice that both [20] and [47, 48] use the geometric formulation introduced in [57].

There are some works concerning properties of the corresponding Dirichlet-Neumann operator in Lipschitz domains, which is a very important ingredient in the research of water waves. Ming and Wang [46] adapts some elliptic theory in corner domains (see for example [25]) to the desired form with boundary dependence. Agrawal and Alazard prove in [2] some refined Rellich inequalities and some sharp inequality for the D-N operator.

Besides, there are also some more related works on models of water waves with contact lines. Lannes and Métivier [43] considers the Green-Naghdi system in a beach-type domain, which is a shallow-water model of the water-waves problem. Lannes [42] works on the floating-body problem and proposes a new formulation which can be easily generalized in order to take into account the presence of a floating body. Lannes and Iguchi [30] considers initial boundary value problem with a free boundary arising in wave-structure interaction, which contains the floating problem in the shallow water regime. In [31] Iguchi and Lannes prove the local well-posedness for a shallow water model with a fixed partially immersed object. For more related works, see [10, 8] etc..

For contact-line problems of Navier-Stokes system, Guo and Tice [26] shows a priori estimates in the case of Stokes equations. Tice and Zheng [62] proves the local well-posedness of the contact-line problem in two-dimensional Stokes flow. Moreover, Guo and Tice [27] proves an a priori estimate for the contact-line problem of two-dimensional Navier-Stokes flow. For Darcy’s flow, one can see [35, 36] etc..

The second kind of problem focuses on crests or cusps on the free surface, i.e. the surface is Lipschitz. Kinsey and Wu, Wu [34, 67] prove the a priori estimate and the local well-posedness for two-dimensional gravity water waves in the half plane, where there is a crest on the free surface. The Taylor’s sign degenerates at the crest point, and the authors flatten the domain with a Riemann mapping and then use some natural weighted norms to finish the energy estimate. The crest angle is less than $\pi/2$ here and doesn’t change with respect to time. Based on these works, Agrawal [1] shows that an initial interface with angled crests remains angled crests in a more general way. Córdoba, Enciso and Grubic [15] considers the case with cusps and crests on the surface in the absence of gravity, where the angles of these crests are less than $\pi/2$ and change in time. Córdoba, Enciso and

Grubic prove in [16] the local well-posedness for gravity water waves in weighted Sobolev spaces which allow for interfaces with corners. In these two works, the Taylor's sign always degenerates at the crest point on the surface. Meanwhile, the weighted norms used in [16] are introduced in [38] for example, including both so-called homogeneous norm and inhomogeneous norm.

We would like to mention the famous Stokes waves which can be dated back to works by G. Stokes [59, 60], where traveling-wave solutions with limit crest angle $2\pi/3$ are discussed.

It is straightforward to see that, the main difference between non-smooth-boundary water waves and smooth-boundary water waves lies in the corners on the boundary. As a result, the analysis involving the corners (i.e. domain singularities) and proper adaptations of the existing results of smooth-boundary water waves become the key points in the non-smooth water waves.

1.1. Main result of the paper. We focus on the case of two dimensional water waves with contact points, where both the surface function ζ and the velocity v have some decay near the corners. The Taylor's sign is also assumed to degenerate at the contact points but it keeps the positive sign with some weight, see Assumption 2. Meanwhile, our problem is considered in the absence of gravity and surface tension.

In fact, it is not clear so far whether the Taylor's sign vanishes (degenerates) at the contact point, so it makes sense for us to consider the special case when the Taylor's sign degenerates at corner points. Moreover, similar weighted conditions for the Taylor's sign can be found in the surface-crest case, see [34, 16].

Our work is inspired by two observations. The first observation is about the singularity decomposition for solutions to elliptic systems (see [25] for example) and the homogeneous norm introduced in [38] for elliptic estimates. On one hand, for an elliptic system in a corner domain, the singular part of the solution usually takes the form of r^λ , where r is the radius to the corner point, and λ is some related eigenvalue of this system.

On the other hand, according to the elliptic theory in corner domains (see for example [38]), proper decay near the corners in boundary conditions can result in corresponding decay near the corners for the solution to an elliptic system. This kind of decay can be described by some homogeneous norms. In fact, for a sector \mathcal{K} with the vertices at the origin, the so-called homogeneous norm is defined as

$$\|u\|_{V_\beta^l(\mathcal{K})} = \left(\sum_{|\alpha| \leq l} \int_{\mathcal{K}} r^{2(\beta-l+|\alpha|)} |\nabla_X^\alpha u|^2 dX \right)^{\frac{1}{2}}$$

with r the radius with respect to the vertices, l the order of derivatives and β the weight. A complete theory of elliptic estimates is developed in [38] using this kind of weighted norms (which is similar as the elliptic theory in Sobolev spaces), and the authors use a conformal mapping to transform the domain into a flat strip sometimes. According to this theory, these weighted norms can describe solutions to elliptic systems with boundary conditions given in the form of r^s with s some real number (there are some more conditions about the power s essentially avoiding the eigenvalues, see [38]). Therefore, as we stated before, *proper decay on the boundary can lead to corresponding decay of the solution.*

The second observation is that one can use a conformal mapping \mathcal{T} to transform a fixed flat strip \mathcal{S}_0 (with complex coordinates $\tilde{Z} = \tilde{x} + i\tilde{z}$, see (6)) to the physical domain Ω at time t (see Figure 2 in Section 2), which behaves like $e^{\tilde{Z}}$ if we focus on the left corner of Ω and a natural weight (see (9)) appears:

$$\alpha(\tilde{x}) \approx |\mathcal{T}'| \approx e^{\tilde{x}} \quad \text{near the left corner where } \tilde{x} \rightarrow -\infty.$$

For more details about \mathcal{T} , one can see Section 2. Consequently, there will be some coefficients in form of \mathcal{T} appearing everywhere in the equation of water waves along with derivatives, but a very good point here is that one can rewrite the D-N operator in Ω into the D-N operator in the flat strip with some coefficient (see (28))! In fact, it is lucky to see that properties for the D-N operator defined in a flat strip are already well-studied, see for example [40, 41].

As a result, it is interesting to see if this kind of homogeneous norms from [38] can be adjusted to our problem, where we transform our equation to an equivalent equation in the flat strip \mathcal{S}_0 with some natural weights. As in our previous works [47, 48, 49], we use the geometric formulation following Shatah and Zeng [57], so the system of water waves (1)-(5) is rewritten into an equation of the mean curvature κ on the free surface, see (23). Using a conformal mapping, this equation of κ is transformed into a weighted equation on the surface the the flat strip.

Based on the homogeneous norms in [38], we introduce some weighted norms in the strip for the surface function ζ , the velocity v etc.. However, there are some difficulties arising even after we identify all the weighted norms and succeed to prove the corresponding weighted elliptic theory.

The first difficulty lies in the estimate for the conformal mapping. The higher-order-derivative estimate is needed in the form of weighted estimate, and the time-dependent estimate is also involved as well. Combining our weighted elliptic theory with some computations in [56], we succeed in finishing the desired estimates (see Section 4.4).

The second difficulty is about the higher-order terms of velocity v and pressure P on the bottom, while we notice that the surface Γ_t and the bottom Γ_b intersect at the contact points with different boundary conditions in the related elliptic systems for v and P etc.. Compared to estimates for smooth-boundary water waves or surface-crest water waves, this is a new part to deal with. Meanwhile, compared to the case of contact points with surface tension, the regularity of the surface Γ_t (i.e. the graph function ζ) is lower in our case without surface tension, and there will be a loss of 1/2-order derivative for the velocity as long as one uses the kinematic condition (4). Therefore, the estimate for v is very tricky. Luckily with our chosen weighted norms in the flat strip, we succeed to prove a delicate higher-order estimate for v (see Section 4.5) combining some tricks in [57] (which considers the case of smooth free surface without any bottom).

Another difficulty lies in dealing with the corner behaviors of the surface, velocity and pressure etc.. In fact, it turns out that we need to trace all the weighted limits of these quantities at the corners. The non-zero weighted limits take place since we assume that the weighted Taylor's sign is bounded below by a positive number, which together with the boundary condition on the bottom of pressure P imply that the weighted limits of pressure at corners are not vanishing. As a result, some more analysis for the corresponding weighted limits is needed. In our settings, we show that the weighted limits for $\partial_t \zeta$, v , P and other quantities are constant with respect to time t .

As a result, one can see that our a priori estimate is different from all the known works mentioned before on well-posedness of water waves. So far there are only two works on the case of contact points without surface tension, see [4] (for a special angle $\pi/2$) and [20] (a priori estimate for general n dimensions), where the Taylor's sign is assumed to be bounded below and Sobolev spaces are used.

Moreover, our result is also very different from the weighted formulations in [34, 15]. The first point is that there are contact points and the bottom Γ_b in our case, which is different from the case of crests on the free surface from the physical settings. The second point is that higher-order derivative terms appear on the bottom when we consider the energy estimate (as mentioned before), while there are no such terms for the case without fixed bottom. The last point is again about the weighted limits, which arise because of the weighted Taylor's sign and the boundary condition of pressure at the bottom.

The main theorem of our paper is stated below in an informal way, while the exact main theorem is Theorem 2 in Section 5. To be more clear, we denote by $\tilde{\zeta} = \zeta \circ \mathcal{T}|_{\Gamma_t}$, $\tilde{\partial}_t \zeta = (\partial_t \zeta) \circ \mathcal{T}|_{\Gamma_t}$ for the surface graph function $z = \zeta$ with a slight abuse of notation (see Section 4). Moreover, we recall the weight function $\alpha = \alpha(\tilde{x})$ introduced by the conformal mapping \mathcal{T} . The energy $\mathcal{E}(t)$ is defined in the beginning of Section 5, which contains weighted norms of $(k + 1/2)$ -th order derivative of the mean curvature $\kappa \circ \mathcal{T}|_{\Gamma_t}$ and k -th order derivative of $(D_t \kappa) \circ \mathcal{T}|_{\Gamma_t}$ and some lower-order terms.

Theorem 1. *(Informal version of Theorem 2) Assume that the contact angles $\omega, \omega_r \in (0, \pi/2)$, the integer k and real γ satisfy*

$$2 \leq k \leq \min\left\{\frac{\pi}{\omega}, \frac{\pi}{\omega_r}\right\}, \quad 0 < \gamma + 1 \leq \min\left\{\frac{2\pi}{\omega}, \frac{2\pi}{\omega_r}\right\}.$$

Let a solution to the water-waves system (1)-(5) be given by the free surface graph function ζ and its time derivative $\partial_t \zeta$ with $\alpha^{-1} \partial_{\tilde{x}} \tilde{\zeta} \in H^{k+3/2}(\mathbb{R})$ and $\alpha^{-\gamma-2} \partial_t \tilde{\zeta} - a_{\zeta_t} \in H^{k+2}(\mathbb{R})$ (a_{ζ_t} is the weighted limit of $\alpha^{-\gamma-2} \partial_t \tilde{\zeta}$ near corners, see (34)) and the initial values are chosen from the set of initial values Λ_0 (see Definition 2 in Section 4.3). Moreover, assume that the weighted Taylor sign condition holds initially for $t = 0$, i.e. there exists a positive constant a_0 such that

$$\alpha^{-2\gamma-3} (-\nabla_{n_t} P) \circ \mathcal{T}|_{\Gamma_t} \geq a_0 > 0 \quad \text{when } t = 0.$$

Then there exists $T^* > 0$ depending only on the corresponding initial values such that the energy estimate for the energy $\mathcal{E}(t)$ is closed for $t \in [0, T]$ with $T \leq T^*$.

To understand better the energy and the a priori estimate performed in \mathcal{S}_0 instead of the physical domain Ω , we use the following remark to see what does the conclusion look like if we transform the weighted norms in \mathcal{S}_0 back into corresponding weighted norms in Ω . To see more details, see Remark 12.

Remark 1. According to our theorem, we set that the mean curvature κ of the surface Γ_t (instead of using the surface directly) and its material derivative $D_t \kappa$ satisfy

$$r^{1/2} \kappa \in L^2, \quad r^{-1/2} (r^{-\gamma} D_t \kappa - a_{w_t}) \in L^2 \quad \text{near the corners,}$$

where $\gamma > -1$, r is the radius with respect to to the corner point and a_{w_t} is some weighted limited defined in Proposition 9. This makes Assumption 1 hold. As a result, we have for the velocity v and the pressure P that

$$r^{-1} (r^{-\gamma-2} v - a_v) \in L^2, \quad r^{-1} (r^{-2\gamma-4} P - a_p) \in L^2 \quad \text{near the corners,}$$

where a_v, a_p are corresponding weighted limits of v, P defined in (49), (63). So one can see that if we set some decay near the corners on the free surface, we will have corresponding decay for the velocity and the pressure etc. in a short time interval. Moreover, there is some faster decay for time derivatives. Therefore, it makes sense that we deal with the special case when the contact points and angles are fixed during this time period.

Remark 2. (1) The weighted Taylor's sign holds for $t \in [0, T]$, see Remark 11 for more details; (2) The surface is set to be tangentially horizontal at contact points because we want to simplify the estimates technically. It is possible to generalize this point. Of course it is also possible to consider more general cases with contact points and angles changing with respect to time.

1.2. Organization of the paper. In Section 2 we introduce the conformal mapping \mathcal{T} from the strip domain \mathcal{S}_0 to the corner domain Ω , and lower-order-derivative estimates of \mathcal{T} are proved. In Section 3 we show some weighted elliptic theory in strip domains and deal with some weighted estimates for the D-N operator. The equation of κ is introduced in Section 4, which is then transformed again using the conformal mapping. Moreover, weighted estimates for \mathcal{T}, v, P etc. are proved and the corresponding weighted limits are discussed. In the end, our a priori energy estimate is proved in Section 5.

1.3. Notations. - The surface Γ_t is parameterized by the graph function $z = \zeta(t, x)$, and Γ_b is parameterized by $z = b(x)$ with b smooth enough.

- we denote by n_t, τ_t the unit outward normal and tangential vectors on the free surface Γ_t . n_b, τ_b are defined in a similar way on the bottom Γ_b .

- Π : the second fundamental form where $\Pi(w) = \nabla_w n_t \in T_X \Gamma_t$ (space of tangential vectors) for $w \in T_X \Gamma_t$.

- \mathcal{S}_0 is the strip domain with height ω defined in (6) with the same notations Γ_t, Γ_b for boundaries.

- $u^\top = (u \cdot \tau_t) \tau_t$ for a vector-valued function u defined on Γ_t , and $u^\perp = u \cdot n_t$ on Γ_t .

- Let $\mathcal{H}(f)$ or $f_{\mathcal{H}}$ be the harmonic extension in Ω (or \mathcal{S}_0) for some function f defined on Γ_t , which is defined by the elliptic system

$$\begin{cases} \Delta \mathcal{H}(f) = 0 & \text{in } \Omega \text{ (or } \mathcal{S}_0), \\ \mathcal{H}(f)|_{\Gamma_t} = f, \quad \nabla_{n_b} \mathcal{H}(f)|_{\Gamma_b} = 0. \end{cases}$$

- We denote by

$$\mathcal{N}f = \nabla_{n_t} f \mathcal{H}|_{\Gamma_t}$$

the Dirichlet-Neumann (D-N) operator without the coefficient $\sqrt{1 + |\partial_x \zeta|^2}$.

- The inner product (\cdot, \cdot) stands for the inner product $(\cdot, \cdot)_{L^2(\mathbb{R})}$.

- The norm $\|\cdot\|_{H^s}$ stands for Sobolev norm in Ω or \mathcal{S}_0 , and $|\cdot|_{H^s}$ denotes the norm in \mathbb{R} (or in Γ_t , Γ_b) when no confusion will be made.

- \tilde{f} stands for $f \circ \mathcal{T}$ and is defined in \mathcal{S}_0 or the surface Γ_t of \mathcal{S}_0 , where f is a function in Ω or the surface Γ_t of Ω .

- χ_l, χ_r are some smooth enough cut-off functions near $\tilde{x} = -\infty, +\infty$:

$$\chi_l(\tilde{x}) = \begin{cases} 1, & \tilde{x} \leq -c_0 - 1, \\ 0, & \tilde{x} \geq -c_0, \end{cases} \quad \chi_r(\tilde{x}) = \begin{cases} 0, & \tilde{x} \leq C_0, \\ 1, & \tilde{x} \geq C_0 + 1, \end{cases}$$

where the constants c_0, C_0 are defined later in the weight function.

- $[s]$ stands for the integer part of $s \in \mathbb{R}$.

- We denote by $a_{f,l}(\tilde{z}) = \lim_{\tilde{x} \rightarrow -\infty} \alpha^{-\beta} f$ the weighted limit when $\tilde{x} \rightarrow -\infty$ for a function $f(\tilde{x}, \tilde{z})$, and we denote by $a_{f,r}$ similar weighted limit when $\tilde{x} \rightarrow +\infty$.

- The index i ($i = l, r$) in the estimates stands for the summation of $i = l, r$.

2. PRELIMINARIES

We first present some Sobolev embedding inequalities, product estimates and a trace theorem.

Lemma 1. (1) *There exists a constant C independent of $f \in H^s(\mathbb{R})$ with $s > 1/2$ such that*

$$|f|_{L^\infty(\mathbb{R})} \leq C|f|_{H^s(\mathbb{R})};$$

(2) *Let \mathcal{G} be an open subset of \mathbb{R}^2 with a Lipschitz boundary. Then there exists a constant C independent of $f \in H^s(\mathcal{G})$ with $s > 1$ such that*

$$|f|_{L^\infty(\mathcal{G})} \leq C|f|_{H^s(\mathcal{G})}.$$

Proof. One can check for example Theorem 1.4.4.1 [25] for a complete embedding statement and proof. \square

Lemma 2. (1) *Let $f \in H^1(\mathbb{R})$ and $g \in H^{1/2}(\mathbb{R})$. Then there exists a constant C independent of f, g such that*

$$|fg|_{H^{1/2}(\mathbb{R})} \leq C|f|_{H^1(\mathbb{R})}|g|_{H^{1/2}(\mathbb{R})};$$

(2) *Let $f \in W^{1,\infty}(\mathbb{R})$ and $g \in H^{1/2}(\mathbb{R})$. Then there exists a constant C independent of f, g such that*

$$|fg|_{H^{1/2}(\mathbb{R})} \leq C|f|_{W^{1,\infty}}|g|_{H^{1/2}(\mathbb{R})};$$

(3) *Let a be a function in $W^{1,\infty}(\mathbb{R})$ such that $a \geq a_0$ for some constant $a_0 > 0$ and $f \in H^{1/2}(\mathbb{R})$. Then there exists a constant C depending on $a_0^{-1}, |a|_{W^{1,\infty}}$ such that*

$$|f|_{H^{1/2}(\mathbb{R})} \leq C(a_0^{-1}, |a|_{W^{1,\infty}})|af|_{H^{1/2}(\mathbb{R})}.$$

Proof. For the first estimate, it follows directly from Lemma 2.2 [47] and Lemma 1. Moreover, the second estimate can be proved by a direct computation.

For the last estimate, one sets $\mathcal{L} = 1/a$ and finds that both $\mathcal{L} : L^2 \rightarrow L^2$ and $\mathcal{L} : H^1 \rightarrow H^1$ are bounded. Therefore, an interpolation with the bounds leads to the desired estimate immediately. \square

Lemma 3. (Traces in a strip domain) *The maps*

$$u \mapsto u|_{\Gamma_j}, \quad j = t, b$$

are continuous from $H^s(\mathcal{S}_0)$ onto $H^{s-1/2}(\Gamma_j)$ for $s > 1/2$ and there hold

$$|u|_{H^{s-1/2}(\Gamma_j)} \leq C\|u\|_{H^s(\mathcal{S}_0)}$$

for some constant C depending only on \mathcal{S}_0 and $j = t, b$.

Proof. The proof follows from the trace theorem for the full space \mathbb{R}^2 (Theorem 1.5.1.1 [25] for example) and standard extension and restriction theorems between \mathbb{R}^2 and \mathcal{S}_0 . \square

We are going to introduce a conformal mapping \mathcal{T} defined from the fixed strip domain \mathcal{S}_0 to our physical domain Ω , which plays a key role in our paper. We show that this conformal mapping has very limited regularity near corners, although it can be smooth enough away from the corner.

We define the fixed strip domain

$$(6) \quad \mathcal{S}_0 = \{(\tilde{x}, \tilde{z}) \mid -\omega_* \leq \tilde{z} \leq 0\}$$

so the depth of \mathcal{S}_0 is ω_* . We choose here the depth

$$\omega_* = \omega,$$

i.e. the depth is equal to the left contact angle. Moreover, we still denote by Γ_t, Γ_b the upper and the lower boundaries of \mathcal{S}_0 when no confusion will be made.

We now define the conformal mapping

$$\mathcal{T} : \tilde{Z} = \tilde{x} + i\tilde{z} \in \mathcal{S}_0 \mapsto Z = x + iz \in \Omega,$$

where we choose that \mathcal{T} maps the left infinity of \mathcal{S}_0 to the left corner point p_l of Ω , and maps the right infinity of \mathcal{S}_0 to the right corner point p_r of Ω .

Remark 3. Notice that we have one more freedom of choosing one point on Γ_t to fix the conformal mapping, which is also postponed to Section 4.3 and is not important. We also notice that since Ω depends on time t , \mathcal{T} varies continuously with respect to t .




Figure 2. The conformal mapping

According to the properties of conformal mappings (see [53] for example), we know directly that \mathcal{T} maps the upper (lower) boundary of \mathcal{S}_0 to the upper (lower) boundary of Ω , and \mathcal{T} can be smooth enough away from the corner. When the surface Γ_t is smooth enough, the limitation for the regularity of \mathcal{T} only comes from the part of the domain near the contact point.

In fact, we see in the following lines that *the conformal mapping only have limited boundary regularity due to the existence of the corner, and its regularity is measured by the angles of the corners.*

The following result focuses on the left contact point and is written in a more general case when the left contact angle ω is different from ω_* (recall that it is the height of the strip \mathcal{S}_0), and the boundary is assumed to be piecewise $C^{k,\sigma}$ for the moment, which is the same as in [53]. The detailed boundary dependence in weighted norms and time dependence estimates are discussed later in Section 4.3.

Lemma 4. (*Behaviear near the left corner*) Let

$$\beta = \frac{\omega}{\omega_*}$$

and the domain Ω be given with piecewise $C^{k,\sigma}$ boundary where k is a nonnegative integer and $\sigma \in (0, 1)$. Then for

$$2 \leq k \leq 1 + \pi/\omega,$$

there exist positive constants c_0 and c_1, C_1 depending on Ω, k such that

$$c_1 e^{\beta \tilde{x}} \leq |\mathcal{T}^{(l)}(\tilde{Z})| \leq C_1 e^{\beta \tilde{x}}, \quad \forall \tilde{x} \leq -c_0, \quad l \leq k.$$

Proof. Step 1. Mapping F from \mathcal{K} (a sector) to Ω . Recalling that p_l represents the left corner point of Ω , we define

$$(7) \quad F(\bar{Z}) = \mathcal{T} \circ [\beta^{-1} L n(\bar{Z} - p_l)] \quad \text{for } \bar{Z} \in \mathcal{K}$$

where

$$\mathcal{K} = \{\bar{Z} \mid \bar{Z} = p_l + r e^{i\theta}, r \geq 0, -\omega \leq \theta \leq 0\}.$$

Notice here that \mathcal{K} is the straightened sector of Ω with the same angle ω of the corner. Therefore, we relate the estimate for \mathcal{T} to the estimate for F .

Step 2. Flattening of domains. Letting

$$\alpha_0 = \frac{\omega}{\pi},$$

we define

$$F_1 = Z^{\alpha_0} + p_l : \mathcal{K}_{\alpha_0} \rightarrow \mathcal{K},$$

which maps the half plane

$$\mathcal{K}_{\alpha_0} = \{Z \mid Z = r e^{i\theta}, r \geq 0, -\frac{\omega}{\alpha_0} = -\pi \leq \theta \leq 0\}$$

to the sector \mathcal{K} .

We also define

$$F_2 = (Z - p_l)^{1/\alpha_0} : \Omega \rightarrow \Omega_{\alpha_0}$$

which flattens Ω in a similar way and move the corner point to $(0, 0)$.

We then set the mapping

$$\tilde{F} = F_2 \circ F \circ F_1 : \mathcal{K}_{\alpha_0} \rightarrow \Omega_{\alpha_0},$$

which now maps between domains without corners. Thanks to the assumption of the boundary regularity, we know that boundaries of \mathcal{K}_{α_0} and Ω_{α_0} are smooth enough near $(0, 0)$ ($k \geq 2$ is needed in the following analysis in fact).

Consequently, we know that \tilde{F} is smooth enough near $(0, 0)$ and its derivatives are bounded by Theorem 3.6 [53] (Kellogg-Warschawski Theorem). Besides, direct computations using conformal mappings to the unit disk and applying Theorem 3.7 [53], we have $\text{Arg} \tilde{F}'(0, 0) = 0$, which implies

$$(8) \quad \tilde{F}'(0, 0) = (c, 0) \quad \text{for some } c > 0.$$

Step 3. Estimates of F and \mathcal{T} . First, we expand F near p_l :

$$\begin{aligned} F(\bar{Z}) &= [\tilde{F}((\bar{Z} - p_l)^{1/\alpha_0})]^{\alpha_0} + p_l \\ &= p_l + \tilde{F}'(0, 0)^{\alpha_0} (\bar{Z} - p_l) + \frac{\tilde{F}''(0, 0)}{2\tilde{F}'(0, 0)^{1-\alpha_0}} (\bar{Z} - p_l)^{1+1/\alpha_0} + o((\bar{Z} - p_l)^{1+1/\alpha_0}). \end{aligned}$$

Notice here that the second term doesn't vanish and the third term makes sense thanks to (8).

Therefore, there exist two constants c_1, C_1 depending on a neighborhood of the corner such that

$$c_1 \leq |F'(\bar{Z})| \leq C_1, \quad \text{when } \bar{Z} \rightarrow p_l.$$

Moreover, for k th-order derivative of F satisfying $k < 1 + 1/\alpha_0 = 1 + \pi/\omega$, we always have $F^{(k)}(0, 0) = 0$ and

$$|F^{(k)}(\bar{Z})| \leq C, \quad \text{when } \bar{Z} \rightarrow p_l$$

for some constant C depending on k and a neighborhood of the corner of Ω .

Now we are ready to deal with derivatives of \mathcal{T} . In fact, we obtain thanks to (7) that

$$\mathcal{T}'(\tilde{Z}) = \frac{F'(\bar{Z})}{[\beta^{-1}Ln(\bar{Z} - p_l)]'} = F'(\bar{Z})[e^{\beta\bar{Z}} + p_l]',$$

where

$$\bar{Z} = \bar{Z}(\tilde{Z}) = e^{\beta\tilde{Z}} + p_l.$$

As a result, using the estimates of F , one has the estimate for \mathcal{T}' when $\tilde{x} \leq -c_0$ for some c_0 .

Moreover, for higher-order derivatives of \mathcal{T} , direct computations show that the main part in $\mathcal{T}^{(l)}$ is always

$$F'(\bar{Z})[e^{\beta\bar{Z}} + p_l]^{(l)},$$

so the desired estimates also follow from the estimate for F' . □

A similar result also holds near the right corner and the proof is omitted.

Lemma 5. (*Behavier near the right corner*) Let

$$\beta_r = \frac{\omega_r}{\omega_*}$$

and the domain Ω satisfy the same assumption as in the previous lemma.

Then for

$$2 \leq k \leq 1 + \pi/\omega_r,$$

there exist positive constants C_0 and c_2, C_2 depending on Ω, l such that

$$c_2 e^{\beta_r \tilde{x}} \leq |\mathcal{T}^{(l)}(\tilde{Z})| \leq C_2 e^{\beta_r \tilde{x}}, \quad \forall \tilde{x} \geq C_0, \quad l \leq k.$$

Based on the estimates near the corner, we now deal with the global estimates for \mathcal{T} in \mathcal{S}_0 and take the time parameter into considerations as well. To begin with, we first introduce the following $C^\infty(\mathbb{R})$ weight function

$$\alpha_\beta(\tilde{x}) = \begin{cases} e^{\beta\tilde{x}}, & \tilde{x} \leq -c_0, \\ e^{-\beta_r\tilde{x}}, & \tilde{x} \geq C_0 \end{cases}$$

with β, β_r defined as above. Since the contact angles ω, ω_r are fixed in our settings, one can see that the weight function $\alpha_\beta(\tilde{x})$ doesn't depend on time t .

Proposition 1. (*Global estimates with time parameter*) Assume that the domain Ω and \mathcal{T} are given by Definition 1 in Section 4.3. Then there exist two positive constants c, C depending on the bound L from Definition 2 such that

$$c\alpha(\tilde{x}) \leq |\mathcal{T}'(t, \tilde{Z})| \leq C\alpha(\tilde{x}), \quad \forall \tilde{Z} = (\tilde{x}, \tilde{z}) \in \mathcal{S}_0, \quad \forall t \in [0, T].$$

Moreover, one also has for all $t \in [0, T]$ that

$$|\mathcal{T}^{(l)}(t, \tilde{Z})| \leq C\alpha(\tilde{x}), \quad \forall \tilde{Z} \in \mathcal{S}_0, \quad l \leq k + 1 \leq 1 + \min\left\{\frac{\pi}{\omega}, \frac{\pi}{\omega_r}\right\}$$

with some constant $C > 0$ depending on the initial values and k .

Proof. In fact, we only need to sum up based on Lemma 4 and Lemma 5 and use the bound Definition 2 to control the estimate with respect to time t .

First of all, according to the settings in Definition 1 and Definition 2, the surface Γ_t lies in a weighted $H^{k+5/2}$ Sobolev space and has uniform bound with respect to the time, so it is locally in $H^{k+5/2}$, which means that the boundary of Ω is piecewise $C^{k+1,\sigma}$ for some $\sigma \in (0, 1)$ by the Sobolev embedding theorem. Therefore, applying Lemma 4 and Lemma 5, we know immediately that the constants in the estimates there are all controlled by the bound from Definition 2.

Second, since we already have $\mathcal{T}^{(l)}$ controlled by $\alpha(\tilde{x})$ near ∞ , what is left for us to deal with is the behavior of $\mathcal{T}^{(l)}$ on the bounded domain $\{(\tilde{x}, \tilde{z}) \in \mathcal{S}_0 \mid -c_0 \leq \tilde{x} \leq C_0\}$. Recalling that the conformal mapping can be a smooth-enough diffeomorphism away from corners due to the regularity of boundary, we know immediately that \mathcal{T}' satisfies

$$c_3 \leq |\mathcal{T}'| \leq C_3$$

for some constants c_3, C_3 depending on the initial values. Besides, we also have

$$|\mathcal{T}^{(l)}| \leq C, \quad l \leq k+1$$

for some positive C depending on the initial values and k .

As a result, summing up all these estimates above, we derive the desired estimates. \square

In the end, we specify the following weight function used throughout this paper, which is a special case of α_β . Recall that we choose the height ω_* of the strip \mathcal{S}_0 to be the left contact angle ω , i.e. $\beta = 1$ for the left corner:

$$(9) \quad \alpha(\tilde{x}) = \begin{cases} e^{\tilde{x}}, & \tilde{x} \leq -c_0 \\ e^{-\beta_r \tilde{x}}, & \tilde{x} \geq C_0 \end{cases}$$

with $\beta_r = \omega_r/\omega$.

We now present a natural and useful corollary, which holds since we set the surface and the bottom to be graphs of functions.

Corollary 1. *Let \mathcal{T} satisfy Definition 1 and Definition 2. Then there exist constants c, C depending on the bound in Definition 2 such that*

$$c\alpha(\tilde{x}) \leq |\partial_{\tilde{x}}x| \leq C\alpha(\tilde{x}) \quad \text{on } \Gamma_t, \Gamma_b$$

holds uniformly with respect to $t \in [0, T]$.

Proof. We only prove the case of Γ_t here, and the case of Γ_b follows in a similar way. Since

$$\mathcal{T}'(Z) = (\partial_{\tilde{x}}x, \partial_{\tilde{x}}z) \quad \text{in } \mathcal{S}_0,$$

we have

$$\mathcal{T}'|_{\Gamma_t: \tilde{z}=0} = (\partial_{\tilde{x}}x|_{\tilde{z}=0}, \partial_{\tilde{x}}\tilde{\zeta})$$

where $\tilde{\zeta} = \zeta \circ \mathcal{T}|_{\Gamma_t} = \zeta(t, x(t, \tilde{x}, 0))$ and

$$\partial_{\tilde{x}}\tilde{\zeta} = \partial_{\tilde{x}}x \widetilde{\partial_x \zeta}.$$

As a result, we obtain

$$\mathcal{T}'|_{\Gamma_t} = \partial_{\tilde{x}}x|_{\tilde{z}=0}(1, \widetilde{\partial_x \zeta}),$$

which implies immediately that

$$|\partial_{\tilde{x}}x|_{\tilde{z}=0}| \leq |\mathcal{T}'|_{\Gamma_t}| \leq (1 + |\widetilde{\partial_x \zeta}|_{L^\infty}^2)^{\frac{1}{2}} |\partial_{\tilde{x}}x|_{\tilde{z}=0}|,$$

where $|\widetilde{\partial_x \zeta}|_{L^\infty} = |(\partial_{\tilde{x}}x)^{-1} \partial_{\tilde{x}}\tilde{\zeta}|_{L^\infty}$ is controlled by the upper bound in Definition 2.

Consequently, applying the previous proposition leads to the desired estimate. \square

Remark 4. The estimates above will be used later only for the lower-order $W^{k+1,\infty}$ estimates of \mathcal{T} due to the Sobolev embedding theorem. There will be weighted higher-order estimates for \mathcal{T} in the Section 4.3, which show the dependence on the boundary, especially for the dependence on the surface function ζ .

3. WEIGHTED ELLIPTIC ESTIMATES RELATED TO D-N OPERATOR

We present some weighted space in this section, which will be used later in the energy estimate. The weighted elliptic estimates are proved here for the mixed-boundary system as well as for the Dirichlet system. Moreover, the corresponding estimates for the D-N operator $\tilde{\mathcal{N}}_0$ in the strip domain \mathcal{S}_0 are proved.

3.1. Weighted elliptic estimates on the strip domain. We consider the following mixed-boundary problem

$$(10) \quad \begin{cases} \Delta u = h & \text{in } \mathcal{S}_0, \\ u|_{\Gamma_t} = f, \quad \nabla_{n_b} u|_{\Gamma_b} = g. \end{cases}$$

Before defining the weighted space, we explain the related eigenvalues of (10), needed in following lines. To begin with, we introduce the Laplace transform with respect to \tilde{x}

$$\check{u}(\lambda, \cdot) = (\mathcal{L}u)(\lambda, \cdot) = \int_{\mathbb{R}} e^{-\lambda\tilde{x}} u(\tilde{x}, \cdot) d\tilde{x}, \quad \forall \lambda \in \mathbb{C}.$$

Performing Laplace Transform, system (10) formally becomes

$$\begin{cases} \lambda^2 \check{u} + \partial_{\tilde{z}}^2 \check{u} = \mathcal{L}(h), & \tilde{z} \in [-\omega, 0] \\ \check{u}|_{\tilde{z}=0} = \mathcal{L}(f), \quad -\partial_{\tilde{z}} \check{u}|_{\tilde{z}=-\omega} = \mathcal{L}(g). \end{cases}$$

As a result, the corresponding eigenvalue problem reads

$$\begin{cases} -\phi''(\tilde{z}) - \lambda^2 \phi(\tilde{z}) = 0, & \tilde{z} \in [-\omega, 0] \\ \phi(0) = 0, \quad -\phi'(-\omega) = 0. \end{cases}$$

Moreover, direct computation shows that the eigenvalues are

$$\lambda_m = \frac{(m+1/2)\pi}{\omega}, \quad m \in \mathbb{Z}.$$

Notice that the first positive eigenvalue is $\lambda_0 = \pi/(2\omega)$ when $m = 0$, which will be used later.

Next, we introduce a simple version of weighted norms in \mathcal{S}_0 . For any $\gamma \in \mathbb{R}$ and an integer $l \geq 0$, recall (5.2.1) [38] the following norms

$$\|u\|_{\mathcal{W}_{2,\gamma}^l(\mathcal{S}_0)} = \|e^{\gamma\tilde{x}} u\|_{H^l(\mathcal{S}_0)}$$

and

$$|u|_{\mathcal{W}_{2,\gamma}^{l-1/2}(\Gamma_t)} = |e^{\gamma\tilde{x}} u|_{H^{l-1/2}(\Gamma_t)},$$

where Γ_t can be replaced by Γ_b .

We present a specialized version of Theorem 5.2.2 from page 160 of [38].

Proposition 2. *Suppose that no eigenvalues λ of the problem (10) lie on the line $Re\lambda = -\gamma$. Then this problem is uniquely solvable in $\mathcal{W}_{2,\gamma}^l(\mathcal{S}_0)$ for every given (h, f, g) from the space $\mathcal{W}_{2,\gamma}^{l-2}(\mathcal{S}_0) \times \mathcal{W}_{2,\gamma}^{l-1/2}(\Gamma_t) \times \mathcal{W}_{2,\gamma}^{l-3/2}(\Gamma_b)$ with $l \geq 2$. Furthermore, the solution satisfies the estimate*

$$\|u\|_{\mathcal{W}_{2,\gamma}^l(\mathcal{S}_0)} \leq C(\|h\|_{\mathcal{W}_{2,\gamma}^{l-2}(\mathcal{S}_0)} + |f|_{\mathcal{W}_{2,\gamma}^{l-1/2}(\Gamma_t)} + |g|_{\mathcal{W}_{2,\gamma}^{l-3/2}(\Gamma_b)})$$

with a constant C depending on l, \mathcal{S}_0 .

Remark 5. Checking the proofs of Theorem 3.6.1 and Theorem 5.2.2 in [38] line by line, one knows that the estimate in Proposition 2 can be extended to the case $l = 1$:

$$\|u\|_{\mathcal{W}_{2,\gamma}^1(\mathcal{S}_0)} \leq C(\|h\|_{\mathcal{W}_{2,\gamma}^1(\mathcal{S}_0)^*} + |f|_{\mathcal{W}_{2,\gamma}^{1/2}(\Gamma_t)} + |g|_{\mathcal{W}_{2,\gamma}^0(\Gamma_b)}),$$

where $\mathcal{W}_{2,\gamma}^1(\mathcal{S}_0)^*$ is the dual space of $\mathcal{W}_{2,\gamma}^1(\mathcal{S}_0)$ and the norm for g is not a sharp one but easier to handle.

In the end, we introduce the modified weighted norm used later in our paper. Let

$$\|u\|_{H_\gamma^l} := \|\alpha^\gamma u\|_{H^l(\mathcal{S}_0)}$$

and the norm on the boundary is defined in a similar way. We will use both of the two notations above for the sake of convenience.

Proposition 3. *Let the integer $l \geq 2$ and real γ satisfy*

$$|\gamma| < \min\left\{\frac{\pi}{2\omega}, \frac{\pi}{2\omega_r}\right\}.$$

Suppose that system (10) admits a solution u such that $\alpha^{\pm\gamma}u \in H^l(\mathcal{S}_0)$ with the right-hand side satisfies

$$(\alpha^\gamma h, \alpha^\gamma f, \alpha^\gamma g) \in H^{l-2}(\mathcal{S}_0) \times H^{l-1/2}(\Gamma_t) \times H^{l-3/2}(\Gamma_b).$$

Then the estimate holds

$$\|\alpha^\gamma u\|_{H^l(\mathcal{S}_0)} \leq C(\|\alpha^\gamma h\|_{H^{l-2}(\mathcal{S}_0)} + |\alpha^\gamma f|_{H^{l-1/2}(\Gamma_t)} + |\alpha^\gamma g|_{H^{l-3/2}(\Gamma_b)})$$

with a constant C depending on $l, \mathcal{S}_0, c_0, \gamma$.

Remark 6. In fact, the unique existence of the solution can also be obtained. We refer to Theorem 2.6.4 [39] where this weighted norm in \mathcal{S}_0 is transformed back to the so-called homogeneous weighted norm in the physical domain Ω . The restriction of the power γ comes directly from the existence part, which takes the symmetric interval with respect to the smallest positive eigenvalue λ_m with $m = 0$ defined above. Therefore, we only focus on the weighted estimate here.

The following lemma is about the unique solvability and estimate of system (10) in H^1 variational formulation.

Lemma 6. *Let the right-hand side of system (10) satisfy*

$$(h, f, g) \in H^{-1}(\mathcal{S}_0) \times H^{1/2}(\Gamma_t) \times H^{-1/2}(\Gamma_b).$$

Then the system admits a unique solution $u \in H^1(\mathcal{S}_0)$ with the estimate

$$\|u\|_{H^1(\mathcal{S}_0)} \leq C(\|h\|_{H^{-1}(\mathcal{S}_0)} + |f|_{H^{1/2}(\Gamma_t)} + |g|_{H^{-1/2}(\Gamma_b)})$$

where the constant C depends on \mathcal{S}_0 .

Proof. This can be proved in a standard way by applying Lax-Milgram theorem, see [40] for example. \square

Proof. (Proof of Proposition 3). In fact, the proof is similar to the analysis in Section 2.6.1 [39], which is proved in Ω with an equivalent weighted norm in Ω . The idea is to start with the existence and estimate in standard variational formula H^1 norm in Ω , and then apply Corollary 6.3.3 [38] to show that the elliptic operator is an isomorphism for $|\gamma| \leq \min\{\pi/(2\omega), \pi/(2\omega_r)\}$, i.e. to stay in the interval bounded by the first positive eigenvalue. In order to be self contained, we provide a direct proof adjust to our case in the following lines.

We first prove the case where the weight is $\alpha^{-\gamma}$ ($\gamma > 0$), which is an easier part. Based on the estimates with weight $\alpha^{-\gamma}$, we deal with the case with weight α^γ by a dual argument. To begin with, we first introduce some localizations to reduce the estimates to the case in Proposition 2.

Step 1. Localization and estimates near ∞ . We only focus on the left infinity, and the proof for the right infinity follows in a similar way.

Recall that χ_l is a $C^\infty(\mathbb{R})$ cut-off function, and let $\bar{\chi}_l(\tilde{x})$ be a similar cut-off function satisfying

$$\chi_l \bar{\chi}_l = \bar{\chi}_l.$$

Defining

$$u_l = \chi_l u, \quad u_r = \chi_r u, \quad u_R = (1 - \chi_l - \chi_r)u,$$

we have immediately

$$u = u_l + u_r + u_R$$

and

$$\alpha^{-\gamma} u_l = \chi_l \alpha^{-\gamma} u = \chi_l e^{-\gamma \tilde{x}} u.$$

As a result, we know that

$$\|\alpha^{-\gamma} u_l\|_{H^l(\mathcal{S}_0)} = \|\chi_l e^{-\gamma \tilde{x}} u\|_{H^l(\mathcal{S}_0)} = \|u_l\|_{\mathcal{W}_{2,-\gamma}^l(\mathcal{S}_0)},$$

and u_l satisfies the system

$$\begin{cases} \Delta u_l = \chi_l h + [\Delta, \chi_l]u := h_l & \text{in } \mathcal{S}_0, \\ u_l|_{\Gamma_t} = \chi_l f := f_l, \quad \nabla_{n_b} u_l|_{\Gamma_b} = \chi_l g + (\nabla_{n_b} \chi_l)u|_{\Gamma_b} := g_l. \end{cases}$$

Applying Proposition 2, we get

$$\|u_l\|_{\mathcal{W}_{2,-\gamma}^l(\mathcal{S}_0)} \leq C(\|h_l\|_{\mathcal{W}_{2,-\gamma}^{l-2}(\mathcal{S}_0)} + |f_l|_{\mathcal{W}_{2,-\gamma}^{l-1/2}(\Gamma_t)} + |g_l|_{\mathcal{W}_{2,-\gamma}^{l-3/2}(\Gamma_b)})$$

where $l \geq 2$ and $C = C(\mathcal{S}_0)$. The right-side terms will be handled one by one. In fact, for h_l term, one has

$$\begin{aligned} \|h_l\|_{\mathcal{W}_{2,-\gamma}^{l-2}(\mathcal{S}_0)} &\leq \|\chi_l h\|_{\mathcal{W}_{2,-\gamma}^{l-2}(\mathcal{S}_0)} + \|[\Delta, \chi_l]u\|_{\mathcal{W}_{2,-\gamma}^{l-2}(\mathcal{S}_0)} \\ &\leq \|\chi_l h\|_{\mathcal{W}_{2,-\gamma}^{l-2}(\mathcal{S}_0)} + \|e^{-\gamma \tilde{x}} [\Delta, \chi_l]u\|_{H^{l-2}(\mathcal{S}_0)} \\ &\leq \|\chi_l h\|_{\mathcal{W}_{2,-\gamma}^{l-2}(\mathcal{S}_0)} + C(c_0, \gamma, l) \|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^{l-1}(\mathcal{S}_0)} \end{aligned}$$

where $\bar{\chi}_r$ is defined in a similar way as $\bar{\chi}_l$.

For g_l term, one has similarly

$$\begin{aligned} |g_l|_{\mathcal{W}_{2,-\gamma}^{l-3/2}(\Gamma_b)} &\leq |\chi_l g|_{\mathcal{W}_{2,-\gamma}^{l-3/2}(\Gamma_b)} + |(\nabla_{n_b} \chi_l)u|_{\mathcal{W}_{2,-\gamma}^{l-2}(\Gamma_b)} \\ &\leq |\chi_l g|_{\mathcal{W}_{2,-\gamma}^{l-3/2}(\Gamma_b)} + C(c_0, \gamma, l) \|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^{l-3/2}(\Gamma_b)}. \end{aligned}$$

Summing up these estimates above, we obtain

(11)

$$\|u_l\|_{\mathcal{W}_{2,-\gamma}^l(\mathcal{S}_0)} \leq C \left(\|\chi_l h\|_{\mathcal{W}_{2,-\gamma}^{l-2}(\mathcal{S}_0)} + |\chi_l f|_{\mathcal{W}_{2,-\gamma}^{l-1/2}(\Gamma_t)} + |\chi_l g|_{\mathcal{W}_{2,-\gamma}^{l-3/2}(\Gamma_b)} + \|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^{l-1}(\mathcal{S}_0)} \right),$$

where the constant $C = C(c_0, \gamma, l, \mathcal{S}_0)$. Moreover, similar estimate can be proved for u_r where $-\gamma$ is replaced by $\gamma\beta_r$.

Step 2. Estimates for u_R . Recalling that $u_R = (1 - \chi_l - \chi_r)u$, we have directly

$$\|\alpha^{-\gamma} u_R\|_{H^l(\mathcal{S}_0)} \leq C \|u_R\|_{H^l(\mathcal{S}_0)}$$

with $C = C(c_0, \gamma, l)$ and u_R satisfies the system

$$\begin{cases} \Delta u_R = (1 - \chi_l - \chi_r)h - [\Delta, \chi_l + \chi_r]u & \text{in } \mathcal{S}_0, \\ u_R|_{\Gamma_t} = (1 - \chi_l - \chi_r)f, \quad \nabla_{n_b} u_R|_{\Gamma_b} = (1 - \chi_l - \chi_r)g - (\nabla_{n_b} \chi_l + \nabla_{n_b} \chi_r)u|_{\Gamma_b}. \end{cases}$$

Standard elliptic estimates in strip domains (see [40]) and similar arguments as for u_l case lead to

$$(12) \quad \begin{aligned} \|u_R\|_{H^l(\mathcal{S}_0)} &\leq C \left(\|(1 - \chi_l - \chi_r)h\|_{H^{l-2}(\mathcal{S}_0)} + |(1 - \chi_l - \chi_r)f|_{H^{l-1/2}(\Gamma_t)} \right. \\ &\quad \left. + |(1 - \chi_l - \chi_r)g|_{H^{l-3/2}(\Gamma_b)} + \|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^{l-1}(\mathcal{S}_0)} \right), \end{aligned}$$

where the constant $C = C(c_0, \gamma, l, \mathcal{S}_0)$.

Summing up (11) and (12), we obtain

$$\begin{aligned} \|\alpha^{-\gamma}u\|_{H^1(\mathcal{S}_0)} &\leq C\left(\|\alpha^{-\gamma}h\|_{H^{l-2}(\mathcal{S}_0)} + |\alpha^{-\gamma}f|_{H^{l-1/2}(\Gamma_t)} + |\alpha^{-\gamma}g|_{H^{l-3/2}(\Gamma_b)}\right. \\ &\quad \left. + \|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^{l-1}(\mathcal{S}_0)}\right). \end{aligned}$$

To finish the proof, we need to deal with the lower-order term on the right side.

Step 3. Lower-order estimate. In fact, a standard interpolation in Sobolev spaces leads to

$$\|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^{l-1}(\mathcal{S}_0)} \leq \epsilon\|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^l(\mathcal{S}_0)} + C_\epsilon\|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{L^2(\mathcal{S}_0)}$$

where $C_\epsilon = C(1/\epsilon, \mathcal{S}_0)$.

For the first term on the right side, thanks to the definition of $\bar{\chi}_l$, one has directly that

$$\|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^l(\mathcal{S}_0)} \leq C(c_0, \gamma, l)\|\alpha^{-\gamma}u\|_{H^l(\mathcal{S}_0)}.$$

On the other hand, applying Lemma 6 and turning Sobolev norms into the weighted norms lead to

$$\begin{aligned} \|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{L^2(\mathcal{S}_0)} &\leq C\|u\|_{L^2(\mathcal{S}_0)} \\ &\leq C(\|h\|_{H^{-1}(\mathcal{S}_0)} + |f|_{H^{1/2}(\Gamma_t)} + |g|_{H^{-1/2}(\Gamma_b)}) \\ &\leq C(\|\alpha^{-\gamma}h\|_{H^{l-2}(\mathcal{S}_0)} + |\alpha^{-\gamma}f|_{H^{l-1/2}(\Gamma_t)} + |\alpha^{-\gamma}g|_{H^{l-3/2}(\Gamma_b)}) \end{aligned}$$

since $\gamma > 0$.

As a result, we conclude that

$$\begin{aligned} \|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^{l-1}(\mathcal{S}_0)} &\leq \epsilon C\|\alpha^{-\gamma}u\|_{H^l(\mathcal{S}_0)} + C(\|\alpha^{-\gamma}h\|_{H^{l-2}(\mathcal{S}_0)} + |\alpha^{-\gamma}f|_{H^{l-1/2}(\Gamma_t)} \\ &\quad + |\alpha^{-\gamma}g|_{H^{l-3/2}(\Gamma_b)}), \end{aligned}$$

which together with Step 2 finishes the proof for the case with weight $\alpha^{-\gamma}$.

Step 4. Estimate for the case α^γ ($\gamma > 0$). To prove this case, we first need to investigate again the variational formula for system (10) with homogeneous boundary conditions and modify the H^1 estimate to obtain the weighted $H_{-\gamma}^1$ estimate. Second, a dual argument is applied here to obtain the estimate for H_γ^1 case. Third, the inhomogeneous-boundary case follows in a natural way by a trace theorem.

To begin with, we consider system (10) with homogeneous boundary conditions:

$$(13) \quad \begin{cases} \Delta u = h & \text{in } \mathcal{S}_0, \\ u|_{\Gamma_t} = 0, \quad \nabla_{n_b} u|_{\Gamma_b} = 0 \end{cases}$$

and we modify the variation formula for $u \in H_0^1(\mathcal{S}_0) = \{v \in H^1(\mathcal{S}_0) \mid v|_{\Gamma_t} = 0, \nabla_{n_b} v|_{\Gamma_b} = 0\}$ (See Section 2.6.1 [39]) now with the right side $h \in (H_\gamma^1)^*$ (the dual space of H_γ^1):

$$\int_{\mathcal{S}_0} \nabla u \cdot \nabla v d\tilde{x}d\tilde{z} = \langle h, v \rangle,$$

where $v \in H_0^1(\mathcal{S}_0)$ and the right side is the dual product on H_γ^1 .

One knows immediately that

$$\langle h, v \rangle \leq \|h\|_{(H_\gamma^1)^*} \|v\|_{H_\gamma^1} \leq C\|h\|_{(H_\gamma^1)^*} \|v\|_{H^1}$$

where C depends on the weight function α .

Consequently, we obtain

$$\|u\|_{H^1} \leq C\|h\|_{(H_\gamma^1)^*}.$$

On the other hand, repeating the proof in Step 1 and Step 2 with $l = 1$, $f, g = 0$ and using Remark 5, we find

$$\|\alpha^{-\gamma}u\|_{H^1} \leq C\left(\|\chi_l h\|_{(\mathcal{W}_{2,\gamma}^1)^*} + \|\chi_r h\|_{(\mathcal{W}_{2,\gamma\beta_r}^1)^*} + \|(1 - \chi_l - \chi_r)h\|_{H^{-1}} + \|(1 - \bar{\chi}_l - \bar{\chi}_r)u\|_{H^1}\right).$$

Moreover, direct proof shows that

$$\|\chi_l h\|_{(\mathcal{W}_{2,\gamma}^1)^*} + \|\chi_r h\|_{(\mathcal{W}_{2,\gamma\beta_r}^1)^*} + \|(1 - \chi_l - \chi_r)h\|_{H^{-1}} \leq C\|h\|_{(H_\gamma^1)^*}$$

with the constant C depending on χ_i, α .

Summing up these estimates above, we conclude that

$$(14) \quad \|u\|_{H_{-\gamma}^1} \leq C \|h\|_{(H_\gamma^1)^*},$$

which shows that the elliptic operator \mathcal{A} for system (13) is an isomorphism from $H_{-\gamma}^1$ to $(H_\gamma^1)^*$ with homogeneous boundary conditions and is consistent with Section 2.6.1 [39].

Next, we consider the adjoint operator \mathcal{A}^* and the elliptic operator \mathcal{A}^+ associated to the formally adjoint problem of (13):

$$\begin{cases} \Delta v = h & \text{in } \mathcal{S}_0, \\ v|_{\Gamma_t} = 0, & -\nabla_{n_b} v|_{\Gamma_b} = 0, \end{cases}$$

where we can see that \mathcal{A}^+ is in fact the same as \mathcal{A} .

We know from Lemma 6.3.4 and (6.3.23) in [38] that $v \in H_{-\gamma}^1$ is a solution of $\mathcal{A}^+v = (h, 0, 0)$ for a given $h \in (H_\gamma^1)^*$ if and only if $\mathcal{A}^*v = h$. As a result, this implies for all $u \in H_\gamma^1$ that

$$\langle h, u \rangle = \langle \mathcal{A}^*v, u \rangle = \langle v, \mathcal{A}u \rangle \leq \|v\|_{H_{-\gamma}^1} \|\mathcal{A}u\|_{(H_{-\gamma}^1)^*}.$$

Besides, we have by (14) that

$$\|v\|_{H_{-\gamma}^1} \leq C \|\mathcal{A}^+v\|_{(H_\gamma^1)^*} = C \|h\|_{(H_\gamma^1)^*},$$

which leads to

$$\langle h, u \rangle \leq C \|h\|_{(H_\gamma^1)^*} \|\mathcal{A}u\|_{(H_{-\gamma}^1)^*}.$$

Therefore, we finally obtain the following variational weighted estimate with weight α^γ :

$$\|u\|_{H_\gamma^1} \leq C \|\mathcal{A}u\|_{(H_{-\gamma}^1)^*}$$

i.e. \mathcal{A} is also an isomorphism from H_γ^1 to $(H_{-\gamma}^1)^*$. Moreover, $\|\mathcal{A}u\|_{(H_{-\gamma}^1)^*}$ can be controlled directly by $\|\alpha^\gamma h\|_{L^2(\mathcal{S}_0)}$.

To close the proof, repeating the proof from Step 1 to Step 3 and using a trace theorem, the weighted estimate for higher-order case can be easily finished. \square

In the end, we also mention the following Dirichlet system:

$$(15) \quad \begin{cases} \Delta u = h & \text{in } \mathcal{S}_0, \\ u|_{\Gamma_t} = f, & u|_{\Gamma_b} = g. \end{cases}$$

A similar weighted estimate holds for this system and the proof is omitted here. Moreover, we also refer to Theorem 3.11 [11] for the corresponding weighted estimate in Ω .

Proposition 4. *Let the integer $l \geq 1$ and the real γ satisfy*

$$|\gamma| < \min\left\{\frac{\pi}{\omega}, \frac{\pi}{\omega_r}\right\}.$$

Suppose that the problem (15) admits a solution u such that $\alpha^\gamma u \in H^l(\mathcal{S}_0)$ with the right-hand side given:

$$(\alpha^\gamma h, \alpha^\gamma f, \alpha^\gamma g) \in H^{l-2}(\mathcal{S}_0) \times H^{l-\frac{1}{2}}(\Gamma_t) \times H^{l-\frac{1}{2}}(\Gamma_b).$$

Then the estimate holds

$$\|\alpha^\gamma u\|_{H^l(\mathcal{S}_0)} \leq C (\|\alpha^\gamma h\|_{H^{l-2}(\mathcal{S}_0)} + |\alpha^\gamma f|_{H^{l-\frac{1}{2}}(\Gamma_t)} + |\alpha^\gamma g|_{H^{l-\frac{1}{2}}(\Gamma_b)})$$

with a constant C depending on $l, \mathcal{S}_0, c_0, \gamma$.

Remark 7. The range of the weight is given in Theorem 3.11 [11] and Theorem 2.6.1 [39], which is also decided by the first eigenvalue $\lambda = \pi/\omega$ for the Dirichlet system.

3.2. Weighted estimates for D-N operator. We first recall some basic properties for the D-N operator $\tilde{\mathcal{N}}_0$ in the strip domain \mathcal{S}_0 from Proposition 3.8 [40]. The Sobolev norm $|\cdot|_{H^s}$ always represents the norm on Γ_t from now on when no confusion will be made.

Lemma 7. *Let f, g be any two functions in Schwartz class $\mathcal{S}(\mathbb{R})$. Then*

(1) *The operator $\tilde{\mathcal{N}}_0$ is self-adjoint:*

$$(\tilde{\mathcal{N}}_0 f, g) = (f, \tilde{\mathcal{N}}_0 g)$$

where (\cdot, \cdot) is the $L^2(\mathbb{R})$ inner product;

(2) *The operator $\tilde{\mathcal{N}}_0$ is positive:*

$$(\tilde{\mathcal{N}}_0 f, f) \geq 0;$$

(3) *There exists a constant C independent of f, g such that the following estimate holds:*

$$|(\tilde{\mathcal{N}}_0 f, g)| \leq C |f|_{H^{1/2}} |g|_{H^{1/2}}.$$

Moreover, there exist some positive μ and c independent of f such that

$$(\tilde{\mathcal{N}}_0 f, f) + \mu |f|_{L^2}^2 \geq c |f|_{H^{1/2}}^2.$$

We now consider weighted estimates involving the D-N operator.

Proposition 5. *Let the integer $l \geq 1$, γ satisfy*

$$|\gamma| < \min\left\{\frac{\pi}{2\omega}, \frac{\pi}{2\omega_r}\right\},$$

and the weight $\alpha = \alpha(\tilde{x})$ be defined by (9). Then the following statements hold true:

(1) *For any f such that $\alpha^\gamma f \in H^{3/2}(\Gamma_t)$, one has*

$$|\alpha^\gamma \tilde{\mathcal{N}}_0 f|_{H^{1/2}} \leq C |\alpha^\gamma f|_{H^{3/2}}$$

with a constant $C = C(c_0, \gamma, \mathcal{S}_0)$.

(2) *For any f such that $\alpha^\gamma f \in H^{l+1}(\Gamma_t)$, one has*

$$|\alpha^\gamma \tilde{\mathcal{N}}_0 f|_{H^l} \leq C |\alpha^\gamma f|_{H^{l+1}}$$

with a constant $C = C(l, c_0, \gamma, \mathcal{S}_0)$.

Proof. (1) The proof is straightforward. In fact, one writes

$$|\alpha^\gamma \tilde{\mathcal{N}}_0 f|_{H^{1/2}} = |\partial_{\tilde{z}}(\alpha^\gamma f_{\mathcal{H}})|_{H^{1/2}},$$

which together with Proposition 3 and Lemma 3 imply immediately that

$$|\alpha^\gamma \tilde{\mathcal{N}}_0 f|_{H^{1/2}} \leq C \|\alpha^\gamma f_{\mathcal{H}}\|_{H^2} \leq C |\alpha^\gamma f|_{H^{3/2}}.$$

(2) Higher-order estimate. Similar arguments lead to the desired conclusion. \square

Moreover, we also need some commutator estimates related to the D-N operator.

Proposition 6. *Let real $\gamma, \gamma_1, \gamma_2$ satisfy*

$$\gamma = \gamma_1 + \gamma_2, \quad |\gamma|, |\gamma_1|, |\gamma_2| < \min\left\{\frac{\pi}{2\omega}, \frac{\pi}{2\omega_r}\right\}$$

and the weight $\alpha = \alpha(\tilde{x})$ be defined by (9). Then the following statements hold true:

(1) *For any two functions f, g such that $\alpha^{\gamma_1} f \in H^2(\Gamma_t)$, $\alpha^{\gamma_2} g \in H^{1/2}(\Gamma_t)$, one has*

$$|\alpha^\gamma [f, \tilde{\mathcal{N}}_0] g|_{L^2} \leq C |\alpha^{\gamma_1} f|_{H^2} |\alpha^{\gamma_2} g|_{H^{1/2}}$$

with some constant $C = C(c_0, \gamma, \mathcal{S}_0)$.

Moreover, when $\alpha^{\gamma_1} f \in W^{2, \infty}$, one has

$$|\alpha^\gamma [f, \tilde{\mathcal{N}}_0] g|_{L^2} \leq C |\alpha^{\gamma_1} f|_{W^{2, \infty}} |\alpha^{\gamma_2} g|_{H^{1/2}}$$

(2) (Higher-order estimate) Let $l \in \mathbb{N}$. For any two functions f, g such that $\alpha^{\gamma_1} f \in H^{l+1}(\Gamma_t)$, $\alpha^{\gamma_2} g \in H^l(\Gamma_t)$, one has

$$|\alpha^\gamma [f, \tilde{\mathcal{N}}_0]g|_{H^l} \leq C |\alpha^{\gamma_1} f|_{H^{l+1}} |\alpha^{\gamma_2} g|_{H^l}$$

with some constant $C = C(l, c_0, \gamma, \mathcal{S}_0)$.

(3) There exists a constant $C = C(c_0, \gamma, \mathcal{S}_0)$ such that

$$|\alpha^{\gamma_1} [\tilde{\mathcal{N}}_0, \alpha^{\gamma_2}]f|_{H^{1/2}} \leq C |\alpha^\gamma f|_{H^{1/2}}.$$

Proof. (1) Recalling that $f_{\mathcal{H}}$ denotes the harmonic extension of f in \mathcal{S}_0 , we start with rewriting the commutator on Γ_t :

$$[f, \tilde{\mathcal{N}}_0]g = f \nabla_{n_t} g_{\mathcal{H}} - \nabla_{n_t} (fg)_{\mathcal{H}} = -(\nabla_{n_t} f_{\mathcal{H}})g - \nabla_{n_t} (f_{\mathcal{H}}g_{\mathcal{H}} - (fg)_{\mathcal{H}}).$$

As a result, denoting by $w = f_{\mathcal{H}}g_{\mathcal{H}} - (fg)_{\mathcal{H}}$, we obtain

$$(16) \quad |\alpha^\gamma [f, \tilde{\mathcal{N}}_0]g|_{L^2} \leq |\alpha^\gamma (\nabla_{n_t} f_{\mathcal{H}})g|_{L^2} + |\alpha^\gamma \nabla_{n_t} w|_{L^2},$$

where w satisfies the system

$$\begin{cases} \Delta w = 2\nabla f_{\mathcal{H}} \cdot \nabla g_{\mathcal{H}} & \text{in } \mathcal{S}_0, \\ w|_{\Gamma_t} = 0, \quad \nabla_{n_b} w|_{\Gamma_b} = 0. \end{cases}$$

For the first term on the right side of (16), we have directly that

$$|\alpha^\gamma (\nabla_{n_t} f_{\mathcal{H}})g|_{L^2} \leq |\alpha^{\gamma_1} \nabla_{n_t} f_{\mathcal{H}}|_{L^\infty} |\alpha^{\gamma_2} g|_{L^2}.$$

Noticing as before that

$$\alpha^\gamma \nabla_{n_t} f_{\mathcal{H}}|_{\Gamma_t} = \partial_{\tilde{z}}(\alpha^\gamma f_{\mathcal{H}})|_{\Gamma_t}$$

and applying Lemma 3 and Proposition 3 on it, we find

$$\begin{aligned} |\alpha^\gamma (\nabla_{n_t} f_{\mathcal{H}})g|_{L^2} &\leq C \|\alpha^{\gamma_1} f_{\mathcal{H}}\|_{H^{5/2}} |\alpha^{\gamma_2} g|_{L^2} \\ &\leq C |\alpha^{\gamma_1} f|_{H^2} |\alpha^{\gamma_2} g|_{L^2}. \end{aligned}$$

For the second term $|\alpha^\gamma \nabla_{n_t} w|_{L^2}$ in (16), we have immediately by Proposition 3 that

$$\begin{aligned} |\alpha^\gamma \nabla_{n_t} w|_{L^2(\Gamma_t)} &\leq |\alpha^\gamma \nabla_{n_t} w|_{H^{1/2}(\Gamma_t)} \leq C \|\alpha^\gamma w\|_{H^2(\mathcal{S}_0)} \\ &\leq C \|\alpha^\gamma \nabla f_{\mathcal{H}} \cdot \nabla g_{\mathcal{H}}\|_{L^2(\mathcal{S}_0)} \\ &\leq C \|\alpha^{\gamma_1} \nabla f_{\mathcal{H}}\|_{L^\infty(\mathcal{S}_0)} \|\alpha^{\gamma_2} \nabla g_{\mathcal{H}}\|_{L^2(\mathcal{S}_0)}. \end{aligned}$$

Using Lemma 1, Lemma 3 and Proposition 3 again, one has

$$\|\alpha^{\gamma_1} \nabla f_{\mathcal{H}}\|_{L^\infty(\mathcal{S}_0)} \leq C |\alpha^{\gamma_1} f|_{H^2(\Gamma_t)}.$$

Moreover, applying Remark 5 to g leads to

$$\|\alpha^{\gamma_2} \nabla g_{\mathcal{H}}\|_{L^2(\mathcal{S}_0)} \leq C \|\alpha^{\gamma_2} g_{\mathcal{H}}\|_{H^1(\mathcal{S}_0)} \leq C |\alpha^{\gamma_2} g|_{H^{1/2}(\Gamma_t)}.$$

Summing up these estimates above, the proof for the first estimate in (1) is finished. For the second one, repeating the proof above while replace $f_{\mathcal{H}}$ by f (constant with respect to \tilde{z}), one can derive the desired estimate, where the second-order derivative of f is reached in the estimate of w .

(2) Higher-order derivative estimates. One has immediate the following estimate in a similar way as above:

$$|\alpha^\gamma [f, \tilde{\mathcal{N}}_0]g|_{H^1} \leq |\alpha^\gamma (\nabla_{n_t} f_{\mathcal{H}})g|_{H^1} + |\alpha^\gamma \nabla_{n_t} w|_{H^1},$$

and the desired estimate follows naturally.

(3) Direct computations lead to

$$[\tilde{\mathcal{N}}_0, \alpha^\gamma]f = \partial_{\tilde{z}} \mathcal{H}(\alpha^\gamma f)|_{\Gamma_t} - \alpha^\gamma \partial_{\tilde{z}} f_{\mathcal{H}}|_{\Gamma_t} = \partial_{\tilde{z}} w_1|_{\Gamma_t},$$

where

$$w_1 = \mathcal{H}(\alpha^\gamma f) - \alpha^\gamma f_{\mathcal{H}}$$

satisfies

$$\begin{cases} \Delta w_1 = -2\partial_{\bar{x}}\alpha^\gamma\partial_{\bar{x}}f_{\mathcal{H}} - \partial_{\bar{x}}^2\alpha^\gamma f_{\mathcal{H}} & \text{in } \mathcal{S}_0, \\ w_1|_{\Gamma_t} = 0, \quad \nabla_{n_b} w_1|_{\Gamma_b} = 0. \end{cases}$$

As a result, one has by Lemma 3, Proposition 3 and Remark 5 that

$$\begin{aligned} |\alpha^{\gamma_1}[\tilde{\mathcal{N}}_0, \alpha^{\gamma_2}]f|_{L^2} &= |\alpha^{\gamma_1}\partial_{\bar{z}}w_1|_{L^2} \leq C\|\alpha^{\gamma_1}w_1\|_{H^2} \\ &\leq C(\|\alpha^{\gamma_1}\partial_{\bar{x}}\alpha^{\gamma_2}\partial_{\bar{x}}f_{\mathcal{H}}\|_{L^2} + \|\alpha^{\gamma_1}\partial_{\bar{x}}^2\alpha^{\gamma_2}f_{\mathcal{H}}\|_{L^2}) \leq C|\alpha^\gamma f|_{H^{1/2}}. \end{aligned}$$

Notice that here the derivatives of α are controlled by $C\alpha$ thanks to the definition of α . Therefore, the proof is finished. \square

4. REFORMULATION OF THE PROBLEM AND PREPARATIONS

Similarly as in previous works [46, 47, 48], we use again the geometric formulation in [57, 58]. We start with recalling and rewriting the equation of the mean curvature κ of the surface in the physical domain Ω , and then transform it into an equivalent equation in the strip domain \mathcal{S}_0 .

4.1. Geometric formulation in the strip domain. To begin with, the mean curvature of Γ_t is expressed by

$$\kappa = \nabla_{\tau_t} n_t \cdot \tau_t.$$

Remembering that the free surface Γ_t is expressed as $z = \zeta(t, x)$ explicitly, we know immediately that κ is parameterized as

$$\kappa = \partial_x \left(\frac{\partial_x \zeta}{\sqrt{1 + |\partial_x \zeta|^2}} \right).$$

In the following we use both the geometric expression and the parameterization of κ for the sake of convenience, and there is a slight abuse of the notation $\Gamma_t = \mathbb{R} \times \{0\}$ of \mathcal{S}_0 when we use the parameterization $z = \zeta$ and consider $\tilde{\zeta}$ in \mathbb{R} indeed.

We also recall some computations from [57, 46]:

$$(17) \quad D_t n_t = -((\nabla v)^T n_t)^\top = -(\nabla_{\tau_t} v \cdot n_t) \tau_t, \quad D_t \tau_t = (\nabla_{\tau_t} v \cdot n_t) n_t \quad \text{on } \Gamma_t.$$

Next, taking the surface tension coefficient $\sigma = 0$ in [57], we find that κ satisfies

$$(18) \quad D_t \kappa = -\Delta_{\Gamma_t} v \cdot n_t - 2\nabla_{\tau_t} n_t \cdot \nabla_{\tau_t} v \quad \text{on } \Gamma_t.$$

Taking D_t on this equation again, one obtains immediately as in [57] the following equation:

$$(19) \quad D_t^2 \kappa = -n_t \cdot \Delta_{\Gamma_t} D_t v + R_1 = n_t \cdot \Delta_{\Gamma_t} (\nabla P) + R_1,$$

where

$$\begin{aligned} R_1 = & 2 \left[\tau_t \nabla_{\tau_t} ((\nabla v)^T n_t)^\top \cdot \tau_t + \nabla_{(\nabla_{\tau_t} v)^\top} n_t \right] \cdot \nabla_{\tau_t} v + \Delta_{\Gamma_t} v \cdot ((\nabla v)^T n_t)^\top + 2\nabla_{\tau_t} n_t \cdot (\nabla v)^2 \tau_t \\ & - 2(\nabla_{\tau_t} v \cdot n_t) \left(\nabla_{\tau_t} n_t \cdot \nabla_{n_t} v + \nabla_{\tau_t} n_t \cdot (\nabla v)^T n_t \right) - 2n_t \cdot D^2 v (\tau_t, (\nabla_{\tau_t} v)^\top) \\ & - 2(\nabla_{n_t} v \cdot n_t) (\nabla_{\tau_t} n_t \cdot \nabla_{\tau_t} v) + n_t \cdot \nabla v ((\Delta_{\Gamma_t} v)^\top) - \kappa |((\nabla v)^T n_t)^\top|^2 + 2\nabla_{\tau_t} n_t \cdot \nabla_{\tau_t} \nabla P. \end{aligned}$$

Moreover, direct computations show that the pressure P satisfies the following system

$$(20) \quad \begin{cases} \Delta P = -tr(\nabla v)^2 & \text{in } \Omega, \\ P|_{\Gamma_t} = 0, \quad \nabla_{n_b} P|_{\Gamma_b} = \nabla_v n_b \cdot v. \end{cases}$$

Similarly as in [57], we rewrite the term $n_t \cdot \Delta_{\Gamma_t} \nabla P$ in (19):

$$n_t \cdot \Delta_{\Gamma_t} \nabla P = -\mathbf{a} \mathcal{N} \kappa + R_2,$$

where

$$\mathbf{a} = -\nabla_{n_t} P$$

is the Taylor's sign and \mathcal{N} is the Dirichlet-Neumann operator. Here R_2 is derived by direct computations related to Δ_{Γ_t} :

$$\begin{aligned}
(21) \quad R_2 &= -n_t \cdot \Delta \nabla P + \kappa n_t \cdot \nabla_{n_t} \nabla P + n_t \cdot D^2(\nabla P)(n_t, n_t) \\
&= -n_t \cdot \nabla(\text{tr}(Dv)^2) + \nabla_{n_t}(\kappa \mathcal{H} \nabla_{n_t} P) - \nabla_{n_t} \kappa \mathcal{H} \nabla_{\bar{n}_t} P - \kappa \nabla_{n_t} \bar{n}_t \cdot \nabla P + \nabla_{n_t}(D^2 P(\bar{n}_t, \bar{n}_t)) \\
&\quad - 2D^2 P(n_t, \nabla_{n_t} \bar{n}_t) \\
&= -n_t \cdot \nabla(\text{tr}(Dv)^2) - \nabla_{n_t} u_p + \kappa \nabla P \cdot \nabla_{n_t} \bar{n}_t + 2D^2 P(n_t, \nabla_{n_t} \bar{n}_t)
\end{aligned}$$

where

$$u_p := \kappa \mathcal{H} \nabla_{\bar{n}_t} P + D^2 P(\bar{n}_t, \bar{n}_t),$$

and \bar{n}_t is an extension of n_t in Ω . For simplicity, we just take

$$\bar{n}_t = n_t \quad \text{in } \Omega$$

since n_t relies on the parametrization $z = \zeta(t, x)$ and depends only on t, x .

Moreover, we need to emphasize that $\nabla_{n_t} u_p$ is in fact a higher-order term about the same order as $\nabla^2 v$. In fact, u_p satisfies

$$(22) \quad \begin{cases} \Delta u_p = 2\nabla \kappa \mathcal{H} \cdot \nabla(\nabla_{n_t} P) + \kappa \mathcal{H} \Delta(\nabla_{n_t} P) + \Delta(D^2 P(n_t, n_t)) & \text{in } \Omega \\ u_p|_{\Gamma_t} = \kappa \nabla_{n_t} P + D^2 P(n_t, n_t) = \Delta P - \Delta_{\Gamma_t} P = -\text{tr}(Dv)^2|_{\Gamma_t}, \\ \nabla_{n_b} u_p|_{\Gamma_b} = \kappa \mathcal{H} \nabla_{n_b}(\nabla_{n_t} P) + \nabla_{n_b} D^2 P(n_t, n_t) + 2D^2 P(\nabla_{n_b} n_t, n_t)|_{\Gamma_b} \end{cases}$$

where one notes the term $\partial^3 P$ on Γ_b .

The boundary dependence of the corresponding elliptic system of P usually leads to a loss of half-order derivative of the boundary, but we will show that in Corollary 4 that this still works since the boundary dependence in this case only lies in the bottom, which can be as smooth as we want.

Substituting these computations into equation (19), we find

$$(23) \quad D_t^2 \kappa + \mathbf{a} \mathcal{N} \kappa = R_1 + R_2 \quad \text{on } \Gamma_t.$$

Now we plan to transform (23) into an equivalent equation in \mathcal{S}_0 by the conformal mapping \mathcal{T} defined before. We always use the notation

$$\tilde{u} = u \circ \mathcal{T}$$

in the following lines, where u is some function defined in Ω .

To begin with, for space derivatives in Ω , we have the following transform

$$(24) \quad \widetilde{\nabla} u = (\nabla u) \circ \mathcal{T} = (\nabla_X \tilde{X}) \circ \mathcal{T} \nabla \tilde{u}$$

where

$$X = (x, z) \in \Omega, \quad \tilde{X} = (\tilde{x}, \tilde{z}) \in \mathcal{S}_0$$

and

$$(25) \quad (\nabla_X \tilde{X}) \circ \mathcal{T} = |\mathcal{T}'|^{-2} \begin{pmatrix} \partial_{\tilde{x}} x & \partial_{\tilde{x}} z \\ \partial_{\tilde{x}} z & -\partial_{\tilde{x}} x \end{pmatrix}, \quad |\mathcal{T}'| = |(\partial_{\tilde{x}} x, \partial_{\tilde{x}} z)| = \sqrt{(\partial_{\tilde{x}} x)^2 + (\partial_{\tilde{x}} z)^2}.$$

Meanwhile, taking $\tilde{z} = 0$ in $z = z(t, \tilde{x}, \tilde{z})$, we obtain directly

$$z = z(t, \tilde{x}, 0) = \zeta(t, x(t, \tilde{x}, 0)) := \tilde{\zeta}(t, \tilde{x}),$$

which is usually denoted by

$$\tilde{\zeta} = \zeta \circ \mathcal{T}|_{\Gamma_t}$$

with a slight abuse of notation. Consequently, we have

$$(26) \quad |\mathcal{T}'|_{\Gamma_t} = |\partial_{\tilde{x}} x| \sqrt{1 + |(\partial_{\tilde{x}} x)^{-1} \partial_{\tilde{x}} \tilde{\zeta}|^2}$$

and

$$(27) \quad \begin{aligned} (\nabla_X \tilde{X}) \circ \mathcal{T}|_{\Gamma_t} &= |\mathcal{T}'|^{-2} \begin{pmatrix} \partial_{\tilde{x}} x & \partial_{\tilde{x}} \tilde{\zeta} \\ \partial_{\tilde{x}} \tilde{\zeta} & -\partial_{\tilde{x}} x \end{pmatrix} \\ &= (\partial_{\tilde{x}} x)^{-1} \frac{1}{1 + |(\partial_{\tilde{x}} x)^{-1} \partial_{\tilde{x}} \tilde{\zeta}|^2} \begin{pmatrix} 1 & (\partial_{\tilde{x}} x)^{-1} \partial_{\tilde{x}} \tilde{\zeta} \\ (\partial_{\tilde{x}} x)^{-1} \partial_{\tilde{x}} \tilde{\zeta} & -1 \end{pmatrix}. \end{aligned}$$

Second, we transform the D-N operator \mathcal{N} in Ω into the D-N operator $\tilde{\mathcal{N}}_0$ in \mathcal{S}_0 , which is well-understood with estimates already. In fact, direct computations show that for a function f defined on Γ_t , the harmonic extension $f_{\mathcal{H}}$ satisfies

$$\begin{cases} \Delta f_{\mathcal{H}} = 0 & \text{in } \mathcal{S}_0, \\ f_{\mathcal{H}}|_{\Gamma_t} = \tilde{f}, \quad \nabla_{n_b} f_{\mathcal{H}}|_{\Gamma_b} = 0. \end{cases}$$

As a result, we obtain

$$(28) \quad (\mathcal{N}f) \circ \mathcal{T} = (\nabla_{n_t} f_{\mathcal{H}}|_{\Gamma_t}) \circ \mathcal{T} = |\mathcal{T}'|^{-1}|_{\Gamma_t} \tilde{\mathcal{N}}_0 \tilde{f}$$

where $\tilde{\mathcal{N}}_0$ is the D-N operator in \mathcal{S}_0 .

On the other hand, we also need to consider the transformation of $D_t u$ for a smooth function u defined in Ω . Taking $\tilde{u} = u \circ \mathcal{T}$ in \mathcal{S}_0 as above, direct computations show that

$$(D_t u) \circ \mathcal{T} = (\partial_t u) \circ \mathcal{T} + (v \cdot \nabla u) \circ \mathcal{T} = (\partial_t u) \circ \mathcal{T} + \tilde{v} \cdot (\nabla_X \tilde{X}) \circ \mathcal{T} \nabla \tilde{u},$$

where

$$(\partial_t u) \circ \mathcal{T} = \partial_t \tilde{u} - \tilde{v}_{\mathcal{T}0} \cdot \tilde{\nabla} u = \partial_t \tilde{u} - \tilde{v}_{\mathcal{T}0} \cdot (\nabla_X \tilde{X}) \circ \mathcal{T} \nabla \tilde{u}$$

with

$$\tilde{v}_{\mathcal{T}0} = (\partial_t x, \partial_t z)^T.$$

Consequently, denoting by

$$(29) \quad \tilde{v}_{\mathcal{T}} = (\nabla_X \tilde{X})^T \circ \mathcal{T} (\tilde{v} - \tilde{v}_{\mathcal{T}0}),$$

we conclude the following formula

$$(D_t u) \circ \mathcal{T} = \partial_t \tilde{u} + \tilde{v}_{\mathcal{T}} \cdot \nabla \tilde{u} := \tilde{D}_t \tilde{u} \quad \text{in } \mathcal{S}_0.$$

Moreover, when we look at a function $f = f(t, x)$ defined by the parametrization of the surface, i.e.

$$f(t, x) = F(t, x, z)|_{\Gamma_t: z=\zeta(t, x)}$$

for some function F , we have

$$\underline{D}_t f = (D_t F)|_{z=\zeta} \quad \text{with} \quad \underline{D}_t = \partial_t + \underline{v}_1 \partial_x$$

where

$$v = (v_1, v_2)^T, \quad \underline{v}_1 = v_1|_{z=\zeta}.$$

To simplify the notations, we also use D_t instead of \underline{D}_t when no confusion will be made.

Similar computations as above show that for

$$\tilde{f} = f \circ \mathcal{T}|_{\tilde{z}=0} = f(t, x(t, \tilde{x}, 0))$$

we have

$$(30) \quad (D_t f) \circ \mathcal{T}|_{\tilde{z}=0} = \partial_t \tilde{f} + u_1 \partial_{\tilde{x}} \tilde{f} := \tilde{D}_t \tilde{f}$$

with

$$u_1 = (\partial_{\tilde{x}} x)^{-1} (\tilde{v}_1 - \partial_t x), \quad \tilde{v}_1 = \underline{v}_1 \circ \mathcal{T}|_{\tilde{z}=0}.$$

We will use the same notation \tilde{D}_t as in the strip domain \mathcal{S}_0 when no confusion will be made in the following text.

Based on the preparations above, we are ready to rewrite (23) in the strip domain \mathcal{S}_0 . First of all, we denote by

$$w := \tilde{\kappa}.$$

Second, using the parameterizing $z = \zeta$ on Γ_t , compositing $\mathcal{T}|_{\Gamma_t: \tilde{z}=0}$ on both sides of (23) and applying (30), we derive

$$(31) \quad \tilde{D}_t^2 w + \tilde{\mathfrak{a}} |\mathcal{T}'|^{-1} \tilde{\mathcal{N}}_0 w = \tilde{R}_1 + \tilde{R}_2$$

where

$$\tilde{R}_i = R_i \circ \mathcal{T}, \quad i = 1, 2.$$

Notice that here we omit the notation $\cdot|_{\Gamma_t}$ for $|\mathcal{T}'|^{-1}$ when no confusion will be made.

4.2. The proper weights. Since a natural weight $|\mathcal{T}'|^{-1}$ appears in (31), we plan to identify the weights for all the quantities. To begin with, recalling the weight function $\alpha(\tilde{x})$ from (9) and setting

$$\gamma \in \mathbb{R} \quad \text{to be fixed later,}$$

$\tilde{D}_t w$ is assumed to be equipped with the weight $\alpha^{-\gamma}$ and we try to use the weighted norm $|\alpha^{-\gamma} \tilde{D}_t w|_{H^k}$.

Remembering that $|\mathcal{T}'|^{-1} \approx \alpha^{-1}$ thanks to Proposition 1, it seems that we should put the weight $\alpha^{-\gamma-\frac{1}{2}}$ before w while notice that there is the Taylor's sign $\tilde{\mathfrak{a}}$ with w .

Besides, recalling (24) shows that for every space derivative in Ω , there is approximately one α^{-1} together with one space derivative in \mathcal{S}_0 . Meanwhile, looking at (26) and (27), we find a natural weight α^{-1} in front of $\partial_{\tilde{x}} \tilde{\zeta} = \partial_{\tilde{x}x} \tilde{\partial}_x \tilde{\zeta}$, which suggests that the weight for $\partial_{\tilde{x}} \tilde{\zeta}$ seems to be α^{-1} .

To have some more clues about weights, we look at the system for $\tilde{P} = P \circ \mathcal{T}$. In fact, \tilde{P} satisfies

$$(32) \quad \begin{cases} \Delta \tilde{P} = -|\mathcal{T}'|^2 \text{tr}((\nabla_X \tilde{X}) \circ \mathcal{T} \nabla \tilde{v})^2 & \text{in } \mathcal{S}_0, \\ \tilde{P}|_{\Gamma_t} = 0, \quad \partial_{\tilde{z}} \tilde{P}|_{\Gamma_b} = |\mathcal{T}'|((\nabla_X \tilde{X})^T \circ \mathcal{T}) \tilde{v} \cdot \nabla \tilde{n}_b \cdot \tilde{v}. \end{cases}$$

Proposition 3 suggests that the terms on the right side of the above system decide the weight of \tilde{P} . Besides, we have

$$|\mathcal{T}'| \approx \alpha, \quad (\nabla_X \tilde{X}) \circ \mathcal{T}|_{\Gamma_t} \approx \alpha^{-1}$$

thanks to Proposition 1, and notice that n_b is assumed to be constant near the corners, which implies immediately that the boundary term $\partial_{\tilde{z}} \tilde{P}|_{\Gamma_b}$ vanishes near $\tilde{x} = \infty$. Consequently, the weight for \tilde{P} should be determined by \tilde{v} -terms on the right side.

Now we are ready to identify all the weights. According to the assumption that $\tilde{D}_t w$ carries the weight $\alpha^{-\gamma}$ and the fact $w \approx \alpha^{-2} \partial^2 \tilde{\zeta}$, we find the weight $\alpha^{-\gamma-2}$ for $\tilde{D}_t \tilde{\zeta}$.

On the other hand, direct computations and applying the kinematic condition (4) show that

$$(33) \quad \begin{aligned} \partial_t \tilde{\zeta} &= \tilde{\partial}_t \tilde{\zeta} + \partial_t x (\partial_{\tilde{x}x})^{-1} \partial_{\tilde{x}} \tilde{\zeta} \\ &= \tilde{v} \cdot \left(-(\partial_{\tilde{x}x})^{-1} \partial_{\tilde{x}} \tilde{\zeta}, 1 \right)^T + \partial_t x (\partial_{\tilde{x}x})^{-1} \partial_{\tilde{x}} \tilde{\zeta}. \end{aligned}$$

As a result, checking (30), (33), (32) and counting the orders of derivatives one by one suggest the following weights:

$$\mathcal{T}' : \alpha^{-1}, \quad \tilde{\partial}_t \tilde{\zeta} : \alpha^{-\gamma-2}, \quad \tilde{v} : \alpha^{-\gamma-2}, \quad \tilde{P} : \alpha^{-2\gamma-4}.$$

The weight for $\partial_t \mathcal{T}$ will be determined later. Moreover, the weight for

$$\tilde{\mathfrak{a}} = -\widetilde{\nabla_{n_t} P} = -|\mathcal{T}'|^{-1} \partial_{\tilde{z}} \tilde{P}|_{\tilde{z}=0}$$

should be $\alpha^{-2\gamma-3}$, which implies that

$$\tilde{\mathfrak{a}}^{1/2} \text{ carries the weight } \alpha^{-\gamma-3/2}.$$

We will assume later that $\tilde{\mathfrak{a}}$ is positive with the weight (the weighted Taylor's sign condition), which together with the system of \tilde{P} suggests that $\alpha^{-2\gamma-4} \tilde{P}$ doesn't go to zero when \tilde{x} goes to ∞ . Consequently, we will have to investigate the weighted limits for \tilde{P} and other quantities as well.

Meanwhile, we assume that

$$w = \tilde{\kappa} \quad \text{carries the weight } \alpha^q,$$

so the weight for $\tilde{\zeta}$ should be α^{q+2} (we use $\partial_{\tilde{x}}\tilde{\zeta}$ indeed).

In the end, the left side of equation (31) suggests that $\tilde{\mathbf{a}}^{1/2}w$ should carry the weight $\alpha^{-\gamma-1/2}$, which means

$$-\gamma - 3/2 + q = -\gamma - 1/2, \quad \text{i.e. } q = 1.$$

Consequently, we have the following weights:

$$w : \alpha, \quad \tilde{\zeta} \quad \text{or } \partial_{\tilde{x}}\tilde{\zeta} : \alpha^{-1},$$

which coincide with our observation before.

4.3. Preparations for the energy estimate. Before introducing the energy, we define the set of solutions which involves the surface function ζ , the conformal mapping \mathcal{T} , the velocity v and the pressure P .

To begin with, we consider the weighted limits of $\tilde{\partial}_t\zeta$, \mathcal{T}' , \tilde{P} , $\partial_{\tilde{z}}\tilde{\Phi}$ which will be used later. We define

$$(34) \quad a_{\zeta_t}(t, \tilde{x}) = \chi_l(\tilde{x})a_{\zeta_t,l}(t) + \chi_r(\tilde{x})a_{\zeta_t,r}(t)$$

where χ_l, χ_r are the cut-off functions near $\mp\infty$, and $a_{\zeta_t,i}$ ($i = l, r$) are the weighted limits of $\tilde{\partial}_t\zeta$:

$$a_{\zeta_t,l} = \lim_{x \rightarrow -\infty} \alpha^{-\gamma-2} \tilde{\partial}_t\zeta, \quad a_{\zeta_t,r} = \lim_{x \rightarrow +\infty} \alpha^{-\gamma-2} \tilde{\partial}_t\zeta.$$

Therefore, a_{ζ_t} is well-defined as long as $\alpha^{-\gamma-2}\tilde{\partial}_t\zeta \in L^\infty(\mathcal{S}_0)$ for each t .

Moreover, we set $T_{1,c}$, a_p and a_{Φ_z} as the corresponding weighted limits for \mathcal{T}' , \tilde{P} and $\partial_{\tilde{z}}\tilde{\Phi}$ defined in (39), (63) and Proposition 10 respectively, which can be well-defined in a similar way and turn out to depend on a_{ζ_t} .

We now are ready to define the set of solutions to (31).

Definition 1. Let $T > 0$, $k \in \mathbb{N}$ and Q_Λ be some positive-coefficient polynomial. Let $\Lambda = \Lambda(T, k)$ be the collection of $(\zeta, \mathcal{T}, \Phi, P)$ satisfying the water-waves system (1)-(5) (where \mathcal{T} is defined by ζ) and the following conditions:

(i) $(\zeta, \mathcal{T}, \Phi, P)$ lies in the following spaces

$$\begin{aligned} \alpha^{-1}\partial_{\tilde{x}}\tilde{\zeta} &\in L_T^\infty(H^{k+3/2}(\mathbb{R})), \quad \alpha^{-\gamma-2}\tilde{\partial}_t\zeta - a_{\zeta_t} \in L_T^\infty(H^{k+2}(\mathbb{R})), \quad \alpha^{-1}\mathcal{T}' - T_{1,c} \in L_T^\infty(H^{k+2}(\mathcal{S}_0)), \\ \alpha^{-\gamma-3}\nabla\tilde{\Phi} - (a_{\Phi_x}, a_{\Phi_z})^T &\in L_T^\infty(H^{k+5/2}(\mathcal{S}_0)), \quad \alpha^{-2\gamma-4}\tilde{P} - a_p \in L_T^\infty(H^{k+7/2}(\mathcal{S}_0)), \end{aligned}$$

where a_{ζ_t} , $T_{1,c}$, a_p and a_{Φ_x}, a_{Φ_z} are well defined such that

$$a_{\zeta_t} \in L^\infty([0, T]), \quad T_{1,c}, a_p, a_{\Phi_x}, a_{\Phi_z} \in L_T^\infty(L^\infty([-\omega, 0]));$$

(ii) $(\zeta, \mathcal{T}, \Phi, P)$ is bounded uniformly for $t \in [0, T]$:

$$\begin{aligned} &|\alpha w|_{H^{k+1/2}}^2 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}^2 + \|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{L^2}^2 + |\alpha^{-\gamma-2}\tilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2}^2 \\ &+ \|\alpha^{-\gamma-3}\nabla\tilde{\Phi} - (a_{\Phi_x}, a_{\Phi_z})^T\|_{L^2}^2 + \|\alpha^{-2\gamma-4}\tilde{P} - a_p\|_{L^2}^2 + |a_{\zeta_t,i}|^2 + |b|^2 + |b_r|^2 + |b_{r2}|^2 \leq Q_0 \end{aligned}$$

where a_{w_t} is well defined in Proposition 9 and

$$\begin{aligned} Q_0 = &Q_\Lambda(|\alpha w(0)|_{H^{k+1/2}}, |\alpha^{-\gamma}\tilde{D}_t w(0) - a_{w_t}(0)|_{H^k}, \|\alpha^{-1}\mathcal{T}'(0) - T_{1,c}(0)\|_{L^2}, |\alpha^{-\gamma-2}\tilde{\partial}_t\zeta(0) - a_{\zeta_t}(0)|_{L^2}, \\ &\|\alpha^{-\gamma-3}\nabla\tilde{\Phi}(0) - (a_{\Phi_x}, a_{\Phi_z})^T(0)\|_{L^2}, \|\alpha^{-2\gamma-4}\tilde{P}(0) - a_p(0)\|_{L^2}, |a_{\zeta_t,i}(0)|, |b(0)|, |b_r(0)|, |b_{r2}(0)|). \end{aligned}$$

Moreover, we define the set for the initial values.

Definition 2. Let $L > 0$ be some fixed constant. Λ_0 is the collection of the initial values of $(\zeta, \mathcal{T}, \Phi, P)$ satisfying the water-waves system (1)-(5) such that

$$\begin{aligned} & |\alpha w(0)|_{H^{k+1/2}}, |\alpha^{-\gamma} \widetilde{D}_t w(0) - a_{w_t}(0)|_{H^k} \leq L, \\ & \|\alpha^{-1} \mathcal{T}(0) - T_{1,c}(0)\|_{L^2}, |\alpha^{-\gamma-2} \widetilde{\partial}_t \zeta(0) - a_{\zeta_t}(0)|_{L^2}, \|\alpha^{-\gamma-3} \nabla \widetilde{\Phi}(0) - (a_{\Phi_x}, a_{\Phi_z})^T(0)\|_{L^2} \leq L, \\ & \|\alpha^{-2\gamma-4} \widetilde{P}(0) - a_p(0)\|_{L^2}, |a_{\zeta_t, i}(0)|, |b(0)|, |b_r(0)|, |b_{r2}(0)| \leq L. \end{aligned}$$

Remark 8. According to the weighted norms for $\widetilde{\zeta}$ and $\widetilde{\partial}_t \widetilde{\zeta}$ in these definitions above, we see directly that Assumption 1 holds for $t \in [0, T]$.

Remark 9. For each t , the third point in Γ_t of \mathcal{S}_0 to fix \mathcal{T} is always chosen such that

$$\lim_{\widetilde{x} \rightarrow -\infty} \alpha^{-1} \partial_{\widetilde{x}} x|_{\Gamma_t: \widetilde{z}=0} = 1.$$

In fact, this choice is not important of course and simplifies a little bit the first condition in (36).

Since the higher-order energy will be based on the curvature w (related to the surface function $\widetilde{\zeta}$), we need to investigate estimates for all the quantities like $\widetilde{v}, \widetilde{P}$ in (31) and express them by w . Meanwhile, we notice that the higher-order space derivatives and the time derivative of \mathcal{T} are involved a lot. As a result, the estimates in our paper will be proved under the choice

$$(\zeta, \mathcal{T}, \Phi, P) \in \Lambda(T, k)$$

with their initial values chosen from Λ_0 and

$$k \leq \min\left\{\frac{\pi}{\omega}, \frac{\pi}{\omega_r}\right\}$$

comes from Proposition 1.

Moreover, we need the following weighted Taylor's sign assumption in our energy estimate.

Assumption 2. There exists a positive constant a_0 such that

$$\alpha^{-2\gamma-3} \widetilde{\mathbf{a}} \geq a_0 > 0 \quad \text{on } \Gamma_t : \widetilde{z} = 0, \quad \text{for } t \in [0, T].$$

Notations:

- (1) We denote by $Q(\cdot)$ some polynomial with positive constant coefficients depending on $k, \gamma, \omega, \omega_r$ and the bound L from Definition 2;
- (2) $Q(\mathcal{T}', \nabla \widetilde{\Phi}, \widetilde{P}, w, s)$ stands for a polynomial $Q(\cdot)$ depending on $|\alpha w|_{H^s}$ and the corresponding weighted L^2 norms of $\mathcal{T}', \nabla \widetilde{\Phi}, \widetilde{P}$ as well as the L^∞ norms of their left and right weighted limits (In fact, all these limits turn out to depend on $T_{1,c}$ and a_{ζ_t} defined in (34) and (39)). The other notations $Q(\cdot, w, s)$ are defined in a similar way;
- (3) Q_b is a polynomial $Q(|b|_{L^\infty}, |b_r|_{L^\infty}, |b_{r2}|_{L^\infty})$;
- (4) C is a positive constant depending on $k, \omega, \omega_r, \gamma$ and the bound L from Definition 2;
- (5) We will denote by $|\cdot|$ the norm with respect to \mathbb{R} (for Γ_t or Γ_b) and by $\|\cdot\|$ the norm with respect to \mathcal{S}_0 when no confusion will be made.

4.4. Weighted estimates for the conformal mapping \mathcal{T} . As the first step, we show detailed boundary-dependence estimates for higher-order space derivatives of \mathcal{T} . To begin with, we investigate the behavior of \mathcal{T}' at infinity, which is already showed to some extent by Proposition 1. Corresponding to the left corner, we denote by

$$(35) \quad a_{x,l}(t, \widetilde{z}) = \lim_{\widetilde{x} \rightarrow -\infty} \alpha^{-1} \partial_{\widetilde{x}} x, \quad a_{z,l}(t, \widetilde{z}) = \lim_{\widetilde{x} \rightarrow -\infty} \alpha^{-1} \partial_{\widetilde{x}} z, \quad \forall \widetilde{z} \in [-\omega, 0]$$

where we know from Proposition 1, Remark 4, Definition 1, Definition 2 and Remark 9 that $a_{x,l}, a_{z,l}$ vary continuously with respect to z with the following end-point values:

$$(36) \quad \begin{cases} a_{x,l}(t, 0) = 1, & a_{x,l}(t, -\omega) = b(t) \\ a_{z,l}(t, 0) = \lim_{\widetilde{x} \rightarrow -\infty} \widetilde{\partial}_x \zeta = 0, & a_{z,l}(t, -\omega) = \lim_{\widetilde{x} \rightarrow -\infty} \widetilde{\partial}_x b = -k_b b(t) \end{cases}$$

where $b(t)$ is some function unknown and to be estimated later.

Similarly, we define for the right corner the following limits:

$$(37) \quad a_{x,r}(t, \tilde{z}) = \lim_{\tilde{x} \rightarrow +\infty} \alpha^{-1} \partial_{\tilde{x}} x, \quad a_{z,r}(t, \tilde{z}) = \lim_{\tilde{x} \rightarrow +\infty} \alpha^{-1} \partial_{\tilde{x}} z, \quad \forall \tilde{z} \in [-\omega, 0]$$

with the values

$$(38) \quad \begin{cases} a_{x,r}(t, 0) = b_r(t), & a_{x,r}(t, -\omega) = b_{r2}(t) \\ a_{z,r}(t, 0) = \lim_{\tilde{x} \rightarrow +\infty} \partial_{\tilde{x}} \zeta = 0, & a_{z,r}(t, -\omega) = \lim_{\tilde{x} \rightarrow +\infty} \partial_{\tilde{x}} \tilde{b} = -k_{b,r} b_{r2}(t). \end{cases}$$

Moreover, we denote by

$$a_x = \chi_l(\tilde{x}) a_{x,l}(t, \tilde{z}) + \chi_r(\tilde{x}) a_{x,r}(t, \tilde{z}), \quad a_z = \chi_l(\tilde{x}) a_{z,l}(t, \tilde{z}) + \chi_r(\tilde{x}) a_{z,r}(t, \tilde{z})$$

where the cut-off functions χ_i are already defined before.

To consider higher-order space derivatives for \mathcal{T} , we denote by

$$(39) \quad T_{1,c} = \chi_l T_{1,l} + \chi_r T_{1,r}$$

where

$$T_{1,l} = (a_{x,l}, a_{z,l}), \quad T_{1,r} = (a_{x,r}, a_{z,r}).$$

We plan to use the norm $\|\alpha^{-1} \mathcal{T}' - T_{1,c}\|_{H^k}$, or equivalently the norms $\|\alpha^{-1} \partial_{\tilde{x}} x - a_x\|_{H^k}$, $\|\alpha^{-1} \partial_{\tilde{x}} z - a_z\|_{H^k}$.

To understand this norm, we need some more computations. In fact, when $\alpha^{-1} \partial_{\tilde{x}} z - a_z \in H^k(\mathcal{S}_0)$ for some integer $k \geq 0$, we have

$$\begin{aligned} \partial_{\tilde{x}}(\alpha^{-1} \partial_{\tilde{x}} z - a_z) &= -\alpha^{-2} \alpha' \partial_{\tilde{x}} z + \alpha^{-1} \partial_{\tilde{x}}^2 z + \chi'_l a_{z,l} + \chi'_r a_{z,r} \\ &= -(\alpha^{-2} \alpha' \partial_{\tilde{x}} z - \chi_{l,1} a_{z,l} + \beta_r \chi_{r,1} a_{z,r}) + (\alpha^{-1} \partial_{\tilde{x}}^2 z - \chi_{l,1} a_{z,l} + \beta_r \chi_{r,1} a_{z,r}) \\ &\quad + \chi'_l a_{z,l} + \chi'_r a_{z,r} \end{aligned}$$

where $\chi_{i,1}$ are cut-off functions similar as χ_i with $\text{supp} \chi_{i,1}$ slightly smaller than $\text{supp} \chi_i$.

As a result, this together with $\alpha^{-1} \partial_{\tilde{x}} z - a_z \in L^2(\mathcal{S}_0)$ implies

$$\alpha^{-1} \partial_{\tilde{x}}^2 z - a_{z,1} \in L^2(\mathcal{S}_0)$$

with

$$a_{z,1} = \chi_{l,1} a_{z,l} - \beta_r \chi_{r,1} a_{z,r},$$

and the following estimate holds

$$\begin{aligned} \|\alpha^{-1} \partial_{\tilde{x}}^2 z - a_{z,1}\|_{L^2} &\leq \|\partial_{\tilde{x}}(\alpha^{-1} \partial_{\tilde{x}} z - a_z)\|_{L^2} + \|\alpha^{-2} \alpha' \partial_{\tilde{x}} z - a_{z,1}\|_{L^2} + \|\chi'_{c,i} a_{z,i}\|_{L^2} \\ &\leq C(\|\alpha^{-1} \partial_{\tilde{x}} z - a_z\|_{H^1} + |a_{z,i}|_{L^\infty}). \end{aligned}$$

Here we denote by $|a_{z,i}|_{L^\infty}$ the summation of $|a_{z,i}|_{L^\infty}$ on $i = l, r$.

Therefore, we also know by an induction that

$$\alpha^{-1} \partial_{\tilde{x}}^m z - a_{z,m} \in L^2(\mathcal{S}_0), \quad 1 \leq m \leq k+1 \quad \text{when} \quad \alpha^{-1} \partial_{\tilde{x}} z - a_z \in H^k(\mathcal{S}_0)$$

with the notation

$$a_{z,m} = \chi_{l,m} a_{z,l} + (-\beta_r)^m \chi_{r,m} a_{z,r}$$

and $\chi_{i,m}$ are cut-off functions similar as χ_i with smaller supports than $\chi_{i,m-1}$. Moreover, $\|\alpha^{-1} \partial_{\tilde{x}}^m z - a_{z,m}\|_{L^2}$ can be controlled by $\|\alpha^{-1} \partial_{\tilde{x}} z - a_z\|_{H^k}$ and $|a_{z,i}|_{L^\infty}$.

Consequently, we summarize these computations in the following lemma with a slightly more general form.

Lemma 8. *Let u be a smooth enough function in \mathcal{S}_0 and*

$$a_u(\tilde{x}, \tilde{z}) = \chi_l a_{u,l} + \chi_r a_{u,r} = \chi_l \lim_{\tilde{x} \rightarrow -\infty} \alpha^{-\gamma} u + \chi_r \lim_{\tilde{x} \rightarrow +\infty} \alpha^{-\gamma} u.$$

Assume that $\alpha^{-\gamma} u - a_u \in H^k(\mathcal{S}_0)$ with γ real.

Then we have for any integer m satisfying $1 \leq m \leq k$ that

$$\alpha^{-\gamma} \partial_{\tilde{x}}^m u - a_{u,m} \in L^2(\mathcal{S}_0)$$

with

$$a_{u,m} = \gamma^m (\chi_{l,m} a_{u,l} + (-\beta_r)^m \chi_{r,m} a_{u,r})$$

and the estimate holds

$$\|\alpha^{-\gamma} \partial_{\bar{x}}^l u - a_{u,m}\|_{L^2} \leq C (\|\alpha^{-\gamma} u - a_u\|_{H^k} + |a_u|_{L^\infty})$$

with the constant C depending on $k, \mathcal{S}_0, \chi_i, \alpha, \beta_r$.

Focusing on \mathcal{T} again and applying C-R equations and Lemma 8, we know immediately that $a_{x,i}$ and $a_{z,i}$ are related and we only write the case $i = l$:

$$\begin{aligned} \partial_{\bar{z}} a_{z,l} &= \lim_{\bar{x} \rightarrow -\infty} \alpha^{-1} \partial_{\bar{x}} \partial_{\bar{z}} z = \lim_{\bar{x} \rightarrow -\infty} \alpha^{-1} \partial_{\bar{x}}^2 x = a_{x,l}, \\ \partial_{\bar{z}} a_{x,l} &= \lim_{\bar{x} \rightarrow -\infty} \alpha^{-1} \partial_{\bar{x}} \partial_{\bar{z}} x = - \lim_{\bar{x} \rightarrow -\infty} \alpha^{-1} \partial_{\bar{x}}^2 z = -a_{z,l}, \end{aligned}$$

where we need $x, z \in C^2(\mathcal{S}_0)$, which is satisfied in Definition 1 already.

Solving these equations above with boundary conditions (36), we obtain

$$(40) \quad a_{x,l} = \cos \tilde{z} + \frac{\cos \omega - b(t)}{\sin \omega} \sin \tilde{z}, \quad a_{z,l} = \frac{k_b b(t)}{\sin \omega} \sin \tilde{z}$$

where k_b is the slope of Γ_b in Ω near the left corner. The expressions for $a_{x,r}$ and $a_{z,r}$ are similar. In fact, these expressions are very natural. When we focus on the left corner of Ω which approximates a sector, the conformal mapping \mathcal{T} is simply like $e^{\tilde{z}}$.

Now we are ready to deal with higher-order space derivatives of \mathcal{T} .

Proposition 7. *Let $T_{1,c}$ be given by (39). Then for any integer $k \geq 1$, there holds*

$$\|\alpha^{-1} \mathcal{T}' - T_{1,c}\|_{H^{k+2}} \leq Q(\mathcal{T}', w, k + 1/2)$$

Moreover, we have

$$|T_{1,c}| \leq C (|b(t)| + |b_r(t)| + |b_{r2}(t)|).$$

Proof. Step 1. We first explore some relationship between \mathcal{T} and the surface curvature κ , which is inspired by Shatah-Walsh-Zeng[56]. According to the definition of \mathcal{T} , the curvature can be expressed as

$$\tilde{\kappa} = \kappa \circ \mathcal{T} = \frac{\partial_{\bar{x}} x \partial_x^2 z - \partial_x^2 x \partial_{\bar{x}} z}{|\mathcal{T}'|^3} \quad \text{on } \Gamma_t,$$

which leads directly to the following expression

$$\operatorname{Re} \frac{\mathcal{T}''}{i \mathcal{T}'} = \tilde{\kappa} |\mathcal{T}'| \quad \text{on } \Gamma_t.$$

Moreover, the same expression can be derived on Γ_b with the curvature κ_b .

Therefore, we express the holomorphic function as

$$(41) \quad \frac{\mathcal{T}''}{i \mathcal{T}'} = u_r + i u_i,$$

where the real component u_r satisfies

$$\begin{cases} \Delta u_r = 0 & \text{in } \mathcal{S}_0, \\ u_r|_{\Gamma_t} = \tilde{\kappa} |\mathcal{T}'| |_{\Gamma_t} = w |\mathcal{T}'| |_{\Gamma_t}, \quad u_r|_{\Gamma_b} = \tilde{\kappa}_b |\mathcal{T}'| |_{\Gamma_b}. \end{cases}$$

Performing standard elliptic estimate in Sobolev spaces, we have

$$\|u_r\|_{H^{k+1}} \leq C (\|w |\mathcal{T}'|\|_{H^{k+1/2}} + \|\tilde{\kappa}_b |\mathcal{T}'|\|_{H^{k+1/2}(\Gamma_b)})$$

where C is a constant depending on k, \mathcal{S}_0 .

We firstly take care of the term $\|w |\mathcal{T}'|\|_{H^{k+1/2}}$. In fact, one derivative on $|\mathcal{T}'|$ in this norm results in terms like $g \partial_{\bar{x}}^2 x w$ and $g \partial_{\bar{x}}^2 z w$, where $\alpha w \in H^{k+1/2}$, and g is some function depending on $\mathcal{T}^{(l)}$ ($l \leq k+1$) and can be bounded by L^∞ norm of $\mathcal{T}^{(l)}$ by Proposition 1. Consequently, we write

$$\partial_{\bar{x}}^2 x w = \alpha^{-1} \partial_{\bar{x}}^2 x \alpha w = (\alpha^{-1} \partial_{\bar{x}}^2 x - a_{x,1}) \alpha w + a_{x,1} \alpha w$$

where $a_{x,1}$ is defined in Lemma 8 where $u = \partial_{\bar{x}}x$. Meanwhile, $\partial_{\bar{x}}^2zw$ can be expressed in a similar way. As a result, thanks to Lemma 8 and (39), we know that the corresponding weighted limit for \mathcal{T}'' can be written in the following form:

$$T_{2,c} = \chi_{l,1}T_{l,1} - \beta_r\chi_{r,1}T_{r,1}.$$

These expressions lead to

$$|w|\mathcal{T}'|_{H^{k+1/2}} \leq C(1 + |\alpha^{-1}\mathcal{T}'' - T_{2,c}|_{H^{k-1/2}} + |a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty})|\alpha w|_{H^{k+1/2}}$$

where C is a constant depending on the bound L given in Definition 2.

Similarly, we can also prove that

$$\begin{aligned} |\tilde{\kappa}_b|\mathcal{T}'|_{H^{k+1/2}(\Gamma_b)} &\leq C(1 + |\alpha^{-1}\mathcal{T}'' - T_{2,c}|_{H^{k-1/2}(\Gamma_b)} + |a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty})|\alpha\tilde{\kappa}_b|_{H^{k+1/2}(\Gamma_b)} \\ &\leq C(1 + |\alpha^{-1}\mathcal{T}'' - T_{2,c}|_{H^{k-1/2}(\Gamma_b)} + |a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty}), \end{aligned}$$

where $|\alpha\tilde{\kappa}_b|_{H^{k+1/2}(\Gamma_b)}$ can be controlled by the the initial bound of in $\Lambda(T, k)$ and the bottom Γ_b .

As a result, going back to the estimate of u_r and applying Lemm 3, we derive

$$\begin{aligned} \|u_r\|_{H^{k+1}} &\leq C(1 + |\alpha^{-1}\mathcal{T}'' - T_{2,c}|_{H^{k-1/2}} + |\alpha^{-1}\mathcal{T}'' - T_{2,c}|_{H^{k-1/2}(\Gamma_b)} + |a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty}) \times \\ &\quad (1 + |\alpha w|_{H^{k+1/2}}) \\ &\leq C(1 + \|\alpha^{-1}\mathcal{T}'' - T_{2,c}\|_{H^k} + |a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty})(1 + |\alpha w|_{H^{k+1/2}}) \end{aligned}$$

Step 2. Retrieve the estimate of u_i from u_r . *Different from the infinite-depth case, direct computations show that the L^2 norm of u_i cannot be bounded by $\|u_r\|_{L^2}$.* But we can still have higher-order estimates for u_i . Applying C-R equations, we have

$$\|\nabla u_i\|_{H^k} = \|\nabla u_r\|_{H^k} \leq C(1 + \|\alpha^{-1}\mathcal{T}'' - T_{2,c}\|_{H^k} + |a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty})(1 + |\alpha w|_{H^{k+1/2}}).$$

As a result, going back to the estimate of the holomorphic function $\frac{\mathcal{T}''}{i\mathcal{T}'}$, we need to take one more derivative and set

$$H := \frac{\mathcal{T}'''}{i\mathcal{T}'} - \frac{\mathcal{T}''i\mathcal{T}''}{(i\mathcal{T}')^2} = \left(\frac{\mathcal{T}''}{i\mathcal{T}'}\right)' = \partial_{\bar{x}}u_r + i\partial_{\bar{x}}u_i.$$

Similarly as before, the corresponding weighted limit for \mathcal{T}''' is denoted by

$$T_{3,c} = \chi_{l,2}T_{l,1} + \beta_r^2\chi_{r,2}T_{r,1}$$

with $\chi_{i,2}$ defined in Lemma 8. We plan to derive the estimate of \mathcal{T}' from $\partial_{\bar{x}}u_r, \partial_{\bar{x}}u_i$. Considering the weighted limits of derivatives of \mathcal{T} as above and rewriting H into the following form

$$H = \frac{\mathcal{T}''' - \alpha T_{3,c}}{i\mathcal{T}'} - \frac{i\mathcal{T}''(\mathcal{T}'' - \alpha T_{2,c})}{(i\mathcal{T}')^2} - \frac{i\alpha(T_{2,c}\mathcal{T}'' - T_{3,c}\mathcal{T}')}{(i\mathcal{T}')^2},$$

we arrive at

$$(42) \quad \begin{aligned} \mathcal{T}''' - \alpha T_{3,c} &= \frac{i\mathcal{T}''(\mathcal{T}'' - \alpha T_{2,c})}{i\mathcal{T}'} + \frac{i\alpha T_{2,c}(\mathcal{T}'' - \alpha T_{2,c})}{i\mathcal{T}'} - \frac{i\alpha T_{3,c}(\mathcal{T}' - \alpha T_{1,c})}{i\mathcal{T}'} \\ &\quad + \frac{i\alpha^2(T_{2,c}^2 - T_{3,c}T_{1,c})}{i\mathcal{T}'} + i\mathcal{T}'(\partial_{\bar{x}}u_r + i\partial_{\bar{x}}u_i). \end{aligned}$$

Noticing that

$$\begin{aligned} T_{2,c}^2 - T_{3,c}T_{1,c} &= (\chi_{l,1}T_{l,1} - \beta_r\chi_{r,1}T_{r,1})^2 - (\chi_{l,2}T_{l,1} + \beta_r^2\chi_{r,2}T_{r,1})(\chi_l T_{l,1} + \chi_r T_{r,1}) \\ &= (\chi_{l,1}^2 - \chi_{l,2}\chi_l)T_{l,1}^2 - \beta_r^2(\chi_{r,1}^2 - \chi_{r,2}\chi_r)T_{r,1}^2, \end{aligned}$$

we conclude

$$\begin{aligned} \|\alpha^{-1}\mathcal{T}''' - T_{1,c}\|_{H^k} &\leq \left\| \frac{(\alpha^{-1}\mathcal{T}'' - T_{2,c})^2}{\mathcal{T}'} \right\|_{H^k} + 2\left\| \frac{T_{2,c}(\mathcal{T}'' - \alpha T_{2,c})}{\mathcal{T}'} \right\|_{H^k} + \left\| \frac{T_{3,c}(\mathcal{T}' - \alpha T_{1,c})}{\mathcal{T}'} \right\|_{H^k} \\ &\quad + \left\| \frac{\alpha(T_{2,c}^2 - T_{3,c}T_{1,c})}{\mathcal{T}'} \right\|_{H^k} + \|\alpha^{-1}\mathcal{T}'(\partial_{\bar{x}}u_r + i\partial_{\bar{x}}u_i)\|_{H^k}. \end{aligned}$$

Applying Proposition 1, Lemma 8 and the estimates of u_r, u_i above leads to

$$\begin{aligned}
& \|\alpha^{-1}\mathcal{T}''' - T_{1,c}\|_{H^k} \\
(43) \quad & \leq C[\|\alpha^{-1}\mathcal{T}'' - T_{2,c}\|_{H^k}^2 + \|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{H^k}^2 + |a_{x,i}|_{L^\infty}^2 + |a_{z,i}|_{L^\infty}^2 + \|\partial_{\bar{x}}u_r\|_{H^k} + \|\partial_{\bar{x}}u_i\|_{H^k}] \\
& \leq C[\|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{H^{k+1}}^2 + |a_{x,i}|_{L^\infty}^2 + |a_{z,i}|_{L^\infty}^2 + (1 + \|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{H^{k+1}} + |a_{x,i}|_{L^\infty} \\
& \quad + |a_{z,i}|_{L^\infty})(1 + |\alpha w|_{H^{k+1/2}})].
\end{aligned}$$

On the other hand, considering H^{k-1} estimate for $\alpha^{-1}\mathcal{T}''' - \chi_l T_{1,c}$ in a similar way as above (where one \mathcal{T}'' from the first term on the right side of (42) is bounded by Proposition 1), we obtain

$$\begin{aligned}
& \|\alpha^{-1}\mathcal{T}''' - T_{3,c}\|_{H^{k-1}} \\
& \leq C[(1 + |a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty})\|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{H^k} + |a_{x,i}|_{L^\infty}^2 + |a_{z,i}|_{L^\infty}^2 + (1 + \|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{H^k} \\
& \quad + |a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty})(1 + |\alpha w|_{H^{k-1/2}})].
\end{aligned}$$

Moreover, applying an interpolation in Sobolev spaces leads to

$$\begin{aligned}
\|\alpha^{-1}\mathcal{T}''' - T_{3,c}\|_{H^{k-1}} & \leq \epsilon(1 + |a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty})\|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{H^{k+1}} \\
& \quad + Q(\epsilon^{-1}, \|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{L^2}, |a_{x,i}|_{L^\infty}, |a_{z,i}|_{L^\infty})(1 + |\alpha w|_{H^{k-1/2}})
\end{aligned}$$

where $\epsilon > 0$ is a small constant. Consequently, we obtain by Lemma 8 that

$$\|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{H^{k+1}} \leq Q(\|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{L^2}, |a_{x,i}|_{L^\infty}, |a_{z,i}|_{L^\infty})(1 + |\alpha w|_{H^{k-1/2}}),$$

where the bound L in Definition 2 is applied to $|a_{x,i}|_{L^\infty} + |a_{z,i}|_{L^\infty}$. As a result, substituting this estimate into (43) and applying Lemma 8 again, we obtain the desired estimate for \mathcal{T} . The estimate for $T_{1,c}$ follows directly from (40). \square

We present a useful estimate which has appeared in the proof above.

Corollary 2. *Let f be a function smooth enough in \mathcal{S}_0 and the integer $k \geq 2$. Then there hold*

$$\|\partial_{\bar{x}}^2 x f\|_{H^k} + \|\partial_{\bar{x}}^2 z f\|_{H^k} \leq Q(\mathcal{T}', w, k - 1/2)\|\alpha f\|_{H^k}$$

and

$$|\partial_{\bar{x}}^2 x f|_{H^k(\Gamma_j)} + |\partial_{\bar{x}}^2 z f|_{H^k(\Gamma_j)} \leq Q(\mathcal{T}', w, k)|\alpha f|_{H^k(\Gamma_j)}$$

where $j = t, b$.

Proof. In fact, similar argument as in the proof above shows that

$$\begin{aligned}
\|\partial_{\bar{x}}^2 x f\|_{H^k} & \leq \|(\alpha^{-1}\partial_{\bar{x}}^2 x - a_{x,1})\alpha f\|_{H^k} + \|a_{x,1}\alpha f\|_{H^k} \\
& \leq C(\|\alpha^{-1}\partial_{\bar{x}}^2 x - a_{x,1}\|_{H^{k+1}} + |a_{x,i}|_{L^\infty})\|\alpha f\|_{H^k},
\end{aligned}$$

where $a_{x,1}$ is defined in Lemma 8 with $u = \partial_{\bar{x}}x$. Applying Proposition 7 leads to the desired estimate immediately. As a result, the remaining estimates can be proved in a similar way. \square

Meanwhile, considering $\partial_{\bar{x}}z$ component of \mathcal{T}' on Γ_t , we obtain immediately the following estimate.

Corollary 3. *Let the integer $k \geq 1$. Then one has*

$$|\alpha^{-1}\partial_{\bar{x}}\tilde{\zeta}|_{H^{k+3/2}} \leq Q(\mathcal{T}, w, k + 1/2).$$

Proof. In fact, noticing that

$$a_z|_{\bar{z}=0} = 0, \quad \partial_{\bar{x}}z|_{\bar{z}=0} = \partial_{\bar{x}}\tilde{\zeta}$$

and applying Lemma 3 lead to

$$|\alpha^{-1}\partial_{\bar{x}}\tilde{\zeta}|_{H^{k+3/2}} \leq C\|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{H^{k+2}}.$$

Therefore, the proof is finished by applying Proposition 7. \square

Besides, we also need another “local” higher-order estimate of \mathcal{T} near the bottom Γ_b , which will be used in the a priori estimate. Notice that these kinds of higher-order estimate work thanks to the separation of Γ_t, Γ_b for \mathcal{S}_0 , which is handled by localization near the bottom as in classical water waves, see [3].

Corollary 4. *Let the integer $k \geq 1$. There holds*

$$|\alpha^{-1}\mathcal{T}' - T_{1,c}|_{H^{k+2}(\Gamma_b)} \leq Q(\mathcal{T}', w, k)$$

Proof. In fact, we only need to modify the proof of Proposition 7. To begin with, we set

$$\chi_b(\tilde{z}) = \begin{cases} 1, & \tilde{z} = -\omega, \\ 0, & \tilde{z} \geq -\omega + \delta \end{cases}$$

for some small constant $\delta > 0$. As a result, we find that $\chi_b u_r$ with u_r from (41) satisfies

$$\begin{cases} \Delta \chi_b u_r = [\Delta, \chi_b] u_r & \text{in } \mathcal{S}_0, \\ \chi_b u_r|_{\Gamma_t} = 0, \quad u_r|_{\Gamma_b} = \tilde{\kappa}_b |\mathcal{T}'| |_{\Gamma_b}. \end{cases}$$

We have by standard elliptic estimates, Proposition 1 and Corollary 2 that

$$\begin{aligned} \|\chi_b u_r\|_{H^{k+3/2}} &\leq C(\|[\Delta, \chi_b] u_r\|_{H^{k-1/2}} + |\tilde{\kappa}_b |\mathcal{T}'||_{H^{k+1}(\Gamma_b)}) \\ &\leq C(\|u_r\|_{H^{k+1/2}} + Q(\mathcal{T}', w, k)), \end{aligned}$$

while notice that the sufficient smoothness of the bottom Γ_b of Ω is used here.

Moreover, applying the estimate of u_r from the proof of Proposition 7 and using an interpolation, we obtain the following estimate

$$\|\chi_b u_r\|_{H^{k+3/2}} \leq Q(\mathcal{T}', w, k),$$

which implies immediately

$$\|u_r\|_{H^{k+3/2}(\mathcal{S}_b)} \leq Q(\mathcal{T}', w, k)$$

where $\mathcal{S}_b \subset \mathcal{S}_0$ is a flat strip near Γ_b with the height smaller than δ .

Besides, we also derive

$$\|\nabla u_i\|_{H^{k+1/2}(\mathcal{S}_b)} = \|\nabla u_r\|_{H^{k+1/2}(\mathcal{S}_b)} \leq Q(\mathcal{T}', w, k).$$

Therefore, repeating estimate (43) from the proof of Proposition 7 and applying the new estimates for u_r, u_i above, we derive a higher-order estimate of \mathcal{T} in \mathcal{S}_b :

$$\begin{aligned} &\|\alpha^{-1}\mathcal{T}''' - T_{3,c}\|_{H^{k+1/2}(\mathcal{S}_b)} \\ &\leq C[\|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{H^{k+3/2}(\mathcal{S}_b)}^2 + |a_{x,i}|_{L^\infty}^2 + |a_{z,i}|_{L^\infty}^2 + \|\partial_{\tilde{x}} u_r\|_{H^{k+1/2}(\mathcal{S}_b)} + \|\partial_{\tilde{x}} u_i\|_{H^{k+1/2}(\mathcal{S}_b)}] \\ &\leq Q(\mathcal{T}', w, k) \end{aligned}$$

In the end, applying Proposition 7 and a trace theorem again, the proof is finished. \square

Next, we investigate the time dependence of the conformal mapping, where a loss of 1/2 derivative appears due to the space derivative in (4).

Proposition 8. *Let γ satisfy*

$$0 < \gamma + 1 \leq \min\left\{\frac{2\pi}{\omega}, \frac{2\pi}{\omega_r}\right\}.$$

Then there holds for $k \geq 0$ that

$$\|\alpha^{-(\gamma+3)/2} \partial_t \mathcal{T}\|_{H^{k+2}} \leq Q(\mathcal{T}', w, k + 1/2) (\|\alpha^{-\gamma-2} \widetilde{\partial}_t \zeta - a_{\zeta_t}\|_{H^{k+3/2}} + |a_{\zeta_t, i}|_{L^\infty}),$$

where a_{ζ_t} is the weighted limit defined in (34).

Proof. The proof here is similar as the proof of the previous proposition. In fact, inspired by [56], we start with computing the following holomorphic function

$$\frac{\partial_t \mathcal{T}}{\mathcal{T}'} = \frac{\partial_t x + i \partial_t z}{\partial_{\bar{x}} x + i \partial_{\bar{x}} z} = \frac{1}{|\mathcal{T}'|^2} [(\partial_{\bar{x}} x \partial_t x + \partial_{\bar{x}} z \partial_t z) + i(\partial_{\bar{x}} x \partial_t z - \partial_{\bar{x}} z \partial_t x)].$$

We set again

$$\frac{\partial_t \mathcal{T}}{\mathcal{T}'} := u_r + i u_i.$$

Step 1. u_i estimate. Since

$$(\partial_t z - (\partial_{\bar{x}} x)^{-1} \partial_{\bar{x}} z \partial_t x)|_{\Gamma_t; \tilde{z}=0} = \widetilde{\partial_t \zeta} \quad \text{and} \quad (\partial_t z - (\partial_{\bar{x}} x)^{-1} \partial_{\bar{x}} z \partial_t x)|_{\Gamma_b; \tilde{z}=-\omega} = 0,$$

we obtain immediately the Dirichlet system of u_i :

$$\begin{cases} \Delta u_i = 0 & \text{in } S_0, \\ u_i|_{\Gamma_t} = |\mathcal{T}'|^{-2} \partial_{\bar{x}} x \widetilde{\partial_t \zeta}, \quad u_i|_{\Gamma_b} = 0. \end{cases}$$

Applying Proposition 4 leads to

$$\|\alpha^{-(\gamma+1)/2} u_i\|_{H^{k+2}} \leq C |\alpha^{-(\gamma+1)/2} |\mathcal{T}'|^{-2} \partial_{\bar{x}} x \widetilde{\partial_t \zeta}|_{H^{k+3/2}},$$

where one requires

$$\frac{|\gamma+1|}{2} \leq \min\left\{\frac{\pi}{\omega}, \frac{\pi}{\omega_r}\right\}.$$

For the right-side term, direct computations and applying Proposition 1 and Corollary 2 lead to

$$|\alpha^{-(\gamma+1)/2} |\mathcal{T}'|^{-2} \partial_{\bar{x}} x \widetilde{\partial_t \zeta}|_{H^{k+3/2}} \leq Q(\mathcal{T}', w, k+1/2) |\alpha^{-(\gamma+1)/2} \alpha^{-1} \widetilde{\partial_t \zeta}|_{H^{k+3/2}},$$

where the right side can be expanded as below following the spirit of Corollary 2:

$$|\alpha^{-(\gamma+1)/2} \alpha^{-1} \widetilde{\partial_t \zeta}|_{H^{k+3/2}} \leq |\alpha^{(\gamma+1)/2} (\alpha^{-\gamma-2} \widetilde{\partial_t \zeta} - a_{\zeta_t})|_{H^{k+3/2}} + |\alpha^{(\gamma+1)/2} a_{\zeta_t}|_{H^{k+3/2}}.$$

Noticing that when $\gamma+1 > 0$, the second term on the right side above is integrable so that we conclude the estimate of u_i :

$$\|\alpha^{-(\gamma+1)/2} u_i\|_{H^{k+2}} \leq Q(\mathcal{T}', w, k-1) (|\alpha^{-\gamma-2} \widetilde{\partial_t \zeta} - a_{\zeta_t}|_{H^{k+3/2}} + |a_{\zeta_t}|_{L^\infty}).$$

Step 2. u_r estimate. In fact, using C-R equations leads directly to the higher-order estimate of u_r :

$$\|\alpha^{-(\gamma+1)/2} \nabla u_r\|_{H^{k+1}} = \|\alpha^{-(\gamma+1)/2} \nabla u_i\|_{H^{k+1}} \leq Q(\mathcal{T}', w, k+1/2) (|\alpha^{-\gamma-2} \widetilde{\partial_t \zeta} - a_{\zeta_t}|_{H^{k+3/2}} + |a_{\zeta_t}|_{L^\infty}).$$

It remains to prove the weighted L^2 estimate of u_r . We plan to recover it by using the Laplace Transform in S_0 . To begin with, C-R equations imply the following system of u_r :

$$\begin{cases} \Delta u_r = 0 & \text{in } S_0, \\ \partial_{\bar{z}} u_r|_{\Gamma_t} = -\partial_{\bar{x}} u_i|_{\Gamma_t}, \quad \partial_{\bar{z}} u_i|_{\Gamma_b} = -\partial_{\bar{x}} u_i|_{\Gamma_b} = 0. \end{cases}$$

Recalling the Laplace Transform on a sufficiently smooth function $u = u(\tilde{x})$:

$$\tilde{u}(\lambda) = \mathcal{L}u(\lambda) = \int_{\mathbb{R}} e^{-\lambda \tilde{x}} u(\tilde{x}) d\tilde{x}, \quad \forall \lambda \in \mathbb{C},$$

we have immediately the following system of $\tilde{u}_r(t, \lambda, \tilde{z})$:

$$\begin{cases} \Delta \tilde{u}_r = 0, & \tilde{z} \in [-\omega, 0], \\ \partial_{\bar{z}} \tilde{u}_r|_{\tilde{z}=0} = -\partial_{\bar{x}} \tilde{u}_i|_{\tilde{z}=0}, \quad \partial_{\bar{z}} \tilde{u}_r|_{\tilde{z}=-\omega} = 0. \end{cases}$$

Solving the system above directly, one has

$$\tilde{u}_r = \frac{\cos(\lambda(\omega + \tilde{z}))}{\sin(\lambda\omega)} \tilde{u}_i|_{\tilde{z}=0} = \frac{e^{i\lambda(\omega + \tilde{z})} + e^{-i\lambda(\omega + \tilde{z})}}{e^{i\lambda\omega} - e^{-i\lambda\omega}} \tilde{u}_i|_{\tilde{z}=0}.$$

Therefore, denoting by

$$\gamma_1 = \frac{\gamma + 1}{2},$$

the weighted L^2 norm of u_r is decomposed as below

$$(44) \quad \begin{aligned} \|\alpha^{-\gamma_1} u_r\|_{L^2} &\leq \|\alpha^{-\gamma_1} \chi_l u_r\|_{L^2} + \|\alpha^{-\gamma_1} \chi_r u_r\|_{L^2} + \|\alpha^{-\gamma_1} (1 - \chi_l - \chi_r) u_r\|_{L^2} \\ &\leq C(\|e^{-\gamma_1 \tilde{x}} u_r\|_{L^2} + \|e^{\gamma_1 \beta_r \tilde{x}} u_r\|_{L^2}), \end{aligned}$$

where both the two weighted terms on the right side above need to be handled.

For the first term, applying the Parseval equality for the Laplace Transform, one obtains

$$\|e^{-\gamma_1 \tilde{x}} u_r\|_{L^2}^2 = \int_{-\omega}^0 \frac{1}{2\pi i} \int_{\operatorname{Re} \lambda = \gamma_1} |\check{u}_r(\lambda, \tilde{z})|^2 d\lambda d\tilde{z}$$

with

$$\lambda = \gamma_1 + i\xi.$$

Substituting the expression of \check{u}_r into this integral, we derive

$$\begin{aligned} \|e^{-\gamma_1 \tilde{x}} u_r\|_{L^2}^2 &= \int_{-\omega}^0 \frac{1}{2\pi i} \int_{\operatorname{Re} \lambda = \gamma_1} \left| \frac{\cos(\lambda(\omega + \tilde{z}))}{\sin(\lambda\omega)} \check{u}_i|_{\tilde{z}=0} \right|^2 d\lambda d\tilde{z} \\ &:= \int_{-\omega}^0 \frac{1}{2\pi i} \int_{\operatorname{Re} \lambda = \gamma_1} B d\lambda d\tilde{z}, \end{aligned}$$

where

$$\begin{aligned} B &= \left| \frac{e^{i(\gamma_1 + i\xi)(\omega + \tilde{z})} + e^{-i(\gamma_1 + i\xi)(\omega + \tilde{z})}}{e^{i(\gamma_1 + i\xi)\omega} - e^{-i(\gamma_1 + i\xi)\omega}} \right|^2 |\check{u}_i|_{\tilde{z}=0}|^2 \\ &= \left| \frac{e^{-\xi(\omega + \tilde{z})} e^{i\gamma_1(\omega + \tilde{z})} + e^{\xi(\omega + \tilde{z})} e^{-i\gamma_1(\omega + \tilde{z})}}{e^{-\xi\omega} e^{i\gamma_1\omega} - e^{\xi\omega} e^{-i\gamma_1\omega}} \right|^2 |\check{u}_i|_{\tilde{z}=0}|^2 \leq C |\check{u}_i|_{\tilde{z}=0}|^2 \end{aligned}$$

with the constant C depending on ω, γ .

Consequently, we arrive at the following estimate

$$\|e^{-\gamma_1 \tilde{x}} u_r\|_{L^2}^2 \leq C \int_{-\omega}^0 \frac{1}{2\pi i} \int_{\operatorname{Re} \lambda = \gamma_1} |\check{u}_i|_{\tilde{z}=0}|^2 d\lambda d\tilde{z} \leq C \|e^{-\gamma_1 \tilde{x}} u_i|_{\tilde{z}=0}\|_{L^2}^2.$$

Meanwhile, for the second term on the right side of (44), similar arguments as above lead to

$$\|e^{\gamma_1 \beta_r \tilde{x}} u_r\|_{L^2}^2 \leq C \|e^{\gamma_1 \beta_r \tilde{x}} u_i|_{\tilde{z}=0}\|_{L^2}^2.$$

Substituting the boundary value

$$u_i|_{\tilde{z}=0} = |\mathcal{T}'|^{-2} \partial_{\tilde{x}} \widetilde{\partial_t \zeta}$$

into the two estimates above, applying Proposition 1 and going back to (44), one derives

$$\begin{aligned} \|\alpha^{-\gamma_1} u_r\|_{L^2} &\leq C(|e^{-\gamma_1 \tilde{x}} \alpha^{-1} \widetilde{\partial_t \zeta}|_{L^2} + |e^{\gamma_1 \beta_r \tilde{x}} \alpha^{-1} \widetilde{\partial_t \zeta}|_{L^2}) \\ &\leq C|\alpha^{-\gamma_1} \alpha^{-1} \widetilde{\partial_t \zeta}|_{L^2} \leq C(|\alpha^{-\gamma-2} \widetilde{\partial_t \zeta} - a_{\zeta_t}|_{L^2} + |a_{\zeta_t}|_{L^\infty}), \end{aligned}$$

where $\gamma_1 = (\gamma + 1)/2 > 0$ is needed again.

As a result, summing up all these estimates of u_r , we finally conclude that

$$\|\alpha^{-(\gamma+1)/2} u_r\|_{H^{k+2}} \leq Q(\mathcal{T}', w, k+1/2) (|\alpha^{-\gamma-2} \widetilde{\partial_t \zeta} - a_{\zeta_t}|_{H^{k+3/2}} + |a_{\zeta_t}|_{L^\infty}).$$

Step 3. End of the proof. Going back to the $\partial_t \mathcal{T}$ expression in the beginning of the proof, we can finish the proof thanks to Step 1-2 and Corollary 2. \square

Remark 10. Combining Proposition 8 with (39), (40) and Proposition 7, we conclude that the weighted limits

$$\partial_t T_{1,i} = 0, \quad (i = l, r), \quad \text{i.e.} \quad b'(t) = b'_r(t) = b'_{r,2}(t) = 0.$$

This happens due to the fact that a higher-order weight is used for $\partial_t \mathcal{T}$ in our settings.

On the other hand, we consider the weighted estimate for $\widetilde{\partial}_t \zeta$, which is a part of the boundary value of $\partial_t \mathcal{T}$ and deals with the relationship between $\widetilde{D}_t w$ and $\widetilde{\partial}_t \zeta$. Notice that a loss of 1/2 order space derivative still exists here.

Proposition 9. *Let the integer $k \geq 0$ and*

$$a_{w_t}(t, \widetilde{x}) = \chi_l a_{w_t, l} + \chi_r a_{w_t, r}$$

with

$$a_{w_t, l}(t) = \lim_{\widetilde{x} \rightarrow -\infty} \alpha^{-\gamma} \widetilde{D}_t w, \quad a_{w_t, r}(t) = \lim_{\widetilde{x} \rightarrow +\infty} \alpha^{-\gamma} \widetilde{D}_t w.$$

Then the following estimate holds:

$$\begin{aligned} |\alpha^{-\gamma-2} \widetilde{\partial}_t \zeta - a_{\zeta_t}|_{H^{k+3/2}} &\leq Q(\mathcal{T}', \widetilde{\Phi}, w, k+1/2) (|\alpha^{-\gamma} \widetilde{D}_t w - a_{w_t}|_{H^{k-1/2}} + |\alpha^{-\gamma-2} \widetilde{v} - a_v|_{H^{k+3/2}} \\ &\quad + |a_v|_{L^\infty} + |\alpha^{-\gamma-2} \widetilde{\partial}_t \zeta - a_{\zeta_t}|_{L^2} + |a_{\zeta_t, i}|_{L^\infty}). \end{aligned}$$

Proof. To begin with, we recall another expression for $D_t \kappa$ from [57]:

$$D_t \kappa = -\Delta_{\Gamma_t}(v \cdot n_t) - |\nabla_{\tau_t} n_t|^2 (v \cdot n_t) + \nabla_{\tau_t} \nabla_{v^\top} n_t \cdot \tau_t - (\nabla_{\tau_t} v^\top)^\top \cdot \nabla n_t \cdot \tau_t \quad \text{on } \Gamma_t.$$

Applying the parametrization $z = \zeta$ and noticing that

$$v \cdot n_t = (1 + |\partial_x \zeta|^2)^{-\frac{1}{2}} \partial_t \zeta$$

by the kinematic condition (4), we rewrite the equation above as

$$D_t \kappa = -\nabla_{\tau_t} \nabla_{\tau_t} ((1 + |\partial_x \zeta|^2)^{-\frac{1}{2}} \partial_t \zeta) + F(\partial v, \partial^2 n_t, \partial \tau_t),$$

where

$$F(\partial v, \partial^2 n_t, \partial \tau_t) = -|\nabla_{\tau_t} n_t|^2 (v \cdot n_t) + \nabla_{\tau_t} \nabla_{v^\top} n_t \cdot \tau_t - (\nabla_{\tau_t} v^\top)^\top \cdot \nabla n_t \cdot \tau_t$$

with at most two derivatives in one term.

Pulling this equation onto the upper boundary of \mathcal{S}_0 by \mathcal{T} and applying (27), (30), we have

$$\widetilde{D}_t w = \widetilde{D}_t \widetilde{\kappa} = -(\partial_{\widetilde{x}} x)^{-1} b_\zeta^{-\frac{1}{2}} \partial_{\widetilde{x}} [(\partial_{\widetilde{x}} x)^{-1} b_\zeta^{-\frac{1}{2}} \partial_{\widetilde{x}} (b_\zeta^{-\frac{1}{2}} \widetilde{\partial}_t \zeta)] + \widetilde{F}$$

where

$$(\nabla_{\tau_t}) \circ \mathcal{T}|_{\Gamma_t} = (\nabla_X \widetilde{X})^T \circ \mathcal{T} \widetilde{\tau}_t \cdot \nabla = (\partial_{\widetilde{x}} x)^{-1} b_\zeta^{-\frac{1}{2}} \partial_{\widetilde{x}}$$

and we write

$$(45) \quad b_\zeta := 1 + |(\partial_{\widetilde{x}} x)^{-1} \partial_{\widetilde{x}} \widetilde{\zeta}|^2.$$

Recalling (36), (38), the definition of a_{ζ_t} in Proposition 8 and the space of $\alpha \widetilde{\zeta}$ and taking the limit of $\alpha^{-\gamma} \widetilde{D}_t w$, we find immediately

$$\begin{aligned} a_{w_t, l} &= \lim_{\widetilde{x} \rightarrow -\infty} \alpha^{-\gamma} \widetilde{D}_t w = -(\gamma+1)(\gamma+2) a_{\zeta_t, l}(t), \\ a_{w_t, r} &= \lim_{\widetilde{x} \rightarrow +\infty} \alpha^{-\gamma} \widetilde{D}_t w = -(\gamma+1)(\gamma+2) \beta_r^2 b_r^2(t) a_{\zeta_t, r}(t). \end{aligned}$$

This means that $\alpha^{-\gamma} \widetilde{D}_t w$ doesn't vanish near ∞ in general.

As a result, denoting by

$$a_{\zeta_t, x} = \chi_l a_{\zeta_t, l}(t) + \chi_r b_r^2(t) a_{\zeta_t, r}(t)$$

and applying $L^2(\mathbb{R})$ inner product on the expression $\alpha^{-\gamma} \widetilde{D}_t w - a_{w_t}$ with $\alpha^{-\gamma} (\partial_{\widetilde{x}} x)^{-2} \widetilde{\partial}_t \zeta - a_{\zeta_t, x}$ leads to

$$\begin{aligned} &\int_{\mathbb{R}} (\alpha^{-\gamma} \widetilde{D}_t w - a_{w_t}) (\alpha^{-\gamma} (\partial_{\widetilde{x}} x)^{-2} \widetilde{\partial}_t \zeta - a_{\zeta_t, x}) d\widetilde{x} \\ (46) \quad &= - \int_{\mathbb{R}} \left(\alpha^{-\gamma} (\partial_{\widetilde{x}} x)^{-1} b_\zeta^{-\frac{1}{2}} \partial_{\widetilde{x}} [(\partial_{\widetilde{x}} x)^{-1} b_\zeta^{-\frac{1}{2}} \partial_{\widetilde{x}} (b_\zeta^{-\frac{1}{2}} \widetilde{\partial}_t \zeta)] + a_{w_t} \right) (\alpha^{-\gamma} (\partial_{\widetilde{x}} x)^{-2} \widetilde{\partial}_t \zeta - a_{\zeta_t, x}) d\widetilde{x} \\ &\quad + \int_{\mathbb{R}} \alpha^{-\gamma} \widetilde{F} (\alpha^{-\gamma} (\partial_{\widetilde{x}} x)^{-2} \widetilde{\partial}_t \zeta - a_{\zeta_t, x}) d\widetilde{x} \\ &:= A_1 + A_2, \end{aligned}$$

and we deal with these two integrals one by one.

Frist, applying integration by parts shows immediately that

$$\begin{aligned}
A_1 &= -\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-1}b_\zeta^{-\frac{1}{2}}(\partial_{\tilde{x}}x)^{-1}b_\zeta^{-\frac{1}{2}}\partial_{\tilde{x}}(b_\zeta^{-\frac{1}{2}}\widetilde{\partial}_t\zeta)(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})\Big|_{-\infty}^{+\infty} \\
&\quad + \int_{\mathbb{R}} \partial_{\tilde{x}}(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-1}b_\zeta^{-\frac{1}{2}})(\partial_{\tilde{x}}x)^{-1}b_\zeta^{-\frac{1}{2}}\partial_{\tilde{x}}(b_\zeta^{-\frac{1}{2}}\widetilde{\partial}_t\zeta)(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\tilde{x} \\
&\quad + \int_{\mathbb{R}} \alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-1}b_\zeta^{-\frac{1}{2}}(\partial_{\tilde{x}}x)^{-1}b_\zeta^{-\frac{1}{2}}\partial_{\tilde{x}}(b_\zeta^{-\frac{1}{2}}\widetilde{\partial}_t\zeta)\partial_{\tilde{x}}(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\tilde{x} \\
&\quad - \int_{\mathbb{R}} a_{w_t}(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\tilde{x} \\
&:= A_{11} + A_{12} + A_{13} + A_{14},
\end{aligned}$$

where direct computations as before lead to

$$A_{11} = 0.$$

- A_{12} part. Expanding the derivative $\partial_{\tilde{x}}(\cdot)$ terms directly leads to

$$\begin{aligned}
A_{12} &= \int_{\mathbb{R}} \partial_{\tilde{x}}(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-1})b_\zeta^{-\frac{1}{2}}(\partial_{\tilde{x}}x)^{-1}b_\zeta^{-\frac{1}{2}}b_\zeta^{-\frac{1}{2}}\partial_{\tilde{x}}\widetilde{\partial}_t\zeta(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\tilde{x} \\
&\quad + \int_{\mathbb{R}} \partial_{\tilde{x}}(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-1})b_\zeta^{-\frac{1}{2}}(\partial_{\tilde{x}}x)^{-1}b_\zeta^{-\frac{1}{2}}(\partial_{\tilde{x}}b_\zeta^{-\frac{1}{2}})\widetilde{\partial}_t\zeta(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\tilde{x} \\
&\quad + \int_{\mathbb{R}} \alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-1}\partial_{\tilde{x}}(b_\zeta^{-\frac{1}{2}})(\partial_{\tilde{x}}x)^{-1}b_\zeta^{-\frac{1}{2}}\partial_{\tilde{x}}(b_\zeta^{-\frac{1}{2}}\widetilde{\partial}_t\zeta)(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\tilde{x} \\
&:= A_{121} + A_{122} + A_{123},
\end{aligned}$$

where one notes that $\partial_{\tilde{x}}(b_\zeta^{-\frac{1}{2}})$ in A_{122}, A_{123} can be estimated directly by some corresponding Sobolev norm of $\alpha^{-1}\widetilde{\zeta}$.

Similarly as before, adding the ‘‘limit part’’ to A_{121} and noticing that

$$\partial_{\tilde{x}}(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-1}) \approx -(\gamma + 1)\alpha^{-\gamma-1}, \quad \tilde{x} \rightarrow -\infty$$

where the computations are similar as those for a_{w_t} , we obtain

$$\begin{aligned}
A_{121} &= \int_{\mathbb{R}} b_\zeta^{-\frac{3}{2}}\left(\partial_{\tilde{x}}(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-1})(\partial_{\tilde{x}}x)^{-1}\partial_{\tilde{x}}\widetilde{\partial}_t\zeta - a_{w_t}\right)(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\tilde{x} \\
&\quad + \int_{\mathbb{R}} b_\zeta^{-\frac{3}{2}}a_{w_t}(\alpha^{-\gamma}(\partial_{\tilde{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\tilde{x} \\
&:= B_1 + B_2.
\end{aligned}$$

As a result, similar arguments and applying Corollary 2 as before lead to

$$|B_1| \leq Q(\mathcal{T}', w, 1)(|\alpha^{-\gamma-2}\partial_{\tilde{x}}\widetilde{\partial}_t\zeta - a_{\zeta_t,1}|_{L^2} + |a_{\zeta_t}|_{L^\infty})(|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2} + |a_{\zeta_t}|_{L^\infty})$$

with $a_{\zeta_t,1}$ defined in Lemma 8.

Moreover, A_{122} can be handled in a similar way as B_1 with the help of Corollary 3 where we take L^2 norm on $\partial_{\tilde{x}}(b_\zeta^{-\frac{1}{2}})$ part:

$$\begin{aligned}
|A_{122}| &\leq Q(\mathcal{T}', w, 1)(|\alpha^{-1}\partial_{\tilde{x}}\widetilde{\zeta}|_{L^2} + |\alpha^{-1}\partial_{\tilde{x}}^2\widetilde{\zeta}|_{L^2})(|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{L^\infty} + |a_{\zeta_t}|_{L^\infty})(|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2} \\
&\quad + |a_{\zeta_t}|_{L^\infty}) \\
&\leq Q(\mathcal{T}', w, 1)(|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{H^1} + |a_{\zeta_t}|_{L^\infty})(|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2} + |a_{\zeta_t}|_{L^\infty})
\end{aligned}$$

For the last term in A_{12} , expanding $\partial_{\bar{x}}$ part and proceeding as above show that

$$\begin{aligned}
A_{123} &= \int_{\mathbb{R}} \partial_{\bar{x}}(b_{\zeta}^{-\frac{1}{2}})b_{\zeta}^{-1} \left(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\partial_{\bar{x}}\widetilde{\partial}_t\zeta - (\gamma+2)\chi_l a_{\zeta_t,l} - (\gamma+2)\beta_r b_r^2(t)a_{\zeta_t,r} \right) \times \\
&\quad (\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x} + \int_{\mathbb{R}} \partial_{\bar{x}}(b_{\zeta}^{-\frac{1}{2}})b_{\zeta}^{-1} \left((\gamma+2)\chi_l a_{\zeta_t,l} + (\gamma+2)\beta_r b_r^2(t)a_{\zeta_t,r} \right) \times \\
&\quad (\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x} + \int_{\mathbb{R}} \alpha^{-\gamma}(\partial_{\bar{x}}x)^{-1}\partial_{\bar{x}}(b_{\zeta}^{-\frac{1}{2}})(\partial_{\bar{x}}x)^{-1}b_{\zeta}^{-\frac{1}{2}}\partial_{\bar{x}}(b_{\zeta}^{-\frac{1}{2}})\widetilde{\partial}_t\zeta \times \\
&\quad (\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x} \\
&:= B_3 + B_4 + B_5,
\end{aligned}$$

where B_3, B_4, B_5 can be handled as above and hence the estimates are omitted here.

- A_{13} part. Expanding the derivative term $\partial_{\bar{x}}(b_{\zeta}^{-\frac{1}{2}}\widetilde{\partial}_t\zeta)$ and adding corresponding limit part lead to

$$\begin{aligned}
A_{13} &= \int_{\mathbb{R}} b_{\zeta}^{-1}\partial_{\bar{x}}(b_{\zeta}^{-\frac{1}{2}})(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})\partial_{\bar{x}}(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x} \\
&\quad + \int_{\mathbb{R}} b_{\zeta}^{-1}\partial_{\bar{x}}(b_{\zeta}^{-\frac{1}{2}})a_{\zeta_t}\partial_{\bar{x}}(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x} \\
&\quad + \int_{\mathbb{R}} b_{\zeta}^{-\frac{3}{2}}\partial_{\bar{x}}(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})\partial_{\bar{x}}(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x} \\
&\quad + \int_{\mathbb{R}} b_{\zeta}^{-\frac{3}{2}}\chi_l' a_{\zeta_t}\partial_{\bar{x}}(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x} \\
&\quad - \int_{\mathbb{R}} b_{\zeta}^{-\frac{3}{2}} \left(\partial_{\bar{x}}(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2})\widetilde{\partial}_t\zeta + (\gamma+2)\chi_l a_{\zeta_t,l} + (\gamma+2)\beta_r b_r^2(t)a_{\zeta_t,r} \right) \partial_{\bar{x}}(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x} \\
&\quad + \int_{\mathbb{R}} b_{\zeta}^{-\frac{3}{2}} \left((\gamma+2)\chi_l a_{\zeta_t,l} + (\gamma+2)\beta_r b_r^2(t)a_{\zeta_t,r} \right) \partial_{\bar{x}}(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x} \\
&:= B_6 + \dots + B_{11},
\end{aligned}$$

where B_6, B_7, B_9, B_{10} can be handled similarly as above and the estimates are omitted again.

Moreover, integrating by parts in B_{11} leads to

$$|B_{11}| \leq Q(\mathcal{T}', w, 1)|a_{\zeta_t}|_{L^\infty} (|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2} + |a_{\zeta_t}|_{L^\infty})$$

For the main non-negative term B_8 , one obtains directly by Corollary 2 and Corollary 3 the following estimate:

$$(47) \quad |\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{H^1}^2 \leq Q(\mathcal{T}', w, 1)(B_8 + |a_{\zeta_t}|_{L^\infty}^2) + |\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2}^2.$$

Consequently, summing up all these estimates, we obtain for A_1 the following inequality

$$A_1 \geq A_{14} + B_2 + B_8 - Q(\mathcal{T}', w, 1)(|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{H^1} + |a_{\zeta_t}|_{L^\infty})(|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2} + |a_{\zeta_t}|_{L^\infty}),$$

where

$$A_{14} + B_2 = \int_{\mathbb{R}} (1 - b_{\zeta}^{-\frac{3}{2}})a_{w_t}(\alpha^{-\gamma}(\partial_{\bar{x}}x)^{-2}\widetilde{\partial}_t\zeta - a_{\zeta_t,x})d\bar{x}$$

and can be handled similarly as above thanks to $1 - b_{\zeta}^{-\frac{3}{2}}$.

Therefore, we conclude that

$$(48) \quad A_1 \geq B_8 - Q(\mathcal{T}', w, 1)(|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{H^1} + |a_{\zeta_t}|_{L^\infty})(|\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2} + |a_{\zeta_t}|_{L^\infty}).$$

- A_2 part. For the $\nabla_{\tau_t}\nabla_{v^+}n_t \cdot \tau_t$ term in F , we perform an integration by parts again (especially for the higher-order estimate) and proceed as before. Meanwhile, notice that each term in \widetilde{F} always

contains $\partial \tilde{n}_t$ when one adds the ‘‘weighted limit’’ to \tilde{v} term and there is at most order-one derivative for \tilde{v} . Here we define the weighted limit of \tilde{v} as

$$(49) \quad a_v = \chi_l \lim_{\tilde{x} \rightarrow -\infty} \alpha^{-\gamma-2} \tilde{v} + \chi_r \lim_{\tilde{x} \rightarrow +\infty} \alpha^{-\gamma-2} \tilde{v}.$$

Consequently, we have the following estimate

$$(50) \quad A_2 \leq Q(\mathcal{T}', w, 1) (|\alpha^{-\gamma-2} \tilde{v} - a_v|_{H^1} + |a_v|_{L^\infty}) (|\alpha^{-\gamma-2} \partial_t \tilde{\zeta} - a_{\zeta_t}|_{L^2} + |a_{\zeta_t}|_{L^\infty})$$

where Corollary 3 and Proposition 10 are applied.

- Weighted H^1 estimate of $\partial_t \tilde{\zeta}$. Summing up all these estimates (47), (48) and (50) above, we finally conclude that

$$\begin{aligned} |\alpha^{-\gamma-2} \partial_t \tilde{\zeta} - a_{\zeta_t}|_{H^1}^2 &\leq Q(\mathcal{T}', w, 1) (|\alpha^{-\gamma} \tilde{D}_t w - a_{w_t}|_{L^2}^2 + |\alpha^{-\gamma-2} \tilde{v} - a_v|_{H^1}^2 + |a_v|_{L^\infty}^2 \\ &\quad + |\alpha^{-\gamma-2} \partial_t \tilde{\zeta} - a_{\zeta_t}|_{L^2}^2 + |a_{\zeta_t}|_{L^\infty}^2). \end{aligned}$$

In the end, for higher-order derivative estimates, one can proceed in a standard way by taking $\partial_{\tilde{x}}$ derivatives and using L^2 inner product from the beginning of this proof. Moreover, counting the order of derivatives for \tilde{v}, ζ , we obtain the desired estimate. \square

4.5. Weighted estimates for the velocity and other quantities. Based on the estimates of \mathcal{T} , we now turn to the estimates for the velocity $\tilde{v} = v \circ \mathcal{T}$, which is also very delicate and involves several steps to reach the desired regularity. Due to the natural loss of 1/2 order space derivative from the kinematic condition (4), we firstly use the potential $\tilde{\Phi} = \Phi \circ \mathcal{T}$ to prove a relatively lower-order estimate for v .

In fact, (4) is rewritten into

$$(51) \quad \partial_t \tilde{\zeta} = (1 + |(\partial_{\tilde{x}} x)^{-1} \partial_{\tilde{x}} \tilde{\zeta}|^2)^{\frac{1}{2}} \tilde{v} \cdot \tilde{n}_t = -(\partial_{\tilde{x}} x)^{-1} \partial_{\tilde{z}} \tilde{\Phi}|_{\tilde{z}=0}.$$

We have the following Neumann system for $\tilde{\Phi}$:

$$(52) \quad \begin{cases} \Delta \tilde{\Phi} = 0 & \text{in } \mathcal{S}_0, \\ \partial_{\tilde{z}} \tilde{\Phi}|_{\Gamma_t} = -(\partial_{\tilde{x}} x) \partial_t \tilde{\zeta}, \quad \partial_{\tilde{z}} \tilde{\Phi}|_{\Gamma_b} = 0 \end{cases}$$

where a loss of 1/2 derivative for Φ takes place.

Proposition 10. *Let real $\gamma > -1$, integer $k \geq 1$, and $\tilde{\Phi}$ satisfy (52).*

(1) *Let*

$$a_{\Phi_z, l} = \lim_{\tilde{x} \rightarrow -\infty} \alpha^{-\gamma-3} \partial_{\tilde{z}} \tilde{\Phi}, \quad a_{\Phi_z, r} = \lim_{\tilde{x} \rightarrow +\infty} \alpha^{-\gamma-3} \partial_{\tilde{z}} \tilde{\Phi}$$

and

$$a_{\Phi_z}(t, \tilde{z}) = \chi_l a_{\Phi_z, l} + \chi_r a_{\Phi_z, r}.$$

Then the following estimates hold:

$$\|\alpha^{-\gamma-3} \partial_{\tilde{z}} \tilde{\Phi} - a_{\Phi_z}\|_{H^{k+2}} \leq Q(\mathcal{T}', \partial_{\tilde{z}} \tilde{\Phi}, w, k+1/2) [|\alpha^{-\gamma-2} \partial_t \tilde{\zeta} - a_{\zeta_t}|_{H^{k+3/2}} + |a_{\zeta_t, i}|_{L^\infty}]$$

and

$$|a_{\Phi_z, i}|_{W_z^{k+2, \infty}} \leq C |a_{\zeta_t, i}|_{L^\infty}.$$

(2) *Let a_v be defined in (49). One also has*

$$\|\alpha^{-\gamma-2} \tilde{v} - a_v\|_{H^{k+2}} \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, w, k+1/2) [|\alpha^{-\gamma-2} \partial_t \tilde{\zeta} - a_{\zeta_t}|_{H^{k+3/2}} + |a_{\zeta_t, i}|_{L^\infty}],$$

where a_v satisfies

$$|a_v|_{W^{k+2, \infty}} \leq Q_b |a_{\zeta_t, i}|_{L^\infty}.$$

Proof. Step 1. Estimate of $\partial_{\tilde{z}}\tilde{\Phi}$. Similarly as in the above estimate for \mathcal{T} , we introduce

$$w_{\Phi} = \alpha^{-\gamma-3}\partial_{\tilde{z}}\tilde{\Phi} - a_{\Phi_z},$$

which satisfies the following system

$$\begin{cases} \Delta w_{\Phi} = [\Delta, \alpha^{-\gamma-3}]\partial_{\tilde{z}}\tilde{\Phi} - \Delta a_{\Phi_z} & \text{in } \mathcal{S}_0, \\ w_{\Phi}|_{\Gamma_t} = -\alpha^{-\gamma-3}(\partial_{\tilde{x}}x)\tilde{\partial}_t\tilde{\zeta} - a_{\Phi_z}|_{\tilde{z}=0}, & w_{\Phi}|_{\Gamma_b} = 0. \end{cases}$$

Here a_{Φ_z} satisfies corresponding boundary conditions for $i = l, r$ naturally:

$$(53) \quad a_{\Phi_z, l}|_{\tilde{z}=0} = -a_{\zeta_t, l}, \quad a_{\Phi_z, r}|_{\tilde{z}=0} = -b_r a_{\zeta_t, r}, \quad a_{\Phi_z, i}|_{\tilde{z}=-\omega} = 0.$$

Moreover, thanks to Lemma 8 and

$$\Delta\partial_{\tilde{z}}\tilde{\Phi} = 0 \quad \text{in } \mathcal{S}_0,$$

direct computations show that $a_{\Phi_z, l}$ satisfies the following equation

$$(54) \quad \partial_{\tilde{z}}^2 a_{\Phi_z, l} = -(\gamma + 3)^2 a_{\Phi_z, l}.$$

As a result, the ODE system (54) and (53) of $a_{\Phi_z, l}$ can be solved similarly as for $a_{x, i}, a_{z, i}$ from (35), and we have immediately

$$(55) \quad a_{\Phi_z, l} = -a_{\zeta_t, l} \cos((\gamma + 3)\tilde{z}) - a_{\zeta_t, l} \cot((\gamma + 3)\omega) \sin((\gamma + 3)\tilde{z})$$

with the constant C depending on γ, m . Moreover, similar expression and estimate hold for $a_{\Phi_z, r}$ depending on a_{ζ_t} and b_r .

On the other hand, when $\tilde{x} \leq -c_0$ (in order to be more clear), recalling the definition of the weight α from (9) and direct computations again lead to

$$\begin{aligned} \Delta w_{\Phi} &= (\alpha^{-\gamma-3})''\partial_{\tilde{z}}\tilde{\Phi} + 2(\alpha^{-\gamma-3})'\partial_{\tilde{x}}\partial_{\tilde{z}}\tilde{\Phi} - \chi_l'' a_{\Phi_z, l} - \chi_l \partial_{\tilde{z}}^2 a_{\Phi_z, l} \\ &= (\gamma + 3)^2 \alpha^{-\gamma-3} \partial_{\tilde{z}}\tilde{\Phi} - 2(\gamma + 3)\alpha^{-\gamma-3} \partial_{\tilde{x}}\partial_{\tilde{z}}\tilde{\Phi} - \chi_l'' a_{\Phi_z, l} - \chi_l \partial_{\tilde{z}}^2 a_{\Phi_z, l} \\ &= (\gamma + 3)^2 (\alpha^{-\gamma-3} \partial_{\tilde{z}}\tilde{\Phi} - \chi_l a_{\Phi_z, l}) - 2(\gamma + 3)(\alpha^{-\gamma-3} \partial_{\tilde{x}}\partial_{\tilde{z}}\tilde{\Phi} - (\gamma + 3)\chi_{l, 1} a_{\Phi_z, l}) - \chi_l'' a_{\Phi_z, l} \\ &\quad + 2(\gamma + 3)^2 (\chi_l - \chi_{l, 1}) a_{\Phi_z, l} \end{aligned}$$

where (54) is applied for a_{Φ_z} and $\chi_{l, 1}$ is defined in Lemma 8.

As a result, applying Proposition 4, Proposition 1, Corollary 1, Corollary 2 and Lemma 8, we obtain

$$\begin{aligned} \|w_{\Phi}\|_{H^{k+2}} &\leq C(\|\alpha^{-\gamma-3}\partial_{\tilde{z}}\tilde{\Phi} - a_{\Phi_z}\|_{H^k} + \|\alpha^{-\gamma-3}\partial_{\tilde{x}}\partial_{\tilde{z}}\tilde{\Phi} - a_{\Phi_z, 1}\|_{H^k} + \|\chi_l'' a_{\Phi_z}\|_{H^k} \\ &\quad + \|(\chi_l - \chi_{l, 1})a_{\Phi_z}\|_{H^k} + |\alpha^{-\gamma-3}(\partial_{\tilde{x}}x)\tilde{\partial}_t\tilde{\zeta} + a_{\Phi_z}|_{H^{k+3/2}}) \\ &\leq C\|w_{\Phi}\|_{H^{k+1}} + Q(\mathcal{T}', w, k + 1/2)[|\alpha^{-\gamma-2}\tilde{\partial}_t\tilde{\zeta} - a_{\zeta_t}|_{H^{k+3/2}} + |a_{\zeta_t, i}|_{L^\infty}]. \end{aligned}$$

Notice that here one has thanks to (53) that

$$\begin{aligned} &|\alpha^{-\gamma-3}(\partial_{\tilde{x}}x)\tilde{\partial}_t\tilde{\zeta} + a_{\Phi_z}|_{H^{k+3/2}} \\ &\leq |\alpha^{-1}\partial_{\tilde{x}}x\chi_l(\alpha^{-\gamma-2}\tilde{\partial}_t\tilde{\zeta} - a_{\zeta_t, l})|_{H^{k+3/2}} + |\alpha^{-1}\partial_{\tilde{x}}x\chi_r(\alpha^{-\gamma-2}\tilde{\partial}_t\tilde{\zeta} - a_{\zeta_t, r})|_{H^{k+3/2}} \\ &\quad + |a_{\zeta_t, l}\chi_l(\alpha^{-1}\partial_{\tilde{x}}x - 1)|_{H^{k+3/2}} + |a_{\zeta_t, r}\chi_r(\alpha^{-1}\partial_{\tilde{x}}x - b_r)|_{H^{k+3/2}} + |(1 - \chi_l - \chi_r)\alpha^{-\gamma-3}(\partial_{\tilde{x}}x)\tilde{\partial}_t\tilde{\zeta}|_{H^{k+3/2}} \\ &\leq Q(\mathcal{T}', w, k + 1/2)[|\alpha^{-\gamma-2}\tilde{\partial}_t\tilde{\zeta} - a_{\zeta_t}|_{H^{k+3/2}} + |a_{\zeta_t, i}|_{L^\infty}], \end{aligned}$$

where we recall (35) and (37) for the limits of $\alpha^{-1}\partial_{\tilde{x}}x$.

Consequently, applying an interpolation to $\|w_{\Phi}\|_{H^k}$, we derive the desired estimate for $\partial_{\tilde{z}}\tilde{\Phi}$.

Step 2. Estimate of $\partial_{\tilde{x}}\tilde{\Phi}$ and \tilde{v} . In fact, using the equation of $\tilde{\Phi}$ and applying the estimate of $\partial_{\tilde{z}}\tilde{\Phi}$, we can have

$$\|\alpha^{-\gamma-3}\partial_{\tilde{x}}\tilde{\Phi} - a_{\Phi_x}\|_{H^{k+2}} \leq Q(\mathcal{T}', \nabla\tilde{\Phi}, w, k + 1/2)[|\alpha^{-\gamma-2}\tilde{\partial}_t\tilde{\zeta} - a_{\zeta_t}|_{H^{k+3/2}} + |a_{\zeta_t, i}|_{L^\infty}]$$

where

$$a_{\Phi_x} = \chi_l a_{\Phi_x} + \chi_r a_{\Phi_x, r}$$

with

$$a_{\Phi_x, l} = \lim_{\tilde{x} \rightarrow -\infty} \alpha^{-\gamma-3} \partial_{\tilde{x}} \tilde{\Phi}, \quad a_{\Phi_x, r} = \lim_{\tilde{x} \rightarrow +\infty} \alpha^{-\gamma-3} \partial_{\tilde{x}} \tilde{\Phi}$$

satisfying

$$a_{\Phi_x, l} = -\frac{1}{\gamma+3} \partial_{\tilde{z}} a_{\Phi_z, l}, \quad a_{\Phi_x, r} = -\frac{1}{(\gamma+3)\beta_r} \partial_{\tilde{z}} a_{\Phi_z, r}.$$

In the end, for the estimate of the velocity v , recalling that $\tilde{v} = \widetilde{\nabla \Phi}$ and applying (25) lead to

$$\tilde{v} = (\nabla_X \tilde{X}) \circ \mathcal{T} \nabla \tilde{\Phi} = |\mathcal{T}'|^{-2} \begin{pmatrix} \partial_{\tilde{x}} x \partial_{\tilde{x}} \tilde{\Phi} + \partial_{\tilde{x}} z \partial_{\tilde{z}} \tilde{\Phi} \\ \partial_{\tilde{x}} z \partial_{\tilde{x}} \tilde{\Phi} - \partial_{\tilde{x}} x \partial_{\tilde{z}} \tilde{\Phi} \end{pmatrix},$$

which implies

(56)

$$\begin{aligned} a_{v, l} &= \lim_{\tilde{x} \rightarrow -\infty} \alpha^{-\gamma-2} \tilde{v} = \lim_{\tilde{x} \rightarrow -\infty} \frac{1}{\alpha^{-2} (\partial_{\tilde{x}} x)^2 + \alpha^{-2} (\partial_{\tilde{x}} z)^2} \begin{pmatrix} \alpha^{-1} \partial_{\tilde{x}} x \alpha^{-\gamma-3} \partial_{\tilde{x}} \tilde{\Phi} + \alpha^{-1} \partial_{\tilde{x}} z \alpha^{-\gamma-3} \partial_{\tilde{z}} \tilde{\Phi} \\ \alpha^{-1} \partial_{\tilde{x}} z \alpha^{-\gamma-3} \partial_{\tilde{x}} \tilde{\Phi} - \alpha^{-1} \partial_{\tilde{x}} x \alpha^{-\gamma-3} \partial_{\tilde{z}} \tilde{\Phi} \end{pmatrix} \\ &= \frac{1}{a_{x, l}^2 + a_{z, l}^2} \begin{pmatrix} a_{x, l} a_{\Phi_x, l} + a_{z, l} a_{\Phi_z, l} \\ a_{z, l} a_{\Phi_x, l} - a_{x, l} a_{\Phi_z, l} \end{pmatrix}. \end{aligned}$$

Moreover, one has a similar expression for $a_{v, r}$.

As a result, we can have the $W^{k, \infty}$ estimate of $a_{v, i}$ thanks to (40) and (55). Applying Corollary 2, we obtain the desired estimate for \tilde{v} . \square

Second, we obtain a higher order estimate for \tilde{v} near the bottom Γ_b thanks to Corollary 4.

Corollary 5. *Under the same assumption of Proposition 10, one has*

$$\|\alpha^{-\gamma-2} \tilde{v} - a_v\|_{H^{k+5/2}(\mathcal{S}_b)} \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, w, k) [\|\alpha^{-\gamma-2} \partial_t \tilde{\zeta} - a_{\zeta_t}\|_{H^{k+1}} + |a_{\zeta_t, i}|_{L^\infty}],$$

where $\mathcal{S}_b \subset \mathcal{S}_0$ is the flat strip near Γ_b .

Proof. The proof is similar as the proof of Corollary (4). We first prove the estimate of $\partial_{\tilde{z}} \tilde{\Phi}$ and use again $w_\Phi = \alpha^{-\gamma-3} \partial_{\tilde{z}} \tilde{\Phi} - a_{\Phi_z}$ from the proof of Proposition 10.

In fact, recalling the cut-off function χ_b near Γ_b from Corollary (4), $\chi_b w_\Phi$ satisfies

$$\begin{cases} \Delta(\chi_b w_\Phi) = [\Delta, \chi_b] w_\Phi - \chi_b \Delta w_\Phi & \text{in } \mathcal{S}_0, \\ \chi_b w_\Phi|_{\Gamma_t} = 0, \quad \chi_b w_\Phi|_{\Gamma_b} = 0. \end{cases}$$

Applying Proposition 10 and the expression of Δw_Φ in that proof, we have

$$\begin{aligned} \|\chi_b w_\Phi\|_{H^{k+5/2}} &\leq C(\|w_\Phi\|_{H^{k+3/2}} + \|\chi_b \Delta w_\Phi\|_{H^{k+1/2}}) \\ &\leq C(\|w_\Phi\|_{H^{k+3/2}} + |a_{\zeta_t, i}|_{L^\infty}) \\ &\leq Q(\mathcal{T}', \partial_{\tilde{z}} \tilde{\Phi}, w, k) [|a_{\zeta_t, i}|_{L^\infty} + \|\alpha^{-\gamma-2} \partial_t \tilde{\zeta} - a_{\zeta_t}\|_{H^{k+1}}]. \end{aligned}$$

Moreover, similar analysis as before, we also derive a similar estimate for $\partial_{\tilde{x}} \tilde{\Phi}$ near Γ_b . Consequently, thanks to the higher order estimate of \mathcal{T}' in Corollary (4) and the expression of \tilde{v} in terms of $\mathcal{T}', \nabla \tilde{\Phi}$ in the proof of Proposition 10, we conclude immediately the desired estimate for \tilde{v} . \square

Third, we present the higher order estimate for \tilde{v} .

Proposition 11. *Under the same assumption of Proposition 10, one has*

$$\|\alpha^{-\gamma-2} \tilde{v} - a_v\|_{H^{k+5/2}} \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \partial_t \tilde{\zeta}, w, k+1/2) [1 + \|\alpha^{-\gamma-2} \tilde{D}_t w - a_{w_t}\|_{H^k} + |a_{\zeta_t, i}|_{L^\infty}].$$

Proof. The proof adopts the idea of the proof for Proposition 4.3 [57], while the key point is the estimate of the boundary condition $\nabla_{n_t} v|_{\Gamma_t}$. As mentioned in the introduction, the key difference compared to the smooth-boundary estimates lies in the estimate for the higher-order terms in the bottom.

Step 1. Decomposition of the boundary condition on Γ_t for v . In fact, remembering that the fluid is irrotational (see (2)), we rewrite

$$(57) \quad \nabla_{n_t} v = (\nabla v)^T n_t := u_v \quad \text{on } \Gamma_t.$$

Decomposing u_v into

$$u_v = u_v^\top + u_v^\perp n_t := (u_v \cdot \tau_t) \tau_t + (u_v \cdot n_t) n_t,$$

we deal with the estimate of u_v^\top and u_v^\perp one by one.

Step 2. The tangential part u_v^\top . Direct computations show that

$$(\Delta_{\Gamma_t} u_v^\top) \cdot \tau_t = (\nabla_{\tau_t} \nabla_{\tau_t} u_v^\top) \cdot \tau_t = \nabla_{\tau_t} (\mathcal{D}_{\tau_t} u_v^\top \cdot \tau_t) = \nabla_{\tau_t} (\mathcal{D} \cdot u_v^\top),$$

where \mathcal{D} is the covariant (tangential) derivative on Γ_t . Therefore, we have

$$-\Delta_{\Gamma_t} u_v^\top = -\nabla_{\tau_t} (\mathcal{D} \cdot u_v^\top) \tau_t \quad \text{on } \Gamma_t.$$

On the other hand, we also know that

$$\begin{aligned} \mathcal{D} \cdot u_v^\top &= \nabla_{\tau_t} u_v^\top \cdot \tau_t = \nabla_{\tau_t} [(n_t \cdot (\nabla v) \tau_t) \tau_t] \cdot \tau_t \\ &= \nabla_{\tau_t} (n_t \cdot (\nabla v) \tau_t) = \nabla_{\tau_t} (n_t \cdot \nabla_{\tau_t} v) = \nabla_{\tau_t} \nabla_{\tau_t} v \cdot n_t + \nabla_{\tau_t} v \cdot \nabla_{\tau_t} n_t. \end{aligned}$$

Since

$$\Delta_{\Gamma_t} = \nabla_{\tau_t} \nabla_{\tau_t} - \nabla_{\mathcal{D}_{\tau_t} \tau_t} = \nabla_{\tau_t} \nabla_{\tau_t}$$

in our 2D settings, substituting (18) into the above expression of $\mathcal{D} \cdot u_v^\top$ leads to

$$\mathcal{D} \cdot u_v^\top = -D_t \kappa - \nabla_{\tau_t} n_t \cdot \nabla_{\tau_t} v \quad \text{on } \Gamma_t.$$

Consequently, we derive the following equation

$$-\Delta_{\Gamma_t} u_v^\top = \nabla_{\tau_t} (D_t \kappa) \tau_t + \nabla_{\tau_t} (\nabla_{\tau_t} n_t \cdot \nabla_{\tau_t} v) \tau_t \quad \text{on } \Gamma_t,$$

which is an elliptic equation on Γ_t and can be handled similar as before.

Performing \mathcal{T} on both sides of the above equation, we arrive at

$$-((\partial_{\tilde{x}} x)^{-1} b_\zeta^{-\frac{1}{2}} \partial_{\tilde{x}})^2 \widetilde{u}_v^\top = (\partial_{\tilde{x}} x)^{-1} b_\zeta^{-\frac{1}{2}} \partial_{\tilde{x}} (\widetilde{D}_t w) \widetilde{\tau}_t + (\partial_{\tilde{x}} x)^{-1} b_\zeta^{-\frac{1}{2}} \partial_{\tilde{x}} (\widetilde{\nabla_{\tau_t} n_t} \cdot \widetilde{\nabla_{\tau_t} v}) \widetilde{\tau}_t$$

on the upper boundary of \mathcal{S}_0 , where b_ζ is defined in (45).

As a result, similar analysis as in the proof of Proposition 9, we have immediately the following elliptic estimate

$$(58) \quad |\alpha^{-\gamma-1} \widetilde{u}_v^\top - a_{u_v, \tau}|_{H^{k+1}} \leq Q(\mathcal{T}', w, k) (|\alpha^{-\gamma} \widetilde{D}_t w - a_{w_t}|_{H^k} + |\alpha^{-\gamma-2} \widetilde{v} - a_v|_{H^{k+1}} + |a_{\zeta_t, i}|_{L^\infty}),$$

where the weighted limit $a_{u_v, \tau}$ is defined for \widetilde{u}_v^\top in a similar way as before and can be handled (depending on $a_{\zeta_t, i}$). Here we omit the analysis for weighted limits since they are decided by the corresponding equations and can always be handled algebraically as before.

Step 3. The normal part u_v^\perp . Since

$$u_v^\perp = u_v \cdot n_t = n_t \cdot \nabla_{n_t} v,$$

we extend n_t as \bar{n}_t in Ω (to be fixed later) and denote by \bar{u}_v^\perp the extension of u_v^\perp with \bar{n}_t .

Direct computations show that

$$\begin{aligned} \nabla \cdot u_v &= \nabla_{\tau_t} u_v \cdot \tau_t + \nabla_{n_t} u_v \cdot n_t \\ &= (\nabla_{\tau_t} \nabla_{\tau_t} v + \nabla_{n_t} \nabla_{\bar{n}_t} v) \cdot n_t + \nabla_{\tau_t} v \cdot \nabla_{\tau_t} n_t + \nabla_{n_t} v \cdot \nabla_{n_t} \bar{n}_t - n_t \cdot \nabla v (\nabla_{\tau_t} \tau_t + \nabla_{n_t} \bar{n}_t) \\ &= (\Delta v) \cdot n_t + r(\partial v, \partial n_t, \partial \tau_t) = r(\partial v, \partial \bar{n}_t, \partial \tau_t), \end{aligned}$$

where the equation $\Delta v = 0$ is applied and

$$r(\partial v, \partial \bar{n}_t, \partial \tau_t) = \nabla_{\tau_t} v \cdot \nabla_{\tau_t} n_t + \nabla_{n_t} v \cdot \nabla_{n_t} \bar{n}_t - 2n_t \cdot \nabla v (\nabla_{\tau_t} \tau_t + \nabla_{n_t} \bar{n}_t)$$

is the lower-order part.

Moreover, we also have on Γ_t that

$$\begin{aligned}\nabla \cdot u_v &= \nabla_{\tau_t} u_v \cdot \tau_t + \nabla_{n_t} u_v \cdot n_t \\ &= \nabla_{\tau_t} u_v^\top \cdot \tau_t + \nabla_{\tau_t} (u^\perp n_t) \cdot \tau_t + \nabla_{n_t} u_v \cdot n_t \\ &= \mathcal{D} \cdot u_v^\top + \kappa u_v^\perp + \nabla_{n_t} u_v \cdot n_t.\end{aligned}$$

Therefore, summing up these expressions above leads to

$$(59) \quad \nabla_{n_t} u_v \cdot n_t = -\mathcal{D} \cdot u_v^\top - \kappa u_v^\perp + r(\partial v, \partial n_t, \partial \tau_t).$$

Next, we plan to derive $\mathcal{N}u_v^\perp$ from the left side of (59) as in [57], which will be adapted to our weighted version in \mathcal{S}_0 .

In fact, we start with rewriting the left side as

$$\nabla_{n_t} u_v \cdot n_t = \nabla_{n_t} (\bar{n}_t \cdot \nabla_{\bar{n}_t} v) - (\nabla v)^T n_t \cdot \nabla_{n_t} \bar{n}_t.$$

Substituting it into (59), we obtain

$$\nabla_{n_t} \bar{u}_v^\perp = \nabla_{n_t} (\bar{n}_t \cdot \nabla_{\bar{n}_t} v) = -\mathcal{D} \cdot u_v^\top + r_1(\partial v, \partial \bar{n}_t, \partial \tau_t),$$

where

$$r_1(\partial v, \partial \bar{n}_t, \partial \tau_t) = (\nabla v)^T n_t \cdot \nabla_{n_t} \bar{n}_t - \kappa u_v^\perp + r(\partial v, \partial \bar{n}_t, \partial \tau_t).$$

Performing \mathcal{T} on both sides of the above equation and noticing that

$$\widetilde{\nabla_{n_t} F} = b_\zeta^{-\frac{1}{2}} (\partial_{\bar{x}x})^{-1} \partial_{\bar{z}} \widetilde{F} \quad \text{on } \Gamma_t,$$

we have

$$(60) \quad b_\zeta^{-\frac{1}{2}} (\partial_{\bar{x}x})^{-1} \partial_{\bar{z}} \widetilde{\bar{u}_v^\perp} = -\widetilde{\mathcal{D} \cdot u_v^\top} + r_1(\widetilde{\partial v}, \widetilde{\partial \bar{n}_t}, \widetilde{\partial \tau_t}) \quad \text{on } \Gamma_t.$$

Denoting by $a_{\bar{u}_v, n}$ the weighted limit of $\widetilde{\bar{u}_v^\perp}$ and $a_{u_v, n}$ the weighted limit of $\widetilde{u_v^\perp}$ on the boundary, the left side above is rewritten into

$$\begin{aligned}b_\zeta^{-\frac{1}{2}} (\partial_{\bar{x}x})^{-1} \partial_{\bar{z}} \widetilde{\bar{u}_v^\perp} &= \alpha^{\gamma+1} b_\zeta^{-\frac{1}{2}} (\partial_{\bar{x}x})^{-1} \partial_{\bar{z}} (\alpha^{-\gamma-1} \widetilde{\bar{u}_v^\perp} - a_{\bar{u}_v, n}) + \alpha^{\gamma+1} b_\zeta^{-\frac{1}{2}} (\partial_{\bar{x}x})^{-1} \partial_{\bar{z}} a_{\bar{u}_v, n} \\ &= \alpha^{\gamma+1} b_\zeta^{-\frac{1}{2}} (\partial_{\bar{x}x})^{-1} [\widetilde{\mathcal{N}_0} (\alpha^{-\gamma-1} \widetilde{\bar{u}_v^\perp} - a_{\bar{u}_v, n}) + \partial_{\bar{z}} w_v] + \alpha^{\gamma+1} b_\zeta^{-\frac{1}{2}} (\partial_{\bar{x}x})^{-1} \partial_{\bar{z}} a_{\bar{u}_v, n}.\end{aligned}$$

Here

$$w_v = (\alpha^{-\gamma-1} \widetilde{\bar{u}_v^\perp} - a_{\bar{u}_v, n}) - \mathcal{H}(\alpha^{-\gamma-1} \widetilde{u_v^\perp} - a_{u_v, n})$$

satisfies

$$\begin{cases} \Delta w_v = \Delta(\alpha^{-\gamma-1} \widetilde{\bar{u}_v^\perp}) - \Delta a_{\bar{u}_v, n} = \Delta(\alpha^{-\gamma-1} \widetilde{\bar{n}_t} \cdot \widetilde{\nabla_{\bar{n}_t} v}) - \Delta a_{\bar{u}_v, n} & \text{in } \mathcal{S}_0, \\ w_v|_{\bar{z}=0} = 0, \\ \partial_{\bar{z}} w_v|_{\bar{z}=-\omega} = \partial_{\bar{z}} (\alpha^{-\gamma-1} \widetilde{\bar{n}_t} \cdot \widetilde{\nabla_{\bar{n}_t} v})|_{\bar{z}=-\omega} - \partial_{\bar{z}} a_{\bar{u}_v, n}|_{\bar{z}=-\omega} := g(\partial^2 \widetilde{v}, \partial \widetilde{\bar{n}_t}). \end{cases}$$

Substituting this expression into (60), we conclude that

$$\widetilde{\mathcal{N}_0} (\alpha^{-\gamma-1} \widetilde{\bar{u}_v^\perp} - a_{\bar{u}_v, n}) = -\partial_{\bar{z}} w_v - \partial_{\bar{z}} a_{\bar{u}_v, n} - \alpha^{-\gamma-1} b_\zeta^{\frac{1}{2}} (\partial_{\bar{x}x}) [\widetilde{\mathcal{D} \cdot u_v^\top} - r_1(\widetilde{\partial v}, \widetilde{\partial \bar{n}_t}, \widetilde{\partial \tau_t})].$$

As a result, we perform the following estimate

$$|\widetilde{\mathcal{N}_0} (\alpha^{-\gamma-1} \widetilde{\bar{u}_v^\perp} - a_{\bar{u}_v, n})|_{H^k} \leq |\partial_{\bar{z}} w_v|_{H^k} + |\partial_{\bar{z}} a_{\bar{u}_v, n} + \alpha^{-\gamma-1} b_\zeta^{\frac{1}{2}} (\partial_{\bar{x}x}) [\widetilde{\mathcal{D} \cdot u_v^\top} - r_1(\widetilde{\partial v}, \widetilde{\partial \bar{n}_t}, \widetilde{\partial \tau_t})]|_{H^k}.$$

Since the corresponding weighted limits for $\widetilde{u_v^\perp}$ and \widetilde{v} can be retrieved from $\partial_{\bar{z}} a_{\bar{u}_v, n}$ on the right side above and the analysis is omitted again, we derive by Lemma 3, Proposition 1 and (58) that

$$(61) \quad \begin{aligned} & |\widetilde{\mathcal{N}_0} (\alpha^{-\gamma-1} \widetilde{\bar{u}_v^\perp} - a_{\bar{u}_v, n})|_{H^k} \\ & \leq C \|w_v\|_{H^{k+3/2}} + Q(\mathcal{T}', w, k) (|\alpha^{-\gamma-1} \widetilde{u_v^\perp} - a_{u_v, \tau}|_{H^{k+1}} + \|\alpha^{-\gamma-2} \widetilde{v} - a_v\|_{H^{k+3/2}}) \\ & \leq C \|w_v\|_{H^{k+3/2}} + Q(\mathcal{T}', w, k) (|\alpha^{-\gamma} \widetilde{D}_t w - a_{w_t}| + \|\alpha^{-\gamma-2} \widetilde{v} - a_v\|_{H^{k+3/2}} + |a_{\zeta_t, i}|_{L^\infty}). \end{aligned}$$

Moreover, similar analysis as in the proof of Proposition 10, one has the following elliptic estimate

$$\begin{aligned} \|w_v\|_{H^{k+3/2}} &\leq C(\|\Delta w_v\|_{H^{k-1/2}} + |\partial_{\bar{z}} w_v|_{H^k(\Gamma_b)}) \\ &\leq Q(\mathcal{T}', w, k+1/2)(\|\alpha^{-\gamma-2}\tilde{v} - a_v\|_{H^{k+3/2}} + |a_{\zeta_t, i}|_{L^\infty} + \|\alpha^{-\gamma-2}\tilde{v} - a_v\|_{H^{k+5/2}(\mathcal{S}_b)}). \end{aligned}$$

Applying the high-order derivative estimate of \tilde{v} in Corollary 5 and lower-order derivative estimate in Proposition 10, we derive

$$\|w_v\|_{H^{k+3/2}} \leq Q(\mathcal{T}', w, k+1/2)(|a_{\zeta_t, i}|_{L^\infty} + |\alpha^{-\gamma-2}\tilde{\partial}_t\zeta - a_{\zeta_t}|_{H^{k+1}}).$$

Substituting this estimate back into (61), we conclude that

$$|\tilde{\mathcal{N}}_0(\alpha^{-\gamma-1}\tilde{u}_v^\perp - a_{\tilde{u}_v, n})|_{H^k} \leq Q(\mathcal{T}', w, k+1/2)(|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} + |a_{\zeta_t, i}|_{L^\infty} + |\alpha^{-\gamma-2}\tilde{\partial}_t\zeta - a_{\zeta_t}|_{H^{k+1}}),$$

which implies immediately the desired estimate for the normal component u_v^\perp :

$$\begin{aligned} (62) \quad |\alpha^{-\gamma-1}\tilde{u}_v^\perp - a_{u_v, n}|_{H^{k+1}} &\leq C(|\tilde{\mathcal{N}}_0(\alpha^{-\gamma-1}\tilde{u}_v^\perp - a_{\tilde{u}_v, n})|_{H^k} + |\alpha^{-\gamma-1}\tilde{u}_v^\perp - a_{u_v, n}|_{L^2}) \\ &\leq Q(\mathcal{T}', w, k+1/2)(|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} + |a_{\zeta_t, i}|_{L^\infty} + |\alpha^{-\gamma-2}\tilde{\partial}_t\zeta - a_{\zeta_t}|_{H^{k+1}}). \end{aligned}$$

Step 4. Elliptic system for \tilde{v} and the estimate. In the end, we are ready to show the desired higher-order derivative estimate for \tilde{v} . In fact, the system for $\alpha^{-\gamma-2}\tilde{v} - a_v$ reads

$$\begin{cases} \Delta(\alpha^{-\gamma-2}\tilde{v} - a_v) = \Delta(\alpha^{-\gamma-2}\tilde{v}) - \Delta a_v & \text{in } \mathcal{S}_0, \\ \partial_{\bar{z}}(\alpha^{-\gamma-2}\tilde{v} - a_v)|_{\bar{z}=0} = \alpha^{-\gamma-2}b_\zeta^{\frac{1}{2}}(\partial_{\bar{x}}x)\tilde{u}_v - \partial_{\bar{z}}a_v|_{\bar{z}=0}, \\ \alpha^{-\gamma-2}\tilde{v} - a_v|_{\bar{z}=-\omega} & \text{is done in Corollary 5.} \end{cases}$$

Meanwhile on the upper surface, we have the following estimate by summing up (58) and (62):

$$|\alpha^{-\gamma-1}\tilde{u}_v - a_{u_v}|_{H^{k+1}} \leq Q(\mathcal{T}', w, k+1/2)(|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} + |a_{\zeta_t, i}|_{L^\infty} + |\alpha^{-\gamma-2}\tilde{\partial}_t\zeta - a_{\zeta_t}|_{H^{k+1}}),$$

where a_{u_v} is the corresponding weighted limit of \tilde{u}_v .

As a result, similarly as in the proof Proposition 10 and thanks to Corollary 5, we conclude estimate for \tilde{v} :

$$\begin{aligned} \|\alpha^{-\gamma-2}\tilde{v} - a_v\|_{H^{k+5/2}} &\leq C(\|\Delta(\alpha^{-\gamma-2}\tilde{v}) - \Delta a_v\|_{H^{k+1/2}} + |\alpha^{-\gamma-2}b_\zeta^{\frac{1}{2}}(\partial_{\bar{x}}x)\tilde{u}_v - \partial_{\bar{z}}a_v|_{\bar{z}=0}|_{H^{k+1}} + |\alpha^{-\gamma-2}\tilde{v} - a_v|_{H^{k+2}(\Gamma_b)}) \\ &\leq Q(\mathcal{T}', w, k)(\|\Delta(\alpha^{-\gamma-2}\tilde{v}) - \Delta a_v\|_{H^{k+1/2}} + |\alpha^{-\gamma-1}\tilde{u}_v - a_{u_v}|_{H^{k+1}} + \|\alpha^{-\gamma-2}\tilde{v} - a_v\|_{H^{k+5/2}(\mathcal{S}_b)}) \\ &\leq Q(\mathcal{T}', w, k+1/2)(|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} + |a_{\zeta_t, i}|_{L^\infty} + |\alpha^{-\gamma-2}\tilde{\partial}_t\zeta - a_{\zeta_t}|_{H^{k+1}}). \end{aligned}$$

Moreover, applying Proposition 9 to the right-side term $|\alpha^{-\gamma-2}\tilde{\partial}_t\zeta - a_{\zeta_t}|_{H^{k+1}}$ and using an interpolation on \tilde{v} , we obtain

$$\begin{aligned} \|\alpha^{-\gamma-2}\tilde{v} - a_v\|_{H^{k+5/2}} &\leq Q(\mathcal{T}', w, k+1/2)(|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} + |a_{\zeta_t, i}|_{L^\infty} + \|\alpha^{-\gamma-2}\tilde{v} - a_v\|_{H^{k+3/2}} \\ &\quad + |\alpha^{-\gamma-2}\tilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2}). \end{aligned}$$

Therefore, the proof is finished. \square

In the end, we deal with \tilde{P} and \mathbf{a} . Recalling Assumption 2 and system (32) for \tilde{P} (especially the boundary conditions), we know immediately that $\alpha^{-2\gamma-4}\tilde{P}$ should not vanish when $\tilde{x} \rightarrow \pm\infty$. Therefore, setting

$$(63) \quad a_p = \chi_l a_{p, l} + \chi_r a_{p, r}$$

with

$$a_{p, l}(t, \tilde{z}) = \lim_{\tilde{x} \rightarrow -\infty} \alpha^{-2\gamma-4}\tilde{P}, \quad a_{p, r}(t, \tilde{z}) = \lim_{\tilde{x} \rightarrow +\infty} \alpha^{-2\gamma-4}\tilde{P},$$

we obtain from (32) the following system for $a_{p,l}$:

$$(64) \quad \begin{cases} \partial_{\bar{z}}^2 a_{p,l} = -(2\gamma + 4)^2 a_{p,l} - a_{Dv^2}, \\ a_{p,l}|_{\bar{z}=0} = 0, \quad \partial_{\bar{z}} a_{p,l}|_{\bar{z}=-\omega} = 0 \end{cases}$$

where

$$a_{Dv^2} = \lim_{\bar{x} \rightarrow -\infty} \alpha^{-2\gamma-4} |\mathcal{T}'|^2 \text{tr}((\nabla_X \tilde{X}) \circ \mathcal{T} \nabla \tilde{v})^2$$

is the combination of weighted limits for $\alpha^{-2\gamma-4} |\mathcal{T}'|^2 \text{tr}((\nabla_X \tilde{X}) \circ \mathcal{T} \nabla \tilde{v})^2$ defined in a similar way as before. The computations in a_{Dv^2} are similar as in (56) and is hence omitted here. Consequently, this limit depends on $T_{1,c}$, a_{Φ_z} . Moreover, we know

$$\partial_{\bar{z}} a_{p,l}|_{\bar{z}=0} < 0$$

thanks to Assumption 2. The analysis for $a_{p,r}$ follows similarly.

Proposition 12. *Let real $\gamma > -1$, integer $k \geq 0$ and \tilde{P} solve (32). Then there holds:*

$$\|\alpha^{-2\gamma-4} \tilde{P} - a_p\|_{H^{k+7/2}} \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \tilde{\partial}_t \zeta, \tilde{P}, w, k+1/2) (1 + |\alpha^{-\gamma} \tilde{D}_t w - a_{w_t}|_{H^k}^2 + |a_{\zeta_t, i}|_{L^\infty}^2)$$

where

$$|a_{p,i}|_{W_{\bar{z}}^{k+2, \infty}} \leq Q_b |a_{\zeta_t, i}(t)|^2.$$

Proof. The proof is similar as before. To begin with, we define

$$w_p = \alpha^{-2\gamma-4} \tilde{P} - a_p,$$

which satisfies

$$\begin{cases} \Delta w_p = [\Delta, \alpha^{-2\gamma-4}] \tilde{P} + \alpha^{-2\gamma-4} \Delta \tilde{P} - \Delta a_p & \text{in } \mathcal{S}_0, \\ w_p|_{\Gamma_t} = 0, \quad \partial_{\bar{z}} w_p|_{\Gamma_b} = \alpha^{-2\gamma-4} \partial_{\bar{z}} \tilde{P}|_{\Gamma_b}. \end{cases}$$

Here the right side above can be computed in a similar way as in the system of w_Φ from the proof of Proposition 10, where we apply (32), (64) and only focus on “the left part”:

$$\begin{aligned} \Delta w_p = & ((\alpha^{-2\gamma-4})'' \tilde{P} - (2\gamma + 4)^2 \chi_l a_{p,l}) - 2(-(\alpha^{-2\gamma-4})' \partial_{\bar{x}} \tilde{P} - (2\gamma + 4)^2 \chi_{l,1} a_{p,l}) \\ & - [\alpha^{-2\gamma-4} |\mathcal{T}'|^2 \text{tr}((\nabla_X \tilde{X}) \circ \mathcal{T} \nabla \tilde{v})^2 + \chi_l (\partial_{\bar{z}}^2 a_{p,l} + (2\gamma + 4)^2 a_{p,l})] \\ & + 2(2\gamma + 4)^2 (\chi_l - \chi_{l,1}) a_{p,l} - \chi_l'' a_{p,l} \end{aligned}$$

Applying standard elliptic estimate, Lemma 8 and (64) leads to

$$\begin{aligned} \|w_p\|_{H^{k+7/2}} &= \|\alpha^{-2\gamma-4} \tilde{P} - a_p\|_{H^{k+7/2}} \\ &\leq C(\|\alpha^{-2\gamma-4} \tilde{P} - a_p\|_{H^{k+5/2}} + |a_{p,i}|_{W_{\bar{z}}^{k+2, \infty}} + \|\alpha^{-2\gamma-4} |\mathcal{T}'|^2 \text{tr}((\nabla_X \tilde{X}) \circ \mathcal{T} \nabla \tilde{v})^2 - a_{Dv^2}\|_{H^{k+3/2}} \\ &\quad + |\alpha^{-2\gamma-4} |\mathcal{T}'| ((\nabla_X \tilde{X})^t \circ \mathcal{T}) \tilde{v} \cdot \nabla \tilde{n}_b \cdot \tilde{v}|_{H^{k+2}(\Gamma_b)}), \end{aligned}$$

where we recall that a_{Dv^2} is defined in (64) and notice that the highest-order derivatives for both \mathcal{T} and \tilde{v} appear in the bottom part.

Applying (25), Proposition 1, Corollary 2 and Corollary 4, one finds

$$\|\alpha^{-2\gamma-4} \tilde{P} - a_p\|_{H^{k+7/2}} \leq C(\|\alpha^{-2\gamma-4} \tilde{P} - a_p\|_{H^{k+5/2}} + |a_{p,i}|_{W_{\bar{z}}^{k+2, \infty}}) + Q(\mathcal{T}', w, k) \|\alpha^{-\gamma-2} \tilde{v} - a_v\|_{H^{k+5/2}}^2$$

As a result, the desired estimate for \tilde{P} follows by applying Proposition 11.

In the end, applying standard one dimensional elliptic estimate to system (64), we have the estimate of $a_{p,l}$ thanks to the estimate (55) for a_{Φ_z} . Therefore, the desired estimate for $a_{p,i}$ follows. \square

Based on the weighted estimate for \tilde{P} , we are able to prove the estimate for $\tilde{\mathbf{a}}$, $\partial_t \tilde{\mathbf{a}}$.

Proposition 13. *Let real $\gamma > -1$ and integer $k \geq 0$. One has*

$$|\alpha^{-2\gamma-3}\tilde{\mathbf{a}} - \partial_{\bar{z}}a_p|_{H^{k+3/2}} \leq Q(\mathcal{T}', \nabla\tilde{\Phi}, \tilde{\partial}_t\tilde{\zeta}, \tilde{P}, w, k+1/2)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^{k-1/2}}^2 + |a_{\zeta_t, i}|_{L^\infty}^2).$$

Moreover, the following estimate for $\partial_t\tilde{\mathbf{a}}$ holds:

$$|\alpha^{-2\gamma-3}\partial_t\tilde{\mathbf{a}}|_{L^\infty} \leq Q(\mathcal{T}', \nabla\tilde{\Phi}, \tilde{\partial}_t\tilde{\zeta}, \tilde{P}, w, 3)Q(|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{L^2}, |a_{\zeta_t, i}|_{L^\infty}).$$

Proof. As the first step, direct computations show that

$$\tilde{\mathbf{a}} = -(\nabla_{n_t}P|_{\Gamma_t}) \circ \mathcal{T} = -\tilde{n}_t \cdot (\nabla_X\tilde{X}) \circ \mathcal{T}\nabla\tilde{P}|_{\tilde{z}=0} = -|\mathcal{T}'|^{-1}\partial_{\tilde{z}}\tilde{P}|_{\tilde{z}=0},$$

so we know immediately from Lemma 3, Proposition 1, Corollary 2 and (27) that

$$\begin{aligned} |\alpha^{-2\gamma-3}\tilde{\mathbf{a}} - \partial_{\bar{z}}a_p|_{H^{k+3/2}} &\leq Q(\mathcal{T}', w, k+1/2)(|\alpha^{-2\gamma-4}\partial_{\tilde{z}}\tilde{P} - \partial_{\bar{z}}a_p|_{H^{k+3/2}} + |a_{p, i}|_{W_{\tilde{z}}^{k+3, \infty}}) \\ &\leq Q(\mathcal{T}', w, k+1/2)(\|\alpha^{-2\gamma-4}\tilde{P} - a_p\|_{H^{k+3}} + |a_{\zeta_t, i}|_{L^\infty}). \end{aligned}$$

Therefore, applying Proposition 12 with the order $k+3$ leads to the desired estimate for $\tilde{\mathbf{a}}$.

Second, we have by (30) that

$$\partial_t\tilde{\mathbf{a}} = -(D_t\nabla_{n_t}P) \circ \mathcal{T}|_{\Gamma_t} - u_1\partial_{\tilde{x}}\tilde{\mathbf{a}}, \quad u_1 = (\partial_{\tilde{x}}x)^{-1}(\tilde{v}_1 - \partial_t x)$$

where direct computations lead to

$$\begin{aligned} D_t\nabla_{n_t}P &= [D_t, \nabla_{n_t}]P + \nabla_{n_t}D_tP \\ &= (D_t n_t \cdot \nabla P - n_t \cdot \nabla v \cdot \nabla P) + \nabla_{n_t}\Delta^{-1}(h_1, g_1) + \nabla_{n_t}\Delta^{-1}(h_2, g_2) \end{aligned}$$

where

$$\begin{aligned} h_1 &= -tr[(-\nabla^2P - (\nabla v)^T\nabla v)\nabla v + \nabla v(-\nabla^2P - (\nabla v)^T\nabla v)], \\ g_1 &= -\nabla_v n_b \cdot (\nabla P + \mathbf{g}) + \nabla_{(\nabla P + \mathbf{g})}n_b \cdot v - v \cdot \nabla v \cdot \nabla n_b \cdot v - \nabla_v((\nabla v)^T n_b)^\top \cdot v, \\ h_2 &= 2\nabla v \cdot \nabla^2P + \Delta v \cdot \nabla P, \\ g_2 &= (\nabla_{n_b}v - \nabla_v n_b) \cdot \nabla P. \end{aligned}$$

As a result, adding corresponding limit terms and proceeding as before, we derive by Lemma 1, Proposition 1, Corollary 2, Proposition 8 that

$$|\alpha^{-2\gamma-3}\partial_t\tilde{\mathbf{a}}|_{L^\infty} \leq Q(\mathcal{T}', w, 1)Q(\|\alpha^{-\gamma-2}\tilde{v} - a_v\|_{H^{5/2}}, \|\alpha^{-2\gamma-4}\tilde{P} - a_p\|_{H^{5/2}}).$$

As a result, the desired estimate follows from Proposition 11 and Proposition 12. \square

4.6. Modification of the equation. Based on all the discussions and estimates above, we find that $\alpha^{-\gamma}\tilde{D}_t w$ doesn't vanish when $\tilde{x} \rightarrow \pm\infty$ in general. As a direct consequence, we need to modify equation (31) of w immediately adding the ‘‘limit part’’ of $\tilde{D}_t w$.

In fact, recalling a_{w_t} from Proposition 9, we rewrite (31) into

$$(65) \quad \tilde{D}_t(\tilde{D}_t w - \alpha^\gamma a_{w_t}) + \tilde{\mathbf{a}}|\mathcal{T}'|^{-1}\tilde{\mathcal{N}}_0 w = \tilde{R},$$

where

$$\tilde{R} = \tilde{R}_1 + \tilde{R}_2 - \tilde{D}_t(\alpha^\gamma a_{w_t}),$$

and direct computations show that

$$\tilde{D}_t(\alpha^\gamma a_{w_t}) = -(\gamma+1)(\gamma+2)\alpha^\gamma(\chi_l a'_{\zeta_t, l} + \chi_r \beta_r^2 (b_r^2 a_{\zeta_t, r})') + (\gamma+1)(\gamma+2)u_1((\alpha^\gamma \chi_l)' a_{\zeta_t, l} + (\alpha^\gamma \chi_r)' \beta_r^2 b_r^2 a_{\zeta_t, r})$$

with u_1 defined in (30).

Naturally, we also need to investigate $a'_{\zeta_t, i}$ ($i = l, r$). Starting from (51) and multiplying the weight $\alpha^{-\gamma-2}$ on both sides, we obtain

$$\begin{aligned} \partial_t(\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta) &= \alpha^{-\gamma-2}\partial_t\widetilde{\partial}_t\zeta = \alpha^{-\gamma-2}\partial_t(b_\zeta^{\frac{1}{2}})\widetilde{v} \cdot \widetilde{n}_t + \alpha^{-\gamma-2}b_\zeta^{\frac{1}{2}}\partial_t\widetilde{v} \cdot \widetilde{n}_t + \alpha^{-\gamma-2}b_\zeta^{\frac{1}{2}}\widetilde{v} \cdot \partial_t\widetilde{n}_t \\ &= -\alpha^{-\gamma-2}b_\zeta^{-\frac{1}{2}}(\partial_{\widetilde{x}x})^{-1}\partial_{\widetilde{x}}\widetilde{\zeta}[-(\partial_{\widetilde{x}x})^{-2}\partial_{\widetilde{x}}\partial_t x \partial_{\widetilde{x}}\widetilde{\zeta} + (\partial_{\widetilde{x}x})^{-1}\partial_{\widetilde{x}}\partial_t\widetilde{\zeta}]\widetilde{v} \cdot \widetilde{n}_t \\ &\quad + \alpha^{-\gamma-2}b_\zeta^{\frac{1}{2}}(\widetilde{D}_t\widetilde{v} - u_1\partial_{\widetilde{x}}\widetilde{v}) \cdot \widetilde{n}_t + \alpha^{-\gamma-2}b_\zeta^{\frac{1}{2}}\widetilde{v} \cdot (\widetilde{D}_t\widetilde{n}_t - u_1\partial_{\widetilde{x}}\widetilde{n}_t). \end{aligned}$$

Substituting (33), the Euler equation (1) and (17) into the equation above, one has

$$\begin{aligned} \alpha^{-\gamma-2}\partial_t\widetilde{\partial}_t\zeta &= -\alpha^{-\gamma-2}b_\zeta^{-\frac{1}{2}}(\partial_{\widetilde{x}x})^{-1}\partial_{\widetilde{x}}\widetilde{\zeta}[(\partial_{\widetilde{x}x})^{-1}\partial_{\widetilde{x}}\widetilde{\partial}_t\zeta - (\partial_{\widetilde{x}x})^{-1}\partial_t x (\partial_{\widetilde{x}x})^{-2}\partial_{\widetilde{x}}^2 x \partial_{\widetilde{x}}\widetilde{\zeta} \\ (66) \quad &\quad + (\partial_{\widetilde{x}x})^{-1}\partial_t x (\partial_{\widetilde{x}x})^{-1}\partial_{\widetilde{x}}^2\widetilde{\zeta}]\widetilde{v} \cdot \widetilde{n}_t + \alpha^{-\gamma-2}b_\zeta^{\frac{1}{2}}(-\widetilde{\nabla}P - u_1\partial_{\widetilde{x}}\widetilde{v}) \cdot \widetilde{n}_t \\ &\quad + \alpha^{-\gamma-2}b_\zeta^{\frac{1}{2}}\widetilde{v} \cdot (-\nabla_{\tau_t} v \cdot \widetilde{n}_t)\tau_t - u_1\partial_{\widetilde{x}}\widetilde{n}_t). \end{aligned}$$

Taking the limit when $\widetilde{x} \rightarrow \pm\infty$ and checking the right side term by term, we conclude

$$a'_{\zeta_t, i} = 0, \quad i = l, r.$$

Notice that when there is gravity, one has

$$a'_{\zeta_t, i} = \infty.$$

Therefore, we conclude that

$$\begin{aligned} (67) \quad \widetilde{D}_t(\alpha^\gamma a_{w_t}) &= -(\gamma+1)(\gamma+2)\beta_r^2\alpha^\gamma\chi_r 2b_r b'_r a_{\zeta_t, l} \\ &\quad + (\gamma+1)(\gamma+2)(\partial_{\widetilde{x}x})^{-1}(\widetilde{v}_1 - \partial_t x)((\alpha^\gamma\chi_l)'a_{\zeta_t, l} + (\alpha^\gamma\chi_r)'\beta_r^2 b_r^2 a_{\zeta_t, r}). \end{aligned}$$

Besides, recalling (55), (56), (64) and Remark 10, we know immediately that

$$(68) \quad \partial_t a_{\Phi_z} = \partial_t a_v = \partial_t a_p = 0.$$

These happen due to our settings of the time derivatives which own higher-order weights since $\gamma+1 > 0$.

5. A PRIORI ESTIMATES

we define the order- k energy ($k \geq 0$) on the surface Γ_t of the strip domain \mathcal{S}_0 :

$$\begin{aligned} (69) \quad E_{k, \gamma}(t) &= |\alpha^{-\gamma}\partial_{\widetilde{x}}^k(\widetilde{D}_t w - \alpha^\gamma a_{w_t})|_{L^2}^2 + |\widetilde{\mathfrak{a}}^{\frac{1}{2}}\alpha^{-\gamma}|\mathcal{T}'|^{-\frac{1}{2}}\partial_{\widetilde{x}}^k w|_{L^2}^2 \\ &\quad + (\widetilde{\mathcal{N}}_0(\widetilde{\mathfrak{a}}^{\frac{1}{2}}\alpha^{-\gamma}|\mathcal{T}'|^{-\frac{1}{2}}\partial_{\widetilde{x}}^k w), \widetilde{\mathfrak{a}}^{\frac{1}{2}}\alpha^{-\gamma}|\mathcal{T}'|^{-\frac{1}{2}}\partial_{\widetilde{x}}^k w)_{L^2}, \end{aligned}$$

and the lower-order energy

$$\begin{aligned} E_0(t) &= \|\alpha^{-1}\mathcal{T}' - T_{1,c}\|_{L^2}^2 + |\alpha^{-\gamma-2}\widetilde{\partial}_t\zeta - a_{\zeta_t}|_{L^2}^2 + \|\alpha^{-\gamma-3}\nabla\Phi - (a_{\Phi_x}, a_{\Phi_z})^T\|_{L^2}^2 + \|\alpha^{-2\gamma-4}\widetilde{P} - a_p\|_{L^2}^2 \\ &\quad + |a_{\zeta_t, i}|_{L^\infty}^2 + |b|_{L^\infty}^2 + |b_r|_{L^\infty}^2 + |b_{r2}|_{L^\infty}^2 \end{aligned}$$

where $T_{1,c}$, a_{ζ_t} , a_{Φ_x} , a_{Φ_z} , and a_p are the corresponding weighted limits given in (39), (34), Proposition 10 and (63) respectively.

The total energy is then defined by

$$\mathcal{E}(t) = \sum_{l \leq k} E_{l, \gamma}(t) + E_0(t).$$

The main theorem of our paper is presented below.

Theorem 2. *Assume that the integer k and real γ satisfy*

$$2 \leq k \leq \min\left\{\frac{\pi}{\omega}, \frac{\pi}{\omega_r}\right\}, \quad 0 < \gamma + 1 \leq \min\left\{\frac{2\pi}{\omega}, \frac{2\pi}{\omega_r}\right\}$$

and $Q_E(\cdot, \cdot, \cdot)$, $P(\cdot)$ are fixed polynomials with positive constant coefficients depending only on $k, \gamma, \omega, \omega_r$ and the bound L from Definition 2.

Let a solution to the water-waves system (1)-(5) be given by the free surface graph function ζ and its time derivative $\partial_t \zeta$ with $\alpha^{-1} \partial_{\tilde{x}} \zeta \in H^{k+3/2}(\mathbb{R})$ and $\alpha^{-\gamma-2} \partial_t \zeta - a_{\zeta_t} \in H^{k+2}(\mathbb{R})$ and the initial values are chosen from Λ_0 . Moreover, assume that Assumption 2 holds initially for $t = 0$.

Then there exists $T^* > 0$ depending only on $Q_E(0) = Q_E(E_0(0), |\alpha w(0)|_{H^{k+1/2}}^2, |\alpha^{-\gamma} \tilde{D}_t w(0) - a_{w_t}(0)|_{H^k}^2)$ such that when $T \leq T^*$, we have $(\zeta, \mathcal{T}, \Phi, P) \in \Lambda(T, k)$ and the following estimate holds

$$\mathcal{E}(t) \leq \mathcal{E}(0) + \int_0^t P(\mathcal{E}(s)) ds, \quad \forall t \in [0, T].$$

Remark 11. Based on the proof of this theorem, Assumption 2 holds for $t \in [0, T]$, i.e. the weighted Taylor's sign holds with a new lower bound $a_0/2$ in stead of a_0 .

Proof. To begin with, direct computations lead to

$$\begin{aligned} (70) \quad \frac{1}{2} \frac{d}{dt} E_{k,\gamma}(t) &= (\alpha^{-\gamma} \partial_{\tilde{x}}^k \partial_t (\tilde{D}_t w - \alpha^\gamma a_{w_t}), \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad + (\partial_t (\tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2}) \partial_{\tilde{x}}^k w, \tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w) \\ &\quad + (\tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k \partial_t w, \tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w) \\ &\quad + (\tilde{\mathcal{N}}_0 (\tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w), \partial_t (\tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2}) \partial_{\tilde{x}}^k w) \\ &\quad + (\tilde{\mathcal{N}}_0 (\tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w), \tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k \partial_t w) \\ &:= I_1 + I_2 + \dots + I_5, \end{aligned}$$

where we take (\cdot, \cdot) for the inner product $(\cdot, \cdot)_{L^2(\mathbb{R})}$.

Step 1. Estimates for $I_1 + I_5$. In fact, one could see that $I_1 + I_5$ is the main part in the energy estimate. Plugging equation (65) into I_1 , one has

$$\begin{aligned} I_1 + I_5 &= -(\alpha^{-\gamma} \partial_{\tilde{x}}^k [u_1 \partial_{\tilde{x}} (\tilde{D}_t w - \alpha^\gamma a_{w_t})], \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad - (\alpha^{-\gamma} \partial_{\tilde{x}}^k [\tilde{\mathbf{a}} |\mathcal{T}'|^{-1} \tilde{\mathcal{N}}_0 w], \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad + (\alpha^{-\gamma} \partial_{\tilde{x}}^k \tilde{R}, \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) + (\tilde{\mathcal{N}}_0 (\tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w), \tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k \partial_t w). \end{aligned}$$

Using commutators to bring some cancellations, one derives

$$\begin{aligned} (71) \quad I_1 + I_5 &= -(\alpha^{-\gamma} \partial_{\tilde{x}}^k [u_1 \partial_{\tilde{x}} (\tilde{D}_t w - \alpha^\gamma a_{w_t})], \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad - (\alpha^{-\gamma} [\partial_{\tilde{x}}^k, \tilde{\mathbf{a}}] |\mathcal{T}'|^{-1} \tilde{\mathcal{N}}_0 w, \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad - (\alpha^{-\gamma} \tilde{\mathbf{a}} [\partial_{\tilde{x}}^k, |\mathcal{T}'|^{-1}] \tilde{\mathcal{N}}_0 w, \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) + (\alpha^{-\gamma} \partial_{\tilde{x}}^k \tilde{R}, \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad - (\tilde{\mathcal{N}}_0 (\tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w), \tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k (u_1 \partial_{\tilde{x}} w)) \\ &\quad + (\tilde{\mathcal{N}}_0 (\tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w), \tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k (\alpha^\gamma a_{w_t})) \\ &\quad + ([\tilde{\mathcal{N}}_0, \tilde{\mathbf{a}}^{1/2}] \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w, \tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad + (\tilde{\mathbf{a}}^{1/2} [\tilde{\mathcal{N}}_0, \alpha^{-\gamma} |\mathcal{T}'|^{-1/2}] \partial_{\tilde{x}}^k w, \tilde{\mathbf{a}}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &:= I_{11} + I_{12} + \dots + I_{18}, \end{aligned}$$

where careful weighted commutators estimates are needed. We firstly deal with all these terms except for I_{14} , and the term I_{14} (which involves \tilde{R}) is handled later.

- I_{11} estimate. Direct computations and integration by parts lead to

$$\begin{aligned} I_{11} &= -(\alpha^{-\gamma} [\partial_{\tilde{x}}^k, u_1] \partial_{\tilde{x}} (\tilde{D}_t w - \alpha^\gamma a_{w_t}), \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad - (\alpha^{-\gamma} u_1 \partial_{\tilde{x}} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t}), \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &= -(\alpha^{-\gamma} [\partial_{\tilde{x}}^k, u_1] \partial_{\tilde{x}} (\tilde{D}_t w - \alpha^\gamma a_{w_t}), \alpha^{-\gamma} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad + \frac{1}{2} (\partial_{\tilde{x}} (\alpha^{-2\gamma} u_1) \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t}), \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t})), \end{aligned}$$

where we recall u_1 from (30). Noticing that

$$\partial_{\bar{x}}\tilde{v} = \partial_{\bar{x}}(v(t, x, \zeta) \circ \mathcal{T}) = \partial_{\bar{x}}x(\partial_{\bar{x}}v|_{\Gamma_t} + (\partial_{\bar{x}}x)^{-1}\partial_{\bar{x}}\tilde{\zeta}\partial_{\bar{x}}v|_{\Gamma_t})$$

where (24) is applied to \tilde{v} . So we count the derivatives and apply Lemma 1, Corollary 1 and Corollary 2 to obtain

$$\begin{aligned} |I_{11}| &\leq C(|u_1|_{W^{k,\infty}} + |\alpha^{2\gamma}\partial_{\bar{x}}(\alpha^{-2\gamma}u_1)|_{L^\infty})|\alpha^{-\gamma}(\tilde{D}_t w - \alpha^\gamma a_{w_t})|_{H^k} \\ &\leq Q(\mathcal{T}', w, k)(|\alpha^{-\gamma-2}\tilde{v} - a_v|_{H^{k+1}} + \|a_v\|_{L^\infty} + |\alpha^{-(\gamma+3)/2}\partial_t x|_{H^{k+1}})|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}^2. \end{aligned}$$

Besides, using Proposition 8 and Proposition 11, we arrive at the following estimate:

$$|I_{11}| \leq Q(\mathcal{T}', \nabla\Phi, \partial_t\tilde{\zeta}, w, k)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^{k-1}} + |a_{\zeta_t, i}|_{L^\infty})|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}^2.$$

- I_{12} estimate. We have by (26), Lemma 1, Corollary 1 and Corollary 2 that

$$\begin{aligned} |I_{12}| &\leq |\alpha^{-\gamma}[\partial_{\bar{x}}^k, \tilde{\mathbf{a}}]|\mathcal{T}'|^{-1}\tilde{\mathcal{N}}_0 w|_{L^2}|\alpha^{-\gamma}\partial_{\bar{x}}^k(\tilde{D}_t w - \alpha^\gamma a_{w_t})|_{L^2} \\ &\leq Q(\mathcal{T}', w, 1)|\alpha^{-\gamma-2}\tilde{\mathbf{a}}|_{W^{k,\infty}}|\alpha\tilde{\mathcal{N}}_0 w|_{H^{k-1}}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}. \end{aligned}$$

Using Proposition 13, we obtain

$$\begin{aligned} |I_{12}| &\leq Q(\mathcal{T}', w, 1)(|\alpha^{-2\gamma-3}\tilde{\mathbf{a}} - \partial_{\bar{z}}a_p|_{H^{k+1}} + |a_{p, i}|_{W_{\bar{z}}^{k+1,\infty}})|\alpha w|_{H^k}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \\ &\leq Q(\mathcal{T}', \nabla\Phi, \tilde{P}, w, k)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^{k-1}} + |a_{\zeta_t, i}|_{L^\infty}^2)|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}. \end{aligned}$$

- I_{13} estimate. Similar arguments as for I_{12} , we arrive at

$$\begin{aligned} I_{13} &\leq Q(\mathcal{T}', w, k-1)|\alpha^{-2\gamma-3}\tilde{\mathbf{a}}|_{L^\infty}|\alpha w|_{H^k}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \\ &\leq Q(\mathcal{T}', \nabla\Phi, \partial_t\tilde{\zeta}, \tilde{P}, w, k)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{L^2} + |a_{\zeta_t, i}|_{L^\infty}^2)|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \end{aligned}$$

where Proposition 13 and Proposition 11 are applied.

- I_{15} estimate. Similarly as for I_{11} , we have

$$\begin{aligned} I_{15} &= -\frac{1}{2}(\tilde{\mathcal{N}}_0(\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_{\bar{x}}^k w), \tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}[\partial_{\bar{x}}^k, u_1]\partial_{\bar{x}} w) \\ &\quad + \frac{1}{2}(\tilde{\mathcal{N}}_0(\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_{\bar{x}}^k w), u_1\partial_{\bar{x}}(\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2})\partial_{\bar{x}}^k w) \\ &\quad + \frac{1}{2}([u_1\partial_{\bar{x}}, \tilde{\mathcal{N}}_0](\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_{\bar{x}}^k w), \tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_{\bar{x}}^k w). \end{aligned}$$

Applying Lemma 2, Lemma 7, Corollary 1, Corollary 2 and Proposition 3.18 [40] (for the standard commutator estimate involving $\tilde{\mathcal{N}}_0$ in the third term), we obtain

$$\begin{aligned} |I_{15}| &\leq C|\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_{\bar{x}}^k w|_{H^{1/2}}(|\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}[\partial_{\bar{x}}^k, u_1]\partial_{\bar{x}} w|_{H^{1/2}} \\ &\quad + |u_1\partial_{\bar{x}}(\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2})\partial_{\bar{x}}^k w|_{H^{1/2}}) + C|u_1|_{W^{1,\infty}}|\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_{\bar{x}}^k w|_{L^2}^2 \\ &\leq C|\alpha^{-\gamma-3/2}\tilde{\mathbf{a}}^{1/2}|_{W^{1,\infty}}^2(1 + |1 - \alpha^{1/2}|\mathcal{T}'|^{-1/2}|_{H^2}^2)|u_1|_{H^{k+1/2}}|\alpha w|_{H^{k+1/2}}^2. \end{aligned}$$

Moreover, direct computations under Assumption 2 and applying Proposition 13, Corollary 1, Corollary 2, one has

$$|\alpha^{-\gamma-3/2}\tilde{\mathbf{a}}^{1/2}|_{W^{1,\infty}} \leq Q(\mathcal{T}', \nabla\Phi, \partial_t\tilde{\zeta}, \tilde{P}, w, 1)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{L^2} + |a_{\zeta_t, i}|_{L^\infty}^2)$$

and

$$|1 - \alpha^{1/2}|\mathcal{T}'|^{-1/2}|_{H^2} \leq Q(\mathcal{T}', w, 1),$$

which together with Proposition 8, Proposition 9 and Proposition 11 results in

$$|I_{15}| \leq Q(\mathcal{T}', \nabla\Phi, \partial_t\tilde{\zeta}, \tilde{P}, w, k+1/2)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^{k-3/2}} + |a_{\zeta_t, i}|_{L^\infty}).$$

- I_{16} estimate. Similarly as in the previous estimate, we obtain

$$\begin{aligned} |I_{16}| &\leq C|\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_x^k w|_{H^{1/2}}|\tilde{\mathbf{a}}^{1/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_x^k(\alpha^\gamma a_{w_t})|_{H^{1/2}} \\ &\leq Q(\mathcal{T}', \nabla\Phi, \tilde{\partial}_t\zeta, \tilde{P}, w, k+1/2)(1|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^{k-1}} + |a_{\zeta_t, i}|_{L^\infty})|a_{\zeta_t, i}|_{L^\infty}. \end{aligned}$$

- I_{17} estimate. We have

$$\begin{aligned} |I_{17}| &\leq C|[\tilde{\mathbf{a}}^{1/2}, \tilde{\mathcal{N}}_0]\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_x^k w|_{L^2}||\mathcal{T}'|^{-1/2}\tilde{\mathbf{a}}^{1/2}|_{L^\infty}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \\ &\leq C|\alpha^{-\gamma-3/2}\tilde{\mathbf{a}}^{1/2}|_{W^{2,\infty}}|\alpha^{\gamma+3/2}\alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_x^k w|_{H^{1/2}}|\alpha^{-1/2}\tilde{\mathbf{a}}^{1/2}|_{L^\infty}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}, \end{aligned}$$

where we use Proposition 6 (1) for the commutator $[\tilde{\mathbf{a}}^{1/2}, \tilde{\mathcal{N}}_0]$ with $\gamma_1 = -\gamma - 3/2$, $\gamma_2 = \gamma + 3/2$ and $g = \alpha^{-\gamma}|\mathcal{T}'|^{-1/2}\partial_x^k w$. Therefore, we require also

$$\gamma + 3/2 = |\gamma + 3/2| < \min\left\{\frac{\pi}{2\omega}, \frac{\pi}{2\omega_r}\right\}.$$

Similar arguments as above lead to the following estimate:

$$|I_{17}| \leq Q(\mathcal{T}', \nabla\Phi, \tilde{\partial}_t\zeta, \tilde{P}, w, 1)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{L^2} + |a_{\zeta_t, i}|_{L^\infty})|\alpha w|_{H^{k+1/2}}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}.$$

- I_{18} estimate. Applying similar analysis as above again, we derive

$$\begin{aligned} |I_{18}| &\leq C|\alpha^{-\gamma-3/2}\tilde{\mathbf{a}}^{1/2}|_{L^\infty}|\alpha^{\gamma+3/2}[\tilde{\mathcal{N}}_0, \alpha^{-\gamma}|\mathcal{T}'|^{-1/2}]\partial_x^k w|_{L^2}|\alpha^{-1/2}\tilde{\mathbf{a}}^{1/2}|_{L^\infty}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \\ &\leq Q(\mathcal{T}', \nabla\Phi, \tilde{\partial}_t\zeta, \tilde{P}, w, 1)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{L^2} + |a_{\zeta_t, i}|_{L^\infty})|\alpha w|_{H^{k+1/2}}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}. \end{aligned}$$

Step 2. The remainder part: Estimates for I_{14} . Recalling from (65) that

$$\tilde{R} = R_1 \circ \mathcal{T} + R_2 \circ \mathcal{T} - \tilde{D}_t(\alpha^\gamma a_{w_t}) \quad \text{on } \Gamma_t$$

where R_1, R_2 and $\tilde{D}_t(\alpha^\gamma a_{w_t})$ are expressed in (19),(21) and (67).

Therefore, we split I_{24} into the following three parts:

$$\begin{aligned} I_{14} &= (\alpha^{-\gamma}\partial_x^k \tilde{R}_1, \alpha^{-\gamma}\partial_x^k(\tilde{D}_t w - \alpha^\gamma a_{w_t})) + (\alpha^{-\gamma}\partial_x^k \tilde{R}_2, \alpha^{-\gamma}\partial_x^k(\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &\quad - (\alpha^{-\gamma}\partial_x^k \tilde{D}_t(\alpha^\gamma a_{w_t}), \alpha^{-\gamma}\partial_x^k(\tilde{D}_t w - \alpha^\gamma a_{w_t})) \\ &:= B_1 + B_2 + B_3. \end{aligned}$$

- Term B_1 . Checking term by term, we know directly from (19) that the terms in R_1 are always like

$$(\partial v)^2 \partial n_t, \partial^2 v \partial v, \partial^2 P \partial n_t.$$

Except $\partial^2 \tilde{\zeta}$ parts with α^{-1} , we know that $R_1 \circ \mathcal{T}|_{\Gamma_t}$ contains terms like

$$\alpha^{-3}(\partial \tilde{v})^2, \alpha^{-3}\partial^2 \tilde{v} \partial \tilde{v}, \alpha^{-3}\partial^2 \tilde{P}.$$

As a result, applying Proposition 1, Corollary 1, Corollary 2 and Corollary 3 similarly as before and remembering that $\gamma > -1$, we obtain for $k \geq 1$ that

$$\begin{aligned} |B_1| &\leq C|\alpha^{-\gamma}\tilde{R}_1|_{H^k}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \\ &\leq Q(\mathcal{T}', w, k)|\alpha^{-1}\tilde{\zeta}|_{H^{k+2}}(|\alpha^{-\gamma-3}(\partial \tilde{v})^2|_{H^k} + |\alpha^{-\gamma-3}\partial^2 \tilde{v} \partial \tilde{v}|_{H^k} + |\alpha^{-\gamma-3}\partial^2 \tilde{P}|_{H^k})|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \\ &\leq Q(\mathcal{T}', w, k)[|\alpha^{-\gamma-2}\tilde{v} - a_v|_{H^{k+2}}^2 + |\alpha^{-2\gamma-4}\tilde{P} - a_p|_{H^{k+2}} + |\alpha^{\gamma+1}\chi_l|_{H^k}(|a_v|_{W^{k+2,\infty}} + |a_{p,i}|_{W^{\frac{k+2}{\tilde{z}},\infty}})] \times \\ &\quad |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}. \end{aligned}$$

Using Proposition 12, Proposition 11 and substituting the estimates above into B_1 estimate, we conclude

$$|B_1| \leq Q(\mathcal{T}', \nabla\Phi, \tilde{\partial}_t\zeta, \tilde{P}, w, k+1/2)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} + |a_{\zeta_t, i}|_{L^\infty})|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}$$

- Term B_2 . Similarly as above, one has from (21) the following estimates for $k \geq 1$:

$$\begin{aligned} |B_2| &\leq C|\alpha^{-\gamma}\tilde{R}_2|_{H^k}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \\ &\leq Q(\mathcal{T}', w, k)(|\alpha^{-\gamma-2}\tilde{v}|_{H^{k+2}}^2 + |\alpha^{-2\gamma-4}\tilde{P}|_{H^{k+2}} + |\alpha^{-\gamma-1}\partial_{\tilde{z}}\tilde{u}_p|_{H^k})|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}. \end{aligned}$$

Now it remains to deal with u_p term, which is also a higher-order-derivative term since system (22) involves $\partial^3 P$ and \mathcal{T}''' at Γ_b . Therefore, we need a close look at this boundary term.

In fact, transforming system (22) directly to an equivalent system in \mathcal{S}_0 as system (32) for \tilde{P} and applying Lemma 3, Proposition 1, Proposition 3 and Corollary 2 as before, we have

$$\begin{aligned} (72) \quad &|\alpha^{-\gamma-1}\partial_{\tilde{z}}\tilde{u}_p|_{H^k} \leq C\|\alpha^{-\gamma-1}\tilde{u}_1\|_{H^{k+3/2}} \\ &\leq Q(\mathcal{T}', w, k)[\|\alpha^{-\gamma-2}\partial\tilde{\kappa}_{\mathcal{H}}\partial\tilde{P}\|_{H^{k-1/2}} + \|\alpha^{-\gamma-2}\partial\tilde{\kappa}_{\mathcal{H}}\partial^2\tilde{P}\|_{H^{k-1/2}} + \|\alpha^{-\gamma-2}\tilde{\kappa}_{\mathcal{H}}\partial^2\tilde{P}\|_{H^{k-1/2}} \\ &\quad + \|\tilde{\kappa}_{\mathcal{H}}\alpha^{-\gamma-2}\partial(\partial\tilde{v}\partial\tilde{v})\|_{H^{k-1/2}} + \|\alpha^{-\gamma-3}\partial\tilde{v}\partial\tilde{v}\|_{H^{k+3/2}} + \|\alpha^{-\gamma-3}\tilde{P}\|_{H^{k+5/2}} + |\alpha^{-\gamma-3}\partial\tilde{v}\partial\tilde{v}|_{H^{k+1}} \\ &\quad + |\alpha^{-\gamma-2}\tilde{\kappa}_{\mathcal{H}}\partial^2\tilde{P}|_{H^k(\Gamma_b)} + |\alpha^{-\gamma-2}\tilde{\kappa}_{\mathcal{H}}\partial\tilde{P}|_{H^k(\Gamma_b)} + (|\alpha^{-1}\mathcal{T}' - T_{1,c}|_{H^{k+2}(\Gamma_b)} + |T_{1,c}|_{L^\infty}) \times \\ &\quad |\alpha^{-\gamma-3}\tilde{P}|_{H^{k+3}(\Gamma_b)}] \end{aligned}$$

where we omit the detailed estimates for n_t and \mathcal{T} and only keep track of the highest-order term of \mathcal{T} on Γ_b .

Recalling the weights for $w = \tilde{\kappa}$, \tilde{v} and \tilde{P} and taking the following estimate of \tilde{v} for example, we obtain

$$\begin{aligned} &\|\alpha^{-\gamma-3}\partial_{\tilde{x}}\tilde{v}\partial_{\tilde{x}}\tilde{v}\|_{H^{k+3/2}} \\ &\leq \|\alpha^{\gamma+1}(\alpha^{-\gamma-2}\partial_{\tilde{x}}\tilde{v} - a_{v,1})(\alpha^{-\gamma-2}\partial_{\tilde{x}}\tilde{v} - a_{v,1})\|_{H^{k+3/2}} + C\|\alpha^{\gamma+1}\alpha^{-\gamma-2}\partial_{\tilde{x}}\tilde{v}a_{v,1}\|_{H^{k+3/2}} \\ &\quad + C\|\alpha^{\gamma+1}(a_{v,1})^2\|_{H^{k+3/2}} \\ &\leq Q(\mathcal{T}', \nabla\tilde{\Phi}, \partial_t\tilde{\zeta}, w, k+1/2)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} + |a_{\zeta_t}|_{L^\infty}^2) \end{aligned}$$

where Proposition 11 is applied and $a_{v,1}$ is defined in Lemma 8.

As a result, thanks to Corollary 4, the estimates for the other terms above in the right side of (72) follow in a similar way.

Consequently, going back to the estimate for B_2 and applying Proposition 12, Proposition 11 and Proposition 9, we obtain

$$|B_2| \leq Q(\mathcal{T}', \nabla\tilde{\Phi}, \partial_t\tilde{\zeta}, \tilde{P}, w, k+1/2)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}^2 + |a_{\zeta_t, i}|_{L^\infty}^2)|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}.$$

Besides, for term B_3 , we have immediately the following estimate by Proposition 1 QQQ:

$$\begin{aligned} |B_3| &\leq C(|\alpha^{-1-(\gamma+1)/2}\tilde{v}|_{W^{k,\infty}} + |\alpha^{-1-(\gamma+1)/2}\partial_t x|_{W^{k,\infty}})|\alpha^{(\gamma+1)/2}a_{\zeta_t}|_{L^2}|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \\ &\leq Q(\mathcal{T}', \nabla\tilde{\Phi}, \tilde{P}, w, k+1/2)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}^2 + |a_{\zeta_t, i}|_{L^\infty}^2)|\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k} \end{aligned}$$

Summing up these estimates above, we finally conclude that

$$\begin{aligned} |I_{14}| &\leq Q(\mathcal{T}', \nabla\tilde{\Phi}, \tilde{P}, w, k+1/2)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^{k-1}}^2 + |\alpha^{-\gamma-2}\partial_t\tilde{\zeta} - a_{\zeta_t}|_{L^2}^2 + |a_{\zeta_t, i}|_{L^\infty}^2) \times \\ &\quad |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^k}. \end{aligned}$$

Step 3. Estimates for I_2, I_3, I_4 . We handle these terms one by one.

To begin with, we have by Proposition 1, Proposition 8, Proposition 13 and Proposition 9 that

$$\begin{aligned} |I_2| &\leq |\alpha^{-\gamma-1/2}|\mathcal{T}'|^{-1/2}\tilde{\mathbf{a}}^{1/2}\partial_t((\alpha^{-2\gamma-3}\tilde{\mathbf{a}})^{1/2}|\mathcal{T}'|^{-1/2})|_{L^\infty}|\alpha\partial_{\tilde{x}}^k w|_{L^2}^2 \\ &\leq C[|\alpha^{-2\gamma-3}\partial_t\tilde{\mathbf{a}}|_{L^\infty} + |\alpha^{-2\gamma-3}\tilde{\mathbf{a}}|_{L^\infty}|\alpha^{1/2}|\mathcal{T}'|^{-5/2}\partial_t(|\mathcal{T}'|^2)|_{L^\infty}]|\alpha w|_{H^k}^2 \\ &\leq Q(\mathcal{T}', \nabla\tilde{\Phi}, \partial_t\tilde{\zeta}, \tilde{P}, w, 1)(1 + |\alpha^{-\gamma}\tilde{D}_t w - a_{w_t}|_{H^2} + |a_{\zeta_t, i}|_{L^\infty})|\alpha w|_{H^k}^2 \end{aligned}$$

Meanwhile, the estimate for I_4 follows similarly as above:

$$|I_4| \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \tilde{\partial}_t \zeta, \tilde{P}, w, 2)(1 + |\alpha^{-\gamma} \tilde{D}_t w - a_{w_t}|_{H^3} + |a_{\zeta_t, i}|_{L^\infty}) |\alpha w|_{H^{k+1/2}}^2.$$

On the other hand, I_3 can be handled in a similar way as I_{11} and I_{16} :

$$\begin{aligned} I_3 = & (\tilde{\alpha}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k (\tilde{D}_t w - \alpha^\gamma a_{w_t}), \tilde{\alpha}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w) \\ & - (\tilde{\alpha}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k (u_1 \partial_{\tilde{x}} w), \tilde{\alpha}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w) \\ & + (\tilde{\alpha}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k (\alpha^\gamma a_{w_t}), \tilde{\alpha}^{1/2} \alpha^{-\gamma} |\mathcal{T}'|^{-1/2} \partial_{\tilde{x}}^k w), \end{aligned}$$

so summing up these estimates above leads to

$$|I_2| + |I_3| + |I_4| \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \tilde{\partial}_t \zeta, \tilde{P}, w, 2)(1 + |\alpha^{-\gamma} \tilde{D}_t w - a_{w_t}|_{H^k} + |a_{\zeta_t, i}|_{L^\infty} + |\alpha w|_{H^k}) |\alpha w|_{H^k}$$

Step 4. The lower-order energy. We still need to deal with the lower-order energy $E_0(t)$.

For the first term of \mathcal{T}' , one has by applying Remark 10, Proposition 8 and Proposition 9 that

$$\frac{1}{2} \frac{d}{dt} |\alpha^{-1} \mathcal{T}' - T_{1,c}|_{L^2}^2 = (\alpha^{-1} \partial_t \mathcal{T}', \alpha^{-1} \mathcal{T}' - T_{1,c}) \leq Q(\mathcal{T}', \tilde{\partial}_t \zeta, a_{\zeta_t}, w, 1).$$

For $\tilde{\partial}_t \zeta$ term, recalling (66) and dealing all the terms as above lead to

$$\frac{1}{2} \frac{d}{dt} |\alpha^{-\gamma-2} \tilde{\partial}_t \zeta - a_{\zeta_t}|_{L^2}^2 = (\partial_t (\alpha^{-\gamma-2} \tilde{\partial}_t \zeta), \alpha^{-\gamma-2} \tilde{\partial}_t \zeta - a_{\zeta_t}) \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \tilde{\partial}_t \zeta, \tilde{P}, w, 1).$$

For $\nabla \tilde{\Phi}$ term, going back directly to system (52) and taking ∂_t , one obtains immediately the following system

$$\begin{cases} \Delta \partial_t \partial_{\tilde{z}} \tilde{\Phi} = 0 & \text{in } \mathcal{S}_0, \\ \partial_t \partial_{\tilde{z}} \tilde{\Phi}|_{\Gamma_t} = -(\partial_{\tilde{x}} \partial_t x) \tilde{\partial}_t \zeta - (\partial_{\tilde{x}} x) \partial_t \tilde{\partial}_t \zeta, & \partial_t \partial_{\tilde{z}} \tilde{\Phi}|_{\Gamma_b} = 0. \end{cases}$$

where one can resort to (66) for the expression of $\partial_t \tilde{\partial}_t \zeta$. Applying Proposition 4 and using similar arguments as above lead to directly

$$\|\alpha^{-\gamma-3} \partial_t \partial_{\tilde{z}} \tilde{\Phi}\|_{L^2} \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \tilde{\partial}_t \zeta, \tilde{P}, w, 1).$$

As a result, one has the following estimate

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\alpha^{-\gamma-3} \partial_{\tilde{z}} \tilde{\Phi} - a_{\Phi_z}\|_{L^2}^2 &= (\alpha^{-\gamma-3} \partial_t \partial_{\tilde{z}} \tilde{\Phi} - \partial_t a_{\Phi_z}, \alpha^{-\gamma-3} \partial_{\tilde{z}} \tilde{\Phi} - a_{\Phi_z}) \\ &\leq \|\alpha^{-\gamma-3} \partial_t \partial_{\tilde{z}} \tilde{\Phi}\|_{L^2} \|\alpha^{-\gamma-3} \partial_{\tilde{z}} \tilde{\Phi} - a_{\Phi_z}\|_{L^2} \\ &\leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \tilde{\partial}_t \zeta, \tilde{P}, w, 1). \end{aligned}$$

where (68) is applied.

Meanwhile, for \tilde{P} term, recalling (68) again and using similar arguments as for $\tilde{\Phi}$ above show that

$$\frac{1}{2} \frac{d}{dt} \|\alpha^{-2\gamma-4} \tilde{P} - a_p\|_{L^2}^2 \leq \|\alpha^{-2\gamma-4} \partial_t \tilde{P}\|_{L^2} \|\alpha^{-2\gamma-4} \tilde{P} - a_p\|_{L^2} \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \tilde{\partial}_t \zeta, \tilde{P}, w, 1).$$

Consequently, using Remark 10 and (68) again we conclude for the lower-order energy part that

$$\frac{1}{2} \frac{d}{dt} E_0(t) \leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \tilde{\partial}_t \zeta, \tilde{P}, w, 1).$$

Step 5. The end of the energy estimate. Summing up all these estimates from Step 1 to Step 4, we arrive at the following energy estimate:

$$(73) \quad \frac{1}{2} \frac{d}{dt} \mathcal{E}(t) \leq Q_k(t),$$

where

$$Q_k(t) = Q(E_0(t), |\alpha w|_{H^{k+1/2}}^2, |\alpha^{-\gamma} \tilde{D}_t w - a_{w_t}|_{H^k}^2).$$

To close the energy estimate, we still need to consider the relationship between $|\alpha w|_{H^{k+1/2}}$ and the higher-order term of w in $E_{k,\gamma}$.

Lemma 9. *Let integer $k \geq 2$. Then there hold*

$$|\alpha w|_{H^{k+1/2}}^2 + |\alpha^{-\gamma} \tilde{D}_t w - a_{w_t}|_{H^k}^2 \leq C\mathcal{E}(t),$$

and

$$\mathcal{E}(t) \leq Q_k(t).$$

Proof. (Proof of the lemma) Comparing all the terms in $Q_k(t)$ with those in $E_{k,\gamma}(t)$, we know that the key point here lies in the estimate of $|\alpha w|_{H^{k+1/2}}$.

We firstly prove the second inequality. In fact, applying Lemma 7 shows directly for $l \leq k$ that

$$\begin{aligned} c_1 |\tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{-\gamma} |\mathcal{T}'|^{-\frac{1}{2}} \partial_{\tilde{x}}^l w|_{H^{1/2}}^2 &\leq |\tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{-\gamma} |\mathcal{T}'|^{-\frac{1}{2}} \partial_{\tilde{x}}^l w|_{L^2}^2 + (\tilde{\mathcal{N}}_0(\tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{-\gamma} |\mathcal{T}'|^{-\frac{1}{2}} \partial_{\tilde{x}}^l w), \tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{-\gamma} |\mathcal{T}'|^{-\frac{1}{2}} \partial_{\tilde{x}}^l w) \\ &\leq C_1 |\tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{-\gamma} |\mathcal{T}'|^{-\frac{1}{2}} \partial_{\tilde{x}}^l w|_{H^{1/2}}^2 \end{aligned}$$

where c_1, C_1 are positive constants depending on \mathcal{S}_0 .

Moreover, one has in a similar way as in the estimates before that

$$\begin{aligned} |\tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{-\gamma} |\mathcal{T}'|^{-\frac{1}{2}} \partial_{\tilde{x}}^l w|_{H^{1/2}} &= |\alpha^{-\gamma-3/2} \tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{3/2} |\mathcal{T}'|^{-\frac{1}{2}} \partial_{\tilde{x}}^l w|_{H^{1/2}} \\ &\leq |\alpha^{-\gamma-3/2} \tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{1/2} |\mathcal{T}'|^{-\frac{1}{2}}|_{W^{1,\infty}} |\alpha w|_{H^{l+1/2}} \\ &\leq Q(\mathcal{T}', \nabla \tilde{\Phi}, \tilde{\partial}_t \zeta, \tilde{P}, w, 1) |\alpha w|_{H^{l+1/2}}. \end{aligned}$$

As a result, the right side of the desired inequality is finished.

For the first inequality, noticing that $\alpha^{-\gamma-3/2} \tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{1/2} |\mathcal{T}'|^{-\frac{1}{2}}$ is also bounded below by Assumption 2 for $t = 0$, Proposition 1 and Proposition 13, we have thanks to Lemma 2 (3) that

$$\begin{aligned} |\alpha \partial_{\tilde{x}}^l w|_{H^{1/2}} &\leq C(a_0^{-1}, |\alpha^{-\gamma-3/2} \tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{1/2} |\mathcal{T}'|^{-\frac{1}{2}}|_{W^{1,\infty}}) |\alpha^{-\gamma-3/2} \tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{1/2} |\mathcal{T}'|^{-\frac{1}{2}} \alpha \partial_{\tilde{x}}^l w|_{H^{1/2}} \\ &\leq C |\tilde{\mathbf{a}}^{\frac{1}{2}} \alpha^{-\gamma} |\mathcal{T}'|^{-\frac{1}{2}} \partial_{\tilde{x}}^l w|_{H^{1/2}}, \end{aligned}$$

where we use the bound in $\Lambda(T, k)$ as well as the uniform bound L from Definition 2 for the lower-order terms in the last inequality. Therefore, the proof is finished. \square

Now we can finally finish the proof of our theorem. In fact, applying Lemma 9 to the right side of (73), we obtain

$$\frac{1}{2} \frac{d}{dt} \mathcal{E}(t) \leq P(\mathcal{E}(t)).$$

As a result, the desired energy estimate is proved. Moreover, we see from this inequality immediately that a small positive T^* exists and depends only on the initial values. \square

Remark 12. The weighted norms can be better understood if we pull them back into the norms in the physical domain Ω . For the sake of simplicity, we only focus on the left corner and omit the details of computations, and the case for the right corner follows similarly. In fact, looking at the part of domain near the left corner, we know that $|\mathcal{T}'| \approx \alpha \approx r$ where r is the radius with respect to the left corner point.

Therefore, we have the following settings:

(1) For $\alpha w = \alpha \tilde{\kappa} \in H^{k+1/2}$, direct transformation shows immediately that

$$r^{1/2} (r \partial_r)^l \tilde{\kappa} \in L^2 \quad \text{near the left corner, } l \leq k,$$

which corresponds to the homogeneous norm defined in [38].

(2) For $\alpha^{-\gamma} \tilde{D}_t w - a_{w_t} \in H^k$, we only look at the L^2 norm for simplicity:

$$r^{-1/2} (r^{-\gamma} D_t \tilde{\kappa} - a_{w_t}) \in L^2 \quad \text{near the left corner.}$$

As a result, we show by the energy estimate that

$$\alpha^{-\gamma-2}\tilde{v} - a_v \in H^{k+5/2}(\mathcal{S}_0), \quad \alpha^{-2\gamma-4}\tilde{P} - a_p \in H^{k+7/2}(\mathcal{S}_0),$$

which means

$$r^{-1}(r^{-\gamma-2}v - a_v) \in L^2, \quad r^{-1}(r^{-2\gamma-4}P - a_p) \in L^2 \quad \text{near the left corner.}$$

Moreover, according to our estimates for time derivatives, we can also see that there are more weights in time derivatives, which means faster decay for time derivatives near the corners.

ACKNOWLEDGEMENTS

This work started from a discussion with Chongchun Zeng in 2019. The author would like to thank Chongchun Zeng, David Lannes and Chao Wang for fruitful discussions and helpful suggestions. This work is supported by National Natural Science Foundation of China no.12071415 and no.12222116.

REFERENCES

- [1] S. Agrawal, Rigidity of singularities of 2D gravity water waves. *J. Differential Equations*, **268** (2020), no.3, 1220–1249.
- [2] S. Agrawal, T. Alazard, Refined Rellich boundary inequalities for the derivatives of a harmonic function. *Proc. Amer. Math. Soc.*, **151** (2023), 2103–2113.
- [3] T. Alazard, N. Burq, C. Zuily, On the water-wave equations with surface tension, *Duke Math. J.* **158** (2011), no.3, 413–499.
- [4] T. Alazard, N. Burq, and C. Zuily, Cauchy theory for the gravity water waves system with non localized initial data. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **33** (2016), no.2, 337–395.
- [5] T. Alazard and J. M. Delort, Global solutions and asymptotic behavior for two dimensional gravity water waves. *Ann. Sci. Ec. Norm. Super.*, **48**(2015) no.5, 1149–1238.
- [6] B. Alvarez-Samaniego and D. Lannes, Large time existence for 3D water-waves and asymptotics. *Invent. Math.*, **171**(2008), 485–541.
- [7] D. M. Ambrose and N. Masmoudi, The zero surface tension limit of two-dimensional water waves. *Comm. Pure Appl. Math.*, **58**(2005), 1287–1315.
- [8] G. Beck, D. Lannes, L. Weynans, A numerical method for wave-structure interactions in the Boussinesq regime. arXiv:2307.01749.
- [9] K. Beyer and M. Günther, On the Cauchy problem for a capillary drop. I. Irrotational motion. *Math. Methods Appl. Sci.*, (12)**21**(1998), 1149–1183.
- [10] E. Bocchi, Compressible-incompressible transitions in fluid mechanics: waves-structures interaction and rotating fluids. Differential Geometry [math.DG]. Université de Bordeaux, 2019. English. NNT: 2019BORD0279. tel-02371058.
- [11] M. Borsuk, V. Kondratiev, Elliptic boundary value problems of second order in piecewise smooth domains, North-Holland Mathematical Library, Volume 69, Elsevier, 2006.
- [12] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo and M. López-Fernández, Rayleigh-Taylor breakdown for the Muskat problem with applications to water waves. *Ann. of Math.*, **175** (2012), 909–948.
- [13] A. Castro, D. Córdoba, C. Fefferman, F. Gancedo and J. Gómez-Serrano, Finite time singularities for the free boundary incompressible Euler equations. *Ann. of Math.*, **178** (2013), 1061–1134.
- [14] D. Christodoulou and H. Lindblad, On the motion of the free surface of a liquid. *Comm. Pure Appl. Math.*, (12)**53**(2000), 1536–1602.
- [15] D. Córdoba, A. Enciso and N. Grubic, Local well-posedness for the free boundary incompressible Euler equations with interfaces that exhibit cusps and corners of nonconstant angle, *Advances in Mathematics*, **433**(2023), 109299.
- [16] D. Córdoba, A. Enciso and N. Grubic, Finite-time singularity formation for angled-crested water waves, arXiv:2303.00027.
- [17] D. Coutand and S. Shkoller, Well-posedness of the free-surface incompressible Euler equations with or without surface tension. *J. Amer. Math. Soc.*, **20** (2007), 829–930.
- [18] D. Coutand and S. Shkoller, On the finite-time splash and splat singularities for the 3-D free-surface Euler equations. *Comm. Math. Phys.* **325** (2014), 143–183.
- [19] W. Craig, An existence theory for water waves and the Boussinesq and Korteweg-de Vries scaling limits. *Comm. Partial Differential Equations*, **10**(1985), 787–1003.
- [20] T. de Poyferré, A priori estimates for water waves with emerging bottom. *Arch. Ration. Mech. Anal.*, **232** (2019), no.2, 763–812.
- [21] Y. Deng, A. D. Ionescu, B. Pausader and F. Pusateri, Global solutions of the gravity-capillary water-wave system in three dimensions, *Acta Math.*, **219** (2017), no.2, 213–402.
- [22] B. Dyda, A fractional order Hardy inequality. *Ill. J. Math.*, 48(2004), no.2, 575–588.
- [23] J.-F. Gerbeau and T. Lelievre, Generalized Navier boundary condition and geometric conservation law for surface tension. *Comput. Methods Appl. Mech. Engrg.*, **198** (2009), 644–656.

- [24] P. Germain, N. Masmoudi and J. Shatah, Global solutions for the gravity surface water waves equation in dimension 3. *Ann. of Math.*, **175**(2012), 691–754.
- [25] P. Grisvard, Elliptic Problems in Non Smooth Domains. *Pitman Advanced Publishing Program, Boston-London-Melbourne*, 1985.
- [26] Y. Guo and I. Tice, Stability of contact lines in fluids: 2D Stokes Flow. *Arch. Ration. Mech. Anal.*, **227** (2018), no.2, 767–854.
- [27] Y. Guo and I. Tice, Stability of contact lines in fluids: 2D Navier-Stokes flow, arXiv:2010.15713v1.
- [28] J. Hunter, M. Ifrim and D. Tataru, Two dimensional water waves in holomorphic coordinates. *Comm. Math. Phys.*, **346** (2016), no.2, 483–552.
- [29] T. Iguchi, Well-posedness of the initial value problem for capillary-gravity waves. *Funkcial. Ekvac.*, **44**(2001) no.2, 219–241.
- [30] T. Iguchi and D. Lannes, Hyperbolic free boundary problems and applications to wave-structure iterations. *Indiana Univ. Math. J.*, **70** (2021), no.1, 353–464.
- [31] T. Iguchi and D. Lannes, The 2D nonlinear shallow water equations with a partially immersed obstacle. arXiv:2306.15285.
- [32] T. Iguchi, N. Tanaka and A. Tani, On a free boundary problem for an incompressible ideal fluid in two space dimensions. *Adv. Math. Sci. Appl.*, **9**(1999), 415–472.
- [33] A. D. Ionescu and F. Pusateri, Global solutions for the gravity water waves system in 2D. *Invent. Math.*, (3)**199**(2015), 653–804.
- [34] R. H. Kinsey and S. Wu, A Priori Estimates for Two-Dimensional Water Waves with Angled Crests. *Camb. J. Math.*, **6**(2018), no.2, 93–181.
- [35] H. Knüpfer and N. Masmoudi, Well-posedness and uniform bounds for a nonlocal third order evolution operator on an infinite wedge. *Comm. Math. Phys.*, **320** (2013), no.2, 395–424.
- [36] H. Knüpfer and N. Masmoudi, Darcy’s flow with prescribed contact angle: well-posedness and lubrication approximation. *Arch. Ration. Mech. Anal.*, **218** (2015), no.2, 589–646.
- [37] V. A. Kondratiev, Boundary value problems for elliptic equations in conical regions, *Soviet Math. Dokl.*, **4**(1963).
- [38] V. A. Kozlov, V. G. Mazya and J. Rossmann, Elliptic Boundary Value Problems in Domains with Point Singularities. *Mathematical Surveys and Monographs*, **52**, American Mathematical Society, Providence, RI, 1997.
- [39] V.A. Kozlov, V.G. Maz’ya, J.Rossmann, Spectral Problems Associated with Corner Singularities of Solutions to Elliptic Equations, *Mathematical Surveys and Monographs*, Volume 85, American Mathematical Society, 2001.
- [40] D. Lannes, Well-posedness of the water-wave equations. *Journal of the American Math. Society*, **18**(2005) no.3, 605–654.
- [41] D. Lannes, The water waves problem. Mathematical analysis and asymptotics. *Mathematical Surveys and Monographs*, Vol. 188. American Mathematical Society, Providence, RI, 2013. xx+321 pp.
- [42] D. Lannes, On the dynamics of floating structures. *Ann. PDE*, **3** (2017) no.1, Art. 11, 81 pp.
- [43] D. Lannes and G. Métivier, The shoreline problem for the one-dimensional shallow water and Green-Naghdi equations. *J. Éc. polytech. Math.*, **5**(2018), 455–518.
- [44] T. Levi-Civita. Détermination rigoureuse des ondes permanentes d’amplitude finie. *Math. Ann.*, **93**(1925), no.1, 264–314.
- [45] H. Lindblad, Well-posedness for the motion of an incompressible liquid with free surface boundary. *Ann. of Math.*, **162**(2005), 109–194.
- [46] M. Ming and C. Wang, Elliptic estimates for D-N operator on corner domains. *Asymptotic analysis*, **104**(2017), 103–166.
- [47] M. Ming and C. Wang, Water waves problem with surface tension in a corner domain I: A priori estimates with constrained contact angle. *SIAM. J. Math. Anal.*, **52**(2020), no.5, 4861–4899.
- [48] M. Ming and C. Wang, Water waves problem with surface tension in a corner domain II: The local well-posedness. *Commun. Pure Appl. Math.*, **74** (2021), no. 2, 225–285.
- [49] M. Ming and C. Wang, Local well-posedness of the capillary-gravity water waves with acute contact angles, arXiv: 2112.14001.
- [50] M. Ming and Z. Zhang, Well-posedness of the water-wave problem with surface tension. *J. Math. Pures Appl.*, **92**(2009), 429–455.
- [51] V. I. Nalimov, The Cauchy-Poisson problem (in Russian). *Dynamika Splosh. Sredy*, **18**(1974), 104–210.
- [52] M. Ogawa and A. Tani, Free boundary problem for an incompressible ideal fluid with surface tension. *Math. Models Methods Appl. Sci.*, **12**(2002), 1725–1740.
- [53] Ch. Pommerenke, Boundary behaviour of conformal maps, Springer-Verlag Berlin Heidelberg 1992.
- [54] W. Ren and W. E, Boundary conditions for the moving contact line problem. *Physics of Fluids*, **19**, 022101(2007), 1–15.
- [55] B. Schweizer, On the three-dimensional Euler equations with a free boundary subject to surface tension. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, (6)**22**(2005), 753–781.
- [56] J. Shatah, S. Walsh, C. Zeng, Travelling water waves with compactly supported vorticity, *Nonlinearity*, **26** (2013), no. 6, 1529–1564.
- [57] J. Shatah, C. Zeng, Geometry and a priori estimates for free boundary problems of the Euler equation. *Commun. Pure Appl. Math.*, **61**(2008), 698–744.

- [58] J. Shatah, C. Zeng, Local well-posedness for the fluid interface problems. *Arch. Ration. Mech. Anal.*, (2)**199**(2011), 653–705.
- [59] G. G. Stokes, On the theory of oscillatory waves. *Trans. Cambridge Philos. Soc.*, **8**(1847), 441–455.
- [60] G. G. Stokes, Considerations relative to the greatest height of oscillatory irrotational waves which can be propagated without change of form. *Mathematical and physical papers*, **1**(1880), 225–228, Cambridge.
- [61] Q. Su, Long time behavior of 2D water waves with point vortices. *Comm. Math. Phys.*, **380**(2020), no.3, 1173–1266.
- [62] I. Tice and Y. Zheng, Local well-posedness of the contact line problem in 2D Stokes flow. *SIAM J. Math. Anal.*,**49** (2017), no.2, 899–953.
- [63] X. Wang, Global infinite energy solutions for the 2D gravity water waves system. *Comm. Pure Appl. Math.*, **71** (2018), no.1, 90–162.
- [64] C. Wang, Z. Zhang, W. Zhao and Y. Zheng, Local well-posedness and break-down criterion of the incompressible Euler equations with free boundary. *Mem. Amer. Math. Soc.*, **270** (2021), no.1318, v+119 pp.
- [65] S. Wu, Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent. Math.*, **1**(1997) no.130, 39–72.
- [66] S. Wu, Well-posedness in Sobolev spaces of the full water wave problem in 3-D. *J. Amer. Math. Soc.*, **12**(1999), 445–495.
- [67] S. Wu, A blow-up criteria and the existence of 2d gravity water waves with angled crests, arXiv:1502.05342.
- [68] S. Wu, Global well-posedness of the 3-D full water wave problem. *Invent. Math.*, **184**(2011), 125–220.
- [69] H. Yosihara, Gravity waves on the free surface of an incompressible perfect fluid of finite depth. *Publ. Res. Inst. Math. Sci.*, **18**(1982), 49–96.
- [70] H. Yosihara, Capillary-gravity waves for an incompressible ideal fluid. *J. Math, Kyoto Univ.*, **23**(1983) no.4, 649–694.
- [71] P. Zhang and Z. Zhang, On the free boundary problem of three-dimensional incompressible Euler equations. *Comm. Pure Appl. Math.*, **61**(2008), 877–940.

SCHOOL OF MATHEMATICS AND STATISTICS, YUNNAN UNIVERSITY, KUNMING 615000, P.R.CHINA
 Email address: mingmei@ynu.edu.cn