

Bound star clusters observed in a lensed galaxy

460 Myr after the Big Bang

Angela Adamo^{1*}, Larry D. Bradley^{2†}, Eros Vanzella^{3†},
Adélaïde Claeysens^{N1}, Brian Welch^{4,5}, Jose M Diego⁶,
Guillaume Mahler^{7,8,9}, Masamune Oguri^{10,11}, Keren Sharon¹²,
Abdurro'uf^{13,2}, Tiger Yu-Yang Hsiao^{13,2}, Xinfeng Xu^{14, 15},
Matteo Messa³, Augusto E. Lassen^{16,1}, Erik Zackrisson^{17,18},
Gabriel Brammer^{19,20}, Dan Coe^{2,13,21}, Vasily Kokorev²²,
Massimo Ricotti⁴, Adi Zitrin²³, Seiji Fujimoto²⁴,
Akio K. Inoue^{25,26}, Tom Resseguier¹³, Jane R. Rigby⁵,
Yolanda Jiménez-Teja^{27,28}, Rogier A. Windhorst²⁹,
Takuya Hashimoto^{30,31}, Yoichi Tamura³²

^{1*}Astronomy Department, Stockholm University & Oskar Klein Centre,
Roslagstullsbacken 21, Stockholm, SE-10691, Sweden.

²Space Telescope Science Institute (STScI), 3700 San Martin Drive,
Baltimore, 21102, MD, USA.

³Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, INAF, via
Gobetti 93/3, Bologna, 40129, Italy.

⁴Department of Astronomy, University of Maryland, 4296 Stadium
Drive, College Park, 20742-2421, USA.

⁵Astrophysics Science Division, Code 660, NASA Goddard Space Flight
Center, 8800 Greenbelt Rd., Greenbelt, MD, 20771, USA.

⁶Instituto de Física de Cantabria , (CSIC-UC), Avda. Los Castros s/n.,
Santander, 39005, Spain.

⁷STAR Institute, Quartier Agora - Allée du six Août, 19c, Liège, B-4000,
Belgium.

⁸Centre for Extragalactic Astronomy, Durham University, South Road,
Durham, DH1 3LE, UK.

⁹Institute for Computational Cosmology, Durham University, South
Road, Durham, DH1 3LE, UK.

- ¹⁰Center for Frontier Science, Chiba University, 1-33 Yayoi-cho, Inage-ku, Chiba, 263-8522, Japan.
- ¹¹Department of Physics, Graduate School of Science, Chiba University, 1-33 Yayoi-cho, Inage-ku, Chiba, 263-8522, Japan.
- ¹²Department of Astronomy, University of Michigan, 1085 S. University Ave, Ann Arbor, MI 48109, USA.
- ¹³Center for Astrophysical Sciences, Department of Physics and Astronomy, The Johns Hopkins University, 3400 N Charles St., Baltimore, MD 21218, USA.
- ¹⁴Department of Physics and Astronomy, Northwestern University, 2145 Sheridan Road, Evanston, IL, 60208, USA.
- ¹⁵Center for Interdisciplinary Exploration and Research in Astrophysics (CIERA), Northwestern University, 1800 Sherman Avenue, Evanston, IL, 60201, USA.
- ¹⁶Instituto de Física, Departamento de Astronomia, Universidade Federal do Rio Grande do Sul, Avenida Bento Gonçalves, 9500, Porto Alegre, 91509-900, Brazil.
- ¹⁷Observational Astrophysics, Department of Physics and Astronomy, Uppsala University, Box 516, Uppsala, SE-751 20, Sweden.
- ¹⁸Swedish Collegium for Advanced Study, Linneanum, Thunbergsvägen 2, Uppsala, SE-752 38 Uppsala, Sweden.
- ¹⁹Cosmic Dawn Center (DAWN), Copenhagen, Denmark.
- ²⁰Niels Bohr Institute, University of Copenhagen, Jagtvej 128, Copenhagen, Denmark.
- ²¹Association of Universities for Research in Astronomy (AURA) for the European Space Agency (ESA), STScI, Baltimore, MD, USA.
- ²²Kapteyn Astronomical Institute, University of Groningen, Landleven 12, Groningen, 9700 AV, Netherlands.
- ²³Department of Physics, Ben-Gurion University of the Negev, P.O. Box 653, Be'er-Sheva, 84105, Israel.
- ²⁴Department of Astronomy, The University of Texas at Austin, Austin, TX 78712, USA.
- ²⁵Department of Physics, School of Advanced Science and Engineering, Faculty of Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku, Tokyo, 169-8555, Japan.
- ²⁶Waseda Research Institute for Science and Engineering, Faculty of Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku, Tokyo, 169-8555, Japan.
- ²⁷Instituto de Astrofísica de Andalucía, (CSIC), Glorieta de la Astronomía s/n., Granada, 18008, Spain.

²⁸Observatório Nacional, (MCTI), Rua Gal. José Cristino 77, São Cristóvão, Rio de Janeiro, 20921-400, Brazil.

²⁹School of Earth and Space Exploration, Arizona State University, Tempe, AZ 85287-1404, USA.

³⁰Division of Physics, Faculty of Pure and Applied Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki, 305-8571, Japan.

³¹Tomonaga Center for the History of the Universe (TCHoU), University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki, 305-8571, Japan.

³²Department of Physics, Graduate School of Science, Nagoya University, Furo, Chikusa, Nagoya, 464-8602, Japan.

*Corresponding author(s). E-mail(s): angela.adamo@astro.su.se;

†These authors contributed equally to this work.

The Cosmic Gems arc is among the brightest and highly magnified galaxies observed at redshift $z \sim 10.2$ [1]. However, it is an intrinsically UV faint galaxy, in the range of those now thought to drive the reionization of the universe [2–4]. Hitherto the smallest features resolved in a galaxy at a comparable redshift are between a few hundreds and a few tens of parsecs [5, 6]. Here we report JWST observations of the Cosmic Gems. The light of the galaxy is resolved into five star clusters located in a region smaller than 70 parsec. They exhibit minimal dust attenuation and low metallicity, ages younger than 50 Myr and intrinsic masses of $\sim 10^6 M_{\odot}$. Their lensing-corrected sizes are approximately 1 pc, resulting in stellar surface densities near $10^5 M_{\odot}/\text{pc}^2$, three orders of magnitude higher than typical young star clusters in the local universe [7]. Despite the uncertainties inherent to the lensing model, they are consistent with being gravitationally bound stellar systems, i.e., proto-globular clusters. We conclude that star cluster formation and feedback likely contributed

to shape the properties of galaxies during the epoch of reionization.

The Cosmic Gems arc (SPT0615-JD1) was initially discovered in HST images obtained by the RELICS survey of the lensing galaxy cluster SPT-CL J0615–5746 at $z = 0.972$ and reported as a redshift $z = 10$ candidate [8].

A recent JWST/NIRCam imaging campaign of this field has observed the Cosmic Gems arc with eight bands covering 0.8 to 5 μm range (Methods). Spectral energy distribution (SED) fitting to the JWST photometry indicates that the Cosmic Gems galaxy has a fairly young stellar population with mass-weighted age < 79 Myr and a lensing-corrected stellar mass in the range $2.4\text{--}5.6 \times 10^7 M_{\odot}$, with low dust extinction ($A_V < 0.15$ mag) and metallicity ($< 1\% Z_{\odot}$) [1].

The FUV-to-optical rest-frame of Cosmic Gems arc reveals bright clumpy structures and extended faint emission over a $5''$ -long arc (Fig. 1). The symmetry between the south-east (hereafter *Img.1*) and the north-west (*Img.2*) part of the arc uncovers two lensed mirror images of the galaxy, implying that the Cosmic Gems arc is observed at very high magnification on the lensing critical curve. Four independent magnification models have been created to account for the galaxy appearance. All the models successfully reproduce the $z = 10.2$ critical line crossing the Cosmic Gems arc (Methods).

Five star cluster candidates are uniquely identified in *Img.1*. In *Img.2*, three sources are clearly distinguishable in the F150W filter (Fig. 1), along with a fourth fainter source (E.2). The appearance of *Img.2* is likely perturbed by further lensing effects due to the northern galaxy at $z \sim 2.6$ (visible in the upper right corner) and possibly by an undetected small scale perturber closer to the arc [1]. Source D.2 is possibly blended with C.2, and is therefore identified for the remaining of the analysis as C.2+D.2. The source E.2 is only detected at a 2σ level and not included in this analysis (Methods). The observed projected distance between the A.1 and E.1 clusters in *Img.1* is about

0.65". Using forward modelling and *Lenstool-A* predictions presented in Methods, we find that their physical distance is 42_{-5}^{+29} pc. The star clusters are all located within this compact region (Fig. 1). The size of this region is similar to that reported for individual stellar clumps observed in the moderately-lensed galaxies at redshift $z \sim 10$ [2, 5].

Table 1: Estimated physical properties of the Cosmic Gems arc star clusters. We report de-convolved observed half-light radii in pixels, $R_{\text{eff,obs}}$, and lensing-corrected R_{eff} and median stellar masses $M_{*,int}$ using magnifications produced by the reference *Lenstool-A* model. M_* are recovered from the BAGPIPES-*exp* reference fit. Errors are estimated from 68 % confidence level of the distributions. These quantities have been used to determine stellar surface density, Σ_* , and dynamical age Π listed in the last columns. Evaluation of magnification uncertainties are discussed in Methods.

ID	$R_{\text{eff,obs}}$ [px]	R_{eff} [pc]	$R_{\text{eff,FM}}$ [pc]	$M_{*,int}$ [$10^6 M_{\odot}$]	Σ_* [$10^5 M_{\odot} / \text{pc}^2$]	$\log(\Pi)$
A.1	$0.6_{-0.1}^{+0.4}$	$1.1_{-0.2}^{+0.7}$	1.1 ± 0.1	$2.45_{-1.56}^{+5.20}$	$1.92_{-1.44}^{+1.60}$	$1.94_{-0.27}^{+0.71}$
B.1	$1.1_{-0.5}^{+0.1}$	$1.1_{-0.5}^{+0.1}$	0.9 ± 0.1	$2.65_{-1.26}^{+1.09}$	$1.93_{-1.11}^{+4.16}$	$2.11_{-0.50}^{+0.83}$
C.1	< 1.25	< 1	0.9 ± 0.2	$1.13_{-0.65}^{+1.77}$	> 1.3	> 1.90
D.1	$1.2_{-1.1}^{+0.2}$	$0.6_{-0.6}^{+0.1}$	0.8 ± 0.2	$1.13_{-0.74}^{+1.23}$	$2.39_{-1.98}^{+7.41}$	$2.17_{-1.03}^{+0.85}$
E.1	$1.5_{-0.5}^{+0.7}$	$0.4_{-0.1}^{+0.2}$	0.7 ± 0.2	$1.01_{-0.36}^{+0.37}$	$6.92_{-4.22}^{+4.90}$	$3.06_{-0.59}^{+0.32}$
A.2	$1.0_{-0.3}^{+0.4}$	$1.7_{-0.4}^{+0.8}$	1.3 ± 0.1	$2.89_{-1.35}^{+1.96}$	$0.88_{-0.46}^{+0.98}$	$1.94_{-0.41}^{+0.52}$
B.2	< 1.25	< 1.4	1.0 ± 0.04	$3.01_{-1.61}^{+3.21}$	> 5.10	> 1.8
C.2+D.2	$2.7_{-2.6}^{+16.1}$	$1.9_{-1.9}^{+11.4}$	0.9 ± 0.1	$4.36_{-1.90}^{+0.98}$	$1.05_{-0.89}^{+7.89}$	$1.99_{-1.49}^{+1.12}$

The intrinsic physical properties of these 5 star clusters are particularly meaningful for probing proto-GC formation mechanisms as well as their potential evolution. As described in Methods, we find that A.1 is only marginally resolved, with an observed effective half-light radius $R_{\text{eff,obs}} = 0.6$ px, while C.1 and B.2 are consistent with being unresolved. For the latter sources, we assumed an upper-limit to their radii coincident with the half-width half maximum (HWHM) of the stellar PSF in the F150W ($0.025'' = 1.25$ px). To derive lensing-corrected R_{eff} , we assumed the predicted *Lenstool-A* tangential magnifications at the location of the star clusters (listed in Extended Data Table 3). The 5 star clusters have intrinsic R_{eff} close to 1 pc (within

uncertainties, Table 1). Using the other lensing models produces similar size ranges (0.3 to 0.9 pc for *Lenstool-B* and 0.3 to 1.2 pc for *Glafic*). Independent intrinsic sizes ($R_{\text{eff,FM}}$ in Table 1) have been derived by projecting the star cluster shapes from the source plane into the image plane. The latter method recovered intrinsic sizes in excellent agreement, within the uncertainties, with those measured in the image plane, strengthening the reliability of the derived values.

The star clusters have been fitted with BAGPIPES [9] and PROSPECTOR [10]. We tested different star formation history (SFH) assumptions that simulate a single burst (inherent to the small sizes of the stellar systems analysed), different high-mass limits of the initial mass function (IMF), and models with stellar binaries (presented in Methods). Despite the assumptions, the resulting cluster physical properties (ages, masses, extinction, metallicities) are in reasonable agreement. In the analysis presented here, we use the physical values derived with a SFH based on a single exponential decline with $\tau = 1$ Myr (listed in Extended Data Table 2).

The recovered ages of the star cluster candidates are between 9 and 36 Myr. The age range suggests that star formation has been propagating within this compact area of the galaxy for a few tens of Myr. The measured rest-frame UV slopes of the star clusters (β between -1.8 and -2.5 , with $F_\lambda \sim \lambda^\beta$, listed in Extended Data Table 1) are similar to those found for more evolved star clusters in the Sunburst arc at redshift 2.37 [11]. While the Cosmic Gems clusters are not extremely young, they have likely delivered large amounts of energy and momentum to their host galaxy.

The lensing-corrected stellar masses range between $1.0 - 2.6 \times 10^6 M_\odot$, for a total combined stellar mass of $8.3 \times 10^6 M_\odot$. The total mass of the clusters is close to 30% of the total stellar mass of the host. Since the mass-weighted age of the galaxy and those of the star clusters are comparable, one can extrapolate the cluster formation efficiency (CFE) [12] to be around 30 %. A caution note is necessary, since the mass estimates (both for the galaxy and star clusters) are subjected to magnification values

and SED fit uncertainties, making the quoted CFE uncertain. A more direct way to establish the CFE is to use the fraction of observed FUV light in star clusters with respect to the host. This quantity is not affected by the same degeneracy as the mass estimates and, thus, is a more reliable indicator of the CFE, under the assumption that the FUV light is produced by stellar populations formed during a similar timescale (as we find here). The analysed star clusters account for $\sim 60\%$ of the total F150W flux of the host extracted within an elliptical Kron aperture ($0.51 \pm 0.01 \mu\text{Jy}$, corresponding to an intrinsic FUV ABmag of -17.8 after lensing correction, [1]), thus reinforcing the conclusion that star formation in star clusters is a major mode for the Cosmic Gems arc and high-redshift galaxies with similar physical properties. This observationally driven conclusion is supported by high-resolution numerical simulations [13] and analytical models [14] that find that compact star clusters with sizes $0.5 - 2$ pc are the dominant star formation mode in the first low-metallicity dwarf galaxies. The compactness of the star clusters appears also to drive the leakage of hydrogen ionizing radiation from their natal molecular cloud [15], making the star clusters observed here potential contributors to cosmic reionization. Massive star clusters like those observed in the Cosmic Gems arc are predicted by the feedback-free starburst model by [16], and could be at the root of the super-Eddington conditions necessary to launch strong outflows in short timescales [17], both models aimed to explain the bright UV luminosity reported for $z > 9$ galaxies.

The resulting stellar surface densities of the clusters are $\sim 10^5 M_{\odot}/\text{pc}^2$ (Table 1). Consistent physical properties have been reported in star clusters detected in the Sunrise arc at $z \sim 6$ [18] and the Sunburst arc at $z = 2.37$ [19] (see Fig. 2). Using the derived ages, masses, and intrinsic sizes, we also determine whether these stellar systems are gravitationally bound. According to the framework introduced by [20], a star cluster is considered bound if its age is greater than the crossing time of the system (where $t_{\text{cross}} = 10\sqrt{R_{\text{eff}}^3/GM}$), or in other words, under the assumption of

virial equilibrium, a cluster is gravitationally bound if $\Pi = \text{Age}/t_{\text{cross}} > 1$. The $\log(\Pi)$ values reported in Table 1 are all significantly larger than unity,

indicating that we are indeed detecting star clusters in an early galaxy, 460 Myr after the Big Bang. This conclusion is valid in spite of the uncertainties inherent to physical quantity estimates as well as lensing models.

The Cosmic Gems arc clusters (Fig. 2) have significantly higher stellar densities and smaller sizes than typical young star clusters observed in the local universe [7] as well as GCs in the Milky Way [21]. The offset with respect to young star clusters in the local universe is expected since the conditions under which star formation operates in reionization-era galaxies are more extreme (e.g, more compact, harbour harder ionising radiation fields, and reach higher electron densities and temperatures,[6, 22]). The offset with respect to local GCs could be explained in terms of dynamical evolution. GCs are hot stellar systems where stars continuously exchange energy and momentum. Three different internal mechanisms contribute to their dynamical evolution over a Hubble time: (i) mass-loss due to stellar evolution; (ii) relaxation due to N-body interactions; (iii) formation and dynamics of stellar black holes (SBHs) [23, 24]. Mass loss due to stellar evolution drives GCs' adiabatic expansion under the condition of virial equilibrium. A typical mass-loss of 50 % will expand the initial radius of the cosmic Gems proto-GCs by a factor of 2, while density will decrease correspondingly by a factor of 8 [23]. Relaxation time scales (shortened by the presence of SBHs [25]) will also contribute to their expansion. Finally, external tidal fields will further affect the dynamical evolution of these bound stellar systems, which appear to be *bona fide* proto-GCs.

Very dense stellar clusters ($\Sigma_* \sim 10^5 \text{ M}_\odot / \text{pc}^2$, which for $R_{\text{eff}} = 1 \text{ pc}$ correspond to $\rho_h \sim 10^5 \text{ M}_\odot / \text{pc}^3$), like those detected in the Cosmic Gems arc, are predicted to form in low metallicity and highly dense gas [26], where radiative pressure cannot counteract the collapse, resulting in extremely high star formation efficiencies ($\sim 80\%$ [27]). The

high stellar densities found in these proto-GCs imply a significant increase in stellar BH mergers in their interiors [28, 29] and therefore pave the way to intermediate mass BH seeds [30]. With stellar masses $> 10^5 M_{\odot}$, these star clusters naturally harbour Wolf-Rayet and very massive stars [31], and because of their elevated stellar densities, satisfy the necessary condition to form supermassive stars in runaway collisions within their cores [32]. These different classes of stars are among the potential polluters that could explain the observed nitrogen enrichment in the ionised gas of high-redshift galaxies [33], possibly linked to the formation of the chemically enriched stellar populations ubiquitously found in Milky Way GCs [34].

Cosmological simulations that focus on Milky Way disk-like assembly find that the majority of its GC population form at redshift $z < 7$ [35–37], suggesting that these star clusters forming at $z \sim 10$ might build up the GC populations of more massive early type galaxies in the local Universe. It is difficult to predict whether the proto-GCs of the Cosmic Gems arc will survive a Hubble time. Their chances would be highly enhanced if they were ejected into their host halo during dynamical interactions (e.g., [12]).

Methods

Size & flux measurements

The JWST NIRCам [38] observations of the SPT-CL J0615–5746 galaxy cluster were obtained in 2023 September (GO 4212: PI Bradley) using four short-wavelength (SW) filters (F090W, F115W, F150W, and F200W) and four long-wavelength (LW) filters (F277W, F356W, F410M, and F444W) spanning $0.8 - 5.0 \mu m$. Each filter had an exposure time of 2920.4 s. The data were reduced with the GRIZLI (version 1.9.5) reduction package [39]. They suffered from strong wisps [40] and required special background subtraction as described in [1]. The final data are in units of 10 nJy. The NIRCам SW (LW) images were drizzled to a pixel scale of 0.02 (0.04)"/pixel. For

further details on the observations and image reduction, please see [1]. We assume throughout the analysis a cosmology with $H_0 = 67.7 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $\Omega_m = 0.31$ [41]. Under these assumptions, 1 SW pixel ($0.02''$) corresponds to 83.7 pc at $z = 10.2$.

We derived the star cluster radii and multiband photometry by applying the method published and tested in [42–44]. We simultaneously fitted for the shape of the light distribution, the flux and the local background (including the galaxy diffuse light) of each of the identified clusters in the reference filter, F150W, which offers the sharpest view of the clusters. Empirical PSFs in all filters used in this work were built by selecting stars in each band. We fitted the PSF out to $0.4''$ with an analytical expression, which was then convolved with varying 2D Gaussians to create a grid of models. Intrinsic sizes (de-convolved by PSF) have then been derived by fitting the observed light distribution of each cluster with this grid of 2D Gaussian models in the F150W. Due to the large shear effects, compact sources might appear resolved in the shear direction (y_{std}) [e.g., 45]. We assumed that the measured major-axis y_{std} of the 2D Gaussian ellipse is the standard deviation of a 2D circular Gaussian which we translated into the observed PSF-deconvolved effective radius, $R_{\text{eff,obs}} = y_{\text{std}} \times \sqrt{2 \ln(2)}$. The derived $R_{\text{eff,obs}}$ are reported in Table 1. The de-lensed intrinsic effective half-light radii, R_{eff} , have been determined by dividing $R_{\text{eff,obs}}$ by the tangential magnification μ_{tan} (reported in Extended Data Table 3).

Extended Data Table 1. Cluster observed properties. JWST photometry and uncertainties in ABmag and measured β slopes of the candidate star clusters.

ID	F150W	F200W	F277W	F356W	F410M	F444W	β
A.1	27.64 ± 0.08	27.83 ± 0.07	27.89 ± 0.07	28.13 ± 0.09	27.85 ± 0.08	28.03 ± 0.10	$-2.36^{+0.19}_{-0.15}$
B.1	27.39 ± 0.11	27.48 ± 0.08	27.66 ± 0.09	27.83 ± 0.10	28.07 ± 0.19	27.90 ± 0.10	$-2.42^{+0.23}_{-0.19}$
C.1	27.81 ± 0.18	27.78 ± 0.17	27.74 ± 0.10	27.86 ± 0.10	27.84 ± 0.16	27.94 ± 0.11	$-1.90^{+0.27}_{-0.34}$
D.1	27.83 ± 0.19	28.02 ± 0.19	27.76 ± 0.11	28.05 ± 0.13	27.80 ± 0.15	28.06 ± 0.12	$-1.80^{+0.35}_{-0.23}$
E.1	28.29 ± 0.28	28.39 ± 0.22	28.21 ± 0.19	28.66 ± 0.24	28.62 ± 0.33	28.24 ± 0.16	$-1.82^{+0.43}_{-0.46}$
A.2	28.08 ± 0.12	28.07 ± 0.09	28.32 ± 0.15	28.87 ± 0.21	28.41 ± 0.19	28.60 ± 0.23	$-2.35^{+0.37}_{-0.27}$
B.2	27.21 ± 0.27	27.19 ± 0.23	27.10 ± 0.10	27.33 ± 0.10	27.16 ± 0.12	27.45 ± 0.14	$-1.80^{+0.31}_{-0.36}$
C.2+D.2	27.03 ± 0.34	27.08 ± 0.20	27.28 ± 0.13	27.45 ± 0.13	27.52 ± 0.17	27.21 ± 0.14	$-2.45^{+0.46}_{-0.55}$

For each star cluster, the flux in the reference filter has been determined by integrating the fitted shape and subtracting the local background. We then measured the fluxes in the other bands by convolving the derived intrinsic shape in F150W with the empirical PSF of the respective bands. Fitting this model to the other bands, we let the centre, normalisation, and local background (including the galaxy diffuse light) as free parameters.

The intrinsic sizes and observed fluxes in the reference filter F150W were derived using a cutout box centered on the source with size of 11×11 px (about 4 times the FWHM) for A.1, A.2. Larger box sizes did not produce noticeable differences in the output sizes and fluxes. Due to their proximity, B.1, C.1, D.1, and E.1 have been fitted simultaneously within a box of 15×15 px. A larger box size, produces consistent values within the uncertainties of measurements for B.1 and C.1, while E.1 gets increasingly elongated, affecting the fit of D.1. To avoid this degeneracy, we fix the box size to 15×15 , which would correspond to fix the source ellipticity of the faint E.1 to 2. Similarly, B.2, C.2, and D.2 were fitted simultaneously within a box of 11×11 (changing the box size does not produce noticeable effects on the recovered parameters). We notice that we do not see two maxima at the location of C.2 and D.2, so we allowed the fit to optimise the centre of a second hidden source. We repeated the fit of this region by assuming only one source. Both approaches produced similar residuals. The flux extracted by assuming only one source is comparable within uncertainties to the flux extracted by fitting for C.2 and D.2. Due to the degeneration in identifying the position of D.2, we extracted physical properties by fitting only one source which we refer to as C.2+D.2. Due to the faintness of E.2 (2σ), our method did not produce significant constraints. We therefore excluded E.2 from our analysis. In the BAGPIPES-*exp* fit we find that C.1 and D.1 have similar ages and a combined total mass of about $2 \times 10^6 M_{\odot}$. C.2+D.2 has a slightly older age (but in agreement within 1σ) than C.1 and D.1.

The total mass and size of C.2+D.2 is a factor of two higher than their counterparts, corroborating the idea that the two star clusters are blended in *Img.2*.

The size uncertainties were derived by bootstrapping the fit of the source taking into account the RMS of the local background. The photometric errors include the latter uncertainties as well as the sum in quadrature of the local background variance estimated within the box where the sources have been fitted. Aperture corrections have been extrapolated up to $0.4''$ in all bands.

In Extended Data Fig. 1, we show the best model of the star clusters and the residual image in the reference filter and two more bands. The extraction of the sources does not produce significant artificial residuals above the RMS of the image.

Independent measurements of intrinsic R_{eff} have been obtained following the forward modeling method of [46]. Briefly, this method creates a model of the galaxy in the source plane, then projects that model into the image plane. After convolving with the measured empirical PSF, the image plane model is compared to the observed data. The source plane model parameters are first optimized using a downhill simplex algorithm, then sampled using an MCMC with the Python package `emcee` [47].

For SPT0615-JD1, the source plane model consisted of five Sersic profiles centered on the five identified clumps A.1–E.1 (Extended Data Fig. 2). No diffuse component of the arc has been included in this analysis, due to the faintness of this component with respect to the clusters. Separately, we modelled clumps A.2–C.2 on the other side of the lensing critical curve. Uncertainties in the *Lenstool-A* lens model resulted in slight offsets between source plane positions of clumps on opposite sides of the critical curve, which prevents simultaneous fitting of the two images of the arc. We found similar results for clump sizes on both sides of the critical curve, with clump radii ranging from 0.7–1.1 pc (see Table 1).

SED fitting analysis

We performed SED fitting with BAGPIPES [9] and test the derived physical properties against different assumptions, as well as with a different software PROSPECTOR [10]. For all the runs we fix the redshift at $z = 10.2$, as measured by [1]. The standard stellar population templates were reprocessed with CLOUDY to generate nebular continuum and line information (see [48] for comparisons of the two code implementations). In both codes we assume a Kroupa IMF, unless otherwise specified. We constrain SFHs to prescriptions that reproduce a short burst in all tests except one where τ is free to vary. The short burst assumption is in agreement with studies of stellar cluster and GC populations in the local universe ([12, 34]). The recovered median of the posterior distributions of age, mass, A_V , metallicity, and associated 68% uncertainties are reported in Extended Data Table 2. We let the ionization parameter, U , to change between -2 and -3.5 . We assumed a Calzetti attenuation [49] but test also the SMC extinction. In the reference set, used to produce results reported in Fig. 2 and Table 1 and referred to as BAGPIPES-*exp*, we assumed an exponential decline with a very short $\tau = 1$ Myr and Calzetti attenuation. The panels in Extended Data Fig. 3 show the observed SEDs of the 5 star clusters identified in *Img.1* (black dots with uncertainties). When available, we include the observed SED of the corresponding clusters in *Img.2* (orange stars with associated errors). The latter have been normalized by the median flux ratio in the 6 bands of the corresponding source in *Img.1* to match the flux level while preserving the intrinsic SED shape. The best spectral and integrated photometry model obtained for the BAGPIPES-*exp* fit is included. The overall shape of the observed SEDs of mirrored clusters in both images are similar within uncertainties, confirming the symmetry.

To check the consistency of the derived cluster physical properties, we test different assumptions. The outputs are summarised in Extended Data Table 2 where we list the median and 68% values produced by the different fits. Changing the attenuation

prescription from Calzetti to SMC, produces noticeable smaller A_v , but all the recovered parameters are still within the 68% uncertainties associated with the recovered values. In BAGPIPES-*burst*, we assumed a single burst. We recovered slightly older ages and larger masses (but notice uncertainties) which would prefer higher stellar surface densities and older dynamical ages, confirming that we are looking at dense and bound star clusters. In a third SED fitting set, BAGPIPES-*BPASS*, we used BPASS v2.2.1 SED templates [50] and the fiducial BPASS IMF with maximum stellar mass of $300 M_\odot$ and a high-mass slope similar to [51]. Also, this model reproduces values which are very close to the reference one, suggesting that the clusters are compatible with being slightly older and therefore less sensitive to presence of very massive stars/binary systems in their light (and the limitation of fitting only 6 broad/medium band covering FUV-blue optical). Letting $\tau = free$ (we report mass-weighted parameters in Extended Data Table 2) produces significantly older ages, but similar masses, thus not affecting the results presented in this article.

PROSPECTOR allows us to test single stellar population (SSP) SFH. In this case, we find that the age of A.1 is slightly younger (but within uncertainties) than those produced by the BAGPIPES, resulting in lower masses. This would result in slightly lower intrinsic mass $M_{*,int} = 0.39 \times 10^6 M_\odot$, $\log(\Sigma_*) = 4.5 M_\odot / \text{pc}^2$, and $\log(\Pi) = 1.2$, but leaving unchanged any of the conclusions of the article.

Finally, given that some of the knots in the Cosmic Gems arc are unresolved (C.1 and B.2) or only marginally resolved (A.1) in our current images, we have also explored scenarios in which these sources are individual, highly magnified stars. Using SED models for stars at high redshifts [52] we find that, while the slopes of the SEDs of these sources would be broadly consistent with individual stars at effective temperatures $\gtrsim 20000$ K, such scenarios would require magnifications well in excess of what our macrolens models predict at the positions of these sources. Even the most massive and luminous stars (initial mass 560–575 M_\odot) described by the stellar evolutionary tracks

of [53] would require magnifications $\mu > 1000$ to explain the observed fluxes of C.1, B.2 or A.1.

Lens Models & uncertainties on the derived star cluster physical properties

Four different lensing models have been created for the SPT-CL J0615–5746 cosmological field. The models are presented in detail in [1]. We include here below a short description.

Lenstool-A, here used as reference model for the analysis presented in this letter, is based on the software LENSTOOL [54] which uses a parametric approach and MCMC sampling of the parameter space to identify the best-fit model and uncertainties. In *Lenstool-A*, we model the cluster lens as a combination of three main halos and cluster member galaxies, all parameterized as pseudo-isothermal mass distributions. The model uses as constraints the positions of 43 multiple images of 14 clumps, belonging to 9 unique source galaxies. The redshifts of three sources are used as constraints (the $z = 10.2$ arc, and sources at $z = 1.358$ and $z = 4.013$ [55]), while the rest of the redshifts are treated as free parameters. Three clumps on each side of the main arc were used as constraints, A, B, and C, assumed to be at $z = 10.2$. The model predicts a counter image at (R.A., Decl.)=(93.9490607, -57.7701814). A possible candidate of this counter image, observed near this location ($\sim 1.8''$), was not used as constraint. The image plane rms of the best-fit model is $0.36''$. All the observed lensed features are well-reproduced by this model.

The second model, here referred to as *Lenstool-B*, uses the same algorithm, but with noticeable different assumptions. This model uses 43 multiple images from 11 unique sources. A secondary cluster scale halo is placed around the location of dusty galaxies nearly 50 arcsec north of the bright centre galaxy and allowed to move within a $20''$ box around this position. As with the previous model, the $z = 10.2$ arc has a

predicted counter image near the possible candidate and is only about $2''$ away from the *Lenstool-A* model. The main differences between those models are the assumptions about the mass distribution of the lens and the addition of constraints. The image plane rms of the best-fit model is $0.68''$.

The third model used in this analysis has been created with `glafic`. The `glafic` [56, 57] mass model is constructed with three elliptical NFW [58] halos, external shear, and cluster member galaxies modeled by pseudo-Jaffe profile. The model parameters are fitted to reproduce the position of 44 multiple images generated from 15 background sources. Spectroscopic redshifts are available for 7 of the 15 sources. We include positions of A.1/A.2 and B.1/B.2 in the Cosmic Gems arc as constraints, with small positional errors of $0.04''$ to accurately predict magnifications of each star cluster images. For the other multiple images, we adopt the positional error of $0.4''$. Our best-fitting model reproduces all the multiple image positions with the root-mean-square of image positions of $0.41''$.

As a consistency check, we excluded the positional constraints from the Cosmic Gems arc to construct the mass model, and confirmed that the critical curve of this mass model still pass through the arc. Our `glafic` best-fitting mass model also predicts a counter-image of the Cosmic Gems arc at around (R.A., Dec.)=(93.9504865, -57.7696559). We find that there is a candidate counter-image at $\sim 2''$ from the predicted position, (R.A., Dec.)=(93.9500002, -57.7702197). Both the consistency check and the presence of the candidate counter-image confirm the validity of this mass model.

A fourth model has also been produced with `WSLAP+` [59, 60]. The `WSLAP+` lens models offer an alternative to parametric models and are free of assumptions made about the distribution of dark matter. When the $z = 10.2$ arc is not included as a constraint, the `WSLAP+` model predicts the critical curve passing at $\approx 1''$ from the $z = 10.2$ arc. This solution predicts a mirrored image of the arc which is not observed, reinforcing

the expectation that the Cosmic Gems is a double image with the critical curve passing through the middle. When the arc is included as a constraint, the predicted critical curve passes between C.1 and D.1, just $0.3''$ from the alleged symmetry point in the arc, and within the uncertainties typical of WSLAP+ models. Also, this model predicts the position of a third counterimage consistent with the previous models. This model is currently under development with the goal of explaining the perturbation seen in *Img.2*, and is therefore not included in this analysis.

The photometric redshift of the candidate counter image is $z_{phot} = 10.8^{+0.6}_{-1.4}$ (95% confidence) [1], in agreement with the expectation.

The total and tangential magnifications at the position of the star clusters, $\mu_{tot} = \mu_{tang} \times \mu_{rad}$, are reported in Extended Data Table 3. For the two Lenstool-based models, we estimated uncertainties following the method presented in [44] based on magnification maps produced from the lenstool MCMC posterior distributions of the lens model. Uncertainties are omitted for the glafic model.

In Extended Data Fig. 4, we show the impact that magnification predictions have in the recovered physical properties (intrinsic half-light radius and mass, black and blue solid lines) and derived quantities (dynamical age and stellar surface density in magenta and orange solid lines). We use logarithmic scales so that all quantities can be included. The colored bands show the level of uncertainties recovered from the analysis. We also include the upper-limits on the R_{eff} (assuming that the source is unresolved and has size smaller than the stellar PSF) and what type of the lower-limits it will translate for the physical quantities that depend on size estimates as dashed lines (notice that these are the reference quantities for C.1 which is unresolved). The magnifications (total in the bottom and tangential in the upper) are reported on the x axis. As we move to lower magnifications, the derived masses and radii become larger, consequently predicting lower stellar surface densities and dynamical ages. However, even in the unlikely case that the magnifications are wrong by one order of magnitude,

stellar surface density will remain above $10^4 M_{\odot}/\text{pc}^2$ and dynamical ages will still be significantly larger than 1 ($\log(\text{II}) > 0$), leaving the main conclusion of this analysis unchanged, i.e., that we are detecting bound proto-GCs within the first 500 Myr of our Universe.

Data availability

The data have been acquired under JWST Program ID 4212, with PI: Bradley. The datasets generated during and/or analysed during the current study may be obtained from the MAST archive at <https://dx.doi.org/10.17909/tcje-1780>. All data generated or analysed during this study are included in this published article (and its supplementary information files).

Code availability

This work made use of NUMPY [61], SCIPY [62], MATPLOTLIB [63] and ASTROPY [64]. SED fit analyses are performed with publicly available software BAGPIPES [9] and PROSPECTOR [10].

Acknowledgments

AA, EV, MM, AC, and MR thank ISSI for sponsoring the ISSI team: "Star Formation within rapidly evolving galaxies" where many ideas discussed in this article have been brainstormed. AA thanks Mark Gieles and Ivan Cabrera-Ziri for discussions on GC physical properties. Based on observations with the NASA/ESA/CSA JWST obtained from the Mikulski Archive for Space Telescopes (MAST) at the Space Telescope Science Institute (STScI), which is operated by the Association of Universities for Research in Astronomy (AURA), Incorporated, under NASA contract NAS5-03127. Support for Program number JWST-GO-04212 was provided through a grant from the STScI under NASA contract NAS5-03127. AA and AC acknowledge support

by the Swedish research council Vetenskapsrådet (2021-05559). J.M.D. acknowledges the support of project PID2022-138896NB-C51 (MCIU/AEI/MINECO/FEDER, UE) Ministerio de Ciencia, Investigación y Universidades. AZ acknowledges support by Grant No. 2020750 from the United States-Israel Binational Science Foundation (BSF) and Grant No. 2109066 from the United States National Science Foundation (NSF); by the Ministry of Science & Technology, Israel; and by the Israel Science Foundation Grant No. 864/23. Y.J-T. acknowledges financial support from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 898633, the MSCA IF Extensions Program of the Spanish National Research Council (CSIC), the State Agency for Research of the Spanish MCIU through the Center of Excellence Severo Ochoa award to the Instituto de Astrofísica de Andalucía (SEV-2017-0709), and grant CEX2021-001131-S funded by MCIN/AEI/ 10.13039/501100011033. EZ acknowledges project grant 2022-03804 from the Swedish Research Council (Vetenskapsrådet) and has also benefited from a sabbatical at the Swedish Collegium for Advanced Study. M.O. acknowledges the support of JSPS KAKENHI Grant Numbers JP22H01260, JP20H05856, JP22K21349. A.K.I. acknowledges the support of JSPS KAKENHI Grant Numbers JP23H00131. EV and MM acknowledge financial support through grants PRIN-MIUR 2020SKSTHZ, the INAF GO Grant 2022 “The revolution is around the corner: JWST will probe globular cluster precursors and Population III stellar clusters at cosmic dawn” and by the European Union – NextGenerationEU within PRIN 2022 project n.20229YBSAN - Globular clusters in cosmological simulations and in lensed fields: from their birth to the present epoch. T.H. is supported by Leading Initiative for Excellent Young Researchers, MEXT, Japan (HJH02007) and by JSPS KAKENHI grant No. 22H01258. Y.T. acknowledges the support of JSPS KAKENHI grant No. 22H04939 and 23K20035.

References

- [1] Bradley, L. D. *et al.* Unveiling the Cosmic Gems Arc at $z \sim 10.2$ with JWST. *arXiv e-prints* arXiv:2404.10770 (2024). [2404.10770](#).
- [2] Roberts-Borsani, G. *et al.* The nature of an ultra-faint galaxy in the cosmic dark ages seen with JWST. *Nature* **618**, 480–483 (2023). [2210.15639](#).
- [3] Atek, H. *et al.* Most of the photons that reionized the Universe came from dwarf galaxies. *Nature* **626**, 975–978 (2024).
- [4] Simmonds, C. *et al.* Low-mass bursty galaxies in JADES efficiently produce ionizing photons and could represent the main drivers of reionization. *Mon. Not. R. Astron. Soc.* **527**, 6139–6157 (2024). [2310.01112](#).
- [5] Hsiao, T. Y.-Y. *et al.* JWST Reveals a Possible $z \sim 11$ Galaxy Merger in Triply Lensed MACS0647-JD. *Astrophys. J. L.* **949**, L34 (2023). [2210.14123](#).
- [6] Morishita, T. *et al.* Enhanced Subkiloparsec-scale Star Formation: Results from a JWST Size Analysis of 341 Galaxies at $5 < z < 14$. *Astrophys. J.* **963**, 9 (2024). [2308.05018](#).
- [7] Brown, G. & Gnedin, O. Y. Radii of young star clusters in nearby galaxies. *Mon. Not. R. Astron. Soc.* **508**, 5935–5953 (2021). [2106.12420](#).
- [8] Salmon, B. *et al.* RELICS: A Candidate $z \sim 10$ Galaxy Strongly Lensed into a Spatially Resolved Arc. *Astrophys. J. L.* **864**, L22 (2018). [1801.03103](#).
- [9] Carnall, A. C., McLure, R. J., Dunlop, J. S. & Davé, R. Inferring the star formation histories of massive quiescent galaxies with BAGPIPES: evidence for multiple quenching mechanisms. *Mon. Not. R. Astron. Soc.* **480**, 4379–4401 (2018). [1712.04452](#).

- [10] Johnson, B. D., Leja, J., Conroy, C. & Speagle, J. S. Stellar Population Inference with Prospector. *Astrophys. J. S.* **254**, 22 (2021). [2012.01426](#).
- [11] Kim, K. J. *et al.* Small Region, Big Impact: Highly Anisotropic Lyman-continuum Escape from a Compact Starburst Region with Extreme Physical Properties. *Astrophys. J. L.* **955**, L17 (2023). [2305.13405](#).
- [12] Adamo, A. *et al.* Star Clusters Near and Far; Tracing Star Formation Across Cosmic Time. *Space Sci. Rev.* **216**, 69 (2020). [2005.06188](#).
- [13] Garcia, F. A. B., Ricotti, M., Sugimura, K. & Park, J. Star cluster formation and survival in the first galaxies. *Mon. Not. R. Astron. Soc.* **522**, 2495–2515 (2023). [2212.13946](#).
- [14] Nebrin, O., Giri, S. K. & Mellema, G. Starbursts in low-mass haloes at Cosmic Dawn. I. The critical halo mass for star formation. *Mon. Not. R. Astron. Soc.* **524**, 2290–2311 (2023). [2303.08024](#).
- [15] He, C.-C., Ricotti, M. & Geen, S. Simulating star clusters across cosmic time - II. Escape fraction of ionizing photons from molecular clouds. *Mon. Not. R. Astron. Soc.* **492**, 4858–4873 (2020). [2001.06109](#).
- [16] Dekel, A., Sarkar, K. C., Birnboim, Y., Mandelker, N. & Li, Z. Efficient formation of massive galaxies at cosmic dawn by feedback-free starbursts. *Mon. Not. R. Astron. Soc.* **523**, 3201–3218 (2023). [2303.04827](#).
- [17] Ferrara, A. Super-early JWST galaxies, outflows, and Ly α visibility in the Epoch of Reionization. *Astron. Astrophys.* **684**, A207 (2024). [2310.12197](#).
- [18] Vanzella, E. *et al.* JWST/NIRCam Probes Young Star Clusters in the Reionization Era Sunrise Arc. *Astrophys. J.* **945**, 53 (2023). [2211.09839](#).

- [19] Vanzella, E. *et al.* High star cluster formation efficiency in the strongly lensed Sunburst Lyman-continuum galaxy at $z = 2.37$. *Astron. Astrophys.* **659**, A2 (2022). [2106.10280](#).
- [20] Gieles, M. & Portegies Zwart, S. F. The distinction between star clusters and associations. *Mon. Not. R. Astron. Soc.* **410**, L6–L7 (2011). [1010.1720](#).
- [21] Baumgardt, H., Hénault-Brunet, V., Dickson, N. & Sollima, A. Evidence for a bottom-light initial mass function in massive star clusters. *Mon. Not. R. Astron. Soc.* **521**, 3991–4008 (2023). [2303.01636](#).
- [22] Roberts-Borsani, G. *et al.* Between the Extremes: A JWST Spectroscopic Benchmark for High Redshift Galaxies Using ~ 500 Confirmed Sources at $z \geq 5$. *arXiv e-prints* arXiv:2403.07103 (2024). [2403.07103](#).
- [23] Gieles, M., Baumgardt, H., Heggie, D. C. & Lamers, H. J. G. L. M. On the mass-radius relation of hot stellar systems. *Mon. Not. R. Astron. Soc.* **408**, L16–L20 (2010). [1007.2333](#).
- [24] Antonini, F. & Gieles, M. Population synthesis of black hole binary mergers from star clusters. *Mon. Not. R. Astron. Soc.* **492**, 2936–2954 (2020). [1906.11855](#).
- [25] Wang, L. The survival of star clusters with black hole subsystems. *Mon. Not. R. Astron. Soc.* **491**, 2413–2423 (2020). [1911.05077](#).
- [26] Fukushima, H. & Yajima, H. The formation of globular clusters with top-heavy initial mass functions. *Mon. Not. R. Astron. Soc.* **524**, 1422–1430 (2023). [2303.12405](#).
- [27] Menon, S. H., Federrath, C. & Krumholz, M. R. Outflows driven by direct and reprocessed radiation pressure in massive star clusters. *Mon. Not. R. Astron.*

- Soc.* **521**, 5160–5176 (2023). [2210.02818](#).
- [28] Choksi, N., Volonteri, M., Colpi, M., Gnedin, O. Y. & Li, H. The Star Clusters That Make Black Hole Binaries across Cosmic Time. *Astrophys. J.* **873**, 100 (2019). [1809.01164](#).
- [29] Antonini, F., Gieles, M., Dosopoulou, F. & Chattopadhyay, D. Coalescing black hole binaries from globular clusters: mass distributions and comparison to gravitational wave data from GWTC-3. *Mon. Not. R. Astron. Soc.* **522**, 466–476 (2023). [2208.01081](#).
- [30] Katz, H., Sijacki, D. & Haehnelt, M. G. Seeding high-redshift QSOs by collisional runaway in primordial star clusters. *Mon. Not. R. Astron. Soc.* **451**, 2352–2369 (2015). [1502.03448](#).
- [31] Crowther, P. A. *et al.* The R136 star cluster dissected with Hubble Space Telescope/STIS. I. Far-ultraviolet spectroscopic census and the origin of He II $\lambda 1640$ in young star clusters. *Mon. Not. R. Astron. Soc.* **458**, 624–659 (2016). [1603.04994](#).
- [32] Charbonnel, C. *et al.* N-enhancement in GN-z11: First evidence for supermassive stars nucleosynthesis in proto-globular clusters-like conditions at high redshift? *Astron. Astrophys.* **673**, L7 (2023). [2303.07955](#).
- [33] Marques-Chaves, R. *et al.* Extreme N-emitters at high redshift: Possible signatures of supermassive stars and globular cluster or black hole formation in action. *Astron. Astrophys.* **681**, A30 (2024). [2307.04234](#).
- [34] Gratton, R. *et al.* What is a globular cluster? An observational perspective. *A&A Rev.* **27**, 8 (2019). [1911.02835](#).

- [35] Reina-Campos, M., Kruijssen, J. M. D., Pfeffer, J. L., Bastian, N. & Crain, R. A. Formation histories of stars, clusters, and globular clusters in the E-MOSAICS simulations. *Mon. Not. R. Astron. Soc.* **486**, 5838–5852 (2019). [1905.02217](#).
- [36] Grudić, M. Y. *et al.* Great balls of FIRE - I. The formation of star clusters across cosmic time in a Milky Way-mass galaxy. *Mon. Not. R. Astron. Soc.* **519**, 1366–1380 (2023). [2203.05732](#).
- [37] Belokurov, V. & Kravtsov, A. In-situ vs accreted Milky Way globular clusters: a new classification method and implications for cluster formation. *Mon. Not. R. Astron. Soc.* (2023). [2309.15902](#).
- [38] Rieke, M. J. *et al.* Performance of NIRCcam on JWST in Flight. *PASP* **135**, 028001 (2023). [2212.12069](#).
- [39] Brammer, G., Strait, V., Matharu, J. & Momcheva, I. grizli (2022). URL <https://doi.org/10.5281/zenodo.6672538>. Please cite this software using these metadata.
- [40] Rigby, J. *et al.* The Science Performance of JWST as Characterized in Commissioning. *PASP* **135**, 048001 (2023). [2207.05632](#).
- [41] Planck Collaboration *et al.* Planck 2018 results. VI. Cosmological parameters. *Astron. Astrophys.* **641**, A6 (2020). [1807.06209](#).
- [42] Messa, M. *et al.* Star-forming clumps in the Lyman Alpha Reference Sample of galaxies - I. Photometric analysis and clumpiness. *Mon. Not. R. Astron. Soc.* **487**, 4238–4260 (2019). [1906.04751](#).
- [43] Messa, M. *et al.* Multiply lensed star forming clumps in the A521-sys1 galaxy at redshift 1. *Mon. Not. R. Astron. Soc.* **516**, 2420–2443 (2022). [2208.02863](#).

- [44] Claeysens, A. *et al.* Star formation at the smallest scales: a JWST study of the clump populations in SMACS0723. *Mon. Not. R. Astron. Soc.* **520**, 2180–2203 (2023). [2208.10450](#).
- [45] Vanzella, E. *et al.* Massive star cluster formation under the microscope at $z = 6$. *Mon. Not. R. Astron. Soc.* **483**, 3618–3635 (2019). [1809.02617](#).
- [46] Welch, B. *et al.* RELICS: Small-scale Star Formation in Lensed Galaxies at $z = 6$ –10. *Astrophys. J.* **943**, 2 (2023). [2207.03532](#).
- [47] Foreman-Mackey, D., Hogg, D. W., Lang, D. & Goodman, J. emcee: The MCMC Hammer. *PASP* **125**, 306 (2013). [1202.3665](#).
- [48] Pacifici, C. *et al.* The Art of Measuring Physical Parameters in Galaxies: A Critical Assessment of Spectral Energy Distribution Fitting Techniques. *Astrophys. J.* **944**, 141 (2023). [2212.01915](#).
- [49] Calzetti, D. *et al.* The Dust Content and Opacity of Actively Star-forming Galaxies. *Astrophys. J.* **533**, 682–695 (2000). [astro-ph/9911459](#).
- [50] Stanway, E. R. & Eldridge, J. J. Re-evaluating old stellar populations. *Mon. Not. R. Astron. Soc.* **479**, 75–93 (2018). [1805.08784](#).
- [51] Kroupa, P. On the variation of the initial mass function. *Mon. Not. R. Astron. Soc.* **322**, 231–246 (2001). [astro-ph/0009005](#).
- [52] Zackrisson, E. *et al.* The detection and characterization of highly magnified stars with JWST: Prospects of finding Population III. *arXiv e-prints* arXiv:2312.09289 (2023). [2312.09289](#).
- [53] Szécsi, D., Agrawal, P., Wünsch, R. & Langer, N. Bonn Optimized Stellar Tracks (BoOST). Simulated populations of massive and very massive stars for

- astrophysical applications. *Astron. Astrophys.* **658**, A125 (2022). [2004.08203](#).
- [54] Jullo, E. *et al.* A Bayesian approach to strong lensing modelling of galaxy clusters. *New Journal of Physics* **9**, 447 (2007). [0706.0048](#).
- [55] Paterno-Mahler, R. *et al.* RELICS: A Strong Lens Model for SPT-CLJ0615-5746, a $z = 0.972$ Cluster. *Astrophys. J.* **863**, 154 (2018). [1805.09834](#).
- [56] Oguri, M. The Mass Distribution of SDSS J1004+4112 Revisited. *PASJ* **62**, 1017 (2010). [1005.3103](#).
- [57] Oguri, M. Fast Calculation of Gravitational Lensing Properties of Elliptical Navarro-Frenk-White and Hernquist Density Profiles. *PASP* **133**, 074504 (2021). [2106.11464](#).
- [58] Navarro, J. F., Frenk, C. S. & White, S. D. M. A Universal Density Profile from Hierarchical Clustering. *Astrophys. J.* **490**, 493–508 (1997). [astro-ph/9611107](#).
- [59] Diego, J. M., Protopapas, P., Sandvik, H. B. & Tegmark, M. Non-parametric inversion of strong lensing systems. *Mon. Not. R. Astron. Soc.* **360**, 477–491 (2005a). [arXiv:astro-ph/0408418](#).
- [60] Diego, J. M., Tegmark, M., Protopapas, P. & Sandvik, H. B. Combined reconstruction of weak and strong lensing data with WSLAP. *Mon. Not. R. Astron. Soc.* **375**, 958–970 (2007). [arXiv:astro-ph/0509103](#).
- [61] van der Walt, S., Colbert, S. C. & Varoquaux, G. The numpy array: A structure for efficient numerical computation. *CSE* **13**, 22–30 (2011).
- [62] Jones, E., Oliphant, T., Peterson, P. *et al.* SciPy: Open source scientific tools for Python (2001–). URL <http://www.scipy.org/>. [Online; accessed 2016-11-22].

- [63] Hunter, J. D. Matplotlib: A 2d graphics environment. *Computing In Science & Engineering* **9**, 90–95 (2007).
- [64] Astropy Collaboration *et al.* Astropy: A community Python package for astronomy. *Astron. Astrophys.* **558**, A33 (2013). [1307.6212](#).




Fig. 1: The Cosmic Gems arc in a JWST color-composite. The filter combination shows the rest-frame UV, blue optical wavelengths (1200–2800 Å). The arc is extended over 5". A foreground galaxy at redshift $z_{\text{phot}} = 2.6$ is visible above and to the right. The field of view is rotated North up. Top right: A zoom inset of the center of the arc where the brightest star clusters are located highlighted by the white square on the left image. Two mirror images are observed due to gravitational lensing. Lensing critical curves based on three models are shown bisecting the arc. Bottom left: each star cluster is labeled in a grayscale FUV rest-frame image of the galaxy. Bottom right: Source plane reconstruction of the core of the galaxy where the star clusters are located showing their relative sizes and positions. The physical distance between A and E and A and D is about 40 pc. Note there is some uncertainty in source positions parallel to the lensing caustic.







Fig. 2: Cluster stellar surface density versus half-light radius R_{eff} . The star clusters in the $z_{\text{phot}} \sim 10.2$ Cosmic Gems arc are shown color-coded by their dynamical age, $\log(\Pi)$. The plotted values have been derived using the reference lensing model. Predictions by other models do not change the observed trends. The error bars do not account for magnification uncertainties. However, we refer the reader to Extended Data Fig. 4, where we show that magnification uncertainties do not affect these results. Other gravitationally-bound star clusters detected in lensed galaxies at $z \sim 6$ and $z = 2.38$ are included, along with the $z = 0$ Milky Way globular clusters and young star clusters properties of star-forming spiral galaxies in the Local Volume (distance < 16 Mpc). Lines of equal mass show the change in density as a function of R_{eff} .



Extended Data Figure 1. Cluster light fit in the image plane. Observed JWST images (left), best-fitted clump shape after removing the local diffuse light, (centre), and residual images (right) in the reference filter F150W (top), and two more bands, the F200W (middle) with similar resolution to the F150W, and the F444W (bottom), with the lowest spatial resolution. We show log-scale images matched in flux in each band.



Extended Data Figure 2. Cluster light fit with forward modeling from the source plane. Observed JWST image of *Img. 1* is shown (far left) along with the best fit image plane from forward modeling (left center) and weighted fit residuals (right). Weighted residuals are calculated as $(\text{data} - \text{model}) / (\text{uncertainty})$.




Extended Data Figure 3. Observed photometry and spectral energy distributions (SEDs) of each star cluster (presented in Methods). We include the observed SEDs of the mirrored image of clusters A, B, C (orange symbols) normalised by the median ratio of the 6 bands, preserving the SED shape.

BAGPIPES $z = 10.2$, $\tau = 1$ Myr, Calzetti attenuation										Prospector SSP, Calzetti attenuation												
ID	Age [Myr]	$10^6 M_*$ [M_\odot]	A_V [mag]	Z/Z_\odot [%]	Age [Myr]	$10^6 M_*$ [M_\odot]	A_V [mag]	Z/Z_\odot [%]	Age [Myr]	$10^6 M_*$ [M_\odot]	A_V [mag]	Z/Z_\odot [%]	Age [Myr]	$10^6 M_*$ [M_\odot]	A_V [mag]	Z/Z_\odot [%]	Age [Myr]	$10^6 M_*$ [M_\odot]	A_V [mag]	Z/Z_\odot [%]	τ [Myr]	
A.1	$9.2^{+13.5}_{-3.2}$	$138.34^{+293.88}_{-51.986}$	$0.22^{+0.15}_{-0.14}$	$6.38^{+2.42}_{-3.08}$	$4.0^{+9.5}_{-1.0}$	$21.94^{+328.83}_{-0.2475}$	$0.00^{+0.27}_{-0.00}$	$16.91^{+0.87}_{-0.14}$														
B.1	$14.0^{+9.2}_{-4.2}$	$289.37^{+187.65}_{-271.70}$	$0.20^{+0.12}_{-0.09}$	$3.42^{+2.49}_{-3.48}$	$51.8^{+21.4}_{-23.3}$	$920.51^{+238.96}_{-268.97}$	$0.04^{+0.03}_{-0.08}$	$0.83^{+0.34}_{-0.82}$														
C.1	$9.1^{+9.2}_{-2.8}$	$172.51^{+100.24}_{-256.24}$	$0.36^{+0.16}_{-0.12}$	$4.56^{+3.49}_{-3.12}$	$14.9^{+4.7}_{-19.6}$	$585.77^{+262.14}_{-337.14}$	$0.49^{+0.29}_{-0.51}$	$0.57^{+1.82}_{-2.17}$														
D.1	$10.9^{+3.9}_{-3.9}$	$235.49^{+154.93}_{-153.99}$	$0.33^{+0.16}_{-0.15}$	$5.27^{+3.03}_{-3.15}$	$16.2^{+5.6}_{-24.2}$	$528.51^{+168.52}_{-254.82}$	$0.11^{+0.27}_{-0.09}$	$0.45^{+2.17}_{-2.08}$														
E.1	$36.8^{+16.4}_{-6.2}$	$421.73^{+151.94}_{-177.77}$	$0.21^{+0.11}_{-0.07}$	$5.05^{+3.14}_{-3.28}$	$51.8^{+18.7}_{-12.1}$	$582.23^{+266.12}_{-160.02}$	$0.11^{+0.23}_{-0.01}$	$2.66^{+2.08}_{-1.580}$														
A.2	$17.2^{+8.8}_{-6.2}$	$166.70^{+90.28}_{-57.14}$	$0.11^{+0.11}_{-0.16}$	$4.39^{+3.86}_{-4.83}$	$8.0^{+13.6}_{-3.1}$	$45.26^{+278.14}_{-216.02}$	$0.40^{+0.15}_{-0.27}$	$1.24^{+3.65}_{-1.36}$														
B.2	$9.3^{+6.6}_{-7.8}$	$294.46^{+137.33}_{-137.33}$	$0.28^{+0.12}_{-0.16}$	$4.83^{+2.83}_{-2.62}$	$13.6^{+11.0}_{-11.0}$	$743.34^{+236.02}_{-136.02}$	$0.13^{+0.13}_{-0.11}$	$3.71^{+6.35}_{-3.25}$														
C.2+D.2	$18.6^{+7.4}_{-7.4}$	$603.09^{+263.11}_{-263.11}$	$0.19^{+0.13}_{-0.13}$	$6.29^{+3.50}_{-3.50}$	$24.5^{+11.0}_{-11.0}$	$794.45^{+308.30}_{-308.30}$	$0.13^{+0.13}_{-0.11}$	$3.71^{+6.35}_{-3.25}$														
BAGPIPES $z = 10.2$, Burst, Calzetti attenuation																						
A.1	$23.5^{+4.2}_{-17.0}$	$503.90^{+106.02}_{-409.27}$	$0.32^{+0.09}_{-0.10}$	$0.74^{+4.93}_{-0.75}$	$16.68^{+10.41}_{-7.65}$	$257.14^{+159.40}_{-171.21}$	$0.09^{+0.10}_{-0.06}$	$3.33^{+3.66}_{-2.19}$														
B.1	$15.4^{+6.0}_{-6.6}$	$421.94^{+229.62}_{-229.56}$	$0.31^{+0.17}_{-0.09}$	$0.17^{+0.14}_{-0.14}$	$18.05^{+5.67}_{-14.20}$	$319.09^{+133.81}_{-275.99}$	$0.09^{+0.08}_{-0.06}$	$0.63^{+0.90}_{-2.63}$														
C.1	$7.0^{+4.7}_{-4.7}$	$149.70^{+434.40}_{-99.12}$	$0.37^{+0.09}_{-0.20}$	$0.30^{+2.71}_{-0.27}$	$14.32^{+14.20}_{-11.11}$	$269.70^{+236.38}_{-166.15}$	$0.17^{+0.15}_{-0.11}$	$1.69^{+2.63}_{-1.09}$														
D.1	$8.4^{+3.7}_{-24.2}$	$173.50^{+391.91}_{-133.47}$	$0.36^{+0.18}_{-0.14}$	$0.57^{+3.84}_{-0.53}$	$20.70^{+10.67}_{-22.97}$	$373.07^{+166.15}_{-196.33}$	$0.20^{+0.12}_{-0.12}$	$1.94^{+3.61}_{-1.31}$														
E.1	$41.0^{+14.4}_{-8.0}$	$439.84^{+133.47}_{-12.90}$	$0.22^{+0.17}_{-0.14}$	$0.22^{+2.52}_{-0.19}$	$31.75^{+22.97}_{-14.31}$	$324.68^{+158.77}_{-127.03}$	$0.17^{+0.16}_{-0.11}$	$3.66^{+4.00}_{-2.55}$														
A.2	$18.6^{+8.0}_{-4.3}$	$222.91^{+73.25}_{-302.06}$	$0.19^{+0.11}_{-0.11}$	$0.28^{+2.20}_{-3.23}$	$18.49^{+9.68}_{-8.04}$	$171.49^{+78.30}_{-286.08}$	$0.06^{+0.08}_{-0.13}$	$1.33^{+2.50}_{-3.46}$														
B.2	$7.7^{+2.3}_{-2.3}$	$263.42^{+125.30}_{-100.62}$	$0.33^{+0.16}_{-0.17}$	$1.04^{+3.98}_{-0.99}$	$12.30^{+8.31}_{-8.00}$	$382.18^{+324.66}_{-201.93}$	$0.14^{+0.13}_{-0.13}$	$2.04^{+4.10}_{-4.10}$														
C.2+D.2	$19.9^{+10.0}_{-10.0}$	$670.79^{+317.92}_{-317.92}$	$0.19^{+0.14}_{-0.14}$	$0.99^{+0.93}_{-0.93}$	$19.18^{+7.78}_{-7.78}$	$521.93^{+248.80}_{-248.80}$	$0.11^{+0.08}_{-0.08}$	$3.02^{+4.10}_{-1.77}$														
BAGPIPES $z = 10.2$, $\tau = 1$ Myr, SMC extinction																						
A.1	$19.3^{+12.9}_{-13.4}$	$278.55^{+166.88}_{-234.09}$	$0.04^{+0.04}_{-0.03}$	$6.67^{+2.39}_{-3.60}$	$27.90^{+21.74}_{-14.82}$	$192.02^{+135.05}_{-107.57}$	$0.08^{+0.08}_{-0.05}$	$6.39^{+2.40}_{-3.13}$														
B.1	$16.4^{+5.7}_{-8.2}$	$272.67^{+115.28}_{-160.50}$	$0.05^{+0.03}_{-0.06}$	$4.62^{+3.17}_{-3.46}$	$19.15^{+7.33}_{-16.10}$	$162.58^{+57.50}_{-137.98}$	$0.04^{+0.03}_{-0.03}$	$2.16^{+2.41}_{-3.36}$														
C.1	$12.6^{+4.8}_{-4.8}$	$259.80^{+157.52}_{-179.19}$	$0.16^{+0.07}_{-0.07}$	$4.10^{+2.76}_{-3.60}$	$15.68^{+9.40}_{-22.90}$	$171.97^{+69.69}_{-176.63}$	$0.20^{+0.12}_{-0.13}$	$3.84^{+2.93}_{-2.93}$														
D.1	$13.6^{+10.7}_{-6.0}$	$249.39^{+168.07}_{-148.91}$	$0.15^{+0.07}_{-0.08}$	$4.53^{+3.00}_{-3.27}$	$18.39^{+10.93}_{-42.68}$	$167.89^{+86.57}_{-176.71}$	$0.19^{+0.11}_{-0.11}$	$5.08^{+2.97}_{-3.12}$														
E.1	$35.3^{+17.9}_{-14.6}$	$357.91^{+148.91}_{-142.66}$	$0.11^{+0.08}_{-0.08}$	$5.38^{+3.66}_{-3.66}$	$54.85^{+30.97}_{-42.68}$	$314.15^{+68.91}_{-68.91}$	$0.19^{+0.15}_{-0.13}$	$4.15^{+3.79}_{-2.70}$														
A.2	$18.3^{+7.8}_{-6.6}$	$157.81^{+78.97}_{-69.06}$	$0.04^{+0.04}_{-0.02}$	$4.02^{+3.60}_{-2.76}$	$27.56^{+22.23}_{-13.42}$	$123.81^{+97.75}_{-35.06}$	$0.05^{+0.03}_{-0.03}$	$3.63^{+3.32}_{-2.34}$														
B.2	$11.1^{+6.6}_{-6.2}$	$350.70^{+271.40}_{-120.36}$	$0.14^{+0.07}_{-0.07}$	$4.44^{+2.56}_{-2.56}$	$15.87^{+15.37}_{-16.40}$	$270.99^{+209.38}_{-173.05}$	$0.16^{+0.12}_{-0.12}$	$4.43^{+3.55}_{-3.72}$														
C.2+D.2	$22.6^{+7.3}_{-7.3}$	$632.09^{+206.55}_{-206.55}$	$0.08^{+0.06}_{-0.06}$	$5.56^{+3.05}_{-3.05}$	$34.25^{+19.41}_{-19.41}$	$474.27^{+48.73}_{-48.73}$	$0.13^{+0.09}_{-0.09}$	$3.79^{+2.60}_{-2.60}$														

Extended Data Table 2. Compilations of different SED fit outputs described in Methods. The reported masses are not corrected for magnification. The output of the fit assuming $\tau = 1$ Myr and Calzetti attenuation is referred to as BAGPIPES-*exp* and used to derive the physical values reported in Table 1. For BAGPIPES fit with $\tau = \text{free}$ we report mass weighted quantities.

Extended Data Table 3. Magnifications and associated uncertainties for 3 different lens models. The model *Lenstool-A* is used as reference in the analysis.

ID	RA	DEC	$\mu_{\text{Lenstool-A}} (\mu_{\text{tan}})$	$\mu_{\text{Lenstool-B}} (\mu_{\text{tan}})$	$\mu_{\text{glafic}} (\mu_{\text{tan}})$
A.1	93.979828	-57.772475	$56.5^{+10.1}_{-8.9}$ (44.8)	$122.0^{+48.8}_{-17.8}$ (81.9)	76.9 (68.2)
B.1	93.979698	-57.772395	$109.2^{+31.2}_{-30.3}$ (84.8)	$212.6^{+152.8}_{-25.4}$ (142.3)	153.8 (125.7)
C.1	93.979660	-57.772373	$153.3^{+91.9}_{-55.8}$ (120.8)	$280.0^{+270.7}_{-27.2}$ (187.9)	200.9 (177.5)
D.1	93.979635	-57.772355	$209.1^{+132.8}_{-93.0}$ (159.8)	$349.5^{+474.6}_{-56.5}$ (234.2)	288.8 (254.7)
E.1	93.979599	-57.772335	$419.3^{+215.7}_{-261.6}$ (341.2)	$527.3^{+708.6}_{-30.1}$ (353.7)	516.4 (455.7)
A.2	93.979316	-57.772180	$57.7^{+27.7}_{-3.7}$ (46.5)	$128.9^{+17.1}_{-17.1}$ (89.5)	76.3 (70.1)
B.2	93.979416	-57.772235	$97.8^{+63.5}_{-10.0}$ (72.0)	$234.3^{+188.7}_{-30.8}$ (157)	133.0 (116.9)
C.2+D.2	93.979460	-57.772256	$138.4^{+263.7}_{-20.4}$ (117)	$372.8^{+527.0}_{-58.2}$ (249.6)	207.7 (182.3)



Extended Data Figure 4. Measured and derived cluster physical properties as a function of their magnification. The most relevant quantities of each cluster in the arc (marked in the central top panel) are expressed as a function of the total magnification (μ_{total} and in case of the R_{eff} as a function of μ_{tan}). The radii (R_{eff}), dynamical ages (Π), stellar mass surface densities (Σ_{mass}) and the stellar masses (M_*) suggest the clumps are bound star clusters even at modest magnification regimes ($\mu_{\text{total}} > 10$). The transparent green region and the vertical lines show the expected magnification from the reference lens model. The dotted horizontal line indicates the region where $\log(\Pi) = 0$. The shaded areas in the plot mark the uncertainties associated with the derived values. The dashed lines show the lower-limits in Σ_* and Π , assuming half the stellar PSF FWHM as upper limit for the R_{eff} of each star cluster.