

Exploring the photoproduction of ρ and ϕ in hadronic heavy-ion collisions

Kaifeng Shen, Xin Wu, Zebo Tang, and Wangmei Zha*
State Key Laboratory of Particle Detection and Electronics,
University of Science and Technology of China, Hefei 230026, China
(Dated: January 17, 2024)

Significant enhancements of J/ψ production have been observed by various experiments at RHIC and LHC for very low transverse momenta in peripheral heavy-ion collisions, which has ignited a surge of investigations into photon-induced processes in hadronic heavy-ion collisions (HHICs). Within this wave of research enthusiasm, the search for more photon induced products in HHICs becomes paramount. In this paper, we perform the calculation of the ρ and ϕ production resulting from photon-nucleus interactions in HHICs, which are crucial probes for studying the properties of Quark-Gluon Plasma (QGP) in HHICs. Our study reveals that, in comparison to hadronic production, the photon-induced production of ρ and ϕ does not reach the same level of significance as that observed in J/ψ production. Nevertheless, it remains substantial, especially in peripheral collisions, holding great promise for experimental verification in the imminent future.

I. INTRODUCTION

Understanding the intricacies of the Quark-Gluon Plasma (QGP) is crucial for unraveling the fundamental nature of matter under extreme conditions, such as those present in the early universe microseconds after the Big Bang. By utilizing the final-state particles resulting from relativistic heavy-ion collisions, notably those conducted at RHIC [1, 2] and LHC [3, 4], valuable insights into the properties and dynamics of the QGP can be gleaned, shedding light on the behavior of quarks and gluons in conditions of extreme temperature and energy density [5]. Among the probes employed to investigate the QGP, vector mesons like ρ and ϕ assume pivotal roles. The measurements of ρ (770) through leptonic decay is sensitive to the chiral symmetry restoration within the hot and dense quantum chromodynamics (QCD) environment, which establishes a connection to the origin of hadronic masses in the universe [6, 7]. Additionally, the study of the ϕ meson, bound state of strange quark and its anti-quark, proves especial insight for deciphering the QGP chemistry [8–11], as strangeness is not present in the valence content of the colliding nuclei and is light enough to be created copiously in the hot medium.

In relativistic heavy-ion collisions, the production of vector mesons primarily arises from the intense strong interactions occurring within the nuclear overlap region; however, it extends beyond the realm of purely hadronic production. Another significant process, photon induced interaction, involving the emission of highly quasireal photons from one of the colliding nucleus [12–14], could also generate vector mesons. These photons could fluctuate into $q\bar{q}$ pairs and subsequently interact with the other nucleus (coherent process) or the individual nucleons in the other nucleus (incoherent process) via Pomeron exchange, a color-neutral two-gluon state, and then emerge as vector mesons. In the coherent process, the final products consist of two intact nuclei, a vector meson with very

low transverse momentum ($p_T < 0.1$ GeV/c) and nothing else. The exclusive photoproduction of a vector meson, where both outgoing nuclei remain intact throughout the process, presents a distinctive opportunity to investigate the gluonic structure of nuclear matter. Notably, coherent interactions are conventionally believed to be only observable in Ultra-Peripheral Collisions (UPCs), where the impact parameter (b) is larger than twice of the nuclear radius. In such collisions, the two colliding nuclei are expected to remain intact, satisfying the conditions necessary for coherent interactions. Therefore, the photoproduction of vector mesons was anticipated exclusively in UPCs, while their hadronic production was expected solely in hadronic heavy-ion collisions (HHICs)..

Is it possible that coherent photonucleus interactions also take place in HHICs, where the nuclei collide and disintegrate? The coherent photoproduction is a electromagnetic process that could occur at the periphery of the colliding nuclei where the influence of the hot medium is less pronounced. Consequently, the vector mesons produced via coherent photonucleus interactions may endure within the fireball and come to dominate in the very low p_T region. The experimental evidence was first revealed by the ALICE [15] and STAR [16] collaborations, showing a significantly excess in the yield of J/ψ at very low p_T (< 0.3 GeV/c). Theoretical models [17–19] based on photoproduction processes have been employed to explain these excesses, since it is evident that these excesses can not be comprehended through known hot and cold medium effects in hadronic J/ψ production. This research [17] explores various photon and Pomeron coupling scenarios and addresses interference between J/ψ photoproduction amplitudes from opposite directions. Besides the photonucleus interactions, STAR and ALICE also proved the coherent photon-photon interactions happened in HHICs by measuring the excesses in dilepton production at very low p_T [20, 21], and the QED based calculations at leading-order [22–24] can describe these measurements in HHICs.

Can we observe additional photon-induced products in HHICs besides J/ψ , such as ρ and ϕ ? If so, what are their

* first@ustc.edu.cn

contributions compared to their hadronic productions? Given that photoproduction has not been considered in previous experimental measurements, there is a question of whether it could impact the derived physical conclusions. Furthermore, can these photon-induced products also function as probes for studying the properties of the QGP? In this paper, we undertake calculations for the p_T spectrum of ρ and ϕ photoproductions across various centrality bins at RHIC and LHC energies. Our objective is to determine the significance of photoproduction over the hadronic background, providing insights into the experimental challenges associated with verifying photon-produced ρ and ϕ in HHICs.

II. METHODOLOGY

The cross section of the ρ and ϕ mesons in the photonuclear process can be calculated by using the similar method in [25, 26]. For the experimental measurements, the production yields are usually performed in the transverse momentum space, especially for coherent photoproduction case where the p_T of the photo-produced vector meson is constrained to be the level of the inverse of the colliding nucleus size [27]. In this case, the photo-produced ρ and ϕ mesons will be dominated in the extremely low p_T region which is about 60 MeV/ c , while the yield of ρ and ϕ mesons from hadronic processes decreases as p_T decreases in such low p_T region. In theoretical calculations, the amplitude distribution of the ρ and ϕ mesons from the photonuclear process in two-dimensional transverse momentum representation can be calculated by performing a Fourier transformation to which in coordinate representation:

$$\vec{A}(\vec{p}_\perp) = \frac{1}{2\pi} \int d^2x_\perp (\vec{A}_1(\vec{x}_\perp) + \vec{A}_2(\vec{x}_\perp)) \times e^{i\vec{p}_\perp \cdot \vec{x}_\perp}, \quad (1)$$

where $\vec{A}_1(\vec{x}_\perp)$ and $\vec{A}_2(\vec{x}_\perp)$ are the amplitudes distributions in the transverse plane from the two colliding nuclei, which can be obtained based on the corresponding $\gamma A \rightarrow VA$ scattering amplitude $\Gamma_{\gamma A \rightarrow VA}$ and the spatial photon flux in the outside the colliding nuclei [28], shown as the following equation:

$$A(\vec{x}_\perp) = \Gamma_{\gamma A \rightarrow VA} \sqrt{N(\omega_\gamma, \vec{x}_\perp)}. \quad (2)$$

The scattering amplitude $\Gamma_{\gamma A \rightarrow VA}$ can be calculated by the Glauber model [29] and vector meson dominance (VMD) approach [30]:

$$\Gamma_{\gamma A \rightarrow VA}(\vec{x}_\perp) = \frac{4\sqrt{\alpha}C}{f_V} \times \frac{f_{\gamma N \rightarrow VN}(0)}{f_{\gamma N \rightarrow VN}} \times 2[1 - \exp(-\frac{\sigma_{VN}}{2} T'(\vec{x}_\perp))], \quad (3)$$

where f_V is the V -photon coupling const and C is a correction factor for the non-diagonal coupling through

higher mass vector mesons [31], and $f_{\gamma N \rightarrow VN}(0)$ is the forward-scattering amplitude for $\gamma + N \rightarrow V + N$, which can be obtained from the well parameterized cross section of the forward-scattering $\frac{d\sigma_{\gamma N \rightarrow VN}}{dt}|_{t=0}$ as shown in Ref. [32]. The modified thickness function, $T'(\vec{x}_\perp)$, is used to accounting for the coherent effect on z direction:

$$T'(\vec{x}_\perp) = \int_{-\infty}^{+\infty} \rho(\sqrt{\vec{x}_\perp^2 + z^2}) e^{iq_L z} dz, \quad (4)$$

$$q_L = \frac{M_{\rho(\phi)} e^y}{2\gamma_c},$$

where q_L is the longitudinal momentum transfer required to produce a real vector meson, γ_c is the Lorentz factor of the nucleus, $M_{\rho(\phi)}$ and y are the mass and rapidity of the ρ (ϕ) meson.

The distribution of photon flux generated by the colliding nuclei can be calculated based on the equivalent photon approximation (EPA):

$$\frac{d^3 N_\gamma(\omega_\gamma, \vec{x}_\perp)}{d\omega_\gamma d\vec{x}_\perp} = \frac{4Z^2\alpha}{\omega_\gamma} \times \left| \int \frac{d^2 k_{\gamma\perp}}{(2\pi)^2} \vec{k}_{\gamma\perp} \frac{F_\gamma(\vec{k}_\gamma)}{|\vec{k}_\gamma|^2} e^{i\vec{x}_\perp \cdot \vec{k}_{\gamma\perp}} \right|^2 \quad (5)$$

$$\vec{k}_\gamma = (\vec{k}_{\gamma\perp}, \frac{\omega_\gamma}{\gamma_c}), \quad \omega_\gamma = \frac{1}{2} M_{\rho(\phi)} e^{\pm y},$$

where Z is the nuclear charge number, α is the electromagnetic coupling constant, ω_γ is the energy of photon, $F_\gamma(\vec{k}_\gamma)$ is the nuclear electromagnetic form factor and the \vec{k}_γ is two-dimensional momentum vectors of the equivalent photons in the transverse momentum space. The form factor is obtained by the Fourier transformation of the charge density in the nucleus, which is described by the parameterized Woods-Saxon distribution:

$$\rho_A(r) = \frac{\rho_0}{1 + \exp[(r - R_{WS})/d]}, \quad (6)$$

where ρ_0 is the normalization factor and the radius R_{WS} and skin depth d are from electron-scattering experiments [33].

In addition, the incoherent photo-nucleus interactions also contribute to the production of ρ and ϕ mesons in the very low p_T region, although the p_T of ρ and ϕ mesons from incoherent photo-nucleus process is constrained to the order of the inverse of the nucleon size, which is about 300 MeV/ c [27]. The cross section of the incoherent ρ and ϕ mesons photoproduction, $\sigma_{\gamma A \rightarrow V A'}$, is scaled from the cross section $\gamma p \rightarrow V p$ by the Glauber plus VMD approach, where A' is the final state of the nucleus after the interaction, which contains the products of the nuclear disintegration. The cross section of incoherent

photo-nucleus interaction can be calculated as:

$$\sigma_{\gamma+A \rightarrow V+A'} = \sigma_{\gamma+p \rightarrow V+p} \times \int d^2 \vec{x}_\perp T(\vec{x}_\perp) e^{-\frac{1}{2} \sigma_{VN}^{\text{in}} T(\vec{x}_\perp)}, \quad (7)$$

$$\sigma_{VN}^{\text{in}} = \sigma_{VN} - \sigma_{VN}^2 / (16\pi B_V),$$

where $T(\vec{x}_\perp)$ is the thickness function of the nucleus, σ_{VN}^{in} is the inelastic vector meson-nucleon cross section and B_V is the slope of the t dependence of the $\gamma + p \rightarrow V + p$ scattering [32].

III. RESULTS




FIG. 1. The calculated invariant yield of ρ meson resulting from photoproduction as a function of p_T in 70-80% Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV (Top) and in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV (Bottom). The magenta solid lines represent the contribution originating from the coherent photon-nucleus process, while the brown solid lines denote the production derived from the incoherent photon-nucleus process. The blue solid lines stand for the total yield of ρ mesons, encompassing contributions from both photoproduction and hadronic production.

Figure 1 illustrates the calculated invariant yield distributions of ρ mesons, $Br_{\rho \rightarrow \pi^+ \pi^-} \frac{d^2 N}{2\pi p_T dp_T dy}$, arising from photoproduction in 70-80% Au+Au collisions at $\sqrt{s_{NN}} =$

200 GeV and Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. The magenta solid lines depict the distribution of photo-produced ρ mesons resulting from coherent interactions, while the brown solid lines represent those resulting from incoherent interactions. The blue solid lines indicate the total yields of ρ mesons, encompassing contributions from both photoproduction and hadronic interactions. As observed in Fig. 1, the p_T distribution of coherent photo-produced ρ mesons is highly concentrated in the extremely low range ($p_T < 0.1$ GeV/c). The decreasing trend as p_T approaches zero is a consequence of interference effects [25]. The peak structures originating from the coherent photon-nucleus process are attributed to the combined effects of interference and diffraction, and noticeable differences between Au+Au and Pb+Pb collisions are evident, stemming from the distinct size and shape of the two nuclei. For the incoherent photoproduction case, the p_T distribution extends to a relatively higher range, as revealed by our calculations. Experimental measurements of ρ meson yields resulting from hadronic interactions in Au+Au at $\sqrt{s_{NN}} = 200$ GeV and in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV are represented by the black points [34, 35].

To estimate the yield of ρ mesons from hadronic production at extremely low transverse momentum (p_T), the experimental data in Fig. 1 is subjected to fitting using the following formula:

$$\frac{1}{2\pi p_T} \frac{d^2 N}{dy dp_T} = A \cdot e^{[-B(\sqrt{p_T^2 + m_V^2} - m_V)]} \quad (8)$$

where A and B are free parameters and m_V is the pole mass of the vector meson. The fitted results are presented as the red band in Fig. 1, where the band conveys the uncertainties associated with the fitting process. Through these extrapolated results, the yield of photo-produced ρ mesons is found to be comparable to that from hadronic interactions at very low p_T . Notably, the ρ meson yield from photoproduction closely aligns with that from hadronic production in 70-80% Au+Au collisions at RHIC and Pb+Pb collisions at LHC. The comparable invariant yield results for ϕ mesons are exhibited in Fig. 2, revealing that the yield fraction of ϕ mesons from photoproduction is lower than that of ρ mesons in both collision systems.

Figure 3 illustrates the ratio of photo-produced ρ mesons to those originating from hadronic production, presenting a dependence on $\langle N_{\text{part}} \rangle$, at very low p_T (< 0.1 GeV/c) and integrated p_T in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. Different colored data points distinguish between various collision systems. The observed trend reveals an increase in the ratio as $\langle N_{\text{part}} \rangle$ decreases, suggesting a more pronounced photo-production of ρ mesons in peripheral nucleus-nucleus collisions. In central collisions, contributions from photo-production appear negligible; however, in peripheral collisions, photo-induced production accounts for approximately 30% of ρ production originating from hadronic interactions at very low p_T ($<$




FIG. 2. The calculated invariant yield of photon produced ϕ meson as a function of p_T in 70-80% Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV (Top) and Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV (Bottom). The magenta solid lines represent the invariant yield originating from the coherent process, while the brown solid lines present similar results derived from incoherent interactions. The blue solid lines denote the total yield of ϕ mesons, encompassing contributions from both coherent and incoherent processes.

0.1 GeV/c). The integrated yield of ρ meson from photoproduction is much smaller than those from hadronic interactions in all centrality classes, which suggests that the additional contribution from photoproduction does not significantly impact the derived physical conclusions from previous experimental measurements. Moving on to the ratios of photo-produced ϕ mesons to those from hadronic interactions, depicted as a function of $\langle N_{part} \rangle$ at $p_T < 0.1$ GeV/c and integrated p_T in Fig. 4, we observe a similar trend to that of ρ mesons. However, the proportion of photo-production for ϕ mesons is relatively lower compared to that of ρ . In the most peripheral Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, the photo-induced production of ϕ mesons reaches up to 15% of the ϕ yield originating from hadronic interactions for very low p_T (< 0.1 GeV/c). While ρ and ϕ photoproduction is not as pronounced as that of J/ψ at very low p_T in peripheral collisions, it remains comparable to that from hadronic interactions on the order of magnitude. This suggests




FIG. 3. The ratio of photo-produced ρ^0 mesons to those originating from hadronic interactions as a function of $\langle N_{part} \rangle$, for $p_T < 0.1$ GeV/c and integrated p_T in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV (depicted by blue markers) and in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV (depicted by red markers).

promising avenues for experimental verification in the near future, necessitating higher statistical requirements. However, it is crucial to note that this study represents an initial step, and more advanced models are required to explore potential hot medium effects on ρ and ϕ photoproduction.




FIG. 4. The ratio of photo-produced ϕ mesons to those originating from hadronic interactions as a function of $\langle N_{part} \rangle$, for $p_T < 0.1$ GeV/c and integrated p_T in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV (depicted by blue markers) and in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV (depicted by red markers).

IV. SUMMARY

In summary, this study delves into the realm of ρ and ϕ photoproduction within in hadronic high-energy heavy-ion collisions at both RHIC and LHC energies. Our calculations compellingly underscore that the yield of ρ and ϕ at very low p_T has significant contribution from co-

herent photon-nucleus process. Methodically considering coherent and incoherent photo-produced ρ and ϕ mesons and juxtaposing them with extrapolated hadronic contributions yields are valuable insights. Compared to its hadronic counterpart, the integrated yield from photoproduction registers as negligible, which suggests that the additional contribution from photoproduction does not wield a significant impact on the established physical conclusions drawn from preceding experimental measurements. However, at $p_T < 0.1$ GeV/ c in peripheral collisions, this yield remains comparable to that from hadronic interactions. This holds promise for experimental verification of ρ and ϕ photoproduction in HHICs in the imminent future, yet it demands augmented statistical requirements. A pivotal query arises: Can these photon-induced products serve as probes for probing the properties of the QGP? This question sparks curiosity

and beckons further investigation. It is paramount to acknowledge that this study constitutes an inaugural exploration, and the pursuit of more advanced models becomes imperative to scrutinize potential hot medium effects on ρ and ϕ photoproduction.

V. ACKNOWLEDGMENTS

This work is supported in part by the National Key Research and Development Program of China under Contract No. 2022YFA1604900 and the National Natural Science Foundation of China (NSFC) under Contract No. 12175223 and 12005220. W. Zha is supported by Anhui Provincial Natural Science Foundation No. 2208085J23 and Youth Innovation Promotion Association of Chinese Academy of Science.

-
- [1] John Adams et al. Experimental and theoretical challenges in the search for the quark gluon plasma: The STAR Collaboration's critical assessment of the evidence from RHIC collisions. *Nucl. Phys. A*, 757:102–183, 2005.
 - [2] Xin Dong, Lijuan Ruan, Ming Shao, Yongjie Sun, Zebo Tang, Zhangbu Xu, Wangmei Zha, and Yifei Zhang. Dilepton Program with Time-of-Flight Detector at the STAR Experiment. *Symmetry*, 15(2):392, 2023.
 - [3] T. Niida and Y. Miake. Signatures of QGP at RHIC and the LHC. *AAPPS Bull.*, 31(1):12, 2021.
 - [4] Renu Bala, Irais Bautista, Jana Bielikova, and Antonio Ortiz. Heavy-ion physics at the LHC: Review of Run I results. *Int. J. Mod. Phys. E*, 25(07):1642006, 2016.
 - [5] Peter Braun-Munzinger and Johanna Stachel. The quest for the quark-gluon plasma. *Nature*, 448:302–309, 2007.
 - [6] C. D. Roberts, M. S. Bhagwat, A. Holl, and S. V. Wright. Aspects of hadron physics. *Eur. Phys. J. ST*, 140:53–116, 2007.
 - [7] Sinya Aoki et al. Review of Lattice Results Concerning Low-Energy Particle Physics. *Eur. Phys. J. C*, 74:2890, 2014.
 - [8] A. Shor. PHI MESON PRODUCTION AS A PROBE OF THE QUARK GLUON PLASMA. *Phys. Rev. Lett.*, 54:1122–1125, 1985.
 - [9] B. Mohanty and N. Xu. Probe the QCD phase diagram with phi-mesons in high energy nuclear collisions. *J. Phys. G*, 36:064022, 2009.
 - [10] J. H. Chen, Y. G. Ma, G. L. Ma, X. Z. Cai, Z. J. He, H. Z. Huang, J. L. Long, W. Q. Shen, C. Zhong, and J. X. Zuo. Elliptic flow of phi mesons and strange quark collectivity. *Phys. Rev. C*, 74:064902, 2006.
 - [11] Md. Nasim, Bedangadas Mohanty, and Nu Xu. Elliptic flow of ϕ mesons as a sensitive probe for the onset of the deconfinement transition in high energy heavy-ion collisions. *Phys. Rev. C*, 87(1):014903, 2013.
 - [12] Carlos A. Bertulani, Spencer R. Klein, and Joakim Nysstrand. Physics of ultra-peripheral nuclear collisions. *Ann. Rev. Nucl. Part. Sci.*, 55:271–310, 2005.
 - [13] C. Bottcher, M. R. Strayer, and M. J. Rhoades-Brown. Approximate analytic formula used to estimate electron capture cross-sections at relativistic energies. *Phys. Rev. A*, 44:4769–4770, 1991.
 - [14] C. F. von Weizsacker. Radiation emitted in collisions of very fast electrons. *Z. Phys.*, 88:612–625, 1934.
 - [15] Jaroslav Adam, Dagmar Adamová, Madan M Aggarwal, G Aglieri Rinella, Maria Agnello, Nikita Agrawal, Zubayer Ahammed, SU Ahn, S Aiola, A Akindinov, et al. Measurement of an excess in the yield of j/ψ at very low p_t in pb–pb collisions at $\sqrt{s_{NN}} = 2.76$ tev. *Physical review letters*, 116(22):222301, 2016.
 - [16] J. Adam et al. Observation of excess J/ψ yield at very low transverse momenta in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and U+U collisions at $\sqrt{s_{NN}} = 193$ GeV. *Phys. Rev. Lett.*, 123(13):132302, 2019.
 - [17] W. Zha, S. R. Klein, R. Ma, L. Ruan, T. Todoroki, Z. Tang, Z. Xu, C. Yang, Q. Yang, and S. Yang. Coherent J/ψ photoproduction in hadronic heavy-ion collisions. *Phys. Rev.*, C97(4):044910, 2018.
 - [18] Mariola Kłusek-Gawenda and Antoni Szczurek. Photoproduction of j/ψ mesons in peripheral and semicentral heavy ion collisions. *Phys. Rev. C*, 93:044912, Apr 2016.
 - [19] James Daniel Brandenburg, Zhangbu Xu, Wangmei Zha, Cheng Zhang, Jian Zhou, and Ya-jin Zhou. Exploring gluon tomography with polarization dependent diffractive J/ψ production. *Phys. Rev. D*, 106(7):074008, 2022.
 - [20] Jaroslav Adam et al. Low- p_T e^+e^- pair production in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and U+U collisions at $\sqrt{s_{NN}} = 193$ GeV at STAR. *Phys. Rev. Lett.*, 121(13):132301, 2018.
 - [21] S. Acharya et al. Dielectron production at midrapidity at low transverse momentum in peripheral and semi-peripheral Pb–Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. *JHEP*, 06:024, 2023.
 - [22] Wangmei Zha, James Daniel Brandenburg, Zebo Tang, and Zhangbu Xu. Initial transverse-momentum broadening of Breit-Wheeler process in relativistic heavy-ion collisions. *Phys. Lett. B*, 800:135089, 2020.
 - [23] James Daniel Brandenburg, Wangmei Zha, and Zhangbu Xu. Mapping the electromagnetic fields of heavy-ion collisions with the Breit-Wheeler process. *Eur. Phys. J. A*, 57(10):299, 2021.

- [24] Jiaxuan. Luo, Xinbai. Li, Zebo. Tang, Xin. Wu, and Wangmei. Zha. Effect of initial nuclear deformation on dielectron photoproduction in hadronic heavy-ion collisions. *Phys. Rev. C*, 108(5):054906, 2023.
- [25] Wangmei Zha, James Daniel Brandenburg, Lijuan Ruan, Zebo Tang, and Zhangbu Xu. Exploring the double-slit interference with linearly polarized photons. *Phys. Rev. D*, 103(3):033007, 2021.
- [26] Xin Wu, Xinbai Li, Zebo Tang, Pengfei Wang, and Wangmei Zha. Reaction plane alignment with linearly polarized photon in heavy-ion collisions. *Phys. Rev. Res.*, 4:L042048, Dec 2022.
- [27] A. Andronic et al. Heavy-flavour and quarkonium production in the LHC era: from proton–proton to heavy-ion collisions. *Eur. Phys. J. C*, 76(3):107, 2016.
- [28] W. Zha, S. R. Klein, R. Ma, L. Ruan, T. Todoroki, Z. Tang, Z. Xu, C. Yang, Q. Yang, and S. Yang. Coherent j/ψ photoproduction in hadronic heavy-ion collisions. *Phys. Rev. C*, 97:044910, Apr 2018.
- [29] Michael L. Miller, Klaus Reygers, Stephen J. Sanders, and Peter Steinberg. Glauber modeling in high energy nuclear collisions. *Ann. Rev. Nucl. Part. Sci.*, 57:205–243, 2007.
- [30] T. H. Bauer, R. D. Spital, D. R. Yennie, and F. M. Pipkin. The Hadronic Properties of the Photon in High-Energy Interactions. *Rev. Mod. Phys.*, 50:261, 1978. [Erratum: *Rev.Mod.Phys.* 51, 407 (1979)].
- [31] Jorg Hufner and Boris Z. Kopeliovich. J / Psi N and Psi-prime N total cross-sections from photoproduction data: Failure of vector dominance. *Phys. Lett. B*, 426:154–160, 1998.
- [32] Spencer R. Klein, Joakim Nystrand, Janet Seger, Yuri Gorbunov, and Joey Butterworth. STARlight: A Monte Carlo simulation program for ultra-peripheral collisions of relativistic ions. *Comput. Phys. Commun.*, 212:258–268, 2017.
- [33] R. C. Barrett, D. F. Jackson, and George W. Greenless. Nuclear Sizes and Structure. *Physics Today*, 31(7):47–48, 07 1978.
- [34] Shreyasi Acharya et al. Production of the $\rho(770)^0$ meson in pp and Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. *Phys. Rev. C*, 99(6):064901, 2019.
- [35] J. Adams et al. Rho0 production and possible modification in Au+Au and p+p collisions at $S(NN)^{1/2} = 200$ -GeV. *Phys. Rev. Lett.*, 92:092301, 2004.