
Upper Limit on the Coronal Cosmic Ray Energy Budget in Seyfert Galaxies

Yoshiyuki INOUE^{1,2,3}, Shinsuke TAKASAO¹ and Dmitry KHANGULYAN^{4, 5, 6}

¹Department of Earth and Space Science, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan

²Interdisciplinary Theoretical & Mathematical Science Program (iTHEMS), RIKEN, 2-1 Hirosawa, Saitama 351-0198, Japan

³Kavli Institute for the Physics and Mathematics of the Universe (WPI), UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

⁴Key Laboratory of Particle Astrophysics, Institute of High Energy Physics, Chinese Academy of Sciences, 100049 Beijing, China

⁵Tianfu Cosmic Ray Research Center, 610000 Chengdu, Sichuan, China

⁶Institute for Cosmic Ray Research, The University of Tokyo, 5-1-5 Kashiwa-no-Ha, Kashiwa City, Chiba, 277-8582, Japan

*E-mail: yoshiyuki.inoue.sci@osaka-u.ac.jp, takasao@astro-osaka.jp, khangulyan@ihep.ac.cn

Received (reception date); Accepted (acceptation date)

Abstract

The IceCube collaboration has reported possible detections of high-energy neutrinos from nearby Seyfert galaxies. While central hot coronae are proposed as the primary neutrino production site, the exact coronal cosmic-ray energy budget has been loosely constrained. In this study, we propose a new stringent upper bound on the coronal cosmic-ray energy budget of Seyfert galaxies, considering both accretion dynamics and observed properties of radio-quiet Seyfert galaxies. Notably, even under the calorimetric condition where cosmic rays lose all their energies, our limit indicates that the coronal neutrino flux of NGC 1068 is about an order of magnitude fainter than the observed levels. This discrepancy suggests the need for further theoretical and observational investigations on the IceCube signals from Seyfert galaxies.

Key words: accretion, accretion disks — black hole physics — galaxies: active — galaxies: jets — cosmic rays — neutrinos

1 Introduction

The IceCube Collaboration has reported a $4.2\text{-}\sigma$ detection of high-energy neutrinos from a nearby Seyfert galaxy, NGC 1068, based on 10 years of survey data (Aartsen et al. 2020; Goswami et al. 2022), where Seyfert galaxies are a subset of active galactic nuclei (AGNs) characterized by a lack of pronounced jet activity. Subsequently, they have also identified $2.9\text{-}\sigma$ signals from two other proximate Seyfert galaxies, NGC 4151 and CGCG 420-015 (Goswami 2023; Glauch et al. 2023). Intriguingly, the neutrino flux

of NGC 1068, as reported, exceeds its GeV gamma-ray flux (Aartsen et al. 2020; IceCube Collaboration et al. 2022; Ajello et al. 2023), which implies a significant attenuation of GeV gamma rays. Although the exact neutrino fluxes of NGC 4151 and CGCG 420-015 have not been determined yet, this discrepancy would also apply to these two Seyferts considering the Fermi observations (Abdollahi et al. 2022; Peretti et al. 2023).

This discrepancy between gamma rays and neutrinos has led to the proposal of various models to account for the neutrino signals from Seyferts, such as accretion-

powered winds from supercritically accreting compact binaries (Sridhar et al. 2022), muon pair production in AGN core (Hooper & Plant 2023), interaction between clouds and the accretion disk (Müller & Romero 2020), interaction between failed winds and the accretion disk (Inoue et al. 2022), jet interaction with interstellar medium (Fang et al. 2023), or stellar-mass black holes (BHs) embedded in the AGN accretion disk (Tagawa et al. 2023). Here, given various known observational properties, the hot plasma surrounding the central supermassive black hole (SMBH) and its accretion disk, namely the corona, emerges as one of the most plausible sites for neutrino production (Inoue et al. 2020; Murase et al. 2020; Gutiérrez et al. 2021; Kheirandish et al. 2021; Eichmann et al. 2022; Murase 2022; Neronov et al. 2023; Blanco et al. 2023; Ajello et al. 2023; Mbarek et al. 2023; Fiorillo et al. 2023).

The influence of high-energy cosmic rays (CRs) in SMBH coronae have been extensively investigated in the literature (see, e.g., Zdziarski 1986; Kazanas & Ellison 1986; Sikora et al. 1987; Begelman et al. 1990; Stecker et al. 1992; Kalashev et al. 2015; Inoue et al. 2019). However, a definitive constraint on the cosmic-ray energy budget within these coronal regions remains elusive, primarily due to the ambiguous non-thermal activity of Seyferts in electromagnetic spectra. To account for the observed neutrino signals, a specific CR energy budget is often postulated without strict constraints. In this paper, we aim to constrain this CR energy budget by considering accretion dynamics and the observed properties of the coronae. Furthermore, we explore potential limits on the high-energy neutrino luminosities characteristic of Seyfert galaxies.

2 Physical Properties of AGN Coronae

Accretion disk coronae serve as the primary sources of AGN X-ray emission through Comptonization of accretion disk photons (Katz 1976; Bisnovatyi-Kogan & Blinnikov 1977; Pozdniakov et al. 1977; Sunyaev & Titarchuk 1980). Through X-ray spectral analyses, we can deduce thermal coronal parameters, specifically the coronal electron temperature T_e and the Thomson scattering optical depth τ_X (Brenneman et al. 2014; Fabian et al. 2015; Ricci et al. 2018). X-ray AGNs typically have $k_B T_e \sim 50$ keV, where k_B is the Boltzmann constant, and $\tau_X \sim 1$ (Fabian et al. 2015; Ricci et al. 2018; Pal et al. 2023).

The coronal size R_c is measured as $R_c \equiv r_c R_s \sim 5 - 40 R_s$, where R_s is the Schwarzschild radius, by optical-X-ray spectral fitting studies (Jin et al. 2012; Kubota & Done 2018), microlensing observations (Morgan et al. 2012), and coronal synchrotron emission studies (Inoue & Doi

2018; Michiyama et al. 2023). In this paper, we adopt $r_c = 10$ as a fiducial value. This size is sufficient to attenuate GeV gamma-ray photons, which is required to match with the IceCube results of NGC 1068 (Inoue et al. 2020; Murase 2022).

Theoretically, it is expected that BH coronae are parts of hot accretion flows, heated through advection (Liu et al. 2002a; Kawanaka et al. 2008; Kawabata & Mineshige 2010), resulting in two-temperature plasma (Narayan & Yi 1994; Manmoto et al. 1997). Such two-temperature plasma has been also confirmed by recent numerical simulations (e.g., Liska et al. 2022). Hereinafter, for the sake of simplicity, we treat coronae as uniform and steady two-temperature plasma surrounding BHs.

The two-temperature plasma model naturally accounts for the observed coronal electron temperature in BH accretion systems (e.g., Kawabata & Mineshige 2010; Inoue et al. 2019). In the hot accretion flow, the ion temperature T_i is close to the virial temperature (Kato et al. 2008; Yuan & Narayan 2014)

$$T_i \approx \frac{GM_{\text{BH}}m_p}{6k_B R_c} = \frac{m_p c^2}{12k_B r_c} \simeq 9.1 \times 10^{10} \text{ K} \left(\frac{r_c}{10}\right)^{-1}, \quad (1)$$

where G is the gravitational constant, M_{BH} is the BH mass, m_p is the proton rest mass, and c is the speed of light. This temperature translates to $k_B T_i \sim 2.6$ MeV.

Through elastic Coulomb (EC) collisions, protons redistribute their heat to electrons, which are in turn cooled via Comptonization. The electron heating rate through EC collisions is given by

$$\frac{dT_e}{dt} \approx \frac{T_i}{t_{\text{EC},pe}} \approx \frac{n_e \sigma_T c \ln \Lambda}{\sqrt{\pi/2}} \left(\frac{m_e}{m_p}\right) T_i \theta_e^{-3/2}, \quad (2)$$

where $t_{\text{EC},pe}$ is the proton-electron Coulomb collision timescale (Spitzer 1962; Stepney 1983), n_e is the coronal electron density, σ_T is the Thomson scattering cross section, $\ln \Lambda \approx 20$ is the Coulomb logarithm, and $\theta_e \equiv k_B T_e / m_e c^2$ is the dimensionless electron temperature. We ignore the term of the dimensionless ion temperature $\theta_i \equiv k_B T_i / m_p c^2$ in $t_{\text{EC},pe}$, as $\theta_i \ll \theta_e$.

The electron cooling rate due to Comptonization is described as

$$\frac{dT_e}{dt} \approx -\frac{4}{3} \frac{\sigma_T u_{\text{ph}} T_e}{m_e c}, \quad (3)$$

where u_{ph} is the accretion disk photon energy density. We define the bolometric disk luminosity as $L_{\text{bol}} = \epsilon L_{\text{Edd}}$, where ϵ is the radiative efficiency and L_{Edd} is the Eddington luminosity. Thus, $u_{\text{ph}} = L_{\text{bol}} / 4\pi R_c^2 c$.

By balancing these electron heating and cooling rates, the coronal electron temperature is estimated to be

$$k_B T_e \approx \left(\frac{3 \ln \Lambda}{4\sqrt{\pi/2}} \frac{m_e}{m_p} \frac{n_e}{u_{\text{ph}}} k_B T_i \right)^{2/5} m_e c^2 \quad (4)$$

$$\simeq 84 \left(\frac{\tau_X}{1} \right)^{2/5} \left(\frac{\epsilon}{0.1} \right)^{-2/5} [\text{keV}],$$

where, given the coronal opacity and the size, the electron number density is expressed as $n_e \approx \tau_X / \sigma_T R_c$. The dependencies on τ_X and ϵ are retained. This temperature aligns closely with the measured coronal temperature ~ 50 keV (Fabian et al. 2015; Pal et al. 2023). Therefore, the two-temperature plasma model provides a coherent explanation for the observed coronal temperatures.

3 Bounds on Coronal CR Properties

3.1 Coronal CR Pressure

The total pressure in the coronal flow p is expressed as

$$p = p_{\text{gas}} + p_{\text{mag}} + p_{\text{cr}}, \quad (5)$$

where p_{gas} is the gas pressure, p_{mag} is the magnetic pressure, and p_{cr} is the cosmic-ray pressure. Here, we assume isotropic momentum distribution for CR particles. We neglect the radiation pressure because we consider the accretion rate regime $\dot{m} \ll 1$, where \dot{m} is the dimensionless mass accretion rate, scaled by the Eddington rate.

Given that the ion temperature T_i exceeds the electron temperature T_e , the coronal thermal gas pressure is dominated by ions, which is

$$p_{\text{gas}} = n_i k_B T_i + n_e k_B T_e \approx n_i k_B T_i, \quad (6)$$

where, assuming the charge neutrality, the ion number density n_i is set equal to n_e . Then, we have the coronal gas pressure as

$$p_{\text{gas}} \approx \frac{m_p c^4}{24 G \sigma_T} \frac{\tau_X}{M_{\text{BH}} r_c^2}. \quad (7)$$

If we consider positron contribution, n_i becomes smaller than n_e , so p_{gas} would decrease.

Magnetic fields accrete onto the central BH together with gas particles. The ratio of the gas pressure against the magnetic pressure is commonly defined by the plasma beta as

$$\beta \equiv \frac{p_{\text{gas}}}{p_{\text{mag}}}, \quad (8)$$

where we consider global coronal magnetic field.

In this paper, we introduce the ratio of the CR pressure against the magnetic pressure as

$$\delta \equiv \frac{p_{\text{cr}}}{p_{\text{mag}}}. \quad (9)$$

Observationally, as described above, coronal gamma-ray fluxes need to be attenuated by coronal X-ray photons. This implies that coronal CR particles should be generated and confined in the coronal region. Therefore, δ , which limits the CR energy budget in the coroneae, is

bounded as

$$\delta \leq 1. \quad (10)$$

This condition is also applicable if CR particles are accelerated by magnetic activity associated with coronal flows as we consider the time-averaged structure. When this CR trapping condition reaches $\delta = 1$, it signifies energy equipartition between magnetic fields and CR particles. It should be noted that in scenarios where $\delta > 1$ and $\beta < 1$, the CR pressure primarily sustains the accretion disk. This leads to distinct accretion dynamics compared to those dominated by gas, radiation, or magnetic pressure, which have been explored in literature.

Incorporating these considerations, the CR pressure is bounded in terms of the gas pressure as (Eqs. 7–9)

$$p_{\text{cr}} = \frac{\delta}{\beta} p_{\text{gas}} \approx \frac{m_p c^4}{24 G \sigma_T} \frac{\delta}{\beta} \frac{\tau_X}{M_{\text{BH}} r_c^2} \quad (11)$$

3.2 Bounds on Coronal Cosmic Ray and Neutrino Emissions

The energy density of cosmic rays, denoted as u_{cr} , is associated with the CR pressure as $u_{\text{cr}} = 3p_{\text{cr}}$. Those CRs generated in the coronal regions are transported to the further inner region while being trapped in turbulent magnetic fields in the accretion flow. As we consider geometrically thick coroneae, we apply spherical approximation here.

Since the energy source is the accretion process, the power of processes in the corona is limited by the rate at which the energy is supplied to the corona by the accretion flow. Using the standard expression for the energy flux (see, e.g., Landau & Lifshitz (1987)) one obtains

$$L_{\text{cr}} \approx 4\pi R_c^2 v_{\text{acc}} u_{\text{cr}}, \quad (12)$$

where v_{acc} represents the accretion speed. Utilizing the self-similar solutions (Narayan & Yi 1994; Yuan et al. 2012; Yuan & Narayan 2014), v_{acc} is approximated $v_{\text{acc}} \simeq 1.1 \times 10^{10} \alpha r_c^{-1/2} \text{ cm s}^{-1}$, where α is the dimensionless parameter representing the kinematic viscosity of the accretion flow, the so-called α parameter, which is typically set to be 0.1.

By combining Eqs. 10, 11 and 12, we can constrain the CR power as

$$L_{\text{cr}} \lesssim 7.3 \times 10^{41} \text{ erg s}^{-1} \left(\frac{\alpha}{0.1} \right) \left(\frac{\beta}{10} \right)^{-1} \times \left(\frac{\delta}{1} \right) \left(\frac{M_{\text{BH}}}{10^7 M_{\odot}} \right) \left(\frac{r_c}{10} \right)^{-1/2} \left(\frac{\tau_X}{1} \right) \quad (13)$$

Here, if energy is injected into CRs from other than accretion processes (see e.g., Blandford & Globus 2022 for the ejection disk model), higher CR power might be realized.

The neutrino production channels are primarily pp

and $p\gamma$ interactions. One needs to consider the efficiency of pp and $p\gamma$ processes. However, for the purpose of this study, we assume the calorimetric limit, where cosmic rays lose all their energies, for both processes to establish bounds on the neutrino luminosity to derive conservative upper limits. The bolometric neutrino luminosity per flavor is approximated as

$$L_\nu \approx f_\nu L_{\text{cr}} \mathcal{C}^{-1}, \quad (14)$$

where f_ν is the neutrino production fraction. $f_\nu \simeq 1/6$ for pp , while $f_\nu \simeq 1/8$ for $p\gamma$. \mathcal{C} reflects the spectral shape effect, which should be larger than unity. To derive the upper bound on the coronal neutrino luminosity, we take $f_\nu = 1/6$. Consequently, the coronal neutrino luminosity is bounded as

$$L_\nu \leq 1.2 \times 10^{41} \text{ erg s}^{-1} \left(\frac{\alpha}{0.1}\right) \left(\frac{\beta}{10}\right)^{-1} \times \left(\frac{\delta}{1}\right) \left(\frac{\mathcal{C}}{1}\right)^{-1} \left(\frac{M_{\text{BH}}}{10^7 M_\odot}\right) \left(\frac{r_c}{10}\right)^{-1/2} \left(\frac{\tau_X}{1}\right). \quad (15)$$

3.3 Coronal Neutrino Emissions of NGC 1068, NGC 4151, and CGCG 420-015

NGC 1068 is a Compton-thick Seyfert (Bauer et al. 2015; Marinucci et al. 2016; Mizumoto et al. 2019) located at the distance of $D_L = 13.97 \pm 2.10$ Mpc (Anand et al. 2021). The mass of the central SMBH of NGC 1068 has been estimated using various methods. Water maser disk studies have reported a mass of $\sim 10^7 M_\odot$ (Greenhill et al. 1996; Huré 2002; Lodato & Bertin 2003; Morishima et al. 2023)¹. We take $1 \times 10^7 M_\odot$ for the SMBH mass of NGC 1068.

Employing Eq. 15, we derive the upper bound for the total neutrino flux of

$$F_{\nu, \text{NGC 1068}} \leq 3.3 \times 10^{-12} \text{ TeV cm}^{-2} \text{ s}^{-1} \times \left(\frac{D_L}{13.97 \text{ Mpc}}\right)^{-2} \left(\frac{M_{\text{BH}}}{10^7 M_\odot}\right) \left(\frac{r_c}{10}\right)^{-1/2} \left(\frac{\tau_X}{1}\right) \times \left(\frac{\alpha}{0.1}\right) \left(\frac{\beta}{10}\right)^{-1} \left(\frac{\delta}{1}\right) \left(\frac{\mathcal{C}}{1}\right)^{-1}. \quad (16)$$

If we assume all the neutrino flux is generated at 1 TeV, the resulting flux bound would be $3.3 \times 10^{-12} \text{ TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$ at 1 TeV. Here, the observed neutrino flux of NGC 1068 is $\Phi_{\nu_\mu + \bar{\nu}_\mu}^{1 \text{ TeV}} = (5.0 \pm 1.5_{\text{stat}} \pm 0.6_{\text{sys}}) \times 10^{-11} \text{ TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$ at 1 TeV with a spectral index of $3.2 \pm 0.2 \pm 0.07$ (IceCube Collaboration et al. 2022) which is about an order of magnitude higher than our bound.

Similar disagreement is seen for the other two reported

Seyferts. NGC 4151 is the X-ray brightest type-1 Seyfert (Oh et al. 2018) located at the distance of $D = 15.8$ Mpc based on the Cepheid distance measurement (Yuan et al. 2020). Recently, GeV gamma-ray emission has been detected from this object (Ajello et al. 2021; Peretti et al. 2023). NGC 4151 has a central SMBH mass of $2.1 \times 10^7 M_\odot$ based on reverberation mapping measurements (De Rosa et al. 2018). CGCG 420-015 is a Compton-thick Seyfert (Tanimoto et al. 2018; Tanimoto et al. 2022) at a distance of 127 Mpc (Shimizu et al. 2016). The central SMBH mass is estimated as $2.0 \times 10^8 M_\odot$ (Koss et al. 2017). The corresponding neutrino flux upper limit is $\leq 5.3 \times 10^{-12} \text{ TeV cm}^{-2} \text{ s}^{-1}$ and $\leq 7.9 \times 10^{-13} \text{ TeV cm}^{-2} \text{ s}^{-1}$ for NGC 4151 and CGCG 420-015, respectively. These upper limits are about an orders of magnitude lower than the 5σ sensitivity of IceCube (Aartsen et al. 2020).

These discrepancies in NGC 1068, NGC 4151, and CGCG 420-015 suggest potential contributions of other mechanisms, other compact sources, or potential background contamination. Lower β would be able to reconcile the discrepancies. However, as described below, achieving low-beta plasma for the entire coronal region would not be straightforward, considering the nature of radio-quiet Seyferts and the dichotomy of AGNs.

4 Discussion

4.1 Challenge of Strongly Magnetized Coronal Models

When strong large-scale poloidal magnetic fields are present near the SMBHs, numerical simulations predict strong magnetic fields in the coronal region with values of $\beta \sim 10^{-2} - 10^{-3}$ (Liska et al. 2022). Such a low- β accreting plasma would be necessary to successfully reproduce powerful jets seen in radio-loud AGNs. In these scenarios, AGN coronae could be magnetically heated by reconnections (Haardt & Maraschi 1991; Liu et al. 2002b; Beloborodov 2017). However, our focus is on radio-quiet Seyfert galaxies, which, by definition, do not exhibit powerful jet activity.

Lack of powerful jets in Seyferts implies that large-scale poloidal fields should not be dynamically important. In such cases, only weakly magnetized coronae will develop (Liska et al. 2022). Given the absence of strong, large-scale poloidal magnetic fields, the Parker instability (Parker 1955; Parker 1966; Matsumoto et al. 1988) could inhibit the magnetic field amplification, keeping $\beta \gtrsim 10$ (see e.g., Takasao et al. 2018; Hogg & Reynolds 2018; Dhang & Sharma 2019). Consequently, this leads to β in the ranges of $\beta \sim 10 - 100$.

¹ Studies focusing on the polarized broad Balmer emission line and the neutral FeK α line have indicated masses of $\sim 7 \times 10^7 M_\odot$ and $\sim 1 \times 10^8 M_\odot$, respectively (Minezaki & Matsushita 2015).

We can relate the observed coronal magnetic field strength, B_c , to β , as the magnetic pressure is given by $p_{\text{mag}} \approx B^2/8\pi$. From observations, the coronal magnetic field B_c of nearby Seyferts with SMBH masses of 10^7 – $10^8 M_\odot$ is estimated to be in the range of $B_c \sim 10$ – 30 G with $r_c \sim 40$ (Inoue & Doi 2018; Michiyama et al. 2023), leading to $\beta \approx 100$. This estimate aligns with the arguments presented earlier. If this estimate holds, the coronal CR power could be reduced by an order of magnitude than our estimate, signifying the discrepancy between our bound and the observed neutrino flux.

The magnetic field estimates are based on the observations of millimeter (mm) emissions from Seyferts, utilizing the unprecedented angular resolution and sensitivity of ALMA. These emissions are interpreted as coronal synchrotron emissions (Inoue & Doi 2018; Michiyama et al. 2023). However, we note that the origin of these mm emissions remains a subject of debate (see, e.g., Raginski & Laor 2016; Baskin & Laor 2021). Nonetheless, the mm–X-ray luminosity correlation (Kawamuro et al. 2022; Ricci et al. 2023; Kawamuro et al. 2023) and the mm–X-ray time variability correlation (Behar et al. 2020; Chen et al. 2022; Panessa et al. 2022; Petrucci et al. 2023) strongly suggest that both X-ray and mm emissions originate from the same compact coronal region.

Low- β plasma might be prevalent in Seyfert coronae if the launch of jets is predominantly influenced by the angular momentum of the BHs. One widely accepted mechanism for the launch of powerful relativistic jets is the Blandford-Znajek process, in which the jet power is extracted by the rotation of SMBHs threaded by magnetic fields (Blandford & Znajek 1977). X-ray reflection spectra studies (e.g., Tanaka et al. 1995) indicate that most AGNs including Seyferts harbor rapidly rotating SMBHs (Vasudevan et al. 2016; Reynolds 2021). This predominance of spinning SMBHs in AGNs is also theoretically anticipated, based on studies of accretion and merger histories of SMBHs (Volonteri et al. 2013). Considering the insights from both theoretical and observational studies on SMBH spins, it is argued that a notable difference in magnetic flux strength, β , is inevitably necessary to explain the dichotomy of radio loudness of radio-loud and radio-quiet AGNs (Sikora et al. 2007; Sikora & Begelman 2013). Consequently, the presence of low- β coronal plasma in radio-quiet Seyferts would be unlikely, given their generally lower levels of jet activity compared to radio-loud populations.

4.2 Particle Acceleration Processes

Recent ALMA and IceCube observations indicate the presence of accelerated particles in AGN coronae (Inoue & Doi 2018; Inoue et al. 2020; IceCube Collaboration et al. 2022; Michiyama et al. 2023). Two-temperature accreting coronae are also characterized as collisionless plasma (e.g., Mahadevan & Quataert 1997), where non-thermal particles would be efficiently accelerated. Yet, the specific particle acceleration mechanisms at play within coronae remain elusive (Inoue et al. 2019; Murase et al. 2020; Kheirandish et al. 2021; Murase 2022; Mbarek et al. 2023; Fiorillo et al. 2023). Inoue et al. (2019) have previously discussed potential acceleration processes in AGN coronae. We expand upon their arguments by combining our coronal CR-bound studies.

Firstly, diffusive shock acceleration (DSA) would be responsible for particle acceleration in coronae (Stecker et al. 1992; Inoue et al. 2019). From the analysis of ALMA mm emission spectra, the injection spectral index of primary electrons at $\gamma_e \sim 10$ – 10^2 is likely around 2 (Inoue et al. 2019; Gutiérrez et al. 2021). This injection index aligns with what is typically expected in a simple DSA scenario for a strong shock (e.g., Drury 1983; Blandford & Eichler 1987). With $\beta \sim 10$ – 100 , DSA could potentially accelerate particles to energies of > 10 TeV (Inoue et al. 2019). Potential triggers for these shock accelerations include falling-back BLR clouds (Müller & Romero 2020; Müller et al. 2022; Sotomayor & Romero 2022) or failed-wind accretion (Inoue et al. 2022).

Secondly, stochastic acceleration might accelerate particles through the scattering by magnetic turbulence (Kimura et al. 2019; Murase et al. 2020; Eichmann et al. 2022). The stochastic acceleration is characterized by Alfvén speed $v_A = B/\sqrt{4\pi m_p n_i}$, turbulence index q , and the ratio of the strength of turbulence against the background (e.g., Dermer et al. 1996). The stochastic acceleration timescale depends on the magnetic field with $t_{\text{sta}} \propto B^{-7/3}$ with the Kolomogorov index $q = 5/3$ (Inoue et al. 2019; Eichmann et al. 2022). In low- β plasma environments (i.e., in radio-loud AGNs), stochastic acceleration is likely to emerge as the predominant process for particle acceleration (Zhdankin et al. 2018; Zhdankin et al. 2019). On the other hand, in radio-quiet AGNs having $\beta \sim 10$ – 100 , stochastic acceleration may not be as effective (but see also Kimura et al. 2015; Lynn et al. 2014; Sun & Bai 2021; Bacchini et al. 2024).

Magnetic reconnection flares are also potential particle accelerators (de Gouveia dal Pino & Lazarian 2005; Singh et al. 2015; Kheirandish et al. 2021; Fiorillo et al. 2023). The specific particle acceleration mechanisms during these

flares remain uncertain, with both DSA and stochastic acceleration being potential candidates (see e.g., Hoshino 2012; Nishizuka & Shibata 2013). It would be beyond the scope of this paper to understand the specific acceleration process occurring in reconnection. However, we can estimate the achievable reconnection flare power. Differential rotation in accretion disks generates shear in magnetic fields, which would trigger magnetic reconnections. The achievable generation rate of magnetic energy in the coronal region can be estimated as

$$P_{\text{mag}} \approx \frac{\Omega_{\text{K}} f_{\text{ani}} B_{\phi}^2 R_c^3}{12\pi}, \quad (17)$$

where Ω_{K} is the Keplerian angular velocity, B_{ϕ} is the azimuthal magnetic field strength, and f_{ani} gives the ratio between radial and azimuthal components of magnetic field strengths. The rotation period is $\approx 2\pi/\Omega_{\text{K}} \simeq 1.7 \text{ days} (M_{\text{BH}}/10^7 M_{\odot}) (r_c/30)^{3/2}$. Considering that a fraction of this magnetic energy f_{flare} will be released by reconnection (Shibata et al. 2013), we estimate the total reconnection power as

$$P_{\text{rec}} \approx f_{\text{flare}} P_{\text{mag}} \quad (18)$$

$$\approx f_{\text{flare}} \frac{\Omega_{\text{K}} f_{\text{ani}} B_{\phi}^2 R_c^3}{12\pi} \quad (19)$$

$$\simeq 2.5 \times 10^{40} \text{ erg s}^{-1} \left(\frac{f_{\text{flare}}}{0.1} \right) \left(\frac{f_{\text{ani}}}{1} \right) \left(\frac{\beta}{10} \right)^{-1} \\ \times \left(\frac{M_{\text{BH}}}{10^7 M_{\odot}} \right) \left(\frac{r_c}{10} \right)^{-1/2} \left(\frac{\tau_{\text{X}}}{1} \right), \quad (20)$$

where we set B_{ϕ} equal to the average B . This power is lower than our CR bound (Eq. 13).

As magnetic reconnections are transient phenomena and considering the radial dependence in Eq. 20, we would expect sporadic giant reconnection flares. Recent numerical simulations found that such giant flare would indeed occur in the inner regions of accretion disks, even in globally high- β plasma cases (Takasao et al. 2019; Porth et al. 2021; Ripperda et al. 2022; Ball et al. 2018). These reconnections could serve as a source of cosmic-ray particles or provide pre-accelerated particles. Detecting flaring variabilities with ALMA or IceCube would be crucial in confirming these phenomena.

4.3 Influence of Coronal Structure and Strong Gravity

In our analysis, we have not explored the inner structure of the corona or the influence of gravity. By adopting a density structure from hot accretion flow without any outflows, $n_i \propto r^{-3/2}$ (Yuan & Narayan 2014), we estimate that the CR power could increase by a factor of 10 in the case of a maximally rotating BH. However, the actual inner

structure of the corona diverges from this simple hot accretion flow model. For example, a more uniform density within $5\text{--}10R_s$ is presented by a recent numerical study (e.g., Liska et al. 2022).

Considering the coronae's proximity to the central BH, the impact of general relativity, especially gravitational redshift, is significant. This effect can dramatically reduce the observable flux from the core region. For rapidly spinning black holes, Hagen & Done (2023) demonstrate that emissions from the central region could be reduced by over 80% based on general relativistic ray-tracing studies. Thus, if neutrinos are produced deep within the coronae, the actual CR energy density would be considerably higher than currently estimated.

5 Summary

In this study, we explore the CR energy budget within the coronae of Seyfert galaxies. Taking into account the accretion dynamics and observed characteristics of Seyfert coronae, we establish a stringent upper limit on the coronal CR power, as denoted by Eq. 13. This constraint is more restrictive than those derived from bolometric luminosities. When considering the calorimetric limit, our established bound yields the neutrino flux that is approximately an order of magnitude lower than observations from the direction of NGC 1068. Such a discrepancy implies potential contributions from other compact regions or the possibility of background event contamination. A more in-depth combination of theoretical and observational analyses of the IceCube signals emanating from Seyfert galaxies is warranted.

Recent numerical simulations indicate a preference for high β plasma ($\beta \gg 1$) in AGNs lacking powerful jet activity (Liska et al. 2022). In such a scenario, the DSA process emerges as the favored particle acceleration mechanism. With a high β value, both stochastic acceleration and magnetic reconnection flares face challenges in efficiently accelerating particles. Nonetheless, sporadic giant magnetic reconnection flares (e.g., Takasao et al. 2019; Porth et al. 2021; Ripperda et al. 2022), whose magnetic fields are locally amplified, would potentially supply the necessary cosmic-ray population.

Acknowledgments

The authors would like to thank Chris Done, Scott, Hagen, and Ellis Owen for useful comments and discussions. YI is supported by NAOJ ALMA Scientific Research Grant Number 2021-17A and JSPS KAKENHI Grant Number JP18H05458, JP19K14772, and JP22K18277. ST is supported by JSPS KAKENHI Grant Number JP21H04487, JP22KK0043, and JP22K14074. This work was sup-

ported by World Premier International Research Center Initiative (WPI), MEXT, Japan.

References

- Aartsen, M. G., Ackermann, M., Adams, J., et al. 2020, *Phys. Rev. Lett.*, 124, 051103
- Abdollahi, S., Acero, F., Baldini, L., et al. 2022, *ApJS*, 260, 53
- Ajello, M., Baldini, L., Ballet, J., et al. 2021, *ApJ*, 921, 144
- Ajello, M., Murase, K., & McDaniel, A. 2023, *ApJL*, 954, L49
- Anand, G. S., Lee, J. C., Van Dyk, S. D., et al. 2021, *MNRAS*, 501, 3621
- Bacchini, F., Zhdankin, V., Gorbunov, E. A., et al. 2024, arXiv e-prints, arXiv:2401.01399
- Ball, D., Özel, F., Psaltis, D., Chan, C.-K., & Sironi, L. 2018, *ApJ*, 853, 184
- Baskin, A. & Laor, A. 2021, *MNRAS*, 508, 680
- Bauer, F. E., Arévalo, P., Walton, D. J., et al. 2015, *ApJ*, 812, 116
- Begelman, M. C., Rudak, B., & Sikora, M. 1990, *ApJ*, 362, 38
- Behar, E., Kaspi, S., Paubert, G., et al. 2020, *MNRAS*, 491, 3523
- Beloborodov, A. M. 2017, *ApJ*, 850, 141
- Bisnovatyi-Kogan, G. S. & Blinnikov, S. I. 1977, *A&A*, 59, 111
- Blanco, C., Hooper, D., Linden, T., & Pinetti, E. 2023, arXiv e-prints, arXiv:2307.03259
- Blandford, R. & Eichler, D. 1987, *Phys. Rep.*, 154, 1
- Blandford, R. & Globus, N. 2022, *MNRAS*, 514, 5141
- Blandford, R. D. & Znajek, R. L. 1977, *MNRAS*, 179, 433
- Brenneman, L. W., Madejski, G., Fuerst, F., et al. 2014, *ApJ*, 788, 61
- Chen, S., Laor, A., & Behar, E. 2022, *MNRAS*, 515, 1723
- de Gouveia dal Pino, E. M. & Lazarian, A. 2005, *A&A*, 441, 845
- De Rosa, G., Fausnaugh, M. M., Grier, C. J., et al. 2018, *ApJ*, 866, 133
- Dermer, C. D., Miller, J. A., & Li, H. 1996, *ApJ*, 456, 106
- Dhang, P. & Sharma, P. 2019, *MNRAS*, 482, 848
- Drury, L. O. 1983, *Reports on Progress in Physics*, 46, 973
- Eichmann, B., Oikonomou, F., Salvatore, S., Dettmar, R.-J., & Tjus, J. B. 2022, *ApJ*, 939, 43
- Fabian, A. C., Lohfink, A., Kara, E., et al. 2015, *MNRAS*, 451, 4375
- Fang, K., Lopez Rodriguez, E., Halzen, F., & Gallagher, J. S. 2023, arXiv e-prints, arXiv:2307.07121
- Fiorillo, D. F. G., Petropoulou, M., Comisso, L., Peretti, E., & Sironi, L. 2023, arXiv e-prints, arXiv:2310.18254
- Glauch, T., Kheirandish, A., Kontrimas, T., Liu, Q., & Niederhausen, H. 2023, arXiv e-prints, arXiv:2308.00024
- Goswami, S. 2023, arXiv e-prints, arXiv:2307.15349
- Goswami, S., Privon, G. C., Santander, M., et al. 2022, in 37th International Cosmic Ray Conference, 1142
- Greenhill, L. J., Gwinn, C. R., Antonucci, R., & Barvainis, R. 1996, *ApJL*, 472, L21
- Gutiérrez, E. M., Vieyro, F. L., & Romero, G. E. 2021, *A&A*, 649, A87
- Haardt, F. & Maraschi, L. 1991, *ApJL*, 380, L51
- Hagen, S. & Done, C. 2023, *MNRAS*, 525, 3455
- Hogg, J. D. & Reynolds, C. S. 2018, *ApJ*, 861, 24
- Hooper, D. & Plant, K. 2023, arXiv e-prints, arXiv:2305.06375
- Hoshino, M. 2012, *Phys. Rev. Lett.*, 108, 135003
- Huré, J. M. 2002, *A&A*, 395, L21
- IceCube Collaboration, Abbasi, R., Ackermann, M., et al. 2022, *Science*, 378, 538
- Inoue, S., Cerruti, M., Murase, K., & Liu, R.-Y. 2022, arXiv e-prints, arXiv:2207.02097
- Inoue, Y. & Doi, A. 2018, *ApJ*, 869, 114
- Inoue, Y., Khangulyan, D., & Doi, A. 2020, *ApJL*, 891, L33
- Inoue, Y., Khangulyan, D., Inoue, S., & Doi, A. 2019, *ApJ*, 880, 40
- Jin, C., Ward, M., Done, C., & Gelbord, J. 2012, *MNRAS*, 420, 1825
- Kalashhev, O., Semikoz, D., & Tkachev, I. 2015, *Soviet Journal of Experimental and Theoretical Physics*, 120, 541
- Kato, S., Fukue, J., & Mineshige, S. 2008, *Black-Hole Accretion Disks — Towards a New Paradigm —* (Kyoto University Press (Kyoto, Japan))
- Katz, J. I. 1976, *ApJ*, 206, 910
- Kawabata, R. & Mineshige, S. 2010, *PASJ*, 62, 621
- Kawamuro, T., Ricci, C., Imanishi, M., et al. 2022, *ApJ*, 938, 87
- Kawamuro, T., Ricci, C., Mushotzky, R. F., et al. 2023, arXiv e-prints, arXiv:2309.02776
- Kawanaka, N., Kato, Y., & Mineshige, S. 2008, *PASJ*, 60, 399
- Kazanas, D. & Ellison, D. C. 1986, *ApJ*, 304, 178
- Kheirandish, A., Murase, K., & Kimura, S. S. 2021, *ApJ*, 922, 45
- Kimura, S. S., Murase, K., & Toma, K. 2015, *ApJ*, 806, 159
- Kimura, S. S., Tomida, K., & Murase, K. 2019, *MNRAS*, 485, 163
- Koss, M., Trakhtenbrot, B., Ricci, C., et al. 2017, *ApJ*, 850, 74
- Kubota, A. & Done, C. 2018, *MNRAS*, 480, 1247
- Landau, L. D. & Lifshitz, E. M. 1987, *Fluid Mechanics*, Second Edition: Volume 6 (Course of Theoretical Physics), 2nd edn., Course of theoretical physics / by L. D. Landau and E. M. Lifshitz, Vol. 6 (Butterworth-Heinemann)
- Liska, M. T. P., Musoke, G., Tchekhovskoy, A., Porth, O., & Beloborodov, A. M. 2022, *ApJL*, 935, L1
- Liu, B. F., Mineshige, S., Meyer, F., Meyer-Hofmeister, E., & Kawaguchi, T. 2002a, *ApJ*, 575, 117
- Liu, B. F., Mineshige, S., & Shibata, K. 2002b, *ApJ*, 572, L173
- Lodato, G. & Bertin, G. 2003, *A&A*, 398, 517
- Lynn, J. W., Quataert, E., Chandran, B. D. G., & Parrish, I. J. 2014, *ApJ*, 791, 71
- Mahadevan, R. & Quataert, E. 1997, *ApJ*, 490, 605
- Manmoto, T., Mineshige, S., & Kusunose, M. 1997, *ApJ*, 489, 791
- Marinucci, A., Bianchi, S., Matt, G., et al. 2016, *MNRAS*, 456, L94
- Matsumoto, R., Horiuchi, T., Shibata, K., & Hanawa, T. 1988, *PASJ*, 40, 171
- Mbarek, R., Philippov, A., Chernoglazov, A., Levinson, A., & Mushotzky, R. 2023, arXiv e-prints, arXiv:2310.15222
- Michiyama, T., Inoue, Y., & Doi, A. 2023, *PASJ*, 75, 874
- Minezaki, T. & Matsushita, K. 2015, *ApJ*, 802, 98
- Mizumoto, M., Izumi, T., & Kohno, K. 2019, *ApJ*, 871, 156
- Morgan, C. W., Hainline, L. J., Chen, B., et al. 2012, *ApJ*, 756, 52
- Morishima, Y., Sudou, H., Yamauchi, A., Taniguchi, Y., & Nakai, N. 2023, *PASJ*, 75, 71
- Müller, A. L., Naddaf, M.-H., Zajaček, M., et al. 2022, *ApJ*, 931, 39
- Müller, A. L. & Romero, G. E. 2020, *A&A*, 636, A92
- Murase, K. 2022, *ApJL*, 941, L17
- Murase, K., Kimura, S. S., & Mészáros, P. 2020, *Phys. Rev. Lett.*, 125, 011101
- Narayan, R. & Yi, I. 1994, *ApJL*, 428, L13
- Neronov, A., Savchenko, D., & Semikoz, D. V. 2023, arXiv e-prints,

- arXiv:2306.09018
- Nishizuka, N. & Shibata, K. 2013, *Physical Review Letters*, 110, 051101
- Oh, K., Koss, M., Markwardt, C. B., et al. 2018, *ApJS*, 235, 4
- Pal, I., Anju, A., Sreehari, H., et al. 2023, arXiv e-prints, arXiv:2310.18196
- Panessa, F., Pérez-Torres, M., Hernández-García, L., et al. 2022, *MNRAS*, 510, 718
- Parker, E. N. 1955, *ApJ*, 121, 491
- Parker, E. N. 1966, *ApJ*, 145, 811
- Peretti, E., Peron, G., Tombesi, F., et al. 2023, arXiv e-prints, arXiv:2303.03298
- Petrucchi, P. O., Piétu, V., Behar, E., et al. 2023, arXiv e-prints, arXiv:2309.01804
- Porth, O., Mizuno, Y., Younsi, Z., & Fromm, C. M. 2021, *MNRAS*, 502, 2023
- Pozdniakov, L. A., Sobol, I. M., & Siuniaev, R. A. 1977, *Soviet Ast.*, 21, 708
- Raginski, I. & Laor, A. 2016, *MNRAS*, 459, 2082
- Reynolds, C. S. 2021, *ARA&A*, 59, 117
- Ricci, C., Chang, C.-S., Kawamuro, T., et al. 2023, *ApJL*, 952, L28
- Ricci, C., Ho, L. C., Fabian, A. C., et al. 2018, *MNRAS*, 480, 1819
- Ripperda, B., Liska, M., Chatterjee, K., et al. 2022, *ApJL*, 924, L32
- Shibata, K., Isobe, H., Hillier, A., et al. 2013, *PASJ*, 65, 49
- Shimizu, T. T., Meléndez, M., Mushotzky, R. F., et al. 2016, *MNRAS*, 456, 3335
- Sikora, M. & Begelman, M. C. 2013, *ApJL*, 764, L24
- Sikora, M., Kirk, J. G., Begelman, M. C., & Schneider, P. 1987, *ApJL*, 320, L81
- Sikora, M., Stawarz, Ł., & Lasota, J.-P. 2007, *ApJ*, 658, 815
- Singh, C. B., de Gouveia Dal Pino, E. M., & Kadowaki, L. H. S. 2015, *ApJL*, 799, L20
- Sotomayor, P. & Romero, G. E. 2022, *A&A*, 664, A178
- Spitzer, L. 1962, *Physics of Fully Ionized Gases* (New York: Interscience)
- Sridhar, N., Metzger, B. D., & Fang, K. 2022, arXiv e-prints, arXiv:2212.11236
- Stecker, F. W., Done, C., Salamon, M. H., & Sommers, P. 1992, *Physical Review Letters*, 69, 2738
- Stepney, S. 1983, *MNRAS*, 202, 467
- Sun, X. & Bai, X.-N. 2021, *MNRAS*, 506, 1128
- Sunyaev, R. A. & Titarchuk, L. G. 1980, *A&A*, 500, 167
- Tagawa, H., Kimura, S. S., & Haiman, Z. 2023, *ApJ*, 955, 23
- Takasao, S., Tomida, K., Iwasaki, K., & Suzuki, T. K. 2018, *ApJ*, 857, 4
- Takasao, S., Tomida, K., Iwasaki, K., & Suzuki, T. K. 2019, *ApJL*, 878, L10
- Tanaka, Y., Nandra, K., Fabian, A. C., et al. 1995, *Nature*, 375, 659
- Tanimoto, A., Ueda, Y., Kawamuro, T., et al. 2018, *ApJ*, 853, 146
- Tanimoto, A., Ueda, Y., Odaka, H., Yamada, S., & Ricci, C. 2022, *ApJS*, 260, 30
- Vasudevan, R. V., Fabian, A. C., Reynolds, C. S., et al. 2016, *MNRAS*, 458, 2012
- Volonteri, M., Sikora, M., Lasota, J. P., & Merloni, A. 2013, *ApJ*, 775, 94
- Yuan, F. & Narayan, R. 2014, *ARA&A*, 52, 529
- Yuan, F., Wu, M., & Bu, D. 2012, *ApJ*, 761, 129
- Yuan, W., Fausnaugh, M. M., Hoffmann, S. L., et al. 2020, *ApJ*, 902, 26
- Zdziarski, A. A. 1986, *ApJ*, 305, 45
- Zhdankin, V., Uzdensky, D. A., Werner, G. R., & Begelman, M. C. 2018, *ApJL*, 867, L18
- Zhdankin, V., Uzdensky, D. A., Werner, G. R., & Begelman, M. C. 2019, *Phys. Rev. Lett.*, 122, 055101