

# GENUINELY RAMIFIED MAPS AND MONODROMY

INDRANIL BISWAS, MANISH KUMAR, AND A. J. PARAMESWARAN

ABSTRACT. For any genuinely ramified morphism  $f : Y \rightarrow X$  between irreducible smooth projective curves we prove that  $(Y \times_X Y) \setminus \Delta$  is connected, where  $\Delta \subset Y \times_X Y$  is the diagonal. Using this result the following are proved:

- (1) If  $f$  is further Morse then the Galois closure is the symmetric group  $S_d$ , where  $d = \text{degree}(f)$ .
- (2) The Galois group of the general projection, to a line, of any smooth curve  $X \subset \mathbb{P}^n$  of degree  $d$ , which is not contained in a hyperplane and contains a non-flex point, is  $S_d$ .

## 1. INTRODUCTION

Let  $f : Y \rightarrow X$  be a generically smooth finite morphism of smooth projective curves over an algebraically closed field  $k$ . It is called genuinely ramified if the corresponding homomorphism of étale fundamental groups is surjective. A map  $Y \rightarrow X$  is genuinely ramified if and only if  $Y \times_X Y$  is connected [BP]. We prove the following improved version of it (see Theorem 3.1):

*For any genuinely ramified morphism  $Y \rightarrow X$ , the closure of*

$$(Y \times_X Y) \setminus \Delta \subset Y \times_X Y$$

*is connected, where  $\Delta$  is the diagonal in  $Y \times_X Y$ .*

Two applications of it are given.

*If a genuinely ramified morphism  $f : Y \rightarrow X$  of degree  $d$  is also Morse (see Section 2 for Morse map), then the Galois group of the Galois closure of  $f$  is  $S_d$ , where  $d = \text{degree}(f)$ . (See Corollary 3.2.)*

*Let  $X \subset \mathbb{P}^n$  be a smooth curve of degree  $d$  and not contained in any hyperplane. Assume that  $X$  contains a non-flex point (see Definition 3.4). Take a line  $L$  in the dual projective space  $\check{\mathbb{P}}^n$  such that it intersects the dual variety  $\check{X}$  transversely, every  $H \in L \cap \check{X}$  is an ordinary tangent to  $X$ , and  $X \cap H_1 \cap H_2$  is empty for  $H_1 \neq H_2 \in L$ . Let  $f : X \rightarrow L$  be the projection of  $X$  from the codimension two linear subspace in  $\mathbb{P}^n$  corresponding to  $L$ . Then the map  $f$  is Morse and its Galois closure has Galois group  $S_d$ . (See Corollary 3.6.)*

Note that the later result (Corollary 3.6) over the field of complex numbers is known and follows from the uniform position principle (see [PS]). Again over  $\mathbb{C}$ , the monodromy group of a degree  $d$  cover  $f : Y \rightarrow \mathbb{P}^1$  has been studied by many people (see [GM, Theorem 2], [Yo, Theorem 1 and 1'], etc). Apart from extending some of these results to

---

2010 *Mathematics Subject Classification.* 14H30, 14G17, 14H60.

*Key words and phrases.* Genuine ramification, fundamental group, monodromy, Morse map.

positive characteristic, our method allows computation of monodromy groups even when the base curve is not  $\mathbb{P}^1$ .

## 2. MORSE MAP

Let  $k$  be an algebraically closed field. Take connected smooth projective curves  $X$  and  $Y$  defined over  $k$ . A generically smooth finite morphism  $f : Y \rightarrow X$  is called a finite covering (or a cover). Since here  $X$  and  $Y$  are integral schemes, the condition that the morphism  $f$  is generically smooth is equivalent to  $f$  being separable. The cover  $f$  is called Galois if  $k(Y)/k(X)$  is a Galois extension. The cover  $f$  is called *Morse* if over every branch point  $x \in X$  the following two conditions hold: there is exactly one ramification point  $y$  over  $x$  and this  $y$  is a double point.

Take a generically smooth finite morphism  $f : Y \rightarrow X$ . Consider the fiber product  $Y \times_X Y$ . For  $i = 1, 2$ , let

$$q_i : Y \times_X Y \rightarrow Y \quad (2.1)$$

be the natural projection to the  $i$ -th factor. Consider the diagonal

$$\Delta \subset Y \times_X Y; \quad (2.2)$$

so  $\Delta$  is the image of the map  $Y \rightarrow Y \times_X Y$  given by the two copies of the identity map of  $Y$ . Let

$$Y' := \overline{(Y \times_X Y) \setminus \Delta} \subset Y \times_X Y \quad (2.3)$$

be the closure of the complement  $(Y \times_X Y) \setminus \Delta$  in  $Y \times_X Y$ .

**Lemma 2.1.** *Let  $f : Y \rightarrow X$  be a Morse map. Then  $Y'$  in (2.3) is smooth. If  $Y'$  is also connected, then the restriction  $q'_1 := q_1|_{Y'} : Y' \rightarrow Y$  of  $q_1$  (see (2.1)) is again a Morse map.*

*Proof.* Take any point  $(y_1, y_2) \in Y'$ , and denote the point  $f(y_1) = f(y_2) \in X$  by  $x$ . If  $y_1 \neq y_2$ , then  $f$  is étale at either  $y_1$  or  $y_2$ . In that case,  $(y_1, y_2)$  is a smooth point of  $Y \times_X Y$ , because the map  $q_i$  in (2.1) is smooth at  $(y_1, y_2)$  if  $f$  is étale at  $y_i$ . Hence  $(y_1, y_2)$  is a smooth point of  $Y'$ . If  $y_1 = y_2$ , and  $f$  is unramified at  $y_1$ , then again  $(y_1, y_2)$  is a smooth point of  $Y \times_X Y$  for the same reason, hence it cannot be a point of  $Y'$ .

Assume that  $y_1 = y_2$ , and that  $f$  is ramified at  $y_1$ . Then by hypothesis  $y_1$  is a double point of  $f$ . Note that  $Y \times_X Y$  has two smooth local branches at  $(y_1, y_1)$  with  $\Delta$  and  $Y'$  being the irreducible components through this point. Therefore, we conclude that  $Y'$  is smooth at  $(y_1, y_1)$ . So  $Y'$  is smooth.

For  $y \in Y$ , if  $f(y)$  is not a branch point of  $f$ , then  $(y, z)$  is not a branch point of  $q'_1$  for every  $z \in f^{-1}(f(y))$ . Assume that  $f(y)$  is a branch point of  $f$ . Let  $z \in f^{-1}(f(y))$  be the unique double point of  $f$  in  $f^{-1}(f(y))$ . Then  $(y, z)$  is the unique double point of  $q'_1$  in  $(q'_1)^{-1}(y)$  if  $y \neq z$  proving that the map  $q'_1$  is Morse. Note that if  $y = z$ , then again  $y$  is not a branch point of  $q'_1$ .  $\square$

**Lemma 2.2.** *Let  $f : Y \rightarrow X$  be a cover of degree  $d$  between irreducible smooth projective curves. If the complement  $(Y \times_X Y) \setminus \Delta$ , where  $\Delta \subset Y \times_X Y$  is the diagonal, is irreducible, then the Galois group of the Galois closure of  $f$  is a two-transitive subgroup of  $S_d$ . Conversely, if the Galois group of the Galois closure of  $f$  is a two-transitive subgroup of  $S_d$ , then  $(Y \times_X Y) \setminus \Delta$  is irreducible.*

*Proof.* Since  $f$  is a covering of degree  $d$ , the function field  $k(Y)$  of  $Y$  satisfies the condition

$$k(Y) = k(X)[z]/F(z)$$

for some irreducible separable polynomial  $F$  of degree  $d$ . As the map  $f$  is finite, every irreducible component of  $Y \times_X Y$  dominates  $X$ . Hence the irreducible components of  $Y \times_X Y$  are in bijection with the generic fiber of  $Y \times_X Y \rightarrow X$ . The generic fiber corresponds to the ring extension  $k(X) \subset k(Y) \otimes_{k(X)} k(Y)$ . But

$$k(Y) \otimes_{k(X)} k(Y) \cong k(Y)[z]/(F(z)).$$

Let  $\alpha$  be a root of  $F(z)$  in  $k(Y)$ . Then  $F(z) = (z - \alpha)F_1(z)$  in  $k(Y)[z]$ .

Note that  $F_1(z)$  is irreducible if and only if  $k(Y)[z]/(F(z))$  has only two closed points. The condition that  $k(Y)[z]/(F(z))$  has only two closed points is evidently equivalent to the condition that  $\Delta \subset Y \times_X Y$  as well as the closure of  $(Y \times_X Y) \setminus \Delta$  in  $Y \times_X Y$  are both irreducible.

Let  $\tilde{Y}$  be the Galois closure of  $f$ . Note that the Galois group  $G = \text{Gal}(\tilde{Y}/X)$  acts on the roots of  $F(z)$  in  $k(\tilde{Y})$ . Since  $F(z)$  is an irreducible separable polynomial of degree  $d$ , the above Galois group  $G$  is a transitive subgroup of  $S_d$ . Let  $\beta$  and  $\gamma$  be roots of  $F_1(z)$ . If  $F_1(z)$  is irreducible, then there is an automorphism  $\sigma$  of  $k(\tilde{Y})$  fixing  $k(X)(\alpha)$  such that  $\sigma(\beta) = \gamma$ . Hence  $G$  is two transitive.

Conversely if  $G$  is two transitive, then there is a  $\sigma \in G$  such that  $\sigma(\alpha) = \alpha$  and  $\sigma(\beta) = \gamma$ . Hence  $\sigma \in \text{Gal}(k(\tilde{Y})/k(X)(\alpha))$ . Since  $\beta$  and  $\gamma$  were arbitrary roots, it follows that  $F_1(z)$  is irreducible.  $\square$

### 3. GENUINELY RAMIFIED MORPHISMS

A covering map  $f : Y \rightarrow X$  between connected smooth projective curves is called *genuinely ramified* if any (all) of the following three equivalent conditions holds:

- (1) the induced homomorphism of étale fundamental groups  $f_* : \pi_1^{\text{ét}}(Y) \rightarrow \pi_1^{\text{ét}}(X)$  is surjective.
- (2) The fiber product  $Y \times_X Y$  is connected.
- (3)  $\dim H^0(Y, f^* f_* \mathcal{O}_Y) = 1$ .

(See [BP, p. 12828, Proposition 2.6] and [BP, p. 12830, Lemma 3.1].)

**Theorem 3.1.** *Let  $f : Y \rightarrow X$  be a genuinely ramified morphism between irreducible smooth projective curves. Then  $Y'$  (defined in (2.3)) is connected.*

*Proof.* We will show this by separating the cases when  $f$  is Galois and when it is not.

**Case 1:** Assume that the map  $f$  is Galois. Let

$$\Gamma := \text{Aut}(Y/X) \subset \text{Aut}(Y)$$

be the Galois group of  $f$ . For each  $\gamma \in \Gamma$ ,

$$Y_\gamma := \{(y, \gamma \cdot y) \mid y \in Y\} \subset Y \times_X Y \tag{3.1}$$

is evidently an irreducible component isomorphic to  $Y$ . It is straightforward to check that these are all the irreducible components of  $Y \times_X Y$ . In other words, the irreducible components of  $Y \times_X Y$  are parametrized by  $\Gamma$ .

Let  $G_f$  denote the dual graph of  $Y \times_X Y$  which is constructed as follows. The vertices of  $G_f$  are the irreducible components of  $Y \times_X Y$ , so the vertices of  $G_f$  are indexed by the elements of  $\Gamma$ . For any two vertices  $v_1$  and  $v_2$  of  $G_f$  there is at most one edge joining them, and there is an edge joining  $v_1$  and  $v_2$  if and only if the irreducible components of  $Y \times_X Y$  corresponding to  $v_1$  and  $v_2$  intersect. It is easy to see that the Galois group  $\Gamma$  acts naturally on  $G_f$ . The action of  $\Gamma$  on the vertices of  $G_f$  coincides with the left-translation action of  $\Gamma$  on itself (recall that the vertices are parametrized by the elements of  $\Gamma$ ). In particular, the action of  $\Gamma$  on the vertices of  $G_f$  is free and transitive.

Since  $Y \times_X Y$  is connected, so is  $G_f$ . The diagonal  $\Delta$  in (2.2) is  $Y_e$  (see (3.1)), where  $e \in \Gamma$  is the identity element. For any element  $g \in \Gamma$ , let  $G_f^g$  be the graph obtained from  $G_f$  by deleting the vertex corresponding to  $g$  together with all the edges that contain this vertex corresponding to  $g$ . So the closure  $\overline{(Y \times_X Y) \setminus \Delta}$  of  $(Y \times_X Y) \setminus \Delta \subset Y \times_X Y$  is connected if and only if the graph  $G_f^e$  is connected.

Assume that  $Y \times_X Y \setminus \Delta = Y \times_X Y \setminus Y_e$  is disconnected. As noted above, this implies that  $G_f^e$  is disconnected. Since the action of  $\Gamma$  on the vertices of  $G_f$  is transitive, we conclude that the graph  $G_f^g$  is disconnected for each  $g \in \Gamma$ .

**Claim 1.** Given a finite connected graph  $\mathbb{G}$ , let  $v$  and  $w$  be two vertices of  $\mathbb{G}$  such that distance between  $v$  and  $w$  is the largest among all pairs of vertices in  $\mathbb{G}$ . Then the graph  $\mathbb{G}_v$  obtained from  $\mathbb{G}$ , by deleting  $v$  together with also all edges containing  $v$ , is connected.

We will prove the claim by showing that all the vertices of  $\mathbb{G}_v$  are connected to  $w$ . If  $w'$  is a vertex of  $\mathbb{G}$  different from  $v$ , then the hypothesis that the distance between  $v$  and  $w$  is the largest implies that there is a path in  $\mathbb{G}$  connecting  $w$  and  $w'$  which does not pass through  $v$ . Hence  $w'$  and  $w$  are connected in  $\mathbb{G}_v$ .

Hence the closure of  $(Y \times_X Y) \setminus \Delta \subset Y \times_X Y$  is connected if  $f$  is Galois.

**Case 2.** Assume that  $f$  is not Galois.

Let

$$\tilde{f} : \tilde{Y} \longrightarrow X \tag{3.2}$$

be the Galois closure of  $f$  with  $h' : \tilde{Y} \longrightarrow Y$  being the map through which  $\tilde{f}$  factors, meaning  $\tilde{f} = f \circ h'$ . It is known that  $f$  is genuinely ramified does not imply  $\tilde{f}$  is genuinely ramified. Let  $g : Z \longrightarrow X$  be the maximal étale cover dominated by  $\tilde{f}$ , and let  $h : \tilde{Y} \longrightarrow Z$  be the factor map; so  $\tilde{f} = g \circ h$ . Let  $W$  be the fiber product  $Z \times_X Y$ , and also denote by  $p_Z$  (respectively,  $p_Y$ ) the projection map from  $W$  to  $Z$  (respectively,  $Y$ ). Note that since  $g : Z \longrightarrow X$  is étale, the map  $p_Y$  is also étale. Moreover, since  $Y$  is smooth, the fiber product  $W$  is also smooth. Consequently, we get the following

commutative diagram:

$$\begin{array}{ccccc}
 & & \tilde{Y} & & \\
 & & \downarrow q & & \\
 & h \swarrow & W & \searrow h' & \\
 & p_Z \swarrow & & \searrow p_Y & \\
 Z & & & & Y \\
 & g \swarrow & & \searrow f & \\
 & & X & & 
 \end{array} \tag{3.3}$$

Since  $f$  is genuinely ramified and  $g$  is étale, we have  $k(Z) \cap k(Y) = k(X)$ . Also note that  $k(Z)$  is Galois over  $k(X)$ . This is because the Galois closure of an étale cover is étale and  $g$  is the maximal intermediate étale cover of the ramified Galois cover  $\tilde{f}$ . So  $W$  is an irreducible smooth curve and the maps  $q$ ,  $p_Y$ ,  $h'$  and  $h$  are all Galois covers. Since  $g$  is the maximal étale cover dominated by  $\tilde{f} : \tilde{Y} \rightarrow X$ , the map  $h$  is genuinely ramified. Consequently,  $p_Z$  — being dominated by  $h$  — is also genuinely ramified.

Since  $p_Z$  is genuinely ramified, the fiber product  $W \times_Z W$  is connected. Let

$$D_{W/Z} \subset W \times_Z W$$

be the diagonal, so  $D_{W/Z}$  is identified with  $W$ . Consider the map

$$\Phi := (p_Y \times p_Y)|_{W \times_Z W} : W \times_Z W \rightarrow Y \times_X Y.$$

**Claim 2.** The inverse image, under this map  $\Phi$ , of  $\Delta \subset Y \times_X Y$  coincides with  $D_{W/Z}$ .

It can be seen that to prove the above claim it suffices to prove that the degree of the map  $\Phi : W \times_Z W \rightarrow Y \times_X Y$  coincides with the degree of  $p_Y : W \rightarrow Y$ . Indeed, as  $D_{W/Z}$  is a component of  $\Phi^{-1}(\Delta)$ , we have  $D_{W/Z} = \Phi^{-1}(\Delta)$  if degrees of  $p_Y$  and  $\Phi$  coincide. Now note that  $W \times_Z W$  is the fiber product of  $Y \times_X Y \rightarrow X$  and  $g : Z \rightarrow X$  with  $\Phi : W \times_Z W \rightarrow Y \times_X Y$  being the first projection. Hence we have  $\text{degree}(\Phi) = \text{degree}(g) = \text{degree}(p_Y)$ . This proves the claim.

From Claim 2 it follows that  $(Y \times_X Y) \setminus \Delta$  is connected if  $W \times_Z W \setminus D_{W/Z}$  is connected (note that  $\Phi$  is surjective).

Let  $G$  (respectively,  $H$ ) be the Galois group of  $h$  (respectively,  $q$ ); see (3.3). Then  $H \leq G$  is a subgroup (need not be normal). Note that the preimage of  $D_{W/Z}$  under the morphism  $\tilde{Y} \times_Z \tilde{Y} \rightarrow W \times_Z W$  is  $\tilde{Y} \times_W \tilde{Y}$ . Hence to show that  $(W \times_Z W) \setminus D_{W/Z}$  is connected it is enough to show that  $(\tilde{Y} \times_Z \tilde{Y}) \setminus \tilde{Y} \times_W \tilde{Y}$  is connected. Let  $\Gamma_G$  (respectively,  $\Gamma_H$ ) be the dual graph of  $\tilde{Y} \times_Z \tilde{Y}$  (respectively,  $\tilde{Y} \times_W \tilde{Y}$ ). Then  $\Gamma_H$  is a subgraph of  $\Gamma_G$ . It will follow that  $\tilde{Y} \times_Z \tilde{Y} \setminus \tilde{Y} \times_W \tilde{Y}$  is connected once we are able to show that  $\Gamma_G \setminus \Gamma_H$  is connected.

As  $h$  is genuinely ramified the graph  $\Gamma_G$  is connected. Also  $G$  acts faithfully and transitively on  $\Gamma_G$ . The cosets of  $H$  define a partition of  $\Gamma_G$  with  $\Gamma_H$  as one of the parts. Let  $\Gamma$  be the quotient graph of  $\Gamma_G$  with respect to this partition. Then the  $H$ -cosets in  $G$  are the vertices of  $\Gamma$ . Since  $\Gamma_G$  is connected, so is  $\Gamma$ . Also, as  $G$  acts on the set of  $H$ -cosets transitively, the action of  $G$  on  $\Gamma$  is transitive. Since  $\Gamma$  is connected there exists a vertex

$gH$  in  $\Gamma$  such that  $\Gamma \setminus \{gH\}$  is connected (see Claim 1). As  $G$  acts transitively on  $\Gamma$ , it now follows that  $\Gamma \setminus \{eH\}$  is connected. But this implies that  $\Gamma_G \setminus \Gamma_H$  is connected.  $\square$

**Corollary 3.2.** *Let  $f : Y \rightarrow X$  be a genuinely ramified cover of degree  $d$  between irreducible smooth projective curves such that  $f$  is a Morse map. Then the Galois group of the Galois closure of  $f$  is full  $S_d$ .*

*Proof.* Note that  $(Y \times_X Y) \setminus \Delta$  is connected by Theorem 3.1. Lemma 2.1 implies that  $(Y \times_X Y) \setminus \Delta$  is smooth and hence it is irreducible. Lemma 2.2 says that the Galois group  $G$  of the Galois closure of  $f$  is a two transitive subgroup of  $S_d$ . Since the ramification type of  $f$  is  $(2, 1, 1, \dots, 1)$ , the group  $G$  contains a transposition ([Abh, Cycle lemma, page 95] or [Ju, Lemma 2.4]). Hence we have  $G = S_d$  [Se, Lemma 4.4.3].  $\square$

**Corollary 3.3.** *Let  $f : Y \rightarrow X$  be a genuinely ramified cover of degree  $d$  between irreducible smooth curves such that  $f$  is a Morse map. Then  $Y'$  in (2.3) is smooth irreducible, and*

$$q'_1 : Y' \rightarrow Y$$

(see Lemma 2.1 for  $q'_1$ ) is a genuinely ramified Morse map of degree  $d - 1$ .

*Proof.* As in (3.2), let  $\tilde{f} : \tilde{Y} \rightarrow X$  be the Galois closure of  $f$ . By Corollary 3.2, this  $\tilde{f}$  is a  $S_d$ -Galois cover. Lemma 2.1 implies that  $Y'$  is smooth and Theorem 3.1 implies  $Y'$  is connected, and hence  $Y'$  is irreducible. Consequently, by Lemma 2.1, the covering morphism  $q'_1$  is a Morse map of degree  $d - 1$ .

Note that  $h' : \tilde{Y} \rightarrow Y$  in (3.3) is a  $S_{d-1}$ -Galois cover which dominates  $q' : Y' \rightarrow Y$ . This follows by noticing that  $\text{Gal}(k(\tilde{Y})/k(Y))$  is an index  $d$  subgroup of  $\text{Gal}(k(\tilde{Y})/k(X))$  and the fact that any subgroup of  $S_d$  of index  $d$  is actually  $S_{d-1}$ . It is standard fact that there is no proper subgroup of  $S_{d-1}$  that properly contains  $S_{d-2}$ . This implies that there is no field in between  $k(Y)$  and  $k(Y')$ . Hence  $q'_1 : Y' \rightarrow Y$  does not admit any intermediate covers. So if we show that  $q'_1$  is ramified then it is genuinely ramified.

Note that if  $d = 2$  then  $q'_1$  is degree 1 and there is nothing more to show. If  $d \geq 3$ , then note that  $h'$  is ramified. Indeed, for a branch point  $x \in X$  of  $f$ , every point of  $\tilde{f}^{-1}(x)$  is a ramification point of  $\tilde{f}$ . Hence every point in  $f^{-1}(x)$  at which  $f$  is unramified is a branch point of  $h'$ . Since  $f$  is a Morse map and the degree  $f$  is at least 3, it follows that  $h'$  is a ramified cover. Also note that  $h'$  is the Galois closure of  $q'_1$  as the degree of  $q'_1$  is  $d - 1$ . Hence  $q'_1$  is also ramified.  $\square$

**Definition 3.4.** Let  $X \subset \mathbb{P}^n$  be a smooth curve of degree  $d$  not contained in any hyperplane. A hyperplane  $H \subset \mathbb{P}^n$  is said to be an *ordinary tangent* of  $X$  at  $x \in X$  if  $x \in H$ , the multiplicity of the intersection of  $H$  and  $X$  at  $x$  is two and their intersection is transversal at points of  $X \cap H \setminus \{x\}$ . We will say a point  $x \in X$  is *non-flex* if there is an hyperplane  $H \subset \mathbb{P}^n$  which is an ordinary tangent of  $X$  at  $x$ .

The dual variety

$$\check{X} := \{H \in \check{\mathbb{P}}^n \mid H \text{ does not intersect } X \text{ transversally}\}$$

is a hypersurface in  $\check{\mathbb{P}}^n$ .

**Lemma 3.5.** *Let  $X \subset \mathbb{P}^n$  be a smooth curve of degree  $d$  not contained in any hyperplane. Assume that  $X$  has a non-flex point. There is an open dense subset  $U$  of  $\check{X}$  such that every  $H$  in  $U$  is an ordinary tangent of  $X$ .*

*Proof.* The inclusion map  $X \hookrightarrow \mathbb{P}^n$  will be denoted by  $\iota_X$ . Let

$$\mathbb{P}^n \times \check{X} \supset \mathcal{H} \xrightarrow{\varpi'} \check{X} \quad (3.4)$$

be the tautological hyperplane bundle over  $\check{X}$ . Let

$$\phi : \mathcal{H} \longrightarrow \mathbb{P}^n \quad (3.5)$$

be the natural projection. Define  $\mathcal{X} := \phi^{-1}(X) \subset \mathcal{H}$ . The inclusion map  $\mathcal{X} \hookrightarrow \mathcal{H}$  will be denoted by  $\iota_{\mathcal{X}}$ . Let

$$\varphi : \mathcal{X} \longrightarrow X \quad (3.6)$$

be the natural projection. Note that  $\varphi$  is given by the restriction, to  $\mathcal{X}$ , of the map  $\phi$  in (3.5). Define

$$\varpi : \mathcal{X} \longrightarrow \check{X} \quad (3.7)$$

to be the restriction of  $\varpi'$  in (3.4). Note that  $\varpi$  is a finite morphism.

Let  $T_{\varpi'} \subset T\mathcal{H}$  be the relative tangent bundle for the projection  $\varpi'$  in (3.4). The natural injective homomorphisms

$$\iota_{\mathcal{X}}^* T_{\varpi'} \longrightarrow (\iota_X \circ \varphi)^* T\mathbb{P}^n \quad \text{and} \quad \varphi^* TX \longrightarrow (\iota_X \circ \varphi)^* T\mathbb{P}^n$$

together produce a homomorphism

$$\Psi : \iota_{\mathcal{X}}^* T_{\varpi'} \oplus \varphi^* TX \longrightarrow (\iota_X \circ \varphi)^* T\mathbb{P}^n.$$

From the assumption that  $X$  has a non-flex point it follows immediately that  $\Psi$  is surjective over some point. Let

$$\mathcal{W} \subset \mathcal{X}$$

be the nonempty Zariski open subset over which  $\Psi$  is an isomorphism. Consider the Zariski open subset

$$\mathcal{U} := \varpi(\mathcal{W}) \subset \check{X}.$$

Take any  $H_0 \in \check{X}$  which is an ordinary tangent of  $X$  at some point  $x_0 \in X$  (such an hyperplane exists by our assumption). Then

$$\varpi^{-1}(H_0) \cap \mathcal{W} = (X \cap H_0) \setminus \{x_0\}.$$

In particular,  $\varpi^{-1}(H_0) \cap \mathcal{W}$  consists of distinct  $d - 2$  reduced points of  $X$ . Note that  $\varpi^{-1}(H_0) = (\varpi^{-1}(H_0) \cap \mathcal{W}) \cup 2x_0 \subset X$ . Also, note that for any  $H \in \mathcal{U}$ , the inverse image  $\varpi^{-1}(H) \cap \mathcal{W}$  is a reduced subscheme of  $X$  whose length is at most  $d - 2$ .

Let  $U \subset \mathcal{U}$  be the locus of all points  $H \in \mathcal{U}$  such that the length of  $\varpi^{-1}(H) \cap \mathcal{W}$  is  $d - 2$  (the maximal possible).

The hyperplane in  $\mathbb{P}^n$  corresponding to each point of  $U$  is an ordinary tangent of  $X$  at some point. Also,  $U$  is an open subset of  $\check{X}$ .  $\square$

Under the hypothesis of the Lemma 3.5, let  $L$  be a line in  $\check{\mathbb{P}}^n$  that intersects  $\check{X}$  transversely,  $L \cap \check{X}$  is contained in  $U$  and  $X \cap H_1 \cap H_2$  is empty for  $H_1 \neq H_2 \in L$ . Note that a general  $L$  in  $\check{\mathbb{P}}^n$  will have this property. We define a map

$$f : X \longrightarrow L \quad (3.8)$$

as follows. For  $x \in X$ , let  $H_x$  be the set of hyperplanes in  $\mathbb{P}^n$  passing through  $x$ . Then  $H_x$  defines a hyperplane in  $\mathbb{P}^n$  not containing  $L$  and hence it intersects  $L$  at a unique point. The map  $f$  sends  $x$  to this point of  $L$ .

The following is a generalization of [Yo] to any smooth irreducible curve in  $\mathbb{P}^n$  and on the other hand to any characteristic.

**Corollary 3.6.** *Let the hypothesis on  $X$  be as in Lemma 3.5 and  $f : X \rightarrow L$  be as in (3.8). The projection of  $X$  from the codimension two linear subspace in  $\mathbb{P}^n$  corresponding to  $L$  coincides with the map  $f$ . The map  $f$  is Morse and has Galois group  $S_d$ .*

*Proof.* Let  $H_1$  and  $H_2$  be two distinct points of  $L$ ; then  $A = H_1 \cap H_2$  is a codimension two linear subspace of  $\mathbb{P}^n$ . The map  $f : X \rightarrow L$  is the projection of  $X$  from  $A$ .

By the description of  $f$ , if  $H \in L$  is not in  $\check{X}$  then  $H$  intersects  $X$  transversally and hence  $f^{-1}(H)$  has  $d$  distinct points. So  $f$  is not branched at  $H$ . If  $H \in L \cap U$  then  $H$  intersects  $X$  transversely at  $d-2$  distinct points and  $H$  intersects  $X$  with multiplicity two at one point (say  $x$ ). Hence  $f$  is étale at all points of  $f^{-1}(H)$  except at  $x$  and  $x$  is a double point of  $f$ . Hence  $f$  is Morse, since  $L \cap \check{X}$  is contained in  $U$ .

Since  $\mathbb{P}^1$  is simply connected  $f$  is also genuinely ramified. Hence by Corollary 3.2, it follows that the Galois group of  $f$  is  $S_d$ .  $\square$

**Remark 3.7.** Note that the hypothesis that  $X \subset \mathbb{P}^2$  contains a non-flex point is equivalent to the dual map from  $X$  to  $\check{X}$  is separable if the characteristic of the base field is odd (see [Par, Proposition 1.5]).

#### ACKNOWLEDGEMENTS

We are very grateful to the referee for comments which helped in removing an error and improving the exposition.

#### REFERENCES

- [Abh] S. S. Abhyankar, Galois theory on the line in nonzero characteristic, *Bull. Amer. Math. Soc.* **27** (1992), 68–133.
- [BP] I. Biswas and A. J. Parameswaran, Ramified covering maps and stability of pulled back bundles, *Int. Math. Res. Not.*, 2022, no. 17, 12821–12851.
- [GM] R. Guralnick and K. Magaard, On the minimal degree of a primitive permutation group, *J. Algebra* **207** (1998), 127–145,
- [Ju] J. Juul, Iterates of generic polynomials and generic rational functions, *Trans. Amer. Math. Soc.* **371** (2019), 809–831.
- [Par] R. Pardini, Some remarks on plane curves over fields of finite characteristic, *Compositio Math.* **60** (1986), 3–17.
- [PS] G. P. Pirola and E. Schlesinger, Monodromy of projective curves, *J. Algebraic Geom.* **14** (2005), 623–642.
- [Se] J.-P. Serre, Topics in Galois theory, *Res. Notes in Math.* **1**, Jones and Bartlett, Boston, MA, 1992
- [Yo] H. Yoshihara, Function field theory of plane curves by dual curves, *Jour. Alg.* **239** (2001), 340–355.



DEPARTMENT OF MATHEMATICS, SHIV NADAR UNIVERSITY, NH91, TEHSIL DADRI, GREATER NOIDA, UTTAR PRADESH 201314, INDIA

*Email address:* `indranil.biswas@snu.edu.in`, `indranil29@gmail.com`

STATISTICS AND MATHEMATICS UNIT, INDIAN STATISTICAL INSTITUTE, BANGALORE 560059, INDIA

*Email address:* `manish@isibang.ac.in`

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400005, INDIA

*Email address:* `param@math.tifr.res.in`