

Large-space and long-time asymptotic behaviors of N_∞ -soliton solutions for the focusing Hirota equation

Weifang Weng^a, Zhenya Yan^{b,c,*}

^aDepartment of Mathematical Sciences, Tsinghua University, Beijing 100084, China

^bKLMM, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China

^cSchool of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

Abstract The Hirota equation is one of the higher-order extensions of the nonlinear Schrödinger equation, and can describe the ultra-short optical pulse propagation in the form $iq_t + \alpha(q_{xx} + 2|q|^2q) + i\beta(q_{xxx} + 6|q|^2q_x) = 0$, $(x, t) \in \mathbb{R}^2$ ($\alpha, \beta \in \mathbb{R}$). In this paper, we analytically explore the large-space and long-time asymptotic behaviors of a soliton gas (the limit $N \rightarrow \infty$ of N -soliton solutions) for the Hirota equation including the complex modified KdV equation by using the Riemann-Hilbert problems with discrete spectra restricted in intervals $(ia, ib) \cup (-ib, -ia)$ ($0 < a < b$). We find that this soliton gas tends slowly to the Jacobian elliptic wave solution with an error $\mathcal{O}(|x|^{-1})$ (zero exponentially quickly) as $x \rightarrow -\infty$ ($x \rightarrow +\infty$). We also present the long-time asymptotics of the soliton gas under the different velocity conditions: $x/t > 4\beta b^2$, $\xi_c < x/t < 4\beta b^2$, $x/t < \xi_c$. Moreover, we analyze the property of the soliton gas of the Hirota equation for the case of the discrete spectra filling uniformly a quadrature domain.

Mathematics Subject Classification: Primary 35Q51 · 35Q15 · 37K15; Secondary 35C20

1 Introduction

Since Zabusky and Kruskal [1] first coined the concept of “soliton” for the study of nonlinear wave propagations of the Korteweg-de Veris (KdV) equation with the periodic initial data in 1965, and Gardner-Greene-Kruskal-Miura [2] first presented the inverse scattering transform (IST) method (alias nonlinear Fourier transform [3]) to analytically solve the N -soliton solution of the KdV equation starting from its linear spectral problem (alias the Lax pair [4]) in 1967, more and more one-soliton and multi-soliton wave phenomena have been found to appear in a variety of physical models including the integrable and nearly-integrable nonlinear wave equations arising from many fields [5–14]. The N -soliton solutions solving integrable nonlinear wave equations via the IST provide us with the very good frameworks of nonlinear superpositions to further analyze their elastic interactions and other related properties. For example, in 1971, Zakharov [15] first introduced the concept of a *soliton gas*, which was defined as the limit of $N \rightarrow \infty$ of the N -soliton solution of the KdV equation. After that, the soliton gas was extended to study the soliton gas, breather gas, dense soliton gas, and hydrodynamic of other nonlinear wave equations [16–22]. More recently, Grava *et al* [23–25] presented a new idea to study the soliton gas, dense soliton gas, and soliton shielding of the KdV, modified KdV, and NLS equations, respectively, starting from the N -soliton solution characterized by a Riemann-Hilbert problem.

Moreover, the larger-order limits of multipole solitons and rogue waves were studied [26–29] by using the robust inverse scattering transform [30] and Riemann-Hilbert problems [31, 32]. The Deift-Zhou’s nonlinear steepest descent method [32] and $\bar{\partial}$ -steepest descent method [33, 34] with Riemann-Hilbert problems [35–37] were used to study the long-time asymptotics of some integrable nonlinear wave systems, for example, the nonlinear Schrödinger (NLS), KdV, Camassa-Holm (CH), Degasperis-Procesi, Fokas-Lenells, Sasa-Satsuma, modified CH, derivative NLS, and short-pulse equations (see [38–52] and references therein).

To study more complicated nonlinear wave phenomena (e.g., the phenomena of oceanic and optical waves), the extensions of the NLS equation were needed. For example, when the optical pulses become shorter (e.g., 100 fs [11, 12]), higher-order dispersive and nonlinear effects such as third-order dispersion, self-frequency shift,

* Email address: zyyan@mmrc.iss.ac.cn (Corresponding author)

and self-steepening arising from the stimulated Raman scattering are significant in the study of ultra-short optical pulse propagation to derive the multi-parameter higher-order NLS equation [53–56]

$$iq_t + \alpha_1 q_{xx} + \alpha_2 |q|^2 q + i\epsilon(\beta_1 q_{xxx} + \beta_2 |q|^2 q_x + \beta_3 q(|q|^2)_x) = 0, \quad (x, t) \in \mathbb{R}^2, \quad (1.1)$$

where $q = q(x, t)$ is complex envelop field, $\alpha_{1,2}, \epsilon, \beta_{1,2,3}$ are real parameters. Generally speaking, Eq. (1.1) may not be completely integrable. However, some special cases of Eq. (1.1) can be solved via the IST if these parameters satisfy some constraint conditions. For example, when $\beta_1 : \beta_2 : \beta_3 = 0 : 0 : 0$, Eq. (1.1) reduces to the integrable focusing ($\alpha_1 \alpha_2 > 0$) or defocusing ($\alpha_1 \alpha_2 < 0$) NLS equation [57]

$$iq_t + \alpha_1 q_{xx} + \alpha_2 |q|^2 q = 0. \quad (1.2)$$

When $\beta_1 : \beta_2 : \beta_3 = 0 : 1 : 1$, Eq. (1.1) reduce to the the modified derivative NLS equation [58]

$$iq_t + \alpha_1 q_{xx} + \alpha_2 |q|^2 q + i\epsilon \beta_2 (|q|^2 q)_x = 0, \quad (1.3)$$

which, using a gauge transform, can be changed into the Kaup-Newell derivative NLS equation [59]

$$iq_t + \alpha_1 q_{xx} + i\epsilon (|q|^2 q)_x = 0. \quad (1.4)$$

When $\beta_1 : \beta_2 : \beta_3 = 0 : 1 : 0$, Eq. (1.1) reduce to the the modified Chen-Lee-Liu derivative NLS equation [60,61]

$$iq_t + \alpha_1 q_{xx} + \alpha_2 |q|^2 q + i\epsilon \beta_2 |q|^2 q_x = 0, \quad (1.5)$$

When $\alpha_1 : \alpha_2 = 1 : 2$, $\beta_1 : \beta_2 : \beta_3 = 1 : 6 : 3$, Eq. (1.1) reduce to the integrable Sasa-Satsuma equation [56]

$$iq_t + \alpha_1 (q_{xx} + 2|q|^2 q) + i\epsilon \beta_1 (q_{xxx} + 6|q|^2 q_x + 3q(|q|^2)_x) = 0, \quad (1.6)$$

When $\alpha_1 : \alpha_2 = 1 : 2$, $\beta_1 : \beta_2 : \beta_3 = 1 : 6 : 0$, Eq. (1.1) reduce to the Hirota equation [62]

$$iq_t + \alpha (q_{xx} + 2|q|^2 q) + i\beta (q_{xxx} + 6|q|^2 q_x) = 0, \quad (1.7)$$

where $q = q(x, t)$ denotes the complex envelope field, the real-valued parameters α, β are used to modulate the second- and third-order dispersion coefficients, respectively. As $\alpha \neq 0, \beta = 0$, Eq. (1.7) reduces to the NLS equation, and at $\alpha = 0, \beta \neq 0$, Eq. (1.7) reduces to the complex modified KdV (cmKdV) equation

$$q_t + \beta (q_{xxx} + 6|q|^2 q_x) = 0. \quad (1.8)$$

Notice that one can take $\beta = 1$ by using the scaling transform $t = \tau/\beta$.

The Hirota equation (1.7) and cmKdV equation (1.8) are the important ones of the above-mentioned higher-order NLS equations, and completely integrable, which can be derived from the AKNS hierarchy [3]. In fact, they can also be found from the reduction of the Maxwell equation (see [54])

$$\frac{\epsilon}{c^2} \frac{\partial^2 \mathbf{E}}{\partial t^2} + \nabla \times \nabla \times \mathbf{E} = 0, \quad (1.9)$$

where \mathbf{E} denotes the electric field intensity, c the speed of light, and ϵ a real-valued parameter.

In this paper, based on the idea of Refs. [23–25], we would like to study the large-space and long-time asymptotic behaviors of a soliton gas of the integrable Hirota equation (1.7) via the Riemann-Hilbert problems (see the following Riemann-Hilbert problem 2 in Sec. 2). The one-soliton solution of the Hirota equation (1.7) is

$$q(x, t) = k \operatorname{sech}[k(x - vt - x_0)] e^{i(\mu x + \omega t + p_0)} \quad (1.10)$$

with

$$v = \beta(k^2 - 3\mu^2) + \frac{8}{3}\alpha\mu + \frac{2\alpha^2\mu}{9\beta\mu - 3\alpha}, \quad \omega = k^2(\alpha - 3\beta\mu) + \mu^2(\beta\mu - \alpha),$$

where x_0 is the initial peak position of the soliton, p_0 the initial phase, v the soliton velocity, $-\omega/\mu$ phase velocity, and $3k^2\beta - 3\beta\mu^2 + 2\alpha\mu$ group velocity, and the Jacobian elliptic double-periodic wave solution of the Hirota equation (1.7) is

$$q(x, t) = k \operatorname{dn}[k(x - v_m t - x_0); m] e^{i(\mu x + \omega_m t + p_0)}, \quad m \in (0, 1) \quad (1.11)$$

with

$$v_m = \beta[k^2(2 - m^2) - 3\mu^2] + \frac{8}{3}\alpha\mu + \frac{2\alpha^2\mu}{9\beta\mu - 3\alpha}, \quad \omega_m = k^2(2 - m^2)(\alpha - 3\beta\mu) + \mu^2(\beta\mu - \alpha).$$

where the elliptic wave velocity is v_m , phase velocity $-\omega_m/\mu$, and group velocity $3(2 - m^2)k^2\beta - 3\beta\mu^2 + 2\alpha\mu$. When $m \rightarrow 1$, the periodic solution (1.11) reduces to the one-soliton solution (1.10).

The Hirota equation and its higher-order extensions have been showed to possess the multi-soliton solutions, breather solutions, and rogue waves [62–68]. Moreover, the long-time asymptotic behaviors of the focusing and defocusing Hirota equations have been studied [69–71]. Recently, Guo *et al* [72] studied nonlinear stability of multi-solitons for the Hirota equation.

It is well known that one can solve the N -soliton solution of the Hirota equation (1.7) by using the inverse scattering transform to study the associated Lax pair [3]

$$\begin{aligned} \frac{\partial \Psi}{\partial x} &= \begin{pmatrix} -iz & q(x, t) \\ -q^*(x, t) & iz \end{pmatrix} \Psi, \\ \frac{\partial \Psi}{\partial t} &= \begin{pmatrix} -i(4\beta z^3 + 2\alpha z^2) + i(2\beta z + \alpha)|q|^2 + 2i\beta \operatorname{Im}(qq_x^*) & 4\beta z^2 q + 2z(\alpha q - i\beta q_x) + T_{01} \\ -4\beta z^2 q^* - 2z(\alpha q^* - i\beta q_x^*) + T_{02} & i(4\beta z^3 + 2\alpha z^2) - i(2\beta z + \alpha)|q|^2 - 2i\beta \operatorname{Im}(qq_x^*) \end{pmatrix} \Psi, \end{aligned} \quad (1.12)$$

where $\Psi = \Psi(x, t; z)$ is a 2×2 matrix eigenfunction, $z \in \mathbb{C}$ a spectral parameter, the star denotes the complex conjugate, and

$$T_{01} = -i\alpha q - \beta(2|q|^2 q + q_{xx}), \quad T_{02} = i\alpha q^* + \beta(2|q|^2 q^* + q_{xx}^*).$$

Let the new matrix function $\Phi(x, t; z)$ be

$$\Phi(x, t; z) = \Psi(x, t; z) e^{i[zx + (4\beta z^3 + 2\alpha z^2)t]\sigma_3}. \quad (1.13)$$

Then based on the boundary-value condition of solitons $\lim_{|x| \rightarrow \infty} q(x, 0) = 0$, it follows from the Lax pair (1.12) that one know that [3]

$$\Psi(x, t; z) \sim e^{-i[zx + (4\beta z^3 + 2\alpha z^2)t]\sigma_3}, \quad \Phi(x, t; z) \sim \mathbb{I}, \quad |x| \rightarrow \infty. \quad (1.14)$$

and the Lax pair (1.12) can be rewritten as

$$\begin{aligned} \frac{\partial \Phi}{\partial x} + ik[\sigma_3, \Phi] &= \begin{pmatrix} 0 & q(x, t) \\ -q^*(x, t) & 0 \end{pmatrix} \Phi, \\ \frac{\partial \Phi}{\partial t} + i(4\beta z^3 + 2\alpha z^2)[\sigma_3, \Phi] &= \begin{pmatrix} i(2\beta z + \alpha)|q|^2 + 2i\beta \operatorname{Im}(qq_x^*) & 4\beta z^2 q + 2z(\alpha q - i\beta q_x) + T_{01} \\ -4\beta z^2 q^* - 2z(\alpha q^* - i\beta q_x^*) + T_{02} & -i(2\beta z + \alpha)|q|^2 - 2i\beta \operatorname{Im}(qq_x^*) \end{pmatrix} \Phi, \end{aligned} \quad (1.15)$$

which can be written as fully differential form such that the Jost solutions $\Phi_+(x, t; z)$ and $\Phi_-(x, t; z)$ can be written as follows:

$$\Phi_{\pm}(x, t; z) = \mathbb{I} - \int_x^{\pm\infty} e^{iz(y-x)\sigma_3} \begin{pmatrix} 0 & q(y, t) \\ -q^*(y, t) & 0 \end{pmatrix} \Phi_{\pm}(y, t; z) e^{-iz(y-x)\sigma_3} dy, \quad (1.16)$$

with $\Phi_{\pm}(x, t; z) = \mathbb{I} + \mathcal{O}(1/z)$, $z \rightarrow \infty$, where σ_3 is the third Pauli matrix.

By using Abel lemma and the zero traces of coefficient matrices of the Lax pair (1.12), one knows that $\det \Psi_{\pm}(k, x, t)$ are independent of variable x and $\det \Psi_{\pm} = 1$. Furthermore, $\Psi_{\pm} = \Phi_{\pm} e^{-i[zx + (4\beta z^3 + 2\alpha z^2)t]\sigma_3}$ are linearly dependent to lead to

$$\Psi_-(x, t; z) = \Psi_+(x, t; z)S(k), \quad \det(S) = 1, \quad (1.17)$$

or

$$\Phi_-(x, t; z) = \Phi_+(x, t; z) e^{-i[zx + (4\beta z^3 + 2\alpha z^2)t]\widehat{\sigma}_3} S(k), \quad \det(S) = 1, \quad (1.18)$$

where $S(z) = (s_{ij}(z))_{2 \times 2}$ is a 2×2 scattering matrix.

$$\begin{aligned} s_{jj}(z) &= (-1)^{j+1} |\Psi_{-j}(x, t; z), \Psi_{+(3-j)}(x, t; z)| = (-1)^{j+1} |\Phi_{-j}(x, t; z), \Phi_{+(3-j)}(x, t; z)|, \quad j = 1, 2, \\ s_{(3-j)j}(z) &= (-1)^j |\Psi_{-j}(x, t; z), \Psi_{+j}(x, t; z)| = (-1)^j |\Phi_{-j}(x, t; z), \Phi_{+j}(x, t; z)|, \quad j = 1, 2, \end{aligned} \quad (1.19)$$

where $\Psi_{\pm} = (\Psi_{\pm 1}(x, t; z), \Psi_{\pm 2}(x, t; z))$, $\Phi_{\pm} = (\Phi_{\pm 1}(x, t; z), \Phi_{\pm 2}(x, t; z))$, where $\Psi_{\pm 1}$ ($\Phi_{\pm 1}$) and $\Psi_{\pm 2}$ ($\Phi_{\pm 2}$) represent two columns of Ψ_{\pm} (Φ_{\pm}), respectively. There are the symmetry: $\Psi_{\pm}(z) = \sigma_2 \Psi_{\pm}^*(z^*) \sigma_2$ and $S(z) = \sigma_2 S^*(z^*) \sigma_2$, where σ_2 is the second Pauli matrix. It follows from Eq. (1.16) that these functions $\Psi_{-1}, \Psi_{+2}, \Phi_{-1}, \Phi_{+2}, s_{11}(z)$ ($\Psi_{+1}, \Psi_{-2}, \Phi_{+1}, \Phi_{-2}, s_{22}(z)$) can be analytically extended to \mathbb{C}_+ (\mathbb{C}_-) and continuous to continuously to $\mathbb{C}_+ \cup \mathbb{R}$ ($\mathbb{C}_- \cup \mathbb{R}$). However another two scattering coefficients $s_{12}(z)$, $s_{21}(z)$ can not be analytically continued away from \mathbb{R} . Thus the reflection coefficient is defined as $\rho(z) = s_{21}(z)/s_{11}(z)$.

The discrete spectra of the Hirota equation are the set of all values $z \in \mathbb{C} \setminus \mathbb{R}$ such that they possess eigenfunctions in $L^2(\mathbb{R})$, and satisfy $s_{11}(z_n) = 0$ for $Z_+ = \{z_n | z_n \in \mathbb{C}_+\}$ and $s_{22}(z_n^*) = 0$ for $Z_- = \{z_n^* | z_n^* \in \mathbb{C}_-\}$. Let the sectionally meromorphic matrix $M^{(0)}(z) = M^{(0)}(x, t; z)$ be

$$M^{(0)}(z) = \begin{cases} \begin{pmatrix} \frac{\Phi_{-1}(x, t; z)}{s_{11}(z)}, & \Phi_{+2}(x, t; z) \\ \Phi_{+1}(x, t; z), & \frac{\Phi_{-2}(x, t; z)}{s_{22}(z)} \end{pmatrix} = \begin{pmatrix} \frac{\Psi_{-1}(x, t; z)}{s_{11}(z)}, & \Psi_{+2}(x, t; z) \\ \Psi_{+1}(x, t; z), & \frac{\Psi_{-2}(x, t; z)}{s_{22}(z)} \end{pmatrix} e^{i\theta(x, t; z)\sigma_3}, & z \in \mathbb{C}_+, \\ \begin{pmatrix} \frac{\Phi_{-1}(x, t; z)}{s_{11}(z)}, & \Phi_{+2}(x, t; z) \\ \Phi_{+1}(x, t; z), & \frac{\Phi_{-2}(x, t; z)}{s_{22}(z)} \end{pmatrix} = \begin{pmatrix} \frac{\Psi_{-1}(x, t; z)}{s_{11}(z)}, & \Psi_{+2}(x, t; z) \\ \Psi_{+1}(x, t; z), & \frac{\Psi_{-2}(x, t; z)}{s_{22}(z)} \end{pmatrix} e^{i\theta(x, t; z)\sigma_3}, & z \in \mathbb{C}_-. \end{cases} \quad (1.20)$$

where the phase function is

$$\theta(x, t; z) = zx + (4\beta z^3 + 2\alpha z^2)t. \quad (1.21)$$

Then based on the scattering relation given by Eq. (1.17) or (1.18), one can introduce the following Riemann-Hilbert problem (RHP) [3, 47, 68]:

RHP 1. Find a 2×2 matrix function $M^{(0)}(x, t; z)$ that satisfies:

- *Analyticity:* $M^{(0)}(x, t; z)$ is meromorphic in $\{z | z \in \mathbb{C} \setminus \mathbb{R}\}$ and takes continuous boundary values on \mathbb{R} ;
- *Jump condition:* The boundary values on the jump contour \mathbb{R} are defined as

$$M_+^{(0)}(x, t; z) = M_-^{(0)}(x, t; z) V_0(x, t; z), \quad z \in \mathbb{R}, \quad (1.22)$$

where the jump matrix is defined as

$$V_0(x, t; z) = \begin{bmatrix} 1 + |\rho(z)|^2 & \rho^*(z) e^{-2i\theta(x, t; z)} \\ \rho(z) e^{2i\theta(x, t; z)} & 1 \end{bmatrix},$$

- *Normalization:* $M^{(0)}(x, t; z) \rightarrow \mathbb{I}, \quad z \rightarrow \infty.$
- *Residue conditions:* $M^{(0)}(x, t; z)$ has simple poles at each discrete spectrum in $Z_+ = \{z_n | z_n \in \mathbb{C}_+\} \cup Z_- = \{z_n^* | z_n^* \in \mathbb{C}_-\}$ with:

$$\begin{aligned} \operatorname{Res}_{z=z_n} M^{(0)}(x, t; z) &= \lim_{z \rightarrow z_n} M^{(0)}(x, t; z) \begin{bmatrix} 0 & 0 \\ c_n e^{2iz_n x + 4i(\alpha z_n^2 + 2\beta z_n^3)t} & 0 \end{bmatrix}, \\ \operatorname{Res}_{z=z_n^*} M^{(0)}(x, t; z) &= \lim_{z \rightarrow z_n^*} M^{(0)}(x, t; z) \begin{bmatrix} 0 & -c_n^* e^{-2iz_n^* x - 4i(\alpha z_n^{*2} + 2\beta z_n^{*3})t} \\ 0 & 0 \end{bmatrix}. \end{aligned} \quad (1.23)$$

where the norming constants $c_n(z_n), c_n^*(z_n^*)$ satisfy

$$c_n(z_n) = \frac{\hat{c}_n(z_n)}{s'_{11}(z_n)}, \quad c_n^*(z_n^*) = \frac{\hat{c}_n^*(z_n^*)}{s'_{22}(z_n^*)} \quad (1.24)$$

with $\hat{c}_n(z_n), \hat{c}_n^*(z_n^*)$ be defined by

$$\Psi_{-1}(x, t; z_n) = \hat{c}_n(z_n) \Psi_{+2}(x, t; z_n), \quad \Psi_{-2}(x, t; z_n) = -\hat{c}_n^*(z_n^*) \Psi_{+1}(x, t; z_n^*), \quad (1.25)$$

Therefore, based on the Plemelj's formulae, one can has

$$\begin{aligned} M^{(0)}(x, t; z) &= \mathbb{I} + \sum_{n=1}^N \left[\frac{\operatorname{Res}_{z=z_n} M^{(0)}(x, t; z)}{z - z_n} + \frac{\operatorname{Res}_{z=z_n^*} M^{(0)}(x, t; z)}{z - z_n^*} \right] \\ &\quad + \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{M_-^{(0)}(x, t; \zeta) (\mathbb{I} - V_0(x, t; \zeta))}{\zeta - z} d\zeta, \quad z \in \mathbb{C} \setminus \mathbb{R}, \end{aligned} \quad (1.26)$$

such that the solution $q(x, t)$ of the Hirota equation is given by using the solution $M^{(0)}(x, t; z)$ given by Eq. (1.26) of the RHP-1

$$\begin{aligned} q(x, t) &= 2i \lim_{z \rightarrow \infty} z (M^{(0)}(x, t; z))_{12} \\ &= 2i \left(\operatorname{Res}_{z=z_n^*} M^{(0)}(x, t; z) \right)_{12} - \frac{1}{\pi} \int_{\mathbb{R}} \left(M_-^{(0)}(x, t; \zeta) (\mathbb{I} - V_0(x, t; \zeta)) \right)_{12} d\zeta. \end{aligned} \quad (1.27)$$

In this paper, we will use the idea of Refs. [23–25] to analyze the asymptotic behaviors of a soliton gas for the Hirota equation. The rest of this paper is arranged as follows. In Sec. 2, based on the Riemann-Hilbert problem 1, we present the Riemann-Hilber problem related to the soliton gas of the Hirota equation, which is regarded as the limit $N \rightarrow \infty$ of its N -soliton solution, where the discrete spectra are chosen as the pure imaginary numbers within the intervals $(ia, ib) \cup (-ib, -ia)$. In Sec. 3, we present the large-space asymptotics of the soliton gas of the potential $q(x, 0)$ of the Hirota equation as $x \rightarrow -\infty$. In Sec. 4, we give the corresponding large-space asymptotics of the soliton gas of the potential $q(x, t_0)$ of the Hirota equation for the fixed $t = t_0$ as $x \rightarrow -\infty$. In Sec. 5, we give the long-time asymptotics of the soliton gas of the potential $q(x, t)$ of the complex mKdV equation ($\alpha = 0, \beta = 1$) as $t \rightarrow +\infty$ at three fundamental spatial domains, $\xi = x/t > 4\beta b^2, \xi_c < \xi = x/t < 4\beta b^2, \xi = x/t < \xi_c$. In Sec. 6, we study the property of the soliton gas with the discrete spectra filling uniformly a quadrature domain. Finally, we give some conclusions and discussions in Sec. 7.

2 Soliton gas: the limit $N \rightarrow \infty$ of N -soliton solution via RHP

In this section, we first of all construct the pure-soliton Riemann-Hilbert problem corresponding to the N -soliton solution with reflectionless potential (i.e., $\rho(z) = 0$) and pure imaginary discrete spectra set $Z_N := \{iz_j, -iz_j\}_{j=1}^N$ with $z_j \in \mathbb{R}_+$ from the RHP 1.

RHP 2. Find a 2×2 matrix function $M(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $M(x, t; z)$ is analytic in $\mathbb{C} \setminus Z_N$, and has simple poles at $z \in Z_N$.
- *Normalization:* $M(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$;
- *Residue conditions:* $M(x, t; z)$ has simple poles at each point in Z_N with:

$$\begin{aligned} \text{Res}_{z=iz_j} M(x, t; z) &= \lim_{z \rightarrow iz_j} M(x, t; z) \begin{bmatrix} 0 & 0 \\ c_j e^{-2z_j x + (8\beta z_j^3 - 4i\alpha z_j^2)t} & 0 \end{bmatrix}, \\ \text{Res}_{z=-iz_j} M(x, t; z) &= \lim_{z \rightarrow -iz_j} M(x, t; z) \begin{bmatrix} 0 & -c_j^* e^{-2z_j x + (8\beta z_j^3 + 4i\alpha z_j^2)t} \\ 0 & 0 \end{bmatrix}. \end{aligned} \quad (2.28)$$

The matrix function solution of the RHP 2 can be written as

$$M(x, t; z) = \mathbb{I} + \sum_{j=1}^N \left(\frac{1}{z - z_j} \begin{bmatrix} u_j(x, t) & 0 \\ v_j(x, t) & 0 \end{bmatrix} + \frac{1}{z - z_j^*} \begin{bmatrix} 0 & -v_j^*(x, t) \\ 0 & u_j(x, t) \end{bmatrix} \right), \quad (2.29)$$

where $u_j(x, t)$, $v_j(x, t)$ are unknown functions to be determined later. Substituting Eq. (2.29) into the first one of Eq. (2.28) yields the system of equations about unknowns u_j , v_j^*

$$\begin{aligned} u_j &= - \sum_{s=1}^N \frac{c_s e^{-2z_s x + (8\beta z_s^3 - 4i\alpha z_s^2)t}}{z_j - z_s^*} v_s^*, \quad j = 1, 2, \dots, N, \\ v_j^* &= \left(1 + \sum_{s=1}^N \frac{c_s e^{-2z_s x + (8\beta z_s^3 - 4i\alpha z_s^2)t}}{z_j^* - z_s} u_j \right) c_j^* e^{-2z_s x + (8\beta z_s^3 + 4i\alpha z_s^2)t}, \quad j = 1, 2, \dots, N. \end{aligned} \quad (2.30)$$

Let $\hat{u} = (\hat{u}_1, \hat{u}_2, \dots, \hat{u}_N)^T$, $\hat{v}^* = (\hat{v}_1^*, \hat{v}_2^*, \dots, \hat{v}_N^*)^T$ and $w = (w_1, w_2, \dots, w_N)^T$, $Q = (Q_{js})_{N \times N}$ with

$$\begin{aligned} \hat{u}_j &= \frac{u_j}{\sqrt{c_j}} e^{z_j x - (4\beta z_j^3 - 2i\alpha z_j^2)t}, \quad \hat{v}_j^* = \frac{v_j^*}{\sqrt{c_j^*}} e^{z_j x - (4\beta z_j^3 + 2i\alpha z_j^2)t}, \\ w_j &= \sqrt{c_j^*} e^{-z_j x + (4\beta z_j^3 + 2i\alpha z_j^2)t}, \quad Q_{js} = \frac{i}{z_s - z_j^*} e^{-(z_j + z_s)x + [4\beta(z_j^3 + z_s^3) + 2i\alpha(z_j^2 - z_s^2)]t}. \end{aligned} \quad (2.31)$$

Then system (2.30) can be rewritten as

$$\begin{bmatrix} \mathbb{I}_N & -iQ^* \\ -iQ & \mathbb{I}_N \end{bmatrix} \begin{bmatrix} \hat{u} \\ \hat{v}^* \end{bmatrix} = \begin{bmatrix} 0 \\ w \end{bmatrix}, \quad (2.32)$$

whose solution can be derived in the form (Note that $\det(\mathbb{I}_N + QQ^*) \neq 0$)

$$\begin{bmatrix} \hat{u} \\ \hat{v}^* \end{bmatrix} = \begin{bmatrix} \mathbb{I}_N - Q^*(\mathbb{I}_N + QQ^*)^{-1}Q & iQ^*(\mathbb{I}_N + QQ^*)^{-1} \\ i(\mathbb{I}_N + QQ^*)^{-1}Q & (\mathbb{I}_N + QQ^*)^{-1} \end{bmatrix} \begin{bmatrix} 0 \\ w \end{bmatrix}. \quad (2.33)$$

According to Eqs. (1.27), (2.29), (2.31), and (2.33), we can recover $q(x, t)$ by the following formula of the N -soliton solution of the Hirota equation (1.7):

$$\begin{aligned} q_N(x, t) &= 2i \lim_{z \rightarrow \infty} (zM(x, t; z))_{12}, \\ &= -2i \sum_{j=1}^N v_j^* \\ &= -2i \sum_{j,s=1}^N ((\mathbb{I}_N + QQ^*)^{-1})_{js} \sqrt{c_j^* c_s^*} e^{-(z_j + z_s)x + [4\beta(z_j^3 + z_s^3) + 2i\alpha(z_j^2 + z_s^2)]t}. \end{aligned} \quad (2.34)$$




Figure 1: Analytical domains of eigenfunctions and distribution of the discrete spectrum Z_N .

Recently, Girotti *et al* [23, 24] investigated the soliton gas phenomena of KdV and mKdV equations with discrete spectra located on the imaginary axis. Similarly, we assume that the discrete spectrum belongs to the interval $(ia, ib) \cup (-ib, -ia)$. We mainly focus on the situation where the limit $N \rightarrow \infty$ of $q_N(x, t)$, under the following conditions [23]:

- The simple poles $\{iz_j\}_{j=1}^N$ are sampled from a smooth density function $p(z)$ so that $\int_a^{z_j} p(\lambda) d\lambda = \frac{j}{N}, j = 1, \dots, N$ (see Fig. 1).
- The norming parameters $\{c_j\}_{j=1}^N$ satisfy

$$c_j = \frac{i(b-a)r(iz_j)}{N\pi}, \quad (2.35)$$

where $r(z)$ is a real-valued, continuous and non-vanishing function for $z \in (ia, ib)$ ($0 < a < b$), with the symmetry $r(z^*) = r(z)$.

Then the residue conditions (2.28) of matrix function $M(x, t; z)$ can be rewrite as:

$$\begin{aligned} \text{Res}_{z=iz_j} M(x, t; z) &= \lim_{z \rightarrow iz_j} M(x, t; z) \begin{bmatrix} 0 & 0 \\ \frac{i(b-a)r(iz_j)}{N\pi} e^{-2z_j x + (8\beta z_j^3 - 4i\alpha z_j^2)t} & 0 \end{bmatrix}, \\ \text{Res}_{z=-iz_j} M(x, t; z) &= \lim_{z \rightarrow -iz_j} M(x, t; z) \begin{bmatrix} 0 & \frac{i(b-a)r(iz_j)}{N\pi} e^{-2z_j x + (8\beta z_j^3 + 4i\alpha z_j^2)t} \\ 0 & 0 \end{bmatrix}. \end{aligned} \quad (2.36)$$

Notice that for the fixed $t = t_0$, as $x \rightarrow +\infty$ the residue conditions (2.36) only have the exponentially small terms such that the potential q is exponentially small by a small norm argument. However, as $x \rightarrow -\infty$ both terms of the residue conditions (2.36) are exponentially large. For this case, one may make the transform

$$\widehat{M}(z) = M(z)T(z), \quad T(z) = \prod_{j=1}^N \left(\frac{z - iz_j}{z + iz_j} \right)^{\sigma_3} \quad (2.37)$$

such that the residue conditions of $\widehat{M}(z)$ are

$$\begin{aligned} \text{Res}_{z=iz_j} \widehat{M}(x, t; z) &= \lim_{z \rightarrow iz_j} M(x, t; z) \begin{bmatrix} 0 & \frac{N\pi}{i(b-a)r(iz_j)T'^2(iz_j)} e^{2z_j x - (8\beta z_j^3 - 4i\alpha z_j^2)t} \\ 0 & 0 \end{bmatrix}, \\ \text{Res}_{z=-iz_j} \widehat{M}(x, t; z) &= \lim_{z \rightarrow -iz_j} M(x, t; z) \begin{bmatrix} 0 & 0 \\ \frac{N\pi T^4(-iz_j)}{i(b-a)r(iz_j)T'^2(-iz_j)} e^{2z_j x - (8\beta z_j^3 + 4i\alpha z_j^2)t} & 0 \end{bmatrix}, \end{aligned} \quad (2.38)$$

in which the moduli of coefficients of two exponential functions are less than $e^{c_1 N + c_2}$ for some positive constants $c_{1,2} > 0$ such that one knows that for the fixed $t = t_0$, these two non-zero terms are exponentially small as $x \rightarrow -\infty$ with $x < (c_1 N + c_2)/(c_3 - 2\min\{z_j\})$ for some positive constant c_3 ($0 < c_3 < 2z_j$). The corresponding potential is

$$q_N(x, t) = 2i \lim_{z \rightarrow \infty} (z \widehat{M}(x, t; z))_{12}, \quad (2.39)$$

Let a closed curve Γ_+ (Γ_-) be a small radius encircling the poles $\{iz_j\}_{j=1}^N$ ($\{-iz_j\}_{j=1}^N$) counterclockwise (clockwise) in the upper (lower) half plane \mathbb{C}_+ (\mathbb{C}_-). Make the following transformation:

$$M^{(1)}(x, t; z) = \begin{cases} M(x, t; z) \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^N \frac{i(b-a)r(iz_j)}{N\pi(z-iz_j)} e^{2i(z_j x + 2\alpha z_j^2 t + 4\beta z_j^3 t)} & 1 \end{bmatrix}, & z \text{ within } \Gamma_+, \\ M(x, t; z) \begin{bmatrix} 1 & -\sum_{j=1}^N \frac{i(b-a)r(iz_j)}{N\pi(z+iz_j)} e^{-2i(z_j x + 2\alpha z_j^2 t + 4\beta z_j^3 t)} \\ 0 & 1 \end{bmatrix}, & z \text{ within } \Gamma_-, \\ M(x, t; z), & \text{otherwise.} \end{cases} \quad (2.40)$$

Then the matrix function $M^{(1)}(x, t; z)$ satisfies the following Riemann-Hilbert problem.

RHP 3. Find a 2×2 matrix function $M^{(1)}(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $M^{(1)}(x, t; z)$ is analytic in $\mathbb{C} \setminus (\Gamma_+ \cup \Gamma_-)$ and takes continuous boundary values on $\Gamma_+ \cup \Gamma_-$.
- *Jump condition:* The boundary values on the jump contour $\Gamma_+ \cup \Gamma_-$ are defined as

$$M_+^{(1)}(x, t; z) = M_-^{(1)}(x, t; z) V_1(x, t; z), \quad z \in \Gamma_+ \cup \Gamma_-, \quad (2.41)$$

where

$$V_1(x, t; z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^N \frac{i(b-a)r(iz_j)}{N\pi(z-iz_j)} e^{2i(z_j x + 2\alpha z_j^2 t + 4\beta z_j^3 t)} & 1 \end{bmatrix}, & z \in \Gamma_+, \\ \begin{bmatrix} 1 & \sum_{j=1}^N \frac{i(b-a)r(iz_j)}{N\pi(z+iz_j)} e^{-2i(z_j x + 2\alpha z_j^2 t + 4\beta z_j^3 t)} \\ 0 & 1 \end{bmatrix}, & z \in \Gamma_-; \end{cases} \quad (2.42)$$

- *Normalization:* $M^{(1)}(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

According to Eq. (2.34), we can recover $q(x, t)$ by the following formula:

$$q(x, t) = 2i \lim_{z \rightarrow \infty} (zM^{(1)}(x, t; z))_{12}. \quad (2.43)$$

For convenience, we can assume that the N poles are equally spaced along (ia, ib) with distance between two poles equal to $s_d = \frac{b-a}{N}$.

Based on Ref. [23], one has the following lemma:

Lemma 2.1. *For any open set A_+ (A_-) containing the interval $[ia, ib]$ ($[-ib, -ia]$), the following identities hold:*

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j e^{2i(z_j x + 2\alpha z_j^2 t + 4\beta z_j^3 t)}}{z \mp iz_j} = \pm \frac{1}{\pi} \int_{\pm ia}^{\pm ib} \frac{r(k) e^{2i(kx + 2\alpha k^2 t + 4\beta k^3 t)}}{z - k} dk, \quad z \in \mathbb{C} \setminus A_{\pm}, \quad (2.44)$$

where the open intervals (ia, ib) and $(-ib, -ia)$ are both oriented upwards.

Proof. For $z \in \mathbb{C} \setminus A_+$, since $c_j = \frac{i(b-a)r(iz_j)}{N\pi}$, one has

$$\begin{aligned} \lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j e^{2i(z_j x + 2\alpha z_j^2 t + 4\beta z_j^3 t)}}{z - iz_j} &= \lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{r(iz_j)(ib - ia)}{N\pi(z - iz_j)} e^{2i(z_j x + 2\alpha z_j^2 t + 4\beta z_j^3 t)} \\ &= \lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{ir(iz_j)s_d}{\pi(z - iz_j)} e^{2i(z_j x + 2\alpha z_j^2 t + 4\beta z_j^3 t)} \\ &= \frac{1}{\pi} \int_{ia}^{ib} \frac{r(k)}{z - k} e^{2i(kx + 2\alpha k^2 t + 4\beta k^3 t)} dk. \end{aligned}$$

Similarly, it can be proven that another identity hold $z \in \mathbb{C} \setminus A_-$. Thus the proof is completed. \square

When $N \rightarrow \infty$, according to Lemma 2.1, the jump matrix $V_1(x, t; z)$ defined by Eq. (2.42) can be rewrite as:

$$V_1(x, t; z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ -\frac{1}{\pi} \int_{ia}^{ib} \frac{r(k)}{z - k} e^{2i(kx + 2\alpha k^2 t + 4\beta k^3 t)} dk & 1 \end{bmatrix}, & z \in \Gamma_+, \\ \begin{bmatrix} 1 & \frac{1}{\pi} \int_{-ib}^{-ia} \frac{r(k)}{z - k} e^{-2i(kx + 2\alpha k^2 t + 4\beta k^3 t)} dk \\ 0 & 1 \end{bmatrix}, & z \in \Gamma_-; \end{cases} \quad (2.45)$$

Make the following transformation:

$$M^{(2)}(x, t; z) = \begin{cases} M^{(1)}(x, t; z) \begin{bmatrix} 1 & 0 \\ \frac{1}{\pi} \int_{ia}^{ib} \frac{r(k)}{z - k} e^{2i(kx + 2\alpha k^2 t + 4\beta k^3 t)} dk & 1 \end{bmatrix}, & z \text{ within } \Gamma_+, \\ M^{(1)}(x, t; z) \begin{bmatrix} 1 & \frac{1}{\pi} \int_{-ib}^{-ia} \frac{r(k)}{z - k} e^{2i(kx + 2\alpha k^2 t + 4\beta k^3 t)} dk \\ 0 & 1 \end{bmatrix}, & z \text{ within } \Gamma_-, \\ M^{(1)}(x, t; z), & \text{otherwise.} \end{cases} \quad (2.46)$$

For convenience, let $\gamma_1 := (a, b)$, $\gamma_2 := (-b, -a)$ and both line segments γ_1 and γ_2 are directed to the right. Then matrix function $M^{(2)}(x, t; z)$ satisfies the following Riemann-Hilbert problem.

RHP 4. Find a 2×2 matrix function $M^{(2)}(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $M^{(2)}(x, t; z)$ is analytic in $\mathbb{C} \setminus (i\gamma_1 \cup i\gamma_2)$ and takes continuous boundary values on $i\gamma_1 \cup i\gamma_2$.
- *Jump condition:* The boundary values on the jump contour $i\gamma_1 \cup i\gamma_2$ are defined as

$$M_+^{(2)}(x, t; z) = M_-^{(2)}(x, t; z)V_2(x, t; z), \quad z \in i\gamma_1 \cup i\gamma_2, \quad (2.47)$$

where

$$V_2(x, t; z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ 2ir(z)e^{2i(xz+2\alpha tz^2+4\beta tz^3)} & 1 \end{bmatrix}, & z \in i\gamma_1, \\ \begin{bmatrix} 1 & 2ir(z)e^{-2i(xz+2\alpha tz^2+4\beta tz^3)} \\ 0 & 1 \end{bmatrix}, & z \in i\gamma_2; \end{cases} \quad (2.48)$$

- *Normalization:* $M^{(2)}(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

According to Eq. (2.43), we can recover $q(x, t)$ by the following formula:

$$q(x, t) = 2i \lim_{z \rightarrow \infty} (zM^{(2)}(x, t; z))_{12}. \quad (2.49)$$

We want to rotate the jump line to the real axis, so we will rewrite Eq. (2.47) as follows:

$$M_+^{(2)}(x, t; iz) = M_-^{(2)}(x, t; iz) \begin{cases} \begin{bmatrix} 1 & 0 \\ 2ir(iz)e^{-2zx-4i\alpha z^2t+8\beta z^3t} & 1 \end{bmatrix}, & z \in \gamma_1, \\ \begin{bmatrix} 1 & 2ir(iz)e^{2zx+4i\alpha z^2t-8\beta z^3t} \\ 0 & 1 \end{bmatrix}, & z \in \gamma_2. \end{cases} \quad (2.50)$$

Make the following transformation:

$$M^{(3)}(x, t; z) = M^{(2)}(x, t; iz), \quad R(z) = 2r(iz), \quad (2.51)$$

Then matrix function $M^{(3)}(x, t; z)$ satisfies the following Riemann-Hilbert problem:

RHP 5. Find a 2×2 matrix function $M^{(3)}(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $M^{(3)}(x, t; z)$ is analytic in $\mathbb{C} \setminus (\gamma_1 \cup \gamma_2)$ and takes continuous boundary values on $\gamma_1 \cup \gamma_2$.
- *Jump condition:* The boundary values on the jump contour $\gamma_1 \cup \gamma_2$ are defined as

$$M_+^{(3)}(x, t; z) = M_-^{(3)}(x, t; z)V_3(x, t; z), \quad z \in \gamma_1 \cup \gamma_2, \quad (2.52)$$

where

$$V_3(x, t; z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ iR(z)e^{-2zx-4i\alpha z^2t+8\beta z^3t} & 1 \end{bmatrix}, & z \in \gamma_1, \\ \begin{bmatrix} 1 & iR(z)e^{2zx+4i\alpha z^2t-8\beta z^3t} \\ 0 & 1 \end{bmatrix}, & z \in \gamma_2. \end{cases} \quad (2.53)$$

- *Normalization:* $M^{(3)}(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

According to Eq. (2.49), we can recover $q(x, t)$ by the following formula:

$$q(x, t) = -2 \lim_{z \rightarrow \infty} (zM^{(3)}(x, t; z))_{12}. \quad (2.54)$$

3 Large-space asymptotics of the potential $q(x, 0)$ as $x \rightarrow -\infty$

Recently, Girotti *et al* [23] studied the asymptotic behavior of the soliton gas of KdV equation when $t = 0$. Similarly, we will study the asymptotic behavior of soliton gas $q(x) := q(x, 0)$ of the Hirota equation at $t = 0$. Let $Y(x; z) = M^{(3)}(x, 0; z)$. Then we construct the following Riemann-Hilbert problem:

RHP 6. Find a 2×2 matrix function $Y(x; z)$ that satisfies the following properties:

- *Analyticity:* $Y(x; z)$ is analytic in $\mathbb{C} \setminus (\gamma_1 \cup \gamma_2)$ and takes continuous boundary values on $\gamma_1 \cup \gamma_2$.
- *Jump condition:* The boundary values on the jump contour $\gamma_1 \cup \gamma_2$ are defined as

$$Y_+(x; z) = Y_-(x; z)V_4(x; z), \quad z \in \gamma_1 \cup \gamma_2, \quad (3.55)$$

where

$$V_4(x; z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ iR(z)e^{-2zx} & 1 \end{bmatrix}, & z \in \gamma_1, \\ \begin{bmatrix} 1 & iR(z)e^{2zx} \\ 0 & 1 \end{bmatrix}, & z \in \gamma_2; \end{cases} \quad (3.56)$$

- *Normalization:* $Y(x; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

According to Eq. (2.54), we can recover $q(x)$ by the following formula:

$$q(x) = -2 \lim_{z \rightarrow \infty} (zY(x; z))_{12}. \quad (3.57)$$

To solve the Riemann-Hilbert problem 6, we make the following transformation:

$$Y_1(x; z) = Y(x; z)[f(z)e^{xg(z)}]^{\sigma_3}, \quad (3.58)$$

where $f(z)$ and $g(z)$ are scalar functions which satisfy the following properties [23]:

- *Analyticity:* $f(z), g(z)$ are analytic in $\mathbb{C} \setminus (-b, b)$ and takes continuous boundary values on $(-b, b)$.
- *Jump condition:* The boundary values of $f(z), g(z)$ on the jump contour are defined as

$$\begin{aligned} f_+(z)f_-(z) &= \begin{cases} R^{-1}(z), & z \in \gamma_1, \\ R(z), & z \in \gamma_2, \end{cases} \\ f_+(z) &= f_-(z)e^{m_1}, \quad z \in [-a, a], \end{aligned} \quad (3.59)$$

and

$$\begin{aligned} g_+(z) + g_-(z) &= 2z, \quad z \in \gamma_1 \cup \gamma_2, \\ g_+(z) - g_-(z) &= m_2, \quad z \in [-a, a], \end{aligned} \quad (3.60)$$

where where

$$m_1 = -\frac{b}{K_1(m)} \int_a^b \frac{\ln R(\lambda)}{C_+(\lambda)} d\lambda, \quad C(z) = \sqrt{(z^2 - a^2)(z^2 - b^2)}, \quad m_2 = -\frac{i\pi b}{K_1(m)},$$

and the first and second kinds of complete elliptic integrals are denoted as:

$$K_1(m) := \int_0^1 \frac{ds}{\sqrt{(1-s^2)(1-m^2s^2)}}, \quad K_2(m) := \int_0^1 \frac{\sqrt{1-m^2s^2}}{\sqrt{1-t^2}} ds. \quad (3.61)$$

with modulus $m = \frac{a}{b}$.




Figure 2: The Riemann-Hilbert problem 8 for matrix function $Y_2(x; z)$. Opening lenses O_1 and O_2 .

- Normalization: $f(z) = 1 + \mathcal{O}(z^{-1})$, $g(z) = \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

The scalar functions $f(z)$ and $g(z)$ can also be accurately solved as [23, 24]:

$$f(z) = \exp \left[\frac{C(z)}{2\pi i} \left(\int_{\gamma_2^- \gamma_1} \frac{\ln R(\lambda)}{C_+(\lambda)(\lambda - z)} d\lambda + \int_{-a}^a \frac{m_1}{C(\lambda)(\lambda - z)} d\lambda \right) \right], \quad (3.62)$$

$$g(z) = z - \int_b^z \frac{\lambda^2 + b^2(K_2(m)/K_1(m) - 1)}{C(\lambda)} d\lambda.$$

Then matrix function $Y_1(x; z)$ satisfies the following Riemann-Hilbert problem.

RHP 7. Find a 2×2 matrix function $Y_1(x; z)$ that satisfies the following properties:

- Analyticity: $Y_1(x; z)$ is analytic in $\mathbb{C} \setminus (-b, b)$ and takes continuous boundary values on $(-b, b)$.
- Jump condition: The boundary values on the jump contour are defined as

$$Y_{1+}(x; z) = Y_{1-}(x; z)V_5(x; z), \quad z \in \gamma_1 \cup [-a, a] \cup \gamma_2, \quad (3.63)$$

where

$$V_5(x; z) = \begin{cases} \begin{bmatrix} \delta f(z)e^{x\Delta g(z)} & 0 \\ i & \delta^{-1}f(z)e^{-x\Delta g(z)} \end{bmatrix}, & z \in \gamma_1, \\ \begin{bmatrix} e^{xm_2+m_1} & 0 \\ 0 & e^{-(xm_2+m_1)} \end{bmatrix}, & z \in [-a, a], \\ \begin{bmatrix} \delta f(z)e^{x\Delta g(z)} & i \\ 0 & \delta^{-1}f(z)e^{-x\Delta g(z)} \end{bmatrix}, & z \in \gamma_2; \end{cases} \quad (3.64)$$

with $\delta f(z) = f_+(z)/f_-(z)$, $\Delta g(z) = g_+(z) - g_-(z)$.

- Normalization: $Y_1(x; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

3.1 Opening lenses: jump matrix decompositions

Based on the idea of Ref. [23], let us open the lens O_1 pass through points $z = a$ and $z = b$ and open the lens O_2 pass through points $z = -b$ and $z = -a$ (see Fig. 2). We define a new function as follows:

$$R_{\pm}(z) := \pm R(z), \quad z \in \gamma_1 \cup \gamma_2. \quad (3.65)$$

Note that the jump matrix $V_5(x; z)|_{\gamma_1}$ has the following decomposition:

$$\begin{aligned} V_5(x; z) &= \begin{bmatrix} \delta f(z)e^{x\Delta g(z)} & 0 \\ i & \delta^{-1}f(z)e^{-x\Delta g(z)} \end{bmatrix} \\ &= \begin{bmatrix} 1 & \frac{ie^{x\Delta g(z)}}{R_-(z)f_-^2(z)} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} \begin{bmatrix} 1 & -\frac{ie^{-x\Delta g(z)}}{R_+(z)f_+^2(z)} \\ 0 & 1 \end{bmatrix}, \quad z \in \gamma_1, \end{aligned} \quad (3.66)$$

and the jump matrix $V_5(x; z)|_{\gamma_2}$ has the following decomposition:

$$\begin{aligned} V_5(x; z) &= \begin{bmatrix} \delta f(z)e^{x\Delta g(z)} & i \\ 0 & \delta^{-1}f(z)e^{-x\Delta g(z)} \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ \frac{if_-^2(z)}{R_-(z)}e^{-x\Delta g(z)} & 1 \end{bmatrix} \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -\frac{if_+^2(z)}{R_+(z)}e^{x\Delta g(z)} & 1 \end{bmatrix}, \quad z \in \gamma_2. \end{aligned} \quad (3.67)$$

According to the above-mentioned decompositions with opening lenses, we make the following transformation:

$$Y_2(x; z) = \begin{cases} Y_1(x; z) \begin{bmatrix} 1 & \frac{ie^{-2x(g(z)-z)}}{R(z)f^2(z)} \\ 0 & 1 \end{bmatrix}, & z \text{ in the upper } (O_1^+)/\text{lower}(O_1^-) \text{ part of lens } O_1, \\ Y_1(x; z) \begin{bmatrix} 1 & 0 \\ \frac{if^2(z)}{R(z)}e^{2x(g(z)-z)} & 1 \end{bmatrix}, & z \text{ in the upper } (O_2^+)/\text{lower}(O_2^-) \text{ part of lens } O_2, \\ Y_1(x; z), & \text{otherwise.} \end{cases} \quad (3.68)$$

Then the matrix function $Y_2(x; z)$ satisfies the following Riemann-Hilbert problem.

RHP 8. Find a 2×2 matrix function $Y_2(x; z)$ that satisfies the following properties:

- *Analyticity:* $Y_2(x; z)$ is analytic in $\mathbb{C} \setminus ((-b, b) \cup O_1 \cup O_2)$ and takes continuous boundary values on $(-b, b) \cup O_1 \cup O_2$.
- *Jump condition:* The boundary values on the jump contour are defined as

$$Y_{2+}(x; z) = Y_{2-}(x; z)V_6(x; z), \quad z \in \gamma_1 \cup [-a, a] \cup \gamma_2 \cup O_1^+ \cup O_1^- \cup O_2^+ \cup O_2^-, \quad (3.69)$$

where the jump matrix is defined as

$$V_6(x; z) = \begin{cases} \begin{bmatrix} 1 & -\frac{ie^{-2x(g(z)-z)}}{R(z)f^2(z)} \\ 0 & 1 \end{bmatrix}, & z \in O_1^+ \cup O_1^-, \\ \begin{bmatrix} 1 & 0 \\ -\frac{if^2(z)}{R(z)}e^{2x(g(z)-z)} & 1 \end{bmatrix}, & z \in O_2^+ \cup O_2^-, \\ \begin{bmatrix} e^{xm_2+m_1} & 0 \\ 0 & e^{-(xm_2+m_1)} \end{bmatrix}, & z \in [-a, a], \\ \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}, & z \in \gamma_1 \cup \gamma_2; \end{cases} \quad (3.70)$$

- *Normalization:* $Y_2(x; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

Lemma 3.1. [23] *The following inequalities hold:*

$$\begin{aligned} \operatorname{Re}(g(z) - z) &< 0, \quad z \in O_1 \setminus \{a, b\}, \\ \operatorname{Re}(z - g(z)) &< 0, \quad z \in O_2 \setminus \{-a, -b\}. \end{aligned} \quad (3.71)$$

According to Lemma 3.1, we know that the off-diagonal entries of the jump matrix defined by Eq. (3.70) along the upper and lower lenses O_1, O_2 are exponentially decay when $x \rightarrow -\infty$.

3.2 The outer matrix parametrix

We construct a new Riemann-Hilbert problem with the matrix function having only jumps on $(-b, b)$.

RHP 9. *Find a 2×2 matrix function $Y^o(x; z)$ that satisfies the following properties:*

- *Analyticity:* $Y^o(x; z)$ is analytic in $\mathbb{C} \setminus (-b, b)$ and takes continuous boundary values on $(-b, b)$.
- *Jump condition:* The boundary values on the jump contour are defined as

$$Y_+^o(x; z) = Y_-^o(x; z)V_7(x; z), \quad z \in \gamma_1 \cup [-a, a] \cup \gamma_2, \quad (3.72)$$

where

$$V_7(x; z) = \begin{cases} \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}, & z \in \gamma_1 \cup \gamma_2, \\ \begin{bmatrix} e^{xm_2+m_1} & 0 \\ 0 & e^{-(xm_2+m_1)} \end{bmatrix}, & z \in [-a, a]. \end{cases} \quad (3.73)$$

- *Normalization:* $Y^o(x; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

Notice that the Riemann-Hilbert problem 9 is similar to the one arising from the long-time asymptotic behavior of KdV equation with step-like initial conditions [76]. In order to solve the Riemann-Hilbert problem 9, we will define a two-sheeted Riemann surface Ω of genus 1 related to the function $C(z)$ (see Fig. 3),

$$\Omega := \{(\zeta_1, \zeta_2) \in \mathbb{C}^2 \mid \zeta_2^2 = C^2(z) = (z^2 - a^2)(z^2 - b^2)\}. \quad (3.74)$$

Note that

$$\begin{aligned} \oint_{A_1} \frac{m_2}{4\pi i C(\lambda)} d\lambda &= \frac{iK_1(\sqrt{1-m^2})}{2K_1(m)} =: \tau, \\ \oint_{A_2} \frac{m_2}{4\pi i C(\lambda)} d\lambda &= 1. \end{aligned} \quad (3.75)$$

Let

$$\begin{aligned} m_3(z) &= \int_b^z \frac{m_2}{4\pi i C(\lambda)} d\lambda, \\ m_4(z) &= \int_b^z \frac{\lambda^2 + b^2 \left(\frac{K_2(m)}{K_1(m)} - 1 \right)}{C(\lambda)} d\lambda, \\ m_5(z) &= \left(\frac{(z-b)(z+a)}{(z-a)(z+b)} \right)^{\frac{1}{4}}, \end{aligned} \quad (3.76)$$




Figure 3: The two-sheeted Riemann surface with genus 1, Ω , given by Eq. (3.74).

Then one has the following jump relations:

$$\begin{aligned}
m_{3+}(z) + m_{3-}(z) &= 0, & z \in \gamma_1, \\
m_{3+}(z) - m_{3-}(z) &= -\tau, & z \in (-a, a), \\
m_{3+}(z) + m_{3-}(z) &= -1, & z \in \gamma_2, \\
m_{3+}(z) - m_{3-}(z) &= 0, & z \in [b, +\infty), \\
m_{5+}(z) &= im_{5-}(z), & z \in \gamma_1 \cup \gamma_2,
\end{aligned} \tag{3.77}$$

Based on Ref. [23], one has the following lemma:

Lemma 3.2. *These functions:*

$$\begin{aligned}
\psi_{11}(x; z) &= (m_5(z) + m_5^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(m_3(z) + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(m_3(z) + \frac{1}{4}; \tau) \vartheta_3(\frac{xm_2+m_1}{2\pi i}; \tau)}, \\
\psi_{21}(x; z) &= (m_5(z) - m_5^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(m_3(z) + \frac{xm_2+m_1}{2\pi i} - \frac{1}{4}; \tau)}{\vartheta_3(m_3(z) - \frac{1}{4}; \tau) \vartheta_3(\frac{xm_2+m_1}{2\pi i}; \tau)}, \\
\psi_{12}(x; z) &= (m_5(z) - m_5^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(-m_3(z) + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(-m_3(z) + \frac{1}{4}; \tau) \vartheta_3(\frac{xm_2+m_1}{2\pi i}; \tau)}, \\
\psi_{22}(x; z) &= (m_5(z) + m_5^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(-m_3(z) + \frac{xm_2+m_1}{2\pi i} - \frac{1}{4}; \tau)}{\vartheta_3(-m_3(z) - \frac{1}{4}; \tau) \vartheta_3(\frac{xm_2+m_1}{2\pi i}; \tau)},
\end{aligned} \tag{3.78}$$

have the following jump relations:

$$\begin{aligned}
\psi_{11+}(x; z) &= i\psi_{12-}(x; z), & \psi_{21+}(x; z) &= i\psi_{22-}(x; z), & z \in \gamma_1 \cup \gamma_2, \\
\psi_{12+}(x; z) &= i\psi_{11-}(x; z), & \psi_{22+}(x; z) &= i\psi_{21-}(x; z), & z \in \gamma_1 \cup \gamma_2, \\
\psi_{11+}(x; z) &= \psi_{11-}(x; z)e^{xm_2+m_1}, & \psi_{12+}(x; z) &= \psi_{12-}(x; z)e^{-(xm_2+m_1)}, & z \in (-a, a), \\
\psi_{21+}(x; z) &= \psi_{21-}(x; z)e^{xm_2+m_1}, & \psi_{22+}(x; z) &= \psi_{22-}(x; z)e^{-(xm_2+m_1)}, & z \in (-a, a).
\end{aligned} \tag{3.79}$$

where the third Jacobi theta function is defined as

$$\vartheta_3(z; \tau) = \sum_{n=-\infty}^{+\infty} e^{i(2\pi n z + \pi n^2 \tau)}, \quad z \in \mathbb{C}, \tag{3.80}$$

which has the periodicity condition:

$$\vartheta_3(z + h\tau + l; \tau) = e^{-(2\pi h z + \pi h^2 \tau)} \vartheta_3(z; \tau), \quad h, l \in \mathbb{Z}. \tag{3.81}$$

Proof. If $z \in \gamma_1 \cup \gamma_2$, we have

$$\begin{aligned}\psi_{11+}(x; z) &= (m_{5+}(z) + m_{5+}^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(m_{3+}(z) + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(m_{3+}(z) + \frac{1}{4}; \tau) \vartheta_3(\frac{xm_2+m_1}{2\pi i}; \tau)} \\ &= i(m_{5-}(z) - m_{5-}^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(-m_{3-}(z) + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(-m_{3-}(z) + \frac{1}{4}; \tau) \vartheta_3(\frac{xm_2+m_1}{2\pi i}; \tau)} \\ &= i\psi_{12-}(x; z).\end{aligned}$$

If $z \in (-a, a)$, we have

$$\begin{aligned}\psi_{11+}(x; z) &= (m_5(z) + m_5^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(m_{3+}(z) + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(m_{3+}(z) + \frac{1}{4}; \tau) \vartheta_3(\frac{xm_2+m_1}{2\pi i}; \tau)} \\ &= (m_5(z) + m_5^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(m_{3-}(z) - \tau + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(m_{3-}(z) - \tau + \frac{1}{4}; \tau) \vartheta_3(\frac{xm_2+m_1}{2\pi i}; \tau)}.\end{aligned}\tag{3.82}$$

According to Eq. (3.81), we have

$$\begin{aligned}& \frac{\vartheta_3(m_{3-}(x; z) - \tau + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(m_{3-}(z) - \tau + \frac{1}{4}; \tau)} \\ &= \frac{e^{-2i\pi\tau + 2\pi i(m_{3-}(z) + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4})} \vartheta_3(m_{3-}(z) + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4}; \tau)}{e^{-2i\pi\tau + 2\pi i(m_{3-}(z) + \frac{1}{4})} \vartheta_3(m_{3-}(z) - \tau + \frac{1}{4}; \tau)} \\ &= \frac{\vartheta_3(m_{3-}(z) + \frac{xm_2+m_1}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(m_{3-}(z) + \frac{1}{4}; \tau)} e^{xm_2+m_1},\end{aligned}\tag{3.83}$$

Substituting Eq. (3.83) into Eq. (3.82), we obtain

$$\psi_{11+}(x; z) = \psi_{11-}(x; z) e^{xm_2+m_1}.\tag{3.84}$$

Similarly, it can be proven that other equations also hold. Thus the proof is completed. \square

Proposition 3.1. *The solution of Riemann-Hilbert problem 9 can be expressed in the following form.*

$$Y^o(x; z) = \frac{1}{2} \begin{bmatrix} \psi_{11}(x; z) & \psi_{12}(x; z) \\ \psi_{21}(x; z) & \psi_{22}(x; z) \end{bmatrix}.\tag{3.85}$$

Proof. According to Lemma 3.2, we know that the matrix function $Y^o(x; z)$ is a solution of Riemann-Hilbert problem 9. \square

3.3 The local matrix parametrix

In this section, we will construct a local matrix parametrix $Y^b(x; z)$ as $z \in \delta^b := \{z \mid |z - b| < \epsilon, \epsilon \text{ is a suitable small, positive parameter}\}$. Similarly, we can define regions δ^a , δ^{-b} , and δ^{-a} . Note that

$$z - g_{\pm}(z) = \mathcal{O}(\sqrt{z - b}), \quad z \rightarrow b.\tag{3.86}$$

Then, we define the following conformal map:

$$\zeta_b := \frac{x^2(g(z) - z)^2}{4}, \quad z \in \delta^b.\tag{3.87}$$

Make the following transformation:

$$Y^{(1)}(x; z) = \begin{cases} Y_2(x; z) \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{R(z)}f(z)} \right)^{\sigma_3} e^{-2\sqrt{\zeta_b}\sigma_3} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, & z \in \mathbb{C}_+ \cap \delta^b, \\ Y_2(x; z) \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{-R(z)}f(z)} \right)^{\sigma_3} e^{-2\sqrt{\zeta_b}\sigma_3} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, & z \in \mathbb{C}_- \cap \delta^b. \end{cases} \quad (3.88)$$

Lemma 3.3. *The matrix function $Y^{(1)}(x; z)$ satisfies the following jump conditions:*

$$Y_+^{(1)}(x; z) = \begin{cases} Y_-^{(1)}(x; z) \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}, & z \in \delta^b \cap \{\text{upper and lower lenses}\}, \\ Y_-^{(1)}(x; z) \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, & z \in \delta^b \cap [0, +\infty). \end{cases} \quad (3.89)$$

Proof. If $z \in \delta^b \cap [0, +\infty)$, we have

$$\begin{aligned} Y_+^{(1)}(x; z) &= Y_{2+}(x; z) \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{R(z)}f(z)} \right)^{\sigma_3} e^{-2\sqrt{\zeta_b}\sigma_3} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ &= Y_{2-}(x; z) \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{R(z)}f(z)} \right)^{\sigma_3} e^{-2\sqrt{\zeta_b}\sigma_3} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ &= Y_-^{(1)}(x; z) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} e^{2\sqrt{\zeta_b}\sigma_3} \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{R(z)}f(z)} \right)^{-\sigma_3} \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{R(z)}f(z)} \right)^{\sigma_3} e^{-2\sqrt{\zeta_b}\sigma_3} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ &= Y_-^{(1)}(x; z) \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}. \end{aligned}$$

If $z \in \delta^b \cap \{\text{upper lens}\}$, we have

$$\begin{aligned} Y_+^{(1)}(x; z) &= Y_{2+}(x; z) \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{R(z)}f(z)} \right)^{\sigma_3} e^{-2\sqrt{\zeta_b}\sigma_3} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ &= Y_{2-}(x; z) \begin{bmatrix} 1 & -\frac{ie^{-2x(g(z)-z)}}{R(z)f^2(z)} \\ 0 & 1 \end{bmatrix} \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{R(z)}f(z)} \right)^{\sigma_3} e^{-2\sqrt{\zeta_b}\sigma_3} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ &= Y_-^{(1)}(x; z) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} e^{2\sqrt{\zeta_b}\sigma_3} \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{R(z)}f(z)} \right)^{-\sigma_3} \begin{bmatrix} 1 & -\frac{ie^{-2x(g(z)-z)}}{R(z)f^2(z)} \\ 0 & 1 \end{bmatrix} \\ &\quad \times \left(\frac{\exp(\frac{i\pi}{4})}{\sqrt{R(z)}f(z)} \right)^{\sigma_3} e^{-2\sqrt{\zeta_b}\sigma_3} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \\ &= Y_-^{(1)}(x; z) \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}. \end{aligned}$$




Figure 4: Regional division for the Bessel model $U_{Bes}(z)$.

As the same way, we can obtain the jump condition of matrix function $Y^{(1)}(x; z)$ as $z \in \delta^b \cap \{\text{lower lens}\}$. Thus the proof is completed. \square

Next, we will introduce the Bessel model $U_{Bes}(z)$ which satisfies the following Riemann-Hilbert problem [23, 73]:

RHP 10. Find a 2×2 matrix function $U_{Bes}(z)$ that satisfies the following properties:

- *Analyticity:* $U_{Bes}(z)$ is analytic for z in the three regions shown in Figure 4, namely, $S_1 := \{z \in \mathbb{C} \mid |\arg(z)| < \frac{2\pi}{3}\}$, $S_2 := \{z \in \mathbb{C} \mid \frac{2\pi}{3} < \arg(z) < \pi\}$ and $S_3 := \{z \in \mathbb{C} \mid -\pi < \arg(z) < -\frac{2\pi}{3}\}$, where $-\pi < \arg(z) \leq \pi$, and takes continuous boundary values on the excluded rays and at the origin from each sector.
- *Jump condition:* The boundary values on the jump contour γ_U where $\gamma_U := \gamma_{\pm} \cup \gamma_0$ with $\gamma_{\pm} = \{\arg(z) = \pm \frac{2\pi}{3}\}$ and $\gamma_0 = \{\arg(z) = \pi\}$ are defined as

$$U_{Bes+}(z) = U_{Bes-}(z)V_{Bes}(z), \quad z \in \gamma_U, \quad (3.90)$$

where

$$V_{Bes}(z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}, & z \in \gamma_{\pm}, \\ \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, & z \in \gamma_0; \end{cases} \quad (3.91)$$

- *Normalization:*

$$U_{Bes}(z) = \mathcal{O}(\ln(|z|)) \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, \quad z \rightarrow 0.$$

The solution of the Riemann-Hilbert Problem 10 can be expressed explicitly according to the Bessel functions [73]. In particular, the solution $U_{Bes}(z)$ satisfies:

$$U_{Bes}(z) = \frac{\sqrt{2}}{2} (2\pi\sqrt{z})^{-\frac{\sigma_3}{2}} \begin{bmatrix} 1 & -i \\ -i & 1 \end{bmatrix} \left(\mathbb{I} + \mathcal{O}(z^{-\frac{1}{2}}) \right) e^{2\sqrt{z}\sigma_3}, \quad z \rightarrow \infty, \quad (3.92)$$

in \mathbb{C} except from the jump curve γ_U .

Then the local matrix parametrix $Y^b(x; z)$ can be expressed as:

$$Y^b(x; z) = Y^{b1}(x; z)U_{Bes}(\zeta_b) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \left(\sqrt{\pm R(z)} f(z) e^{2\sqrt{\zeta_b} - i\pi/4} \right)^{-\sigma_3}, \quad z \in \delta^b \cap \mathbb{C}_{\pm}, \quad (3.93)$$

where

$$Y^{b1}(x; z) = \frac{\sqrt{2}}{2} Y^o(x; z) \left(\frac{\exp(i\pi/4)}{\sqrt{\pm R(z)} f(z)} \right)^{\sigma_3} \begin{bmatrix} i & 1 \\ 1 & i \end{bmatrix} (2\pi\sqrt{\zeta_b})^{\frac{\sigma_3}{2}}.$$

Then, we have

$$Y^b(x; z) (Y^o(x; z))^{-1} = \mathbb{I} + \mathcal{O}(|x|^{-1}), \quad x \rightarrow -\infty, z \in \partial\delta^b \setminus \gamma_U. \quad (3.94)$$

Similarly, we can construct local matrix functions $Y^a(x; z)$, $Y^{-a}(x; z)$ and $Y^{-b}(x; z)$ which satisfy the following properties:

$$\begin{aligned} Y^a(x; z) (Y^o(x; z))^{-1} &= \mathbb{I} + \mathcal{O}(|x|^{-1}), \quad x \rightarrow -\infty, z \in \partial\delta^a \setminus \gamma_U, \\ Y^{-a}(x; z) (Y^o(x; z))^{-1} &= \mathbb{I} + \mathcal{O}(|x|^{-1}), \quad x \rightarrow -\infty, z \in \partial\delta^{-a} \setminus \gamma_U, \\ Y^{-b}(x; z) (Y^o(x; z))^{-1} &= \mathbb{I} + \mathcal{O}(|x|^{-1}), \quad x \rightarrow -\infty, z \in \partial\delta^{-b} \setminus \gamma_U. \end{aligned}$$

Then we construct a matrix function as follows:

$$Y_3(x; z) = \begin{cases} Y^o(x; z), & z \in \mathbb{C} \setminus (\delta^a \cup \delta^b \cup \delta^{-a} \cup \delta^{-b}), \\ Y^{\pm a}(x; z), & z \in \delta^{\pm a}, \\ Y^{\pm b}(x; z), & z \in \delta^{\pm b}, \end{cases} \quad (3.95)$$

which has the following jump condition:

$$Y_{3+}(x; z) = Y_{3-}(x; z)V_8(x; z). \quad (3.96)$$

3.4 The small norm Riemann-Hilbert problem

Define the error matrix function $E(x; z)$ as follow:

$$E(x; z) = Y_2(x; z)Y_3^{-1}(x; z). \quad (3.97)$$

Then error matrix function $E(x; z)$ satisfies the following Riemann-Hilbert problem.

RHP 11. Find a 2×2 matrix function $E(x; z)$ that satisfies the following properties:

- *Analyticity:* $E(x; z)$ is analytic in $\mathbb{C} \setminus \gamma_3$ where $\gamma_3 := O_1 \cup O_2 \cup \partial\delta^a \cup \partial\delta^b \cup \partial\delta^{-a} \cup \partial\delta^{-b} \setminus (\delta^a \cup \delta^b \cup \delta^{-a} \cup \delta^{-b})$ and takes continuous boundary values on γ_3 .
- *Jump condition:* The boundary values on the jump contour are defined as

$$E_+(x; z) = E_-(x; z)V_E(x; z), \quad z \in \gamma_3, \quad (3.98)$$

where

$$V_E(x; z) = Y_{3-}(x; z)V_6(x; z)V_8^{-1}(x; z)Y_{3-}^{-1}(x; z), \quad (3.99)$$

which satisfies the following properties:

$$V_E(x; z) = \begin{cases} \mathbb{I} + \mathcal{O}(e^{-c|x|}), & z \in O_1 \cup O_2 \setminus (\bar{\delta}^a \cup \bar{\delta}^b \cup \bar{\delta}^{-a} \cup \bar{\delta}^{-b}), \\ \mathbb{I} + \mathcal{O}\left(\frac{1}{|x|}\right), & z \in \partial\delta^a \cup \partial\delta^b \cup \partial\delta^{-a} \cup \partial\delta^{-b}, \end{cases} \quad (3.100)$$

with c being a suitable positive, real parameter.

- Normalization: $E(x; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

According to Riemann-Hilbert problem 11, we obtain

$$E(x; z) = \mathbb{I} + \frac{E_1(x)}{xz} + \mathcal{O}(z^{-2}). \quad (3.101)$$

Proposition 3.2. *As $x \rightarrow -\infty$, the potential function $q(x, 0)$ has the following large-space asymptotics*

$$q(x, 0) = (b - a) \frac{\vartheta_3(0; \tau) \vartheta_3\left(\frac{xm_2 + m_1}{2\pi i} + \frac{1}{2}; \tau\right)}{\vartheta_3\left(\frac{1}{2}; \tau\right) \vartheta_3\left(\frac{xm_2 + m_1}{2\pi i}; \tau\right)} + \mathcal{O}(|x|^{-1}). \quad (3.102)$$

Proof. According to Eqs. (3.58), (3.68) and (3.101), we have

$$\begin{aligned} Y(x; z) &= Y_1(x; z) f(z)^{-\sigma_3} e^{-xg(z)\sigma_3} \\ &= Y_2(x; z) f(z)^{-\sigma_3} e^{-xg(z)\sigma_3} \\ &= E(x; z) Y_3(x; z) f(z)^{-\sigma_3} e^{-xg(z)\sigma_3} \\ &= \left(\mathbb{I} + \frac{E_1(x)}{xz} + \mathcal{O}(z^{-2}) \right) Y_3(x; z) f(z)^{-\sigma_3} e^{-xg(z)\sigma_3}, \end{aligned}$$

which leads to

$$\begin{aligned} Y_{12}(x; z) &= \left(Y_{3,12}(x; z) + \frac{E_{1,12}(x)}{xz} + \mathcal{O}(z^{-2}) \right) f(z) e^{xg(z)} \\ &= \left(Y_{12}^o(x; z) + \frac{E_{1,12}(x)}{xz} + \mathcal{O}(z^{-2}) \right) f(z) e^{xg(z)} \\ &= \frac{1}{2} \left(\psi_{12}(x; z) + \frac{E_{1,12}(x)}{xz} + \mathcal{O}(z^{-2}) \right) f(z) e^{xg(z)}. \end{aligned} \quad (3.103)$$

Note that

$$\begin{aligned} f(z) &= 1 + \frac{f_1}{z} + \mathcal{O}(z^{-2}), \\ e^{xg(z)} &= 1 + \frac{x\left(\frac{a^2 + b^2}{2} + b^2\left(\frac{K_2(m)}{K_1(m)} - 1\right)\right)}{z} + \mathcal{O}(z^{-2}), \\ m_5(z) &= 1 + \frac{a - b}{2z} + \mathcal{O}(z^{-2}). \end{aligned} \quad (3.104)$$

According to Eq. (3.57), one can show the large-space asymptotics (3.102) of the solution of the Hirota equation. \square

Note that similarly, we also consider the asymptotic behavior of $q(x, t)$ as $x \rightarrow -\infty$ for each fixed $t = t_0$, where the norming constant c_j of the residue conditions (2.28) of matrix function $M(x, t_0; z)$ is changed as

$$c_j = \frac{i(b - a)r(iz_j)e^{4i\alpha z_j^2 t_0 - 8\beta z_j^3 t_0}}{N\pi}, \quad (3.105)$$

where $r(z)$ is a real-valued, continuous and non-vanishing function for $z \in (ia, ib)$, with the symmetry $r(z^*) = r(z)$. Similarly, as $x \rightarrow -\infty$, we have the same large-space asymptotic behavior of the potential function $q(x, t_0)$ as Eq. (3.102).

4 Long-time asymptotics of the potential $q(x, t)$ as $t \rightarrow +\infty$

We recall the Riemann-Hilbert problem 5 of function $M^{(3)}(x, t; z)$:

$$M_+^{(3)}(x, t; z) = \begin{cases} M_-^{(3)}(x, t; z) \begin{bmatrix} 1 & 0 \\ iR(z)e^{-2zx-4i\alpha z^2 t+8\beta z^3 t} & 1 \end{bmatrix}, & z \in \gamma_1, \\ M_-^{(3)}(x, t; z) \begin{bmatrix} 1 & iR(z)e^{2zx+4i\alpha z^2 t-8\beta z^3 t} \\ 0 & 1 \end{bmatrix}, & z \in \gamma_2, \end{cases} \quad (4.106)$$

$$M^{(3)}(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1}), \quad z \rightarrow \infty.$$

Girotti *et al* [24] studied the long-time asymptotics of the dense soliton gas of modified KdV equation. Similarly, in what follows we will study the long-time asymptotics of the potential $q(x, t)$ as $t \rightarrow +\infty$ based on the Riemann-Hilbert problem. In fact, the following similar results can also be found for $t \rightarrow -\infty$.

4.1 The case $\xi = x/t > 4\beta b^2$

Let $\xi := \frac{x}{t}$, then Eq. (4.106) can be rewrite as:

$$M_+^{(3)}(x, t; z) = \begin{cases} M_-^{(3)}(x, t; z) \begin{bmatrix} 1 & 0 \\ iR(z)e^{-2zt(\xi-4\beta z^2+2i\alpha z)} & 1 \end{bmatrix}, & z \in \gamma_1, \\ M_-^{(3)}(x, t; z) \begin{bmatrix} 1 & iR(z)e^{2zt(\xi-4\beta z^2+2i\alpha z)} \\ 0 & 1 \end{bmatrix}, & z \in \gamma_2 \end{cases} \quad (4.107)$$

$$M^{(3)}(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1}), \quad z \rightarrow \infty.$$

Then we have

$$M^{(3)}(x, t; z) = \mathbb{I} + \mathcal{O}(e^{-2ta(\xi-4\beta b^2)}), \quad \xi > 4\beta b^2, \quad t \rightarrow +\infty. \quad (4.108)$$

and the potential function $q(x, t)$ becomes trivial.

4.2 The case $\xi_c < \xi = x/t < 4\beta b^2$

The jump matrices in Eq. (4.107) contain the two terms $iR(z)e^{\pm 2zt(\xi-4\beta z^2+2i\alpha z)}$. To deal with them conveniently, we only consider them to be of the pure imaginary functions, that is, $\alpha = 0$, $\beta \neq 0$, in which the Hirota equation reduces to the complex modified KdV equation

$$q_t + \beta(q_{xxx} + 6|q|^2 q_x) = 0, \quad (4.109)$$

in which one can take $\beta = 1$ without loss of generality.

Eq. (4.107) with $\alpha = 0$, $\beta = 1$ can be rewritten as:

$$M_+^{(3)}(x, t; z) = \begin{cases} M_-^{(3)}(x, t; z) \begin{bmatrix} 1 & 0 \\ iR(z)e^{-2tz(\xi-4z^2)} & 1 \end{bmatrix}, & z \in \gamma_1, \\ M_-^{(3)}(x, t; z) \begin{bmatrix} 1 & iR(z)e^{2tz(\xi-4z^2)} \\ 0 & 1 \end{bmatrix}, & z \in \gamma_2, \end{cases} \quad (4.110)$$

$$M^{(3)}(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1}), \quad z \rightarrow \infty.$$

In what follows we consider the condition $\xi_c < \xi < b^2$, where ξ_c to be determined below (see Eq. (4.119)).
Let

$$\gamma_{1a_1} := (a_1, b), \quad \gamma_{2a_1} := (-b, -a_1), \quad a_1 \in (a, b).$$

Therefore, we construct the following Riemann-Hilbert problem based on the RHP (4.110).

RHP 12. Find a 2×2 matrix function $Y_{a_1}(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $Y_{a_1}(x, t; z)$ is analytic in $\mathbb{C} \setminus (\gamma_1 \cup \gamma_2)$ and takes continuous boundary values on $\gamma_1 \cup \gamma_2$.
- *Jump condition:* The boundary values on the jump contour $\gamma_1 \cup \gamma_2$ are defined as

$$Y_{a_1+}(x, t; z) = Y_{a_1-}(x, t; z)V_{4a_1}(x, t; z), \quad z \in \gamma_1 \cup \gamma_2, \quad (4.111)$$

where

$$V_{4a_1}(x, t; z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ iR(z)e^{-2tz(\xi-4z^2)} & 1 \end{bmatrix}, & z \in \gamma_1, \\ \begin{bmatrix} 1 & iR(z)e^{2tz(\xi-4z^2)} \\ 0 & 1 \end{bmatrix}, & z \in \gamma_2; \end{cases} \quad (4.112)$$

- *Normalization:* $Y_{a_1}(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

According to Eq. (2.54), we can recover $q(x, t)$ by the following formula:

$$q(x, t) = -2 \lim_{z \rightarrow \infty} (zY_{a_1}(x, t; z))_{12}. \quad (4.113)$$

To solve Riemann-Hilbert problem 12, we make the following transformation:

$$Y_{1a_1}(x, t; z) = Y_{a_1}(x, t; z)[f_{a_1}(z)e^{tg_{a_1}(z)}]^{\sigma_3}, \quad (4.114)$$

where $f_{a_1}(z)$ and $g_{a_1}(z)$ are scalar functions which satisfy the following properties:

- *Analyticity:* $f_{a_1}(z)$, $g_{a_1}(z)$ are analytic in $\mathbb{C} \setminus (-b, b)$ and take continuous boundary values on $(-b, b)$.
- *Jump conditions:* The boundary values on the jump contour are defined as

$$\begin{aligned} f_{a_1+}(z)f_{a_1-}(z) &= R^{-1}(z), & z \in \gamma_{1a_1}, \\ f_{a_1+}(z)f_{a_1-}(z) &= R(z), & z \in \gamma_{2a_1}, \\ f_{a_1+}(z) &= f_{a_1-}(z)e^{m_{1a_1}}, & z \in [-a_1, a_1] \end{aligned} \quad (4.115)$$

and

$$\begin{aligned} g_{a_1+}(z) + g_{a_1-}(z) &= 2\xi z - 8z^3, & z \in \gamma_{1a_1} \cup \gamma_{2a_1}, \\ g_{a_1+}(z) - g_{a_1-}(z) &= m_{2a_1}, & z \in [-a_1, a_1], \end{aligned} \quad (4.116)$$

where

$$m_{1a_1} = \frac{\int_{a_1}^b \frac{2 \ln R(\lambda)}{C_{a_1+}(\lambda)} d\lambda}{\int_{-a_1}^{a_1} \frac{d\lambda}{C_{a_1}(\lambda)}}, \quad C_{a_1}(z) = \sqrt{(z^2 - a_1^2)(z^2 - b^2)},$$

$$m_{2a_1} = -\frac{i\pi b(2a_1^2 + 2b^2 - \xi)}{K_1(m_{a_1})}, \quad m_{a_1} = \frac{a_1}{b}.$$

- Normalization: $f_{a_1}(z) = 1 + \mathcal{O}(z^{-1})$, $g_{a_1}(z) = \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

These scalar functions $f_{a_1}(z)$ and $g_{a_1}(z)$ can also be accurately solved as [23, 24]:

$$f_{a_1}(z) = \exp \left\{ \frac{C_{a_1}(z)}{2\pi i} \left(\int_{\gamma_{2a_1} - \gamma_{1a_1}} \frac{\ln R(\lambda)}{C_{a_1+}(\lambda)(\lambda - z)} d\lambda + \int_{-a_1}^{a_1} \frac{m_{1a_1}}{C_{a_1}(\lambda)(\lambda - z)} d\lambda \right) \right\}, \quad (4.117)$$

$$g_{a_1}(z) = -4z^3 + \xi z + \int_b^z \frac{12z^4 - z^2[\xi + 6(a_1^2 + b^2)] + 4a_1^2 b^2 + b^2[2(a_1^2 + b^2) - \xi] \left(\frac{K_2(m_{a_1})}{K_1(m_{a_1})} - 1 \right)}{C_{a_1}(\lambda)} d\lambda,$$

where the parameter a_1 is determined by the following equation:

$$\xi = 2(a_1^2 + b^2) + \frac{4a_1^2(a_1^2 - b^2)}{a_1^2 + b^2 \left(\frac{K_2(m_{a_1})}{K_1(m_{a_1})} - 1 \right)}, \quad (4.118)$$

such that one has

$$\xi_c = 2(a^2 + b^2) + \frac{4a^2(a^2 - b^2)}{a^2 + b^2 \left(\frac{K_2(m)}{K_1(m)} - 1 \right)}, \quad m = \frac{a}{b}. \quad (4.119)$$

Therefore, the matrix function $Y_{1a_1}(x, t; z)$ satisfies the following Riemann-Hilbert problem.

RHP 13. Find a 2×2 matrix function $Y_{1a_1}(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $Y_{1a_1}(x, t; z)$ is analytic in $\mathbb{C} \setminus (-b, b)$ and takes continuous boundary values on $(-b, b)$.
- *Jump condition:* The boundary values on the jump contour are defined as

$$Y_{1a_1+}(x, t; z) = Y_{1a_1-}(x, t; z) V_{5a_1}(x, t; z), \quad z \in (-b, b), \quad (4.120)$$

where

$$V_{5a_1}(x, t; z) = \begin{cases} \begin{bmatrix} \delta f_{a_1}(z) e^{t\Delta g_{a_1}(z)} & 0 \\ i & \delta^{-1} f_{a_1}(z) e^{-t\Delta g_{a_1}(z)} \end{bmatrix}, & z \in \gamma_{1a_1}, \\ \begin{bmatrix} e^{tm_{2a_1} + m_{1a_1}} & 0 \\ iR(z) \widehat{\delta} f_{a_1}(z) e^{t(\widehat{\Delta} g_{a_1}(z) - 2\xi z + 8z^3)} & e^{-(tm_{2a_1} + m_{1a_1})} \end{bmatrix}, & z \in [a, a_1], \\ \begin{bmatrix} e^{tm_{2a_1} + m_{1a_1}} & 0 \\ 0 & e^{-(tm_{2a_1} + m_{1a_1})} \end{bmatrix}, & z \in [-a, a], \\ \begin{bmatrix} e^{tm_{2a_1} + m_{1a_1}} & \frac{iR(z) e^{-t(\widehat{\Delta} g_{a_1}(z) - 2\xi z + 8z^3)}}{\widehat{\delta} f_{a_1}(z)} \\ 0 & e^{-(tm_{2a_1} + m_{1a_1})} \end{bmatrix}, & z \in [-a_1, -a], \\ \begin{bmatrix} \widehat{\delta} f_{a_1}(z) e^{t\widehat{\Delta} g_{a_1}(z)} & i \\ 0 & \frac{e^{-t\widehat{\Delta} g_{a_1}(z)}}{\widehat{\delta} f_{a_1}(z)} \end{bmatrix}, & z \in \gamma_{2a_1}; \end{cases} \quad (4.121)$$

with $\delta f_{a_1}(z) = f_{a_1+}(z)/f_{a_1-}(z)$, $\widehat{\delta} f_{a_1}(z) = f_{a_1+}(z)f_{a_1-}(z)$, $\Delta g_{a_1}(z) = g_{a_1+}(z) - g_{a_1-}(z)$, $\widehat{\Delta} g_{a_1}(z) = g_{a_1+}(z) + g_{a_1-}(z)$.

- *Normalization:* $Y_{1a_1}(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.




Figure 5: Riemann-Hilbert problem 14 of function $Y_{2a_1}(x, t; z)$. Opening lenses O_1 and O_2 : $t \rightarrow +\infty$.

4.2.1 Opening lenses

Based on the idea [24], we open the lens O_1 to pass through points $z = a_1$ and $z = b$ and lens O_2 to pass through points $z = -b$ and $z = -a_1$ (see Fig. 5). We define a new function as follows:

$$R_{\pm}(z) := \pm R(z), \quad z \in \gamma_{1a_1} \cup \gamma_{2a_1}. \quad (4.122)$$

Note that the jump matrix $V_5(x, t; z)|_{\gamma_{1a_1}}$ has the following decomposition:

$$\begin{aligned} V_{5a_1}(x, t; z) &= \begin{bmatrix} \delta f_{a_1}(z) e^{t\Delta g_{a_1}(z)} & 0 \\ i & \delta^{-1} f_{a_1}(z) e^{-t\Delta g_{a_1}(z)} \end{bmatrix} \\ &= \begin{bmatrix} 1 & \frac{i e^{-2t(g_{a_1}-(z)+4z^3-\xi z)}}{R_-(z) f_{a_1}^2(z)} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} \begin{bmatrix} 1 & -\frac{i e^{-2t(g_{a_1}+(z)+4z^3-\xi z)}}{R_+(z) f_{a_1}^2(z)} \\ 0 & 1 \end{bmatrix}, \quad z \in \gamma_{1a_1}, \end{aligned} \quad (4.123)$$

and the jump matrix $V_5(x, t; z)|_{\gamma_{2a_1}}$ has the following decomposition:

$$\begin{aligned} V_{5a_1}(x, t; z) &= \begin{bmatrix} \widehat{\delta} f_{a_1}(z) e^{t\widehat{\Delta} g_{a_1}(z)} & i \\ 0 & \frac{e^{-t\widehat{\Delta} g_{a_1}(z)}}{\widehat{\delta} f_{a_1}(z)} \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ \frac{i e^{2t(g_{a_1}-(z)+4z^3-\xi z)} f_-^2(z)}{R_-(z)} & 1 \end{bmatrix} \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -\frac{i e^{2t(g_{a_1}+(z)+4z^3-\xi z)} f_+^2(z)}{R_+(z)} & 1 \end{bmatrix}, \quad z \in \gamma_{2a_1}. \end{aligned} \quad (4.124)$$

We can proceed with opening lenses. Make the following transformation:

$$Y_{2a_1}(x, t; z) = \begin{cases} Y_{1a_1}(x, t; z) \begin{bmatrix} 1 & \frac{ie^{-2t(g_{a_1}(z)+4z^3-\xi z)}}{R(z)f^2(z)} \\ 0 & 1 \end{bmatrix}, & \text{in the upper/lower lens } O_1, \\ Y_{1a_1}(x, t; z) \begin{bmatrix} 1 & 0 \\ \frac{ie^{2t(g_{a_1}(z)+4z^3-\xi z)}f^2(z)}{R(z)} & 1 \end{bmatrix}, & \text{in the upper/lower lens } O_2, \\ Y_{1a_1}(x, t; z), & \text{otherwise.} \end{cases}, \quad (4.125)$$

Then matrix function $Y_{2a_1}(x, t; z)$ satisfies the following Riemann-Hilbert problem.

RHP 14. Find a 2×2 matrix function $Y_{2a_1}(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $Y_2(x, t; z)$ is analytic in $\mathbb{C} \setminus ((-b, b) \cup O_1 \cup O_2)$ and takes continuous boundary values on $(-b, b) \cup O_1 \cup O_2$.
- *Jump condition:* The boundary values on the jump contour are defined as

$$Y_{2a_1+}(x, t; z) = Y_{2a_1-}(x, t; z)V_{6a_1}(x, t; z), \quad z \in (-b, b) \cup O_1^+ \cup O_1^- \cup O_2^+ \cup O_2^-, \quad (4.126)$$

where for $z \in O_1^+ \cup O_1^- \cup O_2^+ \cup O_2^-$

$$V_{6a_1}(x; z) = \begin{cases} \begin{bmatrix} 1 & -\frac{ie^{-2t(g_{a_1}(z)+4z^3-\xi z)}}{R(z)f^2(z)} \\ 0 & 1 \end{bmatrix}, & z \in O_1^+ \cup O_1^-, \\ \begin{bmatrix} 1 & 0 \\ -\frac{ie^{2t(g_{a_1}(z)+4z^3-\xi z)}f^2(z)}{R(z)} & 1 \end{bmatrix}, & z \in O_2^+ \cup O_2^-, \end{cases} \quad (4.127)$$

and for $z \in (-b, b)$

$$V_{6a_1}(x; z) = \begin{cases} \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}, & z \in \gamma_{1a_1} \cup \gamma_{2a_1}, \\ \begin{bmatrix} e^{tm_{2a_1}+m_{1a_1}} & 0 \\ iR(z)\widehat{\delta}f_{a_1}(z)e^{t(\widehat{\Delta}g_{a_1}(z)-2\xi z+8z^3)} & e^{-(tm_{2a_1}+m_{1a_1})} \end{bmatrix}, & z \in [a, a_1], \\ \begin{bmatrix} e^{tm_{2a_1}+m_{1a_1}} & 0 \\ 0 & e^{-(tm_{2a_1}+m_{1a_1})} \end{bmatrix}, & z \in [-a, a], \\ \begin{bmatrix} e^{tm_{2a_1}+m_{1a_1}} & \frac{iR(z)e^{-t(\widehat{\Delta}g_{a_1}(z)-2\xi z+8z^3)}}{\widehat{\delta}f_{a_1}(z)} \\ 0 & e^{-(tm_{2a_1}+m_{1a_1})} \end{bmatrix}, & z \in [-a_1, -a], \end{cases} \quad (4.128)$$

- *Normalization:* $Y_{2a_1}(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.




Figure 6: Riemann surface Ω .

Lemma 4.1. [23] *The following inequalities hold:*

$$\begin{aligned}
\operatorname{Re}(g_{a_1}(z) + 4z^3 - \xi z) &> 0, \quad z \in O_1 \setminus \{a_1, b\}, \\
\operatorname{Re}(g_{a_1}(z) + 4z^3 - \xi z) &< 0, \quad z \in O_2 \setminus \{-a_1, -b\}, \\
\operatorname{Re}(g_{a_1+}(z) + g_{a_1-}(z) - 2\xi z + 8z^3) &< 0, \quad z \in [a, a_1), \\
\operatorname{Re}(g_{a_1+}(z) + g_{a_1-}(z) - 2\xi z + 8z^3) &> 0, \quad z \in (-a_1, -a].
\end{aligned} \tag{4.129}$$

According to Lemma 4.1, we know that the off-diagonal entries of the jump matrix defined by Eq. (4.127) along the upper and lower lenses O_1, O_2 are exponentially decay when $t \rightarrow +\infty$.

4.2.2 The outer parametrix

In order to solve the Riemann-Hilbert problem 14, we will introduce a two-sheeted Riemann surface Ω associated to the function function $C_{a_1}(z)$ (see Fig. 6)

$$\Omega := \{(\zeta_1, \zeta_2) \in \mathbb{C}^2 \mid \zeta_2^2 = C_{a_1}^2(z) = (z^2 - a_1^2)(z^2 - b^2)\}. \tag{4.130}$$

We construct a new Riemann-Hilbert problem which only has jumps on $(-b, b)$.

RHP 15. *Find a 2×2 matrix function $Y^o(x, t; z)$ that satisfies the following properties:*

- *Analyticity:* $Y^o(x, t; z)$ is analytic in $\mathbb{C} \setminus (-b, b)$ and takes continuous boundary values on $(-b, b)$.
- *Jump condition:* The boundary values on the jump contour are defined as

$$Y_+^o(x, t; z) = Y_-^o(x, t; z)V_{7a_1}(x, t; z), \quad z \in \gamma_{1a_1} \cup [-a_1, a_1] \cup \gamma_{2a_1}, \tag{4.131}$$

where

$$V_{7a_1}(x, t; z) = \begin{cases} \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}, & z \in \gamma_{1a_1} \cup \gamma_{2a_1}, \\ \begin{bmatrix} e^{tm_{2a_1} + m_{1a_1}} & 0 \\ 0 & e^{-(tm_{2a_1} + m_{1a_1})} \end{bmatrix}, & z \in [-a_1, a_1]. \end{cases} \tag{4.132}$$

- *Normalization:* $Y^o(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

Then, we define the functions:

$$\begin{aligned}
\tau &:= \oint_{A_1} -\frac{b}{4K_1(m_{a_1})C_{a_1}(\lambda)} d\lambda, \\
m_{3a_1}(z) &= \int_b^z -\frac{b}{4K_1(m_{a_1})C_{a_1}(\lambda)} d\lambda, \\
m_{5a_1}(z) &= \left(\frac{(z-b)(z+a_1)}{(z-a_1)(z+b)} \right)^{\frac{1}{4}},
\end{aligned} \tag{4.133}$$

which have the following jump relations:

$$\begin{aligned}
m_{3a_1+}(z) + m_{3a_1-}(z) &= 0, \quad z \in \gamma_{1a_1}, \\
m_{3a_1+}(z) - m_{3a_1-}(z) &= -\tau, \quad z \in (-a_1, a_1), \\
m_{3a_1+}(z) + m_{3a_1-}(z) &= -1, \quad z \in \gamma_{2a_1}, \\
m_{3a_1+}(z) - m_{3a_1-}(z) &= 0, \quad z \in [b, +\infty), \\
m_{5a_1+}(z) &= im_{5a_1-}(z), \quad z \in \gamma_{1a_1} \cup \gamma_{2a_1},
\end{aligned} \tag{4.134}$$

Lemma 4.2. [23] Define the functions:

$$\begin{aligned}
\psi_{11}(z) &= (m_{5a_1}(z) + m_{5a_1}^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(m_{3a_1}(z) + \frac{tm_{2a_1} + m_{1a_1}}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(m_{3a_1}(z) + \frac{1}{4}; \tau) \vartheta_3(\frac{tm_{2a_1} + m_{1a_1}}{2\pi i}; \tau)}, \\
\psi_{21}(z) &= (m_{5a_1}(z) - m_{5a_1}^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(m_{3a_1}(z) + \frac{tm_{2a_1} + m_{1a_1}}{2\pi i} - \frac{1}{4}; \tau)}{\vartheta_3(m_{3a_1}(z) - \frac{1}{4}; \tau) \vartheta_3(\frac{tm_{2a_1} + m_{1a_1}}{2\pi i}; \tau)}, \\
\psi_{12}(z) &= (m_{5a_1}(z) - m_{5a_1}^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(-m_{3a_1}(z) + \frac{tm_{2a_1} + m_{1a_1}}{2\pi i} + \frac{1}{4}; \tau)}{\vartheta_3(-m_{3a_1}(z) + \frac{1}{4}; \tau) \vartheta_3(\frac{tm_{2a_1} + m_{1a_1}}{2\pi i}; \tau)}, \\
\psi_{22}(z) &= (m_{5a_1}(z) + m_{5a_1}^{-1}(z)) \frac{\vartheta_3(0; \tau) \vartheta_3(-m_{3a_1}(z) + \frac{tm_{2a_1} + m_{1a_1}}{2\pi i} - \frac{1}{4}; \tau)}{\vartheta_3(-m_{3a_1}(z) - \frac{1}{4}; \tau) \vartheta_3(\frac{tm_{2a_1} + m_{1a_1}}{2\pi i}; \tau)},
\end{aligned} \tag{4.135}$$

which have the following jump relations:

$$\begin{aligned}
\psi_{11+}(z) &= i\psi_{12-}(z), \quad \psi_{21+}(z) = i\psi_{22-}(z), \quad z \in \gamma_{1a_1} \cup \gamma_{2a_1}, \\
\psi_{12+}(z) &= i\psi_{11-}(z), \quad \psi_{22+}(z) = i\psi_{21-}(z), \quad z \in \gamma_{1a_1} \cup \gamma_{2a_1}, \\
\psi_{11+}(z) &= \psi_{11-}(z) e^{tm_{2a_1} + m_{1a_1}}, \quad \psi_{12+}(z) = \psi_{12-}(z) e^{-(tm_{2a_1} + m_{1a_1})}, \quad z \in (-a_1, a_1), \\
\psi_{21+}(z) &= \psi_{21-}(z) e^{tm_{2a_1} + m_{1a_1}}, \quad \psi_{22+}(z) = \psi_{22-}(z) e^{-(tm_{2a_1} + m_{1a_1})}, \quad z \in (-a_1, a_1).
\end{aligned} \tag{4.136}$$

Proof. According to Lemma 3.2, the above conclusion can be obtained. \square

Proposition 4.1. The solution of Riemann-Hilbert problem 15 can be expressed in the following form.

$$Y^o(x, t; z) = \frac{1}{2} \begin{bmatrix} \psi_{11}(x, t; z) & \psi_{12}(x, t; z) \\ \psi_{21}(x, t; z) & \psi_{22}(x, t; z) \end{bmatrix}. \tag{4.137}$$

Proof. According to Lemma 4.2, we know that the matrix function $Y^o(x, t; z)$ is a solution of Riemann-Hilbert problem 15. \square




Figure 7: Regional division for the Airy model $U_{Ai}(z)$.

4.2.3 The local matrix parametrix

In this section, we will construct a local matrix parametrix $Y^{-a_1}(x, t; z)$ as $z \in \delta^{-a_1} := \{z \mid |z + a_1| < \epsilon, \epsilon \text{ is a suitable small positive parameter}\}$. Similarly, we can define regions δ^{a_1} , δ^b , and δ^{-b} . Then, let the conformal map be [23]:

$$\zeta_{-a_1} := \left(\frac{3t}{4}\right)^{\frac{2}{3}} \left(\int_{-a_1}^z [g'_{a_1+}(\lambda) - g'_{a_1-}(\lambda)] d\lambda\right)^{\frac{2}{3}}, \quad z \in \delta^{-a_1}. \quad (4.138)$$

and make the following transformation:

$$Y^{(1)}(x, t; z) = \begin{cases} Y_{2a_1}(x, t; z) \left(\frac{\sqrt{R(z)}}{f_{a_1}(z)}\right)^{\sigma_3} e^{\left(\frac{i\pi}{4} - \frac{1}{2}(tm_{2a_1} + m_{1a_1}) - \frac{2}{3}\zeta_{-a_1}^{\frac{3}{2}}\right)\sigma_3}, & z \in \mathbb{C}_+ \cap \delta^{-a_1}, \\ Y_{2a_1}(x, t; z) \left(\frac{\sqrt{-R(z)}}{f_{a_1}(z)}\right)^{\sigma_3} e^{\left(\frac{i\pi}{4} + \frac{1}{2}(tm_{2a_1} + m_{1a_1}) - \frac{2}{3}\zeta_{-a_1}^{\frac{3}{2}}\right)\sigma_3}, & z \in \mathbb{C}_- \cap \delta^{-a_1}. \end{cases} \quad (4.139)$$

Lemma 4.3. [23] *Matrix function $Y^{(1)}(x, t; z)$ satisfies the following jump conditions:*

$$Y_+^{(1)}(x, t; z) = \begin{cases} Y_-^{(1)}(x, t; z) \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, & z \in \delta^{-a_1} \cap \{\text{upper and lower lenses}\}, \\ Y_-^{(1)}(x, t; z) \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, & z \in \delta^{-a_1} \cap (0, +\infty), \\ Y_-^{(1)}(x, t; z) \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, & z \in \delta^{-a_1} \cap (-\infty, 0). \end{cases} \quad (4.140)$$

Next, we will introduce the Airy model $U_{Ai}(z)$ [74] which satisfies the following Riemann-Hilbert problem:

RHP 16. *Find a 2×2 matrix function $U_{Ai}(z)$ that satisfies the following properties:*

- *Analyticity:* $U_{Ai}(z)$ is analytic for z in the three regions shown in Figure 7, namely, $S_1 : 0 < \arg(z) < \frac{2\pi}{3}$, $S_2 : \frac{2\pi}{3} < \arg(z) < \pi$, $S_3 : -\pi < \arg(z) < -\frac{2\pi}{3}$ and $S_4 : -\frac{2\pi}{3} < \arg(z) < 0$ where $-\pi < \arg(z) \leq \pi$. It takes continuous boundary values on the excluded rays and at the origin from each sector.
- *Jump condition:* The boundary values on the jump contour γ_A where $\gamma_A := \gamma_{\pm} \cup \gamma_{0\pm}$ with $\gamma_{\pm} = \{\arg(z) = \pm \frac{2\pi}{3}\}$, $\gamma_{0+} = \{\arg(z) = 0\}$ and $\gamma_{0-} = \{\arg(z) = \pi\}$ are defined as:

$$U_{Ai+}(z) = U_{Ai-}(z)V_{Ai}(z), \quad z \in \gamma_A, \quad (4.141)$$

where

$$V_{Ai}(z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, & z \in \gamma_{\pm}, \\ \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, & z \in \gamma_{0+}, \\ \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, & z \in \gamma_{0-}; \end{cases} \quad (4.142)$$

- *Normalization:* $U_{Ai}(z)$ is bounded as $z \rightarrow 0, z \in \mathbb{C} \setminus \gamma_A$.

The solution of the Riemann-Hilbert Problem 16 can be expressed explicitly according to the Airy function $Ai(z)$. In particular, the solution satisfies:

$$U_{Ai}(z) = \frac{\sqrt{2}}{2} z^{-\frac{1}{4}\sigma_3} \begin{bmatrix} 1 & i \\ i & 1 \end{bmatrix} \left(\mathbb{I} + \mathcal{O}(z^{-\frac{3}{2}}) \right) e^{-\frac{2}{3}z^{\frac{3}{2}}\sigma_3}, \quad z \rightarrow \infty, \quad (4.143)$$

in \mathbb{C} except for the jumps.

Then the local matrix parametrix $Y^{-a_1}(x, t; z)$ can be expressed as:

$$Y^{-a_1}(x, t; z) = Y^{-a_{11}}(x, t; z) U_{Ai}(\zeta_{-a_1}) \left(\frac{\sqrt{\pm R(z)}}{f_{a_1}(z)} \right)^{-\sigma_3} e^{-\left(\frac{i\pi}{4} \mp \frac{1}{2}(tm_{2a_1} + m_{1a_1}) - \frac{2}{3}\zeta_{-a_1}^{\frac{3}{2}} \right) \sigma_3}, \quad z \in \delta^{-a_1} \cap \mathbb{C}_{\pm},$$

where

$$Y^{-a_{11}}(x, t; z) = \frac{\sqrt{2}}{2} Y^o(x, t; z) e^{\left(\frac{i\pi}{4} \mp \frac{1}{2}(tm_{2a_1} + m_{1a_1}) \right) \sigma_3} \left(\frac{\sqrt{\pm R(z)}}{f_{a_1}(z)} \right)^{\sigma_3} \begin{bmatrix} 1 & -i \\ -i & 1 \end{bmatrix} (\zeta_{-a_1})^{\frac{1}{4}\sigma_3}.$$

Then, we have

$$Y^{-a_1}(x, t; z) (Y^o(x, t; z))^{-1} = \mathbb{I} + \mathcal{O}(t^{-1}), \quad t \rightarrow +\infty, z \in \partial\delta^{-a_1} \setminus \gamma_A. \quad (4.144)$$

Similarly, we can construct the local matrix function $Y^{a_1}(x, t; z)$ which satisfies the following property:

$$Y^{a_1}(x, t; z) (Y^o(x, t; z))^{-1} = \mathbb{I} + \mathcal{O}(t^{-1}), \quad t \rightarrow +\infty, z \in \partial\delta^{a_1} \setminus \gamma_A. \quad (4.145)$$

Similar to the case of $x \rightarrow -\infty$, we can provide the local function $Y^b(x, t; z)$ when $z \rightarrow b$ and $Y^{-b}(x, t; z)$ when $z \rightarrow -b$ which correspond to the Bessel model $U_{Bes}(z)$. And local matrix functions $Y^b(x, t; z)$ and $Y^{-b}(x, t; z)$ satisfy the following properties:

$$\begin{aligned} Y^b(x, t; z) (Y^o(x, t; z))^{-1} &= \mathbb{I} + \mathcal{O}(t^{-1}), \quad t \rightarrow +\infty, z \in \partial\delta^b \setminus \gamma_A, \\ Y^{-b}(x, t; z) (Y^o(x, t; z))^{-1} &= \mathbb{I} + \mathcal{O}(t^{-1}), \quad t \rightarrow +\infty, z \in \partial\delta^{-b} \setminus \gamma_A. \end{aligned} \quad (4.146)$$

Then we construct a matrix function as follows:

$$Y_{3a_1}(x, t; z) = \begin{cases} Y^o(x; z), & z \in \mathbb{C} \setminus (\delta^{a_1} \cup \delta^b \cup \delta^{-a_1} \cup \delta^{-b}), \\ Y^{\pm a_1}(x, t; z), & z \in \delta^{\pm a_1}, \\ Y^{\pm b}(x, t; z), & z \in \delta^{\pm b}, \end{cases} \quad (4.147)$$

which has the following jump condition:

$$Y_{3a_1+}(x, t; z) = Y_{3a_1-}(x, t; z)V_{8a_1}(x, t; z). \quad (4.148)$$

4.2.4 The small norm Riemann-Hilbert problem

Let the error matrix function $E(x, t; z)$ as follow:

$$E(x, t; z) = Y_{2a_1}(x, t; z)Y_{3a_1}^{-1}(x, t; z). \quad (4.149)$$

Then the error matrix function $E(x, t; z)$ satisfies the following Riemann-Hilbert problem.

RHP 17. Find a 2×2 matrix function $E(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $E(x, t; z)$ is analytic in $\mathbb{C} \setminus \gamma_3$ where $\gamma_3 := (a, a_1) \cup (-a_1, -a) \cup O_1 \cup O_2 \cup \partial\delta^{a_1} \cup \partial\delta^b \cup \partial\delta^{-a_1} \cup \partial\delta^{-b} \setminus (\delta^{a_1} \cup \delta^b \cup \delta^{-a_1} \cup \delta^{-b})$ and takes continuous boundary values on γ_3 .
- *Jump condition:* The boundary values on the jump contour are defined as

$$E_+(x, t; z) = E_-(x, t; z)V_E(x, t; z), \quad z \in \gamma_3, \quad (4.150)$$

where

$$V_E(x, t; z) = Y_{3a_1-}(x, t; z)V_{6a_1}(x, t; z)V_{8a_1}^{-1}(x, t; z)Y_{3a_1-}^{-1}(x, t; z), \quad (4.151)$$

which satisfies the following properties:

$$V_E(x, t; z) = \begin{cases} \mathbb{I} + \mathcal{O}(e^{-ct}), & z \in (a, a_1) \cup (-a_1, -a) \cup O_1 \cup O_2 \setminus (\bar{\delta}^{a_1} \cup \bar{\delta}^b \cup \bar{\delta}^{-a_1} \cup \bar{\delta}^{-b}), \\ \mathbb{I} + \mathcal{O}\left(\frac{1}{t}\right), & z \in \partial\delta^{a_1} \cup \partial\delta^b \cup \partial\delta^{-a_1} \cup \partial\delta^{-b}, \end{cases} \quad (4.152)$$

with c being a suitable positive real parameter.

- *Normalization:* $E(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

According to Riemann-Hilbert problem 17, we obtain

$$E(x, t; z) = \mathbb{I} + \frac{E_1(x, t)}{tz} + \mathcal{O}(z^{-2}). \quad (4.153)$$

Based on Ref. [23], one has the following proposition:

Proposition 4.2. As $t \rightarrow +\infty$, the potential function $q(x, t)$ of the complex mKdV equation has the following asymptotic behaviour:

$$q(x, t) = (b - a_1) \frac{\vartheta_3(0; \tau) \vartheta_3\left(\frac{tm_{2a_1} + m_{1a_1}}{2\pi i} + \frac{1}{2}; \tau\right)}{\vartheta_3\left(\frac{1}{2}; \tau\right) \vartheta_3\left(\frac{tm_{2a_1} + m_{1a_1}}{2\pi i}; \tau\right)} + \mathcal{O}(t^{-1}), \quad t \rightarrow +\infty. \quad (4.154)$$

Proof. According to Eqs. (4.114),(4.125) and (4.153), we have

$$\begin{aligned}
Y_{a_1}(x, t; z) &= Y_{1a_1}(x, t; z)f_{a_1}(z)^{-\sigma_3} e^{-tg_{a_1}(z)\sigma_3} \\
&= Y_{2a_1}(x, t; z)f_{a_1}(z)^{-\sigma_3} e^{-tg_{a_1}(z)\sigma_3} \\
&= E(x, t; z)Y_{3a_1}(x, t; z)f_{a_1}(z)^{-\sigma_3} e^{-tg_{a_1}(z)\sigma_3} \\
&= \left(\mathbb{I} + \frac{E_1(x, t)}{tz} + \mathcal{O}(z^{-2}) \right) Y_{3a_1}(x, t; z)f_{a_1}(z)^{-\sigma_3} e^{-tg_{a_1}(z)\sigma_3},
\end{aligned}$$

which further leads to

$$\begin{aligned}
Y_{a_1,12}(x, t; z) &= \left(Y_{3a_1,12}(x, t; z) + \frac{E_{1,12}(x, t)}{tz} + \mathcal{O}(z^{-2}) \right) f_{a_1}(z)e^{tg_{a_1}(z)} \\
&= \left(Y_{12}^o(x, t; z) + \frac{E_{1,12}(x, t)}{tz} + \mathcal{O}(z^{-2}) \right) f_{a_1}(z)e^{tg_{a_1}(z)} \\
&= \frac{1}{2} \left(\psi_{12}(x, t; z) + \frac{E_{1,12}(x, t)}{tz} + \mathcal{O}(z^{-2}) \right) f_{a_1}(z)e^{tg_{a_1}(z)}.
\end{aligned} \tag{4.155}$$

Note that

$$\begin{aligned}
f_{a_1}(z) &= 1 + \frac{f_1}{z} + \mathcal{O}(z^{-2}), \\
e^{tg_{a_1}(z)} &\rightarrow 1, \quad z \rightarrow \infty, \\
m_{5a_1}(z) &= 1 + \frac{a_1 - b}{2z} + \mathcal{O}(z^{-2}).
\end{aligned} \tag{4.156}$$

According to Eq. (4.113), namely, $q(x, t) = -2 \lim_{z \rightarrow \infty} zY_{a_1,12}(x, t; z)$, one can find that this Theorem holds. \square

4.3 The case $\xi = x/t < \xi_c$

When $\xi < \xi_c$, we need to redefine the function $g(z)$, which satisfies the following properties [23]:

- Analyticity: $g_a(z)$ is analytic in $\mathbb{C} \setminus (-b, b)$ and takes continuous boundary values on $(-b, b)$.
- Jump condition: The boundary values on the jump contour are defined as

$$\begin{aligned}
g_{a+}(z) + g_{a-}(z) &= 2\xi z - 8z^3, \quad z \in \gamma_1 \cup \gamma_2, \\
g_{a+}(z) - g_{a-}(z) &= m_{2a}, \quad z \in (-a, a),
\end{aligned} \tag{4.157}$$

where

$$m_{2a} = \frac{\pi ib(\xi - 2(a^2 + b^2))}{K_1(m)}.$$

- Normalization: $g_a(z) = \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

Then, we have

$$\begin{aligned}
\operatorname{Re}(g_a(z) + 4z^3 - \xi z) &> 0, \quad z \in O_1 \setminus \{a, b\}, \\
\operatorname{Re}(g_a(z) + 4z^3 - \xi z) &< 0, \quad z \in O_2 \setminus \{-a, -b\}.
\end{aligned} \tag{4.158}$$

Based on Ref. [23], one has

Proposition 4.3. *As $t \rightarrow +\infty, \xi < \xi_c$, we have the long-time asymptotic behaviour of the soliton gas of the potential function $q(x, t)$ of the complex mKdV equation:*

$$q(x, t) = (b - a) \frac{\vartheta_3(0; \tau) \vartheta_3\left(\frac{tm_{2a} + m_1}{2\pi i} + \frac{1}{2}; \tau\right)}{\vartheta_3\left(\frac{1}{2}; \tau\right) \vartheta_3\left(\frac{tm_{2a} + m_1}{2\pi i}; \tau\right)} + \mathcal{O}(t^{-1}), \quad t \rightarrow +\infty. \tag{4.159}$$




Figure 8: The domains Ω_1 and Ω_1^* : (a) The parameters are $s_1 = 2i$, $s_2 = \frac{1}{4} + \frac{1}{4}i$, $s_3 = \frac{3}{4}$, $\ell = 1$, (b) The parameters are $s_1 = 2 + 2i$, $s_2 = \frac{1}{2}$, $s_3 = \frac{3}{4}$, $\ell = 2$.

5 Soliton gas in the quadrature domains

Bertola *et al* studied the soliton shielding in the quadrature domains of the focusing nonlinear Schrödinger equation [25]. Similarly, we consider the case that the discrete spectra z_j ($j = 1, \dots, N$) with the norming constants c_j ($j = 1, \dots, N$) filling uniformly compact domain Ω_1 of the complex upper half space \mathbb{C}_+ , for example [25],

$$\Omega_1 := \{z \mid |(z - s_1)^\ell - s_2| < s_3\}, \quad \Omega_1 \subset \mathbb{C}_+, \quad (5.160)$$

where $\ell \in \mathbb{N}_+$, $s_1 \in \mathbb{C}_+$ and $|s_2|, s_3$ are sufficiently small real constants. The boundary of Ω_1^* which is complex conjugate domain of Ω_1 is described by (see Fig. 8)

$$z^* = \left(s_2^* + \frac{s_3^2}{(z - s_1)^\ell - s_2} \right)^{1/\ell} + s_1^*, \quad z \in \Omega_1. \quad (5.161)$$

We assume that the norming constants c_j ($j = 1, \dots, N$) have the following forms:

$$c_j = \frac{|\Omega_1| r(z_j, z_j^*)}{N\pi}. \quad (5.162)$$

where $|\Omega_1|$ means the area of the domain Ω_1 and $r(z, z^*)$ is a smooth function with respect to variables z and z^* . Then we construct the following Riemann-Hilbert problem.

RHP 18. Find a 2×2 matrix function $M_0(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $M_0(x, t; z)$ is meromorphic in \mathbb{C} , with simple poles at $K_1 := \{z_j, z_j^*\}_{j=1}^N$ with $z_j \in \Omega_1$.
- *Normalization:* $M_0(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$;
- *Residue conditions:* $M_0(x, t; z)$ has simple poles at each point in K_1 with:

$$\begin{aligned} \text{Res}_{z=z_j} M_0(x, t; z) &= \lim_{z \rightarrow z_j} M_0(x, t; z) \begin{bmatrix} 0 & 0 \\ c_j e^{2i(zx + (2\alpha z^2 + 4\beta z^3)t)} & 0 \end{bmatrix}, \\ \text{Res}_{z=z_j^*} M_0(x, t; z) &= \lim_{z \rightarrow z_j^*} M_0(x, t; z) \begin{bmatrix} 0 & -c_j^* e^{-2i(zx + (2\alpha z^2 + 4\beta z^3)t)} \\ 0 & 0 \end{bmatrix}. \end{aligned} \quad (5.163)$$

According to Eq. (1.27), we can recover $q(x, t)$ by the following formula:

$$q(x, t) = 2i \lim_{z \rightarrow \infty} (zM_0(x, t; z))_{12}. \quad (5.164)$$

We define a closed curve Γ_{1+} with a very small radius encircling the poles $\{z_j\}_{j=1}^N$ counterclockwise in the upper half plane \mathbb{C}_+ , and a closed curve Γ_{1-} with a very small radius encircling the poles $\{z_j^*\}_{j=1}^N$ counterclockwise in the lower half plane \mathbb{C}_- . Make the following transformation:

$$M_1(x, t; z) = \begin{cases} M_0(x, t; z) \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^N \frac{c_j e^{2i(xz+2\alpha tz^2+4\beta tz^3)}}{z-z_j} & 1 \end{bmatrix}, & z \text{ within } \Gamma_+, \\ M_0(x, t; z) \begin{bmatrix} 1 & \sum_{j=1}^N \frac{c_j^* e^{-2i(xz+2\alpha tz^2+4\beta tz^3)}}{z-z_j^*} \\ 0 & 1 \end{bmatrix}, & z \text{ within } \Gamma_-, \\ M_0(x, t; z), & \text{otherwise.} \end{cases} \quad (5.165)$$

Then matrix function $M_1(x, t; z)$ satisfies the following Riemann-Hilbert problem.

RHP 19. Find a 2×2 matrix function $M_1(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $M_1(x, t; z)$ is analytic in $\mathbb{C} \setminus (\Gamma_{1+} \cup \Gamma_{1-})$ and takes continuous boundary values on $\Gamma_{1+} \cup \Gamma_{1-}$.
- *Jump condition:* The boundary values on the jump contour $\Gamma_{1+} \cup \Gamma_{1-}$ are defined as

$$M_{1+}(x, t; z) = M_{1-}(x, t; z)J_1(x, t; z), \quad z \in \Gamma_{1+} \cup \Gamma_{1-}, \quad (5.166)$$

where

$$J_1(x, t; z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^N \frac{c_j e^{2i(xz_j+2\alpha tz_j^2+4\beta tz_j^3)}}{z-z_j} & 1 \end{bmatrix}, & z \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & \sum_{j=1}^N \frac{c_j^* e^{-2i(xz_j^*+2\alpha tz_j^{*2}+4\beta tz_j^{*3})}}{z-z_j^*} \\ 0 & 1 \end{bmatrix}, & z \in \Gamma_{1-}; \end{cases} \quad (5.167)$$

- *Normalization:* $M_1(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

According to Eq. (5.164), we can recover $q(x, t)$ by the following formula:

$$q(x, t) = 2i \lim_{z \rightarrow \infty} (zM_1(x, t; z))_{12}. \quad (5.168)$$

Based on Ref. [25], one has the following lemma:

Lemma 5.1. For any open set B_+ containing the domain Ω_1 , and any open set B_- containing the domain Ω_1^* , the following identities hold:

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j e^{2i(xz_j+2\alpha tz_j^2+4\beta tz_j^3)}}{z-z_j} = \iint_{\Omega_1} \frac{r(\lambda, \lambda^*)}{2\pi i(z-\lambda)} e^{2i(\lambda x+2\alpha \lambda^2 t+4\beta \lambda^3 t)} d\lambda^* \wedge d\lambda, \quad (5.169)$$

uniformly for all $\mathbb{C} \setminus B_+$,

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j^* e^{-2i(xz_j + 2\alpha t z_j^2 + 4\beta t z_j^3)}}{z - z_j^*} = \iint_{\Omega_1^*} \frac{r^*(\lambda, \lambda^*)}{2\pi i(z - \lambda)} e^{-2i(\lambda x + 2\alpha \lambda^2 t + 4\beta \lambda^3 t)} d\lambda^* \wedge d\lambda, \quad (5.170)$$

uniformly for all $\mathbb{C} \setminus B_-$. The boundary $\partial\Omega_1$ and $\partial\Omega_1^*$ are both counterclockwise.

According to Lemma 5.1, we can obtain the Riemann-Hilbert problem of matrix function $M_1(x, t; z)$ as $N \rightarrow \infty$.

RHP 20. Find a 2×2 matrix function $M_2(x, t; z)$ that satisfies the following properties:

- *Analyticity:* $M_2(x, t; z)$ is analytic in $\mathbb{C} \setminus (\Gamma_{1+} \cup \Gamma_{1-})$ and takes continuous boundary values on $\Gamma_{1+} \cup \Gamma_{1-}$.
- *Jump condition:* The boundary values on the jump contour $\Gamma_{1+} \cup \Gamma_{1-}$ are defined as

$$M_{2+}(x, t; z) = M_{2-}(x, t; z) J_2(x, t; z), \quad z \in \Gamma_{1+} \cup \Gamma_{1-}, \quad (5.171)$$

where

$$J_2(x, t; z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ \iint_{\Omega_1} \frac{r(\lambda, \lambda^*) e^{2i(x\lambda + 2\alpha t \lambda^2 + 4\beta t \lambda^3)}}{2\pi i(\lambda - z)} d\lambda^* \wedge d\lambda & 1 \end{bmatrix}, & z \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & \iint_{\Omega_1^*} \frac{r^*(\lambda, \lambda^*) e^{-2i(x\lambda + 2\alpha t \lambda^2 + 4\beta t \lambda^3)}}{2\pi i(z - \lambda)} d\lambda^* \wedge d\lambda \\ 0 & 1 \end{bmatrix}, & z \in \Gamma_{1-}; \end{cases} \quad (5.172)$$

- *Normalization:* $M_1(x, t; z) = \mathbb{I} + \mathcal{O}(z^{-1})$, $z \rightarrow \infty$.

Let the function $R(z, z^*)$ satisfy the following property:

$$\bar{\partial} R(z, z^*) = r(z, z^*). \quad (5.173)$$

Then the jump matrix $J_2(x, t; z)$ can be written as:

$$J_2(x, t; z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ \int_{\partial\Omega_1} \frac{R(\lambda, \lambda^*) e^{2i(x\lambda + 2\alpha t \lambda^2 + 4\beta t \lambda^3)}}{2\pi i(\lambda - z)} d\lambda & 1 \end{bmatrix}, & z \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & \int_{\partial\Omega_1^*} \frac{R^*(\lambda, \lambda^*) e^{-2i(x\lambda + 2\alpha t \lambda^2 + 4\beta t \lambda^3)}}{2\pi i(z - \lambda)} d\lambda \\ 0 & 1 \end{bmatrix}, & z \in \Gamma_{1-}; \end{cases} \quad (5.174)$$

Based on Ref. [25], one has the following proposition:

Proposition 5.1. If we assume that the function $R(z, z^*) = R(z, s_1^* + \left(s_2^* + \frac{s_3^2}{(z - s_1)^\ell - s_2} \right)^{\frac{1}{\ell}})$ in Ω_1 has n distinct poles $\lambda_j, j = 1, \dots, n$, that is,

$$R\left(z, s_1^* + \left(s_2^* + \frac{s_3^2}{(z - s_1)^\ell - s_2} \right)^{\frac{1}{\ell}}\right) = \widehat{R}(z) \prod_{j=1}^n \frac{1}{z - \lambda_j}, \quad (5.175)$$

where $\widehat{R}(z)$ is some analytic function in Ω_1 , then the solution of the Riemann-Hilbert problem 20 can generate an n -soliton $q_n(x, t)$ with discrete spectrums $\lambda_j, j = 1, \dots, n$ with the norming constants $c_j = \frac{R(\lambda_j)}{\prod_{k \neq j} (\lambda_j - \lambda_k)}, j = 1, \dots, n$.

Proof. Substituting Eq. (5.175) into the integral in Eq. (5.174), we obtain

$$\begin{aligned} & \int_{\partial\Omega_1} \frac{R(\lambda, \lambda^*) e^{2i(x\lambda + 2\alpha t\lambda^2 + 4\beta t\lambda^3)}}{2\pi i(\lambda - z)} d\lambda \\ &= \int_{\partial\Omega_1} \frac{R\left(\lambda, s_1^* + \left(s_2^* + \frac{s_3^2}{(\lambda - s_1)^\ell - s_2}\right)^{\frac{1}{\ell}}\right)}{2\pi i(\lambda - z)} e^{2i(x\lambda + 2\alpha t\lambda^2 + 4\beta t\lambda^3)} d\lambda \\ &= \int_{\partial\Omega_1} \frac{\widehat{R}(\lambda) e^{2i(x\lambda + 2\alpha t\lambda^2 + 4\beta t\lambda^3)}}{2\pi i(z - \lambda) \prod_{j=1}^n (\lambda - \lambda_j)} d\lambda \\ &= \sum_{j=1}^n \frac{\widehat{R}(\lambda_j) e^{2i(x\lambda_j + 2\alpha t\lambda_j^2 + 4\beta t\lambda_j^3)}}{(z - \lambda_j) \prod_{k \neq j} (\lambda_j - \lambda_k)}, \quad z \notin \Omega_1. \end{aligned}$$

Let the norming constants be $c_j = \frac{\widehat{R}(\lambda_j)}{\prod_{k \neq j} (\lambda_j - \lambda_k)}, j = 1, \dots, n$, then, similarly to the Riemann-Hilbert problem 2, the solution of Riemann-Hilbert problem 20 leads to an n -soliton $q_n(x, t)$ of the Hirota equation. Thus the proof is completed. \square

6 Conclusions and discussions

In conclusion, we have used the idea of Refs. [23–25] to investigate the asymptotic behaviors of a soliton gas (the limit $N \rightarrow \infty$ of N -soliton solutions) of the Hirota equation including the complex modified KdV equation with the aid of the Riemann-Hilbert problems with pure imaginary discrete spectra restricted in an interval. We show that that this soliton gas tends slowly to the Jacobian elliptic wave solution (zero exponentially quickly) when $x \rightarrow -\infty$ ($x \rightarrow +\infty$). Moreover, we also give the long-time asymptotics of the soliton gas under the different velocity conditions: $x/t > 4\beta b^2$, $\xi_c < x/t < 4\beta b^2$, $x/t < \xi_c$. Finally, we analyze the property of the soliton gas with the discrete spectra filling uniformly a quadrature domain.

Therefore are some challenging issues, for example, i) when $\xi = x/t < 4\beta b^2$, how will one consider the long-time asymptotics of the potential $q(x, t)$ of the Hirota equation for the case $\alpha\beta \neq 0$? since there is the terms $iR(z)e^{\pm 2zt(\xi - 4\beta z^2 + 2iaz)}$ with the factor $e^{\pm 4iaz^2 t}$ in the jump matrices given by Eq. (4.107); ii) how will one study the asymptotics of soliton gas for the case of the considered discrete spectral intervals localized in the non-imaginary axes?

Acknowledgments

This work was supported by the National Natural Science Foundation of China (Grant No. 11925108).

References

- [1] N. J. Zabusky, M. D. Kruskal, Interaction of “solitons” in a collisionless plasma and the recurrence of initial states, Phys. Rev. Lett. 15, 240 (1965).
- [2] C. S. Gardner, J.M. Greene, M.D. Kruskal, and R.M. Miura, Method for solving the Korteweg-de Vries equation, Phys. Rev Lett. 19, 1095-1097 (1967).

- [3] M.J. Ablowitz, D.J. Kaup, A.C. Newell, and H. Segur, The inverse scattering transform. Fourier analysis for nonlinear problems, *Stud. Appl. Math.* 53, 249-315 (1974).
- [4] P.D. Lax, Integrals of nonlinear evolution equations and solitary waves, *Comm. Pure Appl. Math.* 21, 467-490 (1968).
- [5] S.P. Novikov, S.V. Manakov, L.P. Pitaevskii, and V.E. Zakharov, *Theory of Solitons: The Inverse Scattering-Transform* (Plenum, New York, 1984).
- [6] A.C. Newell, *Solitons in Mathematics and Physics* (SIAM, Philadelphia, 1985).
- [7] L. D. Faddeev, L. A. Takhtajan, *Hamiltonian Methods in the Theory of Solitons* (Springer-Verlag, Berlin, 1987).
- [8] M.J. Ablowitz and P.A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering* (Cambridge University Press, Cambridge, 1991).
- [9] A.C. Scott, *Nonlinear Science: Emergence and Dynamics of Coherent Structures* (Oxford University Press, Oxford, 2003).
- [10] G. P. Agrawal, *Nonlinear Fiber Optics* (5th edn.) (Academic Press, New York, 2012).
- [11] L.F. Mollenauer, R.H. Stolen, J. P. Gordon, Experimental observation of picosecond pulse narrowing and solitons in optical fibers, *Phys. Rev. Lett.* 45, 1095-1112 (1980).
- [12] Y. S. Kivshar and G. P. Agrawal, *Optical Solitons: from Fibers to Photonic Crystals* (Academic Press, New York, 2013).
- [13] B. A. Malomed, D. Mihalache, F. Wise, and L. Torner, Spatiotemporal optical solitons, *J. Opt. B: Quantum Semiclass. Opt.* 7, R53 (2005).
- [14] L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation and Superfluidity* (Oxford University Press, Oxford, 2016).
- [15] V.E. Zakharov, Kinetic equation for solitons. *Sov. Phys. JETP* 33, 538-541 (1971).
- [16] G.A. El, A.M. Kamchantov, Kinetic equation for a dense soliton gas, *Phys. Rev. Lett.* 95, 204101 (2005).
- [17] G.A. El, A.M. Kamchatnov, M.V. Pavlov, S.A. Zykov, Kinetic equation for a soliton gas and its hydrodynamic reductions, *J. Nonlinear Sci.* 21, 151-191 (2011).
- [18] G.A. El, Critical density of a soliton gas, *Chaos* 26, 023105 (2016).
- [19] G.A. El, A. Tovbis, Spectral theory of soliton and breather gases for the focusing nonlinear Schrödinger equation, *Phys. Rev. E* 101, 052207 (2020).
- [20] P. Suret, A. Tikan, F. Bonnefoy, F. Copie, G. Ducrozet, A. Gelash, G. Prabhudesai, G. Michel, A. Cazaubiel, E. Falcon, G. A. El, S. Randoux, Nonlinear spectral synthesis of soliton gas in deep-water surface gravity waves, *Phys. Rev. Lett.* 125, 264101 (2020).
- [21] T. Congy, G. El, G. Roberti, Soliton gas in bidirectional dispersive hydrodynamics, *Phys. Rev. E* 103, 042201 (2021).
- [22] E. G. Shurgalina, E. N. Pelinovsky, Nonlinear dynamics of a soliton gas: Modified Korteweg-de Vries equation framework, *Phys. Lett. A* 380, 2049-2053 (2016).
- [23] M. Girotti, T. Grava, R. Jenkins, K. T. R. McLaughlin, Rigorous asymptotics of a KdV soliton gas, *Commun. Math. Phys.* 384, 733-784 (2021).

- [24] M. Girotti, T. Grava, R. Jenkins, K. T. R. McLaughlin, A. Minakov, Soliton versus the gas: Fredholm determinants, analysis, and the rapid oscillations behind the kinetic equation, *Comm. Pure Appl. Math.* 76, 3233-3299 (2023).
- [25] M. Bertola, T. Grava, G. Orsatti, Soliton shielding of the focusing Nonlinear Schrödinger Equation, *Phys. Rev. Lett.* 130, 127201 (2023).
- [26] G. D. Lyng, P. D. Miller, The N -soliton of the focusing nonlinear Schrödinger equation for N large, *Comm. Pure Appl. Math.* 60, 951-1026 (2007).
- [27] D. Bilman, R. Buckingham, Large-order asymptotics for multiple-pole solitons of the focusing nonlinear Schrödinger equation. *J. Nonlinear Sci.* 29, 2185-2229 (2019).
- [28] D. Bilman, L. Ling, P. D. Miller, Extreme superposition: rogue waves of infinite order and the Painlevé-III hierarchy, *Duke Math. J.* 169, 671-760 (2020).
- [29] D. Bilman, R. Buckingham, D. S. Wang, Far-field asymptotics for multiple-pole solitons in the large-order limit, *J. Differ. Equ.* 297, 320 (2021).
- [30] D. Bilman, P. D. Miller, A robust inverse scattering transform for the focusing nonlinear Schrödinger equation, *Comm. Pure Appl. Math.* 72, 1722 (2019).
- [31] X. Zhou, The Riemann-Hilbert problem and inverse scattering, *SIAM J. Math. Anal.* 20, 966 (1989).
- [32] P. Deift, X. Zhou, A steepest descent method for oscillatory Riemann-Hilbert problems, *Asymptotics for the MKdV equation*, *Ann. of Math.* 137, 295 (1993).
- [33] K.T.R. McLaughlin, P.D. Miller, The $\bar{\partial}$ -steepest descent method and the asymptotic behavior of polynomials orthogonal on the unit circle with fixed and exponentially varying non-analytic weights, *Int. Math. Res. Not.* 2006, 48673 (2006).
- [34] K.T.R. McLaughlin, P.D. Miller, The $\bar{\partial}$ -steepest descent method for orthogonal polynomials on the real line with varying weights, *Int. Math. Res. Not.* 075 (2008).
- [35] R. Beals, R. Coifman, Scattering and inverse scattering for first order systems, *Comm. Pure Appl. Math.* 37, 39-90 (1984).
- [36] R. Beals, P. Deift, C. Tomei, *Direct and Inverse Scattering on the Line* (American Mathematical Society, Providence, 1988).
- [37] T. Trogdon, S. Olver, *Riemann-Hilbert Problems, Their Numerical Solution, and the Computation of Nonlinear Special Functions* (SIAM, Philadelphia, 2016).
- [38] P. Deift, X. Zhou, Long-time asymptotics for solutions of the NLS equation with initial data in a weighted Sobolev space, *Comm. Pure Appl. Math.* 56, 1029-1077 (2003).
- [39] A. H. Vartanian, Long-time asymptotics of solutions to the Cauchy problem for the defocusing nonlinear Schrödinger equation with finite-density initial data. II. Dark solitons on continua. *Math. Phys. Anal. Geom.* 5, 319-413 (2002).
- [40] A. Boutet de Monvel, A. Its, V. Kotlyarov, Long-time asymptotics for the focusing NLS equation with time-periodic boundary condition on the half-line, *Commun. Math. Phys.* 290, 479-522 (2009).
- [41] K. Grunert, G. Teschl, Long-time asymptotics for the Korteweg de Vries equation via nonlinear steepest descent, *Math. Phys. Anal. Geom.* 12, 287-324 (2009).
- [42] A. Boutet de Monvel, J. Lenells, D. Shepelsky, Long-time asymptotics for the Degasperis-Procesi equation on the half-line, *Ann. Inst. Fourier* 69, 171-230 (2019).

- [43] J. Xu, E. Fan, Long-time asymptotics for the Fokas-Lenells equation with decaying initial value problem: without solitons, *J. Differ. Equ.* 259, 1098-1148 (2015).
- [44] H. Liu, X. Geng, B. Xue, The Deift-Zhou steepest descent method to long-time asymptotics for the Sasa-Satsuma equation, *J. Differ. Equ.* 265, 5984-6008 (2018).
- [45] G. Biondini, S. Li, D. Mantzavinos, Long-time asymptotics for the focusing nonlinear Schrödinger equation with nonzero boundary conditions in the presence of a discrete spectrum, *Commun. Math. Phys.* 382, 1495-1577 (2021).
- [46] S. Cuccagna, R. Jenkins, On the asymptotic stability of N-soliton solutions of the defocusing nonlinear Schrödinger equation, *Commun. Math. Phys.* 343, 921-969 (2016).
- [47] M. Borghese, R. Jenkins, K.T.R. McLaughlin, Long-time asymptotic behavior of the focusing nonlinear Schrödinger equation, *Ann. Inst. Henri Poincaré Anal.* 35, 887-920 (2018).
- [48] R. Jenkins, J. Liu, P. Perry, C. Sulem, Soliton resolution for the derivative nonlinear Schrödinger equation, *Commun. Math. Phys.* 363,1003-1049 (2018).
- [49] Y. Yang, E. Fan, Soliton resolution for the short-pulse equation, *J. Differ. Equ.* 280, 644-689 (2021).
- [50] Y. Yang, E. Fan, On the long-time asymptotics of the modified Camassa-Holm equation in space-time solitonic regions. *Adv. Math.* 402, 108340 (2022).
- [51] Z. Wang, E. Fan, The defocusing nonlinear Schrödinger equation with a nonzero background:: Painlevé asymptotics in two transition regions, *Commun. Math. Phys.* 402, 2879-2930 (2023).
- [52] Z. Li, S. Tian, J. Yang, On the asymptotic stability of N-soliton solution for the short pulse equation with weighted Sobolev initial data, *J. Differ. Equ.* 377, 121-187 (2023).
- [53] Y. Kodama, Optical solitons in a monomode fiber, *J. Stat. Phys.* 39, 597 (1985).
- [54] Y. Kodama and A. Hasegawa, Nonlinear pulse propagation in a monomode dielectric guide, *IEEE J. Quantum Electron.* 23, 510 (1987).
- [55] Z. Yan and C. Dai, Optical rogue waves in the generalized inhomogeneous higher-order nonlinear Schrödinger equation with modulating coefficients, *J. Opt.* 15, 064012 (2013).
- [56] N. Sasa, J. Satsuma, New-type of soliton solutions for a high-order nonlinear Schrödinger equation, *J. Phys. Soc. Jpn.* 60, 409-417 (1990).
- [57] V.E. Zakharov, A.B. Shabat, Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media, *Sov. Phys. JETP* 34, 62-69 (1972).
- [58] M. Wadati, K. Konno, Y. H. Ichikawa, A generalization of inverse scattering method, *J. Phys. Soc. Jpn.* 46, 1965 (1979).
- [59] D. J. Kaup, A. C. Newell, An exact solution for a derivative nonlinear Schrödinger equation, *J. Math. Phys.* 19, 798-801 (1978).
- [60] H.H. Chen, Y. C. Lee, C.S. Liu, Integrability of nonlinear Hamiltonian systems by inverse scattering method, *Phys. Scr.* 20, 490 (1979).
- [61] A. Kundu, Landau-Lifshitz and higher-order nonlinear systems gauge generated from nonlinear Schrödinger-type equations, *J. Math. Phys.* 25, 3433 (1984).
- [62] R. Hirota, Exact envelope-soliton solutions of a nonlinear wave equation, *J. Math. Phys.* 14, 805 (1973).

- [63] A. Ankiewicz, J. M. Soto-Crespo, and N. Akhmediev, Rogue waves and rational solutions of the Hirota equation, *Phys. Rev. E* 81, 046602 (2010).
- [64] A. Chowdury, D. J. Kedziora, A. Ankiewicz, and N. Akhmediev, Soliton solutions of an integrable nonlinear Schrödinger equation with quintic terms, *Phys. Rev. E* 90, 032922 (2014).
- [65] Y. Tao, J. He, Multisolitons, breathers, and rogue waves for the Hirota equation generated by the Darboux transformation, *Phys. Rev. E* 85, 026601 (2012).
- [66] Y. Yang, Z. Yan, B. A. Malomed, Rogue waves, rational solitons, and modulational instability in an integrable fifth-order nonlinear Schrödinger equation, *Chaos* 25, 103112 (2015).
- [67] S. Chen, Z. Yan, The Hirota equation: Darboux transform of the Riemann-Hilbert problem and higher-order rogue waves, *Appl. Math. Lett.* 95, 65-71 (2019).
- [68] G. Zhang, S. Chen, Z. Yan, Focusing and defocusing Hirota equations with non-zero boundary conditions: Inverse scattering transforms and soliton solutions, *Commun. Nonlinear Sci. Numer. Simulat.* 80 (2020) 104927.
- [69] L. Huang, J. Xu, E. Fan, Long-time asymptotic for the Hirota equation via nonlinear steepest descent method, *Nonlinear Anal. Real World Appl.* 26, 229-262 (2015).
- [70] B. Guo, N. Liu, Y. Wang, Long-time asymptotics for the Hirota equation on the half-line, *Nonlinear Anal.* 174, 118-140 (2018).
- [71] S. Chen, Z. Yan, B. Guo, Long-time asymptotics for the focusing Hirota equation with non-zero boundary conditions at infinity via the Deift-Zhou approach, *Math. Phys. Anal. Geom.* 24, 17 (2021).
- [72] Y. Xiao, B. Guo, Z. Wang, Nonlinear stability of multi-solitons for the Hirota equation, *J. Differential Equa.* 342, 369-417 (2023).
- [73] A. B. J. Kuijlaars, K. R. McLaughlin, W. Van Assche, M. Vanlessen, The Riemann-Hilbert approach to strong asymptotics for orthogonal polynomials on $[-1,1]$, *Adv. Math.* 188, 337-398 (2004).
- [74] P. Deift, T. Kriecherbauer, K. T. R. McLaughlin, S. Venakides, X. Zhou, Strong asymptotics of orthogonal polynomials with respect to exponential weights, *Comm. Pure Appl. Math.* 52, 1491-1552 (1999).
- [75] D. F. Lawden, *Elliptic Functions and Applications* (Springer, Berlin, 1989).
- [76] I. Egorova, Z. Gladka, V. Kotlyarov, G. Teschl, Long-time asymptotics for the Korteweg-de Vries equation with step-like initial data, *Nonlinearity* 26, 1839-1864 (2013).