

The Assouad dimension of Kakeya sets in \mathbb{R}^3

Hong Wang*

Joshua Zahl †

Abstract

This paper studies the structure of Kakeya sets in \mathbb{R}^3 . We show that for every Kakeya set $K \subset \mathbb{R}^3$, there exist well-separated scales $0 < \delta < \rho \leq 1$ so that the δ neighborhood of K is almost as large as the ρ neighborhood of K . As a consequence, every Kakeya set in \mathbb{R}^3 has Assouad dimension 3 and every Ahlfors–David regular Kakeya set in \mathbb{R}^3 has Hausdorff dimension 3. We also show that every Kakeya set in \mathbb{R}^3 that has “stably equal” Hausdorff and packing dimension (this is a new notion, which is introduced to avoid certain obvious obstructions) must have Hausdorff dimension 3.

The above results follow from certain multi-scale structure theorems for arrangements of tubes and rectangular prisms in three dimensions, and a mild generalization of the sticky Kakeya theorem previously proved by the authors.

1 Introduction

A Kakeya set is a compact subset of \mathbb{R}^n that contains a unit line segment pointing in every direction. The Kakeya set conjecture asserts that every Kakeya set in \mathbb{R}^n has Minkowski and Hausdorff dimension n . This conjecture is proved in the plane [1, 2], and is open in three and higher dimensions. See [9, 18] for a survey of progress on the Kakeya conjecture.

In this paper, we study the structure of Kakeya sets in \mathbb{R}^3 . We show that for every Kakeya set $K \subset \mathbb{R}^3$, there exist well-separated scales $0 < \delta < \rho \leq 1$ so that the δ neighborhood of K is almost as large as the ρ neighborhood of K . The precise statement is given in Theorem 1.5 below. As a consequence, we prove several weaker variants of the Kakeya set conjecture: we prove that every Kakeya set in \mathbb{R}^3 has Assouad dimension 3, and we prove that every Kakeya set in \mathbb{R}^3 with “stably equal” Hausdorff and packing dimension (see Definition 1.3 below) must have Hausdorff and packing dimension 3. In order to explain these statements precisely, we require the following definitions.

Definition 1.1. *Let $E \subset \mathbb{R}^n$ be non-empty. The Assouad dimension of E , denoted $\dim_A(E)$, is the infimum of all $\beta \geq 0$ for which there exist positive constants C and r_0 so that for all $0 < \rho < r \leq r_0$, we have*

$$\sup_{x \in \mathbb{R}^n} \mathcal{E}_\rho(E \cap B(x, r)) \leq C(r/\rho)^\beta.$$

In the above, $\mathcal{E}_\rho(X)$ denotes the ρ -covering number of X .

The Assouad dimension is always at least as large as the Hausdorff and Minkowski dimensions, i.e. if $E \subset \mathbb{R}^n$ is bounded then we have $0 \leq \dim_H(E) \leq \underline{\dim}_M(E) \leq \overline{\dim}_M(E) \leq \dim_A(E) \leq n$. If

*Courant Institute of Mathematical Sciences, New York University. New York, NY, USA.

†Department of Mathematics, The University of British Columbia. Vancouver, BC, Canada.

E is bounded and Ahlfors-David regular, then all of these dimensions are equal. See [5] for further details on Assouad dimension and its properties.

Our first result is the following weak version of theakeya set conjecture in \mathbb{R}^3 .

Theorem 1.2. *Everyakeya set in \mathbb{R}^3 has Assouad dimension 3.*

Remarks

1. Theorem 1.2 holds for a slightly more general class of sets, where the lines satisfy a mild strengthening of the Wolff axioms. See the remarks in Section 8 for details.

2. Fraser, Olson and Robinson [4] proved that every half-extendedakeya set in \mathbb{R}^n (i.e. a subset of \mathbb{R}^n containing a half-infinite line segment in every direction) has Assouad dimension n . However this is a somewhat different question. In brief, Fraser, Olson and Robinson showed that the Assouad dimension does not increase under a “zooming out” rescaling, which transforms an extendedakeya set into one where all lines pass through the origin (the latter type of set has full Assouad dimension).

Next, we considerakeya sets with equal Hausdorff and packing dimension. Recall that for $E \subset \mathbb{R}^n$, the packing dimension is given by

$$\dim_P(E) = \inf\{\sup \overline{\dim}_M E_i\},$$

where $\overline{\dim}_M$ denotes the upper Minkowski dimension, and the infimum is taken over all decompositions $E = \bigcup_i E_i$ into countably many sets. We have $\dim_P(E) \leq \dim_H(E)$ for every set $E \subset \mathbb{R}^n$, and if $\dim_P(E) = \dim_H(E) = \alpha$, then for every $\varepsilon > 0$ there exists a constant C_ε and a measure μ supported on E so that for μ -a.e. $x \in E$ we have the Frostman-type estimate

$$C_\varepsilon^{-1}r^{\alpha+\varepsilon} \leq \mu(B(x, r)) \leq C_\varepsilon r^{\alpha-\varepsilon} \quad \text{for all } 0 < r \leq 1.$$

We would like to say that everyakeya set in \mathbb{R}^3 with equal Hausdorff and packing dimension must have dimension 3. Unfortunately, we cannot prove this statement, and indeed this would imply that everyakeya set in \mathbb{R}^3 has packing dimension 3: let $K \subset \mathbb{R}^3$ be aakeya set, and let $X \subset \mathbb{R}^3$ be a compact set with equal Hausdorff and packing dimension $\dim_H(X) = \dim_P(X) = \dim_P(K)$. Then $K' = K \cup X$ is aakeya set with equal Hausdorff and packing dimension, and this common value is $\dim_P(K)$. In the above example, theakeya set K' has equal Hausdorff and packing dimension for the somewhat trivial reason that it is a union of aakeya set with potentially unequal Hausdorff and packing dimension, and a set of equal Hausdorff and packing dimension¹. In particular, if $\dim_H K < \dim_H K' = \alpha$, then an α -dimensional Frostman measure supported on K' would be null on K . To exclude this type of situation, we introduce the following definition.

Definition 1.3. *We say a set compact set $K \subset \mathbb{R}^n$ is aakeya set with stably equal Hausdorff and packing dimension if for all $\varepsilon > 0$, there is a measure μ supported on K with the following properties.*

- *There is a constant C_ε so that for all $x \in \text{supp}(\mu)$, we have*

$$C_\varepsilon^{-1}r^{\alpha+\varepsilon} \leq \mu(B(x, r)) \leq C_\varepsilon r^{\alpha-\varepsilon}, \quad \text{for all } 0 \leq r \leq 1,$$

where $\alpha = \dim_H(K)$.

¹We could even select X to be a union of unit line segments pointing in different directions (specifically a $\dim X - 1$ dimensional set of directions), and modify K so that K' is a union of unit line segments, with one line segment pointing in each direction.

- There is a positive (two-dimensional Lebesgue) measure set of directions $\Omega \subset S^2$, so that for each direction $e \in \Omega$, there is a line ℓ_e pointing in direction e for which $\ell_e \cap \text{supp}(\mu)$ has positive one-dimensional Lebesgue measure.

In particular, if K is a Kakeya set with unequal Hausdorff and packing dimension, then the set $K' = K \cup X$ described above will not have stably equal Hausdorff and packing dimension².

Theorem 1.4. *Every Kakeya set in \mathbb{R}^3 with stably equal Hausdorff and packing dimension has Hausdorff and packing dimension 3.*

Theorems 1.2 and 1.4 will follow from a common discretized Kakeya estimate, which is the main technical contribution of this paper. In the statement below, $N_\rho(X)$ denotes the ρ -neighborhood of the set X , a δ -tube is the δ -neighborhood of a unit line segment, and say two δ -tubes are essentially distinct if neither tube is contained in the 2-fold dilate of the other; see Section 2 for precise definitions.

Theorem 1.5. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let \mathbb{T} be a set of essentially distinct δ -tubes in $B(0, 1) \subset \mathbb{R}^3$, and suppose that each convex set of volume V contains at most $\delta^{-\eta}V(\#\mathbb{T})$ tubes from \mathbb{T} . For each $T \in \mathbb{T}$, let $Y(T) \subset T$ be a measurable set, and suppose that $\sum_{\mathbb{T}} |Y(T)| \geq \delta^\eta \sum_{\mathbb{T}} |T|$.*

Then there exists $\rho, r \in [\delta, 1]$ with $\rho \leq \delta^\eta r$ and a ball B of radius r so that

$$\left| B \cap N_\rho \left(\bigcup_{T \in \mathbb{T}} Y(T) \right) \right| \geq (\rho/r)^\varepsilon |B|. \quad (1.1)$$

Remark 1.6. In the above theorem, we have required that the tubes be essentially distinct and that every convex set $U \subset \mathbb{R}^3$ contain at most $\delta^{-\eta}|U|(\#\mathbb{T})$ tubes from \mathbb{T} . This condition is satisfied, for example, if $\#\mathbb{T} = \delta^{-2}$ and the tubes in \mathbb{T} point in δ -separated directions. Indeed; for the latter condition we need only consider convex sets $U \subset B(0, 1) \subset \mathbb{R}^3$ of diameter roughly 1, and every set of this form is comparable to an ellipsoid whose axes have lengths $s \leq t \leq w$ with $w \sim 1$. If T is a δ -tube contained in U , then the direction of T lies in a rectangular sector of S^2 of dimensions roughly $s \times t$; if the tubes in \mathbb{T} point in δ -separated directions, we conclude that $\#\{T \in \mathbb{T} : T \subset U\} \lesssim st\delta^{-2} \lesssim |U|(\#\mathbb{T})$. The anti-clustering condition imposed in Theorem 1.5 and will be discussed further in Section 2.2.

Theorem 1.5 can be amplified to obtain the following variant, which says that if \mathbb{T} is a collection of δ -tubes satisfying the anti-clustering condition described above, then there are two well-separated scales $\rho < r$ so that the ρ -neighborhood of $\bigcup_{\mathbb{T}} T$ is almost as large as its r -neighborhood.

Corollary 1.7. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let \mathbb{T} be a set of essentially distinct δ -tubes in $B(0, 1) \subset \mathbb{R}^3$, and suppose that each convex set of volume V contains at most $\delta^{-\eta}V(\#\mathbb{T})$ tubes from \mathbb{T} . For each $T \in \mathbb{T}$, let $Y(T) \subset T$ be a measurable set, and suppose that $\sum_{\mathbb{T}} |Y(T)| \geq \delta^\eta \sum_{\mathbb{T}} |T|$.*

Then there exist sets $Y'(T) \subset Y(T)$, $T \in \mathbb{T}$, with $\sum_{\mathbb{T}} |Y'(T)| \gtrsim |\log \delta|^{-2} \sum_{\mathbb{T}} |Y(T)|$, and there exist scales $\rho, r \in [\delta, 1]$ with $\rho \leq \delta^\eta r$ so that

$$\left| N_\rho \left(\bigcup_{T \in \mathbb{T}} Y'(T) \right) \right| \geq (\rho/r)^\varepsilon \left| N_r \left(\bigcup_{T \in \mathbb{T}} Y'(T) \right) \right|. \quad (1.2)$$

²Unless the set X is itself a Kakeya set with stably equal Hausdorff and packing dimension.

1.1 Main ideas, and a sketch of the proof

Recall that every Kakeya set contains a two-dimensional family of unit line segments pointing in different directions. In the discretized setting, this corresponds to a set of δ -tubes pointing in δ -separated directions. The requirement that the line segments (or δ -tubes) point in different directions can be thought of as a type of anti-clustering condition—without this condition, for example, we could construct a two-dimensional family of coplanar line segments in \mathbb{R}^3 , and the union of these line segments would have dimension at most 2.

In [17], Wolff imposed an anti-clustering condition on his collection of δ -tubes in \mathbb{R}^3 that is slightly weaker than requiring the tubes to point in δ -separated directions. First, Wolff required that for $\delta \leq \rho \leq 1$, each ρ -tube must contain at most $(\rho/\delta)^2$ δ -tubes; if we think of δ -tubes as points in the Grassmannian of (affine) lines in \mathbb{R}^3 , then Wolff's first requirement corresponds to a 2-dimensional Frostman non-concentration condition on the set of tubes. Second, Wolff required that for each $\delta \leq \rho \leq 1$, at most ρ/δ δ -tubes can be contained in any rectangular prism of dimensions $\delta \times \rho \times 1$. This second condition prevents the tubes from clustering into the δ -neighborhood of a plane. These two conditions are now referred to as the *Wolff axioms*.

In our arguments below, we will consider several anti-clustering conditions that are related to the Wolff axioms. The first one already appeared in the statement of Theorem 1.5; it is a mild generalization of the Wolff axioms, which we call the *Convex Wolff Axioms*. Next, we say a set of δ -tubes \mathbb{T} satisfies the *Convex Wolff Axioms at every scale* if for every $\delta \leq \rho \leq 1$, the tubes in \mathbb{T} can be covered by ρ -tubes, so that for each ρ -tube T_ρ in this cover, the portion of \mathbb{T} contained inside T_ρ can be re-scaled to create a collection of (δ/ρ) -tubes that satisfy the convex Wolff axioms. Finally, we say a set of tubes \mathbb{T} satisfies a *Frostman Condition at exponent α* if every ρ -tube contains at most $\rho^\alpha(\#\mathbb{T})$ tubes from \mathbb{T} . Precise versions of these definitions are given in Section 2.2.

In Section 4, we show that if a set of tubes \mathbb{T} satisfies the Convex Wolff Axioms and also satisfies a Frostman condition, then either it must satisfy the Convex Wolff Axioms at every scale, or else the conclusion of Theorem 1.5 must hold for \mathbb{T} (or both). In broad strokes, the argument is as follows. Suppose that \mathbb{T} satisfies a Frostman condition but fails to satisfy the Convex Wolff Axioms at every scale. Then the tubes in \mathbb{T} must cluster into rectangular prisms of dimensions $s \times t \times 1$ for some $\delta \leq s \ll t \leq 1$. Each such prism is essentially the intersection of the s -neighborhood of a plane with the t -neighborhood of a line. First, we consider the case that when two such prisms intersect, the corresponding planes make angle $\theta \gg s/t$. Then the union of the s -neighborhood of these planes will contain a ball of radius $r = \theta t$, which is much larger than s , and hence (1.1) holds. Second, we consider the case that when two such prisms intersect, the corresponding planes make angle $\leq s/t$. This forces the $s \times t \times 1$ prisms cluster into rectangular prisms of dimensions $s' \times t' \times 1$, where $s' \ll t'$ are substantially larger than s and t , respectively. We can then repeat this argument at a larger scale. After finitely many iterations of this argument, we must find ourselves in the First Case. This concludes the argument.

The argument described above is a key geometric input for the proof of Theorem 1.5. With this geometric input, the proof of Theorem 1.5 proceeds by contradiction: suppose there exists a counter-example to Theorem 1.5, i.e. a set of tubes satisfying the Convex Wolff Axioms, for which (1.1) fails for every pair of scales ρ, r and every ball B of radius r . The first step is to find a “worst possible” such counter-example to Theorem 1.5, i.e. a set of δ -tubes \mathbb{T} of largest possible cardinality satisfying the Convex Wolff Axioms, for which the reverse of Inequality (1.1) holds for every pair of scales ρ, r and every ball B of radius r , with the term $(\rho/r)^\varepsilon$ replaced by $(\rho/r)^\omega$, for the largest possible exponent ω .

Since the conclusion of Theorem 1.5 is agnostic to location and scale, if we select $s \in [\delta, 1]$; consider only those tubes from \mathbb{T} that are contained in a s -tube; and re-scale, then the reverse of Inequality (1.1) must continue to hold, with the same exponent ω . But this means that the newly obtained collection of tubes (obtained by restricting \mathbb{T} to a s -tube and rescaling) cannot be larger (relative to the new scale δ/s) than the original set \mathbb{T} , since \mathbb{T} was a set of largest possible cardinality for which the reverse of Inequality (1.1) holds at every location and scale with exponent ω . The above argument shows that \mathbb{T} must satisfy a Frostman condition. We can now apply the geometric input discussed above (and proved in Section 4): \mathbb{T} satisfies the Convex Wolff Axioms (by assumption), and we just established that \mathbb{T} satisfies a Frostman condition. On the other hand, we are assuming that \mathbb{T} does not satisfy the conclusion of Theorem 1.5. We conclude that \mathbb{T} satisfies the Convex Wolff Axioms at every scale.

In summary, we have shown that if Theorem 1.5 is false, then there must exist a “worst possible” counter-example, and this counter-example must satisfy the Convex Wolff Axioms at every scale. Finally, we arrive at a contradiction (and thus complete the proof of Theorem 1.5) using a slight variant of the Sticky Kakeya theorem from [16]. We state an informal version of this result below.

Theorem 1.8. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let \mathbb{T} be a set of essentially distinct δ -tubes in \mathbb{R}^3 that satisfy the Convex Wolff Axioms at every scale. For each $T \in \mathbb{T}$, let $Y(T) \subset T$ be a measurable set, and suppose that $\sum_{\mathbb{T}} |Y(T)| \geq \delta^\eta \sum_{\mathbb{T}} |T|$. Then*

$$\left| \bigcup_{T \in \mathbb{T}} Y(T) \right| \geq \delta^\varepsilon. \quad (1.3)$$

Theorem 1.8 is closely related to previous work by the authors from [16]. In [16], the authors prove a variant of Inequality (1.3), in which the tubes \mathbb{T} satisfy slightly different hypotheses—the tubes are required to point in different directions and satisfy a Frostman condition at exponent 2. There are two places in [16] where one uses the hypothesis that the tubes point in different directions, and this hypothesis can be replaced by the Convex Wolff Axioms with minimal modification. A precise version of Theorem 1.8 is stated in Section 5. In Section 6 we briefly outline the proof of Theorem 1.8, and how it may be obtained by modifying the arguments in [16].

1.2 Thanks

The authors would like to thank Jonathan Fraser and Terence Tao for comments and suggestions on an earlier draft of this manuscript. Hong Wang is supported by NSF CAREER DMS-2238818 and NSF DMS-2055544. Joshua Zahl is supported by a NSERC Discovery Grant.

2 Notation and Definitions

We write $A \lesssim B$ or $A = O(B)$ to mean there is a constant C (which might depend on the ambient dimension n) so that $A \leq CB$. We write $A \sim B$ if $A \lesssim B$ and $B \lesssim A$. If the constant C is allowed to depend on an additional parameter such as ε , we then we denote this by $A \lesssim_\varepsilon B$. Many of our results will involve a small positive parameter, which we will call δ . We write $A \lesssim_\delta B$ to mean that there is a constant C so that $A \leq C(\log 1/\delta)^C B$. As above, if the constant C is allowed to depend on an additional parameter such as ε , we then we denote this by $A \lesssim_{\varepsilon, \delta} B$. Finally, during proof sketches and other informal remarks, we will sometimes write $A \ll B$ to suggest to the reader that the real number A is much smaller than the real number B .

2.1 Tubes and their shadings

Recall that a δ -tube in \mathbb{R}^n is the δ -neighborhood of a unit line segment. We will use T to denote a δ -tube, and $\ell(T)$ to denote its coaxial line. If T is a δ -tube (or more generally, the translation of a centrally symmetric convex set) and $R > 0$, we define RT to be the R -fold dilate of T . We say two tubes T, T' are *essentially distinct* if $T \not\subset 2T'$ and $T' \not\subset 2T$.

We will sometimes be concerned with collections of tubes, plus shadings (i.e. distinguished subsets) of these tubes. To this end, we introduce the following definitions.

Definition 2.1. *Let \mathcal{S} be a collection of subsets of \mathbb{R}^n , and let $\tau \in (0, 1]$. A collection of sets $\{Y(S) \subset S : S \in \mathcal{S}\}$ is called a τ -dense shading of \mathcal{S} if $\sum_{S \in \mathcal{S}} |Y(S)| \geq \tau \sum_{S \in \mathcal{S}} |S|$. $\{Y(S)\}$ is called a uniformly τ -dense shadings of \mathcal{S} if $|Y(S)| \geq \tau |S|$ for each $S \in \mathcal{S}$.*

In practice, \mathcal{S} will either be a collection of δ -tubes, or a collection of $s \times t \times 1$ rectangular prisms. We will use the notation $(\mathbb{T}, Y)_\delta$ to refer to a collection of δ -tubes and an associated shading $\{Y(T) : T \in \mathbb{T}\}$.

Definition 2.2. *Let $S \subset \mathbb{R}^n$, let $Y(S) \subset S$, and let $\delta > 0$. We say the shading $Y(S)$ is regular at scales $\geq \delta$ if for each $x \in Y(S)$ and each $r \in [\delta, 1]$, we have*

$$|Y(S) \cap B(x, r)| \geq (100 \log(1/\delta))^{-1} |Y(S)| \left(\frac{|B(x, r) \cap S|}{|S|} \right). \quad (2.1)$$

If the quantity δ is apparent from context, then we will omit it and say that $Y(S)$ is regular.

The following is Lemma 2.7 from [10].

Lemma 2.3. *Let $S \subset \mathbb{R}^n$ and let $Y(S)$ be a shading. Then there is a regular shading $Y'(S) \subset Y(S)$ with $|Y'(S)| \geq \frac{1}{2} |Y(S)|$.*

The proof in [10] considers the special case where S is a unit line segment (and hence $|B(x, r) \cap S|/|S| \sim r$), but the proof is identical. In brief, we consider the quantity $|Y(S) \cap B(x, r)|$ for each dyadic scale $r = 2^k \delta$ and each ball $B(x, r)$ aligned to the dyadic grid. We delete those balls for which $|Y(S) \cap B(x, r)|$ is too small (i.e. for which (2.1) fails), and we denote the surviving subset of $Y(S)$ by $Y'(S)$. At each dyadic scale we have only deleted a small fraction of $Y(S)$, so at least half of $Y(S)$ survives this process.

2.2 Wolff axioms and their generalizations

In this section we will define several anti-clustering conditions that we can impose on our collection of tubes. In [17] (see also Definition 13.1 from [7]), Wolff introduced what are now called the Wolff axioms; this is a non-concentration condition that forbids a collection of δ -tubes from clustering near a fatter ρ -tube, and also forbids a collection of δ -tubes from concentrating on (affine) 2-planes. The precise definition is as follows.

Definition 2.4. *Let \mathbb{T} be a set of δ -tubes in \mathbb{R}^n . We say \mathbb{T} satisfies the Wolff axioms if:*

- *Every ρ -tube can contain at most $C(\rho/\delta)^{n-1}$ tubes from \mathbb{T} .*
- *Every rectangular prism of dimensions $2\delta \times \rho \times 2$ can contain at most $C(\rho/\delta)$ tubes from \mathbb{T} .*

Remarks

1. In the above definition, the constant C is generally chosen to be a large number depending only on the ambient dimension n .
2. Every set of δ -tubes pointing in δ -separated directions will satisfy the Wolff axioms with constant $C = 100^n$.
3. Every set of δ -tubes contained in $B(0, 1) \subset \mathbb{R}^n$ that satisfies the Wolff axioms has cardinality at most $2C\delta^{1-n}$.

In the arguments that follow, we will need several variants of the Wolff axioms. We define these below.

Definition 2.5. *Let $C \geq 1$. We say a multi-set \mathcal{S} of sets in \mathbb{R}^n satisfies the Convex Wolff Axioms with error C if for all convex sets $W \subset \mathbb{R}^n$, we have*

$$\#\{S \in \mathcal{S}: S \subset W\} \leq C|W|(\#\mathcal{S}). \quad (2.2)$$

In practice we will have $n = 3$, and the sets in \mathcal{S} will be either δ -tubes, or $s \times t \times 1$ rectangular prisms (not necessarily axis-parallel) for some $s, t \in [\delta, 1]$.

Remark 2.6. If the elements of \mathcal{S} are convex, then $\#\mathcal{S} \geq C^{-1}(\inf_{S \in \mathcal{S}} |S|)^{-1}$. In particular, a set of δ -tubes in \mathbb{R}^3 satisfying the Convex Wolff Axioms must have cardinality $\gtrsim \delta^{-2}$.

Remark 2.7. Let \mathcal{S} be a multi-set of subsets of \mathbb{R}^n . For each $S \in \mathcal{S}$, let $T(S) \supset S$, and define the multi-set $\mathcal{S}' = \{T(S): S \in \mathcal{S}\}$, i.e. the multi-sets \mathcal{S} and \mathcal{S}' have the same cardinality. If \mathcal{S} satisfies the Convex Wolff Axioms with error C , then \mathcal{S}' also satisfies the Convex Wolff Axioms with error C .

In our arguments below, we will often consider a Kakeya set (or more accurately a collection of δ -tubes) at many different scales. Typically, for a collection of δ -tubes \mathbb{T} , we will want to examine the coarsening of \mathbb{T} at a larger scale ρ , and also examine the collection of δ -tubes from \mathbb{T} that are contained inside each of these coarser ρ -tubes. As observed in Remark 2.7 above, if \mathbb{T} satisfies the Convex Wolff Axioms, then the coarsening of \mathbb{T} at a larger scale will also satisfy these axioms³. However, the (rescaled) collection of δ -tubes from \mathbb{T} contained inside each coarser ρ -tube might not satisfy the Convex Wolff Axioms. Our next task is to introduce a slightly stricter variant of the Convex Wolff Axioms that addresses this issue.

Definition 2.8. *Let U be a convex set and let $S \subset U$. Let $U^* \supset U$ be the John ellipsoid that circumscribes U , and let ϕ_U be an affine transformation that sends U^* to the unit ball. The unit rescaling of S relative to U is the set $\phi_U(S)$. If \mathcal{S} be a multi-set of subsets of \mathbb{R}^n that are contained in U , then the unit rescaling of \mathcal{S} relative to U is the multi-set $\{\phi_U(S): S \in \mathcal{S}\}$.*

Remark 2.9. In [16], the authors considered a similar definition in the special case where \mathcal{S} is a set of δ -tubes and U is a ρ -tube. In order for this definition to be compatible with the definition from [16] of a tube's shading, the map ϕ_U was defined slightly differently, so that image of U^* was the ellipsoid $\{Cx^2 + Cy^2 + z^2 \leq 1\}$ (for some fixed constant C), rather than the unit ball.

Definition 2.10. *Let \mathcal{A} be a set and \mathcal{B} be a multi-set of convex subsets of \mathbb{R}^n . We say that \mathcal{A} is a cover of \mathcal{B} if each set $B \in \mathcal{B}$ is contained in at least one set from \mathcal{A} . For each $A \in \mathcal{A}$, we write $\mathcal{B}[A]$ to denote the multi-set $\{B \in \mathcal{B}: B \subset A\}$. We say that \mathcal{A} is a K -uniform cover of \mathcal{B} if*

³In contrast, the more traditional condition that the tubes in \mathbb{T} point in δ -separated directions is *not* preserved after coarsening

$\#\mathcal{B}[A] \leq K(\#\mathcal{B}[A'])$ for each pair of sets $A, A' \in \mathcal{A}$. Finally, we say that \mathcal{A} is a partitioning cover of \mathcal{B} if \mathcal{A} is a cover of \mathcal{B} , and for every pair of distinct $A, A' \in \mathcal{A}$, the sets $\mathcal{B}[2A]$ and $\mathcal{B}[2A']$ are disjoint, where $2A$ denotes the 2-fold dilate of A . In practice, \mathcal{A} and \mathcal{B} will be collections of ρ -tubes and δ -tubes respectively, for some $\rho > \delta$.

Remark 2.11. Note that if \mathcal{B} satisfies the Convex Wolff Axioms with error C , and if \mathcal{A} is a K -uniform partitioning cover of \mathcal{B} , then \mathcal{A} satisfies the Convex Wolff Axioms with error KC .

Definition 2.12. Let $C \geq 1, \delta > 0$. We say a set \mathbb{T} of δ -tubes in \mathbb{R}^n satisfies the Convex Wolff Axioms at every scale with error C if the tubes in \mathbb{T} are essentially distinct, and for every $\rho_0 \in [\delta, 1]$, there exists $\rho \in [\rho_0, C\rho_0)$ and a set of ρ -tubes \mathbb{T}_ρ that satisfies the following properties.

- (i) \mathbb{T}_ρ is a C -uniform partitioning cover of \mathbb{T} .
- (ii) For each $T_\rho \in \mathbb{T}_\rho$, the unit re-scaling of $\mathbb{T}[T_\rho]$ relative to T_ρ satisfies the Convex Wolff Axioms with error C .

As the name suggests, Definition 2.12 is a multi-scale property. In particular, if \mathbb{T} satisfies the Convex Wolff Axioms at every scale, then we can cover \mathbb{T} by fatter ρ -tubes; both the ρ -tubes and the (re-scaled) tubes from \mathbb{T} inside each ρ -tube will again satisfy the Convex Wolff Axioms at every scale. The precise statement is as follows.

Lemma 2.13. Let \mathbb{T} be a set of δ -tubes in \mathbb{R}^n that satisfies the Convex Wolff Axioms at every scale with error C . Let $\rho_0 \in [\delta, 1]$. Then there exists $\rho \in [\rho_0, C\rho_0]$ and a set of ρ -tubes \mathbb{T}_ρ , so that \mathbb{T}_ρ satisfies the Convex Wolff Axioms at every scale with error $O(C)$, and for each $T_\rho \in \mathbb{T}_\rho$, the unit rescaling of $\mathbb{T}[T_\rho]$ relative to T_ρ satisfies the Convex Wolff Axioms at every scale with error $O(C)$.

Proof. Let N be the largest integer with $C^{N+1} \leq \delta^{-1}$. For $i = 1, \dots, N$, let \mathbb{T}_{τ_i} be a set of τ_i -tubes, with $\tau_i \in [\delta C^i, \delta C^{i+1})$ that satisfy Items (i) and (ii) from Definition 2.12. We claim that for $i < j$, if $T_{\tau_i} \in \mathbb{T}_{\tau_i}, T_{\tau_j} \in \mathbb{T}_{\tau_j}$, and $\mathbb{T}[T_{\tau_i}] \cap \mathbb{T}[T_{\tau_j}] \neq \emptyset$, then $\mathbb{T}[T_{\tau_i}] \subset \mathbb{T}[T_{\tau_j}]$. To see this, observe that if the intersection contains at least one tube $T \in \mathbb{T}$, then $2T_{\tau_j} \supset N_{\tau_j}(T) \supset T_{\tau_i}$. Since \mathbb{T}_{τ_j} covers \mathbb{T} , and the sets $\{\mathbb{T}[2T'_{\tau_j}]: T'_{\tau_j} \in \mathbb{T}_{\tau_j}\}$ are disjoint, we conclude that $\mathbb{T}[T_{\tau_i}] \subset \mathbb{T}[T_{\tau_j}]$.

The consequence of the above observation is the following: if we partially order the sets $\{\mathbb{T}[T_{\tau_i}]: T_{\tau_i} \in \mathbb{T}_{\tau_i}, i = 1, \dots, N\}$ under inclusion, then we actually obtain a total ordering, and this ordering forms a tree with N levels—the vertices at level i are precisely the sets $\mathbb{T}[T_{\tau_i}]$. By Item (i) of Definition 2.12, for each $i = 1, \dots, N$ the sets $\{\mathbb{T}[T_{\tau_i}]: T_{\tau_i} \in \mathbb{T}_{\tau_i}\}$ have comparable cardinality, up to a multiplicative factor of C . This tree is precisely what is needed to verify Lemma 2.13: given $\rho_0 \in [\delta, 1]$, we select an index i so that $\tau_i \in [\rho_0, C\rho_0)$, and we choose $\rho = \tau_i, \mathbb{T}_\rho = \mathbb{T}_{\tau_i}$; note that since \mathbb{T}_ρ forms a partitioning cover of \mathbb{T} , the ρ -tubes in \mathbb{T}_ρ are essentially distinct. \square

Finally, we will introduce two additional anti-concentration conditions that will play a technical role in the arguments to follow.

Definition 2.14. Let $C \geq 1$ and let \mathbb{T} be a set of δ -tubes in \mathbb{R}^n . We say that \mathbb{T} satisfies a Frostman condition at dimension σ with error C if the tubes in \mathbb{T} are essentially distinct, and for every $\rho \geq \delta$ and every ρ -tube T_ρ , we have $\#\mathbb{T}[T_\rho] \leq C\rho^\sigma(\#\mathbb{T})$.

Definition 2.15. Let $C \geq 1$ and let \mathbb{T} be a set of δ -tubes in \mathbb{R}^n . We say that \mathbb{T} satisfies the Self-Similar Convex Wolff Axioms with error C if the tubes in \mathbb{T} are essentially distinct, and for every $\rho_0 \in [\delta, 1]$, there exists $\rho \in [\rho_0, C\rho_0)$ and a set of ρ -tubes \mathbb{T}_ρ that satisfies the following properties.

- (i) \mathbb{T}_ρ is a C -uniform partitioning cover of \mathbb{T} .
- (ii) For each $T_\rho \in \mathbb{T}_\rho$, the unit re-scaling of $\mathbb{T}[T_\rho]$ relative to T_ρ satisfies the Convex Wolff Axioms with error C .
- (iii) For each $T_\rho \in \mathbb{T}_\rho$, we have

$$C^{-1}(\rho/\delta)^\sigma \leq \#\mathbb{T}[T_\rho] \leq C(\rho/\delta)^\sigma, \quad (2.3)$$

where $\sigma > 0$ is the unique number satisfying $\#\mathbb{T} = \delta^{-\sigma}$.

Note that Items (i) and (ii) above are identical to their counterparts in Definition 2.12. In particular, we have

$$\text{Self-Similar CWA w. err. } C \implies \text{CWA at every scale w. err. } C \implies \text{CWA w/ err. } C. \quad (2.4)$$

2.3 Discretized Assouad dimension

The conclusion of Theorem 1.5 involves a discretized analogue of Assouad dimension. We formalize this as follows.

Definition 2.16. *Let $E \subset \mathbb{R}^n$, let $\beta \in [0, n]$, let $\delta > 0$, and let $A \geq 1$. We say that E has discretized Assouad dimension at least β , at scale δ and scale separation A if there exist scales $\delta \leq \rho \leq r \leq 1$ with $r \geq A\rho$, and a ball B of radius r , so that*

$$|B \cap N_\rho(E)| \geq (\rho/r)^{n-\beta}|B|.$$

If the scale δ is apparent from context, then for brevity we may say “ E has discretized Assouad dimension at least β at scale separation A .”

In this language, the conclusion of Theorem 1.5 says that the set $\bigcup_{T \in \mathbb{T}} Y(T)$ has discretized Assouad dimension at least $3 - \varepsilon$ at scale separation $\delta^{-\eta}$.

3 Assouad-extremal Kakeya sets

In this section, we will suppose that Theorem 1.5 is false, and we seek to construct a “worst possible” counter-example to Theorem 1.5. In this and later sections, we will restrict attention to \mathbb{R}^3 . Our example will be worst possible in two respects. First, Inequality (1.1) will fail as dramatically as possible, i.e. a reverse inequality will hold where the term $(\rho/r)^\varepsilon$ on the RHS of (1.1) will be replaced by $(\rho/r)^\omega$ for the largest possible value of $\omega > 0$. Second, of all collections of tubes for which such a reverse inequality holds for this value of ω , we will choose a set \mathbb{T} of largest possible cardinality. The hypotheses of Theorem 1.5 ensure that \mathbb{T} has size at least δ^{-2} ; we will choose a set of tubes of size $\delta^{-\alpha}$, for the largest possible value of α .

We now turn to the task of carefully defining the quantities ω and α ; this will be done in Sections 3.1 and 3.2, respectively. This task is complicated by the fact that Theorem 1.5 involves a sequence of quantifiers, and we must unwind these quantifiers in the correct order.

3.1 Defining ω

Let $(\mathbb{T}, Y)_\delta$ be a set of tubes and their associated shading. Let $\delta \leq \rho < r \leq 1$, and let B be a ball of radius r whose intersection with $\bigcup Y(T)$ has positive measure. Define $\zeta = \zeta(\mathbb{T}, Y, B, \rho)$ to be the unique number satisfying

$$\left| B \cap N_\rho \left(\bigcup_{T \in \mathbb{T}} Y(T) \right) \right| = \left(\frac{\rho}{r} \right)^\zeta |B|. \quad (3.1)$$

Next, for $\eta > 0$, define

$$\zeta(\mathbb{T}, Y; \eta) = \inf_{B, \rho} \zeta(\mathbb{T}, Y, B, \rho), \quad (3.2)$$

where the infimum is taken over all pairs (B, ρ) , where $\rho \in [\delta, \delta^\eta]$ and B is a ball of radius $r \in [\delta^{-\eta}\rho, 1]$ whose intersection with $\bigcup Y(T)$ has positive measure. Observe that as $\eta \searrow 0$, the set of admissible pairs (B, ρ) becomes larger, and hence $\zeta(\mathbb{T}, Y; \eta)$ weakly decreases, i.e. $\zeta(\mathbb{T}, Y; \eta)$ is a weakly increasing function of η .

For $s, \eta, \delta > 0$ and $u \geq 2$, define

$$\omega(\varepsilon, \eta, \delta; u) = \sup_{(\mathbb{T}, Y)_\delta} \zeta(\mathbb{T}, Y; \eta), \quad (3.3)$$

where the supremum is taken over all choices of $(\mathbb{T}, Y)_\delta$ with the following properties:

- \mathbb{T} is a set of essentially distinct δ -tubes contained in $B(0, 1)$ that satisfies the Convex Wolff Axioms with error $\delta^{-\varepsilon}$, and Y is a δ^ε -dense shading.
- $\#\mathbb{T} \geq \delta^{-u+\varepsilon}$.

Again, $\omega(\varepsilon, \eta, \delta; u)$ is a weakly increasing function of η . It is also a weakly increasing function of ε and weakly decreasing function of u , since as ε become larger or u becomes smaller, the set of admissible pairs $(\mathbb{T}, Y)_\delta$ becomes larger.

Define

$$\omega(u) = \lim_{\varepsilon \rightarrow 0^+} \lim_{\eta \rightarrow 0^+} \limsup_{\delta \rightarrow 0^+} \omega(\varepsilon, \eta, \delta; u) = \inf_{\varepsilon, \eta > 0} \limsup_{\delta \rightarrow 0^+} \omega(\varepsilon, \eta, \delta; u).$$

Definition 3.1. Define $\omega = \omega(2)$.

Unwrapping the above definitions, we have the following.

Lemma 3.2. *Let $\omega \geq 0$ be as in Definition 3.1. For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $(\mathbb{T}, Y)_\delta$ be a set of essentially distinct tubes that satisfy the Convex Wolff Axioms with error $\delta^{-\eta}$, with a δ^η -dense shading.*

Then $\bigcup_{T \in \mathbb{T}} Y(T)$ has discretized Assouad dimension at least $3 - \omega - \varepsilon$ at scale separation $\delta^{-\eta}$.

Proof. Select $\eta > 0$ sufficiently small so that $\limsup_{\delta \rightarrow 0^+} \omega(\eta, \eta, \delta; 2) \leq \omega + \varepsilon/2$. Then there exists $\delta_0 > 0$ so that $\omega(\eta, \eta, \delta; 2) \leq \omega + \varepsilon$ for all $\delta \in (0, \delta_0]$. The conclusion of Lemma 3.2 now holds for this choice of η and δ_0 . \square

In light of Lemma 3.2, in order to prove Theorem 1.5 it suffices to prove that $\omega = 0$.

3.2 Defining α

Note that $\omega(u)$ is weakly monotone decreasing. We also have that $\omega(4) = 0$, since if \mathbb{T} is a set of δ^{-4} essentially distinct δ -tubes in $B(0, 1)$, then $\bigcup_{T \in \mathbb{T}} T$ must have volume $\gtrsim 1$. Define

$$\alpha = \sup\{u \in [2, 4]: \omega(u) = \omega(2)\}. \quad (3.4)$$

The quantity α has the following interpretation. We have defined $3 - \omega$ to be the smallest possible discretized Assouad dimension of a collection of essentially distinct tubes that satisfies the Convex Wolff Axioms. Every such collection of tubes must have cardinality at least δ^{-2} , and in general we might expect that larger collections of tubes should have larger discretized Assouad dimension. Any collection of essentially distinct tubes of cardinality substantially larger than $\delta^{-\alpha}$ that satisfies the Convex Wolff Axioms must have discretized Assouad dimension larger than $3 - \omega$.

3.3 The existence of Assouad-Extremal Kakeya sets

Now that we have defined ω and α , we are ready to construct a (hypothetical) “worst possible” counter-example to Theorem 1.5, in the sense described at the beginning of Section 3.

Definition 3.3. For $\varepsilon > 0$, we say a pair $(\mathbb{T}, Y)_\delta$ is ε Assouad-extremal if it satisfies the following properties.

- $\#\mathbb{T} \geq \delta^{-\alpha+\varepsilon}$.
- The tubes are contained in $B(0, 1)$, essentially distinct, and satisfy the Convex Wolff Axioms with error $\delta^{-\varepsilon}$.
- The shading Y is δ^ε -dense.
- For all $\rho, r \in [\delta, 1]$ with $\rho \leq \delta^\varepsilon r$ and all balls B of radius r , we have

$$\left| B \cap N_\rho \left(\bigcup_{T \in \mathbb{T}} Y(T) \right) \right| \leq (\rho/r)^{\omega-\varepsilon} |B|. \quad (3.5)$$

Note that if \mathbb{T} is ε Assouad-extremal, then it is also ε' Assouad-extremal for all $\varepsilon' \geq \varepsilon$.

Lemma 3.4. For all $\varepsilon, \delta_0 > 0$, there exists $\delta \in (0, \delta_0]$ and an ε Assouad-extremal set of tubes $(\mathbb{T}, Y)_\delta$.

Proof. Fix $u \geq \max(2, \alpha - \varepsilon/2)$ with $\omega(u) = \omega(2) = \omega$. Since $\omega(s, \eta, \delta; u)$ is a weakly increasing function of s and η , we have $\limsup_{\delta \rightarrow 0^+} \omega(\varepsilon/2, \varepsilon, \delta; u) \geq \omega(u)$. Thus there exists $\delta \in (0, \delta_0]$ with $\omega(\varepsilon/2, \varepsilon, \delta; u) \geq \omega - \varepsilon/2$.

In light of (3.3), there exists a set of tubes and their associated shading $(\mathbb{T}, Y)_\delta$, so that (i) $\#\mathbb{T} \geq \delta^{-u+\varepsilon/2} \geq \delta^{-\alpha+\varepsilon}$; (ii) the tubes are contained in $B(0, 1)$, essentially distinct, and satisfy the Convex Wolff Axioms with error $\delta^{-\varepsilon}$; (iii) Y is a δ^ε -dense shading; and (iv) we have $\zeta(\mathbb{T}, Y; \eta) \geq \omega - \varepsilon$. Thus by (3.2), for every pair of scales $\rho, r \in [\delta, 1]$ with $\rho \leq \delta^\varepsilon r$ and every ball B of radius r , we have

$$\zeta(\mathbb{T}, Y, B, \rho) \geq \zeta((\mathbb{T}, Y); \eta) \geq \omega - \varepsilon.$$

But this is precisely (3.5). □

4 Convex Wolff Axioms and multi-scale structure

The purpose of this section is to prove the following result.

Proposition 4.1. *For all $\varepsilon > 0$, there exists $\tau > 0$, so that for all $\eta_1 > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $\beta \geq 2$ and let $(\mathbb{T}, Y)_\delta$ be a set of $\delta^{-\beta}$ essentially distinct δ -tubes that satisfy the Convex Wolff Axioms with error δ^η , with Y a δ^η uniformly dense shading. Then at least one of the following must be true.*

- (A) $\bigcup Y(T)$ has discretized Assouad dimension at least $3 - \varepsilon$ at scale δ , with scale separation $\delta^{-\eta_1}$.
- (B) There exists $\mathbb{T}' \subset \mathbb{T}$ with $\#\mathbb{T}' \geq \delta^{\eta_1}(\#\mathbb{T})$, and \mathbb{T}' satisfies the self-similar Convex Wolff Axioms with error $\delta^{-\varepsilon}$.
- (C) There exists $\rho \in [\delta, \delta^\varepsilon]$ and a set $\tilde{\mathbb{T}}$ of ρ -tubes, with the following properties.
 - (C.i) The tubes in $\tilde{\mathbb{T}}$ are essentially distinct, and $\#\tilde{\mathbb{T}} \geq \rho^{-\beta-\tau}$.
 - (C.ii) $\tilde{\mathbb{T}}$ satisfies the Convex Wolff Axioms with error $\leq \rho^{-\eta_1}$.
 - (C.iii) For each $T_\rho \in \tilde{\mathbb{T}}$, there exists $T \in \mathbb{T}$ with $T \subset T_\rho$.
- (D) There exists $\rho \in [\delta^{1-\varepsilon}, 1]$, a ρ -tube T_ρ , and a set $\tilde{\mathbb{T}} \subset \mathbb{T}[T_\rho]$, so that the following holds.
 - (D.i) $\#\tilde{\mathbb{T}} \geq (\delta/\rho)^{-\beta-\tau}$.
 - (D.ii) The unit rescaling of $\tilde{\mathbb{T}}$ relative to T_ρ satisfies the Convex Wolff Axioms with error $(\delta/\rho)^{-\eta_1}$.

Remark 4.2. Conclusions (A) and (B) are somewhat self-explanatory. Conclusion (C) says that at a coarser scale, we can find a set of tubes that is richer than our original set \mathbb{T} . Conclusion (D) says that we can find a ρ -tube so that the unit-rescaling of $\mathbb{T}[T_\rho]$ is richer than our original set \mathbb{T} .

An important step in proving Proposition 4.1 is to study arrangements of tubes that cluster into rectangular prisms. In the next section, we will show that unions of rectangular prisms of dimensions $s \times t \times 1$ with $s \ll t$ have large Assouad dimension.

4.1 Arrangements of rectangular prisms

The next result is a straightforward consequence of Cauchy-Schwartz; it says that a union of $\delta \times 1 \times 1$ prisms satisfying the Convex Wolff Axioms must have almost maximal volume.

Lemma 4.3. *Let $0 < \delta \leq 1$ and $K \geq 1$. Let \mathcal{R} be a multi-set of $\delta \times 1 \times 1$ rectangular prisms, and suppose that \mathcal{R} satisfies the Convex Wolff Axioms with error K . Let $\{Y(R), R \in \mathcal{R}\}$ be a K^{-1} -dense shading. Then*

$$\left| \bigcup_{R \in \mathcal{R}} Y(R) \right| \gtrsim K^{-3} |\log \delta|^{-1}. \quad (4.1)$$

Proof. For each prism $R \in \mathcal{R}$, there is an associated normal vector $v(R) \in S^2$, which is well-defined up to uncertainty $O(\delta)$. For $R, R' \in \mathcal{R}$, define $\angle(R, R')$ to be the (unsigned) angle between $v(R)$ and $v(R')$.

Fix $R \in \mathcal{R}$. We have

$$\begin{aligned} \sum_{R' \in \mathcal{R}} |R \cap R'| &= \sum_{j=0}^{\log(1/\delta)} \sum_{\substack{R': R \cap R' \neq \emptyset, \\ \angle(R, R') \sim 2^j \delta}} |R \cap R'| \lesssim \sum_{j=0}^{\log(1/\delta)} \left(K(2^j \delta \# \mathcal{R}) \right) \left(\frac{\delta^2}{2^j \delta} \right) \\ &\lesssim K \delta^2 (\# \mathcal{R}) \log(1/\delta). \end{aligned}$$

In the above computation, observe that if $R \cap R'$ is non-empty and $\angle(R, R') = \theta \sim 2^j \delta$, then R' is contained in the $O(\theta) \times 1 \times 1$ rectangular prism concentric with R . Since \mathcal{R} obeys the Convex Wolff Axioms, at most $K(2^j \delta)(\# \mathcal{R})$ slabs $R' \in \mathcal{R}$ can have this property. We conclude that

$$\left\| \sum_{R \in \mathcal{R}} \chi_R \right\|_2^2 \lesssim K \delta^2 (\# \mathcal{R})^2 \log(1/\delta). \quad (4.2)$$

Let $E = \bigcup_{R \in \mathcal{R}} Y(R)$. Since the shading Y is K^{-1} dense, by Cauchy-Schwartz we have

$$K^{-1} \delta (\# \mathcal{R}) \leq \int \chi_E \cdot \sum_{R \in \mathcal{R}} \chi_{Y(R)} \leq \int \chi_E \cdot \sum_{R \in \mathcal{R}} \chi_R \leq |E|^{1/2} \left\| \sum_{R \in \mathcal{R}} \chi_R \right\|_2. \quad (4.3)$$

Rearranging and using (4.2), we obtain (4.1). \square

Next, we will establish a geometric argument showing that a union of $s \times t \times 1$ prisms, $s \ll t$, satisfying the Convex Wolff Axioms must either have nearly maximal discretized Assouad dimension, or the prisms must arrange themselves into $s' \times t' \times 1$ prisms for s' and t' substantially larger than s and t , respectively. The precise statement is as follows.

Lemma 4.4. *For all $\varepsilon, \tau \in (0, 1]$, there exists $\delta_0, \eta > 0$ so that the following is true for all $\delta \in (0, \delta_0]$. Let $\delta \leq s \leq t \leq 1$ with $s \leq \delta^\varepsilon t$. Let \mathcal{R} be a multi-set of $s \times t \times 1$ prisms contained in $B(0, 1)$, and suppose that \mathcal{R} satisfies the Convex Wolff Axioms with error $\delta^{-\eta}$. Let $\{Y(R), R \in \mathcal{R}\}$ be a δ^η -dense shading.*

Then at least one of the following two things must happen.

(A): *There exists $r \in [\delta^{-\eta} s, 1]$ and a r -ball B so that*

$$\left| B \cap \bigcup_{R \in \mathcal{R}} Y(R) \right| \geq (s/r)^\varepsilon |B|. \quad (4.4)$$

(B): *There are scales $s' \in [\delta^{-\varepsilon/2} s, \delta^{\varepsilon/2}]$ and $t' = \min(1, t \frac{s'}{s})$, and a multi-set \mathcal{R}' of $s' \times t' \times 1$ prisms that satisfies the Convex Wolff Axioms with error $\delta^{-\tau}$, so that the shading*

$$Y'(R') = R' \cap N_{s'} \left(\bigcup_{\substack{R \in \mathcal{R} \\ RC2R'}} Y(R) \right) \quad (4.5)$$

is δ^τ -dense.

Remark 4.5. Note that as $\varepsilon \searrow 0$, Conclusion (A) becomes stronger, while conclusion (B) becomes weaker (indeed, if $\varepsilon = 0$ then Conclusion (B) vacuously holds with $s' = s$ and $t' = t$). As $\tau \searrow 0$, Conclusion (A) is unchanged, while Conclusion (B) becomes stronger.

Proof. Step 1.

After dyadic pigeonholing, we can select a multiplicity μ and a $\geq \frac{1}{10}|\log \delta|^{-1}\delta^\eta$ -dense shading $Y_1(R) \subset Y(R)$ so that $\sum_{R \in \mathcal{R}} \chi_{Y_1(R)}(x) \sim \mu$ for all $x \in \bigcup_{R \in \mathcal{R}} Y_1(R)$. Let $Y_2(R) \subset Y_1(R)$ be a regular (in the sense of Definition 2.2) refinement of Y_1 , with $|Y_2(R)| \geq \frac{1}{2}|Y_1(R)|$ for each $R \in \mathcal{R}$.

If $\mu \leq \delta^{-\varepsilon^2+3\eta}(st\#\mathcal{R})$, then

$$\left| \bigcup_{R \in \mathcal{R}} Y(R) \right| \geq \mu^{-1} \sum_{R \in \mathcal{R}} |Y_1(R)| \gtrsim \delta^{\varepsilon^2-3\eta} \delta^\eta \frac{\sum_{R \in \mathcal{R}} |R|}{st(\#\mathcal{R})} \geq \delta^{\varepsilon^2-\eta} \geq \delta^{-\eta} s^\varepsilon,$$

where the second-last inequality used the hypothesis that \mathcal{R} satisfies the Convex Wolff Axioms with error $\delta^{-\eta}$ (and hence $\sum |R| \geq \delta^\eta$) and the final inequality used the hypothesis that $s \leq \delta^\varepsilon t$ and hence $s \leq \delta^\varepsilon$. If δ_0 is sufficiently small depending on η , then Conclusion (A) holds with $r = 1$, and we are done.

Step 2.

Suppose instead that $\mu \geq \delta^{-\varepsilon^2+3\eta}(st\#\mathcal{R})$. Each $s \times t \times 1$ prism $R \in \mathcal{R}$ is the image of the set $[-s/2, s/2] \times [-t/2, t/2] \times [-1/2, 1/2]$ under a rigid motion τ . We define $\ell = \ell(R)$ to be the image of the z -axis under τ , and we define $\Pi(R)$ to be the image of the yz -plane under τ .

For each $x \in \bigcup_{R \in \mathcal{R}} Y_2(R)$, define the ‘‘plane spread’’

$$\text{P-spread}(x) = \max \left\{ \angle(\Pi(R), \Pi(R')) \right\},$$

where the maximum is taken over all pairs R, R' for which $x \in Y_2(R)$ and $x \in Y_2(R')$, and where $\angle(\Pi, \Pi')$ denotes the unsigned angle between the normal vectors of Π and Π' . Since $\Pi(R)$ and $\Pi(R')$ are defined up to uncertainty $O(s/t)$, $\text{P-spread}(x)$ is meaningfully defined if it is substantially larger than s/t . If $x \notin \bigcup_{R \in \mathcal{R}} Y_2(R)$, define $\text{P-spread}(x) = 0$.

Step 3. Suppose there exists a prism $R_0 \in \mathcal{R}$ and a number $\theta \geq \delta^{-4\eta/\varepsilon} \frac{s}{t}$ so that

$$|\{x \in R_0 : \theta \leq \text{P-spread}(x) < 2\theta\}| \geq \frac{1}{100} |\log \delta|^{-2} \delta^\eta |R_0|. \quad (4.6)$$

Since R_0 is a $s \times t \times 1$ prism, for each $x_0 \in R_0$ we have $|R_0 \cap B(x_0, t)| \sim st^2$. By pigeonholing, we can select a point $x_0 \in R_0$ so that

$$|\{x \in R_0 \cap B(x_0, t) : \theta \leq \text{P-spread}(x) < 2\theta\}| \gtrsim \delta^\eta st^2. \quad (4.7)$$

After a harmless translation and rotation, we may suppose that $\Pi(R_0)$ is the yz -plane, i.e. $\text{span}\{\mathbf{e}_2, \mathbf{e}_3\}$, and that $R_0 \cap B(x_0, t)$ is contained in the ‘‘square’’ $S = [-s, s] \times [-t, t] \times [-t, t]$ (note as well that $|S| \sim |R_0 \cap B(x_0, t)|$). Let \tilde{S} denote the thickened square $[-10t\theta, 10t\theta] \times [-10t, 10t] \times [-10t, 10t]$, i.e. \tilde{S} is comparable to the $t\theta$ neighborhood of S .

Denote the set on the LHS of (4.7) by E ; then for each $x \in E$ there is a prism $R_x \in \mathcal{R}$ with $x \in Y_2(R)$ and $\angle(\Pi(R_x), \text{span}\{\mathbf{e}_2, \mathbf{e}_3\}) \in [\theta, 2\theta]$. This implies that $R_x \cap B(x, t) \subset R_x \cap B(x_0, 10t) \subset \tilde{S}$. Since the shading $Y_2(R_x)$ is regular, we also have

$$|Y_2(R_x) \cap B(x_0, 2t)| \gtrsim |Y_2(R_x)| \frac{|R_x \cap B(x_0, 2t)|}{|R_x|} \gtrsim \delta^\eta st^2. \quad (4.8)$$

Define the linear map $A(x, y, z) = \left(\frac{x}{10t\theta}, \frac{y}{10t}, \frac{z}{10t} \right)$, so $A(\tilde{S}) = [-1, 1]^3$. Define

$$\hat{S} = A(S) = \left[-\frac{s}{10t\theta}, \frac{s}{10t\theta} \right] \times [-1/10, 1/10] \times [-1/10, 1/10],$$

and let $\hat{E} = A(E) \subset \hat{S}$. For each $x \in E$, let $\hat{R}_x = A(R_x \cap \tilde{S})$; this is comparable to a rectangular prism of dimensions roughly $\frac{s}{t\theta} \times 1 \times 1$, and this prism makes angle ~ 1 with the yz -plane, i.e. the prism has a normal direction $\mathbf{n}(\hat{R}_x)$ (defined up to uncertainty $O(\frac{s}{t\theta})$) with $|\mathbf{n}(\hat{R}_x) \wedge \mathbf{e}_1| \gtrsim 1$. By (4.8) we have

$$|\hat{Y}_2(\hat{R}_x)| \gtrsim \delta^\eta \frac{s}{t\theta}. \quad (4.9)$$

By pigeonholing, we can find a unit vector $e \in \mathbb{R}^3$ and a set $\hat{E}' \subset \hat{E}$ with $|\hat{E}'| \sim |\hat{E}|$, so that $\angle(e, \mathbf{n}(\hat{R}_x)) \leq \frac{1}{100}$ for each $x \in \hat{E}'$. In particular, this implies $|e \wedge \mathbf{e}_1| \gtrsim 1$, and hence the projection of e to the yz -plane has magnitude ~ 1 ; denote the image of e under this projection by e' .

(4.7) says that $|E| \gtrsim \delta^\eta s t^2$ and hence $|\hat{E}'| \gtrsim |\hat{E}| \gtrsim \delta^\eta \frac{s}{t\theta}$. On the other hand, $\hat{E}' \subset \hat{S}$, and the latter set has volume $\sim \frac{s}{t\theta}$. Since $\hat{S} = \left[-\frac{s}{10t\theta}, \frac{s}{10t\theta} \right] \times [-1/10, 1/10] \times [-1/10, 1/10]$ and e' is contained in the yz plane, by Fubini we can select a line ℓ pointing in direction e' that satisfies $|\ell \cap \hat{E}'| \gtrsim \delta^\eta$.

Select a $\frac{s}{t\theta}$ -separated set of points x_1, \dots, x_N , with $N \gtrsim \delta^\eta \frac{t\theta}{s}$, from $|\ell \cap \hat{E}'|$; we will index these points so that $|x_i - x_j| \geq \frac{s}{t\theta} |i - j|$. Note that $R_{x_i} \cap \ell$ is a line segment of length between $\frac{s}{t\theta}$ and $10\frac{s}{t\theta}$, and hence if i, j are a pair of indices with $|x_i - x_j| \geq 100\frac{s}{t\theta}$, then $\text{dist}(R_{x_i} \cap \ell, R_{x_j} \cap \ell) \gtrsim \frac{s}{t\theta} |i - j|$. Since ℓ points in direction e' , and $|e' \cdot \mathbf{n}(\hat{R}_{x_j})| \sim 1$, we conclude that $\text{dist}(x_i, R_{x_j}) \gtrsim \frac{s}{t\theta} |i - j|$, so in particular R_{x_i} contains a point that has distance $\gtrsim \frac{s}{t\theta} |i - j|$ from R_{x_j} . Thus either R_{x_i} and R_{x_j} do not intersect, or if they do intersect then their normal vectors satisfy $\angle(\mathbf{n}(\hat{R}_{x_i}), \mathbf{n}(\hat{R}_{x_j})) \gtrsim \frac{s}{t\theta} |i - j|$. In either case, we have

$$|\hat{R}_{x_i} \cap \hat{R}_{x_j}| \lesssim \frac{\left(\frac{s}{t\theta}\right)^2}{\frac{s}{t\theta}(1 + |i - j|)} = \frac{s}{t\theta(1 + |i - j|)}. \quad (4.10)$$

The above inequality vacuously holds if i, j are a pair of indices with $|x_i - x_j| < 100\frac{s}{t\theta}$, since in this case $|i - j| \sim 1$ and we can use the trivial bound $|\hat{R}_{x_i} \cap \hat{R}_{x_j}| \leq |\hat{R}_{x_i}|$.

We conclude that

$$\left\| \sum_{i=1}^N \chi_{\hat{Y}_2(\hat{R}_i)} \right\|_2^2 \leq \sum_{i,j=1}^N |\hat{R}_i \cap \hat{R}_j| \lesssim N \frac{s}{t\theta} \left| \log \left(\frac{s}{t\theta} \right) \right|, \quad (4.11)$$

and hence

$$\left| \bigcup_{i=1}^N \hat{Y}_2(\hat{R}_i) \right| \geq \frac{\left(\sum_i |\hat{Y}_2(\hat{R}_i)| \right)^2}{\left\| \sum_i \chi_{\hat{Y}_2(\hat{R}_i)} \right\|_2^2} \gtrsim \frac{\left(N \delta^\eta \frac{s}{t\theta} \right)^2}{N \frac{s}{t\theta} \left| \log \left(\frac{s}{t\theta} \right) \right|} \gtrsim \delta^{2\eta} N \frac{s}{t\theta} \gtrsim \delta^{3\eta},$$

where the second inequality used (4.9) and (4.11). Undoing the scaling transformation A , we conclude that

$$\left| \tilde{S} \cap \bigcup_{R \in \mathcal{R}} Y(R) \right| \gtrsim \delta^{3\eta} |\tilde{S}|.$$

Recall that \tilde{S} is a rectangular prism of dimensions roughly $t\theta \times t \times t$. Define $r = t\theta$; then by pigeonholing, we can find a ball $B \subset \tilde{S}$ of radius r so that

$$\left| B \cap \bigcup_{R \in \mathcal{R}} Y(R) \right| \gtrsim \delta^{3\eta} |B|.$$

By δ_0 sufficiently small depending on η and the implicit constants in the above quasi-inequality, we can ensure that

$$\left| B \cap \bigcup_{R \in \mathcal{R}} Y(R) \right| \geq \delta^{4\eta} |B|, \quad (4.12)$$

and since $\theta \geq \delta^{-4\eta/\varepsilon} \frac{s}{t}$, we have $\delta^{4\eta} \geq (s/r)^\varepsilon$. Thus (4.12) implies (4.4), and hence Conclusion (A) holds.

Step 4. Suppose instead that for every prism $R_0 \in \mathcal{R}$ and every $\theta \geq \delta^{-4\eta/\varepsilon} \frac{s}{t}$, (4.6) fails. Let $s_1 = \delta^{-4\eta/\varepsilon} s$. Since $s \leq \delta^\varepsilon t$, if we select η sufficiently small then $s_1 \leq t$. Select a $\gtrsim \delta^\eta$ dense shading $Y_3(R) \subset Y_2(R)$ so that $\text{P-spread}(x) \leq \frac{s_1}{t}$ for each $x \in Y_3(R)$. Let $\mathcal{R}_0 \subset \mathcal{R}$ consist of those $R \in \mathcal{R}$ with $|Y_3(R)| \gtrsim \delta^\eta |R|$; we will choose the implicit constant and implicit power of $|\log \delta|$ so that $\#\mathcal{R}_0 \gtrsim \delta^\eta (\#\mathcal{R})$.

For each $x \in \bigcup_{R \in \mathcal{R}_0} Y_3(R)$, define the ‘‘line spread’’

$$\text{L-spread}(x) = \max \left\{ \angle(\ell(R), \ell(R')) \right\}, \quad (4.13)$$

where the maximum is taken over all pairs R, R' for which $x \in Y_3(R)$ and $x \in Y_3(R')$, and where $\angle(\ell, \ell')$ denotes the unsigned angle between the lines ℓ and ℓ' . Since $\ell(R)$ and $\ell(R')$ are defined up to uncertainty $O(t)$, $\text{L-spread}(x)$ is meaningfully defined if it is substantially larger than t . If $x \notin \bigcup_{R \in \mathcal{R}_0} Y_3(R)$, define $\text{L-spread}(x) = 0$.

We claim there is a number $\theta \in [\delta^{-\varepsilon^2/2} t, \pi/4]$ so that the shading

$$Y_4(R) = \{x \in Y_3(R) : \text{L-spread}(x) \in [\theta, 2\theta]\} \quad (4.14)$$

is $\gtrsim \delta^\eta$ dense. We argue as follows: if this was not the case, then

$$\begin{aligned} \sum_{R \in \mathcal{R}_0} |\{x \in Y_3(R) : \text{L-spread}(x) < \delta^{-\varepsilon^2/2} t\}| &\gtrsim \sum_{R \in \mathcal{R}_0} |Y_3(R)| \\ &\gtrsim \delta^{2\eta} \sum_{R \in \mathcal{R}} |Y_1(R)| \gtrsim \delta^{2\eta} \mu \left| \bigcup_{R \in \mathcal{R}} Y_1(R) \right| \\ &\geq \delta^{2\eta} \mu \left| \bigcup_{R \in \mathcal{R}_0} Y_3(R) \right|, \end{aligned}$$

and hence there must exist a point $x \in \bigcup_{R \in \mathcal{R}_0} Y_3(R)$ with

$$\#\{R \in \mathcal{R}_0 : x \in Y_3(R)\} \gtrsim \delta^{2\eta} \mu \geq \delta^{-\varepsilon^2+5\eta} (st\#\mathcal{R}), \quad \text{L-spread}(x) \leq \delta^{-\varepsilon^2/2} t, \quad \text{P-spread}(x) \leq \frac{s_1}{t}. \quad (4.15)$$

(the last condition follows from the fact that $x \in \bigcup Y_3(R)$). But this means that the rectangular prisms $\{R \in \mathcal{R}_0 : x \in Y_3(R)\}$ are contained in the $O(1)$ dilate of a rectangular prism of dimensions $s_1 \times \delta^{-\varepsilon^2/2} t \times 1$. Since the prisms in \mathcal{R} satisfy the Convex Wolff Axioms with error $\delta^{-\eta}$, we have

$$\#\{R \in \mathcal{R}_0 : x \in Y_3(R)\} \lesssim \delta^{-\eta} \left(s_1 \cdot \delta^{-\varepsilon^2/2} t \right) (\#\mathcal{R}) = \delta^{-3\eta/\varepsilon - \eta - \varepsilon^2/2} (st\#\mathcal{R}), \quad (4.16)$$

Comparing (4.15) and (4.16), we obtain a contradiction if δ_0 and η are sufficiently small.

Step 5. At this point we can fix $\theta \geq \delta^{-\varepsilon^2/2}t$, and we may suppose that the corresponding shading Y_4 from (4.14) is $\gtrsim \delta^\eta$ dense. For each prism $R \in \mathcal{R}_0$, let \hat{R} be the prism of dimensions $\frac{s_1\theta}{t} \times \theta \times 1$ obtained by anisotropically dilating R . We claim that if $R_0, R \in \mathcal{R}_0$ are prisms and if $x \in Y_4(R_0) \cap Y_3(R)$, then $R \subset 2\hat{R}_0$. This follows from the fact that $\angle(\Pi(R_0), \Pi(R)) \leq \frac{s_1}{t}$, and $\angle(\ell(R_0), \ell(R)) \leq 2\theta$. In fact more is true—if we define $w = \max(\frac{s_1\theta}{t}, t)$, then $N_w(R)$ is comparable to a prism of dimensions $\frac{s_1\theta}{t} \times w \times 1$, and this prism is contained in $2\hat{R}_0$. Furthermore, $N_w(R) \cap R_0$ is comparable to a rectangular prism of dimensions $s \times t \times w/\theta$.

Let $\mathcal{R}_1 \subset \mathcal{R}_0$ consist of those rectangular prisms for which $|Y_4(R)| \gtrsim \delta^{2\eta}|R|$. Let $R_0 \in \mathcal{R}_1$; recall that R_0 is a $s \times t \times 1$ rectangular prism; for notational simplicity we will suppose $R_0 = [-s/2, s/2] \times [-t/2, t/2] \times [-1/2, 1/2]$, and that the intersection of $Y_4(R_0)$ with the z -axis has measure $\gtrsim \delta^\eta$. Define $\hat{R}_0 = [-\frac{s_1\theta}{t}, \frac{s_1\theta}{t}] \times [-\theta, \theta] \times [-1, 1]$.

Select $\frac{w}{\theta}$ -separated points $z_1, \dots, z_N \in Y_4(R_0)$ on the z -axis, with $N \gtrsim \delta^\eta \frac{\theta}{w}$. We will label the points so that $|z_i - z_j| \geq \frac{w}{\theta}|i - j|$. For each index i , select a prism $R_i \in \mathcal{R}_0$ with $z_i \in Y_3(R_i)$ and $\angle((0, 0, 1), \ell(R_i)) \in [\theta, 2\theta]$.

For each index i , let R'_i be the $\frac{s_1\theta}{t}$ neighborhood of R_i , and let $Y'_3(R'_i) = N_{\frac{s_1\theta}{t}}(Y_3(R_i))$. Since $R_i \in \mathcal{R}_0$ (and hence $|Y_3(R_i)| \gtrsim \delta^\eta|R_i|$), we have

$$|Y'_3(R'_i)| \gtrsim \delta^\eta|R'_i|. \quad (4.17)$$

R'_i is comparable to a prism of dimensions $\frac{s_1\theta}{t} \times w \times 1$ whose coaxial line makes angle $\sim \theta$ with the z axis. We have

$$|R'_i \cap R'_j| \lesssim \left(\frac{s_1\theta}{t}\right) \frac{w}{1 + |i - j|},$$

and hence

$$\left\| \sum_{i=1}^N \chi_{Y'_3(R'_i)} \right\|_2^2 \leq \sum_{i,j=1}^N |R'_i \cap R'_j| \lesssim N \frac{s_1\theta w}{t} \left| \log\left(\frac{\theta}{w}\right) \right|.$$

We conclude that

$$\left| \bigcup_{i=1}^N Y'_3(R'_i) \right| \geq \frac{\left(\sum |Y'_3(R'_i)| \right)^2}{\left\| \sum \chi_{Y'_3(R'_i)} \right\|_2^2} \gtrsim \frac{\left(N \delta^\eta \frac{s_1\theta}{t} w \right)^2}{N \frac{s_1\theta w}{t}} = N \delta^{2\eta} \frac{s_1\theta}{t} w \gtrsim \delta^{3\eta} \frac{s_1}{t}. \quad (4.18)$$

On the other hand, each set R'_i is contained in \hat{R}_0 , and thus

$$\left| \hat{R}_0 \cap N_{\frac{s_1\theta}{t}} \left(\bigcup_{\substack{R \in \mathcal{R} \\ R \subset \hat{R}_0}} Y(R) \right) \right| \gtrsim \delta^{3\eta} |\hat{R}_0|. \quad (4.19)$$

Recall that \hat{R}_0 is a prism of dimensions $2\frac{s_1\theta}{t} \times 2\theta \times 2$, and $s_1 = \delta^{-3\eta/\varepsilon}s$. Define $s' = \frac{s_1\theta}{t}$ and $t' = \min(1, t\frac{s'}{s}) = \min(1, \delta^{-3\eta/\varepsilon}\theta)$. Then $s' \in [\delta^{-\varepsilon^2/2}s, \frac{s_1}{t}]$; since $s \leq \delta^\varepsilon t$, we can select η sufficiently small so that $\frac{s_1}{t} \leq \delta^{\varepsilon/2}$, and hence $s' \in [\delta^{-\varepsilon^2/2}s, \delta^{\varepsilon/2}]$.

By (4.19) we conclude that for each $R_0 \in \mathcal{R}_1$, if \hat{R}'_0 is the corresponding $s' \times t' \times 1$ prism and η is selected sufficiently small compared to ε and τ , then the shading $\hat{Y}'(\hat{R}'_0)$ given by (4.5) satisfies

$$|\hat{Y}'(\hat{R}'_0)| \gtrsim \delta^{3\eta} |\hat{R}'_0| \geq \delta^{3\eta+3\eta/\varepsilon} |\hat{R}'_0| \geq \delta^\tau |\hat{R}'_0|. \quad (4.20)$$

To finish the proof, define

$$\mathcal{R}' = \{\hat{R}'_0: \hat{R}_0 \in \mathcal{R}_1\}.$$

(4.20) ensures that $|Y'(R')| \geq \delta^\tau |R'|$ for each $R' \in \mathcal{R}'$. Since \mathcal{R} satisfies the Convex Wolff axioms with error $\delta^{-\eta}$ and $\#\mathcal{R}_1 \gtrsim \delta^{2\eta} \#\mathcal{R}$, we have that \mathcal{R}_1 satisfies the Convex Wolff axioms with error $\lesssim \delta^{-3\eta}$, and hence by Remark 2.7 we have that \mathcal{R}' satisfies the Convex Wolff Axioms with error $\lesssim \delta^{-3\eta}$; if δ_0 and η is selected sufficiently small then this is $\leq \delta^{-\tau}$. \square

The culmination of Section 4.1 is the following result, which is obtained by iterating Lemma 4.4.

Proposition 4.6. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $s, t \in [\delta, 1]$ with $s \leq \delta^\varepsilon t$. Let \mathcal{R} be a multi-set of $s \times t \times 1$ prisms contained in $B(0, 1)$, and suppose that \mathcal{R} satisfies the Convex Wolff Axioms with error $\delta^{-\eta}$. Let $\{Y(R), R \in \mathcal{R}\}$ be a δ^η -dense shading.*

Then $\bigcup_{R \in \mathcal{R}} Y(R)$ has discretized Assouad dimension at least $3 - \varepsilon$ at scale δ and scale separation $\delta^{-\eta}$.

Proof. Fix $\varepsilon > 0$ and let η, δ_0 be small quantities to be determined below. Let $\delta \in (0, \delta_0]$, $s, t \in [\delta, 1]$ with $s \leq \delta^\varepsilon t$; let \mathcal{R} be a set of $s \times t \times 1$ prisms that satisfy the Convex Wolff Axioms with error $\delta^{-\eta}$; and let $Y(R)$ be a δ^η -dense shading.

Define $\mathcal{R}_1 = \mathcal{R}$ and define the shading $Y_1 = Y$. Let $s_1 = s$ and $t_1 = t$. Let $\eta = \tau_1 < \tau_2 < \tau_3 < \dots < \tau_N$, $N = \lfloor 2/\varepsilon^2 \rfloor$ be small quantities to be specified below. We will perform the following iterative process for each $i = 1, 2, \dots, N$. At the beginning of step i , the following things will be true

- We have a set \mathcal{R}_i of $s_i \times t_i \times 1$ prisms, with $s_i \leq \delta^\varepsilon t_i$.
- \mathcal{R}_i satisfies the Convex Wolff axioms with error $\delta^{-\tau_i}$.
- We have a shading $\{Y_i(R): R \in \mathcal{R}_i\}$ that is δ^{τ_i} dense.
- For each $R_i \in \mathcal{R}_i$, we have

$$Y_i(R_i) \subset N_{s_i} \left(\bigcup_{R \in \mathcal{R}} Y(R) \right). \quad (4.21)$$

Apply Lemma 4.4 to (\mathcal{R}_i, Y_i) with ε as above and τ_{i+1} in place of τ . If $\tau_i > 0$ is sufficiently small depending on ε and τ_{i+1} , then the Lemma can be applied (indeed, τ_i must be less than or equal to the quantity “ η ” from the conclusion of Lemma 4.4). The quantity δ_0 (and hence δ) will be chosen sufficiently small so that Lemma 4.4 can be applied at each step.

Case (A). Suppose that Conclusion (A) from Lemma 4.4 holds, i.e. there exists $r \in [\delta^{-\tau_i} s_i, 1]$ and an r -ball B so that

$$\left| B \cap \bigcup_{R \in \mathcal{R}_i} Y_i(R_i) \right| \geq (s_i/r)^\varepsilon |B|.$$

Using (4.21), we conclude that

$$\left| B \cap N_{s_i} \left(\bigcup_{R \in \mathcal{R}} Y(R) \right) \right| \geq (s_i/r)^\varepsilon |B|,$$

i.e. $\bigcup_{R \in \mathcal{R}} Y(R)$ has discretized Assouad dimension $\geq 3 - \varepsilon$ at scale δ and scale separation $\delta^{-\tau_i} \geq \delta^{-\eta}$. If this occurs, then we halt the iteration and conclude the proof.

Case (B). Next, suppose that Conclusion (B) holds. Let $s_{i+1} = s'$; $t_{i+1} = t'_i$; let \mathcal{R}_{i+1} be the set of $s_{i+1} \times t_{i+1} \times 1$ prisms, with shading $Y_{i+1} = Y'_i$, as described by Lemma 4.4. We have that \mathcal{R}_{i+1} satisfies the Convex Wolff Axioms with error $\delta^{-\tau_{i+1}}$, and Y_{i+1} is $\delta^{\tau_{i+1}}$ dense. Furthermore, the shading Y_{i+1} satisfies (4.21) with $i + 1$ in place of i .

Case (B.i). Suppose that Conclusion (B) holds and $t_{i+1} = 1$. Recall that $s_{i+1} \leq \delta^{\varepsilon/2}$. Thus we can apply Lemma 4.3 with s_{i+1} in place of δ . We have

$$\left| N_{s_{i+1}} \left(\bigcup_{R \in \mathcal{R}} Y(R) \right) \right| \geq \left| \bigcup_{R \in \mathcal{R}_{i+1}} Y_{i+1}(R) \right| \gtrsim \delta^{3\tau_{i+1}} |\log s_{i+1}|^{-1} \gtrsim s_{i+1}^{7\tau_{i+1}/\varepsilon}. \quad (4.22)$$

We can select τ_{i+1} and δ_0 sufficiently small so that the LHS of (4.22) has size at least $s_{i+1}^\varepsilon |B(0, 1)|$. Since the set on the LHS of (4.22) is contained in $B(0, 1)$, we conclude that $\bigcup_{R \in \mathcal{R}} Y(R)$ has discretized Assouad dimension $\geq 3 - \varepsilon$ at scale δ and scale separation $\delta^{-\varepsilon/2} \geq \delta^{-\eta}$. If this occurs, then we halt the iteration and conclude the proof.

Case (B.ii). Suppose $t_{i+1} < 1$; then $s_{i+1}/t_{i+1} = s_i/t_i$, so in particular $s_{i+1} \leq \delta^\varepsilon t_{i+1}$. Thus we have verified all of the properties required to continue the iteration, with $i + 1$ in place of i . Note that $s_{i+1} \geq \delta^{-\varepsilon^2/2} s_i$; $t_{i+1} \geq \min(1, \delta^\varepsilon s_{i+1})$, and the iterative process halts if $t_{i+1} = 1$. We conclude that this iterative process will halt after at most $N = \lfloor 2/\varepsilon^2 \rfloor$ steps. \square

4.2 Arrangements of tubes that cluster into rectangular prisms

In Section 4.1, we showed that unions of rectangular prisms that are far from being tubes (i.e. prisms of dimensions $s \times t \times 1$, with s much smaller than t) have large discretized Assouad dimension. In this section, we will use these results to show that if an arrangement of δ -tubes clusters into prisms that are far from being (fatter) tubes, then the union of these tubes must have large discretized Assouad dimension. This is made precise in Lemmas 4.8 and 4.10 below. We begin with a technical lemma.

Lemma 4.7. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $\rho \in [20\delta, 1]$ and let $K \geq 1$. Let $(\mathbb{T}, Y)_\delta$ be a set of essentially distinct δ -tubes, with $\#\mathbb{T} \geq K\rho/\delta$, and Y a uniformly δ^η -dense shading (see Definition 2.1). Let R_0 be a $10\delta \times \rho \times 1$ rectangle, and suppose each $T \in \mathbb{T}$ is contained in R_0 .*

Then there exists a number $K\delta \lesssim \omega \leq \rho$ and a set \mathcal{R} of essentially distinct $10\delta \times \omega \times 1$ prisms, so that

$$\# \left(\bigcup_{R \in \mathcal{R}} \{T \in \mathbb{T} : T \subset R\} \right) \gtrsim \#\mathbb{T}, \quad (4.23)$$

and for each prism $R \in \mathcal{R}$ we have

$$\left| \bigcup_{\substack{T \in \mathbb{T} \\ T \subset R}} Y(T) \right| \geq \delta^\varepsilon |R|. \quad (4.24)$$

Proof. For each $x \in \mathbb{R}^3$, define the ‘‘line spread’’

$$\text{L-spread}(x) = \max \{ \angle(T, T') \},$$

where the maximum is taken over all pairs of tubes T, T' intersecting x . We define $\text{L-spread}(x) = 0$ if at most one tube intersects x . Let C be a large number to be determined below. After dyadic pigeonholing, we can select $\theta \in [\frac{K\delta}{C}, \rho]$ so that the shading

$$Z(T) = \{x \in T : \text{L-spread}(x) \in [\theta, 2\theta]\}$$

satisfies

$$\sum_{T \in \mathbb{T}} |Z(T)| \geq (C |\log \delta|)^{-1} \delta^2 (\#\mathbb{T}). \quad (4.25)$$

(Note that the shading $Z(T)$ is not necessarily a subset of $Y(T)$). Our argument that such a shading $Z(T)$ must exist proceeds as follows: if we could not select a $\theta \in [\frac{K\delta}{C}, \rho]$ so that (4.25) holds, then the shading

$$Z'(T) = \{x \in T : \text{L-spread}(x) \leq \frac{K\delta}{C}\}$$

satisfies $\sum_{T \in \mathbb{T}} |Z'(T)| \geq \frac{1}{100} \delta^2 (\#\mathbb{T})$. But for each $x \in \mathbb{R}^3$, the set of tubes $T \in \mathbb{T}$ with $x \in Z'(T)$ are contained in the 10δ -neighborhood of a plane (since they are contained in R_0), and also make angle $\leq \frac{K\delta}{C}$ with a common unit vector; since the tubes in \mathbb{T} are essentially distinct, we conclude that $O(K/C)$ tubes $T \in \mathbb{T}$ can satisfy $x \in Z'(T)$, and hence

$$\left| \bigcup_{T \in \mathbb{T}} T \right| \geq \left| \bigcup_{T \in \mathbb{T}} Z'(T) \right| \gtrsim (C/K) \sum_{T \in \mathbb{T}} |Z'(T)| \gtrsim C\rho\delta. \quad (4.26)$$

If C is selected sufficiently large then this is impossible, since the set on the LHS of (4.26) is contained in R_0 , which has volume $10\rho\delta$.

At this point we have established the existence of an angle $\theta \in [\frac{K\delta}{C}, \rho]$ so that the associated shading $Z(T)$ satisfies (4.25). In the arguments that follow, the quantity C has been fixed, and all implicit constants are allowed to depend on C . Let

$$\mathbb{T}_1 = \{T \in \mathbb{T} : |Z(T)| \gtrsim |\log \delta|^{-1} |T|\}. \quad (4.27)$$

Selecting the implicit constant in (4.27) appropriately, we can ensure that $\#\mathbb{T}_1 \gtrsim \#\mathbb{T}$. For each $T_1 \in \mathbb{T}_1$, let x_2, \dots, x_N , $N \gtrsim \theta/\delta$ be a set of δ/θ -separated points from $Y_1(T_1)$. For each x_i , select $T_i \in \mathbb{T}$ with $x_i \in T_i$ and $\angle(T_1, T_i) \in [\theta, 2\theta)$. A L^2 argument (see e.g. (4.11)) plus the fact that $|Y(T_i)| \geq \delta^\eta$ shows that

$$\left| \bigcup_{i=2}^N Y(T_i) \right| \gtrsim \delta^{2\eta} \sum_{i=2}^N |T_i| \gtrsim \delta^{2\eta} \delta\theta. \quad (4.28)$$

On the other hand, the set on the LHS of (4.28) is contained in $R_0 \cap N_{2\theta}(T_1)$, which is essentially a rectangular prism of dimensions $10\delta \times 4\theta \times 1$. Denote this prism by $R(T_1)$. We have just shown that

$$\left| \bigcup_{\substack{T \in \mathbb{T} \\ T \subset R(T_1)}} Y(T) \right| \gtrsim \delta^{2\eta} |R(T_1)|. \quad (4.29)$$

To conclude the proof, let $\omega = 4\theta$ and let $\mathcal{R}_1 = \{R(T_1) : T_1 \in \mathbb{T}_1\}$. We claim that (4.23) holds for \mathcal{R}_1 . Indeed, this follows from the fact that $T_1 \subset R(T_1)$ for each $T_1 \in \mathbb{T}_1$, and $\#\mathbb{T}_1 \gtrsim \#\mathbb{T}$. By (4.29) we see that (4.24) holds for each $R \in \mathcal{R}_1$, provided we select η and δ_0 sufficiently small. To conclude the proof, greedily select a set $\mathcal{R} \subset \mathcal{R}_1$ of essentially distinct rectangular prisms; we can perform this selection so that (4.23) continues to hold. \square

Lemma 4.8. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let \mathbb{T} be a set of essentially distinct δ -tubes that satisfies the Convex Wolff Axioms with error $\delta^{-\eta}$. Let Y be a shading with $|Y(T)| \geq \delta^\eta |T|$ for each tube T . Then at least one of the following must occur.*

- (A) *There is a set $\mathbb{T}' \subset \mathbb{T}$ with $\#\mathbb{T}' \geq \frac{1}{2}(\#\mathbb{T})$ so that for every $\delta \leq \rho \leq 1$ and every $\delta \times \rho \times 1$ rectangle R , at most $\delta^{-\varepsilon}(\rho/\delta)$ tubes from \mathbb{T}' are contained in $2R$.*
- (B) *$\bigcup Y(T)$ has discretized Assouad dimension at least $3 - \varepsilon$ at scale δ and scale separation $\delta^{-\eta}$.*

Remark 4.9. Since \mathbb{T} satisfies the Convex Wolff Axioms, we have that at most $\delta^{-\eta}(\rho\delta)(\#\mathbb{T})$ tubes are contained in each $2\delta \times 2\rho \times 2$ rectangle. If $\#\mathbb{T}$ is substantially larger than δ^{-2} , then the estimate from Conclusion (A) is stronger.

Proof. We say a $2\delta \times 2\rho \times 2$ rectangle is “heavy” if it contains more than $\delta^{-\varepsilon}(\rho/\delta)$ tubes from \mathbb{T} . If fewer than $\frac{1}{2}(\#\mathbb{T})$ of the tubes are contained in heavy rectangles, let \mathbb{T}' be the set of tubes that are not contained in heavy rectangles. Then Conclusion (A) holds, and we are done.

Suppose instead that at least $\frac{1}{2}(\#\mathbb{T})$ tubes are contained in heavy rectangles. Since the tubes in \mathbb{T} are essentially distinct, each $2\delta \times 2\rho \times 2$ rectangle can contain at most $O((\rho/\delta)^2)$ tubes from \mathbb{T} . Thus if a $2\delta \times 2\rho \times 2$ rectangle is heavy, then we must have $\rho \gtrsim \delta^{1-\varepsilon}$. Let \mathcal{R}_0 denote this set of heavy rectangles (note that the rectangles in \mathcal{R}_0 can have different dimensions). Let ε_1 be a small quantity to be chosen below. Apply Lemma 4.7 to each rectangle $R_0 \in \mathcal{R}_0$, with ε_1 in place of ε , and $K = \delta^{-\varepsilon}$. We obtain a number $\omega(R_0) \gtrsim \delta^{1-\varepsilon}$ and a set $\mathcal{R}(R_0)$ of $\delta \times \omega(R_0) \times O(1)$ rectangles. We have

$$\#\left(\bigcup_{R_0 \in \mathcal{R}_0} \bigcup_{R \in \mathcal{R}(R_0)} \{T \in \mathbb{T}: T \subset R\}\right) \gtrsim \#\mathbb{T}. \quad (4.30)$$

After dyadic pigeonholing, we can select $\omega_0 \in [\delta^{1-\varepsilon}, 1]$ so that if we define $\mathcal{R}_1 = \{R_0 \in \mathcal{R}_0: \omega(R_0) \in [\omega_0, 2\omega_0]\}$, then (4.30) continues to hold with \mathcal{R}_1 in place of \mathcal{R}_0 .

Define the multi-set

$$\mathcal{R}_2 = \bigsqcup_{R_1 \in \mathcal{R}_1} \mathcal{R}(R_1).$$

(In the above, \bigsqcup is used to suggest that if the same rectangle is present in multiple sets $\mathcal{R}(R_1)$, then this rectangle should occur multiple times in \mathcal{R}_2 ; i.e. the union might create a multi-set). Let

$$\mathbb{T}_2 = \bigcup_{R \in \mathcal{R}_2} \{T \in \mathbb{T}: T \subset R\}.$$

Since \mathbb{T} satisfies the Convex Wolff Axioms with error $\delta^{-\eta}$, by (4.30) we have that the set (i.e. not multi-set) \mathbb{T}_2 satisfies the Convex Wolff Axioms with error $\lesssim \delta^{-\eta}$. Finally, for each $T \in \mathbb{T}_2$, let $R(T)$ be a rectangle in \mathcal{R}_2 with $T \subset R$ (if more than one $R \in \mathcal{R}_2$ contains T , simply select one of them). Define the multi-set $\mathcal{R} = \{R(T): T \in \mathbb{T}_2\}$, i.e. \mathcal{R} is a multi-set with the same cardinality as \mathbb{T}_2 . By Remark 2.7, we conclude that \mathcal{R} satisfies the Convex Wolff Axioms with error $\lesssim \delta^{-\eta}$.

For each $R \in \mathcal{R}$, define the shading

$$Y(R) = R \cap \bigcup_{T \in \mathbb{T}} Y(T).$$

By Lemma 4.7, we have $|Y(R)| \geq \delta^{\varepsilon_1} |R|$ for each $R \in \mathcal{R}$. To conclude the proof, apply Proposition 4.6 with ε as above, $s = \delta$, and $t = \omega_0$; to ensure that Proposition 4.6 can be applied, we must select

ε_1 sufficiently small depending on ε (our choice of ε_1 depends on the quantity “ η ” from Proposition 4.6), and then η sufficiently small depending on ε_1 . Since $\bigcup Y(T) \subset \bigcup Y(R)$, we conclude that Conclusion (B) holds. \square

Lemma 4.8 says that either a union of δ -tubes has large discretized Assouad dimension, or these tubes cannot concentrate into $\delta \times \rho \times 1$ rectangular prisms. The next lemma is similar, except the latter conclusion is strengthened: either a union of δ -tubes has large discretized Assouad dimension, or these tubes cannot concentrate into $s \times t \times 1$ rectangular prisms, for any $\delta \leq s \leq t \leq 1$. To prove this result, we apply Lemma 4.8 to the s -thickening of the tubes in our arrangement, for many different values of s . The precise statement is as follows.

Lemma 4.10. *For all $\varepsilon = 1/N > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $(\mathbb{T}, Y)_\delta$ be a set of essentially distinct δ -tubes that satisfies the Convex Wolff Axioms with error $\delta^{-\eta}$, and let Y be a uniformly δ^η -dense shading. Then at least one of the following must occur.*

(A) *There is a set $\mathbb{T}' \subset \mathbb{T}$ with $\#\mathbb{T}' \geq \delta^\varepsilon(\#\mathbb{T})$ so that the following holds. For every pair of numbers $\delta \leq s \leq t \leq 1$ with s of the form $\delta^{i\varepsilon}$, there is a 1-uniform partitioning cover \mathbb{T}'_s of \mathbb{T}' (see Definition 2.10), where \mathbb{T}'_s is a set of essentially distinct s -tubes. Furthermore, for every $s \times t \times 1$ rectangular prism R , at most $\delta^{-\varepsilon} \frac{t}{s} \frac{(\#\mathbb{T}')}{(\#\mathbb{T}'_s)}$ tubes from \mathbb{T}' are contained in R .*

(B) $\bigcup Y(T)$ has discretized Assouad dimension at least $3 - \varepsilon$ at scale δ and scale separation $\delta^{-\eta}$.

Proof. Step 1.

Our first step is to regularize the set \mathbb{T} at scales of the form $s_i = \delta^{i/N}$, with $i = 1, \dots, N$ (we do not need to consider $s_0 = 1$, since in that case there is nothing to prove). After repeated pigeonholing, we can select a set $\mathbb{T}^\dagger \subset \mathbb{T}$ with $\#\mathbb{T}^\dagger \gtrsim (C|\log \delta|)^N (\#\mathbb{T})$ and sets of s_i -tubes $\mathbb{T}^\dagger_{s_i}$, $i = 0, \dots, N$, so that $\mathbb{T}^\dagger_{s_N} = \mathbb{T}^\dagger$, and each set $\mathbb{T}^\dagger_{s_i}$ is a partitioning cover of $\mathbb{T}^\dagger_{s_{i+1}}$ (and hence also a partitioning cover of $\mathbb{T}^\dagger_{s_j}$ for all $j > i$). See e.g. Lemma 3.4 from [11] for an explicit description of this iterated pigeonholing process. We will choose δ_0 sufficiently small so that $\#\mathbb{T}^\dagger \geq \delta^{\eta/2}(\#\mathbb{T})$. In particular, by Remark 2.11 this means that \mathbb{T}^\dagger and each set $\mathbb{T}^\dagger_{s_i}$, $i = 1, \dots, N$ satisfies the Convex Wolff Axioms with error $\delta^{-2\eta}$.

Step 2.

We would like to apply Lemma 4.8 to each set $\mathbb{T}^\dagger_{s_N}, \dots, \mathbb{T}^\dagger_{s_1}$ in turn. If Conclusion (B) of Lemma 4.8 holds at any step, then we conclude that Conclusion (B) of Lemma 4.10 holds and we halt. If not, then few tubes from $\mathbb{T}^\dagger_{s_i}$ can cluster into a $s_i \times t \times 1$ prism; if each tube from $\mathbb{T}^\dagger_{s_i}$ contains the same number of tubes from \mathbb{T} , then Conclusion (A) of Lemma 4.10 holds, and we are done. The difficulty is that conclusion (A) of Lemma 4.10 only holds for a subset of the tubes in $\mathbb{T}^\dagger_{s_i}$; this complication introduces additional pigeonholing at each step, which in turn forces us to re-label all of the relevant objects. We now turn to the unpleasant details.

Define $\mathbb{T}_{s_N}^{(1)} = \mathbb{T}^\dagger_{s_N}$ and apply Lemma 4.8 with $\varepsilon/2$ in place of ε to $\mathbb{T}_{s_N}^{(1)}$, with the shading Y inherited from \mathbb{T} (recall that this shading is uniformly δ^η dense). If Conclusion (B) of Lemma 4.8 holds, then Conclusion (B) of Lemma 4.10 holds, and we are done.

If Conclusion (A) of Lemma 4.8 holds, then we obtain a set $(\mathbb{T}_{s_N}^{(1)})' \subset \mathbb{T}_{s_N}^{(1)}$ with $\#(\mathbb{T}_{s_N}^{(1)})' \geq \frac{1}{2}(\#\mathbb{T}_{s_N}^{(1)})$, so that for each $t \in [\delta, 1]$ and each $2s_N \times 2t \times 2$ rectangle R (recall $s_N = \delta$), at most $\delta^{-\varepsilon/2}(t/\delta)$ tubes from $(\mathbb{T}_{s_N}^{(1)})'$ are contained in R . After dyadic pigeonholing and refining $(\mathbb{T}_{s_N}^{(1)})'$, we

can select sets $\mathbb{T}_{s_N}^{(2)} \subset (\mathbb{T}_{s_N}^{(1)})'$ and $\mathbb{T}_{s_{N-1}}^{(2)} \subset \mathbb{T}_{s_{N-1}}^\dagger$ so that $\#\mathbb{T}_{s_N}^{(2)} \gtrsim |\log \delta|^{-1}(\mathbb{T}_{s_N}^{(1)})'$, and $\#(\mathbb{T}_{s_N}^{(2)}[T_{s_{N-1}}])$ is the same for each $T_{s_{N-1}} \in \mathbb{T}_{s_{N-1}}^{(2)}$, i.e. $\mathbb{T}_{s_{N-1}}^{(2)}$ is a 1-balanced partitioning cover of $\mathbb{T}_{s_N}^{(2)}$.

Next, we will describe a slightly more elaborate version of the above process. Let $2 \leq k \leq N$, and suppose we have constructed sets $\mathbb{T}_{s_N}^{(k)}, \dots, \mathbb{T}_{s_{N-k+1}}^{(k)}$, that have the following properties:

- (a) For $N - k + 1 \leq i < N$, $\mathbb{T}_{s_i}^{(k)}$ is a 1-balanced partitioning cover of $\mathbb{T}_{s_{i+1}}^{(k)}$ (and hence also a 1-balanced partitioning cover of $\mathbb{T}_{s_j}^{(k)}$ for $j > i$).
- (b) $\#\mathbb{T}_{s_N}^{(k)} \gtrsim_k |\log \delta|^{-k}(\#\mathbb{T}_{s_N}^{(1)})$.
- (c) For $N - k + 2 \leq i \leq N$ and $s_i \leq t \leq 1$, each $s_i \times t \times 1$ rectangle R contains $\lesssim_k |\log \delta|^k \delta^{-\varepsilon/2} \frac{\#\mathbb{T}_{s_N}^{(k)}}{\#\mathbb{T}_{s_i}^{(k)}}$ tubes from $\mathbb{T}_N^{(k)}$.

We construct sets $\mathbb{T}_{s_N}^{(k+1)}, \dots, \mathbb{T}_{s_{N-k}}^{(k+1)}$ as follows. For each $T_{s_{N-k+1}} \in \mathbb{T}_{s_{N-k+1}}^{(k)}$, define the shading

$$Y(T_{s_{N-k+1}}) = T_{s_{N-k+1}} \cap N_{s_{N-k+1}} \left(\bigcup_{T \in \mathbb{T}} Y(T) \right).$$

Since $\mathbb{T}[T_{s_{N-k+1}}]$ is non-empty and each tube $T \in \mathbb{T}$ has a δ^η -dense shading, we conclude that $|Y(T_{s_{N-k+1}})| \gtrsim \delta^\eta |T_{s_{N-k+1}}|$, i.e. the shading is uniformly $\gtrsim \delta^\eta$ -dense. Since $s_{N-k+1} \geq \delta^{1/N}$, this means that the shading is uniformly $s_{N-k+1}^{\eta N}$ dense. We will select η and δ_0 sufficiently small so that Lemma 4.8 can be applied to the pair $(\mathbb{T}_{s_{N-k+1}}^{(k)}, Y)_{s_{N-k+1}}$ with s_{N-k+1} in place of δ , and $\varepsilon/2$ in place of ε . Let $\eta_1 = \eta_1(\varepsilon) > 0$ be the output of that lemma.

If Conclusion (B) of Lemma 4.10 holds, then $\bigcup_{T_{s_{N-k+1}} \in \mathbb{T}_{s_{N-k+1}}^{(k)}} Y(T_{s_{N-k+1}})$ has discretized Assouad dimension at least $3 - \varepsilon/2$ at scale s_{N-k+1} and scale separation $s_{N-k+1}^{-\eta_1}$. But this means that $\bigcup_{T \in \mathbb{T}} Y(T)$ has discretized Assouad dimension at least $3 - \varepsilon/2$ at scale δ and scale separation $\delta^{-\eta_1/N}$; if we select $\eta \leq \eta_1/N$, then Conclusion (B) of Lemma 4.10 holds, and we are done.

Suppose instead that Conclusion (A) of Lemma 4.10 holds. Then there exists a set $(\mathbb{T}_{s_{N-k+1}}^{(k)})' \subset \mathbb{T}_{s_{N-k+1}}^{(k)}$ with $\#(\mathbb{T}_{s_{N-k+1}}^{(k)})' \geq \frac{1}{2}(\#\mathbb{T}_{s_{N-k+1}}^{(k)})$, so that for each $s_{N-k+1} \leq t \leq 1$ and each $2s_{N-k+1} \times 2t \times 2$ rectangle R , at most $s_{N-k+1}^{-\varepsilon/2} \frac{t}{s_{N-k+1}} \leq \delta^{-\varepsilon/2} \frac{t}{s_{N-k+1}}$ tubes from $(\mathbb{T}_{s_{N-k+1}}^{(k)})'$ are contained in R .

After dyadic pigeonholing and refining $(\mathbb{T}_{s_{N-k+1}}^{(k)})'$, we can select sets $\mathbb{T}_{s_{N-k+1}}^{(k+1)} \subset (\mathbb{T}_{s_{N-k+1}}^{(k)})'$ and $\mathbb{T}_{s_{N-k}}^{(k+1)} \subset \mathbb{T}_{s_{N-k}}^\dagger$ so that $\#\mathbb{T}_{s_{N-k+1}}^{(k+1)} \gtrsim |\log \delta|^{-1}(\mathbb{T}_{s_{N-k+1}}^{(k)})'$, and $\mathbb{T}_{s_{N-k}}^{(k+1)}$ is a 1-balanced partitioning cover of $\mathbb{T}_{s_{N-k+1}}^{(k+1)}$. For each $N - k + 1 < i \leq N$, define $\mathbb{T}_{s_i}^{(k+1)}$ to be those tubes from $\mathbb{T}_{s_i}^{(k)}$ that are contained in some tube from $\mathbb{T}_{s_{N-k+1}}^{(k+1)}$; it is still the case that \mathbb{T}_{s_i} is a 1-balanced partitioning cover of $\mathbb{T}_{s_{i+1}}$, for $N - k + 1 \leq i < N$, and by construction it is also the case that $\mathbb{T}_{s_{N-k}}^{(k+1)}$ is a 1-balanced partitioning cover of $\mathbb{T}_{s_{N-k+1}}^{(k+1)}$.

Furthermore, $\#\mathbb{T}_{s_N}^{(k+1)} \gtrsim |\log \delta|^{-1}(\#\mathbb{T}_{s_N}^{(k)})$ (this is because the cardinality of $\mathbb{T}_{s_{N-k+1}}^{(k)}$ was reduced by a factor of $\lesssim |\log \delta|^{-1}$, and $\mathbb{T}_{s_{N-k+1}}^{(k)}$ was a 1-balanced partitioning cover of $\mathbb{T}_{s_N}^{(k)}$), and thus Item (c) remains true for $k+1$ in place of j , and Item (c) remains true for $N - k + 2 \leq i \leq N$, with $k+1$ in place of k (indeed, the cardinality of $\mathbb{T}_{s_N}^{(k+1)}$ has decreased by at most a multiplicative factor of

$O(|\log \delta|)$ compared to the cardinality of $\mathbb{T}_{s_N}^{(k)}$, and the cardinality of $\mathbb{T}_{s_i}^{(k+1)}$ is no larger than that of $\mathbb{T}_{s_i}^{(k)}$

All that remains is to verify Item (c) above for $i = N - k + 1$. Let $s_{N-k+1} \leq t \leq 1$, let R be a $s_{N-k+1} \times t \times 1$ rectangle, and let $R' = 2R$. We already know that at most $\delta^{-\varepsilon/2} \frac{t}{s_{N-k+1}}$ tubes from $\mathbb{T}_{s_{N-k+1}}^{(k+1)}$ can be contained in the R' . On the other hand, if T is a δ -tube that is contained in R and also contained in a s_{N-k+1} tube $T_{s_{N-k+1}}$, then $T_{s_{N-k+1}} \subset 2R$. Since $\mathbb{T}_{s_{N-k+1}}^{(k+1)}$ is a 1-balanced cover of $\mathbb{T}_{s_N}^{(k+1)}$, each tube in $\mathbb{T}_{s_{N-k+1}}^{(k+1)}$ contains $\frac{\#\mathbb{T}_{s_N}^{(k+1)}}{\#\mathbb{T}_{s_{N-k+1}}^{(k+1)}}$ tubes from $\mathbb{T}_{s_N}^{(k+1)}$. We conclude that $\lesssim \delta^{-\varepsilon/2} \frac{t}{s_{N-k+1}} \frac{\#\mathbb{T}_{s_N}^{(k+1)}}{\#\mathbb{T}_{s_{N-k+1}}^{(k+1)}}$ tubes from $\mathbb{T}_{s_N}^{(k+1)}$ are contained in R . This completes the $k \rightarrow k+1$ step of the above iteration.

Step 3.

If the iterative process described above does not halt before the N -th step, then define $\mathbb{T}' = \mathbb{T}_{s_N}^{(N)}$. By selecting δ_0 sufficiently small we ensure that $\#\mathbb{T}' \geq \delta^{-\varepsilon} (\#\mathbb{T})$. For each $s = \delta^{i/N}$, $i = 1, \dots, N$, we have the 1-balanced partitioning cover $\mathbb{T}'_s = \mathbb{T}_{s_i}^{(N)}$. If we select δ_0 sufficiently small depending on ε , then Item (c) above ensures that for every $t \in [s, 1]$ and every $s \times t \times 1$ rectangular prism R , at most $\delta^{-\varepsilon} \frac{t}{s} \frac{(\#\mathbb{T}')}{(\#\mathbb{T}'_s)}$ tubes from \mathbb{T}' are contained in R . \square

4.3 Proof of Proposition 4.1

We are now ready to prove Proposition 4.1. For the reader's convenience, we reproduce it here

Proposition 4.1. *For all $\varepsilon > 0$, there exists $\tau > 0$, so that for all $\eta_1 > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $\beta \geq 2$ and let $(\mathbb{T}, Y)_\delta$ be a set of $\delta^{-\beta}$ essentially distinct δ -tubes that satisfy the Convex Wolff Axioms with error δ^η , with Y a δ^η uniformly dense shading. Then at least one of the following must be true.*

- (A) $\bigcup Y(T)$ has discretized Assouad dimension at least $3 - \varepsilon$ at scale δ , with scale separation $\delta^{-\eta}$.
- (B) There exists $\mathbb{T}' \subset \mathbb{T}$ with $\#\mathbb{T}' \geq \delta^{\eta_1} (\#\mathbb{T})$, and \mathbb{T}' satisfies the self-similar Convex Wolff Axioms with error $\delta^{-\varepsilon}$.
- (C) There exists $\rho \in [\delta, \delta^\varepsilon]$ and a set $\tilde{\mathbb{T}}$ of ρ -tubes, with the following properties.
 - (C.i) The tubes in $\tilde{\mathbb{T}}$ are essentially distinct, and $\#\tilde{\mathbb{T}} \geq \rho^{-\beta-\tau}$.
 - (C.ii) $\tilde{\mathbb{T}}$ satisfies the Convex Wolff Axioms with error $\leq \rho^{-\eta_1}$.
 - (C.iii) For each $T_\rho \in \tilde{\mathbb{T}}$, there exists $T \in \mathbb{T}$ with $T \subset T_\rho$.
- (D) There exists $\rho \in [\delta^{1-\varepsilon}, 1]$, a ρ -tube T_ρ , and a set $\tilde{\mathbb{T}} \subset \mathbb{T}[T_\rho]$, so that the following holds.
 - (D.i) $\#\tilde{\mathbb{T}} \geq (\delta/\rho)^{-\beta-\tau}$.
 - (D.ii) The unit rescaling of $\tilde{\mathbb{T}}$ relative to T_ρ satisfies the Convex Wolff Axioms with error $(\delta/\rho)^{-\eta_1}$.

Proof. Step 1.

Let $\eta_2 = 1/N > 0$ be a small quantity to be chosen below. Apply Lemma 4.10 with η_2 in place of

ε . We conclude that either Conclusion (A) of Proposition 4.1 holds, there exists a set $\mathbb{T}' \subset \mathbb{T}$ that satisfies Conclusion (B) of Lemma 4.10.

Step 2.

Suppose there exists $\rho \in [\delta, \delta^\varepsilon]$ of the form $\rho = \delta^{i/N}$ so that $\#\mathbb{T}'_\rho \geq \delta^{-\beta-\tau}$ (here $\#\mathbb{T}'_\rho$ is a 1-uniform partitioning cover of \mathbb{T}' , coming from Conclusion (B) of Lemma 4.10). Then by Remark 2.11, \mathbb{T}'_ρ satisfies the Convex Wolff Axioms with error $\delta^{-2\eta} \leq \rho^{-2\eta/\varepsilon}$; this is at most $\rho^{-\eta_1}$, provided we choose $\eta \leq \varepsilon\eta_1/2$. Thus Conclusion (C) holds, with $\tilde{\mathbb{T}} = \mathbb{T}'_\rho$.

Step 3. Our goal in this step is to show that either Conclusion (D) of Proposition 4.1 holds, or for every $\rho \in [\delta, \delta^\varepsilon]$ of the form $\rho = \delta^{i/N}$, we have

$$\rho^{-\beta+\varepsilon/3} \leq \#\mathbb{T}'_\rho \leq \rho^{-\beta-\tau}, \quad \rho \in [\delta, \delta^\varepsilon]. \quad (4.31)$$

Note that the RHS of the above inequality was already established in Step 2. First, we consider the case where $\beta \leq 2 + \varepsilon/4$; in this case (4.31) will always hold. Indeed, let $\rho \in [\delta, \delta^\varepsilon]$ be of the form $\rho = \delta^{i/N}$. Then for each $T_\rho \in \mathbb{T}'_\rho$, we have

$$\frac{\#\mathbb{T}'}{\#\mathbb{T}'_\rho} = \#\mathbb{T}'[T_\rho] \leq \delta^{-2\eta} \rho^2 \#\mathbb{T}',$$

where the second inequality follows from the fact that \mathbb{T}' satisfies the Convex Wolff Axioms with error $\delta^{-2\eta}$. We conclude that $\#\mathbb{T}'_\rho \geq \delta^{2\eta} \rho^{-2}$. In particular, if $\beta \leq 2 + \varepsilon/4$, and if we select η sufficiently small, then

$$\#\mathbb{T}'_\rho \geq \delta^{2\eta} \rho^{-2} \geq \rho^{2\eta/\varepsilon+\varepsilon/4} \rho^{-\beta} \geq \rho^{-\beta+\varepsilon/3}, \quad (4.32)$$

and hence (4.31) holds.

Next we consider the case where $\beta > 2 + \varepsilon/4$. Suppose there exists some $\rho \in [\delta, \delta^\varepsilon]$ of the form $\rho = \delta^{i/N}$ for which $\#\mathbb{T}'_\rho < \rho^{-\beta+\varepsilon/4}$. Let $\rho \in [\delta, \delta^\varepsilon]$ be the smallest number of the form $\delta^{i/N}$ for which this is true. Note that $\rho \geq \delta^{1-\varepsilon/20}$, since the tubes in \mathbb{T}' are essentially distinct, and thus $\#\mathbb{T}'_\rho \geq (\delta/\rho)^4 (\#\mathbb{T}') \geq (\delta/\rho)^4 \delta^{-\beta+2\eta}$.

Let $T_\rho \in \mathbb{T}'_\rho$. By Conclusion (B) of Lemma 4.10, for each $s \in [\delta, \rho]$ of the form $\delta^{j/N}$, each $t \in [s, \rho]$ and each $s \times t \times 1$ rectangular prism $R \subset T_\rho$, we have

$$\begin{aligned} \#\{T \in \mathbb{T}'[T_\rho]: T \subset R\} &\leq \delta^{-\eta_2} \frac{t (\#\mathbb{T}')}{s (\#\mathbb{T}'_s)} \leq \delta^{-\eta_2} \frac{t \delta^{-\beta}}{s \rho^{-\beta+\varepsilon/4}} \left(\frac{s}{\rho}\right)^{\beta-\varepsilon/4} \\ &\leq \delta^{-\eta_2} \frac{t \delta^{-\beta}}{s \rho^{-\beta+\varepsilon/4}} \left(\frac{s}{\rho}\right)^2 \leq \delta^{-2\eta_2} \frac{t (\#\mathbb{T}')}{s (\#\mathbb{T}'_\rho)} \left(\frac{s}{\rho}\right)^2 \leq \delta^{-2\eta_2} \frac{st}{\rho^2} (\#\mathbb{T}'[T_\rho]), \end{aligned}$$

where the second inequality used the fact that $\#\mathbb{T}'_s \geq s^{-\beta+\varepsilon/4}$ and the minimality of ρ , the third inequality used the assumption $\beta - \varepsilon/4 \geq 2$, and the fourth inequality used the assumptions $\#\mathbb{T}' \geq \delta^{-\beta+\eta_1}$ and $\#\mathbb{T}'_\rho \leq \rho^{-\beta+\varepsilon/4}$.

Define $\tilde{\mathbb{T}} = \mathbb{T}'[T_\rho]$, and let $W \subset T_\rho$ be a convex set that contains at least one tube from $\tilde{\mathbb{T}}$. Then W is comparable to a $s \times t \times 1$ prism, for some $\delta \leq s \leq t \leq \rho$; increasing s (and possibly t by a factor of at most $\delta^{-1/N}$, we may suppose that s is of the form $\delta^{j/N}$. Thus we have

$$\#\{T \in \tilde{\mathbb{T}}: T \subset W\} \lesssim \delta^{-4\eta_2} \frac{|W|}{|T_\rho|} (\#\tilde{\mathbb{T}}) \leq (\delta/\rho)^{-40\eta_2/\varepsilon} \frac{|W|}{\rho^2} (\#\tilde{\mathbb{T}}).$$

If we select $\eta_1 < 40\eta_2\varepsilon$, then our set $\tilde{\mathbb{T}}$ satisfies Conclusion (D) from Proposition 4.1 and we are done.

Step 4. At this point, we have reduced to the case where (4.31) holds for all $\rho \in [\delta, \delta^\varepsilon]$ of the form $\rho = \delta^{i/N}$. We will show that Conclusion (B) holds. To begin, note that Items (i) and (iii) from Definition 2.15 hold for \mathbb{T}' : for each $\rho_0 \in [\delta, \delta^\varepsilon]$, we can select $\rho \in [\rho_0, \delta^{-\varepsilon}\rho_0]$ of the form $\rho = \delta^{i/N}$; \mathbb{T}_ρ is a partitioning cover of \mathbb{T}' , so Item (i) holds, and Item (iii) holds by (4.31) (if $\rho_0 > \delta^\varepsilon$ then we can select $\rho = 1$ and there is nothing to prove).

It remains to show that Item (ii) holds for \mathbb{T}' ; as explained above it suffices to consider ρ -tubes with $\rho \in [\delta, \delta^\varepsilon]$ of the form $\rho = \delta^{i/N}$, and convex sets R that are $s \times t \times 1$ prisms, with $s \in [\delta, \rho]$ of the form $s = \delta^{j/N}$ and $t \in [s, \rho]$. We would like to estimate the number of tubes from $\mathbb{T}'[T_\rho]$ contained in R .

By Conclusion (B) of Lemma 4.10, we have

$$\begin{aligned} \#\{T \in \mathbb{T}'[T_\rho]: T \subset R\} &\leq \delta^{-\eta_2} \frac{t (\#\mathbb{T}')}{s (\#\mathbb{T}'_s)} \lesssim \delta^{-\eta_2 - \varepsilon/4} t s (s^{\beta-2} \delta^{-\beta}) \\ &\lesssim \delta^{-\eta_2 - \varepsilon/4} t s \rho^{\beta-2} \delta^{-\beta} = \delta^{-\eta_2 - \varepsilon/4} \frac{t s}{\rho^2} \frac{\delta^{-\beta}}{\rho^{-\beta}} \leq \delta^{-2\eta_2 - \varepsilon/4} \frac{t s}{\rho^2} (\#\mathbb{T}''[T_\rho]). \end{aligned}$$

Selecting η_2 sufficiently small, we obtain

$$\#\{T \in \mathbb{T}'[T_\rho]: T \subset R\} \leq \delta^{-\varepsilon} \frac{|R|}{|T_\rho|} (\#\mathbb{T}'[T_\rho]).$$

We conclude that \mathbb{T} satisfies Condition (B) from Definition 2.15, and hence Conclusion (B) of Proposition 4.1 holds. \square

5 Sticky Kakeya and the Self-Similar Wolff Axioms

Our goal in this section is to prove Theorem 1.5. Our first task is to analyze the structure of extremal collections of tubes. If $\omega > 0$, then such collections must satisfy the Self-Similar Convex Wolff Axioms.

5.1 Extremal Kakeya sets satisfy the Self-Similar Convex Wolff Axioms

Proposition 5.1. *Suppose ω from Definition 3.1 is positive. Then for all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $(\mathbb{T}, Y)_\delta$ be η Assouad Extremal. Then there exists $\mathbb{T}' \subset \mathbb{T}$ so that $(\mathbb{T}', Y)_\delta$ is ε Assouad extremal, and \mathbb{T}' satisfies the Self-Similar Convex Wolff axioms with error $\delta^{-\varepsilon}$.*

Proof. Step 1. By decreasing ε if necessary, we can suppose that $\varepsilon < \omega/2$. Let $\tau > 0$ be the value from Proposition 4.1, with this choice of ε . Let η_1 be a small quantity to be chosen below (η_1 will depend on ω and ε), and let η', δ'_0 be the values from Proposition 4.1 for this choice of ε and η_1 .

Let η and δ_0 be small positive quantities to be chosen below. η and δ_0 will depend on the following quantities: $\omega, \varepsilon, \tau, \eta_1, \eta'$. In addition, δ_0 will depend on δ'_0 .

Let $\delta \in (0, \delta_0]$ and let $(\mathbb{T}, Y)_\delta$ be η Assouad Extremal. By pigeonholing, we can select a subset $\mathbb{T}_1 \subset \mathbb{T}$ with $\#\mathbb{T}_1 \gtrsim \delta^\eta (\#\mathbb{T})$ so that $|Y(T)| \geq \frac{1}{2} \delta^\eta |T|$ for each $T \in \mathbb{T}_1$. Then $\#\mathbb{T}_1 = \delta^{-\beta}$ for some

$\beta \geq \alpha - 2\eta$ (recall that α is defined in (3.4)), and \mathbb{T}_1 satisfies the Convex Wolff Axioms with error $\delta^{-2\eta}$. Furthermore, for all ρ_1, r_1 with $\rho_1 \leq \delta^\eta r_1$, and all balls B of radius r_1 , we have

$$\left| B \cap N_{\rho_1} \left(\bigcup_{T \in \mathbb{T}_1} Y(T) \right) \right| \leq (\rho_1/r_1)^{\omega-\eta} |B|. \quad (5.1)$$

Step 2. Apply Proposition 4.1 to $(\mathbb{T}_1, Y)_\delta$. We will show that Conclusion (A) from Proposition 4.1 cannot hold. Suppose to the contrary that Conclusion (A) holds, i.e. there exists $\rho, r \in [\delta, 1]$ with $\rho \leq \delta^{\eta'} r$ and a r -ball B such that

$$\left| B \cap N_\rho \left(\bigcup_{T \in \mathbb{T}_1} Y(T) \right) \right| \geq (\rho/r)^\varepsilon |B|. \quad (5.2)$$

If we select $\eta \leq \min(\eta', \omega_2)$ (recall that $\varepsilon \leq \omega/2$), then (5.2) contradicts (5.1). Hence Conclusion (A) cannot hold.

Step 3. We will show that Conclusion (C) from Proposition 4.1 cannot hold. Suppose to the contrary that Conclusion (C) holds, i.e. there exists $\rho \in [\delta, \delta^\varepsilon]$ and a set of ρ -tubes $\tilde{\mathbb{T}}$ satisfying Items (C.i), (C.ii) and (C.iii) from Conclusion (C). By Item (C.iii), we have that the shading

$$\tilde{Y}(\tilde{T}) = \tilde{T} \cap N_\rho \left(\bigcup_{T \in \mathbb{T}_1} Y(T) \right), \quad \tilde{T} \in \tilde{\mathbb{T}}$$

satisfies $|\tilde{Y}(\tilde{T})| \gtrsim \delta^\eta |\tilde{T}| \gtrsim \rho^{\eta/\varepsilon} |\tilde{T}|$ for each $\tilde{T} \in \tilde{\mathbb{T}}$; we will select $\eta < \eta' \varepsilon$ and δ_0 sufficiently small that $|\tilde{Y}(\tilde{T})| \leq \rho^{\eta'} |\tilde{T}|$ for each $\tilde{T} \in \tilde{\mathbb{T}}$.

Define

$$\gamma = \omega - \omega(\beta + \tau). \quad (5.3)$$

(In the above equation, $\omega(\beta + \tau)$ refers to the function ω , evaluated at the (positive) number $\beta + \tau$). Provided η_1 is selected sufficiently small (depending on τ), we have $\beta + \tau \geq \alpha - 2\eta_1 + \tau > \alpha$, and hence by the definition of α from (3.4), we have $\gamma > 0$. Hence if η_1 is selected sufficiently small, then

$$\limsup_{x \searrow 0} \omega(\eta_1, \eta_1, x, \beta + \tau) \leq \omega - \gamma/2,$$

i.e. there exists $x_0 > 0$ so that

$$\omega(\eta_1, \eta_1, x, \beta + \tau) \leq \omega - \gamma/4 \quad \text{for all } x \in (0, x_0].$$

Select δ_0 sufficiently small so that $\delta_0^\varepsilon \leq x_0$, and thus $\rho \in [\delta, \delta^\varepsilon]$ implies $\rho \leq x_0$. In particular, $\omega(\eta_1, \eta_1, \rho, \beta + \tau) \leq \omega - \gamma/4$, i.e. there exists $\rho_1, r_1 \in [\rho, 1]$ with $\rho_1 \leq \rho^{\eta_1} r_1$, and a r_1 -ball B such that

$$\left| B \cap N_{\rho_1} \left(\bigcup_{\tilde{T} \in \tilde{\mathbb{T}}} \tilde{Y}(\tilde{T}) \right) \right| \geq (\rho_1/r_1)^{\omega-\gamma/8} |B|. \quad (5.4)$$

But if we select $\eta \leq \min(\eta_1 \varepsilon, \gamma/9)$, then $\rho_1 \leq \delta^\eta r_1$, and (5.4) contradicts (5.1). Hence Conclusion (C) cannot hold.

Step 4. We will show that Conclusion (D) from Proposition 4.1 cannot hold. Suppose to the contrary that Conclusion (D) holds, i.e. there exists $\rho \in [\delta^{1-\varepsilon}, 1]$, a ρ -tube T_ρ , and a set $\tilde{\mathbb{T}} \subset \mathbb{T}_1[T_\rho]$ satisfying Items (D.i) and (D.ii) from Conclusion (D). Let $\tilde{\delta} = \delta/\rho \in [\delta, \delta^\varepsilon]$, and let $\tilde{\mathbb{T}}_{\tilde{\delta}}$ be unit

rescaling of $\tilde{\mathbb{T}}$ relative to T_ρ . Let $\tilde{Y}_\delta(\tilde{T}_\delta)$ be the image of $\tilde{Y}(\tilde{T})$ under this same rescaling. We have $|\tilde{Y}_\delta(\tilde{T})| \gtrsim \delta^\eta |\tilde{T}_\delta| \geq \tilde{\delta}^{\eta/\varepsilon} |\tilde{T}_\delta|$, and we will select η and δ_0 sufficiently small so that $|\tilde{Y}_\delta(\tilde{T}_\delta)| \geq \tilde{\delta}^{\eta_1} |\tilde{T}_\delta|$ for all $\tilde{T}_\delta \in \tilde{\mathbb{T}}_\delta$. The tubes in $\tilde{\mathbb{T}}_\delta$ are essentially distinct; by Item (D.i) we have $\#\tilde{\mathbb{T}}_\delta \geq \tilde{\delta}^{-\beta-\tau}$, and by Item (D.ii) we have that $\tilde{\mathbb{T}}_\delta$ satisfies the Convex Wolff Axioms with error $\lesssim \tilde{\delta}^{-\eta_1}$.

Arguing as in Step 3, we conclude that there exists $\rho_2, r_2 \in [\tilde{\delta}, 1]$ with $\rho_2 \leq \tilde{\delta}^{\eta_1} r_2$, and a r_2 -ball B_2 such that

$$\left| B_2 \cap N_{\rho_2} \left(\bigcup_{\tilde{T}_\delta \in \tilde{\mathbb{T}}_\delta} \tilde{Y}_\delta(\tilde{T}_\delta) \right) \right| \geq (\rho_2/r_2)^{\omega-\gamma/8} |B_2|.$$

Undoing the unit rescaling described above, we conclude there exists an ellipsoid E of dimensions $\rho r_2 \times \rho r_2 \times \rho$ so that

$$\left| E \cap N_{\rho r_2} \left(\bigcup_{\tilde{T} \in \tilde{\mathbb{T}}} \tilde{Y}(\tilde{T}) \right) \right| \geq (\rho_2/r_2)^{\omega-\gamma/8} |E|.$$

Indeed, the above inequality follows from the observation that the image of a ρ_2 -ball centered at x under the above rescaling is an ellipse of dimensions $\rho \rho_2 \times \rho \rho_2 \times \rho_2$, and this ellipse contains a $\rho \rho_2$ -ball centered at x . In particular, we can find a ρr_2 -ball B with $|B \cap E| \sim |B|$, so that

$$\left| B \cap N_{\rho r_2} \left(\bigcup_{\tilde{T} \in \tilde{\mathbb{T}}} \tilde{Y}(\tilde{T}) \right) \right| \gtrsim (\rho_2/r_2)^{\omega-\gamma/8} |B|. \quad (5.5)$$

Finally, observe that $\frac{\rho \rho_2}{\rho r_2} = \frac{\rho_2}{r_2} \leq \rho^{\eta_1} \leq \delta^{\eta_1 \varepsilon} \leq \delta^\eta$ (provided we select $\eta \leq \eta_1 \varepsilon$). Thus if we choose $\eta \leq \gamma/9$ and δ_0 sufficiently small, then (5.5) contradicts (5.1). Hence Conclusion (D) cannot hold.

Step 5. At this point, we shown that Conclusions (A), (C), and (D) from Proposition 4.1 cannot hold. Thus Conclusion (B) must hold. This completes the proof of Proposition 5.1. \square

5.2 Proof of Theorem 1.5

Our proof of Theorem 1.5 follows by combining two main ingredients. The first is Proposition 5.1. The second is a variant of the Stickyakeya Theorem, which was proved by the authors in [16]. The precise statement we need is as follows.

Theorem 5.2. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let \mathbb{T} be a set of δ -tubes that satisfy the Convex Wolff Axioms at every scale with error $\delta^{-\eta}$, and let $Y(T)$ be a δ^η dense shading. Then*

$$\left| \bigcup_{T \in \mathbb{T}} Y(T) \right| \geq \delta^\varepsilon. \quad (5.6)$$

As noted above, Theorem 5.2 is similar to the Stickyakeya Theorem proved by the authors in [16]. In Section 6 we will discuss the proof of Theorem 5.2, and how it differs from the arguments in [16]. Assuming this result for now, we can combine these two ingredients to prove Theorem 1.5 as follows

Proof of Theorem 1.5. It suffices to show that $\omega = 0$. Suppose instead that $\omega > 0$. Let η_0, δ_0 be the output from Theorem 5.2, with $\varepsilon = \omega/3$. By Proposition 5.1, there exists $\delta \in (0, \delta_0]$ and a set

$(\mathbb{T}', Y)_\delta$ with discretized Assouad dimension $\leq 3 - \omega/3$ that satisfies the Self-Similar Convex Wolff Axioms (and hence the Convex Wolff Axioms at every scale) with error $\delta^{-\eta_0}$, and for which $\{Y(T)\}$ is a δ^{η_0} dense shading. But then (6.2) says that $\bigcup Y(T)$ has discretized Assouad dimension at least $3 - \omega/3$, which is a contradiction. \square

Thus we have completed the proof of Theorem 1.5, except that it remains to prove Theorem 5.2; we will do so in the next section.

6 Makeya estimates for tubes satisfying the Convex Wolff Axioms at all scales

Our goal in this section is to prove Theorem 5.2. Theorem 5.2 is closely related to the discretized analogue of [16, Theorem 1.1], and specifically the estimate $\sigma_3 = 0$ from that paper; the latter result says that if \mathbb{T} is a collection of δ -tubes in \mathbb{R}^3 of cardinality roughly δ^{-2} that point in δ -separated directions, and if \mathbb{T} satisfies a property called “stickiness,” then the union of the tubes must have volume $\gtrsim 1$. Stickiness is a technical property, which roughly speaking says that for each $\rho \in [\delta, 1]$, the tubes in \mathbb{T} can be covered by a collection of ρ -tubes that point in ρ -separated directions (more accurately, the tubes point in ρ -separated directions up to multiplicity $\lesssim 1$).

Stickiness is a multi-scale property, in the following sense: if \mathbb{T} is sticky, then for each $\rho \geq \delta$ we can cover \mathbb{T} by a sticky collection of ρ -tubes. This multi-scale property was exploited throughout the arguments in [16]. Theorem 5.2 differs from [16, Theorem 1.1], because the stickiness hypothesis has been replaced by the hypothesis that the tubes satisfy the Convex Wolff axioms at every scale. As we will see below, however, this latter hypothesis is sufficient in order to repeat the arguments from [16]. Indeed, with two exceptions that will be explained below, the arguments from [16] can be repeated mutatis mutatis to obtain Theorem 5.2.

The proof of [16, Theorem 1.1] is divided into several sections. The paper begins by supposing that the result is false, and studying the structure of a (hypothetical) counter-example. In Section 2, the authors define what it means for a pair (\mathbb{T}, Y) of tubes and their associated shading to be a worst possible counter-example to [16, Theorem 1.1]; this is called an ε -extremal pair. In our discussion below, we will call it a ε -extremal pair. In [16], Section 3, it is shown that an ε -extremal pair (\mathbb{T}, Y) must look coarsely self similar at every intermediate scale $\rho \in [\delta, 1]$. More precisely, for every such ρ it is possible to cover the tubes in \mathbb{T} by a collection of ρ -tubes, so that the corresponding collection of ρ -tubes is a worst possible counter-example to [16, Theorem 1.1] at scale ρ , and the δ -tubes inside each ρ -tube also form a (re-scaled) worst-possible counter-example at scale δ/ρ . The analogous statement and proof in our setting are nearly identical; we will state the former and briefly sketch the latter.

In Sections 3, 4, and 5 of [16], the authors use the multi-scale self-similarity established in Section 2 of [16] to show that an extremal pair (\mathbb{T}, Y) must have certain structural properties called planiness and graininess, and in particular there must exist local grains that are described by a Lipschitz plane map, and global grains that are described by a C^2 slope function f ; this will be described in greater detail below. The arguments from [16] can be repeated without modification to establish the same conclusions in the present setting. In particular, the hypothesis from [16] that the tubes point in δ -separated directions has not been used up until this step.

In Section 6 of [16], the authors show that the derivative f' of the slope function has magnitude roughly 1 (and in particular, this quantity is bounded away from 0). It is during this step that

the authors use the hypothesis that the tubes point in δ -separated directions. With some small modifications, we can instead use the hypothesis that the tubes satisfy the Convex Wolff axioms at every scale; we will elaborate on this step in Section 6.4.

Finally, in Section 7 of [16], the authors recall an analogue of Wolff's circular maximal function estimate [19, 14], and they show how this estimate contradicts the assumption that there exists a collection of sticky tubes whose union has volume much smaller than 1, thereby completing the proof of the theorem. This section of [16] uses the assumption that the tubes point in different directions, in order to obtain a rich planar arrangement of curves to which Wolff's circular maximal theorem can be applied. With some modifications, the direction-separated assumption can be replaced with a weaker ball condition; we will explain this in detail in Sections 6.5 and 7.

6.1 Relevant definitions from [16]

Before describing the arguments in [16], it will be helpful to recall a few key definitions. First, a minor technical annoyance: tubes and cubes in [16] are defined slightly differently than in the present paper. In [16], a δ -tube in \mathbb{R}^3 is defined to be a set of the form $N_{\delta\delta}(\ell) \cap [-1, 1]^3$, where ℓ is a line in \mathbb{R}^3 . This definition is adopted for technical reasons, and for consistency with the results in [16] we will use this definition throughout Section (6); the distinction between this definition and our previous definition in Section 1 is not important in the arguments that follow. Second, in [16], a shading $Y(T) \subset T$ is defined to be a union of axis-aligned δ -cubes contained in T . Again, the distinction between this definition and our previous one from Section 2.1 is harmless: replace each tube T (in the sense of Section 1) and a each shading $Y(T)$ (in the sense of Section 2.1) with the corresponding tube in the sense of [16] and the the shadings consisting of the union of axis-aligned δ -cubes Q that satisfy $|Q \cap Y(T)| \geq \frac{1}{100}|Y(T)|/|T|$. This gives us a collection of tubes and shadings in the sense of [16], and if the conclusion of Theorem 1.5 holds for this latter collection of tubes and shadings, then it also holds for the original collection of tubes and shadings, except Inequality (1.1) has been weakened by an additional (harmless) factor of δ^η .

Tubes, covers, rescaling. We will often refer to a collection of δ -tubes \mathbb{T} , and a shading $Y(T)$ of these tubes. We will refer to this pair as $(\mathbb{T}, Y)_\delta$. We use $E_{\mathbb{T}}$ to denote the set $\bigcup_{T \in \mathbb{T}} Y(T)$; the shading Y will always be apparent from context. For $p \in \mathbb{R}^3$, we define

$$\mathbb{T}(p) = \{T \in \mathbb{T} : p \in Y(T)\}. \quad (6.1)$$

We say a pair $(\mathbb{T}', Y')_\delta$ is a *refinement* of $(\mathbb{T}, Y)_\delta$ if $\mathbb{T}' \subset \mathbb{T}$; $Y'(T') \subset Y(T')$ for each $T' \in \mathbb{T}'$; and $\sum_{\mathbb{T}'} |Y'(T')| \geq C^{-1} |\log \delta|^{-C} \sum_{\mathbb{T}} |Y(T)|$ for some absolute constant C (in practice, $C = 100$ will always suffice).

Let $0 < \delta \leq \rho \leq 1$. In [16], the authors say that a ρ -tube \tilde{T} *covers* a δ -tube T if their respective coaxial lines satisfy $d(\ell, \tilde{\ell}) \leq \rho/2$, where $d(\cdot, \cdot)$ is an appropriately chosen metric on the affine Grassmannian of lines in \mathbb{R}^3 ; see Section 2 of [16] for details. Note that if \tilde{T} covers T , then $T \subset \tilde{T}$, and hence this definition is consistent with ours from Definition 2.10 (the definition in [16] is slightly stronger than Definition 2.10, but the distinction is harmless).

If $(\mathbb{T}, Y)_\delta$ and $(\tilde{\mathbb{T}}, \tilde{Y})_\rho$ are pairs of δ (resp. ρ) tubes and their associated shadings, then we say $(\tilde{\mathbb{T}}, \tilde{Y})_\rho$ *covers* $(\mathbb{T}, Y)_\delta$ if each $T \in \mathbb{T}$ is covered by at least one $\tilde{T} \in \tilde{\mathbb{T}}$, and for each $\tilde{T} \in \tilde{\mathbb{T}}$ covering T , we have $Y(T) \subset \tilde{Y}(\tilde{T})$. If this is the case, we write $\mathbb{T}[\tilde{\mathbb{T}}]$ to denote the set of tubes in \mathbb{T} covered by $\tilde{\mathbb{T}}$. We say that $(\tilde{\mathbb{T}}, \tilde{Y})_\rho$ is a *balanced cover* of $(\mathbb{T}, Y)_\delta$ if it is a cover, and in addition $|E_{\mathbb{T}} \cap Q|$ is the same for each ρ -cube $Q \subset E_{\tilde{\mathbb{T}}}$.

Let $(\mathbb{T}, Y)_\delta$ be a set of δ -tubes and their associated shading. Let \tilde{T} be a ρ -tube, and suppose each tube in \mathbb{T} is covered by \tilde{T} . We define a new pair $(\hat{\mathbb{T}}, \hat{Y})_{\delta/\rho}$, which we call the *unit rescaling of $(\mathbb{T}, Y)_\delta$ relative to \tilde{T}* . Informally, our definition is as follows: consider the linear transformation that sends the coaxial line of \tilde{T} to the z axis, and dilates the x and y directions by ρ^{-1} ; the image of \tilde{T} under this transformation is comparable to the unit ball. The tubes in $\hat{\mathbb{T}}$ are those (δ/ρ) -tubes whose coaxial lines are the images of the coaxial lines of the tubes in \mathbb{T} under this transformation. The shadings $\hat{Y}(\hat{T})$ consist of the union of axis-aligned δ/ρ -cubes that intersect the image of $Y(T)$ under the above transformation. See Section 3 of [16] for details. As above, the definition of unit rescaling in [16] is slightly different than Definition 2.8, but the distinction is harmless.

The plane map. In the arguments in [16], the authors show that certain collections of tubes possess an important structural property called planiness. Let $(\mathbb{T}, Y)_\delta$ be a set of tubes and their associated shading. We say a function $V: E_{\mathbb{T}} \rightarrow S^2$ is a *plane map* for $(\mathbb{T}, Y)_\delta$ if

$$|\operatorname{dir}(T) \cdot V(p)| \leq \delta \quad \text{for all } (T, p) \in \mathbb{T} \times \mathbb{R}^3 \text{ with } p \in Y(T),$$

where $\operatorname{dir}(T) \in S^2$ is a unit vector parallel to the line coaxial with T .

Discretized Ahlfors-David regular sets. We say a set $E \subset \mathbb{R}$ is a (δ, α, C) -ADset if for all $\rho \geq \delta$, all $r \geq \rho$, and all intervals I of length r , we have $\mathcal{E}_\rho(E \cap I) \leq C(r/\rho)^\alpha$, where $\mathcal{E}_\rho(E \cap I)$ means the ρ -covering number of $E \cap I$. In practice we will have $\alpha \in (0, 1]$, and C will be a small power of $1/\delta$. See Section 4 of [16] for further discussion and motivation for this definition.

6.2 Extremal collection of tubes, and multi-scale structure

We are ready to begin the proof of Theorem 5.2. Suppose to the contrary that the result was false. Then there exists $\sigma' > 0$ so that for all $\eta, \delta_0 > 0$, there exists $\delta \in (0, \delta_0]$; a set \mathbb{T} of δ -tubes that satisfy the Convex Wolff Axioms at every scale with error $\delta^{-\eta}$; and a δ^η -dense shading $Y(T)$, so that

$$\left| \bigcup_{T \in \mathbb{T}} Y(T) \right| \leq \delta^{\sigma'}. \quad (6.2)$$

Let $\sigma > 0$ be the supremum of all numbers σ' with the above property; this supremum exists, since by hypothesis the set of admissible σ' is non-empty, and it is bounded above by 3. Our goal is to show that we must in fact have $\sigma = 0$, and thereby obtain a contradiction.

The following definition is the analogue of [16, Definition 3.1].

Definition 6.1. *Let $\varepsilon, \delta > 0$. We say a pair $(\mathbb{T}, Y)_\delta$ is ε -extremal if*

1. \mathbb{T} satisfies the Convex Wolff Axioms at all scales with error $\delta^{-\varepsilon}$.
2. Y is δ^ε -dense.
3. $\left| \bigcup_{T \in \mathbb{T}} Y(T) \right| \leq \delta^{\sigma-\varepsilon}$.

It is immediate from the above definitions that for every $\varepsilon, \delta_0 > 0$, there exists a ε -extremal collection of tubes $(\mathbb{T}, Y)_\delta$ for some $\delta \in (0, \delta_0]$; cf. [16, Lemma 3.1].

Our next task is to prove an analogue of [16, Proposition 3.2]. The precise statement is as follows.

Proposition 6.2. *For all $\varepsilon > 0$, there exists $\eta > 0$ and $\delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $(\mathbb{T}, Y)_\delta$ be a η -extremal collection of tubes, and let $\rho \in [\delta^{1-\varepsilon}, \delta^\varepsilon]$. Then there is a refinement $(\mathbb{T}', Y')_\delta$ of $(\mathbb{T}, Y)_\delta$ and a balanced cover $(\tilde{\mathbb{T}}, \tilde{Y})_\rho$ of $(\mathbb{T}', Y')_\delta$ with the following properties.*

(i) $(\tilde{\mathbb{T}}, \tilde{Y})_\rho$ is ε -extremal.

(ii) For each $\tilde{T} \in \tilde{\mathbb{T}}$, the unit rescaling of $(\mathbb{T}'[\tilde{T}], Y')_\delta$ relative to \tilde{T} is ε -extremal.

(iii) For each $p \in \mathbb{R}^3$, $\#\tilde{\mathbb{T}}(p) \leq \rho^{2-\sigma-\varepsilon}(\#\tilde{\mathbb{T}})$.

(iv) For each $p \in \mathbb{R}^3$ and each $\tilde{T} \in \tilde{\mathbb{T}}$, $\#\mathbb{T}'[\tilde{T}](p) \leq (\delta/\rho)^{2-\sigma-\varepsilon}(\#\mathbb{T}'[\tilde{T}])$.

Proof. The proof is nearly identical to the proof of the corresponding statement in [16, Proposition 3.2]. In [16], the first major step was to establish the existence of a cover \mathbb{T}_ρ of \mathbb{T} (or more accurately, a large subset $\mathbb{T}' \subset \mathbb{T}$), with the property that both \mathbb{T}_ρ and the re-scaled sets $\mathbb{T}'[T_\rho]$ are sticky. In our case, this step was handled by Lemma 2.13. The second major step was to establish a shading on \mathbb{T}_ρ and a refinement of the shadings on the sets $\mathbb{T}'[T_\rho]$ so that not too many tubes pass through each point (and as a consequence, the corresponding collections are ε -extremal).

Next we will give a slightly more detailed sketch, though we refer the reader to the proof of [16, Proposition 3.2] for full details. We begin by reducing to the case where each $T \in \mathbb{T}$ satisfies $|Y(T)| \geq \frac{1}{2}\delta^\eta|T|$. This can be done by discarding those tubes with $|Y(T)| < \frac{1}{2}\delta^\eta|T|$ and then repeatedly pigeonholing to re-establish a collection of tubes that satisfies the Convex Wolff Axioms at every scale (with a slightly worse constant, say $\delta^{-2\eta}$); after this process, the cardinality of our collection of tubes has been reduced by at most a factor of δ^η .

Next, we apply Lemma 2.13 with $\rho_0 = \delta^{2\eta}\rho$; we get a number $\rho' \in [\delta^{2\eta}\rho, \rho]$; and a collection $\mathbb{T}_{\rho'}$ of ρ' -tubes, as described by that Lemma. Of course, we may replace each ρ' tube by corresponding ρ -tube, and denote the resulting collection by $\tilde{\mathbb{T}}$. After pigeonholing to re-establish the property that the tubes are essentially distinct, it is still the case that $\tilde{\mathbb{T}}$ satisfies the Convex Wolff Axioms at every scale with error $\delta^{-O(\eta)}$, and each of the re-scaled collections $\mathbb{T}[\tilde{T}]$ satisfies the Convex Wolff Axioms at every scale with error $\delta^{-O(\eta)}$.

After refining the shadings $Y(T)$, we may suppose that for each collection $\mathbb{T}[\tilde{T}]$, the same number of tubes pass through each δ -cube in $E_{\mathbb{T}[\tilde{T}]}$ in the sense of (6.1); after further pigeonholing we may suppose that this number is the same (up to a multiplicative factor of 2) for each set $\mathbb{T}[\tilde{T}]$; call this number μ_{fine} . Since $\mathbb{T}[\tilde{T}]$ is a (re-scaled) collection of tubes satisfying the Convex Wolff Axioms at every scale, if η and δ_0 are chosen sufficiently small then we must have $\mu_{\text{fine}} \leq (\delta/\rho)^{2-\sigma-\varepsilon/10}(\#\mathbb{T}[\tilde{T}])$; this is Conclusion (iv) above.

Our next task is to define a shading \tilde{Y} on the tubes of $\tilde{\mathbb{T}}$. To begin, for each \tilde{T} we consider the union of all ρ -cubes $\tilde{Q} \subset \tilde{T}$ that intersect $E_{\mathbb{T}[\tilde{T}]}$. After pigeonholing, we can refine this shading so that each ρ -cube intersects $E_{\mathbb{T}[\tilde{T}]}$ in a set of roughly the same size, and the same number of tubes from $\tilde{\mathbb{T}}$ pass through each cube of $E_{\tilde{\mathbb{T}}}$; call this number μ_{coarse} . We refine the corresponding shadings $Y(T)$, $T \in \mathbb{T}$ so that $Y(T) \subset \tilde{Y}(\tilde{T})$ whenever $T \in \mathbb{T}[\tilde{T}]$; in particular $(\tilde{\mathbb{T}}, \tilde{Y})_\rho$ is now a balanced cover of $(\mathbb{T}, Y)_\delta$. Note that Conclusion (iv) remains true after this procedure. Since $\tilde{\mathbb{T}}$ satisfies the Convex Wolff Axioms with error $\delta^{-O(\eta)}$, if η and δ_0 are chosen sufficiently small then we must have $\mu_{\text{coarse}} \leq \rho^{2-\sigma-\varepsilon/10}(\#\tilde{\mathbb{T}})$; this is Conclusion (iii) above.

Finally, observe that $\mu_{\text{fine}}\mu_{\text{coarse}} \leq \delta^{2-\sigma-\varepsilon/5}(\#\mathbb{T})$, but on the other hand since \mathbb{T} was η -extremal,

we must have $\mu_{\text{fine}}\mu_{\text{coarse}} \gtrsim \delta^{2-\sigma+\eta}(\#\mathbb{T})$. We conclude that

$$\begin{aligned} (\delta/\rho)^{2-\sigma+\varepsilon/10}(\#\mathbb{T}'[\tilde{T}]) &\leq \mu_{\text{fine}} \leq (\delta/\rho)^{2-\sigma-\varepsilon/10}(\#\mathbb{T}'[\tilde{T}]), \quad \tilde{T} \in \tilde{\mathbb{T}}, \\ \rho^{2-\sigma+\varepsilon/10}(\#\tilde{\mathbb{T}}) &\leq \mu_{\text{coarse}} \leq \rho^{2-\sigma-\varepsilon/10}(\#\tilde{\mathbb{T}}), \end{aligned}$$

and in particular Conclusions (i) and (ii) are satisfied. We refer the reader to [16] for complete details. \square

6.3 Local and global grains

Using Proposition 6.2 in place of [16, Proposition 3.2], we can follow an identical argument as in [16] to obtain the analogue of [16, Proposition 4.1], which we recall here.

Proposition 6.3. *For all $\varepsilon, \delta_0 > 0$, there exists $\delta \in (0, \delta_0]$ and a ε -extremal set of tubes $(\mathbb{T}, Y)_\delta$ with the following two properties.*

1. $E_{\mathbb{T}}$ is a union of global grains with Lipschitz slope function.

There is a 1-Lipschitz function $f : [-1, 1] \rightarrow \mathbb{R}$ so that $(E_{\mathbb{T}} \cap \{z = z_0\}) \cdot (1, f(z_0), 0)$ is a $(\delta, 1 - \sigma, \delta^{-\varepsilon})_1$ -ADset for each $z_0 \in [-1, 1]$.

2. $E_{\mathbb{T}}$ is a union of local grains with Lipschitz plane map.

$(\mathbb{T}, Y)_\delta$ has a 1-Lipschitz plane map V . For all $\rho \in [\delta, 1]$ and all $p \in \mathbb{R}^3$, $V(p) \cdot (B(p, \rho^{1/2}) \cap E_{\mathbb{T}})$ is a $(\rho, 1 - \sigma, \delta^{-\varepsilon})_1$ -ADset.

Following the same proof as [16, Section 5] using radial projections, we can upgrade the global grains slope function f in Proposition 6.3 from Lipschitz to C^2 . This is the analogue of [16, Proposition 5.1]. The precise statement is as follows.

Proposition 6.4. *For all $\varepsilon, \delta_0 > 0$, there exists $\delta \in (0, \delta_0]$ and a ε -extremal set of tubes $(\mathbb{T}, Y)_\delta$ with the following properties.*

- 1' **Global grains with C^2 -slope function.**

There is a function $f : [-1, 1] \rightarrow \mathbb{R}$ with $\|f\|_{C^2} \leq 1$ so that $(E_{\mathbb{T}} \cap \{z = z_0\}) \cdot (1, f(z_0), 0)$ is a $(\delta, 1 - \sigma, \delta^{-\varepsilon})_1$ -ADset for each $z_0 \in [-1, 1]$.

- 2 **Local grains with Lipschitz plane map.**

$(\mathbb{T}, Y)_\delta$ has a 1-Lipschitz plane map V . For all $\rho \in [\delta, 1]$ and all $p \in \mathbb{R}^3$, $V(p) \cdot (B(p, \rho^{1/2}) \cap E_{\mathbb{T}})$ is a $(\rho, 1 - \sigma, \delta^{-\varepsilon})_1$ -ADset.

6.4 The slope function has large slope

Our next task is to show that the global grains slope function f in Proposition 6.3 has derivative f' with magnitude bounded away from 0. As discussed above, the proof of the analogous statement [16, Proposition 6.1] used the property that \mathbb{T} contains tubes pointing in many different directions. Here, we will use the property that \mathbb{T} satisfies the Convex Wolff Axioms at every scale. The precise statement we need is as follows.

Proposition 6.5. For all $\varepsilon, \delta_0 > 0$, there exists $\delta \in (0, \delta_0]$ and a ε -extremal set of tubes $(\mathbb{T}, Y)_\delta$ with the following properties:

1" **Global grains with nonsingular C^2 -slope function.** There is a function $f : [-1, 1] \rightarrow \mathbb{R}$ with $1 \leq |f'(z)| \leq 2$ and $|f''(z)| \leq 1/100$ for all $z \in [-1, 1]$, so that $(E_{\mathbb{T}} \cap \{z = z_0\}) \cdot (1, f(z_0), 0)$ is a $(\delta, 1 - \sigma, \delta^{-\varepsilon})_1$ -ADset for each $z_0 \in [-1, 1]$.

2 **Local grains with Lipschitz plane map.**

$(\mathbb{T}, Y)_\delta$ has a 1-Lipschitz plane map V . For all $\rho \in [\delta, 1]$ and all $p \in \mathbb{R}^3$, $V(p) \cdot (B(p, \rho^{1/2}) \cap E_{\mathbb{T}})$ is a $(\rho, 1 - \sigma, \delta^{-\varepsilon})_1$ -ADset.

Before proving 6.5, we will establish the following technical result, which is the analogue of [16, Lemma 6.3].

Lemma 6.6. For all $\varepsilon > 0$, there exists $\eta > 0, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $(\mathbb{T}, Y)_\delta$ be a set of tubes that satisfies the conclusions of Proposition 6.4, with η in place of ε . Let $J \subset [-1, 1]$ be an interval, with $\delta^{1/2} \leq |J| \leq \delta^\varepsilon$. Suppose that

$$\sum_{T \in \mathbb{T}} |Y(T) \cap (\mathbb{R}^2 \times J)| \geq \delta^\eta |J|, \quad (6.3)$$

and that $(\mathbb{R}^2 \times J) \cap \bigcup_{T \in \mathbb{T}} Y(T)$ can be covered by a union of $\leq \delta^{-\eta} |J|^{-2+\sigma}$ cubes of side-length J .

Then the slope function $f : [-1, 1] \rightarrow \mathbb{R}$ from Proposition 6.4 satisfies

$$|f'(z)| \geq |J| \text{ for all } z \in J.$$

Proof. The proof is the same as [16, Lemma 6.3] until **Step 4: \mathbb{T} contains parallel tubes.**

Here is a summary of Steps 1-3. After a refinement, we may suppose that for each point $p \in F := (\mathbb{R}^2 \times J) \cap \bigcup_{T \in \mathbb{T}} Y(T)$,

$$\delta^{2-\sigma+2\eta} (\#\mathbb{T}) \leq \#\mathbb{T}(p) \leq \delta^{2-\sigma-2\eta} (\#\mathbb{T}). \quad (6.4)$$

After a translation and a rotation, we may suppose that $J = [0, \rho^{1/2}]$ and $f(0) = 0$. Since $|f''(z)| \lesssim 1$, if $|f'(z)| \leq |J|$ for some $z \in J$, then $|f'(z)| \lesssim |J|$ for all $z \in J$, and so $|f(z)| \lesssim |J|^2$. We identify a subset $F_2 \subset F$ with $|F_2| \geq \delta^{4\eta} |F|$ so that

1. for each $z_1 \in J$ and each $r \in (F \cap \{z = z_1\}) \cdot (1, f(z_1), 0)$, the $1 \times \delta \times \delta$ -rectangular prism

$$G_{r, z_1} = \{|z - z_1| \leq \delta\} \cap \{|(x, y) \cdot (1, f(z_1)) - r| \leq \delta\} \quad (6.5)$$

has a large intersection with F_2 :

$$|G_{r, z_1} \cap F_2| \geq \delta^{4\eta} |G_{r, z_1}| \geq \delta^{2+4\eta}. \quad (6.6)$$

2. Let $\Pi_{x, z} : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be the projection to the first and third coordinates, then

$$|N_{\rho^{1/2}} \Pi_{x, z}(F_2)| \leq \delta^{-7\eta} \rho^{1/2+\sigma/2}.$$

3. There is a $y_0 \in [-1, 1]$ such that

$$|\{y = y_0\} \cap F_2| \geq |F_2|/2$$

where the $|\cdot|$ on the LHS denote 2-dimensional measure and $|\cdot|$ on the RHS denote 3-dimensional measure, and each G_{r,z_1} defined in (6.5) satisfies

$$G_{r,z_1} \cap \{y = y_0\} \cap F_2 \neq \emptyset.$$

Cover $[-1, 1] \times \{y = y_0\} \times J$ by a set \mathcal{S} of interior-disjoint squares of side-length $\rho^{1/2} = |J|$. Let $S \in \mathcal{S}$ be a square with center $p_S = (x_S, y_0, \rho^{1/2}/2)$. By Item 2 from Proposition 6.4, $S \cap F_2 \cdot V(p_S)$ is a $(\rho, 1 - \sigma, \delta^{-\eta})_1$ -ADset. Write $V(p_S) = (V_x(p_S), V_y(p_S), V_z(p_S))$ and define $\tilde{V}(p_S) = (V_x(p_S), 0, V_z(p_S))$. Since $S \cap F_2 \subset \{y = y_0\}$,

$$(S \cap F_2) \cdot \tilde{V}(p_S) \text{ is a } (\rho, 1 - \sigma, \delta^{-\eta})_1\text{-ADset.} \quad (6.7)$$

Define $P_S = \{(x, y, z) : (x, z) \in S\}$, then

$$(F_2 \cap P_S) \cdot \tilde{V}(p_S) \text{ is a } (\rho, 1 - \sigma, 2\delta^{-\eta})_1\text{-ADset.}$$

This finishes the summary of Steps 1-3 and we have introduced the necessary geometric objects and notations.

In Step 4, we will show that the constraints from the above steps will force many tubes from \mathbb{T} to be contained in a $\delta^{-2C_0\eta}\rho^{1/2} \times 1 \times 1$ rectangular prism (henceforth called a *slab*, to match the terminology in [16]) for some constant $C_0 = C_0(\sigma)$, which contradicts the assumption that \mathbb{T} satisfies the Convex Wolff Axioms (see Definition 6.1 and Definition 2.5).

For each $T \in \mathbb{T}$, define $F_2(T) = Y(T) \cap F_2$. Recalling (6.3) and (6.4), we have

$$\sum_{T \in \mathbb{T}} |F_2(T)| \geq \delta^{10\eta} |J| = \delta^{10\eta} \rho^{1/2}. \quad (6.8)$$

Recall that each $T \in \mathbb{T}$ intersects at most 3 prisms P_S , $S \in \mathcal{S}$ and for each such prism we have $|F_2(T) \cap P_S| \leq \rho^{1/2}\delta^2$. By pigeonholing, there exists a square $S \in \mathcal{S}$ and a set $\mathbb{T}_S = \{T \in \mathbb{T} : T \cap P_S \neq \emptyset\}$ with

$$\#\mathbb{T}_S \gtrsim (\#\mathbb{T})(\#\mathcal{S})^{-1} \gtrsim \delta^\eta \rho^{-\sigma/2+1/2} (\#\mathbb{T}). \quad (6.9)$$

But for each $T \in \mathbb{T}_S$, we have

$$F_2(T) \cdot \tilde{V}(p_S) \subset (F_2 \cap P_S) \cdot \tilde{V}(p_S) \text{ is a } (\rho, 1 - \sigma, 2\delta^{-\eta})_1\text{-ADset.}$$

By [16, Lemma 6.2], we have

$$|\text{dir}(T) \cdot \tilde{V}(p_S)| \leq \delta^{-C_0\eta} \rho^{1/2}. \quad (6.10)$$

It follows that \mathbb{T}_S is contained in the slab

$$\Sigma_S = \{(x, y, z) \in [0, 1]^3 : |(x - x_S, 0, z) \cdot \tilde{V}(p_S)| \leq \delta^{-2C_0\eta} \rho^{1/2}\}.$$

By the Convex Wolff Axioms,

$$\#\mathbb{T}_S \leq |\Sigma_S| \delta^{-\eta} (\#\mathbb{T}) \leq \delta^{-(2C+1)\eta} \rho^{1/2} (\#\mathbb{T}),$$

which yields a contradiction to (6.9). \square

The proof of Proposition 6.5 is the same as [16, Proposition 6.1] using Lemma 6.6 and [16, Lemma 6.4].

6.5 Twisted projections, and the completion of the proof

Recalling Definition 6.1 from [16], for $f: [-1, 1] \rightarrow \mathbb{R}$ we define the twisted projection $\pi_f(x, y, z) = (x + f(z)y, z)$. The consequence of Proposition 6.5 is the following analogue of [16, Corollary 6.5]

Lemma 6.7. *For all $\varepsilon, \delta_0 > 0$, there exists $\delta \in (0, \delta_0]$; a set pair (\mathbb{T}, Y) of δ -tubes satisfying the Convex Wolff Axioms with error $\delta^{-\varepsilon}$ and an δ^ε -dense shading; and a function $f: [-1, 1] \rightarrow \mathbb{R}$ with $1 \leq |f'(z)| \leq 2$ and $|f''(z)| \leq 1/100$ for all $z \in [-1, 1]$, so that*

$$\left| \bigcup_{T \in \mathbb{T}} \pi_f(Y(T)) \right| \leq \delta^{\sigma + \varepsilon}. \quad (6.11)$$

To complete the proof of Theorem 5.2, it suffices to establish the following estimate, which is the analogue of [16, Proposition 7.1]

Proposition 6.8. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $(\mathbb{T}, Y)_\delta$ be a set of δ -tubes satisfying the Convex Wolff Axioms with error $\delta^{-\eta}$, and let Y be a δ^η -dense shading. Let $f: [-1, 1] \rightarrow \mathbb{R}$ with $1 \leq |f'(z)| \leq 2$ and $|f''(z)| \leq 1/100$ for all $z \in [-1, 1]$. Then*

$$\left| \bigcup_{T \in \mathbb{T}} \pi_f(Y(T)) \right| \geq \delta^\varepsilon.$$

Indeed, comparing Lemma 6.7 and Proposition 6.8, we conclude that $\sigma = 0$, which completes the proof of Theorem 5.2. It remains to prove Proposition 6.8; we defer this to the next section.

7 Twisted projections of tubes satisfying the Convex Wolff Axioms

Our goal in this section is to prove Proposition 6.8. In fact, we will prove a slightly stronger statement. We say a set of tubes \mathbb{T} satisfies the *Tube Wolff Axioms* with error C , if for every ρ -tube T_ρ , at most $C|T_\rho|(\#\mathbb{T})$ tubes from \mathbb{T} are contained in T_ρ ; since ρ -tubes are convex, this is a special case of the Convex Wolff Axioms. We will prove the following mild generalization of Proposition 6.8

Proposition 7.1. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $(\mathbb{T}, Y)_\delta$ be a set of δ -tubes satisfying the Tube Wolff Axioms with error $\delta^{-\eta}$, and let Y be a δ^η -dense shading. Let $f: [-1, 1] \rightarrow \mathbb{R}$ with $1 \leq |f'(z)| \leq 2$ and $|f''(z)| \leq 1/100$ for all $z \in [-1, 1]$. Then*

$$\left| \bigcup_{T \in \mathbb{T}} \pi_f(Y(T)) \right| \geq \delta^\varepsilon.$$

Before proving Proposition 7.1, we will need to introduce two closely related non-concentration conditions for subsets of \mathbb{R}^n , which arise from discretizing fractal sets. The first was proposed by Katz and Tao [8], while the second was explored by Orponen and Shmerkin [12].

Definition 7.2. *Let $\delta > 0$, $s \in (0, n]$, and $C > 0$. A non-empty bounded set $A \subset \mathbb{R}^n$ is called a (δ, s, C) -Katz-Tao set if*

$$\mathcal{E}_\delta(A \cap B(x, r)) \leq C(\delta/r)^s, \quad \forall x \in \mathbb{R}^n, r \in [\delta, 1].$$

Definition 7.3. Let $\delta > 0$, $s \in (0, n]$, and $C > 0$. A non-empty bounded set $A \subset \mathbb{R}^n$ is called a (δ, s, C) -Frostman set if

$$\mathcal{E}_\delta(A \cap B(x, r)) \leq Cr^s \mathcal{E}_\delta(A), \quad \forall x \in \mathbb{R}^n, r \in [\delta, 1].$$

Remark 7.4. If \mathbb{T} is a set of δ -tubes contained in a bounded set such as $B(0, 1)$, then we can identify the coaxial line of each tube with a point in \mathbb{R}^4 . If \mathbb{T} satisfies the Tube Wolff Axioms, then the corresponding set of points in \mathbb{R}^4 forms a $(\delta, 2, O(1))$ -Frostman set (technically the corresponding set of points might be a multi-set, but this is harmless—we can either extend the definition of a Orponen-Shmerkin-set to include this possibility, or infinitesimally perturb the tubes to avoid this situation).

Remark 7.5. If $A \subset \mathbb{R}^n$ is a (δ, s, C) -Katz-Tao set (resp. (δ, s, C) -Frostman set), then $N_\delta(A)$ is a (δ, s, C') -Katz-Tao set (resp. (δ, s, C') -Frostman set), where $C' = O_n(C)$, and conversely.

Lemma 7.6. Let $A \subset \mathbb{R}^n$ be a (δ, s, C) -Frostman set. Then there exists a $(\delta, s, 100)$ -Katz-Tao set $A' \subset A$, with $\mathcal{E}_\delta(A') \gtrsim_{n,s} \delta^{-s} |\log \delta|^{-1} C^{-1}$.

Proof. Recall that the s -dimensional Hausdorff content is defined as

$$\mathcal{H}_\infty^s(A) := \inf \left\{ \sum_i r(B_i)^s : A \subset \cup_i B_i \right\},$$

where each B_i in the covering is a cube and $r(B_i)$ is the side-length of B_i . By [3, Lemma 3.13], if $\mathcal{H}_\infty^s(A) = \kappa > 0$, then there exists a $(\delta, s, 100)$ -Katz-Tao set $A' \subset A$ with $\mathcal{E}_\delta(A') \gtrsim \kappa \delta^{-s}$. Thus in order to prove the lemma, it suffices to obtain the bound

$$\mathcal{H}_\infty^s(A) \gtrsim_{n,s} |\log \delta|^{-1} C^{-1}. \quad (7.1)$$

The remainder of the proof is devoted to establishing (7.1). By Remark 7.5, we can assume A is a union of δ -balls (this introduces a harmless $O_n(1)$ loss). Let $\{B_i\}$ be a cover of A by balls. Sort the balls according to their side-length; it suffices to consider balls of side-length $\geq \delta$ since A is a union of δ -balls. By dyadic pigeonholing, there exists a dyadic number $\lambda \in [\delta, 1]$ such that

$$\mathcal{E}_\delta \left(\bigcup_{r(B_i) \sim \lambda} B_i \cap A \right) \gtrsim |\log \delta|^{-1} \mathcal{E}_\delta(A).$$

Since A is a (δ, s, C) -Frostman set, $\mathcal{E}_\delta(A \cap B_i) \leq C \lambda^s \mathcal{E}_\delta(A)$. The number of B_i in the cover with $r(B_i) \sim \lambda$ is $\gtrsim_{n,s} \lambda^{-s} |\log \delta|^{-1} C^{-1}$. This establishes (7.1) and concludes the proof. \square

The following definition describes when a discretized set is homogeneous at many different scales.

Definition 7.7 (η -uniform set up to scale δ). Let $\eta > 0$ be a small parameter. A set $A \subset [0, 1]^n$ is called an η -uniform set up to scale δ if for any $\rho \in 2^{-\eta^{-1}\mathbb{N}} \cap [\delta, 1]$, if two ρ -cubes $Q_1, Q_2 \subset N_\rho(A)$, then the δ -covering numbers are comparable, in the sense that

$$\frac{1}{100} \mathcal{E}_\delta(Q_1 \cap A) \leq \mathcal{E}_\delta(Q_2 \cap A) \leq 100 \mathcal{E}_\delta(Q_1 \cap A).$$

Let $\eta > 0$ be a small parameter and $\delta \in 2^{-\eta^{-1}\mathbb{N}}$. Given an arbitrary set $A \subset [0, 1]^n$, [13, Lemma 2.15] says there exists an η uniform set $A' \subset A$ up to scale δ and $\mathcal{E}_\delta(A') \gtrsim |\log \delta|^{-\eta^{-1}} \mathcal{E}_\delta(A)$. We call A' an η -uniform refinement (usually δ is clear from context and we omit it from notation).

Even though [13, Lemma 2.15] is for dyadic cubes, the same argument works for any set A and the relevant ρ -balls $Q \subset N_\rho(A)$, which we restate in our language below.

Lemma 7.8. *Let $A \subset [0, 1]^n$ and let $\eta > 0, T \in \mathbb{N}$, $\delta = 2^{-\eta^{-1}T}$. Then there exists an η -uniform refinement $A' \subset A$ such that*

$$\mathcal{E}_\delta(A') \geq (2T)^{-\eta^{-1}} \mathcal{E}_\delta(A).$$

In particular, if $\varepsilon > 0$ and $T^{-1} \log(2T) \leq \varepsilon$, then $\mathcal{E}_\delta(A') \geq \delta^\varepsilon \mathcal{E}_\delta(A)$.

Given an η -uniform set A with δ -covering number δ^{-s} , the next lemma finds an Orponen-Shmerkin-set if one zooms in at sufficiently small scale.

Lemma 7.9. *For any $\varepsilon > 0$, there exists $\eta > 0$ such that for any $\delta \in 2^{-\eta^{-1}\mathbb{N}}$, the following holds. Let $A \subset [0, 1]^n$ be an η -uniform set with $\mathcal{E}_\delta(A) \geq \delta^{-s}$. Then there exists $\rho \in (\delta^{1-\varepsilon}, 1]$ such for any ρ -cube $Q \in N_\rho(A)$,*

$$A_Q := S_Q(A \cap Q) \text{ is a } (\delta/\rho, s, (\delta/\rho)^{-4d\varepsilon})\text{-Frostman set,}$$

where $S_Q : Q \rightarrow [0, 1]^n$ is an affine transformation that maps Q to the unit cube.

Proof. Let $\eta \ll \varepsilon$ to be determined later. We shall apply the multi-scale decomposition techniques developed by Keleti and Shmerkin [11]. By [15, Lemma 5.21], there exists a decomposition of scales such that

- $a_1 = \eta^{1/2}, a_{J+1} = 1, a_{j+1} - a_j \geq \tau$ where $\tau = \tau(\varepsilon) > 0$ depends only on ε .
- For each δ^{a_j} -cube $Q \subset N_{\delta^{a_j}}(A)$, $A_Q := S_Q(A \cap Q)$ is a $(\delta^{a_{j+1}-a_j}, t_j, \delta^{-\eta})$ -set, where $S_Q : Q \rightarrow [0, 1]^3$ is an affine transformation that maps Q to the unit cube.
- $\sum t_j(a_{j+1} - a_j) \geq s - \varepsilon^2$.

In principle there is no control on the values of t_j . However, we can fix this as follows. The ‘‘Merging Lemma,’’ [15, Lemma 5.20] says that if $t_j \geq t_{j+1}$, then for each δ^{a_j} -cube $Q \subset N_{\delta^{a_j}}(A)$, $A_Q := S_Q(A \cap Q)$ is a $(\delta^{a_{j+2}-a_j}, t'_j, \delta^{-\eta})$ -set, where $t'_j = \frac{(a_{j+1}-a_j)t_j + (a_{j+2}-a_{j+1})t_{j+1}}{a_{j+2}-a_j}$.

Start with the output of [15, Lemma 5.21] applying to A , if there exists $t_j \geq t_{j+1}$, apply [15, Lemma 5.20] to merge the two intervals $[a_j, a_{j+1}]$ and $[a_{j+1}, a_{j+2}]$ to $[a_j, a_{j+2}]$ and the two numbers t_j, t_{j+1} becomes t'_j . Rename the set of intervals and t_j s and replace J by $J - 1$. Iterate the process until $t_j < t_{j+1}$ for all j . Now we have obtained a decomposition of scales $\{[a_j, a_{j+1}]\}_{j=1}^J$ such that

- $a_1 = \eta^{1/2}, a_{J+1} = 1, a_{j+1} - a_j \geq \tau$ where $\tau = \tau(\varepsilon) > 0$ depends only on ε .
- For each δ^{a_j} -cube $Q \subset N_{\delta^{a_j}}(A)$, $A_Q := S_Q(A \cap Q)$ is a $(\delta^{a_{j+1}-a_j}, t_j, \delta^{-\eta})$ -set, where $S_Q : Q \rightarrow [0, 1]^3$ is an affine transformation that maps Q to the unit cube.
- $t_j < t_{j+1}$ and $\sum t_j(a_{j+1} - a_j) \geq s - \varepsilon^2$.

Let J_0 be the smallest j such that $t_j \geq s - \varepsilon$. It follows from

$$s - \varepsilon^2 \leq \sum_j t_j (a_{j+1} - a_j) \leq (s - \varepsilon) a_{J_0} + d(1 - a_{J_0})$$

that $a_{J_0} \leq 1 - \frac{\varepsilon}{2d}$.

Define $\rho = \delta^{a_{J_0}}$. For any ρ -cube $Q \subset N_\rho(A)$ and $A_Q = S_Q(A \cap Q)$. Since $s - \varepsilon \leq t_{J_0} < t_{J_0+1} < \dots < t_J$, A_Q is a union of δ/ρ -balls that is a $(\delta/\rho, s, (\delta/\rho)^{-2\varepsilon})$ -set if $\eta \ll \varepsilon^2$. The conclusion holds with ε replaced by $2d\varepsilon$. \square

Lemma 7.10. *Let $s \in (0, n]$. Suppose $Q_0 \subset [0, 1]^n$ is a cube of side-length ρ and $A_0 \subset Q_0$ is a (δ, s, C) -Katz-Tao set with $\mathcal{E}_\delta(A_0) \gtrsim (\rho/\delta)^s$. Then there exists a $(\delta, s, O(C))$ -Katz-Tao set A with $\mathcal{E}_\delta(A) \gtrsim \delta^{-s}$ so that for each axis-aligned ρ -cube Q that intersects A , we have that $Q \cap A$ is a translation of A_0 .*

Proof. Divide the unit cube $[0, 1]^n$ into a union of cubes \tilde{Q} of side length $\rho^{s/d} \leq \rho$. For each \tilde{Q} , let $A_{\tilde{Q}}$ denote the translation that maps the center of Q_0 to the center of \tilde{Q} . Define $A = \sqcup A_{\tilde{Q}}(A_0)$. Since the side-length of \tilde{Q} is greater than the side-length of Q_0 , $A_{\tilde{Q}}(A_0)$ are disjoint, and so $\mathcal{E}_\delta(A) \gtrsim \delta^{-s}$.

It remains to verify that A is a (δ, s, C) -Katz-Tao set. By the construction of A , it suffices to check that for each $r \geq \rho$,

$$\mathcal{E}_\delta(A \cap B_r) \lesssim C(r/\delta)^s.$$

To see this, when $r \in (\rho, \rho^{s/n})$, we have $\mathcal{E}_\delta(A \cap B_r) \sim \mathcal{E}_\delta(A_0) \lesssim C(\rho/\delta)^s$. On the other hand, when $r \geq \rho^{s/n}$, we have

$$\mathcal{E}_\delta(A \cap B_r) \lesssim C\left(\frac{r}{\rho^{s/n}}\right)^n \cdot \left(\frac{\rho}{\delta}\right)^s \lesssim C\left(\frac{r}{\delta}\right)^s. \quad \square$$

We prove a single scale version of Proposition 7.1.

Lemma 7.11. *For all $\varepsilon > 0$, there exists $\eta, \delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $(\mathbb{T}, Y)_\delta$ be a set of δ -tubes satisfying the Tube Wolff Axioms with error $\delta^{-\eta}$, and let Y be a δ^η -dense shading. Let $f: [-1, 1] \rightarrow \mathbb{R}$ with $1 \leq |f'(z)| \leq 2$ and $|f''(z)| \leq 1/100$ for all $z \in [-1, 1]$. Then there exists $\rho \in (\delta^{1-\varepsilon^2}, 1)$ and $\delta^{4\eta}$ -shading $Y' \subset Y$ such that*

$$\left| \bigcup_{T \in \mathbb{T}} \pi_f(Y'(T)) \right| \geq (\delta/\rho)^\varepsilon \left| N_\rho\left(\bigcup_{T \in \mathbb{T}} \pi_f(Y'(T)) \right) \right|. \quad (7.2)$$

The quantity on the RHS of (7.2) will be closely related to the graph of cinematic functions (see [16, Definition 7.1]). To estimate it, we recall [16, Theorem 7.2].

Theorem 7.12. *Let $I \subset \mathbb{R}$ be a compact interval and let $M \subset C^2(I)$ be a family of cinematic functions (see [16, Definition 1.6]). Then for all $\varepsilon > 0$, there exists $\delta_0 > 0$ so that the following holds for all $\delta \in (0, \delta_0]$. Let $G \subset M$ be a δ -separated set that satisfies the Frostman non-concentration condition*

$$\#(G \cap B) \leq \delta^{-\varepsilon} (r/\delta) \text{ for all balls } B \subset C^2(I) \text{ of diameter } r. \quad (7.3)$$

Then

$$\left\| \sum_{g \in G} \chi_{g^\delta} \right\|_{3/2} \leq \delta^{-\varepsilon},$$

where g^δ is the δ -neighborhood of $\text{graph}(g)$.

Proof of Lemma 7.11. Let $X_0 = \bigcup_{T \in \mathbb{T}} \pi_f(Y(T))$ and $0 < \eta \ll \varepsilon$ be a small parameter to be chosen later. For each dyadic number $1 < \lambda < \delta^{-4}$, let X_λ be the set of $x \in X_0$ such that $\sum_{p \in \pi_f^{-1}(x)} \#\mathbb{T}(p) \sim \lambda$, where the sum is over a δ -separated set of $p \in \pi_f^{-1}(x)$ and $\#\mathbb{T}(p)$ was defined in (6.1). By pigeonholing, there exists λ such that $Y_\lambda(T) := Y(T) \cap \pi_f^{-1}(X_\lambda)$ is a $|\log \delta|^{-2} \delta^\eta$ -dense shading.

Apply Lemma 7.8 to find an η -uniform subset $X \subset X_\lambda$ such that $|X| \geq \delta^{\eta/2} |X_\lambda|$. By double counting, this implies that for a $\geq \delta^{2\eta}$ -fraction of \mathbb{T} , $|Y(T) \cap \pi_f^{-1}(X)| \geq \delta^{2\eta} |T|$. Define $Y'(T) = Y_\lambda(T) \cap \pi_f^{-1}(X)$.

Observe that the image of the line $l_{a,b,c,d} := \{(a + ct, b + dt, t) : t \in \mathbb{R}\}$ under the twisted projection π_f is the curve

$$\pi_f(l_{a,b,c,d}) = \{(a + ct + f(t)(b + dt), t)\}.$$

After possibly refinement and rotation, each tube $T \in \mathbb{T}$ is δ -neighborhood of a line of the form $l_{a,b,c,d}$ intersecting the unit cube. Let $A(\mathbb{T})$ denote set of parameters of these lines, i.e.

$$A(\mathbb{T}) := \{(a, b, c, d) : T_{a,b,c,d} := N_\delta(l_{a,b,c,d}) \cap [0, 1]^3 \in \mathbb{T}\}.$$

Since \mathbb{T} satisfies the Tube Wolff axioms, if $\rho \in [\delta, 1]$ then every ρ -tube T_ρ contains at most $\delta^{-\eta} |T_\rho| (\#\mathbb{T})$ tubes from \mathbb{T} , and hence the point set $A(\mathbb{T}) \subset \mathbb{R}^4$ of parameters is a $(\delta, 2, \delta^{-\eta})$ -Frostman set.

Let $A_c = \{(a, b, d) : (a, b, c, d) \in A(\mathbb{T})\}$. By Fubini, there exists a $c \in [0, 1]$ such that $\mathcal{E}_\delta(A_c) := \delta^{-s} \geq \delta^{-1+\eta} \geq \delta^{-1+\varepsilon^2}$. Apply Lemma 7.8 to refine A_c to be an η -uniform subset.

So far we only know the δ -covering number of A_c but not the spacing condition. To apply Theorem 7.12, we need to know the spacing of A_c . By Lemma 7.9, there exists $\rho' \in (\delta^{1-\varepsilon^2}, 1]$ such that for any ρ -cube $Q \subset N_{\rho'}(A_c)$, $A_Q := S_Q(A_c \cap Q)$ is a $(\rho, 1 - \varepsilon^2, \rho^{-12\varepsilon^2})$ -set with $\rho = \delta/\rho'$.

By Lemma 7.6, since A_Q is a $(\rho, 1 - \varepsilon^2, \rho^{-12\varepsilon^2})$ -set, there exists a $(\rho, 1 - \varepsilon^2, 100)$ -Katz-Tao set $A'_Q \subset A_Q$ with $\mathcal{E}_\rho(A'_Q) \gtrsim \rho^{14\varepsilon^2-1}$. Then $S_Q^{-1}(A'_Q)$ is a $(\delta, 1 - \varepsilon^2, 100)$ -Katz-Tao set contained in Q , a cube of side-length $\rho' = \delta/\rho$. By Lemma 7.10, there exist $\sim \rho'^{-1}$ translations A_i such that the set $A'_c := \sqcup_i A_i(S_Q^{-1}(A'_Q))$ is a $(\rho, 1 - \varepsilon^2, 100)$ -Katz-Tao set with $\mathcal{E}_\delta(A'_c) \gtrsim \rho^{14\varepsilon^2} \delta^{-1}$. For each A_i and $(a, b, d) \in S_Q^{-1}(A'_Q)$, define the shading on the tube $T' = T_{a',b',c,d'}$, where $(a', b', d') = A_i(a, b, d)$, as

$$(a' + ct, b' + d't, t) \in Y'(T') \text{ if and only if } (a + ct, b + dt, t) \in Y'(T).$$

For each t , define $\pi_{f,t} = (x + f(t)y)$. Then for each fixed pair (A_i, A_j) ,

$$\left| \bigcup_{A_i} \pi_{f,t}(Y'(T)) \right| \sim \left| \bigcup_{A_j} \pi_{f,t}(Y'(T)) \right|, \quad (7.4)$$

where the union \bigcup_{A_i} is taken over all $(a, b, d) \in A_i(S_Q^{-1}(A'_Q))$ and the corresponding $T = T_{a,b,c,d}$. To see that (7.4) holds, let $A_j \circ A_i^{-1}(x) = x + (a_0, b_0, d_0)$. Then $a + ct + f(t)(b + dt) \in \Omega_{i,t} := \bigcup_{A_i} \pi_{f,t}(Y'(T))$ if and only if

$$a + a_0 + ct + f(t)(b + b_0 + (d + d_0)t) \in \Omega_{j,t} := \bigcup_{A_j} \pi_{f,t}(Y'(T)).$$

Therefore,

$$\Omega_{i,t} = \Omega_{j,t} + a_0 + b_0 f(t) + d_0 f(t)t,$$

which establishes (7.4). Finally, integrating (7.4) in t yields

$$\left| \bigcup_{A_i} \pi_f(Y'(T)) \right| \sim \left| \bigcup_{A_j} \pi_f(Y'(T)) \right|. \quad (7.5)$$

By [16, Lemma 7.3], the set of functions

$$M_f := \{g_{a,b,d}(t) = a + bf(t) + dtf(t) : a, b, d \in [-1, 1]\}$$

is a family of cinematic functions (see [16, Definition 7.1]). Apply Theorem 7.12 with $I = [-1, 1]$, $M = M_f$ and $G = \{g_{a,b,d} : (a, b, d) \in A'_c\}$. We conclude that

$$\left\| \sum_{g \in G} \chi_{g^{2\delta}} \right\|_{3/2} \lesssim \delta^{-\varepsilon^2}.$$

If $g = g_{a,b,d}$ then $\pi_f(T) \subset g^{2\delta}$ with $T = T_{a,b,c,d}$. Let $\mathbb{T}_0 = \{T_{a,b,c,d} : (a, b, d) \in A'_c\}$. Therefore,

$$\begin{aligned} \rho^{15\varepsilon^2} &\leq \sum_{T \in \mathbb{T}_0} |Y'(T)| \leq \left| \bigcup_{T \in \mathbb{T}_0} \pi_f(Y'(T)) \right|^{1/3} \left\| \sum_{T \in \mathbb{T}_0} \chi_{\pi_f(Y'(T))} \right\|_{3/2} \\ &\leq \left| \bigcup_{T \in \mathbb{T}_0} \pi_f(Y'(T)) \right|^{1/3} \left\| \sum_{T \in \mathbb{T}_0} \chi_{\pi_f(T)} \right\|_{3/2} \\ &\leq \left| \bigcup_{T \in \mathbb{T}_0} \pi_f(Y'(T)) \right|^{1/3} \left\| \sum_{g \in G} \chi_{g^{2\delta}} \right\|_{3/2}. \end{aligned}$$

Therefore,

$$\left| \bigcup_{T \in \mathbb{T}_0} \pi_f(Y'(T)) \right| \gtrsim \rho^{45\varepsilon^2}.$$

Since A'_c consists of $\lesssim \rho/\delta$ translated copies of $S_Q^{-1}(A'_Q)$, and the latter is a subset of $A_c \cap Q$, (7.5) implies

$$\left| \bigcup_{(a,b,c,d) \in A_c \cap Q} \pi_f(Y'(T_{a,b,c,d})) \right| \gtrsim \rho^{45\varepsilon^2} \delta / \rho. \quad (7.6)$$

Note that the LHS of (7.6) is contained in the $10\delta/\rho$ -neighborhood of a curve $\pi_f(l_{a,b,c,d})$, $(a, b, d) \in Q$, whose volume is comparable to the RHS of (7.6), up to a factor of $\rho^{-45\varepsilon^2}$. We conclude that the η -uniform set $X = \bigcup_{T \in \mathbb{T}} \pi_f(Y'(T))$ satisfies

$$|N_{\delta/\rho} X| \leq \rho^{-50\varepsilon^2} |X|. \quad \square$$

We prove Proposition 7.1 by iterating Lemma 7.11. The details are as follows.

Proof of Proposition 7.1. Let $0 < \eta \ll \varepsilon$ to be determined. Let $\mathbb{T}' \subset \mathbb{T}$ be the set of tubes with $|Y(T)| \geq \delta^{2\eta}|T|$, then $\#\mathbb{T}' \geq \delta^\eta \#\mathbb{T}$. Apply Lemma 7.8 to find an η -uniform subset $\mathbb{T}'' \subset \mathbb{T}'$ with $\#\mathbb{T}'' \geq \delta^\eta \#\mathbb{T}'$. Then $(\mathbb{T}'', Y)_\delta$ becomes a set of δ -tubes satisfying the Tube Wolff Axioms with error $\delta^{-3\eta}$ and Y is a $\delta^{2\eta}$ -dense shading. To ease the notation, still let \mathbb{T} denote \mathbb{T}'' .

By Lemma 7.11, we can find a scale $\rho_1 \in (\delta^{1-\varepsilon^2}, 1)$ and a $\delta^{4\eta}$ -shading $Y_1 \subset Y$ such that

$$\left| \bigcup_{T \in \mathbb{T}} \pi_f(Y(T)) \right| \geq \left| \bigcup_{T \in \mathbb{T}} \pi_f(Y_1(T)) \right| \geq (\delta/\rho_1)^\varepsilon \left| N_{\rho_1} \left(\bigcup_{T \in \mathbb{T}} \pi_f(Y_1(T)) \right) \right|.$$

Consider \mathbb{T}_{ρ_1} as the set of distinct ρ_1 -tubes containing at least one tube in \mathbb{T} . Since \mathbb{T} is η -uniform, \mathbb{T}_{ρ_1} satisfies the Tube Wolff Axioms with error $\delta^{-3\eta}$. Define Y_{ρ_1} be a shading on $\tilde{T} \in \mathbb{T}_{\rho_1}$ as $Y_{\rho_1} = N_{\rho_1}(\cup_{T \in \mathbb{T}[\tilde{T}]} Y_1(T))$. Therefore, Y_{ρ_1} is a $\delta^{2\eta}$ -dense shading. Apply Lemma 7.11 on $(\mathbb{T}_{\rho_1}, Y_{\rho_1})_{\rho_1}$ and find $\rho_2 \in (\rho_1^{1-\varepsilon^2}, 1)$ and corresponding $Y_2 \subset Y_{\rho_1}$ such that

$$\left| \bigcup_{\tilde{T} \in \mathbb{T}_{\rho_1}} \pi_f(Y_2(\tilde{T})) \right| \geq (\rho_1/\rho_2)^\varepsilon \left| N_{\rho_2} \left(\bigcup_{\tilde{T} \in \mathbb{T}_{\rho_1}} \pi_f(Y_2(\tilde{T})) \right) \right|.$$

Since for any $\tilde{T} \in \mathbb{T}_{\rho_1}$, $Y_2 \subset Y_{\rho_1} \subset N_{\rho_1}(\cup_{T \in \mathbb{T}[\tilde{T}]} \pi_f(Y_1(T)))$, we have

$$\left| \bigcup_{T \in \mathbb{T}} \pi_f(Y(T)) \right| \geq (\delta/\rho_2)^\varepsilon \left| N_{\rho_2} \left(\bigcup_{\tilde{T} \in \mathbb{T}_{\rho_1}} \pi_f(Y_2(\tilde{T})) \right) \right|.$$

Iterate the above process and obtain correspondingly scales ρ_3, \dots, ρ_N until $\rho_N \geq \delta^{\varepsilon^2}$. Since $\rho_{k+1} \geq \rho_k^{1-\varepsilon^2}$, $N \leq \varepsilon^{-2}$. When $\eta > 0$ is sufficiently small depending on ε ,

$$\left| \bigcup_{T \in \mathbb{T}} \pi_f(Y(T)) \right| \geq (\delta/\rho_2)^\varepsilon \left| N_{\rho_N} \left(\bigcup_{\tilde{T} \in \mathbb{T}_{\rho_{N-1}}} \pi_f(Y_N(\tilde{T})) \right) \right| \geq \delta^\varepsilon \rho_N^2 \geq \delta^{2\varepsilon}.$$

We conclude with ε replaced by $\varepsilon/2$. □

8 The consequences of Theorem 1.5

In this section we will show how Theorem 1.5 implies Theorem 1.2, Theorem 1.4, and Corollary 1.7.

Proof of Theorem 1.2. Suppose to the contrary that that there exists a Kakeya set $K \subset \mathbb{R}^3$ with $\dim_A(K) = 3 - \beta$ for some $\beta > 0$, i.e. there exists $C, r_0 > 0$ so that

$$\sup_{x \in \mathbb{R}^3} \mathcal{E}_\rho(K \cap B(x, r)) \leq C(r/\rho)^{3-\beta} \quad \text{for all } x \in \mathbb{R}^3, 0 < \rho < r \leq r_0. \quad (8.1)$$

Replacing C by $(1 + 100r_0^{-3})C$, we may suppose that (8.1) holds with $r_0 = 1$.

Let η, δ_0 be the values from Theorem 1.5, with $\varepsilon = \beta/2$. Let $\delta \in (0, \delta_0]$ be chosen sufficiently small (we will describe the choice of δ in further detail below), and let \mathbb{T} be a set of δ^{-2} δ -tubes contained in $N_\delta(K)$ that point in δ separated directions; in particular the tubes in \mathbb{T} are essentially distinct and satisfy the Convex Wolff Axioms with error 100; we will select $\delta \in (0, \delta_0]$ sufficiently small so that $100 \leq \delta^{-\eta}$. For each $T \in \mathbb{T}$, let $Y(T) = T$; this is a 1-dense shading.

Apply Theorem 1.5 with $\varepsilon = \beta/2$. We conclude that there exists $\rho, r \in [\delta, 1]$ with $\rho \leq \delta^\eta r$, and a ball B of radius r so that (1.1) holds, and hence

$$\mathcal{E}_\rho(K \cap B) \gtrsim \rho^{-3} \left| B \cap N_\rho \left(\bigcup_{T \in \mathbb{T}} Y(T) \right) \right| \gtrsim (\rho/r)^\varepsilon (r/\rho)^3 \geq \delta^{-\beta\eta/2} (r/\rho)^{3-\beta}. \quad (8.2)$$

Comparing (8.1) and (8.2), we obtain a contradiction, provided $\delta > 0$ is selected sufficiently small depending on C, β, η , and the implicit constant in (8.2). We conclude that every Kakeya set in \mathbb{R}^3 has Assouad dimension 3. □

Remark 8.1. As noted in the introduction, the conclusion of Theorem 1.2 continues to hold if we remove the hypotheses that the lines point in different directions, and replace this with the requirement that the lines satisfy a variant of the Wolff axioms. This is most easily stated in the discretized setting: For all $\varepsilon > 0$, there exists $\eta > 0$ so that the following holds. Let \mathbb{T} be a set of δ -tubes that satisfy the Convex Wolff Axioms with error $\delta^{-\eta}$. Then $\bigcup T$ has discretized Assouad dimension $\geq 3 - \varepsilon$ at scale δ and scale separation $\delta^{-\eta}$.

Proof of Theorem 1.4. Suppose to the contrary that there exists a Keakeya set K with essentially equal Hausdorff and packing dimension $\alpha < 3$. Without loss of generality, we can assume $K \subset B(0, 1)$. Let $\varepsilon = (3 - \alpha)/4$; in what follows, all implicit constant may depend on ε . Let $\eta > 0$ and $\delta_0 > 0$ be the quantities from Theorem 1.5 for this choice of ε . Let $\eta_1 = (3 - \varepsilon)\eta/6$. Since K has essentially equal Hausdorff and packing dimension, we can find a measure μ supported on K so that

$$r^{\alpha+\eta_1} \lesssim \mu(B(x, r)) \lesssim r^{\alpha-\eta_1}, \quad x \in \text{supp}(\mu), \quad 0 \leq r \leq 1, \quad (8.3)$$

and there exists $\tau > 0$ and a set $\Omega \subset S^2$ of positive measure, so that for each $e \in \Omega$ there is a line ℓ_e pointing in direction e , with $|\ell_e \cap \text{supp}(\mu)| \geq \tau$.

Let Ω_δ be a δ -separated subset of Ω of cardinality $\#\Omega_\delta \gtrsim \delta^{-2}|\Omega|$. For each $e \in \Omega_\delta$, let T_e be a δ -tube with coaxial line ℓ_e , which satisfies $|T_e \cap \ell_e \cap \text{supp}(\mu)| \geq \tau/2$. Define the shading

$$Y(T_e) = \bigcup_{x \in T_e \cap \ell_e \cap \text{supp}(\mu)} B(x, \delta).$$

We have $|Y(T_e)| \gtrsim \tau|T|$ for each tube T_e ; we will select $\delta \in (0, \delta_0]$ sufficiently small so that $|Y(T_e)| \geq \delta^\eta|T|$. Let $\mathbb{T} = \{T_e : e \in \Omega_\delta\}$. Since the tubes in \mathbb{T} point in δ -separated directions, the tubes are essentially distinct and satisfy the Convex Wolff Axioms with error $O(\delta^{-2}/(\#\mathbb{T})) = O(|\Omega|^{-1})$; we will select $\delta \in (0, \delta_0]$ sufficiently small so that \mathbb{T} satisfies the convex Wolff Axioms with error at most $\delta^{-\eta}$. Thus we can apply Theorem 1.5 to conclude there exist scales $\rho, r \in [\delta, 1]$ with $\rho \leq \delta^\eta r$, and a ball B of radius r so that (1.1) holds.

Let $E \subset B$ be a ρ -separated subset of $B \cap N_\rho\left(\bigcup_{T \in \mathbb{T}} Y(T)\right)$ of cardinality $\gtrsim (r/\rho)^{3-\varepsilon}$. Then by (8.3) we have

$$r^{\alpha-\eta_1} \gtrsim \mu(B) \gtrsim \sum_{x \in E} \mu(B(x, \rho)) \gtrsim (r/\rho)^{3-\varepsilon} \rho^{\alpha+\eta_1}.$$

Re-arranging and using the fact that $\rho/r \leq \delta^\eta$, as well as the definition of ε and η_1 , we conclude that

$$\delta^{-\eta(3-\varepsilon)/2} = \delta^{-\eta(3-\varepsilon-\alpha)} \leq (r/\rho)^{-\eta(3-\varepsilon-\alpha)} \lesssim (r\rho)^{-\eta_1} \lesssim \delta^{-2\eta_1} = \delta^{-\eta(3-\varepsilon)/3}.$$

If $\delta \in (0, \delta_0]$ is selected sufficiently small, then this is impossible. We conclude that every Keakeya set in \mathbb{R}^3 with essentially equal Hausdorff and packing dimension must have dimension 3. \square

Proof of Corollary 1.7. Let $\varepsilon > 0$. Let $\eta_0 > 0$ and $\delta_0 > 0$ be the quantities from Theorem 1.5 with $\varepsilon/2$ in place of ε . Let $\eta = \eta_0/2$, let \mathbb{T} be a set of essentially distinct δ -tubes in $B(0, 1)$ that satisfies the Convex Wolff Axioms with error $\delta^{-\eta}$, and let $Y(T) \subset T$ be a shading with $\sum_{T \in \mathbb{T}} |Y(T)| \geq \delta^\eta \sum_{T \in \mathbb{T}} |T|$.

Let $Y_0(T) = Y(T)$. For each $i = 1, \dots$, apply Theorem 1.5 to \mathbb{T} with the shading Y_{i-1} ; let r_i, ρ_i and B_i be the output from this theorem. Define $Y_i(T) = Y_{i-1}(T) \setminus B_i$. We repeat this process until

$\sum_{\mathbb{T}} |Y_i(T)| < \frac{1}{2} \delta^\eta \sum_{\mathbb{T}} |T|$, at which point we have

$$\sum_{T \in \mathbb{T}} \left| Y(T) \cap \bigcup_i B_i \right| \geq \frac{1}{2} \sum_{T \in \mathbb{T}} |Y(T)|.$$

After dyadic pigeonholing, we can select $\rho, r \in [\delta, 1]$ with $\rho \leq \delta^\eta r$, and a set of indices I with the following properties.

- For each index $i \in I$ we have $\rho_i \in [\rho, 2\rho)$ and $r_i \in [r, 2r)$
- $\sum_{T \in \mathbb{T}} \left| Y(T) \cap \bigcup_{i \in I} B_i \right| \gtrsim |\log \delta|^{-2} \sum_{T \in \mathbb{T}} |Y(T)|$.

Next, we can greedily select a set of indices $I' \subset I$ so that the above properties continue to hold (with the implicit constant in the second item worsened slightly), so that the sets $\{B_i : i \in I'\}$ are pairwise disjoint.

We claim that the values of ρ and r described above, plus the shading $Y'(T) = Y(T) \cap \bigcup_{i \in I'} B_i$ satisfies the conclusions of Corollary 1.7. Indeed, we can compute

$$\begin{aligned} \left| N_\rho \left(\bigcup_{T \in \mathbb{T}} Y'(T) \right) \right| &= \left| N_\rho \left(\bigcup_{i \in I'} B_i \cap \bigcup_{T \in \mathbb{T}} Y'(T) \right) \right| \\ &\gtrsim \sum_{i \in I'} \left| B_i \cap N_\rho \left(\bigcup_{T \in \mathbb{T}} Y'(T) \right) \right| \\ &\gtrsim \sum_{i \in I'} (r/\rho)^{\varepsilon/2} |B_i| \\ &\gtrsim (r/\rho)^{\varepsilon/2} \left| N_r \left(\bigcup_{T \in \mathbb{T}} Y'(T) \right) \right| \\ &\geq \delta^{-\eta_0 \varepsilon/2} (r/\rho)^\varepsilon \left| N_r \left(\bigcup_{T \in \mathbb{T}} Y'(T) \right) \right|. \end{aligned}$$

After decreasing δ_0 (and hence δ) if necessary, the quantity $\delta^{-\eta_0 \varepsilon/2}$ dominates the implicit constants in the above inequality, and we obtain (1.2). \square

References

- [1] A. Cordoba. The Keakeya maximal function and the spherical summation multipliers. *Am. J. Math.* 99:1–22, 1977.
- [2] R. Davies. Some remarks on the Keakeya problem. *Math. Proc. Cambridge Philos. Soc.* 69:417–421, 1971.
- [3] K. Fässler and T. Orponen. On restricted families of projections in \mathbb{R}^3 . *Proc. Lond. Math. Soc.* (3) 109 (2014), no. 2, 353–381.
- [4] J. Fraser, E. Olson, and J. Robinson. Some results in support of the Keakeya conjecture. *Real Anal. Exch.* 42(2): 253–268, 2017.
- [5] J. Fraser. *Assouad Dimension and Fractal Geometry*. Cambridge University Press. Cambridge UK. 2020.

- [6] S. Gan, L. Guth, and D. Maldague. An exceptional set estimate for restricted projections to lines in \mathbb{R}^3 . *J. Geom. Anal.* 34:15, 2024.
- [7] N.H. Katz, I. Laba, and T. Tao. An improved bound on the Minkowski dimension of Besicovitch sets in \mathbb{R}^3 . *Ann. of Math.* 152, 383–446, 2000.
- [8] N.H. Katz and T. Tao. Some connections between Falconer’s distance set conjecture, and sets of Furstenberg type. *New York J. Math.* 7, 149–187, 2001.
- [9] N.H. Katz and T. Tao. Recent progress on the Kakeya conjecture. *Publicacions Matemàtiques*, Proceedings of the 6th International Conference on Harmonic Analysis and Partial Differential Equations, U. Barcelona, pp. 161–180, 2002.
- [10] N.H. Katz, S. Wu, and J. Zahl. Kakeya sets from lines in SL_2 . *Ars Inven. Anal.* paper no 6, 1–23, 2023.
- [11] T. Keleti and P. Shmerkin. New bounds on the dimensions of planar distance sets. *Geom. Funct. Anal.*, 29(6):1886–1948, 2019.
- [12] T. Orponen and P. Shmerkin. On the Hausdorff dimension of Furstenberg sets and orthogonal projections in the plane. To appear, *Duke Math. J.* [arXiv:2106.03338](https://arxiv.org/abs/2106.03338), 2021.
- [13] T. Orponen and P. Shmerkin. Projections, Furstenberg sets, and the ABC sum-product problem. [arXiv:2301.10199](https://arxiv.org/abs/2301.10199), 2023
- [14] M. Pramanik, T. Yang, J. Zahl. A Furstenberg-type problem for circles, and a Kaufman-type restricted projection theorem in \mathbb{R}^3 . [arXiv:2207.02259](https://arxiv.org/abs/2207.02259), 2022.
- [15] P. Shmerkin and H. Wang. On the distance sets spanned by sets of dimension $d/2$ in \mathbb{R}^d . [arXiv:2112.09044](https://arxiv.org/abs/2112.09044), 2021.
- [16] H. Wang and J. Zahl. Sticky Kakeya sets and the sticky Kakeya conjecture. [arXiv:2210.09581](https://arxiv.org/abs/2210.09581), 2022.
- [17] T. Wolff. An improved bound for Kakeya type maximal functions. *Rev. Mat. Iberoam.* 11(3):651–674, 1995.
- [18] T. Wolff. Recent work connected with the Kakeya problem, in *Prospects in Mathematics* (Princeton, NJ, 1996), pp. 129–162, Amer. Math. Soc, Providence, RI, 1999.
- [19] T. Wolff. A Kakeya-type problem for circles. *Am. J. Math.* 119(5):985–1026, 1997.