

ON THE METRIC OF THE JET BUNDLE AND SIMILARITY ON WEIGHTED DIRICHLET SPACES

KUI JI, SHANSHAN JI, HYUN-KYOUNG KWON, AND JING XU

ABSTRACT. In general, it is more difficult to formulate a sufficient condition for similarity than a necessary condition. We give sufficient conditions for a Cowen-Douglas operator with a positivity condition to be similar to the backward shift operator on weighted Dirichlet space. This condition involves the holomorphic jet bundle of the eigenvector bundle of the operator.

1. INTRODUCTION

Let $\mathcal{L}(\mathcal{H})$ denote the algebra of bounded linear operators on a complex separable Hilbert space \mathcal{H} . Recall that operators T and S in $\mathcal{L}(\mathcal{H})$ are *unitarily equivalent* ($T \sim_u S$) if there is a unitary operator U such that $T = U^*SU$. An operator T is *similar* to S ($T \sim_s S$) if there is a bounded invertible operator X satisfying $T = X^{-1}SX$. The problem of finding unitary or similarity invariants on an infinite-dimensional space \mathcal{H} is no simple task. An effective approach to study the problem is to make use of operator models given in [1, 2, 4–6, 18, 21, 43, 47, 48, 51], where the shift operators on certain reproducing kernel Hilbert spaces are regarded as universal operators.

Let \mathbb{D} denote the open unit disk in the complex plane \mathbb{C} and let $f(z) = \sum_{n=0}^{\infty} \hat{f}(n)z^n$ be the Taylor expansion of a holomorphic function f on \mathbb{D} . For each real number k , we consider a corresponding reproducing kernel Hilbert space \mathcal{H}_k defined by

$$\mathcal{H}_k := \left\{ f \in \text{Hol}(\mathbb{D}) : \|f\|_{\mathcal{H}_k}^2 = \sum_{n=0}^{\infty} (n+1)^k |\hat{f}(n)|^2 < \infty \right\}.$$

Note that we obtain well-known reproducing kernel Hilbert spaces by substituting different values for k : $k = 1$ gives the Dirichlet space \mathcal{D} (or \mathcal{D}_1), for $k \in (0, 1)$, we have the weighted Dirichlet spaces \mathcal{D}_k , $k = 0$ yields the Hardy space $H^2(\mathbb{D})$, and $k = -1$ results in the Bergman space $L^2(\mathbb{D})$. The backward shift operator M_z^* on a reproducing kernel Hilbert space \mathcal{H}_k belongs to the class $\mathcal{B}_1(\mathbb{D})$ introduced by Cowen and Douglas in [13, 14]. Given an open connected subset Ω of \mathbb{C} , the Cowen-Douglas class $\mathcal{B}_n(\Omega)$ of rank n , is defined as

$$\begin{aligned} \mathcal{B}_n(\Omega) = \{T \in \mathcal{L}(\mathcal{H}) : & (1) \Omega \subset \sigma(T) = \{w \in \mathbb{C} : T - w \text{ is not invertible}\}, \\ & (2) \dim \ker (T - w) = n \text{ for } w \in \Omega, \\ & (3) \bigvee_{w \in \Omega} \ker (T - w) = \mathcal{H}, \text{ for a closed linear span } \bigvee, \\ & (4) \text{ran } (T - w) = \mathcal{H} \text{ for } w \in \Omega\}. \end{aligned}$$

2000 *Mathematics Subject Classification.* Primary 47A45; Secondary 46E22, 46J20.

Key words and phrases. Dirichlet Space, Model Theorem, Multipliers, Weights, Backward Shift Operator, Similarity.

The authors except for HK were supported by the National Natural Science Foundation of China, Grant No. 12371129 and 12471123. The second and the fifth authors appreciate the hospitality of the Department of Mathematics and Statistics at the University at Albany during their visit from Aug. 2023 to Feb. 2024.

Every operator in $\mathcal{B}_n(\Omega)$ can be realized as the adjoint of the multiplication operator on a reproducing kernel Hilbert space consisting of holomorphic functions on $\Omega^* = \{w \in \mathbb{C} : \bar{w} \in \Omega\}$. Moreover, the study of a Cowen-Douglas operator $T \in \mathcal{B}_n(\Omega)$ involves the Hermitian holomorphic vector bundle \mathcal{E}_T and the curvature function $\mathcal{K}_{\mathcal{E}_T}$ defined to be

$$\mathcal{E}_T = \{(w, x) \in \Omega \times \mathcal{H} : x \in \ker(T - w)\}, \quad \pi(w, x) = w,$$

and

$$\mathcal{K}_{\mathcal{E}_T}(w) = -\frac{\partial}{\partial \bar{w}} \left(h_{\mathcal{E}_T}^{-1}(w) \frac{\partial h_{\mathcal{E}_T}(w)}{\partial w} \right),$$

where $h_{\mathcal{E}_T}(w) = (\langle \gamma_j(w), \gamma_i(w) \rangle)_{i,j=1}^n$ denotes the metric associated with a holomorphic frame $\{\gamma_1, \dots, \gamma_n\}$ for \mathcal{E}_T . It is shown in [13] that the operators T and S in $\mathcal{B}_n(\Omega)$ are unitarily equivalent if and only if their complex bundles \mathcal{E}_T and \mathcal{E}_S are equivalent as Hermitian holomorphic vector bundles. They also proved that the curvature and its covariant derivatives form a complete set of unitary invariants. However, the similarity classification problem is much more complex and has not yet been fully resolved.

For operators T and S in $\mathcal{B}_1(\mathbb{D})$, each having $\overline{\mathbb{D}}$ as a K -spectral set, the Cowen-Douglas conjecture ([13], §4.35) states that T is similar to S if and only if $\lim_{|w| \rightarrow 1} \mathcal{K}_T(w)/\mathcal{K}_S(w) = 1$, where \mathcal{K}_T denotes the curvature of \mathcal{E}_T . Clark and Misra gave a positive result in [10], confirming that the conjecture is sufficient for the class of Toeplitz operators considered in [12]. Subsequently, they in [11] introduced a weighted backward shift operator to show that the conjecture is unnecessary, and found that the ratio of the metrics associated with Hermitian holomorphic line bundles is a better similarity invariant than the ratio of the curvatures, which is the main point of the note. In addition, Clark and Misra showed that the backward shift operator T , with $\|T\| \leq 1$, is similar to the backward shift operator S_α with weight sequence $\{[(n+1)/(n+2)]^{\alpha/2}\}_{n \geq 0}$ if and only if the ratio $a_w := \frac{h_{\mathcal{E}_T}(w)}{h_{\mathcal{E}_{S_\alpha}}(w)}$ of the metric is bounded and bounded from 0 ([11], Theorem 1). In the third section of this note, we will use the ratio of metrics and the operator realization of holomorphic vector bundles to provide necessary and sufficient conditions for the similarity of the backward shift operators on the Hardy space and the weighted Bergman space.

There have been some new developments in the past four decades on the similarity problem for the weighted backward shift operators on the spaces \mathcal{H}_k [15, 26, 27, 30, 33, 34, 39, 42, 57]. Uchiyama used a holomorphic frame of Hermitian holomorphic vector bundles to show the similarity between a contraction and the backward shift operator on the Hardy space [55]. Douglas, Treil and the third author gave a necessary and sufficient condition for an n -hypercontraction to be similar to the backward shift operator on a weighted Bergman spaces by using the trace of the curvature [16, 38]. Sarkar and the first author described the similarity of backward shift operators on weighted Bergman spaces using Hilbert modules [29]. The third author gave a necessary condition for the similarity of the backward shift on the Dirichlet space [37]. Note that the original Cowen-Douglas conjecture does not include the backward shift operator in the Dirichlet space and operators similar to it, see Remark 2 in [11]. In this note, we introduce holomorphic jet bundle, which is closely related to holomorphic vector bundles, to consider the similarity of the backward shift operators on the weighted Dirichlet spaces.

In brief, there have been three different approaches to dealing with the similarity problem of operators. The first is a geometric approach used in [16, 24, 37, 38] which gives an explicit formulation of similarity in terms of the trace of the curvature. The second approach is algebraic and it describes the similarity of the direct sum of strongly irreducible operators [8, 31, 32]. The third approach is due to Shields in [49] in which a necessary and sufficient condition for the similarity of weighted shift operators is presented using weight sequences.

In [40], Lin considered the operator realization problem of when a function $f \in H^\infty(\mathbb{D})$ gives rise to a Cowen-Douglas operator and proved that the tensor product $\mathcal{E}_T \otimes \mathcal{E}_S$ from operators $T \in \mathcal{B}_m(\mathbb{D})$ and $S \in \mathcal{B}_n(\mathbb{D})$ is indeed a Hermitian holomorphic eigenvector bundle over \mathbb{D} of some Cowen-Douglas operator in $\mathcal{B}_{nm}(\mathbb{D})$. We are interested in using the geometric quantities of the Hermitian holomorphic eigenvector bundle associated with a Cowen-Douglas operator to characterize similarity. In particular, we will use the metric of the Hermitian holomorphic eigenvector bundle and associated holomorphic jet bundles to give a similarity criteria.

The paper is organized as following. Section 2 fixes the notation and executes some preliminaries. Section 3 deals with some extension results, Theorem 3.4 and Proposition 3.6, which are given in terms of the tensor product of holomorphic vector bundles and the holomorphic vector bundles induced by the defect operator corresponding to the n -hypercontraction. Section 4 shows two sufficient conditions for operator $T \in \mathcal{L}(\mathcal{H})$ to be similar to the backward shift operators D_α^* on the weighted Dirichlet space \mathcal{D}_α , namely Theorem 4.3 and Theorem 4.9, the techniques used for these results are the jet bundles of holomorphic complex bundles and the properties of multiplier algebras in weighted Dirichlet spaces.

2. PRELIMINARIES

2.1. Multipliers. Given a reproducing kernel Hilbert space \mathcal{H}_K of holomorphic functions on \mathbb{D} with reproducing kernel K , the *multiplier algebra* of \mathcal{H}_K is the set of $\phi \in \text{Hol}(\mathbb{D})$ defined as

$$\text{Mult}(\mathcal{H}_K) = \{\phi \in \text{Hol}(\mathbb{D}) : \phi f \in \mathcal{H}_K \text{ whenever } f \in \mathcal{H}_K\}.$$

It follows from the closed graph theorem that if ϕ is a multiplier of \mathcal{H}_K , then M_ϕ , the multiplication operator by ϕ , is a bounded linear operator on \mathcal{H}_K and

$$\|\phi\|_{\text{Mult}(\mathcal{H}_K)} = \|M_\phi\| = \sup_{\|f\|_{\mathcal{H}_K}=1} \|M_\phi f\|_{\mathcal{H}_K} = \sup_{\|f\|_{\mathcal{H}_K}=1} \|\phi f\|_{\mathcal{H}_K}.$$

Since $\langle f, M_\phi^* K(\cdot, w) \rangle_{\mathcal{H}_K} = \langle M_\phi f, K(\cdot, w) \rangle_{\mathcal{H}_K} = \phi(w)f(w) = \langle f, \overline{\phi(w)} K(\cdot, w) \rangle_{\mathcal{H}_K}$ for any $f \in \mathcal{H}_K$,

$$M_\phi^* K(\cdot, w) = \overline{\phi(w)} K(\cdot, w), \quad w \in \mathbb{D},$$

and $\|\phi\|_\infty \leq \|M_\phi\| = \|\phi\|_{\text{Mult}(\mathcal{H}_K)}$.

For the Hardy space $H^2(\mathbb{D})$ and the Bergman space $L^2(\mathbb{D})$,

$$\text{Mult}(H^2(\mathbb{D})) = \text{Mult}(L^2(\mathbb{D})) = H^\infty(\mathbb{D}).$$

For the multiplier algebra of the weighted Dirichlet space \mathcal{D}_α for $\alpha \in (0, 1]$, it is well-known that $\text{Mult}(\mathcal{D}_\alpha) \subsetneq H^\infty(\mathbb{D})$. In fact, it is even proven in [17, 50] that $\text{Mult}(\mathcal{D}_\alpha) \subsetneq H^\infty(\mathbb{D}) \cap \mathcal{D}_\alpha$.

Given separable Hilbert spaces E_1 and E_2 , the space $\mathcal{H}_K \otimes E_1$ can be seen as a space of holomorphic functions $f : \mathbb{D} \rightarrow E_1$ with Taylor series $f(z) = \sum_{n=0}^{\infty} \hat{f}(n)z^n$, where $\{\hat{f}(n)\}_{n=0}^{\infty} \subset E_1$ and

$\sum_{n=0}^{\infty} \|\hat{f}(n)\|_{E_1}^2 \|z^n\|_{\mathcal{H}_K}^2 < \infty$. A multiplier from $\mathcal{H}_K \otimes E_1$ to $\mathcal{H}_K \otimes E_2$ is an operator-valued function $\Phi : \mathbb{D} \rightarrow \mathcal{L}(E_1, E_2)$ such that for every $f \in \mathcal{H}_K \otimes E_1$, $\Phi f \in \mathcal{H}_K \otimes E_2$. The multiplication operator M_Φ is also one such that given $f \in \mathcal{H}_K \otimes E_1$, $M_\Phi f \in \mathcal{H}_K \otimes E_2$. Then for every $f \in \mathcal{H}_K \otimes E_1$ and $g \in E_2$,

$$\begin{aligned} \langle f, M_\Phi^* K(\cdot, w) \otimes g \rangle_{\mathcal{H}_K \otimes E_1} &= \langle M_\Phi f, K(\cdot, w) \otimes g \rangle_{\mathcal{H}_K \otimes E_2} \\ &= \langle \Phi(w) f(w), g \rangle_{E_2} \\ &= \langle f(w), \Phi(w)^* g \rangle_{E_1} \\ &= \langle f, K(\cdot, w) \otimes \Phi(w)^* g \rangle_{\mathcal{H}_K \otimes E_1}. \end{aligned}$$

The multiplication operator has the following important property given in [3]:

Theorem 2.1. *Let $\Phi : \mathbb{D} \rightarrow \mathcal{L}(E_1, E_2)$ be an operator-valued function. If $\Phi \in \text{Mult}(\mathcal{H}_K \otimes E_1, \mathcal{H}_K \otimes E_2)$, then*

$$M_\Phi^*(K(\cdot, w) \otimes g) = K(\cdot, w) \otimes \Phi(w)^* g, \quad w \in \mathbb{D}, g \in E_2.$$

Conversely, if $\Phi : \mathbb{D} \rightarrow \mathcal{L}(E_1, E_2)$ and the mapping $K(\cdot, w) \otimes g \mapsto K(\cdot, w) \otimes \Phi(w)^ g$ extends to a bounded operator $X \in \mathcal{L}(\mathcal{H}_K \otimes E_2, \mathcal{H}_K \otimes E_1)$, then $\Phi \in \text{Mult}(\mathcal{H}_K \otimes E_1, \mathcal{H}_K \otimes E_2)$ and $X = M_\Phi^*$.*

It is clear now that any multiplier Φ is bounded and holomorphic, that is, $\Phi \in H_{E_1 \rightarrow E_2}^\infty(\mathbb{D})$, where $H_{E_1 \rightarrow E_2}^\infty(\mathbb{D})$ is the space of bounded analytic functions defined on \mathbb{D} whose function values are bounded linear operators from a Hilbert space E_1 to another one E_2 . When $\text{Mult}(\mathcal{H}_K) = H^\infty(\mathbb{D})$, then $\text{Mult}(\mathcal{H} \otimes E_1, \mathcal{H}_K \otimes E_2) = H_{E_1 \rightarrow E_2}^\infty(\mathbb{D})$.

Since a bounded linear operator on the Hardy space commuting with the operator of multiplication by z is given by the multiplication operator by a function $\varphi \in H^\infty(\mathbb{D})$ with norm $\|\varphi\|_\infty$, the corona problem is closely related to the multipliers on the reproducing kernel Hilbert space of analytic functions (see [7, 56]). The original corona problem posed by Kakutani in 1941 asks whether the unit disk \mathbb{D} is dense in the maximal ideal space of $H^\infty(\mathbb{D})$. The question was later answered in the affirmative by Carleson in [9]. In [23], Hörmander introduced a new and a much simpler method for solving the problem which was popularized by Wolff (see [20, 45]). There have been many versions of the operator corona theorem given by Fuhrmann, Rosenblum, Tolokonnikov, Uchiyama, and Vasyunin [19, 46, 53, 54]. In particular, we have the following result by Fuhrmann [53]:

Theorem 2.2. *For an operator-valued function $F \in H_{E_1 \rightarrow E_2}^\infty(\mathbb{D})$ with $\dim E_1 < \infty$ and $\dim E_2 = \infty$ such that*

$$F^*(w)F(w) \geq \delta, \quad w \in \mathbb{D},$$

for some $\delta > 0$, there exists a function $G \in H_{E_2 \rightarrow E_1}^\infty(\mathbb{D})$ satisfying

$$G(w)F(w) \equiv I_{E_1}, \quad w \in \mathbb{D}.$$

One can consider other multiplier algebras. For a weighted Dirichlet space \mathcal{D}_α for $\alpha \in (0, 1]$, $\text{Mult}(\mathcal{D}_\alpha)$ is a Banach algebra under the norm $\|\phi\|_{\text{Mult}(\mathcal{D}_\alpha)} = \|M_\phi\|$ for $f \in \text{Mult}(\mathcal{D}_\alpha)$. In the work [36], Kidane and Trent proved the corona theorem for $\text{Mult}(\mathcal{D}_\alpha)$ for $\alpha \in (0, 1]$. Let $\{f_i\}_{i=1}^\infty \subset \text{Mult}(\mathcal{D}_\alpha)$

and $F = (f_1, f_2, \dots)$. Define a row operator $M_F^R : \bigoplus_1^\infty \mathcal{D}_\alpha \rightarrow \mathcal{D}_\alpha$ by

$$M_F^R((h_1, h_2, \dots)^T) = \sum_{i=1}^{\infty} f_i h_i,$$

where $(h_1, h_2, \dots)^T \in \bigoplus_1^\infty \mathcal{D}_\alpha$. Similarly, define a column operator $M_F^C : \mathcal{D}_\alpha \rightarrow \bigoplus_1^\infty \mathcal{D}_\alpha$ by

$$M_F^C(h) = (f_1 h, f_2 h, \dots)^T, \quad h \in \mathcal{D}_\alpha.$$

A result that we will later need concerning these row and column operators is as follows and is due to Kidane and Trent [36]:

Theorem 2.3. *Let $M_F^C \in \mathcal{L}\left(\mathcal{D}_\alpha, \bigoplus_1^\infty \mathcal{D}_\alpha\right)$. Then $M_F^R \in \mathcal{L}\left(\bigoplus_1^\infty \mathcal{D}_\alpha, \mathcal{D}_\alpha\right)$ and $\|M_F^R\| \leq \sqrt{10} \|M_F^C\|$.*

The following corona theorem for $\text{Mult}(\mathcal{D}_\alpha)$ is also proven in [36]:

Theorem 2.4. *Let $\{f_i\}_{i=1}^\infty \subset \text{Mult}(\mathcal{D}_\alpha)$. Assume that $\|M_F^C\| \leq 1$ and $0 < \delta^2 \leq \sum_{i=1}^\infty |f_i(w)|^2$ for all $w \in \mathbb{D}$. Then there exists a positive number $C_\alpha < \infty$ and $\{g_i\}_{i=1}^\infty \subset \text{Mult}(\mathcal{D}_\alpha)$ such that*

$$\sum_{i=1}^{\infty} f_i g_i = 1 \quad \text{and} \quad \|M_G^C\| \leq \frac{C_\alpha}{\delta^4}.$$

2.2. Model theorem. The original model theorem states that a contraction $T \in \mathcal{L}(\mathcal{H})$ with $\lim_{j \rightarrow \infty} \|T^j h\| = 0$ for all $h \in \mathcal{H}$ is unitarily equivalent to the backward shift operator restricted to some co-invariant subspace of a vector-valued Hardy space [18, 45, 48, 51]. Numerous analogues have been proposed in [1, 2, 4–6, 43, 44]. In particular, we mention the model involving the backward shift on the Hardy space, weighted Bergman space, and weighted Dirichlet space given by Sz.-Nagy-Foiaş [51], Agler [1, 2], and Müller [43], respectively. The result has been previously used by Douglas, Treil and the third author to give a necessary and sufficient condition for the similarity to the backward shift on the Hardy space [38] and the weighted Bergman space [16]. Subsequently, the third author to give a necessary condition for the similarity to the backward shift on the Dirichlet space [37].

If n is a positive integer, then an n -hypercontraction is an operator T with $\sum_{j=0}^k (-1)^j \binom{k}{j} (T^*)^j T^j \geq 0$ for all $1 \leq k \leq n$. In addition, we denote $D_T := \left(\sum_{j=0}^n (-1)^j \binom{n}{j} (T^*)^j T^j \right)^{1/2}$ as the defect operator of the n -hypercontraction T .

We denote \mathcal{M}_n as a Hilbert space with the reproducing kernel $K_n(z, w) = \frac{1}{(1-\bar{w}z)^n}$ for $z, w \in \mathbb{D}$, where n is a positive integer. As can be easily checked, different function spaces correspond to different n 's: the Hardy space for $n = 1$ and the weighted Bergman spaces for $n \geq 2$. The vector-valued space $\mathcal{M}_{n,E}$ for a separable Hilbert space E can be defined as previously. The forward shift $S_{n,E}$ on $\mathcal{M}_{n,E}$ is defined as $S_{n,E}f(z) = zf(z)$, and the backward shift operator $S_{n,E}^*$ is its adjoint. We are ready to state the following theorem by Agler [1, 2].

Theorem 2.5. *For $T \in \mathcal{L}(\mathcal{H})$, there exist a Hilbert space E and an $S_{n,E}^*$ -invariant subspace $\mathcal{N} \subseteq \mathcal{M}_{n,E}$ such that $T \sim_u S_{n,E}^*|_{\mathcal{N}}$ if and only if T is an n -hypercontraction and $\lim_{j \rightarrow \infty} \|T^j x\| = 0$ for all $x \in \mathcal{H}$.*

Consider the vector-valued l^2 space with components from a Hilbert space E with orthonormal basis $\{e_i\}_{i=0}^\infty$. The backward shift $S_{\alpha,E}^* \in \mathcal{L}(l^2 \otimes E)$ with a weight sequence $\{\alpha_i\}_{i \geq 1}$ is defined by

$$S_{\alpha,E}^* e_0 = 0 \quad \text{and} \quad S_{\alpha,E}^* e_i = \alpha_i e_{i-1}, \quad i \geq 1.$$

Setting $b_i = \prod_{j=1}^i \alpha_j^{-2}$ for $i \geq 1$, we have the following theorem by Müller [43], where we assume $\alpha_1 \geq \alpha_2 \geq \dots > 0$:

Theorem 2.6. *Let $T \in \mathcal{L}(\mathcal{H})$ satisfy $\sum_{s=1}^\infty \|T^s\|^2 b_s \leq 1$. Then there exist a Hilbert space E and a subspace $\mathcal{N} \subset l^2 \otimes E$ with $S_{\alpha,E}^* \mathcal{N} \subset \mathcal{N}$ such that $T \sim_u S_{\alpha,E}^*|_{\mathcal{N}}$.*

For general unilateral shift operators, the work of Shields given in [49] is quite extensive. The result that we need the most in this note is the following, which easily shows that the backward shift operators on the Hardy space, the weighted Bergman spaces, and the weighted Dirichlet spaces are not similar.

Theorem 2.7. *Let T_1 and T_2 be unilateral weighted shifts with weight sequences $\{\alpha_i\}_{i \geq 1}$ and $\{\beta_i\}_{i \geq 1}$, respectively. Then $T_1 \sim_s T_2$ if and only if there exist constants C_1 and C_2 such that for every positive integer l ,*

$$0 < C_1 \leq \left| \frac{\alpha_k \alpha_{k+1} \cdots \alpha_l}{\beta_k \beta_{k+1} \cdots \beta_l} \right| \leq C_2$$

for all $1 \leq k \leq l$.

3. SIMILARITY IN THE CLASS $\mathcal{B}_m(\mathbb{D})$

In this section, we study Cowen-Douglas operators whose associated Hermitian holomorphic eigenvector bundles possess a tensor structure. A relationship between the corresponding metric matrix and similarity will be given. Let \mathcal{E} be a Hermitian holomorphic vector bundle over \mathbb{D} of rank n . One can form the metric matrix $h(w) = (\langle \gamma_j(w), \gamma_i(w) \rangle)_{i,j=1}^n$ using a holomorphic frame $\{\gamma_1, \dots, \gamma_n\}$ of \mathcal{E} . Denote by $\{\sigma_i\}_{i=1}^n$ an orthonormal basis for \mathbb{C}^n and let

$$E = \bigvee_{w \in \mathbb{D}} \{\gamma_i(w) : 1 \leq i \leq n\}.$$

Now define an operator-valued function $F \in H_{\mathbb{C}^n \rightarrow E}^\infty(\mathbb{D})$ such that

$$F(w)\sigma_i = \gamma_i(w), \quad 1 \leq i \leq n.$$

Lemma 3.1. *The following statements are equivalent:*

- (1) F is left-invertible in $H_{\mathbb{C}^n \rightarrow E}^\infty(\mathbb{D})$.
- (2) There exist constants C_1 and C_2 such that

$$0 < C_1 I_n \leq h(w) \leq C_2 I_n, \quad w \in \mathbb{D}.$$

- (3) For every $1 \leq i \leq n$, $\|\gamma_i(w)\|$ is uniformly bounded on \mathbb{D} and there exist constants A_1 and A_2 such that

$$0 < A_1 \leq \det h(w) \leq A_2, \quad w \in \mathbb{D}.$$

Proof. In proving the implications, we use the well-known fact that for the eigenvalues $\{\lambda_i(w)\}_{i=1}^n$ of the metric matrix $h(w) = (\langle \gamma_j(w), \gamma_i(w) \rangle)_{i,j=1}^n$,

$$\sum_{i=1}^n \lambda_i(w) = \sum_{i=1}^n \|\gamma_i(w)\|^2, \quad w \in \mathbb{D}.$$

(1) \Rightarrow (2): Since F is left-invertible in $H_{\mathbb{C}^n \rightarrow E}^\infty(\mathbb{D})$, there is a $G \in H_{E \rightarrow \mathbb{C}^n}^\infty(\mathbb{D})$ such that

$$G(w)F(w) = I_{\mathbb{C}^n}, \quad w \in \mathbb{D}.$$

It is then obvious that $C_1 = 1/\|G\|^2$. Next, since $\|\gamma_i(w)\|, 1 \leq i \leq n$, are uniformly bounded on \mathbb{D} , $C_2 = n \sup\{\|\gamma_i(w)\|, 1 \leq i \leq n\}$.

(2) \Rightarrow (1): Since $0 < C_1 I_n \leq h(w) \leq C_2 I_n$,

$$C_1 \leq \lambda_i(w) \leq C_2 \quad \text{and} \quad 0 < nC_1 \leq \sum_{i=1}^n \|\gamma_i(w)\|^2 = \sum_{i=1}^n \lambda_i(w) \leq nC_2,$$

so that the function $F \in H_{\mathbb{C}^n \rightarrow E}^\infty(\mathbb{D})$ satisfies

$$F(w)^* F(w) = (\langle \gamma_j(w), \gamma_i(w) \rangle)_{i,j=1}^n \geq C_1, \quad w \in \mathbb{D}.$$

Then by Theorem 2.2, there exists a function $G \in H_{E \rightarrow \mathbb{C}^n}^\infty(\mathbb{D})$ such that

$$G(w)F(w) = I_{\mathbb{C}^n}, \quad w \in \mathbb{D}.$$

(2) \Rightarrow (3): Since $0 < C_1 I_n \leq h(w) \leq C_2 I_n$,

$$C_1 \leq \lambda_i(w) \leq C_2 \quad \text{and} \quad \sum_{i=1}^n \|\gamma_i(w)\|^2 = \sum_{i=1}^n \lambda_i(w) \leq nC_2,$$

so that each $\|\gamma_i(w)\|, 1 \leq i \leq n$, is uniformly bounded on \mathbb{D} and

$$0 < C_1^n \leq \det h(w) = \prod_{i=1}^n \lambda_i(w) \leq (nC_2)^n, \quad w \in \mathbb{D}.$$

(3) \Rightarrow (2): Since $h(w)$ is positive definite and there exists some $A > 0$ such that $\|\gamma_i(w)\|^2 \leq A$ for every $1 \leq i \leq n$ and $w \in \mathbb{D}$,

$$\lambda_i(w) \geq 0, \quad \prod_{i=1}^n \lambda_i(w) \leq (nA)^n, \quad \text{and} \quad h(w) \leq nAI_n.$$

We claim that $C := \inf_{w \in \mathbb{D}} \{\lambda_i(w) : 1 \leq i \leq n\} > 0$. If not, we assume, without loss of generality that $\inf_{w \in \mathbb{D}} \lambda_1(w) = 0$. Then there is a Cauchy sequence $\{w_k\}_{k=1}^\infty \subset \mathbb{D}$ such that $\lim_{k \rightarrow \infty} \lambda_1(w_k) = 0$. This implies that $\prod_{i=2}^n \lambda_i(w_k) \rightarrow \infty$ as $k \rightarrow \infty$, which is obviously contradictory to $\|\gamma_i(w)\|^2 \leq A$. Thus, $0 < CI_n \leq h(w) \leq nAI_n$ for every $w \in \mathbb{D}$. \square

For the Cowen-Douglas conjecture, Clark and Misra noted in [11] that the ratio a_w of the metrics is more suitable as a similarity invariant for the Cowen-Douglas operator. For operators $T \in \mathcal{B}_m(\mathbb{D})$ and $S \in \mathcal{B}_m(\mathbb{D})$, if $\mathcal{E}_T = \mathcal{E}_S \otimes \mathcal{E}$ for some Hermitian holomorphic vector bundle \mathcal{E} over \mathbb{D} , then the metrics of these holomorphic vector bundles satisfy $h_{\mathcal{E}_T} = h_{\mathcal{E}_S} \otimes h_{\mathcal{E}}$. In particular, when \mathcal{E}_T and \mathcal{E}_S are holomorphic line bundles, the ratio of their metrics is the metric of holomorphic line bundle \mathcal{E} , and thus $a_w := h_{\mathcal{E}}(w) = \frac{h_{\mathcal{E}_T}(w)}{h_{\mathcal{E}_S}(w)}$. From the model theorems given in [1, 2, 4, 43–45, 52] and

the operator theoretic realization in [40], we know many operators whose associated holomorphic eigenvector bundles possess a tensor structure. Now provide detailed proof of the Cowen-Douglas operator with n -hypercontraction possessing this property, which was proposed in [2].

Lemma 3.2. *If $T \in \mathcal{B}_m(\mathbb{D})$ is an n -hypercontraction, then there exists a Hermitian holomorphic vector bundle \mathcal{E} over \mathbb{D} of rank m such that \mathcal{E}_T and $\mathcal{E}_{S_n^*} \otimes \mathcal{E}$ are equivalent as Hermitian holomorphic vector bundles, where S_n^* is the backward shift operator on \mathcal{M}_n .*

Proof. Since $T \in \mathcal{L}(\mathcal{H})$ is an n -hypercontraction and $\bigvee_{w \in \mathbb{D}} \ker(T-w) = \mathcal{H}$, we have that $\lim_{j \rightarrow \infty} \|T^j x\| = 0$ for all $x \in \mathcal{H}$. From Theorem 2.5, T is unitarily equivalent to the restriction of $S_{n,E}^*$ to an invariant subspace \mathcal{N} of $\mathcal{M}_{n,E}$ for some Hilbert space E , that is, there exists a unitary operator $V : \mathcal{H} \rightarrow \mathcal{N}$ defined as

$$Vx = \sum_{k=0}^{\infty} \frac{z^k}{\|z^k\|_{\mathcal{M}_n}^2} \otimes D_T T^k x$$

such that $\mathcal{N} = \overline{\text{ran} V}$, $E = \overline{\text{ran} D_T}$ and $VT = (S_{n,E}^*|_{\mathcal{N}})V$, where D_T is the defect operator of the n -hypercontraction T . One of the main results in [13], Theorem 1.14, shows that two Cowen-Douglas operators are unitarily equivalent if and only if their induced holomorphic complex bundles are equivalent as Hermitian holomorphic vector bundles. Therefore, for a holomorphic frame $\gamma = \{\gamma_1, \dots, \gamma_m\}$ of \mathcal{E}_T , we have that

$$V\gamma_i(w) = \sum_{k=0}^{\infty} \frac{z^k}{\|z^k\|_{\mathcal{M}_n}^2} \otimes D_T T^k \gamma_i(w) = \sum_{k=0}^{\infty} \frac{z^k}{\|z^k\|_{\mathcal{M}_n}^2} \otimes D_T w^k \gamma_i(w) = K_n(z, \bar{w}) \otimes D_T \gamma_i(w), \quad 1 \leq i \leq m,$$

and the eigenvector bundle $\ker(S_{n,E}^*|_{\mathcal{N}} - w) = \bigvee \{K_n(\cdot, \bar{w}) D_T \gamma_i(w) : 1 \leq i \leq m\}$. From the eigenvector subspaces

$$\ker(T - w) = \bigvee \{\gamma_1(w), \dots, \gamma_m(w)\} \quad \text{and} \quad \ker(S_n^* - w) = \bigvee \{K_n(\cdot, \bar{w})\}$$

are analytically dependent on $w \in \mathbb{D}$, we can say the same for the subspace $\mathcal{E}_{D_T}(w) = \bigvee \{D_T \gamma_i(w) : 1 \leq i \leq m\}$. So $\mathcal{E}_{D_T} = \{(w, x) \in \mathbb{D} \times \mathcal{E}(w) : x \in \mathcal{E}_{D_T}(w)\}$ is a m -dimensional Hermitian holomorphic vector bundle over \mathbb{D} with the natural projection $\pi, \pi(w, x) = w$, and $\{D_T \gamma_1, \dots, D_T \gamma_m\}$ is a holomorphic frame of \mathcal{E}_{D_T} . Hence, \mathcal{E}_T and $\mathcal{E}_{S_n^*} \otimes \mathcal{E}_{D_T}$ are equivalent as Hermitian holomorphic vector bundles. \square

For the n -hypercontraction $T \in \mathcal{B}_m(\mathbb{D})$, let \mathcal{E}_{D_T} be the holomorphic complex bundle induced by the defect operator D_T of T , which has the following definition.

Definition 3.3. *If $T \in \mathcal{B}_m(\mathbb{D})$ is an n -hypercontraction and $\gamma = \{\gamma_1, \dots, \gamma_m\}$ is a holomorphic frame of \mathcal{E}_T , we said $\mathcal{E}_{D_T} = \{(w, x) \in \mathbb{D} \times \mathcal{E}_{D_T}(w) : x \in \mathcal{E}_{D_T}(w)\}$ is a Hermitian holomorphic vector bundle over \mathbb{D} of rank m induced by the defect operator D_T of T , where $\mathcal{E}_{D_T}(w) = \bigvee \{D_T \gamma_i(w) : 1 \leq i \leq m\}$.*

For the n -hypercontraction in $\mathcal{B}_1(\mathbb{D})$, Theorem 1 in [11] and Theorem 3.4 in [25] show that the ratio of the metrics of holomorphic line bundles is a similarity invariant for the backward shift operators on the Hardy space and the weighted Bergman spaces. More generally, for the similarity of operators in $\mathcal{B}_m(\mathbb{D})$, we use the multiplier algebra and Lemma 3.2 to give the following theorem, the main theorem of this section, concerns a similarity criteria for n -hypercontraction $T \in \mathcal{B}_m(\mathbb{D})$ by using the metric $h_{\mathcal{E}_{D_T}}$ of the holomorphic complex bundle \mathcal{E}_{D_T} :

Theorem 3.4. *Let $T \in \mathcal{B}_m(\mathbb{D})$ be an n -hypercontraction, then the following statements are equivalent:*

- (1) *The operator T is similar to the backward shift operator S_{n, \mathbb{C}^m}^* on $\mathcal{M}_{n, \mathbb{C}^m}$.*
- (2) *There exist constants C_1 and C_2 such that the metric matrix $h_{\mathcal{E}_{D_T}}(w)$ of \mathcal{E}_{D_T} in some holomorphic frame satisfies*

$$0 < C_1 I_m \leq h_{\mathcal{E}_{D_T}}(w) \leq C_2 I_m, \quad w \in \mathbb{D}.$$

- (3) *There exists a holomorphic frame $\tilde{\gamma} = \{\tilde{\gamma}_1, \dots, \tilde{\gamma}_m\}$ of \mathcal{E}_{D_T} such that each $\|\tilde{\gamma}_i(w)\|$ is uniformly bounded on \mathbb{D} , and there are constants A_1 and A_2 such that the metric matrix $h_{\mathcal{E}_{D_T}}(w)$ under holomorphic frame $\tilde{\gamma}$ satisfies*

$$0 < A_1 \leq \det h_{\mathcal{E}_{D_T}}(w) \leq A_2 \quad w \in \mathbb{D}.$$

Proof. Since $T \in \mathcal{B}_m(\mathbb{D})$ is an n -hypercontraction, by Lemma 3.2, we know that

$$\mathcal{E}_{D_T} = \{(w, x) \in \mathbb{D} \times \mathcal{E}_{D_T}(w) : x \in \mathcal{E}_{D_T}(w)\}$$

is a Hermitian holomorphic vector bundle over \mathbb{D} of rank m , $\{D_T \gamma_1, \dots, D_T \gamma_m\}$ is a holomorphic frame of \mathcal{E}_{D_T} for every holomorphic frame $\gamma = \{\gamma_1, \dots, \gamma_m\}$ of \mathcal{E}_T , and $\mathcal{E}_T \sim_u \mathcal{E}_{S_n^*} \otimes \mathcal{E}_{D_T}$. Since the equivalence (2) \Leftrightarrow (3) was already explained above in Lemma 3.1, it remains to show (3) \Rightarrow (1) \Rightarrow (2) in order to complete the proof.

(3) \Rightarrow (1): Using the assumption that $\|D_T \gamma_i(w)\|, 1 \leq i \leq m$, are uniformly bounded on \mathbb{D} , we first define a function $F \in H_{\mathbb{C}^m \rightarrow E}^\infty(\mathbb{D})$ by

$$F(w)(\sigma_i) = D_T \gamma_i(w), \quad 1 \leq i \leq m,$$

where $\{\sigma_i\}_{i=1}^m$ is an orthonormal basis of \mathbb{C}^m and $E = \bigvee_{w \in \mathbb{D}} \{D_T \gamma_i(w) : 1 \leq i \leq m\}$. Then by Lemma 3.1, there exists a function $G \in H_{E \rightarrow \mathbb{C}^m}^\infty(\mathbb{D})$ such that

$$G(w)F(w) = I_{\mathbb{C}^m} \quad w \in \mathbb{D}.$$

Next we let $F^\#(w) := F(\bar{w})$ and $G^\#(w) := G(\bar{w})$ and note that

$$(F^\#)^* \in H_{E \rightarrow \mathbb{C}^m}^\infty(\mathbb{D}) = \text{Mult}(\mathcal{M}_{n, E}, \mathcal{M}_{n, \mathbb{C}^m}) \quad \text{and} \quad (G^\#)^* \in H_{\mathbb{C}^m \rightarrow E}^\infty(\mathbb{D}) = \text{Mult}(\mathcal{M}_{n, \mathbb{C}^m}, \mathcal{M}_{n, E}).$$

Then from Theorem 2.1, we know that for $1 \leq i \leq m$,

$$M_{(F^\#)^*}^*(K_n(\cdot, \bar{w}) \otimes \sigma_i) = K_n(\cdot, \bar{w}) \otimes F^\#(\bar{w})(\sigma_i) = K_n(\cdot, \bar{w}) \otimes F(w)(\sigma_i) = K_n(\cdot, \bar{w}) \otimes D_T \gamma_i(w),$$

and

$$M_{(G^\#)^*}^* M_{(F^\#)^*}^*(K_n(\cdot, \bar{w}) \otimes \sigma_i) = M_{(G^\#)^*}^*(K_n(\cdot, \bar{w}) \otimes F(w)(\sigma_i)) = K_n(\cdot, \bar{w}) \otimes G(w)F(w)(\sigma_i) = K_n(\cdot, \bar{w}) \otimes \sigma_i.$$

Hence,

$$M_{(G^\#)^*}^* M_{(F^\#)^*}^* = I_{\mathcal{M}_{n, \mathbb{C}^m}}.$$

Since $\mathcal{M}_{n, \mathbb{C}^m} = \bigvee_{w \in \mathbb{D}} \{K_n(\cdot, \bar{w}) \otimes \sigma_i : 1 \leq i \leq m\}$ and the operator $M_{(F^\#)^*}^*$ is left-invertible, we conclude that $\text{ran } M_{(F^\#)^*}^* = \bigvee_{w \in \mathbb{D}} \{K_n(\cdot, \bar{w}) \otimes D_T \gamma_i(w) : 1 \leq i \leq m\}$. Therefore, the operator

$$X := M_{(F^\#)^*}^* : \mathcal{M}_{n, \mathbb{C}^m} \rightarrow \bigvee_{w \in \mathbb{D}} \{K_n(\cdot, \bar{w}) \otimes D_T \gamma_i(w) : 1 \leq i \leq m\}$$

is invertible and for every $w \in \mathbb{D}$,

$$X(\mathcal{E}_{S_{n,\mathbb{C}^m}^*}(w)) = (\mathcal{E}_{S_n^*} \otimes \mathcal{E}_{D_T})(w).$$

Considering the composite of the unitary operator V , shown Lemma 3.2, with X , we see that the invertible operator $X^{-1}V$ satisfies $(X^{-1}V)\mathcal{E}_T(w) = \mathcal{E}_{S_{n,\mathbb{C}^m}^*}(w)$ for all $w \in \mathbb{D}$. It follows that T is similar to S_{n,\mathbb{C}^m}^* .

(1) \Rightarrow (2): Since $T \sim_s S_{n,\mathbb{C}^m}^*$ and $\mathcal{E}_T \sim_u \mathcal{E}_{S_n^*} \otimes \mathcal{E}_{D_T}$, there exists a bounded invertible operator X such that

$$X(\mathcal{E}_{S_{n,\mathbb{C}^m}^*}(w)) = (\mathcal{E}_{S_n^*} \otimes \mathcal{E}_{D_T})(w), \quad w \in \mathbb{D}.$$

Thus, for some holomorphic frame $\{\tilde{\gamma}_1, \dots, \tilde{\gamma}_m\}$ of \mathcal{E}_{D_T} , we have

$$(3.1) \quad X(K_n(\cdot, \bar{w}) \otimes \sigma_i) = K_n(\cdot, \bar{w}) \otimes \tilde{\gamma}_i(w), \quad 1 \leq i \leq m,$$

where $\{\sigma_i\}_{i=1}^m$ is an orthonormal basis of \mathbb{C}^m . Since X is bounded, $\|\tilde{\gamma}_i(w)\|$ is uniformly bounded on \mathbb{D} for each $1 \leq i \leq m$, and then

$$h_{\mathcal{E}_{D_T}}(w) = (\langle \tilde{\gamma}_j(w), \tilde{\gamma}_i(w) \rangle)_{i,j=1}^m \leq C_2 I_m, \quad w \in \mathbb{D},$$

for some $C_2 > 0$.

Now define a function $F \in H_{\mathbb{C}^m \rightarrow E}^\infty(\mathbb{D})$ as

$$F(w)\sigma_i = \tilde{\gamma}_i(w), \quad 1 \leq i \leq m,$$

where $E = \bigvee_{w \in \mathbb{D}} \{\tilde{\gamma}_i(w) : 1 \leq i \leq m\}$. Obviously, the function $F^\#$ defined on \mathbb{D} as $F^\#(w) := F(\bar{w})$ is such that $(F^\#)^* \in \text{Mult}(\mathcal{M}_{n,E}, \mathcal{M}_{n,\mathbb{C}^m})$. By Theorem 2.1,

$$(3.2) \quad M_{(F^\#)^*}^*(K_n(\cdot, \bar{w}) \otimes \sigma_i) = K_n(\cdot, \bar{w}) \otimes F^\#(\bar{w})(\sigma_i) = K_n(\cdot, \bar{w}) \otimes F(w)(\sigma_i) = K_n(\cdot, \bar{w}) \otimes \tilde{\gamma}_i(w),$$

and $\text{ran } M_{(F^\#)^*}^* = \bigvee_{w \in \mathbb{D}} \{K_n(\cdot, \bar{w}) \otimes \tilde{\gamma}_i(w) : 1 \leq i \leq m\}$. From $\mathcal{M}_n = \bigvee_{w \in \mathbb{D}} K_n(\cdot, \bar{w})$, (3.1) and (3.2), we see that $X = M_{(F^\#)^*}^*$. Since X is invertible, for every $h = \lambda K_n(\cdot, \bar{w}) \in \mathcal{M}_n$, $\lambda \in \mathbb{C}$, $g \in \mathbb{C}^m$, and $w \in \mathbb{D}$, there exists a constant $\delta > 0$ such that

$$\langle X^*X(h \otimes g), h \otimes g \rangle = \left\langle M_{(F^\#)^*}^* M_{(F^\#)^*}^*(h \otimes g), h \otimes g \right\rangle = \|h\|^2 \langle F^*(w)F(w)g, g \rangle \geq \delta^2 \|g\|^2 \|h\|^2.$$

It follows that $h_{\mathcal{E}_{D_T}}(w) = F^*(w)F(w) = (\langle \tilde{\gamma}_j(w), \tilde{\gamma}_i(w) \rangle)_{i,j=1}^m \geq C_1 I_m$ for all $w \in \mathbb{D}$ and $C_1 := \delta^2$. \square

Theorem 2.4 in [16] and Theorem 2.3 in [24] use the curvature of the holomorphic complex bundles to characterize the similarity of n -hypercontraction. Next, we give the following brief and concise proof using Theorem 3.4:

Corollary 3.5. *Let $T \in \mathcal{B}_1(\mathbb{D})$ be an n -hypercontraction. Then T is similar to the backward shift operator S_n^* on \mathcal{M}_n if and only if there exists a bounded subharmonic function φ over \mathbb{D} such that*

$$\mathcal{K}_{S_n^*}(w) - \mathcal{K}_T(w) = \frac{\partial^2}{\partial w \partial \bar{w}} \varphi(w), \quad w \in \mathbb{D}.$$

Proof. Since $T \in \mathcal{B}_1(\mathbb{D})$ is an n -hypercontraction, by Lemma 3.2, there exists a Hermitian holomorphic line bundle \mathcal{E}_{D_T} over \mathbb{D} such that $\mathcal{E}_T \sim_u \mathcal{E}_{S_n^*} \otimes \mathcal{E}_{D_T}$.

If T is similar to S_n^* , there is a bounded invertible operator X such that

$$X(\mathcal{E}_{S_n^*}(w)) = (\mathcal{E}_{S_n^*} \otimes \mathcal{E}_{D_T})(w), \quad w \in \mathbb{D}.$$

Therefore, there exist constants C_1 , C_2 , and a non-vanishing holomorphic cross-section γ of \mathcal{E}_{D_T} satisfying

$$XK_n(\cdot, \bar{w}) = K_n(\cdot, \bar{w}) \otimes \gamma(w) \quad \text{and} \quad 0 < C_1 \leq \|\gamma(w)\|^2 \leq C_2, \quad w \in \mathbb{D}.$$

Hence,

$$\mathcal{K}_{S_n^*}(w) - \mathcal{K}_T(w) = \frac{\partial^2}{\partial w \partial \bar{w}} \log \|\gamma(w)\|^2 = \frac{\|\gamma(w)\|^2 \frac{\partial^2}{\partial w \partial \bar{w}} \|\gamma(w)\|^2 - \frac{\partial}{\partial w} \|\gamma(w)\|^2 \frac{\partial}{\partial \bar{w}} \|\gamma(w)\|^2}{\|\gamma(w)\|^4} \geq 0.$$

To complete the proof, one sets $\varphi(w) := \log \|\gamma(w)\|^2$ on \mathbb{D} .

For the other implication, we first note that since $\mathcal{E}_T \sim_u \mathcal{E}_{S_n^*} \otimes \mathcal{E}_{D_T}$ and $\mathcal{K}_T = \mathcal{K}_{S_n^*} + \mathcal{K}_{\mathcal{E}_{D_T}}$. Then

$$\frac{\partial^2}{\partial w \partial \bar{w}} \varphi(w) = \frac{\partial^2}{\partial w \partial \bar{w}} \log \|\gamma(w)\|^2$$

for bounded subharmonic function φ on \mathbb{D} and a non-vanishing holomorphic cross-section γ of \mathcal{E}_{D_T} .

Now, $u(w) := \log \left(\frac{\|\gamma(w)\|^2}{e^{\varphi(w)}} \right)^{\frac{1}{2}}$ is a real-valued harmonic function, and letting $\phi := e^{u+iv} \in \text{Hol}(\mathbb{D})$, where v is the harmonic conjugate of u , we see that

$$|\phi(w)| = e^{u(w)} = \left(\frac{\|\gamma(w)\|^2}{e^{\varphi(w)}} \right)^{\frac{1}{2}}.$$

Then $e^{\varphi(w)} = \left\| \frac{1}{\phi(w)} \gamma(w) \right\|^2$ and $\frac{1}{\phi} \gamma$ is a holomorphic cross-section of \mathcal{E}_{D_T} . Since φ is bounded on \mathbb{D} , there are constants C_1 and C_2 such that

$$0 < C_1 \leq e^{\varphi(w)} = \left\| \frac{1}{\phi(w)} \gamma(w) \right\|^2 \leq C_2, \quad w \in \mathbb{D},$$

and therefore by Theorem 3.4, $T \sim_s S_n^*$. □

Two Cowen-Douglas operators are unitarily equivalent if and only if their corresponding holomorphic complex bundles are equivalent as Hermitian holomorphic vector bundles. It is known from [40] that many holomorphic complex bundles with tensor products correspond to the Cowen-Douglas operator. Therefore, we use holomorphic complex bundles to describe the similar classification of the Cowen-Douglas operators and obtain the following proposition, whose proof is similar to Theorem 3.4, so the proof is omitted.

Proposition 3.6. *Let $T \in \mathcal{B}_m(\mathbb{D})$ and M_z^* be the backward shift operator on reproducing kernel Hilbert space \mathcal{H}_K with $\text{Mult}(\mathcal{H}_K) = H^\infty(\mathbb{D})$. If $\mathcal{E}_T \sim_u \mathcal{E}_{M_z^*} \otimes \mathcal{E}$ for some Hermitian holomorphic vector bundle \mathcal{E} over \mathbb{D} of rank m , then the following statements are equivalent:*

- (1) *The operator T is similar to the backward shift operator M_{z, \mathbb{C}^m}^* on $\mathcal{H}_{K, \mathbb{C}^m}$.*
- (2) *There exist constants C_1 and C_2 such that the metric $h_{\mathcal{E}}(w)$ of \mathcal{E} in some holomorphic frame satisfies $0 < C_1 I_m \leq h_{\mathcal{E}}(w) \leq C_2 I_m$ for any $w \in \mathbb{D}$.*
- (3) *There exists a holomorphic frame $\gamma = \{\gamma_1, \dots, \gamma_m\}$ of \mathcal{E} such that each $\|\gamma_i(w)\|$, $1 \leq i \leq m$, is uniformly bounded on \mathbb{D} , and there are constants A_1 and A_2 such that the metric $h_{\mathcal{E}}(w)$ under the holomorphic frame γ satisfies $0 < A_1 \leq \det h_{\mathcal{E}}(w) \leq A_2$ for any $w \in \mathbb{D}$.*

In Proposition 3.6, if the metric of the holomorphic complex bundle \mathcal{E} under a holomorphic frame is a polynomial for $|w|^2$, with each coefficient being a non-negative real number, then we can construct

many contractive operators similar to the backward shift operators on the Hardy space or the weighted Bergman spaces. This provides positive support for Theorem 1 in [11].

4. SIMILARITY ON WEIGHTED DIRICHLET SPACES

Shields, in [49], provided a necessary and sufficient condition for the similarity of backward shift operators on Hilbert space using weight sequences. From [28, 40], we know when a tensor product of Hermitian holomorphic vector bundles corresponds to a bounded linear operator. The goal of this section is to use the holomorphic jet bundle to describe the similarity of the backward shift operator on weighted Dirichlet spaces.

Let \mathcal{E} be a Hermitian holomorphic line bundle over \mathbb{D} and γ be a non-vanishing holomorphic cross-section of \mathcal{E} . For $k = 0, 1, 2, \dots$, we associate to \mathcal{E} a $(k+1)$ -dimensional Hermitian holomorphic vector bundle $\mathcal{J}_k(\mathcal{E})$ called the holomorphic k -jet bundle of \mathcal{E} . The work [13, 22, 35] are good references for k -jet bundles.

Suppose that for each w in \mathbb{D} , $\gamma(w), \gamma'(w), \gamma^{(2)}(w), \dots, \gamma^{(k)}(w)$ are independent. The k -jet bundle $\mathcal{J}_k(\mathcal{E})$ of the bundle \mathcal{E} has an associated holomorphic frame $\mathcal{J}_k(\gamma) = \{\gamma, \gamma', \gamma^{(2)}, \dots, \gamma^{(k)}\}$. The metric $h_{\mathcal{E}}(w) = \langle \gamma(w), \gamma(w) \rangle$ for \mathcal{E} induces a metric $\mathcal{J}_k(h_{\mathcal{E}})(w)$ for $\mathcal{J}_k(\mathcal{E})$ of the form

$$\mathcal{J}_k(h_{\mathcal{E}})(w) = \begin{pmatrix} h_{\mathcal{E}}(w) & \frac{\partial}{\partial w} h_{\mathcal{E}}(w) & \cdots & \frac{\partial^k}{\partial w^k} h_{\mathcal{E}}(w) \\ \frac{\partial}{\partial \bar{w}} h_{\mathcal{E}}(w) & \frac{\partial^2}{\partial w \partial \bar{w}} h_{\mathcal{E}}(w) & \cdots & \frac{\partial^{k+1}}{\partial w^k \partial \bar{w}} h_{\mathcal{E}}(w) \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^k}{\partial \bar{w}^k} h_{\mathcal{E}}(w) & \frac{\partial^{k+1}}{\partial w \partial \bar{w}^k} h_{\mathcal{E}}(w) & \cdots & \frac{\partial^{2k}}{\partial w^k \partial \bar{w}^k} h_{\mathcal{E}}(w) \end{pmatrix}.$$

Throughout the section, we denote by D_{α}^* and $D_{\alpha, E}^*$ the backward shift operators on weighted Dirichlet space \mathcal{D}_{α} and the space $\mathcal{D}_{\alpha, E}$, respectively, where $\alpha \in (0, 1]$ and E is a separable Hilbert space. The reproducing kernel of \mathcal{D}_{α} is $K_{\alpha}(z, w) = \sum_{i=0}^{\infty} \frac{z^i \bar{w}^i}{(i+1)^{\alpha}}$, $z, w \in \mathbb{D}$. In particular, D_1^* and \mathcal{D}_1 are also denoted as D^* and \mathcal{D} , respectively. Following the notations and ideas of Müller in [43], the Theorem 2.6, we use the following lemma to demonstrate how to construct a defect operator corresponding to operator $T \in \mathcal{L}(\mathcal{H})$ with $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{(s+1)^{\alpha}} \leq 1$ such that an operator T is unitary equivalent to the restriction of $D_{\alpha, E}^*$ on some invariant subspace.

Lemma 4.1. *Let $T \in \mathcal{L}(\mathcal{H})$ and $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{(s+1)^{\alpha}} \leq 1$ for some constant $\alpha \in (0, 1]$. Then there exist a Hilbert space E and an $D_{\alpha, E}^*$ -invariant subspace $\mathcal{N} \subseteq \mathcal{D}_{\alpha, E}$ such that $T \sim_u D_{\alpha, E}^*|_{\mathcal{N}}$.*

Proof. Let $b_0 = 1$, $c_0 = 1$, and for $k \geq 1$, $b_k = \frac{1}{(k+1)^{\alpha}}$, and $c_k = -\sum_{j=0}^{k-1} \frac{c_j}{(k-j+1)^{\alpha}}$, then for any fixed positive integer $l \geq 1$, $\sum_{i=0}^l b_i c_{l-i} = 0$. Obviously, $-b_1 = c_1 < 0$. Without losing generality, assume that $-b_i \leq c_i < 0$ for $1 \leq i \leq k-1$. Then by the assumption $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{(s+1)^{\alpha}} \leq 1$ and Lemma 2.1 in [43], we know that $0 \geq c_k \geq -b_k$ for every $k \geq 1$, which gives

$$\left\langle \left(\sum_{k=0}^{\infty} c_k (T^k)^* T^k \right) x, x \right\rangle \geq \|x\|^2 + \sum_{k=1}^{\infty} c_k \|T^k\|^2 \|x\|^2 \geq \|x\|^2 - \sum_{k=1}^{\infty} \frac{\|T^k\|^2}{(k+1)^{\alpha}} \|x\|^2 \geq 0, \quad x \in \mathcal{H}.$$

Hence, $\sum_{k=0}^{\infty} c_k (T^*)^k T^k$ is a positive operator.

Next, we define an operator $V : \mathcal{H} \rightarrow \mathcal{N}$ by

$$Vx = \sum_{n=0}^{\infty} \frac{z^n}{\|z^n\|_{\mathcal{D}_\alpha}^2} \otimes D_T T^n x, \quad x \in \mathcal{H},$$

where $\mathcal{N} = \overline{\text{ran } V}$ and $D_T = \left(\sum_{k=0}^{\infty} c_k (T^k)^* T^k \right)^{1/2}$. Since the orthonormal basis for the space \mathcal{D}_α is $\left\{ e_n = \frac{z^n}{\sqrt{(n+1)^\alpha}} \right\}_{n \geq 0}$, setting $E := \overline{\text{ran } D_T}$, we have for every $x \in \mathcal{H}$,

$$\|Vx\|^2 = \sum_{n=0}^{\infty} b_n \left\langle \left(\sum_{k=0}^{\infty} c_k (T^k)^* T^k \right) T^n x, T^n x \right\rangle = \sum_{l=0}^{\infty} \left(\|T^l x\|^2 \sum_{n+k=l} b_n c_k \right) = \|x\|^2$$

and

$$VTx = \sum_{n=0}^{\infty} \frac{z^n}{\|z^n\|_{\mathcal{D}_\alpha}^2} \otimes D_T T^{n+1} x = \sum_{n=1}^{\infty} \frac{z^{n-1}}{\|z^{n-1}\|_{\mathcal{D}_\alpha}^2} \otimes D_T T^n x = D_{\alpha, E}^* \left(\sum_{n=0}^{\infty} \frac{z^n}{\|z^n\|_{\mathcal{D}_\alpha}^2} \otimes D_T T^n x \right) = D_{\alpha, E}^* Vx.$$

Hence, $T \sim_u D_{\alpha, E}^*|_{\mathcal{N}}$ for the $D_{\alpha, E}^*$ -invariant subspace $\mathcal{N} \subset \mathcal{D}_{\alpha, E}$. \square

Similar to the definition of the defect operator of n -hypercontraction in Section 2, to prevent confusion, we still denote $D_T = \left(\sum_{k=0}^{\infty} c_k (T^k)^* T^k \right)^{1/2}$ as the *defect operator* of the operator T with $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{(s+1)^\alpha} \leq 1$. Then we have the following lemma from Lemma 4.1, whose proof is omitted.

Lemma 4.2. *Let $T \in \mathcal{B}_1(\mathbb{D})$ be an operator with $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{(s+1)^\alpha} \leq 1$ for some constant $\alpha \in (0, 1]$, and let γ be a non-vanishing holomorphic cross-section of \mathcal{E}_T . Then $\mathcal{E}_{D_T} = \{(w, x) \in \mathbb{D} \times \mathcal{E}_{D_T}(w) : x \in \mathcal{E}_{D_T}(w)\}$ is a Hermitian holomorphic line bundle over \mathbb{D} with the natural projection $\pi, \pi(w, x) = w$, where $\mathcal{E}_{D_T}(w) = \vee \{D_T \gamma(w)\}$.*

Moreover, $\mathcal{E}_T \sim_u \mathcal{E}_{D_\alpha}^ \otimes \mathcal{E}_{D_T}$ and $D_T \gamma$ is a non-vanishing holomorphic cross-section of \mathcal{E}_{D_T} .*

The ratio of the metrics on holomorphic line bundles, as derived in the previous section, serves as a similarity invariant for the backward shift operators on the Hardy space and the weighted Bergman space. However, whether this extends to weighted Dirichlet remains unclear. In what follows, we give a sufficient condition for certain bounded linear operators to be similar to backward shift operator D_α^* on weighted Dirichlet space \mathcal{D}_α , derived through the holomorphic jet bundle.

Theorem 4.3. *Let $T \in \mathcal{B}_1(\mathbb{D})$ and $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{(s+1)^\alpha} \leq 1$ for some constant $\alpha \in (0, 1]$. If there exist constants $c_2 > c_1 > 0$ satisfy*

$$c_1 \leq h_{\mathcal{E}_{D_T}}(w) < \text{trace } \mathcal{J}_1(h_{\mathcal{E}_{D_T}})(w) \leq c_2, \quad w \in \mathbb{D},$$

for the metric $h_{\mathcal{E}_{D_T}}$ of \mathcal{E}_{D_T} and $\mathcal{E}_{D_T}(w) \subset \mathbb{C}^n$ for some $n > 0$, then $T \sim_s D_\alpha^$.*

Proof. Since $T \in \mathcal{B}_1(\mathbb{D})$ satisfies $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{(s+1)^\alpha} \leq 1$ for some constant $\alpha \in (0, 1]$, by Lemma 4.1 and Lemma 4.2, we obtain that $\mathcal{E}_T \sim_u \mathcal{E}_{D_\alpha}^* \otimes \mathcal{E}_{D_T}$ for the holomorphic line bundle \mathcal{E}_{D_T} induced by the defect operator D_T .

Now, a holomorphic function $f \in \mathcal{D}_\alpha$ can be represented as a power series

$$f(z) = \sum_{i=0}^{\infty} a_i z^i, \quad \text{where} \quad \|f\|_{\mathcal{D}_\alpha}^2 = \sum_{i=0}^{\infty} (i+1)^\alpha |a_i|^2 < \infty.$$

Since $\|zf\|_{\mathcal{D}_\alpha}^2 = \sum_{i=0}^{\infty} (i+2)^\alpha |a_i|^2 \leq 2 \sum_{i=0}^{\infty} (i+1)^\alpha |a_i|^2 < \infty$, $z \in \text{Mult}(\mathcal{D}_\alpha)$. From [41], $\phi \in H^\infty(\mathbb{D})$ is in $\text{Mult}(\mathcal{D}_\alpha)$ if and only if there is a constant $d > 0$ such that

$$\int_{\mathbb{D}} |f(z)|^2 |\phi'(z)|^2 (1-|z|^2)^{1-\alpha} dA(z) \leq d \|f\|_{\mathcal{D}_\alpha}^2, \quad f \in \mathcal{D}_\alpha,$$

where dA is the normalized area measure. Setting $\phi(z) = z$, we have

$$(4.1) \quad \int_{\mathbb{D}} |f(z)|^2 (1-|z|^2)^{1-\alpha} dA(z) \leq d \|f\|_{\mathcal{D}_\alpha}^2, \quad f \in \mathcal{D}_\alpha.$$

By assumption, the metric $\mathcal{J}_1(h_{\mathcal{E}_{D_T}})$ of the 1-jet bundle $\mathcal{J}_1(\mathcal{E}_{D_T})$ satisfies

$$(4.2) \quad c_1 \leq h_{\mathcal{E}_{D_T}}(w) < \text{trace } \mathcal{J}_1(h_{\mathcal{E}_{D_T}})(w) \leq c_2, \quad w \in \mathbb{D},$$

for some constants $c_2 > c_1 > 0$, where $h_{\mathcal{E}_{D_T}}$ denotes the metric obtained from a non-vanishing holomorphic cross-section γ of \mathcal{E}_{D_T} . Furthermore, since $\mathcal{E}_{D_T}(w) \subset \mathbb{C}^n$ for some $n > 0$, there is an orthonormal basis $\{e_i\}_{i=1}^n$ of the Hilbert space $E = \bigvee_{w \in \mathbb{D}} \mathcal{E}_{D_T}(w)$ such that $\gamma(w) = \sum_{i=1}^n \gamma_i(w) e_i$. From

(4.2),

(4.3)

$$c_1 \leq h_{\mathcal{E}_{D_T}}(w) = \sum_{i=1}^n |\gamma_i(w)|^2 < \sum_{i=1}^n |\gamma_i(w)|^2 + \sum_{i=1}^n |\gamma_i'(w)|^2 = h_{\mathcal{E}_{D_T}}(w) + \frac{\partial^2}{\partial w \partial \bar{w}} h_{\mathcal{E}_{D_T}}(w) \leq 2c_2, \quad w \in \mathbb{D}.$$

By (4.1) and (4.3), we have for every $f \in \mathcal{D}_\alpha$,

$$(4.4) \quad \int_{\mathbb{D}} |f(z)|^2 |\gamma_i'(z)|^2 (1-|z|^2)^{1-\alpha} dA(z) \leq 2c_2 \int_{\mathbb{D}} |f(z)|^2 (1-|z|^2)^{1-\alpha} dA(z) \leq 2dc_2 \|f\|_{\mathcal{D}_\alpha}^2.$$

This shows that $\{\gamma_1, \dots, \gamma_n\} \subset \text{Mult}(\mathcal{D}_\alpha)$ and that $\|M_{\gamma_i}\| = \|\gamma_i\|_{\text{Mult}(\mathcal{D}_\alpha)} < \infty$ for $1 \leq i \leq n$.

We next define an operator-valued function $F : \mathbb{D} \rightarrow \mathcal{L}(\mathbb{C}, E)$ by

$$F(w)(1) = \gamma(w).$$

Then for every $f \in \mathcal{D}_\alpha$, $M_F f = f\gamma$ and

$$\|M_F\|^2 = \sup_{\|f\|_{\mathcal{D}_\alpha} \neq 0} \frac{\|M_F f\|_{\mathcal{D}_\alpha, E}^2}{\|f\|_{\mathcal{D}_\alpha}^2} = \sup_{\|f\|_{\mathcal{D}_\alpha} \neq 0} \frac{\sum_{i=1}^n \|f\gamma_i\|_{\mathcal{D}_\alpha}^2}{\|f\|_{\mathcal{D}_\alpha}^2} \leq \sum_{i=1}^n \|M_{\gamma_i}\|^2 < \infty,$$

that is, $F \in \text{Mult}(\mathcal{D}_\alpha, \mathcal{D}_\alpha, E)$. For $\gamma_i \in H^\infty(\mathbb{D})$, $1 \leq i \leq n$, let $\gamma_i(z) := \sum_{j=0}^{\infty} a_{i,j} z^j$ and $\tilde{\gamma}_i(z) := \sum_{j=0}^{\infty} \overline{a_{i,j}} z^j$.

Then using (4.3) and (4.4), we deduce that for every $w \in \mathbb{D}$, $|\tilde{\gamma}_i(w)|^2, |\tilde{\gamma}_i'(w)|^2 \leq 2c_2$ and

$$\int_{\mathbb{D}} |f(z)|^2 |\tilde{\gamma}_i'(z)|^2 (1-|z|^2)^{1-\alpha} dA(z) \leq 2c_2 \int_{\mathbb{D}} |f(z)|^2 (1-|z|^2)^{1-\alpha} dA(z) \leq 2dc_2 \|f\|_{\mathcal{D}_\alpha}^2, \quad f \in \mathcal{D}_\alpha, 1 \leq i \leq n.$$

Thus $\{\tilde{\gamma}_1, \dots, \tilde{\gamma}_n\} \subset \text{Mult}(\mathcal{D}_\alpha)$ and $(F^\#)^* \in \text{Mult}(\mathcal{D}_{\alpha,E}, \mathcal{D}_\alpha)$ for $F^\#(w) = F(\bar{w})$. Since $1 \in \text{Mult}(\mathcal{D}_\alpha)$, $\gamma_i \in \text{Mult}(\mathcal{D}_\alpha)$, and $\sum_{i=1}^n |\gamma_i(w)|^2 \geq c_1 > 0$ for every $w \in \mathbb{D}$, by Theorem 2.4, there are functions $\{g_1, \dots, g_n\} \subset \text{Mult}(\mathcal{D}_\alpha)$ such that $\sum_{i=1}^n g_i(w)\gamma_i(w) = 1$ for every $w \in \mathbb{D}$.

We now define a function $G : \mathbb{D} \rightarrow \mathcal{L}(E, \mathbb{C})$ by

$$G(w)h := \sum_{i=1}^n g_i(w)h_i,$$

where $h = \sum_{i=1}^n h_i e_i \in E$. It is easy to see that $G \in \text{Mult}(\mathcal{D}_{\alpha,E}, \mathcal{D}_\alpha)$ and $(G^\#)^* \in \text{Mult}(\mathcal{D}_\alpha, \mathcal{D}_{\alpha,E})$ for $G^\#(w) = G(\bar{w})$. Hence, by Theorem 2.1, for the reproducing kernel $K_\alpha(z, w) = \sum_{n=0}^{\infty} \frac{z^n \bar{w}^n}{(n+1)^\alpha}$ of \mathcal{D}_α ,

$$M_{(F^\#)^*}^*(K_\alpha(\cdot, \bar{w}) \otimes 1) = K_\alpha(\cdot, \bar{w}) \otimes F^\#(\bar{w})(1) = K_\alpha(\cdot, \bar{w}) \otimes F(w)(1) = K_\alpha(\cdot, \bar{w}) \otimes \gamma(w),$$

and therefore,

$$M_{(G^\#)^*}^* M_{(F^\#)^*}^*(K_\alpha(\cdot, \bar{w}) \otimes 1) = M_{(G^\#)^*}^*(K_\alpha(\cdot, \bar{w}) \otimes F(w)(1)) = K_\alpha(\cdot, \bar{w}) \otimes \sum_{i=1}^n g_i(w)\gamma_i(w) = K_\alpha(\cdot, \bar{w}) \otimes 1,$$

so that using the fact that $\mathcal{D}_\alpha = \bigvee_{w \in \mathbb{D}} K_\alpha(\cdot, \bar{w})$, we conclude that $M_{(G^\#)^*}^* M_{(F^\#)^*}^* \equiv I_{\mathcal{D}_\alpha}$. Moreover, if we let $\mathcal{N} := \bigvee_{w \in \mathbb{D}} \{K_\alpha(\cdot, \bar{w}) \otimes \gamma(w)\}$, then since the operator $M_{(F^\#)^*}^*$ is left-invertible, $\text{ran } M_{(F^\#)^*}^* = \mathcal{N}$. Hence, the operator $X := M_{(F^\#)^*}^* : \mathcal{D}_\alpha \rightarrow \mathcal{N}$ is a bounded invertible operator satisfying

$$X(\mathcal{E}_{D_\alpha^*}(w)) = (\mathcal{E}_{D_\alpha^*} \otimes \mathcal{E}_{D_T})(w), \quad w \in \mathbb{D}.$$

Finally, by combining $\mathcal{E}_T \sim_u \mathcal{E}_{D_\alpha^*} \otimes \mathcal{E}_{D_T}$, we conclude that $T \sim_s D_\alpha^*$. \square

In Theorem 4.3, the ratio of the metrics of holomorphic line bundles \mathcal{E}_T and $\mathcal{E}_{D_\alpha^*}$ is the metric of the holomorphic line bundle \mathcal{E}_{D_T} induced by the defect operator D_T , that is, $h_{\mathcal{E}_{D_T}} = \frac{h_{\mathcal{E}_T}}{h_{D_\alpha^*}}$. Letting \mathcal{H}_K is a Hilbert space with diagonal reproducing kernel $K(z, w) = \sum_{m,n=0}^{\infty} a_{m,n} z^m \bar{w}^n$, which means that for all $n \neq m$, $a_{m,n} = 0$. From this, the similarity classification of holomorphic complex bundles and the similarity of backward shift operators on non-diagonal reproducing kernel Hilbert space can be obtained.

Corollary 4.4. *Let \mathcal{E}_1 and \mathcal{E}_2 be Hermitian holomorphic line bundle over \mathbb{D} and $\mathcal{E}_1 \sim_u \mathcal{E}_{D_\alpha^*} \otimes \mathcal{E}_2$. If there exist constants $c_2 > c_1 > 0$ and a metric $h_{\mathcal{E}_2}$ of \mathcal{E}_2 satisfy*

$$c_1 \leq h_{\mathcal{E}_2}(w) < \text{trace } \mathcal{J}_1(h_{\mathcal{E}_2})(w) \leq c_2$$

and $\mathcal{E}_2(w) \subset \mathbb{C}^n$ for some positive integer $n > 0$, then $\mathcal{E}_1 \sim_s \mathcal{E}_{D_\alpha^*}$.

Example 4.5. Let $T \in \mathcal{B}_1(\mathbb{D})$ and $\mathcal{E}_T \sim_u \mathcal{E}_{D_\alpha^*} \otimes \mathcal{E}$ for some Hermitian holomorphic line bundle \mathcal{E} over \mathbb{D} . If the metric $h_{\mathcal{E}}$ of \mathcal{E} satisfies $h_{\mathcal{E}}(w) = 1 + \sum_{i=0}^n |\phi_i(w)|^2$ for polynomials ϕ_1, \dots, ϕ_n , and positive integer n , then $\mathcal{E}_T \sim_s \mathcal{E}_{D_\alpha^*}$, that is, $T \sim_s D_\alpha^*$.

Given the discussion in this section, we use holomorphic line bundles to provide a necessary and sufficient for certain operators to be similar to backward shift operators on weighted Dirichlet spaces.

Proposition 4.6. *For an operator $T \in \mathcal{B}_1(\mathbb{D})$, if there exists a Hermitian holomorphic line bundle \mathcal{E} over \mathbb{D} such that $\mathcal{E}_T \sim_u \mathcal{E}_{D_\alpha^*} \otimes \mathcal{E}$, then $T \sim_s D_\alpha^*$ if and only if $\mathcal{E}_{D_\alpha^*} \otimes \mathcal{E} \sim_s \mathcal{E}_{D_\alpha^*}$.*

If the reproducing kernel of a Hilbert space \mathcal{H}_K is of the form $K(z, w) = \sum_{m,n=0}^{\infty} a_{m,n} z^m \bar{w}^n$, where, for some $n \neq m$, $a_{m,n} \neq 0$, then Theorem 2.7 cannot be used to determine similarity. The following example is a good illustration of how one can use Theorem 4.3 to conclude similarity:

Example 4.7. Let M_z^* be the backward shift operator on Hilbert space \mathcal{H}_K with reproducing kernel

$$K(z, w) = 2 + z + \bar{w} + 2z\bar{w} + (z + \bar{w}) \sum_{n=1}^{\infty} \frac{2n+1}{n(n+1)} z^n \bar{w}^n \\ + (z^2 + \bar{w}^2) \sum_{n=0}^{\infty} \frac{z^n \bar{w}^n}{n+1} + \sum_{n=0}^{\infty} \frac{4n^2 + 15n + 13}{(n+1)(n+2)(n+3)} z^{n+2} \bar{w}^{n+2}, \quad z, w \in \mathbb{D}.$$

Then $M_z^* \sim_s D^*$.

Proof. Since the reproducing kernel of Dirichlet space \mathcal{D} is $K_1(z, w) = \sum_{i=0}^{\infty} \frac{z^i \bar{w}^i}{i+1}$, $z, w \in \mathbb{D}$, we obtain that $K(z, w) = K_1(z, w) \cdot [1 + (1 + z + z^2)(1 + \bar{w} + \bar{w}^2)]$. Then there exists a Hermitian holomorphic line bundle \mathcal{E} over \mathbb{D} such that $\mathcal{E}_{M_z^*} = \mathcal{E}_{D^*} \otimes \mathcal{E}$, and a non-vanishing holomorphic cross-section γ of \mathcal{E} satisfies $\|\gamma(w)\|^2 = 1 + |1 + w + w^2|^2$. Therefore, the metric $h_{\mathcal{E}}(w) = \|\gamma(w)\|^2$ of \mathcal{E} satisfies

$$1 \leq h_{\mathcal{E}}(w) < \text{trace} \mathcal{J}_1(h_{\mathcal{E}})(w) \leq 19, \quad w \in \mathbb{D}.$$

Hence the conclusion can be derived from Theorem 4.3. \square

In Theorem 4.3, the condition that $\mathcal{E}_{D_T}(w) \subset \mathbb{C}^n$ for some integer $n > 0$ was necessary. A natural question is whether the result also exists when $n = \infty$. The following example answers the question in the affirmative:

Example 4.8. Let M_z^* be the backward shift operator on Hilbert space \mathcal{H}_K with reproducing kernel

$$K(z, w) = \sum_{n=0}^{\infty} \left(\sum_{i=0}^n \frac{1}{(i+1)(n-i+1)^2} \right) z^n \bar{w}^n, \quad z, w \in \mathbb{D}.$$

Then there is a Hermitian holomorphic line bundle $\mathcal{E} = \{(w, \gamma) : w \in \mathbb{D}, \gamma(w) \in \mathcal{E}(w)\}$ with the natural projection $\pi, \pi(w, \gamma) = w$, such that $\mathcal{E}_{M_z^*} = \mathcal{E}_{D^*} \otimes \mathcal{E}$ and $\|\gamma(w)\|^2 = \sum_{n=0}^{\infty} \frac{1}{(n+1)^2} |w|^{2n}$, where $\mathcal{E}(w) = \bigvee \gamma(w)$.

From the relationship between the weight sequence and the reproducing kernel of Hilbert space, we obtain that the weight sequence of M_z^* is $\left\{ \alpha_n = \sqrt{\frac{a_{n-1}}{a_n}} \right\}_{n \geq 1}$, where $a_n = \sum_{i=0}^n \frac{1}{(i+1)(n-i+1)^2}$. Since the weight sequence of D^* is $\left\{ \beta_n = \sqrt{\frac{n+1}{n}} \right\}_{n \geq 1}$, we have that $\prod_{i=1}^n \beta_i / \prod_{i=1}^n \alpha_i = \sqrt{(n+1)a_n} \geq 1$.

Now, to prove from Theorem 2.7 that M_z^* is similar to D^* , we only need to prove that $\lim_{n \rightarrow \infty} (n+1)a_n = c$ for some constant c .

From $\lim_{n \rightarrow \infty} \left[\sum_{k=1}^n \frac{1}{k} - \ln n \right] = \Upsilon$ for the Euler-Mascheroni constant Υ , we obtain that

$$\lim_{n \rightarrow \infty} \frac{n+1}{(n+2)^2} \sum_{k=1}^{n+1} \frac{1}{k} = 0, \quad \lim_{n \rightarrow \infty} \frac{n+1}{(n+2)^2} \sum_{k=1}^{n+1} \frac{1}{n+2-k} = 0, \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{n+1}{n+2} \sum_{k=1}^{n+1} \frac{1}{(n+2-k)^2} = \frac{\pi^2}{6}.$$

Therefore,

$$\begin{aligned} \lim_{n \rightarrow \infty} (n+1)a_n &= \lim_{n \rightarrow \infty} (n+1) \left(\sum_{k=1}^{n+1} \frac{1}{k(n+2-k)^2} \right) \\ &= \lim_{n \rightarrow \infty} (n+1) \sum_{k=1}^{n+1} \left[\frac{1}{(n+2)^2} \frac{1}{k} + \frac{1}{(n+2)^2} \frac{1}{n+2-k} + \frac{1}{n+2} \frac{1}{(n+2-k)^2} \right] \\ &= \frac{\pi^2}{6}. \end{aligned}$$

Next is another major theorem in this section, as Theorem 4.3, we provide another criterion for the similarity of backward shift operators in the weighted Dirichlet space when $n = \infty$.

Theorem 4.9. *Let $T \in \mathcal{B}_1(\mathbb{D})$ and $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{(s+1)^\alpha} \leq 1$ for some constant $\alpha \in (0, 1]$. If there is a non-vanishing holomorphic cross-section γ of \mathcal{E}_{D_T} such that for some $\delta > 0$,*

$$(4.5) \quad \delta \leq \|\gamma(w)\|_E^2 \leq \sum_{i=1}^{\infty} \|\gamma_i\|_{\text{Mult}(\mathcal{D}_\alpha)}^2 < \infty, \quad w \in \mathbb{D},$$

where $\gamma(w) = \sum_{i=1}^{\infty} \gamma_i(w)e_i$, $E = \bigvee_{w \in \mathbb{D}} \gamma(w)$, and $\{e_i\}_{i=1}^{\infty}$ is an orthonormal basis of E , then $T \sim_s D_\alpha^*$.

Proof. Denoting by $K_\alpha(z, w) = \sum_{i=0}^{\infty} \frac{z^i \bar{w}^i}{(i+1)^\alpha}$, $z, w \in \mathbb{D}$, the reproducing kernel for the space \mathcal{D}_α , we have for every $f \in \mathcal{D}_\alpha$ and $\phi \in \text{Mult}(\mathcal{D}_\alpha)$,

$$\langle f, M_\phi^* K_\alpha(\cdot, w) \rangle = \langle M_\phi f, K_\alpha(\cdot, w) \rangle = \phi(w) f(w) = \langle f, \overline{\phi(w)} K_\alpha(\cdot, w) \rangle,$$

and therefore, $\|\phi\|_\infty \leq \|M_\phi\| = \|\phi\|_{\text{Mult}(\mathcal{D}_\alpha)}$. Let γ be a non-vanishing holomorphic cross-section of \mathcal{E}_{D_T} satisfying (4.5). If we assume without loss of generality that $\sum_{i=1}^{\infty} \|\gamma_i\|_{\text{Mult}(\mathcal{D}_\alpha)}^2 \leq 1$, then

$$\|\gamma(w)\|_E^2 = \sum_{i=1}^{\infty} |\gamma_i(w)|^2 \leq \sum_{i=1}^{\infty} \|\gamma_i\|_\infty^2 \leq \sum_{i=1}^{\infty} \|\gamma_i\|_{\text{Mult}(\mathcal{D}_\alpha)}^2 \leq 1, \quad w \in \mathbb{D}.$$

We next define an operator-valued function $F : \mathbb{D} \rightarrow \mathcal{L}(\mathbb{C}, E)$ by

$$F(w)(1) = \gamma(w).$$

Then $F \in H_{\mathbb{C} \rightarrow E}^\infty(\mathbb{D})$ and the multiplication operator $M_F : \mathcal{D}_\alpha \rightarrow \mathcal{D}_{\alpha, E}$ satisfies

$$(M_F f)(z) = F(z) f(z) = f(z) \otimes \gamma(z).$$

Moreover, for every $f \in \mathcal{D}_\alpha$,

$$\|f \otimes \gamma\|_{\mathcal{D}_\alpha \otimes E}^2 = \sum_{i=1}^{\infty} \|f \gamma_i\|_{\mathcal{D}_\alpha}^2 = \sum_{i=1}^{\infty} \|M_{\gamma_i} f\|_{\mathcal{D}_\alpha}^2 \leq \|f\|_{\mathcal{D}_\alpha}^2 \sum_{i=1}^{\infty} \|M_{\gamma_i}\|^2 = \|f\|_{\mathcal{D}_\alpha}^2 \sum_{i=1}^{\infty} \|\gamma_i\|_{\text{Mult}(\mathcal{D}_\alpha)}^2 \leq \|f\|_{\mathcal{D}_\alpha}^2.$$

It follows that $F \in \text{Mult}(\mathcal{D}_\alpha, \mathcal{D}_{\alpha, E})$.

Now let $F^\#(w) := F(\bar{w})$ and hence, $(F^\#)^* \in \text{Mult}(\mathcal{D}_{\alpha, E}, \mathcal{D}_\alpha)$. If we denote by M_γ^C the column operator from \mathcal{D}_α to $\bigoplus_1^\infty \mathcal{D}_\alpha$ defined by

$$M_\gamma^C(f) = (\gamma_1 f, \gamma_2 f, \dots)^T = f \otimes \gamma, \quad f \in \mathcal{D}_\alpha,$$

then $M_F = M_\gamma^C$ and

$$\|M_\gamma^C\| = \|M_F\| = \sup_{\|f\|_{\mathcal{D}_\alpha} \neq 0} \frac{\|M_F f\|_{\mathcal{D}_{\alpha,E}}}{\|f\|_{\mathcal{D}_\alpha}} = \sup_{\|f\|_{\mathcal{D}_\alpha} \neq 0} \frac{\|f \otimes \gamma\|_{\mathcal{D}_{\alpha,E}}}{\|f\|_{\mathcal{D}_\alpha}} \leq 1.$$

Since $\|\gamma(w)\|_E^2 = \sum_{i=1}^{\infty} |\gamma_i(w)|^2 \geq \delta > 0$, by Theorem 2.4, we obtain functions $\{g_i\}_{i=1}^{\infty} \subset \text{Mult}(\mathcal{D}_\alpha)$ such that $\sum_{i=1}^{\infty} g_i(w)\gamma_i(w) = 1$ for any $w \in \mathbb{D}$. Moreover, the column operator $M_G^C : \mathcal{D}_\alpha \rightarrow \bigoplus_1^{\infty} \mathcal{D}_\alpha$ defined by

$$M_G^C(f) = (g_1 f, g_2 f, \dots)^T, \quad f \in \mathcal{D}_\alpha,$$

is bounded and again from Theorem 2.3, we know that the row operator $M_G^R : \bigoplus_1^{\infty} \mathcal{D}_\alpha \rightarrow \mathcal{D}_\alpha$ is bounded as well.

Finally, we define an operator-valued function $\tilde{G} : \mathbb{D} \rightarrow \mathcal{L}(E, \mathbb{C})$ by

$$\tilde{G}(w)(h) = \sum_{i=1}^{\infty} g_i(w)h_i, \quad h = \sum_{i=1}^{\infty} h_i e_i \in E.$$

Then the operator $M_{\tilde{G}} : \mathcal{D}_{\alpha,E} \rightarrow \mathcal{D}_\alpha$ defined by

$$M_{\tilde{G}} f = \tilde{G} f, \quad f \in \mathcal{D}_{\alpha,E},$$

is such that $\|M_{\tilde{G}}\| \leq \|M_G^R\| < \infty$, so that $M_{\tilde{G}}$ is bounded and $\tilde{G} \in \text{Mult}(\mathcal{D}_{\alpha,E}, \mathcal{D}_\alpha)$. Since M_G^C is bounded, $(\tilde{G}^\#)^* \in \text{Mult}(\mathcal{D}_\alpha, \mathcal{D}_{\alpha,E})$, where $\tilde{G}^\#(w) = \tilde{G}(\bar{w})$. Proceeding similarly as in the proof of Theorem 4.3, we obtain an invertible operator X such that $X(\mathcal{E}_{D_\alpha^*}(w)) = (\mathcal{E}_{D_\alpha^*} \otimes \mathcal{E}_{D_T})(w)$, and since $\mathcal{E}_T \sim_u \mathcal{E}_{D_\alpha^*} \otimes \mathcal{E}_{D_T}$, $T \sim_s D_\alpha^*$ as claimed. \square

The condition $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{(s+1)^\alpha} \leq 1$ in Theorem 4.9 ensures that the Hermitian holomorphic line bundle \mathcal{E}_T acquires a tensor product structure. The natural question arises, what type of operator corresponds to a Hermitian holomorphic complex bundle with a tensor product structure in the unitarily equivalent sense? We will give a precise answer, distinct from the condition $\sum_{s=1}^{\infty} \frac{\|T^s\|^2}{s+1} \leq 1$, through the following example.

Example 4.10. Let M_z^* be the backward shift operator on the Hilbert space \mathcal{H}_K with reproducing kernel $K(z, w) = \sum_{n=0}^{\infty} a_n z^n \bar{w}^n$, $z, w \in \mathbb{D}$, where $a_n = \sum_{\substack{i+j^5=n \\ i, j \in \mathbb{N}}} \frac{1}{(i+1)(j+1)^8}$ for $n \in \mathbb{N}$. We have that

$\{\sqrt{a_n} z^n\}_{n \geq 0}$ is an orthonormal basis of \mathcal{H}_K and $\left\{\sqrt{\frac{a_{n-1}}{a_n}}\right\}_{n \geq 1}$ is the weight sequence of M_z^* . After calculations, we obtain that

$$(4.6) \quad K(z, w) = \sum_{n=0}^{\infty} \left(\sum_{\substack{i+j^5=n \\ i, j \in \mathbb{N}}} \frac{1}{(i+1)(j+1)^8} \right) z^n \bar{w}^n = \left(\sum_{i=0}^{\infty} \frac{z^i \bar{w}^i}{i+1} \right) \left(\sum_{j=0}^{\infty} \frac{z^{j^5} \bar{w}^{j^5}}{(j+1)^8} \right).$$

If M_z^* satisfies $\sum_{s=1}^{\infty} \frac{\|M_z^{*s}\|^2}{s+1} \leq 1$, then

$$\left\langle \left(I - \sum_{s=1}^{\infty} \frac{M_z^s M_z^{*s}}{s+1} \right) x, x \right\rangle = \|x\|^2 - \sum_{s=1}^{\infty} \frac{\|M_z^{*s} x\|^2}{s+1} \geq \left(1 - \sum_{s=1}^{\infty} \frac{\|M_z^{*s}\|^2}{s+1} \right) \|x\|^2 \geq 0, \quad x \in \mathcal{H}_K.$$

It follows that $\sum_{s=1}^{\infty} \frac{M_z^s M_z^{*s}}{s+1} \leq I$, and then $a_n - \frac{1}{2}a_{n-1} - \dots - \frac{1}{n+1}a_0 \geq 0$ for any positive integer n . However, for $a_2 = \frac{515}{1536}$, $a_1 = \frac{129}{256}$ and $a_0 = 1$, we have that $a_2 - \frac{1}{2}a_1 - \frac{1}{3}a_0 = \frac{515}{1536} - \frac{1}{2} \times \frac{129}{256} - \frac{1}{3} = -\frac{1}{4} < 0$. This means that M_z^* does not satisfy $\sum_{s=1}^{\infty} \frac{\|M_z^{*s}\|^2}{s+1} \leq 1$.

Setting \mathcal{E} is a Hermitian holomorphic line bundle over \mathbb{D} with a non-vanishing holomorphic cross-section γ satisfying $\|\gamma(w)\|^2 = \sum_{j=0}^{\infty} \frac{|w|^{2j^5}}{(j+1)^8}$, then from (4.6), $\mathcal{E}_{M_z^*} \sim_u \mathcal{E}_{D^*} \otimes \mathcal{E}$ for the backward shift operator D^* on the Dirichlet space \mathcal{D} . Let $E = \bigvee_{w \in \mathbb{D}} \gamma(w)$ and $\{e_k\}_{k=0}^{\infty}$ be an orthonormal basis of E , then $\gamma(w) = \sum_{k=0}^{\infty} \gamma_k(w) e_k$ for $\gamma_k(w) = \frac{w^{k^5}}{(k+1)^4}$. For any holomorphic function $f \in \mathcal{D}$ with Taylor expansion $f(z) = \sum_{n=0}^{\infty} \hat{f}(n) z^n$, there is $\|f\|_{\mathcal{D}}^2 = \sum_{n=0}^{\infty} (n+1) |\hat{f}(n)|^2 < \infty$. Therefore, for $\gamma_k f$, $k \geq 0$, we have that $\gamma_k(z) f(z) = \sum_{n=0}^{\infty} \frac{\hat{f}(n)}{(k+1)^4} z^{n+k^5}$, and then

$$\begin{aligned} \|\gamma_k f\|_{\mathcal{D}}^2 &= \sum_{n=0}^{\infty} (n + k^5 + 1) \left| \frac{\hat{f}(n)}{(k+1)^4} \right|^2 \\ &= \frac{1}{(k+1)^8} \left[\sum_{n=0}^{\infty} (n+1) |\hat{f}(n)|^2 + \sum_{n=0}^{\infty} k^5 |\hat{f}(n)|^2 \right] \\ &\leq \left[\frac{1}{(k+1)^8} + \frac{1}{(k+1)^5} \right] \|f\|_{\mathcal{D}}^2. \end{aligned}$$

It follows that $\|\gamma_k\|_{\text{Mult}(\mathcal{D})}^2 \leq \frac{1}{(k+1)^8} + \frac{1}{(k+1)^5} \leq \frac{2}{(k+1)^2}$, Hence,

$$1 \leq \|\gamma(w)\|_E^2 = \sum_{k=0}^{\infty} |\gamma_k(w)|^2 \leq \sum_{k=0}^{\infty} \|\gamma_k\|_{\infty}^2 \leq \sum_{k=0}^{\infty} \|\gamma_k\|_{\text{Mult}(\mathcal{D})}^2 \leq \frac{\pi^2}{3}, \quad w \in \mathbb{D}.$$

Using the proof of Theorem 4.9, we obtain that $M_z^* \sim_s D^*$.

From Example 4.10, we demonstrate that the weighted backward shift operator M_z^* does not satisfy the condition $\sum_{s=1}^{\infty} \frac{\|M_z^{*s}\|^2}{s+1} \leq 1$. Furthermore, using Shields' result, Theorem 2.7, it is not straightforward to verify that M_z^* is similar to D^* . Now, let's consider an example of a Hilbert space with a non-diagonal reproducing kernel. The proof, being similar to that of Example 4.10, is omitted here.

Example 4.11. Let M_z^* be the backward shift operator on Hilbert space \mathcal{H}_K with reproducing kernel

$$K(z, w) = 1 + \frac{5}{8} z \bar{w} + (z + \bar{w}) \sum_{n=1}^{\infty} \frac{1}{8n} z^n \bar{w}^n + \sum_{n=2}^{\infty} \left(\sum_{i=1}^{n+1} \frac{1}{i(n+2-i)^3} + \frac{1}{8(n-1)} \right) z^n \bar{w}^n, \quad z, w \in \mathbb{D}.$$

Then $M_z^* \sim_s D^*$.

We close this section with a proposition that shows that if a Hermitian holomorphic complex bundle with a tensor product satisfies the given conditions, there exists a corresponding bounded linear operator in $\mathcal{B}_1(\mathbb{D})$. This establishes the operator realization of the tensor product of holomorphic complex bundles.

Proposition 4.12. *Let \mathcal{E} be a Hermitian holomorphic line bundle over \mathbb{D} , and let γ be a non-vanishing holomorphic cross-section of \mathcal{E} . Suppose that $E = \bigvee_{w \in \mathbb{D}} \gamma(w)$, and $\{e_i\}_{i=0}^\infty$ is an orthonormal basis of E . If*

$$\delta \leq \|\gamma(w)\|_E^2 \leq \sum_{i=0}^{\infty} \|\gamma_i\|_{Mult(\mathcal{D}_\alpha)}^2 < \infty, \quad w \in \mathbb{D},$$

for $\gamma(w) = \sum_{i=0}^{\infty} \gamma_i(w)e_i$ and some constant $\delta > 0$, then there exists operator $T \in \mathcal{B}_1(\mathbb{D})$ such that $\mathcal{E}_T = \mathcal{E}_{D_\alpha^*} \otimes \mathcal{E}$ and $T \sim_s D_\alpha^*$.

CONCLUDING REMARKS

Recall the Cowen-Douglas conjecture is that for operators T and S in $\mathcal{B}_1(\mathbb{D})$ having $\overline{\mathbb{D}}$ as a K -spectral set, T is similar to S if and only if $\lim_{|w| \rightarrow 1} \mathcal{K}_T(w)/\mathcal{K}_S(w) = 1$. In [10, 11], Clark and Misra constructed examples to show that the Cowen-Douglas conjecture is false and claimed that the ratio of the metrics can be a better indication of similarity than the ratio of the curvatures. Meanwhile, supporting example are provided in Theorem 1 of [11]. Theorem 3.4 and Proposition 3.6 in this note are positive results supporting the claim made by Clark and Misra. Then a natural question is, which Cowen-Douglas operator's similarity invariant is the ratio of the metrics?

Every Cowen-Douglas operator is unitary equivalent to the adjoint of a multiplication operator on a reproducing kernel Hilbert space consisting of holomorphic functions. For Hardy space and weighted Bergman spaces, Theorem 1 in [11], Theorem 3.8 in [55], Theorem 3.4 in [25], Theorem 3.4 and Proposition 3.6 suggest that, under certain conditions, the ratio of the metrics is a similarity invariant of the backward shift operator on these spaces. In particular, for the n -hypercontraction $T \in \mathcal{B}_1(\mathbb{D})$, from Theorem 0.1 in [38], Theorem 2.4 in [16], Theorem 2.3 in [24], and Theorem 3.4, we obtain that the operator T is similar to S_n^* on reproducing kernel Hilbert space \mathcal{M}_n if and only if $a_w := h_{\mathcal{E}_T}(w)/h_{\mathcal{E}_{S_n^*}}(w)$ is bonded and bounded from 0 if and only if the curvature $\mathcal{K}_{\mathcal{E}_{D_T}}$ of the holomorphic line bundle \mathcal{E}_{D_T} induced by the defect operator D_T satisfies $\mathcal{K}_{\mathcal{E}_{D_T}} = -\frac{\partial^2}{\partial w \partial \bar{w}} \varphi(w)$ for some bounded subharmonic function φ defined on \mathbb{D} .

Clark and Misra constructed two contraction backward shift operators in 2.5 of [11], such that the ratio of the metrics of the holomorphic line bundles associated with operators is bounded and bounded from 0, but they are not similar. This indicates that the ratio of metrics is not a similarity invariant for all backward shift operators on the Hilbert space of diagonal reproducing kernels. However, the following question is unknown:

Question 1. Is the ratio of the metrics of holomorphic line bundles a similarity invariant of the backward shift operator on the weighted Dirichlet space?

The similarity invariant of backward shift operators on weighted Dirichlet spaces cannot be obtained from the previous statements about Hardy space and weighted Bergman spaces. The main reason

weighted Dirichlet spaces differ from Hardy space and weighted Bergman spaces in Question 1 is that the multiplier algebra of weighted Dirichlet spaces is not equal to $H^\infty(\mathbb{D})$. For the similarity of backward shift operators on weighted Dirichlet spaces, assuming that the ratio of the metrics is bounded and bounded from 0, we introduce new tools such as holomorphic jet bundles and provide two sufficient conditions in Theorem 4.3 and Theorem 4.9, respectively. Meanwhile, as shown in Example 4.8 and Example 4.10, it is challenging to construct a backward shift operator that satisfies the following two conditions: (1) the ratio of the metric of the holomorphic line bundle induced by the operator to the metric of the holomorphic line bundle associated with the backward shift operator on Dirichlet space is bounded (or unbounded or tends to 0 as $|w| \rightarrow 1$); (2) the operator is not similar to the backward shift operator on Dirichlet space.

ACKNOWLEDGMENT

We gratefully acknowledge Y. H. Zhang and J. M. Yang for their valuable input and many helpful conversations regarding this paper.

REFERENCES

- [1] J. Agler, *The Arveson extension theorem and coanalytic models*, Integral Equ. Oper. Theory, **5** (1982), 608-631.
- [2] J. Agler, *Hypercontractions and subnormality*, J. Operator Theory, **13** (1985), 203-217.
- [3] J. Agler and J. E. McCarthy, *Pick Interpolation and Hilbert Function Spaces*, Vol. 44 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2002.
- [4] C.-G. Ambrozie, M. Enliš and V. Müller, *Operator tuples and analytic models over general domains in \mathbb{C}^n* , J. Operator Theory, **47** (2002), no. 2, 287-302.
- [5] J. Arazy and M. Enliš, *Analytic models for commuting operator tuples on bounded symmetric domains*, Trans. Amer. Math. Soc. **355** (2003), no. 2, 837-864.
- [6] A. Athavale, *Model theory on the unit ball in \mathbb{C}^m* , J. Operator Theory, **27** (1992), no. 2, 347-358.
- [7] J. A. Ball, T. T. Trent and V. Vinnikov, *Interpolation and commutant lifting for multipliers on reproducing kernel Hilbert spaces* in Operator Theory and Analysis, Oper. Theory Adv. Appl., 122, Birkhäuser, Basel, 2001, 89-138.
- [8] Y. Cao, J. S. Fang and C. L. Jiang, *K-groups of Banach algebras and strongly irreducible decomposition of operators*, J. Operator Theory, **48** (2002), no. 1, 235-253.
- [9] L. Carleson, *Interpolations by bounded analytic functions and the corona problem*, Ann. of Math. **76** (1962), no. 2, 547-559.
- [10] D. N. Clark and G. Misra, *On curvature and similarity*, Michigan Math. J. **30** (1983), no. 3, 361-367.
- [11] D. N. Clark and G. Misra, *On weighted shifts, curvature, and similarity*, J. Lond. Math. Soc. **31** (1985), no. 2, 357-368.
- [12] D. N. Clark and J. H. Morrel, *On Toeplitz Operators and Similarity*, Amer. J. Math. **100** (1978), no. 5, 973-986.
- [13] M. J. Cowen and R. G. Douglas, *Complex geometry and operator theory*, Acta. Math. **141** (1978), no. 1, 187-261.

- [14] M. J. Cowen and R. G. Douglas, *Equivalence of connections*, Acta. Math. **56** (1985), no. 1, 39-91.
- [15] R. E. Curto and N. Salinas, *Generalized Bergman kernels and the Cowen-Douglas theory*, Am. J. Math. **106** (1984), no. 2, 447-488.
- [16] R. G. Douglas, H. Kwon and S. Treil, *Similarity of n -hypercontractions and backward Bergman shifts*, J. Lond. Math. Soc. **88** (2013), no. 3, 637-648.
- [17] O. El-Fallah, K. Kellay, J. Mashreghi and T. Ransford, *A primer on the Dirichlet space* in Cambridge Tracts in Mathematics, 203, Cambridge University Press, Cambridge, 2014.
- [18] C. Foiaş, *A remark on the universal model for contractions of G* . C. Rota, Com. Acad. R. P. Romîne, **13** (1963), 349-352.
- [19] P. A. Fuhrmann, *On the corona theorem and its application to spectral problems in Hilbert space*, Trans. Amer. Math. Soc. **132** (1968), 55-66.
- [20] J. B. Garnett and P. W. Jones, *The corona theorem for Denjoy domains*, Acta Math. **155** (1985), no. 1-2, 27-40.
- [21] S. Ghara, S. Kumar and P. Pramanick, *\mathbb{K} -homogeneous tuple of operators on bounded symmetric domains*, Isr. J. Math. **247** (2022), no. 1, 331-360.
- [22] M. Golubitsky and V. Guillemin, *Stable mappings and their singularities*, Springer-Verlag, New York, 1973.
- [23] L. Hörmander, *Generators for some rings of analytic functions*, Bull. Amer. Math. Soc. **73** (1967), 943-949.
- [24] Y. L. Hou, K. Ji and H. Kwon, *The trace of the curvature determines similarity*, Studia Math. **236** (2017), no. 2, 193-200.
- [25] K. Ji and S. S. Ji, *The metrics of Hermitian holomorphic vector bundles and the similarity of Cowen-Douglas operators*, Indian J. Pure Appl. Math. **53** (2022), 736-749.
- [26] K. Ji, S. S. Ji, H. Kwon and J. Xu, *The Cowen-Douglas Theory for operator tuples and similarity*, <https://arxiv.org/abs/2210.00209v2>.
- [27] K. Ji, C. L. Jiang, D. K. Keshari and G. Misra, *Rigidity of the flag structure for a class of Cowen-Douglas operators*, J. Funct. Anal. **272** (2017), no. 7, 2899-2932.
- [28] K. Ji, H. Kwon, J. Sarkar and J. Xu, *A subclass of the Cowen-Douglas class and similarity*, Math. Nachr. **295** (2022), no. 11, 2197-2222.
- [29] K. Ji and J. Sarkar, *Similarity of quotient Hilbert modules in the Cowen-Douglas class*, Eur. J. Math. **5** (2019), no. 4, 1331-1351.
- [30] S. S. Ji and J. Xu, *On the trace of curvature of irreducible Cowen-Douglas operators*, J. Math. Anal. Appl. **512** (2022), 126181.
- [31] C. L. Jiang, *Similarity classification of Cowen-Douglas operators*, Can. J. Math. **56** (2004), no. 4, 742-775.
- [32] C. L. Jiang, X. Z. Guo and K. Ji, *K -group and similarity classification of operators*, J. Funct. Anal. **225** (2005), no. 1, 167-192.
- [33] C. L. Jiang and K. Ji, *Similarity classification of holomorphic curves*, Adv. Math. **215** (2007), 446-468.
- [34] C. L. Jiang and Y. C. Li, *The commutant and similarity invariant of analytic Toeplitz operators on Bergman space*, Sci. China Ser. A **50** (2007), no. 5, 651-664.

- [35] D. K. Keshari, *Trace formulae for curvature of jet bundles over planar domains*, Complex Anal. Oper. Theory, **8** (2014), no. 8, 1723-1740.
- [36] B. Kidane and T. T. Trent, *The corona theorem for multiplier algebras on weighted Dirichlet spaces*, Rocky Mt. J. Math. **43** (2013), no. 1, 241-271.
- [37] H. Kwon, *Similarity to the backward shift operator on the Dirichlet space*, J. Operator Theory, **76** (2016), no. 1, 133-140.
- [38] H. Kwon and S. Treil, *Similarity of operators and geometry of eigenvector bundles*, Publ. Mat. **53** (2009), no. 2, 417-438.
- [39] Y. C. Li and P. Ma, *On similarity and commutant of a class of multiplication operators on the Dirichlet space*, Banach J. Math. Anal. **17** (2023), no. 1, Paper No. 9.
- [40] Q. Lin, *Operator theoretical realization of some geometric notions*, Trans. Amer. Math. Soc. **305** (1988), no. 1, 353-367.
- [41] S. Luo, *Corona Theorem for the Dirichlet-Type Space*, J. Geom. Anal. **32** (2022), no. 3, 1-20.
- [42] G. Misra, *Curvature inequalities and extremal properties of bundle shifts*, J. Operator Theory, **11** (1984), no. 2, 305-317.
- [43] V. Müller, *Models for operators using weighted shifts*, J. Operator Theory, **20** (1988), no. 1, 3-20.
- [44] V. Müller and F. H. Vasilescu, *Standard models for some commuting multioperators*, Proc. Am. Math. Soc. **117** (1993), no. 4, 979-989.
- [45] N. K. Nikolskii, *Treatise on the shift operator. Spectral function theory*, With an appendix by S. V. Hruščev and V. V. Peller, Translated from the Russian by Jaak Peetre. Grundlehren der mathematischen Wissenschaften **273**, Springer-Verlag, Berlin, 1986.
- [46] M. Rosenblum, *A corona theorem for countably many functions*, Integral Equ. Oper. Theory, **3** (1980), no. 1, 125-137.
- [47] G. C. Rota, *Note on the invariant subspaces of linear operators*, Rend. Circ. Mat. Palermo, **8** (1959), no. 2, 182-184.
- [48] G. C. Rota, *On models for linear operators*, Comm. Pure Appl. Math. **13** (1960), 469-472.
- [49] A. L. Shields, *Weighted shift operators and analytic function theory*, in Math. Surveys, Amer. Math. Soc. **13** Providence, RI, 1974, 49-128.
- [50] D. Stegenga, *Multipliers of the Dirichlet space*, Illinois J. Math., **24** (1980), no. 1, 113-139.
- [51] B. Sz.-Nagy and C. Foiaş, *Sur les contractions de l'espace de Hilbert. VIII.*, Acta Sci. Math. **25** (1964), 38-71.
- [52] B. Sz.-Nagy and C. Foiaş, *Harmonic analysis of operators on Hilbert space*, North-Holland Publi. Co., Amsterdam 1970.
- [53] V. A. Tolokonnikov, *Estimates in the Carleson corona theorem, ideals of the algebra H^∞ , a problem of Sz.-Nagy*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **113** (1981), 178-198.
- [54] M. Uchiyama, *Corona theorems for countably many functions and estimates for their solutions*, preprint (1980).
- [55] M. Uchiyama, *Curvatures and similarity of operators with holomorphic eigenvectors*, Trans. Amer. Math. Soc. **319** (1990), no. 1, 405-415.

- [56] J. Xiao, *The $\bar{\partial}$ -problem for multipliers of the Sobolev space*, Manuscripta Math. **97** (1998), no. 2, 217-232.
- [57] K. Zhu, *Operators in Cowen-Douglas classes*, Ill. J. Math. **44** (2000), no. 4, 767-783.

Current address, Kui Ji and Shanshan Ji: School of Mathematics, Hebei Normal University, Shijiazhuang, Hebei 050016, China

Current address, Hyun-Kyoung Kwon: Department of Mathematics and Statistics, University at Albany-State University of New York, Albany, NY, 12222, USA

Current address, Jing Xu: School of Mathematics and Science, Hebei GEO University, Shijiazhuang 050031, China

Email address, K. Ji: jikui@hebtu.edu.cn

Email address, S. Ji: jishanshan15@outlook.com

Email address, H. Kwon: hkwon6@albany.edu

Email address, J. Xu: xujingmath@outlook.com