

Almost global existence and radiative decay of three dimensional spherical gas bubble inside inviscid compressible liquid

Liangchen Zou

School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui, 230026, PR China, zlc0601@mail.ustc.edu.cn

Abstract

The present paper considers the model of a homogeneous bubble inside an unbounded isentropic compressible inviscid liquid. The exterior liquid is governed by the Euler equation while the free bubble surface is determined by the kinematic and dynamic boundary conditions on the bubble-liquid interface. This classical model is of great concern in physics, and has been studied in a variety of approximations [11], [25], [16]. In this paper, we study this model in a first attempt from a mathematical point of view. We first proved the local existence and uniqueness of the complete nonlinear system using energy methods under an iteration scheme. Then we proved the almost global existence of the solution and the radiative decay of bubble oscillation through a bootstrap argument. Except for the energy estimate, this bootstrap argument encompasses a generalized KSS (Keel-Smith-Sogge) estimate and the analysis of backward pressure wave using the method of characteristics, which are the novelty of the present paper.

We developed a generalized weighted $L_t^2 H_x^j$ -estimate, or the so-called KSS estimate, which extends the KSS estimate [9] to nonlinear wave equations in exterior domains regardless of the boundary conditions, at the cost of only the appearance of a L_t^2 norm of the boundary value. To handle this boundary value, we establish a method of characteristics to study the backward pressure wave, which is then used to decouple the ODE of the boundary value from the hyperbolic system of backward and forward pressure wave. The analysis of backward pressure wave takes advantage of a change of variable between the backward and forward characteristics generated by the sound speed field in a geometric way. These two methods can not only be used for the bubble-liquid model studied in this paper, but are expected to be applied on other questions regarding nonlinear wave equations with complex boundary conditions.

Keywords: compressible Euler equation, free boundary, almost global existence, radiative decay, pressure wave

1. Introduction

1.1. Equations governing the liquid-bubble system

We consider a spherical gas bubble immersed in a compressible inviscid liquid. This physical relevant model plays an important role in a large variety of fields including underwater explosion [10], ultrasound imaging [2], medicine delivery [13], acoustic blocking [5], acoustic communication across the water-air interface [6], in which cases acoustic wave generated by compressibility and bubble oscillation are essential. For other applications of bubble dynamics, one can refer to [15], [14], [27],

and the references therein. The liquid under consideration is assumed inviscid and compressible with spherical symmetry, so the dynamics of liquid is governed by the compressible Euler equations, which, written in radial coordinates, is

$$\begin{cases} \partial_t \rho + r^{-2} \partial_r (r^2 \rho u) = 0, & r > R(t), t > 0, \\ \rho \partial_t u + \rho u \partial_r u + \partial_r p = 0, & r > R(t), t > 0. \end{cases} \quad (1.1)$$

Here ρ , u , p are the density, velocity and pressure of liquid respectively, and $R(t)$ is the bubble radius. We have assumed in (1.1) that the bubble center is on the origin, which otherwise can be achieved by transforming to the center-of-mass coordinates. On the other hand, the bubble surface is determined by the kinematic condition and the stress balance on the interface:

$$\begin{cases} \frac{dR}{dt} = u|_{r=R(t)}, & t > 0, \\ p|_{r=R(t)} = p_b - 2\sigma R^{-1}, & t > 0, \end{cases} \quad (1.2)$$

where p_b is the pressure at the inner bubble surface, and σ denotes the surface tension.

We further make the assumptions that the liquid is isentropic, so $p = C_0 \rho^\gamma$ for some positive constants C_0 and $\gamma > 1$, and that the bubble is homogeneous and satisfies a polytropic gas law, which says that the bubble pressure is uniform in space and is proportional to a power of bubble volume, say $p_b(t) = C_1 |R(t)|^{-3\gamma_0}$ for positive constant C_1 and $\gamma_0 > 1$. By nondimensionalization, we can assume without loss of generality that the liquid pressure $p = \frac{Ca}{2} \rho^\gamma$, the bubble pressure $p_b = \left(\frac{Ca}{2} + \frac{2}{We}\right) R^{-3\gamma_0}$, and the surface tension $\sigma = \frac{1}{We}$, where Ca and We are dimensionless constants, and namely, the Cavitation number and Weber number respectively. Therefore, (1.1)(1.2) admit an equilibrium state $(u, \rho, R) = (0, 1, 0)$. We assume in addition that the pressure at infinity is equal to the pressure at this equilibrium state, or equivalently, $\rho = 1$ at infinity.

To deal with this free boundary problem, a natural way is to introduce the Lagrangian coordinate $x := \int_{R(t)}^r \rho(t, s) s^2 dx$, which corresponds to the mass of liquid within a spherical domain of radius r and exterior to the bubble. Using the continuity equation and that $\frac{dR}{dt} = u|_{r=R(t)}$, there holds $\partial_t x(r, t) = -(r^2 \rho u)(r, t)$, therefore the Jacobian $\frac{\partial(x, t)}{\partial(r, t)}$ as well as its inverse reads

$$\begin{bmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial t} \\ \frac{\partial t}{\partial r} & \frac{\partial t}{\partial t} \end{bmatrix} = \begin{bmatrix} \rho r^2 & -\rho r^2 u \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} \frac{\partial r}{\partial x} & \frac{\partial r}{\partial t} \\ \frac{\partial t}{\partial x} & \frac{\partial t}{\partial t} \end{bmatrix} = \begin{bmatrix} (\rho r^2)^{-1} & u \\ 0 & 1 \end{bmatrix}. \quad (1.3)$$

By a change of variable using (1.3), equations (1.1)(1.2) are transformed into

$$\begin{cases} \partial_t \rho + \rho^2 \partial_x (r^2 u) = 0, & x > 0, t > 0, & (1.4) \\ \partial_t u + \frac{Ca}{2} r^2 \partial_x (\rho^\gamma) = 0, & x > 0, t > 0, & (1.5) \end{cases}$$

$$\begin{cases} \frac{dR}{dt} = u|_{x=0}, & t > 0, & (1.6) \end{cases}$$

$$\begin{cases} \frac{Ca}{2} \rho^\gamma|_{x=0} = \left(\frac{Ca}{2} + \frac{2}{We}\right) R^{-3\gamma_0} - \frac{2}{We} R^{-1} =: \tilde{f}(R), & t > 0, & (1.7) \end{cases}$$

$$\begin{cases} r = \left(R(t)^3 + 3 \int_0^x \rho^{-1}(y, t) dy \right)^{\frac{1}{3}} = r(x, 0) + \int_0^t u(y, \tau) d\tau, & x > 0, t > 0. & (1.8) \end{cases}$$

Next, we define $q = \rho^{-1} - 1$, and denote the specific sound speed by $c = \left(\frac{Ca\gamma}{2}\rho^{\gamma+1}\right)^{\frac{1}{2}}$ and the value of c at equilibrium state by $c_0 := \left(\frac{Ca\gamma}{2}\right)^{\frac{1}{2}}$. Then the fourth equation implies

$$q|_{x=0} = \left[\left(1 + \frac{2}{Ca} \frac{2}{We}\right) R^{-3\gamma_0} - \frac{2}{Ca} \frac{2}{We} R^{-1} \right]^{-\frac{1}{\gamma}} - 1 =: f(R).$$

$f(R)$ is smooth and strictly increasing for $R \in (0, \bar{R})$ with $\bar{R} = \left(1 + \frac{Ca}{2} \frac{We}{2}\right)^{\frac{1}{3\gamma_0-1}}$, and a simple calculation gives $c_0^2 f'(1) = -\tilde{f}'(1)$. Hence f admits a smooth inverse function f^{-1} defined on the range $f((0, \bar{R}))$. Replacing ρ by $(1+q)^{-1}$ in the above equations yields

$$\begin{cases} \partial_t q = \partial_x(r^2 u), & x > 0, t > 0, & (1.9) \\ \partial_t u = c^2 r^2 \partial_x q, & x > 0, t > 0, & (1.10) \end{cases}$$

$$\begin{cases} \frac{dR}{dt} = u|_{x=0}, & t > 0, & (1.11) \\ q|_{x=0} = f(R), & t > 0, & (1.12) \end{cases}$$

$$\begin{cases} r = \left(R(t)^3 + 3x + \int_0^x q(y, t) dy\right)^{\frac{1}{3}} = r(x, 0) + \int_0^t u(y, \tau) d\tau, & x > 0, t > 0. & (1.13) \end{cases}$$

We now impose the initial data at $t = 0$ that

$$(u, q, R)|_{t=0} = (u_{in}, q_{in}, R_{in}), \quad (1.14)$$

and compatibly $r(x, 0) = \left(R_{in}^3 + 3x + \int_0^x q_{in}(y) dy\right)^{\frac{1}{3}}$. The bubble-liquid system is then determined by equations (1.9)-(1.14).

In addition to the above equations, for later use we hereby derive the equation governing the velocity potential φ which is defined by $\rho r^2 \partial_x \varphi = u$ and $\varphi = 0$ at infinity. Substituting u by $\rho r^2 \partial_x \varphi$ in (1.5) and using (1.3)(1.4), there holds

$$\begin{aligned} & -\frac{Ca\gamma}{2(\gamma-1)} \rho r^2 \partial_x (\rho^{\gamma-1}) = \partial_t u \\ & = \rho r^2 \partial_x \partial_t \varphi + \partial_t (\rho r^2) \partial_x \varphi = \rho r^2 \partial_x \partial_t \varphi - \rho^2 \partial_x (r^2 u) r^2 \partial_x \varphi + 2r \rho u \partial_x \varphi \\ & = \rho r^2 \partial_x \partial_t \varphi - \rho^2 r^4 \partial_x u \partial_x \varphi = \rho r^2 \partial_x \left(\partial_t \varphi - \frac{u^2}{2} \right). \end{aligned}$$

Hence by cancelling $\rho r^2 \partial_x$ on each side, we obtain the relation between ρ and φ , namely the Bernoulli equation:

$$-\frac{Ca}{2} \frac{\gamma}{\gamma-1} (\rho^{\gamma-1} - 1) = \partial_t \varphi - \frac{u^2}{2}. \quad (1.15)$$

Multiplying (1.4) by $\rho^{\gamma-2}$ and replacing ρ using (1.15) yields

$$0 = \partial_t \left(\partial_t \varphi - \frac{u^2}{2} \right) - \frac{Ca\gamma}{2} \rho^\gamma \partial_x (\rho r^4 \partial_x \varphi).$$

Since $u \cdot \partial_t u = -\rho r^2 \partial_x \varphi \cdot \frac{Ca\gamma}{2} \rho^{\gamma-1} r^2 \partial_x \rho$, φ satisfies the following quasilinear wave equation on $(x, t) \in \mathbb{R}_+ \times \mathbb{R}_+$,

$$\partial_t^2 \varphi - c^2 \partial_x (r^4 \partial_x \varphi) = 0. \quad (1.16)$$

1.2. Observation from linear approximation

To illustrate the damping mechanism of bubble radius, we look at the linearisation of (1.16) at the equilibrium state. Denote by $\xi(x) = (1 + 3x)^{\frac{1}{3}}$ the radius function and by $c_0 = \left(\frac{Ca\gamma}{2}\right)^{\frac{1}{2}}$ the sound speed at equilibrium state, then the linearised equation reads

$$\partial_t^2 \varphi - c_0^2 (\partial_\xi^2 + \frac{2}{\xi} \partial_\xi) \varphi = 0.$$

In view of (1.15), the boundary conditions (1.11)(1.12) under linearisation become $\frac{dR}{dt} = \partial_\xi \varphi|_{\xi=1}$, and $\partial_t \varphi|_{\xi=1} = c_0^2 f'(1)(R - 1)$. Eliminating R by taking t -derivative in the latter yields

$$\partial_t^2 \varphi|_{\xi=1} - c_0^2 f'(1) \partial_\xi \varphi|_{\xi=1} = 0.$$

Let $\psi = \frac{\varphi}{\xi}$, then ψ satisfies the one-dimensional wave equation $\partial_t^2 \psi - c_0^2 \partial_\xi^2 \psi = 0$, while the above equation of boundary value becomes

$$\partial_t^2 \psi|_{\xi=1} + c_0 f'(1) \partial_t \psi|_{\xi=1} + c_0^2 f'(1) \psi|_{\xi=1} = c_0 f'(1) (\partial_t \psi + c_0 \partial_\xi \psi)|_{\xi=1}.$$

This is a second order inhomogeneous ODE of $\psi|_{\xi=1}$ with source term $c_0 f'(1) (\partial_t \psi + c_0 \partial_\xi \psi)|_{\xi=1}$, which corresponds to the backward pressure wave. Since ψ satisfies the one dimensional wave equation, we have

$$(\partial_t \psi + c_0 \partial_\xi \psi)|_{\xi=1}(t) = (\partial_t \psi + c_0 \partial_\xi \psi)(1 + c_0 t, 0).$$

The two eigenvalues, denoted by Λ_1, Λ_2 , have negative real part since f is strictly increasing, and therefore $(\psi|_{\xi=1}, \partial_t \psi|_{\xi=1})$ as well as $R - 1$ enjoy an exponential decay in time if we put the source term aside. In fact, solving the above ODE gives

$$\begin{aligned} \partial_t \psi|_{\xi=1}(t) &= \frac{\Lambda_1}{\Lambda_1 - \Lambda_2} e^{\Lambda_1 t} (\partial_t \psi - \Lambda_2 \psi)|_{\xi=1}(0) - \frac{\Lambda_2}{\Lambda_1 - \Lambda_2} e^{\Lambda_2 t} (\partial_t \psi - \Lambda_1 \psi)|_{\xi=1}(0) \\ &\quad + c_0 f'(1) \int_0^t \left(\frac{\Lambda_1}{\Lambda_1 - \Lambda_2} e^{\Lambda_1 s} - \frac{\Lambda_2}{\Lambda_1 - \Lambda_2} e^{\Lambda_2 s} \right) (\partial_t \psi + c_0 \partial_\xi \psi)(1 + c_0 s, 0) ds. \end{aligned}$$

Heuristically, for the nonlinear equations, $(\partial_t \psi + c_0 \partial_\xi \psi)$ will be replaced by

$$w_B := \partial_t \psi + cr^2 \partial_x \psi$$

since it stands for the backward pressure wave, and $(1 + c_0 s, 0)$ should be replaced by $(\xi_0(s), 0)$, where $\xi_0(s)$ is defined such that the characteristic starting from $(\xi_0(s), 0)$ intersects $\xi = 1$ at $(1, s)$. This leads to define

$$\begin{aligned} \mathcal{R}(t) &:= \frac{\Lambda_1}{\Lambda_1 - \Lambda_2} e^{\Lambda_1 t} (\partial_t \psi - \Lambda_2 \psi)|_{\{x=0, t=0\}} - \frac{\Lambda_2}{\Lambda_1 - \Lambda_2} e^{\Lambda_2 t} (\partial_t \psi - \Lambda_1 \psi)|_{\{x=0, t=0\}} \\ &\quad + c_0 f'(1) \int_0^t \left(\frac{\Lambda_1}{\Lambda_1 - \Lambda_2} e^{\Lambda_1 s} - \frac{\Lambda_2}{\Lambda_1 - \Lambda_2} e^{\Lambda_2 s} \right) w_B(\xi_0(s), 0) ds, \end{aligned} \tag{1.17}$$

where we also modify the definition of ψ to

$$\psi = r\varphi.$$

In the course of later proof, $\mathcal{R}(t)$ will be regarded as the principle part of $\partial_t \psi|_{\xi=1}(t)$. Meanwhile, we denote the forward pressure wave by

$$w_F := \partial_t \psi - cr^2 \partial_x \psi.$$

We will derive the precise nonlinear boundary ODE in Section 5 and specify the source term thereby, while the control of this source term is given in Section 6.

1.3. Review on established approaches describing bubble dynamics

The bubble dynamics was first studied by L. Rayleigh [26] in 1917, which describes how the liquid pressure develops in the process of collapse of a vacuum cavity in water. This first attempt considered the most simplified model. The cavity and liquid are assumed spherical, and other factors including compressibility, viscosity, thermal effects are omitted, while the pressure at infinity is assumed constant. Plesset (1949) [23] extended Rayleigh's work to the case where the pressure at infinity is time-dependent, which also established the well-known Plesset-Rayleigh equation. Plesset-Rayleigh equation have been widely applied in numerical simulations of bubble related models in which compressibility is less important.

To describe the pressure wave transmitted from a underwater explosion into the surrounding liquid, and to explain the damping of the bubble oscillation which Plesset-Rayleigh model failed to predict, Keller (1956) [10] modified Plesset's theory by taking compressibility into consideration and assumed the velocity potential satisfies a wave equation, which, as explained in a later work (1980) [11], is derived from linear approximation of Euler equations. It was also in [11] that Keller and Miksis established another frequently used ordinary differential equation (Keller-Miksis equation) regarding the bubble dynamics when the pressure wave emission has to be taken into consideration. Numerical simulations using Keller-Miksis model indicate the radiation decay due to acoustic wave emission by the bubble oscillation [31], [30], which also agrees well with experiment results. Prosperetti and Lezzi (1987) [25], [16] developed a second order theory in terms of the bubble-wall Mach number, which significantly increased the determinacy compared with the previous first order models. There are also many other generalizations of Plesset-Rayleigh equation regarding additional physical factors, including viscosity [3], the loss of spherical symmetry [12], the presence of mass exchange between bubble and liquid [24], thermal effects [1]. For a history of the understanding of bubbles, readers can refer to the review by Prosperetti in the book [4, Page 735].

To describe non-symmetry and compressibility simultaneously, numerical simulations have been carried out to identify mechanisms and features of non-spherical dynamics of acoustic cavitation bubbles [32], [33]. It is notable that Shapiro and Weinstein (2011) [27] use a new approach incorporating spherical harmonic decomposition and Laplace transform to describe in the linear level the dynamics of a non-spherical gas bubble in an inviscid, compressible liquid, and obtained the frequency of each mode in the spherical harmonic decomposition series. In our previous work [34], we also studied the model of a spherical gas bubble in a compressible viscous fluid, and proved the viscous damping for bubble oscillation.

1.4. Statement of the main result

To the best of our knowledge, our results in the present paper is a first attempt to rigorously study the complete nonlinear partial differential equations without any approximation or asymptotic expansion describing the bubble dynamics in a compressible, inviscid liquid. From (1.9)-(1.14) and

(1.16), we see that this question corresponds to a quasilinear wave equation in the exterior domain with a free boundary and nonlinear boundary conditions. Obviously, the approach of Laplace transform is no longer applicable as such to the nonlinear system (1.9)-(1.14). Hence we restricted the present study in the case with spherical symmetry.

The following is the statement of results in this work. Since we will work with the (x, t) variables instead of the usual Euclidean coordinates, we first write down the definition of function spaces we will work in.

Definition 1.1. Write $\xi(x) = (1 + 3x)^{\frac{1}{3}}$, and thus $x = \frac{\xi^3 - 1}{3}$, $\partial_\xi = \xi^2 \partial_x$. For positive integer j , define $H^j := \{v \in L^2(0, +\infty) : (\xi^2 \partial_x)^i v \in L^2(0, +\infty) \text{ for } i \in [0, j] \cap \mathbb{Z}\}$. For each $M \in (1, +\infty]$, denote by $L_\xi^2(1, M) := \{\tilde{v} \in L_{loc}^1(1, M) : \int_1^M |\tilde{v}(\xi)|^2 \xi^2 d\xi < +\infty\}$ the space of 3-dimensional radial square integrable functions on $B(0, M) \setminus \overline{B(0, 1)}$, and $H_\xi^j(1, M) := \{\tilde{v} \in L_\xi^2(1, M) : \partial_\xi^i \tilde{v} \in L_\xi^2(1, M) \text{ for } i \in [0, j] \cap \mathbb{Z}\}$ the corresponding Sobolev spaces.

From the definition we see immediately that if $v \in L^2(0, +\infty)$, then $\tilde{v}(\xi) := v(\frac{\xi^3 - 1}{3}) \in L_\xi^2(1, +\infty)$, and $v \in H^j$ is equivalent to $\tilde{v} \in H_\xi^j(1, +\infty)$. The first result is the local existence of system (1.9)-(1.14).

Theorem 1.2 (Local existence and uniqueness). *Let the initial value $(u_{in}, q_{in}, R_{in}) \in H^2 \times H^2 \times \mathbb{R}_+$ be such that $-1 < \underline{q} \leq q_{in} \leq \bar{q} < +\infty$ for all $x > 0$ and constants \underline{q}, \bar{q} , and that $0 < R_{in} < \bar{R}$. Then there exists a time $T > 0$ and a unique solution*

$$(u, q, R) \in L^\infty([0, T]; H^2) \cap C([0, T]; H^1) \times L^\infty([0, T]; H^2) \cap C([0, T]; H^1) \times C^1[0, T]$$

to (1.9)-(1.14) with initial value (u_{in}, q_{in}, R_{in}) . Moreover, a lower bound of T can be given in terms of $\|u_{in}\|_{H^2}$, $\|q_{in}\|_{H^2}$, \underline{q}, \bar{q} and $|f(R_{in})|$.

Theorem 1.3 (Almost global existence and radiative decay). *There exist constants $\epsilon_0, \tilde{\epsilon}_0, \kappa_0 > 0$ such that if the initial data $(u_{in}, q_{in}, R_{in}) \in H^2 \times H^2 \times \mathbb{R}_+$ satisfies*

$$\begin{aligned} \epsilon^2 &:= \sum_{j=0}^2 \|L_0^j q_{in}\|_{L^2}^2 + \sum_{j=0}^2 \|\tilde{L}_0^j u_{in}\|_{L^2}^2 + |f(R_{in})|^2 \leq \epsilon_0^2, \\ \tilde{\epsilon} &:= \|\xi^2 u_{in}\|_{L^2} + \|\xi^2 q_{in}\|_{L^2} \leq \tilde{\epsilon}_0, \end{aligned} \tag{1.18}$$

where the operators L_0^j, \tilde{L}_0^j are defined as

$$L_0^1 q := c_0^2 \partial_\xi q, L_0^2 q := c_0^2 \left(\partial_\xi^2 + \frac{2}{\xi} \partial_\xi \right) q, \tilde{L}_0^1 u = \left(\partial_\xi + \frac{2}{\xi} \partial_\xi \right) u, \tilde{L}_0^2 u = c_0^2 \left(\partial_\xi^2 + \frac{2}{\xi} \partial_\xi - \frac{2}{\xi^2} \right) u,$$

then there exist constants \underline{c}, \bar{c} such that $-\epsilon \lesssim \underline{c} - c_0 \leq c - c_0 \leq \bar{c} - c_0 \lesssim \epsilon$, and (1.9)-(1.14) has a unique solution

$$(u, q, R) \in L^\infty([0, T_\epsilon]; H^2) \cap C([0, T_\epsilon]; H^1) \times L^\infty([0, T_\epsilon]; H^2) \cap C([0, T_\epsilon]; H^1) \times C^1[0, T_\epsilon]$$

with $1 + \bar{c}T_\epsilon \geq \exp\left(\frac{\kappa_0}{\epsilon}\right)$. Moreover, for t satisfying $1 + \bar{c}t \leq \exp\left(\frac{\kappa_0}{\epsilon}\right)$, the bubble radius R satisfies the following estimate

$$|F(R) - \mathcal{R}(t)| \lesssim \frac{\log(1 + \bar{c}t)}{1 + \bar{c}t} \epsilon^{\frac{3}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \frac{\epsilon^{\frac{3}{2}}}{1 + \bar{c}t} \left(\epsilon^{\frac{1}{2}} + \tilde{\epsilon}^{\frac{1}{2}} \right) + \frac{\epsilon}{(1 + \bar{c}t)^2} (\epsilon + \tilde{\epsilon}),$$

where F is an explicit smooth function defined around $R = 1$, cf. (7.11)(7.13), and $\mathcal{R}(t)$ is as in (1.17). If assume additionally $w_B|_{t=0} = 0$, then the above estimate can be improved to

$$|F(R) - \mathcal{R}(t)| \lesssim \frac{\epsilon^{\frac{3}{2}}}{1 + \underline{c}t} \left(\epsilon^{\frac{1}{2}} + \tilde{\epsilon}^{\frac{1}{2}} \right).$$

Corollary 1.4 (Pointwise decay near the surface). *Let (u_{in}, q_{in}, R_{in}) be as in Theorem 1.3 with $\epsilon \ll \kappa_0$ and denote $T_0 := \bar{c}^{-1} \left(\exp\left(\frac{\kappa_0}{\epsilon}\right) - 1 \right)$. Assume additionally u_{in}, q_{in} are compactly supported in $[0, x_b]$, where x_b satisfies $\xi_b := (1 + 3x_b)^{\frac{1}{3}} \ll \exp\left(\frac{\kappa_0}{\epsilon}\right)$. Then for each point (x, t) with*

$$\xi := (1 + 3x)^{\frac{1}{3}} < 1 + \frac{1 + \underline{c}T_0 - \xi_b}{2}, \quad (1.19)$$

$$\frac{\xi + \xi_b - 2}{\underline{c}} \leq t \leq T_0 - \frac{\xi - 1}{\underline{c}}, \quad (1.20)$$

the solution (u, q, R) given by Theorem 1.3 with the initial data (u_{in}, q_{in}, R_{in}) satisfies the following pointwise decay for $t \gtrsim 1$:

$$\begin{aligned} |u(x, t)| + |q(x, t)| &\lesssim \xi^{-1} e^{\Lambda(t - \frac{\xi-1}{\underline{c}})} \left(|Y(0)| + \frac{|\tilde{f}'(1)|}{c_0} \int_0^{\frac{\xi_b-1}{\underline{c}}} e^{-\Lambda s} |w_B(\xi_0(s), 0)| ds \right) \\ &\quad + \xi^{-1} (2 + \underline{c}t - \xi)^{-1} \log \left(1 + \bar{c} \left(t + \frac{\xi - 1}{\underline{c}} \right) \right) \epsilon^2 \xi_b, \end{aligned}$$

where $\Lambda := \max\{Re\Lambda_1, Re\Lambda_2\}$. If assume additionally $w_B|_{t=0} = 0$, then the above bound can be improved to

$$\begin{aligned} |u(x, t)| + |q(x, t)| &\lesssim \xi^{-1} e^{\Lambda(t - \frac{\xi-1}{\underline{c}})} \left(|Y(0)| + \frac{|\tilde{f}'(1)|}{c_0} \int_0^{\frac{\xi_b-1}{\underline{c}}} e^{-\Lambda s} |w_B(\xi_0(s), 0)| ds \right) \\ &\quad + \xi^{-1} (2 + \underline{c}t - \xi)^{-1} \epsilon^2 \xi_b. \end{aligned}$$

1.5. Comments on the results

1. The maximal lifespan. The bound on the maximal lifespan given in Theorem 1.3 is sharp in three dimensions, as shown by finite propagation speed and the counterexample of Sideris [29] on singularities formation of boundaryless compressible fluids, which shows that a general small data may lead to blow up after a time of size $\exp\frac{\kappa}{\epsilon}$. Meanwhile, as shown in (1.16), the equations (1.9)-(1.14) can be written as a quasilinear wave equation, and the bound on the maximal lifespan in Theorem 1.3 also agrees with the results of Keel, Smith, and Sogge's studies on nonlinear wave equations in exterior domains [8], [9], [20]. Counterexamples of Sideris [28] and John [7] on nonlinear wave equations also indicate this optimal lifespan.

2. The KSS type estimate (4.1). Our techniques in Section 4 is in some way an extension to the KSS (Keel-Smith-Sogge) estimate developed in [9] for wave equations in exterior domains with Dirichlet condition, and can be applied on general quasilinear wave equations in exterior domains regardless of

boundary conditions. In fact, our calculation in Section 4 does not rely on any boundary conditions over the velocity potential φ which satisfies the quasilinear wave equation (1.16) in the exterior domain. The main feature of this extension can be illustrated by (4.22). We have the boundary values of the higher order space-time derivatives on the left-hand side of (4.22) and the boundary value of a lower order time derivative on the right. Hence (4.22) actually provides an approach to control higher regularities in terms of the lower ones on the boundary, which is unusual in any analysis. Moreover, by taking a delicate linear combination, all the boundary values of derivatives of φ can be controlled by barely the boundary value of φ on the L_t^2 level, cf. (4.1). We also remark that the spherical symmetry assumption is not necessary for this extension. In fact, the only assumption regarding the boundary we need is a uniform distance from the origin, which we used in (4.6) and (4.7).

3. The method of characteristics. The estimate of $w_B|_{\xi=1}$ in Proposition 6.1 is obtained by solving the hyperbolic system (5.1)(5.2) using characteristics. We note that the treatment in Section 6 can be extended to binary hyperbolic systems with variable coefficients on $(\xi, t) \in [1, +\infty) \times [0, +\infty)$ of the form

$$\begin{aligned}(\partial_t - c(x, t)\partial_\xi)w_B + \epsilon f_1(\xi, t)w_F + \epsilon g_1(\xi, t)w_B &= 0, \\(\partial_t + c(x, t)\partial_\xi)w_F + \epsilon f_2(\xi, t)w_B + \epsilon g_2(\xi, t)w_F &= 0,\end{aligned}$$

where $|c - c_0| \lesssim \epsilon \xi^{-1}$, $|f_i(\xi, t)| \lesssim \xi^{-1}$, $|g_i(\xi, t)| \leq g(\xi)$ for a given $g \in L^1(1, +\infty)$, and ϵ is sufficiently small. This will result in a Volterra inequality which holds for $t \lesssim \exp(\frac{\kappa}{\epsilon})$ and $\kappa \lesssim 1$:

$$\begin{aligned}v_B(\xi_0(t), 0) &\leq |w_B(\xi_0(t), 0)| + C\epsilon \mathcal{V}(\xi_0(t)) + C\epsilon \int_0^t \left(1 + \frac{\underline{c}}{2}(t-s)\right)^{-1} |w_F(1, t)| dt \\ &\quad + C\epsilon \kappa \int_1^{\xi_0(t)} (2\mu + \xi_0(t) - \eta)^{-1} v_B(\eta, 0) d\eta, \\ \mathcal{V}(\xi) &= |w_B(\xi, 0)| + \xi^{-1} \int_1^\xi |w_F(\eta, 0)| d\eta.\end{aligned}$$

Here $\xi_0(t)$ is the ξ coordinate of the intersection of $t = 0$ with the backward characteristic starting from $(1, t)$, and $v_B(\xi_0(t))$ is the maximum of $|w_B|$ along this characteristic. This Volterra inequality shall give a bound on $|w_B|$, and thus can play a role in decoupling the above binary hyperbolic system.

4. The decay rate of second order terms. The difference between $F(R)$ and its linear part $\mathcal{R}(t)$ is second order in terms of the size of initial data, and decays at a rate of $(1 + \underline{c}t)^{-1} \log(1 + \bar{c}t)$ or $(1 + \underline{c}t)^{-1}$ depending on whether backward pressure wave w_B exists at the initial time. This second order error comes from the reflection of pressure waves due to the variation of sound speed, or by formula from the terms

$$\left(\frac{\partial_t c + cr^2 \partial_x c}{2c}\right) \frac{w_B}{c^{\frac{1}{2}}}, \quad \left(\frac{\partial_t c - cr^2 \partial_x c}{2c}\right) \frac{w_F}{c^{\frac{1}{2}}}$$

in (5.1)(5.2). As proved in Section 6, these two terms will result in the error estimate (6.41). We remark that the decay rate $(1 + \underline{c}t)^{-1}$ in (6.41) of second order terms is the optimal one which can be deduced from the Volterra inequality (6.26). In fact, if we regard $w_B|_{t=0} \neq 0$ as first order, then

$\epsilon\mathcal{V}(\xi_0(t_*))$ and $\epsilon\Psi(t_*)$ should be viewed as second order perturbations, so we may extract the main part of (6.26) as

$$v_B(\xi_0(t), 0) \leq \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t), 0) \right| + C\epsilon \int_1^{\xi_0(t)} (2\mu + \xi_0(t) - \eta)^{-1} v_B(\eta, 0) d\eta.$$

Even if we assume $(c^{-\frac{1}{2}}w_B)(\xi, 0) = \mathbb{1}_{[1, \xi_b]}(\xi)$ for a constant $\xi_b \simeq 1$ so that it is compactly supported, solving the above Volterra equation in the standard way gives that

$$\begin{aligned} v_B(\xi, 0) &= \mathbb{1}_{[1, \xi_b]}(\xi) + \sum_{k=1}^{\infty} (C\epsilon)^k \int_1^{\xi} \cdots \int_1^{\xi} \prod_{i=1}^k \mathbb{1}_{\{\xi_{i-1} > \xi_i\}} (2\mu + \xi_{i-1} - \xi_i)^{-1} \mathbb{1}_{[1, \xi_b]}(\xi_n) d\xi_n \cdots d\xi_1 \\ &\geq \mathbb{1}_{[1, \xi_b]}(\xi) + C\epsilon \log \left(1 + \frac{\xi_b - 1}{2\mu + \xi - \xi_b} \right) \end{aligned}$$

with $\xi_0 := \xi$. Hence for large ξ we see that $v_B(\xi, 0) \geq C\epsilon \frac{\xi_b - 1}{2\mu + \xi - \xi_b}$, and one can not expect a faster decay rate than $(1 + c_t)^{-1}$ in the second order part of $v_B(\xi_0(t), 0)$.

5. The condition (1.18). Generally, the norms of the form in (1.18) will grow quadratic in time even for linear wave equations, so it may seem unnatural to impose (1.18) as a condition on initial data. However, we will only use this condition at $t = 0$. In fact, (1.18) is only used in Lemma 7.1 to deduce the controls of $w_B|_{t=0}$, and $w_F|_{t=0}$ at the initial time. While the norms in (1.18) may increase quadratic in time, the controls in Lemma 7.1 can propagate uniformly to $t > 0$, although we do not need this throughout our proofs. In other words, condition (1.18) is not a necessary but convenient condition to deduce the bounds in Lemma 7.1.

6. Expected results in higher dimensions or with null conditions. In contrast to the current three-dimensional isentropic setting, the solutions to quasilinear wave equations in exterior domains are global for small initial data in higher dimensions ($n \geq 4$) [19], [22], or with additional null conditions in three dimensions [21], [18]. In particular, if we consider a Chaplygin fluid (in which case the pressure p is linear in ρ^{-1} , or linear in q in our notations) outside the bubble instead of an isentropic fluid, our techniques can be easily modified to yield a global solution. In that case, the specific sound speed c is a constant, and our analysis would be greatly simplified.

1.6. Strategy of the proof

We provide in this section an overview of our proof.

1. Local existence. We construct an approximate solution sequence by iteration. In each step of this iteration, we define the approximate solution $(u_k, q_k, f(R_k))$ by solving the linear system (2.4)-(2.8) in which the parameters $r_{k-1}, c_{k-1}, f'(R_{k-1})$ are determined by $(u_{k-1}, q_{k-1}, f(R_{k-1}))$ from the previous step. It is worth noting that we choose $f(R_k)$ as the unknown in the linear equations (2.7) instead of naively linearising $f(R)$ at $R = 1$ and setting $q_k|_{x=0} = f'(R_{k-1})(R_k - 1)$. The latter choice will result in unclosure in the energy identity due to the appearance of the terms involving

$$\partial_t^{j+1} R_k \cdot \partial_t^i R_{k-1}.$$

This can also be observed from Lemma 2.2 where we do not allow an inhomogeneity in $q|_{x=0} = f$. We prove by induction local-in-time uniform L^2 bounds on all the space-time derivatives of order no

larger than two of each term in the approximate solution sequence, cf. Proposition 2.4. Due to the presence of the boundary, the higher order energy functionals e_k^j are defined by taking time derivatives which preserve the boundary. The method is to prove the local-in-time bounds on e_k^j , then convert these bounds on time derivatives to all space-time derivatives using equations (2.4)(2.5), cf. Lemma 2.6. Using the uniform bounds obtained in Proposition 2.4, Aubin-Lions compactness theorem is employed to yield a convergent subsequence and a limit, which is verified to be the solution of (1.9)-(1.14).

2. Bootstrap scheme. The almost global existence is proved under a bootstrap setting. We consider an interval $[0, T]$ on which the energy functionals satisfy the bound (3.1). It also follows that the parameters r, c and the solution itself satisfy the pointwise control (3.6)-(3.10). We aim at strictly improving (3.1) on the interval $[0, T]$, so the solution can be continued beyond $[0, T]$ until T reaches the bound (3.11).

3. Control of nonlinearities via KSS estimate. To improve the bound (3.1), the first step is to control the nonlinearities in the energy identity (2.34) (removing the iteration subscripts k) or (7.1). In view of the pointwise controls (3.6)-(3.10), the integrands in these nonlinearities are of the size $\epsilon \xi^{-1} \left((\partial_t^j u)^2 + (\partial_t^j q)^2 \right)$. Since the equations (1.9)-(1.14) can be transformed to the quasilinear wave equation (1.16), the estimate of these nonlinearities can hopefully be achieved by the said KSS (Keel-Smith-Sogge) estimate, which has been applied to control such nonlinearities for nonlinear wave equations in exterior domains with Dirichlet condition in [8], [9]. The difference is that we need an additional control over $\int_0^T \varphi^2|_{x=0} dt$ due to the absence of a vanishing boundary condition.

4. Hyperbolic system (5.1)(5.2) coupling boundary ODE (5.3). To control $\int_0^T \varphi^2|_{x=0} dt$, we begin by looking at how $\varphi|_{x=0}$ is determined. By taking time derivative in (1.7) and substituting ρ using (1.15), we arrive at the ODE (5.3) of $\psi|_{x=0}$ with $\psi = r\varphi$. On one hand, the backward pressure wave $w_B|_{x=0}$ acts as an inhomogeneity in the ODE (5.3), so we have to carry out the characteristic method to estimate $w_B|_{x=0}$, which is initialized in (5.1)(5.2), or equivalently (6.1)(6.2). On the other hand, to solve the hyperbolic system (5.1)(5.2) one needs the data of w_F on the boundary $\xi = 1$, which is in fact given by the ODE (5.3) and $w_F = 2\partial_t\psi - w_B$.

5. Decoupling (5.1)(5.2)(5.3). We first integrate (5.1)(5.2) on each characteristic to obtain inequalities (6.6)(6.7)(6.8) of v_B and v_F , which are the maximum of $|c^{-\frac{1}{2}}w_B|$ along the backward characteristic and the maximum of $|c^{-\frac{1}{2}}w_F|$ along the forward characteristic. To combine these inequalities, we study the regularities of the change of variables between the backward and forward characteristics in different regions through a geometric approach and simplify these inequalities to (6.18)(6.19)(6.20). Then from (6.18)(6.19)(6.20), we derive a Volterra inequality (6.26) on v_B which then deduces an estimate (6.32) of $w_B|_{\xi=1}$. With the estimate (6.32) we are able to decoupling (5.3) from (5.1)(5.2) and obtain the estimate (6.41) for $\psi|_{x=0}$ and $\partial_t\psi|_{x=0}$.

6. Closing the bootstrap. Combing the KSS estimate (4.1) and (6.41), the bound (3.1) is then strictly improved on $[0, T]$ for all T satisfying (3.11), which finishes the bootstrap argument. A by-products of (6.41) is an error estimate of the bubble radius R from the solution of linearised equation as shown in Theorem 1.3. Moreover, a pointwise decay-in-time estimate of (u, q) as shown in Corollary 1.4 is recovered from the estimates of w_F and w_B obtained in Section 6.

1.7. Organization of the sections

In Section 2, we prove the local existence and uniqueness under the iteration scheme. In Section 3, we state the bootstrap assumptions and deduce the resulting pointwise bounds on c , r and the solution itself. In Section 4, we derive the generalized KSS estimate without boundary conditions. In Section 5, we derive the boundary ODE determining $\psi|_{x=0}$ and the binary hyperbolic system of the forward and backward pressure waves w_F, w_B . In Section 6, we first study the regularities of the change of variables between the forward and backward characteristics. Using these change of variables, we prove the Volterra inequality (6.26), and use (6.26) to obtain the control on $\psi|_{x=0}$. In Section 7, we close the bootstrap argument by combing the KSS estimate and the control on $\psi|_{x=0}$, and prove the pointwise estimate of (u, q, R) .

Acknowledgments. To do this study has been caused by helpful discussions with Associate Professor Lifeng Zhao (University of Science of Technology of China). This work was supported by NSFC Grant of China No. 12271497.

2. Energy estimate and Local existence

In this section, we shall prove Theorem 1.2 by means of an iterative scheme to solve (1.9)-(1.14) and to show a uniform bound for each iterative, then proceed by compactness argument. Before the analysis of iteration, we look at an elliptic estimate and a model question, which will provide the energy identity.

Lemma 2.1 (Weighted elliptic estimate). *Let $v \in H^1$ and $r \in AC[0, +\infty)$ be such that $r(0) > 0$ and $+\infty > \bar{\theta} \geq r^2 \partial_x r \geq \underline{\theta} > 0$ for some constant $\underline{\theta}$ and $\bar{\theta}$. Then we have the identity*

$$\int_0^\infty (r^2 \partial_x r)^{-1} (\partial_x (r^2 v))^2 dx + 2rv^2|_{x=0} = \int_0^\infty (r^2 \partial_x r)^{-1} (r^2 \partial_x v)^2 dx + 2 \int_0^\infty (r^2 \partial_x r) r^{-2} v^2 dx, \quad (2.1)$$

and the pointwise control

$$\begin{aligned} |v(x)|^2 &\leq r^{-2}(x) (\|v\|_{L^2}^2 + \|r^2 \partial_x v\|_{L^2}^2) \\ |v(x)|^2 &\leq r^{-2}(x) ((8r(0)^{-2} \bar{\theta}^2 + 1) \|v\|_{L^2}^2 + 2 \|\partial_x (r^2 v)\|_{L^2}^2) \end{aligned} \quad (2.2)$$

Proof. The assumption of r gives $r(0)^3 + 3\bar{\theta}x \geq r(x)^3 \geq r(0)^3 + 3\underline{\theta}x$, so $r^2 \partial_x v \in L^2$. Note that $(\partial_x (r^2 v))^2 = (r^2 \partial_x v)^2 + 2(r^2 \partial_x r)^2 r^{-2} v^2 + 2(r^2 \partial_x r) \partial_x (r v^2)$, therefore (2.1) follows from integration by parts. For any $y > x > 0$, there holds $v^2(x) - v^2(y) = \int_x^y 2v(z) \partial_x v(z) dz$. Integrating in y on $I := [x, \tilde{x}]$ yields $|I|v^2(x) = \int_I v^2(y) dy + 2 \int_I \int_x^y v(z) \partial_x v(z) dz dy$. Then by Cauchy-Schwartz

$$v^2(x) \leq (\tilde{x} - x)^{-1} \int_I v^2(y) dy + r^{-2}(x) \int_I v^2(z) dz + r^{-2}(x) \int_I (r^2 \partial_x v(z))^2 dz$$

The first inequality of (2.2) follows by forcing \tilde{x} to infinity. Note that $|r^2 \partial_x v| \leq |\partial_x (r^2 v)| + 2|r \partial_x r| |v|$ and $|r \partial_x r| \leq r(0)^{-1} \bar{\theta}$, therefore the second line in (2.2) follows from Cauchy-Schwartz. \square

Lemma 2.2 (Model question). *Let (u, q, f) satisfy the following equations*

$$\begin{cases} \partial_t u - c^2 r^2 \partial_x q = F_1, & x > 0, t > 0, \\ \partial_t q - \partial_x(r^2 u) = F_2, & x > 0, t > 0, \\ u|_{x=0} = g \frac{d}{dt} f + F_3, & t > 0, \\ q|_{x=0} = f, & t > 0. \end{cases}$$

Then the energy identity reads

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left[\int_0^\infty (u^2 + c^2 q^2) dx + (c^2 r^2)|_{x=0} g f^2 \right] \\ &= \int_0^\infty \left(\frac{1}{2} (\partial_t c^2) q^2 - (r^2 \partial_x c^2) u q + F_1 u + c^2 F_2 q \right) dx + \frac{1}{2} \partial_t [(c^2 r^2)|_{x=0} g] f^2 - (c^2 r^2)|_{x=0} F_3 f. \end{aligned} \quad (2.3)$$

The proof is straightforward by multiplying u on the first equation and integration by parts. We now begin with the construction of iteration sequence. Set $u_0 = 0, q_0 = 0, R_0 = 1, c_0^2 = \frac{Ca\gamma}{2}, r_0(x) = (1 + 3x)^{\frac{1}{2}}$ and thus $f(R_0) = 0$. Then define $(u_k, q_k, R_k), k \geq 1$ inductively by solving the linear equations of $(u_k, q_k, f(R_k))$

$$\begin{cases} \partial_t q_k = \partial_x(r_{k-1}^2 u_k), & x > 0, t > 0, \\ \partial_t u_k = c_{k-1}^2 r_{k-1}^2 \partial_x q_k, & x > 0, t > 0, \end{cases} \quad (2.4)$$

$$\begin{cases} u_k|_{x=0} = (f'(R_{k-1}))^{-1} \frac{d}{dt} f(R_k), & t > 0, \end{cases} \quad (2.6)$$

$$\begin{cases} q_k|_{x=0} = f(R_k), & t > 0, \end{cases} \quad (2.7)$$

$$\begin{cases} (u_k, q_k, f(R_k))|_{t=0} = (u_{in}, q_{in}, f(R_{in})). \end{cases} \quad (2.8)$$

Here the parameters r_{k-1} and c_{k-1} are determined by the previous iterative that

$$r_{k-1}^3 = R_{k-1}^3 + 3x + 3 \int_0^x q_{k-1}(y) dy, \quad c_{k-1}^2 = \frac{Ca\gamma}{2} (1 + q_{k-1})^{-1-\gamma}.$$

Introduce the time dependent operators

$$\begin{aligned} L_{k-1}^{2j} &:= [\partial_x(r_{k-1}^4 c_{k-1}^2 \partial_x)]^j, & L_{k-1}^{2j+1} &:= c_{k-1}^2 r_{k-1}^2 \partial_x L_{k-1}^{2j}, \\ \tilde{L}_{k-1}^{2j} &:= [c_{k-1}^2 r_{k-1}^2 \partial_x^2 r_{k-1}^2]^j, & \tilde{L}_{k-1}^{2j+1} &:= \partial_x(r_{k-1}^2 \tilde{L}_{k-1}^{2j}). \end{aligned}$$

Equations (2.4)(2.5) can then be written as $\partial_t q_k = \tilde{L}_{k-1}^1 u_k$ and $\partial_t u_k = L_{k-1}^1 q_k$. Moreover, for each j , there holds

$$\tilde{L}_{k-1}^{j+1} = L_{k-1}^j \tilde{L}_{k-1}^1, \quad L_{k-1}^{j+1} = \tilde{L}_{k-1}^j L_{k-1}^1.$$

Since L_0^1 and \tilde{L}_0^1 commute with ∂_t , applying ∂_t^j to (2.4)-(2.7) yields

$$\begin{cases} \partial_t \partial_t^j u_1 - L_0^1 \partial_t^j q_1 = 0, & x > 0, t > 0, \\ \partial_t \partial_t^j q_1 - \tilde{L}_0^1 \partial_t^j u_1 = 0, & x > 0, t > 0, \\ \partial_t^j q_1|_{x=0} = \partial_t^j f(R_1), & t > 0, \\ \partial_t^j u_1|_{x=0} = f'(1)^{-1} \frac{d}{dt} \left(\frac{d}{dt} \right)^j f(R_1), & t > 0. \end{cases}$$

Using (2.3), we obtain

$$0 = \frac{1}{2} \frac{d}{dt} \left[\int_0^\infty ((\partial_t^j u_1)^2 + c_0^2 (\partial_t^j q_1)^2) dx + f'(1)^{-1} c_0^2 (\partial_t^j f(R_1))^2 \right] \quad (2.9)$$

In order to obtain the estimate for H^2 norms of u_1 and q_1 , the next step is to exchange t -derivatives to x -derivatives.

Proposition 2.3 (Bound of mixed derivatives for initial iterative).

$$\begin{aligned} & \sum_{j=0}^2 |\partial_t^j f(R_1)|^2 + \sum_{j_1+j_2 \leq 2} \left(\|L_0^{j_2} \partial_t^{j_1} q_1\|_{L^2}^2 + \|\tilde{L}_0^{j_2} \partial_t^{j_1} u_1\|_{L^2}^2 \right) \\ & \lesssim \sum_{j=0}^2 \|L_0^j q_{in}\|_{L^2}^2 + \sum_{j=0}^2 \|\tilde{L}_0^j u_{in}\|_{L^2}^2 + |f(R_{in})|^2. \end{aligned} \quad (2.10)$$

Proof. By (2.4)(2.5) and that L_0^1, \tilde{L}_0^1 commute with ∂_t , for each (j_1, j_2) , there holds

$$\begin{aligned} L_0^{j_1} \partial_t^{j_2} q_1 &= \partial_t^{j_1+j_2} q_1, & \tilde{L}_0^{j_1} \partial_t^{j_2} u_1 &= \partial_t^{j_1+j_2} u_1, & \text{for } j_1 \text{ even,} \\ L_0^{j_1} \partial_t^{j_2} q_1 &= \partial_t^{j_1+j_2} q_1, & \tilde{L}_0^{j_1} \partial_t^{j_2} u_1 &= \partial_t^{j_1+j_2} u_1, & \text{for } j_1 \text{ odd.} \end{aligned}$$

Hence in view of (2.9) the left hand side of (2.10) is bounded by multiple of $\sum_{j=0}^2 \|\partial_t^j q_1|_{t=0}\|_{L^2}^2 + \sum_{j=0}^2 \|\partial_t^j u_1|_{t=0}\|_{L^2}^2 + \sum_{j=0}^2 |\partial_t^j f(R_1)|_{t=0}|^2$, and for the same reason the right hand side is exactly $\sum_{j=0}^2 \|\partial_t^j q_1|_{t=0}\|_{L^2}^2 + \sum_{j=0}^2 \|\partial_t^j u_1|_{t=0}\|_{L^2}^2 + |f(R_1)|_{t=0}|^2$. It remains to bound $\partial_t f(R_1)|_{t=0}$ and $\partial_t^2 f(R_1)|_{t=0}$. By (2.6), $\partial_t^{j+1} f(R_1) = f'(1) \partial_t^j u_1|_{x=0}$. The elliptic estimate (2.2) gives that for $j = 0, 1$,

$$|\partial_t^j u|_{x=0} \lesssim \|\partial_t^j u_1\|_{L^2} + \|\tilde{L}_0^1 \partial_t^j u_1\|_{L^2} \leq \|\tilde{L}_0^j u_1\|_{L^2} + \|\tilde{L}_0^{j+1} u_1\|_{L^2} + \|L_0^j q_1\|_{L^2} + \|L_0^{j+1} q_1\|_{L^2}$$

Restricting t to 0 shows $|\partial_t f(R_1)|_{t=0}|^2 + |\partial_t^2 f(R_1)|_{t=0}|^2 \lesssim \sum_{j=0}^2 \|L_0^j q_{in}\|_{L^2}^2 + \sum_{j=0}^2 \|\tilde{L}_0^j u_{in}\|_{L^2}^2$. \square

The center of construction of local solution is to prove the same bound as in Proposition 2.3 for each iterative $(u_k, q_k, f(R_k))$ on a common interval $[0, T]$. That is

Proposition 2.4 (Uniform bound of mixed derivatives). *There exists $T > 0$ and a constant $A > 0$ such that for each $k \geq 0$,*

$$\begin{aligned} & \sum_{j=0}^2 |\partial_t^j f(R_k)|^2 + \sum_{j_1+j_2 \leq 2} \left(\|L_{k-1}^{j_2} \partial_t^{j_1} q_k\|_{L^2}^2 + \|\tilde{L}_0^{j_2} \partial_t^{j_1} u_k\|_{L^2}^2 \right) \\ & \leq A \left(\sum_{j=0}^2 \|L_0^j q_{in}\|_{L^2}^2 + \sum_{j=0}^2 \|\tilde{L}_0^j u_{in}\|_{L^2}^2 + |f(R_{in})|^2 \right), \end{aligned} \quad (2.11)$$

holds on $t \in [0, T]$.

Define for $j = 0, 1, 2$ and $k \geq 0$ that

$$e_k^j(t) := \frac{1}{2} \int_0^\infty \left((\partial_t^j u_k(t))^2 + c_{k-1}^2(t) (\partial_t^j q_k(t))^2 \right) dx + \frac{1}{2} c_{k-1}(t)|_{x=0}^2 f'(R_{k-1}(t))^{-1} \left(\partial_t^j f(R_k(t)) \right)^2,$$

$$E_k(t) := \sum_{j=0}^2 |\partial_t^j f(R_k)(t)|^2 + \sum_{j_1+j_2 \leq 2} \left(\|L_{k-1}^{j_2} \partial_t^{j_1} q_k(t)\|_{L^2}^2 + \|\tilde{L}_{k-1}^{j_2} \partial_t^{j_1} u_k(t)\|_{L^2}^2 \right),$$

$$\epsilon^2 := \sum_{j=0}^2 \|L_0^j q_{in}\|_{L^2}^2 + \sum_{j=0}^2 \|\tilde{L}_0^j u_{in}\|_{L^2}^2 + |f(R_{in})|^2.$$

The proof will be completed by induction in k with the assistance of the following lemmas.

Lemma 2.5 (Control of parameters). *There exists $T > 0$ and constant $C > 0$ dependent on q, \bar{q} and R_{in} such that for each $k \geq 0$ and $t \in [0, T]$*

$$\frac{q-1}{2} \leq q_k(x, t) \leq 2\bar{q} + 1, \quad x \in [0, +\infty), \quad (2.12)$$

$$\frac{Ca\gamma}{2} (2 + 2\bar{q})^{-\gamma-1} \leq c_k^2(x, t) \leq \frac{Ca\gamma}{2} \left(\frac{q+1}{2} \right)^{-\gamma-1}, \quad x \in [0, +\infty), \quad (2.13)$$

$$\frac{f(R_{in})-1}{2} \leq f(R_k) \leq 2f(R_{in}) + 1, \quad f^{-1} \left(\frac{f(R_{in})-1}{2} \right) \leq R_k \leq f^{-1} (2f(R_{in}) + 1), \quad (2.14)$$

$$R_k^3 + \frac{3}{2}(q+1)x \leq r_k^3 \leq R_k^3 + 6(\bar{q}+1)x, \quad x \in [0, \infty), \quad (2.15)$$

$$\|v\|_{L^\infty}^2 \leq C (\|v\|_{L^2}^2 + \|L_k v\|_{L^2}^2), \quad \|v\|_{L^\infty} \leq C (\|v\|_{L^2}^2 + \|\tilde{L}_k^1 v\|_{L^2}^2). \quad (2.16)$$

Lemma 2.6. *There exists constant $B > 0$ and $\tilde{B} > 0$ such that for each $k \geq 0$*

$$E_k(t) \leq B \sum_{j=0}^2 e_k^j(t), \quad t \in [0, T], \quad (2.17)$$

$$\sum_{j=0}^2 e_k^j(0) \leq \tilde{B} \epsilon^2. \quad (2.18)$$

From Lemma 2.6 we see that if $\sum_{j=0}^2 e_k^j(t) \leq 2 \sum_{j=0}^2 e_k^j(0)$ holds on $t \in [0, T]$ for some T independent on k , then we obtain $E_k(t) \leq 2B\tilde{B}\epsilon^2$, i.e. (2.11) holds for $A = 2B\tilde{B}$. Now we suppose that Proposition 2.4, Lemma 2.5, Lemma 2.6 hold for all $k' \leq k-1$.

Proof of Lemma 2.6. Equations (2.4)(2.5) is equivalent to $\partial_t u_k = L_{k-1}^1 q_k$ and $\partial_t q_k = \tilde{L}_{k-1}^1 u_k$, and also yield that

$$L_{k-1}^2 q_k = \tilde{L}_{k-1}^1 \partial_t u_k, \quad \tilde{L}_{k-1}^2 u_k = L_{k-1}^1 \partial_t q_k. \quad (2.19)$$

Hence according to the construction of E_k , in order to show (2.17) it suffices to control $\|L_{k-1}^1 \partial_t q_k\|_{L^2}$ and $\|\tilde{L}_{k-1}^1 \partial_t u_k\|_{L^2}$. By differentiating (2.4)(2.5) we obtain for $t \in [0, T]$

$$\begin{aligned}
\partial_t^2 q_k - \tilde{L}_{k-1}^1 \partial_t u_k &= 2\partial_x(r_{k-1}u_{k-1}u_k) \\
&= 2r_{k-1}^{-1}u_{k-1}\tilde{L}_{k-1}^1 u_k + 2r_{k-1}^2 \partial_x(r_{k-1}^{-1}u_{k-1})u_k \\
&= 2r_{k-1}^{-1}u_{k-1}\tilde{L}_{k-1}^1 u_k + 2r_{k-1}^2 \partial_x((r_{k-1}r_{k-2}^2)^{-1}r_{k-2}^2 u_{k-1})u_k \\
&= 2r_{k-1}^{-1}u_{k-1}\tilde{L}_{k-1}^1 u_k + 2r_{k-2}^{-2}r_{k-1}u_k\tilde{L}_{k-2}^1 u_{k-1} \\
&\quad + 2r_{k-1}^2 r_{k-2}^2 u_k u_{k-1} \partial_x(r_{k-1}r_{k-2}^2)^{-1} \\
&= 2r_{k-1}^{-1}u_{k-1}\tilde{L}_{k-1}^1 u_k + 2r_{k-2}^{-2}r_{k-1}u_k\tilde{L}_{k-2}^1 u_{k-1} \\
&\quad - 2r_{k-1}^{-2}(1+q_{k-1})u_k u_{k-1} - 4r_{k-2}^{-3}r_{k-1}(1+q_{k-2})u_k u_{k-1} \\
&= 2r_{k-1}^{-1}u_{k-1}\partial_t q_k + 2r_{k-2}^{-2}r_{k-1}u_k \partial_t q_{k-1} \\
&\quad - 2r_{k-1}^{-2}(1+q_{k-1})u_k u_{k-1} - 4r_{k-2}^{-3}r_{k-1}(1+q_{k-2})u_k u_{k-1},
\end{aligned} \tag{2.20}$$

$$\begin{aligned}
\partial_t^2 u_k - L_{k-1}^1 \partial_t q_k &= \frac{\partial_t(c_{k-1}^2 r_{k-1}^2)}{c_{k-1}^2 r_{k-1}^2} L_{k-1} q_k \\
&= \left(-(\gamma+1) \frac{\partial_t q_{k-1}}{1+q_{k-1}} + 2 \frac{u_{k-1}}{r_{k-1}} \right) \partial_t u_k \\
&= \left(-(\gamma+1) \frac{\tilde{L}_{k-2}^1 u_{k-1}}{1+q_{k-1}} + 2 \frac{u_{k-1}}{r_{k-1}} \right) L_{k-1}^1 q_k.
\end{aligned} \tag{2.21}$$

Therefore by Lemma 2.5 and Proposition 2.4 for $t \in [0, T]$ there holds

$$\begin{aligned}
&\|\partial_t^2 q_k - \tilde{L}_{k-1}^1 \partial_t u_k\|_{L^2} \\
&\leq C(\|u_{k-1}\|_{L^\infty} + \|\partial_t q_{k-1}\|_{L^\infty})(\|\partial_t q_k\|_{L^2} + \|u_k\|_{L^2}) \\
&\leq C\left(\|u_{k-1}\|_{L^2} + \|\tilde{L}_{k-2}^1 u_{k-1}\|_{L^2} + \|\partial_t q_{k-1}\|_{L^2} + \|L_{k-2}^1 \partial_t q_{k-1}\|_{L^2}\right)(\|\partial_t q_k\|_{L^2} + \|u_k\|_{L^2}) \\
&\leq CA^{\frac{1}{2}}\epsilon(\|\partial_t q_k\|_{L^2} + \|u_k\|_{L^2}) \\
&= CA^{\frac{1}{2}}\epsilon\left(\|\tilde{L}_{k-1}^1 u_k\|_{L^2} + \|u_k\|_{L^2}\right),
\end{aligned} \tag{2.22}$$

$$\begin{aligned}
\|\partial_t^2 u_k - L_{k-1}^1 \partial_t q_k\|_{L^2} &\leq C(\|u_{k-1}\|_{L^\infty} + \|\partial_t q_{k-1}\|_{L^\infty})\|\partial_t u_k\|_{L^2} \\
&\leq CA^{\frac{1}{2}}\epsilon\|\partial_t u_k\|_{L^2} \\
&= CA^{\frac{1}{2}}\epsilon\|L_{k-1}^1 q_k\|_{L^2},
\end{aligned} \tag{2.23}$$

which complete the proof of (2.17) for $B = C(1 + A^{\frac{1}{2}}\epsilon)$.

To show (2.18), restricting t to 0 in (2.20)(2.21) and using (2.19) gives

$$\begin{aligned}
\partial_t^2 q_k|_{t=0} - L_{k-1}^2 q_{in} &= 4r_{in}^{-1}u_{in}\partial_x(r_{in}^2 u_{in}) - 6r_{in}^{-2}(1+q_{in})u_{in}^2 \\
&= 4r_{in}^{-1}u_{in}\tilde{L}_{k-1}^1 u_{in},
\end{aligned} \tag{2.24}$$

$$\begin{aligned}
\partial_t^2 u_k|_{t=0} - \tilde{L}_{k-1}^2 u_{in} &= (-(\gamma+1)(1+q_{in})^{-1}\partial_x(r_{in}^2 u_{in}) + 2r_{in}^{-1}u_{in})c_{in}^2 r_{in}^2 \partial_x q_{in} \\
&= \left(-(\gamma+1)(1+q_{in})^{-1}\tilde{L}_{k-1}^1 u_{in} + 2r_{in}^{-1}u_{in} \right) L_{k-1}^1 q_{in}.
\end{aligned} \tag{2.25}$$

Moreover, we can re-express L_{k-1} and \tilde{L}_{k-1} by L_0 and \tilde{L}_0 as the following.

$$\tilde{L}_{k-1}^1 u_k = \frac{r_{k-1}^2}{r_0^2} \tilde{L}_0^1 u_k + 2r_{k-1}^{-1} (1 + q_{k-1} - r_0^{-3} r_{k-1}^3) u_k, \quad (2.26)$$

$$L_{k-1}^1 q_k = \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^2}{r_0^2} L_0^1 q_k, \quad (2.27)$$

$$\begin{aligned} & \tilde{L}_{k-1}^2 u_k \\ &= \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^4}{r_0^4} \tilde{L}_0^2 u_k + \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^2}{r_0^2} L_0^1 \left(\frac{r_{k-1}^2}{r_0^2} \right) \tilde{L}_0^1 u_k + 2 \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^{k-1}}{r_0^2} (1 + q_{k-1} - r_0^{-3} r_{k-1}^3) L_0^1 u_k \\ & \quad + 2 \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^2}{r_0^2} L_0^1 (r_{k-1}^{-1} (1 + q_{k-1} - r_0^{-3} r_{k-1}^3)) u_k \\ &= \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^4}{r_0^4} \tilde{L}_0^2 u_k + 4c_{k-1}^2 r_{k-1} r_0^{-2} (1 + q_{k-1} - r_{k-1}^3 r_0^{-3}) \tilde{L}_0^1 u_k \\ & \quad - 4c_{k-1}^2 r_{k-1} r_0^{-3} (1 + q_{k-1} - r_0^{-3} r_{k-1}^3) u_k - 2c_{k-1}^2 r_{k-1}^{-2} (1 + q_{k-1}) (1 + q_{k-1} - r_0^{-3} r_{k-1}^3) u_k \\ & \quad + 6c_{k-1}^2 r_{k-1} r_0^{-3} (r_0^{-3} r_{k-1}^3 - 1 - q_{k-1}) u_k + 2c_{k-1}^2 r_{k-1} \partial_x q_{k-1} u_k \\ &= \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^4}{r_0^4} \tilde{L}_0^2 u_k + 4c_{k-1}^2 r_{k-1} r_0^{-2} (1 + q_{k-1} - r_0^{-3} r_{k-1}^3) \tilde{L}_0^1 u_k + 2 \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^{k-1}}{r_0^2} L_0^1 q_{k-1} u_k \\ & \quad - c_{k-1}^2 (1 + q_{k-1} - r_0^{-3} r_{k-1}^3) (10r_0^{-3} r_{k-1} - 2r_{k-1}^{-2} (1 + q_{k-1})) u_k. \end{aligned} \quad (2.28)$$

$$\begin{aligned} L_{k-1}^2 q_k &= \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^4}{r_0^4} L_0^2 q_k + r_0^2 \partial_x \left(\frac{r_{k-1}^4}{r_0^4} \frac{c_{k-1}^2}{c_0^2} \right) L_0^1 q_k \\ &= \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^4}{r_0^4} L_0^2 q_k - (\gamma + 1) \frac{r_{k-1}^4}{r_0^2} \frac{(1 + q_{k-1})^{-\gamma-2}}{c_0^2} \partial_x q_{k-1} L_0^1 q_k \\ & \quad + 4 \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^{k-1}}{r_0^2} (1 + q_{k-1} - r_0^{-3} r_{k-1}^3) L_0^1 q_k, \end{aligned} \quad (2.29)$$

Hence by (2.26)-(2.29) and (2.12)-(2.15) for $k-1$ there holds

$$C^{-1} \|L_0^1 q_k\|_{L^2} \leq \|L_{k-1}^1 q_k\|_{L^2} \leq C \|L_0^1 q_k\|_{L^2}, \quad (2.30)$$

$$\|\tilde{L}_{k-1}^1 u_k - r_0^{-2} r_{k-1}^2 \tilde{L}_0^1 u_k\|_{L^2} \leq C \|u_k\|_{L^2}, \quad (2.31)$$

$$\left\| L_{k-1}^2 q_k - \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^4}{r_0^4} L_0^2 q_k \right\|_{L^2} \leq C (1 + \|L_0^1 q_{k-1}\|_{L^\infty}) \|L_0^1 q_k\|_{L^2}, \quad (2.32)$$

$$\left\| \tilde{L}_{k-1}^2 u_k - \frac{c_{k-1}^2}{c_0^2} \frac{r_{k-1}^4}{r_0^4} \tilde{L}_0^2 u_k \right\|_{L^2} \leq C (1 + \|L_0^1 q_{k-1}\|_{L^\infty}) \left(\|u\|_{L^2} + \|\tilde{L}_0^1 u_k\|_{L^2} \right). \quad (2.33)$$

Then using (2.30)-(2.33) in (2.24)(2.25) yields

$$\begin{aligned} & \|\partial_t^2 q_k|_{t=0}\|_{L^2} + \|\partial_t^2 u_k|_{t=0}\|_{L^2} \\ & \leq \|L_{k-1}^2 q_{in}\|_{L^2} + \|\tilde{L}_{k-1}^2 u_{in}\|_{L^2} + C (\|u_{in}\|_{L^\infty} + \|L_{k-1}^1 q_{in}\|_{L^\infty}) \left(\|\tilde{L}_{k-1}^1 u_{in}\|_{L^2} + \|L_{k-1}^1 q_{in}\|_{L^2} \right) \\ & \leq C (1 + \|L_0^1 q_{in}\|_{L^\infty} + \|u_{in}\|_{L^\infty}) \sum_{j=0}^2 \left(\|L_0^j q_{in}\|_{L^2} + \|\tilde{L}_0^j u_{in}\|_{L^2} \right). \end{aligned}$$

(2.30)(2.31) also shows the lower t -derivatives can be controlled by the same bound. It remains to bound $\partial_t^j f(R_k)|_{t=0}$, for which we have

$$\begin{aligned}
|\partial_t f(R_k)|_{t=0} &= f'(R_{k-1})|_{t=0} |u_k|_{(x=0,t=0)} = f'(R_{in}) |u_{in}|_{x=0} \leq C \left(\|\tilde{L}_0^1 u_{in}\|_{L^2} + \|u_{in}\|_{L^2} \right), \\
|\partial_t^2 f(R_k)|_{t=0} &= |f'(R_{k-1}) \partial_t u_k|_{(x=0,t=0)} + \partial_t (f'(R_{k-1})) u_k|_{(x=0,t=0)}| \\
&= \left| f'(R_{k-1}) L_{k-1}^1 q_k|_{(x=0,t=0)} + (f' \circ f^{-1})' (f(R_{k-1})) \partial_t f(R_{k-1}) u_k|_{(x=0,t=0)} \right| \\
&= \left| f'(R_{in}) \frac{c_{in}^2}{c_0^2} \frac{r_{in}^2}{r_0^2} L_0^1 q_{in} + (f' \circ f^{-1})' (f(R_{in})) f'(R_{in}) u_{in}^2|_{x=0} \right| \\
&\leq C \left(\|L_0^2 q_{in}\|_{L^2} + \|L_0^1 q_{in}\|_{L^2} \right) + C \|u_{in}\|_{L^\infty} \left(\|u_{in}\|_{L^2} + \|\tilde{L}_0^1 u_{in}\| \right).
\end{aligned}$$

Collecting the above bounds gives

$$\begin{aligned}
\sum_{j=0}^2 e_k^j(0) &\leq C \sum_{j=0}^2 \left(\|\partial_t^j q_k|_{t=0}\|_{L^2}^2 + \|\partial_t^j u_k|_{t=0}\|_{L^2}^2 + |\partial_t^j f(R_k)|_{t=0}|^2 \right) \\
&\leq C(1 + \|L_0^1 q_{in}\|_{L^\infty} + \|u_{in}\|_{L^\infty}) \left[\sum_{j=0}^2 \left(\|L_0^j q_k\|_{L^2}^2 + \|\tilde{L}_0^j u_k\|_{L^2}^2 \right) + |f(R_{in})|^2 \right].
\end{aligned}$$

Note that by the embedding (2.16)

$$\|L_0^1 q_{in}\|_{L^\infty} + \|u_{in}\|_{L^\infty} \leq C \left(\|L_0^2 q_{in}\|_{L^2} + \|L_0^1 q_{in}\|_{L^2} + \|u_{in}\|_{L^2} + \|\tilde{L}_0^1 u_{in}\|_{L^2} \right) \leq C\epsilon,$$

therefore, by choosing $\tilde{B} = C(1 + \epsilon)$ we arrive at (2.18). \square

The next step is to prove Proposition 2.4 for k , which is achieved by propagating e_k^j through an energy method and applying Lemma 2.6.

Proof of Proposition 2.4. Applying ∂_t^j to (2.4)-(2.7) yields

$$\begin{cases} \partial_t \partial_t^j u_k - c_{k-1}^2 r_{k-1}^2 \partial_x \partial_t^j q_k = [\partial_t^j, c_{k-1}^2 r_{k-1}^2 \partial_x] q_k, & x > 0, t > 0, \\ \partial_t \partial_t^j q_k - \partial_x (r_{k-1}^2 \partial_t^j u_k) = [\partial_t^j, \partial_x r_{k-1}^2] u_k, & x > 0, t > 0, \\ \partial_t^j q_k|_{x=0} = \partial_t^j f(R_k), & t > 0, \\ \partial_t^j u_k|_{x=0} = f'(R_{k-1})^{-1} \partial_t^{j+1} f(R_k) + [\partial_t^j, f'(R_{k-1})^{-1}] \partial_t f(R_k), & t > 0. \end{cases}$$

(2.3) gives

$$\begin{aligned}
&\frac{1}{2} \frac{d}{dt} \left[\int_0^\infty \left((\partial_t^j u_k)^2 + c_{k-1}^2 (\partial_t^j q_k)^2 \right) dx + (c_{k-1}^2 r_{k-1}^2)|_{x=0} f'(R_{k-1})^{-1} |\partial_t^j f(R_k)|^2 \right] \\
&= \int_0^\infty \left(\frac{1}{2} \partial_t c_{k-1}^2 (\partial_t^j q_k)^2 - r_{k-1}^2 \partial_x c_{k-1}^2 \partial_t^j u_k \partial_t^j q_k \right) dx \\
&\quad + \frac{1}{2} \partial_t \left[(c_{k-1}^2 r_{k-1}^2)|_{x=0} f'(R_{k-1})^{-1} \right] |\partial_t^j f(R_k)|^2 \\
&\quad + \int_0^\infty \left([\partial_t^j, c_{k-1}^2 r_{k-1}^2 \partial_x] q_k \partial_t^j u_k + c_{k-1}^2 [\partial_t^j, \partial_x r_{k-1}^2] u_k \partial_t^j q_k \right) dx \\
&\quad - (c_{k-1}^2 r_{k-1}^2)|_{x=0} [\partial_t^j, f'(R_{k-1})^{-1}] \partial_t f(R_k) \partial_t^j f(R_k).
\end{aligned} \tag{2.34}$$

For the first term on the right we compute that

$$\begin{aligned}
\left\| \frac{\partial_t c_{k-1}^2}{c_{k-1}^2} \right\|_{L^\infty} &\leq (\gamma + 1) \|(1 + q_{k-1})^{-1}\|_{L^\infty} \|\partial_t q_{k-1}\|_{L^\infty} \\
&\leq C (\|\partial_t q_{k-1}\|_{L^2} + \|L_{k-2}^1 \partial_t q_{k-1}\|_{L^2}) \\
&\leq CA^{\frac{1}{2}} \epsilon,
\end{aligned} \tag{2.35}$$

where we used $k-1$ version of Lemma 2.5 in the second and last inequalities, and in a same manner that

$$\begin{aligned}
\|c_{k-1}^{-1} r_{k-1}^2 \partial_x c_{k-1}^2\|_{L^\infty} &= \left(\frac{Ca\gamma}{2}\right)^{\frac{1}{2}} (\gamma + 1) \|(1 + q_{k-1})^{-\frac{\gamma+3}{2}} r_{k-1}^2 \partial_x q_{k-1}\|_{L^\infty} \\
&\leq \left(\frac{Ca\gamma}{2}\right)^{\frac{1}{2}} (\gamma + 1) \|(1 + q_{k-1})^{-\frac{\gamma+3}{2}} c_{k-2}^{-2} r_{k-2}^{-2} r_{k-1}^2\|_{L^\infty} \|L_{k-2}^1 q_{k-1}\|_{L^\infty} \\
&\leq C (\|L_{k-2}^1 q_{k-1}\|_{L^2} + \|L_{k-2}^2 q_{k-1}\|_{L^2}) \\
&\leq CA^{\frac{1}{2}} \epsilon,
\end{aligned} \tag{2.36}$$

For the second term, there holds

$$\begin{aligned}
|\partial_t [(c_{k-1}^2 r_{k-1}^2)|_{x=0} f'(R_{k-1})^{-1}]| &= \frac{Ca\gamma}{2} |\partial_t [R_{k-1}^2 (q_{k-1}|_{x=0} + 1)^{-\gamma-1} f'(R_{k-1})^{-1}]| \\
&= |\partial_t [R_{k-1}^2 (f(R_{k-1}) + 1)^{-\gamma-1} f'(R_{k-1})^{-1}]| \\
&\leq C |\partial_t R_{k-1}| \\
&\leq CA^{\frac{1}{2}} \epsilon.
\end{aligned} \tag{2.37}$$

The first inequality holds since $R^2(f(R) + 1)^{-\gamma-1} f'(R)^{-1}$ is smooth for R in the range given by (2.14). Therefore, the first and second terms are bounded by $CA^{\frac{1}{2}} \epsilon e_k^j(t)$. For the commutators we compute and introduce the notations $NL_{\alpha,k}^j$, $\alpha = 1, 2, 3, 4, 5$ as follows

$$\begin{aligned}
[\partial_t^j, c_{k-1}^2 r_{k-1}^2 \partial_x] q_k \partial_t^j u_k &= \sum_{i=1}^j \binom{j}{i} \partial_t^i (r_{k-1}^2 c_{k-1}^2) \partial_t^{j-i} \partial_x q_k \partial_t^j u_k \\
&= \sum_{i=1}^j \binom{j}{i} \partial_t^i (r_{k-1}^2 c_{k-1}^2) \partial_t^{j-i} (c_{k-1}^{-2} r_{k-1}^{-2} \partial_t u_k) \partial_t^j u_k \\
&= \sum_{i=1}^j \sum_{l=0}^{j-i} \binom{j}{i} \binom{j-i}{l} \partial_t^i (r_{k-1}^2 c_{k-1}^2) \partial_t^l (c_{k-1}^{-2} r_{k-1}^{-2}) \partial_t^{j+1-i-l} u_k \partial_t^j u_k \\
&=: NL_{1,k}^j,
\end{aligned}$$

$$\begin{aligned}
c_{k-1}^2[\partial_t^j, \partial_x r_{k-1}^2]u_k \partial_t^j q_k &= c_{k-1}^2 \sum_{i=1}^j \binom{j}{i} \left(\partial_x \partial_t^i r_{k-1}^2 \partial_t^{j-i} u_k \partial_t^j q_k + \partial_t^i r_{k-1}^2 \partial_t^{j-i} \partial_x u_k \partial_t^j q_k \right) \\
&= c_{k-1}^2 \sum_{i=1}^j \binom{j}{i} \partial_x \partial_t^i r_{k-1}^2 \partial_t^{j-i} u_k \partial_t^j q_k \\
&\quad + c_{k-1}^2 \sum_{i=1}^j \binom{j}{i} \partial_t^i r_{k-1}^2 \partial_t^{j-i} (r_{k-1}^{-2} \partial_t q_k + \partial_x (r_{k-1}^{-2}) r_{k-1}^2 u_k) \partial_t^j q_k \\
&= c_{k-1}^2 \sum_{i=1}^j \binom{j}{i} \partial_x \partial_t^i r_{k-1}^2 \partial_t^{j-i} u_k \partial_t^j q_k \\
&\quad + c_{k-1}^2 \sum_{i=1}^j \binom{j}{i} \partial_t^i r_{k-1}^2 \sum_{l=0}^{j-i} \binom{j-i}{l} \partial_t^l (r_{k-1}^{-2}) \partial_t^{j+1-i-l} q_k \partial_t^j q_k \\
&\quad + c_{k-1}^2 \sum_{i=1}^j \binom{j}{i} \partial_t^i r_{k-1}^2 \sum_{l=0}^{j-i} \binom{j-i}{l} \partial_t^l (r_{k-1}^2 \partial_x (r_{k-1}^{-2})) \partial_t^{j-i-l} u_k \partial_t^j q_k \\
&=: NL_{2,k}^j + NL_{3,k}^j + NL_{4,k}^j, \\
NL_{5,k}^j &:= (c_{k-1}^2 r_{k-1}^2) |_{x=0} [\partial_t^j, f'(R_{k-1})^{-1}] \partial_t f(R_k) \partial_t^j f(R_k) \\
&= \sum_{i=1}^j \binom{j}{i} (c_{k-1}^2 r_{k-1}^2) |_{x=0} \partial_t^i (f'(R_{k-1})^{-1}) \partial_t^{j+1-i} f(R_k) \partial_t^j f(R_k).
\end{aligned}$$

• $NL_{1,k}^j$ **terms.** We expand the coefficients as

$$\begin{aligned}
\partial_t (r_{k-1}^2 c_{k-1}^2) &= 2c_{k-1}^2 r_{k-1} u_{k-1} - (\gamma + 1) c_{k-1}^2 (1 + q_{k-1})^{-1} r_{k-1}^2 \partial_t q_{k-1}, \\
\partial_t^2 (r_{k-1}^2 c_{k-1}^2) &= 2c_{k-1}^2 r_{k-1} \partial_t u_{k-1} + 2c_{k-1}^2 u_{k-1}^2 + (\gamma + 1)(\gamma + 2) c_{k-1}^2 (1 + q_{k-1})^{-2} r_{k-1}^2 (\partial_t q_{k-1})^2 \\
&\quad - (\gamma + 1) c_{k-1}^2 (1 + q_{k-1})^{-1} r_{k-1}^2 \partial_t^2 q_{k-1} - 4(\gamma + 1) c_{k-1}^2 (1 + q_{k-1})^{-1} r_{k-1} u_{k-1} \partial_t q_{k-1}, \\
\partial_t (c_{k-1}^{-2} r_{k-1}^{-2}) &= -2c_{k-1}^{-2} r_{k-1}^{-3} u_{k-1} + (\gamma + 1) c_{k-1}^{-2} (1 + q_{k-1})^{-1} r_{k-1}^{-2} \partial_t q_{k-1}.
\end{aligned}$$

Hence except for the case $(i, l) = (2, 0)$ there holds

$$\begin{aligned}
|\partial_t^i (r_{k-1}^2 c_{k-1}^2) \partial_t^l (c_{k-1}^{-2} r_{k-1}^{-2})| &\leq C (1 + |u_{k-1}| + |\partial_t q_{k-1}|) (|u_{k-1}| + |\partial_t q_{k-1}|) \\
&\leq CA^{\frac{1}{2}} \epsilon (1 + A^{\frac{1}{2}} \epsilon),
\end{aligned}$$

while for $(i, l) = (2, 0)$

$$\begin{aligned}
&|c_{k-1}^{-2} r_{k-1}^{-2} (\partial_t^2 (r_{k-1}^2 c_{k-1}^2) + (\gamma + 1) c_{k-1}^2 (1 + q_{k-1})^{-1} r_{k-1}^2 \partial_t^2 q_{k-1})| \\
&\leq C (|\partial_t u_{k-1}| + |u_{k-1}|^2 + |\partial_t q_{k-1}|^2) \\
&\leq CA^{\frac{1}{2}} \epsilon (1 + A^{\frac{1}{2}} \epsilon),
\end{aligned}$$

and that

$$\begin{aligned}
& \left| \int_0^\infty (\gamma + 1)(1 + q_{k-1})^{-1} \partial_t^2 q_{k-1} \partial_t u_k \partial_t^2 u_k dx \right| \\
& \leq C \|\partial_t^2 q_{k-1}\|_{L^2} \|\partial_t^2 u_k\|_{L^2} \|\partial_t u_k\|_{L^\infty} \\
& \leq CA^{\frac{1}{2}} \epsilon \|\partial_t^2 u_k\|_{L^2} \left(\|\partial_t u_k\|_{L^2} + \|\tilde{L}_0^1 \partial_t u_k\|_{L^2} \right) \\
& \leq CA^{\frac{1}{2}} \epsilon \|\partial_t^2 u_k\|_{L^2} \left(\|\partial_t u_k\|_{L^2} + \|\partial_t^2 q_k\|_{L^2} + CA^{\frac{1}{2}} \epsilon (\|\partial_t q_k\|_{L^2} + \|u_k\|_{L^2}) \right),
\end{aligned}$$

where in the last step we used (2.22). Collecting the above bounds for all possible pairs (i, l) we obtain

$$\int_0^\infty |NL_{1,k}^1| dx \leq CA^{\frac{1}{2}} \epsilon \|\partial_t u_k\|_{L^2}^2, \quad (2.38)$$

$$\begin{aligned}
\int_0^\infty |NL_{1,k}^2| dx & \leq CA^{\frac{1}{2}} \epsilon (1 + A^{\frac{1}{2}} \epsilon) \|\partial_t^2 u_k\|_{L^2} \left(\|\partial_t u_k\|_{L^2} + \|\partial_t^2 u_k\|_{L^2} + \|\partial_t^2 q_k\|_{L^2} \right) \\
& \quad + CA \epsilon^2 \|\partial_t^2 u_k\|_{L^2} \left(\|\partial_t q_k\|_{L^2} + \|u_k\|_{L^2} \right).
\end{aligned} \quad (2.39)$$

- $NL_{2,k}^j$ **terms.** Expand the coefficients $\partial_x \partial_t^i r_{k-1}^2$ as follows.

$$\begin{aligned}
\partial_x \partial_t r_{k-1}^2 & = 2r_{k-1}^{-1} \partial_t q_{k-1} - 2(1 + q_{k-1}) r_{k-1}^{-2} u_{k-1}, \\
\partial_x \partial_t^2 r_{k-1}^2 & = 2\partial_t \left(r_{k-1}^{-1} \partial_t q_{k-1} - (1 + q_{k-1}) r_{k-1}^{-2} u_{k-1} \right) \\
& = 2r_{k-1}^{-1} \partial_t^2 q_{k-1} - 4r_{k-1}^{-2} u_{k-1} \partial_t q_{k-1} \\
& \quad - 2(1 + q_{k-1}) r_{k-1}^{-2} \partial_t u_{k-1} + 4(1 + q_{k-1}) r_{k-1}^{-3} u_{k-1}^2.
\end{aligned}$$

Using the bounds in Lemma 2.5 and the induction hypothesis we obtain

$$\begin{aligned}
\int_0^\infty |NL_{2,k}^1| dx & \leq C \left(\|\partial_t q_{k-1}\|_{L^\infty} + \|u_{k-1}\|_{L^\infty} \right) \|u_k\|_{L^2} \|\partial_t q_k\|_{L^2} \\
& \leq CA^{\frac{1}{2}} \epsilon \|u_k\|_{L^2} \|\partial_t q_k\|_{L^2},
\end{aligned} \quad (2.40)$$

$$\begin{aligned}
\int_0^\infty |NL_{2,k}^2| dx & \leq C \left(\|\partial_t q_{k-1}\|_{L^\infty} + \|\partial_t u_{k-1}\|_{L^\infty} \right) \|\partial_t u_k\|_{L^2} \|\partial_t^2 q_k\|_{L^2} \\
& \quad + C \|\partial_t^2 q_{k-1}\|_{L^2} \|u_k\|_{L^\infty} \|\partial_t^2 q_k\|_{L^2} \\
& \quad + C \left(\|u_{k-1}\|_{L^\infty} \|\partial_t q_{k-1}\|_{L^\infty} + \|\partial_t u_{k-1}\|_{L^\infty} + \|u_{k-1}\|_{L^2} \right) \|u_k\|_{L^2} \|\partial_t^2 q_k\|_{L^2} \\
& \leq CA^{\frac{1}{2}} \epsilon (1 + A^{\frac{1}{2}} \epsilon) \left(\|\partial_t u_k\|_{L^2} + \|u_k\|_{L^\infty} + \|u_k\|_{L^2} \right) \|\partial_t^2 q_k\|_{L^2} \\
& \leq CA^{\frac{1}{2}} \epsilon (1 + A^{\frac{1}{2}} \epsilon) \left(\|\partial_t u_k\|_{L^2} + \|\partial_t q_k\|_{L^2} + \|u_k\|_{L^2} \right) \|\partial_t^2 q_k\|_{L^2}.
\end{aligned} \quad (2.41)$$

- $NL_{3,k}^j$ **terms.** Note that

$$\begin{aligned}
|\partial_t(r_{k-1}^{-2})| & = |-2r_{k-1}^{-3} u_{k-1}| \leq CA^{\frac{1}{2}} \epsilon r_{k-1}^{-3}, \\
|\partial_t(r_{k-1}^2)| & = 2|r_{k-1} u_{k-1}| \leq CA^{\frac{1}{2}} \epsilon r_{k-1}, \\
|\partial_t^2(r_{k-1}^2)| & = |2u_{k-1}^2 + 2r_{k-1} \partial_t u_{k-1}| \leq CA^{\frac{1}{2}} \epsilon (1 + A^{\frac{1}{2}} \epsilon) r_{k-1},
\end{aligned}$$

therefore collecting the bounds of $|\partial_t^i(r_{k-1}^2)\partial_t^l(r_{k-1}^{-2})|$ for each possible pair (i, l) gives

$$\int_0^\infty |NL_{3,k}^1| dx \leq CA^{\frac{1}{2}}\epsilon \|\partial_t q_k\|_{L^2}^2, \quad (2.42)$$

$$\int_0^\infty |NL_{3,k}^2| dx \leq CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon) (\|\partial_t q_k\|_{L^2} + \|\partial_t^2 q_k\|_{L^2}) \|\partial_t^2 q_k\|_{L^2}. \quad (2.43)$$

• **$NL_{4,k}^j$ terms.** The bounds of $|\partial_t^i r_{k-1}^2|$ has been obtained in the previous step. We compute for $\partial_t^l(r_{k-1}^2 \partial_x(r_{k-1}^{-2}))$ that

$$|r_{k-1}^2 \partial_x(r_{k-1}^{-2})| = |-2r_{k-1}^{-3}(1 + q_{k-1})| \leq Cr_{k-1}^{-3},$$

$$|\partial_t(r_{k-1}^2 \partial_x(r_{k-1}^{-2}))| = |-2r_{k-1}^{-3} \partial_t q_{k-1} + 6r_{k-1}^{-4} u_{k-1}(1 + q_{k-1})| \leq CA^{\frac{1}{2}}\epsilon r_{k-1}^{-3}.$$

Hence

$$\int_0^\infty |NL_{4,k}^1| dx \leq CA^{\frac{1}{2}}\epsilon \|u_k\|_{L^2} \|\partial_t q_k\|_{L^2}, \quad (2.44)$$

$$\int_0^\infty |NL_{4,k}^2| dx \leq CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon) (\|u_k\|_{L^2} + \|\partial_t u_k\|_{L^2}) \|\partial_t^2 q_k\|_{L^2}. \quad (2.45)$$

• **$NL_{5,k}^j$ terms.** Write $(f'(R_{k-1}))^{-1}$ as $\left(\frac{1}{f'} \circ f^{-1}\right) \circ f(R_{k-1})$, then

$$\partial_t (f'(R_{k-1})^{-1}) = \left(\frac{1}{f'} \circ f^{-1}\right)' (f(R_{k-1})) \partial_t f(R_{k-1}),$$

$$\partial_t^2 (f'(R_{k-1})^{-1}) = \left(\frac{1}{f'} \circ f^{-1}\right)'' (f(R_{k-1})) (\partial_t f(R_{k-1}))^2 + \left(\frac{1}{f'} \circ f^{-1}\right)' (f(R_{k-1})) \partial_t^2 f(R_{k-1}).$$

Since f is smooth for R within the range given by (2.15), the last term can be estimate as follows.

$$|NL_{5,k}^1| \leq CA^{\frac{1}{2}}\epsilon |\partial_t f(R_k)|^2, \quad (2.46)$$

$$|NL_{5,k}^2| \leq CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon) (|\partial_t f(R_k)| + |\partial_t^2 f(R_k)|) |\partial_t^2 f(R_k)|. \quad (2.47)$$

• **Final estimate of e_k^j .** Recall the energy estimate (2.34), and we collect the bounds (2.35)-(2.47) for each term on the right hand side and obtain

$$\frac{d}{dt} e_k^0(t) \leq CA^{\frac{1}{2}}\epsilon e_k^0(t),$$

$$\frac{d}{dt} e_k^1(t) \leq CA^{\frac{1}{2}}\epsilon (e_k^0(t) + e_k^1(t)),$$

$$\frac{d}{dt} e_k^2(t) \leq CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon) (e_k^0(t) + e_k^1(t) + e_k^2(t)).$$

Hence $\sum_{j=0}^2 e_k^j(t) \leq \sum_{j=0}^2 e_k^j(0) \exp\left\{CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon)t\right\}$, and by Lemma 2.6

$$E_k(t) \leq B\tilde{B} \exp\left\{CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon)t\right\} \epsilon^2 = C(1 + \epsilon)(1 + A^{\frac{1}{2}}\epsilon) \exp\left\{CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon)t\right\} \epsilon^2.$$

Now we choose A large enough such that

$$A \geq 2C(1 + \epsilon)(1 + A^{\frac{1}{2}}\epsilon),$$

and choose a small T satisfying

$$\exp \left\{ CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon)t \right\} = 2,$$

then for all $t \in [0, T]$ there holds $E_k(t) \leq A\epsilon^2$, which is (2.11). \square

Remark 2.7. From the proof above we see that the restriction on T are

$$\exp \left\{ CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon)T \right\} \leq 2, \quad CA^{\frac{1}{2}}T \leq \min \left\{ \frac{q+1}{2}, \frac{f(R_{in})+1}{2} \right\}.$$

Hence the maximal lifespan admits a lower bound in forms of

$$\min \left\{ \left(CA^{\frac{1}{2}}\epsilon(1 + A^{\frac{1}{2}}\epsilon) \right)^{-1}, \frac{q+1}{CA^{\frac{1}{2}}}, \frac{f(R_{in})+1}{CA^{\frac{1}{2}}} \right\}.$$

The last step of closing the induction is to prove Lemma 2.5 for k .

Proof of Lemma 2.5. By (2.34), the following holds:

$$\begin{aligned} |q_k(x, t) - q_k(x, 0)| &\leq \int_0^t \|\partial_t q_k(s)\|_{L^\infty} ds \\ &\leq C \int_0^t (\|\partial_t q_k(s)\|_{L^2} + \|L_{k-1}^1 \partial_t q_k(s)\|_{L^2}) ds \\ &\leq CA^{\frac{1}{2}}\epsilon t. \end{aligned}$$

Hence by choosing T small, we have for $t \in [0, T]$

$$\frac{q-1}{2} = \underline{q} - \frac{q+1}{2} \leq q_k(x, 0) - CA^{\frac{1}{2}}\epsilon t \leq q_k(x, t) \leq q_k(x, 0) + CA^{\frac{1}{2}}\epsilon t \leq 2\bar{q} + 1,$$

which is (2.12). (2.13) is a consequence of (2.12) by the construction of c_k . In a same manner,

$$|f(R_k)(t) - f(R_{in})| \leq \int_0^t |\partial_t f(R_k)(s)| ds \leq A^{\frac{1}{2}}\epsilon t,$$

$$\frac{f(R_{in})-1}{2} = f(R_{in}) - \frac{f(R_{in})+1}{2} \leq f(R_{in}) - A^{\frac{1}{2}}\epsilon t \leq f(R_k)(t) \leq f(R_{in}) + A^{\frac{1}{2}}\epsilon t \leq 2f(R_{in}) + 1.$$

Hence we obtain the first part of (2.14), and the other part as well as (2.15) is a consequence by applying f^{-1} and the construction of r_k . (2.16) is given by Lemma 2.1 since $r_k(x, t) \geq R_k(x, t) \geq f^{-1} \left(\frac{f(R_{in})-1}{2} \right)$ and $r_k^2 \partial_x r_k = 1 + q_k \in [\frac{1}{2}(\underline{q} + 1), 2(\bar{q} + 1)]$. \square

At this point the k step of induction on Proposition 2.4, Lemma 2.5, Lemma 2.6 has been finished, and therefore their proof is complete. A direct corollary is the uniform bound of $(\partial_t^j u_k, \partial_t^j q_k)$ in $L^\infty([0, T]; H^{2-j})$.

Corollary 2.8 (Uniform bounds on H^2). *There exists $T > 0$ such that for all $k \geq 0$ and $t \in [0, T]$, $j = 0, 1, 2$ that*

$$\|\partial_t^j u_k(t)\|_{H^{2-j}} + \|\partial_t^j q_k(t)\|_{H^{2-j}} \lesssim A^{\frac{1}{2}}(1 + A^{\frac{1}{2}}\epsilon)\epsilon,$$

and $\tilde{u}_k(\xi, t) := u_k\left(\frac{\xi^3-1}{3}, t\right)$, $\tilde{q}_k(\xi, t) := q_k\left(\frac{\xi^3-1}{3}, t\right)$ satisfy

$$\|\partial_t^j \tilde{u}_k(t)\|_{H_\xi^{2-j}(1, +\infty)} + \|\partial_t^j \tilde{q}_k(t)\|_{H_\xi^{2-j}(1, +\infty)} \lesssim A^{\frac{1}{2}}(1 + A^{\frac{1}{2}}\epsilon)\epsilon.$$

Proof. The case $j = 2$ is obvious. Using inequalities (2.30)-(2.33), (2.11) and the bounds in Lemma 2.5 for c_k and r_k , we obtain

$$\|q_k(t)\|_{H^2} + \|u_k(t)\|_{H^2} \lesssim \sum_{j=0}^2 \left(\|L_0^j q_k(t)\|_{L^2} + \|\tilde{L}_0^j u_k(t)\|_{L^2} \right) \lesssim A^{\frac{1}{2}}(1 + A^{\frac{1}{2}}\epsilon)\epsilon.$$

Since $L_0^1 \partial_t q_k = \frac{c_0^2}{c_{k-1}^2} \frac{r_0^2}{r_{k-1}^2} L_{k-1}^1 \partial_t q_k$ and $\frac{r_{k-1}^2}{r_0^2} \tilde{L}_0^1 \partial_t u_k = \tilde{L}_{k-1}^1 \partial_t u_k + 2r_{k-1}^{-1}(r_0^{-3} r_{k-1}^3 - 1 - q_{k-1})u_k$, there holds

$$\|\partial_t q_k(t)\|_{H^1} + \|\partial_t u_k(t)\|_{H^1} \lesssim \sum_{j=0}^1 \left(\|L_0^j \partial_t q_k(t)\|_{L^2} + \|\tilde{L}_0^j \partial_t u_k(t)\|_{L^2} \right) \lesssim A^{\frac{1}{2}}\epsilon.$$

The bounds on \tilde{u}_k and \tilde{q}_k follow by a change of variable. \square

Next, we construct the solution to (1.9)-(1.14). For arbitrary $M > 1$, since we have the compact embedding $H_\xi^2(1, M) \subset\subset H_\xi^1(1, M) \subset\subset L_\xi^2(1, M)$, applying Aubin-Lions compactness theorem twice gives that there exists a subsequence, still denoted by $\{\tilde{u}_k\}$, $\{\tilde{q}_k\}$, and limit points \tilde{u} , $\tilde{q} \in L^\infty([0, T], H_\xi^2(1, M))$ such that

$$\tilde{u}_k \rightharpoonup \tilde{u} \text{ weakly* in } L^\infty([0, T]; H_\xi^2(1, M)), \quad \tilde{u}_k \rightarrow \tilde{u} \text{ strongly in } C([0, T]; H_\xi^1(1, M)), \quad (2.48)$$

$$\tilde{q}_k \rightharpoonup \tilde{q} \text{ weakly* in } L^\infty([0, T]; H_\xi^2(1, M)), \quad \tilde{q}_k \rightarrow \tilde{q} \text{ strongly in } C([0, T]; H_\xi^1(1, M)), \quad (2.49)$$

$$\partial_t \tilde{u}_k \rightharpoonup \partial_t \tilde{u} \text{ weakly* in } L^\infty([0, T]; H_\xi^1(1, M)), \quad \partial_t \tilde{u}_k \rightarrow \partial_t \tilde{u} \text{ strongly in } C([0, T]; L_\xi^2(1, M)), \quad (2.50)$$

$$\partial_t \tilde{q}_k \rightharpoonup \partial_t \tilde{q} \text{ weakly* in } L^\infty([0, T]; H_\xi^1(1, M)), \quad \partial_t \tilde{q}_k \rightarrow \partial_t \tilde{q} \text{ strongly in } C([0, T]; L_\xi^2(1, M)). \quad (2.51)$$

Moreover, using a diagonal argument and further passing to a subsequence, we can assume that (2.48)-(2.51) hold for each $M > 1$. Meanwhile, applying Azela-Ascoli theorem on $\{f(R_k)\}$ and $\{\partial_t f(R_k)\}$ yields the existence of a subsequence of $\{f(R_k)\}$, still denoted by $\{f(R_k)\}$, and a limit point \tilde{f} such that

$$f(R_k) \rightarrow \tilde{f} \text{ in } C^1[0, T].$$

Then we construct the solution (u, q, R) as follows.

$$R(t) := f^{-1} \tilde{f}(t), \quad r^3(x, t) := R^3(t) + 3x + 3 \int_0^x q(y, t) dy,$$

$$u(x, t) := \tilde{u}((1 + 3x)^{\frac{1}{2}}, t), \quad q(x, t) := \tilde{q}((1 + 3x)^{\frac{1}{3}}, t).$$

To verify (u, q, R) is a solution to (1.9)-(1.14), we denote $\tilde{r}_k(\xi, t) := r_k\left(\frac{\xi^3-1}{3}, t\right)$, and $\tilde{r}(\xi, t) := r\left(\frac{\xi^3-1}{3}, t\right)$. First note that the construction of r is exactly (1.13). From Sobolev embedding and the following inequality

$$\left|\frac{\tilde{r}_k^3 - \tilde{r}^3}{\xi^3}\right| \leq \xi^{-3}|R_k^3 - R^3| + 3\xi^{-3} \int_1^\xi |\tilde{q}_k - \tilde{q}|\eta^2 d\eta \leq \xi^{-3}|R_k^3 - R^3| + (1 - \xi^{-3})\|\tilde{q}_k - \tilde{q}\|_{L^\infty},$$

it follows that $\xi^{-2}\tilde{r}_k^2$ converges strongly to $\xi^{-2}\tilde{r}^2$ in L_ξ^∞ due to (2.49), and (2.49) also implies $(1 + \tilde{q}_k)^{-\gamma-1}$ converges strongly to $(1 + \tilde{q})^{-\gamma-1}$ in L_ξ^∞ . By the equation (2.5), we see that \tilde{u}_k, \tilde{q}_k satisfy

$$\partial_t \tilde{u}_k = c_0^2(1 + \tilde{q}_{k-1})^{-\gamma-1} \frac{\tilde{r}_{k-1}^2}{\xi^2} \partial_\xi \tilde{q}_k.$$

Forcing k to infinity and using (2.49)(2.50) yields (1.10). Next, (2.4) is equivalent to

$$\partial_t \tilde{q}_k = \xi^{-2} \partial_\xi (\tilde{r}_{k-1}^2 \tilde{u}_k) = \frac{\tilde{r}_{k-1}^2}{\xi^2} \partial_\xi \tilde{u}_k + \frac{2}{\tilde{r}_{k-1}} (1 + \tilde{q}_{k-1}) \tilde{u}_k.$$

Forcing k to infinity and using (2.48)(2.51) yields (1.9). For boundary conditions, forcing k to infinity in the equation

$$(\tilde{u}_k, \tilde{q}_k)|_{\xi=1} = (u_k, q_k)|_{x=0} = (f'(R_{k-1}) \frac{d}{dt} f(R_k), f(R_k))$$

and applying trace theorem yields $(u, q)|_{x=0} = (\tilde{u}, \tilde{q})|_{\xi=1} = (\frac{dR}{dt}, f(R))$, which is (1.11)(1.12). Last, (1.14) holds since $\tilde{u}_k(0), \tilde{q}_k(0), f(R_k)(0)$ converge to $\tilde{u}(0), \tilde{q}(0), f(R)(0)$ in $H_\xi^1(1, M)$ or \mathbb{R} as $k \rightarrow +\infty$ and $(\tilde{u}_k(0), \tilde{q}_k(0), f(R_k)(0)) = (\tilde{u}_{in}, \tilde{q}_{in}, f(R_{in}))$ independent of k . Therefore, we conclude that (u, q, R) is a solution to (1.9)-(1.14) with the wanted regularity of Theorem 1.2. Last, the uniqueness can be shown in a standard way by taking difference.

Proposition 2.9 (Uniqueness). *Let (u_i, q, R_i) , $i = 1, 2$ be two solutions to (1.9)-(1.14) with $u_i, q_i \in L^\infty([0, T]; H^2) \cap C([0, T]; H^1)$, $R_i \in C^1([0, T])$ and $-1 < \underline{q} \leq q_i \leq \bar{q} < +\infty$, $0 < R_i < \bar{R}$, then $(u_1, q_1, R_1) = (u_2, q_2, R_2)$.*

Proof. By taking difference, we obtain

$$\partial_t(u_1 - u_2) = c_0^2(1 + q_1)^{-\gamma-1} r_1^2 \partial_x(q_1 - q_2) + c_0^2[(1 + q_{k-1})^{-\gamma-1} r_1^2 - (1 + q_2)^{-\gamma-1} r_2^2] \partial_x q_2,$$

$$\partial_t(q_1 - q_2) = \partial_x[r_1^2(u_1 - u_2) + (r_1^2 - r_2^2)u_2],$$

$$\frac{d}{dt}(f(R_1) - f(R_2)) = f'(R_1)(u_1 - u_2)|_{x=0} + (f'(R_1) - f'(R_2))u_2|_{x=0},$$

$$(q_1 - q_2)|_{x=0} = f(R_1) - f(R_2).$$

Multiplying the first two equations by $(u_1 - u_2)$ and $(q_1 - q_2)$ respectively and integrating by parts

yields

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \left[\int_0^\infty ((u_1 - u_2)^2 + c_0^2 (1 + q_1)^{-\gamma-1} (q_1 - q_2)^2) dx \right] \\
& + \frac{1}{2} \frac{d}{dt} [c_0^2 R_1^2 (1 + q_1)^{-\gamma-1} |_{x=0} (f'(R_1))^{-1} (f(R_1) - f(R_2))^2] \\
= & \int_0^\infty \left[\frac{c_0^2}{2} \partial_t (1 + q_1)^{-\gamma-1} (q_1 - q_2)^2 - c_0^2 r_1^2 \partial_x (1 + q_1)^{-\gamma-1} (u_1 - u_2) (q_1 - q_2) \right] dx \\
& + \int_0^\infty c_0^2 [(1 + q_1)^{-\gamma-1} r_1^2 - (1 + q_2)^{-\gamma-1} r_2^2] \partial_x q_2 (u_1 - u_2) dx \\
& + \int_0^\infty c_0^2 (1 + q_1)^{-\gamma-1} \partial_x ((r_1^2 - r_2^2) u_2) (q_1 - q_2) dx \\
& + \frac{c_0^2}{2} \partial_t [(1 + q_1)^{-\gamma-1} |_{x=0} R_1^2 (f'(R_1))^{-1}] (f(R_1) - f(R_2))^2 \\
& + c_0^2 (1 + q_1)^{-\gamma-1} |_{x=0} R_1^2 (f'(R_1))^{-1} (f'(R_1) - f'(R_2)) u_2 |_{x=0} (f(R_1) - f(R_2)).
\end{aligned} \tag{2.52}$$

To control the source terms on the right, write

$$\frac{r_2}{r_1} - 1 = \frac{r_2^3 - r_1^3}{r_1(r_2^2 + r_1 r_2 + r_1^2)} = [r_1(r_2^2 + r_2 r_1 + r_1^2)]^{-1} \left[R_2^3 - R_1^3 + \int_0^x (q_2 - q_1)(y, t) dy \right].$$

From the $L^2 - L^2$ boundness of the maximal functional, it follows

$$\begin{aligned}
\left\| \frac{r_2}{r_1} - 1 \right\|_{L^2} & \leq C |f(R_2) - f(R_1)| + \left\| \frac{x}{r_1(r_2^2 + r_2 r_1 + r_1^2)} \right\|_{L^\infty} \left\| \frac{1}{x} \int_0^x (q_2 - q_1)(y, t) dy \right\|_{L^2} \\
& \leq C (|f(R_2) - f(R_1)| + \|q_2 - q_1\|_{L^2}),
\end{aligned}$$

and similarly

$$\left\| \frac{r_1}{r_2} - 1 \right\|_{L^2} \leq C (|f(R_2) - f(R_1)| + \|q_1 - q_2\|_{L^2}).$$

The first term is bounded by multiple of $\|q_1 - q_2\|_{L^2} + \|u_1 - u_2\|_{L^2}$ since $\partial_t q_1 = \partial_x (r_1^2 u_1)$ and $\|\partial_x (r_1^2 u_1)\|_{L^\infty} + \|r_1^2 \partial_x q_1\|_{L^\infty} \leq \|u_1\|_{H^2} + \|q_1\|_{H^2} \leq C$. For the second term on the right, we compute

$$\begin{aligned}
& (1 + q_1)^{-\gamma-1} \frac{r_1^2}{r_2^2} - (1 + q_2)^{-\gamma-1} \\
= & (q_1 - q_2) \frac{(1 + q_1)^{-\gamma-1} - (1 + q_2)^{-\gamma-1}}{q_1 - q_2} + (1 + q_1)^{-\gamma-1} \left(\frac{r_1}{r_2} - 1 \right) \left(\frac{r_1}{r_2} + 1 \right),
\end{aligned}$$

and thus by Sobolev embedding

$$\begin{aligned}
\left\| (1 + q_1)^{-\gamma-1} \frac{r_1^2}{r_2^2} - (1 + q_2)^{-\gamma-1} \right\|_{L^2} & \leq C (|f(R_1) - f(R_2)| + \|q_1 - q_2\|_{L^2}), \\
\left| \int_0^\infty c_0^2 [(1 + q_1)^{-\gamma-1} r_1^2 - (1 + q_2)^{-\gamma-1} r_2^2] \partial_x q_2 (u_1 - u_2) dx \right| \\
& \leq C \|r_2^2 \partial_x q_2\|_{L^\infty} (|f(R_1) - f(R_2)| + \|q_1 - q_2\|_{L^2}) \|u_1 - u_2\|_{L^2} \\
& \leq C (|f(R_1) - f(R_2)| + \|q_1 - q_2\|_{L^2}) \|u_1 - u_2\|_{L^2}.
\end{aligned}$$

For the third term, there holds

$$\partial_x[(r_1^2 - r_2^2)u_2] = 2r_1^{-1}(q_1 - q_2)u_2 + 2(r_1^{-1} - r_2^{-1})(1 + q_2)u_2 + \left(1 - \frac{r_1^2}{r_2^2}\right)r_2^2\partial_x u_2,$$

and thus

$$\begin{aligned} & \|\partial_x[(r_1^2 - r_2^2)u_2]\|_{L^2} \\ & \leq C \left(\left\| \frac{u_2}{r_1} \right\|_{L^\infty} (1 + \|q_2\|_{L^\infty}) + (1 + \|r_2^{-1}r_1\|_{L^\infty}) \|r_2^2\partial_x u_2\|_{L^\infty} \right) (|f(R_1) - f(R_2)| + \|q_1 - q_2\|_{L^2}) \\ & \leq C (|f(R_1) - f(R_2)| + \|q_1 - q_2\|_{L^2}). \end{aligned}$$

Since $q_1|_{x=0} = f(R_1)$ is a C^1 function and f is invertible, the last two boundary terms can be controlled by multiple of $|f(R_1) - f(R_2)|^2$. Therefore, we conclude from (2.52) that

$$\frac{d}{dt}e \leq C (\|u_1 - u_2\|_{L^2}^2 + \|q_1 - q_2\|_{L^2}^2 + |f(R_1) - f(R_2)|^2) \leq Ce$$

where we denote

$$\begin{aligned} e := & \int_0^\infty ((u_1 - u_2)^2 + c_0^2(1 + q_1)^{-\gamma-1}(q_1 - q_2)^2) dx \\ & + c_0^2 R_1^2 (1 + q_1)^{-\gamma-1}|_{x=0} (f'(R_1))^{-1} (f(R_1) - f(R_2))^2. \end{aligned}$$

Gronwall's inequality then gives $e = 0$ on $[0, T]$, which shows $(u_1, q_1, R_1) = (u_2, q_2, R_2)$. \square

3. Setup of Bootstrap

The following several sections are devoted to the proof of Theorem 1.3. The strategy is to use a bootstrap argument encompassing the norms in energy spaces and parameters. In this section we detail the set up of the bootstrap.

We inherit the notations from Section 2 and drop the subscript k :

$$e^j(t) := \frac{1}{2} \int_0^\infty \left(\left(\partial_t^j u(t) \right)^2 + c^2(t) \left(\partial_t^j q(t) \right)^2 \right) dx + \frac{1}{2} c(t)|_{x=0}^2 f'(R(t))^{-1} \left(\partial_t^j f(R)(t) \right)^2,$$

$$E(t) := \sum_{j=0}^2 |\partial_t^j f(R)(t)|^2 + \sum_{j_1+j_2 \leq 2} \left(\|L^{j_2} \partial_t^{j_1} q(t)\|_{L^2}^2 + \|\tilde{L}^{j_2} \partial_t^{j_1} u(t)\|_{L^2}^2 \right),$$

$$L^1 q = c^2 r^2 \partial_x q, \quad L^2 q = \partial_x (c^2 r^4 \partial_x q), \quad \tilde{L}^1 u = \partial_x (r^2 u), \quad \tilde{L}^2 u = c^2 r^2 \partial_x^2 (r^2 u).$$

Assume the initial data (u_{in}, q_{in}, R_{in}) satisfies the assumption of Theorem 1.3:

$$\epsilon^2 := \sum_{j=0}^2 \|L_0^j q_{in}\|_{L^2}^2 + \sum_{j=0}^2 \|\tilde{L}_0^j u_{in}\|_{L^2}^2 + |f(R_{in})|^2 \leq \epsilon_0^2,$$

where the constant ϵ_0^2 will be determined later. Taking $r = \xi(x) = (1 + 3x)^{\frac{1}{3}}$ in Lemma 2.1 yields $\|q_{in}\|_{L^\infty} \leq C\epsilon$, and by choosing ϵ_0 small there holds

$$-\frac{1}{4} < -C\epsilon \leq q_{in} \leq C\epsilon \leq \frac{1}{4}.$$

Meanwhile, since f is smooth and strictly increasing near $R = 1$, we have $|R_{in} - 1| \leq C\epsilon$ and thus $0 < 1 - C\epsilon \leq R_{in} \leq 1 + C\epsilon < \bar{R}$. Then Theorem 1.2 gives the existence of a local solution. Note that the inequality (2.18) is essentially independent of k , therefore $\sum_{j=0}^2 e^j(0) \leq \tilde{B}\epsilon^2$ for some constant \tilde{B} . We now consider the time interval $[0, T]$ such that the following bounds hold on $[0, T]$:

$$\sum_{j=0}^2 e^j(t) \leq 4 \sum_{j=0}^2 e^j(0) \leq 4\tilde{B}\epsilon^2, \quad (3.1)$$

$$-\frac{1}{2} \leq q(x, t) \leq \frac{1}{2}, \quad x > 0, \quad (3.2)$$

and consequently for $x > 0, t \in [0, T]$

$$-\frac{1}{2} \leq f(R) \leq \frac{1}{2},$$

$$\frac{Ca\gamma}{2} \left(\frac{2}{3}\right)^{\gamma+1} \leq c^2(x, t) = \frac{Ca\gamma}{2} (1 + q(x, t))^{-\gamma-1} \leq \frac{Ca\gamma}{2} 2^{\gamma+1},$$

$$R^3(t) + \frac{3}{2}x \leq r^3(x, t) = R^3(t) + 3 \int_0^x (1 + q(y, t)) dy \leq R^3(t) + \frac{9}{2}x.$$

Such an interval exists due to continuity and by choosing ϵ_0 further small. By dropping the subscript in (2.20)(2.21) we obtain

$$\partial_t^2 q - \tilde{L}^1 \partial_t u = 4r^{-1} u \partial_t q - 6r^{-2} (1 + q) u^2,$$

$$\partial_t^2 u - L^1 \partial_t q = (-\gamma - 1)(1 + q)^{-1} \partial_t q + 2r^{-1} u \partial_t u.$$

Hence by Lemma 2.1 the following bounds hold on $[0, T]$:

$$\begin{aligned} \|L^2 q\|_{L^2} &= \|\tilde{L}^1 \partial_t u\|_{L^2} \leq \|\partial_t^2 q\|_{L^2} + C \left(\|u\|_{L^2} + \|\tilde{L}^1 u\|_{L^2} \right) (\|\partial_t q\|_{L^2} + \|u\|_{L^2}) \\ &= \|\partial_t^2 q\|_{L^2} + C (\|u\|_{L^2} + \|\partial_t q\|_{L^2})^2 \\ &\leq C \sum_{j=0}^2 e^j(t)^{\frac{1}{2}}, \end{aligned} \quad (3.3)$$

$$\begin{aligned} \|\tilde{L}^2 u\|_{L^2} &= \|L^1 \partial_t q\|_{L^2} \leq \|\partial_t^2 u\|_{L^2} + C \left(\|\partial_t u\|_{L^2} + \|\tilde{L}^1 \partial_t u\|_{L^2} \right) (\|u\|_{L^2} + \|\partial_t q\|_{L^2}) \\ &\leq \|\partial_t^2 q\|_{L^2} + C\epsilon (\|u\|_{L^2} + \|\partial_t q\|_{L^2}) \\ &\leq C \sum_{j=0}^2 e^j(t)^{\frac{1}{2}}, \end{aligned} \quad (3.4)$$

where we have used that $\sum_{j=0}^2 e^j(t) \leq 4\tilde{B}\epsilon^2$ on $[0, T]$ and $\epsilon < \epsilon_0 \lesssim 1$. Therefore, there exists a constant $A > 0$ such that on $[0, T]$

$$E(t) \leq A \sum_{j=0}^2 e^j(t). \quad (3.5)$$

Moreover, applying Lemma 2.1 again gives that for $x > 0$, $t \in [0, T]$

$$\begin{aligned} |q(x, t)| + |\partial_\xi q(x, t)| + |\partial_t q(x, t)| &\lesssim \epsilon \xi^{-1}, \\ |u(x, t)| + |\partial_\xi u(x, t)| + |\partial_t u(x, t)| &\lesssim \epsilon \xi^{-1}, \end{aligned} \quad (3.6)$$

where we used that $\xi^{-1} \lesssim r^{-1} \lesssim \xi^{-1}$ since $R(t)$ is bounded from below and above, and consequently

$$\begin{aligned} |c(x, t) - c_0| + |\partial_\xi c(x, t)| + |\partial_t c(x, t)| &\lesssim \epsilon \xi^{-1}, \\ |R-1| &\lesssim \epsilon. \end{aligned} \quad (3.7)$$

By Hölder and (1.13), we also compute

$$\begin{aligned} \left| \frac{r(x, t)^2}{\xi^2} - 1 \right| &= \frac{\xi r(x, t) + \xi^2}{r(x, t)^2 + \xi r(x, t) + \xi^2} \xi^{-3} \left| R(t)^3 - 1 + 3 \int_0^x q(y, t) dy \right| \\ &\lesssim \xi^{-3} \left(|R(t)^3 - 1| + x^{\frac{1}{2}} \|q\|_{L^2} \right) \\ &\lesssim \epsilon \xi^{-\frac{3}{2}}. \end{aligned} \quad (3.8)$$

Combining (3.7)(3.8) yields

$$|cr^2\xi^{-2} - c_0| \lesssim \epsilon \xi^{-1}, \quad (3.9)$$

and thus there exist constants \bar{c} and \underline{c} such that

$$-\epsilon \lesssim \underline{c} - c_0 \leq cr^2\xi^{-2} - c_0 \leq \bar{c} - c_0 \lesssim \epsilon. \quad (3.10)$$

If T is large enough so that $1 + \bar{c}T \geq \exp\left(\frac{\kappa_0}{\epsilon}\right)$, then we have obtained the almost global existence. Hence we assume with out of generality that

$$1 + \bar{c}T \leq \exp\left(\frac{\kappa}{\epsilon}\right) \quad (3.11)$$

holds for a constant $\kappa \leq \kappa_0$ with κ_0 to be determined. The heart of the proof of almost global existence is the following:

Proposition 3.1. *Assume that (3.1)(3.2) and consequently (3.3)-(3.10) hold on $[0, T]$ with T satisfying (3.11). Then the following holds:*

1. *The bounds (3.1)(3.2) can be strictly improved on $[0, T]$. To be specific, we will prove*

$$\sum_{j=0}^2 e^j(t) \leq 3\tilde{B}\epsilon^2, \quad t \in [0, T], \quad (3.12)$$

and thus $|q(x, t)| \lesssim \epsilon$ by (3.6), which improves (3.2) in turn for suitable ϵ_0 .

2. *(Control of error). The following estimate holds on $[0, T]$:*

$$|F(R) - \mathcal{R}(t)| \lesssim \frac{\log(1 + \bar{c}t)}{1 + \underline{c}t} \epsilon^{\frac{3}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \frac{\epsilon^{\frac{3}{2}}}{1 + \underline{c}t} \left(\epsilon^{\frac{1}{2}} + \tilde{\epsilon}^{\frac{1}{2}} \right) + \frac{\epsilon}{(1 + \underline{c}t)^2} (\epsilon + \tilde{\epsilon}).$$

Remark 3.2. In fact, system (1.4)-(1.8) admits a conserved energy. We have by (1.4) and (1.5) that

$$\begin{aligned} 0 &= u \left(\partial_t u + \frac{Ca}{2} r^2 \partial_x (\rho^\gamma) \right) + \frac{Ca}{2} \rho^{-2} (\rho^\gamma - 1) (\partial_t \rho + \rho^2 \partial_x (r^2 u)) \\ &= \partial_t \left(\frac{1}{2} u^2 + \frac{Ca}{2} \left(\frac{1}{\gamma-1} \rho^{\gamma-1} - \frac{\gamma}{\gamma-1} + \rho^{-1} \right) \right) + \frac{Ca}{2} \partial_x (r^2 u (\rho^\gamma - 1)). \end{aligned}$$

Integrating the above identity on the strip $\mathbb{R}_+ \times [0, T]$ and using the boundary conditions gives

$$\begin{aligned} 0 &= \int_0^\infty \left(\frac{1}{2} u^2 + H(\rho) \right) (T) dx - \int_0^\infty \left(\frac{1}{2} u^2 + H(\rho) \right) (0) dx \\ &\quad - \int_0^T \left(R^2 \left(\left(\frac{Ca}{2} + \frac{2}{We} \right) R^{-3\gamma_0} - \frac{2}{We} R^{-1} - \frac{Ca}{2} \right) \frac{dR}{dt} \right) dt, \end{aligned}$$

where $H(\rho) := \frac{Ca}{2(\gamma-1)} (\rho^{\gamma-1} - \gamma + (\gamma-1)\rho^{-1})$ denotes the entropy. Note that

$$\begin{aligned} &- R^2 \left(\left(\frac{Ca}{2} + \frac{2}{We} \right) R^{-3\gamma_0} - \frac{2}{We} R^{-1} - \frac{Ca}{2} \right) \frac{dR}{dt} \\ &= \frac{d}{dt} \left\{ \frac{1}{3\gamma_0 - 3} \left(\frac{Ca}{2} + \frac{2}{We} \right) (R^{-3\gamma_0+3} - 1) + \frac{1}{We} (R^2 - 1) + \frac{Ca}{6} (R^3 - 1) \right\} \\ &=: \frac{d}{dt} W(R). \end{aligned}$$

$W(R)$ is actually the work done by the surface tension and the pressure at both the bubble surface and the infinity. Since $W(R)$ is convex in R , we obtain the conserved energy:

$$\int_0^\infty \left(\frac{1}{2} u^2 + H(\rho) \right) dx + W(R).$$

This conserved energy can give a uniform in time the upper and lower bound of R as well as the control of L^2 norms of u and q . However, since L^2 is sub-critical compared with the existence space, such controls are of less importance, and we will not use this in the proceeding proof.

4. KSS type estimate with boundary terms

In this section we establish a KSS (Keel-Smith-Sogge) type estimate regarding the quasilinear wave equation (1.16). This will provide a long time control upon the weighted norms of the solution, which will be used later to improve (3.1). However, due to the non-vanishing boundary value, we will obtain boundary terms in this KSS type estimate. Fortunately, these boundary terms have positive sign in higher regularity and negative sign in lower regularity. Therefore, by taking a delicate linear combination, the only left bad term is the one with the lowest regularity, whose treatment will be given in the next two sections. To be specific, the goal of this section is to prove the estimate below.

Proposition 4.1. *Let φ be a solution to (1.16) on $[0, T]$, and assume that (3.1)(3.2) hold on $[0, T]$, then the following bound holds:*

$$\begin{aligned} & \sum_{j=0}^2 \left\{ \int_0^T \int_0^\infty \xi^{-1} \left((\partial_t^{1+j} \varphi)^2 + (cr^2 \partial_x \partial_t^j \varphi)^2 \right) dx dt + \int_0^T \int_0^\infty \xi^{-3} (\partial_t^j \varphi)^2 dx dt \right. \\ & \left. + \int_0^T (\partial_t^{1+j} \varphi)^2 \Big|_{x=0} dt + \int_0^T \left(cr^2 \partial_x \partial_t^j \varphi + \frac{c}{\rho r} \partial_t^j \varphi \right)^2 \Big|_{x=0} dt \right\} \\ & \lesssim \max \{ \log_2 T, 1 \} \sum_{j=0}^2 \max_{0 \leq t \leq T} \int_0^\infty \left((\partial_t^{1+j} \varphi)^2 + (cr^2 \partial_x \partial_t^j \varphi)^2 \right) dx + \int_0^T \varphi^2|_{x=0} dt. \end{aligned} \quad (4.1)$$

Proof. Step 1. We begin with defining a sequence of modified momentum density for $j = 0, 1, 2$ and $k \in \mathbb{N} \cup \{0\}$:

$$\begin{aligned} P_{0,k}^j & := M_k \partial_t^{1+j} \varphi \cdot cr^2 \partial_x \partial_t^j \varphi + \frac{1}{2} N_k \partial_t^j \varphi \partial_t^{1+j} \varphi, \\ P_{1,k}^j & := \frac{1}{2} M_k \left[(\partial_t^{1+j} \varphi)^2 + (cr^2 \partial_x \partial_t^j \varphi)^2 \right] + \frac{1}{2} N_k \partial_t^j \varphi \cdot cr^2 \partial_x \partial_t^j \varphi - \frac{1}{4} cr^2 (\partial_x N_k) (\partial_t^j \varphi)^2, \end{aligned}$$

where M_k, N_k are undetermined weight functions. By taking time derivative in (1.16), the following holds:

$$\partial_t^2 \partial_t^j \varphi - c^2 \partial_x (r^4 \partial_x \partial_t^j \varphi) = [\partial_t^j, c^2 \partial_x r^4 \partial_x] \varphi. \quad (4.2)$$

We compute by brute force and (4.2) that

$$\begin{aligned} & \int_0^T \int_0^\infty \left(-\partial_t P_{0,k}^j + \partial_x (cr^2 P_{1,k}^j) \right) dx dt \\ & = \int_0^T \int_0^\infty \left(\frac{r^2 \partial_x (cM_k)}{2} + \frac{c \partial_x r^2 M_k}{2} - \frac{N_k}{2} \right) (\partial_t^{1+j} \varphi)^2 dx dt \\ & \quad + \int_0^T \int_0^\infty \left(\frac{r^2 \partial_x (cM_k)}{2} + \frac{N_k}{2} - \frac{c \partial_x r^2 M_k}{2} \right) (cr^2 \partial_x \partial_t^j \varphi)^2 dx dt \\ & \quad + \int_0^T \int_0^\infty -\frac{1}{4} \partial_x (c^2 r^4 \partial_x N_k) (\partial_t^j \varphi)^2 dx dt \\ & \quad + \int_0^T \int_0^\infty M_k (r^2 \partial_x c) (cr^2 \partial_x \partial_t^j \varphi)^2 dx dt - \int_0^T \int_0^\infty M_k \frac{\partial_t (cr^2)}{cr^2} \partial_t^{1+j} \varphi \cdot cr^2 \partial_x \partial_t^j \varphi dx dt \\ & \quad + \int_0^T \int_0^\infty N_k (r^2 \partial_x c) \partial_t^j \varphi \cdot cr^2 \partial_x \partial_t^j \varphi dx dt \\ & \quad - \int_0^T \int_0^\infty \left(M_k cr^2 \partial_x \partial_t^j \varphi + \frac{1}{2} N_k \partial_t^j \varphi \right) [\partial_t^j, c^2 \partial_x r^4 \partial_x] \varphi dx dt \\ & \quad - \int_0^T \int_0^\infty \partial_t M_k \partial_t^{1+j} \varphi \cdot cr^2 \partial_x \partial_t^j \varphi dx dt - \int_0^T \int_0^\infty \frac{\partial_t N_k}{2} \partial_t^j \varphi \partial_t^{1+j} \varphi dx dt. \end{aligned} \quad (4.3)$$

The construction of P_0 's and P_1 's as well as the calculation is an analogue to Section 5 of [20], which also considers a more general case without symmetry, see also [9], but for integrity, we write

down the computation process of the present context in Appendix A. Now select $M_k := \frac{1}{c} \frac{\xi}{2^k + \xi}$, $N_k := c \partial_x r^2 \cdot M_k = \frac{2}{\rho r} \frac{\xi}{2^k + \xi}$, and thus the coefficients of the leading order quadratic terms are

$$\frac{r^2 \partial_x (c M_k)}{2} + \frac{c \partial_x r^2 M_k}{2} - \frac{N_k}{2} = \frac{r^2 \partial_x (c M_k)}{2} - \frac{c \partial_x r^2 M_k}{2} + \frac{N_k}{2} = \frac{r^2}{2 \xi^2} \frac{2^k}{(2^k + \xi)^2}.$$

The coefficient of the third term is

$$\begin{aligned} & -\frac{1}{4} \partial_x \left(c^2 r^4 \partial_x \left(\frac{2}{\rho r} \frac{\xi}{2^k + \xi} \right) \right) \\ &= -\frac{1}{2} \partial_x \left(\frac{c^2 r^3}{\rho \xi^2} \frac{2^k}{(2^k + \xi)^2} \right) + \frac{1}{2} \partial_x \left(\frac{c^2}{\rho^2} \frac{\xi}{2^k + \xi} \right) + \frac{1}{2} \partial_x \left(\frac{c^2 r^3 \partial_x \rho}{\rho^2} \frac{\xi}{2^k + \xi} \right) \\ &= -\frac{1}{2} \partial_x \left(\frac{c^2 r^3}{\rho \xi^3} \right) \frac{2^k \xi}{(2^k + \xi)^2} - \frac{1}{2} \frac{c^2 r^3}{\rho \xi^5} \frac{2^k (2^k - \xi)}{(2^k + \xi)^3} + \frac{1}{2} \partial_x \left(\frac{c^2}{\rho^2} \right) \frac{\xi}{2^k + \xi} \\ &\quad + \frac{1}{2} \frac{c^2}{\rho^2 \xi^2} \frac{2^k}{(2^k + \xi)^2} + \frac{1}{2} \partial_x \left(\frac{c^2 r^3 \partial_x \rho}{\rho^2} \right) \frac{\xi}{2^k + \xi} + \frac{1}{2} \frac{c^2 r^3 \partial_x \rho}{\rho^2 \xi^2} \frac{2^k}{(2^k + \xi)^2} \\ &= \frac{1}{2} \frac{c^2}{\rho^2 \xi^2} \frac{2^k}{(2^k + \xi)^2} \left(1 + \frac{\rho r^3}{\xi^3} \frac{\xi - 2^k}{2^k + \xi} \right) - \frac{1}{2} \partial_x \left(\frac{c^2 r^3}{\rho \xi^3} \right) \frac{2^k \xi}{(2^k + \xi)^2} \\ &\quad + \frac{1}{2} \partial_x \left(\frac{c^2}{\rho^2} \right) \frac{\xi}{2^k + \xi} + \frac{1}{2} \partial_x \left(\frac{c^2 r^3 \partial_x \rho}{\rho^2} \right) \frac{\xi}{2^k + \xi} + \frac{1}{2} \frac{c^2 r^3 \partial_x \rho}{\rho^2 \xi^2} \frac{2^k}{(2^k + \xi)^2} \\ &= \frac{1}{2} \frac{c^2}{\rho^2} \left(1 + \frac{\rho r^3}{\xi^3} \right) \frac{2^k}{\xi (2^k + \xi)^3} + \frac{1}{2} \frac{c^2}{\rho^2} \left(1 - \frac{\rho r^3}{\xi^3} \right) \frac{2^{2k}}{\xi^2 (2^k + \xi)^3} - \frac{1}{2} \partial_x \left(\frac{c^2 r^3}{\rho \xi^3} \right) \frac{2^k \xi}{(2^k + \xi)^2} \\ &\quad + \frac{1}{2} \partial_x \left(\frac{c^2}{\rho^2} \right) \frac{\xi}{2^k + \xi} + \frac{1}{2} \partial_x \left(\frac{c^2 r^3 \partial_x \rho}{\rho^2} \right) \frac{\xi}{2^k + \xi} + \frac{1}{2} \frac{c^2 r^3 \partial_x \rho}{\rho^2 \xi^2} \frac{2^k}{(2^k + \xi)^2}. \end{aligned}$$

Hence we obtain the main part $\frac{1}{2} \frac{c^2}{\rho^2} \left(1 + \frac{\rho r^3}{\xi^3} \right) \frac{2^k}{\xi (2^k + \xi)^3}$, while the left will be regarded as nonlinear terms. Next, by divergence theorem there holds

$$\begin{aligned} & \int_0^\infty P_{0,k}^j(x, 0) dx - \int_0^\infty P_{0,k}^j(x, T) dx - \int_0^T (c r^2 P_{1,k}^j)(0, t) dt \\ &= \int_0^T \int_0^\infty \left(-\partial_t P_{0,k}^j + \partial_x (c r^2 P_{1,k}^j) \right) dx dt. \end{aligned} \tag{4.4}$$

Moreover, we expand the boundary term $P_{1,k}^j(0, t)$ as

$$\begin{aligned} & P_{1,k}^j(0, t) \\ &= \left[\frac{1}{2} M_k \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(c r^2 \partial_x \partial_t^j \varphi \right)^2 \right) + \frac{1}{2} N_k \partial_t^j \cdot c r^2 \partial_x \partial_t^j \varphi - \frac{1}{4} c r^2 \partial_x N_k \left(\partial_t^j \varphi \right)^2 \right] \Big|_{x=0} \\ &= \left[\frac{1}{2} M_k \left(\partial_t^{1+j} \varphi \right)^2 + \frac{1}{2} M_k \left(c r^2 \partial_x \partial_t^j \varphi + \frac{N_k}{2 M_k} \partial_t^j \varphi \right)^2 - \left(\frac{N_k^2}{8 M_k} + \frac{1}{4} c r^2 \partial_x N_k \right) \left(\partial_t^j \varphi \right)^2 \right] \Big|_{x=0}, \end{aligned}$$

and compute the coefficients explicitly:

$$\frac{1}{2} M_k(0, t) = \frac{1}{2c(0, t)} \frac{1}{2^k + 1}, \quad \frac{N_k(0, t)}{2 M_k(0, t)} = \left(\frac{c}{\rho r} \right) (0, t),$$

$$\begin{aligned}
\frac{N_k(0,t)^2}{8M_k(0,t)} &= \frac{1}{2} \left(\frac{c}{\rho^2 r^2}(0,t) \right) \frac{1}{2^k + 1}, \\
&\frac{1}{4} (cr^2 \partial_x N_k)(0,t) \\
&= \frac{1}{4} \left[cr^2 \partial_x \left(\frac{2}{\rho r} \frac{\xi}{2^k + \xi} \right) \right] \Big|_{x=0} \\
&= \frac{1}{2} \left[\frac{cr}{\rho} \frac{2^k}{\xi^2(2^k + \xi)^2} \right] \Big|_{x=0} - \frac{1}{2} \left[\frac{c}{\rho^2} (r \partial_x \rho + r^{-2}) \frac{\xi}{2^k + \xi} \right] \Big|_{x=0} \\
&= \frac{1}{2} \left(\frac{cr}{\rho} \right) \Big|_{x=0} \frac{2^k}{(2^k + 1)^2} - \frac{1}{2} \left(\frac{c}{\rho^2 r^2} \right) \Big|_{x=0} \frac{1}{2^k + 1} - \frac{1}{2} \left(\frac{c}{\rho^2} r \partial_x \rho \right) \Big|_{x=0} \frac{1}{2^k + 1}, \\
\frac{N_k(0,t)^2}{8M_k(0,t)} + \frac{1}{4} (cr^2 \partial_x N_k)(0,t) &= \frac{1}{2} \left(\frac{cr}{\rho} \right) \Big|_{x=0} \frac{2^k}{(2^k + 1)^2} - \frac{1}{2} \left(\frac{c}{\rho^2} r \partial_x \rho \right) \Big|_{x=0} \frac{1}{2^k + 1}.
\end{aligned}$$

Combining (4.3)(4.4) yields the identity

$$\begin{aligned}
&\int_0^T \int_0^\infty \frac{r^2}{2\xi^2} \frac{2^k}{(2^k + \xi)^2} \left((\partial_t^{1+j} \varphi)^2 + (cr^2 \partial_x \partial_t^j \varphi)^2 \right) dx dt \\
&+ \int_0^T \int_0^\infty \frac{c^2}{2\rho^2} \left(1 + \frac{\rho r^3}{\xi^3} \right) \frac{2^k}{\xi(2^k + \xi)^3} (\partial_t^j \varphi)^2 dx dt \\
&+ \int_0^T \frac{R^2}{2} \frac{1}{2^k + 1} (\partial_t^{1+j} \varphi)^2 \Big|_{x=0} dt + \int_0^T \frac{R^2}{2} \frac{1}{2^k + 1} \left(cr^2 \partial_x \partial_t^j \varphi + \frac{c}{\rho r} \partial_t^j \varphi \right)^2 \Big|_{x=0} dt \quad (4.5) \\
&= \int_0^\infty P_{0,k}^j(x,0) dx - \int_0^\infty P_{0,k}^j(x,T) dx + \int_0^T \frac{1}{2} \left(\frac{c^2 r^3}{\rho} \right) \Big|_{x=0} \frac{2^k}{(2^k + 1)^2} (\partial_t^j \varphi)^2 \Big|_{x=0} dt \\
&+ \sum_{l=1}^8 NL_{k,l}^j,
\end{aligned}$$

with the nonlinearities given by

$$\begin{aligned}
NL_{k,1}^j &:= - \int_0^T \int_0^\infty \left[\frac{1}{2} \frac{c^2}{\rho^2} \left(1 - \frac{\rho r^3}{\xi^3} \right) \frac{2^{2k}}{\xi^2(2^k + \xi)^3} - \frac{1}{2} \partial_x \left(\frac{c^2 r^3}{\rho \xi^3} \right) \frac{2^k \xi}{(2^k + \xi)^2} + \frac{1}{2} \partial_x \left(\frac{c^2}{\rho^2} \right) \frac{\xi}{2^k + \xi} \right. \\
&\quad \left. + \frac{1}{2} \partial_x \left(\frac{c^2 r^3 \partial_x \rho}{\rho^2} \right) \frac{\xi}{2^k + \xi} + \frac{1}{2} \frac{c^2 r^3 \partial_x \rho}{\rho^2 \xi^2} \frac{2^k}{(2^k + \xi)^2} \right] (\partial_t^j \varphi)^2 dx dt, \\
NL_{k,2}^j &= - \int_0^T \int_0^\infty (r^2 \partial_x c) M_k (cr^2 \partial_x \partial_t^j \varphi)^2 dx dt, \\
NL_{k,3}^j &= - \int_0^T \int_0^\infty M_k \frac{\partial_t(cr^2)}{cr^2} \partial_t^{1+j} \varphi \cdot cr^2 \partial_x \partial_t^j \varphi dx dt, \\
NL_{k,4}^j &= - \int_0^T \int_0^\infty N_k (r^2 \partial_x c) \partial_t^j \varphi \cdot cr^2 \partial_x \partial_t^j \varphi dx dt, \\
NL_{k,5}^j &= \int_0^T \int_0^\infty \left(M_k cr^2 \partial_x \partial_t^j \varphi + \frac{1}{2} N_k \partial_t^j \varphi \right) [\partial_t^j, c^2 \partial_x r^4 \partial_x] \varphi dx dt, \\
NL_{k,6}^j &= \int_0^T \int_0^\infty \partial_t M_k \partial_t^{1+j} \varphi \cdot cr^2 \partial_x \partial_t^j \varphi dx dt,
\end{aligned}$$

$$\begin{aligned}
NL_{k,7}^j &= \int_0^T \int_0^\infty \frac{\partial_t N_k}{2} \partial_t^j \varphi \partial_t^{1+j} \varphi dx dt, \\
NL_{k,8}^j &= - \int_0^T \frac{1}{2} \left(\frac{c^2 r^3}{\rho^2} \partial_x \rho \right) \Big|_{x=0} \frac{1}{2^k + 1} \left(\partial_t^j \varphi \right)^2 \Big|_{x=0} dt.
\end{aligned}$$

Note that $\frac{2^k}{(2^k + \xi)^2} \simeq \frac{1}{\xi}$ and $\frac{2^k}{\xi(2^k + \xi)^3} \simeq \frac{1}{\xi^3}$ for $\xi \in [2^k, 2^{k+1}]$. Summing in k with $2^k \leq T$, or equivalently, for $k \leq K$ with $K := \max\{-1, \lfloor \log_2 T \rfloor\}$ gives

$$\begin{aligned}
& \int_0^T \int_0^\infty \frac{1}{\xi} \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx dt \\
& \leq \sum_{k=0}^K \int_0^T \int_{2^k \leq \xi < 2^{k+1}} \frac{1}{\xi} \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx dt \\
& \quad + \int_0^T \int_{\xi \geq 2^{K+1}} \frac{1}{\xi} \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx dt \tag{4.6} \\
& \lesssim \sum_{k=0}^K \int_0^T \int_0^\infty \frac{r^2}{2\xi^2} \frac{2^k}{(2^k + \xi)^2} \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx dt \\
& \quad + \max_{0 \leq t \leq T} \int_0^\infty \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx,
\end{aligned}$$

where we used $2^{-(K+1)} \leq T^{-1}$ and (3.2), and similarly,

$$\begin{aligned}
& \int_0^T \int_0^\infty \frac{1}{\xi^3} \left(\partial_t^j \varphi \right)^2 dx dt \\
& \leq \sum_{k=0}^K \int_0^T \int_{2^k \leq \xi < 2^{k+1}} \frac{1}{\xi^3} \left(\partial_t^j \varphi \right)^2 dx dt + \int_0^T \int_{\xi \geq 2^{K+1}} \frac{1}{\xi^3} \left(\partial_t^j \varphi \right)^2 dx dt \tag{4.7} \\
& \lesssim \sum_{k=0}^K \int_0^T \int_0^\infty \frac{1}{2} \frac{c^2}{\rho^2} \left(1 + \frac{\rho r^3}{\xi^3} \right) \frac{2^k}{\xi(2^k + \xi)^3} \left(\partial_t^j \varphi \right)^2 dx dt + \max_{0 \leq t \leq T} \int_0^\infty \frac{1}{\xi^2} \left(\partial_t^j \varphi \right)^2 dx.
\end{aligned}$$

Step 2. In this part we shall bound the nonlinearities by the left hand of (4.6) and (4.7) with the help of (3.1)-(3.10). Since we are using the new unknown φ , it is necessary to build a bridge between φ and u, ρ . Recall $\frac{c}{\rho} u = cr^2 \partial_x \varphi$ and (1.15) that $\partial_t \varphi + \frac{C\alpha\gamma}{2(\gamma-1)} (\rho^{\gamma-1} - 1) = \frac{1}{2} u^2$. By taking time derivatives, the following hold:

$$\partial_t^2 \varphi - \frac{c^2}{\rho} \partial_t q = u \partial_t u,$$

$$\partial_t^3 \varphi - \frac{c^2}{\rho} \partial_t^2 q = \partial_t \left(\frac{c^2}{\rho} \right) \partial_t q + (\partial_t u)^2 + u \partial_t^2 u.$$

Similarly, taking time derivatives on $\partial_x \varphi = \frac{u}{\rho r^2}$ yields

$$cr^2 \partial_x \partial_t \varphi = \frac{c}{\rho} \partial_t u + cu \left(\partial_t q - \frac{2u}{\rho r} \right),$$

$$cr^2\partial_x\partial_t^2\varphi = \frac{c}{\rho}\partial_t^2u + 2c\left(\partial_tq - \frac{2u}{\rho r}\right)\partial_tu + cu\left(-\frac{2}{\rho}\frac{\partial_tu}{r} + \frac{6u^2}{\rho r^2} - \frac{4u\partial_tq}{r} + \partial_t^2q\right).$$

Hence by (3.6)(3.7)

$$\left|\partial_t\varphi - \frac{c^2}{\rho}(\rho^{\gamma-1} - 1)\right| \lesssim \epsilon\xi^{-1}|u| \simeq \epsilon\xi^{-1}|cr^2\partial_x\varphi|,$$

$$\left|\partial_t^2\varphi - \frac{c^2}{\rho}\partial_tq\right| \lesssim \epsilon\xi^{-1}|u| \simeq \epsilon\xi^{-1}|cr^2\partial_x\varphi|, \quad (4.8)$$

$$\left|\partial_t^3\varphi - \frac{c^2}{\rho}\partial_t^2q\right| \lesssim \epsilon\xi^{-1}(|\partial_tq| + |\partial_tu| + |\partial_t^2u|),$$

$$|cr^2\partial_x\partial_t\varphi - \frac{c}{\rho}\partial_tu| \lesssim \epsilon\xi^{-1}|u| \simeq \epsilon\xi^{-1}|cr^2\partial_x\varphi|,$$

$$|cr^2\partial_x\partial_t^2\varphi - \frac{c}{\rho}\partial_t^2u| \lesssim \epsilon\xi^{-1}(|u| + |\partial_tu| + |\partial_t^2q|). \quad (4.9)$$

Moreover, for sufficiently small ϵ_0 and $\epsilon < \epsilon_0$, (4.8)(4.9) further implies that

$$\begin{aligned} |\partial_t^2u| + |\partial_t^2q| &\lesssim |cr^2\partial_x\partial_t^2\varphi| + |\partial_t^3\varphi| + \epsilon\xi^{-1}(|u| + |\partial_tq| + |\partial_tu|) \\ &\lesssim |cr^2\partial_x\partial_t^2\varphi| + |\partial_t^3\varphi| + \epsilon\xi^{-1}(|cr^2\partial_x\varphi| + |cr^2\partial_x\partial_t\varphi| + |\partial_t^2\varphi|), \end{aligned} \quad (4.10)$$

and consequently

$$|\partial_t^2u - \rho r^2\partial_x\partial_t^2\varphi| \lesssim \epsilon\xi^{-1} \sum_{j=0}^2 \left(|cr^2\partial_x\partial_t^j\varphi| + |\partial_t^{j+1}\varphi|\right), \quad (4.11)$$

$$|\partial_t^2q - \frac{\rho}{c^2}\partial_t^3\varphi| \lesssim \epsilon\xi^{-1} \sum_{j=0}^2 \left(|cr^2\partial_x\partial_t^j\varphi| + |\partial_t^{j+1}\varphi|\right). \quad (4.12)$$

• $NL_{1,k}^j$ **terms.** We use the bootstrap bounds (3.6)(3.8) to compute

$$\begin{aligned} \left|\frac{c^2}{\rho^2}\left(1 - \frac{\rho r^3}{\xi^3}\right)\frac{2^{2k}}{\xi^2(2^k + \xi)^3}\right| &\lesssim \left(|\rho - 1| + \rho\left|\frac{r^3}{\xi^3} - 1\right|\right)\frac{2^{2k}}{\xi^2(2^k + \xi)^3} \lesssim \epsilon\frac{2^{2k}}{\xi^3(2^k + \xi)^3}, \\ \left|\partial_x\left(\frac{c^2r^3}{\rho\xi^3}\right)\frac{2^k\xi}{(2^k + \xi)^2}\right| &\lesssim \left|\partial_x\left(\rho^\gamma\frac{r^3}{\xi^3}\right)\right|\frac{2^k\xi}{(2^k + \xi)^2} \lesssim \epsilon\frac{2^k}{\xi^2(2^k + \xi)^2}, \\ \left|\partial_x\left(\frac{c^2}{\rho^2}\right)\frac{\xi}{2^k + \xi}\right| &\lesssim |\partial_x(\rho^{\gamma-1})|\frac{\xi}{2^k + \xi} \lesssim \epsilon\frac{1}{\xi^2(2^k + \xi)}, \\ \left|\frac{c^2r^3\partial_x\rho}{\rho^2\xi^2}\frac{2^k}{(2^k + \xi)^2}\right| &= |c^2r^3\partial_xq|\frac{2^k}{\xi^2(2^k + \xi)^2} \lesssim \epsilon\frac{2^k}{\xi^2(2^k + \xi)^2}. \end{aligned}$$

Noting that $L^2q = \partial_x(c^2r^4\partial_xq)$, it follows that

$$\begin{aligned} \left|\partial_x\left(\frac{c^2r^3\partial_x\rho}{\rho^2}\right)\frac{\xi}{2^k + \xi}\right| &= |\partial_x(c^2r^3\partial_xq)|\frac{\xi}{2^k + \xi} \leq \left|\frac{c^2}{\rho}\partial_xq\right|\frac{\xi}{2^k + \xi} + \frac{1}{r}|L^2q|\frac{\xi}{2^k + \xi} \\ &\lesssim \epsilon\frac{1}{\xi^2(2^k + \xi)} + |L^2q|\frac{1}{2^k + \xi}. \end{aligned}$$

Using the above bounds we estimate the $NL_{1,k}^j$ terms except $\partial_x \left(\frac{c^2 r^3 \partial_x \rho}{\rho^2} \right) \frac{\xi}{2^k + \xi} \left(\partial_t^j \varphi \right)^2$:

$$\begin{aligned}
& \sum_{k=0}^K \left| \int_0^T \int_0^\infty \left[\frac{1}{2} \frac{c^2}{\rho^2} \left(1 - \frac{\rho r^3}{\xi^3} \right) \frac{2^{2k}}{\xi^2 (2^k + \xi)^3} - \frac{1}{2} \partial_x \left(\frac{c^2 r^3}{\rho \xi^3} \right) \frac{2^k \xi}{(2^k + \xi)^2} + \frac{1}{2} \partial_x \left(\frac{c^2}{\rho^2} \right) \frac{\xi}{2^k + \xi} \right. \right. \\
& \quad \left. \left. + \frac{1}{2} \frac{c^2 r^3 \partial_x \rho}{\rho^2 \xi^2} \frac{2^k}{(2^k + \xi)^2} \right] \left(\partial_t^j \varphi \right)^2 dx dt \right| \\
& \lesssim \epsilon \sum_{k=0}^K \int_0^T \int_0^\infty \left(\frac{2^{2k}}{\xi^3 (2^k + \xi)^3} + \frac{2^k}{\xi^2 (2^k + \xi)^2} + \frac{1}{\xi^2 (2^k + \xi)} \right) \left| \partial_t^j \varphi \right|^2 dx dt \\
& \lesssim \epsilon (1 + K) \int_0^T \int_0^\infty \xi^{-3} \left| \partial_t^j \varphi \right|^2 dx dt, \\
& \lesssim \kappa \int_0^T \int_0^\infty \xi^{-3} \left| \partial_t^j \varphi \right|^2 dx dt.
\end{aligned}$$

The last inequality holds due to the assumption (3.11) and that

$$\epsilon(1 + K) \leq \epsilon \max\{0, 1 + \lfloor \log_2 T \rfloor\} \lesssim \kappa. \quad (4.13)$$

Next, we integrate by parts and obtain

$$\begin{aligned}
& \left| \int_0^T \int_0^\infty \frac{1}{2} \partial_x \left(\frac{c^2 r^3 \partial_x \rho}{\rho^2} \right) \frac{\xi}{2^k + \xi} \left(\partial_t^j \varphi \right)^2 dx dt \right| \\
& \leq \left| \int_0^T \frac{1}{2} \left(c^2 r^3 \partial_x q \frac{\xi}{2^k + \xi} \left(\partial_t^j \varphi \right)^2 \right) \Big|_{x=0} dt \right| \\
& \quad + \left| \int_0^T \int_0^\infty \frac{1}{2} c r^2 r^3 \partial_x q \left(\frac{2^k}{\xi^2 (2^k + \xi)^2} \left(\partial_t^j \varphi \right)^2 + \frac{2\xi}{2^k + \xi} \partial_t^j \varphi \partial_x \partial_t^j \varphi \right) dx dt \right| \\
& \lesssim \epsilon \left(\int_0^T \int_0^\infty \frac{2^k}{\xi^2 (2^k + \xi)^2} \left(\partial_t^j \varphi \right)^2 dx dt + \int_0^T \int_0^\infty \frac{1}{2^k + \xi} \left(c r^2 \partial_x \partial_t^j \varphi \right)^2 dx dt \right) \\
& \quad + \epsilon \frac{1}{2^k + 1} \int_0^T \left(\partial_t^j \varphi \right)^2 \Big|_{x=0} dt.
\end{aligned}$$

Then sum in k and use (4.13) again:

$$\begin{aligned}
& \sum_{k=0}^K \left| \int_0^T \int_0^\infty \frac{1}{2} \partial_x \left(\frac{c^2 r^3 \partial_x \rho}{\rho^2} \right) \frac{\xi}{2^k + \xi} \left(\partial_t^j \varphi \right)^2 dx dt \right| \\
& \lesssim \kappa \left(\int_0^T \int_0^\infty \xi^{-3} \left(\partial_t^j \varphi \right)^2 dx dt + \int_0^T \int_0^\infty \xi^{-1} \left(c r^2 \partial_x \partial_t^j \varphi \right)^2 dx dt \right) + \epsilon \int_0^T \left(\partial_t^j \varphi \right)^2 \Big|_{x=0} dt.
\end{aligned}$$

Therefore, we conclude with the control of $NL_{1,k}^j$:

$$\begin{aligned} & \sum_{k=0}^K \left| NL_{1,k}^j \right| \\ & \lesssim \kappa \left(\int_0^T \int_0^\infty \xi^{-3} \left(\partial_t^j \varphi \right)^2 dxdt + \int_0^T \int_0^\infty \xi^{-1} \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 dxdt \right) + \epsilon \int_0^T \left(\partial_t^j \varphi \right)^2 \Big|_{x=0} dt. \end{aligned} \quad (4.14)$$

• $NL_{2,k}^j, NL_{3,k}^j, NL_{4,k}^j, NL_{8,k}^j$ **terms.** These terms can be bound easily by using (3.6)(3.7) and (4.13):

$$\begin{aligned} \sum_{k=0}^K \left| NL_{2,k}^j \right| & \leq \sum_{k=0}^K \int_0^T \int_0^\infty \left| \frac{r^2 \partial_x c}{c} \right| \frac{\xi}{2^k + \xi} \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 dxdt \\ & \lesssim \epsilon \sum_{k=0}^K \int_0^T \int_0^\infty \frac{1}{2^k + \xi} \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 dxdt \\ & \lesssim \kappa \int_0^T \int_0^\infty \xi^{-1} \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 dxdt, \end{aligned} \quad (4.15)$$

$$\begin{aligned} \sum_{k=0}^K \left| NL_{3,k}^j \right| & \lesssim \sum_{k=0}^K \int_0^T \int_0^\infty \left| \frac{\partial_t(cr^2)}{c^2 r^2} \right| \frac{\xi}{2^k + \xi} \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dxdt \\ & \lesssim \epsilon \sum_{k=0}^K \int_0^T \int_0^\infty \frac{1}{2^k + \xi} \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dxdt \\ & \lesssim \kappa \left(\int_0^T \int_0^\infty \xi^{-1} \left(\partial_t^{1+j} \varphi \right)^2 dxdt + \int_0^T \int_0^\infty \xi^{-1} \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 dxdt \right), \end{aligned} \quad (4.16)$$

$$\begin{aligned} \sum_{k=0}^K \left| NL_{4,k}^j \right| & \lesssim \sum_{k=0}^K \int_0^T \int_0^\infty \frac{2}{\rho r} \frac{1}{2^k + \xi} |r^2 \partial_x c| \left(\xi^{-2} \left(\partial_t^j \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dxdt, \\ & \lesssim \epsilon \sum_{k=0}^K \int_0^T \int_0^\infty \frac{1}{2^k + \xi} \left(\xi^{-2} \left(\partial_t^j \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dxdt \\ & \lesssim \kappa \left(\int_0^T \int_0^\infty \xi^{-3} \left(\partial_t^j \varphi \right)^2 dxdt + \int_0^T \int_0^\infty \xi^{-1} \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 dxdt \right), \end{aligned} \quad (4.17)$$

$$\begin{aligned} \sum_{k=0}^K \left| NL_{8,k}^j \right| & \leq \sum_{k=0}^K \int_0^T \frac{1}{2} |c^2 r^3 \partial_x q|_{x=0} \frac{1}{2^k + 1} \left(\partial_t^j \varphi \right)^2 \Big|_{x=0} dt \\ & \lesssim \epsilon \int_0^T \left(\partial_t^j \varphi \right)^2 \Big|_{x=0} dt. \end{aligned} \quad (4.18)$$

- $NL_{5,k}^j$ **term.** We expand the commutators and use (1.16) to obtain:

$$\begin{aligned}
[\partial_t^j, c^2 \partial_x r^4 \partial_x] \varphi &= [\partial_t^j, c^2] (\partial_x (r^4 \partial_x \varphi)) + c^2 \partial_x ([\partial_t^j, r^4] \partial_x \varphi) \\
&= [\partial_t^j, c^2] (c^{-2} \partial_t^2 \varphi) + c^2 \partial_x ([\partial_t^j, r^4] r^{-4} r^4 \partial_x \varphi) \\
&= [\partial_t^j, c^2] (c^{-2} \partial_t^2 \varphi) + c^2 [\partial_t^j, r^4] r^{-4} (c^{-2} \partial_t^2 \varphi) + c^2 [\partial_x, [\partial_t^j, r^4] r^{-4}] r^4 \partial_x \varphi.
\end{aligned}$$

For $j = 1$, we compute:

$$[\partial_t, c^2 \partial_x r^4 \partial_x] \varphi = \left(\frac{\partial_t c^2}{c^2} + \frac{\partial_t r^4}{r^4} \right) \partial_t^2 \varphi + c^2 \partial_x \left(\frac{\partial_t r^4}{r^4} \right) r^4 \partial_x \varphi$$

and using (3.6):

$$|[\partial_t, c^2 \partial_x r^4 \partial_x] \varphi| \lesssim \epsilon \xi^{-1} |\partial_t^2 \varphi| + \epsilon \xi^{-2} |c r^2 \partial_x \varphi|.$$

For $j = 2$, the commutator reads

$$\begin{aligned}
&[\partial_t^2, c^2 \partial_x r^4 \partial_x] \varphi \\
&= (\partial_t^2 c^2 + 2 \partial_t c^2 \partial_t) (c^{-2} \partial_t^2 \varphi) + c^2 \left(\frac{\partial_t^2 r^4}{r^4} + 2 \frac{\partial_t r^4}{r^4} \partial_t + 2 \partial_t r^4 \partial_t r^{-4} \right) (c^{-2} \partial_t^2 \varphi) \\
&\quad + c^2 \left[\partial_x, \frac{\partial_t^2 r^4}{r^4} + 2 \frac{\partial_t r^4}{r^4} \partial_t + 2 \partial_t r^4 \partial_t r^{-4} \right] r^4 \partial_x \varphi \\
&= \left(2 \frac{\partial_t c^2}{c^2} + 2 \frac{\partial_t r^4}{r^4} \right) \partial_t^3 \varphi + \left(\frac{\partial_t^2 c^2}{c^2} + 2 \partial_t c^2 \partial_t c^{-2} + \frac{\partial_t^2 r^4}{r^4} + 2 \partial_t r^4 \partial_t r^{-4} + 2 \frac{\partial_t r^4}{r^4} c^2 \partial_t c^{-2} \right) \partial_t^2 \varphi \\
&\quad + c^2 r^2 \left(\partial_x \left(\frac{\partial_t^2 r^4}{r^4} \right) + 2 \partial_x (\partial_t r^4 \partial_t r^{-4}) \right) r^2 \partial_x \varphi + 2 c^2 r^2 \partial_x \left(\frac{\partial_t r^4}{r^4} \right) \left(r^2 \partial_x \partial_t \varphi + \frac{\partial_t r^4}{r^4} r^2 \partial_x \varphi \right).
\end{aligned}$$

We also compute:

$$\begin{aligned}
\frac{\partial_t^2 c^2}{c^2} &= -(\gamma + 1) \frac{\partial_t (c^2 \rho \partial_t q)}{c^2} = -(\gamma + 1) \rho \partial_t^2 q + (\gamma + 1) (\gamma + 2) c^2 \rho^2 (\partial_t q)^2, \\
\frac{\partial_t^2 r^4}{r^4} &= 4 \frac{\partial_t u}{r} + 12 \frac{u^2}{r^2}, \\
\partial_x \left(\frac{\partial_t r^4}{r^4} \right) &= 4 \partial_x \left(\frac{u}{r} \right) = 4 r^{-3} (\partial_t q - 3 \rho^{-1} u), \\
\partial_x \left(\frac{\partial_t^2 r^4}{r^4} \right) &= 4 \partial_x \left(\frac{\partial_t (r^2 u)}{r^3} + \frac{u^2}{r^2} \right) = 4 r^{-3} \left(\partial_t^2 q - \frac{3}{\rho r^3} (2 r u^2 + r^2 \partial_t u) - \frac{6 u^2}{\rho r^2} + \frac{2 u}{r} \partial_t q \right),
\end{aligned}$$

where we used that $\partial_t \partial_x (r^2 u) = \partial_t^2 q$ to avoid the presence of mixed derivative. Therefore, it follows from (3.6)(3.7) that

$$\begin{aligned}
|[\partial_t^2, c^2 \partial_x r^4 \partial_x] \varphi| &\lesssim \epsilon \xi^{-1} |\partial_t^3 \varphi| + \epsilon \xi^{-2} (|\partial_t^2 \varphi| + |c r^2 \partial_x \partial_t \varphi| + \xi^{-1} |c r^2 \partial_x \varphi|) \\
&\quad + |\partial_t^2 \varphi| (|\partial_t^2 \varphi| + \xi^{-1} |c r^2 \partial_x \varphi|).
\end{aligned}$$

Then we conclude in view of (4.8)(4.10) that

$$\begin{aligned}
& \sum_{k=0}^K (|NL_{5,k}^1| + |NL_{5,k}^2|) \\
& \lesssim \epsilon \sum_{k=0}^K \int_0^T \int_0^\infty \frac{1}{2^k + \xi} \sum_{i=1}^2 (|cr^2 \partial_x \partial_t^i \varphi| + \xi^{-1} |\partial_t^i \varphi|) (|\partial_t^3 \varphi| + |\partial_t^2 \varphi| + |cr^2 \partial_x \partial_t \varphi| + |cr^2 \partial_x \varphi|) \quad (4.19) \\
& \lesssim \kappa \sum_{i=0}^2 \int_0^T \int_0^\infty \xi^{-1} \left((\partial_t^{1+j} \varphi)^2 + (cr^2 \partial_x \partial_t^j \varphi)^2 \right) dx dt.
\end{aligned}$$

• $NL_{6,k}^j, NL_{7,k}^j$ terms. We estimate by (3.6)(3.7) that

$$\begin{aligned}
\sum_{k=0}^K |NL_{6,k}^j| & \leq \sum_{k=0}^K \int_0^T \int_0^\infty |\partial_t c^{-1}| \frac{\xi}{2^k + \xi} |\partial_t^{1+j} \varphi| |cr^2 \partial_x \partial_t^j \varphi| dx dt \\
& \lesssim \epsilon \sum_{k=0}^K \int_0^T \int_0^\infty \frac{1}{2^k + \xi} |\partial_t^{1+j} \varphi| |cr^2 \partial_x \partial_t^j \varphi| dx dt \quad (4.20) \\
& \lesssim \kappa \left(\int_0^T \int_0^\infty \xi^{-1} (\partial_t^{1+j} \varphi)^2 dx dt + \int_0^T \int_0^\infty \xi^{-1} (cr^2 \partial_x \partial_t^j \varphi)^2 dx dt \right),
\end{aligned}$$

$$\begin{aligned}
\sum_{k=0}^K |NL_{7,k}^j| & \leq \sum_{k=0}^K \int_0^T \int_0^\infty \left| \partial_t \left(\frac{1}{\rho r} \right) \right| \frac{\xi}{2^k + \xi} |\partial_t^j \varphi| |\partial_t^{1+j} \varphi| dx dt \\
& \lesssim \epsilon \sum_{k=0}^K \int_0^T \int_0^\infty \xi^{-1} \frac{1}{2^k + \xi} |\partial_t^j \varphi| |\partial_t^{1+j} \varphi| dx dt \quad (4.21) \\
& \lesssim \kappa \left(\int_0^T \int_0^\infty \xi^{-3} (\partial_t^j \varphi)^2 dx dt + \int_0^T \int_0^\infty \xi^{-1} (\partial_t^{1+j} \varphi)^2 dx dt \right).
\end{aligned}$$

Step 3. Now we combine (4.5)(4.6)(4.7) to obtain:

$$\begin{aligned}
& \int_0^T \int_0^\infty \xi^{-1} \left((\partial_t^{1+j} \varphi)^2 + (cr^2 \partial_x \partial_t^j \varphi)^2 \right) dx dt + \int_0^T \int_0^\infty \xi^{-3} (\partial_t^j \varphi)^2 dx dt \\
& + \int_0^T \left(\partial_t^{1+j} \varphi \right)^2 \Big|_{x=0} dt + \int_0^T \left(cr^2 \partial_x \partial_t^j \varphi + \frac{c}{\rho r} \partial_t^j \varphi \right)^2 \Big|_{x=0} dt \\
& \lesssim \sum_{k=0}^K \left\{ \left| \int_0^\infty P_{0,k}^j(x, 0) dx \right| + \left| \int_0^\infty P_{0,k}^j(x, T) dx \right| + \sum_{l=1}^8 |NL_{l,k}^j| \right\} \quad (4.22) \\
& + \int_0^T \left(\partial_t^j \varphi \right)^2 \Big|_{x=0} dt + \max_{0 \leq t \leq T} \int_0^\infty \left((\partial_t^{1+j} \varphi)^2 + (cr^2 \partial_x \partial_t^j \varphi)^2 + \xi^{-2} (\partial_t^j \varphi)^2 \right) dx.
\end{aligned}$$

By the explicit expression of $P_{0,k}^j$ and Hardy's inequality it follows that

$$\begin{aligned} \sum_{k=0}^K \left| \int_0^\infty P_{0,k}^j(x,t) dx \right| &\lesssim (1+K) \int_0^\infty \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 + \xi^{-2} \left(\partial_t^j \varphi \right)^2 \right) dx \\ &\lesssim \max \{ \log_2 T, 1 \} \int_0^\infty \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx. \end{aligned}$$

Moreover, by taking a suitable linear combination of (4.22) of $j = 0, 1, 2$, we can cancel the boundary term on the right hand side except the one with the lowest order. Then we conclude by using the bounds (4.14)-(4.21) of nonlinearities:

$$\begin{aligned} &\sum_{j=0}^2 \left\{ \int_0^T \int_0^\infty \xi^{-1} \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx dt + \int_0^T \int_0^\infty \xi^{-3} \left(\partial_t^j \varphi \right)^2 dx dt \right. \\ &\quad \left. + \int_0^T \left(\partial_t^{1+j} \varphi \right)^2 \Big|_{x=0} dt + \int_0^T \left(cr^2 \partial_x \partial_t^j \varphi + \frac{c}{\rho r} \partial_t^j \varphi \right)^2 \Big|_{x=0} dt \right\} \\ &\lesssim \max \{ \log_2 T, 1 \} \sum_{j=0}^2 \max_{0 \leq t \leq T} \int_0^\infty \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx + \int_0^T \varphi^2|_{x=0} dt \\ &\quad + \sum_{j=0}^2 \sum_{k=0}^K \sum_{l=1}^8 \left| NL_{l,k}^j \right| \\ &\lesssim \max \{ \log_2 T, 1 \} \sum_{j=0}^2 \max_{0 \leq t \leq T} \int_0^\infty \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx + \int_0^T \varphi^2|_{x=0} dt \\ &\quad + \kappa \sum_{j=0}^2 \left\{ \int_0^T \int_0^\infty \xi^{-1} \left(\left(\partial_t^{1+j} \varphi \right)^2 + \left(cr^2 \partial_x \partial_t^j \varphi \right)^2 \right) dx dt + \int_0^T \int_0^\infty \xi^{-3} \left(\partial_t^j \varphi \right)^2 dx dt \right\} \\ &\quad + \epsilon \sum_{j=0}^2 \int_0^T \left(\partial_t^j \varphi \right)^2 \Big|_{x=0} dt. \end{aligned}$$

Hence for sufficiently small κ_0 and ϵ_0 we arrive at (4.1). \square

5. Derivation of precise boundary ODE

In order to handle the boundary value of φ present on the right hand side of (4.1), we derive in this section the equation satisfied by $\varphi|_{x=0}$. Introduce $\psi := r\varphi$, and substitute φ by $\frac{\psi}{r}$ in (1.16) yields

$$\begin{aligned} 0 &= r^{-1} \partial_t^2 \psi + 2\partial_t r^{-1} \partial_t \psi + \partial_t^2 r^{-1} \psi - c^2 \partial_x (r^3 \partial_x \psi + r^4 \partial_x r^{-1} \psi) \\ &= r^{-1} \left(\partial_t^2 \psi + 2r \partial_t r^{-1} \partial_t \psi - c^2 r^2 \partial_x (r^2 \partial_x \psi) \right) + \left(\partial_t^2 r^{-1} - c^2 \partial_x (r^4 \partial_x r^{-1}) \right) \psi. \end{aligned}$$

On the other hand, from (1.5)

$$0 = \partial_t u + \frac{Ca}{2} r^2 \partial_x (\rho^\gamma) = \partial_t^2 r - \frac{Ca\gamma}{2} \rho^{\gamma+1} r^2 \partial_x \rho^{-1} = \partial_t r^2 - c^2 r^2 \partial_x (r^2 \partial_x r),$$

and thus

$$\partial_t^2 r^{-1} - c^2 \partial_x (r^4 \partial_x r^{-1}) = \partial_t^2 r^{-1} + c^2 \partial_x (r^2 \partial_x r) = \partial_t^2 r^{-1} + r^{-2} \partial_t^2 r = 2r^{-3} (\partial_t r)^2.$$

Henceforth

$$0 = \partial_t^2 \psi - c^2 r^2 \partial_x (r^2 \partial_x \psi) - 2 \frac{u}{r} \partial_t \psi + 2 \frac{u^2}{r^2} \psi.$$

We rewrite the above equation to a hyperbolic system of the form

$$\begin{aligned} 0 &= (\partial_t + cr^2 \partial_x) (\partial_t - cr^2 \partial_x) \psi + \left(\frac{\partial_t c + cr^2 \partial_x c}{c} + \frac{2u}{r} \right) cr^2 \partial_x \psi - \frac{2u}{r} \partial_t \psi + \frac{2u^2}{r^2} \psi \\ &= c^{\frac{1}{2}} (\partial_t + cr^2 \partial_x) \left(c^{-\frac{1}{2}} (\partial_t - cr^2 \partial_x) \psi \right) + \left(\frac{\partial_t c + cr^2 \partial_x c}{2c} \right) (\partial_t + cr^2 \partial_x) \psi \\ &\quad - \frac{2u}{r} (\partial_t - cr^2 \partial_x) \psi + \frac{2u^2}{r^2} \psi, \end{aligned}$$

and similarly,

$$\begin{aligned} 0 &= c^{\frac{1}{2}} (\partial_t - cr^2 \partial_x) \left(c^{-\frac{1}{2}} (\partial_t + cr^2 \partial_x) \psi \right) + \left(\frac{\partial_t c - cr^2 \partial_x c}{2c} \right) (\partial_t - cr^2 \partial_x) \psi \\ &\quad - \frac{2u}{r} (\partial_t + cr^2 \partial_x) \psi + \frac{2u^2}{r^2} \psi. \end{aligned}$$

By introducing $w_F = (\partial_t - cr^2 \partial_x) \psi$ and $w_B = (\partial_t + cr^2 \partial_x) \psi$, which in fact stand for the forward and backward pressure wave, we see that w_F, w_B satisfy the transport equations for $(x, t) \in \mathbb{R}_+ \times [0, T]$:

$$0 = (\partial_t + cr^2 \partial_x) \frac{w_F}{c^{\frac{1}{2}}} + \left(\frac{\partial_t c + cr^2 \partial_x c}{2c} \right) \frac{w_B}{c^{\frac{1}{2}}} - \frac{2u}{r} \frac{w_F}{c^{\frac{1}{2}}} + \frac{2u^2}{r^2} \frac{\psi}{c^{\frac{1}{2}}}, \quad (5.1)$$

$$0 = (\partial_t - cr^2 \partial_x) \frac{w_B}{c^{\frac{1}{2}}} + \left(\frac{\partial_t c - cr^2 \partial_x c}{2c} \right) \frac{w_F}{c^{\frac{1}{2}}} - \frac{2u}{r} \frac{w_B}{c^{\frac{1}{2}}} + \frac{2u^2}{r^2} \frac{\psi}{c^{\frac{1}{2}}}. \quad (5.2)$$

One can check that (5.1)(5.2) is genuinely nonlinear in w_F, w_B , and this is actually a main feature of isentropic compressible Euler equations. In view of Majda's conjecture [17, Page 89], it is reasonable not to expect a global solution. To solve (5.1)(5.2) as a hyperbolic system on the half line, we shall need the boundary data of w_F . Recalling (1.7) and taking time derivative yields

$$\frac{C\alpha\gamma}{2} \rho^{\gamma-1} \partial_t \rho \Big|_{x=0} = \tilde{f}'(R) (\rho r^2 \partial_x \varphi) \Big|_{x=0} = \tilde{f}'(R) (\rho r \partial_x \psi - r^{-2} \psi) \Big|_{x=0}.$$

Then substituting $\rho^\gamma - 1$ by (1.15) gives

$$- \partial_t \left(\partial_t \varphi - \frac{1}{2} u^2 \right) \Big|_{x=0} = \tilde{f}'(R) (r \partial_x \psi) \Big|_{x=0} - \tilde{f}'(R) (\rho^{-1} r^{-2} \psi) \Big|_{x=0}.$$

In order to obtain an equation of $\psi|_{x=0}$, we compute

$$\partial_t^2 \varphi = r^{-1} \partial_t^2 \psi + 2 \partial_t r^{-1} \partial_t \psi + \partial_t^2 r^{-1} \psi = r^{-1} \partial_t^2 \psi - 2 \frac{u}{r^2} \partial_t \psi - \left(\frac{\partial_t u}{r^2} - 2 \frac{u^2}{r^3} \right) \psi,$$

$$r^2 \partial_x \psi = \frac{1}{c} (w_B - \partial_t \psi),$$

$$u = \rho r^2 \partial_x \varphi = \rho r \partial_x \psi - r^{-2} \psi = \frac{\rho}{rc} (w_B - \partial_t \psi) - r^{-2} \psi.$$

Therefore by writing $r|_{x=0} = R$ we obtain

$$\begin{aligned} 0 &= \left\{ \partial_t^2 \psi - Ru \partial_t u - \left(\frac{\tilde{f}'(R)}{c} + 2R^{-1}u \right) \partial_t \psi \right. \\ &\quad \left. - \left(\tilde{f}'(R) \rho^{-1} R^{-1} - 2R^{-2}u^2 + R^{-1} \partial_t u \right) \psi + \frac{\tilde{f}'(R)}{c} w_B \right\} \Big|_{x=0} \\ &= \left\{ \partial_t^2 \psi - \left(\frac{\tilde{f}'(R)}{c} + 2R^{-1}u - \frac{\rho \partial_t u}{c} \right) \partial_t \psi \right. \\ &\quad \left. - \left(\tilde{f}'(R) \rho^{-1} R^{-1} - 2R^{-2}u^2 - R^{-1} \partial_t u \right) \psi + \left(\frac{\tilde{f}'(R)}{c} - \frac{\rho \partial_t u}{c} \right) w_B \right\} \Big|_{x=0}. \end{aligned}$$

In short, there holds

$$\partial_t^2 \psi|_{x=0} + b_1 \partial_t \psi|_{x=0} + b_0 \psi|_{x=0} + a w_B|_{x=0} = 0, \quad (5.3)$$

with

$$\begin{aligned} b_1(t) &:= - \left(\frac{\tilde{f}'(R)}{c} + 2R^{-1}u - \frac{\rho \partial_t u}{c} \right) \Big|_{x=0}, \\ b_0(t) &:= - \left(\tilde{f}'(R) \rho^{-1} R^{-1} - 2R^{-2}u^2 - R^{-1} \partial_t u \right) \Big|_{x=0}, \\ a(t) &:= \left(\frac{\tilde{f}'(R)}{c} - \frac{\rho \partial_t u}{c} \right) \Big|_{x=0}. \end{aligned}$$

Moreover, we estimate these coefficients by using (3.6)(3.7) and the smoothness of \tilde{f} near 1:

$$|b_1(t) + \frac{\tilde{f}'(1)}{c_0}| \lesssim \epsilon, \quad |b_0(t) + \tilde{f}'(1)| \lesssim \epsilon, \quad |a(t) - \frac{\tilde{f}'(1)}{c_0}| \lesssim \epsilon. \quad (5.4)$$

Let Λ_1, Λ_2 denote the two eigenvalues of the unperturbed equation, namely, the two roots of $\lambda^2 - \frac{\tilde{f}'(1)}{c_0} \lambda - \tilde{f}'(1)$. Then we rewrite (5.3) in the form of column vectors and diagonalize the resulted equation:

$$\partial_t Y - \begin{bmatrix} \Lambda_1 & \\ & \Lambda_2 \end{bmatrix} Y + \Delta Y + W = 0, \quad (5.5)$$

$$Y := \begin{bmatrix} 1 & -\Lambda_2 \\ 1 & -\Lambda_1 \end{bmatrix} \begin{bmatrix} \partial_t \psi|_{x=0} \\ \psi|_{x=0} \end{bmatrix}, \quad \Delta = \frac{1}{\Lambda_1 - \Lambda_2} \begin{bmatrix} \delta_1 & \delta_2 \\ \delta_1 & \delta_2 \end{bmatrix}, \quad W = \begin{bmatrix} a w_B|_{x=0} \\ a w_B|_{x=0} \end{bmatrix},$$

$$\delta_1 = \Lambda_1 \left(b_1 + \frac{\tilde{f}'(1)}{c_0} \right) + (b_0 + \tilde{f}'(1)), \quad \delta_2 = -\Lambda_2 \left(b_1 + \frac{\tilde{f}'(1)}{c_0} \right) - (b_0 + \tilde{f}'(1)).$$

By solving (5.5) it follows

$$Y(t) = \exp \begin{bmatrix} \Lambda_1 t & \\ & \Lambda_2 t \end{bmatrix} Y(0) - \int_0^t \exp \begin{bmatrix} \Lambda_1(t-s) & \\ & \Lambda_2(t-s) \end{bmatrix} (\Delta Y)(s) ds \\ - \int_0^t \exp \begin{bmatrix} \Lambda_1(t-s) & \\ & \Lambda_2(t-s) \end{bmatrix} W(s) ds. \quad (5.6)$$

We remark here that from (1.7) there holds $\tilde{f}'(1) = -3\gamma \left(\frac{Ca}{2} + \frac{2}{We} \right) + \frac{2}{We} < 0$, which shows that $\text{Re}\Lambda_i < 0$, $i = 1, 2$.

6. The analysis of backward pressure wave

Recall the KSS type estimate (4.1). To control the boundary term $\int_0^T \varphi^2|_{x=0} dt$, we have obtained in the previous section the integral equation (5.6) satisfied by $\psi := r^{-1}\varphi$ with the source term containing the backward pressure wave w_B . To close this argument, we shall estimate w_B in this section by using the equations (5.1)(5.2), which form a hyperbolic system on the domain $\{(x, t) : x \geq 0, t \geq 0\}$. Generally, in view of the signs of characteristic speeds we will need the data of w_B on $\{t = 0\}$ and the data of w_F on $\{t = 0\} \cup \{x = 0\}$ to solve the Cauchy problem associated with system (5.1)(5.2), but in the present context the data of w_F on $\{x = 0\}$ is instead given implicitly by equation (5.6) by the relation $w_F = (\partial_t - cr^2\partial_x)\psi = 2\partial_t\psi - w_B$.

Recall that $\xi = (1 + 3x)^{\frac{1}{3}}$ and we rewrite (5.1)(5.2) as

$$0 = (\partial_t + cr^2\xi^{-2}\partial_\xi) \frac{w_F}{c^{\frac{1}{2}}} + \left(\frac{\partial_t c + cr^2\xi^{-2}\partial_\xi c}{2c} \right) \frac{w_B}{c^{\frac{1}{2}}} - \frac{2u}{r} \frac{w_F}{c^{\frac{1}{2}}} + \frac{2u^2}{r^2} \frac{\psi}{c^{\frac{1}{2}}}, \quad (6.1)$$

$$0 = (\partial_t - cr^2\xi^{-2}\partial_\xi) \frac{w_B}{c^{\frac{1}{2}}} + \left(\frac{\partial_t c - cr^2\xi^{-2}\partial_\xi c}{2c} \right) \frac{w_F}{c^{\frac{1}{2}}} - \frac{2u}{r} \frac{w_B}{c^{\frac{1}{2}}} + \frac{2u^2}{r^2} \frac{\psi}{c^{\frac{1}{2}}}. \quad (6.2)$$

Now we begin the analysis by introducing the characteristics. Note that the bootstrap bounds (3.6)(3.7) imply the Lipschitz continuity of $cr^2\xi^{-2}$ in both ξ and t argument. In fact, we have

$$|\partial_\xi(cr^2\xi^{-2})| + |\partial_t(cr^2\xi^{-2})| \lesssim \epsilon\xi^{-1}. \quad (6.3)$$

For each $\xi \in [1, +\infty)$, $t \in [0, T]$, define X_B as the unique solution (figure 1) to

$$\begin{cases} X_B(\xi, t; 0) = (\xi, t), \\ \frac{d}{ds} X_B(\xi, t; s) = -(cr^2\xi^{-2})(X_B(\xi, t; s)), 1, \end{cases}$$

and X_F to be the unique solution to

$$\begin{cases} X_F(\xi, t; 0) = (\xi, t), \\ \frac{d}{ds} X_F(\xi, t; s) = ((cr^2\xi^{-2})(X_F(\xi, t; s)), 1). \end{cases}$$

We then obtain the integral equation corresponding to (6.1)(6.2):

$$\begin{aligned}
& \frac{w_F}{c^{\frac{1}{2}}}(X_F(\xi, t; s)) \\
&= \exp \left\{ \int_0^s \frac{2u}{r}(X_F(\xi, t; \sigma)) d\sigma \right\} \frac{w_F}{c^{\frac{1}{2}}}(X_F(\xi, t; 0)) \\
& \quad - \int_0^s \exp \left\{ \int_0^{s-\sigma} \frac{2u}{r}(X_F(\xi, t; \tau)) d\tau \right\} \left(\frac{\partial_t c + cr^2 \xi^{-2} c}{2c} \right) \frac{w_B}{c^{\frac{1}{2}}}(X_F(\xi, t; \sigma)) d\sigma \\
& \quad - \int_0^s \exp \left\{ \int_0^{s-\sigma} \frac{2u}{r}(X_F(\xi, t; \tau)) d\tau \right\} 2r^{-2} u^2 \frac{\psi}{c^{\frac{1}{2}}}(X_F(\xi, t; \sigma)) d\sigma,
\end{aligned} \tag{6.4}$$

$$\begin{aligned}
& \frac{w_B}{c^{\frac{1}{2}}}(X_B(\xi, t; s)) \\
&= \exp \left\{ \int_0^s \frac{2u}{r}(X_B(\xi, t; \sigma)) d\sigma \right\} \frac{w_B}{c^{\frac{1}{2}}}(X_B(\xi, t; 0)) \\
& \quad - \int_0^s \exp \left\{ \int_0^{s-\sigma} \frac{2u}{r}(X_B(\xi, t; \tau)) d\tau \right\} \left(\frac{\partial_t c + cr^2 \xi^{-2} c}{2c} \right) \frac{w_F}{c^{\frac{1}{2}}}(X_B(\xi, t; \sigma)) d\sigma \\
& \quad - \int_0^s \exp \left\{ \int_0^{s-\sigma} \frac{2u}{r}(X_B(\xi, t; \tau)) d\tau \right\} 2r^{-2} u^2 \frac{\psi}{c^{\frac{1}{2}}}(X_B(\xi, t; \sigma)) d\sigma.
\end{aligned} \tag{6.5}$$

For each $t \in [0, T]$, we denote $\xi_0(t) := X_B(1, t; -t)$, and thus $X_B(\xi_0(t), 0; t) = (1, t)$, and for each $t_* \in [0, T]$ denote by $\Omega_B(t_*)$ the backward acoustic cone $\{(\xi, t) : 0 \leq t \leq t_*, 1 \leq \xi \leq X_B(\xi(t_*), 0; t)\}$. Introduce the new unknowns $v_F(\xi, t, t_*)$, $v_B(\xi, t, t_*)$ by

$$\begin{aligned}
v_F(\xi, t, t_*) &:= \sup \left\{ \left| \frac{w_F}{c^{\frac{1}{2}}}(X_F(\xi, t; s)) \right| : X_F(\xi, t; s) \in \Omega_B(t_*) \right\}, \\
v_B(\xi, t, t_*) &:= \sup \left\{ \left| \frac{w_B}{c^{\frac{1}{2}}}(X_B(\xi, t; s)) \right| : X_B(\xi, t; s) \in \Omega_B(t_*) \right\}.
\end{aligned}$$




Figure 1

$X_B(\xi, t; s) \in \Omega_B(t_*)$ is automatically fulfilled for each $(\xi, t) \in \Omega_B(t_*)$ due to the uniqueness of integral curves. Therefore we abbreviate $v_B(\xi, t; t_*)$ to $v_B(\xi, t)$. Moreover, from the definition we have

$$v_F(X_F(\xi, t; \tau); t_*) = v_F(\xi, t; t_*), \quad v_B(X_B(\xi, t; \tau)) = v_B(\xi, t).$$

It then follows from (6.4)(6.5) that

$$\begin{aligned} v_F(1, t; t_*) &\leq \exp \left\{ \sup_s \int_0^s \left| \frac{2u}{r}(X_F(1, t; \sigma)) \right| d\sigma \right\} \left\{ \left| \frac{w_F}{c^{\frac{1}{2}}}(1, t) \right| \right. \\ &\quad + \sup_s \int_0^s \left| \left(\frac{\partial_t c + cr^2 \xi^{-2} \partial_\xi c}{2c} \right) (X_F(1, t; \sigma)) \right| v_B(X_F(1, t; \sigma)) d\sigma \\ &\quad \left. + \sup_s \int_0^s \left| \left(2r^{-2} u^2 \frac{\psi}{c^{\frac{1}{2}}} \right) (X_F(1, t; \sigma)) \right| d\sigma \right\}, \end{aligned} \quad (6.6)$$

$$\begin{aligned} v_F(\xi, 0; t_*) &\leq \exp \left\{ \sup_s \int_0^s \left| \frac{2u}{r}(X_F(\xi, 0; \sigma)) \right| d\sigma \right\} \left\{ \left| \frac{w_F}{c^{\frac{1}{2}}}(\xi, 0) \right| \right. \\ &\quad + \sup_s \int_0^s \left| \left(\frac{\partial_t c + cr^2 \xi^{-2} \partial_\xi c}{2c} \right) (X_F(\xi, 0; \sigma)) \right| v_B(X_F(\xi, 0; \sigma)) d\sigma \\ &\quad \left. + \sup_s \int_0^s \left| \left(2r^{-2} u^2 \frac{\psi}{c^{\frac{1}{2}}} \right) (X_F(\xi, 0; \sigma)) \right| d\sigma \right\}, \end{aligned} \quad (6.7)$$

$$\begin{aligned} v_B(\xi, 0) &\leq \exp \left\{ \sup_s \int_0^s \left| \frac{2u}{r}(X_B(\xi, 0; \sigma)) \right| d\sigma \right\} \left\{ \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi, 0) \right| \right. \\ &\quad + \sup_s \int_0^s \left| \left(\frac{\partial_t c - cr^2 \xi^{-2} \partial_\xi c}{2c} \right) (X_B(\xi, 0; \sigma)) \right| v_F(X_B(\xi, 0; \sigma); t_*) d\sigma \\ &\quad \left. + \sup_s \int_0^s \left| \left(2r^{-2} u^2 \frac{\psi}{c^{\frac{1}{2}}} \right) (X_B(\xi, 0; \sigma)) \right| d\sigma \right\}. \end{aligned} \quad (6.8)$$

Each supremum above is taken over all $s \geq 0$ such that $X_F(1, t; \sigma)$ (or $X_F(\xi, 0; \sigma)$, $X_B(\xi, 0; \sigma)$ respectively) remains in $\Omega_B(t_*)$ for all $\sigma \in [0, s]$.

The goal of this section is to obtain a decay estimate of $\psi(1, \cdot)$. To this end, we shall need an estimate of the source term W in (5.6). We begin with the estimate of the nonlinearities in (6.6)-(6.8). These estimates rely on a change of variable between the forward and backward characteristics, which will be specified in the following analysis. Then by collecting these estimates we will be able to measure the difference between $w_B(1, t)$ and $w_B(\xi_0(t), 0)$, which further implies the estimate of W . Hereby we state the expected estimate of w_B and ψ :

Proposition 6.1. *Introduce the notations:*

$$\begin{aligned} \mathcal{V}(\xi) &:= |w_B(\xi, 0)| + \xi^{-1} \int_1^\xi |w_F(\eta, 0)| d\eta + \epsilon \xi^{-3} |\psi(\xi, 0)| + \epsilon^2 \xi^{-1} \int_1^\xi \eta^{-3} |\psi(\eta, 0)| d\eta, \\ \Psi(t_*) &:= \int_0^{t_*} \left(1 + \frac{c}{2}(t_* - t) \right)^{-1} |\partial_t \psi(1, t)| dt + \epsilon \int_0^{t_*} \left(1 + \frac{c}{2}(t_* - t) \right)^{-4} |\psi(1, t)| dt \\ &\quad + \epsilon^2 \int_0^{t_*} \left(1 + \frac{c}{2}(t_* - t) \right)^{-1} |\psi(1, t)| dt, \end{aligned}$$

then w_B satisfies the following bound:

$$\begin{aligned} & \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left(\left| \frac{w_B}{c^{\frac{1}{2}}} (1, t_*) - \frac{w_B}{c^{\frac{1}{2}}} (\xi_0(t_*), 0) \right| \right) \right\} \\ & \lesssim \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \mathcal{V}(\xi_0(t_*)) + \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \Psi(t_*) + \kappa \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left| \frac{w_B}{c^{\frac{1}{2}}} (\xi_0(t_*), 0) \right| \right\}. \end{aligned}$$

Proposition 6.2. *The following estimate of $Y(t)$ holds for $t \in [0, T]$:*

$$|Y(t) - Y_0(t)| \lesssim \frac{1}{1 + \underline{c}t} \left\{ \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}} (\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon |Y(0)| \right\},$$

where

$$Y_0(t) := \exp \begin{bmatrix} \Lambda_1 t & \\ & \Lambda_2 t \end{bmatrix} Y(0) - \frac{\tilde{f}'(1)}{c_0} \int_0^t \exp \begin{bmatrix} \Lambda_1(t-s) & \\ & \Lambda_2(t-s) \end{bmatrix} \begin{bmatrix} w_B(\xi_0(s), 0) \\ w_B(\xi_0(s), 0) \end{bmatrix} ds.$$

Since $\psi^2|_{x=0} \simeq \varphi^2|_{x=0}$, it follows

$$\int_0^T \varphi^2|_{x=0} dt \lesssim \int_0^T |Y_0(t)|^2 dt + \left(\kappa \max_{\eta \in [1, \xi_0(T)]} \left\{ \eta \left| \frac{w_B}{c^{\frac{1}{2}}} (\eta, 0) \right| \right\} + \epsilon \max_{\eta \in [1, \xi_0(T)]} \eta \mathcal{V}(\eta) + \epsilon |Y(0)| \right)^2.$$

6.1. Estimate of (6.6)

Let $X_F^{(\xi)}$, $X_B^{(\xi)}$ denote the space argument of X_F , X_B respectively. By continuity, for each $\eta \in [\xi_0(t), \xi_0(T)]$, there exists a unique $\sigma_1(\eta, t) \in [0, T-t]$ such that (figure 2)

$$X_F(1, t; \sigma_1(\eta, t)) = X_B(\eta, 0; t + \sigma_1(\eta, t)), \quad \sigma_1(\xi_0(t), t) = 0.$$

In particular, for each $t_* \in [t, T]$, $s_1(t, t_*) := \sigma_1(\xi_0(t_*), t)$ satisfies

$$X_F(1, t; s_1(t, t_*)) = X_B(\xi_0(t_*), 0; t + s_1(t, t_*)),$$

and thus the integrals in (6.6) is actually taken over $\sigma \in (0, s_1(t, t_*))$. Using the bootstrap bounds (3.1)-(3.11), we have the following lemma for σ_1 :

Lemma 6.3. *For each $t \in [0, T]$, $\sigma_1(\cdot, t)$ is a strictly increasing C^1 function on $(\xi_0(t), \xi_0(T))$. Moreover, for $\eta \in (\xi_0(t), \xi_0(T))$, there holds*

$$\begin{aligned} (2\bar{c})^{-1} e^{-C\kappa} & \leq \frac{d\sigma_1(\eta, t)}{d\eta} \leq (2\underline{c})^{-1} e^{C\kappa}, \\ 1 + \frac{\underline{c}e^{-C\kappa}}{\underline{c} + \bar{c}} (\eta - \xi_0(t_*)) & \leq X_F^{(\xi)}(1, t; \sigma_1(\eta, t)) \leq 1 + \frac{\bar{c}e^{C\kappa}}{\underline{c} + \bar{c}} (\eta - \xi_0(t_*)). \end{aligned}$$

Proof. The C^1 regularity of $\sigma_1(\cdot, t)$ is given by the implicit function theorem and the H^2 regularity of c . First, we claim that $\sigma(\eta, t)$ is strictly increasing in η . If assume that $\eta_2 > \eta_1$ and $\sigma_1(\eta_2, t) \leq \sigma_1(\eta_1, t)$, we find

$$\begin{aligned} X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_2, t)) &= X_F^{(\xi)}(1, t; \sigma_1(\eta_2, t)) \leq X_F^{(\xi)}(1, t; \sigma_1(\eta_1, t)) \\ &= X_B^{(\xi)}(\eta_1, 0; t + \sigma_1(\eta_1, t)) \leq X_B^{(\xi)}(\eta_1, 0; t + \sigma_1(\eta_2, t)), \end{aligned}$$

while $X_B^{(\xi)}(\eta_2, 0; 0) > X_B^{(\xi)}(\eta_1, 0; 0)$, which leads to contradiction with the uniqueness of integral curves. Hence $\sigma_1(\eta_2, t) > \sigma_1(\eta_1, t)$ and for the same reason $X_B^{(\xi)}(\eta_2, 0; \tau) > X_B^{(\xi)}(\eta_1, 0; \tau)$ for τ in the common interval of $X_B(\eta_1, 0; \cdot)$ and $X_B(\eta_2, 0; \cdot)$ if $\eta_2 > \eta_1$. By mean value formula and (6.3) we have for $\eta_2, \eta_1 \in [\xi_0(t), \xi_0(T)]$ with $\eta_2 > \eta_1$ that

$$\begin{aligned} \left| \frac{d}{d\tau} \left(X_B^{(\xi)}(\eta_2, 0; \tau) - X_B^{(\xi)}(\eta_1, 0; \tau) \right) \right| &= |(cr^2\xi^{-2})(X_B(\eta_1, 0; \tau)) - (cr^2\xi^{-2})(X_B(\eta_2, 0; \tau))| \\ &\leq C\epsilon X_B^{(\xi)}(\eta_1, 0; \tau)^{-1} \left(X_B^{(\xi)}(\eta_2, 0; \tau) - X_B^{(\xi)}(\eta_1, 0; \tau) \right), \end{aligned}$$

and thus by Gronwall's inequality

$$\begin{aligned} &(\eta_2 - \eta_1) \exp \left\{ - \int_0^{t+\sigma_1(\eta_1, t)} C\epsilon X_B^{(\xi)}(\eta_1, 0; \tau)^{-1} d\tau \right\} \\ &\leq X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_1, t)) - X_B^{(\xi)}(\eta_1, 0; t + \sigma_1(\eta_1, t)) \\ &\leq (\eta_2 - \eta_1) \exp \left\{ \int_0^{t+\sigma_1(\eta_1, t)} C\epsilon X_B^{(\xi)}(\eta_1, 0; \tau)^{-1} d\tau \right\}. \end{aligned}$$

Using the rough bound (3.10) $cr^2\xi^{-2} \geq \underline{c}$, we further obtain

$$\begin{aligned} &\int_0^{t+\sigma_1(\eta_1, t)} X_B^{(\xi)}(\eta_1, 0; \tau)^{-1} d\tau \\ &\leq \int_0^{t+\sigma_1(\eta_1, t)} \left(X_B^{(\xi)}(\eta_1, 0; t + \sigma_1(\eta_1, t)) + \underline{c}(t + \sigma_1(\eta_1, t) - \tau) \right)^{-1} d\tau \\ &\leq \frac{1}{\underline{c}} \log(1 + \underline{c}(t + \sigma_1(\eta_1, t))). \end{aligned}$$

Hence by the assumption (3.11) and that $\sigma_1(\eta_1, t) + t \leq T$, it follows that

$$\int_0^{t+\sigma_1(\eta_1, t)} C\epsilon X_B^{(\xi)}(\eta_1, 0; \tau)^{-1} d\tau \leq C\kappa,$$

therefore

$$e^{-C\kappa}(\eta_2 - \eta_1) \leq X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_1, t)) - X_B^{(\xi)}(\eta_1, 0; t + \sigma_1(\eta_1, t)) \leq e^{C\kappa}(\eta_2 - \eta_1). \quad (6.9)$$




Figure 2

By the rough bound (3.10) again, there holds (figure 2)

$$\begin{aligned}
& X_F^{(\xi)}(1, t; \sigma_1(\eta_1, t)) + \underline{c}(\sigma_1(\eta_2, t) - \sigma_1(\eta_1, t)) \leq X_F^{(\xi)}(1, t; \sigma_1(\eta_2, t)) \\
& = X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_2, t)) \leq X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_1, t)) - \underline{c}(\sigma_1(\eta_2, t) - \sigma_1(\eta_1, t)), \\
& X_F^{(\xi)}(1, t; \sigma_1(\eta_1, t)) + \bar{c}(\sigma_1(\eta_2, t) - \sigma_1(\eta_1, t)) \geq X_F^{(\xi)}(1, t; \sigma_1(\eta_2, t)) \\
& = X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_2, t)) \geq X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_1, t)) - \bar{c}(\sigma_1(\eta_2, t) - \sigma_1(\eta_1, t)).
\end{aligned}$$

Combining the above inequalities yields

$$\begin{aligned}
2\underline{c}(\sigma_1(\eta_2, t) - \sigma_1(\eta_1, t)) & \leq X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_1, t)) - X_F^{(\xi)}(1, t; \sigma_1(\eta_1, t)) \\
& = X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_1, t)) - X_B^{(\xi)}(\eta_1, 0; t + \sigma_1(\eta_1, t)) \\
& \leq e^{C\kappa}(\eta_2 - \eta_1), \\
2\bar{c}(\sigma_1(\eta_2, t) - \sigma_1(\eta_1, t)) & \geq X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_1, t)) - X_F^{(\xi)}(1, t; \sigma_1(\eta_1, t)) \\
& = X_B^{(\xi)}(\eta_2, 0; t + \sigma_1(\eta_1, t)) - X_B^{(\xi)}(\eta_1, 0; t + \sigma_1(\eta_1, t)) \\
& \geq e^{-C\kappa}(\eta_2 - \eta_1),
\end{aligned}$$

which gives the estimate of $\frac{d\sigma_1(\eta, t)}{d\eta}$ since η_2, η_1 can be arbitrary. Note that

$$\begin{aligned}
& \max \left\{ 1 + \underline{c}\sigma_1(\eta, t), X_B^{(\xi)}(\eta, 0; t) - \bar{c}\sigma_1(\eta, t) \right\} \\
& \leq X_F^{(\xi)}(1, t; \sigma_1(\eta, t)) = X_B^{(\xi)}(\eta, 0; t + \sigma_1(\eta, t)) \\
& \leq \min \left\{ 1 + \bar{c}\sigma_1(\eta, t), X_B^{(\xi)}(\eta, 0; t) - \underline{c}\sigma_1(\eta, t) \right\},
\end{aligned}$$

then taking $\eta_2 = \eta$, $\eta_1 = \xi_0(t)$ in (6.9) yields

$$e^{-C\kappa}(\eta - \xi_0(t)) \leq X_B^{(\xi)}(\eta, 0; t) - 1 \leq e^{C\kappa}(\eta - \xi_0(t)).$$

It follows

$$\begin{aligned} X_F^{(\xi)}(1, t; \sigma_1(\eta, t)) &\geq (\bar{c} + \underline{c})^{-1} \left(\bar{c} + \underline{c} X_B^{(\xi)}(\eta, 0; t) \right) = 1 + \frac{\underline{c} e^{-C\kappa}}{\underline{c} + \bar{c}} (\eta - \xi_0(t_*)), \\ X_F^{(\xi)}(1, t; \sigma_1(\eta, t)) &\leq (\bar{c} + \underline{c})^{-1} \left(\underline{c} + \bar{c} X_B^{(\xi)}(\eta, 0; t) \right) = 1 + \frac{\bar{c} e^{C\kappa}}{\underline{c} + \bar{c}} (\eta - \xi_0(t_*)). \end{aligned}$$

□

Using the bound (3.7) and Lemma 6.3, we estimate the second line in (6.6):

$$\begin{aligned} &\sup_s \int_0^s \left| \left(\frac{\partial_t c + cr^2 \xi^{-2} \partial_\xi c}{2c} \right) (X_F(1, t; \sigma)) \right| v_B(X_F(1, t; \sigma)) d\sigma \\ &\leq C\epsilon \int_0^{s_1(t, t_*)} X_F^{(\xi)}(1, t; \sigma)^{-1} v_B(X_F(1, t; \sigma)) d\sigma \\ &\leq \frac{C e^{C\kappa} \epsilon}{\underline{c}} \int_{\xi_0(t)}^{\xi_0(t_*)} X_F^{(\xi)}(1, t; \sigma_1(\eta, t))^{-1} v_B(X_B(\eta, 0; t + \sigma_1(\eta, t))) d\eta \\ &\leq \frac{C e^{C\kappa} \epsilon}{\underline{c}} \int_{\xi_0(t)}^{\xi_0(t_*)} \left(1 + \underline{c} (\bar{c} + \underline{c})^{-1} e^{-C\kappa} (\eta - \xi_0(t)) \right)^{-1} v_B(\eta, 0) d\eta. \end{aligned} \tag{6.10}$$

Recall that $w_B = (\partial_t + cr^2 \partial_x) \psi$. We then estimate the third line in (6.6) using (3.6):

$$\begin{aligned} &\sup_s \int_0^s \left| \left(2r^{-2} u^2 \frac{\psi}{c^{\frac{1}{2}}} \right) (X_F(1, t; \sigma)) \right| d\sigma \\ &\leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_1(t, t_*)} X_F^{(\xi)}(1, t; \sigma)^{-4} |\psi(X_F(1, t; \sigma))| d\sigma \\ &\leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_1(t, t_*)} X_F^{(\xi)}(1, t; \sigma)^{-4} \left| \psi(1, t) + \int_0^\sigma w_B(X_F(1, t; \tau)) d\tau \right| d\sigma \\ &\leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_1(t, t_*)} X_F^{(\xi)}(1, t; \sigma)^{-4} d\sigma |\psi(1, t)| \\ &\quad + \frac{C\epsilon^2 \bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{1}{2}}} \int_0^{s_1(t, t_*)} \int_0^\sigma X_F^{(\xi)}(1, t; \sigma)^{-4} v_B(X_F(1, t; \tau)) d\tau d\sigma, \end{aligned}$$

and exchange the order of integration:

$$\begin{aligned} &\int_0^{s_1(t, t_*)} \int_0^\sigma X_F^{(\xi)}(1, t; \sigma)^{-4} v_B(X_F(1, t; \tau)) d\tau d\sigma \\ &= \int_0^{s_1(t, t_*)} \left(\int_\tau^{s_1(t, t_*)} X_F^{(\xi)}(1, t; \sigma)^{-4} d\sigma \right) v_B(X_F(1, t; \tau)) d\tau. \end{aligned}$$

Use the rough bound (3.10):

$$\int_{\tau}^{s_1(t, t_*)} X_F^{(\xi)}(1, t; \sigma)^{-4} d\sigma \leq \int_{\tau}^{s_1(t, t_*)} \left(X_F^{(\xi)}(1, t; \tau) + \underline{c}(\sigma - \tau) \right)^{-4} d\sigma \leq \frac{1}{3\underline{c}} X_F^{(\xi)}(1, t; \tau)^{-3}.$$

We proceed the estimate by Lemma 6.3:

$$\begin{aligned} & \sup_s \int_0^s \left| \left(2r^{-2} u^2 \frac{\psi}{c^{\frac{1}{2}}} \right) (X_F(1, t; \sigma)) \right| d\sigma \\ & \leq \frac{C\epsilon^2}{\underline{c}^{\frac{3}{2}}} |\psi(1, t)| + \frac{C\epsilon^2 \bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{3}{2}}} \int_0^{s_1(t, t_*)} X_F^{(\xi)}(1, t; \tau)^{-3} v_B(X_F(1, t; \tau)) d\tau \\ & \leq \frac{C\epsilon^2}{\underline{c}^{\frac{3}{2}}} |\psi(1, t)| + \frac{C\epsilon^2 \bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{3}{2}}} (2\underline{c})^{-1} e^{C\kappa} \int_{\xi_0(t)}^{\xi_0(t_*)} X_F^{(\xi)}(1, t; \sigma_1(\eta, t))^{-3} v_B(X_B(\eta, 0; \sigma_1(\eta, t))) d\eta \\ & \leq \frac{C\epsilon^2}{\underline{c}^{\frac{3}{2}}} |\psi(1, t)| + \frac{C\epsilon^2 \bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{5}{2}}} e^{C\kappa} \int_{\xi_0(t)}^{\xi_0(t_*)} \left(1 + \frac{\underline{c}}{\bar{c} + \underline{c}} e^{-C\kappa} (\eta - \xi_0(t)) \right)^{-3} v_B(\eta, 0) d\eta. \end{aligned} \tag{6.11}$$

6.2. Estimate of (6.7)

By continuity, for each $\eta \in [\xi, \xi_0(T)]$, there exists a unique $\sigma_2(\eta, \xi) \in [0, T - t]$ such that (figure 3)

$$X_F(\xi, 0; \sigma_2(\eta, \xi)) = X_B(\eta, 0; \sigma_2(\eta, \xi)), \quad \sigma_2(\xi, \xi) = 0.$$

In particular, for each $t_* \leq T$ such that $\xi_0(t_*) \geq \xi$, $s_2(\xi, t_*) := \sigma_2(\xi_0(t_*), \xi)$ satisfies

$$X_F(\xi, 0; s_2(\xi, t_*)) = X_B(\xi_0(t_*), 0; s_2(\xi, t_*)),$$

so the integrals in (6.7) are taken over $\sigma \in (0, s_2(\xi, t_*))$.

Lemma 6.4. *For each $\xi \in [1, \xi_0(T)]$, $\sigma_2(\cdot, \xi)$ is a strictly increasing C^1 function on $(\xi, \xi_0(T))$. Moreover, for $\eta \in (\xi, \xi_0(T))$, there holds*

$$\begin{aligned} (2\bar{c})^{-1} e^{-C\kappa} & \leq \frac{d\sigma_2(\eta, \xi)}{d\eta} \leq (2\underline{c})^{-1} e^{C\kappa}, \\ \xi + \frac{\underline{c}}{\bar{c} + \underline{c}} (\eta - \xi) & \leq X_F^{(\xi)}(\xi, 0; \sigma_2(\eta, \xi)) \leq \xi + \frac{\bar{c}}{\bar{c} + \underline{c}} (\eta - \xi). \end{aligned}$$

Proof. The implicit function theorem and the H^2 regularity of c give the C^1 regularity of $\sigma_2(\cdot, \xi)$. Assume otherwise $\sigma_2(\eta_2, \xi) \leq \sigma_2(\eta_1, \xi)$ for some $\eta_2 > \eta_1$. Then

$$\begin{aligned} X_B^{(\xi)}(\eta_2, 0; \sigma_2(\eta_2, \xi)) & = X_F^{(\xi)}(\xi, 0; \sigma_2(\eta_2, \xi)) \leq X_F^{(\xi)}(\xi, 0; \sigma_2(\eta_1, \xi)) \\ & = X_B^{(\xi)}(\eta_1, 0; \sigma_2(\eta_1, \xi)) \leq X_B^{(\xi)}(\eta_1, 0; \sigma_2(\eta_2, \xi)), \end{aligned}$$

while $X_B^{(\xi)}(\eta_2, 0; 0) > X_B^{(\xi)}(\eta_1, 0; 0)$, which contradicts with the uniqueness of integral curves. Hence $\sigma_2(\eta_2, t) > \sigma_2(\eta_1, t)$ and for the same reason $X_B^{(\xi)}(\eta_2, 0; \tau) > X_B^{(\xi)}(\eta_1, 0; \tau)$ for τ in the common

The estimate of $X_F^{(\xi)}(\xi, 0; \sigma_2(\eta, \xi))$ follows from the the following inequalities:

$$\begin{aligned}
& \max \{ \xi + \underline{c}\sigma_2(\eta, \xi), \eta - \bar{c}\sigma_2(\eta, \xi) \} \\
& \leq X_F^{(\xi)}(\xi, 0; \sigma_2(\eta, \xi)) = X_B^{(\xi)}(\eta, 0; \sigma_2(\eta, \xi)) \\
& \leq \min \{ \xi + \bar{c}\sigma_2(\eta, \xi), \eta - \underline{c}\sigma_2(\eta, \xi) \}, \\
& \frac{\bar{c}}{\bar{c} + \underline{c}} (\xi + \underline{c}\sigma_2(\eta, \xi)) + \frac{\underline{c}}{\bar{c} + \underline{c}} (\eta - \bar{c}\sigma_2(\eta, \xi)) = \xi + \frac{\underline{c}}{\bar{c} + \underline{c}} (\eta - \xi), \\
& \frac{\underline{c}}{\bar{c} + \underline{c}} (\xi + \bar{c}\sigma_2(\eta, \xi)) + \frac{\bar{c}}{\bar{c} + \underline{c}} (\eta - \underline{c}\sigma_2(\eta, \xi)) = \xi + \frac{\bar{c}}{\bar{c} + \underline{c}} (\eta - \xi).
\end{aligned}$$

□

By the bound (3.7) and Lemma 6.4, the second line in (6.7) can be controlled by

$$\begin{aligned}
& \sup_s \int_0^s \left| \left(\frac{\partial_t c + cr^2 \xi^{-2} \partial_\xi c}{2c} \right) (X_F(\xi, 0; \sigma)) \right| v_B(X_F(\xi, 0; \sigma)) d\sigma \\
& \leq C\epsilon \int_0^{s_2(\xi, t_*)} X_F^{(\xi)}(\xi, 0; \sigma)^{-1} v_B(X_F(\xi, 0; \sigma)) d\sigma \\
& \leq \frac{C\epsilon^{C\kappa\epsilon}}{\underline{c}} \int_\xi^{\xi_0(T)} X_F^{(\xi)}(\xi, 0; \sigma_2(\eta, \xi))^{-1} v_B(X_B(\eta, 0; \sigma_2(\eta, \xi))) d\eta \\
& \leq \frac{C\epsilon^{C\kappa\epsilon}}{\underline{c}} \int_\xi^{\xi_0(T)} \left(\xi + \frac{\underline{c}}{\bar{c} + \underline{c}} (\eta - \xi) \right)^{-1} v_B(\eta, 0) d\eta.
\end{aligned} \tag{6.12}$$

Use the bound (3.6) and that $(\partial_t + cr^2 \partial_x) \psi = w_B$ to estimate the third line in (6.7):

$$\begin{aligned}
& \sup_s \int_0^s \left| \left(2r^{-2} u^2 \frac{\psi}{c^{\frac{3}{2}}} \right) (X_F(\xi, 0; \sigma)) \right| d\sigma \\
& \leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_2(\xi, t_*)} X_F^{(\xi)}(\xi, 0; \sigma)^{-4} |\psi(X_F(\xi, 0; \sigma))| d\sigma \\
& \leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_2(\xi, t_*)} X_F^{(\xi)}(\xi, 0; \sigma)^{-4} \left| \psi(\xi, 0) + \int_0^\sigma w_B(X_F(\xi, 0; \tau)) d\tau \right| d\sigma \\
& \leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_2(\xi, t_*)} X_F^{(\xi)}(\xi, 0; \sigma)^{-4} d\sigma |\psi(\xi, 0)| \\
& \quad + \frac{C\epsilon^2 \bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{1}{2}}} \int_0^{s_2(\xi, t_*)} \int_0^\sigma X_F^{(\xi)}(\xi, 0; \sigma)^{-4} v_B(X_F(\xi, 0; \tau)) d\tau d\sigma \\
& = \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_2(\xi, t_*)} X_F^{(\xi)}(\xi, 0; \sigma)^{-4} d\sigma |\psi(\xi, 0)| \\
& \quad + \frac{C\epsilon^2 \bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{1}{2}}} \int_0^{s_2(\xi, t_*)} \left(\int_\tau^{s_2(\xi, t_*)} X_F^{(\xi)}(\xi, 0; \sigma)^{-4} d\sigma \right) v_B(X_F(\xi, 0; \tau)) d\tau.
\end{aligned}$$

Note that the rough bound (3.10) yields

$$\int_{\tau}^{s_2(\xi, t_*)} X_F^{(\xi)}(\xi, 0; \sigma)^{-4} d\sigma \leq \int_{\tau}^{s_2(\xi, t_*)} \left(X_F^{(\xi)}(\xi, 0; \tau) + \underline{c}(\sigma - \tau) \right)^{-4} d\sigma \leq \frac{1}{3\underline{c}} X_F^{(\xi)}(\xi, 0; \tau)^{-3}.$$

Then we proceed by Lemma 6.4 and obtain

$$\begin{aligned} & \sup_s \int_0^s \left| \left(2r^{-2} u^2 \frac{\psi}{c^{\frac{1}{2}}} \right) (X_F(\xi, 0; \sigma)) \right| d\sigma \\ & \leq \frac{C\epsilon^2}{\underline{c}^{\frac{3}{2}}} \xi^{-3} |\psi(\xi, 0)| + \frac{C\epsilon^2 \bar{c}}{\underline{c}^{\frac{3}{2}}} \int_0^{s_2(\xi, t_*)} X_F^{(\xi)}(\xi, 0; \tau)^{-3} v_B(X_F(\xi, 0; \tau)) d\tau \\ & \leq \frac{C\epsilon^2}{\underline{c}^{\frac{3}{2}}} \xi^{-3} |\psi(\xi, 0)| + \frac{C\epsilon^2 \bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{3}{2}}} (2\underline{c})^{-1} e^{C\kappa} \int_{\xi}^{\xi_0(t_*)} X_F^{(\xi)}(\eta, 0; \sigma_2(\eta, \xi))^{-3} v_B(X_B(\eta, 0; \sigma_2(\eta, \xi))) d\eta \\ & \leq \frac{C\epsilon^2}{\underline{c}^{\frac{3}{2}}} \xi^{-3} |\psi(\xi, 0)| + \frac{C\epsilon^2 \bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{5}{2}}} e^{C\kappa} \int_{\xi}^{\xi_0(t_*)} \left(\xi + \underline{c}(\bar{c} + \underline{c})^{-1}(\eta - \xi) \right)^{-3} v_B(\eta, 0) d\eta. \end{aligned} \tag{6.13}$$

6.3. Estimate of (6.8)

In fact, we will take $\xi = \xi_0(t_*)$ in (6.8) for the later analysis, and thus it is convenient to establish the estimate with ξ replaced by $\xi_0(t_*)$ in (6.8) with $t_* \in [0, T]$, in which case the integrals in (6.8) is in fact taken over $\sigma \in (0, t_*)$ by the construction of $\xi_0(\cdot)$. For each $\eta \in [1, \xi_0(t_*)]$, by continuity there exists a unique $\sigma_3(\eta, t_*) \in [0, t_*]$ such that (figure 4)

$$X_F(\eta, 0; \sigma_3(\eta, t_*)) = X_B(\xi_0(t_*), 0; \sigma_3(\eta, t_*)), \quad \sigma_3(\xi_0(t_*), t_*) = 0.$$

In particular, $s_3(t_*) := \sigma_3(1, t_*)$ satisfies

$$X_F(1, 0; s_3(t_*)) = X_B(\xi_0(t_*), 0; s_3(t_*)).$$

Analogously, for each $t \in [0, t_*]$ there exists a unique $\sigma_4(t, t_*) \in [t, t_*]$ such that (figure 5)

$$X_F(1, t; \sigma_4(t, t_*) - t) = X_B(\xi_0(t_*), 0; \sigma_4(t, t_*)), \quad \sigma_4(t_*, t_*) = t_*, \quad \sigma_4(0, t_*) = s_3(t_*).$$

Recall that $X_F(1, t; \sigma_1(\eta, t)) = X_B(\eta, 0; t + \sigma_1(\eta, t))$ for $t \in [0, t]$ and $\eta \in [\xi_0(t), \xi_0(t_*)]$. Hence by taking $\eta = \xi_0(t_*)$ we have in particular

$$\sigma_1(\xi_0(t_*), t) = \sigma_4(t, t_*) - t.$$

Lemma 6.5. *For each $t_* \in [0, T]$, $\sigma_3(\cdot, t_*)$ is a strictly decreasing C^1 function on $(1, \xi_0(t_*))$, and $\sigma_4(\cdot, t_*)$ is a strictly decreasing C^1 function on $(0, t_*)$. Moreover, the following estimates hold:*

$$\begin{aligned} (2\bar{c})^{-1} e^{-C\kappa} & \leq -\frac{d\sigma_3(\eta, t_*)}{d\eta} \leq (2\underline{c})^{-1} e^{C\kappa}, \quad \eta \in (1, \xi_0(t_*)), \\ (2\bar{c})^{-1} \underline{c} e^{-C\kappa} & \leq \frac{d\sigma_4(t, t_*)}{dt} \leq (2\underline{c})^{-1} \bar{c} e^{C\kappa}, \quad t \in (0, t_*), \\ 1 + \frac{\underline{c}}{\bar{c} + \underline{c}} (\xi_0(t_*) - 1) & \leq \eta + \frac{\underline{c}}{\bar{c} + \underline{c}} (\xi_0(t_*) - \eta) \leq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta, t_*)) \leq \eta + \frac{\underline{c}}{\bar{c} + \underline{c}} (\xi_0(t_*) - \eta), \\ 1 + \frac{\underline{c}}{2} (t_* - t) & \leq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*)) \leq 1 + \frac{\bar{c}}{2} (t_* - t). \end{aligned}$$

Proof. The C^1 regularity of $\sigma_3(\cdot, t_*)$ and $\sigma_4(\cdot, t_*)$ follows from the implicit function theorem and the H^2 regularity of c . Let $\eta_1, \eta_2 \in [1, \xi_0(t_*)]$ with $\eta_2 > \eta_1$ and $t_1, t_2 \in [0, t_*]$ with $t_2 > t_1$. To show the monotonicity, assume on the contrary that $\sigma_3(\eta_2, t_*) \geq \sigma_3(\eta_1, t_*)$. Then

$$\begin{aligned} X_F^{(\xi)}(\eta_1, 0; \sigma_3(\eta_1, t_*)) &= X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta_1, t_*)) \geq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta_2, t_*)) \\ &= X_F^{(\xi)}(\eta_2, 0; \sigma_3(\eta_2, t_*)) \geq X_F^{(\xi)}(\eta_2, 0; \sigma_3(\eta_1, t_*)), \end{aligned}$$

while $X_F(\eta_1, 0; 0) < X_F(\eta_2, 0; 0)$, which is impossible by the uniqueness of integral curves. Similarly, if we assume $\sigma_4(t_1, t_*) \geq \sigma_4(t_2, t_*)$, then the inequalities

$$\begin{aligned} X_F^{(\xi)}(1, t_2; \sigma_4(t_2, t_*) - t_2) &= X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t_2, t_*)) \geq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t_1, t_*)) \\ &= X_F^{(\xi)}(1, t_1; \sigma_4(t_1, t_*) - t_1) \geq X_F^{(\xi)}(1, t_1; \sigma_4(t_2, t_*) - t_1) = X_F^{(\xi)}(X_F(1, t_1; t_2 - t_1); \sigma_4(t_2, t_*) - t_2), \\ 1 &= X_F^{(\xi)}(1, t_2; 0) < X_F^{(\xi)}(1, t_1; t_2 - t_1) = X_F^{(\xi)}(X_F(1, t_1; t_2 - t_1); 0) \end{aligned}$$

lead to contradiction. By recalling (3.6)(3.7) we have

$$\begin{aligned} \left| \frac{d}{d\tau} \left(X_F^{(\xi)}(\eta_2, 0; \tau) - X_F^{(\xi)}(\eta_1, 0; \tau) \right) \right| &= \left| (cr^2\xi^{-2})(X_F(\eta_2, 0; \tau)) - (cr^2\xi^{-2})(X_F(\eta_1, 0; \tau)) \right| \\ &\leq C\epsilon X_F^{(\xi)}(\eta_1, 0; \tau)^{-1} \left(X_F^{(\xi)}(\eta_2, 0; \tau) - X_F^{(\xi)}(\eta_1, 0; \tau) \right). \end{aligned}$$




Figure 4

By integrating the above inequality and noting that (3.11) gives

$$\begin{aligned} C\epsilon \int_0^{\sigma_3(\eta_2, t_*)} X_F^{(\xi)}(\eta_1, 0; \tau)^{-1} d\tau &\leq C\epsilon \int_0^{\sigma_3(\eta_2, t_*)} \left(X_F^{(\xi)}(\eta_1, 0; 0) + \underline{c}\tau \right)^{-1} d\tau \\ &\leq \frac{C\epsilon}{\underline{c}} \log(1 + \eta_1^{-1} \underline{c}\sigma_3(\eta_2, t_*)) \\ &\leq C\kappa, \end{aligned}$$

we obtain

$$e^{-C\kappa}(\eta_2 - \eta_1) \leq X_F^{(\xi)}(\eta_2, 0; \sigma_3(\eta_2, t_*)) - X_F^{(\xi)}(\eta_1, 0; \sigma_3(\eta_1, t_*)) \leq e^{C\kappa}(\eta_2 - \eta_1).$$

Use the rough bound (3.10) (figure 4):

$$\begin{aligned} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta_2, t_*)) - \bar{c}(\sigma_3(\eta_1, t_*) - \sigma_3(\eta_2, t_*)) &\leq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta_1, t_*)) \\ = X_F^{(\xi)}(\eta_1, 0; \sigma_3(\eta_1, t_*)) &\leq X_F^{(\xi)}(\eta_1, 0; \sigma_3(\eta_2)) + \bar{c}(\sigma_3(\eta_1, t_*) - \sigma_3(\eta_2, t_*)), \\ X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta_2, t_*)) - \underline{c}(\sigma_3(\eta_1, t_*) - \sigma_3(\eta_2, t_*)) &\geq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta_1, t_*)) \\ = X_F^{(\xi)}(\eta_1, 0; \sigma_3(\eta_1, t_*)) &\geq X_F^{(\xi)}(\eta_1, 0; \sigma_3(\eta_2)) + \underline{c}(\sigma_3(\eta_1, t_*) - \sigma_3(\eta_2, t_*)). \end{aligned}$$




Figure 5

Combining the above inequalities and recalling $X_B(\xi_0(t_*), 0; \sigma_3(\eta_2, t_*)) = X_F(\eta_2, 0; \sigma_3(\eta_2, t_*))$, it follows

$$(2\bar{c})^{-1} e^{-C\kappa}(\eta_2 - \eta_1) \leq \sigma_3(\eta_1, t_*) - \sigma_3(\eta_2, t_*) \leq (2\underline{c})^{-1} e^{C\kappa}(\eta_2 - \eta_1),$$

which in turn gives the expected estimate of $\frac{d\sigma_3(\eta, t_*)}{d\eta}$. To obtain the estimate of $\frac{d\sigma_4(t, t_*)}{dt}$, using the relation

$$X_F^{(\xi)}(1, t_1; \sigma_4(t_1, t_*) - t_1) = X_F^{(\xi)}(X_F(1, t_1; t_2 - t_1); \sigma_4(t_1, t_*) - t_2),$$

it can be shown in a similar way that

$$\begin{aligned} e^{-C\kappa} \left(X_F^{(\xi)}(1, t_1; t_2 - t_1) - 1 \right) &\leq X_F^{(\xi)}(1, t_1; \sigma_4(t_1, t_*) - t_1) - X_F^{(\xi)}(1, t_2; \sigma_4(t_1, t_*) - t_2) \\ &\leq e^{C\kappa} \left(X_F^{(\xi)}(1, t_1; t_2 - t_1) - 1 \right). \end{aligned}$$

The rough bound (3.10) yields (figure 5)

$$\begin{aligned} \underline{c}(t_2 - t_1) &\leq X_F^{(\xi)}(1, t_1; t_2 - t_1) - 1 \leq \bar{c}(t_2 - t_1), \\ X_F^{(\xi)}(1, t_2; \sigma_4(t_1, t_*) - t_2) + \underline{c}(\sigma_4(t_2, t_*) - \sigma_4(t_1, t_*)) &\leq X_F^{(\xi)}(1, t_2; \sigma_4(t_2, t_*)) \\ = X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t_2, t_*)) &\leq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t_1, t_*)) - \underline{c}(\sigma_4(t_2, t_*) - \sigma_4(t_1, t_*)), \\ X_F^{(\xi)}(1, t_2; \sigma_4(t_1, t_*) - t_2) + \bar{c}(\sigma_4(t_2, t_*) - \sigma_4(t_1, t_*)) &\geq X_F^{(\xi)}(1, t_2; \sigma_4(t_2, t_*)) \\ = X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t_2, t_*)) &\geq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t_1, t_*)) - \bar{c}(\sigma_4(t_2, t_*) - \sigma_4(t_1, t_*)). \end{aligned}$$

Combining the above inequalities and using the relation

$$X_B(\xi_0(t_*), 0; \sigma_4(t_1, t_*)) = X_F(1, t_1; \sigma_4(t_1, t_*) - t_1),$$

we obtain

$$(2\bar{c})^{-1} \underline{c} e^{-C\kappa} (t_2 - t_1) \leq \sigma_4(t_2, t_*) - \sigma_4(t_1, t_*) \leq (2\underline{c})^{-1} \bar{c} e^{C\kappa},$$

which gives the claimed estimate of $\frac{d\sigma_4(t, t_*)}{dt}$. Next, we have by the rough bound (3.10)

$$\begin{aligned} &\max \{ \xi_0(t_*) - \bar{c}\sigma_3(\eta, t_*), \eta + \underline{c}\sigma_3(\eta, t_*) \} \\ &\leq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta, t_*)) = X_F^{(\xi)}(\eta, 0; \sigma_3(\eta, t_*)) \\ &\leq \min \{ \xi_0(t_*) - \underline{c}\sigma_3(\eta, t_*), \eta + \bar{c}\sigma_3(\eta, t_*) \}, \end{aligned}$$

and thus

$$1 + \frac{\underline{c}}{\bar{c} + \underline{c}} (\xi_0(t_*) - 1) \leq \eta + \frac{\underline{c}}{\bar{c} + \underline{c}} (\xi_0(t_*) - \eta) \leq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta, t_*)) \leq \eta + \frac{\bar{c}}{\bar{c} + \underline{c}} (\xi_0(t_*) - \eta).$$

For $X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*))$, recall that $X_B(\xi_0(t_*), 0; t_*) = (1, t_*)$, and thus

$$1 + \underline{c}(t_* - \sigma_4(t, t_*)) \leq X_B(\xi_0(t_*), 0; \sigma_4(t, t_*)) = X_B(1, t_*; \sigma_4(t, t_*) - t_*) \leq 1 + \bar{c}(t_* - \sigma_4(t, t_*)).$$

It follows that

$$\begin{aligned} &\max \{ 1 + \underline{c}(t_* - \sigma_4(t, t_*)), 1 + \underline{c}(\sigma_4(t, t_*) - t) \} \\ &\leq X_F^{(\xi)}(1, t; \sigma_4(t, t_*)) = X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*)) \\ &\leq \min \{ 1 + \bar{c}(t_* - \sigma_4(t, t_*)), 1 + \bar{c}(\sigma_4(t, t_*) - t) \}. \end{aligned}$$

Therefore

$$1 + \frac{\underline{c}}{2}(t_* - t) \leq X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*)) \leq 1 + \frac{\bar{c}}{2}(t_* - t).$$

□

Now we estimate the second line in (6.8) by using (3.7), Lemma 6.5 and splitting the integral into two parts in which $\sigma \in (0, s_3(t_*))$ and $\sigma \in (s_3(t_*), t_*)$:

$$\begin{aligned}
& \int_0^{s_3(t_*)} \left| \frac{(\partial_t - cr^2\xi^{-2}\partial_\xi)c}{2c} (X_B(\xi_0(t_*), 0; \sigma)) \right| v_F(X_B(\xi_0(t_*), 0; \sigma); t_*) d\sigma \\
& \leq C\epsilon \int_0^{s_3(t_*)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma)^{-1} v_F(X_B(\xi_0(t_*), 0; \sigma); t_*) d\sigma \\
& \leq \frac{C\epsilon^{C\kappa}\epsilon}{\underline{c}} \int_1^{\xi_0(t_*)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_3(\eta, t_*))^{-1} v_F(X_F(\eta, 0; \sigma_3(\eta, t_*)); t_*) d\eta \\
& \leq \frac{C\epsilon^{C\kappa}\epsilon}{\underline{c}} \int_1^{\xi_0(t_*)} \left(1 + \frac{\underline{c}}{\bar{c} + \underline{c}}(\xi_0(t_*) - 1)\right)^{-1} v_F(\eta, 0; t_*) d\eta,
\end{aligned} \tag{6.14}$$

$$\begin{aligned}
& \int_{s_3(t_*)}^{t_*} \left| \frac{(\partial_t - cr^2\xi^{-2}\partial_\xi)c}{2c} (X_B(\xi_0(t_*), 0; \sigma)) \right| v_F(X_B(\xi_0(t_*), 0; \sigma); t_*) d\sigma \\
& \leq C\epsilon \int_{s_3(t_*)}^{t_*} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma)^{-1} v_F(X_B(\xi_0(t_*), 0; \sigma); t_*) d\sigma \\
& \leq \frac{C\bar{c}\epsilon^{C\kappa}\epsilon}{\underline{c}} \int_0^{t_*} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*))^{-1} v_F(X_F(\eta, 0; \sigma_4(t, t_*)); t_*) dt \\
& \leq \frac{C\bar{c}\epsilon^{C\kappa}\epsilon}{\underline{c}} \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-1} v_F(\eta, 0; t_*) dt.
\end{aligned} \tag{6.15}$$

We also split the integral in the third line of (6.8) into two parts:

$$\begin{aligned}
& \int_0^{t_*} \left| 2r^{-2}u^2 \frac{\psi}{c^{\frac{1}{2}}} (X_B(\xi_0(t_*), 0; \sigma)) \right| d\sigma \\
& = \int_0^{s_3(t_*)} \left| 2r^{-2}u^2 \frac{\psi}{c^{\frac{1}{2}}} (X_B(\xi_0(t_*), 0; \sigma)) \right| d\sigma + \int_{s_3(t_*)}^{t_*} \left| 2r^{-2}u^2 \frac{\psi}{c^{\frac{1}{2}}} (X_B(\xi_0(t_*), 0; \sigma)) \right| d\sigma.
\end{aligned}$$

We use $(\partial_t - cr^2\partial_x)\psi = w_F$ and (3.6)(3.10) to estimate the first part:

$$\begin{aligned}
& \int_0^{s_3(t_*)} \left| 2r^{-2}u^2 \frac{\psi}{c^{\frac{1}{2}}} (X_B(\xi_0(t_*), 0; \sigma)) \right| d\sigma \\
& \leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_3(t_*)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma)^{-4} \left| \psi(\xi_0(t_*)) + \int_0^\sigma w_F(X_B(\xi_0(t_*), 0; \tau)) d\tau \right| d\sigma \\
& \leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_3(t_*)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma)^{-4} |\psi(\xi_0(t_*))| d\sigma \\
& \quad + \frac{C\epsilon^2\bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{1}{2}}} \int_0^{s_3(t_*)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma)^{-4} \int_0^\sigma v_F(X_B(\xi_0(t_*), 0; \tau); t_*) d\tau d\sigma \\
& \leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_0^{s_3(t_*)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma)^{-4} |\psi(\xi_0(t_*))| d\sigma \\
& \quad + \frac{C\epsilon^2\bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{1}{2}}} \int_0^{s_3(t_*)} \left(\int_\tau^{s_3(t_*)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma)^{-4} d\sigma \right) v_F(X_B(\xi_0(t_*), 0; \tau); t_*) d\tau.
\end{aligned}$$

Use the rough bound (3.10) and Lemma 6.5:

$$\begin{aligned} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma) &\geq X_B^{(\xi)}(\xi_0(t_*), 0; s_3(t_*)) + \underline{c}(s_3(t_*) - \sigma), \\ \int_{\tau}^{s_3(t_*)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma)^{-4} d\sigma &\leq \frac{1}{3\underline{c}} X_B^{(\xi)}(\xi_0(t_*), 0; s_3(t_*))^{-3} \\ &\leq \frac{1}{3\underline{c}} \left(1 + \frac{\underline{c}}{\bar{c} + \underline{c}}(\xi_0(t_*) - 1)\right)^{-3}. \end{aligned}$$

Therefore

$$\begin{aligned} &\int_0^{s_3(t_*)} \left| 2r^{-2}u^2 \frac{\psi}{c^{\frac{1}{2}}} (X_B(\xi_0(t_*), 0; \sigma)) \right| d\sigma \\ &\leq \frac{C\epsilon^2}{\underline{c}^{\frac{3}{2}}} \left(1 + \frac{\underline{c}}{\bar{c} + \underline{c}}(\xi_0(t_*) - 1)\right)^{-3} |\psi(\xi_0(t_*))| \\ &\quad + \frac{C\epsilon^2\bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{3}{2}}} \left(1 + \frac{\underline{c}}{\bar{c} + \underline{c}}(\xi_0(t_*) - 1)\right)^{-3} \int_0^{s_3(t_*)} v_F(X_B(\xi_0(t_*), 0; \tau); t_*) d\tau \quad (6.16) \\ &\leq \frac{C\epsilon^2}{\underline{c}^{\frac{3}{2}}} \left(1 + \frac{\underline{c}}{\bar{c} + \underline{c}}(\xi_0(t_*) - 1)\right)^{-3} |\psi(\xi_0(t_*))| \\ &\quad + \frac{C\epsilon^2\bar{c}^{\frac{1}{2}}}{\underline{c}^{\frac{5}{2}}} e^{C\kappa} \left(1 + \frac{\underline{c}}{\bar{c} + \underline{c}}(\xi_0(t_*) - 1)\right)^{-3} \int_1^{\xi_0(t_*)} v_F(\eta, 0; t_*) d\eta. \end{aligned}$$

For the second part, we first use $(\partial_t + cr^2\partial_x)\psi = w_B$ (3.6) and Lemma 6.5:

$$\begin{aligned} &\int_{s_3(t_*)}^{t_*} \left| 2r^{-2}u^2 \frac{\psi}{c^{\frac{1}{2}}} (X_B(\xi_0(t_*), 0; \sigma)) \right| d\sigma \\ &\leq \frac{C\epsilon^2}{\underline{c}^{\frac{1}{2}}} \int_{s_3(t_*)}^{t_*} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma)^{-4} |\psi(X_B(\xi_0(t_*), 0; \sigma))| d\sigma \\ &\leq \frac{C\epsilon^2\bar{c}}{\underline{c}^{\frac{3}{2}}} e^{C\kappa} \int_0^{t_*} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*))^{-4} |\psi(X_F(1, t; \sigma_4(t, t_*) - t))| dt \\ &\leq \frac{C\epsilon^2\bar{c}}{\underline{c}^{\frac{3}{2}}} e^{C\kappa} \int_0^{t_*} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*))^{-4} \left| \psi(1, t) + \int_0^{\sigma_4(t, t_*) - t} w_B(X_F(1, t; \tau)) d\tau \right| dt \\ &\leq \frac{C\epsilon^2\bar{c}}{\underline{c}^{\frac{3}{2}}} e^{C\kappa} \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-4} |\psi(1, t)| dt \\ &\quad + \frac{C\epsilon^2\bar{c}^{\frac{3}{2}}}{\underline{c}^{\frac{3}{2}}} e^{C\kappa} \int_0^{t_*} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*))^{-4} \int_0^{\sigma_4(t, t_*) - t} v_B(X_F(1, t; \tau)) d\tau dt. \end{aligned}$$

We further apply Lemma 6.3 and recall $\sigma_4(t, t_*) - t = \sigma_1(\xi_0(t_*), t)$:

$$\begin{aligned} \int_0^{\sigma_4(t, t_*) - t} v_B(X_F(1, t; \tau)) d\tau dt &\leq \frac{e^{C\kappa}}{2\underline{c}} \int_{\xi_0(t)}^{\xi_0(t_*)} v_B(X_B(\eta, 0; t + \sigma_1(\eta, t))) d\eta \\ &= \frac{e^{C\kappa}}{2\underline{c}} \int_{\xi_0(t)}^{\xi_0(t_*)} v_B(\eta, 0) d\eta, \end{aligned}$$

$$\begin{aligned}
& \int_0^{t_*} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*))^{-4} \int_0^{\sigma_4(t, t_*)-t} v_B(X_F(1, t; \tau)) d\tau dt \\
& \leq \frac{e^{C\kappa}}{2\underline{c}} \int_0^{t_*} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*))^{-4} \int_{\xi_0(t)}^{\xi_0(t_*)} v_B(\eta, 0) d\eta dt \\
& = \frac{e^{C\kappa}}{2\underline{c}} \int_1^{\xi_0(t_*)} \left(\int_0^{\xi_0^{-1}(\eta)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*))^{-4} dt \right) v_B(\eta, 0) d\eta,
\end{aligned}$$

where ξ_0^{-1} denotes the inverse function of ξ_0 . Use Lemma 6.5 to estimate the integral in the bracket:

$$\begin{aligned}
\int_0^{\xi_0^{-1}(\eta)} X_B^{(\xi)}(\xi_0(t_*), 0; \sigma_4(t, t_*))^{-4} dt & \leq \int_0^{\xi_0^{-1}(\eta)} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-4} dt \\
& \leq \frac{2}{3\underline{c}} \left(1 + \frac{\underline{c}}{2}(t_* - \xi_0^{-1}(\eta))\right)^{-3}.
\end{aligned}$$

Combining the above inequalities yields

$$\begin{aligned}
& \int_{s_3(t_*)}^{t_*} \left| 2r^{-2}u^2 \frac{\psi}{c^{\frac{1}{2}}}(X_B(\xi_0(t_*), 0; \sigma)) \right| d\sigma \\
& \leq \frac{C\epsilon^2\bar{c}}{\underline{c}^{\frac{3}{2}}} e^{C\kappa} \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-4} |\psi(1, t)| dt \\
& \quad + \frac{C\epsilon^2\bar{c}^{\frac{3}{2}}}{\underline{c}^{\frac{7}{2}}} e^{C\kappa} \int_1^{\xi_0(t_*)} \left(1 + \frac{\underline{c}}{2}(t_* - \xi_0^{-1}(\eta))\right)^{-3} v_B(\eta, 0) d\eta.
\end{aligned} \tag{6.17}$$

6.4. Collection of (6.6)-(6.17) and the estimate of $w_B(1, t)$

We now collect all the bounds obtained in this section. First we note in addition that from the bound (3.6)(3.10) and the smallness $\epsilon \leq \epsilon_0 \lesssim 1$ it follows

$$\begin{aligned}
& \exp \left\{ \int_0^s \left| \frac{2u}{r}(X_F(1, t; \sigma)) \right| d\sigma \right\} \leq \exp \left\{ C\epsilon \int_0^s (1 + \underline{c}\sigma)^{-2} d\sigma \right\} \leq e^{C\epsilon} \leq 1 + C\epsilon, \\
& \exp \left\{ \int_0^s \left| \frac{2u}{r}(X_F(\xi, 0; \sigma)) \right| d\sigma \right\} \leq \exp \left\{ C\epsilon \int_0^s (\xi + \underline{c}\sigma)^{-2} d\sigma \right\} \leq e^{C\epsilon} \leq 1 + C\epsilon, \\
& \exp \left\{ \int_0^{t_*} \left| \frac{2u}{r}(X_B(\xi_0(t_*), 0; \sigma)) \right| d\sigma \right\} \leq \exp \left\{ C\epsilon \int_0^{t_*} (1 + \underline{c}\sigma)^{-2} d\sigma \right\} \leq e^{C\epsilon} \leq 1 + C\epsilon.
\end{aligned}$$

For simplicity we write $\mu := \frac{\bar{c}}{\underline{c}} e^{C\kappa} > 1$. Combining (6.6)(6.10)(6.11) gives

$$v_F(1, t; t_*) \leq (1 + C\epsilon) \left| \frac{w_F}{c^{\frac{1}{2}}}(1, t) \right| + C\epsilon \int_{\xi_0(t)}^{\xi_0(t_*)} \left(1 + \frac{1}{2\mu}(\eta - \xi_0(t))\right)^{-1} v_B(\eta, 0) d\eta + C\epsilon^2 |\psi(1, t)|. \tag{6.18}$$

Analogously, from (6.7)(6.12)(6.13) there holds

$$v_F(\xi, 0; t_*) \leq (1 + C\epsilon) \left| \frac{w_F}{c^{\frac{1}{2}}}(\xi, 0) \right| + C\epsilon \int_{\xi}^{\xi_0(t_*)} \left(\xi + \frac{1}{2\mu}(\eta - \xi)\right)^{-1} v_B(\eta, 0) d\eta + C\epsilon^2 \xi^{-3} |\psi(\xi, 0)|, \tag{6.19}$$

Since $1 + \frac{\underline{c}}{\bar{c} + \underline{c}}(\xi_0(t_*) - 1) \geq \frac{\underline{c}}{\bar{c} + \underline{c}}\xi_0(t_*)$, from (6.8), (6.14)-(6.17) it follows

$$\begin{aligned}
& v_B(\xi_0(t_*), 0) \\
& \leq (1 + C\epsilon) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| + C\epsilon^2 \xi_0(t_*)^{-3} |\psi(\xi_0(t_*))| + C\epsilon^2 \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-4} |\psi(1, t)| dt \\
& \quad + C\epsilon \xi_0(t_*)^{-1} \int_1^{\xi_0(t_*)} v_F(\xi, 0; t_*) d\xi + C\epsilon \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-1} v_F(1, t; t_*) dt \\
& \quad + C\epsilon^2 \int_1^{\xi_0(t_*)} \left(1 + \frac{\underline{c}}{2}(t_* - \xi_0^{-1}(\eta))\right)^{-3} v_B(\eta, 0) d\eta.
\end{aligned} \tag{6.20}$$

We further call upon (6.5) to obtain

$$\begin{aligned}
& \left| \frac{w_B}{c^{\frac{1}{2}}}(1, t_*) - \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \\
& \leq C\epsilon |w_B(\xi_0(t_*), 0)| + C\epsilon^2 \xi_0(t_*)^{-3} |\psi(\xi_0(t_*))| + C\epsilon^2 \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-4} |\psi(1, t)| dt \\
& \quad + C\epsilon \xi_0(t_*)^{-1} \int_1^{\xi_0(t_*)} v_F(\xi, 0; t_*) d\xi + C\epsilon \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-1} v_F(1, t; t_*) dt \\
& \quad + C\epsilon^2 \int_1^{\xi_0(t_*)} \left(1 + \frac{\underline{c}}{2}(t_* - \xi_0^{-1}(\eta))\right)^{-3} v_B(\eta, 0) d\eta.
\end{aligned} \tag{6.21}$$

To obtain an explicit control of the last terms in (6.20) and (6.21), we have the following control of ξ_0^{-1} :

Lemma 6.6. *For each $\eta_1, \eta_2 \in [1, \xi_0(T)]$ with $\eta_1 < \eta_2$, the following inequality holds:*

$$\frac{e^{C\kappa}}{\underline{c}}(\eta_2 - \eta_1) \geq \xi_0^{-1}(\eta_2) - \xi_0^{-1}(\eta_1) \geq \frac{e^{-C\kappa}}{\bar{c}}(\eta_2 - \eta_1).$$

In particular, it follows that

$$\begin{aligned}
\frac{e^{C\kappa}}{\underline{c}} & \geq \frac{d\xi_0^{-1}}{d\eta} \geq \frac{e^{-C\kappa}}{\bar{c}}, \\
t_* - \xi_0^{-1}(\eta) & \geq \frac{e^{-C\kappa}}{\bar{c}}(\xi_0(t_*) - \eta).
\end{aligned}$$

Proof. By simulating the proof of Lemma 6.3 we have

$$e^{C\kappa}(\eta_2 - \eta_1) \geq X_B^{(\xi)}(\eta_2, 0; \xi_0^{-1}(\eta_1)) - X_B^{(\xi)}(\eta_1, 0; \xi_0^{-1}(\eta_1)) \geq e^{-C\kappa}(\eta_2 - \eta_1).$$

Then use the rough bound (3.10) (figure 6):

$$\begin{aligned}
\xi_0^{-1}(\eta_2) - \xi_0^{-1}(\eta_1) & \geq \frac{1}{\underline{c}} \left(X_B^{(\xi)}(\eta_2, 0; \xi_0^{-1}(\eta_1)) - X_B^{(\xi)}(\eta_1, 0; \xi_0^{-1}(\eta_1)) \right) \geq \frac{e^{-C\kappa}}{\underline{c}}(\eta_2 - \eta_1), \\
\xi_0^{-1}(\eta_2) - \xi_0^{-1}(\eta_1) & \leq \frac{1}{\underline{c}} \left(X_B^{(\xi)}(\eta_2, 0; \xi_0^{-1}(\eta_1)) - X_B^{(\xi)}(\eta_1, 0; \xi_0^{-1}(\eta_1)) \right) \leq \frac{e^{C\kappa}}{\underline{c}}(\eta_2 - \eta_1).
\end{aligned}$$

□




Figure 6

From the above lemma we can write $(1 + \frac{c}{2}(t_* - \xi_0^{-1}(\eta)))^{-3} \geq (1 + \frac{1}{2\mu}(\xi_0(t_*) - \eta))^{-3}$. Then substituting $v_F(\eta, 0; t_*)$ and $v_F(1, t; t_*)$ in (6.20) by the bounds (6.18)(6.19) yields

$$\begin{aligned}
& v_B(\xi_0(t_*), 0) \\
& \leq (1 + C\epsilon) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| + C\epsilon^2 \xi_0(t_*)^{-3} |\psi(\xi_0(t_*))| + C\epsilon^2 \int_0^{t_*} \left(1 + \frac{c}{2}(t_* - t)\right)^{-4} |\psi(1, t)| dt \\
& \quad + C\epsilon^3 \xi_0(t_*)^{-1} \int_1^{\xi_0(t_*)} \eta^{-3} |\psi(\eta, 0)| d\eta + C\epsilon^3 \int_0^{t_*} \left(1 + \frac{c}{2}(t_* - t)\right)^{-1} |\psi(1, t)| dt \\
& \quad + C\epsilon \xi_0(t_*)^{-1} \int_1^{\xi_0(t_*)} |w_F(\eta, 0)| d\eta + C\epsilon \int_0^{t_*} \left(1 + \frac{c}{2}(t_* - t)\right)^{-1} |w_F(1, t)| dt \\
& \quad + C\epsilon^2 \int_1^{\xi_0(t_*)} \left(1 + \frac{1}{2\mu}(\xi_0(t_*) - \eta)\right)^{-3} v_B(\eta, 0) d\eta \\
& \quad + C\epsilon^2 \xi_0(t_*)^{-1} \int_1^{\xi_0(t_*)} \int_{\xi}^{\xi_0(t_*)} \left(\xi + \frac{1}{2\mu}(\eta - \xi)\right)^{-1} v_B(\eta, 0) d\eta d\xi \\
& \quad + C\epsilon^2 \int_0^{t_*} \int_{\xi_0(t)}^{\xi_0(t_*)} \left(1 + \frac{c}{2}(t_* - t)\right)^{-1} \left(1 + \frac{1}{2\mu}(\eta - \xi_0(t))\right)^{-1} v_B(\eta, 0) d\eta dt.
\end{aligned} \tag{6.22}$$

Since $w_F = (\partial_t - cr^2\xi^{-2}\partial_\xi)\psi = 2\partial_t\psi - w_B$, there holds

$$\begin{aligned}
& \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-1} |w_F(1, t)| dt \\
& \leq 2 \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-1} |\partial_t\psi(1, t)| dt + \bar{c}^{\frac{1}{2}} \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-1} v_B(\xi_0(t), 0) dt \\
& \leq 2 \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-1} |\partial_t\psi(1, t)| dt + \frac{e^{C\kappa}\bar{c}^{\frac{1}{2}}}{\underline{c}} \int_1^{\xi_0(t_*)} \left(1 + \frac{1}{2\mu}(\xi_0(t_*) - \eta)\right)^{-1} v_B(\eta, 0) d\eta.
\end{aligned} \tag{6.23}$$

The last inequality follows from Lemma 6.6. To simplify the last two terms in (6.22) we compute

$$\begin{aligned}
& \int_1^{\xi_0(t_*)} \int_\xi^{\xi_0(t_*)} \left(\xi + \frac{1}{2\mu}(\eta - \xi)\right)^{-1} v_B(\eta, 0) d\eta d\xi \\
& = \int_1^{\xi_0(t_*)} \left(\int_1^\eta \left(\xi + \frac{1}{2\mu}(\eta - \xi)\right)^{-1} d\xi\right) v_B(\eta, 0) d\eta \\
& = \int_1^{\xi_0(t_*)} \frac{1}{1 - \frac{1}{2\mu}} \log\left(\frac{\eta}{1 + \frac{1}{2\mu}(\eta - 1)}\right) v_B(\eta, 0) d\eta \\
& \leq C \int_1^{\xi_0(t_*)} v_B(\eta, 0) d\eta,
\end{aligned} \tag{6.24}$$

and using Lemma 6.6 to obtain

$$\begin{aligned}
& \int_0^{t_*} \int_{\xi_0(t)}^{\xi_0(t_*)} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-1} \left(1 + \frac{1}{2\mu}(\eta - \xi_0(t))\right)^{-1} v_B(\eta, 0) d\eta dt \\
& = \int_1^{\xi_0(t_*)} \left(\int_0^{\xi_0^{-1}(\eta)} \left(1 + \frac{\underline{c}}{2}(t_* - t)\right)^{-1} \left(1 + \frac{1}{2\mu}(\eta - \xi_0(t))\right)^{-1} dt\right) v_B(\eta, 0) d\eta \\
& \leq \frac{e^{C\kappa}}{\underline{c}} \int_1^{\xi_0(t_*)} \left(\int_1^\eta \left(1 + \frac{1}{2\mu}(\xi_0(t_*) - \xi)\right)^{-1} \left(1 + \frac{1}{2\mu}(\eta - \xi)\right)^{-1} d\xi\right) v_B(\eta, 0) d\eta \\
& = \frac{e^{C\kappa}}{\underline{c}} \int_1^{\xi_0(t_*)} \frac{4\mu^2}{\xi_0(t_*) - \eta} \left(\log \frac{2\mu + \eta - 1}{2\mu} - \log \frac{2\mu + \xi_0(t_*) - 1}{2\mu + \xi_0(t_*) - \eta}\right) v_B(\eta, 0) d\eta \\
& = \frac{e^{C\kappa}}{\underline{c}} \int_1^{\xi_0(t_*)} \frac{4\mu^2}{\xi_0(t_*) - \eta} \log\left(1 + \frac{(\eta - 1)(\xi_0(t_*) - \eta)}{4\mu^2 + 2\mu(\xi_0(t_*) - 1)}\right) v_B(\eta, 0) d\eta.
\end{aligned}$$

If $\eta \in (1, \xi_0(t_*) - 1)$, using the bound (3.10)(3.11) the following holds:

$$\begin{aligned}
& \frac{1}{\xi_0(t_*) - \eta} \leq \frac{1 + 2\mu}{\xi_0(t_*) - \eta + 2\mu}, \\
& \log\left(1 + \frac{(\eta - 1)(\xi_0(t_*) - \eta)}{2\mu(2\mu + \xi_0(t_*) - 1)}\right) \leq \log \xi_0(t_*) \leq \log(1 + \bar{c}t_*) \leq \frac{\kappa}{\epsilon},
\end{aligned}$$

and thus

$$\frac{4\mu^2}{\xi_0(t_*) - \eta} \log\left(1 + \frac{(\eta - 1)(\xi_0(t_*) - \eta)}{4\mu^2 + 2\mu(\xi_0(t_*) - 1)}\right) \leq \frac{\kappa}{\epsilon} \frac{4\mu^2(1 + 2\mu)}{\xi_0(t_*) - \eta + 2\mu}.$$

If $\eta \in [\xi_0(t_*) - 1, \xi_0(t_*)]$, we have

$$\frac{4\mu^2}{\xi_0(t_*) - \eta} \log \left(1 + \frac{(\eta - 1)(\xi_0(t_*) - \eta)}{4\mu^2 + 2\mu(\xi_0(t_*) - 1)} \right) \leq \frac{2\mu(\eta - 1)}{2\mu + \xi_0(t_*) - 1} \leq 2\mu \leq \frac{\kappa}{\epsilon} \frac{4\mu^2}{\xi_0(t_*) - \eta + 2\mu}.$$

Hence in each case we obtain

$$\frac{4\mu^2}{\xi_0(t_*) - \eta} \log \left(1 + \frac{(\eta - 1)(\xi_0(t_*) - \eta)}{4\mu^2 + 2\mu(\xi_0(t_*) - 1)} \right) \lesssim \frac{\kappa}{\epsilon} \frac{1}{\xi_0(t_*) - \eta + 2\mu},$$

and thus

$$\begin{aligned} & \int_0^{t_*} \int_{\xi_0(t)}^{\xi_0(t_*)} \left(1 + \frac{\underline{c}}{2}(t_* - t) \right)^{-1} \left(1 + \frac{1}{2\mu}(\eta - \xi_0(t)) \right)^{-1} v_B(\eta, 0) d\eta dt \\ & \leq \frac{C e^{C\kappa}}{\underline{c}} \frac{\kappa}{\epsilon} \int_1^{\xi_0(t_*)} \frac{1}{\xi_0(t_*) - \eta + 2\mu} v_B(\eta, 0) d\eta. \end{aligned} \quad (6.25)$$

From (6.23)(6.24)(6.25) we see that all the terms on the right-hand side of (6.22) involving v_B can be controlled by

$$C\epsilon(\kappa + 1) \int_1^{\xi_0(t_*)} (2\mu + \xi_0(t_*) - \eta)^{-1} v_B(\eta, 0) d\eta.$$

Recall that

$$\begin{aligned} \mathcal{V}(\xi) & := |w_B(\xi, 0)| + \xi^{-1} \int_1^\xi |w_F(\eta, 0)| d\eta + \epsilon \xi^{-3} |\psi(\xi, 0)| + \epsilon^2 \xi^{-1} \int_1^\xi \eta^{-3} |\psi(\eta, 0)| d\eta, \\ \Psi(t_*) & := \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t) \right)^{-1} |\partial_t \psi(1, t)| dt + \epsilon \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t) \right)^{-4} |\psi(1, t)| dt \\ & \quad + \epsilon^2 \int_0^{t_*} \left(1 + \frac{\underline{c}}{2}(t_* - t) \right)^{-1} |\psi(1, t)| dt, \end{aligned}$$

and classify the terms on the right-hand side of (6.22) in view of $\kappa \leq \kappa_0 \lesssim 1$:

$$\begin{aligned} v_B(\xi_0(t_*), 0) & \leq \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| + C\epsilon \mathcal{V}(\xi_0(t_*)) + C\epsilon \Psi(t_*) \\ & \quad + C\epsilon \int_1^{\xi_0(t_*)} (2\mu + \xi_0(t_*) - \eta)^{-1} v_B(\eta, 0) d\eta. \end{aligned} \quad (6.26)$$

From (6.21) and in a same manner, there is also

$$\begin{aligned} & \left| \frac{w_B}{c^{\frac{1}{2}}}(1, t_*) - \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \\ & \leq C\epsilon \mathcal{V}(\xi_0(t_*)) + C\epsilon \Psi(t_*) + C\epsilon \int_1^{\xi_0(t_*)} (2\mu + \xi_0(t_*) - \eta)^{-1} v_B(\eta, 0) d\eta. \end{aligned} \quad (6.27)$$

Multiplying (6.26) by $\xi_0(t_*)$ yields

$$\begin{aligned}
& \xi_0(t_*) \left(v_B(\xi_0(t_*), 0) - \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right) \\
& \lesssim \epsilon \xi_0(t_*) (\mathcal{V}(\xi_0(t_*)) + \Psi(t_*)) \\
& \quad + \epsilon \xi_0(t_*) \int_1^{\xi_0(t_*)} (2\mu + \xi_0(t_*) - \eta)^{-1} \left(v_B(\eta, 0) - \left| \frac{w_B}{c^{\frac{1}{2}}}(\eta, 0) \right| \right) d\eta \\
& \quad + \epsilon \xi_0(t_*) \int_1^{\xi_0(t_*)} (2\mu + \xi_0(t_*) - \eta)^{-1} \left| \frac{w_B}{c^{\frac{1}{2}}}(\eta, 0) \right| d\eta.
\end{aligned} \tag{6.28}$$

By recalling that (3.10)(3.11), we obtain

$$\log \xi_0(t_*) \leq \log \xi_0(T) \leq \log(1 + \bar{c}T) \leq \frac{\kappa}{\epsilon}, \tag{6.29}$$

and thus

$$\epsilon \xi_0(t_*) \int_1^{\xi_0(t_*)} (2\mu + \xi_0(t_*) - \eta)^{-1} \eta^{-1} d\eta = \frac{\epsilon \xi_0(t_*)}{\xi_0(t_*) + 2\mu} \left(\log \xi_0(t_*) + \log \frac{2\mu - 1 + \xi_0(t_*)}{2\mu} \right) \lesssim \kappa.$$

Therefore, taking maximum in (6.28) over $t_* \in [0, T]$ yields

$$\begin{aligned}
& \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left(v_B(\xi_0(t_*), 0) - \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right) \right\} \\
& \lesssim \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \mathcal{V}(\xi_0(t_*)) + \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \Psi(t_*) + \kappa \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right\} \\
& \quad + \kappa \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left(v_B(\xi_0(t_*), 0) - \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right) \right\}.
\end{aligned}$$

We remark here that since ξ_0 is a bijection from $[0, T]$ to $[1, \xi_0(T)]$ there holds

$$\begin{aligned}
& \max_{t_* \in [0, T]} \xi_0(t_*) \mathcal{V}(\xi_0(t_*)) = \max_{\eta \in [1, \xi(T)]} \eta \mathcal{V}(\eta), \\
& \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right\} = \max_{\eta \in [1, \xi(T)]} \left\{ \eta \left| \frac{w_B}{c^{\frac{1}{2}}}(\eta, 0) \right| \right\}.
\end{aligned}$$

By choosing κ_0 small, we arrive at

$$\begin{aligned}
& \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left(v_B(\xi_0(t_*), 0) - \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right) \right\} \\
& \lesssim \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \mathcal{V}(\xi_0(t_*)) + \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \Psi(t_*) + \kappa \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right\},
\end{aligned} \tag{6.30}$$

and it follows

$$\begin{aligned}
& \max_{t_* \in [0, T]} \xi_0(t_*) v_B(\xi_0(t_*), 0) \\
& \lesssim \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \mathcal{V}(\xi_0(t_*)) + \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \Psi(t_*) + \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right\}.
\end{aligned} \tag{6.31}$$

Furthermore, multiplying (6.27) by $\xi_0(t_*)$, taking maximum over $t_* \in [0, T]$ and applying the bound (6.31) to the v_B term gives

$$\begin{aligned} & \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left(\left| \frac{w_B}{c^{\frac{1}{2}}}(1, t_*) - \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right) \right\} \\ & \lesssim \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \mathcal{V}(\xi_0(t_*)) + \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \Psi(t_*) + \kappa \max_{t_* \in [0, T]} \left\{ \xi_0(t_*) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(t_*), 0) \right| \right\}, \end{aligned} \quad (6.32)$$

which proves Proposition 6.1.

6.5. The decay estimate of $\psi(1, t)$.

Recall from (5.4)(5.5) that $W(s) = [aw_B|_{x=0}, aw_B|_{x=0}]^T(s)$ with $a := \left(\frac{\tilde{f}'(R)}{c} - \frac{\rho \partial_t u}{c} \right) \Big|_{x=0}$ and satisfying $\left| a(t) - \frac{\tilde{f}'(1)}{c_0} \right| \lesssim \epsilon$. We write

$$aw_B|_{x=0}(s) = \frac{\tilde{f}'(1)}{c_0} w_B(\xi_0(s), 0) + \frac{\tilde{f}'(1)}{c_0} (w_B(1, s) - w_B(\xi_0(s), 0)) + \left(a(s) - \frac{\tilde{f}'(1)}{c_0} \right) w_B(1, s).$$

The last two terms on the right-hand side are regarded as perturbation, and from (6.31)(6.32) they satisfy

$$\begin{aligned} & \left| \int_0^t \exp(\Lambda_i(t-s)) (w_B(1, s) - w_B(\xi_0(s), 0)) ds \right| \\ & \lesssim \left(\int_0^t \exp(\operatorname{Re}\Lambda_i(t-s)) \xi_0(s)^{-1} ds \right) \left(\kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} \right. \\ & \quad \left. + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon \max_{s \in [0, T]} \xi_0(s) \Psi(s) \right) \\ & \lesssim \frac{1}{1 + \underline{c}t} \left(\kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon \max_{s \in [0, T]} \xi_0(s) \Psi(s) \right), \end{aligned} \quad (6.33)$$

$$\begin{aligned} & \left| \int_0^t \exp(\Lambda_i(t-s)) \left(a(s) - \frac{\tilde{f}'(1)}{c_0} \right) w_B(1, s) ds \right| \\ & \lesssim \epsilon \left(\int_0^t \exp(\operatorname{Re}\Lambda_i(t-s)) \xi_0(s)^{-1} ds \right) \left(\max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} \right. \\ & \quad \left. + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon \max_{s \in [0, T]} \xi_0(s) \Psi(s) \right) \\ & \lesssim \frac{1}{1 + \underline{c}t} \left(\epsilon \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon^2 \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon^2 \max_{s \in [0, T]} \xi_0(s) \Psi(s) \right), \end{aligned} \quad (6.34)$$

where $i = 1, 2$ and we used (3.10) and $\operatorname{Re}\Lambda_i < 0$ to deduce the inequality

$$\int_0^t \exp(\Lambda_i(t-s)) \xi_0(s)^{-1} ds \leq \int_0^t \exp(\operatorname{Re}\Lambda_i(t-s)) (1 + \underline{c}s)^{-1} ds \lesssim \frac{1}{1 + \underline{c}t}.$$

Use the above inequality again, we control the second term on the right-hand side of (5.6) by

$$\begin{aligned}
& \left| \int_0^t \exp \begin{bmatrix} \Lambda_1(t-s) & \\ & \Lambda_2(t-s) \end{bmatrix} (\Delta Y)(s) ds \right| \\
& \lesssim \epsilon \left(\int_0^t \exp \left[\max_{i=1,2} \{\operatorname{Re} \Lambda_i\} (t-s) \right] (1 + \underline{c}s)^{-1} (s) ds \right) \left(\max_{s \in [0,t]} (1 + \underline{c}s) (|\partial_t \psi(1, s)| + |\psi(1, s)|) \right) \\
& \lesssim \frac{\epsilon}{1 + \underline{c}t} \left(\max_{s \in [0,t]} (1 + \underline{c}s) (|\partial_t \psi(1, s)| + |\psi(1, s)|) \right)
\end{aligned} \tag{6.35}$$

To eliminate the $\Psi(s)$ terms in (6.33)(6.34) by $\psi(1, t)$ and $\partial_t \psi(1, t)$, we recall the notation (1.17) and compute

$$\begin{aligned}
\xi_0(t) \Psi(t) & \leq \xi_0(t) \int_0^t \left(1 + \frac{\underline{c}}{2}(t-s)\right)^{-1} |\mathcal{R}(s)| ds \\
& \quad + \xi_0(t) \left(\int_0^t \left(1 + \frac{\underline{c}}{2}(t-s)\right)^{-1} (1 + \underline{c}s)^{-1} ds \right) \max_{s \in [0,t]} (1 + \underline{c}s) |\partial_t \psi(1, s) - \mathcal{R}(s)| \\
& \quad + \epsilon \xi_0(t) \left(\int_0^t \left(1 + \frac{\underline{c}}{2}(t-s)\right)^{-4} (1 + \underline{c}s)^{-1} ds \right) \max_{s \in [0,t]} (1 + \underline{c}s) |\psi(1, s)| \\
& \quad + \epsilon^2 \xi_0(t) \left(\int_0^t \left(1 + \frac{\underline{c}}{2}(t-s)\right)^{-1} (1 + \underline{c}s)^{-1} ds \right) \max_{s \in [0,t]} (1 + \underline{c}s) |\psi(1, s)| \\
& \lesssim \xi_0(t) \int_0^t \left(1 + \frac{\underline{c}}{2}(t-s)\right)^{-1} |\mathcal{R}(s)| ds + \frac{\kappa}{\epsilon} \max_{s \in [0,t]} (1 + \underline{c}s) |\partial_t \psi(1, s) - \mathcal{R}(s)| \\
& \quad + \epsilon \max_{s \in [0,t]} (1 + \underline{c}s) |\psi(1, s)| + \epsilon \kappa \max_{s \in [0,t]} (1 + \underline{c}s) |\psi(1, s)|.
\end{aligned} \tag{6.36}$$

Now applying (6.33)-(6.36) to (5.6) yields

$$\begin{aligned}
& \left| Y(t) - \exp \begin{bmatrix} \Lambda_1 t & \\ & \Lambda_2 t \end{bmatrix} Y(0) + \frac{\tilde{f}'(1)}{c_0} \int_0^t \exp \begin{bmatrix} \Lambda_1(t-s) & \\ & \Lambda_2(t-s) \end{bmatrix} \begin{bmatrix} w_B(\xi_0(s), 0) \\ w_B(\xi_0(s), 0) \end{bmatrix} ds \right| \\
& \lesssim \frac{1}{1 + \underline{c}t} \left\{ \kappa \max_{s \in [0,T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0,T]} \xi_0(s) \mathcal{V}(\xi_0(s)) \right. \\
& \quad + \epsilon \xi_0(t) \int_0^t \left(1 + \frac{\underline{c}}{2}(t-s)\right)^{-1} |\mathcal{R}(s)| ds + \kappa \max_{s \in [0,t]} (1 + \underline{c}s) |\partial_t \psi(s) - \mathcal{R}(s)| \\
& \quad \left. + \epsilon^2 \max_{s \in [0,t]} (1 + \underline{c}s) |\psi(s)| \right\}.
\end{aligned}$$

Denote the principle part of $Y(t)$ by $Y_0(t)$, namely

$$Y_0(t) := \exp \begin{bmatrix} \Lambda_1 t & \\ & \Lambda_2 t \end{bmatrix} Y(0) - \frac{\tilde{f}'(1)}{c_0} \int_0^t \exp \begin{bmatrix} \Lambda_1(t-s) & \\ & \Lambda_2(t-s) \end{bmatrix} \begin{bmatrix} w_B(\xi_0(s), 0) \\ w_B(\xi_0(s), 0) \end{bmatrix} ds.$$

This automatically yields

$$\mathcal{R}(t) = \begin{bmatrix} \frac{\Lambda_1}{\Lambda_1 - \Lambda_2} & -\frac{\Lambda_2}{\Lambda_1 - \Lambda_2} \end{bmatrix} Y_0(t), \quad \partial_t \psi(t) - \mathcal{R}(t) = \begin{bmatrix} \frac{\Lambda_1}{\Lambda_1 - \Lambda_2} & -\frac{\Lambda_2}{\Lambda_1 - \Lambda_2} \end{bmatrix} (Y(t) - Y_0(t)).$$

Then the above inequality can be rewritten as

$$\begin{aligned}
& |Y(t) - Y_0(t)| \\
& \lesssim \frac{1}{1 + \underline{c}t} \left\{ \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) \right. \\
& \quad \left. + \epsilon \xi_0(t) \int_0^t \left(1 + \frac{\underline{c}}{2}(t-s) \right)^{-1} |Y_0(s)| ds + \epsilon^2 \max_{s \in [0, t]} (1 + \underline{c}s) |Y_0(s)| \right. \\
& \quad \left. + (\kappa + \epsilon^2) \max_{s \in [0, t]} (1 + \underline{c}s) |Y(s) - Y_0(s)| \right\}. \tag{6.37}
\end{aligned}$$

Multiplying (6.37) by $1 + \underline{c}t$ and taking maximum over $[0, T]$ yields

$$\begin{aligned}
& \max_{t \in [0, T]} (1 + \underline{c}t) |Y(t) - Y_0(t)| \\
& \lesssim \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon^2 \max_{s \in [0, T]} (1 + \underline{c}s) |Y_0(s)| \\
& \quad + \epsilon \max_{t \in [0, T]} \left\{ \xi_0(t) \int_0^t \left(1 + \frac{\underline{c}}{2}(t-s) \right)^{-1} |Y_0(s)| ds \right\} + (\kappa + \epsilon^2) \max_{t \in [0, T]} (1 + \underline{c}t) |Y(t) - Y_0(t)|.
\end{aligned}$$

Hence for small enough κ_0 and ϵ_0 , this shows

$$\begin{aligned}
& \max_{t \in [0, T]} (1 + \underline{c}t) |Y(t) - Y_0(t)| \\
& \lesssim \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) \\
& \quad + \epsilon \max_{t \in [0, T]} \left\{ \xi_0(t) \int_0^t \left(1 + \frac{\underline{c}}{2}(t-s) \right)^{-1} |Y_0(s)| ds \right\} + \epsilon^2 \max_{s \in [0, T]} (1 + \underline{c}s) |Y_0(s)|.
\end{aligned}$$

For the last two terms, we have

$$\begin{aligned}
|Y_0(t)| & \lesssim \exp \left(\max_{i=1,2} \{\operatorname{Re} \Lambda_i\} t \right) |Y(0)| \\
& \quad + \int_0^t \exp \left(\max_{i=1,2} \{\operatorname{Re} \Lambda_i\} (t-s) \right) \xi_0(s)^{-1} ds \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} \\
& \lesssim \exp \left(\max_{i=1,2} \{\operatorname{Re} \Lambda_i\} t \right) |Y(0)| + \frac{1}{1 + \underline{c}t} \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\}, \tag{6.38}
\end{aligned}$$

and thus

$$\begin{aligned}
& \epsilon \max_{t \in [0, T]} \left\{ \xi_0(t) \int_0^t \left(1 + \frac{\underline{c}}{2}(t-s) \right)^{-1} |Y_0(s)| ds \right\} + \epsilon^2 \max_{s \in [0, T]} (1 + \underline{c}s) |Y_0(s)| \\
& \lesssim \epsilon |Y(0)| + \epsilon^2 \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} \\
& \quad + \epsilon \max_{t \in [0, T]} \left\{ \xi_0(t) \int_0^t \left(1 + \frac{\underline{c}}{2}(t-s) \right)^{-1} (1 + \underline{c}s)^{-1} ds \right\} \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} \\
& \lesssim \epsilon |Y(0)| + (\kappa + \epsilon^2) \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\}. \tag{6.39}
\end{aligned}$$

Here we used that

$$\xi_0(t) \int_0^t \left(1 + \frac{c}{2}(t-s)\right)^{-1} (1 + \underline{c}s)^{-1} ds = \xi_0(t) \frac{2}{3\underline{c} + \underline{c}^2 t} \left(\log(1 + \underline{c}t) + \log\left(1 + \frac{c}{2}t\right) \right) \lesssim \frac{\kappa}{\epsilon}.$$

Therefore by noting that $|w_B(\eta)| \leq \mathcal{V}(\eta)$,

$$\begin{aligned} & \max_{t \in [0, T]} (1 + \underline{c}t) |Y(t) - Y_0(t)| \\ & \lesssim (\kappa + \epsilon^2) \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon |Y(0)| \\ & \lesssim \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon |Y(0)|. \end{aligned} \quad (6.40)$$

By substituting the last third terms on the right-hand side of (6.37) by the bounds (6.39)(6.40), we obtain

$$\begin{aligned} & |Y(t) - Y_0(t)| \\ & \lesssim \frac{1}{1 + \underline{c}t} \left\{ \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon |Y(0)| \right\}, \end{aligned} \quad (6.41)$$

which proves Proposition 6.2.

7. Close the bootstrap

With the help of the bounds obtained in the previous sections, we shall complete the bootstrap argument in this section.

7.1. Proof of almost global existence

To prove the bound (3.12), recall the energy identity (2.34), and by dropping the iteration subscripts we obtain

$$\frac{1}{2} \frac{d}{dt} \left[\int_0^\infty ((\partial_t^j u)^2 + c^2 (\partial_t^j q)^2) dx + (c^2 r^2)|_{x=0} f'(R)^{-1} |\partial_t^j f(R)|^2 \right] = \sum_{l=1}^5 nl_l^j(t), \quad (7.1)$$

where we denote

$$\begin{aligned} nl_1^j(t) &:= \int_0^\infty \left(\frac{1}{2} \partial_t c^2 (\partial_t^j q)^2 - r^2 \partial_x c^2 \partial_t^j u \partial_t^j q \right) dx, \\ nl_2^j(t) &:= \int_0^\infty [\partial_t^j, c^2 r^2 \partial_x] q \partial_t^j u dx, \quad nl_3^j(t) := \int_0^\infty c^2 [\partial_t^j, \partial_x r^2] u \partial_t^j q dx, \\ nl_4^j(t) &:= \frac{1}{2} \partial_t [(c^2 r^2)|_{x=0} f'(R)^{-1}] |\partial_t^j f(R)|^2, \\ nl_5^j(t) &:= -(c^2 r^2)|_{x=0} [\partial_t^j, f'(R)^{-1}] \partial_t f(R) \partial_t^j f(R). \end{aligned}$$

From (4.8)-(4.12) and (3.7)-(3.9), it follows

$$\sum_{j=0}^2 \left(|\partial_t^j u|^2 + |\partial_t^j q|^2 \right) \simeq \sum_{j=0}^2 \left(|\partial_t^{j+1} \varphi|^2 + |c r^2 \partial_x \partial_t^j \varphi|^2 \right), \quad (7.2)$$

and thus by Hardy inequality

$$\begin{aligned} & \sum_{j=0}^2 \left\{ \int_0^T \int_0^\infty \xi^{-1} \left((\partial_t^{1+j} \varphi)^2 + (cr^2 \partial_x \partial_t^j \varphi)^2 \right) dx dt + \int_0^T \int_0^\infty \xi^{-3} \left(\partial_t^j \varphi \right)^2 dx dt \right\} \\ & \simeq \sum_{j=0}^2 \int_0^T \int_0^\infty \xi^{-1} \left((\partial_t^j u)^2 + (\partial_t^j q)^2 \right) dx dt, \end{aligned}$$

with which we further deduce from (4.1) that

$$\sum_{j=0}^2 \int_0^T \int_0^\infty \xi^{-1} \left((\partial_t^j u)^2 + (\partial_t^j q)^2 \right) dx dt \lesssim \max \{ \log_2 T, 1 \} \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \int_0^T \varphi^2|_{x=0} dt \quad (7.3)$$

Using the above inequality with (3.7)(3.11), we estimate

$$\begin{aligned} \int_0^T |nl_1^j(t)| dt & \lesssim (\|\xi \partial_t c^2\|_{L^\infty} + \|\xi r^2 \partial_x c^2\|_{L^\infty}) \int_0^T \int_0^\infty \xi^{-1} \left((\partial_t^j u)^2 + (\partial_t^j q)^2 \right) dx dt \\ & \lesssim \epsilon \left(\max \{ \log_2 T, 1 \} \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \int_0^T \varphi^2|_{x=0} dt \right) \\ & \lesssim \kappa \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \epsilon \int_0^T \varphi^2|_{x=0} dt. \end{aligned} \quad (7.4)$$

Next, we compute

$$\begin{aligned} [\partial_t, r^2 c^2 \partial_x] q & = \partial_t (r^2 c^2) \partial_x q = \frac{\partial_t (c^2 r^2)}{c^2 r^2} \partial_t u, \\ [\partial_t^2, r^2 c^2 \partial_x] q & = [\partial_t^2, r^2 c^2] \partial_x q = [\partial_t^2, r^2 c^2] (r^{-2} c^{-2} \partial_t u) \\ & = 2 \frac{\partial_t (r^2 c^2)}{r^2 c^2} \partial_t^2 u + \frac{\partial_t^2 (r^2 c^2)}{r^2 c^2} \partial_t u + 2 \partial_t (r^2 c^2) \partial_t (r^{-2} c^{-2}) \partial_t u, \\ \frac{\partial_t^2 (r^2 c^2)}{r^2 c^2} & = \frac{\partial_t^2 c^2}{c^2} + 2 \frac{\partial_t r^2 \partial_t c^2}{r^2 c^2} + \frac{\partial_t^2 r^2}{r^2} \\ & = -(\gamma + 1) \rho \partial_t^2 q + (\gamma + 1)(\gamma + 2) \rho^2 (\partial_t q)^2 + 2 \frac{\partial_t u}{r} + 2 \frac{u^2}{r^2} + 8 \frac{u \partial_t c}{rc}. \end{aligned}$$

Hence by (3.6)(3.7),

$$|[\partial_t^2, r^2 c^2 \partial_x] q| \lesssim \epsilon \xi^{-1} (|\partial_t^2 u| + |\partial_t^2 q| + |\partial_t u|).$$

From (7.3)(3.11), it follows

$$\int_0^T |nl_2^j(t)| dt \lesssim \epsilon \sum_{i=0}^j \int_0^T \int_0^\infty \xi^{-1} \left((\partial_t^i u)^2 + (\partial_t^i q)^2 \right) dx dt \lesssim \kappa \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \epsilon \int_0^T \varphi^2|_{x=0} dt. \quad (7.5)$$

Similarly, to control nl_3^j , we compute

$$[\partial_t, \partial_x r^2] u = \partial_x [\partial_t, r^2] u = \partial_x (2ru^2) = 4 \frac{u}{r} \partial_t q - 6 \frac{u^2}{\rho r^2},$$

$$\begin{aligned}
[\partial_t^2, \partial_x r^2]u &= \partial_x [\partial_t^2, r^2]u = \partial_x [2u^3 + 4ru\partial_t u] = \partial_x \left[4\frac{u}{r}\partial_t(r^2u) - 6u^3 \right], \\
\partial_x \left[\frac{u}{r}\partial_t(r^2u) \right] &= \frac{u}{r}\partial_t^2 q + 4\partial_x \left(\frac{u}{r} \right) \partial_t(r^2u) = \frac{u}{r}\partial_t^2 q + r^{-1} (\partial_t q - 3\rho^{-1}r^{-1}u) (\partial_t u + 2r^{-1}u^2), \\
\partial_x u^3 &= 3\frac{u^2}{r^2} (\partial_t q - 2\rho^{-1}r^{-1}u).
\end{aligned}$$

Then by (7.3)(3.11), we obtain

$$\int_0^T |nl_3^j(t)| dt \lesssim \epsilon \sum_{i=0}^2 \int_0^T \int_0^\infty \xi^{-1} \left((\partial_t^j u)^2 + (\partial_t^j q)^2 \right) dx dt \lesssim \kappa \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \epsilon \int_0^T \varphi^2|_{x=0} dt. \quad (7.6)$$

Recall that $f(R) = q|_{x=0}$ and $\rho^{\gamma-1} - 1 = -\left(\frac{Ca\gamma}{2(\gamma-1)}\right)^{-1} (\partial_x \varphi - \frac{1}{2}u^2)$, from which we write

$$q = \rho^{-1} - 1 = -\frac{\rho^{-1} - 1}{\rho^{\gamma-1} - 1} \left(\frac{Ca\gamma}{2(\gamma-1)} \right)^{-1} \left(\partial_x \varphi - \frac{1}{2}u^2 \right).$$

In view of (3.6)(3.7), this shows

$$\begin{aligned}
|f(R)| &\lesssim \left| \left(\partial_t \varphi - \frac{1}{2}u^2 \right) \Big|_{x=0} \right| \leq |\partial_t \varphi|_{x=0} + C\epsilon |(cr^2 \partial_x \varphi)|_{x=0} \\
&\leq |\partial_t \varphi|_{x=0} + C\epsilon \left| \left(cr^2 \varphi + \frac{c}{\rho r} \varphi \right) \Big|_{x=0} \right| + C\epsilon |\varphi|_{x=0}.
\end{aligned}$$

For derivatives of $f(R)$, we have

$$\begin{aligned}
\partial_t q &= -(\gamma-1)^{-1} \rho^{-\gamma} \partial_t (\rho^{\gamma-1} - 1) = \left(\frac{Ca\gamma}{2} \right)^{-1} \rho^{-\gamma} (\partial_t^2 \varphi - u \partial_t u), \\
|\partial_t f(R)| &\lesssim |\partial_t^2 \varphi|_{x=0} + C\epsilon |\partial_t u|_{x=0} \\
&= |\partial_t^2 \varphi|_{x=0} + C\epsilon |\partial_t (\rho r^2 \partial_x \varphi)|_{x=0} \\
&\leq |\partial_t^2 \varphi|_{x=0} + C\epsilon |\rho r^2 \partial_x \partial_t \varphi|_{x=0} + C\epsilon \left| \frac{\partial_t (\rho r^2)}{\rho r^2} \Big|_{x=0} \right| \cdot |(\rho r^2 \partial_x \varphi)|_{x=0} \\
&\leq |\partial_t^2 \varphi|_{x=0} + C\epsilon \left| \left(cr^2 \partial_x \partial_t \varphi + \frac{c}{\rho r} \partial_t \varphi \right) \Big|_{x=0} \right| + C\epsilon |\partial_t \varphi|_{x=0} \\
&\quad + C\epsilon \left| \left(cr^2 \partial_x \varphi + \frac{c}{\rho r} \varphi \right) \Big|_{x=0} \right| + C\epsilon |\varphi|_{x=0}.
\end{aligned}$$

Similarly, by separating the top derivatives and using (3.6)(3.7), we continue and obtain

$$\begin{aligned}
|\partial_t^2 f(R)| &= |\partial_t^2 q|_{x=0} = \left(\frac{Ca\gamma}{2}\right)^{-1} |\partial_t (\rho^{-\gamma} (\partial_t^2 \varphi - u \partial_t u))|_{x=0} \\
&\lesssim |(\partial_t^3 \varphi - (\partial_t u)^2 - u \partial_t^2 u)|_{x=0} + |\partial_t \rho^{-\gamma}|_{x=0} \cdot |(\partial_t^2 \varphi - u \partial_t u)|_{x=0} \\
&\lesssim |(\partial_t^3 \varphi - (\partial_t u)^2 - u \partial_t^2 u)|_{x=0} + \epsilon \left(\sum_{j=0}^2 |\partial_t^j \varphi|_{x=0} + \sum_{j=0}^1 \left| \left(cr^2 \partial_x \partial_t^j \varphi + \frac{c}{\rho r} \partial_t^j \varphi \right) \right|_{x=0} \right) \\
&\lesssim |\partial_t^3 \varphi|_{x=0} + \epsilon |\partial_t^2 u|_{x=0} + \epsilon \left(\sum_{j=0}^2 |\partial_t^j \varphi|_{x=0} + \sum_{j=0}^1 \left| \left(cr^2 \partial_x \partial_t^j \varphi + \frac{c}{\rho r} \partial_t^j \varphi \right) \right|_{x=0} \right),
\end{aligned}$$

$$\begin{aligned}
|\partial_t^2 u|_{x=0} &= |\partial_t^2 (\rho r^2 \partial_x \varphi)|_{x=0} \\
&= |(\rho r^2 \partial_x \partial_t^2 \varphi + 2 \partial_t (\rho r^2) \partial_x \partial_t \varphi + \partial_t^2 (\rho r^2) \partial_x \varphi)|_{x=0} \\
&\leq |(\rho r^2 \partial_x \partial_t^2 \varphi)|_{x=0} + C \epsilon |(\rho r^2 \partial_x \partial_t \varphi)|_{x=0} \\
&\quad + \left| \left(\rho^{-1} \partial_t^2 \rho + 4 \frac{u}{\rho r} \partial_t \rho + \frac{2u^2}{r^2} + \frac{2 \partial_t u}{r} \right) \right|_{x=0} \cdot |(\rho r^2 \partial_x \varphi)|_{x=0} \\
&\lesssim |(\rho r^2 \partial_x \partial_t^2 \varphi)|_{x=0} + \epsilon (|(\rho r^2 \partial_x \partial_t \varphi)|_{x=0} + |(\rho r^2 \partial_x \varphi)|_{x=0} + |\partial_t^2 \rho|_{x=0} + |\partial_t \rho|_{x=0}) \\
&\lesssim |(\rho r^2 \partial_x \partial_t^2 \varphi)|_{x=0} + \epsilon (|(\rho r^2 \partial_x \partial_t \varphi)|_{x=0} + |(\rho r^2 \partial_x \varphi)|_{x=0} + |\partial_t^2 q|_{x=0} + |\partial_t q|_{x=0}) \\
&\lesssim \sum_{j=0}^2 \left| \left(cr^2 \partial_x \partial_t^j \varphi + \frac{c}{\rho r} \partial_t^j \varphi \right) \right|_{x=0} + \sum_{j=0}^2 |\partial_t^j \varphi|_{x=0} + \epsilon |\partial_t^2 q|_{x=0}.
\end{aligned}$$

Combining the above two inequalities yields

$$|\partial_t^2 f(R)| \lesssim |\partial_t^3 \varphi|_{x=0} + \epsilon \sum_{j=0}^2 \left| \left(cr^2 \partial_x \partial_t^j \varphi + \frac{c}{\rho r} \partial_t^j \varphi \right) \right|_{x=0} + \epsilon \sum_{j=0}^2 |\partial_t^j \varphi|_{x=0} + \epsilon^2 |\partial_t^2 f(R)|.$$

Hence for small enough ϵ this gives

$$|\partial_t^2 f(R)| \lesssim |\partial_t^3 \varphi|_{x=0} + \epsilon \sum_{j=0}^2 \left| \left(cr^2 \partial_x \partial_t^j \varphi + \frac{c}{\rho r} \partial_t^j \varphi \right) \right|_{x=0} + \epsilon \sum_{j=0}^2 |\partial_t^j \varphi|_{x=0},$$

and thus for $j = 0, 1, 2$

$$|\partial_t^j f(R)| \lesssim |\partial_t^{1+j} \varphi|_{x=0} + \epsilon \sum_{i=0}^j \left| \left(cr^2 \partial_x \partial_t^i \varphi + \frac{c}{\rho r} \partial_t^i \varphi \right) \right|_{x=0} + \epsilon \sum_{i=0}^j |\partial_t^i \varphi|_{x=0} \quad (7.7)$$

We return to the estimate of nl_4^j . Since $c^2 = c_0^2(1+q)^{-\gamma-1}$ and $q|_{x=0} = f(R)$, there exists a function f_1 which is smooth near $R = 1$ such that $(cr^2)|_{x=0} f'(R)^{-1} = f_1(R)$, and thus

$$|\partial_t [(cr^2)|_{x=0} f'(R)^{-1}]| = |\partial_t f_1(R)| = |f_1'(R)| |u|_{x=0} \lesssim \epsilon.$$

It then follows from (7.7)(4.1)(7.2)(3.11) that

$$\begin{aligned} \int_0^T |nl_4^j(t)|dt &\lesssim \epsilon \left(\max\{\log_2 T, 1\} \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \int_0^T \varphi^2|_{x=0} dt \right) \\ &\lesssim \kappa \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \epsilon \int_0^T \varphi^2|_{x=0} dt. \end{aligned} \quad (7.8)$$

The nl_5^j terms can also be treated by direct computation.

$$|[\partial_t, f'(R)^{-1}]\partial_t f(R) \cdot \partial_t f(R)| = |\partial_t(f'(R)^{-1})| \cdot |\partial_t f(R)|^2 \lesssim \epsilon |\partial_t f(R)|^2,$$

$$[\partial_t^2, f'(R)^{-1}]\partial_t f(R) \cdot \partial_t^2 f(R) = 2\partial_t(f'(R)^{-1})|\partial_t^2 f(R)|^2 + \partial_t^2(f'(R)^{-1})\partial_t f(R)\partial_t^2 f(R),$$

By Leibniz rule,

$$\begin{aligned} \partial_t^2(f'(R)^{-1}) &= \partial_t^2 \left(\frac{1}{f'} \circ f^{-1} \circ f(R) \right) = \partial_t \left[\left(\frac{1}{f'} \circ f^{-1} \right)' \circ f(R) \cdot \partial_t f(R) \right] \\ &= \left(\frac{1}{f'} \circ f^{-1} \right)' \circ f(R) \cdot \partial_t^2 f(R) + \left(\frac{1}{f'} \circ f^{-1} \right)'' \circ f(R) \cdot (\partial_t f(R))^2. \end{aligned}$$

Since $|\partial_t R| = |u|_{x=0} \lesssim \epsilon$, it follows

$$|[\partial_t^2, f'(R)^{-1}]\partial_t f(R) \cdot \partial_t^2 f(R)| \lesssim \epsilon ((\partial_t^2 f(R))^2 + (\partial_t f(R))^2),$$

and by (7.7)(4.1)(3.11)

$$\begin{aligned} \int_0^T |nl_5^j(t)|dt &\lesssim \epsilon \left(\left| \partial_t^{1+j} \varphi|_{x=0} \right| + \epsilon \sum_{i=0}^j \left| \left(cr^2 \partial_x \partial_t^i \varphi + \frac{c}{\rho r} \partial_t^i \varphi \right) \Big|_{x=0} \right| + \epsilon \sum_{i=0}^j \left| \partial_t^i \varphi|_{x=0} \right| \right) \\ &\lesssim \epsilon \left(\max\{\log_2 T, 1\} \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \int_0^T \varphi^2|_{x=0} dt \right) \\ &\lesssim \kappa \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \epsilon \int_0^T \varphi^2|_{x=0} dt. \end{aligned} \quad (7.9)$$

Integrating (7.1) over $[0, T]$ and using (7.4)(7.5)(7.6)(7.8)(7.9) yields

$$e^j(t) - e^j(0) \lesssim \kappa \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + \epsilon \int_0^T \varphi^2|_{x=0} dt,$$

and it follows by taking maximum

$$\sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) \leq \sum_{j=0}^2 e^j(0) + C\kappa \sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) + C\epsilon \int_0^T \varphi^2|_{x=0} dt.$$

Hence for small enough κ_0 and $\kappa \leq \kappa_0$ such that $C\kappa_0 \leq \frac{1}{2}$, we obtain

$$\sum_{j=0}^2 \max_{0 \leq t \leq T} e^j(t) \leq 2 \sum_{j=0}^2 e^j(0) + C\epsilon \int_0^T \varphi^2|_{x=0} dt. \quad (7.10)$$

To prove (3.12), it remains to bound $\int_0^T \varphi^2|_{x=0} dt$. To this end, we call upon Proposition 6.2 and use the following lemma.

Lemma 7.1. *Under the condition (1.18) and for $\xi \geq 1$, we have the bounds*

$$\begin{aligned} |Y(0)| &\lesssim \epsilon, \quad \int_1^\infty |w_B(\eta, 0)|^2 d\eta \lesssim \epsilon^2, \\ \max_{\eta \in [1, \xi_0(T)]} \{\eta |w_B(\eta, 0)|\} &\lesssim \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}}, \quad \max_{\xi \in [1, \xi_0(T)]} \int_1^\xi |w_F(\eta, 0)| d\eta \lesssim \epsilon^{\frac{1}{2}} \left(\epsilon^{\frac{1}{2}} + \tilde{\epsilon}^{\frac{1}{2}} \right), \\ \xi^{-2} |\psi(\xi, 0)| &\lesssim \epsilon \xi^{-1}, \quad \int_1^\xi \eta^{-3} |\psi(\eta, 0)| d\eta \lesssim \epsilon. \end{aligned}$$

In particular,

$$\epsilon \max_{\eta \in [1, \xi_0(T)]} \eta \mathcal{V}(\eta) \lesssim \epsilon^{\frac{3}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \epsilon^2.$$

Proof. Recall that $\rho r^2 \partial_x \varphi = u$. By the bootstrap bound (3.2)(3.8) at $t = 0$ and a Hölder inequality, there holds

$$|\varphi(x, 0)| \leq \int_x^\infty \left| (1 + q_{in}) \frac{u_{in}}{r_{in}^2}(y) \right| dy \lesssim \|u_{in}\|_{L^2} \lesssim \epsilon.$$

From (1.15), we also have

$$|\partial_t \varphi(x, 0)| \lesssim \|u_{in}\|_{L^\infty}^2 + \|q_{in}\|_{L^\infty} \lesssim \epsilon.$$

From $\psi = r\varphi$, $\partial_t \psi = r\partial_t \varphi + u\varphi$, and $r_{in}|_{\xi=1} = R_{in} \simeq 1$, it follows that

$$\begin{aligned} |Y(0)| &\lesssim |\partial_t \psi|_{\{\xi=1, t=0\}} + |\psi|_{\{\xi=1, t=0\}} \lesssim |\partial_t \varphi|_{\{x=0, t=0\}} + |\varphi|_{\{x=0, t=0\}} \lesssim \epsilon, \\ \xi^{-2} |\psi(\xi, 0)| &= \xi^{-2} r_{in}(x) |\varphi(x, 0)| \lesssim \epsilon \xi^{-1}, \\ \int_1^\xi \eta^{-3} |\psi(\eta, 0)| d\eta &\lesssim \epsilon \int_1^\xi \eta^{-2} d\eta \lesssim \epsilon. \end{aligned}$$

Using (3.2)(3.8) again, we compute

$$\begin{aligned} \xi |w_B(\xi, 0)| &= \xi |(\partial_t + cr^2 \partial_x)(r\varphi)(x, 0)| \\ &\leq \xi r_{in}(x) (|\partial_t \varphi(x, 0)| + |cr^2 \partial_x \varphi(x, 0)|) + \xi \left| \left(u_{in}(x) + \frac{c(x, 0)}{\rho_{in}(x)} \right) \varphi(x, 0) \right| \\ &\lesssim \xi^2 (|q_{in}(x)| + |u_{in}(x)|) + \xi |\varphi(x, 0)|. \end{aligned}$$

Similarly,

$$\xi |w_F(\xi, 0)| \lesssim \xi^2 (|q_{in}(x)| + |u_{in}(x)|) + \xi |\varphi(x, 0)|.$$

Then by condition (1.18),

$$\begin{aligned}\xi^2|u_{in}(x)| &\leq \left(2 \int_0^\infty \xi^2|u_{in}(\xi)| |\partial_x(\xi^2 u_{in})| dx\right)^{\frac{1}{2}} \lesssim \|\tilde{L}_0^1 u_{in}\|_{L^2}^{\frac{1}{2}} \|\xi^2 u_{in}\|_{L^2}^{\frac{1}{2}} \lesssim \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}}, \\ \xi^2|q_{in}(x)| &\leq \left(2 \int_0^\infty \xi^2|q_{in}(\xi)| |\partial_x(\xi^2 q_{in})| dx\right)^{\frac{1}{2}} \lesssim (\|L_0^1 q_{in}\|_{L^2} + \|q_{in}\|_{L^2})^{\frac{1}{2}} \|\xi^2 q_{in}\|_{L^2}^{\frac{1}{2}} \lesssim \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}}, \\ \xi|\varphi(x,0)| &\leq \xi \int_x^\infty (1+q_{in}) \frac{|u_{in}|}{r_{in}^2}(y) dy \lesssim \xi \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}} \int_x^\infty (1+3y)^{-\frac{4}{3}} dy \lesssim \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}}.\end{aligned}$$

Hence we obtain $\xi|w_B(\xi,0)| \lesssim \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}}$. Next, for the bound on $\int_1^\infty |w_B(\eta,0)|^2 d\eta$, we use a change of variable and Hardy inequality to obtain

$$\begin{aligned}\int_1^\infty |w_B(\eta,0)|^2 d\eta &\lesssim \int_1^\infty (|\eta q_{in}(x(\eta))|^2 + |\eta u_{in}(x(\eta))|^2 + |\varphi(x(\eta),0)|^2) d\eta \\ &\lesssim \int_0^\infty (|q_{in}(x)|^2 + |u_{in}(x)|^2 + |\xi^{-1}\varphi(x,0)|^2) dx \\ &\lesssim \|u_{in}\|_{L^2}^2 + \|q_{in}\|_{L^2}^2 \\ &\lesssim \epsilon^2.\end{aligned}$$

It remains to control $\max_{\xi \in [1, \xi_0(T)]} \int_1^\xi |w_F(\eta,0)| d\eta$, for which we use

$$\begin{aligned}\max_{\xi \in [1, \xi_0(T)]} \int_1^\xi |w_F(\eta,0)| d\eta &\lesssim \left(\int_1^\infty |\eta w_F(\eta,0)|^2 d\eta\right)^{\frac{1}{2}} \\ &\lesssim \left(\int_1^\infty |\eta^2 q_{in}(x(\eta))|^2 d\eta\right)^{\frac{1}{2}} + \left(\int_1^\infty |\eta^2 u_{in}(x(\eta))|^2 d\eta\right)^{\frac{1}{2}} + \left(\int_1^\infty |\eta \varphi(x(\eta),0)|^2 d\eta\right)^{\frac{1}{2}}.\end{aligned}$$

By a change of variable and Hölder,

$$\left(\int_1^\infty |\eta^2 q_{in}(x(\eta))|^2 d\eta\right)^{\frac{1}{2}} = \|\xi q_{in}\|_{L^2} \leq \|q_{in}\|_{L^2}^{\frac{1}{2}} \|\xi^2 q_{in}\|_{L^2}^{\frac{1}{2}} \leq \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}},$$

and similarly,

$$\left(\int_1^\infty |\eta^2 u_{in}(x(\eta))|^2 d\eta\right)^{\frac{1}{2}} \leq \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}}.$$

For the last term, integration by parts gives

$$\begin{aligned}&\int_1^\infty |\eta \varphi(x(\eta),0)|^2 d\eta \\ &\lesssim |\varphi(0,0)|^2 + \int_1^\infty |\eta \varphi(x(\eta),0)| |\eta^2 \partial_\eta \varphi(x(\eta),0)| d\eta \\ &\lesssim |\varphi(0,0)|^2 + \left(\int_1^\infty |\eta \varphi(x(\eta),0)|^2 d\eta\right)^{\frac{1}{2}} \left(\int_1^\infty |\eta^2 u_{in}(x(\eta),0)|^2 d\eta\right)^{\frac{1}{2}},\end{aligned}$$

where we used that $\rho r^2 \simeq \xi^2$ by (3.2)(3.8). Hence by Cauchy-Schwartz

$$\int_1^\infty |\eta\varphi(x(\eta), 0)|^2 d\eta \lesssim \epsilon^2 + \epsilon\tilde{\epsilon}.$$

Collecting the above bounds yields

$$\max_{\xi \in [1, \xi_0(T)]} \int_1^\xi |w_F(\eta, 0)| d\eta \lesssim \left(\int_1^\infty |\eta w_F(\eta, 0)|^2 d\eta \right)^{\frac{1}{2}} \lesssim \epsilon^{\frac{1}{2}} \left(\epsilon^{\frac{1}{2}} + \tilde{\epsilon}^{\frac{1}{2}} \right).$$

□

Now we combine Proposition 6.2 with Lemma 7.1 in view of $\epsilon \leq \epsilon_0$, $\tilde{\epsilon} \leq \tilde{\epsilon}_0$, $\kappa \leq \kappa_0$ to deduce

$$\begin{aligned} & C\epsilon \int_0^T \varphi^2|_{x=0} dt \\ & \leq C\epsilon \int_0^T |Y_0(t)|^2 dt + C\epsilon \left(\epsilon |Y(0)| + \kappa \max_{\eta \in [1, \xi_0(T)]} \left\{ \eta \left| \frac{w_B}{c^{\frac{1}{2}}}(\eta, 0) \right| \right\} + \epsilon \max_{\eta \in [1, \xi_0(T)]} \eta \mathcal{V}(\eta) \right)^2 \\ & \leq C\epsilon \int_0^T |Y_0(t)|^2 dt + C\epsilon \left(\epsilon^2 + \kappa \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \epsilon^{\frac{3}{2}} \tilde{\epsilon}^{\frac{1}{2}} \right)^2 \\ & \leq C\epsilon \int_0^T |Y_0(t)|^2 dt + C\epsilon^2 \left(\epsilon_0^{\frac{3}{2}} + \kappa_0 \tilde{\epsilon}_0^{\frac{1}{2}} + \epsilon_0 \tilde{\epsilon}_0^{\frac{1}{2}} \right)^2. \end{aligned}$$

Meanwhile, by Young inequality, Lemma 6.6 and Lemma 7.1 again we obtain

$$\begin{aligned} & \left\| \int_0^t \exp\left(\max_{i=1,2} \{\operatorname{Re}\Lambda_i\}(t-s)\right) |w_B(\xi_0(s), 0)| ds \right\|_{L^2[0,T]}^2 \\ & \lesssim \int_0^T |w_B(\xi_0(t), 0)|^2 dt \lesssim \int_0^{\xi_0(T)} |w_B(\eta, 0)|^2 d\eta \lesssim \epsilon^2, \end{aligned}$$

and thus $\int_0^T |Y_0(t)|^2 dt \lesssim \epsilon^2$, which further gives

$$C\epsilon \int_0^T \varphi^2|_{x=0} dt \leq C\epsilon^2 \left(\epsilon_0 + \left(\epsilon_0^{\frac{3}{2}} + \kappa_0 \tilde{\epsilon}_0^{\frac{1}{2}} + \epsilon_0 \tilde{\epsilon}_0^{\frac{1}{2}} \right)^2 \right).$$

Hence for sufficient small $\epsilon_0, \tilde{\epsilon}_0, \kappa_0$ such that $C \left(\epsilon_0 + \left(\epsilon_0^{\frac{3}{2}} + \kappa_0 \tilde{\epsilon}_0^{\frac{1}{2}} + \epsilon_0 \tilde{\epsilon}_0^{\frac{1}{2}} \right)^2 \right) \leq \tilde{B}$, we obtain

$$C\epsilon \int_0^T \varphi^2|_{x=0} dt \leq \tilde{B}\epsilon^2,$$

which together with (7.10) yields (3.12). Therefore the solution can be continued up to $[0, T_\epsilon]$ with $1 + \bar{c}T_\epsilon \geq \exp\left(\frac{\kappa}{\epsilon}\right)$ for all $\kappa \leq \kappa_0$. In particular, choosing $\kappa = \kappa_0$ gives $1 + \bar{c}T_\epsilon \geq \exp\left(\frac{\kappa_0}{\epsilon}\right)$.

7.2. Estimate of bubble radius

Now assume that $1 + \bar{c}T = \exp\left(\frac{\kappa}{\epsilon}\right)$ for $\kappa \leq \kappa_0$ and $t \in [0, T]$. To retrieve an estimate of $R(t)$, recall the relation (1.15). For simplicity, denote

$$g(q) := -\frac{Ca\gamma}{2(\gamma-1)} \left((1+q)^{1-\gamma} - 1 \right) \quad (7.11)$$

so that

$$g(q) = \partial_t \varphi - \frac{1}{2}u^2 = r^{-1}\partial_t \psi - \frac{u}{r^2}\psi - \frac{1}{2}u^2.$$

From the boundary condition (1.12), it follows that

$$F(R) = \partial_t \psi|_{x=0} - \left(\frac{u}{r}\psi + \frac{1}{2}ru^2 \right) \Big|_{x=0}, \quad (7.12)$$

where

$$F(R) = R \cdot g \circ f(R). \quad (7.13)$$

Write

$$\partial_t \psi|_{x=0} = \frac{\Lambda_1}{\Lambda_1 - \Lambda_2} (\partial_t \psi - \Lambda_2 \psi)|_{x=0} - \frac{\Lambda_2}{\Lambda_1 - \Lambda_2} (\partial_t \psi - \Lambda_1 \psi)|_{x=0},$$

and from (6.41) we obtain

$$\begin{aligned} & |\partial_t \psi|_{x=0} - \mathcal{R}(t)| \\ & \lesssim \frac{1}{1 + \bar{c}t} \left\{ \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon |Y(0)| \right\} \\ & \lesssim \frac{1}{1 + \bar{c}t} \left(\kappa \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \epsilon^2 + \epsilon^{\frac{3}{2}} \tilde{\epsilon}^{\frac{1}{2}} \right), \end{aligned} \quad (7.14)$$

where we used Lemma 7.1 and recall that

$$\begin{aligned} \mathcal{R}(t) & := \frac{\Lambda_1}{\Lambda_1 - \Lambda_2} e^{\Lambda_1 t} (\partial_t \psi - \Lambda_2 \psi)|_{\{x=0, t=0\}} - \frac{\Lambda_2}{\Lambda_1 - \Lambda_2} e^{\Lambda_2 t} (\partial_t \psi - \Lambda_1 \psi)|_{\{x=0, t=0\}} \\ & \quad - \frac{\tilde{f}'(1)}{c_0} \int_0^t \left(\frac{\Lambda_1}{\Lambda_1 - \Lambda_2} e^{\Lambda_1 s} - \frac{\Lambda_2}{\Lambda_1 - \Lambda_2} e^{\Lambda_2 s} \right) w_B(\xi_0(s), 0) ds. \end{aligned}$$

To control the nonlinearity $\left(\frac{u}{r}\psi + \frac{1}{2}ru^2 \right) \Big|_{x=0}$, we use

$$u = \rho r^2 \partial_x \varphi = \rho r \partial_x \psi - r^{-2} \psi = \frac{\rho}{cr} \left(w_B - \partial_t \psi - \frac{c}{\rho r} \psi \right)$$

to obtain

$$\frac{u}{r} \psi + \frac{1}{2} r u^2 = \frac{\rho^2}{2c^2 r} \left(w_B - \partial_t \psi - \frac{c}{\rho r} \psi \right) \left(w_B - \partial_t \psi + \frac{c}{\rho r} \psi \right).$$

In view of (3.2)(3.10)(6.38)(6.41) and Lemma 7.1, there holds

$$\begin{aligned}
& \left| \partial_t \psi|_{x=0} + \left| \left(\frac{c}{\rho r} \psi \right) \Big|_{x=0} \right| \\
& \lesssim |Y(t) - Y_0(t)| + |Y_0(t)| \\
& \lesssim \frac{1}{1 + \underline{c}t} \left\{ \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon |Y(0)| \right\} + |Y_0(t)| \quad (7.15) \\
& \lesssim \frac{1}{1 + \underline{c}t} \left(\kappa \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \epsilon^{\frac{3}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \epsilon^2 \right) + \frac{1}{1 + \underline{c}t} \left(\epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \epsilon \right) \\
& \lesssim \frac{\epsilon^{\frac{1}{2}}}{1 + \underline{c}t} \left(\epsilon^{\frac{1}{2}} + \tilde{\epsilon}^{\frac{1}{2}} \right).
\end{aligned}$$

Meanwhile, by (6.31)(6.32)

$$|w_B(1, t)| \lesssim \frac{1}{1 + \underline{c}t} \left(\epsilon \max_{\eta \in [1, \xi_0(T)]} \eta \mathcal{V}(\eta) + \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \Psi(t_*) + \max_{\eta \in [1, \xi_0(T)]} \left\{ \eta \left| \frac{w_B}{c^{\frac{1}{2}}}(\eta, 0) \right| \right\} \right).$$

From (6.36)(6.39)(6.40), we obtain

$$\begin{aligned}
& \epsilon \max_{t_* \in [0, T]} \xi_0(t_*) \Psi(t_*) \\
& \lesssim \epsilon |Y(0)| + (\kappa + \epsilon^2) \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + (\kappa + \epsilon^2) \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)). \quad (7.16)
\end{aligned}$$

In view of $\epsilon, \kappa \lesssim 1$, it follows

$$\begin{aligned}
|w_B(1, t)| & \lesssim \frac{1}{1 + \underline{c}t} \left(\epsilon \max_{\eta \in [1, \xi_0(T)]} \eta \mathcal{V}(\eta) + \epsilon |Y(0)| + \max_{\eta \in [1, \xi_0(T)]} \left\{ \eta \left| \frac{w_B}{c^{\frac{1}{2}}}(\eta, 0) \right| \right\} \right) \\
& \lesssim \frac{1}{1 + \underline{c}t} \left(\epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \epsilon^2 \right). \quad (7.17)
\end{aligned}$$

Combing (7.15)(7.17) yields

$$\left| \left(\frac{u}{r} \psi + \frac{1}{2} r u^2 \right) \Big|_{x=0} \right| \lesssim \frac{\epsilon}{(1 + \underline{c}t)^2} (\epsilon + \tilde{\epsilon}).$$

Therefore (7.12)(7.14) give

$$|F(R) - \mathcal{R}(t)| \lesssim \frac{\kappa}{1 + \underline{c}t} \epsilon^{\frac{1}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \frac{\epsilon^{\frac{3}{2}}}{1 + \underline{c}t} \left(\epsilon^{\frac{1}{2}} + \tilde{\epsilon}^{\frac{1}{2}} \right) + \frac{\epsilon}{(1 + \underline{c}t)^2} (\epsilon + \tilde{\epsilon}). \quad (7.18)$$

Now we set $T = t$ so that $\kappa = \epsilon \log(1 + \bar{c}t)$. Then (7.18) becomes

$$|F(R) - \mathcal{R}(t)| \lesssim \frac{\log(1 + \bar{c}t)}{1 + \underline{c}t} \epsilon^{\frac{3}{2}} \tilde{\epsilon}^{\frac{1}{2}} + \frac{\epsilon^{\frac{3}{2}}}{1 + \underline{c}t} \left(\epsilon^{\frac{1}{2}} + \tilde{\epsilon}^{\frac{1}{2}} \right) + \frac{\epsilon}{(1 + \underline{c}t)^2} (\epsilon + \tilde{\epsilon}),$$

which is exactly the desired estimate of Theorem 1.3. If we assume additionally $w_B|_{t=0} = 0$, then (7.14)(7.15)(7.17) can be improved to

$$|\partial_t \psi|_{x=0} - \mathcal{R}(t)| \lesssim \frac{1}{1 + \underline{c}t} \left(\epsilon^2 + \epsilon^{\frac{3}{2}} \tilde{\epsilon}^{\frac{1}{2}} \right),$$

$$|\partial_t \psi|_{x=0} + \left| \left(\frac{c}{\rho r} \psi \right) \Big|_{x=0} \right| \lesssim \frac{1}{1 + \underline{c}t} \left(\epsilon^{\frac{3}{2}} \tilde{c}^{\frac{1}{2}} + \epsilon^2 \right) + \exp \left(\max_{i=1,2} \{\operatorname{Re} \Lambda_i\} t \right) \epsilon,$$

$$|w_B(1, t)| \lesssim \frac{1}{1 + \underline{c}t} \left(\epsilon^{\frac{3}{2}} \tilde{c}^{\frac{1}{2}} + \epsilon^2 \right).$$

Hence

$$\left| \left(\frac{u}{r} \psi + \frac{1}{2} r u^2 \right) \Big|_{x=0} \right| \lesssim \left(\frac{1}{1 + \underline{c}t} \left(\epsilon^2 + \epsilon^{\frac{3}{2}} \tilde{c}^{\frac{1}{2}} \right) + \exp \left(\max_{i=1,2} \{\operatorname{Re} \Lambda_i\} t \right) \epsilon \right)^2,$$

$$|F(R) - \mathcal{R}(t)| \lesssim \frac{\epsilon^{\frac{3}{2}}}{1 + \underline{c}t} \left(\epsilon^{\frac{1}{2}} + \tilde{c}^{\frac{1}{2}} \right) + \exp \left(2 \max_{i=1,2} \{\operatorname{Re} \Lambda_i\} t \right) \epsilon^2 \lesssim \frac{\epsilon^{\frac{3}{2}}}{1 + \underline{c}t} \left(\epsilon^{\frac{1}{2}} + \tilde{c}^{\frac{1}{2}} \right).$$

7.3. Proof of Corollary 1.4

Now we assume additionally u_{in}, q_{in} are compactly supported in $[0, x_b)$. From the relations $u = \rho r^2 \partial_x \varphi$, $\psi = r \varphi$ and (1.15), it follows $w_B(\cdot, 0) = (\partial_t + cr^2 \xi^{-2} \partial_\xi) \psi|_{t=0}$ is compactly supported in $\xi \in [1, \xi_b)$ with $\xi_b = (1 + x_b)^{\frac{1}{2}}$. Suppose (x, t) satisfies (1.19)(1.20). Define T such that $T = t + \frac{\xi - 1}{\underline{c}}$ if $t \geq \frac{1}{\underline{c}}$ and $T = \frac{\xi}{\underline{c}}$ if $t < \frac{1}{\underline{c}}$, so by (1.19)(1.20) we have $T \leq T_0$. Let $\kappa \leq \kappa_0$ be such that $1 + \bar{c}T = \exp\left(\frac{\kappa}{\epsilon}\right)$, so we always have $\kappa \gtrsim \epsilon$. Note that ξ_0 is a bijection from $[0, T]$ to $[1, \xi_0(T)]$. Hence taking (7.16) back to (6.30) yields that for $\eta \in [1, \xi_0(T)]$

$$v_B(\eta, 0) - \left| \frac{w_B}{c^{\frac{1}{2}}}(\eta, 0) \right| \lesssim \eta^{-1} \left\{ \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \right\}. \quad (7.19)$$

For any $t' \leq T$ with $\xi_0(t') \geq \xi_b$, using (7.19) in (6.18) gives that

$$v_F(1, t'; T) - \left| \frac{w_F}{c^{\frac{1}{2}}}(1, t') \right| \lesssim \epsilon \left| \frac{w_F}{c^{\frac{1}{2}}}(1, t') \right| + \epsilon^2 |\psi(1, t')| + \epsilon \max_{\eta \in [\xi_0(t'), \xi_0(T)]} \eta v_B(\eta, 0) \int_{\xi_0(t')}^{\xi_0(T)} \left(1 + \frac{1}{2\mu} (\eta - \xi_0(t')) \right)^{-1} \eta^{-1} d\eta$$

$$\lesssim \epsilon \left| \frac{w_F}{c^{\frac{1}{2}}}(1, t') \right| + \epsilon^2 |\psi(1, t')| + \epsilon \int_{\xi_0(t')}^{\xi_0(T)} \left(1 + \frac{1}{2\mu} (\eta - \xi_0(t')) \right)^{-1} \eta^{-1} d\eta$$

$$\cdot \left\{ \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \right\}.$$

We compute for $\xi_0(t') > 2\mu$ that

$$\int_{\xi_0(t')}^{\xi_0(T)} \left(1 + \frac{1}{2\mu} (\eta - \xi_0(t')) \right)^{-1} \eta^{-1} d\eta$$

$$= \frac{2\mu}{\xi_0(t') - 2\mu} \int_{\xi_0(t')}^{\xi_0(T)} \left(\frac{1}{\eta - \xi_0(t') + 2\mu} - \frac{1}{\eta} \right) d\eta$$

$$= \frac{2\mu}{\xi_0(t') - 2\mu} \log \frac{(\xi_0(T) - \xi_0(t') + 2\mu) \xi_0(t')}{2\mu \xi_0(T)}$$

$$= \frac{2\mu}{\xi_0(t') - 2\mu} \log \left(1 + \frac{1}{2\mu \xi_0(T)} (\xi_0(T) - \xi_0(t')) (\xi_0(t') - 2\mu) \right).$$

If $\xi_0(t') \in (2\mu, 2\mu + 1]$,

$$\frac{2\mu}{\xi_0(t') - 2\mu} \log \left(1 + \frac{1}{2\mu\xi_0(T)} (\xi_0(T) - \xi_0(t')) (\xi_0(t') - 2\mu) \right) \leq \frac{\xi_0(T) - \xi_0(t')}{\xi_0(T)} \leq 1 \lesssim \frac{1}{\xi_0(t')}.$$

If $\xi_0(t') > 2\mu + 1$, then

$$\begin{aligned} & \frac{2\mu}{\xi_0(t') - 2\mu} \log \left(1 + \frac{1}{2\mu\xi_0(T)} (\xi_0(T) - \xi_0(t')) (\xi_0(t') - 2\mu) \right) \\ & \leq \frac{2\mu}{\xi_0(t') - 2\mu} \log \left(1 + \frac{(\xi_0(T) - 2\mu)^2}{8\mu\xi_0(T)} \right) \\ & \lesssim \frac{\kappa}{\epsilon} \frac{1}{\xi_0(t')}. \end{aligned}$$

For $\xi_0(t') \leq 2\mu$, the estimate is much simpler:

$$\int_{\xi_0(t')}^{\xi_0(T)} \left(1 + \frac{1}{2\mu} (\eta - \xi_0(t')) \right)^{-1} \eta^{-1} d\eta \leq 2\mu \int_{\xi_0(t')}^{\xi_0(T)} \eta^{-2} d\eta \leq 2\mu\xi_0(t')^{-1}.$$

Hence in each case we have

$$\int_{\xi_0(t')}^{\xi_0(T)} \left(1 + \frac{1}{2\mu} (\eta - \xi_0(t')) \right)^{-1} \eta^{-1} d\eta \lesssim \frac{\kappa}{\epsilon} \xi_0(t')^{-1},$$

and it follows

$$\begin{aligned} & \left| v_F(1, t'; T) - \left| \frac{w_F}{c^{\frac{1}{2}}}(1, t') \right| \right| \\ & \lesssim \epsilon \left| \frac{w_F}{c^{\frac{1}{2}}}(1, t') \right| + \epsilon^2 |\psi(1, t')| \\ & \quad + \kappa \xi_0(t')^{-1} \left\{ \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \right\}. \end{aligned} \tag{7.20}$$

Since $w_F = -w_B + 2\partial_t \psi$, there holds $|w_F(1, t')| \leq |w_B(1, t')| + 2|\partial_t \psi(1, t')|$, and thus

$$\left| \frac{w_F}{c^{\frac{1}{2}}}(1, t') \right| \lesssim v_B(1, t') + |Y(t')| = v_B(\xi_0(t'), 0) + |Y(t')|. \tag{7.21}$$

Denote $\Lambda := \max \{\text{Re}\Lambda_1, \text{Re}\Lambda_2\}$. Since w_B is compactly supported in $[1, \xi_b)$ and $\xi_0(t') \geq 1 + \underline{c}t'$, we have by (6.41) that

$$\begin{aligned} |Y(t')| & \leq e^{\Lambda t'} \left(|Y(0)| + \frac{|\tilde{f}'(1)|}{c_0} \int_0^{\frac{\xi_b - 1}{\underline{c}}} e^{-\Lambda s} |w_B(\xi_0(s), 0)| ds \right) \\ & \quad + \frac{1}{1 + \underline{c}t'} \left\{ \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \right\}. \end{aligned} \tag{7.22}$$

Using $T_\xi(t) \geq t - \frac{\xi-1}{\underline{c}}$ and (1.20), there holds

$$\xi_0(T_\xi(t)) \geq 1 + \underline{c}T_\xi(t) \geq \underline{c}t + 2 - \xi \geq \xi_b.$$

Hence it follows from (7.19) and $w_B(\eta, 0) = 0$, $\eta \geq \xi_b$ that

$$\begin{aligned} |\psi(\xi, t)| &\lesssim |Y(T_\xi(t))| + \int_{\xi_0(T_\xi(t))}^{X_B^{(\xi)}(\xi, t; -t)} \eta^{-1} d\eta \\ &\cdot \left\{ \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \right\}. \end{aligned}$$

We claim that for all $s \in [0, T_\xi(t)]$,

$$X_B^{(\xi)}(\xi, t; T_\xi(t) - t - s) - X_B^{(\xi)}(1, T_\xi(t); -s) \leq e^{C\kappa} \left(X_B^{(\xi)}(\xi, t; T_\xi(t) - t) - 1 \right).$$

In particular, taking $s = T_\xi(t)$ yields

$$\begin{aligned} X_B^{(\xi)}(\xi, t; -t) - \xi_0(T_\xi(t)) &\leq e^{C\kappa} \left(X_B^{(\xi)}(\xi, t; T_\xi(t) - t) - 1 \right) \\ &\leq e^{C\kappa} (\xi - 1 + \bar{c}(t - T_\xi(t))) \leq e^{C\kappa} \frac{\bar{c} + \underline{c}}{\underline{c}} (\xi - 1). \end{aligned}$$

Note that $X_B^{(\xi)}(\xi, t; T_\xi(t) - t - \tau) = X_B^{(\xi)}(X_B(\xi, t; T_\xi(t) - t); -\tau)$. Using (3.6)(3.7) and a mean value formula, we obtain

$$\begin{aligned} &\left| \frac{d}{d\tau} \left(X_B^{(\xi)}(\xi, t; T_\xi(t) - t - \tau) - X_B^{(\xi)}(1, T_\xi(t); -\tau) \right) \right| \\ &\leq C\epsilon X_B^{(\xi)}(1, T_\xi(t); -\tau)^{-1} \left(X_B^{(\xi)}(\xi, t; T_\xi(t) - t - \tau) - X_B^{(\xi)}(1, T_\xi(t); -\tau) \right) \\ &\leq C\epsilon(1 + \underline{c}\tau)^{-1} \left(X_B^{(\xi)}(\xi, t; T_\xi(t) - t - \tau) - X_B^{(\xi)}(1, T_\xi(t); -\tau) \right). \end{aligned}$$

Gronwall's inequality then yields

$$\begin{aligned} &X_B^{(\xi)}(\xi, t; T_\xi(t) - t - s) - X_B^{(\xi)}(1, T_\xi(t); -s) \\ &\leq \exp \left\{ \int_0^s C\epsilon(1 + \underline{c}\tau)^{-1} d\tau \right\} \left(X_B^{(\xi)}(\xi, t; T_\xi(t) - t) - 1 \right) \\ &\leq e^{C\kappa} \left(X_B^{(\xi)}(\xi, t; T_\xi(t) - t) - 1 \right), \end{aligned}$$

which proves the claim. Therefore

$$\int_{\xi_0(T_\xi(t))}^{X_B^{(\xi)}(\xi, t; -t)} \eta^{-1} d\eta \leq \log \left(1 + \xi_0(T_\xi(t))^{-1} e^{C\kappa} \frac{\bar{c} + \underline{c}}{\underline{c}} (\xi - 1) \right) \lesssim \xi_0(T_\xi(t))^{-1} (\xi - 1),$$

and

$$\begin{aligned} |\psi(\xi, t)| &\lesssim |Y(T_\xi(t))| + \xi_0(T_\xi(t))^{-1} (\xi - 1) \\ &\cdot \left\{ \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \right\}. \end{aligned} \quad (7.24)$$

Now we take $\eta = X_B(\xi, t; -t)$ in (7.19), $t' = T_\xi(t)$ in (7.20), and collect the resulted bounds as well as (7.24). It then follows from (7.23) that

$$\begin{aligned} |u(x, t)| &\lesssim \xi^{-1} \left| \frac{w_F}{c^{\frac{1}{2}}}(1, T_\xi(t)) \right| + \xi^{-2} |Y(T_\xi(t))| + \epsilon^2 \xi^{-1} |\psi(1, T_\xi(t))| \\ &\quad + \left(\xi^{-1} X_B^{(\xi)}(\xi, t; -t)^{-1} + \xi^{-2} (\xi - 1) \xi_0(T_\xi(t))^{-1} + \kappa \xi^{-1} \xi_0(T_\xi(t))^{-1} \right) \\ &\quad \cdot \left\{ \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \right\}. \end{aligned}$$

Next, we proceed by (7.21) and using (7.19) again:

$$\begin{aligned} |u(x, t)| &\lesssim \xi^{-1} |Y(T_\xi(t))| + \left(\xi^{-1} X_B^{(\xi)}(\xi, t; -t)^{-1} + \xi^{-2} (\xi - 1) \xi_0(T_\xi(t))^{-1} + \xi^{-1} \xi_0(T_\xi(t))^{-1} \right) \\ &\quad \cdot \left\{ \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \right\} \\ &\lesssim \xi^{-1} |Y(T_\xi(t))| + \xi^{-1} \xi_0(T_\xi(t))^{-1} \\ &\quad \cdot \left\{ \epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \kappa \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \right\}, \end{aligned} \tag{7.25}$$

where we used that $\xi^{-2}(\xi - 1) \leq \xi^{-1}$ and $\xi_0(T_\xi(t)) \leq X_B^{(\xi)}(\xi, t; -t)$. Since u_{in}, q_{in} are compactly supported in $[1, x_b)$, we see that $\tilde{\epsilon} \lesssim \xi_b^2 \epsilon$. Hence from Lemma 7.1 it follows

$$\epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \lesssim \kappa \epsilon \xi_b + \epsilon^2 \xi_b \lesssim \kappa \epsilon \xi_b.$$

Using (7.22) and that $T_\xi(t) \geq t - \frac{\xi-1}{\underline{c}}$, $\xi_0(T_\xi(t)) \geq 1 + \underline{c}T_\xi(t) \geq 2 + \underline{c}t - \xi$, (7.25) gives further that

$$|u(x, t)| \lesssim \xi^{-1} e^{\Lambda(t - \frac{\xi-1}{\underline{c}})} \left(|Y(0)| + \frac{|\tilde{f}'(1)|}{c_0} \int_0^{\frac{\xi_b-1}{\underline{c}}} e^{-\Lambda s} |w_B(\xi_0(s), 0)| ds \right) + \frac{\kappa \epsilon \xi_b}{\xi(2 + \underline{c}t - \xi)}.$$

Hence for $t \geq \frac{1}{\underline{c}}$ we obtain the desired bound of u

$$\begin{aligned} |u(x, t)| &\lesssim \xi^{-1} e^{\Lambda(t - \frac{\xi-1}{\underline{c}})} \left(|Y(0)| + \frac{|\tilde{f}'(1)|}{c_0} \int_0^{\frac{\xi_b-1}{\underline{c}}} e^{-\Lambda s} |w_B(\xi_0(s), 0)| ds \right) \\ &\quad + \xi^{-1} (2 + \underline{c}t - \xi)^{-1} \log \left(1 + \bar{c} \left(t + \frac{\xi-1}{\underline{c}} \right) \right) \epsilon^2 \xi_b. \end{aligned} \tag{7.26}$$

If assume additionally $w_B|_{t=0} = 0$, then

$$\epsilon \max_{s \in [0, T]} \xi_0(s) \mathcal{V}(\xi_0(s)) + \epsilon \max_{s \in [0, T]} \left\{ \xi_0(s) \left| \frac{w_B}{c^{\frac{1}{2}}}(\xi_0(s), 0) \right| \right\} + \epsilon |Y(0)| \lesssim \epsilon^2 \xi_b,$$

and thus

$$\begin{aligned} |u(x, t)| &\lesssim \xi^{-1} e^{\Lambda(t - \frac{\xi-1}{\underline{c}})} \left(|Y(0)| + \frac{|\tilde{f}'(1)|}{c_0} \int_0^{\frac{\xi_b-1}{\underline{c}}} e^{-\Lambda s} |w_B(\xi_0(s), 0)| ds \right) \\ &\quad + \xi^{-1} (2 + \underline{c}t - \xi)^{-1} \epsilon^2 \xi_b. \end{aligned} \tag{7.27}$$

In view of $\rho \simeq 1$, $|u| \lesssim \epsilon$, relation (1.15) and that $\partial_t \varphi = r^{-1} \partial_t \psi - \frac{u}{r^2} \psi$, we can bound q by

$$\begin{aligned} |q(x, t)| &\lesssim \xi^{-1} |\partial_t \psi(\xi, t)| + |u(x, t)| (|u(x, t)| + \xi^{-2} |\psi(\xi, t)|) \\ &\lesssim \xi^{-1} (|w_B(\xi, t)| + |w_F(\xi, t)|) + |u(x, t)| (|u(x, t)| + \xi^{-2} |\psi(\xi, t)|) \\ &\lesssim \xi^{-1} (v_B(X_B(\xi, t; -t)) + v_F(1, \xi_0(T_\xi(t); T))) + \xi^{-2} |\psi(\xi, t)|. \end{aligned}$$

Hence $|q(x, t)|$ can be controlled by the same bound as in (7.26) or (7.27).

Appendix A.

In this appendix we write down the explicit calculation of (4.3), for which we consider the following model question.

Assume that φ satisfies for $x > 0$, $t \in [0, T]$ that

$$\partial_t^2 \varphi - c^2 \partial_x (r^4 \partial_x \varphi) = F, \quad (\text{A.1})$$

and the modified momentum density are defined as

$$P_0 = A \partial_t \varphi \cdot cr^2 \partial_x \varphi + \frac{1}{2} B \varphi \partial_t \varphi,$$

$$P_1 = \frac{1}{2} A \left[(\partial_t \varphi)^2 + (cr^2 \partial_x \varphi)^2 \right] + \frac{1}{2} B \varphi \cdot cr^2 \partial_x \varphi - \frac{1}{4} cr^2 \partial_x B \varphi^2,$$

where A and B are undetermined weight functions. We compute:

$$\begin{aligned} \partial_x (cr^2 P_1) &= \frac{1}{2} \partial_x \left[Acr^2 \left((\partial_t \varphi)^2 + (cr^2 \partial_x \varphi)^2 \right) \right] + \frac{1}{2} B (cr^2 \partial_x \varphi)^2 + \frac{1}{2} B \varphi \partial_x (c^2 r^4 \partial_x \varphi) \\ &\quad + \frac{1}{2} cr^2 \partial_x B \cdot \varphi \cdot cr^2 \partial_x \varphi - \frac{1}{4} \partial_x (c^2 r^4 \partial_x B) \varphi^2 - \frac{1}{2} cr^2 \partial_x B \cdot \varphi \cdot cr^2 \partial_x \varphi \\ &= \frac{1}{2} \partial_x (cr^2 A) \left((\partial_t \varphi)^2 + (cr^2 \partial_x \varphi)^2 \right) + \frac{1}{2} Acr^2 \partial_x \left((\partial_t \varphi)^2 + (cr^2 \partial_x \varphi)^2 \right) \\ &\quad + \frac{1}{2} B (cr^2 \partial_x \varphi)^2 + \frac{1}{2} B \varphi \partial_x (c^2 r^4 \partial_x \varphi) - \frac{1}{4} \partial_x (c^2 r^4 \partial_x B) \varphi^2, \end{aligned}$$

$$\begin{aligned} -\partial_t P_0 &= -\partial_t (A \partial_t \varphi \cdot cr^2 \partial_x \varphi) - \frac{1}{2} (\partial_t B) \varphi \partial_t \varphi - \frac{1}{2} B (\partial_t \varphi)^2 - \frac{1}{2} B \varphi \partial_t^2 \varphi \\ &= -(\partial_t A) \partial_t \varphi \cdot cr^2 \partial_x \varphi - A \partial_t (\partial_t \varphi \cdot cr^2 \partial_x \varphi) - \frac{1}{2} B (\partial_t \varphi)^2 - \frac{1}{2} B \varphi \partial_t^2 \varphi - \frac{1}{2} (\partial_t B) \varphi \partial_t \varphi. \end{aligned}$$

Note by A.1 that

$$\begin{aligned} &\frac{1}{2} cr^2 \partial_x \left((\partial_t \varphi)^2 + (cr^2 \partial_x \varphi)^2 \right) - \partial_t (\partial_t \varphi \cdot cr^2 \partial_x \varphi) \\ &= \partial_t \varphi (cr^2 \partial_x \partial_t \varphi - \partial_t (cr^2 \partial_x \varphi)) + cr^2 \partial_x \varphi (cr^2 \partial_x (cr^2 \partial_x \varphi) - \partial_t^2 \varphi) \\ &= -\frac{\partial_t (cr^2)}{cr^2} \partial_t \varphi \cdot cr^2 \partial_x \varphi + cr^2 \partial_x \varphi \left(cr^2 \partial_x \left(\frac{c}{r^2} \right) r^4 \partial_x \varphi - F \right) \\ &= -c \partial_x r^2 (cr^2 \partial_x \varphi)^2 + r^2 \partial_x c (cr^2 \partial_x \varphi)^2 - \frac{\partial_t (cr^2)}{cr^2} \partial_t \varphi \cdot cr^2 \partial_x \varphi - cr^2 \partial_x \cdot F, \end{aligned}$$

and

$$\partial_x (c^2 r^4 \partial_x \varphi) - \partial_t^2 \varphi = 2 (r^2 \partial_x c) c r^2 \partial_x \varphi - F.$$

Hence

$$\begin{aligned} & \partial_x (c r^2 P_1) - \partial_t P_0 \\ &= \left(\frac{\partial_x (c r^2 A)}{2} - \frac{B}{2} \right) (\partial_t \varphi)^2 + \left(\frac{\partial_x (c r^2 A)}{2} - A c \partial_x r^2 + \frac{B}{2} \right) (c r^2 \partial_x \varphi)^2 \\ & \quad - \frac{1}{4} \partial_x (c^2 r^4 \partial_x B) \varphi^2 + r^2 \partial_x c \cdot A (c r^2 \partial_x \varphi)^2 - A \frac{\partial_t (c r^2)}{c r^2} \partial_t \varphi \cdot c r^2 \partial_x \varphi \\ & \quad + B r^2 \partial_x c \cdot \varphi \cdot c r^2 \partial_x \varphi - \left(A c r^2 \partial_x \varphi + \frac{1}{2} B \varphi \right) F - \partial_t A \cdot \partial_t \varphi \cdot c r^2 \partial_x \varphi - \frac{\partial_t B}{2} \varphi \partial_t \varphi. \end{aligned}$$

Therefore we obtain (4.3) by substituting φ by $\partial_t^j \varphi$, F by $[\partial_t^j, c^2 \partial_x r^4 \partial_x] \varphi$, A by M_k and B by N_k in view of (4.2).

References

- [1] J.-P. Franc. *The Rayleigh-Plesset equation: a simple and powerful tool to understand various aspects of cavitation*, pages 1–41. Springer Vienna, Vienna, 2007.
- [2] P. Frinking, T. Segers, Y. Luan, and F. Tranquart. Three decades of ultrasound contrast agents: A review of the past, present and future improvements. *Ultrasound in Medicine & Biology*, 46(4):892–908, 2020.
- [3] F. R. Gilmore. The collapse and growth of a spherical bubble in a viscous compressible liquid. *california institute of technology*, 1952.
- [4] N. J. Higham, M. R. Dennis, P. Glendinning, P. A. Martin, F. Santosa, and J. Tanner, editors. *The Princeton Companion to Applied Mathematics*. Princeton University Press, Princeton, 2016.
- [5] Z. Huang, S. Zhao, M. Su, Q. Yang, Z. Li, Z. Cai, H. Zhao, X. Hu, H. Zhou, F. Li, J. Yang, Y. Wang, and Y. Song. Bioinspired patterned bubbles for broad and low-frequency acoustic blocking. *ACS Applied Materials & Interfaces*, 12(1):1757–1764, 2020.
- [6] Z. Huang, S. Zhao, Y. Zhang, Z. Cai, Z. Li, J. Xiao, M. Su, Q. Guo, C. Zhang, Y. Pan, X. Cai, Y. Song, and J. Yang. Tunable fluid-type metasurface for wide-angle and multifrequency water-air acoustic transmission. *Research*, 2021, 2021.
- [7] F. John. *Blow-Up for Quasi-Linear Wave Equations in Three Space Dimensions*, pages 566–588. Birkhäuser Boston, Boston, MA, 1985.
- [8] M. Keel, H. Smith, and C. D. Sogge. Almost global existence for some semilinear wave equations. *J. Anal. Math.*, 87, 09 2001.
- [9] M. Keel, H. Smith, and C. D. Sogge. Almost global existence for quasilinear wave equations in three space dimensions. *J. Amer. Math. Soc.*, 17(1):109–153, 2004.

- [10] J. B. Keller and I. I. Kolodner. Damping of underwater explosion bubble oscillations. *Journal of Applied Physics*, 27(10):1152–1161, 1956.
- [11] J. B. Keller and M. Miksis. Bubble oscillations of large amplitude. *The Journal of the Acoustical Society of America*, 68(2):628–633, 08 1980.
- [12] E. Klaseboer and B.C. Khoo. A modified rayleigh–plesset model for a non-spherically symmetric oscillating bubble with applications to boundary integral methods. *Engineering Analysis with Boundary Elements*, 30(1):59–71, 2006.
- [13] K. Kooiman, H. J. Vos, M. Versluis, and N. de Jong. Acoustic behavior of microbubbles and implications for drug delivery. *Advanced Drug Delivery Reviews*, 72:28–48, 2014. Ultrasound triggered drug delivery.
- [14] C.-C. Lai and M. I. Weinstein. Free boundary problem for a gas bubble in a liquid, and asymptotic stability of the manifold of spherically symmetric equilibria. *arXiv e-prints*, page arXiv:2207.04079, July 2022.
- [15] T. G. Leighton. From seas to surgeries, from babbling brooks to baby scans: The acoustics of gas bubbles in liquids. *International Journal of Modern Physics B*, 18(25):3267–3314, 2004.
- [16] A. Lezzi and A. Prosperetti. Bubble dynamics in a compressible liquid. part 2. second-order theory. *Journal of Fluid Mechanics*, 185:289–321, 1987.
- [17] A. Majda. *Compressible fluid flow and systems of conservation laws in several space variables*. Applied Mathematical Sciences. Springer, 1984.
- [18] J. Metcalfe, M. Nakamura, and C. D. Sogge. Global existence of quasilinear, nonrelativistic wave equations satisfying the null condition. *Japan. J. Math. (N.S.)*, 31(2):391–472, 2005.
- [19] J. Metcalfe and C. D. Sogge. Global existence for Dirichlet-wave equations with quadratic nonlinearities in high dimensions. *Math. Ann.*, 336(2):391–420, 2006.
- [20] J. Metcalfe and C. D. Sogge. Long-time existence of quasilinear wave equations exterior to star-shaped obstacles via energy methods. *SIAM J. Math. Anal.*, 38(1):188–209, 2006.
- [21] J. Metcalfe and C. D. Sogge. Global existence of null-form wave equations in exterior domains. *Math. Z.*, 256(3):521–549, 2007.
- [22] J. Metcalfe and C. D. Sogge. Global existence for high dimensional quasilinear wave equations exterior to star-shaped obstacles. *Discrete Contin. Dyn. Syst.*, 28(4):1589–1601, 2010.
- [23] M. S. Plesset. The Dynamics of Cavitation Bubbles. *Journal of Applied Mechanics*, 16(3):277–282, 09 1949.
- [24] A. Prosperetti. A generalization of the Rayleigh–Plesset equation of bubble dynamics. *The Physics of Fluids*, 25(3):409–410, 03 1982.
- [25] A. Prosperetti and A. Lezzi. Bubble dynamics in a compressible liquid. part 1. first-order theory. *Journal of Fluid Mechanics*, 168:457–478, 1986.

- [26] L. Rayleigh. Viii. On the pressure developed in a liquid during the collapse of a spherical cavity. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 34(200):94–98, 1917.
- [27] A. M. Shapiro and M. I. Weinstein. Radiative decay of bubble oscillations in a compressible fluid. *SIAM Journal on Mathematical Analysis*, 43(2):828–876, 2011.
- [28] T. C. Sideris. Global behavior of solutions to nonlinear wave equations in three dimensions. *Communications in Partial Differential Equations*, 8(12):1291–1323, 1983.
- [29] T. C. Sideris. Formation of singularities in three-dimensional compressible fluids. *Communications in Mathematical Physics*, 101(4):475 – 485, 1985. Cited by: 359; All Open Access, Green Open Access.
- [30] W. R. Smith and Q. X. Wang. Radiative decay of the nonlinear oscillations of an adiabatic spherical bubble at small mach number. *Journal of Fluid Mechanics*, 837:1–18, 2018.
- [31] Q. X. Wang. Local energy of a bubble system and its loss due to acoustic radiation. *Journal of Fluid Mechanics*, 797:201–230, 2016.
- [32] Q. X. Wang and J. R. Blake. Non-spherical bubble dynamics in a compressible liquid. part 1. travelling acoustic wave. *Journal of Fluid Mechanics*, 659:191–224, 2010.
- [33] Q. X. Wang and J. R. Blake. Non-spherical bubble dynamics in a compressible liquid. part 2. acoustic standing wave. *Journal of Fluid Mechanics*, 679:559–581, 2011.
- [34] Lifeng Zhao and Liangchen Zou. The viscous damping of three dimensional spherical gas bubble inside unbounded compressible liquid. *arXiv e-prints*, page arXiv:2212.00299, December 2022.