

# A synthetic approach to detecting $v_1$ -periodic families

Christian Carrick\* and Jack Morgan Davies†

January 31, 2024

## Abstract

We provide a simple proof that the unit map from the sphere spectrum to the connective image-of- $J$  spectrum  $j$  is surjective on homotopy groups. This is achieved using a novel  $t$ -structure on the category of  $E$ -synthetic spectra and a specific construction of  $\mathbf{F}_p$ - and BP-synthetic lifts of  $j$ . These synthetic lifts then easily produce modified Adams and Adams–Novikov spectral sequences for  $j$  which we use to prove the above detection statement, all without ever calculating  $\mathbf{F}_p$ - or BP-homology nor the associated Ext groups.

## Contents

<b>Introduction</b>	<b>2</b>
<b>1 The signature of a synthetic spectrum</b>	<b>6</b>
<b>2 Linear <math>t</math>-structures on synthetic spectra</b>	<b>11</b>
<b>3 Synthetic analogues of topological <math>K</math>-theory</b>	<b>17</b>
<b>4 Two synthetic image-of-<math>J</math> spectra</b>	<b>24</b>
<b>5 Adams <math>v_1</math> self-maps and detection for <math>ko</math></b>	<b>29</b>
<b>6 Detection results for <math>j</math></b>	<b>32</b>
<b>References</b>	<b>35</b>
<b>A Spectral sequence diagrams</b>	<b>37</b>

---

\*c.d.carrick@uu.nl

†davies@math.uni-bonn.de

## Introduction

New synthetic models of the connective image-of- $J$  spectrum  $\mathbf{j}$  are constructed inside both  $\mathbf{F}_p$ - and BP-synthetic spectra for each prime  $p$ . These models are then used to produce modified Adams and Adams–Novikov spectral sequences for  $\mathbf{j}$  without ever considering its  $\mathbf{F}_p$ - or BP-homology—these are drastic simplifications to the previous methods to calculate related spectral sequences for  $\mathbf{j}$ . Indeed, the classical Adams spectral sequence (ASS) for  $\mathbf{j}$  requires difficult and subtle computations, going back to Davis [Dav75] and completed only recently by Bruner–Rognes [BR22]. As far as the authors are aware, there is no computation of an Adams–Novikov spectral sequence (ANSS) of  $\mathbf{j}$  in the literature nor of the comodule  $\mathrm{BP}_*\mathbf{j}$ . In contrast, using synthetic methods we can easily produce a modified ASS and a modified ANSS for  $\mathbf{j}$  and compute them completely using only minimal input, such as the effect of the Adams operations on the homotopy groups of  $\mathrm{KU}$ . To demonstrate the utility of these modified spectral sequences, we reprove the following classical detection statement.

**Theorem A.** *Fix a prime  $p$ . The unit map  $\mathbf{S} \rightarrow \mathbf{j}$  induces a surjection on homotopy groups.*

Since the homotopy groups of  $\mathbf{j}$  are readily calculated, this theorem recovers well-known collections of infinite families in the stable homotopy groups of the sphere  $\mathbf{S}$ . Moreover, as the above unit map is known to induce Adams  $e$ -invariant on homotopy groups, such a detection statement provides a splitting of the inclusion of the image of the  $J$ -homomorphism and the  $\mu$ -family. Our modified ANSS also provides a filtration for  $\pi_*\mathbf{j}$  such that this classical splitting of abelian groups refines to a splitting of filtered abelian groups; see Th.C.

## History and motivation

The  $J$ -homomorphism is a nontrivial homomorphism from the homotopy groups of the infinite special orthogonal group to the stable homotopy groups of spheres  $\pi_*\mathbf{S}$ . In [Ada66], Adams uses real topological  $K$ -theory  $\mathrm{KO}$  to study the  $J$ -homomorphism, and showed two facts regarding  $\pi_*\mathbf{S}$  at the prime 2:

- The image of the  $J$ -homomorphism detects a certain cyclic subgroup of  $\pi_k\mathbf{S}$  for  $k \geq 1$  congruent to 0, 1, 3, or 7 modulo 8. The order of this subgroup was determined by Adams in [Ada66] up to a factor of 2 in degrees congruent to 7 modulo 8; this ambiguity is known as the *Adams conjecture* and was confirmed by Quillen [Qui71].
- The unit map  $\mathbf{S} \rightarrow \mathrm{KO}$ , arising from the fact that  $\mathrm{KO}$  is a ring spectrum, detects certain cyclic subgroups of order 2 inside  $\pi_k\mathbf{S}$  for  $k \geq 1$  congruent to 1 or 2 modulo 8. These subgroups are generated by the so-called  *$\mu$ -family*.

It was not long before a spectrum was discovered that detects both the image of the  $J$ -homomorphism and the  $\mu$ -family at once. The (connective) *image-of- $J$  spectrum*  $\mathbf{j}$  is defined at the prime 2 using the fibre sequence

$$\mathbf{j} \rightarrow \mathrm{ko} \xrightarrow{\psi^3-1} \tau_{\geq 4} \mathrm{ko}$$

where  $ko = \tau_{\geq 0} KO$  is connective real topological  $K$ -theory and  $\psi^3$  is the third stable Adams operation—the 4-connective cover  $\tau_{\geq 4} ko$  can be seen as making up for the fact that the image of the  $J$ -homomorphism and the  $\mu$ -family agree in degree  $k = 1$ . The homotopy groups of  $ko$  are well-known by Bott periodicity, as is the action of the Adams operations  $\psi^k$  on  $\pi_* ko$ , so the homotopy groups of  $j$  are readily calculated from the above fibre sequence. The classical detection theorem for  $j$  (Th.A) then states that all of these homotopy groups detect elements in  $\pi_* \mathbf{S}$ , which can then be shown to be the image of the  $J$ -homomorphism and the  $\mu$ -family; in degrees  $k \geq 3$ , these are the red and blue classes, respectively, found in either Fig.4 or Fig.6.

We do not know where or when Th.A was originally stated—it is mentioned in [Koc90, Th.4.2.1]. One method to prove this detection statement for  $j$  would be to calculate its ASS or ANSS, but this proves to be rather subtle. Indeed,  $H^*(j, \mathbf{F}_2)$  itself is only free as an  $\mathcal{A} // \mathcal{A}(3)$ -module, and this calculation is then fed into an Ext-calculator to find the  $E_2$ -page, and finally there are differentials of arbitrary length left to calculate. The  $E_2$ -page of the ASS for  $j$  was computed by Davis [Dav75] and Bruner–Rognes [BR22] recently determined the rest of this spectral sequence. We cannot find any account of an ANSS for  $j$  in the literature.

## Synthetic techniques

The aim of this article is to show how *synthetic spectra* can be used to produce a simple proof of Th.A. The  $\infty$ -category of synthetic spectra  $\text{Syn}_E$  is a categorification of the  $E$ -based Adams spectral sequence ( $E$ -ASS) for some Adams-type homology theory  $E$ . Each synthetic spectrum has an associated spectral sequence, which we call its *signature*. In particular, for each spectrum  $X$  there is a synthetic spectrum  $\nu X$  whose *signature* is the standard  $E$ -ASS for  $X$ . There can be, however, many other synthetic spectra  $Y$  not of the form  $\nu X$  whose signature still abuts to the homotopy groups of  $X$ —we call such a spectral sequence a *modified  $E$ -ASS* for  $X$ ; see Df.1.5.

In this article, we construct synthetic spectra  $j_E$  for  $E = \mathbf{F}_p$  and BP and for every prime  $p$ , whose associated spectral sequences are almost as easy to calculate as the homotopy groups of  $j$  itself—this is in stark contrast to the classical ASS or ANSS for  $j$ . With minimal effort we can calculate the modified ASSs and ANSSs associated with  $j_E$  without ever referring to the  $\mathbf{F}_p$ - or BP-homology of  $j$  nor any Ext-calculations. As an additional curiosity, these synthetic methods also recover the ASS and ANSS for  $ko$  without calculating its  $\mathbf{F}_p$ - or BP-homology. As evidence for the usefulness of these modified spectral sequences and the relevance of  $j_E$ , we provide a much-simplified proof of Th.A as well as demonstrate a detection statement about the synthetic unit map  $\mathbf{1} \rightarrow j_{\text{BP}}$ .

**Theorem B.** *For odd primes  $p$ , the unit map  $\mathbf{1} \rightarrow j_{\text{BP}}$  induces a split surjection on synthetic homotopy groups, and at  $p = 2$ , it induces a split surjection on  $E_\infty$ -pages of the associated spectral sequences.*

Although it is not true that  $\mathbf{1} \rightarrow j_{\text{BP}}$  induces a surjection on bigraded homotopy groups at the prime 2, see Rmk.6.9, the modified ANSS associated with  $j_{\text{BP}}$  does allow us to strengthen the classical statement that  $\pi_* \mathbf{S} \rightarrow \pi_* j$  is a split surjection of abelian groups.

**Theorem C.** *Fix  $p = 2$ . Consider  $\pi_*\mathbf{S}$  and  $\pi_*\mathbf{j}$  as filtered abelian groups using the (modified) ANSSs associated with the BP-synthetic spectra  $\mathbf{1}$  and  $\mathbf{j}_{\text{BP}}$ . Then the map  $\pi_*\mathbf{S} \rightarrow \pi_*\mathbf{j}$  is a split surjection of filtered abelian groups.*

Again, we would like to emphasise that the computational input for these calculations is only slightly more involved than the usual computation of  $\pi_*\mathbf{j}$  using the homotopy groups of  $\mathbf{ko}$  and the action of the stable Adams operations  $\psi^k$ .

The framework of synthetic spectra allows us to also construct modified ASSs and ANSSs for the periodic spectra  $\mathbf{KO}$  and  $\mathbf{J}$ , and we again compute these spectral sequences in their entirety. This is particularly interesting for the ASS, as the classical ASSs for all of the periodic spectra  $\mathbf{KU}$ ,  $\mathbf{KO}$ , and  $\mathbf{J}$  are trivial (and, in particular, do not converge); a consequence of the fact that the  $\mathbf{F}_p$ -homology of these spectra is zero. This modified ASS for  $\mathbf{J}$  behaves like a nontrivial ASS converging to the  $\mathbf{K}(1)$ -local sphere; see §3 and §4.4.

Our construction and study of  $\mathbf{j}_E$  is done using the usual techniques of synthetic spectra together with a new  $t$ -structure on  $\text{Syn}_E$ . In a little more detail, we construct *linear  $t$ -structures* on the stable  $\infty$ -category  $\text{Syn}_E$  depending on a line  $L$  and calculate their hearts. To simplify our exposition, we will restrict ourselves to the vertical line defined by the equation  $x = 0$  in the following statement; see Th.2.2 for a more general statement.

**Theorem D.** *Let  $E$  be either  $\text{BP}$  or  $\mathbf{F}_p$ . There exists a left and right complete monoidal  $t$ -structure on  $\text{Syn}_E$ , called the vertical  $t$ -structure, whose connective objects are given by those synthetic spectra  $X$  with  $\pi_{a,b}X = 0$  for  $a < 0$ . Moreover, the heart of this  $t$ -structure is given by the abelian category of graded  $\pi_{0,*}\mathbf{1}$ -modules.*

We then define  $\mathbf{j}_E$  as the fibre of  $\psi^3 - 1: \nu\mathbf{ko} \rightarrow \tau_{\geq 4}^{\uparrow}\nu\mathbf{ko}$ , where the 4-connective cover is taken with respect to the vertical  $t$ -structure on  $\text{Syn}_E$ ; see §4. The synthetic homotopy groups of  $\mathbf{j}_E$  and the associated modified  $E$ -ASSs are then readily computed from the long exact sequence of the above fibre sequence; see §A for many spectral sequence diagrams demonstrating this.

## Outline

This article is divided into three main sections: in §1 and 2, we provide some clarifications and refinements to concepts concerning the  $\infty$ -category  $\text{Syn}$ , then in §3 and 4, we construct various lifts of topological  $K$ -theories and image-of- $J$ -spectra in  $\text{Syn}$ , and finally in §6, we prove detection results using these synthetic constructions. Experts on synthetic spectra may comfortably skip §1, and experts on topological  $K$ -theory and the Hurwicz image of  $\mathbf{ko}$  may skip §3 and 5—these latter two sections serve to provide motivation and notation for our study of  $\mathbf{j}$ . In some more detail:

- In §1, the connection between synthetic spectra, filtered spectra, and their associated spectral sequences is discussed. In particular, this provides us with a language to discuss *modified  $E$ -ASSs*, reproducing  $v_n$ -localised ASSs as an example.

- In §2, we construct linear  $t$ -structures on synthetic spectra and prove generalisations of Th.D for a variety of lines  $L$  through the origin. The heart of these  $t$ -structures is calculated as well as some basic properties of the associated truncation and connective cover functors.
- In §3, various  $E$ -synthetic lifts of topological  $K$ -theory are constructed, first for  $E = \mathbf{F}_p$  and then for  $E = \mathbf{BP}$ . It is shown how the classical ASSs and ANSSs can be recovered from a version of Wood's theorem internal to  $\mathbf{Syn}$ , using only simple long exact sequence arguments. These constructions are a warm-up to our synthetic construction of various image-of- $J$ -spectra.
- In §4, the main characters of this article are introduced: the synthetic lifts for  $\mathbf{j}$ . Similar to the classical calculation of the homotopy groups  $\mathbf{j}$ , the synthetic homotopy groups of these lifts are then easily computed simultaneously giving us modified ASSs and ANSSs for  $\mathbf{j}$ . We also show that each  $\mathbf{j}_E$  has a chosen  $\mathbf{E}_\infty$ -structure in  $\mathbf{Syn}$  and produce periodic variants.
- In §5, we reprove the classical statements of Adams concerning the existence of  $v_1$  self-maps on mod  $p$  Moore spectra and how these can be used to prove detection results concerning real topological  $K$ -theory. These arguments are a necessary step towards the detection results concerning image-of- $J$ -spectra.
- In §6, we prove Th.A, stating the surjectivity of the unit map  $\mathbf{S} \rightarrow \mathbf{j}$  on homotopy groups, using the calculations from §4 together with the detection arguments for  $\mathbf{ko}$  from §5. We also prove two more refined statements: Th.B, which states that the synthetic unit map  $\mathbf{1} \rightarrow \mathbf{j}_{\mathbf{BP}}$  induces a **split** surjection on synthetic homotopy groups at odd primes, and Th.C, which states that the surjection of Th.A refines to a split surjection of filtered abelian groups at the prime 2.
- In §A, we provide many diagrams of modified  $E$ -ASS used throughout this article.

## Future work

Our proof of Th.A does not use any homology calculations of  $\mathbf{j}$ , however, one could still take a more computational approach to prove this theorem using direct ASS calculations. There are other situations though, where these direction ASS computations become unmanageable. The authors' initial motivation for these synthetic constructions was to try to calculate the image of the unit map from  $\mathbf{S}$  into the fibre of a map  $F - \lambda: \mathbf{tmf} \rightarrow \tau_{\geq 8} \mathbf{tmf}$ , where  $F$  is an Adams operation  $\psi^k$  or a Hecke operator  $T_n$  and  $\lambda$  is the eigenvalue of the unit. Computing directly with the ASS for  $\mathbf{tmf}$  is very difficult, as is determining the homology of  $\tau_{\geq 8} \mathbf{tmf}$ , let alone its ASS. These computations can hopefully lead to more computations of connective variants of Behrens'  $\mathbf{Q}(N)$ -spectra. We will return to the question above regarding the Hurewicz image of such spectra in future work, where we can immediately apply the techniques of this article.

## Notation

Through this article,  $p$  will be a fixed prime and our spectra are implicitly  $p$ -complete. If  $E$  is an Adams-type homology theory, we will write  $E$ -ASS for the  $E$ -based Adams spectral sequence. In particular,  $\mathbf{F}_p$ -ASS will be written as ASS, and BP-ASS as ANSS. We will denote the category of  $E$ -synthetic spectra  $\text{Syn}_E$  of [Pst23] by  $\text{Syn}$ . We use the “stem–filtration” grading for synthetic spectra, meaning that  $\Sigma^{a,b}\mathbf{1} = \Sigma^{-b}\nu\mathbf{S}^{a+b}$ .

## Logical assumptions

The basic inputs surrounding topological  $K$ -theory and the spectrum  $j$  we are taking for granted in this article are the following:

- The  $\mathbf{E}_\infty$ -ring  $ku$  exists, whose homotopy groups are given by the ring

$$\pi_* ku \simeq \mathbf{Z}[u] \quad |u| = 2.$$

As we will be working in  $p$ -complete spectra for each prime  $p$ , we can take  $ku = \tau_{\geq 0} \text{KU}$ , where  $\text{KU}$  is the Morava  $E$ -theory of the formal multiplicative group  $\mathbf{G}_m$  over  $\mathbf{F}_p$ ; see [Lur18, §5].

- By [Lur18, Rmk.5.0.6], there is an action of the automorphisms of  $\widehat{\mathbf{G}}_m$  over  $\mathbf{F}_p$  on  $\text{KU}$ . This group of automorphisms is given by  $\mathbf{Z}_p^\times$ , and for each  $k \in \mathbf{Z}_p^\times$ , we denote the action of  $k$  on  $\text{KU}$  by the automorphism of  $\mathbf{E}_\infty$ -rings  $\psi^k$ ; we will also use this notation for the action of  $\mathbf{Z}_p^\times$  on  $ku$ . By [Dav20, Prs.5.2.8 & 5.2.10], the map  $\psi^k$  induces multiplication by  $k^d$  on  $\pi_{2d} ku$  and deserves to be called a *stable Adams operation*.
- Writing  $F$  for the maximal finite subgroup of  $\mathbf{Z}_p^\times$  at odd primes, then we define the  $\mathbf{E}_\infty$ -ring  $L = \text{KU}^{hF}$ , called the *periodic Adams summand*, as well as its connective cover  $\ell = \tau_{\geq 0} L$ .

Using these inputs, we define the  $\mathbf{E}_\infty$ -ring  $ko = \tau_{\geq 0}(ku^{hC_2})$ .

To reiterate: we are not presupposing any calculations about the  $E$ -homology of  $ko$  or  $j$ , for  $E = \mathbf{F}_p$  or BP.

## Acknowledgements

We would like to thank Shaul Barkan, Lennart Meier, Sven van Nigtevecht, and Lucas Piesseaux for their helpful conversations and suggestions on various drafts.

## 1 The signature of a synthetic spectrum

Associated with a synthetic spectrum  $X$  are two spectral sequences: the  $\tau$ -Bockstein spectral sequence (BSS) and what we call the  $\Gamma$ -spectral sequence ( $\Gamma$ -SS). These concepts will be used

to concretely define what we mean by *modified E-ASSs*. Recall that we are using the “stem-filtration” grading for synthetic spectra, meaning that  $\Sigma^{a,b}\mathbf{S} = \Sigma^{-b}\nu\mathbf{S}^{a+b}$ . In particular,  $\tau$  has degree  $(0, -1)$  and suspension as a stable  $\infty$ -category has degree  $(1, -1)$ .

This section uses filtered spectra as a homotopical model for spectral sequences as done in [Lur17, §1.2.2] and [Hed20, §II.1], for example.

## 1.1 Filtered spectra and spectral sequences

Spectral sequences come from filtrations. In stable homotopy theory, this is encoded as saying that spectral sequences come from filtered spectra. A filtered spectrum is an object in  $\text{Fil}(\text{Sp}) := \text{Fun}(\mathbf{Z}^{\text{op}}, \text{Sp})$ , the  $\infty$ -category of filtered spectra, where  $\mathbf{Z}$  is the poset consisting of the integers with the usual ordering. Thus a filtered spectrum consists of a tower  $X^\bullet$  of spectra

$$\dots \rightarrow X^{n+1} \rightarrow X^n \rightarrow X^{n-1} \rightarrow \dots$$

where  $X^i \in \text{Sp}$ . One extracts a spectral sequence from  $X^\bullet$  by forming the exact couple

$$\begin{array}{ccc} \bigoplus_{s,t} \pi_{t-s} X^{t+1} & \xrightarrow{\quad\quad\quad} & \bigoplus_{s,t} \pi_{t-s} X^t \\ & \swarrow \text{---} & \searrow \text{---} \\ & \bigoplus_{s,t} \pi_{t-s} X_t^t & \end{array}$$

giving a spectral sequence with signature

$$E_2^{t-s,s} = \pi_{t-s} X_t^t \implies \pi_{t-s} X^{-\infty}.$$

Here we write  $X_t^t := \text{cofib}(X^{t+1} \rightarrow X^t)$  and  $X^{-\infty} = \text{colim } X^n$ , and the (descending) filtration on  $\pi_{t-s} X^{-\infty}$  is defined by

$$F^s \pi_{t-s} X^{-\infty} = \text{image}(\pi_{t-s} X^t \rightarrow \pi_{t-s} X^{-\infty})$$

for all  $t \in \mathbf{Z}$ . If  $\lim X^n = X^\infty = 0$ , then the spectral sequences converge in the sense that the  $F^s$  filtration is exhaustive and Hausdorff, and that  $E_\infty \simeq F^s/F^{s+1}$ ; see [Boa99, §5].

If one knows the homotopy groups  $\pi_* X^n$  and the effect of the maps  $\pi_* X^n \rightarrow \pi_* X^{n-1}$  for every  $n$ , then one knows the  $E_2$  page and the behaviour of all the differentials via the exact couple. It is helpful then to package the homotopy groups  $\pi_* X^n$  as  $n$  varies into one bigraded group.

**Definition 1.1.** For  $X^\bullet \in \text{Fil}(\text{Sp})$ , we define the bigraded homotopy group  $\pi_{n,s} X^\bullet = \pi_n X^{n+s}$ .

*Remark 1.2.* This indexing convention does not agree with many sources on bigraded homotopy groups. We have chosen them so that elements in  $\pi_{n,s} X^\bullet$  correspond to classes in  $\pi_n X^{-\infty}$  detected in filtration  $s$  of the spectral sequence associated with  $X^\bullet$ .

There are bigraded spheres in  $\text{Fil}(\text{Sp})$  that corepresent these bigraded homotopy groups; one simply sets

$$\mathbf{S}^{n,s}(k) := \begin{cases} \mathbf{S}^n & k \leq n + s \\ 0 & k > n + s \end{cases}$$

where all the transition maps are either zero or the identity. The transition maps in a filtered spectrum  $\pi_* X^n \rightarrow \pi_* X^{n-1}$  give  $\pi_{*,*} X^\bullet$  the structure of a bigraded module over the bigraded ring  $\mathbf{Z}[\tau]$  where  $|\tau| = (0, -1)$ . This is encoded in the category  $\text{Fil}(\text{Sp})$  by the bigraded self-map  $\tau : \mathbf{S}^{0,-1} \rightarrow \mathbf{S}^{0,0}$  of the unit  $\mathbf{1} = \mathbf{S}^{0,0}$ , defined in the obvious way. Since multiplication by  $\tau$  in  $\pi_{*,*} X^\bullet$  corresponds to the transition maps in the exact couple above, we see that the  $\mathbf{Z}[\tau]$ -module  $\pi_{*,*} X^\bullet$  encodes the data of the spectral sequence associated with  $X^\bullet$ . This sentiment is captured in the following statement, which is proven by the standard arguments for exact couples; see [BHS19, Th.A.1], for example.

**Theorem 1.3.** *We have the following facts about the relationship between the spectral sequence associated with  $X^\bullet \in \text{Fil}(\text{Sp})$  and the  $\mathbf{Z}[\tau]$ -module  $\pi_{*,*} X^\bullet$ .*

1. *There is a short exact sequence*

$$0 \rightarrow (\pi_{n,s} X^\bullet)/\tau \rightarrow E_2^{n,s} \xrightarrow{\partial} (\pi_{n-1,s+2} X^\bullet)[\tau] \rightarrow 0$$

where  $(\pi_{n-1,s+2} X^\bullet)[\tau]$  is the kernel of multiplication by  $\tau$  on  $\pi_{n-1,s+2} X^\bullet$ .

2. *The subgroup  $(\pi_{n,s} X^\bullet)/\tau \simeq Z_\infty^{n,s}$  of  $E_2^{n,s}$  consists of permanent cycles in stem  $n$  and filtration  $s$ .*
3. *The subgroup  $B_r^{n,s} \subset Z_\infty^{n,s}$  of classes that are hit by a  $d_r$ -differential is precisely the set of  $x \in Z_\infty^{n,s}$  admitting a lift  $\tilde{x} \in \pi_{n,s} X^\bullet$  such that  $\tau^{r-1} x = 0$ .*
4. *Suppose the map  $\partial : E_2^{n,s} \rightarrow (\pi_{n-1,s+2} X^\bullet)[\tau]$  sends  $x \mapsto \tau^{r-2} y \neq 0$  where  $y$  is not divisible by  $\tau$ . Then  $d_2, \dots, d_{r-1}$  vanish on  $x$  and  $d_r(x)$  is represented by the image of  $y$  along the map*

$$\pi_{n-1,s+r} X^\bullet \twoheadrightarrow (\pi_{n-1,s+r} X^\bullet)/\tau \hookrightarrow E_2^{n-1,s+r}$$

Conversely, if  $d_2, \dots, d_{r-1}$  vanish on  $x$  and  $d_r(x) = y \neq 0$ , then  $y$  admits a lift to  $E_2$  that admits a lift  $\tilde{y}$  to  $\pi_{n-1,s+r} X$  such that  $\partial(x) = \tau^{r-2} \tilde{y}$ .

## 1.2 Synthetic spectra and modified Adams spectral sequences

Let  $E$  be an Adams-type spectrum, then for any  $X \in \text{Sp}$ , one has the  $E$ -ASS of  $X$

$$\text{Ext}_{E_* E}(E_*, E_* X) \implies \pi_*(X_E^\wedge)$$

converging to the  $E$ -nilpotent completion of  $X$ . For some applications, this particular spectral sequence is ineffective and could be replaced by a variety of other spectral sequences with similar formal behaviour. These are called *modified  $E$ -ASSs* for  $X$ . Such spectral sequences

have a rich history; for example, one often considers localised ASSs, such as the  $v_n$ -localised ASS

$$v_n^{-1} \operatorname{Ext}_{\mathcal{A}_*}(\mathbf{F}_p, H_*(Y)) \implies \pi_*(v_n^{-1}Y)$$

for a spectrum  $Y$  with a  $v_n$  self-map; for example, see [AKQ20, §2.2]. This spectral sequence converges under certain conditions on  $Y$ , but it is not the same as the  $\mathbf{F}_p$ -ASS of  $v_n^{-1}Y$ , as  $H_*(v_n^{-1}Y) = 0$ . Another common example of a modified ASS is the one used to compute the  $X$  homology of a Moore spectrum  $\mathbf{S}/(p^{i_0}, v_1^{i_1}, \dots, v_n^{i_n})$  for some spectrum  $X$ , where one has a spectral sequence

$$\operatorname{Ext}_{\mathcal{A}_*}(\mathbf{F}_p, H(i_0, \dots, i_n) \otimes H_*(X; \mathbf{F}_p)) \implies X_*(\mathbf{S}/(p^{i_0}, v_1^{i_1}, \dots, v_n^{i_n}))$$

where  $H(i_0, \dots, i_n)$  is a Moore object  $\mathbf{1}/(q_0^{i_0}, \dots, q_n^{i_n})$  formed inside the derived category  $\mathcal{D}(\operatorname{Comod}(\mathcal{A}_*))$ ; for example, see [BHHM08, §8]. This  $E_2$ -page is much easier to work with, but is different to the  $E_2$ -page of the  $\mathbf{F}_p$ -ASS of  $X \otimes \mathbf{S}/(p^{i_0}, v_1^{i_1}, \dots, v_n^{i_n})$  if any  $i_j > 1$ .

Classically, the construction of these modified ASSs is quite complicated and *ad hoc*. The category  $\operatorname{Syn}$  of  $E$ -based synthetic spectra makes it easy to construct such spectral sequences and we also find it is convenient to think about objects in  $\operatorname{Syn}$  in this way. Very roughly speaking, objects in  $\operatorname{Syn}$  are pairs  $(X, \mathcal{E})$ , where  $X$  is a spectrum, and  $\mathcal{E}$  is a choice of modified  $E$ -based Adams spectral sequence for  $X$ . This point of view can be made into a rigorous description of  $\operatorname{Syn}$  (up to technical issues in  $\operatorname{Syn}$  like cellularity and  $\tau$ -completeness of the unit); see the *filtered model* of [BHS22, Pr.C.22]. More categorically,  $\operatorname{Syn}$  can be considered as a bundle of spectral sequences living over  $\operatorname{Sp}$

$$\begin{array}{c} \operatorname{Syn} \\ \tau^{-1} \downarrow \overset{\lambda}{\curvearrowright} \nu \\ \operatorname{Sp} \end{array}$$

where the synthetic analog functor  $\nu$  gives a canonical section of this fibration, sending  $X \mapsto (X, E\text{-ASS}(X))$  where  $E\text{-ASS}(X)$  is the standard  $E$ -ASS of  $X$ . From this perspective, the fibration  $\tau^{-1}: \operatorname{Syn} \rightarrow \operatorname{Sp}$  is projection onto the first factor, and we devote the rest of the section to describing projection onto the second factor, namely how to extract a modified  $E$ -ASS for  $\tau^{-1}X$  from a synthetic spectrum  $X$ . We refer the reader again to [BHS22, Appendix C] for a more categorical point of view on this construction.

The unit in  $\operatorname{Syn}$  has a bigraded self-map  $\tau: \Sigma^{0,-1}\mathbf{1} \rightarrow \mathbf{1}$ , so for any  $X \in \operatorname{Syn}$ , one has a filtered synthetic spectrum

$$X^\bullet = \dots \rightarrow \Sigma^{0,-1}X \xrightarrow{\tau} X \xrightarrow{\tau} \Sigma^{0,1}X \rightarrow \dots$$

where  $X^0 = X$ . Since  $\operatorname{Syn}$  is a stable  $\infty$ -category, we may apply the mapping spectrum functor  $\operatorname{map}_{\operatorname{Syn}}(\mathbf{1}, -)$  to this filtered synthetic spectrum to get a filtered spectrum we will call  $\Gamma(X)$ .

**Theorem 1.4.** *The functor  $\Gamma: \operatorname{Syn} \rightarrow \operatorname{Fil}(\operatorname{Sp})$  has the following properties*

1. For any  $X \in \text{Syn}$ , the map  $\pi_{*,*}X \rightarrow \pi_{*,*}\Gamma(X)$  is an isomorphism of  $\mathbf{Z}[\tau]$ -modules.
2. The spectral sequence associated with the filtered spectrum  $\Gamma(X)$  has signature

$$E_2 = \pi_{t-s,s}(X/\tau) \implies \pi_{t-s}\tau^{-1}X.$$

This is the  $\Gamma$ -SS for  $X$ .

3. If  $X$  is  $\tau$ -complete, the spectral sequence converges in the sense that the corresponding filtration on  $\pi_*\tau^{-1}X$  is exhaustive and Hausdorff.
4. Considering  $\text{Fil}(\text{Sp})$  as a symmetric monoidal  $\infty$ -category with the Day convolution monoidal structure, then  $\Gamma$  has the canonical structure of a lax symmetric monoidal functor.

*Proof.* The second and third claims follow from the first by definition of  $\Gamma$  along with the discussion in the previous section. For the first claim, one has

$$\begin{aligned} \pi_{n,s}\Gamma(X) &= \pi_n\Gamma(X)^{n+s} \\ &= \pi_n\text{map}_{\text{Syn}}(\mathbf{1}, \Sigma^{0,-n-s}X) \\ &= [\mathbf{S}^{n,-n}, \Sigma^{0,-n-s}X]_{\text{Syn}} \\ &= \pi_{n,s}X \end{aligned}$$

using that the stable  $\infty$ -categorical suspension functor on  $\text{Syn}$  coincides with the bigraded suspension  $\Sigma^{1,-1}$ . The lax monoidality of  $\Gamma$  comes from the natural maps

$$\begin{aligned} (\Gamma X \otimes \Gamma Y)^n &\simeq \bigoplus_{i+j=n} \text{map}_{\text{Syn}}(\mathbf{1}, \Sigma^{0,-i}X) \otimes \text{map}_{\text{Syn}}(\mathbf{1}, \Sigma^{0,-j}Y) \\ &\rightarrow \bigoplus_{i+j=n} \text{map}_{\text{Syn}}(\mathbf{1}, \Sigma^{0,-i-j}X \otimes Y) \rightarrow \Gamma(X \otimes Y)^n \end{aligned}$$

and the unit map for  $n \geq 0$

$$\mathbf{S}^{0,0}(n) = \mathbf{S} \rightarrow \text{map}_{\text{Syn}}(\Sigma^{0,n}\mathbf{1}, \mathbf{1}) \simeq \Gamma(\mathbf{1})^n. \quad \square$$

**Definition 1.5.** Given a synthetic spectrum  $X$ , we say the *signature* of  $X$  is the spectral sequence associated with the filtered spectrum  $\Gamma(X)$  converging to  $\tau^{-1}X$ . In particular, given a filtered spectrum  $\mathcal{E}$  whose associated spectral sequence converges to the homotopy groups of a spectrum  $Y$ , as well as an equivalence of filtered spectra  $\mathcal{E} \simeq \Gamma(X)$  and an equivalence of spectra  $\tau^{-1}X \simeq Y$ , we say that  $X$  has signature  $\mathcal{E}$ . We will also call the  $\Gamma$ -SS of  $X$  a *modified  $E$ -ASS* for  $Y$  (from  $X$ ).

*Example 1.6.* When  $Y = \nu X$ , then [BHS19, Th.A.1] shows that the signature of  $Y$  is the standard  $E$ -ASS of  $X$ —this is also implicit in [Pst23]. Combining Ths.1.3 and 1.4, we now have a precise meaning to the sense that the bigraded synthetic homotopy groups of  $\nu X$  encode the data of the  $E$ -ASS of  $X$ . Another perspective on this fact will be provided in [CDN24], where the authors together with Sven van Nigtevecht show that there is a natural equivalence of filtered spectra  $\Gamma(\nu X)$  and (the décalage of) the  $E$ -ASS for  $X$ . In [CDN24], we will also construct various synthetic spectra whose signatures are *descent spectral sequences* associated to a global sections of some even periodic spectral Deligne–Mumford stack.

*Example 1.7.* Suppose  $Y$  is a spectrum with a  $v_n$  self-map  $v$ . Then using [BHS19, Th.A.1], one can show that there is a bigraded self-map  $\tilde{v}$  of  $\nu(Y) \in \text{Syn}_{\mathbf{F}_p}$  such that  $\tau^{-1}(\tilde{v}) = v$  and reduction mod  $\tau$  sends  $\tilde{v}$  to the class in  $\text{Ext}_{\mathcal{A}_*}(\mathbf{F}_p, H_*Y)$  detecting  $v$ . It follows that the telescope  $\tilde{v}^{-1}Y$  formed in  $\text{Syn}_{\mathbf{F}_p}$  has the property that

$$\pi_{*,*}(\tilde{v}^{-1}Y/\tau) = v_n^{-1} \text{Ext}_{\mathcal{A}_*}(\mathbf{F}_p, H_*(Y))$$

so that the signature of  $\tilde{v}^{-1}Y$  is the  $v_n$ -localised  $\mathbf{F}_p$ -ASS of  $Y$ .

*Example 1.8.* Let  $X \in \text{Sp}$ . Set  $Y_{-1} = \mathbf{1} \in \text{Syn}_{\mathbf{F}_p}$  and assume by induction that we have defined  $Y_{j-1} =: \mathbf{1}/(q_0^{i_0}, \dots, q_{j-1}^{i_{j-1}}) \in \text{Syn}_{\mathbf{F}_p}$  so that  $\tau^{-1}Y_{j-1} = \mathbf{S}/(p^{i_0}, \dots, v_{j-1}^{i_{j-1}})$ . Since everything is implicitly  $p$ -complete,  $\mathbf{1} \in \text{Syn}_{\mathbf{F}_p}$  is  $\tau$ -complete, hence so is  $\text{End}(Y_{j-1})$ , and thus the corresponding  $\Gamma$ -SS

$$\pi_{*,*}(\text{End}(Y_{j-1})/\tau) \implies \pi_* \text{End}(\mathbf{S}/(p^{i_0}, \dots, v_{j-1}^{i_{j-1}}))$$

converges. The self-map  $v_j^{i_j} \in \pi_* \text{End}(\mathbf{S}/(p^{i_0}, \dots, v_{j-1}^{i_{j-1}}))$  is therefore detected by an endomorphism  $q_j^{i_j}$  of  $Y_{j-1}/\tau$ . Since  $q_j^{i_j}$  is a permanent cycle in the  $\Gamma$ -SS of  $\text{End}(Y_{j-1})$ , it is also a permanent cycle in the  $\tau$ -BSS of  $\text{End}(Y_{j-1})$ , and it therefore admits a lift to a self-map of  $Y_{j-1}$  that  $\tau$ -inverts to  $v_j^{i_j}$ .

Denoting this self-map of  $Y_{j-1}$  also by  $q_j^{i_j}$ , we set  $Y_j = Y_{j-1}/q_j^{i_j}$ . It follows that the synthetic spectrum

$$\mathbf{1}/(q_0^{i_0}, \dots, q_n^{i_n}) := Y_n$$

has the property that

$$\pi_{*,*}(\nu X \otimes \mathbf{1}/(q_0^{i_0}, \dots, q_n^{i_n}, \tau)) = \text{Ext}_{\mathcal{A}_*}(\mathbf{F}_p, H(i_0, \dots, i_n) \otimes H_*(X; \mathbf{F}_p))$$

so that the signature of  $\nu X \otimes \mathbf{1}/(q_0^{i_0}, \dots, q_n^{i_n})$  is the modified  $\mathbf{F}_p$ -ASS of  $X \otimes \mathbf{1}/(p^{i_0}, \dots, v_n^{i_n})$ .

## 2 Linear $t$ -structures on synthetic spectra

Let  $E$  be a spectrum of Adams-type, such that  $\text{Syn}_E = \text{Syn}$  is defined. In this section, we construct  $t$ -structures on  $\text{Syn}$  coming from lines in  $\mathbf{R}^2$  through the origin. When restricted to the line defined by the equation  $x = 0$ , we call the resulting  $t$ -structure the *vertical  $t$ -structure*.

### 2.1 Construction of the linear $t$ -structure

**Definition 2.1.** Let  $L$  be a line in  $\mathbf{R}^2$  through the origin not containing  $(1, -1)$ . Write  $\mathbf{R}^2 - L = L_{>} \cup L_{<}$  as the union of two connected subspaces where the region containing  $(1, -1)$  is  $L_{>}$ . Let us also write  $L_{\geq} = L \cup L_{>}$  and  $L_{\leq} = L \cup L_{<}$ . For any  $n \in \mathbf{Z}$ , define the  $\infty$ -subcategory  $\text{Syn}_{\geq n}^L$  of  $\text{Syn}$  to be spanned by those synthetic spectra  $X$  such that  $\pi_{a+n, b-n}X = \pi_{a, b}X[-n] = 0$  if  $(a, b) \in L_{<}$ , so such that  $X[-n]$  has homotopy groups supported in  $L_{\geq}$ . Likewise, define  $\text{Syn}_{\leq n}^L$  as the  $\infty$ -subcategory of  $\text{Syn}$  spanned by synthetic spectra  $X$  with  $\pi_{a-n, b+n}X = 0$  for  $(a, b) \in L_{>}$ .<sup>1</sup>

<sup>1</sup>If  $L$  is defined by  $y = -x$ , then there are two possible choices for  $\text{Syn}^{\leq L}$ , and both could potentially produce  $t$ -structures on  $\text{Syn}$ —we will not need either of them.

**Theorem 2.2.** *Let  $L$  be a line in  $\mathbf{R}^2$  through the origin not containing  $(1, -1)$ ,  $E$  be a spectrum such that  $\mathrm{Syn}$  is cellular, and suppose that the unit  $\mathbf{1} \in \mathrm{Syn}$  lies in  $\mathrm{Syn}_{\geq 0}^L$ . Then the  $\infty$ -subcategories  $\mathrm{Syn}_{\geq}^L$  and  $\mathrm{Syn}_{\leq}^L$  determine an accessible  $t$ -structure  $\mathrm{Syn}$  which is both right and left complete and compatible with the monoidal structure on  $\mathrm{Syn}$ .*

Note that Pstragowski's homological  $t$ -structure on  $\mathrm{Syn}$  of [Pst23, §4.2] is not an instance of a linear  $t$ -structure, but there is a connection: by [Pst23, Cor.4.19], we see that a synthetic spectrum  $X$  is connective in Pstragowski's  $t$ -structure if and only if  $\nu E \otimes X$  lies in  $\mathrm{Syn}_{\geq 0}^L$  for the line  $L$  defined by  $y = 0$ .<sup>2</sup>

*Proof.* Consider the collection of objects  $S = \{\mathbf{S}^{a,b}\}$  where  $(a, b) \in L_{\geq}$ . Define  $C_{\geq}$  as the full  $\infty$ -subcategory of  $\mathrm{Syn}$  generated by the objects in  $S$  by small colimits and extensions. By [Lur17, Pr.1.4.4.11(2)],  $C_{\geq}$  is a presentable  $\infty$ -category, as  $\mathrm{Syn}$  itself is presentable and stable [Pst23, Pr.4.2], and [Lur17, Pr.1.4.4.11(1)] then states that  $\mathrm{Syn}$  has an accessible  $t$ -structure with with connective objects  $C_{\geq}$ . Notice that by definition  $\mathrm{Syn}_{\geq 0}^L$  contains the spheres in  $S$ . Moreover, it is clearly closed under extensions as this subcategory is defined by a vanishing condition on homotopy groups. Likewise,  $\mathrm{Syn}_{\leq 0}^L$  is also closed under colimits, as coproducts and filtered colimits commute with homotopy groups and for pushouts we use the associated Mayer–Vietoris sequence on homotopy groups. It follows that  $C_{\geq} \subseteq \mathrm{Syn}_{\geq 0}^L$ . The fact that  $\mathrm{Syn}$  is cellular implies this inclusion is an equivalence of subcategories of  $\mathrm{Syn}$ .

To see this  $t$ -structure is left and right complete, if  $\pi_{a,b}X = 0$  for all  $(a, b)$ , then the synthetic spectrum  $X = 0$  from our cellularity hypothesis. In particular, we see that both of the  $\infty$ -subcategories  $\bigcap \mathrm{Syn}_{\leq -n}^L$  and  $\bigcap \mathrm{Syn}_{\geq n}^L$  contain only the zero synthetic spectrum. Just as in the proof of completeness in [Lur17, Pr.1.4.3.6], referring to [Lur17, Pr.1.2.1.19] and its dual, it now suffices to show that  $\mathrm{Syn}_{\leq 0}^L$  and  $\mathrm{Syn}_{\geq 0}^L$  are closed under products and coproducts. This is clear though, as these categories are defined by a condition on synthetic homotopy groups, which commute with products, as well as finite coproducts and filtered colimits.

Finally, we want to check compatibility with the monoidal structure on  $\mathrm{Syn}$ . The unit lies in  $\mathrm{Syn}_{\geq 0}^L$  by assumption, so it suffices to see that the objects of  $S$  are closed under the tensor product of synthetic spectra. This is clear, as  $\mathbf{S}^{a,b} \otimes \mathbf{S}^{c,d} \simeq \mathbf{S}^{a+c,b+d}$ , and if  $(a, b), (c, d) \in L_{\geq}$ , then also  $(a + c, b + d) \in L_{\geq}$ .  $\square$

The condition about the unit  $\mathbf{1}$  above depends on both  $L$  and  $E$ . For example, when  $E = \mathrm{BP}$  or  $\mathbf{F}_p$  and  $L$  is defined by  $x = 0$ , which we call the vertical line, then this unit condition is satisfied as the  $E$ -ASS for the  $E$ -local sphere are concentrated in nonnegative stems. A more general sufficient condition is that the  $E_2$ -page of the  $E$ -ASS is concentrated in nonnegative stems. For other  $E$  and  $L$ , this relates to deep questions concerning vanishing lines in the  $E$ -ASS for the  $E$ -local sphere. The cellularity condition holds for  $E = \mathrm{BP}$  by a variant of [Pst23, Th.6.2], which proves the  $E = \mathrm{MU}$ -case, and for  $E = \mathbf{F}_p$  by the following argument.

---

<sup>2</sup>There are not many interesting  $E$  where the line defined by  $y = 0$  produces a  $t$ -structure using Th.2.2, as this would imply the  $E$ -ASS for  $\mathbf{S}$  is concentrated in filtration 0.

**Lemma 2.3.** *The category  $\text{Syn}_{\mathbf{F}_p}$  is compactly generated by the bigraded spheres  $S^{a,b}$  for any prime  $p$ .*

*Proof.* By [Pst23, Rmk.4.14],  $\text{Syn}_{\mathbf{F}_p}$  is compactly generated by the set of bigraded suspensions of  $\nu P$  where  $P$  ranges through finite  $\mathbf{F}_p$ -projectives, so it remains to show  $\nu P$  is in the subcategory  $\mathcal{C}$  of  $\text{Syn}_{\mathbf{F}_p}$  generated by the bigraded spheres under colimits. Let  $X$  be a bounded below spectrum, shifted so that its bottom homotopy group is in degree zero. We may form the following tower

$$\begin{array}{ccccc}
 \bigoplus_{\alpha_0} \mathbf{S}^0 & & \bigoplus_{\alpha_1} \mathbf{S}^1 & & \bigoplus_{\alpha_2} \mathbf{S}^2 \\
 \downarrow q_0 & & \downarrow q_1 & & \downarrow q_2 \\
 X & \longrightarrow & X_{(0)} & \longrightarrow & X_{(1)} \longrightarrow \dots
 \end{array} \tag{2.4}$$

where  $q_i$  is an isomorphism in  $H_i(-; \mathbf{Z})$ , and  $X_{(i)}$  is the cofibre of  $q_i$ , by use of the Hurewicz theorem.

First, suppose  $H_*(X; \mathbf{F}_p) = 0$ . It follows that  $q_0$  induces the zero map in  $H_*(-; \mathbf{F}_p)$  and  $q_i$  induces an iso in  $H_*(-; \mathbf{F}_p)$  for all  $i > 0$ . Using [Pst23, Lm.4.23], the fact that  $\text{colim } X_{(i)} = 0$ , and that  $\nu$  commutes with filtered colimits, one sees then that  $\nu X \in \mathcal{C}$ . If  $P$  is any finite  $\mathbf{F}_p$ -projective, one has a localisation cofibre sequence

$$Q \rightarrow P \rightarrow P_{(p)}$$

Since  $H_*(Q; \mathbf{F}_p) = 0$ , it follows again from [Pst23, Lm.4.23] that

$$\nu Q \rightarrow \nu P \rightarrow \nu P_{(p)}$$

is a cofibre sequence in  $\text{Syn}_{\mathbf{F}_p}$ . We may therefore assume that  $P$  is a  $p$ -local finite spectrum.

Consider now a variation on (2.4), where we set  $X = P$  and instead  $q_i$  is chosen to be an isomorphism on  $H_i(-; \mathbf{F}_p)$ . The same argument now shows that  $\nu X \in \mathcal{C}$  since  $q_i$  induces a monomorphism on  $H_*(-; \mathbf{F}_p)$  for all  $i$ .  $\square$

*Remark 2.5.* Similar  $t$ -structures on filtered objects in a stable  $\infty$ -category were explored by Lee–Levy in [LL23, §2.2]. There is no moral difference between our construction and theirs—both are designed to cut out certain regions of spectral sequences. However, their construction takes place in filtered objects of a stable  $\infty$ -category, and ours takes place in synthetic spectra; we have not attempted to use the filtered model for  $\text{Syn}$  ([BHS22, Pr.C.22]) to relate our constructions. Moreover, we could consider  $t$ -structures on  $\text{Syn}$  defined with arbitrary pairs of subsets covering  $\mathbf{Z} \times \mathbf{Z}$  satisfying certain axioms, similar to how Lee–Levy study the graphs of functions  $\mathbf{Z} \rightarrow \mathbf{Z}$ . For simplicity, we will ignore these generalisations here.

**Definition 2.6.** Let  $L$  and  $E$  be as in Th.2.2 and consider the  $t$ -structure associated with this line on  $\text{Syn}$ . Write  $\tau_{\geq n}^L: \text{Syn} \rightarrow \text{Syn}_{\geq n}^L$  for the  $n$ th *connective cover* functor and  $\tau_{\leq n}^L: \text{Syn} \rightarrow \text{Syn}_{\leq n}^L$  for the  $n$ th *truncation* functor. By [Lur17, 1.4.4.13], which states accessible  $t$ -structures lead to accessible connective cover and truncation functors, the adjoint functor theorem [Lur09,

Cor.5.5.2.9], and the presentability of  $\text{Syn}$  [Pst23, Pr.4.2], the functor  $\tau_{\geq n}^L$  is right adjoint and  $\tau_{\leq n}^L$  is left adjoint to their respective inclusions. If  $L$  is the vertical line, then we write

$$\text{Syn}_{\geq n}^{\uparrow} = \text{Syn}_{\geq n}^L, \quad \text{Syn}_{\leq n}^{\uparrow} = \text{Syn}_{\geq n}^L, \quad \tau_{\geq n}^L = \tau_{\geq n}^{\uparrow}, \quad \tau_{\leq n}^L = \tau_{\leq n}^{\uparrow}$$

and refer to the associated  $t$ -structure as the *vertical  $t$ -structure*.

## 2.2 The heart of the linear $t$ -structure

To calculate the heart of these  $t$ -structures, we need the following algebraic construction.

**Definition 2.7.** Let  $L$  and  $E$  be as in Th.2.2. For  $X \in \text{Syn}$ , write  $\pi_*^L X$  for the bigraded homotopy groups  $\pi_{a,b} X$  where  $(a, b) \in L$ . As  $L$  is a line through the origin and the  $t$ -structure defined by  $L$  is monoidal, then  $\pi_*^L \mathbf{1}$  is a graded ring.

**Proposition 2.8.** *Let  $L$  and  $E$  be as in Th.2.2. Then the functor*

$$\pi_*^L: (\text{Syn}^L)^{\heartsuit} \xrightarrow{\simeq} \text{Mod}_{\pi_*^L \mathbf{1}}^{\text{gr}}$$

*is an equivalence of  $\infty$ -categories between the heart of the  $t$ -structure associated with  $L$  and the abelian category of graded modules over  $\pi_*^L \mathbf{1}$ .*

*Proof.* The functor

$$\pi_*^L: (\text{Syn}^L)^{\heartsuit} \rightarrow \text{Mod}_{\pi_*^L \mathbf{1}}^{\text{gr}}$$

is conservative as  $\text{Syn}_{\geq 0}^L$  is generated by the bigraded spheres  $\mathbf{S}^{a,b}$  with  $(a, b) \in L_{\geq}$  by our cellularity assumption. Moreover, it preserves cokernels, as for any  $(a, b) \in L$  and any  $Y \in \text{Syn}$ ,

$$[\mathbf{S}^{a,b}, Y] = [\mathbf{S}^{a,b}, \tau_{\leq 0}^L Y]$$

and if  $X \in \text{Syn}_{\geq 0}^L$ , then

$$[\mathbf{S}^{a,b}, \Sigma X] = [\mathbf{S}^{a-1, b+1}, X] = 0$$

as  $(a-1, b+1) \in L_{<}$ . A similar argument shows that  $\pi_*^L$  preserves kernels and products, and it follows that  $\pi_*^L$  has a left adjoint  $H(-)$ .

The classical Barr–Beck theorem implies that the adjunction is monadic, and it remains to show the monad  $\pi_*^L H(-)$  is the identity functor. Using that  $\pi_*^L$  commutes with filtered colimits, finite coproducts, and cokernels, it suffices to show that the counit map  $\epsilon: H(\pi_*^L \mathbf{1}) \rightarrow \tau_{\leq 0}^L \mathbf{1}$  is an equivalence. For any  $X \in (\text{Syn}^L)^{\heartsuit}$ , this induces a map

$$[\tau_{\leq 0}^L \mathbf{1}, X] \rightarrow [H(\pi_*^L \mathbf{1}), X]$$

Both sides are isomorphic to  $\pi_{0,0} X$ , on the left because  $X \simeq \tau_{\leq 0}^L X$  and on the right by adjunction. Using generation by bigraded spheres, it suffices to check this map is an isomorphism when  $X = \tau_{\leq 0}^L \mathbf{S}^{a,b}$  for all  $(a, b) \in L$ , which together give a map of  $\pi_*^L \mathbf{1}$ -modules  $\pi_*^L \mathbf{1} \rightarrow \pi_*^L \mathbf{1}$ , which is an isomorphism because it sends 1 to the map adjoint to the counit  $\epsilon$ , in other words, the identity.  $\square$

*Remark 2.9.* Although two different lines  $L$  and  $L'$  produce different  $t$ -structures on  $\text{Syn}$ , it follows from Pr.2.8, that many  $t$ -structures give the same heart. For example, all lines  $L$  defined by the equation  $y = ax$ , where  $a$  is irrational, have heart equivalent to graded modules over  $\pi_*^L \mathbf{1} \simeq \pi_{0,0} \mathbf{1}$ .

### 2.3 Homotopy groups of linear connective covers and truncations

Let us end this section by calculating some  $L$ -connective covers and  $L$ -truncations, as well as the compatibility of these  $t$ -structures with the usual  $t$ -structure on spectra.

**Proposition 2.10.** *Let  $L$  and  $E$  be as in Th.2.2 and  $X \in \text{Syn}$ . Furthermore, suppose that  $L$  has slope  $m$  with  $-1 < m \leq \infty$ . Then we have natural isomorphisms of abelian groups*

$$\pi_{a,b} \tau_{\geq n}^L X \simeq \begin{cases} \pi_{a,b} X & (a-n, b+n) \in L_{\geq} \\ 0 & (a-n, b+n) \in L_{<} \end{cases}$$

If  $m = \infty$ , then we have natural isomorphisms of abelian groups

$$\pi_{a,b}(\tau_{\geq n}^L(X)/\tau) \simeq \begin{cases} \pi_{a,b}(X/\tau) & (a-n, b+n) \in L_{>} \\ Z_L^\infty & (a-n, b+n) \in L \\ 0 & (a-n, b+n) \in L_{<} \end{cases}$$

where  $Z_L^\infty$  is the subgroup of  $\pi_*^L(X/\tau)$  consisting of permanent cycles in the  $\tau$ -BSS for  $X$ . If  $-1 < m < \infty$ , we have

$$\pi_{a,b}(\tau_{\geq n}^L(X)/\tau) \simeq \begin{cases} \pi_{a,b}(X/\tau) & (a-n, b+n) \in L_{>} \\ \pi_{a,b} X & (a-n, b+n) \in L \\ 0 & (a-n, b+n) \in L_{<} \end{cases}$$

*Proof.* The first claim follows from the definitions of  $\text{Syn}_{\geq n}^L$  and  $\text{Syn}_{\leq n-1}^L$  and the long exact sequence associated with

$$\tau_{\geq n}^L X \rightarrow X \rightarrow \tau_{\leq n-1}^L X \quad (2.11)$$

The claim about homotopy groups mod  $\tau$  follows from investigating the diagram of long exact sequences

$$\begin{array}{ccccccccc} \pi_{a,b+1} \tau_{\geq n}^L X & \xrightarrow{\tau} & \pi_{a,b} \tau_{\geq n}^L X & \longrightarrow & \pi_{a,b}(\tau_{\geq n}^L(X)/\tau) & \xrightarrow{\partial} & \pi_{a-1,b+2} \tau_{\geq n}^L X & \xrightarrow{\tau} & \pi_{a-1,b+1} \tau_{\geq n}^L X \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \pi_{a,b+1} X & \xrightarrow{\tau} & \pi_{a,b} X & \longrightarrow & \pi_{a,b}(X/\tau) & \xrightarrow{\partial} & \pi_{a-1,b+2} X & \xrightarrow{\tau} & \pi_{a-1,b+1} X \end{array} \quad (2.12)$$

If  $(a-n, b+n) \in L$ , then the second vertical map from the left is an isomorphism, and the last two terms in the top row are zero. If  $L$  is not vertical, then the top left term is also zero, and the claim follows in this case. If  $L$  is vertical, the leftmost vertical map is an isomorphism, and this identifies  $\pi_{a,b}(\tau_{\geq n}^L(X)/\tau)$  with the kernel of  $\partial$ , which is the subgroup of permanent cycles in the  $\tau$ -BSS.  $\square$

*Remark 2.13.* Similar claims hold for the  $L$ -truncations using (2.11) and (2.12).

*Remark 2.14.* One can make a more general statement when the line  $L$  has slope  $m$  with  $-\infty < m < -1$ . For example, the proof of Pr.2.10 shows that if  $m = -r$  for some integer  $2 \leq r < \infty$ , then the  $Z_L^\infty$  term in the case of  $(a-n, b+n) \in L$ , needs to be replaced with  $Z_L^{r-1}$ , the subgroup of  $\pi_*^L(X/\tau)$  consisting of  $d_{\leq r-1}$ -cycles in the  $\tau$ -BSS for  $X$ .

**Corollary 2.15.** *Let  $L$  and  $E$  be as in Pr.2.10,  $X$  be a  $\tau$ -complete synthetic spectrum, and  $n$  be an integer. Then both  $\tau_{\geq n}^L X$  and  $\tau_{\leq n}^L X$  are also  $\tau$ -complete.*

*Proof.* We will prove that  $\tau_{\geq n}^L X$  is  $\tau$ -complete for all  $n$ ; the same follows for  $\tau_{\leq n}^L X$  since being  $\tau$ -complete is closed under taking cofibres. Consider the diagram of Milnor sequences

$$\begin{array}{ccccccc}
0 & \longrightarrow & \lim_m^1 \pi_{a+1, b-1}(X_n/\tau^m) & \longrightarrow & \pi_{a,b}(\lim_m X_n/\tau^m) & \longrightarrow & \lim_m \pi_{a,b}(X_n/\tau^m) \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \lim_m^1 \pi_{a+1, b-1}(X/\tau^m) & \longrightarrow & \pi_{a,b}X & \longrightarrow & \lim_m \pi_{a,b}(X/\tau^m) \longrightarrow 0
\end{array} \tag{2.16}$$

where  $X_n = \tau_{\geq n}^L(X)$ . Applying Pr.2.10, we see that if  $(a-n, b+n) \in L_{>}$ , then there is an isomorphism  $\pi_{a,b}\tau_{\geq n}^L(X) \simeq \pi_{a,b}X$  and  $\pi_{a+1, b-1}(\tau_{\geq n}^L(X)) \simeq \pi_{a+1, b-1}X$ . The diagram above then shows that

$$\pi_{a,b}\tau_{\geq n}^L(X) \rightarrow \pi_{a,b}(\lim_m \tau_{\geq n}^L(X)/\tau^m)$$

is an isomorphism. If  $(a-n, b+n) \in L_{<}$ , then

$$\pi_{a,b}(\lim_m \tau_{\geq n}^L(X)/\tau^m) = \lim_m \pi_{a,b}(\tau_{\geq n}^L(X)/\tau^m) = \pi_{a,b}\tau_{\geq n}^L(X) = 0$$

so it remains to show that  $\lim_m^1 \pi_{a+1, b-1}(\tau_{\geq n}^L(X)/\tau^m) = 0$ . This is automatic if  $(a+1-n, b-1+n) \in L_{<}$ , and if  $(a+1-n, b-1+n) \in L_{\geq}$ , this follows as one has  $\pi_{s,t}(\tau_{\geq n}^L(X)/\tau^m) = \pi_{s,t}(\tau_{\geq n}^L(X))/\tau^m$  whenever  $(s-n, t+n) \in L_{\geq}$  and  $(s-1-n, t+1+n) \in L_{<}$ , so that the inverse system is Mittag-Leffler.

If  $(a-n, b+n) \in L$ , the lefthand vertical map is an isomorphism as before, and the result follows from the claim that the map

$$\lim_m \pi_{a,b}(\tau_{\geq n}^L(X)/\tau^m) \rightarrow \lim_m \pi_{a,b}(X/\tau^m)$$

is an isomorphism. To see this, first note that for each  $m$

$$\pi_{a,b}(\tau_{\geq n}^L(X)/\tau^m) \rightarrow \pi_{a,b}(X/\tau^m)$$

is an injection as it is identified with the map  $\pi_{a,b}(X)/\tau^m \rightarrow \pi_{a,b}(X/\tau^m)$ , which implies the map given by taken limits is also an injection. The map also becomes a surjection in the limit because, by  $\tau$ -completeness of  $X$ , each class on the right-hand side admits a lift to  $\pi_{a,b}X$ , whose projection to  $\pi_{a,b}(X/\tau^m)$  must factor through  $\pi_{a,b}(X)/\tau^m$  since the composite  $\pi_{a,b}X \rightarrow \pi_{a,b}(X/\tau^m) \xrightarrow{\hat{\partial}} \pi_{a-1, b+1}X$  is zero.  $\square$

Although we have worked mostly with a general line through the origin in this section, there are some statements that behave best with respect to vertical  $t$ -structure. For example, the following interaction between vertical connective covers and truncations and  $\tau$ -inversion is an immediate consequence of Pr.2.10.

**Corollary 2.17.** *Let  $L$  be the vertical line,  $E$  be as in Th.2.2,  $X \in \text{Syn}$ , and  $n$  be an integer. Then there are natural equivalences of spectra*

$$\tau^{-1}\tau_{\geq n}^{\uparrow}X \simeq \tau_{\geq n}\tau^{-1}X \qquad \tau^{-1}\tau_{\leq n}^{\uparrow}X \simeq \tau_{\leq n}\tau^{-1}X.$$

*In particular, if  $Y \in \text{Sp}$  is connective, then  $\nu Y$  is vertically connective.*

*Proof.* Since  $\tau$  has bidegree  $(0, -1)$ , we have that

$$\tau^{-1}\pi_{k,*}\tau_{\geq n}^{\uparrow}X = \pi_k(\tau^{-1}X)[\tau^{\pm}]$$

for all  $k$ , so that in particular the homotopy groups of  $\tau^{-1}\tau_{\geq n}^{\uparrow}X$  are concentrated in degrees  $\geq n$ . In particular, the natural map  $\tau^{-1}\tau_{\geq n}^L X \rightarrow \tau^{-1}X$  factors through a map  $\tau^{-1}\tau_{\geq n}^L X \rightarrow \tau_{\geq n}\tau^{-1}X$ , which is an equivalence by Pr.2.10. For truncations, combine the same argument with Rmk.2.13.

The “in particular” statement follows this together with Th.1.3 applied to  $\Gamma\nu Y$ .  $\square$

*Remark 2.18.* To see the necessity of  $L$  being the vertical line, consider  $E = \text{BP}$  at the prime 2,  $X = \text{j}_{\text{BP}}$  as defined in Df.4.2 to come, and let  $L$  be defined by the equation  $x = y$ . Then the connective cover of  $\tau^{-1}\text{j}_{\text{BP}}$  is equivalent to  $\text{j}$ , and in particular,  $\pi_4\tau^{-1}\text{j}_{\text{BP}} = 0$ . On the other hand, from Fig.5, we see that  $\pi_4\tau^{-1}\tau_{\geq 0}^L\text{j}_{\text{BP}} \neq 0$ . Indeed, the connective cover will cut off exactly the top-diagonal line of red circles in the  $\Gamma$ -SS associated with  $\tau_{\geq 0}^L\text{j}_{\text{BP}}$ , meaning the class in degree  $(4, 2)$  on the  $E_2$ -page cannot support any differentials, so it survives to a nonzero class in  $\pi_4(\tau^{-1}\tau_{\geq 0}^L\text{j}_{\text{BP}})$ . In general, if  $X$  is an  $E$ -nilpotent complete spectrum with the property that the  $E_2$ -page of its  $E$ -ASS vanishes in  $L_{<}$ , then  $\tau^{-1}\tau_{\geq 0}^L\nu X \simeq \tau_{\geq 0}\tau^{-1}\nu X$ . Similar statements hold for certain vanishing *curves* with associated  $t$ -structure given by the graph of said curve; we will come back to this again in future work.

### 3 Synthetic analogues of topological $K$ -theory

As a warm-up to our synthetic constructions and calculations of various image-of- $J$  spectra, let us show that calculations of the ASSs and ANSSs of  $\text{ko}$  and  $\text{KO}$ , and their odd primary counterparts  $\ell$  and  $\text{L}$ , can easily be obtained in a purely synthetic manner. We remind the reader that we are implicitly working in the  $p$ -complete setting.

#### 3.1 $\nu \text{ku}$ and $\nu \ell$ in $\mathbf{F}_p$ -synthetic spectra

A standard calculation using the cofibre sequences of spectra

$$\Sigma^{2p-2}\text{BP}\langle 1 \rangle \xrightarrow{v_1} \text{BP}\langle 1 \rangle \rightarrow \mathbf{Z}_{(p)} \qquad \mathbf{Z}_{(p)} \xrightarrow{p} \mathbf{Z}_{(p)} \rightarrow \mathbf{F}_p$$

allows one to compute the mod  $p$  homology of any form of  $\mathrm{BP}\langle 1 \rangle$  (such as  $\mathrm{ku}$  at  $p = 2$  and  $\ell$  at  $p > 2$ )—this is the only homology calculation we will require as input for this article. The result is an isomorphism of comodules

$$H_*(\mathrm{BP}\langle 1 \rangle; \mathbf{F}_p) \simeq \mathcal{A}_{*\square_{\mathcal{E}(1)}} \mathbf{F}_p \quad (3.1)$$

The resulting Adams  $E_2$ -page is polynomial on even generators, implying collapse, and we deduce the following.

**Proposition 3.2.** *In  $\mathrm{Syn}_{\mathbf{F}_2}$ , we have*

$$\pi_{*,*} \nu \mathrm{ku} = \mathbf{Z}[h_0, q_1, \tau]/(\tau h_0 - 2)$$

where  $|h_0| = (0, 1)$ ,  $|q_1| = (2, 1)$ , and  $|\tau| = (0, -1)$ . In  $\mathrm{Syn}_{\mathbf{F}_p}$  for  $p > 2$ , we have

$$\pi_{*,*} \nu \ell = \mathbf{Z}[h_0, q_1, \tau]/(\tau h_0 - p)$$

where  $|h_0| = (0, 1)$ ,  $|q_1| = (2p - 2, 1)$ , and  $|\tau| = (0, -1)$ .

This computation allows us to define lifts of  $\mathrm{KU}$  and  $L$  to  $\mathrm{Syn}_{\mathbf{F}_p}$  with nonvanishing Adams spectral sequences, in contrast to the fact that  $\nu \mathrm{KU}/\tau$  and  $\nu L/\tau$  are contractible as  $\mathrm{KU} \otimes \mathbf{F}_p$  vanishes.

**Definition 3.3.** Set  $\mathrm{KU}_{\mathbf{F}_2} := \nu \mathrm{ku}[q_1^{-1}]$  and  $L_{\mathbf{F}_p} := \nu \ell[q_1^{-1}]$ .

It is easy to check that  $\tau^{-1} \mathrm{KU}_{\mathbf{F}_2} \simeq \mathrm{KU}$  and  $\tau^{-1} L_{\mathbf{F}_p} \simeq L$ , as  $\tau^{-1}$  commutes with colimits and  $\tau^{-1}(q_1) = v_1$ . Moreover, from Pr.2.10, we see there are equivalences of  $\mathbf{F}_p$ -synthetic spectra  $\tau_{\geq 0}^{\uparrow} \mathrm{KU}_{\mathbf{F}_p} \simeq \nu \mathrm{ku}$  and similarly that  $\tau_{\geq 0}^{\uparrow} L_{\mathbf{F}_p} \simeq \nu \ell$ .

### 3.2 Synthetic Wood's theorem

Recall the following standard way to compute  $\pi_* \mathrm{ko}$ . One starts by proving Wood's theorem, which states that there is an equivalence  $\mathrm{KO} \otimes C(\eta) \simeq \mathrm{KU}$  or, in other words, a cofibre sequence

$$\Sigma \mathrm{KO} \xrightarrow{\eta} \mathrm{KO} \rightarrow \mathrm{KU}. \quad (3.4)$$

One can prove this using the identification  $C(\eta) \simeq \Sigma^{-2} \mathbf{CP}^2$  and the fact that  $\mathrm{KU}_*(\mathbf{CP}^2)$  is dual to  $\mathrm{KU}^*(\mathbf{CP}^2)$ ; see §3.2 in the arXiv version 4 of [Mat16]. From Wood's theorem, it follows that  $\mathrm{KO}$  detects  $\eta \in \pi_1 \mathbf{S}$ . Indeed, if not, then we would have a splitting  $\mathrm{KU} \simeq \mathrm{KO} \otimes C(\eta) \simeq \mathrm{KO} \oplus \mathrm{KO}[2]$ , however, the existence of the Möbius bundle on  $S^1$ , which represents a 2-torsion element in  $\pi_1 \mathrm{KO}$ , shows this to be impossible.

The complexification map  $c: \mathrm{KO} \rightarrow \mathrm{KU}$  induces an isomorphism on  $\pi_0$ , and it follows from the Wood sequence (3.4) and the evenness of  $\mathrm{KU}$  that  $\pi_k \mathrm{KO} = 0$  for  $k = -1, -2$ . As  $C(\eta)$  is a finite spectrum with cells only in dimensions 0 and 2, it follows that  $\mathrm{ko} \otimes C(\eta) \simeq \mathrm{ku}$ . Using (3.4), one can now compute all of the homotopy groups of  $\mathrm{ko}$  using only that  $\eta$  is nilpotent. One needs only to determine the effect of the boundary map in the Wood sequence on homotopy

groups, which can be determined from the simple facts that  $\mathrm{ko} \rightarrow \mathrm{ku}$  is an isomorphism on  $\pi_0$  and that  $\pi_1 \mathrm{ko}$  contains 2-torsion. One then obtains the calculation

$$\partial(u^n) = \begin{cases} 0 & n = 4k \\ 2\beta^k & n = 4k + 1 \\ \eta^2 \beta^k & n = 4k + 2 \\ \alpha \beta^k & n = 4k + 3 \end{cases} \quad (3.5)$$

where  $\alpha$  and  $\beta$  are generators of  $\pi_4 \mathrm{ko}$  and  $\pi_8 \mathrm{ko}$ , respectively. We will now observe that this argument can be run with little modification in  $\mathrm{Syn}_{\mathbf{F}_2}$  and  $\mathrm{Syn}_{\mathrm{BP}}$  to compute the bigraded synthetic homotopy groups of  $\nu \mathrm{ko}$ . We begin with a synthetic version of the Wood sequence.

**Proposition 3.6.** *There is a cofibre sequence in  $\mathrm{Syn}_E$*

$$\Sigma^{1,1} \nu \mathrm{ko} \xrightarrow{h_1} \nu \mathrm{ko} \xrightarrow{\nu c} \nu \mathrm{ku}$$

for  $E = \mathbf{F}_2$  and  $E = \mathrm{BP}$ , which is a synthetic lift of the connective cover of (3.4), where  $h_1$  is the unique class in  $\pi_{1,1} \mathbf{S}$  such that  $\tau^{-1}(h_1) = \eta$ .

*Proof.* The cofibre sequence of spectra

$$\mathbf{S}^0 \rightarrow C(\eta) \rightarrow \mathbf{S}^2$$

induces a short exact sequence on  $\mathbf{F}_2$ - and BP-homology, so by [Pst23, Lm.4.23], it follows that one has a cofibre sequence

$$\mathbf{S}^{0,0} \rightarrow \nu C(\eta) \rightarrow \mathbf{S}^{2,0}$$

in  $\mathrm{Syn}_E$ . The  $\tau$ -inversion is an exact functor, so we see that the boundary map  $\partial: \mathbf{S}^{1,1} \rightarrow \mathbf{S}^{0,0}$  in this cofibre sequence must  $\tau$ -invert to  $\eta$ . This  $\partial$  then represents the desired  $h_1 \in \pi_{1,1} \mathbf{S}$ . In particular, we have an equivalence of synthetic spectra  $\nu C(\eta) \simeq C(h_1)$ . As  $C(\eta)$  is both an  $\mathbf{F}_2$ - and BP-finite projective, we use [Pst23, Lm.4.24] to see that

$$\nu \mathrm{ku} \simeq \nu(\mathrm{ko} \otimes C(\eta)) \simeq \nu \mathrm{ko} \otimes \nu C(\eta) \simeq \nu \mathrm{ko} \otimes C(h_1). \quad \square$$

*Warning 3.7.* According to the last line in the above proof, there is an equivalence of  $\mathbf{F}_2$ -synthetic spectra  $\nu \mathrm{ku} \simeq \nu \mathrm{ko} \otimes C(h_1)$ . Not all classical relationships between  $\mathrm{ku}$  and  $\mathrm{ko}$  lift to synthetic spectra though. For example,  $\tau_{\geq 0}^\uparrow(\nu \mathrm{ku})^{hC_2}$  is **not** equivalent to  $\nu \mathrm{ko}$ ; see Warn.3.16.

Using the calculation of  $\pi_{*,*} \nu \mathrm{ku}$  from Pr.3.2, we can now calculate  $\pi_{*,*} \nu \mathrm{ko}$  from the long exact sequence

$$\cdots \rightarrow \pi_{n+1,*-1} \nu \mathrm{ku} \rightarrow \pi_{n-1,*-1} \nu \mathrm{ko} \xrightarrow{h_1} \pi_{n,*} \nu \mathrm{ko} \rightarrow \pi_{n,*} \nu \mathrm{ku} \xrightarrow{\partial} \pi_{n-2,*} \nu \mathrm{ko} \rightarrow \cdots \quad (3.8)$$

We begin with the  $\mathbf{F}_2$ -synthetic case.

### 3.3 $\nu \text{ko}$ inside $\text{Syn}_{\mathbf{F}_2}$

In this subsection, we compute the bigraded  $\mathbf{F}_2$ -synthetic homotopy groups of  $\nu \text{ko}$  using only the synthetic version of Wood's theorem and straightforward long exact sequence arguments. This, by Th.1.3, records the data of the (standard) ASS for  $\text{ko}$  at the prime 2. By contrast, the usual recipe to compute the ASS of  $\text{ko}$  involves first determining the comodule  $H_*(\text{ko}; \mathbf{F}_2)$  and then computing  $\text{Ext}_{\mathcal{A}(1)_*}(\mathbf{F}_2, \mathbf{F}_2)$ . The latter may be done by hand using minimal resolutions, but is a significantly more tedious calculation than what we will see here.

**Corollary 3.9.** *The  $\mathbf{F}_2$ -synthetic homotopy groups of  $\nu \text{ko}$  are given by*

$$\pi_{*,*} \nu \text{ko} = \mathbf{Z}[h_0, h_1, \alpha, \beta, \tau] / (h_0\tau - 2, h_0h_1, h_1^3, h_1\alpha, \alpha^2 - h_0^2\beta)$$

where  $|h_0| = (0, 1)$ ,  $|h_1| = (1, 1)$ ,  $|\alpha| = (4, 3)$ ,  $|\beta| = (8, 4)$ , and  $|\tau| = (0, -1)$ . The associated spectral sequence is given as Fig.1.

*Proof.* For notational simplicity, write  $A$  for the graded ring  $\mathbf{Z}[\tau, h_0]/(\tau h_0 - 2)$ . We will show by induction that, when  $k \geq 0$ , there are isomorphisms of  $\pi_{0,*} \nu \text{ko}$ -modules

$$\pi_{8k+r,*} \nu \text{ko} \simeq \begin{cases} A\{\beta^k\} & r = 0 \\ (A/h_0)\{h_1^r \beta^k\} & r = 1, 2 \\ A\{\alpha\beta^k\} & r = 4 \\ 0 & \text{else,} \end{cases} \quad |\beta| = (8, 4), |\alpha| = (4, 3),$$

where the degrees of the generators help us track the  $\tau$ -degree of these modules. Simultaneously, we will show that the boundary maps are given by

$$\partial(q_1^n) = \begin{cases} 0 & n = 4k \\ h_0\beta^k & n = 4k + 1 \\ h_1^2\beta^k & n = 4k + 2 \\ \alpha\beta^k & n = 4k + 3. \end{cases}$$

using that, in each case, the claimed value of  $\partial(q_1^n)$  is the unique class in the correct bidegree which  $\tau$ -inverts to the known value of  $\partial(u^n)$  in (3.5). The claimed multiplicative structure then follows from the  $h_1$ -multiplications and the fact that  $\nu \text{ko} \rightarrow \nu \text{ku}$  is a ring map.

In fact, we only need to show that  $\pi_{8k,*} \nu \text{ko} \rightarrow \pi_{8k,*} \nu \text{ku}$  is an isomorphism to determine  $\pi_{8k+l,*} \nu \text{ko}$  for  $0 < l \leq 8$ , and the effect of the boundary map in this range. For the base case, consider (3.8) when  $n = 0$ ; the fact that  $\text{ko}$  is connective combined with Cor.2.17 shows that  $\nu \text{ko}$  is connective in the vertical  $t$ -structure, implying that  $\pi_{0,*} \nu \text{ko} \rightarrow \pi_{0,*} \nu \text{ku} = A$  is an isomorphism by (3.8).

Assume now that  $\pi_{8k,*} \nu \text{ko} \rightarrow \pi_{8k,*} \nu \text{ku}$  is an isomorphism, for  $k \geq 0$ . Then  $\pi_{8k,*} \nu \text{ko}$  is  $\tau$ -torsion free, so the boundary map  $\pi_{8k+2,*} \nu \text{ku} \rightarrow \pi_{8k,*} \nu \text{ko}$  is determined by its effect after inverting  $\tau$ . It follows that it sends  $q_1^{4k+1} \mapsto h_0\beta^k$ . Using (3.8) along with the fact

that  $\pi_{\text{odd},*}\nu\text{ku} = 0$  we see that  $\pi_{8k+1,*}\nu\text{ko} = (A/h_0)\{h_1\beta^k\}$ , and that  $h_1 : \pi_{8k+1,*}\nu\text{ko} \rightarrow \pi_{8k+2,*+1}\nu\text{ko}$  is an isomorphism.

In particular, we see that  $\pi_{8k+2,*}\nu\text{ko}$  is  $\tau$ -torsion free, so as before we see that  $\partial$  sends  $q_1^{4k+2} \mapsto h_1^2\beta^k$ . By (3.8) we see that  $\pi_{8k+3,*}\nu\text{ko} = 0$ , which implies  $\pi_{8k+4,*}\nu\text{ko} \rightarrow \pi_{8k+4,*}\nu\text{ku} = A$  is an injection with image the ideal generated by  $h_0$  (note there is an  $A$ -module isomorphism  $(h_0) \cong \Sigma^{0,1}A$ ).

Then  $\pi_{8k+4,*}\nu\text{ko}$  is  $\tau$ -torsion free, so as before we see that  $\partial$  sends  $q_1^{4k+3} \mapsto \alpha\beta^k$ . Similar arguments to the above now imply that  $\pi_{8k+l,*}\nu\text{ko} = 0$  for  $4 < l < 8$  and that  $\pi_{8k+8,*}\nu\text{ko} \rightarrow \pi_{8k+8,*}\nu\text{ku}$  is an isomorphism. This completes the induction.

The computation of the associated spectral sequence (Fig.1) follows immediately from Th.1.3; since  $\pi_{*,*}\nu\text{ko}$  is  $\tau$ -torsion free, we have  $\pi_{*,*}(\nu\text{ko}/\tau) = (\pi_{*,*}\nu\text{ko})/\tau$ , and there are no nonzero differentials.  $\square$

We consider the above computation as only a slight complication of the classical LES used to calculate  $\pi_*\text{ko}$  from  $\pi_*\text{ku}$ . In particular, we consider this computation of  $\pi_{*,*}\nu\text{ko}$  and its signature Fig.1 as a vast simplification of the usual process of first computing the comodule  $H_*(\text{ko}; \mathbf{F}_2) \simeq \mathcal{A}_* \square_{\mathcal{A}(1)_*} \mathbf{F}_2$ , which also follows from (3.1) and Wood's theorem, and then calculating  $\text{Ext}_{\mathcal{A}(1)_*}(\mathbf{F}_2, \mathbf{F}_2)$ .

The above calculation also leads to a definition of  $\text{KO}_{\mathbf{F}_2}$ , an object which produces a nontrivial modified  $\mathbf{F}_2$ -ASS for  $\text{KO}$ .

**Definition 3.10.** In  $\text{Syn}_{\mathbf{F}_2}$ , we set  $\text{KO}_{\mathbf{F}_2} := \nu\text{ko}[\beta^{-1}]$ .

**Corollary 3.11.** *The  $\mathbf{F}_p$ -synthetic spectra  $\nu\text{ku}, \text{KU}_{\mathbf{F}_p}, \nu\ell, \text{L}_{\mathbf{F}_p}, \nu\text{ko}, \text{KO}_{\mathbf{F}_2}$  are all  $\tau$ -complete, and applying  $\tau^{-1}$  gives the spectra  $\text{ku}, \text{KU}, \ell, \text{L}, \text{ko}, \text{KO}$ , respectively.*

*Proof.* By [BHS19, Pr.A.13], the fact that all spectra in sight are implicitly  $p$ -complete, hence  $\mathbf{F}_p$ -complete, lead to  $\nu\mathbf{F}_p$ - and  $\tau$ -complete synthetic analogues. In the periodic cases, one checks directly that the  $\tau$ -adic towers are all Mittag-Leffler on bigraded homotopy groups and converge to the bigraded homotopy groups of the synthetic spectrum in question. The claim about  $\tau$ -inversion follows from the natural equivalence  $\tau^{-1} \circ \nu \simeq \text{id}$ , the fact that  $\tau^{-1}(q_1) = v_1$ , and the fact that  $\tau^{-1}$  commutes with colimits.  $\square$

### 3.4 In BP-synthetic spectra

These arguments simplify further if we take  $E = \text{BP}$ . Indeed, in this setting, we require *no* BP-homology calculations, only the knowledge of  $\text{KU}_* = \mathbf{Z}[u^{\pm}]$  and the fact that the Adams–Novikov spectral sequence of a complex-oriented spectrum  $E$  degenerates at the  $E_2$ -page. This, in turn, corresponds to the fact that the coaugmented cosimplicial object

$$E \rightarrow E \otimes \text{BP} \rightrightarrows E \otimes \text{BP} \otimes \text{BP} \rightrightarrows \dots$$

from which the ANSS of  $E$  is constructed has a  $(-1)$ -st codegeneracy coming from the ring map  $\text{BP} \rightarrow E$ . This tells us that the bigraded homotopy groups of  $\nu E$  in  $\text{Syn}_{\text{BP}}$  are  $\tau$ -torsion free; for example, we immediately obtain the calculations

$$\pi_{*,*}\nu \text{ku} \simeq \mathbf{Z}[u, \tau], \quad \pi_{*,*}\nu \text{KU} \simeq \mathbf{Z}[u^\pm, \tau], \quad \pi_{*,*}\nu \ell \simeq \mathbf{Z}[v_1, \tau], \quad \pi_{*,*}\nu L \simeq \mathbf{Z}[v_1^\pm, \tau]$$

where  $|u| = (2, 0)$  and  $|v_1| = (2p - 2, 0)$ . As in the previous subsection, we use Pr.3.6 and the long exact sequence (3.8) to compute  $\pi_{*,*}\nu \text{ko}$  in  $\text{Syn}_{\text{BP}}$ . This determines the entire data of the ANSS of  $\text{ko}$ .

**Proposition 3.12.** *The BP-synthetic homotopy groups of  $\nu \text{ko}$  are given by*

$$\pi_{*,*}\nu \text{ko} = \mathbf{Z}[h_1, \alpha, \beta, \tau]/(2h_1, \tau^2 h_1^3, h_1 \alpha, \alpha^2 - 4\beta)$$

where  $|h_1| = (1, 1)$ ,  $|\alpha| = (4, 0)$ ,  $|\beta| = (8, 0)$ , and  $|\tau| = (0, -1)$ , The associated spectral sequence is given as Fig.2.

*Proof.* The argument closely resembles the  $\text{Syn}_{\mathbf{F}_2}$ -case, so we will be brief. The crucial point is to compute the effect of the boundary map in (3.8) in  $\text{Syn}_{\text{BP}}$ ; the computation then follows as usual by computing the kernel and cokernel. Note that the boundary maps  $\pi_{2k,*}\nu \text{ku} \rightarrow \pi_{2k-2,*}\nu \text{ko}$  are determined by their effect on  $u^k \in \pi_{2k,0}\nu \text{ku}$  as maps of  $\mathbf{Z}[\tau]$ -modules. The boundary maps are therefore determined by their effect after inverting  $\tau$ , as  $\pi_{n,0}\nu \text{ko}$  is always  $\tau$ -torsion free. Indeed, a nonzero  $\tau$ -torsion class in  $\pi_{n,0}\nu \text{ko}$  would imply the existence of a nonzero differential in  $\text{ANSS}(\text{ko})$  with target in bidegree  $(n, 0)$ ; this is impossible as standard ANSSs are concentrated in nonnegative filtrations. In particular, we find that

$$\partial(u^n) = \begin{cases} 0 & n = 4k \\ 2\beta^k & n = 4k + 1 \\ \tau^2 h_1^2 \beta^k & n = 4k + 2 \\ \alpha \beta^k & n = 4k + 3 \end{cases}$$

as, in each case, the claimed value of  $\partial(u^n)$  is the unique class in the correct bidegree which  $\tau$ -inverts to the known value of  $\partial(u^n)$  in (3.5).  $\square$

Either by inverting  $\beta$  or using a similar argument as above with  $\nu \text{KO}$ , one may also compute  $\pi_{*,*}\nu \text{KO}$ .

In the category  $\text{Syn}_{\text{BP}}$ , it is interesting to know that one may construct synthetic lifts of  $\text{KO}$  and  $\text{ko}$  directly from  $\nu \text{KU}$ , and these turn out to recover the synthetic analogues  $\nu \text{KO}$  and  $\nu \text{ko}$ . The rest of this section is not necessary for the main purposes of this article, we only include it to provide further intuition and warnings for both the reader and ourselves.

**Definition 3.13.** In  $\text{Syn}_{\text{BP}}$ , we set

$$\begin{aligned} \text{KO}_{\text{BP}} &:= (\nu \text{KU})^{hC_2} \\ \text{ko}_{\text{BP}} &:= \tau_{\geq 0}^\uparrow \text{KO}_{\text{BP}} \end{aligned}$$

**Proposition 3.14.** *There are equivalences of BP-synthetic spectra  $\text{Syn}_{\text{BP}}, \nu \text{KO} \simeq \text{KO}_{\text{BP}}$  and  $\nu \text{ko} \simeq \text{ko}_{\text{BP}}$ .*

*Proof.* There are natural maps  $\nu \text{KO} \rightarrow \text{KO}_{\text{BP}}$  and  $\nu \text{ko} \rightarrow \text{ko}_{\text{BP}}$  by the universal property of fixed-points, and it suffices to show they induce isomorphisms on bigraded homotopy groups. We may compute the bigraded homotopy groups of  $\text{KO}_{\text{BP}}$  (and thereby of  $\text{ko}_{\text{BP}}$ ) by running the homotopy fixed point spectral sequence (HFPSS) internal to  $\text{Syn}_{\text{BP}}$ . This HFPSS has signature

$$E_2 = H^s(C_2; \pi_{n,f} \nu \text{KU}) \implies \pi_{n-s, f+s} \text{KO}_{\text{BP}}$$

drawn in Adams trigrading  $(n-s, s+f, f)$ , the differential  $d_r$  has signature  $(-1, +1, +r)$ .

The synthetic homotopy groups  $\pi_{*,*} \nu \text{KU} = \mathbf{Z}[u, \tau]$  are  $\tau$ -torsion free, so it follows that the generator of  $C_2$  fixes  $\tau$  and sends  $u \mapsto -u$ , as this is true after inverting  $\tau$ . Standard group cohomology calculations then give

$$E_2 = \mathbf{Z}[h_1, u^{\pm 2}, \tau]/(2h_1)$$

This is, in particular,  $\tau$ -torsion free, and one recovers the classical HFPSS for  $\text{KU}$  upon  $\tau$ -inversion, since applying  $\tau^{-1}$  to the cosimplicial object

$$\nu \text{KU} \implies F(C_{2+}, \nu \text{KU}) \cong F(C_2^{\times 2}_{+}, \nu \text{KU}) \cdots$$

gives the cobar resolution for  $\text{KU}$ . In the classical HFPSS for  $\text{KU}$ , there is a differential  $d_3(u^2) = \eta^3$ , which may be deduced from the fact that  $\eta$  is nilpotent. Therefore  $\tau^{-1}(d_3(u^2)) = \eta^3$ , and synthetically,  $\tau^2 h_1^3$  is the unique class in the tridegree of  $d_3(u^2)$  that  $\tau$ -inverts to  $\eta^3$ , so we deduce  $d_3(u^2) = \tau^2 h_1^3$ . The HFPSS collapses on the  $E_4$  page for degree reasons, and the  $E_\infty$  page is isomorphic to the bigraded homotopy groups of  $\nu \text{KO}$  as in Pr.3.12. There are no nontrivial extensions as each relation holds in or above the highest nonzero filtration in its respective stem on the  $E_\infty$  page of the HFPSS.  $\square$

*Remark 3.15.* One can directly show that  $\text{KO}_{\text{BP}}$  and  $\text{ko}_{\text{BP}}$  are  $\tau$ -complete and that  $\tau^{-1} \text{KO}_{\text{BP}} = \text{KO}$  and  $\tau^{-1} \text{ko}_{\text{BP}} = \text{ko}$  without referring to Pr.3.14. From the point of view of this article, it is unnecessary to consider  $\nu \text{KO}$  and  $\nu \text{ko}$  in  $\text{Syn}_{\text{BP}}$  and rather to focus on the purely synthetic constructions, even in this case they happen to agree. Regardless, the calculations in this subsection show that the bigraded homotopy groups of a synthetic lift of  $\text{KO}$  and  $\text{ko}$ , and therefore also the associated modified ANSSs, may be calculated without ever computing BP-homology.

*Warning 3.16.* It is not possible to construct  $\mathbf{F}_2$ -synthetic lifts of  $\text{KO}$  and  $\text{ko}$  using homotopy fixed points internal to  $\text{Syn}_{\mathbf{F}_2}$ . Indeed, from the HFPSS internal to  $\text{Syn}_{\mathbf{F}_2}$  with signature

$$E_2 = H^s(C_2; \pi_{n,f} \text{KU}_{\mathbf{F}_2}) \implies \pi_{n-s, f+s} (\text{KU}_{\mathbf{F}_2})^{hC_2}.$$

the nonzero class in  $H^1(C_2; \pi_{2,1} \text{KU}_{\mathbf{F}_2})$  detects a class  $x \in \pi_{1,2} (\text{KU}_{\mathbf{F}_2})^{hC_2}$  with  $h_0 x \neq 0$  and  $2x = 0$ . This implies that  $h_0 x$  is a nonzero  $\tau$ -torsion class. However, it follows from Cor.3.9 that  $\pi_{*,*} \text{KO}_{\mathbf{F}_2}$  is  $\tau$ -torsion free, so  $\text{KO}_{\mathbf{F}_2} \not\cong (\text{KU}_{\mathbf{F}_2})^{hC_2}$  and similarly  $\tau_{\geq 0}^{\uparrow} (\nu \text{ku})^{hC_2} \not\cong \nu \text{ko}$ . In fact, the computation shows that  $(\text{KU}_{\mathbf{F}_2})^{hC_2}$  is not even a synthetic lift of  $\text{KO}$ , that is,  $\tau^{-1}((\text{KU}_{\mathbf{F}_2})^{hC_2}) \not\cong \text{KO}$ .

The following is simply [BHS19, Pr.A.13] and the natural equivalence  $\tau^{-1} \circ \nu \simeq \text{id}$ .

**Corollary 3.17.** *The BP-synthetic spectra  $\nu \text{ku}, \nu \text{KU}, \nu \ell, \nu L, \nu \text{ko}, \nu \text{KO}$  are all  $\tau$ -complete, and applying  $\tau^{-1}$  gives the spectra  $\text{ku}, \text{KU}, \text{KO}, \text{ko}, \ell, L$ , respectively.*

## 4 Two synthetic image-of- $J$ spectra

Classically, one defines the connective image-of- $J$  spectrum  $j$  at the prime 2 as follows: first, one shows that the map of spectra  $\psi^3 - 1: \text{ko} \rightarrow \text{ko}$  factors through  $\tau_{\geq 4} \text{ko}$ , and then one defines  $j$  as the fibre of this factorisation. To construct our synthetic lifts of  $j$ , we need to show that the map of synthetic spectra  $\psi^3 - 1: \nu \text{ko} \rightarrow \nu \text{ko}$  factors through a connective cover with respect to our vertical  $t$ -structure.

### 4.1 Construction of synthetic lifts of $j$

**Proposition 4.1.** *Let  $E$  be either  $\mathbf{F}_p$  or BP. Then there is a factorization of the maps*

$$\psi^3 - 1 : \nu \text{ko} \rightarrow \nu \text{ko} \quad \psi^{p+1} - 1 : \nu \ell \rightarrow \nu \ell$$

through  $\tau_{\geq 4}^{\uparrow} \nu \text{ko}$  and  $\tau_{\geq 2p-2}^{\uparrow} \nu \ell$  at  $p = 2$  and odd primes, respectively.

*Proof.* There is a fibre sequence of synthetic spectra  $\tau_{\geq 3}^{\uparrow} \nu \text{ko} \rightarrow \nu \text{ko} \rightarrow \tau_{\leq 2}^{\uparrow} \nu \text{ko}$ . Notice that the composite

$$\nu \text{ko} \xrightarrow{\psi^3 - 1} \nu \text{ko} \rightarrow \tau_{\leq 2}^{\uparrow} \nu \text{ko},$$

or its adjunct  $\tau_{\leq 2}^{\uparrow} \nu \text{ko} \xrightarrow{\tau_{\leq 2}^{\uparrow}(\psi^3 - 1)} \tau_{\leq 2}^{\uparrow} \nu \text{ko}$ , is null. Indeed, as the truncation of the unit  $\tau_{\leq 2}^{\uparrow} \mathbf{1} \rightarrow \tau_{\leq 2}^{\uparrow} \nu \text{ko}$  is an equivalence for both  $E = \text{BP}$  and  $\mathbf{F}_p$ , then this adjunct is equivalent to a map  $\tau_{\leq 2}^{\uparrow} \mathbf{1} \rightarrow \tau_{\leq 2}^{\uparrow} \nu \text{ko}$ , which is in turn adjoint to the map

$$(\psi^3 - 1)(1): \mathbf{1} \rightarrow \tau_{\leq 2}^{\uparrow} \nu \text{ko}.$$

This map is null as it represents zero in  $\pi_{0,0} \tau_{\leq 2}^{\uparrow} \nu \text{ko}$ . This shows the desired map factors through  $\tau_{\geq 3}^{\uparrow} \nu \text{ko}$ . As  $\pi_{3,*} \nu \text{ko} = 0$  for  $E = \mathbf{F}_2$ , see Fig.1, it follows that  $\tau_{\geq 3}^{\uparrow} \nu \text{ko} = \tau_{\geq 4}^{\uparrow} \nu \text{ko}$ , so the  $E = \mathbf{F}_2$ -case is done. Similarly, for an odd prime  $p$ , we have  $\pi_{2p-3,*} \nu \ell = 0$  for both  $E = \mathbf{F}_p$  and BP, so these cases are also covered. For  $E = \text{BP}$  at  $p = 2$ , we have the following solid diagram of synthetic spectra

$$\begin{array}{ccc} & \tau_{\geq 4}^{\uparrow} \nu \text{ko} & \\ & \nearrow & \downarrow \\ \nu \text{ko} & \xrightarrow{\psi^3 - 1} \tau_{\geq 3}^{\uparrow} \nu \text{ko} & \longrightarrow \Sigma^{3,3} \mathbf{F}_2[\tau]/\tau^2 \end{array}$$

where the composite of the vertical and the right map is a fibre sequence; see Fig.2. To obtain the dashed arrow, we want to show that the lower composite is null. This follows from another

series of formal manipulations. This composite precomposed with the unit  $\mathbf{1} \rightarrow \nu \text{ko}$  vanishes for degree reasons, so the above composite factors through the cofibre of the unit, denoted by  $\nu \text{ko}/\mathbf{1}$ . As this unit induces an isomorphism on  $\pi_{n,*}$  for  $n \leq 2$  and a surjection on  $\pi_{3,*}$ , we see that  $\tau_{\leq 3}^\uparrow(\nu \text{ko}/\mathbf{1}) = 0$ . In particular, we see the factorisation of the above composite is adjunct to the zero map

$$0 = \tau_{\leq 3}^\uparrow \nu \text{ko}/\mathbf{1} \xrightarrow{\tau_{\leq 3}^\uparrow(\psi^3-1)} \Sigma^{3,3} \mathbf{F}_2[\tau]/\tau^2. \quad \square$$

Our synthetic lifts of  $j$  are then defined just as in the classical case, now using the vertical  $t$ -structure of Th.2.2.

**Definition 4.2.** Let  $E$  be either  $\mathbf{F}_p$  or  $\text{BP}$ . By Pr.4.1, we have maps of synthetic spectra

$$\psi^3 - 1: \nu \text{ko} \rightarrow \tau_{\geq 4}^\uparrow \nu \text{ko} \quad \psi^{p+1} - 1: \nu \ell \rightarrow \tau_{\geq 2p-2}^\uparrow \nu \ell$$

where  $p$  is an odd prime. Define  $j_E$  as the fibre of the above maps. As the above maps are zero on  $\pi_{0,0}$ , we obtain a map  $\mathbf{1} \rightarrow j_E$  which we call the *unit map*; in Pr.4.9 we will see these synthetic spectra have preferred  $\mathbf{E}_\infty$ -structures.

First, let us show  $j_E$  are indeed lifts of the spectra  $j$ .

**Proposition 4.3.** *For all primes  $p$ , the synthetic spectra  $j_{\mathbf{F}_p}$  and  $j_{\text{BP}}$  are  $\tau$ -complete synthetic lifts of the connective image-of- $J$  spectrum  $j$ .*

*Proof.* By Cors.2.15, 3.11 and 3.17, and the fact that  $\tau$ -inversion is exact, we see that  $j_E$  is  $\tau$ -complete, and the fibre sequences of synthetic spectra defining  $j_E$  are mapped precisely to the fibre sequences of spectra defining  $j$ .  $\square$

## 4.2 Synthetic homotopy groups of $j_E$

To highlight the usefulness of our definition of  $j_E$ , let us calculate  $\pi_{a,b}j_E$ —by Pr.4.3, these synthetic homotopy groups are the signature of a modified  $E$ -ASS for the spectrum  $j$ . The proofs of all of the following calculations are the same, so we will only prove the first two statements at the prime 2.

**Proposition 4.4.** *Write  $A$  for the graded ring  $\mathbf{Z}[\tau, h_0]/(\tau h_0 - 2)$ . For  $k \in \mathbf{Z}$  and  $0 \leq r \leq 7$ , there are isomorphisms of graded  $A$ -modules*

$$\pi_{8k+r,*}(j_{\mathbf{F}_2}) \simeq \begin{cases} A\{1\} & k = r = 0 \\ (A/h_0)\{P^{k-1}(h_1 h_3)\} & k > 0, r = 0 \\ (A/h_0)\{h_1^r\} & k = 0, r = 1, 2 \\ (A/h_0)\{P^{k-1}(h_1^2 h_3), P^k(h_1)\} & k > 0, r = 1 \\ (A/h_0)\{P^k(h_1^2)\} & k > 0, r = 2 \\ (A/(\tau h_0)^3)\{P^k(h_2)\} & k \geq 0, r = 3 \\ (A/(\tau h_0)^{\text{ord}_2(k+1)+4})\{\alpha_k\} & k \geq 0, r = 7 \\ 0 & \text{otherwise} \end{cases}$$

with bidegrees on multiplicative generators given by  $|h_i| = (2^i - 1, 1)$ ,  $|\alpha_k| = (7 + 8k, 5 + 4k)$ , and  $P^k(x) = |x| + (8k, 4k)$ .

The notation  $P^k(-)$  is supposed to suggest the existence of a “synthetic Adams periodicity operator”; see Rmk.6.7.

*Proof.* The computation follows by determining the effect of the map  $\psi^3 - 1 : \nu \mathbf{ko} \rightarrow \tau_{\geq 4}^{\uparrow} \nu \mathbf{ko}$  on bigraded homotopy groups. This is easy, as the source and target are  $\tau$ -torsion free, and we know the effect after inverting  $\tau$ ; we see that the map is zero in  $\pi_{8k+r,*}$  when  $r \neq 0, 4$ , and it is given by multiplication by  $8 = \tau^3 h_0^3$  when  $r = 4$  and multiplication by  $2^{\text{ord}_2(k)+4} = (\tau h_0)^{\text{ord}_2(k)+4}$  when  $r = 0$  and  $k > 0$ , each of these holding up to multiplication by an odd number, in particular a 2-adic unit. This finishes our calculations up to extensions—the rest of the proof is simply a resolution of a classical extension problem calculating  $\pi_d \mathbf{j}$  for  $d \equiv_8 1$ .

For degree reasons, the only possible nontrivial extensions in the long exact sequence computing  $\pi_{*,*} \mathbf{j}_{\mathbf{F}_2}$  appear in stems of the form  $8k + 1$ . For a contradiction, suppose this extension was not trivial so that after inverting  $\tau$  we have  $\pi_{8k+1} \mathbf{j} \simeq \mathbf{Z}/4\mathbf{Z}$ . A quick calculation from the long exact sequence on homotopy groups associated with  $\mathbf{S}/2$  would show that  $\pi_{8k+1} \mathbf{j}/2$  has 4-elements. However, the long exact sequence on homotopy groups associated with the cofibre sequence

$$\mathbf{j}/2 \rightarrow \mathbf{ko}/2 \xrightarrow{(\psi^3-1)/2} \tau_{\geq 4} \mathbf{ko}/2$$

produces the exact sequence

$$\pi_{8k+2} \mathbf{ko}/2 \xrightarrow{\psi^3-1} \pi_{8k+2} \tau_{\geq 4} \mathbf{ko}/2 \rightarrow \pi_{8k+1} \mathbf{j}/2 \rightarrow \pi_{8k+1} \mathbf{ko} \xrightarrow{\psi^3-1=0} \pi_{8k+1} \tau_{\geq 4} \mathbf{ko}. \quad (4.5)$$

If there is an isomorphism  $\pi_{8k+2} \mathbf{ko}/2 \simeq \mathbf{Z}/4\mathbf{Z}$  and  $\psi^3 - 1$  acts trivially on this group, then we are done. Indeed, as  $\pi_{8k+1} \mathbf{ko}/2 \simeq \mathbf{Z}/2\mathbf{Z}$ , which easily follows from the long exact sequence on homotopy groups, we see that  $\pi_{8k+1} \mathbf{j}/2$  must have 8 elements from (4.5), hence  $\pi_{8k+1} \mathbf{j}$  must be isomorphic to  $\mathbf{Z}/2\mathbf{Z} \oplus \mathbf{Z}/2\mathbf{Z}$ , as desired. To see that  $\pi_{8k+2} \mathbf{ko}/2 \simeq \mathbf{Z}/4\mathbf{Z}$ , we use 8-fold Bott periodicity, combined with the facts that  $\mathbf{S} \rightarrow \mathbf{ko}$  induces an isomorphism on  $\pi_s$  for  $s = 1, 2$  and the classical calculation  $\pi_2 \mathbf{S}/2 \simeq \mathbf{Z}/4\mathbf{Z}$ ; see Fig.15. To see that  $\psi^3 - 1$  acts trivially on  $\pi_{8k+2} \mathbf{ko}/2$ , it suffices to show that  $\psi^3$  acts by the identity. To see this, take a generator  $\bar{\beta}^k x$  of this group, where  $x$  is a generator of  $\pi_2 \mathbf{ko}/2 \simeq \mathbf{Z}/4\mathbf{Z}$ . Then  $\psi^3(\bar{\beta}^k x) = \psi^3(\bar{\beta})^k \psi^3(x)$  by multiplicativity, and as  $\psi^3(\bar{\beta}) = \bar{\beta}$ , we are reduced to the  $\pi_2$ -case again. However,  $\psi^3$  acts trivially on  $\pi_2 \mathbf{ko}/2$  as the unit  $\mathbf{S} \rightarrow \mathbf{ko}$  induces an isomorphism on  $\pi_s$  for  $s = 1, 2$ , and  $\psi^k$  commutes with this unit as a map of spectra, hence its reduction mod 2 also commutes with this isomorphism  $\pi_2 \mathbf{S}/2 \simeq \pi_2 \mathbf{ko}/2$ .  $\square$

In the classical ASS for  $\mathbf{j}$ , there are no such extension problems in  $\pi_{8k+1} \mathbf{j}$  to consider for degree reasons.

**Corollary 4.6.** *The MASS for  $\mathbf{j}$  associated with the  $\mathbf{F}_2$ -synthetic spectrum  $\mathbf{j}_{\mathbf{F}_2}$  has  $E_2$ -page given by  $\pi_{*,*}(\nu \mathbf{ko}/\tau) \oplus \pi_{*,*}(\tau_{\geq 4}^{\uparrow}(\nu \mathbf{ko})/\tau)[-1]$ . There are  $d_4$ 's leaving stems of the form  $8k + 4$  and  $d_{\text{ord}_2(k)+5}$ 's leaving the  $8k$  stem, as shown in Fig.3. Moreover, there are exotic  $\eta$ -extensions indicated by the orange lines of slope 4 in Fig.3.*

*Proof.* We see from Pr.4.4 that  $\psi^3 - 1$  induces the zero map on  $\pi_{*,*}(-/\tau)$  from which the claim about the  $E_2$  page follows. The differentials follow immediately from the  $\tau$ -torsion described in Pr.4.4. For the  $\eta$ -extensions, we need to use the modified ANSS for  $j$ , which is easy to describe from Pr.4.7; also see Fig.5. In particular, in this modified ANSS for  $j$  is clear that in  $\pi_{*,j}$  we have  $\eta P^k(\eta^2) = 4P^k(\nu)$ , where  $P^k(\eta^2)$  is the class in  $\pi_{8k+2j}$  detected by  $P^k(h_1^2)$  and  $P^k(\nu)$  the class in  $\pi_{8k+3j}$  detected by  $P^k(h_2)$ . This necessitates the exotic  $\eta$ -extensions of Fig.3.  $\square$

Then the  $E = \text{BP}$ -case at  $p = 2$ . Because of the  $\eta$ -towers in  $\pi_{*,*}\nu\text{ko}$ , it is difficult to describe  $\pi_{*,*}j_{\text{BP}}$  stem-by-stem. We make the following statement instead:

**Proposition 4.7.** *Let  $R$  be the ring  $\mathbf{Z}_2[\tau, \eta]/(2\eta, \tau^2\eta^4)$ , and for  $k \geq 0$ , let  $B_k$  be the ring  $\mathbf{Z}/2^{\text{ord}_2(k+1)+4}[\tau, \eta]/(2\eta, \tau^2\eta^3)$ , where  $|\eta| = (1, 1)$  and  $|\tau| = (0, -1)$ . Then there is an isomorphism of  $\mathbf{Z}_2[\tau]$ -modules*

$$\pi_{*,*}j_{\text{BP}} = R\{1, \theta_{3,5}, \nu\}/(2\theta_{3,5}, \tau^2\eta^3\theta_{3,5}, \eta\nu, 4\nu - \tau^2\eta^3) \oplus \bigoplus_{k \geq 0} B_k\{\alpha_k, x_k, y_k\}/(2x_k, \eta y_k, 4y_k - \tau^2\eta^2 x_k)$$

where  $|\theta_{3,5}| = (3, 5)$ ,  $|\nu| = (3, 1)$ ,  $|\alpha_k| = (8k + 7, 1)$ ,  $|x_k| = (8k + 9, 1)$ ,  $|y_k| = (8k + 11, 1)$ .

*Proof.* We again proceed by calculating the long exact sequence in bigraded homotopy groups associated with the fibre sequence defining  $j_{\text{BP}}$ . Here  $\pi_{*,*}\text{ko}_{\text{BP}}$  is no longer  $\tau$ -torsion free, so it is easier to calculate the effect of the ring map  $\psi^3 : \text{ko}_{\text{BP}} \rightarrow \text{ko}_{\text{BP}}$  by checking on the generators, use that  $\tau_{\geq 4}^{\uparrow} \text{ko}_{\text{BP}} \rightarrow \text{ko}_{\text{BP}}$  induces an injection on bigraded homotopy groups, and then calculate  $\psi^3 - 1$ . Since  $\eta$  and  $\tau$  are in the image of the unit map to  $\text{ko}_{\text{BP}}$ , we see that  $\psi^3(\eta) = \eta$  and  $\psi^3(\tau) = \tau$ . It follows from Pr.3.12 that  $\pi_{4,0}\text{ko}_{\text{BP}}$  and  $\pi_{8,0}\text{ko}_{\text{BP}}$  are  $\tau$ -torsion free, so we see that  $\psi^3(\alpha) = 9\alpha$  and  $\psi^3(\beta) = 81\beta$  since this holds after inverting  $\tau$ .

The calculation now follows from Pr.3.12; it is best to think of the calculation of these homotopy groups in terms of the spectral sequence they correspond to, which is depicted in Fig.5 (here the red dots come from  $\pi_{*,*}(\tau_{\geq 4}^{\uparrow}(\nu\text{ko})/\tau)$  via the boundary map, and the blue dots are sent to the corresponding classes in  $\pi_{*,*}(\nu\text{ko}/\tau)$ ). The only nontrivial point is the relations  $4\nu = \tau^2\eta^3$  and  $4y_k = \tau^2\eta^2 x_k$ , which follow from the fact that the corresponding stem must  $\tau$ -invert to known homotopy groups of  $j$  which are isomorphic to  $\mathbf{Z}/8$ .  $\square$

As per usual, the odd prime case for  $E = \mathbf{F}_p$  and  $\text{BP}$  are much easier; we omit proofs.

**Proposition 4.8.** *Let  $p$  be an odd prime and write  $q = 2p - 2$ . Write  $C$  for the graded ring  $\mathbf{Z}[\tau, h_0]/(h_0\tau - p)$  and  $D$  for the graded ring  $\mathbf{Z}[\tau]$ . For  $n \in \mathbf{Z}$  we have isomorphisms of  $C$ - and  $D$ -modules*

$$\pi_{n,*}(j_{\mathbf{F}_p}) \simeq \begin{cases} C\{1\} & n = 0 \\ C/h_0^{\text{ord}_p(n)+2}\{\alpha_{n/\mathbf{F}_p}\} & n \geq 0, n \equiv -1 \text{ modulo } q \\ 0 & \text{otherwise} \end{cases}$$

$$\pi_{n,*}(j_{\text{BP}}) \simeq \begin{cases} D\{1\} & n = 0 \\ D/p^{\text{ord}_p(n)+2}\{\alpha_{n/\text{BP}}\} & n \geq 0, n \equiv -1 \text{ modulo } q \\ 0 & \text{otherwise} \end{cases}$$

respectively, where  $|\alpha_{n/\mathbf{F}_p}| = (nq - 1, \frac{nq}{2} + 1)$  and  $|\alpha_{n/\text{BP}}| = (nq - 1, 1)$ .

Note that the calculation of the synthetic homotopy groups of  $\mathbf{j}_{\mathbf{F}_p}$  is only slightly more complicated for  $p = 2$ , in stark contrast to the classical situation, where the ASS for  $\mathbf{j}$  at odd primes is easy to produce, but that for  $p = 2$  has only recently appeared in the literature; see [BR22]. Comparing our calculation above to these calculations of Bruner–Rognes, we also see that  $\mathbf{j}_{\mathbf{F}_p}$  is **not** equivalent to  $\nu\mathbf{j}$ .

### 4.3 Multiplicative structure on $\mathbf{j}_E$

Using these synthetic homotopy groups, we can show  $\mathbf{j}_E$  can be given an  $\mathbf{E}_\infty$ -structure.

**Proposition 4.9.** *Let  $E$  be either  $\mathbf{F}_p$  or BP. Write  $\mathbf{j}'_E$  for the synthetic  $\mathbf{E}_\infty$ -ring defined by taking the homotopy fixed points of the  $\mathbf{Z}$ -action on  $\nu\mathbf{ku}$  generated by  $\psi^3$  at the prime 2, and the  $\mathbf{Z}$ -action on  $\nu\ell$  generated by  $\psi^{p+1}$  at odd primes. Then there is a Cartesian diagram of synthetic spectra*

$$\begin{array}{ccc} \mathbf{j}_E & \longrightarrow & \mathbf{j}'_E \\ \downarrow & & \downarrow \\ \tau_{\leq 2}^\uparrow \mathbf{1} & \longrightarrow & \tau_{\leq 2}^\uparrow \mathbf{j}'_E. \end{array}$$

*In particular, the lower-horizontal and right-vertical maps are maps of synthetic  $\mathbf{E}_\infty$ -rings, so  $\mathbf{j}_E$  can be endowed with the structure of a synthetic  $\mathbf{E}_\infty$ -ring.*

*Proof.* Suppose that  $p = 2$  for simplicity. One calculates the synthetic homotopy groups of  $\mathbf{j}'_E$  as it can be written as the fibre of the map  $\psi^3 - 1: \nu\mathbf{ko} \rightarrow \nu\mathbf{ko}$ . One can then calculate the synthetic homotopy groups of the actual pullback, which we will denote by  $P$ , using a Mayer–Vietoris sequence. In total, we find that the synthetic homotopy groups of  $P$  match those of  $\mathbf{j}_E$ . Moreover, at the prime 2, as the composite

$$P \rightarrow \mathbf{j}'_E \rightarrow \nu\mathbf{ko} \xrightarrow{\psi^3 - 1} \tau_{\geq 4}^\uparrow \nu\mathbf{ko}$$

is null by construction, we obtain a map of synthetic spectra  $\mathbf{j}_E \rightarrow P$  that recognises the above isomorphism on synthetic homotopy groups. The argument at odd primes follows *mutatis mutandis*.  $\square$

### 4.4 Periodic variants

Just as in §3, we can also define periodic versions of  $\mathbf{j}_E$ , denoted by  $\mathbf{J}_E$ , which behave both as synthetic lifts of the classical periodic image-of- $J$  spectrum and produce a nonzero modified  $E$ -ASS.

**Definition 4.10.** Let  $p$  be a prime and  $E = \mathbf{F}_p$  or BP. Then we define  $\mathbf{J}_E$  as the fibre of

$$\mathbf{KO}_E \xrightarrow{\psi^3 - 1} \mathbf{KO}_E \qquad \mathbf{L}_E \xrightarrow{\psi^{p+1} - 1} \mathbf{L}_E$$

for  $p = 2$  and odd primes, respectively.

The following two statements are proven just as Prs.4.3 and 4.4; we omit the proofs. Calculations of  $\pi_{*,*}\mathbf{J}_{\mathbf{F}_p}$  at other primes follow similarly.

**Proposition 4.11.** *The synthetic spectra  $\mathbf{J}_E$  are  $\tau$ -complete synthetic lifts of  $J$ .*

**Proposition 4.12.** *At the prime 2, the synthetic homotopy groups of  $\mathbf{J}_{\mathbf{F}_2}$  and  $\mathbf{J}_{\mathbf{BP}}$  are given by Figs.10 and 11 and Figs.12 and 13, respectively.*

Of course, the odd primary cases are simpler, so we will omit them here.

As  $J$  is equivalent to the  $K(1)$ -local sphere, we see that Figs.10 and 11 provide a modified  $\mathbf{F}_p$ -ASS for  $\mathbf{S}_{K(1)}$ . A similar conclusion can be made about a modified ANSS for  $\mathbf{S}_{K(1)}$  using the synthetic homotopy groups of  $\mathbf{J}_{\mathbf{BP}}$ . In this case, however, we claim that  $\nu J \simeq \mathbf{J}_{\mathbf{BP}}$ , so this simply recovers the usual ANSS for the  $K(1)$ -local sphere. This calculation is well-known, and a modern reference and diagram can be found in [Mor23, Fig.11].

## 5 Adams $v_1$ self-maps and detection for $\mathbf{ko}$

This section follows from the classical work of Adams [Ada66], however, the our proofs below will both act as a warm-up for §6 and allow us to solidify our notation.

### 5.1 Construction of $v_1$ self-maps

To show that the unit map  $\mathbf{S} \rightarrow j$  is surjective on  $\pi_{8k-1}$ , we will use Adams'  $v_1$  self-map  $v_1^4: \mathbf{S}/2[8] \rightarrow \mathbf{S}/2$ ; see [Ada66]. Let us reprove this classical result with the tools we have at hand—this will help us set up the following detection statements as well. Let us write  $u$  for the generator of  $\pi_2 \mathbf{ku}/2$ .

**Theorem 5.1** ([Ada66]). *There is a map of spectra  $v_1^4: \mathbf{S}/2[8] \rightarrow \mathbf{S}/2$  inducing multiplication by  $u^4$  on  $\mathbf{ku}$ -homology.*

*Proof.* We grant ourselves an Adams chart and an Adams–Novikov chart for  $\mathbf{S}$  at the prime 2 in stems  $\leq 8$ ; for example, see [Rav78, p.412 & 429], respectively. By [Pst23, Lm.4.23], one has a cofibre sequence in  $\text{Syn}_{\mathbf{F}_2}$

$$\mathbf{S}^{0,1} \xrightarrow{h_0} \nu(\mathbf{S}) \rightarrow \nu(\mathbf{S}/2) \rightarrow \nu(\mathbf{S}^1) \xrightarrow{h_0} \mathbf{S}^{1,0}$$

from which an Adams chart for  $\mathbf{S}/2$  in stems  $\leq 8$  easily follows; see Fig.14. Similarly, one has a cofibre sequence in  $\text{Syn}_{\mathbf{BP}}$

$$\nu(\mathbf{S}) \xrightarrow{2} \nu(\mathbf{S}) \rightarrow \nu(\mathbf{S}/2) \rightarrow \mathbf{S}^{1,-1} \xrightarrow{2} \mathbf{S}^{1,-1}$$

which yields an Adams–Novikov chart for  $\mathbf{S}/2$  in stems  $\leq 8$ ; see Figs.15 and 16.

From Fig.14, we see that  $2 \cdot \pi_8(\mathbf{S}/2) = 0$ . In fact, the  $h_0$ -multiplications therein show that this could only fail to happen if 2 times the class detected in filtration 2 is detected by the

class in filtration 4. However, the long exact sequence associated with the cofibre sequence in  $\text{Syn}_{\mathbf{F}_2}$  above shows that the class in filtration 2 is in the image of the map  $\pi_8 \mathbf{S} \rightarrow \pi_8 \mathbf{S}/2$ , and  $2 \cdot \pi_8 \mathbf{S} = 0$ ; this fact is clear from the  $h_0$ -multiplications in the Adams chart for  $\mathbf{S}$ . It follows that any map  $\mathbf{S}^8 \rightarrow \mathbf{S}/2$  factors over  $\mathbf{S}/2[8]$ .

It remains to use the Adams–Novikov chart (Figs.15 and 16) to show that there is some class  $v_1^4 \in \pi_8 \mathbf{S}/2$  which is sent to  $u^4 \in \pi_8 \text{ku}/2$ . There is such a class on the  $E_2$ -page of the ANSS for  $\mathbf{S}/2$  from the calculation

$$\text{Ext}_{\text{BP}_* \text{BP}}^{0,*}(\text{BP}_*, \text{BP}_*/2) = \mathbf{F}_2[\tilde{v}_1];$$

see [Rav04, Th.4.3.2]. Let us check that the class  $\tilde{v}_1^4$  here is a permanent cycle, but this follows for degree reasons. Indeed, the first differential  $v_1^4$  could support would be a  $d_3$ , but the only class in bidegree  $(7, 3)$  supports a  $d_3$  and thus can't be a boundary. Everything in higher filtration vanishes on the  $E_4$ -page.

It remains to show that  $\tilde{v}_1^4$  is sent to  $u^4 \in \pi_8 \text{ku}/2$ . As  $\text{ku}/2$  is complex oriented, then the Ext-group

$$\text{Ext}_{\text{BP}_* \text{BP}}^{0,8}(\text{BP}_*, \text{BP}_* \text{ku}/2) \simeq \mathbf{F}_2$$

generated by a class that detects  $u^4$ . The image of  $\tilde{v}_1^4$  under the left unit  $\bar{\eta}_L: \text{BP}_*/2 \rightarrow \text{BP}_* \text{ku}/2$  is precisely  $\tilde{v}_1^4$  by construction. This agrees with its image under the right unit  $\bar{\eta}_R$ , as in general  $v_n \equiv \eta_R(v_n)$  modulo  $I_n$ ; see [Rav04, (4.3.1)]. Hence  $\tilde{v}_1^4$  is sent to  $u^4$  in  $\pi_8 \text{ku}/2$ .  $\square$

*Remark 5.2.* The generator  $\tilde{v}_1$  of  $\pi_2 \mathbf{S}/2$  is also detected in  $\pi_2 \text{ku}/2$ . In fact, copying the argument above that  $\tilde{v}_1^4$  hits  $u^4$ , we see that the image of  $\tilde{v}_1$  is precisely  $u$ . However, this does not factor into a self-map of  $\mathbf{S}/2$  as this class  $\tilde{v}_1$  is **not** 2-torsion.

Using the notation from the above proof, we obtain an infinite family of elements  $\partial \tilde{v}_1^{4k} \in \pi_{8k-1} \mathbf{S}$  as the composite

$$\mathbf{S}^{8k} \rightarrow \mathbf{S}^{8k}/2 \xrightarrow{v_1^{4k}} \mathbf{S}/2 \xrightarrow{\partial} \mathbf{S}^1. \quad (5.3)$$

These elements are nonzero as they project to nonzero classes mod  $\tau$  in filtration 1; this is because the boundary map

$$\mathbf{F}_2[\tilde{v}_1] = \pi_{*,0} \nu \mathbf{S}/2, \tau \rightarrow \pi_{*-1,1} \nu \mathbf{S}/\tau$$

is an injection when  $* > 0$ , as  $\pi_{*,0} \nu \mathbf{S} = \text{Ext}_{\text{BP}_* \text{BP}}^{0,*}(\text{BP}_*, \text{BP}_*) = 0$  for  $* > 0$ . The class  $\partial \tilde{v}_1^{4k}$  is then detected by a nonzero class mod  $\tau$ , and it cannot be hit by a differential because it is in filtration 1. The work of Adams [Ada66] and Quillen [Qui71] show this 2-torsion element is divisible by  $2^{\text{ord}_2(k)+2}$ . Let us write  $\alpha_k$  for any element in  $\pi_{8k-1} \mathbf{S}$  such that  $2^{\text{ord}_2(k)+2} \alpha_k = \partial \tilde{v}_1^{4k}$ .

The odd primary case is simpler. Let  $p$  be an odd prime, and write  $q = 2p - 2$  and  $v$  for the generator of  $\pi_q \ell/p$ .

**Theorem 5.4** ([Ada66]). *For an odd prime  $p$ , there is a map of spectra  $v_1: \mathbf{S}/p[q] \rightarrow \mathbf{S}/p$  inducing multiplication by  $v$  on  $\ell$ -homology.*

*Proof.* The proof is similar to Th.5.1, however, many steps are simplified. Indeed, the cofibre sequence  $\mathbf{S} \xrightarrow{p} \mathbf{S} \rightarrow \mathbf{S}/p$  can be used to show that  $\pi_{2p-2}\mathbf{S}/p \simeq \mathbf{Z}/p\mathbf{Z}$ . We write  $\tilde{v}_1$  for a generator of this group. To see this class is sent to  $v \in \pi_q\ell$ , we make the same arguments with the ANSS for  $\mathbf{S}$  using the left and right units.  $\square$

## 5.2 The Hurewicz image of $\mathbf{ko}$

**Theorem 5.5** ([Ada66]). *The unit  $f: \mathbf{S} \rightarrow \mathbf{ko}$  is surjective on  $\pi_{8k+r}$  for all integers  $k$  and  $r = 1, 2$ .*

We will use synthetic spectra in the following proof, both to act as a warm-up for §6 and to highlight that although such results can also be obtained using the classical ASS or ANSS for  $\mathbf{ko}$ , we have still avoided calculations of the comodules  $H_*(\mathbf{ko}; \mathbf{F}_2)$  and  $\mathbf{BP}_*\mathbf{ko}$ , as well as their associated Ext-groups.

*Proof.* There is nothing to show for  $k < 0$  and  $k = 0$  we are also done by Wood's theorem as discussed in §3.2. Let us then set  $k > 0$  and  $r = 1$  and consider the following commutative diagram of BP-synthetic homotopy groups

$$\begin{array}{ccccc}
\pi_{8k+2,0}(\nu\mathbf{S}/\tau) & \longrightarrow & \pi_{8k+2,0}((\nu\mathbf{S})/2, \tau) & \xrightarrow{\partial} & \pi_{8k+1,1}(\nu\mathbf{S}/\tau) \\
\downarrow & & \downarrow & & \downarrow \\
\pi_{8k+2,0}(\nu\mathbf{ko}/\tau) & \longrightarrow & \pi_{8k+2,0}((\nu\mathbf{ko})/2, \tau) & \xrightarrow{\partial} & \pi_{8k+1,1}(\nu\mathbf{ko}/\tau) \\
\downarrow & & \downarrow & & \downarrow \\
\pi_{8k+2,0}(\nu\mathbf{ku}/\tau) & \longrightarrow & \pi_{8k+2,0}((\nu\mathbf{ku})/2, \tau) & \xrightarrow{\partial} & \pi_{8k+1,1}(\nu\mathbf{ku}/\tau) = 0
\end{array} \tag{5.6}$$

from the maps of  $\mathbf{E}_\infty$ -rings  $\mathbf{S} \rightarrow \mathbf{ko} \rightarrow \mathbf{ku}$  tensored with the cone of  $\tau$  and the defining cofibre sequence for  $\mathbf{S}/2$ .

Let us write  $\tilde{v}_1 \in \pi_2\mathbf{S}/2 \simeq \mathbf{Z}/4\mathbf{Z}$  for a generator, see Fig.15, which is detected in  $\pi_{2,0}$  of  $(\nu\mathbf{S})/(2, \tau)$  by a generator which we also denote by  $\tilde{v}_1$ ; see Rmk.5.2. Let us write  $x \in \pi_{8k+2}\mathbf{S}/2$  for the element  $v_1^{4k}\tilde{\nu}$ , and  $\mu = \partial(x) \in \pi_{8k+1+c}\mathbf{S}$  for the image of  $x$  under the boundary map for the cofibre sequence defining  $\mathbf{S}/2$ . From the proof of Th.5.1, we see that the class  $x$  is detected by an element  $v_1^{4k+1}$  inside  $\pi_{8k+2,0}((\nu\mathbf{S})/2, \tau)$ .

We claim that the  $f(\mu)$  is nonzero in the homotopy groups of  $\mathbf{ko}$ . To see this, notice that the class  $v_1^{4k+1}$  defined above is nonzero, as its image in  $\pi_{8k+2,0}((\nu\mathbf{ku})/2, \tau)$  is precisely the class  $\bar{u}^{4k+1}$ , where  $\bar{u}$  is the only class on the  $E_2$ -page of the  $\Gamma$ -SS for  $\nu\mathbf{ku}/2$  which could detect  $u$ . Indeed, this follows from the fact that the generator  $\tilde{v}_1$  of  $\pi_2\mathbf{S}/2 \simeq \mathbf{Z}/4\mathbf{Z}$  is sent to  $u$ , which is precisely Rmk.5.2. As the unit  $\mathbf{1} \rightarrow \nu\mathbf{ku}$  factors through  $f$ , we see that  $f(v_1^{4k+1})$  is also nonzero. As  $\pi_{8k+2,0}(\nu\mathbf{ko}/\tau) = 0$ , see Fig.5, we see that  $\partial f(v_1^{4k+1}) = f(\partial(v_1^{4k+1}))$  must be nonzero in  $\pi_{8k+1,1}(\nu\mathbf{ko})/\tau$ . We now claim that this nonzero element on the  $E_2$ -page of

the  $\Gamma$ -SS from  $\nu \text{ko}$  is a permanent cycle. This follows from the fact that we have an explicit geometric representative for this class,  $f(\mu) = f(\partial(x))$ . Therefore,  $f(\mu)$  is nonzero.

The case for  $r = 2$  follows from the above argument by multiplication by  $\eta$ .  $\square$

Notice that our arguments above actually show that  $\mu \in \pi_{8k+1}\mathbf{S}$  and  $\mu\eta \in \pi_{8k+2}\mathbf{S}$  are also both nonzero.

## 6 Detection results for $j$

Extending the ideas used in the proof of Th.5.5 from  $\nu \text{ko}$  to  $j_E$ , we will prove Th.A that the unit map  $\mathbf{S} \rightarrow j$  induces a surjection on homotopy groups.

### 6.1 Hurewicz image of $j$ at $p = 2$

Implicitly complete all spectra at the prime 2.

**Proposition 6.1.** *The unit map  $\mathbf{S} \rightarrow j$  induces an isomorphism on  $\pi_k$  for  $k \leq 3$ .*

*Proof.* The orders of the homotopy groups in question match, so it suffices to show that  $j$  detects both  $\eta \in \pi_1\mathbf{S}$  and  $\nu \in \pi_3\mathbf{S}$ . By assumption, we know that  $\eta$  is detected in  $\pi_1 \text{ko}$ , and since the unit  $\mathbf{S} \rightarrow \text{ko}$  factors through  $j$ , we see  $j$  also detects  $\eta$ . For  $\nu$ , we look at  $j_{\text{BP}}$ ; see Fig.6. In particular, the 2-extension from  $2h_2$  to  $h_1^3$ , comes from the fact that we know  $\pi_3 j \simeq \mathbf{Z}/8\mathbf{Z}$ . As  $\eta^3$  lies in  $\pi_3 j$  and  $\eta^3 = 4\nu$  in  $\pi_3\mathbf{S}$ , we see that the image of  $\nu$  on  $\pi_3 j$  must be nonzero.  $\square$

**Proposition 6.2.** *The unit map  $\mathbf{S} \rightarrow j$  induces a surjection on  $\pi_{8k-1}$  for all integers  $k$ .*

*Proof.* The case of  $k \leq 0$  is uninteresting, so let  $k > 0$ . Recall that  $\alpha_k$  is any element in  $\pi_{8k-1}\mathbf{S}$  such that  $2^{\text{ord}_2(k)+2}\alpha_k = \partial\tilde{v}_1^{4k}$  using the notation from the proof of Th.5.1. As the 2-power-torsion of these classes  $\alpha_k$  is precisely the size of the cyclic group  $\pi_{8k-1}j$ , which we obtained purely from the action of  $\psi^3 - 1$  on the homotopy groups of  $\text{ko}$ , the surjectivity of  $\mathbf{S} \rightarrow j$  will follow if the image of  $\partial\tilde{v}_1^{4k}$  are nonzero in  $\pi_*j$ .

Consider the following commutative diagram of synthetic homotopy groups, induced from the maps of synthetic spectra  $\nu\mathbf{S} \rightarrow j_{\text{BP}} \rightarrow \text{ku}$  tensored with the cofibre sequence defining  $(\nu\mathbf{S})/2$  and  $C(\tau)$ :

$$\begin{array}{ccccc}
\pi_{8k,0}(\nu\mathbf{S}/\tau) & \longrightarrow & \pi_{8k,0}((\nu\mathbf{S})/2, \tau) & \xrightarrow{\partial} & \pi_{8k-1,1}(\nu\mathbf{S}/\tau) \\
\downarrow & & \downarrow & & \downarrow \\
\pi_{8k,0}(j_{\text{BP}}/\tau) & \longrightarrow & \pi_{8k,0}(j_{\text{BP}}/2, \tau) & \xrightarrow{\partial} & \pi_{8k-1,1}(j_{\text{BP}}/\tau) \\
\downarrow & & \downarrow & & \downarrow \\
\pi_{8k,0}(\nu \text{ku}/\tau) & \longrightarrow & \pi_{8k,0}((\nu \text{ku})/2, \tau) & \xrightarrow{\partial} & \pi_{8k-1,1}(\nu \text{ku}/\tau) = 0
\end{array}$$

We claim, that the image of  $\partial\tilde{v}_1^{4k}$  inside  $\pi_{8k-1,1}(\mathbf{jBP}/2, \tau)$  is nonzero. Indeed, by construction,  $\tilde{v}_1^{4k}$  is nonzero in  $\pi_{8k,0}((\nu\mathbf{ku})/2, \tau)$ , hence this class is also nonzero in  $\pi_{8k,0}(\mathbf{jBP}/2, \tau)$ . It then follows that  $\partial\tilde{v}_1^{4k}$  is nonzero in  $\pi_{8k-1,1}(\mathbf{jBP}/2, \tau)$  as we can easily calculate  $\pi_{8k,0}(\mathbf{jBP}/\tau) = 0$  by construction; see Pr.4.7 and Fig.6. This shows there is a nonzero class in the  $E_2$ -page of a modified ANSS for  $\mathbf{j}$  detecting the desired element from  $\pi_{8k-1}\mathbf{S}$ , and again, this class is a permanent cycle as it has an explicit geometric representation; simply post-compose Equation (5.3) with the unit  $\mathbf{S} \rightarrow \mathbf{j}$ .  $\square$

**Proposition 6.3.** *The unit map  $\mathbf{S} \rightarrow \mathbf{j}$  induces a surjection on  $\pi_{8k}$  for all  $k \in \mathbf{Z}$ .*

*Proof.* For  $k \leq 0$ , this is uninteresting or covered by Pr.6.1, so let  $k \geq 1$ . The class generating  $\pi_{8k-1}\mathbf{j}$  supports multiplication by  $\eta$ . Therefore, the generator of  $\pi_{8k}\mathbf{j} \simeq \mathbf{Z}/2\mathbf{Z}$  is hit by the element  $\eta\alpha_k$  from the sphere—again, this also proves that  $\eta\alpha_k$  is nonzero.  $\square$

In degrees  $8k + r$  for  $r = 1, 2$ , we can deduce the surjectivity of  $\mathbf{S} \rightarrow \mathbf{j}$  using  $\mathbf{ko}$ .

**Proposition 6.4.** *The unit map  $\mathbf{S} \rightarrow \mathbf{j}$  induces a surjection on  $\pi_{8k+r}$  for all  $k \in \mathbf{Z}$  and  $r = 1, 2$ .*

*Proof.* By construction, the unit map  $\mathbf{S} \rightarrow \mathbf{ko}$  factors through  $\mathbf{j}$ . By Th.5.5, we see that the blue classes in our modified ASS for  $\mathbf{j}$ , see Fig.4, are hit by elements from the sphere. Combining this with Pr.6.3, we know the red class in  $\pi_{8k+1}\mathbf{j}$  is hit by  $\eta\alpha_k$ , which gives us our desired surjectivity.  $\square$

**Proposition 6.5.** *The unit map  $\mathbf{S} \rightarrow \mathbf{j}$  induces a surjection on  $\pi_{8k+3}$  for all  $k \in \mathbf{Z}$ .*

*Proof.* The orange extensions by  $\eta$  in our modified ASS for  $\mathbf{j}$  (Fig.4) combined with Pr.6.1 and Pr.6.4 show that  $\mathbf{j}$  detects  $\eta^3\alpha_k$ . This implies that  $\mathbf{j}$  also detects  $\nu\alpha_k$ , as we have  $4\nu = \eta^3$  in the homotopy groups of both  $\mathbf{S}$  and  $\mathbf{j}$ .  $\square$

## 6.2 Hurewicz image of $\mathbf{j}$ at odd primes

The odd primary case can be proven in one go.

**Proposition 6.6.** *For an odd prime  $p$ , the unit map  $\mathbf{S} \rightarrow \mathbf{j}$  induces a surjection on homotopy groups.*

*Proof.* The proof is similar to Pr.6.2, so we omit some details. First, we note that  $\pi_n\mathbf{j} = 0$  unless  $n = 0$  or  $n = kq - 1$  for some  $k \geq 1$ . The key step is now to show that  $\partial v_1^k$  is detected in  $\pi_{kq-1}\mathbf{j}$ . To see this, we use the commutative diagram of synthetic homotopy groups

$$\begin{array}{ccccc}
\pi_{kq,0}(\nu\mathbf{S}/\tau) & \longrightarrow & \pi_{kq,0}((\nu\mathbf{S})/p, \tau) & \xrightarrow{\partial} & \pi_{kq-1,1}(\nu\mathbf{S}/\tau) \\
\downarrow & & \downarrow & & \downarrow \\
\pi_{kq,0}(\mathbf{jBP}/\tau) & \longrightarrow & \pi_{kq,0}(\mathbf{jBP}/p, \tau) & \xrightarrow{\partial} & \pi_{kq-1,1}(\mathbf{jBP}/\tau) \\
\downarrow & & \downarrow & & \downarrow \\
\pi_{kq,0}(\nu\mathbf{ku}/\tau) & \longrightarrow & \pi_{kq,0}((\nu\mathbf{ku})/p, \tau) & \xrightarrow{\partial} & \pi_{kq-1,1}(\nu\mathbf{ku}/\tau) = 0
\end{array}$$

and the fact that  $\pi_{a,b}j_{\text{BP}}/\tau$  is supported in bidegrees  $a = kq - 1$  and  $b = 1$  by construction; see Pr.4.8.  $\square$

Combining these results yields our main theorem.

*Proof of Th.A.* At odd primes this is Pr.6.6, and at the prime 2 combine Prs.6.2 to 6.5 with the fact that  $\pi_{8k+r}j = 0$  for  $r \in \{4, 5, 6\}$ .  $\square$

*Remark 6.7.* Much of the surjectivity of  $\mathbf{S} \rightarrow j$  is concluded from the classical  $\mathbf{F}_2$ -ASS for  $j$  using Adams periodicity operator  $P = \langle 16, \sigma, - \rangle$ . This approach can also be applied here, using the synthetic Toda bracket  $\langle h_0^4, h_3, - \rangle$ , where  $h_3$  is considered as an element in  $\pi_{7,1}\mathbf{1}$ , rather than in  $\pi_{7,5}j_{\mathbf{F}_2}$ . This is the reason we named the classes in Pr.4.4 with  $P^k(-)$  decorations. This operator also produces an alternative proof to many of the statements leading to our proof of Th.A. We have chosen to give the BP-synthetic proofs here instead as they are simple arguments with long exact sequences.

### 6.3 Split surjectivity on filtered homotopy groups

These detection statements imply a split surjectivity statement for our modified ANSS for  $j$ .

**Corollary 6.8.** *Let  $p$  be a prime. The unit map  $\mathbf{1} \rightarrow j_{\text{BP}}$  induces a split surjection on  $E_\infty$ -pages of  $\Gamma$ -SSs.*

*Proof.* By Th.A, we see the associated map is surjective. Combined with the calculation of the  $\alpha$ -family in the ANSS for  $\mathbf{S}$ , see [Rav04, Th.5.3.7], we see that the  $E_\infty$ -page of the  $\Gamma$ -SS for  $j_{\text{BP}}$  detects exactly the  $\alpha$ -family in  $\mathbf{S}$ , a split summand of the  $E_\infty$ -page of the ANSS for  $\mathbf{S}$ .  $\square$

*Remark 6.9.* Although  $\mathbf{1} \rightarrow j_{\text{BP}}$  induces a surjection on synthetic homotopy groups, this map is not a split surjection. Indeed, notice that  $\pi_{3,5}\mathbf{1} = 0$ , however,  $\pi_{3,5}j \neq 0$  as it contains  $\tau$ -torsion classes; this is evident from Pr.4.7 and Fig.5. One might try to clear out some of these  $\tau$ -torsion classes by using something such as  $\tau_{\geq 4}^L \text{ko}$  where  $L$  is defined by the equation  $x = y$ , however, then the line of slope 1 of red classes supporting the differentials which hit the line we just cut out would survive and not be hit by synthetic unit.

On the other hand, it is easy to see that  $\mathbf{1} \rightarrow j_{\mathbf{F}_2}$  cannot be split surjective on synthetic homotopy groups for degree reasons.

*Remark 6.10.* In [Ada66], Adams shows that the image of the  $J$ -homomorphism is a split summand of  $\pi_*\mathbf{S}$ . In fact, more is shown, as it is further shown that the  $\mu$ -family is also a split summand of  $\pi_*\mathbf{S}$ . Adams uses the *e-invariant* to construct a splitting of the image of the  $J$ -homomorphism and the  $\mu$ -family, and according to [Koc90, §4.1], Mahowald first defined the connective image-of- $J$  spectrum  $j$  in [Mah75] such that the unit map  $\mathbf{S} \rightarrow j$  lifts this *e-invariant*. To be explicit about Adams splitting now, let us write  $i: \pi_*j \rightarrow \pi_*\mathbf{S}$  for the Adams map of graded abelian groups which splits the surjective unit map  $\pi_*\mathbf{S} \rightarrow \pi_*j$ .

From these considerations, Th.C follows rather formally from an algebraic fact. Recall that for a filtered abelian group

$$G = G_0 \supseteq G_1 \supseteq \cdots$$

with  $\bigcap G_i = \{0\}$  there is a function  $\pi_G: G \rightarrow \text{gr}G$  from the filtered abelian group to its associated graded abelian groups, which sends  $0 \mapsto 0$  and for  $0 \neq x \in G_i \setminus G_{i+1}$ ,  $x$  is sent to  $[x] \in G_i/G_{i+1}$ . For two such filtered abelian groups  $H, G$ , given a map of graded abelian groups  $j: \text{gr}H \rightarrow \text{gr}G$ , we say a homomorphism of abelian groups  $i: H \rightarrow G$  is a *lift* of  $j$  if the following diagram of sets commutes:

$$\begin{array}{ccc} H & \xrightarrow{i} & G \\ \downarrow \pi_H & & \downarrow \pi_G \\ \text{gr}H & \xrightarrow{j} & \text{gr}G \end{array} \quad (6.11)$$

**Lemma 6.12.** *With notation as above, if  $i$  is a lift of  $j$ , then  $i$  is a map of filtered abelian groups and  $\text{gr}(i) = j$ .*

*Proof.* It suffices to show that if  $0 \neq x \in H_i \setminus H_{i+1}$ , then  $i(x) \in G_i$ . This follows immediately from the definition of  $\pi_G$ , since the diagram (6.11) commutes.  $\square$

*Proof of Th.C.* The unit map  $\pi_*\mathbf{S} \rightarrow \pi_*j$  is a morphism of filtered abelian groups where  $\pi_*\mathbf{S}$  and  $\pi_*j$  have filtrations defined by  $\Gamma$ -SS of the BP-synthetic spectra  $\mathbf{1}$  and  $j_{\text{BP}}$ . It suffices to show that the classical splitting map  $i$  of Rmk.6.10 is a morphism of filtered abelian groups. To this end, we apply Lm.6.12, where  $i$  to be the splitting of Rmk.6.10 and  $j$  as the splitting on  $E_\infty$ -pages from Cor.6.8. One checks directly that the diagram (6.11) commutes: if  $i(x) = y$ , then  $j$  sends the element on  $E_\infty$  detecting  $x$  to the element detecting  $y$ .  $\square$

## References

- [Ada66] John Frank Adams. On the groups  $J(X)$ . (IV). *Topology*, 5:21–71, 1966.
- [AKQ20] Gabriel Angelini-Knoll and J. D. Quigley. Chromatic complexity of the algebraic  $k$ -theory of  $y(n)$ , 2020.
- [BHHM08] M. Behrens, M. Hill, M. J. Hopkins, and M. Mahowald. On the existence of a  $v_2^{32}$ -self map on  $M(1, 4)$  at the prime 2. *Homology Homotopy Appl.*, 10(3):45–84, 2008.
- [BHS19] Robert Burklund, Jeremy Hahn, and Andrew Senger. On the boundaries of highly connected, almost closed manifolds, 2019.
- [BHS22] Robert Burklund, Jeremy Hahn, and Andrew Senger. Galois reconstruction of artin-tate  $\mathbb{R}$ -motivic spectra, 2022.

- [Boa99] J. Michael Boardman. Conditionally convergent spectral sequences. In *Homotopy invariant algebraic structures. A conference in honor of J. Michael Boardman. AMS special session on homotopy theory, Baltimore, MD, USA, January 7–10, 1998*, pages 49–84. Providence, RI: American Mathematical Society, 1999.
- [BR22] Robert R. Bruner and John Rognes. The Adams spectral sequence for the image-of- $J$  spectrum. *Trans. Am. Math. Soc.*, 375(8):5803–5827, 2022.
- [CDN24] Christian Carrick, Jack Morgan Davies, and Sven van Nigtevecht. Descent spectral sequences through synthetic spectra. In preparation, 2024.
- [Dav75] Donald M. Davis. The cohomology of the spectrum  $bJ$ . *Bol. Soc. Mat. Mex., II. Ser.*, 20:6–11, 1975.
- [Dav20] Jack Morgan Davies. On Lurie’s theorem and applications, 2020. arXiv preprint, arXiv:2007.00482.
- [Hed20] Alice Petronella Hedenlund. *Multiplicative Tate spectral sequences*. PhD thesis, University of Oslo, Oslo, Norway, 2020.
- [Koc90] Stanley O. Kochman. *Stable homotopy groups of spheres. A computer-assisted approach*, volume 1423 of *Lect. Notes Math.* Berlin etc.: Springer-Verlag, 1990.
- [LL23] David Lee and Ishan Levi. Topological hochschild homology of the image of  $j$ . arXiv preprint arXiv:2307.04248, July 2023.
- [Lur09] Jacob Lurie. *Higher topos theory*, volume 170. Princeton, NJ: Princeton University Press, 2009.
- [Lur17] Jacob Lurie. Higher algebra. Available at <https://www.math.ias.edu/~lurie/>, 2017.
- [Lur18] Jacob Lurie. Elliptic Cohomology II: Orientations. Available at <https://www.math.ias.edu/~lurie/>, April 2018 version, 2018.
- [Mah75] Mark Mahowald. Description homotopy of the elements in the image of the  $J$ -homomorphism. *Manifolds, Proc. int. Conf. Manifolds relat. Top. Topol.*, Tokyo 1973, 255-263 (1975)., 1975.
- [Mat16] Akhil Mathew. The homology of  $tmf$ . *Homology Homotopy Appl.*, 18(2):1–29, 2016.
- [Mor23] Itamar Mor. Picard and brauer groups of  $k(n)$ -local spectra via profinite galois descent, 2023.
- [Pst23] Piotr Pstragowski. Synthetic spectra and the cellular motivic category. *Invent. Math.*, 232(2):553–681, 2023.
- [Qui71] D. Quillen. The Adams conjecture. *Topology*, 10:67–80, 1971.

- [Rav78] Douglas C. Ravenel. A novice's guide to the Adams-Novikov spectral sequence. *Geom. Appl. Homotopy Theory II, Proc. Conf., Evanston 1977, Lect. Notes Math.* 658, 404-475 (1978)., 1978.
- [Rav04] Douglas C. Ravenel. *Complex cobordism and stable homotopy groups of spheres. 2nd ed.* Providence, RI: AMS Chelsea Publishing, 2nd ed. edition, 2004.

## A Spectral sequence diagrams

The spectral sequences diagrams in this section use the following conventions:

- Each dot represents a single copy of  $\mathbf{F}_p$ . Squares represent a copy of the integers  $\mathbf{Z}$ .
- If the spectral sequence is the  $\Gamma$ -SS of a synthetic spectrum  $Y$ , which itself sits in an evident cofibre sequence of synthetic spectra  $X \rightarrow Y \rightarrow Z$ , then we write in blue those classes detected in  $Z$ , and in red those classes detected in  $X$ . Examples of this include the image-of- $J$  spectra  $j_E$  and  $J_E$  as well as the mod 2 Moore spectrum  $\mathbf{S}/2$ .
- Orange lines indicate exotic extensions.




Figure 1: ASS for  $ko$  at the prime 2 in the range  $0 \leq s \leq 20$ .




Figure 2: ANSS for  $ko$  at the prime  $2$  in the range  $0 \leq s \leq 20$ .




Figure 3: MASS for  $j$  at the prime  $2$  (from  $j_{\mathbb{F}_2}$ ) in the range  $0 \leq s \leq 36$  with all differentials.




Figure 4: MASS for  $j$  at the prime 2 (from  $j_{\mathbf{F}_2}$ ) in the range  $0 \leq s \leq 36$  on the  $E_\infty$ -page.




Figure 5: MANSS for  $j$  at the prime 2 (from  $j_{BP}$ ) in the range  $0 \leq s \leq 36$  with all differentials.




Figure 6: MANSS for  $j$  at the prime 2 (from  $j_{BP}$ ) in the range  $0 \leq s \leq 36$  on the  $E_\infty$ -page.




Figure 7: MASS for  $j$  at the prime 3 (from  $j_{\mathbf{F}_3}$ ) in the range  $0 \leq s \leq 36$  with all differentials.




Figure 8: MASS for  $j$  at the prime 3 (from  $j_{\mathbf{F}_3}$ ) in the range  $0 \leq s \leq 36$  on the  $E_\infty$ -page.




Figure 9: MANSS for  $j$  at the prime 3 (from  $j_{BP}$ ) in the range  $0 \leq s \leq 36$ .




Figure 10: MASS for  $J$  at the prime 2 (from  $J_{\mathbf{F}_2}$ ) in the range  $-18 \leq s \leq 18$  with all differentials.




Figure 11: MASS for  $J$  at the prime 2 (from  $J_{\mathbf{F}_2}$ ) in the range  $-18 \leq s \leq 18$  on the  $E_\infty$ -page.




Figure 12: MANSS for  $J$  at the prime 2 (from  $J_{\mathbf{BP}}$ ) in the range  $-18 \leq s \leq 18$  with all differentials.




Figure 13: MASS for  $J$  at the prime 2 (from  $J_{\mathbf{BP}}$ ) in the range  $-18 \leq s \leq 18$  on the  $E_\infty$ -page.




Figure 14: ASS for  $\mathbf{S}/2$  at the prime 2 in the range  $0 \leq s \leq 8$ .




Figure 15: ANSS for  $\mathbf{S}/2$  at the prime 2 in the range  $0 \leq s \leq 8$  with all differentials.




Figure 16: ANSS for  $\mathbf{S}/2$  at the prime 2 in the range  $0 \leq s \leq 8$  on the  $E_\infty$ -page.