arXiv:2402.01210v1 [astro-ph.CO] 2 Feb 2024

Big Bang Nucleosynthesis constraints on the Energy-Momentum Squared Gravity: The \mathbb{T}^2 model

Dukjae Jang,^{1,*} Mayukh R. Gangopadhyay,^{2,†} Myung-Ki Cheoun,^{3,‡} Toshitaka Kajino,^{4,§} and M. Sami^{5,¶}

¹School of Physics and International Research Center for Big-Bang

Cosmology and Element Genesis, Beihang University, Beijing 100083, China

²Centre For Cosmology and Science Popularization (CCSP),

SGT University, Gurugram, Delhi- NCR, Haryana- 122505, India

³Department of Physics and OMEG Institute, Soongsil University, Seoul 156-743, Republic of Korea

⁴School of Physics and International Research Center for Big-Bang

Cosmology and Element Genesis, Beihang University, Beijing 100083. China.

The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan,

National Astronomical Observatory of Japan 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

⁵Centre For Cosmology and Science Popularization (CCSP),

SGT University, Gurugram, Delhi- NCR, Haryana- 122505, India

Center for Theoretical Physics, Eurasian National University, Astana 010008, Kazakhstan

Chinese Academy of Sciences, 52 Sanlihe Rd, Xicheng District, Beijing.

A scale-independent energy-momentum squared gravity (EMSG) allows different gravitational couplings for different types of sources and has been proven to have interesting implications in cosmology. In this paper, the Big Bang Nucleosynthesis (BBN) formalism and the latest observational constraints are being used in order to extract constraints on this class of modified gravity models. The model has been constrained using the light element reaction rates. Using the tight constraint from BBN on the correction term in the Friedman equation due to EMSG, we find a significant deviation in the matter power spectrum of the cosmic microwave background (CMB) even for a very small allowed value of the modification parameter, thus pointing out the sensitivity of the matter power spectrum at the small scales.

I. INTRODUCTION

The idea of modification of Albert Einstein's General Relativity (GR)[1, 2] dates back to the first few months after the seminal paper published by Einstein. The proposals were made to extend GR and incorporate it into a larger, more unified theory. A few examples are Eddington's theory of connections, Weyl's scale-independent theory, and the higher dimensional theories of Kaluza and Klein. But almost after 108 years, the field equations proposed by Einstein remain still the best description of how space-time behaves on macroscopic scales. Einstein's equations govern everything that happens in our universe, starting from its expansion, structure formation, and black holes to the propagation of gravitational waves. Yet, the efforts to extend or modify GR never stopped, simply with the aim of understanding the dynamics of dark energy (DE) and dark matter (DM). To make a comprehensive list of such models, readers can go through the following to understand the motives and development of such theories: [3–9]. The recent curve ball thrown to us by the universe dubbed the H_0 tension, and the S_8 tension also points towards the requirement of some modification in the GR or some extension.

There exists a specific class of modified theories that permit the presence of scalars constructed from the energymomentum tensor $T_{\mu\nu}$ in the action. One can see it in the f(R,T) gravity, where the action involves the scalar $T = g^{\mu\nu}T_{\mu\nu}$, which is the trace of $T_{\mu\nu}$ [10]. The $f(R, \mathbb{T}^2)$ model has, $\mathbb{T}^2 \equiv T^{\mu\nu}T_{\mu\nu}$ in the action [11–14]. This model inspired by phenomenological considerations is coined as Energy-Momentum-Squared-Gravity (EMSG). A similar term is induced on RS brane as high energy correction to Einstein equations [15-19]. It should be noted that most of the modifications of gravity, in particular scale modifications, involve extra degrees of freedom, which need to be screened out locally by involving mechanisms such as chameleons or Vainstein. Theories that involve chameleon screening have great potential for late-time cosmology. However, proper screening consistent with local gravity constraints leaves no scope for late-time acceleration caused by large-scale modifications in this scenario. One of the interesting features of EMSG is that, unlike most of the modified theories of gravity, it does not involve extra degrees of freedom.

Implications of EMSG for late time acceleration have been studied in Refs.[13, 14]. The model can be constrained

^{*} havevirtue2@gmail.com

[†] mayukh_ccsp@sgtuniversity.org

[‡] cheoun@ssu.ac.kr

[§] kajino@buaa.edu.cn

[¶] sami_ccsp@sgtuniversity.org,

using observations [20–31]. Chaotic inflation [32, 33] has recently been examined in the framework of EMSG[34] and then the Primordial Black Hole and Gravitational Wave production is being studied in [35]. It has been reported that in this case, the model like chaotic inflation (excluded in standard cosmology by observation) in the larger umbrella theory of EMSG falls well within the allowed limits of Planck'18 [36].

Let us note that the stringent constraints on modified gravity models in the early universe come from the era of Big Bang Nucleosynthesis (BBN)[37–43]. Thus, it is necessary to go for the litmus test of the BBN on the EMSG model to make an astute comment on primordial cosmology and its phenomenology within the framework of the given modification of gravity.

The paper is organized as follows: In section II, we have given a short synopsis of the EMSG model and finally introduce the modified Friedmann Equation which plays the main role in this analysis. There also the stability criterion is also discussed. In section III, we constrained the model of EMSG using the latest observations of BBN. Interestingly, we have shown, BBN itself demands the negative value of the model parameter α which is required for the stable solution in this framework. With the given constraints imposed on EMSG, we calculated the matter power spectrum and reported in IV. Finally, we conclude with our findings and future directions in this context of modified gravity models in the last section V.




Figure 1. The cosmic expansion rate in Eq. (2) during the BBN epoch. The red-solid, blue-dotted and black-dashed lines indicate the H for $\alpha = -10^{-12} \text{ cm}^6/\text{s}^2/\text{g}^2$, $\alpha = -10^{-13} \text{ cm}^6/\text{s}^2/\text{g}^2$, and $\alpha = 0$ (standard), respectively.

II. BACKGROUND EQUATIONS AND THE MODEL

THe EMSG model actio is given by [13, 14]:

$$S = \frac{8\pi G}{2} \int d^4x \sqrt{-g} \left[M_{\rm p}^2 R - \alpha \, M_{\rm p}^{4(1-2\beta)} (\mathbb{T}^2)^\beta + 2\mathcal{L}_{\rm m} \right],\tag{1}$$

where $M_{\rm p}$ is the reduced Planck mass, R is the Ricci scalar associated with the spacetime metric $g_{\mu\nu}$, $\mathcal{L}_{\rm m}$ is the Lagrangian density corresponding to the matter source described by the energy-momentum tensor $T_{\mu\nu}$. In addition, $\mathbb{T}^2 \equiv T_{\mu\nu}T^{\mu\nu}$ is a scalar and α is a dimensionless constant that determines the coupling strength of the EMSG modification.

The details of the development of the background theory of this model can be found in [34]. The main consequence of this action is the modification of the Friedmann equation which becomes:

$$H^{2} = \frac{8\pi G}{3}\rho - \alpha \left(\frac{1}{2}p^{2} + \frac{4}{3}\rho p + \frac{1}{6}\rho^{2}\right)$$
(2)

The modification term with the constant α in front, remains non-negligible deep into the radiation domination and thus can impact the BBN. Thus, the BBN constraints become very important to look into while making any claim in this domain. Finally, one important fact is that $\alpha < 0$ is simply from the stability criterion (taking $\beta = 1$). Otherwise, the model faces gradient instability as discussed in [34]. Not only that, from the point of view of the ghost and gradient instability for $\beta = 1$, one needs $-2 \le \alpha \le 0$ Interestingly, this condition also becomes very important as we will see in the next section, BBN also demands a negative value of α to match the current bounds.

III. CONSTRAINTS FROM BBN

To constrain the EMSG model on BBN, we employ the BBN calculation code [44, 45] with updated reaction rates from the JINA REACLIB Database [46]. As input parameters, we adopt the central value of the neutron mean lifetime provided by the Particle Data Group, $\tau_n =$ 878.6±0.6 [47], and the lower limit of the baryon-to-photon ratio, $\eta = (6.104 \pm 0.058) \times 10^{-10}$, which corresponds to the baryon density based on the Λ CDM model (TT, TE, EE+lowE) from Planck observations of the cosmic microwave background, $\Omega_b h^2 = 0.02230 \pm 0.0021$ [48].

Fig. 1 illustrates the cosmic expansion rate (H) given by Eq. (2) for $\alpha = -10^{-12} \text{ cm}^6/\text{s}^2/\text{g}^2$ and $\alpha = -10^{-13} \text{ cm}^6/\text{s}^2/\text{g}^2$, comparing it with the standard formula for $\alpha = 0$. A notable deviation in the H is observed in the high-temperature region. However, as the temperature decreases, both the energy and pressure squared terms




Figure 2. Final abundances of D and Y_p as a function of α . We adopted the observational data of D abundance from the metal-poor Lyman- α absorption, D/H = 2.527 ± 0.030 [49], and mass fraction of ⁴He from metal-poor extra-galactic H II regions, $Y_p = 0.2448 \pm 0.0033$ [50]. In both panels, the red and blue boxes indicate constrained regions by the observational data within 2σ and 4σ range, respectively.

in the correction term rapidly decrease, leading to a reduction in the change of H. Consequently, in this model, H becomes similar to the standard case for temperatures $T \lesssim 10^9$ K.

Adopting the modified H given by Eq. (2), we proceed with BBN calculation. Fig. 1 depicts the final abundances as a function of α . The negative α values result in an increased H, leading nuclear reaction rates to freeze out earlier. The substantial change in H during the early stages leads to a higher neutron-to-proton ratio. The increased relic neutron abundance, in turn, increases the D abundance through the enhancement of the ${}^{1}\text{H}(n, \gamma){}^{2}\text{H}$ reaction. The increased D abundance further contributes to the ${}^{4}\text{He}$ abundance via the ${}^{3}\text{H}(d, n){}^{4}\text{He}$ reaction. Consequently, both D and ${}^{4}\text{He}$ abundances increase as α decreases. This trend aligns with findings from other studies on the effects of a modified expansion rate on primordial abundances [51–55]. From the results presented in Fig. 2, we can constrain the range of α . For the D/H abundance, we obtain lower limits on α of $-1.73 \times 10^{-12} \text{ cm}^6/\text{s}^2/\text{g}^2$ (2σ) and $-2.93 \times 10^{-12} \text{ cm}^6/\text{s}^2/\text{g}^2$ (4σ). For Y_p , the constrained region is narrower, with values of $-3.50 \times 10^{-13} \text{ cm}^6/\text{s}^2/\text{g}^2$ (2σ) and $-9.50 \times 10^{-13} \text{ cm}^6/\text{s}^2/\text{g}^2$ (4σ). For ⁷Li, we note that there is no consistent parameter space to explain the observational data: ⁷Li/H = 1.58 ± 0.31 [56].




Figure 3. Theoretical prediction for the matter power spectrum in the presence of the central values of α for the most stringent bound from Y_p (Green), allowed by D/H abundance (Red) and the Λ CDM model (Black).

IV. OBSERVATIONAL CONSEQUENCES AT CMB

The correction term with constant α creates a deviation of the matter power spectrum P(k, z = 0) from the standard Λ CDM. This is depicted in figure 3. Here, we have considered one massive neutrino with a fixed mass at 0.06eV. It is known that on large scales where $k < k_{nr}$, the matter power spectrum P(k, z = 0) only depends on the matter density fraction Ω_m today. Here k_{nr} means the free-streaming scale where the neutrinos become nonrelativistic. Whereas at the small scales $k > k_{nr}$, even the slightest presence of the term with α , can have a significant effect. Thus, the matter power spectrum can be written as:

$$P(k, z = 0) = \left\langle \left| \frac{\delta \rho_{CDM} + \delta \rho_b + \delta \rho_{SP}}{\rho_{CDM} + \rho_b + \rho_{SP}} \right| \right\rangle$$
$$= \Omega_m^{-2} \left\langle \left| \Omega_{CDM} \delta_{CDM} + \Omega_b \delta_b + \Omega_{SP} \delta_{SP} \right|^2 \right\rangle \quad , \qquad (3)$$

where, ρ_{CDM} , ρ_b , ρ_{SP} represents density fluctuation of the cold dark matter, baryon, and the sterile particle respectively. Whereas, Ω_{CDM} , Ω_b , Ω_{SP} represent the fractional energy density of the cold dark matter, baryon, and the

sterile particle respectively. For the most stringent bound from Y_p , α takes the value of $-3.50 \times 10^{-13} \,\mathrm{cm}^6/\mathrm{s}^2/\mathrm{g}^2$ (2σ) . For the D/H abundance, we obtain lower limits on α of $-1.73 \times 10^{-12} \,\mathrm{cm}^6/\mathrm{s}^2/\mathrm{g}^2$ (2 σ). For this value of α , the matter power spectrum is depicted in Fig. 3 with the green line. At large scales, one can see there is not much deviation from ACDM which is expected as the linear effects are dominant there. However, at a small scale, one can find that we observe significant deviations even for such a small α . Thus, keeping the BBN constraints in mind, one can see that the presence of α will affect the small-scale observations and can be studied to check the issues in Λ CDM. The suppression we see here is a combined effect of the massive neutrino and the negative α value correction on δ_{CDM} . One can compare the result with the findings of [34] where only the CMB+BAO bounds were accounted for. Interestingly enough even with the much smaller value of α compared to [34], the suppression remains significant. On the other hand, in this analysis, we did not consider any positive value of α , simply from the stability conditions demand which matches the demand from the BBN bounds as well.

V. CONCLUSION

We have explored the impact of the EMSG model on BBN and its implications for the matter power spectrum. For negative values of α , the correction term in the EMSG model enhances the cosmic expansion rate, depending on the squared energy density and pressure. Constraints on the parameter α were obtained earlier using CMB and BAO data; here we have focused on implications due to nucleosynthesis. Given the radiation domination due to relativistic species during the BBN epoch, the correction term rapidly decays over cosmic time, proportional to T^8 . Consequently, the EMSG correction specifically affects the initial stage of the BBN epoch, which leads to an increase in the primordial abundances of D and ⁴He for larger α .

Using the BBN observations, we have shown that the lower limit of α is constrained to $-3.50 \times 10^{-13} \,\mathrm{cm}^6/\mathrm{s}^2/\mathrm{g}^2$ and $-9.50 \times 10^{-13} \,\mathrm{cm}^6/\mathrm{s}^2/\mathrm{g}^2$ within 2σ and 4σ ranges, respectively.

One can find that the BBN bounds are much more stringent than the combined CMB+BBO bounds, as reported in [22]. Of course, one can expect the bounds of BBN to always be much more stringent, as explained. Interestingly though, even with the tightest bounds on α (10⁻¹²), the effect is quite prominent on the small scales of the matter power spectrum. Of course, one path to explore in this regard is to consider if the suppression can indicate some part of Dark Matter to be hot, following [57]. One important aspect that came of the analysis is that BBN bounds prefer the negative value of α which is also the theoretical demand to avoid ghost and gradient instability in this model with $\beta = 1$.

Finally, one more time, we would argue that any modification to Einstein's General Relativity, which can have effects at the scales of the BBN, has to satisfy BBN bounds, to go further, and be cultivated to be useful at a later time. In that case, BBN will provide the tightest constraints on the model.

VI. ACKNOWLEDGEMENT

Work of MRG is supported by the Department of Science and Technology(DST), Government of India under the Grant Agreement number IF18-PH-228 (INSPIRE Faculty Award) and by the Science and Engineering Research Board(SERB), DST, Government of India under the Grant Agreement number CRG/2020/004347(Core Research Grant).

MS is supported by the Science and Engineering Research Board (SERB), DST, Government of India under the Grant Agreement number CRG/2022/004120 (Core Research Grant). MS is also partially supported by the Ministry of Education and Science of the Republic of Kazakhstan, Grant No. AP14870191 and CAS President's International Fellowship Initiative(PIFI).

TK and DJ are partly supported by the National Natural Science Foundation of China (No. 12335009), and the National Key R&D Program of China (2022YFA1602401). TK is also supported in part by Grants-in-Aid for Scientific Research of Japan Society for the Promotion of Science (20K03958).

MKC's work was supported by the Basic Science Research Program of the National Research Foundation of Korea (NRF) under Grants No. 2021R1A6A1A03043957 and No. 2020R1A2C3006177.

MRG also wants to thank the organizers of OECE,2023, Beihang University for the hospitality received when the project was initiated.

- A. Einstein, The foundation of the general theory of relativity, Annalen Phys. 49, no.7, 769-822 (1916).
- [2] A. Einstein, Explanation of the Perihelion Motion of Mer-

cury from the General Theory of Relativity, Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.) **1915**, 831-839 (1915).

- [3] C. Brans and R. H. Dicke, Mach's Principle and a Relativistic Theory of Gravitation, Physical Review 124 (1961) 925–935.
- [4] A. H. Chamseddine and V. Mukhanov, Mimetic Dark Matter, JHEP 11 (2013) 135, [1308.5410].
- [5] J. D. Bekenstein, Relativistic gravitation theory for the MOND paradigm, Phys. Rev. D 70 (2004) 083509, [astroph/0403694].
- [6] J. W. Moffat, Scalar-tensor-vector gravity theory, JCAP 03 (2006) 004, [gr-qc/0506021].
- [7] C. Skordis and T. Zlosnik, New Relativistic Theory for Modified Newtonian Dynamics, Phys. Rev. Lett. 127 (2021) 161302, [2007.00082].
- [8] T. P. Sotiriou and V. Faraoni, f(R) Theories Of Gravity, Rev. Mod. Phys. 82 (2010) 451–497, [0805.1726].
- [9] F. W. Hehl and B. Mashhoon, A Formal framework for a nonlocal generalization of Einstein's theory of gravitation, Phys. Rev. D 79 (2009) 064028, [0902.0560].
- T. Harko, F. S. N. Lobo, S. Nojiri and S. D. Odintsov, f (R, T) gravity, Phys. Rev. D 84 (2011) 024020, [1104.2669].
- [11] N. Katirci and M. Kavuk, $f(R, T_{\mu\nu}T^{\mu\nu})$ gravity and Cardassian-like expansion as one of its consequences, Eur. Phys. J. Plus 129 (2014) 163, [1302.4300].
- [12] M. Roshan and F. Shojai, Energy-Momentum Squared Gravity, Phys. Rev. D 94 (2016) 044002, [1607.06049].
- [13] O. Akarsu, N. Katırcı and S. Kumar, Cosmic acceleration in a dust only universe via energy-momentum powered gravity, Phys. Rev. D 97 (2018) 024011, [1709.02367].
- [14] C. V. R. Board and J. D. Barrow, Cosmological Models in Energy-Momentum-Squared Gravity, Phys. Rev. D 96 (2017) 123517, [1709.09501].
- [15] L. Randall and R. Sundrum, Phys. Rev. Lett. 83, 3370-3373 (1999) [arXiv:hep-ph/9905221 [hep-ph]].
- [16] L. Randall and R. Sundrum, Phys. Rev. Lett. 83, 4690-4693
 (1999) [arXiv:hep-th/9906064 [hep-th]].
- [17] M. Bennai, H. Chakir and Z. Sakhi, Eur. J. Phys. 9, 84-93 (2006) [arXiv:0806.1137 [gr-qc]].
- [18] M. R. Gangopadhyay and G. J. Mathews, JCAP 03, 028 (2018) [arXiv:1611.05123 [astro-ph.CO]].
- [19] G. Calcagni, Phys. Rev. D 69, 103508 (2004) [arXiv:hepph/0402126 [hep-ph]].
- [20] E. Nazari, F. Sarvi and M. Roshan, Generalized Energy-Momentum-Squared Gravity in the Palatini Formalism, Phys. Rev. D 102 (2020) 064016, [2008.06681].
- [21] S. Bahamonde, M. Marciu and P. Rudra, Dynamical system analysis of generalized energy-momentum-squared gravity, Phys. Rev. D 100 (2019) 083511, [1906.00027].
- [22] O. Akarsu, N. Katirci, S. Kumar, R. C. Nunes and M. Sami, Cosmological implications of scale-independent energymomentum squared gravity: Pseudo nonminimal interact Phys. Rev. D 98 (2018) 063522, [1807.01588].
- [23] O. Akarsu and N. M. Uzun, Cosmological models in scale-independent energy-momentum squared gravity, Phys. Dark Univ. 40 (2023) 101194, [2301.11204].
- [24] E. Nazari, Light bending and gravitational lensing in energy-momentum-squared gravity, Phys. Rev. D 105 (2022) 104026, [2204.11003].
- [25] N. Nari and M. Roshan, Compact stars in Energy-Momentum Squared Gravity, Phys. Rev. D 98 (2018)

024031, [1802.02399].

- [26] A. Kazemi, M. Roshan, I. De Martino and M. De Laurentis, Jeans analysis in energy–momentum-squared gravity, Eur. Phys. J. C 80 (2020) 150, [2001.04702].
- [27] E. Nazari, M. Roshan and I. De Martino, Constraining energy-momentum-squared gravity by binary pulsar observations, Phys. Rev. D 105 (2022) 044014, [2201.08578].
- [28] O. Akarsu, E. Nazari and M. Roshan, Relativistic binary systems in scale-independent energy-momentum squared gravity, 2302.04682.
- [29] O. Akarsu, J. D. Barrow, S. C, ikinto glu, K. Y. Ek,si and N. Katirci, Constraint on energy-momentum squared gravity from neutron stars and its cosmological implications, Phys. Rev. D 97 (2018) 124017, [1802.02093].
- [30] M. Faraji, N. Rashidi and K. Nozari, Inflation in energymomentum squared gravity in light of Planck2018, Eur. Phys. J. Plus 137 (2022) 593, [2107.13547].
- [31] C. Ranjit, P. Rudra and S. Kundu, Constraints on Energy–Momentum Squared Gravity from cosmic chronometers and Supernovae Type Ia data, Annals Phys. 428 (2021) 168432, [2010.02753]
- [32] A. D. Linde, Chaotic inflation, Physics Letters B 129 (1983) 177–181.
- [33] A. D. Linde, A new inflationary universe scenario: a possible solution of the horizon, flatness, homogeneity, isotropy, and primordial monopole problems, Physics Letters B 108 (1982) 389–393.
- [34] S. A. Hosseini Mansoori, F. Felegary, M. Roshan, O. Akarsu and M. Sami, T²- inflation: Sourced by energy-momentum squared gravity., Phys. Dark Univ. 42, 101360 (2023), [arXiv:2306.09181 [gr-qc]].
- [35] S. A. Hosseini Mansoori, F. Felegray, A. Talebian and M. Sami, JCAP 08, 067 (2023) [arXiv:2307.06757 [astroph.CO]].
- [36] BICEP, Keck collaboration, P. A. R. Ade et al., Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations Phys. Rev. Lett. 127 (2021) 151301, [2110.00483
- [37] J. Bernstein, L. S. Brown, and G. Feinberg, COSMOLOG-ICAL HELIUM PRODUCTION SIMPLIFIED., Rev. Mod. Phys. 61 (1989) 25. I, IV [44] E. W. Kolb and M. S. Turner, The Early Universe, vol. 69. 1990.
- [38] K. A. Olive, G. Steigman, and T. P. Walker, Primordial nucleosynthesis: Theory and observations., Phys. Rept. 333 (2000) 389–407, arXiv:astro-ph/9905320.
- [39] R. H. Cyburt, B. D. Fields, K. A. Olive, and T.-H. Yeh, Big Bang Nucleosynthesis: 2015., Rev. Mod. Phys. 88 (2016) 015004, arXiv:1505.01076 [astro-ph.CO].
- [40] J. D. Barrow, S. Basilakos, and E. N. Saridakis, Big Bang Nucleosynthesis constraints on Barrow entropy., Phys. Lett. B 815 (2021) 136134, arXiv:2010.00986 [gr-qc].
- [41] P. Asimakis, S. Basilakos, N. E. Mavromatos, and E. N. Saridakis, Big bang nucleosynthesis constraints on higherorder modified gravities., Phys. Rev. D 105 (2022) no. 8, 084010, arXiv:2112.10863 [gr-qc].
- [42] N. Sasankan, M. R. Gangopadhyay, G. J. Mathews and M. Kusakabe, New observational limits on dark radiation in braneworld cosmology., Phys. Rev. D 95, no.8, 083516 (2017), [arXiv:1607.06858 [astro-ph.CO]].

- [43] K. Ichiki, M. Yahiro, T. Kajino, M. Orito and G. J. Mathews, Observational constraints on dark radiation in brane cosmology., Phys. Rev. D 66, 043521 (2002), [arXiv:astroph/0203272 [astro-ph]].
- [44] L. Kawano, "Let's go: Early universe. 2. Primordial nucleosynthesis: The Computer way," FERMILAB-PUB-92-004-A.
- [45] M. S. Smith, L. H. Kawano and R. A. Malaney, Astrophys. J. Suppl. 85, 219-247 (1993).
- [46] R. H. Cyburt *et al.*, Astrophys. J. Suppl. Ser. **189**, 240 (2010).
- [47] R. L. Workman *et al.* [Particle Data Group], PTEP **2022**, 083C01 (2022).
- [48] N. Aghanim *et al.* [Planck], Astron. Astrophys. **641**, A6 (2020) [erratum: Astron. Astrophys. **652**, C4 (2021)].
- [49] R. J. Cooke, M. Pettini and C. C. Steidel, Astrophys. J. 855, no.2, 102 (2018).

- [50] E. Aver et al., Mon. Not. Roy. Astr. Soc. 510, 1, 373 (2022).
- [51] M. Kusakabe, S. Koh, K. S. Kim and M. K. Cheoun, Phys. Rev. D 93, no.4, 043511 (2016).
- [52] D. Jang, M. Kusakabe and M. K. Cheoun, Phys. Rev. D 97, no.4, 043005 (2018).
- [53] N. Sasankan, M. R. Gangopadhyay, G. J. Mathews and M. Kusakabe, Int. J. Mod. Phys. E 26, no.08, 1741007 (2017) [arXiv:1706.03630 [astro-ph.CO]]
- [54] G. J. Mathews, M. Gangopadhyay, N. Sasankan, K. Ichiki and T. Kajino, AIP Conf. Proc. **1947**, no.1, 020014 (2018) [arXiv:1711.04873 [astro-ph.CO]].
- [55] G. Mathews, M. Kusakabe, M. Gangopadhyay, T. Kajino and N. Sasankan, EPJ Web Conf. 184, 01011 (2018)
- [56] L. Sbordone et al., Astron. Astrophys. 522, A26 (2010).
- [57] S. Bhattacharya, S. Das, K. Dutta, M. R. Gangopadhyay, R. Mahanta and A. Maharana, Phys. Rev. D 103, no.6, 063503 (2021) [arXiv:2009.05987 [astro-ph.CO]].