

# SRF Cavity as Galactic Dark Photon Telescope

Yifan Chen,<sup>1</sup> Chunlong Li,<sup>2</sup> Yuxiang Liu,<sup>3,4</sup> Yuxin Liu,<sup>5,6</sup> Jing Shu,<sup>3,4,7</sup> and Yanjie Zeng<sup>2,6</sup>

<sup>1</sup>*Niels Bohr International Academy, Niels Bohr Institute, Blegdamsvej 17, 2100 Copenhagen, Denmark*

<sup>2</sup>*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China*

<sup>3</sup>*School of Physics and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, China*

<sup>4</sup>*Center for High Energy Physics, Peking University, Beijing 100871, China*

<sup>5</sup>*International Center for Theoretical Physics Asia-Pacific, Beijing/Hangzhou, China*

<sup>6</sup>*School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China*

<sup>7</sup>*Beijing Laser Acceleration Innovation Center, Huairou, Beijing, 101400, China*

(Dated: February 7, 2024)

Dark photons, aside from constituting non-relativistic dark matter, can also be generated relativistically through the decay or annihilation of other dark matter candidates, contributing to a galactic dark photon background. The production of dark photons tends to favor specific polarization modes, determined by the microscopic coupling between dark matter and dark photons. We leverage data obtained from previous searches for dark photon dark matter using a superconducting radio-frequency cavity to explore galactic dark photon fluxes. The interplay of anisotropic directions and Earth's rotation introduces a diurnal modulation of signals within the cavities, manifesting distinct variation patterns for longitudinal and transverse modes. Our findings highlight the efficacy of superconducting radio-frequency cavities, characterized by significantly high-quality factors, as powerful telescopes for detecting galactic dark photons, unveiling a novel avenue in the indirect search for dark matter through multi-messenger astronomy.

**Introduction** — Dark photon, posited as a minimal extension to the Standard Model (SM) via an additional U(1) gauge group, arise from speculative new physics theories such as extra dimensions or grand unification [1–4]. They are considered a substantial candidate for dark matter, particularly distinguishing themselves as coherent vector waves at sub-eV mass scales [5, 6]. This results in an effective electric current due to kinetic mixing with standard model photons. To probe this intricate interaction, various experimental approaches have been undertaken [7–36]. Notably, superconducting radio-frequency (SRF) cavities have shown exceptional promise, offering the most stringent constraints thus far [26, 30, 32] due to their extremely high quality factor.

Beyond merely serving as a component of dark matter in a standard halo distribution, dark photons may manifest across various cosmic scenarios [37], such as an isotropic cosmological background. Additionally, dark photons generated from dark matter decay or annihilation contribute to a galactic background with a distinct directionality emanating from the galactic center. The detection of these galactic dark photons thus serves as an indirect detection of dark matter. Their anisotropic distribution on Earth results in diurnal signal modulation in direction-sensitive detectors [38], which can be utilized as distinctive indicators of their galactic origins. Moreover, analyzing the polarization and spectral characteristics of dark photons offers in-depth insights into the nature of dark matter interactions, a subject further explored in a companion theoretical paper.

In this work, we initiate the search for galactic dark photons using an SRF cavity. We demonstrate that the projection of dark photon wavefunctions onto cav-

ity modes will result in a polarization-dependent modulation of signatures. Utilizing datasets initially assembled for dark photon dark matter (DPDM) searches [32], we establish pioneering constraints for galactic dark photons across two distinct polarization modes. Our analysis considers the spectra of galactic dark photons from two models: a flat spectrum emanating from four-body cascade decay [38] and a narrow spectrum (1/500 of central frequency) associated with two-body parametric decay. This exploration expands the capabilities and reach of current dark matter research methodologies, utilizing cavities for the multi-messenger, indirect detection of ultralight dark matter. We navigate a vast, previously untapped parameter space, capitalizing on the exceptional quality factor of the SRF cavity, which offers unique advantages compared with traditional copper cavities [38]. **Galactic Dark Photons from Dark Matter Decay** — We begin by characterizing the wavefunctions and correlations inherent to a general dark photon background. The wavefunction  $X^\mu$  is formulated in the frequency domain as:

$$X^\mu(t, \vec{x}) = \int \sum_P X_P(\omega, \hat{k}) e_P^\mu e^{i(\omega t - \vec{k} \cdot \vec{x})} \frac{d\omega}{2\pi} d^2\hat{k}, \quad (1)$$

where the frequency  $\omega$  and spatial momentum  $\vec{k}$  adhere to the dispersion relation  $\omega^2 = |\vec{k}|^2 + m_X^2$  for the dark photon mass  $m_X$ .  $X_P$  signifies the amplitude for each polarization mode  $P$ , incorporating one longitudinal (L) and two transverse (T) components, while  $e_P^\mu$  is the associated polarization vector basis.

Considering the non-coherent nature of dark matter decay, galactic dark photons are expected to manifest

as a stochastic Gaussian field. This is quantified by the two-point correlation function:

$$\langle X_P(\omega, \hat{k}) X_{P'}(\omega', \hat{k}')^* \rangle = \delta(\omega - \omega') \delta^2(\hat{k} - \hat{k}') \delta_{PP'} \mathcal{S}_P(\omega, \hat{k}). \quad (2)$$

Here,  $\langle \dots \rangle$  denotes the ensemble average and  $\mathcal{S}_P(\omega, \hat{k})$  is the power spectral density (PSD) of the  $P$ -mode dark photon from direction  $\hat{k}$ . The total energy density of the dark photon is obtained by integrating  $\omega^2 \mathcal{S}_P(\omega, \hat{k})$  over the frequency spectrum.

The polarization states of the dark photon are contingent upon its interactions with dark matter fields. We examine scenarios where either longitudinal or transverse modes are predominant. Specifically, the longitudinal modes often arise in dark Higgs models, where a scalar field  $\Phi$  decays into two relativistic dark photons through an interaction Lagrangian proportional to  $\Phi X^\mu X_\mu$ . Conversely, transverse modes are typically generated through axion-photon-like interactions, denoted by a term proportional to  $\Phi X_{\mu\nu} \tilde{X}^{\mu\nu}$ , where  $X_{\mu\nu}$  represents the dark photon field tensor and  $\tilde{X}^{\mu\nu}$  its dual. Comprehensive derivations of these models and the dependencies of their couplings are elaborated in the Supplementary Material.

In the interaction models under consideration, a massive scalar field's decay into two dark photons may induce parametric instability [39–60], rapidly diminishing the scalar field through Bose-enhanced dark photon production [61]. In conventional virialized scalar halos, such instabilities might be moderated by gravitational redshift when couplings are sufficiently weak [49], albeit at the cost of reducing the decay products to near undetectable levels. To address this, we consider that a fraction of the scalar dark matter aggregates into dense clumps nested within broader, more diffuse miniclusters [62–76]. Within these dense regions, an equilibrium between dark photon production and escape, limited by the clumps' size, culminates in a saturated state of dark photons. The clumps achieve stability via a balance of parametric decay and accretion, resulting in a steady and substantial dark photon flux that eclipses that from mere perturbative decay, akin to saturated superradiant clouds in axion-photon systems [41, 55]. The parametric instability band predominantly shapes the resulting dark photon spectrum, related to the scalar clumps' core size [42, 50]. Assuming a virialized velocity distribution with a dispersion of  $10^{-3}c$ , the produced dark photons exhibit a bandwidth approximately 1/500 of the central frequency. An alternative production mechanism involves a scalar dark matter field decaying into two intermediate states,  $\Phi$ , which then undergo cascade decay into four dark photons [38]. Here, the dark photon spectrum is broadly flat, confined within a frequency range dictated by the dark matter mass. We categorize the two production mechanisms as follows:

- Two-body parametric decay;
- Four-body cascade decay.




FIG. 1: Top: Schematic of the single-cell 1.3 GHz elliptical cavity utilized in this study with incoming dark photon fields from direction  $\hat{k}$  (orange). The angle between  $\hat{k}$  and the cavity's symmetry axis  $\hat{l}_{\text{cav}}$  is denoted as  $\Psi$ . The electric fields of the targeted cavity mode  $\text{TM}_{010}$  are illustrated with red arrows, their length proportional to the field strength. Bottom: The overlap factor between the  $\text{TM}_{010}$  mode and incoming dark photon fields of different polarization modes, as defined in Eq. (3), assuming  $|\vec{k}| = \omega_0$  for a relativistic dark photon.

In both scenarios, local dark photon flux densities may soar to  $1000 \rho_\gamma$ , compatible with cosmological constraints on dark matter decay [38, 61], where  $\rho_\gamma$  signifies the local electromagnetic photon density [77]. The angular distribution of the incoming dark photon flux adheres to the line-of-sight integral of the dark matter density, often referred to as the J-factor in dark matter indirect detection literature [78].

**Cavity as a Dark Photon Telescope** — When dark photon fluxes transit through a cavity, they can potentially excite a cavity mode if their frequency aligns with the resonant frequency of the cavity, a process facilitated by kinetic mixing with electromagnetic photons [32]. In the interaction picture, dark photons contribute to an effective electric current  $\vec{J}_{\text{eff}} = \epsilon m_X^2 \vec{X}$ , where  $\epsilon$  denotes the kinetic mixing coefficient, and  $\vec{X}$  represents the spatial component of the dark photon wavefunction. As elucidated in the Supplementary Material, the rate of excitation depends on the overlap between  $\vec{X}$  and the cavity mode's electric field  $\vec{E}_0$ , quantified by the parameter:

$$C_P(\hat{k}) \equiv \frac{1}{V} \left| \int_V e^{-i\vec{k}\cdot\vec{x}} \hat{e}_P(\hat{k}) \cdot \vec{E}_0(\vec{x}) d^3\vec{x} \right|^2, \quad (3)$$

where  $\hat{e}_P$  is the spatial unit directional vector of the polarization basis  $e_P^\mu$ , and the integral is taken over the cavity's volume  $V$ . Typically, the phase factor  $e^{-i\vec{k}\cdot\vec{x}}$  is approximated as 1 for the overlap function for DPDM [32], due to the de-Broglie wavelength  $1/|\vec{k}|$  of the non-relativistic field being considerably longer than the cavity scale. For axisymmetric cavity modes, the angular dependence reduces to the polar angle  $\Psi \equiv -\hat{k}\cdot\hat{l}_{\text{cav}}$ , with  $\hat{l}_{\text{cav}}$  representing the cavity's symmetry axis, as depicted in the top panel of Fig. 1. We numerically evaluate the overlap factor  $C_P(\Psi)$  for the  $\text{TM}_{010}$  mode of a single-cell 1.3 GHz elliptical cavity utilized in this study, illustrated in the bottom panel of Fig. 1. The longitudinal mode, with  $\hat{e}_L(\hat{k})$  aligned with  $\hat{k}$ , achieves maximum overlap at  $\Psi = 0$  and diminishes to zero at  $\Psi = \pi/2$ . The two transverse circular polarized modes, indistinct from one another, display an inverse variation compared to the longitudinal mode. These evaluations assume a relativistic limit for the dark photon, where  $|\vec{k}|$  is nearly equal to the mode's resonant frequency  $\omega_0$ .

The PSD absorbed by the cavity from dark photon fields bears resemblance to that from DPDM [32]:

$$S_{\text{sig}}^P(\omega, t) = \frac{\epsilon^2 m_X^4 V |\vec{e}_P|^2}{4(\omega - \omega_0)^2 + \omega_0^2/Q_L^2} \int S_P(\omega, \hat{k}) C_P(\Psi(t)) d^2\hat{k}. \quad (4)$$

The distinction comes with the factor  $|\vec{e}_P|^2$  denotes the magnitude square of the spatial component of the polarization basis  $e_P^\mu$ . This factor equates to 1 for transverse modes and escalates to  $\omega^2/m_X^2$  for the longitudinal mode, suggesting that a relativistic longitudinal mode could significantly enhance signal power compared to transverse modes, given the same spectra. Here,  $Q_L$  indicates the cavity's loaded quality factor, determining the resonance bandwidth. In the Galactic frame, the Earth's rotation leads to temporal variations in the cavity's symmetry axis  $\hat{l}_{\text{cav}}(t)$  and the associated  $\Psi(t)$ . Consequently, these changes result in polarization-dependent modulations in the PSD, which can be leveraged as distinct signatures for the detection of galactic dark photons.

**Constraining Galactic Dark Photons through Diurnal Modulation Analysis** — We now embark on a search for the diurnal modulation of signals stemming from galactic dark photons, leveraging data from a previous DPDM search employing a single-cell elliptical niobium SRF cavity [32]. The cavity was equipped with a frequency tuning mechanism [79, 80] and an amplifier circuit, and maintained in liquid helium at a temperature of approximately 2 K. The experimental platform [81–86] facilitated the calibration of relevant parameters, including the loaded quality factor  $Q_L$ , coupling factor  $\beta$ , net amplification factor  $G_{\text{net}}$ , and the resonant frequency of the  $\text{TM}_{010}$  mode, centered around  $\omega_0 \approx 2\pi 1.3$  GHz, along with its stability range for each scanning process. Each scan step was approximately 5 minutes, with 100 seconds dedicated to data acquisi-




FIG. 2: Normalized power excess from experimental data (black dots) for each scan step as a function of measurement time, and the predicted modulation signals from galactic dark photons for two polarization modes (blue and red lines, respectively). The right panel illustrates the distribution of normalized power excess, fitting a Gaussian distribution with no deviations exceeding 4-sigma. The experiment's initial time,  $t_0$ , was UTC 22:23:52, April 12, 2023. The two runs are delineated by the gray region. The modulation signals are customized for the detector's location ( $116.4^\circ\text{E}$ ,  $39.9^\circ\text{N}$ , Beijing) and orientation (cavity's symmetry axis pointing towards Earth's center), with the signal normalization set to the 90% exclusion bound.

tion and the remainder to frequency tuning and parameter calibration. During two continuous operational periods from April 12-14 and 23-26, 2023, we completed  $N_{\text{bin}} = 350$  and 800 scan steps respectively, covering a total frequency range of  $2\pi 1.37$  MHz.

In our analysis, only the received power at the resonant frequency bin with a bandwidth of  $2\pi 11.5$  Hz [32], denoted by  $\mathcal{P}^i$  for the  $i$ -th step, was considered. The dominant noise contribution typically comes from the amplifier. For each run, we computed the average and standard deviation of  $\mathcal{P}^i$ , denoted as  $\bar{\mathcal{P}}$  and  $\sigma_P$ , respectively, and determined the normalized power excess  $\delta^i \equiv (\mathcal{P}^i - \bar{\mathcal{P}})/\sigma_P$ . These are depicted as black dots in Fig. 2, fitting well with a standard normal distribution without 4-sigma excess. Unlike Ref. [32], where constant fit and normalized power excess were calculated every 50 contiguous bins for environmental stability, the current search aggregates all bins from the two runs without suppressing potential diurnal variations.

The expected power signal induced by galactic dark photons at scan step  $i$  is given by:

$$\mathcal{P}_{\text{sig}}^i(t) = G_{\text{net}} \eta_{\text{bin}} \frac{\beta}{\beta + 1} \frac{(\omega_0^i)^2}{8\pi Q_L^2} \sum_P S_{\text{sig}}^P(\omega_0^i, t). \quad (5)$$

Here,  $\omega_0^i$  is the cavity's resonant frequency, and  $\eta_{\text{bin}} \simeq 84\%$  accounts for the signal leakage outside the bin due

to variations in the resonant frequency. As the total scan range of  $2\pi 1.37$  MHz is less than the considered galactic dark photons' bandwidth, we assume the spectrum  $\mathcal{S}_P(\omega_0^i, \hat{k})$  in Eq. (4) remains constant with respect to  $\omega_0^i$ . The diurnal signal variation is calculated by considering the detector's location ( $116.4^\circ\text{E}$ ,  $39.9^\circ\text{N}$ , Beijing) and orientation (symmetry axis pointing towards Earth's center), along with Earth's rotational changes in the Galactic frame, as detailed in the Supplementary Materials. The time variation during each scan is negligible compared to the diurnal variation timescale. Analogous to  $\delta^i$ , we define  $\delta_{\text{sig}}^i \equiv (\mathcal{P}_{\text{sig}}^i - \Sigma_i \mathcal{P}_{\text{sig}}^i / N_{\text{bin}}) / \sigma_{\mathcal{P}}$  for the signal variations, exemplified in Fig. 2 for the two polarization modes.

We evaluate the hypothesis of galactic dark photons characterized by mass  $m_X$  and kinetic mixing coefficient  $\epsilon$  utilizing the following probability function:

$$\Pr(\delta^i | \epsilon, m_X) \propto \prod_i \exp\left(\frac{-(\delta^i - \delta_{\text{sig}}^i)^2}{2 + 2(\delta^i)^2 \sum_j F_j^2}\right), \quad (6)$$

where  $F_j^2$  quantifies the uncertainties of the involved parameters, collectively accounting for less than 5% [32]. The 90% exclusion limit derived from this probability function is depicted in Fig. 3, showcasing constraints for the two benchmark spectra: a flat spectrum from four-body cascade decay, and a  $1/500$  bandwidth around the central frequency of approximately  $\omega_0$  from two-body parametric decay. Both spectra are normalized to a dark photon energy density of  $\rho_X = 1000 \rho_\gamma$ . We constrain our analysis to regions where dark photons are relativistic, specifically where  $m_X \leq \omega/10$ , as non-relativistic regions lead to a mix of polarization modes and diminish the diurnal modulation. Notably, the longitudinal mode is more significantly constrained than the transverse modes, attributed to the distinct factors  $|\vec{e}_L|^2 = \omega^2/m_X^2$  and  $|\vec{e}_T|^2 = 1$  in the signal power expression (Eq. (4)). The constraints for the four-body cascade decay also depend on the mass of the original dark matter particle, which influences the bandwidth of  $\mathcal{S}_P$  and thus the amplitude at the cavity's resonant frequency. The gradient color region in the figure indicates the variation in exclusion for dark matter masses ranging from  $4\omega_0$  (bottom) to  $100\omega_0$  (top). Alternatively, the two-body parametric decay leads to dark photons with energy half that of the dark matter particle mass, requiring the latter to be around  $2\omega_0$  for our analysis. We also juxtapose our findings with established constraints on kinetic-mixing dark photons from various sources. This includes light-shining-through-walls (LSW) experiments such as Dark SRF [30], which use SRF cavities similar to ours, alongside a spectrum of previous results [87–92], combined constraints from fifth-force searches [93–97], and the effects on the cosmic microwave background (CMB) from photon-dark photon oscillations [98, 99].

**Conclusion** — In this study, we have investigated galac-




FIG. 3: The 90% exclusion limits on the kinetic mixing coefficient  $\epsilon$  for galactic dark photons with mass  $m_X$ . The top and bottom panels correspond to the two production mechanisms: four-body cascade decay and two-body parametric decay, respectively. Exclusions for the longitudinal and transverse polarization modes are shown in blue and red. For context, we include established constraints from light-shining-through-walls (LSW) experiments like Dark SRF [30] using similar SRF cavities, and earlier results [87–92], fifth-force searches [93–97], and CMB distortions from photon-dark photon oscillations [98, 99]. The gradient color in the four-body cascade decay panel indicates variations in the dark matter particle mass from  $4\omega_0$  (bottom) to  $100\omega_0$  (top). For the two-body parametric decay, the dark matter mass is set at approximately  $2\omega_0$ . All exclusions assume the energy density of galactic dark photons is  $1000 \rho_\gamma$ .

tic dark photons as decay products from scalar dark matter, demonstrating the SRF cavity's sensitivity to their largely unexplored parameter space in terms of mass and kinetic mixing coefficient. The two production mechanisms we analyzed allow for ultralight bosonic dark matter decay without exponential depletion of the background, thus providing a novel avenue for the indirect detection of scalar dark matter via dark photons. Owing to their anisotropic distribution, the induced signal in a cavity at rest with respect to the Earth undergoes a diur-

nal modulation. This modulation pattern is sensitive to the polarization mode of dark photons, with constraints being particularly stringent for the longitudinal polarization mode. The exceptional sensitivity of our study is largely attributed to the extremely high quality factor of the SRF cavity. This high quality factor results in a narrow bandwidth, which inherently supports low noise levels, thus enabling the exclusion of galactic dark photons across a spectrum bandwidth substantially broader than the cavity's own bandwidth. This approach mirrors the methodology employed in DPDM searches [26, 30, 32], showcasing the effectiveness of SRF cavities in probing wide spectral ranges of dark photon signatures.

Enhancements to the SRF cavity's sensitivity to dark photons are possible through the integration of advanced readout systems. Innovations such as quantum non-demolition photon counting [16, 100], squeezing techniques [101–106], and multi-mode couplings [107–111] offer pathways to significantly mitigate noise, which is predominantly amplifier-based in current configurations.

Looking ahead, as more cavities become operational, the establishment of a network of detectors [37, 112, 113] with extensive long-baseline separation [31] will enable a detailed analysis of the general properties of dark photons. Long-baseline interferometry and analyzing the relative amplitude from the projection of the dark photon wavefunction will allow for the precise localization and characterization of incoming dark photons, including their polarization modes [37]. Furthermore, an array of strategically positioned and oriented detectors can investigate the presence of chiral components in the dark photon sector [37]. This multifaceted approach not only provides insights into the macroscopic behavior of dark photons but also delves into their microscopic nature, enhancing our understanding of their origins. Additionally, integrating these cavity networks with other astrophysical observations [114], such as gravitational waves, paves the way for new frontiers in multi-messenger astronomy and cosmology.

The versatility of SRF cavities extends well beyond the realm of dark photon detection. They also hold potential for axion searches and the detection of high-frequency gravitational waves through heterodyne upconversion, a process where a bosonic field facilitates a transition from an injected pump mode to a signal mode [111, 115–123]. This unique capability positions SRF cavities as instrumental in searches for galactic axions [38] at frequencies significantly lower than their resonant frequencies, showcasing their broad applicability in the field of particle physics and astrophysics.

**Acknowledgements.** We are grateful to Nicholas L. Rodd and Hyeonseok Seong for useful discussions. We acknowledge the utilization of the Platform of Advanced Photon Source Technology R&D. This work is supported by the National Key Research and Development Program

of China under Grant No. 2020YFC2201501. Y.C. is supported by VILLUM FONDEN (grant no. 37766), by the Danish Research Foundation, and under the European Union's H2020 ERC Advanced Grant "Black holes: gravitational engines of discovery" grant agreement no. Gravitas-101052587, and the Munich Institute for Astro-, Particle and BioPhysics (MIAPbP) which is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2094 – 390783311, and by the European Consortium for Astroparticle Theory in the form of an Exchange Travel Grant, and by FCT (Fundação para a Ciência e Tecnologia I.P, Portugal) under project No. 2022.01324.PTDC. J.S. is supported by Peking University under startup Grant No. 7101302974 and the National Natural Science Foundation of China under Grants No. 12025507, No.12150015; and is supported by the Key Research Program of Frontier Science of the Chinese Academy of Sciences (CAS) under Grants No. ZDBS-LY-7003.

- 
- [1] Peter Svrcek and Edward Witten, "Axions In String Theory," *JHEP* **06**, 051 (2006), [arXiv:hep-th/0605206](#).
  - [2] S. A. Abel, M. D. Goodsell, J. Jaeckel, V. V. Khoze, and A. Ringwald, "Kinetic Mixing of the Photon with Hidden U(1)s in String Phenomenology," *JHEP* **07**, 124 (2008), [arXiv:0803.1449 \[hep-ph\]](#).
  - [3] Asimina Arvanitaki, Savvas Dimopoulos, Sergei Dubovsky, Nemanja Kaloper, and John March-Russell, "String Axiverse," *Phys. Rev. D* **81**, 123530 (2010), [arXiv:0905.4720 \[hep-th\]](#).
  - [4] Mark Goodsell, Joerg Jaeckel, Javier Redondo, and Andreas Ringwald, "Naturally Light Hidden Photons in LARGE Volume String Compactifications," *JHEP* **11**, 027 (2009), [arXiv:0909.0515 \[hep-ph\]](#).
  - [5] Ann E. Nelson and Jakub Scholtz, "Dark Light, Dark Matter and the Misalignment Mechanism," *Phys. Rev. D* **84**, 103501 (2011), [arXiv:1105.2812 \[hep-ph\]](#).
  - [6] Paola Arias, Davide Cadamuro, Mark Goodsell, Joerg Jaeckel, Javier Redondo, and Andreas Ringwald, "WISPy Cold Dark Matter," *JCAP* **06**, 013 (2012), [arXiv:1201.5902 \[hep-ph\]](#).
  - [7] Dieter Horns, Joerg Jaeckel, Axel Lindner, Andrei Lobanov, Javier Redondo, and Andreas Ringwald, "Searching for WISPy Cold Dark Matter with a Dish Antenna," *JCAP* **04**, 016 (2013), [arXiv:1212.2970 \[hep-ph\]](#).
  - [8] Saptarshi Chaudhuri, Peter W. Graham, Kent Irwin, Jeremy Mardon, Surjeet Rajendran, and Yue Zhao, "Radio for hidden-photon dark matter detection," *Phys. Rev. D* **92**, 075012 (2015), [arXiv:1411.7382 \[hep-ph\]](#).
  - [9] J. Suzuki, T. Horie, Y. Inoue, and M. Minowa, "Experimental Search for Hidden Photon CDM in the eV mass range with a Dish Antenna," *JCAP* **09**, 042 (2015), [arXiv:1504.00118 \[hep-ex\]](#).
  - [10] Stefan Knirck, Takayuki Yamazaki, Yoshiki Okesaku, Shoji Asai, Toshitaka Idehara, and Toshiaki Inada, "First results from a hidden photon dark matter search

- in the meV sector using a plane-parabolic mirror system,” *JCAP* **11**, 031 (2018), [arXiv:1806.05120 \[hep-ex\]](#).
- [11] Pierre Brun, Laurent Chevalier, and Christophe Flouzat, “Direct Searches for Hidden-Photon Dark Matter with the SHUKET Experiment,” *Phys. Rev. Lett.* **122**, 201801 (2019), [arXiv:1905.05579 \[hep-ex\]](#).
- [12] A. Phipps *et al.*, “Exclusion Limits on Hidden-Photon Dark Matter near 2 neV from a Fixed-Frequency Superconducting Lumped-Element Resonator,” *Springer Proc. Phys.* **245**, 139–145 (2020), [arXiv:1906.08814 \[astro-ph.CO\]](#).
- [13] Le Hoang Nguyen, Andrei Lobanov, and Dieter Horns, “First results from the WISPDMM radio frequency cavity searches for hidden photon dark matter,” *JCAP* **10**, 014 (2019), [arXiv:1907.12449 \[hep-ex\]](#).
- [14] A. Andrianavalomahefa *et al.* (FUNK Experiment), “Limits from the Funk Experiment on the Mixing Strength of Hidden-Photon Dark Matter in the Visible and Near-Ultraviolet Wavelength Range,” *Phys. Rev. D* **102**, 042001 (2020), [arXiv:2003.13144 \[astro-ph.CO\]](#).
- [15] Nozomu Tomita, Shugo Oguri, Yoshizumi Inoue, Makoto Minowa, Taketo Nagasaki, Jun’ya Suzuki, and Osamu Tajima, “Search for hidden-photon cold dark matter using a K-band cryogenic receiver,” *JCAP* **09**, 012 (2020), [arXiv:2006.02828 \[hep-ex\]](#).
- [16] Akash V. Dixit, Srivatsan Chakram, Kevin He, Ankur Agrawal, Ravi K. Naik, David I. Schuster, and Aaron Chou, “Searching for Dark Matter with a Superconducting Qubit,” *Phys. Rev. Lett.* **126**, 141302 (2021), [arXiv:2008.12231 \[hep-ex\]](#).
- [17] Benjamin Godfrey *et al.*, “Search for dark photon dark matter: Dark E field radio pilot experiment,” *Phys. Rev. D* **104**, 012013 (2021), [arXiv:2101.02805 \[physics.ins-det\]](#).
- [18] Sumita Ghosh, E. P. Ruddy, Michael J. Jewell, Alexander F. Leder, and Reina H. Maruyama, “Searching for dark photons with existing haloscope data,” *Phys. Rev. D* **104**, 092016 (2021), [arXiv:2104.09334 \[hep-ph\]](#).
- [19] Andrea Caputo, Alexander J. Millar, Ciaran A. J. O’Hare, and Edoardo Vitagliano, “Dark photon limits: A handbook,” *Phys. Rev. D* **104**, 095029 (2021), [arXiv:2105.04565 \[hep-ph\]](#).
- [20] Michael A. Fedderke, Peter W. Graham, Derek F. Jackson Kimball, and Saarik Kalia, “Earth as a transducer for dark-photon dark-matter detection,” *Phys. Rev. D* **104**, 075023 (2021), [arXiv:2106.00022 \[hep-ph\]](#).
- [21] Michael A. Fedderke, Peter W. Graham, Derek F. Jackson Kimball, and Saarik Kalia, “Search for dark-photon dark matter in the SuperMAG geomagnetic field dataset,” *Phys. Rev. D* **104**, 095032 (2021), [arXiv:2108.08852 \[hep-ph\]](#).
- [22] R. Cervantes *et al.*, “Search for 70  $\mu\text{eV}$  Dark Photon Dark Matter with a Dielectrically Loaded Multiwavelength Microwave Cavity,” *Phys. Rev. Lett.* **129**, 201301 (2022), [arXiv:2204.03818 \[hep-ex\]](#).
- [23] Shumpei Kotaka *et al.* (DOSUE-RR), “Search for Dark Photon Dark Matter in the Mass Range 74–110  $\mu\text{eV}$  with a Cryogenic Millimeter-Wave Receiver,” *Phys. Rev. Lett.* **130**, 071805 (2023), [arXiv:2205.03679 \[hep-ex\]](#).
- [24] Joshua Berger and Amit Bhoonah, “First Laboratory Bounds on Ultralight Dark Photon Dark Matter from Precision Atomic Spectroscopy,” (2022), [arXiv:2206.06364 \[hep-ph\]](#).
- [25] Haipeng An, Shuailiang Ge, Wen-Qing Guo, Xiaoyuan Huang, Jia Liu, and Zhiyao Lu, “Direct Detection of Dark Photon Dark Matter Using Radio Telescopes,” *Phys. Rev. Lett.* **130**, 181001 (2023), [arXiv:2207.05767 \[hep-ph\]](#).
- [26] Raphael Cervantes, Caterina Braggio, Bianca Giaccone, Daniil Frolov, Anna Grassellino, Roni Harnik, Oleksandr Melnychuk, Roman Pilipenko, Sam Posen, and Alexander Romanenko, “Deepest Sensitivity to Wavelike Dark Photon Dark Matter with SRF Cavities,” (2022), [arXiv:2208.03183 \[hep-ex\]](#).
- [27] Xing Fan, Gerald Gabrielse, Peter W. Graham, Roni Harnik, Thomas G. Myers, Harikrishnan Ramani, Benedict A. D. Sukra, Samuel S. Y. Wong, and Yawen Xiao, “One-Electron Quantum Cyclotron as a Milli-eV Dark-Photon Detector,” *Phys. Rev. Lett.* **129**, 261801 (2022), [arXiv:2208.06519 \[hep-ex\]](#).
- [28] Karthik Ramanathan, Nikita Klimovich, Ritoban Basu Thakur, Byeong Ho Eom, Henry G. LeDuc, Shibo Shu, Andrew D. Beyer, and Peter K. Day, “Wideband Direct Detection Constraints on Hidden Photon Dark Matter with the QUALIPHIDE Experiment,” *Phys. Rev. Lett.* **130**, 231001 (2023), [arXiv:2209.03419 \[astro-ph.CO\]](#).
- [29] Haipeng An, Xingyao Chen, Shuailiang Ge, Jia Liu, and Yan Luo, “Searching for Ultralight Dark Matter Conversion in Solar Corona using LOFAR Data,” (2023), [arXiv:2301.03622 \[hep-ph\]](#).
- [30] A. Romanenko *et al.*, “Search for Dark Photons with Superconducting Radio Frequency Cavities,” *Phys. Rev. Lett.* **130**, 261801 (2023), [arXiv:2301.11512 \[hep-ex\]](#).
- [31] Min Jiang, Taizhou Hong, Dongdong Hu, Yifan Chen, Fengwei Yang, Tao Hu, Xiaodong Yang, Jing Shu, Yue Zhao, and Xinhua Peng, “Search for dark photons with synchronized quantum sensor network,” (2023), [arXiv:2305.00890 \[quant-ph\]](#).
- [32] Zhenxing Tang *et al.*, “SRF Cavity Searches for Dark Photon Dark Matter: First Scan Results,” (2023), [arXiv:2305.09711 \[hep-ex\]](#).
- [33] Faye Bajjali *et al.*, “First results from BRASS-p broadband searches for hidden photon dark matter,” *JCAP* **08**, 077 (2023), [arXiv:2306.05934 \[hep-ex\]](#).
- [34] Ibrahim A. Sulai *et al.*, “Hunt for magnetic signatures of hidden-photon and axion dark matter in the wilderness,” *Phys. Rev. D* **108**, 096026 (2023), [arXiv:2306.11575 \[hep-ph\]](#).
- [35] Tim Schneemann, Kristof Schmieden, and Matthias Schott, “First results of the SUPAX Experiment: Probing Dark Photons,” (2023), [arXiv:2308.08337 \[hep-ex\]](#).
- [36] Stefan Knirck *et al.*, “First Results from a Broadband Search for Dark Photon Dark Matter in the 44 to 52  $\mu\text{eV}$  range with a coaxial dish antenna,” (2023), [arXiv:2310.13891 \[hep-ex\]](#).
- [37] Yifan Chen, Min Jiang, Jing Shu, Xiao Xue, and Yanjie Zeng, “Dissecting axion and dark photon with a network of vector sensors,” *Phys. Rev. Res.* **4**, 033080 (2022), [arXiv:2111.06732 \[hep-ph\]](#).
- [38] T. Nitta *et al.* (ADMX), “Search for a Dark-Matter-Induced Cosmic Axion Background with ADMX,” *Phys. Rev. Lett.* **131**, 101002 (2023), [arXiv:2303.06282 \[hep-ex\]](#).
- [39] I. I. Tkachev, “An Axionic Laser in the Center of a Galaxy?” *Phys. Lett. B* **191**, 41–45 (1987).
- [40] Eric Braaten, Abhishek Mohapatra, and Hong Zhang,

- “Emission of Photons and Relativistic Axions from Axion Stars,” *Phys. Rev. D* **96**, 031901 (2017), arXiv:1609.05182 [hep-ph].
- [41] João G. Rosa and Thomas W. Kephart, “Stimulated Axion Decay in Superradiant Clouds around Primordial Black Holes,” *Phys. Rev. Lett.* **120**, 231102 (2018), arXiv:1709.06581 [gr-qc].
- [42] Mark P. Hertzberg and Enrico D. Schiappacasse, “Dark Matter Axion Clump Resonance of Photons,” *JCAP* **11**, 004 (2018), arXiv:1805.00430 [hep-ph].
- [43] Mateja Boskovic, Richard Brito, Vitor Cardoso, Taishi Ikeda, and Helvi Witek, “Axionic instabilities and new black hole solutions,” *Phys. Rev. D* **99**, 035006 (2019), arXiv:1811.04945 [gr-qc].
- [44] Taishi Ikeda, Richard Brito, and Vitor Cardoso, “Blasts of Light from Axions,” *Phys. Rev. Lett.* **122**, 081101 (2019), arXiv:1811.04950 [gr-qc].
- [45] Andrea Caputo, Marco Regis, Marco Taoso, and Samuel J. Witte, “Detecting the Stimulated Decay of Axions at RadioFrequencies,” *JCAP* **03**, 027 (2019), arXiv:1811.08436 [hep-ph].
- [46] Pierluca Carenza, Alessandro Mirizzi, and Günter Sigl, “Dynamical evolution of axion condensates under stimulated decays into photons,” *Phys. Rev. D* **101**, 103016 (2020), arXiv:1911.07838 [hep-ph].
- [47] Gonzalo Alonso-Álvarez, Rick S. Gupta, Joerg Jaeckel, and Michael Spannowsky, “On the Wondrous Stability of ALP Dark Matter,” *JCAP* **03**, 052 (2020), arXiv:1911.07885 [hep-ph].
- [48] Z. Wang, L. Shao, and L. X. Li, “Resonant instability of axionic dark matter clumps,” *JCAP* **07**, 038 (2020), arXiv:2002.09144 [hep-ph].
- [49] Ariel Arza, Thomas Schwetz, and Elisa Todarello, “How to suppress exponential growth—on the parametric resonance of photons in an axion background,” *JCAP* **10**, 013 (2020), arXiv:2004.01669 [hep-ph].
- [50] D. G. Levkov, A. G. Panin, and I. I. Tkachev, “Radio-emission of axion stars,” *Phys. Rev. D* **102**, 023501 (2020), arXiv:2004.05179 [astro-ph.CO].
- [51] Liang Chen and Thomas W. Kephart, “Stimulated radiation from axion cluster evolution in static spacetimes,” *JCAP* **09**, 034 (2021), arXiv:2007.12337 [hep-ph].
- [52] Xiaolong Du, David J. E. Marsh, Miguel Escudero, Andrew Benson, Diego Blas, Charis Kaur Pooni, and Malcolm Fairbairn, “Soliton Merger Rates and Enhanced Axion Dark Matter Decay,” (2023), arXiv:2301.09769 [astro-ph.CO].
- [53] Miguel Escudero, Charis Kaur Pooni, Malcolm Fairbairn, Diego Blas, Xiaolong Du, and David J. E. Marsh, “Axion Star Explosions: A New Source for Axion Indirect Detection,” (2023), arXiv:2302.10206 [hep-ph].
- [54] Liina M. Chung-Jukko, Eugene A. Lim, David J. E. Marsh, Josu C. Aurrekoetxea, Eloy de Jong, and Bo-Xuan Ge, “Electromagnetic instability of compact axion stars,” *Phys. Rev. D* **108**, L061302 (2023), arXiv:2302.10100 [hep-ph].
- [55] Thomas F. M. Spieksma, Enrico Cannizzaro, Taishi Ikeda, Vitor Cardoso, and Yifan Chen, “Superradiance: Axionic couplings and plasma effects,” *Phys. Rev. D* **108**, 063013 (2023), arXiv:2306.16447 [gr-qc].
- [56] Guenter Sigl, “Radio signatures from conversion of axion-like particle to photons,” <https://agenda.infn.it/event/34125/contributions/206291/> (2023).
- [57] Liang Chen, Da Huang, and Chao-Qiang Geng, “Effects of stimulated emission and superradiant growth of non-spherical axion cluster,” (2023), arXiv:2311.01819 [hep-ph].
- [58] Haoran Di, “Stimulated Decay of Collapsing Axion Stars and Fast Radio Bursts,” (2023), arXiv:2311.06860 [hep-ph].
- [59] P. S. Bhupal Dev, Francesc Ferrer, and Takuya Okawa, “On the Galactic radio signal from stimulated decay of axion dark matter,” (2023), arXiv:2311.13653 [hep-ph].
- [60] Liang Chen and Thomas W. Kephart, “A Review of Axion Lasing in Astrophysics,” (2023), arXiv:2311.16453 [hep-ph].
- [61] Jeff A. Dror, Hitoshi Murayama, and Nicholas L. Rodd, “Cosmic axion background,” *Phys. Rev. D* **103**, 115004 (2021), [Erratum: *Phys.Rev.D* 106, 119902 (2022)], arXiv:2101.09287 [hep-ph].
- [62] C.J. Hogan and M.J. Rees, “Axion miniclusters,” *Physics Letters B* **205**, 228–230 (1988).
- [63] Edward W. Kolb and Igor I. Tkachev, “Nonlinear axion dynamics and formation of cosmological pseudosolitons,” *Phys. Rev. D* **49**, 5040–5051 (1994), arXiv:astro-ph/9311037.
- [64] Edward W. Kolb and Igor I. Tkachev, “Axion miniclusters and Bose stars,” *Phys. Rev. Lett.* **71**, 3051–3054 (1993), arXiv:hep-ph/9303313.
- [65] Kathryn M. Zurek, Craig J. Hogan, and Thomas R. Quinn, “Astrophysical Effects of Scalar Dark Matter Miniclusters,” *Phys. Rev. D* **75**, 043511 (2007), arXiv:astro-ph/0607341.
- [66] I. I. Tkachev, “Fast Radio Bursts and Axion Miniclusters,” *JETP Lett.* **101**, 1–6 (2015), arXiv:1411.3900 [astro-ph.HE].
- [67] Peter Tinyakov, Igor Tkachev, and Konstantin Zioutas, “Tidal streams from axion miniclusters and direct axion searches,” *JCAP* **01**, 035 (2016), arXiv:1512.02884 [astro-ph.CO].
- [68] Luca Visinelli, Sebastian Baum, Javier Redondo, Katherine Freese, and Frank Wilczek, “Dilute and dense axion stars,” *Phys. Lett. B* **777**, 64–72 (2018), arXiv:1710.08910 [astro-ph.CO].
- [69] D. G. Levkov, A. G. Panin, and I. I. Tkachev, “Gravitational Bose-Einstein condensation in the kinetic regime,” *Phys. Rev. Lett.* **121**, 151301 (2018), arXiv:1804.05857 [astro-ph.CO].
- [70] Alejandro Vaquero, Javier Redondo, and Julia Stadler, “Early seeds of axion miniclusters,” *JCAP* **04**, 012 (2019), arXiv:1809.09241 [astro-ph.CO].
- [71] Malte Buschmann, Joshua W. Foster, and Benjamin R. Safdi, “Early-Universe Simulations of the Cosmological Axion,” *Phys. Rev. Lett.* **124**, 161103 (2020), arXiv:1906.00967 [astro-ph.CO].
- [72] Asimina Arvanitaki, Savvas Dimopoulos, Marios Galanis, Luis Lehner, Jedidiah O. Thompson, and Ken Van Tilburg, “Large-misalignment mechanism for the formation of compact axion structures: Signatures from the QCD axion to fuzzy dark matter,” *Phys. Rev. D* **101**, 083014 (2020), arXiv:1909.11665 [astro-ph.CO].
- [73] Benedikt Eggemeier, Javier Redondo, Klaus Dolag, Jens C. Niemeyer, and Alejandro Vaquero, “First Simulations of Axion Minicluster Halos,” *Phys. Rev. Lett.* **125**, 041301 (2020), arXiv:1911.09417 [astro-ph.CO].
- [74] David Ellis, David J. E. Marsh, and Christoph Behrens, “Axion Miniclusters Made Easy,” *Phys. Rev. D* **103**,

- 083525 (2021), arXiv:2006.08637 [astro-ph.CO].
- [75] Huangyu Xiao, Ian Williams, and Matthew McQuinn, “Simulations of axion minihalos,” *Phys. Rev. D* **104**, 023515 (2021), arXiv:2101.04177 [astro-ph.CO].
- [76] David Ellis, David J. E. Marsh, Benedikt Eggemeier, Jens Niemeyer, Javier Redondo, and Klaus Dolag, “Structure of axion miniclusters,” *Phys. Rev. D* **106**, 103514 (2022), arXiv:2204.13187 [hep-ph].
- [77] R. L. Workman and Others (Particle Data Group), “Review of Particle Physics,” *PTEP* **2022**, 083C01 (2022).
- [78] Marco Cirelli, Gennaro Corcella, Andi Hektor, Gert Hutsi, Mario Kadastik, Paolo Panci, Martti Raidal, Filippo Sala, and Alessandro Strumia, “PPPC 4 DM ID: A Poor Particle Physicist Cookbook for Dark Matter Indirect Detection,” *JCAP* **03**, 051 (2011), [Erratum: *JCAP* **10**, E01 (2012)], arXiv:1012.4515 [hep-ph].
- [79] Yuriy Pischalnikov, Evgueni Borissov, Ivan Gonin, Jeremiah Holzbauer, Timergali Khabiboulline, Warren Schappert, Samuel Smith, and Jae-Chul Yun, “Design and Test of the Compact Tuner for Narrow Bandwidth SRF Cavities,” in *6th International Particle Accelerator Conference* (2015) p. WEPTY035.
- [80] Y. Pischalnikov, D. Bice, A. Grassellino, T. Khabiboulline, O. Melnychuk, R. Pilipenko, S. Posen, O. Pronichev, and A. Romanenko, “Operation of an SRF Cavity Tuner Submerged into Liquid Helium,” (2019).
- [81] Zheng-Hui Mi *et al.*, “Design and test of frequency tuner for CAEP high power THz free-electron laser,” *Chin. Phys. C* **39**, 028102 (2015), arXiv:1404.3487 [physics.acc-ph].
- [82] Na Liu, Yi Sun, Guang-Wei Wang, Zheng-Hui Mi, Hai-Ying Lin, Qun-Yao Wang, Rong Liu, and Xin-Peng Ma, “Tuner control system of Spoke012 SRF cavity for C-ADS injector I,” *Chin. Phys. C* **40**, 097001 (2016), arXiv:1602.02125 [physics.acc-ph].
- [83] Dejun Zhou *et al.*, “Vertical Test System for Superconducting RF Cavities at Peking University,” in *18th International Conference on RF Superconductivity* (2018) p. TUPB099.
- [84] Xinying Zhang *et al.*, “The mechanical design, fabrication and tests of dressed 650 MHz 2-cell superconducting cavities for CEPC,” *Nucl. Instrum. Meth. A* **1031**, 166590 (2022).
- [85] Zhitao Yang, Jiankui Hao, Shengwen Quan, Lin Lin, Fang Wang, Fei Jiao, and Kexin Liu, “Effective medium temperature baking of 1.3 ghz single cell srf cavities,” *Physica C: Superconductivity and its Applications* **599**, 1354092 (2022).
- [86] Gai Wang, Shengwen Quan, Lin Lin, Manqian Ren, Jiankui Hao, Fang Wang, Fei Jiao, Feng Zhu, Senlin Huang, Xueqing Yan, and Kun Zhu, “Nb3Sn cavities coated by tin vapor diffusion method at peking university,” *Applied Sciences* **13** (2023), 10.3390/app13158618.
- [87] Klaus Ehret *et al.*, “New ALPS Results on Hidden-Sector Lightweights,” *Phys. Lett. B* **689**, 149–155 (2010), arXiv:1004.1313 [hep-ex].
- [88] A. Wagner *et al.* (ADMX), “A Search for Hidden Sector Photons with ADMX,” *Phys. Rev. Lett.* **105**, 171801 (2010), arXiv:1007.3766 [hep-ex].
- [89] M. Betz and F. Caspers, “A microwave paraphoton and axion detection experiment with 300 db electromagnetic shielding at 3 ghz,” (2012), arXiv:1207.3275 [physics.ins-det].
- [90] T. Inada, T. Namba, S. Asai, T. Kobayashi, Y. Tanaka, K. Tamasaku, K. Sawada, and T. Ishikawa, “Results of a Search for Paraphotons with Intense X-ray Beams at SPring-8,” *Phys. Lett. B* **722**, 301–304 (2013), arXiv:1301.6557 [physics.ins-det].
- [91] M. Betz, F. Caspers, M. Gasior, M. Thumm, and S. W. Rieger, “First results of the CERN Resonant Weakly Interacting sub-eV Particle Search (CROWS),” *Phys. Rev. D* **88**, 075014 (2013), arXiv:1310.8098 [physics.ins-det].
- [92] Stephen R. Parker, John G. Hartnett, Rhys G. Povey, and Michael E. Tobar, “Cryogenic resonant microwave cavity searches for hidden sector photons,” *Phys. Rev. D* **88**, 112004 (2013), arXiv:1410.5244 [hep-ex].
- [93] S. J. Plimpton and W. E. Lawton, “A Very Accurate Test of Coulomb’s Law of Force Between Charges,” *Phys. Rev.* **50**, 1066 (1936).
- [94] D. F. Bartlett and S. Loegl, “LIMITS ON AN ELECTROMAGNETIC FIFTH FORCE,” *Phys. Rev. Lett.* **61**, 2285–2287 (1988).
- [95] Liang-Cheng Tu, Jun Luo, and G. T. Gillies, “The mass of the photon,” *Rept. Prog. Phys.* **68**, 77–130 (2005).
- [96] Joerg Jaeckel and Sabyasachi Roy, “Spectroscopy as a test of Coulomb’s law: A Probe of the hidden sector,” *Phys. Rev. D* **82**, 125020 (2010), arXiv:1008.3536 [hep-ph].
- [97] D. Kroff and P. C. Malta, “Constraining hidden photons via atomic force microscope measurements and the Plimpton-Lawton experiment,” *Phys. Rev. D* **102**, 095015 (2020), arXiv:2008.02209 [hep-ph].
- [98] Samuel D. McDermott and Samuel J. Witte, “Cosmological evolution of light dark photon dark matter,” *Phys. Rev. D* **101**, 063030 (2020), arXiv:1911.05086 [hep-ph].
- [99] Andrea Caputo, Hongwan Liu, Siddharth Mishra-Sharma, and Joshua T. Ruderman, “Dark Photon Oscillations in Our Inhomogeneous Universe,” *Phys. Rev. Lett.* **125**, 221303 (2020), arXiv:2002.05165 [astro-ph.CO].
- [100] Ankur Agrawal, Akash V. Dixit, Tanay Roy, Srivatsan Chakram, Kevin He, Ravi K. Naik, David I. Schuster, and Aaron Chou, “Stimulated emission of signal photons from dark matter waves,” (2023), arXiv:2305.03700 [quant-ph].
- [101] Huaixiu Zheng, Matti Silveri, R.T. Brierley, S.M. Girvin, and K.W. Lehnert, “Accelerating dark-matter axion searches with quantum measurement technology,” (2016), arXiv:1607.02529 [hep-ph].
- [102] M. Malnou, D. A. Palken, B. M. Brubaker, Leila R. Vale, Gene C. Hilton, and K. W. Lehnert, “Squeezed vacuum used to accelerate the search for a weak classical signal,” *Phys. Rev. X* **9**, 021023 (2019), [Erratum: *Phys.Rev.X* **10**, 039902 (2020)], arXiv:1809.06470 [quant-ph].
- [103] K. M. Backes *et al.* (HAYSTAC), “A quantum-enhanced search for dark matter axions,” *Nature* **590**, 238–242 (2021), arXiv:2008.01853 [quant-ph].
- [104] K. W. Lehnert, “Quantum enhanced metrology in the search for fundamental physical phenomena,” (2021) arXiv:2110.04912 [quant-ph].
- [105] Stephen E. Kuenstner *et al.*, “Quantum metrology of low frequency electromagnetic modes with frequency upconverters,” (2022), arXiv:2210.05576 [quant-ph].

- [106] M. J. Jewell *et al.* (HAYSTAC), “New results from HAYSTAC’s phase II operation with a squeezed state receiver,” *Phys. Rev. D* **107**, 072007 (2023), [arXiv:2301.09721 \[hep-ex\]](#).
- [107] Xiang Li, Maxim Goryachev, Yiqiu Ma, Michael E. Tobar, Chunnong Zhao, Rana X. Adhikari, and Yanbei Chen, “Broadband sensitivity improvement via coherent quantum feedback with PT symmetry,” (2020), [arXiv:2012.00836 \[quant-ph\]](#).
- [108] Yifan Chen, Minyuan Jiang, Yiqiu Ma, Jing Shu, and Yuting Yang, “Axion haloscope array with PT symmetry,” *Phys. Rev. Res.* **4**, 023015 (2022), [arXiv:2103.12085 \[hep-ph\]](#).
- [109] K. Wurtz, B. M. Brubaker, Y. Jiang, E. P. Ruddy, D. A. Palken, and K. W. Lehnert, “Cavity Entanglement and State Swapping to Accelerate the Search for Axion Dark Matter,” *PRX Quantum* **2**, 040350 (2021), [arXiv:2107.04147 \[quant-ph\]](#).
- [110] Yue Jiang, Elizabeth P. Ruddy, Kyle O. Quinlan, Maxime Malnou, Nicholas E. Frattini, and Konrad W. Lehnert, “Accelerated Weak Signal Search Using Mode Entanglement and State Swapping,” *PRX Quantum* **4**, 020302 (2023), [arXiv:2211.10403 \[quant-ph\]](#).
- [111] Yifan Chen, Chunlong Li, Yuxin Liu, Jing Shu, Yuting Yang, and Yanjie Zeng, “Simultaneous Resonant and Broadband Detection for Dark Sectors,” (2023), [arXiv:2309.12387 \[hep-ph\]](#).
- [112] Joshua W. Foster, Yonatan Kahn, Rachel Nguyen, Nicholas L. Rodd, and Benjamin R. Safdi, “Dark Matter Interferometry,” *Phys. Rev. D* **103**, 076018 (2021), [arXiv:2009.14201 \[hep-ph\]](#).
- [113] S. Afach *et al.* (GNOME), “What can a GNOME do? Search targets for the Global Network of Optical Magnetometers for Exotic physics searches,” *Annalen Phys.* **2023**, 2300083 (2023), [arXiv:2305.01785 \[hep-ph\]](#).
- [114] Conner Dailey, Colin Bradley, Derek F. Jackson Kimball, Ibrahim A. Sulai, Szymon Pustelny, Arne Wickensbrock, and Andrei Derevianko, “Quantum sensor networks as exotic field telescopes for multimessenger astronomy,” *Nature Astron.* **5**, 150–158 (2021), [arXiv:2002.04352 \[astro-ph.IM\]](#).
- [115] Maxim Goryachev, Ben McAllister, and Michael E. Tobar, “Axion detection with precision frequency metrology,” *Phys. Dark Univ.* **26**, 100345 (2019), [Erratum: *Phys. Dark Univ.* **32**, 100787 (2021)], [arXiv:1806.07141 \[physics.ins-det\]](#).
- [116] Asher Berlin, Raffaele Tito D’Agnolo, Sebastian A.R. Ellis, Christopher Nantista, Jeffrey Neilson, Philip Schuster, Sami Tantawi, Natalia Toro, and Kevin Zhou, “Axion Dark Matter Detection by Superconducting Resonant Frequency Conversion,” *JHEP* **07**, 088 (2020), [arXiv:1912.11048 \[hep-ph\]](#).
- [117] Robert Lasenby, “Microwave cavity searches for low-frequency axion dark matter,” *Phys. Rev. D* **102**, 015008 (2020), [arXiv:1912.11056 \[hep-ph\]](#).
- [118] Catriona A. Thomson, Ben T. McAllister, Maxim Goryachev, Eugene N. Ivanov, and Michael E. Tobar, “Upconversion Loop Oscillator Axion Detection Experiment: A Precision Frequency Interferometric Axion Dark Matter Search with a Cylindrical Microwave Cavity,” *Phys. Rev. Lett.* **126**, 081803 (2021), [Erratum: *Phys. Rev. Lett.* **127**, 019901 (2021)], [arXiv:1912.07751 \[hep-ex\]](#).
- [119] Asher Berlin, Raffaele Tito D’Agnolo, Sebastian A. R. Ellis, and Kevin Zhou, “Heterodyne broadband detection of axion dark matter,” *Phys. Rev. D* **104**, L111701 (2021), [arXiv:2007.15656 \[hep-ph\]](#).
- [120] Michael E. Tobar, Catriona A. Thomson, William M. Campbell, Aaron Quiskamp, Jeremy F. Bourhill, Benjamin T. McAllister, Eugene N. Ivanov, and Maxim Goryachev, “Comparing Instrument Spectral Sensitivity of Dissimilar Electromagnetic Haloscopes to Axion Dark Matter and High Frequency Gravitational Waves,” *Symmetry* **14**, 2165 (2022), [arXiv:2209.03004 \[physics.ins-det\]](#).
- [121] J. F. Bourhill, E. C. I. Paterson, M. Goryachev, and M. E. Tobar, “Searching for ultralight axions with twisted cavity resonators of anyon rotational symmetry with bulk modes of nonzero helicity,” *Phys. Rev. D* **108**, 052014 (2023), [arXiv:2208.01640 \[hep-ph\]](#).
- [122] Catriona A. Thomson, Maxim Goryachev, Ben T. McAllister, Eugene N. Ivanov, Paul Altin, and Michael E. Tobar, “Searching for low-mass axions using resonant upconversion,” *Phys. Rev. D* **107**, 112003 (2023), [arXiv:2301.06778 \[hep-ex\]](#).
- [123] Asher Berlin, Diego Blas, Raffaele Tito D’Agnolo, Sebastian A. R. Ellis, Roni Harnik, Yonatan Kahn, Jan Schütte-Engel, and Michael Wentzel, “Electromagnetic cavities as mechanical bars for gravitational waves,” *Phys. Rev. D* **108**, 084058 (2023), [arXiv:2303.01518 \[hep-ph\]](#).
- [124] T. Damour and Alexander M. Polyakov, “String theory and gravity,” *Gen. Rel. Grav.* **26**, 1171–1176 (1994), [arXiv:gr-qc/9411069](#).
- [125] Ignatios Antoniadis, Savvas Dimopoulos, and Amit Giveon, “Little string theory at a TeV,” *JHEP* **05**, 055 (2001), [arXiv:hep-th/0103033](#).
- [126] Naomi Gendler, David J. E. Marsh, Liam McAllister, and Jakob Moritz, “Glimmers from the Axiverse,” (2023), [arXiv:2309.13145 \[hep-th\]](#).
- [127] Emile Mathieu, “Memoire sur le mouvement vibratoire d’une membrane de forme elliptique.” *Journal de Mathematiques Pures et Appliquees* **13**, 137–203 (1868).
- [128] Ivana Kovacic, Richard Rand, and Si Mohamed Sah, “Mathieu’s equation and its generalizations: overview of stability charts and their features,” *Applied Mechanics Reviews* **70**, 020802 (2018).
- [129] James Johnston Stoker, “Nonlinear vibrations in mechanical and electrical systems,” (No Title) (1950).
- [130] Gizem Acar and Brian F Feeny, “Floquet-based analysis of general responses of the mathieu equation,” *Journal of Vibration and Acoustics* **138**, 041017 (2016).
- [131] Yifan Chen, Xiao Xue, and Vitor Cardoso, “Black Holes as Neutrino Factories,” (2023), [arXiv:2308.00741 \[hep-ph\]](#).
- [132] I. I. Tkachev, “On the possibility of Bose star formation,” *Phys. Lett. B* **261**, 289–293 (1991).
- [133] Jan Veltmaat, Bodo Schwabe, and Jens C. Niemeyer, “Baryon-driven growth of solitonic cores in fuzzy dark matter halos,” *Phys. Rev. D* **101**, 083518 (2020), [arXiv:1911.09614 \[astro-ph.CO\]](#).
- [134] Jiajun Chen, Xiaolong Du, Erik W. Lentz, David J. E. Marsh, and Jens C. Niemeyer, “New insights into the formation and growth of boson stars in dark matter halos,” *Phys. Rev. D* **104**, 083022 (2021), [arXiv:2011.01333 \[astro-ph.CO\]](#).
- [135] Jiajun Chen, Xiaolong Du, Erik W. Lentz, and David J. E. Marsh, “Relaxation times for Bose-Einstein con-

- densation by self-interaction and gravity,” *Phys. Rev. D* **106**, 023009 (2022), arXiv:2109.11474 [astro-ph.CO].
- [136] Dmitry Budker, Joshua Eby, Marco Gorghetto, Minyuan Jiang, and Gilad Perez, “A generic formation mechanism of ultralight dark matter solar halos,” *JCAP* **12**, 021 (2023), arXiv:2306.12477 [hep-ph].
- [137] Joshua Eby, Madelyn Leembruggen, Peter Suranyi, and L. C. R. Wijewardhana, “Collapse of Axion Stars,” *JHEP* **12**, 066 (2016), arXiv:1608.06911 [astro-ph.CO].
- [138] D. G. Levkov, A. G. Panin, and I. I. Tkachev, “Relativistic axions from collapsing Bose stars,” *Phys. Rev. Lett.* **118**, 011301 (2017), arXiv:1609.03611 [astro-ph.CO].
- [139] Gilly Elor, Nicholas L. Rodd, and Tracy R. Slatyer, “Multistep cascade annihilations of dark matter and the Galactic Center excess,” *Phys. Rev. D* **91**, 103531 (2015), arXiv:1503.01773 [hep-ph].
- [140] Gilly Elor, Nicholas L. Rodd, Tracy R. Slatyer, and Wei Xue, “Model-Independent Indirect Detection Constraints on Hidden Sector Dark Matter,” *JCAP* **06**, 024 (2016), arXiv:1511.08787 [hep-ph].
- [141] Julio F. Navarro, Carlos S. Frenk, and Simon D. M. White, “The Structure of cold dark matter halos,” *Astrophys. J.* **462**, 563–575 (1996), arXiv:astro-ph/9508025.
- [142] A. Chen *et al.* (DES), “Constraints on dark matter to dark radiation conversion in the late universe with DES-Y1 and external data,” *Phys. Rev. D* **103**, 123528 (2021), arXiv:2011.04606 [astro-ph.CO].
- [143] IAU SOFA Board, “Iau sofa software collection,” .

---

## Supplemental Materials: SRF Cavity as Galactic Dark Photon Telescope

### I: CAVITY RESPONSE TO GENERAL DARK PHOTON BACKGROUND

This section delves into a comprehensive calculation of the signal generated in a cavity by a general background of kinetically mixing dark photons. The background is characterized by an angular-dependent power spectral density (PSD) function. Unlike non-relativistic dark photon dark matter (DPDM), a relativistic dark photon can exhibit a position-dependent phase factor in the cavity signal, owing to its de Broglie wavelength being comparable to the cavity size.

#### General Dark Photon Background

We examine a dark photon  $X^\mu$  with mass  $m_X$  and momentum  $\vec{k}$  in the Galactic frame. As a massive vector boson, it encompasses three degrees of freedom: the longitudinal (L) and two transverse modes (left/right-hand circularly polarized states,  $T_l$  and  $T_r$ , respectively). Polarization (P) basis vectors  $e_P^\mu$  are introduced, conforming to the unitary gauge and normalization condition  $e_P^\mu e_\mu^P = -1$ , and takes the forms:

$$\begin{aligned} e_{T_l}^\mu &= (0, (\hat{e}_{T_1} - i\hat{e}_{T_2})/\sqrt{2}), \\ e_{T_r}^\mu &= (0, (\hat{e}_{T_1} + i\hat{e}_{T_2})/\sqrt{2}), \\ e_L^\mu &= (k/m_X, \omega \hat{k}/m_X). \end{aligned} \tag{S1}$$

Here,  $\hat{k}$ ,  $\hat{e}_{T_1}$  and  $\hat{e}_{T_2}$  form a Cartesian orthogonal basis parameterized in the Galactic frame as:

$$\begin{aligned} \hat{k} &\equiv (\sin \theta_k \cos \phi_k, \sin \theta_k \sin \phi_k, \cos \theta_k), \\ \hat{e}_{T_1} &\equiv (\cos \theta_k \cos \phi_k, \cos \theta_k \sin \phi_k, -\sin \theta_k), \\ \hat{e}_{T_2} &\equiv (-\sin \phi_k, \cos \phi_k, 0), \end{aligned} \tag{S2}$$

using spherical coordinate  $(\theta_k, \phi_k)$ . The dark photon energy  $\omega$  satisfies the dispersion relation  $\omega^2 = |\vec{k}|^2 + m_X^2$ . It’s notable that the spatial components of the transverse basis,  $\vec{e}_{T_{l/r}} \equiv (\hat{e}_{T_1} \mp i\hat{e}_{T_2})/\sqrt{2}$ , are unit vectors, while the longitudinal one is  $|\vec{e}_L| = \omega/m_X$ . Therefore, for relativistic dark photons, the longitudinal mode can produce significantly larger signals in a cavity, as will be discussed further.

A general time-dependent dark photon field is a sum of dark photons with various momenta:

$$X^\mu(t, \vec{x}) = \int \sum_P X_P(\omega, \hat{k}) e_P^\mu e^{i(\omega t - \vec{k} \cdot \vec{x})} \frac{d\omega}{2\pi} d^2\hat{k}, \tag{S3}$$

where  $X_P(\omega, \hat{k})$  represents the corresponding field amplitude. As the production mechanisms considered in this study are based on incoherent decay of dark matter from different locations in the galaxy, the dark photon field should be

a sum of incoherent waves. Thus, we treat  $X_P(\omega, \hat{k})$  as a Gaussian stochastic field, characterized by its two-point correlation function:

$$\langle X_P(\omega, \hat{k}) X_{P'}(\omega', \hat{k}')^* \rangle = \delta(\omega - \omega') \delta^2(\hat{k} - \hat{k}') \delta_{PP'} \mathcal{S}_P(\omega, \hat{k}). \quad (\text{S4})$$

In this expression,  $\langle \dots \rangle$  denotes the ensemble average.  $\mathcal{S}_P(\omega, \hat{k})$  is the dark photon PSD, reflecting the dark photon spectrum from the direction  $\hat{k}$  of polarization mode  $P$ . The normalization of the PSD leads to the dark photon energy density

$$\rho_X = \frac{1}{4\pi^2} \int \omega^2 \sum_P \mathcal{S}_P(\omega, \hat{k}) d\omega d^2\hat{k}. \quad (\text{S5})$$

### Cavity Response to Dark Photons

We analyze the behavior of a cavity during the transit of a dark photon, particularly in the context of kinetic mixing with electromagnetic photons  $A^\mu$ . The interaction Lagrangian in the interaction picture is given by  $\epsilon m_X^2 A^\mu X_\mu$ , where  $\epsilon$  represents the kinetic mixing coefficient. In this scenario, the dark photon functions as an effective current, altering the electromagnetic field equation:

$$\square \vec{E}(t, \vec{x}) = \epsilon m_X^2 \partial_t \vec{X}. \quad (\text{S6})$$

Here, the electric field  $\vec{E}$  can be expressed as a sum of orthogonal eigenmodes within the cavity:

$$\vec{E}(t, \vec{x}) = \sum_n \xi_n(t) \vec{E}_n(\vec{x}). \quad (\text{S7})$$

The mode basis functions  $\vec{E}_n$  comply with:

$$(\nabla^2 + \omega_n^2) \vec{E}_n(\vec{x}) = 0, \quad \int_V \vec{E}_n \cdot \vec{E}_m^* d^3\vec{x} = \delta_{mn}, \quad (\text{S8})$$

where  $\omega_n$  are the resonant frequencies, and  $\xi_n(t)$  describe their temporal evolutions. Integrating Eq. (S6) and (S7) into Eq. (S6) and considering cavity dissipation, we can solve for  $\xi_0$  in the frequency domain as detailed in [32]:

$$\xi_0(\omega) = \frac{i \epsilon m_X^2 \omega}{\omega^2 - \omega_0^2 + i\omega_0\omega/Q_L} \sum_P \int X_P(\omega, \hat{k}) d^2\hat{k} \int_V e^{-i\vec{k}\cdot\vec{x}} \vec{e}_P \cdot \vec{E}_0(\vec{x}) d^3\vec{x}. \quad (\text{S9})$$

Here,  $Q_L$  is the loaded quality factor of the  $\vec{E}_0$  mode. The two-point correlation function of  $\xi_i(\omega)$  contributes to the signal power in the cavity as  $\langle \xi_0^P(\omega) \xi_0^{P'}(\omega')^* \rangle = \delta_{PP'} \delta(\omega - \omega') S_{\text{sig}}^P(\omega)$ . Utilizing Eq. (S4), we obtain the expression for  $S_{\text{sig}}^P$ :

$$S_{\text{sig}}^P(\omega) = \frac{\epsilon^2 m_X^4 V |\vec{e}_P|^2}{4(\omega - \omega_0)^2 + \omega_0^2/Q_L^2} \int \mathcal{S}_P(\omega, \hat{k}) C_P(\hat{k}) d^2\hat{k}, \quad (\text{S10})$$

where we apply  $Q_L \gg 1$  to simplify the Lorentz resonant factor. The dark photon overlapping form factor  $C_P(\hat{k})$  is defined as:

$$C_P(\hat{k}) \equiv \frac{1}{V} \left| \int_V e^{-i\vec{k}\cdot\vec{x}} \hat{e}_P(\hat{k}) \cdot \vec{E}_0(\vec{x}) d^3\vec{x} \right|^2. \quad (\text{S11})$$

As previously discussed, the factor  $|\vec{e}_P|^2$  in Eq. (S10) enhances the longitudinal mode's contribution to the signal for relativistic dark photons. The form factor is maximized when the dark photon wavefunction, proportional to  $\hat{e}_P$ , aligns with the electric field  $\vec{E}_0$ . Compared to non-relativistic DPDM [32], the form factor here includes an additional phase factor  $e^{-i\vec{k}\cdot\vec{x}}$ , which could potentially suppress the signal. As the  $\text{TM}_{010}$  mode of the axis-symmetric cavity in our study lacks an azimuthal component, the two transverse modes,  $T_l$  and  $T_r$ , have identical form factors.

## II: GALACTIC DARK PHOTONS FROM DARK MATTER DECAY

This section delves into the production of galactic dark photons as a result of dark matter decay. We initially explore various coupling types, each leading to distinct dominant polarization modes. When the coupling strength is significant, two-body decay can exponentially deplete the scalar dark matter background; otherwise, it may yield a dark photon flux too weak for detection. We then introduce two alternative production mechanisms, detailing how they can generate observable galactic dark photons.

### Perturbative Decay Widths

We examine three interaction types between a scalar field  $\Phi$  and dark photons: dark Higgs-like  $g_H \Phi X_\mu X^\mu$ , dilaton-like  $g_d \Phi X_{\mu\nu} X^{\mu\nu}$ , and axion-like  $g_a \Phi X_{\mu\nu} \tilde{X}^{\mu\nu}$ . Assuming the scalar mass  $m_\Phi$  is much larger than the dark photon mass  $m_X$ , the corresponding decay widths for either longitudinal or transverse modes are summarized in Table S1. The dark Higgs-like interaction, originating from spontaneous symmetry breaking and characterized by  $g_H = \sqrt{2}em_X$  (with  $e$  as the dark charge of the dark Higgs), predominantly decays into longitudinal modes. Both dilaton-like and axion-like interactions, frequently proposed in theories featuring extra dimensions [1, 3, 124–126], mainly produce transverse modes of dark photons. In our analysis, we consider purely longitudinal or transverse modes of dark photons following dark matter decay.

	$g_H \Phi X_\mu X^\mu$	$g_d \Phi X_{\mu\nu} X^{\mu\nu}$	$g_a \Phi X_{\mu\nu} \tilde{X}^{\mu\nu}$
$\Gamma(\Phi \rightarrow X_T X_T)$	$\frac{g_H^2}{4\pi m_\Phi}$	$\frac{m_\Phi^4 g_d^2 m_X^4}{2m_X^4 2\pi m_\Phi}$	$\frac{g_a^2 m_\Phi^3}{4\pi}$
$\Gamma(\Phi \rightarrow X_L X_L)$	$\frac{m_\Phi^4 g_H^2}{8m_X^4 4\pi m_\Phi}$	$\frac{g_d^2 m_X^4}{2\pi m_\Phi}$	/

TABLE S1: Perturbative decay widths for three distinct interactions between a scalar particle  $\Phi$  and dark photons  $X^\mu$ , focusing on the relativistic limit where the final-state dark photons are significantly lighter than the scalar ( $m_X \ll m_\Phi$ ). The total decay widths are categorized based on the polarization modes of the dark photons, distinguishing between transverse and longitudinal modes.

### Two-body Parametric Decay

In scenarios with a coherent oscillating scalar background  $\Phi = \Phi_0 \cos m_\Phi t$ , the decay rate of a scalar into dark photons with the same momentum as previously generated particles is enhanced due to Bose enhancement in the final state phase space, fostering exponential growth in dark photon density. This phenomenon is evident from the dark photon's equation of motion:

$$\ddot{X} + (k^2 + m_X^2 + q \cos m_\Phi t) X = 0, \quad (\text{S12})$$

where  $q$  is a parameter dependent on the coupling between  $\Phi$  and the dark photon. For example,  $q = g_a \Phi_0 k m_\Phi$  in the case of axion-photon coupling, and  $q = g_H \Phi_0$  or  $g_H \Phi_0 m_\Phi^2 / (2m_X^2)$  for dark Higgs-like coupling with transverse and longitudinal modes of dark photon, respectively. The above equation (S12) is analogous to the Mathieu equation, which predicts several instability bands in momentum space [127, 128]. Within these bands, one can expect a significant exponential increase in occupation number, a process typically referred to as parametric resonance. The first band typically has the broadest bandwidth and can be analyzed either numerically or through semi-analytic methods like Floquet theory [42, 129, 130] or non-adiabatic condition checks [131], yielding the range:

$$\frac{m_\Phi^2}{4} - q \leq k^2 + m_X^2 \leq \frac{m_\Phi^2}{4} + q. \quad (\text{S13})$$

Under the assumption that  $m_\Phi \gg m_X$ , the first instability band centers around  $m_\Phi$  with a bandwidth  $\Delta k = g_a \Phi_0 m_\Phi$  for axion-photon coupling, and  $\Delta k = 2g_H \Phi_0 / m_\Phi$  and  $g_H \Phi_0 m_\Phi / m_X^2$  for the transverse and longitudinal modes in dark Higgs coupling. Both numerical and Floquet theory analyses suggest that the exponential growth rate approximately

equals the bandwidth  $\Delta k$ . Consequently, the production of dark photons is predominantly in the longitudinal mode from dark-Higgs coupling, consistent with perturbative analysis.

Alternatively, this exponential growth can be effectively described in kinetic theory, where the decay rate is significantly amplified by the dark photon occupation number  $f_X$  [39, 45–47, 59]:

$$\frac{dn_X}{dt} = 2n_\Phi(1 + 2f_X)\Gamma(\Phi \rightarrow XX) + \dots \quad (\text{S14})$$

Here,  $n_X$  and  $n_\Phi$  represent the number densities of dark photons and the scalar dark matter background, respectively, with  $\dots$  denoting other terms influencing the density. In the narrow bandwidth limit, where  $\Delta k \ll m_\Phi/2$ , the occupation number can be approximated as  $f_X \approx 2\pi^2 n_X / (k^2 \Delta k)$ . The exponential growth factor, derived directly from Eq. (S14), is  $8\pi^2 n_\Phi \Gamma(\Phi \rightarrow XX) / (k^2 \Delta k)$ . This factor approximates to  $\Delta k$  when considering  $n_\Phi \approx \rho_\Phi / m_\Phi \approx m_\Phi \Phi_0^2 / 2$  and  $k \approx m_\Phi / 2$ .

We can estimate the critical energy density of dark photons,  $\rho_{\text{crit}}$ , at which Bose-enhanced decay becomes significant, by considering the occupation number  $f_X$  to be of the order of 1. This leads to the following expression:

$$\rho_{\text{crit}} = \int \frac{4\pi k^2 dk}{(2\pi)^3} \sqrt{k^2 + m_X^2} \approx 2.1 \times 10^{-11} \rho_\gamma. \quad (\text{S15})$$

where we assume the bandwidth of the dark photon,  $\Delta k$ , to be  $10^{-3} k$  and  $k \approx m_\Phi/2 \approx 2\pi 1.3 \text{ GHz}$ . Here,  $\rho_\gamma \approx 0.26 \text{ eV/cm}^3$  represents the local electromagnetic photon density [77]. When the density of dark photons surpasses  $\rho_{\text{crit}}$ , their population begins to exhibit exponential growth, which could potentially disrupt the stability of the dark matter halo. To mitigate this, a suitably weak coupling constant must be chosen. For example, in the case of axion-dark photon coupling, the resulting local dark photon density from perturbative decay can be approximated as  $\rho_X \sim \Gamma(\Phi \rightarrow XX) \rho_\odot^{\text{DM}} r_G \propto g_a^2$ . In this approximation,  $\rho_\odot^{\text{DM}} = 0.3 \text{ GeV/cm}^3$  is the local dark matter density, and we use  $r_G = 10 \text{ kpc}$  as a typical galactic scale. This suggests that  $g_a$  should be less than approximately  $10^{-6} \text{ GeV}^{-1}$  to keep  $\rho_X$  below  $\rho_{\text{crit}}$ . However, it's still possible for an initially small dark photon density to grow beyond  $\rho_{\text{crit}}$  at a later stage. To suppress such growth, one must consider dissipative effects on the dark photon density, such as an escape rate due to the finite size of the boson background or gravitational redshift from the dark matter profile [49], both of which impart an exponential decay rate with a factor of  $1/R$ , where  $R$  denotes the typical length scale of the boson background or gravitational potential. By comparing the exponential growth rate, roughly  $\Delta k$ , with the decay rate of  $1/R$ , a similar condition is deduced for axion-photon coupling when considering gravitational redshift in the Milky Way:  $g_a \ll 10^{-6} \text{ GeV}^{-1}$  [49]. Nonetheless, within this parameter range, the energy density of the decay products of dark photons is too low to be detectable.

To address the paradox of achieving sufficiently large dark photon fluxes without depleting the dark matter halo, we consider a scenario where a fraction of the dark matter profile consists of scalar clumps. Formed due to early universe's small-scale density perturbations [64, 68, 69, 71], these clumps are considerably denser than the average dark matter halo. The inner region of these clumps, often termed as a boson star, is surrounded by a minicluster [62–76], with densities ranging between that of the boson star and the ambient halo profile. The production of dark photons through Bose-enhancement predominantly occurs within the boson star region, eventually attaining a saturation phase for both the dark photon field and the scalar background [55, 56]. The dark photon field's equilibrium is upheld by equilibrating the parametric decay rate, approximately  $\Delta k$ , with the escape rate from the boson star, about  $1/R_{\text{BS}}$ , where  $R_{\text{BS}} \approx 1/(m_\Phi v_0)$  denotes the boson star radius:

$$\text{Saturation of dark photon:} \quad g_a \Phi_0^{\text{sat}} m_\Phi = m_\Phi v_0. \quad (\text{S16})$$

Here,  $\Phi_0^{\text{sat}}$  symbolizes the saturated boson star field value. The dark photon's bandwidth is characterized by a quality factor  $Q_X = m_\Phi / (2\Delta k) = 1/(2v_0)$ . Assuming  $v_0$  approximates the typical virial velocity of the Milky Way ( $\sim 10^{-3} c$ ), we derive  $Q_X \approx 500$ . The boson star's equilibrium hinges on balancing energy acquisition from accretion in the surrounding minicluster and energy loss due to parametric decay into dark photons. The accretion rate, derived from simulations, is approximately  $1/(0.3 \tau_{\text{rel}})$ , where  $\tau_{\text{rel}} \approx 2\sqrt{2} m_\Phi^3 v_0^2 / (3\sigma_\Phi n_{\text{MC}}^2 \pi^2)$  is the relaxation time from the minicluster [69, 132–136], characterized by number density  $n_{\text{MC}}$  and self-interaction cross-section  $\sigma_\Phi^{\text{SI}}$ . For a quartic self-interaction,  $\sigma_\Phi = m_\Phi^2 / (8\pi f_\Phi^4)$ , with  $f_\Phi$  being the symmetry breaking scale or decay constant for axion-like particles. The parametric decay rate is given by  $2\Gamma(\Phi \rightarrow XX) f_X$ . Consequently, the boson star's equilibrium condition is:

$$\text{Saturation of boson star:} \quad 1/(0.3 \tau_{\text{rel}}) = 2\Gamma(\Phi \rightarrow XX) f_X^{\text{sat}}, \quad (\text{S17})$$

where  $f_X^{\text{sat}}$  is the saturated dark photon occupation number. These two saturation conditions, as outlined in Eq. (S16) and (S17), can dynamically be achieved from an initially low-density boson star. The relaxation process exponentially amplifies the boson star field value, and upon reaching  $\Phi_0^{\text{sat}}$  as per Eq. (S16), the dark photon starts accumulating, rapidly saturating  $f_X^{\text{sat}}$  and subsequently quenching the exponential growth of the boson star. During the saturation phase, the boson star continuously feeds from the minicluster, converting it into steady dark photon fluxes. To contrast with decay from a smoothly distributed dark matter halo, we introduce an effective decay width for the parametric decay process originating from the boson star:

$$\Gamma_{\text{PD}} = \frac{\eta}{0.3\tau_{\text{rel}}}, \quad (\text{S18})$$

where  $\eta$  represents the mass fraction of boson stars in the overall dark matter halo. The effective decay's number spectrum is presumed to be flat within the bandwidth  $\Delta k = m_\Phi/(2Q_X)$ :

$$\frac{dN_{\text{PD}}}{d\omega} = \frac{4Q_X}{m_\Phi} \Theta\left(\frac{m_\Phi}{4Q_X} - \left|\omega - \frac{m_\Phi}{2}\right|\right), \quad (\text{S19})$$

where  $\Theta(\dots)$  represents the Heaviside step function.

The feasibility of the saturated boson star model can be evaluated using a set of realistic benchmark parameters. Consider, for example, a dark matter particle with mass  $m_\Phi = 4\pi 1.3 \text{ GHz}$ , a decay constant  $f_\Phi = 10^9 \text{ GeV}$ , and a minicluster number density  $n_{\text{MC}} = 10^{30} \text{ cm}^{-3}$  [64, 66, 67]. Given these parameters, the relaxation time  $\tau_{\text{rel}}$  is calculated to be approximately  $10^7$  years, significantly shorter than the age of the universe,  $t_U \approx 1.37 \times 10^9$  years [77]. Setting the boson star mass fraction of the total dark matter mass to  $\eta = 0.1\%$  aligns the effective decay width with the inverse of  $t_U$ , which is crucial for normalizing the local energy density of dark photons in the main text. If we adopt a coupling constant  $g_a = 10^{-11} \text{ GeV}^{-1}$ , the saturated boson star field value,  $\Phi_0^{\text{sat}}$ , is predicted to be around  $f_\Phi/100$ , as inferred from the condition (S16). This ensures the stability of the boson star, preventing its collapse due to self-interactions [137, 138]. Furthermore, this choice of  $g_a$  effectively avoids triggering parametric resonance within miniclusters, assuming they typically have a radius of about  $10^7 \text{ m}$  [64, 66, 67, 76].

#### Four-body Cascade Decay

An alternative approach to circumvent the rapid depletion of the scalar dark matter background is by treating the scalar  $\Phi$  as an intermediate state in the decay chain from dark matter  $\chi$  to dark photon  $X^\mu$ . This transforms the decay process into a cascade sequence, represented as

$$\chi \rightarrow \Phi\Phi \rightarrow X^\mu X^\mu X^\mu X^\mu, \quad (\text{S20})$$

which is analogous to the galactic axion production explored in Ref. [38]. The kinematics inherent in this four-body decay process significantly expand the final state phase space. Specifically, in the regime where  $m_\Phi \ll m_{\text{DM}}$  (with  $m_{\text{DM}}$  denoting the dark matter particle  $\chi$ 's mass), the spectrum of the dark photons generated in each decay instance can be approximated as a uniform distribution up to a limit of  $m_{\text{DM}}/2$  [139, 140]:

$$\frac{dN_{\text{CD}}}{d\omega} = \frac{8}{m_{\text{DM}}} \Theta\left(\frac{m_{\text{DM}}}{2} - \omega\right). \quad (\text{S21})$$

The decay width of the  $\chi$  is primarily governed by the initial decay process  $\chi \rightarrow \Phi\Phi$ , denoted as  $\Gamma_{\text{CD}}$ .

#### Angular Spectral Density

Both the two-body parametric decay and the four-body cascade decay processes occur throughout the galaxy. To calculate the flux at Earth, it's necessary to integrate along the line of sight (l.o.s.) in a specific direction, resulting in the differential flux [78]:

$$\frac{d\Phi_X}{d^2\hat{k} d\omega} = \frac{r_\odot \rho_\odot^{\text{DM}}}{4\pi m_{\text{DM}}} J(\theta_k) \Gamma_{\text{PD/CD}} \frac{dN_{\text{PD/CD}}}{d\omega}, \quad J(\theta_k) \equiv \int_{\text{l.o.s.}} \left( \frac{\rho^{\text{DM}}(r(s, \theta_k))}{\rho_\odot^{\text{DM}}} \right) \frac{ds}{r_\odot}. \quad (\text{S22})$$

Note that  $m_{\text{DM}} = m_\Phi$  for parametric decay, while  $m_{\text{DM}} = m_\chi$  for cascade decay. We employ spherical coordinates in the Galactic frame, with Earth at the origin and the Galactic Center (GC) as the  $-\hat{z}$  axis, defining  $\hat{k} \equiv (\theta_k, \phi_k)$ .

We assume a spherically symmetric dark matter halo density  $\rho^{\text{DM}}(r)$  that varies with distance  $r$  from the GC. The parameter  $r_{\odot} = 8.33 \text{ kpc}$  is the distance from Earth to the GC. The factor  $J(\theta_k)$  quantifies the line of sight integral of the dark matter density, with  $r(s, \theta_k) \equiv (r_{\odot}^2 + s^2 - 2r_{\odot}s \cos \theta_k)^{1/2}$ . From the differential flux (S22), we derive the local dark photon energy density:

$$\rho_X = \iint \frac{d\Phi_X}{d^2\hat{k} d\omega} \omega 2\pi \sin \theta_k d\theta_k d\omega = \frac{1}{2} r_{\odot} \rho_{\odot}^{\text{DM}} \Gamma_{\text{PD/CD}} \int J(\theta_k) \sin \theta_k d\theta_k. \quad (\text{S23})$$

The dark matter distribution is modeled using the Navarro, Frenk, and White (NFW) profile [141]:

$$\rho^{\text{DM}}(r) = \rho_s \frac{r_s}{r} \left(1 + \frac{r}{r_s}\right)^{-2}, \quad (\text{S24})$$

with scale parameters  $r_s = 24.42 \text{ kpc}$  and  $\rho_s = 0.184 \text{ GeV/cm}^3$ . For the scenario involving parametric decay from boson stars, it is assumed that the distribution of the boson stars adheres to the NFW profile. We adopt the assumption that the effective decay rate  $\Gamma_{\text{PD/CD}}$  attains its maximum value consistent with existing cosmological constraints [77, 142]:

$$\Gamma_{\text{PD/CD}}^{-1} \approx 3.6 t_U \approx 4.97 \times 10^{10} \text{ years}. \quad (\text{S25})$$

Incorporating Eq. (S24) and (S25) into Eq. (S23) yields a maximally allowed flux of dark photons, approximately  $\rho_X \approx 1000 \rho_{\gamma}$  [38].

Comparing Eq. (S5) and (S23), we derive the relationship between the angular spectral density and differential flux:

$$\mathcal{S}_P(\omega, \hat{k}) = \frac{4\pi^2}{\omega} \frac{d\Phi_X}{d^2\hat{k} d\omega}. \quad (\text{S26})$$

This relationship forms the basis for two benchmark angular spectral densities discussed in the maintext:

$$\begin{aligned} \mathcal{S}_P^{\text{PD}}(\omega, \hat{k}) &= \frac{4\pi Q \rho_X}{\omega m_{\text{DM}}^2 \bar{J}} J(\theta_k) \Theta\left(\frac{m_{\text{DM}}}{4Q} - \left|\omega - \frac{m_{\text{DM}}}{2}\right|\right), \\ \mathcal{S}_P^{\text{CD}}(\omega, \hat{k}) &= \frac{8\pi \rho_X}{\omega m_{\text{DM}}^2 \bar{J}} J(\theta_k) \Theta\left(\frac{m_{\text{DM}}}{2} - \omega\right), \end{aligned} \quad (\text{S27})$$

where  $\bar{J} \equiv \int J(\hat{k}) d^2\hat{k} / (4\pi) \approx 2.19$ .

To calculate the modulation of galactic dark photon signals, it is necessary to consider the variation of the cavity's symmetry axis in the Galactic frame, denoted as  $\hat{l}_{\text{cav}}(t) \equiv (\theta_{\text{cav}}(t), \phi_{\text{cav}}(t))$ . For simplicity, we set the azimuthal angle of the Earth's rotation axis at 0, leading to the following geometric relationships:

$$\begin{aligned} \cos \theta_{\text{cav}}(t) &= \cos \delta_{\text{GC}} \cos \varphi_{\text{cav}} \cos(2\pi t/24 + \lambda_{\text{cav}} - \alpha_{\text{GC}}) + \sin \delta_{\text{GC}} \sin \varphi_{\text{cav}}, \\ \sin \theta_{\text{cav}}(t) \sin \phi_{\text{cav}}(t) &= \cos \varphi_{\text{cav}} \sin(2\pi t/24 + \lambda_{\text{cav}} - \alpha_{\text{GC}}). \end{aligned} \quad (\text{S28})$$

In this context,  $(\varphi_{\text{cav}}, \lambda_{\text{cav}}) = (39.9^\circ, 116.4^\circ)$  represent the latitude and longitude of the cavity, which is oriented towards the Earth's center throughout the experiment. The coordinates  $(\delta_{\text{GC}}, \alpha_{\text{GC}}) = (-29.0^\circ, 266.4^\circ)$  specify the declination and right ascension of the GC in the celestial sphere. The time variable  $t$  is expressed in hours, based on the J2000 standard time system [143]. Practically, due to the axial symmetry of the dark photon angular spectrum, the modulation of the cavity signal does not exhibit sensitivity to variations in  $\phi_{\text{cav}}(t)$ .

### III: DATA ANALYSIS

The data for this study were sourced from DPDM searches as detailed in Ref. [32]. The relevant parameters, along with their associated uncertainties, are systematically cataloged in Table S2.

Given the Gaussian nature of the measured data, we utilize the following probability function:

$$\text{Pr}(\delta^i | \epsilon, m_X) \propto \prod_i \exp\left(-\frac{\left(\delta^i / \tilde{\delta}_{\text{sig}}^i - (\epsilon / \epsilon_{\text{ref}})^2\right)^2}{2(\sigma_{\text{tot}}^i)^2}\right), \quad (\text{S29})$$

	Value	Fractional Uncertainty
$V$	3900 mL	< 1%
$\beta$	$0.634 \pm 0.014$	1.4%
$G_{\text{net}}$	$(57.30 \pm 0.14)$ dB	3.1%
$Q_L$	$(9.092 \pm 0.081) \times 10^9$	/

TABLE S2: Calibrated parameters for the SRF and amplifier used in the study, including average values, uncertainties, and fractional uncertainties on the dark photon-induced signal, denoted by  $F_j$ .

where  $\tilde{\delta}_{\text{sig}}^i \equiv \delta_{\text{sig}}^i |_{\epsilon=\epsilon_{\text{ref}}}$  is introduced alongside an arbitrary reference value  $\epsilon_{\text{ref}}$  to account for both measurement error and uncertainties from calibrated parameters. Employing the error propagation formula, we deduce the combined uncertainty for  $\delta^i / \tilde{\delta}_{\text{sig}}^i$ , labeled as  $\sigma_{\text{tot}}^i$ :

$$(\sigma_{\text{tot}}^i)^2 = \frac{1}{(\tilde{\delta}_{\text{sig}}^i)^2} + \frac{(\delta^i)^2}{(\tilde{\delta}_{\text{sig}}^i)^4} \sum_j \left( \frac{\partial \tilde{\delta}_{\text{sig}}^i}{\partial j} \sigma_j \right)^2. \quad (\text{S30})$$

Here,  $j$  represents various calibrated parameters with  $\sigma_j$  as their respective uncertainties. The first term in Eq. (S30) signifies the measurement error, assuming a standard deviation of 1 for  $\delta^i$ . Substituting Eq. (S30) into Eq. (S29) simplifies the probability function to:

$$\Pr(\delta^i | \epsilon, m_X) \propto \prod_i \exp\left(\frac{-(\delta^i - \delta_{\text{sig}}^i)^2}{2 + 2(\delta^i)^2 \sum_j F_j^2}\right), \quad (\text{S31})$$

as utilized in the main text. The fractional uncertainty is defined as:

$$F_j \equiv \frac{\partial \tilde{\delta}_{\text{sig}}^i}{\partial j} \frac{\sigma_j}{\tilde{\delta}_{\text{sig}}^i}. \quad (\text{S32})$$

The relevant contributions include  $j = G_{\text{net}}, \beta$ , and  $V$ :

$$\sum_j F_j^2 = \frac{\sigma_V^2}{V^2} + \left(\frac{1}{\beta} - \frac{1}{1+\beta}\right)^2 \sigma_\beta^2 + \frac{\sigma_{G_{\text{net}}}^2}{G_{\text{net}}^2} \quad (\text{S33})$$

with mean values and uncertainties presented in Table S2. Note that  $G_{\text{net}}$  is evaluated as  $(57.30 \pm 0.14)$  dB  $\rightarrow 10^{(5.730 \pm 0.014)}$ .

Using the probability function (S31), we can determine the 90% exclusion limit for the kinetic mixing parameter, denoted as  $\epsilon_{90\%}$ , by solving:

$$\frac{\int_0^{\epsilon_{90\%}} \Pr(\delta^i | \epsilon, m_X) d\epsilon^2}{\int_0^\infty \Pr(\delta^i | \epsilon, m_X) d\epsilon^2} = 90\%. \quad (\text{S34})$$