Reactor neutrino background in third-generation dark matter detectors

D. Aristizabal Sierra,^{1,*} Valentina De Romeri,^{2,†} and Christoph A. Ternes^{3,‡}

¹Universidad Técnica Federico Santa María - Departamento de Física

Casilla 110-V, Avda. España 1680, Valparaíso, Chile

²Instituto de Física Corpuscular (CSIC-Universitat de València),

Parc Científic UV C/ Catedrático José Beltrán, 2 E-46980 Paterna (Valencia) - Spain

³Istituto Nazionale di Fisica Nucleare (INFN), Laboratori Nazionali del Gran Sasso,

67100 Assergi, L'Aquila (AQ), Italy

Third-generation dark matter detectors will be fully sensitive to the boron-8 solar neutrino flux. Because of this, the characterization of such a background has been the subject of extensive analyses over the last few years. In contrast, little is known about the impact of reactor neutrinos. In this letter we report on the implications of such a flux for dark matter direct detection searches. We consider five potential detector deployment sites envisioned by the recently established XLZD consortium: SURF, SNOLAB, Kamioka, LNGS and Boulby. By using public reactor data we construct five reactor clusters—involving about 100 currently operating commercial nuclear reactors each—and determine the net neutrino flux at each detector site. Assuming a xenon-based detector and a 50 tonne-year exposure, we show that in all cases the neutrino event rate may be sizable, depending on energy recoil thresholds. Of all possible detector sites, SURF and LNGS are those with the smallest reactor neutrino fluxes, with Kamioka being subject to a more moderate background. Our findings demonstrate that reactor neutrino fluxes should be taken into account in the next round of dark matter searches. We argue that this background may be particularly relevant for directional detectors, provided they meet the requirements we have employed in this analysis.

INTRODUCTION

A wealth of cosmological and astrophysical data supports the idea that the dominant form of matter in the Universe has feeble or none electromagnetic interactions. The conventional wisdom is that this new form of matter-dubbed dark matter (DM)—is of microscopical origin and its abundance is determined by fast-scattering processes with Standard Model (SM) particles at very early epochs, much before the onset of cosmic neutrino decoupling and primordial nucleosynthesis (for a review see e.g. Ref. [1]). Although at high temperatures DM is thermalized, as the temperature decreases-because of the expansion of the Universe --- these scattering processes are unable to keep the species in thermodynamic equilibrium and so its abundance freezes out. This weakly interacting massive particle (WIMP) is a rather generic candidate appearing in a large class of particle physics models. It is a dominant paradigm that has driven DM searches.

DM direct detection is a subject that dates back to the mid 80's, when Goodman and Witten pointed out that WIMPs could be searched for by using the same detectors proposed by Drukier and Stodolsky for coherent elastic neutrino-nucleus scattering (CEvNS) measurements [2, 3]. Since then, and because of the lack of a signal, detector technologies as well as fiducial volumes have dramatically evolved. At present, DM searches in direct detection experiments are led by liquid xenon (LXe) dual-phase time projection chambers (secondgeneration DM detectors). Detectors at the INFN "Laboratori Nazionali del Gran Sasso" (LNGS) in Italy (XENONnT), at the Sanford Underground Research Facility (SURF) in South Dakota in the US (LZ) and at the China Jinping Underground Laboratory in Sichuan, China (PandaX-4T) are using active volumes of the order of 5 tonne [4–6]. With their high capabilities for background rejection, along with low nuclear recoil energy thresholds, these second-generation DM detectors are sensitive to spin-independent WIMP-nucleon total cross sections of the order of 10^{-48} cm² [7]. Indeed, XENONnT and LZ have recently published results where sensitivities of the order of $\sigma_{WIMP-nuc} \sim 10^{-47}$ cm² have been reported [4, 5]. PandaX-4T has set the most stringent upper limit in the low WIMP mass region (≤ 10 GeV), $\sigma_{WIMP-nuc} \sim 10^{-44}$ cm² [6].

A new generation of LXe detectors—third-generation DM detectors—is expected to pave the way for a discovery ¹. Recently the XENONnT, LZ and DARWIN collaborations have united forces and created the XLZD consortium [8]. Their goal is the construction of a 40-100 tonne detector with unprecedented sensitivities. With such active volume, a detector of this kind will be subject to an irreducible neutrino background dominated by ⁸B solar neutrinos (for nuclear-channel signals) and by *pp* neutrinos (for electron-channel signals) [9].

The morphology and size of this background have been the subject of different analyses in recent years, first identified as the so-called "neutrino floor" [9–13] and its more recent redefinition, the "neutrino fog" [14], where a first estimation of the reactor neutrino background at LNGS was addressed. It is well known that the impact of the neutrino background on a WIMP discovery signal is mainly dominated by neutrino flux uncertainties, with uncertainties on the weak mixing angle and on the root-mean-square radii of the neutron distribu-

¹ Note that if a discovery takes place in second-generation detectors, the experimental environment provided by their third-generation follow-ups will allow precise measurements of WIMP properties.

tions playing a rather subdominant role [15]. The presence of a neutrino background, however, does not mean that an identification of a WIMP signal is impossible. First of all, improvements in the determination of solar neutrino flux uncertainties are expected. Secondly, WIMP and neutrino spectra in general do not fully degenerate in most regions of parameter space. Even in regions where they strongly do, an identification is possible with sufficiently large data sets [12]. Furthermore, even if data is not abundant, directionality will potentially—enable a distinction between WIMP and neutrino nuclear recoil spectra [16], if they turn out to be strongly degenerate.

Given this landscape, and the fact that DM direct detection will soon enter the third-generation detector phase, one should wonder whether other neutrino sources might contribute to the background and hence should be taken into account. This is a rather relevant question to raise, aiming to leverage the full discovery power of these types of detectors. Motivated by this question, in this Letter we assess the impact of nuclear reactor neutrinos. Since the reactor neutrino flux strongly depends on the geographical position of the detector—for definitiveness—we use LNGS, SURF, Boulby (UK), Kamioka (Japan) and SNOLAB (Canada) as possible deployment sites ².

NUCLEAR REACTOR SOURCES: LOCATIONS AND EVENT RATES

The data sets we employ follow from data provided on the Geoneutrinos.org website [17, 18]. We consider only commercial power plants (that involve the most powerful reactors) for which a non-zero operating power is reported. Reactors for which the thermal capacity is known but have zero operating power and those that have been permanently shut down are not included. Depending on the baseline, each detector site that we consider is "surrounded" by a cluster of nuclear reactor power plants, at a certain distance L_i . Table I shows the minimum and maximum baseline and power for each cluster, along with the number of reactors involved. For each detector site, we do not include reactors located at distances beyond L_{max} , as their contribution to the event rates would be negligible.

Location	NR	L _{min} [km]	L _{max} [km]	P_{\min} [GW]	P_{\max} [GW]
SURF	111	790	2951	0.34	3.9
SNOLAB	104	239	2874	0.92	3.9
Kamioka	86	146	2895	0.15	3.9
LNGS	146	417	4027	0.42	3.7
Boulby	141	26	3654	0.51	3.7

TABLE I. Minimum and maximum baselines (L_{min} and L_{min}) along with minimum and maximum reactor powers (P_{min} and P_{max}) for the SURF, SNOLAB, Kamioka, LNGS and Boulby reactor clusters. The number of reactors in each cluster (NR) is also shown. Data has been extracted from the Geoneutrinos.org website.

The largest clusters are those around the LNGS and Boulby detector sites (as expected, given that for these two cases the radius defining the cluster exceeds by about 1000 km the radius at the other sites). However, this does not necessarily mean that the largest flux is obtained for these two positions, as we now discuss. The reactor neutrino flux decreases rapidly with increasing baseline. So, a rather fair assumption is that the flux is dominated by the sub-cluster defined by all reactors included in a radius $\lesssim 1000$ km. For the SURF and LNGS locations one finds that these sub-clusters involve only 5 reactors with a 2.1 GW and 1.8 GW average power, respectively. For the Kamioka, SNOLAB and Boulby locations, the subclusters are composed instead of 35, 59 and 49 reactors. The average power in each case (and in that order) is: 2.1 GW, 4.9 GW and 1.9 GW. Thus, already from these numbers one expects the SURF and LNGS location sites to involve a less intense reactor neutrino flux.

Fig. 1 shows the distribution of nuclear reactors in terms of baseline and power for the five different clusters we consider. The distributions involve the full data sets. From the graph, one can see that for the Boulby and SNOLAB clusters the reactor density for baselines below 1000 km is high, with a few of those reactors having powers above 3 GW. The distribution for the Kamioka cluster is somewhat different. Although below 1000 km there are a few reactors, their density is lower as well as their power. For the SURF and LNGS clusters, the reactor density for baselines below 1000 km is, instead, rather moderate. For these clusters, most reactors are at baselines above 1000 km. So, even without a dedicated calculation of the event rate, expectations are that in terms of increasing reactor neutrino fluxes the clusters can be sorted into three groups: SURF/LNGS, Kamioka, SNOLAB/Boulby.

The calculation of the differential nuclear recoil spectrum at each cluster (C) requires the convolution of the differential CEvNS cross section [3, 19] with the reactor neutrino flux, namely

$$\frac{dR_{\rm C}}{dE_r} = \frac{m_{\rm det}N_A\,\mathcal{T}\,\eta_{\rm C}}{m_{\rm mol}^{\rm Xe}} \int_{E_{\rm v}^{\rm min}}^{E_{\rm v}^{\rm max}} \frac{d\Phi_{\overline{\nu}_e}}{dE_{\rm v}} \frac{d\sigma}{dE_r} F_{\rm H}^2(E_r)dE_{\rm v} \,. \tag{1}$$

Here, m_{det} refers to the detector active volume mass, $m_{\text{mol}}^{\text{Xe}}$ to the xenon molar mass, \mathcal{T} to the exposure time, $E_{v}^{\text{min}} = \sqrt{m_{N}E_{r}/2}$ (E_{r} and m_{N} refer to nuclear recoil energy and

² These underground facilities are considered as potential locations for detector deployment by the XLZD Consortium [8].



FIG. 1. Location of the different reactors within the SURF, SNOLAB, Kamioka, LNGS and Boulby clusters and their corresponding operating power.

mass), and E_v^{max} to the neutrino spectrum kinematic "highenergy" tail taken at 8 MeV. The average nuclear mass is $\langle m_{\text{Xe}} \rangle / \text{GeV} = 0.93 \times \langle A \rangle$, $\langle A \rangle = \sum_i X_i A_i = 131.4$ being the mass number averaged over the nine stable xenon isotopes. We include—for completeness—the weak-charge nuclear form factor, $F_{\text{H}}(E_r)$, parametrized à la Helm [20]. Note that if not included results would deviate from those presented here at most by ~ 2%, because of the process occurring deep in the full coherent regime.

Regarding the electron antineutrino spectrum, we proceed as follows. For the ²³⁵U and ²³⁸U emission spectra we use results from Ref. [21]. For ²³⁹Pu and ²⁴¹Pu we use instead results from Ref. [22]. The full electron antineutrino differential flux is then calculated according to

$$\frac{d\Phi_{\overline{\mathbf{v}}_e}}{dE_{\mathbf{v}}} = \sum_{i=\text{Isotopes}} f_i \frac{d\Phi_{\overline{\mathbf{v}}_e}^i}{dE_{\mathbf{v}}} , \qquad (2)$$

where $f_i = \{f_{235_{\rm U}}, f_{238_{\rm U}}, f_{239_{\rm Pu}}, f_{241_{\rm Pu}}\} = \{5.5, 0.7, 3.2, 0.6\} \times 10^{-1}$ are the uranium and plutonium fission fractions [23]. Note that we do not include electron antineutrinos produced in neutron capture by ²³⁸U. The reason is that the spectra for those neutrinos dominate at energies below ~ 1.5 MeV, hence in a LXe detector would produce nuclear recoils below 0.04 keV (much below any realistic operation threshold). We assume the spectral function in Eq. (2) to be universal for all the reactors within the clusters ³. Thus, the difference among clusters is determined only by the normalization factor, which we calculate assuming that in each fission process an energy of $\varepsilon = 205.24$ MeV is released and that neutrinos are emitted

isotropically. Explicitly, each normalization factor is given by

$$\eta_{\rm C} = \sum_j \frac{P_j}{4\pi L_j^2 \varepsilon} \,, \tag{3}$$

where *j* runs over all reactors relevant for cluster C and P_j and L_j are the operating power and distance for reactor *j*. Their values are displayed in Tab. II, showing that SURF is subject to the least abundant neutrino flux, whereas Boulby to the most severe.

Cluster	SURF	SNOLAB	Kamioka	LNGS	Boulby
$\eta_{C}[cm^{-2}sec^{-1}]$	20422	156630	103903	56677	932874

TABLE II. Neutrino flux normalization factors for the five reactor clusters.

With these results at hand, we are now in a position to calculate the differential event rate as well as the total event rate for each detector site. We assume a 50-tonne active volume LXe detector and 100% efficiency ⁴. Since current realistic thresholds amount to 0.3 keV [25], we use $E_r^{\text{th,min}} = 0.1 \text{ keV}$ as a value envisioned for future detector operations. Results are displayed in Fig. 2. The left (right) graph shows the differential event rate (total event rate) as a function of the recoil energy (recoil energy threshold) for the five different reactor clusters we have considered. The inset plot in the right panel is meant to zoom in on the bottom left corner. Inline with expectations, the differential and total event rates at the SURF (Boulby) detector site are the smallest (largest). The event rate at the LNGS detector location is slightly higher, followed by Kamioka and SNOLAB.

³ Each reactor has its own fission fractions, but variations are at the permille level (see e.g. Tab. 4 in Ref. [24]).

⁴ The XLZD consortium aims at masses from 40 to 100 tonne. So this value is used just as a proxy of what the actual detector will use.



FIG. 2. Left graph: Reactor neutrino differential event rate for the five detector sites considered in this work: SURF, SNOLAB, Kamioka, LNGS, and Boulby as a function of nuclear recoil energy. Shown as well is the ⁸B differential event rate. **Right graph**: Reactor neutrino total event rate for the same detector locations.

DISCUSSION

Naively one would expect the reactor neutrino flux to be suppressed and of little relevance. This expectation is mainly based on the fact that most reactors are far away from the detector sites. However, the fact that the clusters around each detector site involve a large number of active nuclear power plants (with in some cases powerful reactors), combined with a large active volume produces a non-zero event rate in all cases.

Ideally one would like a very low threshold to explore the small WIMP mass window and increase the WIMP-nucleus event rate. At 0.1 keV, we find that the total neutrino-nucleus event rate per year is: 16 (SURF), 44 (LNGS), 82 (Kamioka), 124 (SNOLAB) and 733 (Boulby). If that operation threshold is not achieved and instead the detector is operated at 0.3 keV, these numbers will be degraded by about a factor 7. In such an experimental scenario the reactor neutrino background becomes, of course, less severe. Thus, the question of whether the reactor neutrino background matters is—as anticipated—strongly linked to operation thresholds.

It is worth emphasizing that variations of these estimated numbers are expected in the future, depending on the exact number of reactors that enter in either operation phase or are decommissioned. However, these results demonstrate that the reactor neutrino flux should be seriously taken into account in decision making as well as in data taken, contrary to expectations.

Finally, one might wonder how much this neutrino background matters compared to the boron-8 solar neutrino flux. For the detector configurations we have considered, with a 0.1 keV operation threshold, the number of boron-8 nuclear recoil induced events is overwhelming, 36500 events/year. So, of course, this will be the dominant background source. All the efforts to understand the morphology of this background are indeed motivated by this fact. The question is then whether one should be concerned with the reactor neutrino background whatsoever.

It is well known that the boron-8 background can be to a certain degree circumvented. As we have already stressed, large data sets might enable differentiating neutrino from WIMP signals, if the WIMP parameters are such that the neutrino and WIMP event rates strongly degenerate. In general, however, directional detectors seem to be the most promising avenue [16, 26]⁵. For these detectors it seems that the reactor neutrino background might even become the most dominant background source. Therefore, if the boron-8 nuclear recoil-induced events can be efficiently discriminated, there will be yet another background source that will require careful identification and proper treatment, depending on statistics and operation capabilities.

CONCLUSIONS

With the advent of third-generation DM direct detection detectors, the quantification of reactor neutrino fluxes becomes of pivotal importance. In this work we have quantified the size of the neutrino flux produced by clusters of reactors surrounding five potential detector deployment sites. For definitiveness we have considered the locations envisioned by the recently established XLZD consortium: SURF, SNOLAB, Kamioka, LNGS, and Boulby.

Our findings show that detectors with active volumes of the order of 50 tonne and recoil energy thresholds of the or-

⁵ They have been as well recently considered for CEvNS measurements and beyond the SM searches using neutrino beamlines at Fermilab [27–29]

der of 0.1 keV, will be sensitive to a certain amount of reactor neutrino-induced events. The exact amount depends, to a large degree, on the energy threshold at which the detector is operated. However, even assuming a realistic threshold of 0.3 keV, the event rate turns out to be sizable in all cases. We find that the site with the smallest reactor neutrino background is SURF followed by LNGS, Kamioka, SNOLAB, and Boulby (in that order).

Although subdominant compared to the solar boron-8 neutrino background, we point out that the reactor neutrino background (and its corresponding events) should be—in principle—considered during data taken. Reactor neutrinoinduced events should be taken into account in background discrimination, regardless of the detector technique employed. This result will be particularly relevant for directional detection, if future detectors meet the requirements we have used here.

ACKNOWLEDGMENTS

We thank P. Martínez-Miravé for pointing out to us the Geoneutrinos.org website. The work of D.A.S. is funded by ANID under grant "Fondecyt Regular" 1221445. He thanks "Le Service de Physique Théorique (Université Libre de Bruxelles)" and "Instituto de Física Corpuscular (CSIC y Universidad de Valencia)" for their kind hospitality and their stimulating research environment during the completion of this work. V.D.R. acknowledges financial support from the CIDEXG/2022/20 grant (project "D'AMAGAT") funded by Generalitat Valenciana and by the Spanish grant PID2020-113775GB-I00

(MCIN/AEI/10.13039/501100011033). C.A.T. is very thankful for the hospitality at Universidad Técnica Federico Santa María, where this work was initiated.

* daristizabal@uliege.be

- [†] deromeri@ific.uv.es
- [‡] christoph.ternes@lngs.infn.it
- [1] G. Arcadi, M. Dutra, P. Ghosh, M. Lindner, Y. Mambrini, M. Pierre, S. Profumo, and F. S. Queiroz, "The waning of the WIMP? A review of models, searches, and constraints," *Eur. Phys. J. C* 78 no. 3, (2018) 203, arXiv:1703.07364 [hep-ph].
- [2] M. W. Goodman and E. Witten, "Detectability of Certain Dark Matter Candidates," *Phys. Rev. D* 31 (1985) 3059.
- [3] A. Drukier and L. Stodolsky, "Principles and Applications of a Neutral Current Detector for Neutrino Physics and Astronomy," *Phys. Rev. D* 30 (1984) 2295.
- [4] XENON Collaboration, E. Aprile *et al.*, "First Dark Matter Search with Nuclear Recoils from the XENONnT Experiment," *Phys. Rev. Lett.* **131** no. 4, (2023) 041003, arXiv:2303.14729 [hep-ex].
- [5] LZ Collaboration, J. Aalbers *et al.*, "First Dark Matter Search Results from the LUX-ZEPLIN (LZ) Experiment," *Phys. Rev.*

Lett. 131 no. 4, (2023) 041002, arXiv:2207.03764 [hep-ex].

- [6] PandaX Collaboration, S. Li *et al.*, "Search for Light Dark Matter with Ionization Signals in the PandaX-4T Experiment," *Phys. Rev. Lett.* **130** no. 26, (2023) 261001, arXiv:2212.10067 [hep-ex].
- [7] J. Billard *et al.*, "Direct detection of dark matter—APPEC committee report*," *Rept. Prog. Phys.* 85 no. 5, (2022) 056201, arXiv:2104.07634 [hep-ex].
- [8] "The XLZD Consortium." https://xlzd.org/.
- [9] L. E. Strigari, "Neutrino Coherent Scattering Rates at Direct Dark Matter Detectors," *New J. Phys.* 11 (2009) 105011, arXiv:0903.3630 [astro-ph.CO].
- [10] J. Monroe and P. Fisher, "Neutrino Backgrounds to Dark Matter Searches," *Phys. Rev. D* 76 (2007) 033007, arXiv:0706.3019 [astro-ph].
- [11] J. D. Vergados and H. Ejiri, "Can Solar Neutrinos be a Serious Background in Direct Dark Matter Searches?," *Nucl. Phys. B* 804 (2008) 144–159, arXiv:0805.2583 [hep-ph].
- [12] J. Billard, L. Strigari, and E. Figueroa-Feliciano, "Implication of neutrino backgrounds on the reach of next generation dark matter direct detection experiments," *Phys. Rev. D* 89 no. 2, (2014) 023524, arXiv:1307.5458 [hep-ph].
- [13] C. A. J. O'Hare, "Dark matter astrophysical uncertainties and the neutrino floor," *Phys. Rev. D* 94 no. 6, (2016) 063527, arXiv:1604.03858 [astro-ph.CO].
- [14] C. A. J. O'Hare, "New Definition of the Neutrino Floor for Direct Dark Matter Searches," *Phys. Rev. Lett.* **127** no. 25, (2021) 251802, arXiv:2109.03116 [hep-ph].
- [15] D. Aristizabal Sierra, V. De Romeri, L. J. Flores, and D. K. Papoulias, "Impact of COHERENT measurements, cross section uncertainties and new interactions on the neutrino floor," *JCAP* 01 no. 01, (2022) 055, arXiv:2109.03247 [hep-ph].
- [16] S. E. Vahsen, C. A. J. O'Hare, and D. Loomba, "Directional Recoil Detection," Ann. Rev. Nucl. Part. Sci. 71 (2021) 189–224, arXiv:2102.04596 [physics.ins-det].
- [17] S. Dye and A. Barna, "Global Antineutrino Modeling for a Web Application," arXiv:1510.05633 [physics.ins-det].
- [18] "Geoneutrinos website." https://reactors.geoneutrinos.org/.
 [19] D. Z. Freedman, "Coherent Neutrino Nucleus Scattering as a Probe of the Weak Neutral Current," *Phys. Rev. D* 9 (1974)
- [20] R. H. Helm, "Inelastic and Elastic Scattering of 187-Mev
- Electrons from Selected Even-Even Nuclei," *Phys. Rev.* 104 (1956) 1466–1475.
- [21] V. Kopeikin, M. Skorokhvatov, and O. Titov, "Reevaluating reactor antineutrino spectra with new measurements of the ratio between U235 and Pu239 β spectra," *Phys. Rev. D* 104 no. 7, (2021) L071301, arXiv:2103.01684 [nucl-ex].
- [22] P. Huber, "On the determination of anti-neutrino spectra from nuclear reactors," *Phys. Rev. C* 84 (2011) 024617, arXiv:1106.0687 [hep-ph]. [Erratum: Phys.Rev.C 85, 029901 (2012)].
- [23] TEXONO Collaboration, H. T. Wong *et al.*, "A Search of Neutrino Magnetic Moments with a High-Purity Germanium Detector at the Kuo-Sheng Nuclear Power Station," *Phys. Rev.* D 75 (2007) 012001, arXiv:hep-ex/0605006.
- [24] C. Giunti, Y. F. Li, C. A. Ternes, and Z. Xin, "Reactor antineutrino anomaly in light of recent flux model refinements," *Phys. Lett. B* 829 (2022) 137054, arXiv:2110.06820 [hep-ph].

- [25] B. Lenardo *et al.*, "Measurement of the ionization yield from nuclear recoils in liquid xenon between 0.3 - 6 keV with single-ionization-electron sensitivity," arXiv:1908.00518 [physics.ins-det].
- [26] S. E. Vahsen *et al.*, "CYGNUS: Feasibility of a nuclear recoil observatory with directional sensitivity to dark matter and neutrinos," arXiv:2008.12587 [physics.ins-det].
- [27] M. Abdullah, D. Aristizabal Sierra, B. Dutta, and L. E. Strigari, "Coherent Elastic Neutrino-Nucleus Scattering with directional detectors," *Phys. Rev. D* 102 no. 1, (2020) 015009, arXiv:2003.11510 [hep-ph].
- [28] D. Aristizabal Sierra, B. Dutta, D. Kim, D. Snowden-Ifft, and L. E. Strigari, "Coherent elastic neutrino-nucleus scattering with the vBDX-DRIFT directional detector at next generation neutrino facilities," *Phys. Rev. D* 104 no. 3, (2021) 033004, arXiv:2103.10857 [hep-ph].
- [29] vBDX-DRIFT Collaboration, D. Aristizabal Sierra, J. L. Barrow, B. Dutta, D. Kim, D. Snowden-Ifft, L. Strigari, and M. H. Wood, "Rock neutron backgrounds from FNAL neutrino beamlines in the vBDX-DRIFT detector," *Phys. Rev. D* 107 no. 1, (2023) 013003, arXiv:2210.08612 [hep-ex].