

# HOLOMORPHIC GEOMETRIC STRUCTURES ON OELJEKLAUS–TOMA MANIFOLDS

INDRANIL BISWAS AND SORIN DUMITRESCU

ABSTRACT. We prove that any holomorphic geometric structure of affine type on an Oeljeklaus–Toma manifold is locally homogeneous. For locally conformally Kähler Oeljeklaus–Toma manifolds we prove that all holomorphic geometric structures, and also all holomorphic Cartan geometries, on them are locally homogeneous.

## 1. INTRODUCTION

The Oeljeklaus–Toma manifolds (which are simply referred to as OT–manifolds) form an important class of compact complex non-Kähler manifolds. They were constructed by Oeljeklaus and Toma in [OT] using tools from algebraic number theory and generalizing the construction of Inoue surfaces in class VII<sub>0</sub> [In]. Oeljeklaus–Toma manifolds provide a counterexample to Vaisman’s conjecture [Va2], which is a conjecture on locally conformally Kähler geometry that was open for 25 years.

The geometry of Oeljeklaus–Toma manifolds was systematically studied in the last two decades (see, for instance, [APV, BO, Ka, Ka2, OV, OVV, PV, Ve]). In particular, it was proved that OT–manifolds share many of the geometric features of Inoue surfaces of class VII<sub>0</sub> : they are non-Kähler flat affine manifolds [OT], they are of algebraic dimension zero (meaning that all meromorphic functions are constants) [BO], they contain no holomorphic curves [Ve] and they are known to be solvmanifolds admitting no Vaisman metrics [Ka2]. Moreover, locally conformally Kähler OT–manifolds do not admit proper complex subvarieties [OV].

Let us also mention that, among other results, it was proved in [OT] that the canonical bundle of an OT–manifold and its tensor powers are flat. This result was extended in [APV] to all line bundles.

Our article studies holomorphic geometric structures, and also holomorphic Cartan geometries, on Oeljeklaus–Toma manifolds.

By construction, any OT–manifold  $M$  admits a standard complex affine structure [OT], which endows  $M$  with a torsion-free holomorphic flat affine connection [Ko]. This is an example of holomorphic geometric structure in Gromov’s sense [Gr, DG] and can also be seen as a flat holomorphic Cartan geometry [Sh].

Our main results in this context are the following:

---

2020 *Mathematics Subject Classification.* 53A15, 53C23, 57S20, 14D10.

*Key words and phrases.* Oeljeklaus–Toma manifold, geometric structure, Cartan geometry, Killing vector field.

**Theorem 1.1.** *Let  $M$  be an OT–manifold, and let  $\phi$  be a holomorphic geometric structure of affine type on  $M$ . Then  $\phi$  is locally homogeneous. Moreover, the Lie algebra of local holomorphic affine vector fields on  $M$  preserving  $\phi$  is transitive.*

In particular, the above theorem asserts that any holomorphic tensor on an OT–manifold is locally homogeneous. Moreover, the Lie algebra of all local holomorphic affine vector fields preserving the holomorphic tensor is transitive on  $M$ .

**Theorem 1.2.** *Let  $M$  be a locally conformally Kähler OT–manifold. Then the following statements hold:*

- (i) *Any holomorphic geometric structure  $\phi$  on  $M$  is locally homogeneous. Moreover, the Lie algebra of all local holomorphic affine vector fields on  $M$  that preserve the geometric structure  $\phi$  is transitive.*
- (ii) *Any holomorphic Cartan geometry of algebraic type on  $M$  is locally homogeneous. Moreover, the Lie algebra of local holomorphic affine vector fields on  $M$  preserving the Cartan geometry is transitive.*

The article is organized as follows. Section 2 introduces the two concepts of geometric structures which are studied in the paper: geometric structures in Gromov’s sense [Gr, DG] and Cartan geometries [Sh]. Section 3 briefly presents the construction of OT–manifolds as discovered in [OT] and also recall some of their main geometric features with focus on the properties used in our proofs. Section 4 provides the proofs of our results, in particular that of Theorem 1.1 (see Theorem 4.1) and of Theorem 1.2 (see Theorem 4.2 and Corollary 4.7).

## 2. HOLOMORPHIC GEOMETRIC STRUCTURES

Let  $M$  be a complex manifold of complex dimension  $n$ . For any integer  $k \geq 1$ , we associate the principal bundle on it of  $k$ -frames

$$R^k(M) \longrightarrow M,$$

which is the bundle of  $k$ -jets of local holomorphic coordinates on  $M$ . The structure group of this principal bundle is  $D^k$ : the group of all  $k$ -jets of local biholomorphisms of  $\mathbb{C}^n$  fixing the origin. It may be mentioned that  $D^k$  is a complex affine algebraic group [Gr, DG, Ho, Hu].

We define a geometric structure on  $M$  as in [Gr] (see also the expository survey [DG]).

**Definition 2.1.** *A holomorphic geometric structure of order  $k$  on  $M$  is a holomorphic  $D^k$ -equivariant map  $\phi$  from  $R^k(M)$  to a complex algebraic manifold  $Z$  endowed with an algebraic action of the group  $D^k$ .*

Such a geometric structure  $\phi$  is said to be *of affine type* if the above algebraic manifold  $Z$  is a complex affine variety.

Holomorphic tensors are holomorphic geometric structures of affine type of order one. Indeed, a holomorphic tensor on  $M$  is a holomorphic  $\mathrm{GL}(n, \mathbb{C})$ -equivariant map from the frame bundle  $R^1(M)$  to a linear complex algebraic representation  $W$  of  $\mathrm{GL}(n, \mathbb{C})$ . More

precisely, for any integers  $p, q \geq 0$ , a holomorphic section of  $(TM)^{\otimes p} \otimes (T^*M)^{\otimes q}$  is a  $\mathrm{GL}(n, \mathbb{C})$ -equivariant holomorphic map from  $R^1(M)$  to the representation  $(\mathbb{C})^{\otimes p} \otimes (\mathbb{C}^*)^{\otimes q}$ .

Holomorphic affine connections are holomorphic geometric structures of affine type of order two [Gr, DG]. Holomorphic foliations and holomorphic projective connections are holomorphic geometric structure of non-affine type.

The natural notion of (local) symmetry of a holomorphic geometric structure is classically defined as follows.

A (local) biholomorphism of  $M$  preserves a holomorphic geometric structure  $\phi$  if its canonical lift to the associated  $k$ -frame  $R^k(M)$  fixes each fiber of the map  $\phi$ . Such a local biholomorphism is called a *local isometry* of  $\phi$ .

A (local) holomorphic vector field on  $M$  is called a *Killing vector field* with respect to  $\phi$  if its (local) flow acts on  $M$  by local isometries.

**Definition 2.2.** A holomorphic geometric structure  $\phi$  is called *rigid* of order  $l$  in Gromov's sense if any local biholomorphism preserving  $\phi$  is uniquely determined by its  $l$ -jet at any given point.

Holomorphic affine connections are rigid of order one in Gromov's sense (see [Gr] and [DG]). The rigidity comes from the fact that local biholomorphisms fixing a point and preserving a connection linearize in exponential coordinates, so they are indeed completely determined by their differential at the fixed point.

Holomorphic Riemannian metrics, holomorphic projective connections and holomorphic conformal structures of dimension at least three are all rigid holomorphic geometric structures, while holomorphic symplectic structures and holomorphic foliations are non-rigid geometric structures [DG].

By a theorem of Amores [Am], analytic local Killing fields on simply connected manifolds extend to global holomorphic Killing fields. This implies that the sheaf of local Killing fields of a holomorphic rigid geometric structure  $\phi$  is locally constant. Its fiber is a finite dimensional Lie algebra called *the Killing Lie algebra* of  $\phi$  [DG, Gr]. If the Killing Lie algebra is transitive on an open dense subset  $U$  in  $M$ , we say that that  $\phi$  is *locally homogeneous* on  $U$ .

Let us now recall the classical definition of a *holomorphic Cartan geometry*.

Let  $G$  be a connected complex Lie group and  $H \subset G$  a connected complex Lie subgroup. We denote by  $\mathfrak{h}$  and  $\mathfrak{g}$  the Lie algebras of  $H$  and  $G$  respectively.

**Definition 2.3.** A holomorphic Cartan geometry of type  $(G, H)$  on a complex manifold  $M$  is a holomorphic principal  $H$ -bundle  $\pi : E_H \rightarrow M$  endowed with a  $\mathfrak{g}$ -valued holomorphic 1-form  $\eta$  on  $E_H$  satisfying the following conditions:

- (1)  $\eta : TE_H \rightarrow E_H \times \mathfrak{g}$  is an isomorphism;
- (2)  $\eta$  is  $H$ -equivariant with  $H$  acting on  $\mathfrak{g}$  via the adjoint representation;
- (3) The restriction of  $\eta$  to each fiber of  $\pi$  coincides with the Maurer–Cartan form associated to the action of  $H$  on  $E_H$ .

The above definition is the infinitesimal version of the canonical fibration  $G \rightarrow G/H$ , seen as a holomorphic principal  $H$ -bundle equipped with the left-invariant Maurer-Cartan form of  $G$  (see [Sh]).

The Cartan geometry is called *flat* if its curvature  $\Omega(\eta) = d\eta + \frac{1}{2}[\eta, \eta]$  vanishes identically.

By a classical result of Cartan, a flat Cartan geometry of type  $(G, H)$  given by a principal  $H$ -bundle  $E_H$  and a form  $\eta$  is flat if and only if  $(E_H, \eta)$  is locally isomorphic to the fibration  $G \rightarrow G/H$  endowed with its canonical Maurer-Cartan form. As a consequence, Ehresmann proved that a complex manifold  $M$  admits a flat Cartan geometry of type  $(G, H)$  if and only if  $M$  admits a holomorphic  $(G, G/H)$ -structure, which is a holomorphic atlas with values in the model space  $G/H$  with the transition maps being given by the restriction of left-translation action, on  $G/H$ , of elements of  $G$ .

The Cartan geometry  $(E_H, \eta)$  of type  $(G, H)$  will be called of *algebraic type* if the image of  $H$  through the adjoint representation of  $G$  is an algebraic subgroup of  $\text{Aut}(\mathfrak{g})$  [Ho, Hu].

Holomorphic affine and projective connections, holomorphic conformal structures and holomorphic Riemannian metrics can be defined as holomorphic *Cartan geometries of algebraic type* [Sh]. In particular, for  $G = \text{GL}(n, \mathbb{C}) \times \mathbb{C}^n$  and  $H = \text{GL}(n, \mathbb{C})$  one gets the definition of a Cartan geometry with model the complex affine space. In this case the bundle  $E_H$  is necessarily isomorphic to the bundle of 1-frames  $R^1(M)$ . This data is equivalent to the existence of a holomorphic affine connection  $\nabla$  on  $M$ . The Cartan curvature is flat if and only if  $\nabla$  is torsion-free and its curvature vanishes. By Ehresmann result, this is equivalent to the data of a holomorphic  $(G = \text{GL}(n, \mathbb{C}) \times \mathbb{C}^n, \mathbb{C}^n)$ -structure on  $M$ . Such an equivalence class of atlases with transition maps which are restrictions of affine transformations is classically termed as a *complex affine structure*.

There is a natural notion of (local) symmetry for holomorphic Cartan geometries. A (local) Killing field of the Cartan geometry  $(E_H, \eta)$  is a local holomorphic vector field on  $M$  which lifts to a local holomorphic vector field on  $E_H$  whose local flow acts by bundle automorphisms on  $E_H$  that preserve  $\eta$ . A local Killing field  $X$  of a Cartan geometry extends to any simply connected open subset in  $M$  containing the domain of definition of  $X$  [Am]. This implies that the sheaf of local Killing vector fields is a locally constant sheaf on  $M$  with fiber a finite dimensional Lie algebra; this Lie algebra is called the *Killing Lie algebra* of the Cartan geometry. If the Killing Lie algebra is transitive on an open dense subset  $U$  in  $M$ , we say that  $(E_H, \eta)$  is locally homogeneous on  $U$ .

A flat Cartan geometry with model  $(G, H)$  is locally homogeneous on  $M$ ; its Killing Lie algebra is the Lie algebra  $\mathfrak{g}$  of  $G$ . In particular, a complex affine structure on  $M$  is locally homogeneous, and its local Killing Lie algebra is that of holomorphic affine vector fields (isomorphic to the Lie algebra of the complex affine group  $G = \text{GL}(n, \mathbb{C}) \times \mathbb{C}^n$ ).

We will see in Section 3 that all OT-manifolds admit, by construction, a standard complex affine structure.

## 3. OELJEKLAUS–TOMA MANIFOLDS

An important new class of non-Kähler complex manifolds, generalizing Inoue surfaces in class VII<sub>0</sub> [In], was constructed by Oeljeklaus and Toma, [OT], using ingredients from algebraic number theory. In this section we briefly describe the construction of those Oeljeklaus–Toma manifolds (which will be also denoted as OT–manifold) highlighting their geometric properties which will be useful for our study.

Let us follow [OT] (see also [PV]) and consider an algebraic number field  $F$  of degree  $n$  over  $\mathbb{Q}$ . We denote by  $\mathcal{O}_F$  the ring of algebraic integers of  $F$ , and by  $\mathcal{O}_F^*$  its multiplicative group of units. We assume that  $F$  admits  $n = s + 2t$  embeddings into  $\mathbb{C}$ , with  $s > 0$  of them being real and  $2t > 0$  being complex non-real. Let us denote by  $\sigma_i$ , with  $i \in \{1, \dots, s\}$ , the real embeddings and by  $\sigma_{s+i} = \bar{\sigma}_{s+i+t}$ , where  $i \in \{1, \dots, t\}$ , the complex non-real ones.

Using the above embeddings one defines a natural map

$$\sigma : F \longrightarrow \mathbb{C}^{s+t}, \quad x \longmapsto (\sigma_1(x), \dots, \sigma_{s+t}(x)).$$

The image of the algebraic integers  $\mathcal{O}_F$  through  $\sigma$  is known to be a lattice of rank  $n = s + 2t$  in  $\mathbb{C}^{s+t}$ . Therefore there is a natural action of  $\mathcal{O}_F$  on  $\mathbb{C}^{s+t}$  by translations, defined as

$$T(a, (z_1, \dots, z_{s+t})) = (\sigma_1(a) + z_1, \dots, \sigma_{s+t}(a) + z_{s+t}),$$

for all  $a \in \mathcal{O}_F$ .

There is as well a natural multiplicative action  $M$  of  $\mathcal{O}_F^*$  on  $\mathbb{C}^{s+t}$  given by the formula

$$M(u, (z_1, \dots, z_{s+t})) = (\sigma_1(u)z_1, \dots, \sigma_{s+t}(u)z_{s+t}),$$

for all  $u \in \mathcal{O}_F^*$ .

The above actions  $M$  and  $T$  together define an action of  $\mathcal{O}_F^* \times \mathcal{O}_F$  on  $\mathbb{C}^{s+t}$  by affine transformations. More precisely, an element  $(u, a) \in \mathcal{O}_F^* \times \mathcal{O}_F$  acts on  $\mathbb{C}^{s+t}$  as  $M_u \circ T_a$ . The locus of fixed points of this  $\mathcal{O}_F^* \times \mathcal{O}_F$ -action on  $\mathbb{C}^{s+t}$  is contained in  $\mathbb{R}^s \times \mathbb{C}^t$ . Moreover, if one denotes by  $\mathcal{O}_F^{*,+}$  the subgroup of positive units, meaning those units which are positive in all real embeddings of  $F$ , the above  $\mathcal{O}_F^* \times \mathcal{O}_F$  action induces an  $\mathcal{O}_F^{*,+} \times \mathcal{O}_F$  action on  $\mathbb{C}^{s+t}$  preserving  $\mathbb{H}^s \times \mathbb{C}^t$ , where  $\mathbb{H}$  is the upper-half space. Therefore, this  $\mathcal{O}_F^{*,+} \times \mathcal{O}_F$ -action on  $\mathbb{H}^s \times \mathbb{C}^t$  is free. In general this action is not properly discontinuous, but it is proved in [OT] that one can find for any  $F$  an *admissible* subgroup  $U \subset \mathcal{O}_F^{*,+}$  such that the induced action of  $U \times \mathcal{O}_F$  on  $\mathbb{H}^s \times \mathbb{C}^t$  is properly discontinuous with compact quotient. Moreover, for  $t = 1$ , any finite index subgroup  $U \subset \mathcal{O}_F^{*,+}$  is admissible.

**Definition 3.1.** For any algebraic field extension  $[F, \mathbb{Q}] = n = s + 2t$ , with  $s > 0$  and  $t > 0$ , and any admissible subgroup  $U \subset \mathcal{O}_F^{*,+}$ , the compact complex manifold constructed as quotient of  $\mathbb{H}^s \times \mathbb{C}^t$  by the above  $U \times \mathcal{O}_F$ -action is called an *Oeljeklaus–Toma manifold* (abbreviated as *OT–manifold*).

It was proved in [OT] that all Oeljeklaus–Toma manifolds are non-Kähler.

Since the above  $U \times \mathcal{O}_F$ -action on  $\mathbb{H}^s \times \mathbb{C}^t$  is by complex affine transformations, the affine structure of  $\mathbb{H}^s \times \mathbb{C}^t$ , induced by its inclusion in the standard complex affine space  $\mathbb{C}^{s+t}$ ,

descends on the quotient by  $U \ltimes \mathcal{O}_F$  and endows any quotient OT–manifold with a standard *complex affine structure*.

A very useful result dealing with the algebraic dimension of the OT–manifolds was obtained in [BO] where the authors prove that all OT–manifolds are of algebraic dimension zero. This means that there are no nonconstant meromorphic functions on a OT–manifold.

Therefore, OT–manifolds share with Inoue surfaces of class VII<sub>0</sub> the properties that they are non-Kähler manifolds with algebraic dimension zero and admit a standard complex affine structure.

The *locally conformally Kähler geometry* of OT–manifolds was studied in [OV, PV]. Let us recall that a *locally conformally Kähler metric* on a complex manifold  $M$  is a conformal class of Hermitian metrics on  $M$  which admits a local representative given by a Kähler metric. As it was first observed in [Va],  $M$  admits a locally conformally Kähler metric if and only if  $M$  has a Galois covering which admits a Kähler metric satisfying the condition that the Galois group of the covering acts by holomorphic homotheties with respect to the Kähler metric.

In [OT] its authors proved that for  $t = 1$  (number of complex non-real embeddings) all corresponding OT–manifolds admit a locally conformally Kähler metric.

Following [OV], we will say that a OT–manifold is locally conformally Kähler if it is constructed from a number field  $F$  with  $t = 1$  (meaning it admits exactly 2 complex non-real embeddings).

Ornea and Verbitsky proved in [OV] that all locally conformally Kähler OT–manifolds are *simple*, meaning they do not admit closed proper subvarieties of positive dimension.

For general OT–manifolds, it was proved in [OVV, Theorem 1.2] that all their irreducible subvarieties  $Z$  of smallest possible positive dimension are complex affine in the sense that at any smooth point of  $Z$ , the tangent sub-bundle to  $Z$  is preserved by the standard flat affine connection of the ambient OT–manifold.

#### 4. HOLOMORPHIC GEOMETRIC STRUCTURES ON OELJEKLAUS–TOMA MANIFOLDS

**Theorem 4.1.** *Let  $M$  be an OT–manifold, and let  $\phi$  be a holomorphic geometric structure of affine type on  $M$ . Then  $\phi$  is locally homogeneous. Moreover, the Lie algebra of local holomorphic affine vector fields preserving  $\phi$  is transitive on  $M$ .*

*Proof.* We will prove that any OT–manifold  $M$  satisfies the following property: Any non-zero holomorphic tensor has an empty vanishing set, namely for any integers  $p, q \geq 0$  a non-zero holomorphic section of the holomorphic vector bundle  $(TM)^{\otimes p} \otimes (T^*M)^{\otimes q}$  does not vanish anywhere on  $M$ . Therefore, all holomorphic geometric structures of affine type on  $M$  are locally homogeneous as a consequence of the criteria proved in Lemma 3.2 in [Du1].

It was recently proved in [Ka] (see Remark 3.2 of [Ka]) that for any nonnegative integers  $p, q$ , the holomorphic vector bundle  $(TM)^{\otimes p} \otimes (T^*M)^{\otimes q}$  over the OT–manifold splits as a sum of holomorphic line bundles over  $M$ . By Theorem 2.1 in [APV], any holomorphic line bundle

over an OT–manifold is flat. Now a flat line bundle over an OT–manifold admits a non-zero holomorphic section if and only if it is trivial; consequently, the non-zero holomorphic section is a holomorphic section of the trivial bundle and it does not vanish anywhere. Indeed, this was first prove in [In] (see Proposition 2, i) on page 274) for Inoue surfaces with the consequence that Inoue surfaces do not contain curves. The same argument generalizes to OT–manifolds thanks to Lemma 2.4 (on page 167) in [OT], with the consequence that OT–manifolds do not contain divisors (see Proposition 2.5 in [APV] and Theorem 3.5 in [BO]).

This finishes the proof of the above mentioned property, and, as mentioned before Lemma 3.2 in [Du1] applies.

Notice that the juxtaposition of  $\phi$  with the standard complex affine structure on  $M$  provides a holomorphic geometric structure  $\phi'$  on  $M$  of affine type [Gr, DG]. Since  $\phi'$  is locally homogeneous, its Killing Lie algebra is transitive  $M$ . The Killing Lie algebra of  $\phi'$  is the intersection of the Killing Lie algebra of  $\phi$  with the Killing Lie algebra of the standard complex affine structure. It is the Lie algebra of affine vector fields preserving  $\phi$ ; so this last Lie algebra is transitive on  $M$ .  $\square$

**Theorem 4.2.** *Let  $M$  be an OT–manifold which is locally conformally Kähler. Suppose  $M$  is endowed with a holomorphic geometric structure  $\phi$ . Then  $\phi$  is locally homogeneous. Moreover, the Lie algebra of local holomorphic affine vector fields preserving  $\phi$  is transitive on  $M$ .*

*Proof.* The starting point of the proof is the property, proved by Ornea and Verbitsky in [OV, Corollary 1.7], that locally conformally Kähler OT–manifolds are of algebraic dimension zero, meaning they do not admit non-constant meromorphic functions. Therefore, Theorem 3 and its Corollary 4 in [Du1] (see also Theorem 2.1 and its Corollary 2.2 in [Du3]) imply that any holomorphic rigid geometric structures on  $M$  is locally homogeneous on an open dense subset  $M \setminus S$ , where  $S$  is a nowhere dense closed analytic subset of  $M$  of positive codimension.

Let us now consider a (possibly non-rigid) holomorphic geometric structure  $\phi$  on  $M$ . Add together the holomorphic affine structure of the OT–manifold with  $\phi$  in order to obtain a holomorphic rigid geometric structure in Gromov’s sense (the juxtaposition of a geometric structure with a rigid one is a rigid geometric structure) [Gr, DG]; denote the rigid geometric structure thus obtained by  $\phi'$ . The Killing Lie algebra of  $\phi'$  is the intersection of the Killing Lie algebra of  $\phi$  with the sheaf of the affine vector fields (with respect to the affine structure of the OT–manifold). Applying the above mentioned result of [Du1] to  $\phi'$  one gets that the Killing Lie algebra of  $\phi'$  is transitive on  $M \setminus S$ , with  $S$  being the closed analytic subset of  $M$  where the Killing Lie algebra of  $\phi'$  drops rank.

Moreover, the main result in [OV] (Theorem 1.3 of [OV]) proves that locally conformally Kähler OT–manifolds do not admit non-trivial complex submanifolds. This implies that  $S$  is of dimension zero: it is a finite number of points.

To prove by contradiction, assume that  $S$  is non-empty.

Let us consider the universal cover  $\widetilde{M}$  of  $M$  which identifies with the open set  $\mathbb{H}^s \times \mathbb{C}^t$  in the standard complex affine space  $\mathbb{C}^{s+t}$ . The pull-back of the sheaf of Killing fields of  $\phi'$  to the universal cover  $\widetilde{M}$  is generated by global section [Am]: it identifies with the Lie subalgebra  $\mathfrak{g}'$  of complex affine vector fields in  $\mathfrak{gl}(s+t, \mathbb{C}) \ltimes \mathbb{C}^{s+t}$  which preserve the pull-back of  $\phi$  to  $\widetilde{M}$  (it is the Killing Lie algebra of the pull-back of  $\phi'$  to  $\widetilde{M}$ ). Let us consider a basis  $(k_1, \dots, k_l)$  of  $\mathfrak{g}'$ . The inverse image of  $S$  in universal cover is the subset  $\widetilde{S} \subset \widetilde{M}$  which identifies with the common zeros of the affine vector fields  $k_i$ . Therefore  $\widetilde{S}$  is an algebraically Zariski closed set in  $\mathbb{H}^s \times \mathbb{C}^t$  (it coincides with the intersection of  $\mathbb{H}^s \times \mathbb{C}^t$  with a linear subvariety in  $\mathbb{C}^{s+t}$ ). On the other hand we have seen that  $\widetilde{S}$  has dimension zero. Moreover, since the fundamental group of  $M$  is infinite, we know that  $\widetilde{S}$  is an infinite discrete subset: it cannot be an algebraically Zariski closed subset in  $\mathbb{H}^s \times \mathbb{C}^t$ ; a contradiction. This completes the proof.  $\square$

In the case of a general OT-manifold the above proof gives the following weaker result:

**Proposition 4.3.** *A holomorphic geometric structure  $\phi$  on a OT-manifold is locally homogeneous on  $M \setminus S$ , with  $S$  a nowhere dense analytic subset of positive codimension. Moreover, the Lie algebra of local holomorphic affine vector fields preserving  $\phi$  is transitive on  $M \setminus S$ .*

*Proof.* As in the proof of Theorem 4.2 one considers the holomorphic rigid geometric structure  $\phi'$  which is the juxtaposition of the geometric structure  $\phi$  with the standard complex affine structure of the OT-manifold  $M$ . The main result in [BO] proves that all OT-manifolds have algebraic dimension zero. Therefore, Theorem 3 and its Corollary 4 in [Du1] (see also Theorem 2.1 and its Corollary 2.2 in [Du3]) imply that  $\phi'$  is locally homogeneous on an open dense subset  $M \setminus S$ , where  $S$  is a closed subset in  $M$ . This implies that the Killing Lie algebra of  $\phi'$ , which coincides with the Lie algebra of affine vector fields preserving  $\phi$ , is transitive on  $M \setminus S$ . Moreover,  $S$  coincides with the locus where this Lie algebra of affine vector fields drops rank. Consequently,  $S$  is a nowhere dense analytic subset of positive codimension in  $M$ .  $\square$

**Corollary 4.4.** *Let  $M$  be an OT-manifold associated to an admissible subgroup of units  $U$  such that any element  $u \in U \setminus \{1\}$  is a primitive element for the number field  $F$ . Then any holomorphic geometric structure  $\phi$  on  $M$  is locally homogeneous. Moreover, the intersection of the Killing Lie algebra of  $\phi$  with the Killing Lie algebra of the affine structure is transitive on  $M$ .*

*Proof.* The same proof of the previous Proposition, together with the fact that, under the assumptions,  $M$  does not admit nontrivial complex subvarieties by [OVV, Theorem 1.1], imply that the Killing Lie subalgebra of the affine structure preserving the holomorphic geometric structure is transitive away from a finite set of points  $S$ . We prove that  $S$  is empty as in the proof of Theorem 4.2.  $\square$

Let us now state and prove the analogous results for Cartan geometries.

**Theorem 4.5.** *Let  $M$  be a complex manifold of algebraic dimension zero, bearing a holomorphic torsion-free affine connection  $\nabla$ . Then the following statements hold:*



- (i) *Any holomorphic Cartan geometry of algebraic type on  $M$  is locally homogeneous on an open dense subset  $M \setminus S$ , with  $S$  a nowhere dense closed analytic subset of positive codimension.*
- (ii) *The intersection of the Killing Lie algebra of the Cartan geometry with the Killing Lie algebra of  $\nabla$  is transitive on  $M \setminus S$ .*

*Proof.* Let  $M$  be a complex manifold of complex dimension  $n$ . Let  $(E_H, \eta)$  be a Cartan geometry on  $M$  of type  $(G, H)$ . There is a canonical isomorphism of vector bundles between  $TM$  and  $E_H \times_H \mathfrak{g}/\mathfrak{h}$ , where  $H$  acts on  $\mathfrak{g}/\mathfrak{h}$  via the adjoint representation  $\text{Ad}$ . Moreover, each point  $p \in E_H$  lying above  $b \in M$  defines an isomorphism  $i_p : T_b M \rightarrow \mathfrak{g}/\mathfrak{h}$  such that  $i_{ph} = \text{Ad}(h^{-1})i_p$ , for all  $h \in H$  (see [Sh, p. 188, Theorem 3.15]).

Denote by  $K$  the kernel the adjoint representation  $\text{Ad} : H \rightarrow \text{GL}(\mathfrak{g}/\mathfrak{h})$ , and also denote by  $F_H$  the quotient of  $E_H$  by  $K$ . It follows that the principal  $H/K$ -bundle  $F_H$  get identified with a principal subbundle of the first jet bundle  $R^1(M)$ . This defines a quotient map

$$i : E_H \rightarrow F_H \simeq R^1(M).$$

Notice that the form  $\eta$  trivializes the holomorphic tangent vector bundle of  $E_H$  and gives an isomorphism of vector bundles:  $TE_H \simeq E_H \times \mathfrak{g}$ . The curvature  $\Omega$  of the Cartan connection is completely determined by an  $H$ -equivariant map  $\Omega : E_H \rightarrow V$ , with  $V = \Lambda^2(\mathfrak{g}/\mathfrak{h}) \otimes \mathfrak{g}$ , with  $H$  acting linearly on  $V$  by

$$h \cdot l(u, v) = (\text{Ad}(p) \circ l)(\widetilde{\text{Ad}}(h^{-1})u, \widetilde{\text{Ad}}(h^{-1})v)$$

for all  $h \in H$ , where  $\widetilde{\text{Ad}}$  is the induced adjoint  $H$ -action on  $\mathfrak{g}/\mathfrak{h}$  (see [Sh, Chapter 5, Lemma 3.23]).

Following [Me, Pe] we define the  $m$ -jet of the Cartan curvature  $\Omega$  of  $(E_H, \eta)$  as a map:

$$J^m \Omega : E_H \rightarrow \text{Hom}(\otimes^m \mathfrak{g}, V)$$

defined as  $J^m \Omega(bh^{-1}) = h \cdot (J^m \Omega(b) \circ \text{Ad}^m h^{-1})$ , for all  $b \in E_H$ , with  $\text{Ad}^m$  being the  $m$ -th tensor power  $\otimes^m \mathfrak{g}$  of the adjoint representation restricted of  $H$  (see [Me, Proposition 3.1]).

Consider now, as in [Me, Pe], the map

$$\mathcal{D}^m : E_H \rightarrow V \oplus \text{Hom}(\mathfrak{g}, V) \oplus \cdots \oplus \text{Hom}(\otimes^m \mathfrak{g}, V)$$

defined as  $\mathcal{D}^m = (\Omega, J^1 \Omega, \dots, J^m \Omega)$ .

It was proved in [Me, Theorem 4.1] and in [Pe, Theorem 1.1] that, for  $m = \dim(H)$ , the image of the map  $\mathcal{D}^m$  is exactly a  $H$ -orbit if and only if the Cartan geometry  $(E_H, \eta)$  is locally homogeneous.

Notice that  $(E_H, \eta)$  being of algebraic type the  $H$ -action on

$$V \oplus \text{Hom}(\mathfrak{g}, V) \oplus \cdots \oplus \text{Hom}(\otimes^m \mathfrak{g}, V)$$

is algebraic. For algebraic actions, Rosenlicht's Theorem, [Ro], implies that  $H$ -orbits in general position in  $V \oplus \text{Hom}(\mathfrak{g}, V) \oplus \cdots \oplus \text{Hom}(\otimes^m \mathfrak{g}, V)$  are separated by rational functions. In other words, any  $H$ -invariant rational function  $\mathcal{R}$  on  $V \oplus \text{Hom}(\mathfrak{g}, V) \oplus \cdots \oplus \text{Hom}(\otimes^m \mathfrak{g}, V)$  defines an  $H$ -invariant meromorphic map  $\mathcal{R} \circ \mathcal{D}^m$  on  $E_H$ . Consequently, the meromorphic

map  $\mathcal{R} \circ \mathcal{D}^m$  descends on  $M$  as a constant function, since the algebraic dimension of  $M$  is zero. This implies that  $\mathcal{R}$  is constant on  $\mathcal{D}^m(E_H)$ . From this it follows that there exists an open dense set  $U$  in  $M$  such that  $\mathcal{D}^m(E_H|_U)$  is exactly an  $H$ -orbit. Therefore,  $(E_H, \eta)$  is locally homogeneous over  $U$ ; more details can be found in [Du2, Corollary 1.3].

Since the Killing Lie algebra for  $(E_H, \eta)$  is transitive on an open dense subset, it follows that  $U = M \setminus S$ , with  $S$  being the closed subset of  $M$  where the Killing Lie algebra drops rank. Consequently,  $S$  is a nowhere dense analytic subset of positive codimension in  $M$ . This proves point (i) of Theorem 4.5.

Proof of (ii). Let us now consider the geometric structure which is the juxtaposition of the Cartan geometry  $(E_H, \eta)$  and the holomorphic affine connection  $\nabla$ . This juxtaposition is not in general a Cartan geometry, but we will see that this can be handled as an extended Cartan geometry in the sense of [Pe].

Consider a point  $(b, l)$  in the frame bundle  $R^1(M)$  above  $b \in M$ . This point defines an isomorphism  $l$  between  $\mathbb{C}^n$  and  $T_b M$ . The point  $(b, l) \in R^1(M)$  defines a unique system of local coordinates at  $b \in M$  which are exponential with respect to  $\nabla$  (meaning that the Christoffel coefficients of  $\nabla$  vanishes at the origin for this system of local coordinates). Then we consider the  $m$ -jet of the curvature of  $\nabla$  at the origin with respect to this exponential coordinates. This gives us a  $\mathrm{GL}(n, \mathbb{C})$ -equivariant map  $R^1(M) \rightarrow W^m$ , with  $W^m$  being the vector space of  $m$ -jets of curvature. Replacing the point  $(b, l) \in R^1(M)$  by  $(b, l') \in R^1(M)$ , where  $l' = lh$  with  $h \in \mathrm{GL}(n, \mathbb{C})$ , gives another system of local exponential coordinates at  $b$ , and taking the  $m$ -jet of the curvature in those coordinates defines the action of  $\mathrm{GL}(n, \mathbb{C})$  on  $W^m$ . This action is algebraic (see [DG] for more details).

Therefore the  $m$ -jet of the curvature of  $\nabla$  is given by a  $\mathrm{GL}(n, \mathbb{C})$ -equivariant map

$$\mathcal{K}^m : R^1(M) \rightarrow W^m.$$

Now consider the  $H$ -equivariant map

$$E_H \rightarrow V \oplus \mathrm{Hom}(\mathfrak{g}, V) \oplus \dots \oplus \mathrm{Hom}(\otimes^m \mathfrak{g}, V) \oplus W^m$$

defined by

$$p \mapsto \mathcal{D}^m(p) \oplus (\mathcal{K}^m \circ i)(p),$$

where  $i : E_H \rightarrow F_H \simeq R^1(M)$  is the quotient bundle map described above.

As before, Rosenlicht's Theorem, [Ro], implies that there exists an open dense set  $U$  in  $M$  such that  $(\mathcal{D}^m \oplus (\mathcal{K}^m \circ i))(E_H|_U)$  is exactly an  $H$ -orbit. Therefore,  $(E_H, \eta)$  together with  $\nabla$  form an extended Cartan geometry which is locally homogeneous over  $U$  [Pe, Section 4.4.2 and Section 4.4.3]. Hence the intersection between the Killing Lie algebra of  $(E_H, \eta)$  and the Killing Lie algebra of  $\nabla$  is transitive on  $U$ . It follows that  $U = M \setminus S$ , with  $S$  being the closed subset of  $M$  where the Killing Lie algebra drops rank. Therefore,  $S$  is a nowhere dense analytic subset of positive codimension in  $M$ .  $\square$

**Corollary 4.6.** *A holomorphic Cartan geometry of algebraic type on an OT-manifold is locally homogeneous on  $M \setminus S$ , where  $S$  is a nowhere dense analytic subset of positive codimension in  $M$ . Moreover, the Lie algebra of local holomorphic affine vector fields on  $M$  preserving the Cartan geometry is transitive on  $M \setminus S$ .*

*Proof.* Let  $(E_H, \eta)$  be a holomorphic Cartan geometry of algebraic type on the OT–manifold  $M$ . Since OT–manifolds are known to be of algebraic dimension zero [BO] and are endowed, by construction, with a standard affine structure, [OT], Theorem 4.5 applies to  $(E_H, \eta)$  and to the flat torsion-free affine connection  $\nabla$  associated to the affine structure. Hence the Lie algebra of affine vector fields preserving the Cartan geometry  $(E_H, \eta)$  is transitive on  $M \setminus S$ . Since  $S$  is the subset in  $M$  where the Lie algebra of affine vector fields preserving  $(E_H, \eta)$  drops rank, it follows that  $S$  is a nowhere dense analytic subset of positive codimension in  $M$ .  $\square$

**Corollary 4.7.** *A holomorphic Cartan geometry of algebraic type on a locally conformally Kähler OT–manifold is locally homogeneous. Moreover, the intersection of the Killing Lie algebra of the Cartan geometry with the Killing Lie algebra of the affine structure is transitive.*

*Proof.* By Corollary 4.6, the intersection of the Killing Lie algebra of the Cartan geometry with the Killing Lie algebra of the affine structure is transitive away from the analytic subset  $S$ . By [OV, Theorem 3.1], locally conformally Kähler OT–manifolds do not admit nontrivial complex submanifolds. Therefore  $S$  is a finite union of points. We conclude as in Theorem 4.2 that  $S$  is the empty set.  $\square$

**Corollary 4.8.** *Let  $M$  be an OT–manifold associated to an admissible subgroup of units  $U$  such that any element  $u \in U \setminus \{1\}$  is a primitive element for the number field  $F$ . Then any holomorphic Cartan geometry of algebraic type on  $M$  is locally homogeneous. Moreover, the intersection of the Killing Lie algebra of the Cartan geometry with the Killing Lie algebra of the affine structure is transitive.*

*Proof.* By Corollary 4.6, the intersection of the Killing Lie algebra of the Cartan geometry with the Killing Lie algebra of the affine structure is transitive away from the analytic subset of positive codimension  $S$ . By [OVV, Theorem 1.1]  $M$  does not admit nontrivial complex submanifolds. Therefore  $S$  is a finite union of points. We conclude as in Theorem 4.2 that  $S$  is the empty set.  $\square$

#### ACKNOWLEDGEMENTS

We thank the referee for showing how to correct an argument in the proof of Theorem 1.1. The authors are grateful to Vincent Pecastaing for helpful and friendly discussions and, in particular, for his suggestion to use his results about symmetries of extended Cartan geometries, [Pe], in the proof of Theorem 4.5 and its Corollary 4.7. The authors are also grateful to Hisashi Kasuya, Liviu Ornea, Misha Verbitsky and Victor Vuletescu for helpful comments. The first author is partially supported by a J. C. Bose Fellowship (JBR/2023/000003).

Declaration of interests. The authors do not work for, advise, own shares in, or receive funds from any organisation that could benefit from this article, and have declared no affiliation other than their research organisations.

Data availability statement. No data were generated or used.

## REFERENCES

- [Am] A. M. Amores, Vector fields of a finite type  $G$ -structure, *Jour. Diff. Geom.* **14** (1980), 1–6.
- [APV] D. Angella, M. Parton and V. Vuletescu, Rigidity of Oeljeklaus-Toma manifolds, *Ann. Inst. Fourier* **70** (2020), 2409–2423.
- [BO] L. Battisti and K. Oeljeklaus, Holomorphic line bundles over domains in Cousin groups and the algebraic dimension of OT-manifolds, *Proc. Edin. Math. Soc.* **58**, (2015), 273–285.
- [DG] G. D’Ambra and M. Gromov, *Lectures on transformations groups: geometry and dynamics*, Surveys in Differential Geometry, Cambridge MA, (1991).
- [Du1] S. Dumitrescu, Structures géométriques holomorphes sur les variétés complexes compactes, *Ann. Scient. Éc. Norm. Sup.* **34** (2001), 557–571.
- [Du2] S. Dumitrescu, Killing fields of holomorphic Cartan geometries, *Monat. Math.* **161**, (2010), 145–154.
- [Du3] S. Dumitrescu, Meromorphic almost rigid geometric structures, *Geometry, Rigidity and Group Actions*, Editors Benson Farb and David Fisher, Chicago Lectures in Mathematics Series, 2011, 32–58.
- [Gr] M. Gromov, Rigid Transformations Groups, Editors D. Bernard and Y. Choquet Bruhat, *Géométrie Différentielle*, **33**, Hermann, (1988), 65–139.
- [Ho] G. Hochschild, *Basic theory of algebraic groups and Lie algebras*, Graduate Texts in Mathematics, 75, Springer-Verlag, New York-Berlin, 1981.
- [Hu] J. E. Humphreys, *Linear algebraic groups*, Grad. Texts in Math., No. 21 Springer-Verlag, New York-Heidelberg, 1975.
- [Ka] H. Kasuya, Cohomology of holomorphic line bundles and Hodge symmetry on Oeljeklaus-Toma manifolds, *Euro. J. Math.* **9** (2023), 15 pp.
- [Ka2] H. Kasuya, Vaisman metrics on solvmanifolds and Oeljeklaus-Toma manifolds, *Bull. London Math. Soc.* **45** (2013), 15–26.
- [Ko] S. Kobayashi, *Differential geometry of complex vector bundles*, Princeton University Press, Princeton, NJ, Iwanami Shoten, Tokyo, 1987.
- [In] M. Inoue, On surfaces of class  $VII_0$ , *Invent. Math.* **24**, (1974), 269–310.
- [Me] K. Melnick, A Frobenius theorem for Cartan geometries, with applications, *Enseignement Math.* **57** (2011), 57–89.
- [OT] K. Oeljeklaus and M. Toma, Non-Kähler compact complex manifolds associated to number fields, *Ann. Inst. Fourier* **55** (2005), 1291–1300.
- [OV] L. Ornea and M. Verbitsky, Oeljeklaus-Toma manifolds admitting no complex subvarieties, *Math. Res. Lett.* **18** (2011), 747–754.
- [OVV] L. Ornea, M. Verbitsky and V. Vuletescu, Flat affine subvarieties in Oeljeklaus-Toma manifolds, *Math. Zeit.* **292** (2019), 839–847.
- [PV] M. Parton and V. Vuletescu, Examples of non-trivial rank in locally conformal Kähler geometry, *Math. Zeit.* **270** (2012), 179–187.
- [Pe] V. Pecastaing, On two theorems about local automorphisms of geometric structures, *Ann. Inst. Fourier* **66** (2016), 175–208.
- [Ro] M. Rosenlicht, A remark on quotient spaces, *An. Acad. Sci. Brazil Ci.* **35**, (1963), 487–489.
- [Sh] R. Sharpe, *Differential Geometry: Cartan’s generalization of Klein’s Erlangen program*, New York: Springer, (1996).
- [Va] I. Vaisman, Locally conformal Kähler manifolds with parallel Lee form, *Rend. Math. Roma* **12** (1979), 263–284.
- [Va2] I. Vaisman, On locally and globally conformal Kähler manifolds, *Trans. Amer. Math. Soc.* **262**, (1980), 533–542.
- [Ve] S. M. Verbitskaya, Curves in Oeljeklaus-Toma manifolds, *Funct. Anal. Appl.* **48**, 223–226.

DEPARTMENT OF MATHEMATICS, SHIV NADAR UNIVERSITY, NH91, TEHSIL DADRI, GREATER NOIDA, UTTAR PRADESH 201314, INDIA

*Email address:* indranil.biswas@snu.edu.in, indranil29@gmail.com

UNIVERSITÉ CÔTE D’AZUR, CNRS, LJAD, FRANCE

*Email address:* dumitres@unice.fr