

Countably Colorful Hyperplane Transversal

Sutanoya Chakraborty*

Arijit Ghosh*

Soumi Nandi*

Abstract

Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be an infinite sequence of families of compact connected sets in \mathbb{R}^d . An infinite sequence of compact connected sets $\{B_n\}_{n \in \mathbb{N}}$ is called *heterochromatic sequence* from $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ if there exists an infinite sequence $\{i_n\}_{n \in \mathbb{N}}$ of natural numbers satisfying the following two properties

- *Property A*: $\{i_n\}_{n \in \mathbb{N}}$ is a monotonically increasing sequence, and
- *Property B*: for all $n \in \mathbb{N}$, we have $B_n \in \mathcal{F}_{i_n}$

We show that if every heterochromatic sequence from $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ contains $d + 1$ sets that can be pierced by a single hyperplane then there exists a finite collection \mathcal{H} of hyperplanes from \mathbb{R}^d that pierces all but finitely many families from $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$. As a direct consequence of our result, we get that if every countable subcollection from an infinite family \mathcal{F} of compact connected sets in \mathbb{R}^d contains $d + 1$ sets that can be pierced by a single hyperplane then \mathcal{F} can be pierced by finitely many hyperplanes. This is an infinite and colorful generalization of the (p, q) -Theorem for hyperplanes by Alon and Kalai (Discrete & Computational Geometry 1995). To establish the optimality of our result we show, for all $d \in \mathbb{N}$, there exists an infinite sequence $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ of families of compact connected sets satisfying the following two conditions

- *Condition 1*: for all $n \in \mathbb{N}$, \mathcal{F}_n is not pierceable by finitely many hyperplanes, and
- *Condition 2*: for any $m \in \mathbb{N}$ and every sequence $\{B_n\}_{n=m}^{\infty}$ of compact connected sets in \mathbb{R}^d , where $B_i \in \mathcal{F}_i$ for all $i \geq m$, there exists a hyperplane in \mathbb{R}^d that pierces at least $d + 1$ sets in the sequence.

1 Introduction

Helly's Theorem, proved more than a century ago has been one of the most fundamental results in Convex and Discrete Geometry [ADLS17, BK22]. In its original form, Helly's Theorem, asserts that for a family \mathcal{F} of compact convex sets in \mathbb{R}^d if any $d + 1$ members from \mathcal{F} have a nonempty intersection then the intersection of all the members of \mathcal{F} is also nonempty. Note that the compactness condition can be removed if \mathcal{F} is finite, however, if \mathcal{F} is not finite then the compactness condition becomes necessary.

Geometric transversal theory, an area that studies extensions of the Helly Theorem and its different relatives, has been an active area of research in Convex and Discrete Geometry. In this paper, we will study an infinite variant of geometric transversal.

We will first give some required definitions from geometric transversal theory. A k -flat H is said to *hit* or *pierce* a set $F \subseteq \mathbb{R}^d$ if intersection of F and H is nonempty. A set \mathcal{T} of k -flats is defined as a k -*transversal* for a family \mathcal{F} of subsets in \mathbb{R}^d if, for every $F \in \mathcal{F}$, there exists a k -flat in \mathcal{T} that pierces F . Moreover, when $k = 0$, $k = 1$, or $k = d - 1$, \mathcal{T} is denoted as *point transversal*, *line transversal*, and *hyperplane transversal* of \mathcal{F} , respectively. We also state that a family \mathcal{F} has a k -transversal of size m (or at most m) if there exists a k -transversal \mathcal{T} of \mathcal{F} with $|\mathcal{T}| = m$ (or $|\mathcal{T}| \leq m$). For any two natural numbers p and q , a family (finite or infinite) of subsets of \mathbb{R}^d is said to satisfy the (p, q) -*property with respect to k -flats* if, for any p members

*Indian Statistical Institute, India

of \mathcal{F} , there exists a k -flat that intersects at least q members out of the p members. Hadwiger and Debrunner [HD57] were the first to introduce the (p, q) -property, and they also proved the following important generalization of Helly's Theorem.

Theorem 1.1 (Hadwiger and Debrunner [HD57]). *For all $p \geq q \geq d + 1$ satisfying $q > p(1 - \frac{1}{d}) + 1$, if \mathcal{F} is a family of compact convex sets in \mathbb{R}^d satisfying the (p, q) -property with respect to points, then \mathcal{F} has a point transversal of size at most $p - q + 1$.*

Hadwiger and Debrunner [HD57] asked the following fundamental question.

Question 1.2 (Hadwiger and Debrunner (p, q) -problem [HD57]). *Given $p \geq q \geq d + 1$, does there exist a constant $c(p, q, d) > 0$ such that if a family \mathcal{F} of compact convex sets in \mathbb{R}^d satisfies the (p, q) -property for point transversals, then \mathcal{F} has a point transversal of size at most $c(p, q, d)$?*

Notably, the above problem remained open for 35 years. In a breakthrough paper, Alon and Kleitman [AK92] proved the following fundamental theorem answering the above question.

Theorem 1.3 (Alon-Kleitman [AK92]: (p, q) -theorem). *For any three natural numbers $p \geq q \geq d + 1$, $\exists c = c(p, q, d)$ such that if \mathcal{F} is a finite collection of compact convex sets satisfying the (p, q) -property in \mathbb{R}^d then there exists a point transversal for \mathcal{F} of size at most c .*

Alon and Kleitman's (p, q) -theorem has been extremely influential and since then, this area has seen a lot of interest. Alon and Kalai [AK95] proved the hyperplane variant of the above theorem.

Theorem 1.4 (Alon-Kalai [AK95]: (p, q) -theorem for hyperplane transversal). *For any three natural numbers $p \geq q \geq d + 1$, $\exists c' = c'(p, q, d)$ such that if \mathcal{F} is a finite collection of compact convex sets in \mathbb{R}^d satisfying the (p, q) -property with respect to hyperplanes then there exists a hyperplane transversal for \mathcal{F} of size at most c' .*

Unlike point and hyperplane transversal, Alon, Kalai, Matoušek, and Meshulam [AKMM02] established the impossibility of getting a (p, q) -theorem for k -transversal when $0 < k < d - 1$.

Bárány, Fodor, Montejano, Oliveros, and Pór [BFM⁺14] gave a heterochromatic generalization of Theorem 1.3. Let $\mathcal{F}_1, \dots, \mathcal{F}_p$ be p families of convex sets in \mathbb{R}^d . A *heterochromatic p -tuple* of $\{\mathcal{F}_i \mid i \in [p]\}$ is the collection of p sets C_1, \dots, C_p , where $C_i \in \mathcal{F}_i$ for every $i \in [p]$. The collection $\{\mathcal{F}_i \mid i \in [p]\}$ is said to satisfy *heterochromatic (p, q) -property* if every heterochromatic p -tuple of $\{\mathcal{F}_i \mid i \in [p]\}$ contains q sets which have a non-empty intersection.

Theorem 1.5 (Bárány et al. [BFM⁺14]: Heterochromatic (p, q) -theorem). *Let p, q, d be positive integers with $p \geq q \geq d + 1$. Then there exists $M = M(p, q, d) \in \mathbb{N}$ such that the following holds. Given finite families $\mathcal{F}_1, \dots, \mathcal{F}_p$ of convex sets in \mathbb{R}^d satisfying heterochromatic (p, q) -property, there are $q - d$ indices $i \in [p]$ such that there exists a point transversal for \mathcal{F}_i of size at most M .*

Keller and Perles [KP22] studied infinite variants of (p, q) -type properties with a primary focus on establishing conditions that determine the existence of a finite-sized k -transversal for infinite families of convex sets. As part of this investigation, Keller and Perles [KP22] modified the classical (p, q) -property, referred to as $(\aleph_0, q(k))$ -property with respect to k -flats. This modification involves considering an infinite sequence: within every infinite sequence of convex sets from a family \mathcal{F} , there must exist $q(k)$ sets that can be pierced by a k -flat, where $q(k)$ is an appropriate function of k . Using this new property, Keller and Perles proved the following important theorem.

Theorem 1.6 (Keller-Perles [KP22]: $(\aleph_0, k + 2)$ -theorem for k -transversal). *Let \mathcal{F} be a family of compact convex sets in \mathbb{R}^d , $0 < r < R$, and $0 \leq k \leq d - 1$.*

- (i) *If each convex set C in \mathcal{F} contains a ball of radius $r > 0$ and is contained in a ball of radius R , and \mathcal{F} satisfies $(\aleph_0, k + 2)$ -property with respect to k -flats then the whole family \mathcal{F} has a k -transversal of finite size.*

(ii) If \mathcal{F} is a family of closed balls and \mathcal{F} satisfies $(\aleph_0, 2)$ -property with respect to points then the whole family \mathcal{F} has a point transversal of finite size.

In an updated version [KP23] Keller and Perles have extended Theorem 1.6 to families of *near-balls*. Note that near balls are a generalization of closed balls, and also the convex sets considered in Theorem 1.6. Refer to [KP23] for the exact definition of near-balls.

For all k satisfying $0 < k < d - 1$, Keller and Perles observed the impossibility of getting a $(\aleph_0, k + 2)$ -theorem for k -transversal similar to the above theorem for infinite collections of arbitrary compact convex sets. They also gave an explicit example showing that the compactness assumption in Theorem 1.6 can not be weakened.

Chakraborty, Ghosh, and Nandi [CGN23a] gave heterochromatic generalizations of Theorem 1.6. They also showed that for even non-axis parallel rectangles in \mathbb{R}^2 , $(\aleph_0, 2)$ -Theorem does not hold.

Our results

In this paper, we give the $(\aleph_0, d + 1)$ -Theorem for hyperplane transversal in \mathbb{R}^d .

Theorem 1.7 ($(\aleph_0, d + 1)$ -Theorem for hyperplane transversal). *Let \mathcal{F} be a family of compact connected sets in \mathbb{R}^d such that any infinite subcollection contains $d + 1$ members that can be pierced by a single hyperplane. Then the family \mathcal{F} has a finite hyperplane transversal.*

We get the following corollary directly from the above theorem.

Corollary 1.8 ($(\aleph_0, d + 1)$ -Theorem for hyperplane transversal of convex sets). *Let \mathcal{F} be a family of compact convex sets in \mathbb{R}^d such that any infinite subcollection contains $d + 1$ members that can be pierced by a single hyperplane. Then the family \mathcal{F} has a finite hyperplane transversal.*

Observe that $d + 1$ in Theorem 1.7 is optimal since any nonempty d sets in \mathbb{R}^d can be pierced by a single hyperplane in \mathbb{R}^d . Also, unlike for $k < d - 1$, observe that the above theorem gives a $(\aleph_0, d + 1)$ -Theorem for hyperplane transversal for arbitrary compact connected sets.

Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a sequence of families of compact connected sets in \mathbb{R}^d . A *heterochromatic sequence* with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ is defined as a sequence $\{B_n\}_{n \in \mathbb{N}}$ where we have a strictly increasing sequence $\{m_n\}_{n \in \mathbb{N}}$ in \mathbb{N} such that $\forall n \in \mathbb{N}$, we have $B_n \in \mathcal{F}_{m_n}$. A *strongly heterochromatic sequence* with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ is defined as a sequence $\{B_n\}_{n \in \mathbb{N}}$ where $\forall n \in \mathbb{N}$, we have $B_n \in \mathcal{F}_n$. We will also prove the following heterochromatic generalization of Theorem 1.7.

Theorem 1.9 (Heterochromatic $(\aleph_0, d + 1)$ -Theorem for hyperplane transversal). *Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a sequence of families of compact connected sets such that every heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ has $d + 1$ sets that are pierceable by a single hyperplane. Then all but finitely many \mathcal{F}_n 's are pierceable by finitely many hyperplanes.*

Additionally, from Theorem 1.7 and Theorem 1.9, we can also show the existence of a finite-size collection of hyperplanes for the whole sequence $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ that is a transversal for all but finitely many families from the sequence $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$.

Theorem 1.10 (Stronger variant of Theorem 1.9). *Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a sequence of families of compact connected sets such that every heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ has $d + 1$ sets that are pierceable by a single hyperplane. Then there exists a finite-sized collection \mathcal{H} of hyperplanes in \mathbb{R}^d such that \mathcal{H} pierces all but finitely many families from $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$.*

We complement the above result by showing the following no-go theorem which establishes the optimality of our generalization of Theorem 1.7.

Theorem 1.11 (No-go theorem for strongly heterochromatic generalization). *There exists a sequence $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ of families of compact convex sets in \mathbb{R}^d such that*




Figure 1: Smallest box containing F .

- for all $n \in \mathbb{N}$, \mathcal{F}_n is not pierceable by finitely many hyperplanes, and
- every strongly heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ contains $d + 1$ balls which can be pierced by a single hyperplane.

The proof of the above no-go theorem shows the impossibility of getting a strongly heterochromatic $(\aleph_0, d + 1)$ -Theorem for hyperplanes even when the compact connected sets in question have a nice shape when $d > 1$ as closed unit balls do. Additionally, we can prove the following stronger impossibility result.

Theorem 1.12 (Stronger variant of Theorem 1.11). *There exists a sequence $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ of compact convex sets in \mathbb{R}^d satisfying the following two conditions:*

- for all $n \in \mathbb{N}$, \mathcal{F}_n is not pierceable by finitely many hyperplanes, and
- for any $m \in \mathbb{N}$ and every sequence $\{B_n\}_{n=m}^{\infty}$ of compact connected sets in \mathbb{R}^d , where $B_i \in \mathcal{F}_i$ for all $i \geq m$, there exists a hyperplane in \mathbb{R}^d that pierces at least $d + 1$ sets in the sequence.

Comparison with some recent result

Chakraborty, Ghosh, and Nandi [CGN23b] proved the following related theorem about *axis-parallel boxes*¹.

Theorem 1.13 (Chakraborty, Ghosh, and Nandi [CGN23b]). *Let $\{\mathcal{G}_n\}_{n \in \mathbb{N}}$ be an infinite sequence of collections of compact axis-parallel boxes in \mathbb{R}^d such that every heterochromatic sequence with respect to $\{\mathcal{G}_n\}_{n \in \mathbb{N}}$ has 2 sets that can be pierced by a k -dimensional axis-parallel k -flat. Then there exists a finite-sized collection \mathcal{H}_k of axis-parallel k -flats such that \mathcal{H}_k pierces all but finitely many families from $\{\mathcal{G}_n\}_{n \in \mathbb{N}}$.*

Observe that using the above theorem we get the following corollary.

Corollary 1.14 (Corollary of Theorem 1.13). *Let $\{\mathcal{S}_n\}_{n \in \mathbb{N}}$ be an infinite sequence of collections of compact connected sets in \mathbb{R}^d such that every heterochromatic sequence with respect to $\{\mathcal{S}_n\}_{n \in \mathbb{N}}$ has 2 sets that can be pierced by a axis-parallel hyperplane. Then there exists a finite-sized collection \mathcal{H} of axis-parallel hyperplane such that \mathcal{H} pierces all but finitely many families from $\{\mathcal{S}_n\}_{n \in \mathbb{N}}$.*

Given a connected set $F \in \mathcal{F}$ we consider the box

$$B_F := \pi_{x_1}(F) \times \pi_{x_2}(F) \times \cdots \times \pi_{x_d}(F),$$

¹Defined in Section 2.

see Figure 1. Note that for all $1 \leq i \leq d$, $\pi_{x_i}(F)$ denotes the *orthogonal projection* of F on the i -th axis. Observe that if F is connected then if a axis-parallel hyperplane H pierces B_F then H also pierces F . Using Theorem 1.13, with $k = d - 1$, on the sequence $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$, where $\mathcal{B}_n := \{B_F \mid F \in \mathcal{F}_n\}$ for all $n \in \mathbb{N}$, we get that there exists a constant size family \mathcal{H} of axis-parallel hyperplanes in \mathbb{R}^d that is a hyperplane transversal of all but finitely many families from the sequence $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$. Again as the sets in \mathcal{S}_n are connected for all $n \in \mathbb{N}$, \mathcal{H} is also a hyperplane transversal of all but finitely many families from the sequence $\{\mathcal{S}_n\}_{n \in \mathbb{N}}$.

Jung and Pálvölgyi [JP23] have developed a general framework, using fractional Helly Theorem and (p, q) -theorem, for proving similar $(\aleph_0, k + 2)$ -Theorems. They have given an alternative proofs of Theorem 1.6, Corollary 1.8, and the following result which is a special case of Corollary 1.14.

Theorem 1.15 (Jung and Pálvölgyi [JP23]). *Let \mathcal{F} be a family of compact convex sets in \mathbb{R}^d . If among every infinite sequence of sets from \mathcal{F} contains at least two sets that can be pierced by an axis-parallel hyperplane, then \mathcal{F} has a constant size hyperplane transversal.*

Observe that Theorem 1.13 holds for axis-parallel k -flats (and not just axis-parallel hyperplanes) and it also directly implies a stronger colorful result, see Corollary 1.14.

2 Definitions and notations

- For all $n \in \mathbb{N}$, $[n] := \{1, \dots, n\}$, and $\mathbb{S}^{d-1} := \{x \in \mathbb{R}^d \mid \|x\| = 1\}$.
- An *axis-parallel box* B in \mathbb{R}^d is a set of the form $[a_1, b_1] \times [a_2, b_2] \times \dots \times [a_d, b_d]$, where $\forall i \in [d]$, $a_i, b_i \in \mathbb{R}$ with $a_i \leq b_i$.
- For subsets A and B of \mathbb{R}^d , $\text{CH}(A, B)$ denotes the convex hull of the set $A \cup B$.
- Given a line ℓ in \mathbb{R}^d and $\varepsilon \in (0, \infty)$, define

$$M(\ell, \varepsilon) := \left\{ p \in \mathbb{R}^d \mid d(p, \ell) \leq \varepsilon \right\}.$$

- Given $p \in \mathbb{R}^d$ and $\varepsilon \in (0, \infty)$, define

$$\text{box}(p, \varepsilon) := \left\{ q \in \mathbb{R}^d \mid \forall i \in [d], |q_i - p_i| \leq \varepsilon \right\}.$$

- For a subset B of \mathbb{R}^d and a k -flat S_k , $0 \leq k \leq d$, $\pi_{S_k}(B)$ denotes the *projection* of B on S_k . If $S_k = \{(x_1, \dots, x_d) \in \mathbb{R}^d \mid x_i = 0 \forall k < i \leq d\}$, then we denote $\pi_{S_k}(B)$ by $\pi_k(B)$.
- Let S_k be a k -flat and H be a λ -flat in \mathbb{R}^d . We say H is *perpendicular* to S_k if $H \cap S_k = \pi_{S_k}(H)$.
- Let $A \subseteq \mathbb{R}^d$. We will denote by A° and \bar{A} *interior* and *closure*, respectively, of A . Also, $\text{bd}(A)$ denotes the *boundary* of A , that is, $\text{bd}(A) := \bar{A} \setminus A^\circ$.
- For $c \in \mathbb{R}^d$ and $r \in (0, \infty)$, $B(c, r)$ (and $B^\circ(c, r)$) denotes the *closed ball* (and *open ball*) centered at c with radius r .
- A subset A of \mathbb{R}^d is said to be a *polyhedron* if A is the intersection of finitely many halfspaces. If in addition, A is bounded, then A is called a *polytope*.
- A sequence $\{A_n\}_{n \in \mathbb{N}}$ of sets is called a *nested sequence* if $\forall n \in \mathbb{N}$ we have $A_n \supseteq A_{n+1}$.




Figure 2: Progressively smaller regions containing a 2-set.

- $\text{cone}(\vec{r}, \varepsilon)$: Suppose $\varepsilon > 0$ and \vec{r} is a ray in \mathbb{R}^d from $p \in \mathbb{R}^d$ given by $\vec{r} = \{p + tv \mid t \geq 0\}$ for some $v \in \mathbb{S}^{d-1}$. We define

$$\text{cone}(\vec{r}, \varepsilon) := \{p + tx \mid t \geq 0, x \in B(v, \varepsilon)\}.$$

- Given a ray $\vec{r} = \{p + tv \mid t \geq 0\}$, $v \in \mathbb{S}^{d-1}$ and $n \in \mathbb{N}$, we write

$$R(\vec{r}, 1/n) := \text{cone}(\vec{r}, 1/n) \cap B(p, 1/n)$$

and

$$Q(\vec{r}, n) := \text{cone}(\vec{r}, 1/n) \setminus B(p, n).$$

- *d-set*: \mathcal{F} is a *d-set* if \mathcal{F} is a collection of compact connected sets in \mathbb{R}^d which does not have a finite hyperplane transversal. We say $A \subseteq \mathbb{R}^d$ contains a *d-set* \mathcal{B} or \mathcal{B} is contained in A if $\forall B \in \mathcal{B}, B \subseteq A$.
- *d-point*: Let \mathcal{F} be a *d-set*. A point $p \in \mathbb{R}^d$ is said to be a *d-point* with respect to \mathcal{F} if for every open set A with $p \in A$, \exists a subcollection $\mathcal{F}_A \subseteq \mathcal{F}$ such that \mathcal{F}_A is a *d-set* and for all $B \in \mathcal{F}_A, B \subseteq A$.
- Given sets A_1, A_2, \dots, A_k in S_k where S_k is a k -flat, let Λ be the set of all $(k-1)$ -flats in S_k that intersect A_i for all $i \in [k]$. We define $C_k(A_1, \dots, A_k)$ as $\bigcup_{X \in \Lambda} X$.

3 Outline of the proofs

3.1 $(\aleph_0, d+1)$ -Theorem for Hyperplanes

We will be proving the following contrapositive version of Theorem 1.7.

Theorem 3.1 (Contraposition of Theorem 1.7). *Let \mathcal{F} be a family of compact connected sets in \mathbb{R}^d that cannot be pierced by finitely many hyperplanes. Then there exists an infinite sequence $\{B_n\}_{n \in \mathbb{N}} \subseteq \mathcal{F}$ such that no $d+1$ distinct sets from the sequence can be hit by a hyperplane.*

To highlight the key concepts underpinning the proof of Theorem 3.1, we will discuss two specific scenarios: one where $d = 1$ and another where $d = 2$. The proof for the general case will follow similar lines with certain adjustments, which we will elucidate while sketching out the proofs for the aforementioned instances. The core ideas behind the proof, coupled with some of the details of the construction, can be discerned from this section.

3.1.1 Finite point transversal in \mathbb{R}

If we have a family \mathcal{F} in \mathbb{R} of closed and bounded intervals, which are not pierceable by finitely many points, then there are two possibilities: either for every $n \in \mathbb{N}$ there is an interval $B_n \in \mathcal{F}$ that lies completely outside $[-n, n]$, or there is an $N \in \mathbb{N}$ such that $\forall B \in \mathcal{F}, B \cap [-N, N] \neq \emptyset$. In the first case, it is evident that we can get a sequence of mutually non-intersecting intervals. In the second case, we claim that there is a point $p \in [-N, N]$ such that any neighborhood of p contains a subfamily of \mathcal{F} which cannot be hit by finitely many points. We can prove this claim using the compactness of $[-N, N]$ and the fact that every open cover of a compact set has a finite subcover. For a rigorous proof of the general version of this claim, see Theorem 4.4. Observe that p is the 1-dimensional version of a d -point in \mathbb{R}^d . We will use this point to build the required sequence by first picking a $B_1 \in \mathcal{F}$ that does not contain p . Then there is a neighbourhood $[p - \delta_1, p + \delta_1]$ that does not intersect B_1 . Now pick a $B_2 \in \mathcal{F}$ that is contained in $[p - \delta_1, p + \delta_1]$ but does not contain p . Continuing this way, we get an infinite sequence $\{B_n\}_{n \in \mathbb{N}}$ of non-intersecting intervals in \mathcal{F} as p is a 1-point of \mathcal{F} .

3.1.2 Finite line transversal in \mathbb{R}^2

Let \mathcal{F} be a family of compact connected sets in \mathbb{R}^2 that cannot be pierced by finitely many lines. Just like we saw for \mathbb{R} , we either get a 2-set contained in a bounded region, or there are no 2-sets contained in a bounded region. Recall from the Definitions and notations section that a 2-set contained in a bounded region A means that the sets in the 2-sets are subsets of A . In the first case, as before, we get a 2-point (the proof for the general d -dimension is seen in Theorem 4.4), and in the second case, there are no 2-points corresponding to \mathcal{F} .




Figure 3: Progressively smaller regions about \vec{r}_2 containing a d -set.

Case-I: There is a 2-point $p \in \mathbb{R}^2$ for \mathcal{F}

In this case, any ball $B(p, \delta)$, $\delta > 0$, contains a subfamily of \mathcal{F} that cannot be hit by finitely many lines. For d -dimension, such subfamilies are called d -sets, and here we would call them 2-sets (see Figure 2). We can show that there is a ray \vec{r}_2 whose endpoint is p such that the intersection of any cone around \vec{r}_2 and any ball $B(p, \delta)$ contains a 2-set that is a subset of \mathcal{F} (see Figure 3). This we can show by choosing progressively smaller regions $R(\vec{r}_2, 1/n)$ that contain a 2-set as shown in Figure 3, and the general case is worked out in Theorem 4.5. Now, since for every $n \in \mathbb{N}$, $R(\vec{r}_2, 1/n)$ (as shown in Figure 3) contains a 2-set, we can construct a nested sequence of 2-sets $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$ that are subsets of \mathcal{F} and the sets in each \mathcal{B}_n lie in $R(\vec{r}_2, 1/n)$. Further, we can assume that for every $n \in \mathbb{N}$ and $\forall B \in \mathcal{B}_n$, we have $\vec{r}_2 \cap B = \emptyset$, since the collection of all sets in

\mathcal{F} that intersect \vec{r}_2 cannot be a 2-set, which means that if we drop those sets from \mathcal{F} , it will still be a 2-set.




Figure 4: Obtaining an appropriate $R(\vec{r}_2, \frac{1}{t_2})$.

Now we pick $B_1 \in \mathcal{B}_1$. As per our construction, B_1 does not intersect \vec{r}_2 . Then the lines through B_1 and p , whose union is denoted by $C_2(\{p\}, B_1)$, intersect \vec{r}_2 at only p . So we can pick a $t_2 \in \mathbb{N}$ for which there is an $R(\vec{r}_2, \frac{1}{t_2})$ that does not intersect $C_2(\{p\}, B_1)$. Choose $B_2 \in \mathcal{B}_{t_2}$. Now there is a $t'_2 \in \mathbb{N}$ for which there is an $R(\vec{r}_2, \frac{1}{t'_2})$ that does not intersect $C_2(\{p\}, B_i)$ for $i = 1, 2$ as well as $C_2(B_1, B_2)$. We pick $B_3 \in \mathcal{B}_{t'_2}$, as shown in Figure 4. Proceeding this way, we obtain our sequence. The general construction is described in Section 4.1.1.




Figure 5: Progressively smaller unbounded regions containing a 2-set.

Case-II: There are no 2-points for \mathcal{F}

In this case, we get a ray \vec{r}_2 such that for every $n \in \mathbb{N}$, the region $Q(\vec{r}_2, n)$ contains all sets of a 2-set that is a subset of \mathcal{F} . In other words, we get a nested sequence $\{\mathcal{B}'_n\}_{n \in \mathbb{N}}$ of 2-sets with $\mathcal{B}'_n \subseteq \mathcal{F}$, such that $\forall n \in \mathbb{N}, \forall B \in \mathcal{B}'_n$ we have $B \subset Q(\vec{r}_2, n)$ (see Figure 5, and Theorem 4.15 for proof). Without loss of generality, we can assume that \vec{r}_2 is the nonnegative x_2 -axis and the line passing through its endpoint and perpendicular to it is the x_1 -axis. Let, for all $B \in \mathcal{F}$, $\pi_1(B)$ denote the projection of B onto the x_1 -axis. Since our 2-sets are contained in $Q(\vec{r}_2, n)$'s, we would like to choose sets A, B from \mathcal{F} for our sequence for which $\pi_1(A) \cap \pi_1(B) = \emptyset$, because otherwise for all $n \in \mathbb{N}$, we would have $Q(\vec{r}_2, n) \cap C_2(A, B) \neq \emptyset$, where $C_2(A, B)$ denotes the




Figure 6: $Q(\vec{q}, n)$ containing $Q(\vec{r}_2, t_n)$.

union of all lines through A and B , as defined in Section 2. Let ℓ denote the x_2 -axis, which is also the straight line on which \vec{r}_2 lies. Now, there are two distinct possibilities:

Subcase (1): either there exists an $\varepsilon > 0$ such that $\forall n \in \mathbb{N}$, the subset of \mathcal{B}'_n that is contained in $M(\ell, \varepsilon)$ is a 2-set

Subcase (2): or for every $\varepsilon > 0$, there is an $n_\varepsilon \in \mathbb{N}$ for which the subset of $\mathcal{B}'_{n_\varepsilon}$ that is contained in $M(\ell, \varepsilon)$ is not a 2-set.

In **Subcase (1)**, we can show, as we prove later in Lemma 4.14, that there exists a point $p = (p_1, 0)$ on the x_1 -axis, and $\ell_p := \{p + x \mid x \in \ell\}$ such that for every $n \in \mathbb{N}$, $M(\ell_p, \frac{1}{n})$ contains a 2-set $\mathcal{F}_n \subseteq \mathcal{B}'_n$.

In this subcase, note that if $\vec{q} = \{p + x \mid x \in \vec{r}_2\}$, then there is a strictly increasing sequence $\{t_n\}_{n \in \mathbb{N}} \subseteq \mathbb{N}$ for which $Q(\vec{r}_2, t_n) \subseteq Q(\vec{q}, n)$ for all $n \in \mathbb{N}$ (see Figure 6, and Lemma 4.12 for proof). For every $n \in \mathbb{N}$, set $\mathcal{B}_n = \bigcup_{i=n}^{\infty} \mathcal{F}_i \cap \mathcal{B}'_{t_i}$. For convenience, let us rename \vec{q} as \vec{r}_2 .

In **Subcase (2)**, choose any $\varepsilon > 0$. Then for all $n \in \mathbb{N}$, there exists a $B \in \mathcal{B}'_n$ such that $B \cap M(\ell, \varepsilon) = \emptyset$. In this case, for each $n \in \mathbb{N}$, rename \mathcal{B}'_n as \mathcal{B}_n .

To summarize the notation changes, in order to avoid switching between two subcases, we write \vec{q} as \vec{r}_2 with endpoint at O for **Subcase (1)**, and \mathcal{B}'_n as \mathcal{B}_n for **Subcase (2)**. For the rest of this section, any statement we make about \vec{r}_2 and \mathcal{B}_n for all $n \in \mathbb{N}$ applies for both subcases.

Now note that in both **Subcase (1)** and **Subcase (2)**, if $B_1, \dots, B_m \in \mathcal{B}_1$, then for all $n \in \mathbb{N}$, there is a $B \in \mathcal{B}_n$ such that $\pi_1(B_i) \cap \pi_1(B) = \emptyset$ for all $i \in [m]$. Now we describe the construction of the sequence, the general case of which is covered in Section 4.2.1. Pick any $B_1 \in \mathcal{B}_1$ which does not intersect \vec{r}_2 . So, we can choose a $B_2 \in \mathcal{B}_2$ that does not intersect \vec{r}_2 and $\pi_1(B_1) \cap \pi_1(B_2) = \emptyset$. Then the set consisting of the union of all lines through B_1 and B_2 , denoted by $C_2(B_1, B_2)$, does not contain \vec{r}_2 and $\vec{r}_2 \cap C_2(B_1, B_2)$ is contained in a bounded region (see Lemma 4.17 for proof). So, we can pick a $t_2 \in \mathbb{N}$ for which $Q(\vec{r}_2, n) \cap C_2(B_1, B_2) = \emptyset$ for all $n \geq t_2$. Then we have an $n_2 \in \mathbb{N}$ for which the sets of \mathcal{B}_{n_2} are contained in $Q(\vec{r}_2, t_2)$ (see Figure 7). Pick $B_3 \in \mathcal{B}_{n_2}$ such that B_3 does not intersect \vec{r}_2 and $\pi_1(B_3) \cap \pi_1(B_i) = \emptyset$ for $i = 1, 2$. Continuing this way, we obtain our sequence. For the general construction, see Section 4.2.1.

3.2 Heterochromatic $(\aleph_0, d + 1)$ -Theorem for Hyperplanes

The goal of this section is to give the proof of the following heterochromatic $(\aleph_0, d + 1)$ -Theorem for hyperplanes and (arbitrary) compact connected sets.

Theorem 1.10 (Stronger variant of Theorem 1.9). *Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a sequence of families of compact connected sets such that every heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ has*




Figure 7: Choosing an appropriate $Q(\vec{r}_2, t_n)$.

$d + 1$ sets that are pierceable by a single hyperplane. Then there exists a finite-sized collection \mathcal{H} of hyperplanes in \mathbb{R}^d such that \mathcal{H} pierces all but finitely many families from $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$.

Note that Theorem 1.9 is a direct consequence of the above result.

Theorem 1.9 (Heterochromatic $(\aleph_0, d + 1)$ -Theorem for hyperplane transversal). *Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a sequence of families of compact connected sets such that every heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ has $d + 1$ sets that are pierceable by a single hyperplane. Then all but finitely many \mathcal{F}_n 's are pierceable by finitely many hyperplanes.*

Now consider the following *weaker-looking* heterochromatic theorem compared to Theorem 1.9 and Theorem 1.10.

Theorem 3.2. *Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a sequence of families of compact connected sets in \mathbb{R}^d such that every heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ has $d + 1$ sets that are pierceable by a single hyperplane. Then there exists an $n \in \mathbb{N}$ for which \mathcal{F}_n is pierceable by finitely many hyperplanes.*

We will first deduce Theorem 1.9 from Theorem 3.2. Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a sequence of families of compact connected sets in \mathbb{R}^d such that every heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ has $d + 1$ sets that are pierceable by a single hyperplane. To reach a contradiction assume $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ contains an infinite sequence $\{\mathcal{F}_{i_n}\}_{n \in \mathbb{N}}$ such that no \mathcal{F}_{i_n} can be pierceable by finitely many hyperplanes, where $\{i_n\}_{n \in \mathbb{N}}$ is a strictly increasing sequence in \mathbb{N} . Observe that we will reach a contradiction using Theorem 3.2, since every heterochromatic sequence with respect to $\{\mathcal{F}_{i_n}\}_{n \in \mathbb{N}}$ contains $d + 1$ sets which can be pierceable by a single hyperplane and, from Theorem 3.2, there is an $n \in \mathbb{N}$ for which \mathcal{F}_{i_n} can be pierceable by finitely many hyperplanes.

Using Theorem 1.7 and Theorem 1.9, we can prove that if $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ satisfies the conditions given in Theorem 1.9 then there exists a fixed collection of hyperplanes of constant size that pierces all but finitely many families from the infinite sequence.

Theorem 1.10 (Stronger variant of Theorem 1.9). *Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a sequence of families of compact connected sets such that every heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ has $d + 1$ sets that are pierceable by a single hyperplane. Then there exists a finite-sized collection \mathcal{H} of hyperplanes in \mathbb{R}^d such that \mathcal{H} pierces all but finitely many families from $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$.*

To see how Theorem 1.10 follows from Theorem 1.7 and Theorem 1.9, first observe that from Theorem 1.9 we get that there exists an $m \in \mathbb{N}$ such that for each $n \geq m$, \mathcal{F}_n is pierceable by finitely many hyperplanes. Pick a countably infinite subset \mathcal{S} of $\bigcup_{n \geq m} \mathcal{F}_n$. Then either \mathcal{S}

contains a heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n=m}^\infty$, or \mathcal{S} contains infinitely many sets from \mathcal{F}_r for some $r \geq m$. In either case, by Theorem 1.9 and Theorem 1.7, \mathcal{S} contains $d + 1$ sets that are pierceable by a single hyperplane. Then, by Theorem 1.7, we get that $\bigcup_{n \geq m} \mathcal{F}_n$ is finitely pierceable.

We will give the proof of Theorem 3.2. We will be using the following proposition of Keller and Perles [KP23].

Proposition 3.3 (Keller and Perles [KP23]). *Let \mathcal{F} be a family of compact sets in \mathbb{R}^d , $0 \leq k \leq d - 1$ and $m \in \mathbb{N}$. If any finite subfamily of \mathcal{F} has a k -transversal of size at most m , then \mathcal{F} also has a k -transversal of size at most m .*

Since none of the families in the sequence $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ is finitely pierceable by hyperplanes, from Proposition 3.3 we get that for all $n \in \mathbb{N}$ there exists $\mathcal{S}_n \subseteq \mathcal{F}_n$ satisfying the following properties:

- *Property A:* \mathcal{S}_n does not have a hyperplane transversal of size at most n , and
- *Property B:* \mathcal{S}_n is a finite sized family.

Now consider the following family of compact connected sets:

$$\mathcal{S} := \bigcup_{n \in \mathbb{N}} \mathcal{S}_n.$$

Observe that the family \mathcal{S} by construction does not have a finite-sized hyperplane transversal. Therefore, using Theorem 3.1, we can show that there exists an infinite sequence $\{B_n\}_{n \in \mathbb{N}}$ of compact sets from \mathcal{S} and $\{B_n\}_{n \in \mathbb{N}}$ does not contain $d + 1$ sets that can be pierced by a single hyperplane in \mathbb{R}^d . Since each family \mathcal{S}_n is of finite size, we can extract an infinite subsequence from $\{B_n\}_{n \in \mathbb{N}}$ which is a heterochromatic sequence with respect to $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$. This completes the proof of Theorem 3.2.

3.3 Impossibility of a strongly colorful variant

In this section, we prove Theorem 1.11. Chakraborty, Ghosh and Nandi [CGN23a] proved the following theorem:

Theorem 3.4 (Chakraborty, Ghosh and Nandi [CGN23a]). *For all $d \in \mathbb{N}$ and $0 \leq k < d$, there exists an infinite sequence $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ of families of boxes in \mathbb{R}^d satisfying the following properties:*

- *for all $n \in \mathbb{N}$, \mathcal{F}_n does not have a finite size axis-parallel k -transversal, and*
- *every infinite sequence $\{B_n\}_{n \in \mathbb{N}}$, where $B_n \in \mathcal{F}_n$ for all $n \in \mathbb{N}$, contains two boxes B_i and B_j with $i \neq j$ and both B_i and B_j can be pierced by a single axis-parallel k -flat.*

Putting $d = 1$, we get Theorem 1.11 for \mathbb{R} . It is worth noting that in the above theorem, if for $d = 1$ we substitute families of boxes with families of unit-length intervals, then the theorem is not true. However, for $d > 1$, Theorem 1.11 is true even for unit balls, as the construction below shows. For $d > 1$, pick d d -sets $\mathcal{F}_1, \dots, \mathcal{F}_d$, where $\mathcal{F}_i = \{B_{n,i}\}_{n \in \mathbb{N}}$ and every $B_{n,i}$ is a unit ball. For every d -tuple $(n_1, \dots, n_d) \in \mathbb{N}^d$, we *tightly pack* the region $C_d(B_{n_1,1}, \dots, B_{n_d,d})$ with unit balls, and denote this family of unit balls by $\mathcal{F}_{(n_1, \dots, n_d)}$. It is easy to check that for every $\alpha \in \mathbb{N}^d$, \mathcal{F}_α is a d -set. Now, every strongly heterochromatic sequence chosen from the sequence $\{\mathcal{F}_1, \dots, \mathcal{F}_d\} \cup \{\mathcal{F}_\alpha \mid \alpha \in \mathbb{N}^d\}$ has $d + 1$ sets that are pierced by a single hyperplane, because for any choice of $B_i \in \mathcal{F}_i$ for $i \in [d]$, there is an $\alpha \in \mathbb{N}^d$ for which every set in \mathcal{F}_α pierces a hyperplane that passes through B_1, \dots, B_d . This shows that there cannot be a strongly heterochromatic theorem for any value of $d > 0$.

In the above construction, by taking infinitely many copies of each \mathcal{F}_i for $i \in [d]$ along with \mathcal{F}_α for all $\alpha \in \mathbb{N}^d$, we can prove something stronger than Theorem 1.11.

Theorem 3.5. *Let $d > 1$. There exists a sequence $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ of closed unit balls in \mathbb{R}^d satisfying the following two conditions:*

- *for all $n \in \mathbb{N}$, \mathcal{F}_n is not pierceable by finitely many hyperplanes, and*
- *for any $m \in \mathbb{N}$ and every sequence $\{B_n\}_{n=m}^\infty$ of compact connected sets in \mathbb{R}^d , where $B_i \in \mathcal{F}_i$ for all $i \geq m$, there exists a hyperplane in \mathbb{R}^d that pierces at least $d+1$ sets in the sequence.*

For $d = 1$, the construction given in [CGN23a] can be adapted to satisfy the above theorem when the condition of taking families of unit balls (unit-length intervals) is removed. Combining these results, Theorem 1.12 is satisfied.

4 Proof of Theorem 1.7

We have already proved Theorem 1.7 when $d = 1$ in Section 3.1, and in this section we will prove Theorem 1.7 for $d \geq 2$. First, we present a few observations that are straightforward.

Observation 4.1. *Suppose S_k is a k -flat with $A_1, \dots, A_{k+1} \subseteq S_k$ such that*

$$A_1 \cap C_k(A_2, \dots, A_{k+1}) = \emptyset.$$

Then for any $j \in \{2, \dots, k+1\}$, we have

$$A_j \cap C_k(A_1, \dots, A_{j-1}, A_{j+1}, \dots, A_{k+1}) = \emptyset.$$

Observation 4.2. *Let S_k be a k -flat in \mathbb{R}^d and $P \subseteq \mathbb{R}^d$ be any polyhedron such that if $A = S_k \cap P$, then $\text{bd}(A)$ is nonempty. Then there are finitely many hyperplanes H_1, \dots, H_m such that $\forall i \in [m]$, we have*

$$H_i \cap S_k = \pi_{S_k}(H_i), \text{ and } \text{bd}(A) = \left(\bigcup_{i \in [m]} H_i \right) \cap \bar{A}.$$

Observation 4.3. *Let p be a d -point with respect to a collection of compact connected sets \mathcal{F} in \mathbb{R}^d and A be any open set with $O \in A$. Then there exists a sequence of d -sets $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ with $\mathcal{F} \supseteq \mathcal{F}_n \supseteq \mathcal{F}_{n+1}$, $\forall n \in \mathbb{N}$, such that if $B \in \mathcal{F}_n$, then $B \subseteq \{p + \frac{1}{n}x \mid x \in A\}$.*

4.1 Case-I: When there is a d -set $\mathcal{F}' \subseteq \mathcal{F}$ contained in a bounded region

In this subsection we shall discuss how to get the required infinite sequence when there is a d -set $\mathcal{F}' \subseteq \mathcal{F}$ contained in a bounded region. Note that, as we have defined in Section 2, a d -set \mathcal{B} contained in a set $A \subset \mathbb{R}^d$ means that for all $B \in \mathcal{B}$, we have $B \subseteq A$.

Theorem 4.4. *Let $\mathcal{F}' \subseteq \mathcal{F}$ be a d -set such that there is an $N \in \mathbb{N}$ for which every $B \in \mathcal{F}'$ is contained in $B(O, N)$. Then there is a d -point $p \in \mathbb{R}^d$ with respect to \mathcal{F} .*

Proof. Without loss of generality, we assume that \mathcal{F}' is contained in an axis parallel box

$$A = [a_{1,1}, a_{1,2}] \times \dots \times [a_{d,1}, a_{d,2}].$$

If A contains no d -point, then for each $a \in A$, there is a polytope P_a with $a \in P_a^o$ and P_a^o contains no d -set. As A is compact, there are finitely many points a_1, \dots, a_t such that $A \subseteq \bigcup_{i=1}^t P_{a_i}^o$. Let $\mathcal{F}'' \subseteq \mathcal{F}'$ be the family of all sets in \mathcal{F}' that intersect the boundary of some P_{a_i} , $i \in [t]$. Since the boundary of each P_{a_i} consists of finitely many hyperplanes, \mathcal{F}'' cannot be a d -set. Then $\mathcal{F}' \setminus \mathcal{F}''$ is a d -set such that for every $B \in \mathcal{F}' \setminus \mathcal{F}''$, we have $B \subseteq P_{a_i}^o$, for some $i \in [t]$. But this is a contradiction, since $P_{a_i}^o$ contains no d -set for every $i \in [t]$. Therefore, there must be a d -point in A with respect to \mathcal{F}' , and hence, with respect to \mathcal{F} . \square

Theorem 4.5. Let $\mathcal{F}' \subseteq \mathcal{F}$ be a d -set which is contained in a bounded region and $p \in \mathbb{R}^d$ be a d -point with respect to \mathcal{F}' . Then there exists

(a) a nested sequence of d -sets $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$ such that for all $n \in \mathbb{N}$, $\mathcal{B}_n \subseteq \mathcal{F}$

(b) for every $k \in \{2, \dots, d\}$ a k -flat S_k containing a ray \vec{r}_k whose endpoint is p , and for all $k \in \{2, \dots, d-1\}$, we have $\vec{r}_{k+1} \perp S_k$ and $S_{k+1} \supseteq S_k$

such that for every $n \in \mathbb{N}$ and $B \in \mathcal{B}_n$, we have $\pi_{S_k}(B) \subseteq R(\vec{r}_k, 1/n)$ and $\pi_{S_k}(B) \cap \vec{r}_k = \emptyset$ for all $k \in \{2, \dots, d\}$.

Proof. Without loss of generality, let p be the origin O . We prove Theorem 4.5 with the help of the following claims:

1. There is a nested sequence $\{\mathcal{F}_{n,d}\}_{n \in \mathbb{N}}$, $\mathcal{F}_{n,d} \subseteq \mathcal{F}'$ for all $n \in \mathbb{N}$, with the property that given any $n \in \mathbb{N}$, if $B \in \mathcal{F}_{n,d}$, then $B \subseteq B(O, 1/n)$.
2. Let $k \in \{2, \dots, d\}$ and S_k be a k -flat passing through O . Let there be a nested sequence $\{\mathcal{F}_{n,k}\}_{n \in \mathbb{N}}$ of d -sets, $\mathcal{F}_{n,k} \subseteq \mathcal{F}'$ for all $n \in \mathbb{N}$, such that for all $n \in \mathbb{N}$, if $B \in \mathcal{F}_{n,k}$, then $\pi_{S_k}(B) \subseteq B(O, 1/n) \cap S_k$. Then there is a ray \vec{r}_k in S_k with endpoint at O and a nested sequence $\{\mathcal{B}_{n,k}\}_{n \in \mathbb{N}}$ of d -sets such that $B_{n,k} \subseteq \mathcal{F}_{n,k} \forall n \in \mathbb{N}$, and $\forall B \in \mathcal{B}_{n,k}$, we have $\pi_{S_k}(B) \subseteq R(\vec{r}_k, 1/n)$ and $\pi_{S_k}(B) \cap \vec{r}_k = \emptyset$.

Claim 1 is a simple consequence of Observation 4.3. We will prove Claim 2 in Lemma 4.6. For now, we will prove Theorem 4.5 assuming that Lemma 4.6 is true. Putting $k = d$ in Lemma 4.6, we get a nested sequence $\{\mathcal{B}_{n,d}\}_{n \in \mathbb{N}}$ of d -sets and a ray \vec{r}_d with endpoint at O such that for all $n \in \mathbb{N}$, $\mathcal{B}_{n,d} \subseteq \mathcal{F}_{n,d}$ and $\forall B \in \mathcal{B}_{n,d}$, we have $B \subseteq R(\vec{r}_d, 1/n)$ and $B \cap \vec{r}_d = \emptyset$.

Let S_{d-1} be the $(d-1)$ -flat that passes through O and is perpendicular to \vec{r}_d . Now note that for all $n \in \mathbb{N}$ and $\forall B \in \mathcal{B}_{n,d}$, we have $\pi_{S_{d-1}}(B) \subseteq B(O, 1/n)$. Therefore, Lemma 4.6 applies and we get a ray \vec{r}_{d-1} in S_{d-1} and a nested sequence $\{\mathcal{B}_{n,d-1}\}_{n \in \mathbb{N}}$ of d -sets such that for all $n \in \mathbb{N}$ and $\forall B \in \mathcal{B}_{n,d-1}$, we have $\pi_{S_{d-1}}(B) \subseteq R(\vec{r}_{d-1}, 1/n)$ and $\pi_{S_{d-1}}(B) \cap \vec{r}_{d-1} = \emptyset$.

Proceeding this way, we obtain

- (a) for every $k \in \{2, \dots, d\}$ a nested sequence $\{\mathcal{B}_{n,k}\}_{n \in \mathbb{N}}$ of d -sets such that for every $k \in \{2, \dots, d-1\}$ and $\forall n \in \mathbb{N}$, we have $\mathcal{F} \supseteq \mathcal{B}_{n,k+1} \supseteq \mathcal{B}_{n,k}$
- (b) for every $k \in \{2, \dots, d\}$, a k -flat S_k containing a ray \vec{r}_k whose endpoint is O , and for all $k \in \{2, \dots, d-1\}$, we have $\vec{r}_{k+1} \perp S_k$, $S_{k+1} \supseteq S_k$

such that for every $n \in \mathbb{N}$ and $\forall B \in \mathcal{B}_{n,k}$, we have $\pi_{S_k}(B) \subseteq R(\vec{r}_k, 1/n)$ and $\pi_{S_k}(B) \cap \vec{r}_k = \emptyset$ for all $k \in \{2, \dots, d\}$. Now since for all $n \in \mathbb{N}$ and for all $k \in \{2, \dots, d\}$, we have $\mathcal{B}_{n,2} \subseteq \mathcal{B}_{n,k}$, it follows that for all $B \in \mathcal{B}_{n,2}$, we have $\pi_{S_k}(B) \cap \vec{r}_k = \emptyset$ and $\pi_{S_k}(B) \subseteq R(\vec{r}_k, 1/n)$ for all $k \in \{2, \dots, d\}$ and for all $n \in \mathbb{N}$. We set, for every $n \in \mathbb{N}$, $\mathcal{B}_n = \mathcal{B}_{n,2}$, which gives us the sequence described in the Theorem. \square

Now we prove Lemma 4.6.

Lemma 4.6. Let $p \in \mathbb{R}^d$, $k \in \{2, \dots, d\}$ and S_k be a k -flat through p . Let there be a nested sequence $\{\mathcal{F}_{n,k}\}_{n \in \mathbb{N}}$ of d -sets such that for all $n \in \mathbb{N}$ and $\forall B \in \mathcal{F}_{n,k}$, we have $\pi_{S_k}(B) \subseteq B(p, 1/n)$. Then there is a ray \vec{r}_k in S_k with endpoint at p and a nested sequence $\{\mathcal{B}_{n,k}\}_{n \in \mathbb{N}}$ of d -sets such that $\mathcal{B}_{n,k} \subseteq \mathcal{F}_{n,k} \forall n \in \mathbb{N}$, and $\forall B \in \mathcal{B}_{n,k}$, we have $\pi_{S_k}(B) \subseteq R(\vec{r}_k, 1/n)$ and $\pi_{S_k}(B) \cap \vec{r}_k = \emptyset$.

Proof. Without loss of generality, let $p = O$ and $S_k = \{x \in \mathbb{R}^d \mid x_i = 0 \forall k < i \leq d\}$. For any set $A \subseteq \mathbb{R}^d$, we shall denote its projection onto S_k by $\pi_k(A)$. For any $n \in \mathbb{N}$, we can get a finite sequence $v_{n_1}, \dots, v_{n_{t_n}} \in S_k \cap \mathbb{S}^{d-1}$ such that

$$S_k \cap \mathbb{S}^{d-1} \subseteq \bigcup_{j=1}^{t_n} \left(\text{box} \left(v_{n_j}, \frac{1}{d^2 n^2} \right) \right)^o.$$

If $A \in \mathcal{F}_{n,k}$, then $\pi_k(A) \subseteq B(O, \frac{1}{n})$. Every $CH(\{O\}, \text{box}(\frac{1}{n}v_{n_j}, \frac{1}{d^2n^3}))$ is a polytope, so by Observation 4.2, for every $n \in \mathbb{N}$, the collection of sets in $\mathcal{F}_{n,k}$ whose projections intersect the boundaries of

$$S_k \cap CH\left(\{O\}, \text{box}\left(\frac{1}{n}v_{n_j}, \frac{1}{d^2n^3}\right)\right),$$

where $j \in [t_n]$, is not a d -set. Also, there must be at least one v_{n_j} such that the interior of $CH(\{O\}, \text{box}(\frac{1}{n}v_{n_j}, \frac{1}{d^2n^3})) \cap S_k$ contains the projection of a d -set that is a subset of $\mathcal{F}_{n^2,k}$, since

$$S_k \cap B(O, 1/n) \subseteq \bigcup_{j=1}^{t_n} CH\left(\{O\}, \text{box}\left(\frac{1}{n}v_{n_j}, \frac{1}{d^2n^3}\right)\right) \cap S_k.$$

Let

$$V_n = \bigcup_{j \in J_n} \left(S^{d-1} \cap S_k \cap \text{box}\left(v_{n_j}, \frac{1}{d^2n^2}\right) \right),$$

where $J_n \subseteq \{n_1, \dots, n_{t_n}\}$, such that $CH(\{O\}, \text{box}(\frac{1}{n}v_{n_j}, \frac{1}{d^2n^3}))$ contains the projection of a d -set in its interior that is a subset of $\mathcal{F}_{n^2,k}$ if and only if $j \in J_n$. V_n is compact for each $n \in \mathbb{N}$ and since $\mathcal{F}_{n,k} \supseteq \mathcal{F}_{n+1,k}$ for all $n \in \mathbb{N}$, we have $\bigcap_{i=1}^n V_i \neq \emptyset$ for all $n \in \mathbb{N}$. Therefore, we have $\bigcap_{n \in \mathbb{N}} V_n \neq \emptyset$ and there is a $v \in \bigcap_{n \in \mathbb{N}} V_n$.

Define $\vec{r}_k = \{tv \mid t \in \mathbb{R}^+ \cup \{0\}\}$. Then, due to the way we have chosen v , for every $n \in \mathbb{N}$ there exists a $v_n \in S^{d-1} \cap S_k$ such that $v \in \text{box}(v_n, \frac{1}{d^2n^2})$, and $CH(\{O\}, \text{box}(\frac{1}{n}v_n, \frac{1}{d^2n^3})) \cap S_k$ contains the projection of a d -set $\mathcal{B}'_{n,k} \subseteq \mathcal{F}_{n^2,k}$ onto S_k . Observe that as

$$CH(\{O\}, \text{box}(v_n, d^{-2}n^{-2})) \cap \mathcal{B}(O, 1/n^2) \cap S_k \subseteq R(\vec{r}_k, 1/n) \cap S_k,$$

therefore for every $B \in \mathcal{B}'_{n,k}$ we have

$$\pi_k(B) \subseteq R(\vec{r}_k, 1/n) \cap S_k.$$

For all $n \in \mathbb{N}$, let

$$\mathcal{B}''_{n,k} = \{B \in \mathcal{B}'_{n,k} \mid \pi_k(B) \cap \vec{r}_k = \emptyset\}.$$

Clearly, for all $n \in \mathbb{N}$, $\mathcal{B}''_{n,k}$ is a d -set. Set

$$\mathcal{B}_{n,k} = \bigcup_{i=n}^{\infty} \mathcal{B}''_{i,k}.$$

This is the required nested sequence. □

We shall now prove a few lemmas which would establish some properties that will be used to construct our required sequence. From Theorem 4.5, we have obtained a sequence $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$ of d -sets, and for each $k \in \{2, \dots, d\}$, a k -flat S_k and a ray \vec{r}_k .

We assume, without loss of generality, that for every $k \in [d]$, $S_k = \text{Span}(e_1, \dots, e_k)$ and $\vec{r}_k = \{Re_k \mid R > 0\}$, where $e_k = (x_1, \dots, x_n)$ with $x_i = 0$ if and only if $i \neq k$, and $x_k = 1$.

Lemma 4.7. *Let B_1, B_2, \dots, B_m be such that $B_i \in \mathcal{B}_1$ and $m \in \mathbb{N}$. Then $\exists t_2 \in \mathbb{N}$, $t_2 > m$ such that $R(\vec{r}_2, 1/t_2) \cap \pi_2(B_i) = \emptyset \forall i \in [m]$.*

Proof. Since $\vec{r}_2 \cap \pi_2(B_i) = \emptyset$ for all $i \in [m]$ by our construction of \mathcal{B}_i 's, we have our $t_2 \in \mathbb{N}$ that satisfies $R(\vec{r}_2, 1/t_2) \cap \pi_2(B_i) = \emptyset$ for all $i \in [m]$. □

Lemma 4.8. *Let $B \in \mathcal{B}_n$ where $n \in \mathbb{N}$. Then $\vec{r}_2 \cap C_2(\{O\}, \pi_2(B)) = \{O\}$.*

Proof. If $x \in \vec{r}_2 \cap C_2(\{O\}, \pi_2(B))$ and $x \neq O$, then \vec{r}_2 must intersect $\pi_2(B)$, but that is a contradiction. □

Lemma 4.9. Let $k \in \{2, \dots, d-1\}$ and B_1, \dots, B_m be such that for all $i \in [m]$ we have $B_i \in \mathcal{B}_1$ and $O \notin C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k}))$, where i_j 's are distinct integers in $[m]$, $j \in [k]$. Then we have

$$\vec{r}_{k+1} \cap C_{k+1}(\pi_{k+1}(B_{i_1}), \dots, \pi_{k+1}(B_{i_k}), \{O\}) = \{O\}.$$

Proof. If not, then there is an $x \in \vec{r}_{k+1} \setminus \{O\}$ such that

$$x \in C_{k+1}(\pi_{k+1}(B_{i_1}), \dots, \pi_{k+1}(B_{i_k}), \{O\}).$$

This implies that there is a k -flat that contains \vec{r}_{k+1} and passes through each $\pi_{k+1}(B_{i_j})$. Therefore, this k -flat is perpendicular to S_k and hence its projection onto S_k is a $(k-1)$ -flat that passes through O and $\pi_k(B_{i_j})$ for every $j \in [k]$. But this is a contradiction, since $O \notin C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k}))$. \square

Lemma 4.10. Let $k \in \{2, \dots, d\}$ and B_1, \dots, B_m be such that $B_i \in \mathcal{B}_1$ for all $i \in [m]$ and

$$\vec{r}_k \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \{O\}.$$

Then there exists $t_k \in \mathbb{N}$, with $t_k > m$, such that

$$R(\vec{r}_k, 1/t_k) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \{O\}.$$

Proof. Since $C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\})$ is closed, there is a $t_k \in \mathbb{N}$, $t_k > m$, for which

$$B(e_k, 1/t_k) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \emptyset.$$

Hence the lemma follows. \square

4.1.1 Construction of the sequence

Now we are ready to construct the sequence $\{B_n\}_{n \in \mathbb{N}} \subseteq \mathcal{F}$ so that no $d+1$ distinct sets of the sequence are intersected by a hyperplane. Let $\{B_n\}_{n \in \mathbb{N}}$, S_k and \vec{r}_k for all $k \in \{2, \dots, d\}$ be as defined in Theorem 4.5, and assume, as before, that for every $k \in [d]$, $S_k = \text{Span}(e_1, \dots, e_k)$ and $\vec{r}_k = \{Re_k \mid R > 0\}$, where $e_k = (x_1, \dots, x_n)$ with $x_i = 0$ if and only if $i \neq k$, and $x_k = 1$. Pick $B_1 \in \mathcal{B}_1$. Then there is a $t_2 \in \mathbb{N}$, $t_2 > 1$, for which

$$C_2(\{O\}, \pi_2(B_1)) \cap R(\vec{r}_2, 1/t_2) = \{O\},$$

following Lemma 4.8 and Lemma 4.10. Pick $B_2 \in \mathcal{B}_{t_2}$. Then $O \notin C_2(\pi_2(B_1), \pi_2(B_2))$, since $\pi_2(B_2) \subseteq R(\vec{r}_2, 1/t_2)$ and $O \notin \pi_2(B_2)$. By Lemma 4.9, we have

$$\vec{r}_3 \cap C_3(\pi_3(B_1), \pi_3(B_2), \{O\}) = \{O\}.$$

Therefore, by Lemma 4.10, we have a $t_3 > 2$ for which

$$R(\vec{r}_3, 1/t_3) \cap C_3(\pi_3(B_1), \pi_3(B_2), \{O\}) = \{O\}.$$

Also, since $O \notin C_2(\pi_2(B_1), \pi_2(B_2))$ and $\vec{r}_2 \cap C_2(\pi_2(B_i), \{O\}) = \{O\}$ for $i = 1, 2$, we have a $t'_2 \in \mathbb{N}$ for which

$$R(\vec{r}_2, 1/t'_2) \cap C_2(\pi_2(B_i), \{O\}) = \{O\}$$

and

$$R(\vec{r}_2, 1/t'_2) \cap C_2(\pi_2(B_1), \pi_2(B_2)) = \emptyset.$$

We choose B_3 from \mathcal{B}_t , where $t = \max\{t'_2, t_3\}$. Now observe that

$$\pi_2(B_3) \cap C_2(\pi_2(B_1), \pi_2(B_2)) = \emptyset.$$

Proceeding this way, we construct our sequence.

In general, let us have the sequence B_1, B_2, \dots, B_m where $B_i \in \mathcal{B}_i$ that we have constructed as above. So, for all $k \in \{2, \dots, \min\{m, d\}\}$, we have

- $\vec{r}_k \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \{O\}$,
- $O \notin C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k}))$, and

where i_1, \dots, i_k denote distinct integers in $[m]$. Further, for all $k \in \{2, 3, \dots, \min\{m-1, d\}\}$, we have $\pi_k(B_{i_{k+1}}) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset$, where each i_j is distinct and chosen from $[m]$. We have shown that we can obtain such a sequence for $m = 3$. We shall extend this sequence now. Using Lemma 4.9 and 4.10, we can obtain a $t \in \mathbb{N}$ such that for every $k \in \{2, \dots, \min\{m+1, d\}\}$, we have

$$R(\vec{r}_k, 1/t) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \{O\},$$

and for every $k \in \{2, \dots, \min\{m, d\}\}$, we have

$$R(\vec{r}_k, 1/t) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset$$

for distinct i_j 's. Now, we can easily check that for any $B_{m+1} \in \mathcal{B}_t$ and all $k \in \{2, \dots, \min\{m+1, d\}\}$, we have

$$\pi_k(B_{m+1}) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset$$

for distinct i_j 's in $[m]$. Further, for all $k \in \{2, \dots, \min\{m+1, d\}\}$, we have

- $O \notin C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k}))$,
- $\vec{r}_k \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \{O\}$ and
- $\pi_k(B_{i_k}) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \emptyset$,

where each i_j is distinct and chosen from $[m+1]$. So, for all $k \in \{2, \dots, \min\{m+1, d\}\}$, we have

- $\vec{r}_k \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \{O\}$,
- $O \notin C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k}))$ and
- $\pi_k(B_{i_k}) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \emptyset$,

where i_1, \dots, i_k denote distinct integers in $[m+1]$. For all $k \in \{2, 3, \dots, \min\{m, d\}\}$, we have

$$\pi_k(B_{i_{k+1}}) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset$$

where each i_j is distinct and chosen from $[m+1]$. Therefore, we can extend our sequence. This concludes the proof of **Case-I**.

4.2 Case-II: when there are no d -sets in a bounded region that is a subset of \mathcal{F}

Next we discuss how to get the required infinite sequence when there are no d -sets in a bounded region that is a subset of \mathcal{F} . Before that we will need the following technical lemmas.

Lemma 4.11. *Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a nested sequence of d -sets and S_k be a k -flat with $p' \in S_k$. Then exactly one of the following holds:*

- For every $n \in \mathbb{N}$, $S_k \setminus B(p', n)$ contains the projection of a d -set $\mathcal{B}'_{n,k} \subseteq \mathcal{F}_n$. Then there is a ray \vec{r}_k on S_k such that every $Q(\vec{r}_k, n) \cap S_k$ contains the projection of a d -set $\mathcal{B}_{n,k} \subseteq \mathcal{F}_n$ such that $\mathcal{B}_{n,k} \supseteq \mathcal{B}_{n+1,k}$, for all $n \in \mathbb{N}$.
- There is an $N \in \mathbb{N}$ and an $n' \in \mathbb{N}$ for which the subset of $\mathcal{F}_{n'}$ consisting of all sets whose projections onto S_k are not contained in $S_k \cap B(p', N)$ is not a d -set. In this case, there is a point $p \in S_k$ such that there is a ray \vec{r}_k on S_k with endpoint at p , and for every $n \in \mathbb{N}$, $R(\vec{r}_k, 1/n) \cap S_k$ contains the projection of a d -set $\mathcal{B}_{n,k} \subseteq \mathcal{F}_n$ such that $\mathcal{B}_{n,k} \supseteq \mathcal{B}_{n+1,k}$, for all $n \in \mathbb{N}$.

Lemma 4.12. Let \vec{r}' be a ray in \mathbb{R}^d and let \vec{r} be a ray in \mathbb{R}^d such that there is a $p \in \mathbb{R}^d$ for which $\vec{r}' = p + \vec{r}$. Then there exists a strictly increasing sequence $\{t_n\}_{n \in \mathbb{N}} \subseteq \mathbb{N}$ for which $Q(\vec{r}', t_n) \subseteq Q(\vec{r}, n)$.

Lemma 4.13. Let there be a $k' \in \{2, \dots, d\}$ for which there exists a nested sequence of d -sets $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ and a point $p \in \mathbb{R}^d$ such that for every $k \in \{k', \dots, d\}$ there is a k -flat S_k containing a ray \vec{r}_k with endpoint at p satisfying the property that

- $\vec{r}_{k+1} \perp S_k$ and $S_{k+1} \supsetneq S_k$ when $k < d$
- $\forall n \in \mathbb{N}$ and $\forall B \in \mathcal{F}_n$, we have $\pi_{S_k}(B) \subseteq Q(\vec{r}_k, n) \cap S_k$

Pick any $p' \in S_{k'}$ and for every $k \in \{k', \dots, d\}$, let \vec{r}'_k denote the ray \vec{r}_k translated so that its endpoint is at p' . Then there is a strictly increasing sequence $\{t_n\}_{n \in \mathbb{N}} \subseteq \mathbb{N}$ with the property that

- $\vec{r}'_{k+1} \perp S_k$ and $S_{k+1} \supsetneq S_k$ when $k < d$
- $\forall n \in \mathbb{N}$ and $\forall B \in \mathcal{F}_{t_n}$, we have $\pi_{S_k}(B) \subseteq Q(\vec{r}'_k, n)$

Lemma 4.14. Let $\{\mathcal{F}_n\}_{n \in \mathbb{N}}$ be a nested sequence of d -sets and S_2 be a 2-flat. Further, let, for all $p \in S_2$, $\forall n \in \mathbb{N}$ and for every $R > 0$, $\mathcal{F}'_n := \{B \in \mathcal{F}_n \mid \pi_{S_2}(B) \cap B(p, R) = \emptyset\}$ be a d -set. Then we have a nested sequence $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$ of d -sets with $\mathcal{B}_n \subseteq \mathcal{F}$ $\forall n \in \mathbb{N}$ and a ray \vec{r}_2 on S_2 such that for every $n \in \mathbb{N}$ and $B \in \mathcal{B}_n$, we have $\pi_{S_2}(B) \subseteq Q(\vec{r}_2, n) \cap S_2$ and $\pi_{S_2}(B) \cap \vec{r}_2 = \emptyset$. Moreover, we can choose the \mathcal{B}_n 's in such a way that for any $n \in \mathbb{N}$ and any $B_1, \dots, B_m \in \mathcal{B}_1$, we can choose a $B' \in \mathcal{B}_n$ for which $\pi_{S_1}(B_j) \cap \pi_{S_1}(B') = \emptyset$ for all $i \in [m]$, where S_1 is the straight line that passes through the endpoint of \vec{r}_2 and is perpendicular to \vec{r}_2 .

The proofs of the above four lemmas can be found in Appendix A.

Theorem 4.15. Suppose \mathcal{F} is a family of d -sets in \mathbb{R}^d such that there are no d -sets in \mathcal{F} contained in a bounded region. Then there is a $k' \in \{2, \dots, d\}$ and there exists a nested sequence of d -sets $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$ with $\mathcal{B}_n \subseteq \mathcal{F}$ for all $n \in \mathbb{N}$ and a point $p \in \mathbb{R}^d$ such that for every $k \in \{2, \dots, d\}$ there is a k -flat S_k containing a ray \vec{r}_k with endpoint at p and when $k < d$, we have $\vec{r}_{k+1} \perp S_k$ and $S_{k+1} \supsetneq S_k$. Moreover,

- (a) for all $k \in \{k', \dots, d\}$ and $\forall n \in \mathbb{N}$, $\forall B \in \mathcal{B}_n$, we have $\pi_{S_k}(B) \subseteq Q(\vec{r}_k, n)$ and $\pi_{S_k}(B) \cap \vec{r}_k = \emptyset$, and
- (b) for all $k \in \{2, \dots, k' - 1\}$ and $\forall n \in \mathbb{N}$, $\forall B \in \mathcal{B}_n$, we have $\pi_{S_k}(B) \subseteq R(\vec{r}_k, \frac{1}{n})$ and $\pi_{S_k}(B) \cap \vec{r}_k = \emptyset$. Note that $\{2, \dots, k' - 1\}$ is the empty set if $k' = 2$.

Further, if $k' = 2$, then we can choose $\{\mathcal{B}_n\}_{n \in \mathbb{N}}$ so that for any $B_1, \dots, B_m \in \mathcal{B}_1$, for any $n \in \mathbb{N}$ we can pick a $B' \in \mathcal{B}_n$ such that if S_1 denotes the straight line on S_2 that passes through p and is perpendicular to \vec{r}_2 , then $\pi_{S_1}(B') \cap \pi_{S_1}(B_i) = \emptyset$ for all $i \in [m]$.

Proof. $\mathcal{F}_{n,d} = \{A \in \mathcal{F} \mid A \cap B(O, n) = \emptyset\}$ is a d -set for all $n \in \mathbb{N}$. Applying Lemma 4.11(a), we obtain a nested sequence $\{\mathcal{B}_{n,d}\}_{n \in \mathbb{N}}$ of d -sets such that for all $n \in \mathbb{N}$, $\mathcal{B}_{n,d} \subseteq \mathcal{F}_{n,d}$. Also, we get a ray \vec{r}_d with endpoint at O in \mathbb{R}^d such that for every $n \in \mathbb{N}$, $\forall B \in \mathcal{B}_{n,d}$, we have $B \in Q(\vec{r}_d, n)$.

Let S_{d-1} be the $(d-1)$ -flat that is perpendicular to \vec{r}_d and passes through O . We apply Lemma 4.11 for this $(d-1)$ -flat. Either (a) or (b) of Lemma 4.11 applies. If (a) applies, then we would obtain a ray \vec{r}'_{d-1} with endpoint at O and a nested sequence $\{\mathcal{B}_{n,d-1}\}_{n \in \mathbb{N}}$ of d -sets such that $\forall n \in \mathbb{N}$, we have $\mathcal{B}_{n,d-1} \subseteq \mathcal{B}_{n,d}$ and $\forall B \in \mathcal{B}_{n,d-1}$, $\pi_{S_{d-1}}(B) \subseteq Q(\vec{r}'_{d-1}, n)$. If (a) does not apply, then (b) applies. As we move from S_d to S_{d-1} and so on as above, let k' be the smallest integer in $\{d, d-1, \dots, 2\}$ for which Case (a) of Lemma 4.11 applies to $\{d, d-1, \dots, k'\}$.

If $k' = 2$, then we apply Lemma 4.14 and we are done. If $k' > 2$, then we obtain, for each $k \in \{k', \dots, d\}$, a k -flat S_k containing O and a ray \vec{r}'_k in S_k with endpoint at O , and a sequence

$\{\mathcal{B}_{n,k'}\}_{n \in \mathbb{N}}$ of d -sets with $\mathcal{B}_{n,k'} \subsetneq \mathcal{F}_{n,d}$ and $\forall B \in \mathcal{B}_{n,k'}, \pi_{S_k}(B) \subsetneq Q(\vec{r}_k, n)$ for all $n \in \mathbb{N}$. Now, let $S_{k'-1}$ denote the $(k'-1)$ -flat contained in $S_{k'}$, passing through O and perpendicular to $\vec{r}_{k'}$. Now, Case (b) of Lemma 4.11 applies, and we obtain a point p on $S_{k'-1}$, a ray $\vec{r}_{k'-1}$ in $S_{k'-1}$ from p and a nested sequence $\{\mathcal{B}_{n,k'-1}\}_{n \in \mathbb{N}}$ of d -sets such that $\mathcal{B}_{n,k'-1} \subseteq \mathcal{B}_{n,k'}$ and $\forall B \in \mathcal{B}_{n,k'-1}$, we have $\pi_{S_{k'-1}}(B) \subseteq R(\vec{r}_{k'-1}, \frac{1}{n})$ for all $n \in \mathbb{N}$. Now let for each $k \in \{k', \dots, d\}$, $\vec{r}_k = p + \vec{r}_{k'}$. Then, using Lemma 4.13, we get a strictly increasing sequence $\{t_n\}_{n \in \mathbb{N}}$ for which we have, $\forall n \in \mathbb{N}$ and $\forall B \in \mathcal{B}_{t_n,k'}$, $\pi_{S_k}(B) \subsetneq Q(\vec{r}_k, n)$. Since for every $n \in \mathbb{N}$ and any $k \in \{k', \dots, d\}$, the collection of sets in $\mathcal{B}_{t_n,k'}$ whose projections onto S_k intersect \vec{r}_k cannot be a d -set due to Observation 4.2, we can assume, without loss of generality, that for every $n \in \mathbb{N}$ and $\forall B \in \mathcal{B}_{t_n,k'}$, we have $\vec{r}_k \cap \pi_{S_k}(B) = \emptyset$ for all $k \in \{k', \dots, d\}$. For all $n \in \mathbb{N}$, set

$$\mathcal{F}_{n,k'-1} := \{B \in \mathcal{B}_{t_n,k'-1} \mid \pi_{S_k}(B) \cap \vec{r}_k = \emptyset, \forall k \in \{k'-1, \dots, d\}\}.$$

Now, by repeated application of Lemma 4.6, we obtain, for all $k \in \{2, \dots, k'-1\}$, a k -flat S_k , a ray \vec{r}_k from p and a nested sequence $\{B_n\}$ of d -sets with $B_n \subseteq \mathcal{F}_{n,k'-1}$ such that for $k < k'-1$, $S_k \perp \vec{r}_{k+1}$ and $S_k \subsetneq S_{k+1}$, and $\forall B \in B_n$, $\vec{r}_k \cap \pi_{S_k}(B) = \emptyset$ and $\pi_{S_k}(B) \subseteq R(\vec{r}_k, 1/n)$. This concludes the proof. \square

Again we shall prove a few lemmas before we proceed to construct the sequence. Let for every $k \in \{1, \dots, d\}$, $e_k \in \mathbb{R}^d$ denote the point $e_k = (x_1, \dots, x_d)$ where $x_i = 0$ for all $i \in [d] \setminus \{k\}$ and $x_k = 1$. Without loss of generality, we assume that p is the origin O and $\forall k \in \{2, \dots, d\}$, $S_k = \text{Span}(e_1, \dots, e_k)$ and $\vec{r}_k = \{Re_k \mid R > 0\}$.

Lemma 4.16. *Let B_1, \dots, B_m with $B_i \in \mathcal{B}_1 \forall i \in [m]$ and $k \in \{1, \dots, d-1\}$ be such that we have*

$$C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) \cap \pi_k(B_{i_{k+1}}) = \emptyset$$

for distinct i_j 's in $[m]$. Then $C_{k+1}(\pi_{k+1}(B_{i_1}), \dots, \pi_{k+1}(B_{i_{k+1}}))$ does not contain any straight lines parallel to \vec{r}_{k+1} .

Proof. It suffices to show that for any choice of $a_j \in \pi_{k+1}(B_{i_j})$, for each $j \in \{1, 2, \dots, k+1\}$, $C_{k+1}(\{a_1\}, \dots, \{a_{k+1}\})$ is a single k -flat which does not contain a straight line parallel to \vec{r}_{k+1} . Now if there is no unique k -flat in S_{k+1} that passes through a_1, \dots, a_{k+1} , then there is a λ -flat L that passes through a_1, \dots, a_{k+1} , where $\lambda < k$. Then the projection of L onto S_k passes through $\pi_k(a_1), \dots, \pi_k(a_{k+1})$. But this contradicts the assumption that

$$C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) \cap \pi_k(B_{i_{k+1}}) = \emptyset.$$

Therefore, there must be a unique k -flat \mathcal{K} that passes through a_1, \dots, a_{k+1} . Now if \mathcal{K} contains a straight line parallel to \vec{r}_{k+1} , then the projection of \mathcal{K} onto S_k is a $(k-1)$ -flat, which contradicts the assumption that

$$C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) \cap \pi_k(B_{i_{k+1}}) = \emptyset.$$

\square

Lemma 4.17. *Let B_1, \dots, B_m , $B_i \in \mathcal{B}_1 \forall i \in [m]$ and $k \in \{k', \dots, d\}$ be such that we have*

$$C_{k-1}(\pi_{k-1}(B_{i_1}), \dots, \pi_{k-1}(B_{i_{k-1}})) \cap \pi_{k-1}(B_{i_k}) = \emptyset$$

for distinct i_j 's in $[m]$. Then there is a $t_k \in \mathbb{N}$, $t_k > m$ for which we have the following: $\forall n \geq t_k$ and distinct $i_1, \dots, i_k \in [m]$, we have

$$Q(\vec{r}_k, n) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset.$$

Proof. For distinct B_{i_1}, \dots, B_{i_k} , define

$$f : B_{i_1} \times \dots \times B_{i_k} \rightarrow \mathbb{R}$$

such that $f(b_1, \dots, b_k) = R$ if and only if the $(k-1)$ -flat through b_1, \dots, b_k intersects $\{ce_k \mid c \in \mathbb{R}\}$ at the point Re_k , where $b_j \in B_{i_j}$ for all $j \in [k]$. Note that we have already shown in Lemma 4.16 that such a $(k-1)$ -flat is unique, and since no such $(k-1)$ -flat contains a straight line parallel to \vec{r}_k , the function f is well-defined. Also, it is easy to see that f is continuous, and as such, its image in \mathbb{R} must be compact. Therefore, we have a $t_k \in \mathbb{N}$, $t_k > m$ for which

$$Q(\vec{r}_k, t_k) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset$$

for every choice of distinct i_1, \dots, i_k . Hence the result follows. \square

4.2.1 Construction of the sequence

Now we are ready to construct the sequence $\{B_n\}_{n \in \mathbb{N}}$ such that $B_n \in \mathcal{B}_n$ for all $n \in \mathbb{N}$ and no single hyperplane can pierce $d+1$ distinct B_n 's. If $k' > 2$, then pick $B_1, \dots, B_{k'}$ the way described in subsection 4.1.1 of Case-I, so that for all $k \in \{2, \dots, k' - 1\}$, they satisfy

$$(\{O\} \cup \pi_k(B_{i_{k+1}})) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset,$$

and

$$(\vec{r}_k \setminus \{O\}) \cap C_k(\{O\}, \pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}})) = \emptyset$$

for all distinct i_j 's in $[k']$.

If $k' = 2$, then choose $B_1 \in \mathcal{B}_1$. Due to our choice of \vec{r}_2 as described in Theorem 4.15, there is a $B_2 \in \mathcal{B}_2$ for which $\pi_1(B_2) \cap \pi_1(B_1) = \emptyset$.

Now we have the sequence $B_1, \dots, B_{k'}$. We show how to extend a sequence of length at least k' . Let B_1, \dots, B_m , $m \geq k'$ be a sequence such that for all $k \in \{2, \dots, k' - 1\}$, we have

$$\begin{aligned} & [\{O\} \cup \pi_k(B_{i_{k+1}})] \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset \text{ and} \\ & (\vec{r}_k \setminus \{O\}) \cap C_k(\{O\}, \pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}})) = \emptyset. \end{aligned}$$

for distinct i_j 's in $[m]$. Also, for all $k \in \{k', \dots, \min\{d, m-1\}\}$, we have

$$\pi_k(B_{i_{k+1}}) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset$$

for distinct i_j 's in $[m]$. Then, if $k' > 2$, we can obtain a $t' \in \mathbb{N}$, $t' > m$ for which

$$\begin{aligned} & R(\vec{r}_k, 1/t') \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}}), \{O\}) = \{O\} \text{ and} \\ & R(\vec{r}_k, 1/t') \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset \end{aligned}$$

for distinct i_j 's in $[m]$ for all $k \in \{2, \dots, k' - 1\}$, using Lemma 4.10. For $k \in \{k', \dots, \min\{d, m\}\}$, we use Lemma 4.17 to obtain a $t'' > m$ such that

$$Q(\vec{r}_k, n) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset$$

for distinct i_j 's in $[m]$ whenever $n \geq t''$.

If $k' = 2$, then by Theorem 4.15, we can pick a suitable B_{m+1} from $B_{t''}$ such that $\pi_1(B_{m+1}) \cap \pi_1(B_i) = \emptyset$ for any $i \in [m]$. Therefore, by repeated application of Lemma 4.17, we get that $\forall k \in \{k', \dots, \min\{m, d\}\}$, we have

$$\pi_k(B_{i_{k+1}}) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset$$

for distinct i_j 's in $[m+1]$.

If $k' > 2$, then we set $t = \max\{t', t''\}$ and pick B_{m+1} from \mathcal{B}_t . Then for every $k \in \{2, \dots, k'-1\}$, we have

$$\begin{aligned} [\{O\} \cup \pi_k(B_{i_{k+1}})] \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) &= \emptyset \text{ and} \\ (\vec{r}_k \setminus \{O\}) \cap C_k(\{O\}, \pi_k(B_{i_1}), \dots, \pi_k(B_{i_{k-1}})) &= \emptyset, \end{aligned}$$

for distinct i_j 's in $[m+1]$. At this point, by repeatedly applying Lemma 4.17, we get that $\forall k \in \{k', \dots, \min\{m, d\}\}$, we have

$$\pi_k(B_{i_{k+1}}) \cap C_k(\pi_k(B_{i_1}), \dots, \pi_k(B_{i_k})) = \emptyset$$

for distinct i_j 's in $[m+1]$.

A Missing proofs

Proof of Lemma 4.11. Without loss of generality, let us assume that $p' = O$.

(a) For every $n \in \mathbb{N}$, set

$$\mathcal{F}_{n,k} := \bigcup_{i=n}^{\infty} \mathcal{B}'_{i,k}.$$

If $A \in \mathcal{F}_{n,k}$, then we have $\pi_{S_k}(A) \subseteq S_k \setminus B(O, n)$. For any $n \in \mathbb{N}$, there are finitely many points $v_{n_1}, \dots, v_{n_{t_n}} \in S_k \cap \mathbb{S}^{d-1}$ such that

$$\bigcup_{i=1}^{t_n} \text{box}^o \left(v_{n_i}, \frac{1}{d^2 n^2} \right) \supseteq S_k \cap \mathbb{S}^{d-1}.$$

Then, following similar reasoning as we used in the proof of Lemma 4.3, we can say that there is at least one $i \in \{1, \dots, t_n\}$ such that there is a projection of a d -set that is a subset of $\mathcal{F}_{n^2,k}$ in the interior of

$$S_k \cap \left(\bigcup_{t \geq 0} \text{box} \left(tv_{n_i}, \frac{t}{d^2 n^2} \right) \setminus B(O, n) \right).$$

Let $J_n \subseteq \{1, \dots, t_n\}$ be such that there is a projection of a d -set that is a subset of $\mathcal{F}_{n^2,k}$ in the interior of

$$S_k \cap \left(\bigcup_{t \geq 0} \text{box} \left(tv_{n_i}, \frac{t}{d^2 n^2} \right) \setminus B(O, n) \right)$$

if and only if $i \in J_n$. Let

$$V_n = \bigcup_{i \in J_n} CH \left(\{O\}, \text{box} \left(v_{n_i}, \frac{1}{d^2 n^2} \right) \right) \cap S^{d-1} \cap S_k.$$

Since $\bigcap_{i=1}^n V_i \neq \emptyset$ for all $n \in \mathbb{N}$, we have a $v_k \in \bigcap_{i=1}^{\infty} V_i$. Set $\vec{r}_k = \{tv_k \mid t \geq 0\}$. Then for every $n \in \mathbb{N}$, $Q(\vec{r}_k, n)$ contains the projection of a d -set $\mathcal{F}'_{n,k} \subseteq \mathcal{F}_{n^2,k}$. Setting

$$\mathcal{B}_{n,k} = \bigcup_{i=n}^{\infty} \mathcal{F}'_{i,k},$$

we get the required nested sequence $\{\mathcal{B}_{n,k}\}_{n \in \mathbb{N}}$ of d -sets satisfying the conditions in (a).

(b) Let, for all $n \in \mathbb{N}$,

$$\mathcal{F}'_n = \{A \in \mathcal{F}_n \mid \pi_{S_k}(A) \subseteq B(O, N)\}.$$

Then for all $n \in \mathbb{N}$, \mathcal{F}'_n is a d -set. We claim that there is a point p in $S_k \cap B(O, N)$ such that if P is a polytope with $p \in P^\circ$ then $S_k \cap P^\circ$ also contains the projection of a d -set which is a subset of \mathcal{F}'_n , for all $n \in \mathbb{N}$. If not, then for every $x \in S_k \cap B(O, N)$, there is a polytope P_x on S_k with $x \in P_x^\circ$ and $n_x \in \mathbb{N}$ such that the collection of all sets in \mathcal{F}'_{n_x} whose projections lie in the interior of P_x does not form a d -set. Since $S_k \cap B(O, N)$ is compact, there are finitely many $x_1, \dots, x_s \in S_k \cap B(O, N)$ such that

$$S_k \cap B(O, N) \subseteq \bigcup_{i=1}^s P_{x_i}^\circ.$$

Let $a = \max\{n_{x_1}, \dots, n_{x_s}\}$. The projection of any set in \mathcal{F}'_a on S_k lies either in some P_{x_i} or hits the boundary of some P_{x_i} , $i \in \{1, \dots, s\}$. We can ignore the sets whose projections on S_k hit the boundary of some P_{x_i} because those sets are already hit by finitely many hyperplanes, namely the hyperplanes H_1, \dots, H_t that contain the boundaries of the P_{x_i} 's and $\forall i \in [t]$, $H_i \cap S_k = \pi_{S_k}(H_i)$. This means that the sets whose projections are in the interior of the P_{x_i} 's form a d -set. But this is a contradiction. Therefore, there must be a point p such that for every polytope P with $p \in P^\circ$ and every $n \in \mathbb{N}$, all sets in \mathcal{F}'_n such that their projections lie within $P \cap S_k$ form a d -set. Then by Lemma 4.6, there exists a ray \vec{r}_k on S_k with endpoint at p and a nested sequence $\{\mathcal{B}_{n,k}\}_{n \in \mathbb{N}}$ of d -sets such that for every $n \in \mathbb{N}$, $R(\vec{r}_k, 1/n) \cap S_k$ contains the projection of a d -set $\mathcal{B}_{n,k}$, and we have $\mathcal{B}_{n,k} \subseteq \mathcal{F}_n$. □

Proof of Lemma 4.12. Without loss of generality, let the origin O be the endpoint of \vec{r} and $\vec{r} = \{te_1 \mid t \geq 0\}$. Choose an arbitrary $n \in \mathbb{N}$. There is a point q on \vec{r} such that

$$q \in Q(\vec{r}, n) \setminus [-2n, 2n]^d.$$

So, we have $Q(\vec{r}_q, n) \subseteq Q(\vec{r}, n)$, where $\vec{r}_q = q + \vec{r}$. Now, since there exists a $t_n \in \mathbb{N}$ for which $Q(\vec{r}_q, n) \supseteq Q(\vec{r}', m)$ for all $m \geq t_n$, we have $Q(\vec{r}, n) \supseteq Q(\vec{r}', m)$ for all $m \geq t_n$. □

Proof of Lemma 4.13. Without loss of generality, let for every $k \in \{k', \dots, d\}$,

$$\begin{aligned} S_k &= \text{Span}(e_1, \dots, e_k), \\ \vec{r}_k &= \{te_k \mid t \geq 0\}, \text{ and} \\ p' &= (p_1, \dots, p_{k'}, 0, \dots, 0). \end{aligned}$$

Then for every $k \in \{k', \dots, d\}$, we have $\vec{r}'_k = p' + \vec{r}_k$. We set $S_d := \mathbb{R}^d$. Note that for each $k \in \{k', \dots, d\}$, as S_k is a linear space containing both \vec{r}_k and p' , S_k must contain \vec{r}'_k . When $k' \leq k < d$, suppose S'_k is the k -flat contained in S_{k+1} such that S'_k is perpendicular to \vec{r}'_{k+1} at its endpoint. Then S'_k is parallel to S_k since \vec{r}_k is parallel to \vec{r}'_k . But since $p' \in S'_k \cap S_k$, we must have $S_k = S'_k$, and therefore we have $\vec{r}'_{k+1} \perp S_k$. Now by repeated application of Lemma 4.12, we get an increasing sequence $\{t_n\}_{n \in \mathbb{N}}$ for which we have $\forall n \in \mathbb{N}$ and $\forall B \in \mathcal{F}_{t_n}$, we have

$$\pi_{S_k}(B) \subseteq Q(\vec{r}'_k, n).$$
□

Proof of Lemma 4.14. Exactly one of the two cases below holds:

- (i) either there is a straight line ℓ_p on S_2 and a ray \vec{r}_2 on ℓ_p such that for all $n \in \mathbb{N}$, there is a d -set $\mathcal{B}_n \subseteq \mathcal{F}_n$ such that for every $B \in \mathcal{B}_n$, we have $\pi_{S_2}(B) \subseteq M(\ell_p, 1/n) \cap Q(\vec{r}_2, n)$ and $\pi_{S_2}(B) \cap \ell = \emptyset$,
- (ii) or for no straight line $\ell' \subsetneq S_2$ and $\varepsilon > 0$, there is a sequence of d -sets $\{\mathcal{F}'_n\}_{n \in \mathbb{N}}$ with $\mathcal{F}'_n \subseteq \mathcal{F}_n$ for all $n \in \mathbb{N}$, whose projection on S_2 is contained in the set $M(\ell', \varepsilon)$.

Clearly, if (ii) holds, then (i) does not hold, and by using similar techniques as in Lemma 4.11, we can obtain a sequence $\{B_n\}_{n \in \mathbb{N}}$ and a ray \vec{r}_2 satisfying the properties described in the Lemma. Now, pick any B_1, \dots, B_m from \mathcal{B}_1 . Then there exist straight lines ℓ_1, \dots, ℓ_m in S_2 for which we have for all $i \in [m]$, $\pi_{S_2}(B_i) \subsetneq M(\ell_i, \varepsilon)$ for some $\varepsilon > 0$. Since we can find an $n \in \mathbb{N}$ for which no $M(\ell_i, \varepsilon)$ contains the projection of a d -set that is a subset of \mathcal{F}_n , we can always find a $B \in \mathcal{B}_n$ whose projection onto S_2 does not intersect any $M(\ell_i, \varepsilon)$ (note that the collection of sets in \mathcal{B}_n whose projections onto S_2 intersects the boundaries of $M(\ell_i, \varepsilon)$'s cannot be a d -set). So, we have shown that the lemma holds for (ii).

Now we show that if (ii) does not hold, then (i) holds. So let us assume that (ii) does not hold. Then there is a straight line ℓ on S_2 and $\varepsilon > 0$, and a sequence of d -sets $\{\mathcal{F}''_n\}_{n \in \mathbb{N}}$ with $\mathcal{F}''_n \subseteq \mathcal{F}_n$ for all $n \in \mathbb{N}$, whose projection is contained in the set $M(\ell, \varepsilon)$. We can assume the sequence $\{\mathcal{F}''_n\}_{n \in \mathbb{N}}$ to be nested, because we can easily set, for each $n \in \mathbb{N}$, $\mathcal{B}'_n = \bigcup_{i=n}^{\infty} \mathcal{F}''_i$. Without loss of generality, let $S_2 = \text{span}(e_1, e_2)$, $\ell = \{te_2 \mid t \in \mathbb{R}\}$, and $\varepsilon = 1$. Then we claim that there is a $p = (p_1, 0, \dots, 0) \in \mathbb{R}^d$ such that if ℓ_p is a line on S_2 parallel to ℓ and ℓ_p passes through p , then for every $n \in \mathbb{N}$, $M(\ell_p, 1/n)$ contains the projection of a d -set $\mathcal{B}'_n \subseteq \mathcal{F}''_n$ on S_2 . If not, then for every $x \in [-1, 1] \times \{0\}^{d-1}$, there is an $n_x \in \mathbb{N}$ for which $M\left(\ell_x, \frac{1}{n_x}\right)$ does not contain the projection of a d -set in \mathcal{F}''_{n_x} onto S_2 . Since $[-1, 1]$ is compact, we have finitely many $x_1, \dots, x_t \in [-1, 1] \times \{0\}^{d-1}$ for which

$$M(\ell, 1) \cap S_2 \subseteq \bigcup_{i=1}^t M\left(\ell_{x_i}, \frac{1}{n_{x_i}}\right).$$

Let $N = \max\{n_{x_1}, \dots, n_{x_t}\}$. Then since $\mathcal{F}''_N \supseteq \mathcal{F}''_{N+1}$, we have that the family of the sets of \mathcal{F}''_N whose projections lie in $M\left(\ell_{x_i}, \frac{1}{n_{x_i}}\right)$ for any $i \in [t]$ cannot be a d -set. Further, the collection of sets in \mathcal{F}''_N whose projections onto S_2 intersect the boundaries of $M\left(\ell_{x_i}, \frac{1}{n_{x_i}}\right)$'s cannot form a d -set, as evident from Observation 4.2. But this is a contradiction since \mathcal{F}''_N is a d -set and the projections of all sets in it on S_2 lie in $M(\ell, 1)$. Therefore, (i) holds. Now, S_1 divides S_2 into two half-planes, one of which contains, for every $n \in \mathbb{N}$, the projection of the sets of a d -set that is a subset of \mathcal{F}''_n . Let \vec{r}_2 be the ray on ℓ_p that lies in the half-plane, and let

$$\mathcal{B}_n := \{B \in \mathcal{F}''_n \mid \pi_{S_2}(B) \subsetneq M(\ell_p, 1/n) \text{ and } \pi_{S_2}(B) \cap B(p, n) = \emptyset\}.$$

□

References

- [ADLS17] N. Amenta, J. A. De Loera, and P. Soberón. Helly's Theorem: New Variations and Applications. *Algebraic and Geometric Methods in Discrete Mathematics*, 685:55–95, 2017.
- [AK92] N. Alon and D. J. Kleitman. Piercing Convex Sets and the Hadwiger-Debrunner (p, q) -Problem. *Advances in Mathematics*, 96(1):103–112, 1992.
- [AK95] N. Alon and G. Kalai. Bounding the Piercing Number. *Discrete & Computational Geometry*, 13:245–256, 1995.

- [AKMM02] N. Alon, G. Kalai, J. Matoušek, and R. Meshulam. Transversal numbers for hypergraphs arising in geometry. *Advances in Applied Mathematics*, 29(1):79–101, 2002.
- [BFM⁺14] I. Bárány, F. Fodor, L. P. Montejano, D. Oliveros, and A. Pór. Colourful and Fractional (p, q) -Theorems. *Discrete & Computational Geometry*, 51:628–642, 2014.
- [BK22] I. Bárány and G. Kalai. Helly-type Problems. *Bulletin of the American Mathematical Society*, 59(4):471–502, 2022.
- [CGN23a] S. Chakraborty, A. Ghosh, and S. Nandi. Heterochromatic Geometric Transversals for Convex Sets. *CoRR*, abs/2212.14091, 2023.
- [CGN23b] S. Chakraborty, A. Ghosh, and S. Nandi. Stabbing boxes with finitely many axis-parallel lines and flats. *CoRR*, abs/2308.10479, 2023.
- [HD57] H. Hadwiger and H. Debrunner. Über eine Variante zum Hellyschen Satz. *Archiv der Mathematik*, 8(4):309–313, 1957.
- [JP23] A. Jung and D. Pálvölgyi. k -dimensional transversals for balls. *CoRR*, abs/2311.15646, 2023.
- [KP22] C. Keller and M. A. Perles. An $(\aleph_0, k+2)$ -Theorem for k -Transversals. In *Proceedings of the 38th International Symposium on Computational Geometry, SoCG*, volume 224, pages 50:1–50:14, 2022.
- [KP23] C. Keller and M. A. Perles. An $(\aleph_0, k+2)$ -Theorem for k -Transversals. *CoRR*, abs/2306.02181, 2023.