

# KHOVANOV HOMOLOGY AND EXOTIC 4-MANIFOLDS

QIUYU REN AND MICHAEL WILLIS

ABSTRACT. We show that the Khovanov-Rozansky  $\mathfrak{gl}_2$  skein lasagna module distinguishes the exotic pair of knot traces  $X_{-1}(-5_2)$  and  $X_{-1}(P(3, -3, -8))$ , an example first discovered by Akbulut. This gives the first analysis-free proof of the existence of exotic compact orientable 4-manifolds. We also present a family of exotic knot traces that seem not directly recoverable from gauge/Floer-theoretic methods. Along the way, we present new explicit calculations of the Khovanov skein lasagna modules, and we define lasagna generalizations of the Lee homology and Rasmussen  $s$ -invariant, which are of independent interest. Other consequences of our work include a slice obstruction of knots in 4-manifolds with nonvanishing skein lasagna module, a sharp shake genus bound for some knots from the lasagna  $s$ -invariant, and a construction of induced maps on Khovanov homology for cobordisms in  $k\mathbb{C}\mathbb{P}^2$ .

## 1. INTRODUCTION

The revolutionary work of Donaldson and Freedman in the 1980s revealed a drastic dichotomy between the topological and smooth category of manifolds in dimension 4. Since then, constant efforts are made to find exotic pairs of 4-manifolds.

A pair of smooth 4-manifolds  $X_1$  and  $X_2$  is said to be *exotic* if they are homeomorphic but not diffeomorphic. In preferable cases, showing two 4-manifolds are homeomorphic boils down to checking some algebraic topological invariants agree, thanks to Freedman's fundamental theorems [Fre82]. Showing that they are non-diffeomorphic usually involves showing some gauge-theoretic or Floer-theoretic smooth invariants (Donaldson invariants [Don90], Seiberg-Witten invariants [SW94], Heegaard Floer type invariants [OS04], and various variations/generalizations) differ. To date, all detection of compact orientable exotic 4-manifolds depends heavily on analysis, mostly elliptic PDE. We present the first analysis-free detection result.<sup>1,2</sup>


The relevant smooth invariant in our story is the *Khovanov-Rozansky  $\mathfrak{gl}_2$  skein lasagna module* (Khovanov skein lasagna module for short) defined by Morrison-Walker-Wedrich [MWW22], which is an extension of a suitable variation of the classical Khovanov homology [Kho00] via a skein-theoretic construction. It assigns a tri-graded (by homological degree  $h$ , quantum degree  $q$ , and homology class degree  $\alpha$  below) abelian group

$$\mathcal{S}_{0,h,q}^2(X; L) = \bigoplus_{\alpha \in H_2(X, L)} \mathcal{S}_{0,h,q}^2(X; L; \alpha)$$

to every pair  $(X, L)$  of compact oriented smooth 4-manifold and framed oriented link on its boundary. When  $L$  is the empty link we simply write  $\mathcal{S}_0^2(X)$  for the skein lasagna module.

<sup>1</sup>For a noncompact example, an exotic  $\mathbb{R}^4$  can be constructed from a topologically slice but not smoothly slice knot. Such knots can be detected by Freedman's result [Fre82] and Rasmussen's  $s$ -invariant [Ras10] from Khovanov homology. For nonorientable examples, Kreck [Kre06] showed  $K3\#\mathbb{R}\mathbb{P}^4$  and  $11(S^2 \times S^2)\#\mathbb{R}\mathbb{P}^4$  form an exotic pair using Rochlin's Theorem, and indeed exotic  $\mathbb{R}\mathbb{P}^4$ 's were constructed by Cappell-Shaneson [CS76] (shown to be homeomorphic to  $\mathbb{R}\mathbb{P}^4$  later in [HKT94]).

<sup>2</sup>Exotic surfaces in 4-manifolds were also known to be detectable by Khovanov homology [HS21].

FIGURE 1. Kirby diagrams of  $X_1, X_2$ .

The  $n$ -trace on a knot  $K$ , denoted  $X_n(K)$ , is the 4-manifold obtained by attaching an  $n$ -framed 2-handle to  $B^4$  along  $K \subset S^3 = \partial B^4$ . Our main theorem shows that the  $(-1)$ -traces on the mirror image of the knot  $5_2$  and the Pretzel knot  $P(3, -3, -8)$ , namely

$$X_1 := X_{-1}(-5_2), \quad X_2 := X_{-1}(P(3, -3, -8)),$$

have non-isomorphic skein lasagna module.

**Theorem 1.1.** *The graded modules  $\mathcal{S}_0^2(X_1)$  and  $\mathcal{S}_0^2(X_2)$  are non-isomorphic over  $\mathbb{Q}$ . Therefore,  $X_1$  and  $X_2$  form an exotic pair of 4-manifolds. In fact, even their interiors are exotic.*

The exotic pair  $X_1, X_2$  was first discovered by Akbulut [Akb91b; Akb91a]. The fact that  $X_1, X_2$  are homeomorphic is more or less standard: a sequence of Kirby moves shows  $\partial X_1$  and  $\partial X_2$  are homeomorphic and a suitable extension of Freedman's Theorem [Fre82; Boy86] then implies  $X_1$  and  $X_2$  are homeomorphic. On the other hand, the fact that  $X_1, X_2$  are not diffeomorphic was detected by an intricate calculation of Donaldson invariants. Later, the result was reproved by Akbulut-Matveyev [AM97, Theorem 3.2] using a simplified argument which essentially says  $X_{-1}(-5_2)$  is Stein (since the maximal Thurston-Bennequin number of  $-5_2$  is 1) while  $X_{-1}(P(3, -3, -8))$  is not (since  $P(3, -3, -8)$  is slice). This latter proof is elegant, yet still depends heavily on calculation of Seiberg-Witten invariants of Stein surfaces, as well as deep results of Eliashberg [Eli90] and Lisca-Matić [LM97]. In comparison, our proof of Theorem 1.1 depends on the much lighter package of Khovanov homology, which is combinatorial in nature.

Since the Khovanov skein lasagna module over a field of a connected sum or boundary connected sum of 4-manifolds is equal to the tensor product of the Khovanov skein lasagna modules of the components, a fact proved in [MN22],  $X_1$  and  $X_2$  remain exotic after (boundary) connected sums with any 4-manifolds  $W$  with nonvanishing skein lasagna module (at least when the support of  $\mathcal{S}_0^2(W)$  is small in a suitable sense). Such tricks lead to many new exotic pairs. We write down some explicit examples in the following corollary to give the reader a flavor of what can be proven. For more choices of such candidates of connected summands one may look into Section 6.

**Corollary 1.2.** *The 4-manifolds  $X_1^{\natural a} \natural X_2^{\natural b}$  and  $X_1^{\natural a'} \natural X_2^{\natural b'}$  form an exotic pair if  $a + b = a' + b'$  and  $(a, b) \neq (a', b')$ . The exotica remain under connected sums or boundary sums with any copies of  $S^1 \times S^3$ ,  $S^2 \times D^2$ ,  $\mathbb{C}\mathbb{P}^2$ , and the negative  $E8$  manifold. The exotica also remains upon further taking the interiors of the pair.*

Anubhav Mukherjee informed us that most of these variations seem also achievable via gauge/Floer-theoretic methods and topological arguments. However, below we present a family of exotic knot traces generalizing Theorem 1.1 that seems not directly recoverable from other methods. Let  $P_{n,m}$

and  $Q_{n,m}$  be satellite patterns drawn as in Figure 2, and  $P_{n,m}(K)$ ,  $Q_{n,m}(K)$  be the corresponding satellite knots with companion knot  $K$ . These satellite patterns were first studied by Yasui [Yas15]. Let  $s(K)$  denote the Rasmussen  $s$ -invariant of  $K$ .

**Theorem 1.3.** *If a knot  $K \subset S^3$  has a slice disk  $\Sigma$  in  $k\mathbb{C}\mathbb{P}^2 \setminus \text{int}(B^4)$  and  $n, m$  are two integers satisfying*

$$s(K) = |[\Sigma]| - [\Sigma]^2, \quad n < -[\Sigma]^2, \quad m \geq 0,$$

*then  $X_n(P_{n,m}(K))$  and  $X_n(Q_{n,m}(K))$  form an exotic pair. Their interiors are also exotic.*

In the special case when  $K = U$  is the unknot,  $k = 0$ ,  $\Sigma$  is the trivial slice disk of  $K$  in  $B^4$ , and  $n = -1$ ,  $m = 0$ , Theorem 1.3 recovers Theorem 1.1. One can also construct more exotic pairs in the spirit of Corollary 1.2 by taking connected sums of these examples and other standard manifolds appearing in that corollary.

Now we turn to some explicit calculations of  $\mathcal{S}_0^2$  and other applications. Some topological applications we derived can indeed be proved in other ways, but our approach provides the first analysis-free proofs. We also give a proof of Theorem 1.1 in Section 1.5, assuming some results stated.

Throughout this paper, all 4-manifolds are assumed to be smooth, compact, and oriented. All surfaces are oriented and smoothly and properly embedded. In the current section, many statements are presented in a specialized form. In such cases, we indicate the corresponding generalizations in the parentheses after the statement numbers.

**1.1. A vanishing result and slice obstruction.** For a knot  $K \subset S^3$ , let  $TB(K)$  denote its maximal Thurston-Bennequin number. We have the following vanishing criterion for the Khovanov skein lasagna module.

**Theorem 1.4.** *If the knot trace  $X_n(K)$  for some knot  $K \subset S^3$  and framing  $n \geq -TB(-K)$  embeds into a 4-manifold  $X$ , then  $\mathcal{S}_0^2(X; L) = 0$  for any  $L \subset \partial X$ .*

For example, since  $S^2 \times S^2$  contains an embedded sphere with self-intersection 2, we see the 2-trace of the unknot embeds into  $S^2 \times S^2$ , implying  $\mathcal{S}_0^2(S^2 \times S^2) = 0$ . This answers a question of Manolescu (see [SZ24, Question 2]), generalizing one main Theorem of Sullivan-Zhang [SZ24, Theorem 1.3] and answering [SZ24, Question 3] in the affirmative.

The folklore trace embedding lemma says that  $X_n(K)$  embeds into  $X$  if and only if  $-K$  is  $(-n)$ -slice in  $X$ , i.e. there is a framed slice disk in  $X \setminus \text{int}(B^4)$  of the  $(-n)$ -framed knot  $-K \subset S^3 = \partial B^4$ . Therefore, the contrapositive of Theorem 1.4 can be phrased as follows.

**Corollary 1.5.** *Suppose  $\mathcal{S}_0^2(X; L) \neq 0$  for some  $L \subset \partial X$ . If a knot  $K \subset S^3$  is  $n$ -slice in  $X$ , then  $n > TB(K)$ .  $\square$*

**1.2. Lee skein lasagna modules and lasagna  $s$ -invariants.** In the construction of the Khovanov skein lasagna module, the relevant TQFT (the Khovanov-Rozansky  $\mathfrak{gl}_2$  homology) enters only at the very last step. We could replace this TQFT by its Lee deformation [Lee05], and define an analogous notion of a Lee skein lasagna module of a 4-manifold  $X$  with link  $L \subset \partial X$ , denoted  $\mathcal{S}_0^{Lee}(X; L)$ . It is a  $\mathbb{Q}$ -vector space equipped with a homological  $\mathbb{Z}$ -grading, a quantum  $\mathbb{Z}/4$ -grading, a homology class grading by  $H_2(X, L)$ , and a quantum  $\mathbb{Z}$ -filtration (where 0 has filtration degree  $-\infty$ ), with the caveat that a nonzero vector can have quantum filtration degree  $-\infty$ .

We show in Section 4 that the structure of  $\mathcal{S}_0^{Lee}(X; L)$ , except for the quantum filtration, is determined by simple algebraic data of  $(X, L)$ . Moreover, we extract numerical invariants from the

quantum filtration structure<sup>3</sup>. For the purpose of exposition, in the current section we assume  $L$  is the empty link. Let  $q: \mathcal{S}_0^{Lee}(X) \rightarrow \mathbb{Z} \cup \{-\infty\}$  denote the quantum filtration function.

**Theorem 1.6.** (Theorem 4.1) *The Lee skein lasagna module of a 4-manifold  $X$  is  $\mathcal{S}_0^{Lee}(X) \cong \mathbb{Q}^{H_2(X)^2}$ , with a basis consisting of canonical generators  $x_{\alpha_+, \alpha_-}$ , one for each pair  $\alpha_+, \alpha_- \in H_2(X)$ . Moreover,*

- (1)  $x_{\alpha_+, \alpha_-}$  has homological degree  $-2\alpha_+ \cdot \alpha_-$ ;
- (2)  $x_{\alpha_+, \alpha_-}$  has homology class degree  $\alpha := \alpha_+ + \alpha_-$ ;
- (3)  $x_{\alpha_+, \alpha_-} + x_{\alpha_-, \alpha_+}$ ,  $x_{\alpha_+, \alpha_-} - x_{\alpha_-, \alpha_+}$  have quantum  $\mathbb{Z}/4$  degrees  $-\alpha^2$ ,  $-\alpha^2 + 2$ , respectively;
- (4)  $q(x_{\alpha_+, \alpha_-}) = q(x_{\alpha_-, \alpha_+}) = \max(q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+}))$ , which also equals  $\min(q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+})) + 2$  if  $\alpha_+ \neq \alpha_-$  in  $H_2(X; \mathbb{Q})$ .

**Definition 1.7.** (Definition 4.6) The *lasagna  $s$ -invariant* of  $X$  at a homology class  $\alpha \in H_2(X)$  is

$$s(X; \alpha) := q(x_{\alpha, 0}).$$

The lasagna  $s$ -invariants of  $X$  thus define a smooth invariant  $s(X; \bullet): H_2(X) \rightarrow \mathbb{Z} \cup \{-\infty\}$ .

The genus function of a 4-manifold  $X$ , defined by

$$g(X, \alpha) := \min\{g(\Sigma): \Sigma \subset X, [\Sigma] = \alpha\},$$

captures subtle information about the smooth topology of  $X$ . Like the classical  $s$ -invariant [Ras10; BW08], which provides a lower bound on the slice genus of links, the lasagna  $s$ -invariants provide a lower bound on the genus function of 4-manifolds.

**Theorem 1.8.** (Theorem 4.7(6) and Corollaries after) *The genus function of a 4-manifold  $X$  has a lower bound given by*

$$g(X; \alpha) \geq \frac{s(X; \alpha) + \alpha^2}{2}.$$

Unlike the classical  $s$ -invariant, the lasagna  $s$ -invariants are not directly computable from link diagrams. However, we are able to compute them in some special cases. We indicate one special case where Theorem 1.8 does provide useful information.

The  $n$ -shake genus of a knot  $K$  is defined by  $g_{sh}^n(K) := g(X_n(K); 1)$ , where 1 is a generator of  $H_2(X_n(K)) \cong \mathbb{Z}$ . Clearly,  $g_{sh}^n(K) \leq g_4(K)$ , but the inequality may not be strict [Akb77; Pic19]. Theorem 1.8 specializes to a shake genus bound

$$g_{sh}^n(K) \geq \frac{s(X_n(K); 1) + n}{2}. \tag{1}$$

**Theorem 1.9.** (1) *If  $K$  is concordant to a positive knot, then for  $n \leq 0$  we have  $s(X_n(K); 1) = s(K) - n$ , where  $s(K)$  is the Rasmussen  $s$ -invariant of  $K$ . Thus*

$$g_{sh}^n(K) = g_4(K) = \frac{s(K)}{2}, \quad n \leq 0.$$

(2) *If  $\Sigma \subset \overline{k\mathbb{C}\mathbb{P}^2} \setminus (\text{int}(B^4) \sqcup \text{int}(B^4))$  is a concordance from  $K$  to a positive knot  $K'$ , then for  $n \leq [\Sigma]^2$  we have  $s(X_n(K); 1) \geq s(K') + [\Sigma]^2 + |[\Sigma]| - n$ . Thus*

$$g_{sh}^n(K) \geq \frac{s(K') + [\Sigma]^2 + |[\Sigma]|}{2}, \quad n \leq [\Sigma]^2.$$

---

<sup>3</sup>The authors are aware that the recent paper [MWW24] by Morrison-Walker-Wedrich has an independent proof of a part of this structural result (except addenda (3) and (4)), stated in a more general form. They extract slightly different invariants from the filtration structure, which gives somewhat less information than ours in the  $\mathfrak{gl}_2$  case. Moreover, our invariants generalize Rasmussen  $s$ -invariants for links and enjoy nice functorial properties.

*Remark 1.10.* The computation of shake genus in Theorem 1.9(1) also follows from the following two facts:

- (i) The adjunction inequality for Stein manifolds [LM97], which implies

$$g_{sh}^n(K) \geq (tb(\mathcal{K}) + |\text{rot}(\mathcal{K})| + 1)/2 \quad (2)$$

for any Legendrian representative  $\mathcal{K}$  of  $K$  with  $tb(\mathcal{K}) = n + 1$ . When  $n < TB(K)$ , take  $\mathcal{K}$  to be a stabilization that maximize  $|\text{rot}|$ , of a max  $tb$  representative of  $K$ . Then the right hand side agrees with  $s(K)/2$  for positive knots [Tan99, Theorem 2].

- (ii) Two concordant knots have the same shake genus (by an easy topological argument).

Lasagna  $s$ -invariants also obstruct embeddings between 4-manifolds in the following sense.

**Proposition 1.11.** *If  $i: X \hookrightarrow X'$  is an inclusion between two 4-manifolds, then for any class  $\alpha \in H_2(X)$ ,*

$$s(X'; i_*\alpha) \leq s(X; \alpha).$$

**1.3. Nonvanishing result for 2-handlebodies from lasagna  $s$ -invariants.** The Khovanov homology of a link admits a spectral sequence to its Lee homology. In particular, in every bidegree, the rank of the Khovanov homology is bounded below by the dimension of the associated graded vector space of the Lee homology. Although we are not able to define a spectral sequence from  $\mathcal{S}_0^2$  to  $\mathcal{S}_0^{Lee}$ , we do have a rank inequality when  $X$  is a 2-handlebody. Let  $gr_\bullet V$  denote the associated graded vector space of a filtered vector space  $V$ .

**Theorem 1.12.** *If  $X$  is a 2-handlebody, then in any tri-degree  $(h, q, \alpha) \in \mathbb{Z}^2 \times H_2(X, L)$ ,*

$$\text{rank}(\mathcal{S}_{0,h,q}^2(X; L; \alpha)) \geq \dim(\text{gr}_q(\mathcal{S}_{0,h}^{Lee}(X; L; \alpha))).$$

*In particular, if the quantum filtration on  $\mathcal{S}_0^{Lee}(X; L)$  is not identically  $-\infty$ , then  $\mathcal{S}_0^2(X; L) \neq 0$ .*

We state a comparison result for Lee skein lasagna modules of 2-handlebodies, which can sometimes be used as a nonvanishing criterion for Khovanov skein lasagna modules in view of Theorem 1.12.

**Proposition 1.13.** *(Proposition 5.3) Suppose  $X$  (resp.  $X'$ ) is a 2-handlebody obtained by attaching 2-handles to  $B^4$  along a framed oriented link  $K \subset S^3$  (resp.  $K' \subset S^3$ ). If  $K$  is framed concordant to  $K'$ , then  $\mathcal{S}_0^{Lee}(X) \cong \mathcal{S}_0^{Lee}(X')$  as tri-graded, quantum filtered vector spaces. Under the identification  $H_2(X) \cong H_2(X')$  induced by a link concordance,  $X$  and  $X'$  have equal lasagna  $s$ -invariants.*

**1.4. Nonvanishing result for 2-handlebodies from link diagram.** Using a diagrammatic argument, in Section 5.4, we prove a nonvanishing result for 2-handlebodies with sufficiently negative framings. For the ease of exposition, here we only state the case for knot traces. Let  $n_-(K)$  be the minimal number of negative crossings among all diagrams of a knot  $K$ .

**Theorem 1.14.** *(Theorem 5.5) Let  $K$  be a knot and  $X = X_n(K)$  be its  $n$ -trace.*

- (1) *If  $n < -2n_-(K)$ , then  $s(X; 0) = 0$  and  $\mathcal{S}_{0,0,q}^2(X; 0) \otimes \mathbb{Q} = \text{gr}_q(\mathcal{S}_{0,0}^{Lee}(X; 0)) = \begin{cases} \mathbb{Q}, & q = 0 \\ 0, & q \neq 0. \end{cases}$*

- (2) *If  $K$  is positive, then for  $n \leq 0$ ,  $s(X; 0) = 0$ ,  $s(X; 1) = s(K) - n$ . For  $n < 0$ ,*

$$\mathcal{S}_{0,0,q}^2(X; 1) \otimes \mathbb{Q} = \text{gr}_q(\mathcal{S}_{0,0}^{Lee}(X; 1)) = \begin{cases} \mathbb{Q}, & q = s(X; 1), s(X; 1) - 2 \\ 0, & \text{otherwise.} \end{cases}$$

**1.5. Proof of Theorem 1.1.** Since  $-5_2$  is a positive knot with  $s(-5_2) = 2$ , by Theorem 1.9(1) we have  $g_{sh}^{-1}(-5_2) = 1$ . On the other hand,  $P(3, -3, -8)$  is slice, thus  $g_{sh}^{-1}(P(3, -3, 8)) = 0$ . This shows  $int X_1$  and  $int X_2$  are non-diffeomorphic, without actually calculating  $\mathcal{S}_0^\bullet(X_i)$ .

We still need to argue the Khovanov (and indeed Lee) skein lasagna modules of  $X_1, X_2$  actually differ. Apply Theorem 1.14(2) to  $-5_2$ , we obtain  $s(X_1; 1) = 3$ ,

$$\mathcal{S}_{0,0,q}^2(X_1; 1) \otimes \mathbb{Q} = gr_q(\mathcal{S}_{0,0}^{Lee}(X_1; 1)) = \begin{cases} \mathbb{Q}, & q = 1, 3 \\ 0, & \text{otherwise.} \end{cases}$$

Apply Theorem 1.14(2) to the unknot, and then Proposition 1.13 to  $P(3, -3, -8)$  and the unknot, we obtain  $s(X_2; 1) = 1$ , and

$$\mathcal{S}_{0,0,q}^2(X_2; 1) \otimes \mathbb{Q} \supset gr_q(\mathcal{S}_{0,0}^{Lee}(X_2; 1)) = \begin{cases} \mathbb{Q}, & q = \pm 1 \\ 0, & \text{otherwise.} \end{cases}$$

where the first containment follows from Theorem 1.12. In particular,  $\mathcal{S}_{0,0,-1}^2(X_1; 1) \otimes \mathbb{Q} \neq \mathcal{S}_{0,0,-1}^2(X_2; 1) \otimes \mathbb{Q}$ .  $\square$

**1.6. Example: Khovanov and Lee skein lasagna module of  $\overline{\mathbb{C}\mathbb{P}^2}$ .** As an example, we note that the recent work by the first author [Ren23] has the following implications on the Lee and Khovanov skein lasagna modules for  $\overline{\mathbb{C}\mathbb{P}^2}$ .

**Proposition 1.15.** *The lasagna  $s$ -invariants of  $\overline{\mathbb{C}\mathbb{P}^2}$  are given by  $s(\overline{\mathbb{C}\mathbb{P}^2}; \alpha) = |\alpha|$ . The associated graded vector space of the Lee skein lasagna module of  $\overline{\mathbb{C}\mathbb{P}^2}$  is given by  $gr_q \mathcal{S}_{0,(\beta^2-\alpha^2)/2}^{Lee}(\overline{\mathbb{C}\mathbb{P}^2}; \alpha) = \mathbb{Q}$  for*

$$\beta \equiv \alpha \pmod{2}, \quad q = \begin{cases} \frac{\alpha^2 - \beta^2}{2} + |\beta| - 1 \pm 1, & \beta \neq 0 \\ \frac{\alpha^2}{2}, & \beta = 0, \end{cases}$$

and 0 in other tri-degrees. Moreover, for  $\beta \equiv \alpha \pmod{2}$ ,

$$\mathcal{S}_{0,(\beta^2-\alpha^2)/2,(\alpha^2-\beta^2)/2+|\beta|}^2(\overline{\mathbb{C}\mathbb{P}^2}; \alpha) = \mathbb{Z}.$$

A consequence of this result, together with some properties of lasagna  $s$ -invariants, is the following.

**Corollary 1.16.** *If a 4-manifold  $X$  admits an embedding  $i: X \hookrightarrow k\overline{\mathbb{C}\mathbb{P}^2}$  for some  $k$ , then  $s(X; \alpha) \geq |i_*\alpha|$ , with equality achieved if  $X$  is a 2-handlebody and  $\alpha = 0$ . In particular,  $\mathcal{S}_{0,0,q}^2(X; \alpha) \neq 0$  for  $q = s(X; \alpha)$ , and for  $q = s(X; \alpha) - 2$  if  $\alpha \neq 0 \in H_2(X; \mathbb{Q})$ .*

It is also worth pointing out that Conjecture 6.1 in [Ren23] is equivalent to a complete determination of the Khovanov skein lasagna module of  $\overline{\mathbb{C}\mathbb{P}^2}$  (see Conjecture 6.14).

If  $\Sigma \subset \mathbb{C}\mathbb{P}^2 \setminus (int(B^4) \sqcup int(\overline{B^4}))$  is an oriented cobordism in  $\mathbb{C}\mathbb{P}^2$  between two oriented links  $L_0, L_1$ , the last part of Proposition 1.15 enables us to define an induced map

$$Kh(\Sigma): Kh(L_0) \rightarrow Kh(L_1)$$

between ordinary Khovanov homology of  $L_0$  and  $L_1$ , at least over a field and up to sign, with bidegree  $(0, \chi(\Sigma) - [\Sigma]^2 + |[\Sigma]|)$ . We carry out this construction in more details in Section 6.11.

**1.7. A conjecture for Stein manifolds.** Many similarities between skein lasagna modules and other gauge/Floer-theoretic smooth invariants appear. However, a serious constraint for our computation is that we were not able to produce any 4-manifold with  $b_2^+ > 0$  that has finite lasagna  $s$ -invariants or nonvanishing Khovanov skein lasagna module. In fact, if one were able to find a simply-connected closed 4-manifold  $X$  with  $b_2^+(X) > 0$  that has  $\mathcal{S}_0^2(X) \neq 0$ , then  $X \# \overline{\mathbb{C}\mathbb{P}^2}$  is an exotic closed 4-manifold, as it has nonvanishing  $\mathcal{S}_0^2$  while its standard counterpart (some  $k\mathbb{C}\mathbb{P}^2 \# \overline{\ell\mathbb{C}\mathbb{P}^2}$ ) has vanishing  $\mathcal{S}_0^2$ .

To pursue this thread, we propose the following conjecture, which seems plausible and desirable.

**Conjecture 1.17.** *If  $K$  is a knot and  $X = X_n(K)$  for some  $n < TB(K)$ , then  $s(X; 0) = 0$  and  $s(X; 1) = s(K) - n$ . In particular,  $\mathcal{S}_0^2(X) \neq 0$ . More generally, any Stein 2-handlebody  $X$  has  $s(X; 0) > -\infty$  and nonvanishing Khovanov skein lasagna module.*

The part concerning  $s(X; 1)$  in Conjecture 1.17 implies the  $n$ -shake genus of  $K$  for  $n < TB(K)$  (or more generally  $n < TB(K')$  for some  $K'$  concordant to  $K$ ) is bounded below by  $s(K)/2$ , generalizing Theorem 1.9(1). This seems to be the proper straightening of the bound from the Stein adjunction inequality (2).

As will be clear later, Conjecture 1.17 is equivalent to an assertion on the  $s$ -invariants of cables of  $K$  (or some Legendrian link in the 2-handlebody case) (cf. Proposition 5.4).

**1.8. Organization of the paper.** In Section 2, we review the theory of Khovanov skein lasagna modules, expanding on a few properties, as well as making some new comments on the skein-theoretic construction. In the short Section 3, we prove Theorem 1.4 by applying a bound on Khovanov homology by Ng [Ng05], and we give some examples of 4-manifolds with vanishing Khovanov skein lasagna module.

In Section 4, we develop the theory of Lee skein lasagna modules and lasagna  $s$ -invariants, which can be thought of as a lasagna generalization of the classical theory by Lee [Lee05] and Rasmussen [Ras10] for links in  $S^3$ . A generalization of Theorem 1.6, in particular the definition of canonical Lee lasagna generators, is presented in Section 4.1. The behavior of canonical Lee lasagna generators under morphisms is given in Theorem 4.5 in Section 4.2. Building on these results, in Section 4.3 we define and examine the lasagna  $s$ -invariants, generalizing Definition 1.7, Theorem 1.8, and Proposition 1.11.

In Section 5, we make use of a 2-handlebody formula by Manolescu-Neithalath [MN22] to prove nonvanishing results for Khovanov and Lee skein lasagna modules of 2-handlebodies. In Section 5.1, we prove Theorem 1.12 which shows that finiteness results for the quantum filtration of Lee skein lasagna modules imply nonvanishing results for Khovanov skein lasagna modules. In Section 5.2 we prove a generalization of Proposition 1.13 and in Section 5.3 we show that lasagna  $s$ -invariants can be expressed in terms of classical  $s$ -invariants of some cable links. Finally, in the somewhat technical Section 5.4, we prove a vast generalization of the diagrammatic nonvanishing criterion Theorem 1.14.

Finally, in Section 6, we give examples and applications, mostly to the theory and tools developed in Section 4 and Section 5. This includes proofs of Corollary 1.2, Theorem 1.3, Theorem 1.9, and an expanded discussion of Section 1.6. Other discussions include a proof that  $s$ -invariants of cables of the Conway knot can obstruct its sliceness, and that the Khovanov skein lasagna module over  $\mathbb{Q}$  does not detect potential exotica arising from Gluck twists in  $S^4$ .

Most of our developments are independent to proving Theorem 1.1. The readers only interested in Theorem 1.1 may want to look at its proof in Section 1.5 and delve into the corresponding sections. In particular, to distinguish the smooth structures of the pair  $X_1, X_2$  in Theorem 1.1, one can read the first half of Section 2, certain statements in Section 4 (mainly Definition 4.6 and Corollary 4.13), Section 5.4, and Section 6.4. The minimal amount of theory needed to actually distinguish their Khovanov skein lasagna modules depends moreover on Section 5.1 and Section 5.2 (but not really on Corollary 4.13 and Section 6.4). Still, the statements in those sections are often much more general than needed for Theorem 1.1, which at times complicate the proofs.

**Acknowledgement.** We thank Ian Agol, Daren Chen, Kyle Hayden, Sungkyung Kang, Ciprian Manolescu, Marco Marengon, Anubhav Mukherjee, Ikshu Neithalath, Lisa Piccirillo, Mark Powell, Yikai Teng, Joshua Wang, Paul Wedrich, Hongjian Yang for helpful discussions. This work was supported in part by the Simons grant “new structures in low-dimensional topology” through the workshop on exotic 4-manifolds held at Stanford University December 16-18, 2023.

## 2. KHOVANOV AND LEE SKEIN LASAGNA MODULES

The Khovanov-Rozansky  $\mathfrak{gl}_2$  skein lasagna modules were defined by Morrison-Walker-Wedrich [MWW22]. In the first part of this section we review its construction, cast in a slightly different way, while defining its Lee deformation under the same framework. In Section 2.1, we begin with a general skein-theoretic construction of skein lasagna modules, using an arbitrary (stratified) TQFT for links in  $S^3$ . In Section 2.2, we state a few properties of skein lasagna modules, most of which are essentially proved in [MN22]. Only then, in Section 2.3, do we fix the TQFT in consideration to be either the  $\mathfrak{gl}_2$  Khovanov-Rozansky homology  $KhR_2$  [MWW22] or its Lee deformation  $KhR_{Lee}$  [Lee05]. We state a few more computational results for these settings in Section 2.4.

The second part of the section is new, and can be skipped on a first reading. In Section 2.5, we identify the structure of the underlying space of skeins in the construction of Section 2.1. The proof will eventually be useful in Section 4.1 where we calculate the Lee skein lasagna module of a pair  $(X, L)$ . Finally, in Section 2.6, we suggest a few variations of the skein-theoretic construction.

**2.1. Construction of skein lasagna modules.** <sup>4</sup> Let  $X$  be a 4-manifold and  $L \subset \partial X$  be a framed oriented link on its boundary. A *skein* in  $(X, L)$  or in  $X \text{ rel } L$  is a properly embedded framed oriented surface in  $X$  with the interiors of finitely many disjoint 4-balls deleted, whose boundary on  $\partial X$  is  $L$ . That is, a skein in  $(X, L)$  can be written as

$$\Sigma \subset X \setminus \sqcup_{i=1}^k \text{int}(B_i), \text{ with } \partial \Sigma|_{\partial X} = L, \quad (3)$$

The deleted balls  $B_i$  are called *input balls* of the skein  $\Sigma$ .

We define the *category of skeins* in  $(X, L)$ , denoted  $\mathcal{C}(X; L)$ , as follows. The objects consist of all skeins in  $X \text{ rel } L$ . Suppose  $\Sigma$  is an object as in (3) and  $\Sigma'$  is another object with input balls  $B'_j$ ,  $j = 1, \dots, k'$ . If  $\sqcup B_i \subset \sqcup B'_j$  and each  $B'_j$  either coincides with some  $B_i$  or does not intersect any  $B_i$  on its boundary, then the set of morphisms from  $\Sigma$  to  $\Sigma'$  consists of pairs of an isotopy class of surfaces<sup>5</sup>  $S \subset \sqcup B'_j \setminus \sqcup \text{int}(B_i)$  rel boundary with  $S|_{\partial B_i} = \Sigma|_{\partial B_i}$ ,  $S|_{\partial B'_j} = \Sigma'|_{\partial B'_j}$ , and an isotopy class of isotopies from  $S \cup \Sigma'^6$  to  $\Sigma$  rel boundary. We will suppress the isotopy and write  $[S]: \Sigma \rightarrow \Sigma'$  for

<sup>4</sup>The formulation of our construction was motivated by a conversation with Sungkyung Kang.

<sup>5</sup>We formally allow some components of  $S$  to be curves on the boundaries of those  $B_i = B'_j$ .

<sup>6</sup>In order for the surfaces to glue smoothly, one may assume throughout that all embedded surfaces have a standard collar near each boundary, and all isotopies rel boundary respect these collars.



this morphism. If the input balls  $B_i, B'_j$  do not satisfy the condition above, there is no morphism from  $\Sigma$  to  $\Sigma'$ . The identity morphism at  $\Sigma$  is given by  $[\Sigma|_{\sqcup \partial B_i}]$ . The composition of two morphisms  $[S]:\Sigma \rightarrow \Sigma'$  and  $[S']:\Sigma' \rightarrow \Sigma''$  is  $[S \cup S']$  (together with the composition of isotopies).

Next we introduce an arbitrary (stratified) TQFT of framed oriented links in  $S^3$ , by which we mean a symmetric monoidal functor  $Z$  from the category of framed oriented links in  $S^3$  and link cobordisms in  $S^3 \times I$  up to isotopy rel boundary, to the category of modules over a commutative ring  $R$ , such that  $Z(\emptyset) = R$  and  $Z(L)$  is finitely generated for all links  $L$ . Equivalently,  $Z$  is a functorial link invariant in  $\mathbb{R}^3$  satisfying the conditions in Theorem 2.1 of [MWW24]. At times we also assume  $R$  is a field and  $Z$  is involutive, meaning that  $Z(-L) \cong Z(L)^*$  for a link  $L$  and  $Z(-\Sigma) \cong Z(\Sigma)^*$  for a cobordism  $\Sigma: L_0 \rightarrow L_1$ , where  $-\Sigma: -L_1 \rightarrow -L_0$  is the reverse cobordism.

Assign to every object  $\Sigma$  of  $\mathcal{C}(X; L)$  as in the notations before the  $R$ -module  $\otimes_{i=1}^k Z(K_i)$  which we denote by an abuse of notation  $Z(\Sigma)$ , where each  $K_i = \partial \Sigma|_{\partial B_i}$  is a framed link. Assign to every morphism  $[S]:\Sigma \rightarrow \Sigma'$  the morphism  $Z([S]):Z(\Sigma) \rightarrow Z(\Sigma')$  induced by the cobordism  $S: \sqcup K_i \rightarrow \sqcup K'_j$ . Then  $Z$  defines a functor from the category of skeins in  $(X, L)$  to the category of  $R$ -modules.

The *skein lasagna module* of  $(X, L)$  with respect to  $Z$ , denoted  $\mathcal{S}_0^Z(X; L)$ , can be now defined as

$$\mathcal{S}_0^Z(X; L) := \text{colim}(Z: \mathcal{C}(X; L) \rightarrow \mathbf{Mod}_R). \quad (4)$$

More explicitly,

$$\mathcal{S}_0^Z(X; L) = \{(\Sigma, v): \Sigma \subset X \setminus \sqcup_{i=1}^k \text{int}(B_i), v \in Z(\Sigma)\} / \sim \quad (5)$$

where  $\sim$  is the equivalence relation generated by linearity in  $v$ , and  $(\Sigma, v) \sim (\Sigma', Z([S])(v))$  for  $[S]:\Sigma \rightarrow \Sigma'$ . A pair  $(\Sigma, v)$  (or just  $v$  if there's no confusion) as in (5) is called a *lasagna filling* of  $(X, L)$ .

The skein lasagna module  $\mathcal{S}_0^Z(X; L)$  is an invariant of the pair  $(X, L)$  up to diffeomorphism. When  $L = \emptyset$ , we write for short  $\mathcal{S}_0^Z(X)$  for  $\mathcal{S}_0^Z(X; L)$ . In this case, since all lasagna fillings and their relations appear in the interior of  $X$ ,  $\mathcal{S}_0^Z(X)$  is an invariant of  $\text{int}(X)$ .

**2.2. Properties of skein lasagna modules.** We state some properties of the skein lasagna modules, mostly following Manolescu-Neithalath [MN22]. Although their paper dealt with the case when the TQFT  $Z$  is the Khovanov-Rozansky  $\mathfrak{gl}_N$  homology, the proofs remain valid in the more general setup. We remark that the more recent paper [MWW23] gives a formula for  $\mathcal{S}_0^Z(X; L)$  in terms of a handlebody decomposition of  $X$ . However, we will not be using this result, so we refer interested readers to their paper.

Every object  $\Sigma$  of  $\mathcal{C}(X; L)$  represents a homology class in  $H_2(X, L)$ , whose image under the boundary homomorphism  $\partial$  in the long exact sequence

$$0 \rightarrow H_2(X) \rightarrow H_2(X, L) \xrightarrow{\partial} H_1(L) \rightarrow H_1(X) \rightarrow \dots$$

is  $[L]$ , the fundamental class of  $L$ . If  $[S]:\Sigma \rightarrow \Sigma'$  is a morphism, then  $[\Sigma] = [\Sigma'] \in H_2(X, L)$ . This means  $\mathcal{S}_0^Z(X; L)$  admits a grading by  $H_2^L(X) := \partial^{-1}([L])$ , which we write as

$$\mathcal{S}_0^Z(X; L) = \bigoplus_{\alpha \in H_2^L(X)} \mathcal{S}_0^Z(X; L; \alpha).$$

Of course,  $H_2^L(X)$  is nonempty if and only if the image of  $[L]$  in  $H_1(X)$  is zero, i.e.  $L$  is null-homologous in  $X$ . From now on this will be assumed, since otherwise  $\mathcal{S}_0^Z(X; L)$  is trivially zero as there is no lasagna filling.

**Proposition 2.1 (3, 4-handle attachments, [MN22, Proposition 2.1]).** *If  $(X', L)$  is obtained from  $(X, L)$  by attaching a 3-handle (resp. 4-handle) away from  $L$ , then there is a natural surjection (resp. isomorphism)  $\mathcal{S}_0^Z(X; L) \rightarrow \mathcal{S}_0^Z(X'; L)$ .*  $\square$

**Proposition 2.2 (Connected sum formula, [MN22, Theorem 1.4, Corollary 7.3]).** *If the base ring  $R$  is a field and  $Z$  is involutive, then there is a natural isomorphism*

$$\mathcal{S}_0^Z(X \natural X'; L \sqcup L') \cong \mathcal{S}_0^Z(X; L) \otimes \mathcal{S}_0^Z(X'; L').$$

*The same formula holds with the boundary sum replaced by the connected sum or the disjoint union.*  $\square$

Suppose  $(X, L)$  is a pair and  $Y \subset X$  is a properly embedded separating oriented 3-manifold, possibly with transverse intersections with  $L$  in some point set  $P \subset \partial Y$ . Then  $Y$  cuts  $(X, L)$  into two pairs  $(X_1, T_1)$  and  $(X_2, T_2)$ , where  $\partial X_1$  induces the orientation on  $Y$  and  $\partial X_2$  induces the reverse orientation, and  $T_1, T_2$  are framed oriented tangles with endpoints  $P$ . The framing and the orientation of  $L$  restrict to a framing and an orientation of  $P \subset \partial Y$ .

If  $T_0 \subset Y$  is a framed oriented tangle with  $\partial T_0 = P$  in the framed oriented sense, then  $T_1 \cup T_0$  is a framed oriented link in  $\partial X_1$  and  $-T_0 \cup T_2$  is one in  $\partial X_2$ . A lasagna filling of  $(X_1, T_1 \cup T_0)$  and a lasagna filling of  $(X_2, -T_0 \cup T_2)$  glue to a lasagna filling of  $(X, L)$  in a bilinear way, inducing a map

$$\mathcal{S}_0^Z(X_1; T_1 \cup T_0) \otimes \mathcal{S}_0^Z(X_2; -T_0 \cup T_2) \rightarrow \mathcal{S}_0^Z(X; L) \quad (6)$$

**Proposition 2.3 (Gluing).** *The gluing map*

$$\bigoplus_{\substack{T_0 \subset Y \\ \partial T_0 = P}} \mathcal{S}_0^Z(X_1; T_1 \cup T_0) \otimes \mathcal{S}_0^Z(X_2; -T_0 \cup T_2) \rightarrow \mathcal{S}_0^Z(X; L)$$

*is a surjection.*

*Proof.* Every lasagna filling of  $(X, L)$  can be isotoped to intersect  $Y$  transversely in some tangle  $T_0$ .  $\square$

Write  $L_1 = T_1 \cup T_0$ . A framed surface  $S \subset X_2$  with  $\partial S = -T_0 \cup T_2$  is a lasagna filling of  $(X_2; -T_0 \cup T_2)$  (without input balls), thus putting its class in the second tensor summand of (6) defines a gluing map

$$\mathcal{S}_0^Z(X_2; S): \mathcal{S}_0^Z(X_1; L_1) \rightarrow \mathcal{S}_0^Z(X; L). \quad (7)$$

The most useful cases of gluing might be when  $Y$  is closed, where  $P = \emptyset$ , and  $T_0, T_1, T_2$  are links. Still more specially, if  $X \subset \text{int}(X')$  is an inclusion of 4-manifolds, there is an induced map  $\mathcal{S}_0^Z(X) \rightarrow \mathcal{S}_0^Z(X')$  by gluing the exterior of  $X \subset X'$  to  $X$ . For example, the map in Proposition 2.1 is defined this way.

The skein lasagna module with respect to  $Z$  refines the TQFT  $Z$  in the following sense.

**Proposition 2.4 (Recover TQFT).** *For a framed oriented link  $L \subset S^3$  we have a natural isomorphism  $\mathcal{S}_0^Z(B^4; L) \cong Z(L)$ . If  $S \subset S^3 \times I$  is a cobordism from  $L$  to  $L'$ , then the gluing map  $\mathcal{S}_0^Z(S^3 \times I; S): \mathcal{S}_0^Z(B^4; L) \rightarrow \mathcal{S}_0^Z(B^4; L')$  agrees with  $Z(S): Z(L) \rightarrow Z(L')$  under the natural isomorphisms.*

*Proof.* The statement on objects is proved in [MWW22, Example 5.6] and the statement on morphisms is a simple exercise.  $\square$

Finally, we remark that extra structures on the TQFT  $Z$  usually induce structures on the skein lasagna module  $\mathcal{S}_0^Z$ . For example, suppose  $Z$  takes values in  $R$ -modules with a grading by an abelian group  $A$ , so that for a link cobordism  $S$ ,  $Z(S)$  is homogeneous with degree  $\chi(S)a$  for some  $a \in A$  independent of  $S$ . Then the lasagna module  $\mathcal{S}_0^Z(X; L)$  inherits an  $A$ -grading, once we impose an extra grading shift of  $\chi(\Sigma)a$  in the definition of  $Z(\Sigma)$  for a skein  $\Sigma$ . Analogously, a filtration structure on  $Z$  induces a filtration structure on  $\mathcal{S}_0^Z$ . The isomorphisms or morphisms stated or constructed in this section all respect the extra grading or filtration structure, with a warning that the gluing morphism  $\mathcal{S}_0^Z(X_2; S)$  in (7) has degree or filtration degree  $(\chi(S) - \chi(T_0))a$ .

*Remark 2.5.* In view of the gluing morphism (7) (in the special case  $\partial Y = \emptyset$  and  $T_1 = \emptyset$ ), one is tempted to define a TQFT for framed oriented links in oriented 3-manifolds and cobordisms between them. Namely, for a 3-manifold  $Y$  and link  $L \subset Y$ , define

$$Z(Y, L) := \bigoplus_{\partial X=Y} \mathcal{S}_0^Z(X; L).$$

For a cobordism  $(W, S): (Y, L) \rightarrow (Y', L')$ , define  $Z(W, S): Z(Y, L) \rightarrow Z(Y', L')$  by taking the direct sum of the gluing maps  $\mathcal{S}_0^Z(X; L) \rightarrow \mathcal{S}_0^Z(X \cup_Y W; L')$ .

However, the resulting modules  $Z(Y, L)$  are usually infinite dimensional. To obtain a finite theory, one has to either restrict the domain category (one extreme of which is to recover back the TQFT  $Z$  for links in  $S^3$  again), or modulo suitable skein-theoretic relations on  $Z(Y, L)$ . We leave the exploration of these possibilities to future works.

**2.3. Khovanov-Rozansky  $\mathfrak{gl}_2$  homology and its Lee deformation.** In this paper we will take the TQFT  $Z$  in the skein lasagna construction to be the Khovanov-Rozansky  $\mathfrak{gl}_2$  homology or its Lee deformation. In this section we review some features of these TQFTs, partly also to fix the renormalization conventions and do some bookkeeping. We assume the reader is familiar with the classical Khovanov homology as well as its Lee deformation. See [Kho00; Lee05; Ras10] for relevant backgrounds and [Bar02] for an accessible overview.

We follow the convention in [MWW22]. Khovanov-Rozansky  $\mathfrak{gl}_2$  homology is a renormalization of the Khovanov-Rozansky  $\mathfrak{sl}_2$  homology defined in [KR08] (and extended to  $\mathbb{Z}$ -coefficient in [Bla10]) that is sensitive to the framing of a link. Since conventionally most calculations have been done using the classical Khovanov homology [Kho00], we write down explicitly the renormalization differences between it and the  $\mathfrak{gl}_2$  theory.

Let  $KhR_2$  and  $Kh$  denote the Khovanov-Rozansky  $\mathfrak{gl}_2$  homology and the usual Khovanov homology, respectively. Then for a framed oriented link  $L \subset S^3$ , we have

$$KhR_2^{h,q}(L) \cong Kh^{h,-q-w(L)}(-L), \quad (8)$$

where  $w(L)$  is the writhe of  $L$ , and  $-L$  is the mirror image of  $L$ . When we want to emphasize that we are working over a commutative ring  $R$ , write  $Kh(L; R)$  and  $KhR_2(L; R)$  for the corresponding homology theories.

A dotted framed oriented cobordism  $S: L_0 \rightarrow L_1$  in  $S^3 \times I$  induces a map  $KhR_2(S): KhR_2(L_0) \rightarrow KhR_2(L_1)$  of bidegree  $(0, -\chi(S) + 2\#(\text{dots}))$  which only depends on the isotopy class of  $S$  rel boundary. Let  $\bar{S}: -L_0 \rightarrow -L_1$  denote the mirror image of  $S$ . Then the identification (8) is functorial in the sense that  $Kh(\bar{S}): Kh(-L_0) \rightarrow Kh(-L_1)$  is equal to  $KhR_2(S)$  under (8) up to sign. Since the classical Khovanov homology is only known to be functorial in  $\mathbb{R}^3$  up to sign,  $KhR_2$  can be thought of as a sign fix and a functorial upgrade of  $Kh$ .

Both the constructions of  $KhR_2$  and  $Kh$  depend on the input of the Frobenius algebra  $V = \mathbb{Z}[X]/(X^2)^7$  with comultiplication and counit given by

$$\Delta 1 = X \otimes 1 + 1 \otimes X, \quad \Delta X = X \otimes X, \quad \epsilon 1 = 0, \quad \epsilon X = 1.$$

Here  $X$  has quantum degree 2 in the  $KhR_2$  case and  $-2$  in the  $Kh$  case.

The Lee deformation of  $V$  over  $\mathbb{Q}$  [Lee05] is the Frobenius algebra  $V_{Lee} = \mathbb{Q}[X]/(X^2 - 1)$  with

$$\Delta 1 = X \otimes 1 + 1 \otimes X, \quad \Delta X = X \otimes X + 1 \otimes 1, \quad \epsilon 1 = 0, \quad \epsilon X = 1.$$

Using  $V_{Lee}$  instead of  $V$  defines the Khovanov-Rozansky  $\mathfrak{gl}_2$  Lee homology and the classical Lee homology, denoted  $KhR_{Lee}$  and  $Kh_{Lee}$ , respectively. Since  $V_{Lee}$  is no longer graded but only filtered, the resulting homology theories carry quantum filtration structures, instead of quantum gradings (although a compatible quantum  $\mathbb{Z}/4$ -grading is still present). We remark that  $0 \in KhR_{Lee}$  has quantum filtration degree  $-\infty$  while  $0 \in Kh_{Lee}$  has quantum filtration degree  $+\infty$ , since the two filtrations go in different directions.

Either  $KhR_2$  or  $KhR_{Lee}$  changes only by a renormalization upon changing the orientation of the link  $L$ . More precisely, if  $(L, \mathfrak{o})$  denote the link  $L$  equipped with a possibly different orientation  $\mathfrak{o}$ , then for  $\bullet \in \{2, Lee\}$ ,

$$KhR_{\bullet}(L, \mathfrak{o}) \cong KhR_{\bullet}(L) \left[ \frac{w(L) - w(L, \mathfrak{o})}{2} \right] \left\{ \frac{w(L, \mathfrak{o}) - w(L)}{2} \right\}. \quad (9)$$

It is sometimes more convenient to change from the basis  $\{1, X\}$  of  $V_{Lee}$  to the basis  $\{\mathbf{a} = (1 + X)/2, \mathbf{b} = (1 - X)/2\}$ . In this new basis,

$$\mathbf{a}\mathbf{b} = 0, \quad \mathbf{a}^2 = \mathbf{a}, \quad \mathbf{b}^2 = \mathbf{b}, \quad 1 = \mathbf{a} + \mathbf{b}, \quad X = \mathbf{a} - \mathbf{b}, \quad \Delta \mathbf{a} = 2\mathbf{a} \otimes \mathbf{a}, \quad \Delta \mathbf{b} = -2\mathbf{b} \otimes \mathbf{b}, \quad \epsilon \mathbf{a} = \frac{1}{2}, \quad \epsilon \mathbf{b} = -\frac{1}{2}.$$

Like the classical story, the Khovanov-Rozansky  $\mathfrak{gl}_2$  Lee homology of a framed oriented link  $L$  has a basis consisting of “(rescaled) canonical generators”  $x_{\mathfrak{o}}$  [Ras10, Section 2.3]<sup>8</sup>, one for each orientation  $\mathfrak{o}$  of  $L$  as an unoriented link. Having fixed  $\mathfrak{o}$  (and thus the generator  $x_{\mathfrak{o}}$ ), we write  $L = L_+(\mathfrak{o}) \cup L_-(\mathfrak{o})$  as a union of sublinks on which  $\mathfrak{o}$  agrees (resp. differs) with the orientation of  $L$ . Then  $x_{\mathfrak{o}}$  has homological degree  $-2\ell k(L_+(\mathfrak{o}), L_-(\mathfrak{o})) = (w(L, \mathfrak{o}) - w(L))/2$ , where  $w(L, \mathfrak{o})$  is the writhe of  $L$  with orientation  $\mathfrak{o}$ . Moreover,  $x_{\mathfrak{o}} + x_{\bar{\mathfrak{o}}}, x_{\mathfrak{o}} - x_{\bar{\mathfrak{o}}}$  have quantum  $\mathbb{Z}/4$  degrees  $-w(L) + \#L, -w(L) + \#L + 2$ , respectively, where  $\bar{\mathfrak{o}}$  is the reverse orientation of  $\mathfrak{o}$  (This statement is preserved under disjoint union, cobordisms (in view of (10)), so it suffices to be checked for the two Hopf links).

The unique canonical generator of the empty link is  $1 \in \mathbb{Q}$ . For the 0-framed unknot  $(U, \mathfrak{o})$  we have  $KhR_{Lee}(U) = V_{Lee}\{-1\}$  as a filtered vector space sitting in homological degree 0, and the canonical generators are given by  $x_{\mathfrak{o}} = 2\mathbf{a}, x_{\bar{\mathfrak{o}}} = -2\mathbf{b}$ . Via Reidemeister I induced maps, for any  $n$ , the  $n$ -framed unknot  $(U^n, \mathfrak{o})$  has  $KhR_{Lee}(U^n) = V_{Lee}\{-1 - n\}$  naturally, and the canonical generators are again given by  $x_{\mathfrak{o}} = 2\mathbf{a}, x_{\bar{\mathfrak{o}}} = -2\mathbf{b}$ .

The generators  $x_{\mathfrak{o}}$  behave nicely under cobordisms (cf. Footnote 8 and proof of Theorem 3.3 in [MWW24] about twisting; see also [Ras05, Proposition 3.2] for the classical version with sign

<sup>7</sup>Strictly speaking, the construction of  $KhR_2$  uses the formalism of foams, but for simplicity, our discussion stays in the classical Khovanov formalism throughout.

<sup>8</sup>Here  $x_{\mathfrak{o}}$ 's are defined to be the (grading renormalized) canonical generators on the right hand side of equation (1) in [MWW24] for  $\Sigma = \{\pm 1\}$  (see Proposition 3.12 there), rescaled by  $2^{\#L/2}(-1)^{\#L - (\mathfrak{o})/2}$ . More classically,  $x_{\mathfrak{o}}$  up to sign is Rasmussen's generator  $[\mathfrak{s}_{-\mathfrak{o}}]$  of  $-L$  renormalized to the  $\mathfrak{gl}_2$  theory, rescaled by  $2^{(w(D, -\mathfrak{o}) - r(D, -\mathfrak{o}) + \#L)/2}$  where  $w(D, -\mathfrak{o}), r(D, -\mathfrak{o})$  are the writhe and the number of Seifert circles of a framed diagram  $D$  of  $-L$  equipped with orientation  $-\mathfrak{o}$  used to define  $[\mathfrak{s}_{-\mathfrak{o}}]$ .

ambiguity<sup>9</sup>). If  $S: L \rightarrow L'$  is a framed oriented link cobordism and  $\mathfrak{o}$  is an orientation of  $L$  as an unoriented link, then

$$KhR_{Lee}(S)(x_{\mathfrak{o}}) = 2^{n(S)} \sum_{\mathfrak{D}|_{L=\mathfrak{o}}} (-1)^{n(S_-(\mathfrak{D}))} x_{\mathfrak{D}|_{L'}} \quad (10)$$

where  $\mathfrak{D}$  runs over orientations on  $S$  that are compatible with the orientation  $\mathfrak{o}$  (thus each  $(S, \mathfrak{D})$  is an oriented cobordism from  $(L, \mathfrak{o})$  to  $(L', \mathfrak{D}|_{L'})$ ),  $S_-(\mathfrak{D})$  is the union of components of  $S$  whose orientation disagree with  $\mathfrak{D}$ , and

$$n(\Sigma) = -\frac{\chi(\Sigma) + \#\partial_+\Sigma - \#\partial_-\Sigma}{2} \in \mathbb{Z} \quad (11)$$

for a surface  $\Sigma$  with positive boundary  $\partial_+\Sigma$  and negative boundary  $\partial_-\Sigma$ . In particular,  $n(\cdot)$  is additive under gluing surfaces.

The  $\mathfrak{gl}_2$   $s$ -invariant of  $L$  is defined to be

$$s_{\mathfrak{gl}_2}(L) := q(x_{\mathfrak{o}_L}) - 1$$

where  $\mathfrak{o}_L$  is the orientation of  $L$  and  $q: KhR_{Lee}(L) \rightarrow \mathbb{Z} \sqcup \{-\infty\}$  is the quantum filtration function. It relates to the classical  $s$ -invariant of oriented links (as defined in [BW08], after [Ras10]) by

$$s_{\mathfrak{gl}_2}(L) = -s(-L) - w(L). \quad (12)$$

**2.4. Properties for Khovanov and Lee skein lasagna modules.** Letting  $Z = KhR_2$  or  $KhR_{Lee}$  as the TQFT for the skein lasagna module, we recover the Khovanov and Lee skein lasagna modules, denoted  $\mathcal{S}_0^2$  and  $\mathcal{S}_0^{Lee}$ , respectively, the first of which agrees with the construction in [MWW22] (or more concisely in [MN22]) for the Khovanov-Rozansky  $\mathfrak{gl}_2$  skein lasagna module. Since  $KhR_2$  and  $KhR_{Lee}$  allow dots on cobordisms, in the definition of lasagna fillings we may also allow dots on skeins, which does not change the structure of  $\mathcal{S}_0^2$  or  $\mathcal{S}_0^{Lee}$  but makes the description of certain elements easier. One can think of a dot as an  $X$ -decoration on the unknot in the boundary of an extra deleted local 4-ball near the dot.

It follows from our general discussion in Section 2.1 and Section 2.2 that for a pair  $(X, L)$  of 4-manifold  $X$  and framed oriented link  $L \subset \partial X$ , the Khovanov skein lasagna module

$$\mathcal{S}_0^2(X; L) = \bigoplus \mathcal{S}_{0,h,q}^2(X; L; \alpha)$$

is an abelian group with a homological grading by  $h \in \mathbb{Z}$ , a quantum grading by  $q \in \mathbb{Z}$ , and a homology class grading by  $\alpha \in H_2^L(X)$ . Similarly, the Lee skein lasagna module  $\mathcal{S}_0^{Lee}(X; L)$  is a  $\mathbb{Q}$ -vector space with a homological  $\mathbb{Z}$ -grading, a quantum filtration, a compatible quantum  $\mathbb{Z}/4$ -grading, and a homology class grading by  $H_2^L(X)$ . A Khovanov/Lee lasagna filling of  $(X, L)$  is of the form  $(\Sigma, v)$ , where  $\Sigma$  is a skein in  $X$  rel  $L$  with some input balls  $B_i$ , and  $v \in \otimes_i KhR_{\bullet}(\partial\Sigma|_{\partial B_i})$ ,  $\bullet \in \{2, Lee\}$ . If  $v$  is homogeneous with homological degree  $h$  and quantum degree (or filtration degree)  $q$ , then the class represented by  $(\Sigma, v)$  in  $\mathcal{S}_0^{\bullet}(X; L)$  has homological degree  $h$ , quantum degree (or filtration degree at most)  $q - \chi(\Sigma) + 2\#(\text{dots})$ , and homology class degree  $[\Sigma]$ . When we work over a ring  $R$ , write  $\mathcal{S}_0^2(X; L; R) = \oplus_{\alpha} \mathcal{S}_0^2(X; L; \alpha; R)$  for the corresponding Khovanov skein lasagna module.

Next we describe a formula of Khovanov and Lee skein lasagna modules for 2-handlebodies, developed in [MN22]. From now on till the end of this section, suppose  $X$  is a 2-handlebody obtained by attaching 2-handles along a framed oriented link  $K = K_1 \cup \dots \cup K_m \subset S^3 = \partial B^4$  disjoint from a framed oriented link  $L \subset S^3$ . Then  $L$  can be regarded as a link in  $\partial X$ . A class  $\alpha \in H_2^L(X)$  intersects the cocore of the 2-handle on  $K_i$  algebraically some  $\alpha_i$  times, and the assignment  $\alpha \mapsto (\alpha_i)_i$  gives

<sup>9</sup>The equation in [Ras05, Proposition 3.2] missed a  $\frac{1}{2}$  factor on the exponent on the right hand side.

an isomorphism  $H_2^L(X) \cong \mathbb{Z}^m$ . Write  $\alpha_i^+ = \max(0, \alpha_i)$  and  $\alpha_i^- = \max(0, -\alpha_i)$ . Write  $|\cdot|$  for the  $L^1$ -norm on  $\mathbb{Z}^m$ .

For  $k^+, k^- \in \mathbb{Z}_{\geq 0}^m$ , let  $K(k^+, k^-) \cup L \subset S^3$  denote the framed oriented link obtained from  $K \cup L$  by replacing each  $K_i$  by  $k_i^+ + k_i^-$  parallel copies of itself, with orientations on  $k_i^-$  of them reversed.

**Proposition 2.6 (2-handlebody formula).** *Let  $(X, L)$  be given as above. Then*

$$\mathcal{S}_0^\bullet(X; L; \alpha) \cong \left( \bigoplus_{r \in \mathbb{Z}_{\geq 0}^m} \text{KhR}_\bullet(K(\alpha^+ + r, \alpha^- + r) \cup L) \{ -|\alpha| - 2|r| \} \right) / \sim \quad (13)$$

where  $\{m\}$  denotes the shifting of the quantum degree by  $m$ ,  $\bullet \in \{2, \text{Lee}\}$ , and  $\sim$  is the linear equivalence relation generated by

- (i)  $\sigma_i \cdot x \sim x$  for  $x \in \text{KhR}_\bullet(K(\alpha^+ + r, \alpha^- + r) \cup L) \{ -|\alpha| - 2|r| \}$  and  $\sigma_i \in S_{|\alpha_i| + 2r_i}$ , where  $\sigma_i$  acts by permuting the cables for  $K_i$  (cf. [GLW18])<sup>10</sup>;
- (ii)  $\phi_i(x) \sim x$  for  $x \in \text{KhR}_\bullet(K(\alpha^+ + r, \alpha^- + r) \cup L) \{ -|\alpha| - 2|r| \}$  where

$$\begin{aligned} \phi_i: \text{KhR}_\bullet(K(\alpha^+ + r, \alpha^- + r) \cup L) \{ -|\alpha| - 2|r| \} \rightarrow \\ \text{KhR}_\bullet(K(\alpha^+ + r + e_i, \alpha^- + r + e_i) \cup L) \{ -|\alpha| - 2|r| - 2 \} \end{aligned}$$

is the map induced by the dotted annular cobordism that creates two parallel, oppositely oriented cables of  $K_i$  (which is bidegree-preserving), and where  $e_i$  is the  $i^{\text{th}}$  coordinate vector.

Here, when  $\bullet = 2$ , we assume to work over a base ring where 2 is invertible; otherwise,  $\mathcal{S}_0^2(X; L; \alpha)$  is given by a further quotient of the right hand side of (13).

*Proof.* The  $\mathcal{S}_0^2$  version follows from a slight generalization of Theorem 1.1 and Proposition 3.8<sup>11</sup> in [MN22], stated as Theorem 3.2 in [MWW23]. The claim for the case when 2 is not invertible follows from the proof of Proposition 3.8 in [MN22]. The version for  $\mathcal{S}_0^{\text{Lee}}$  has an identical proof.  $\square$

Changing the orientation of a link only affects  $\text{KhR}_\bullet$  by a grading shift, given by (9). Therefore, by (13),  $\mathcal{S}_0^\bullet(X; L)$  is independent of the orientation of  $L$  (as long as it is null-homologous) up to renormalizations, and  $\mathcal{S}_0^\bullet(X; L; \alpha)$  only depends on the mod 2 reduction of  $\alpha \in H_2^L(X) \subset H_2(X, L)$  up to renormalizations. Explicitly, suppose  $\beta \in H_2(X, L)$  has the same mod 2 reduction as  $\alpha \in H_2^L(X)$  such that  $\partial\beta \in H_1(L)$  is the fundamental class of  $L$  with a possibly different orientation. Let  $(L, \partial\beta)$  denote the link  $L$  with orientation given by  $\partial\beta$ . Then  $\beta \in H_2^{(L, \partial\beta)}(X)$ , and

$$\mathcal{S}_0^\bullet(X; (L, \partial\beta); \beta) \cong \mathcal{S}_0^\bullet(X; L; \alpha) \left[ \frac{\alpha^2 - \beta^2}{2} \right] \left\{ \frac{\beta^2 - \alpha^2}{2} \right\}. \quad (14)$$

Here when  $\bullet = 2$  we assume 2 is invertible in the base ring.

The bilinear product on  $H_2^L(X)$  in (14) (similarly for  $H_2^{(L, \partial\beta)}(X)$ ) is defined by counting intersections of two surfaces in  $X$  with boundary  $L$ , one copy of which is pushed off  $L$  via its framing. In particular, it is independent of the choice of  $K \cup L \subset S^3$ .

**Example 2.7.** For the 2-handlebody  $X = S^2 \times D^2$ ,  $K$  is the 0-framed unknot. It follows from Proposition 2.6 that  $\mathcal{S}_0^\bullet(S^2 \times D^2)$  can be computed from the Khovanov or Lee homology of unlinks

<sup>10</sup>Strictly speaking, elements in  $S_{|\alpha_i| + 2r_i}$  may permute cables of  $K_i$  with different orientations, so it does not act on  $\text{KhR}_\bullet(K(\alpha^+ + r, \alpha^- + r) \cup L) \{ -|\alpha| - 2|r| \}$ . Nevertheless, since changing orientation only affects  $\text{KhR}_\bullet$  by a renormalization, we can first work with  $K(\alpha^+ + \alpha^- + 2r, 0) \cup L$  and then renormalize.

<sup>11</sup>The  $\text{Kh}$  in Proposition 3.8 in [MN22] should be  $\text{KhR}_2$  instead.

and dotted cobordism maps between them. Explicitly, [MN22, Theorem 1.2] calculated that for every  $\alpha \in H_2(S^2 \times D^2)$ ,

$$\mathcal{S}_{0,h,q}^2(S^2 \times D^2; \alpha) = \begin{cases} \mathbb{Z}, & h = 0, q = -2k \leq 0 \\ 0, & \text{otherwise.} \end{cases}$$

Similarly,  $gr_q \mathcal{S}_{0,h}^{Lee}(S^2 \times D^2; \alpha)$  is given by the same formula with  $\mathbb{Z}$  replaced by  $\mathbb{Q}$ .

Using Proposition 2.6, [MN22, Theorem 1.3] also obtained partial information on Khovanov skein lasagna modules for  $D^2$ -bundles over  $S^2$  with euler number  $n \neq 0$ , denoted  $D(n)$ , namely that

$$\mathcal{S}_{0,0,*}^2(D(n); 0) = \begin{cases} \mathbb{Z}, & n < 0, * = 0 \\ 0, & \text{otherwise.} \end{cases}$$

For  $n = \pm 1$ , this also yields the corresponding calculation for  $\mathbb{CP}^2, \overline{\mathbb{CP}^2}$ , in view of Proposition 2.1. These results will be improved for  $n > 0$  in Example 3.2, and for  $n < 0$  in Example 6.1, Section 6.8, and Section 6.9.

The set of framings of a link  $L$  in an oriented 3-manifold is affine over  $\mathbb{Z}^{\pi_0(L)}$ . If  $L, L'$  are two framed links with the same underlying link, the writhe difference of them is defined to be  $w(L', L) := \epsilon(L' - L)$  where  $L' - L$  is thought of as an element in  $\mathbb{Z}^{\pi_0(L)}$  and  $\epsilon: \mathbb{Z}^{\pi_0(L)} \rightarrow \mathbb{Z}$  is the augmentation map, which is defined by taking the sum over all coordinates.

**Proposition 2.8 (Framing change).** *If  $L' \subset \partial X$  is a framed oriented link which agrees with  $L \subset \partial X$  upon forgetting the framing, then for  $\bullet \in \{2, Lee\}$ ,*

$$\mathcal{S}_0^\bullet(X; L') \cong \mathcal{S}_0^\bullet(X; L) \{-w(L', L)\}.$$

*Proof.* For a given lasagna filling of  $(X, L)$ , create new input balls near each component of  $L$  that intersect the skein in disks. Change the framing of  $L$  to  $L'$  and push the framing change into the intersection disks. Then the sum of framings on the input unknots is  $w(L', L)$ . Assigning these unknots the Khovanov or Lee class 1 (the element corresponds to  $1 \in KhR_\bullet(U)$  under the Reidemeister I induced isomorphisms, where  $U$  is the 0-framed unknot) produces a lasagna filling of  $(X, L')$ . This assignment induces the claimed isomorphism.  $\square$

Finally, in view of the trick for framing changes in the proof of Proposition 2.8, we can state a slight improvement of the gluing construction (7). In the general setup when gluing  $(X_2, S)$  to  $(X_1, L_1)$  to obtain  $(X, L)$ , in order to get an induced map on skein lasagna modules we needed to assume  $S$  to be framed so that it can be regarded as a skein in  $X_2 \text{ rel } L_2$ . In the Khovanov or Lee skein lasagna module setup, however, we can drop the framing assumption, at the expense of deleting extra input balls to compensate. Thus, for any (unframed) surface  $S \subset X_2$  with the previous assumptions, there is a gluing morphism

$$\mathcal{S}_0^\bullet(X_2; S): \mathcal{S}_0^\bullet(X_1; L_1) \rightarrow \mathcal{S}_0^\bullet(X; L), \tag{15}$$

which has homological degree 0 and quantum (filtration) degree  $-\chi(S) + \chi(T_0) - [S]^2$ . If  $S$  is already framed, this map agrees with the previous construction.

**2.5. Structure of space of skeins.** The skein lasagna module  $\mathcal{S}_0^Z(X; L)$  decomposes as a direct sum over  $\Omega_{\geq 0}(X; L) := \pi_0(\mathcal{C}(X; L))$ , the set of connected components of  $\mathcal{C}(X; L)$ . We call  $\Omega_{\geq 0}(X; L)$  the *space of skeins* in  $X \text{ rel } L$ . In this section we show that  $\Omega_{\geq 0}(X; L)$  is no finer than  $H_2^L(X)$ .

**Theorem 2.9.** *Taking homology defines an isomorphism*

$$\Omega_{\geq 0}(X; L) \xrightarrow{\cong} H_2^L(X). \tag{16}$$

We recall that  $\Omega_{\geq 0}(X; L)$  is the set of (properly embedded, oriented) framed surfaces  $\Sigma \subset X \setminus \sqcup_i \text{int}(B_i)$  with  $\partial\Sigma|_{\partial X} = L$  up to

- (a) isotopy rel boundary;
- (b) enclosing some small balls by some large balls and deleting the interior of the large balls, or its converse.

Although

- (c) isotopy of input balls in  $X$  (that drags surfaces along)

is not directly allowed in the relations, it can be recovered using a combination of (a)(b).

**Lemma 2.10.** *The relations (a),(b) between skeins are equivalent to the relation of framed cobordisms in  $X \times I$  between such skeins away from the tubular neighborhood of a 1-dimensional complex, relative to  $L \subset \partial X$ .*

*Proof.* One direction is clear: (a) is clearly contained in the relation of framed cobordisms, and (b) is realized by taking the identity cobordism but gradually enlarging the small deleted balls so that they merge with each other and become the large balls.

For the converse, after an isotopy we can decompose every framed cobordism of surfaces rel  $L \subset \partial X$  away from a 1-complex into the following elementary moves:

- (i) Isotopy of the surface and deleted balls rel  $\partial X$ ;
- (ii) Morse move away from deleted balls;
- (iii) Pushing a Morse critical point into one deleted ball;
- (iv) Deleting a local ball away from the surface, or its converse;
- (v) Choosing a path  $\gamma$  between two deleted balls  $B_1, B_2$  disjoint from the surface and replacing  $B_1, B_2$  by a small tubular neighborhood of  $B_1 \cup B_2 \cup \gamma$ , or its converse.

We see immediately that (i) is realized by (a) and (c), while (iii),(iv), and (v) are realized by (b). Since a Morse move is local, we can use (b) twice to delete a local ball and reglue it back to realize (ii).  $\square$

*Proof of Theorem 2.9.* First we show the map is surjective. Any class  $\alpha \in H_2^L(X)$  can be represented by an oriented embedded surface  $\Sigma \subset X$  with  $\partial\Sigma = L$ . The obstruction to putting a framing on  $\Sigma$  lies in  $H^2(\Sigma, L)$ . Once we delete local 4-balls at points on each component of  $\Sigma$ , this obstruction vanishes and we get a skein  $\Sigma$  with  $[\Sigma] = \alpha$ .

Next we show the map is injective. Suppose two skeins  $\Sigma_0, \Sigma_1$  have the same homology class. By enclosing all input balls by one large ball, we may assume  $\Sigma_0, \Sigma_1 \in X^\circ := X \setminus \text{int}(B^4)$  for some fixed ball  $B^4$  in  $X$ . Since  $[\Sigma_1] - [\Sigma_0] = 0 \in H_2(X^\circ \times I, \partial X \times I)$ , we can find a properly embedded immersed oriented 3-manifold  $Y \subset X^\circ \times I$  that cobounds  $\Sigma_0, \Sigma_1$ . By transversality we may assume the self-intersection of  $Y$  is a properly embedded 1-manifold in  $X^\circ \times I$ . Deleting a tubular neighborhood along the self intersection makes  $Y$  an embedded cobordism between  $\Sigma_0$  and  $\Sigma_1$  in  $X \times I$  with the tubular neighborhood of a 1-complex deleted. The obstruction of putting a compatible framing on  $Y$  lies in  $H^2(Y, \partial Y|_{\partial(X \times I)}) \cong H_1(Y, \partial Y \setminus \partial Y|_{\partial(X \times I)})$ , which can be made zero upon a further deletion of the tubular neighborhood of a 1-complex in  $X^\circ \times I$ .  $\square$

*Remark 2.11.* An alternative proof of Theorem 2.9 can be deduced from a relative and noncompact version of [KMT12, Theorem 2] applied to  $(X^\circ, L)$  together with the fact that a framed link translation (as defined in their paper) can be undone in  $\Omega_{\geq 0}(X; L)$  by a neck-cutting. However, for our purpose in Section 4, we need a similar result for the space of a pair of skeins, which can



be proved in the same way as we demonstrated, but is inaccessible by directly applying [KMT12, Theorem 2].

**2.6. Variations of the construction.** In this section we assume  $X$  to be connected. The skein lasagna module  $\mathcal{S}_0^Z(X; L)$  is the colimit of  $Z$  along the category  $\mathcal{C}_{\geq 0}(X; L) := \mathcal{C}(X; L)$  of skeins. One can restrict to the subcategory  $\mathcal{C}_1(X; L) \subset \mathcal{C}_{\geq 0}(X; L)$  of skeins with exactly 1 input ball. The colimit of  $Z$  along  $\mathcal{C}_1(X; L)$  produces a slightly different version of skein lasagna module, say  $\mathcal{S}'_0^Z(X; L)$ . It comes with a map

$$\mathcal{S}'_0^Z(X; L) \twoheadrightarrow \mathcal{S}_0^Z(X; L) \tag{17}$$

which is surjective since every object in  $\mathcal{C}_{\geq 0}(X; L)$  has a successor in  $\mathcal{C}_1(X; L)$ .

In fact, (17) is an isomorphism whenever  $X$  has no 1-handles. This is because the relevant formulas of  $\mathcal{S}_0^Z$  for attaching 2, 3, 4-handles (generalized appropriately for 2-handles when  $Z \neq KhR_\bullet$ ), developed by Manolescu-Walker-Wedrich [MWW23], still hold for  $\mathcal{S}'_0^Z$ . In particular, most calculations done in this paper, including the proof of Theorem 1.1, remain valid with  $\mathcal{S}_0^\bullet$  replaced by  $\mathcal{S}'_0^\bullet$ ,  $\bullet \in \{2, Lee\}$ .

However, when  $H_1(X) \neq 0$ , (17) is usually not an injection. In fact, a relative and noncompact version of [KMT12, Theorem 2] shows that the relevant space of skeins  $\Omega_1(X; L) := \pi_0(\mathcal{C}_1(X; L))$  admits a surjection onto  $H_2^L(X) \cong \Omega_{\geq 0}(X; L)$  by taking homology, whose fiber is usually nontrivial.

We illustrate the problem in the following example. Take  $Z = KhR_2$  and let  $K$  be a knot. Then the lasagna filling  $(K \times S^1, 1)$  of  $S^3 \times S^1$  (with any framing on  $K \times S^1$ ) is equivalent to the lasagna filling  $(\emptyset, 2)$  by a neck-cutting relation in  $\mathcal{S}_0^2$  [MN22, Lemma 7.2]. However, the neck-cutting relation no longer holds for  $\mathcal{S}'_0^2$ , so  $(K \times S^1, 1)$  cannot be simplified there in general. More precisely, if the framing on  $K \times S^1$  has twists along the  $K$  factor, then  $K \times S^1$  and  $\emptyset$  do not represent the same element in  $\Omega_1(S^3 \times S^1)$  [KMT12], so  $(K \times S^1, 1)$  is not equivalent to  $(\emptyset, 2)$  in  $\mathcal{S}'_0^2(S^3 \times S^1)$ , as the latter represents a class that is nonzero in the quotient  $\mathcal{S}_0^2(S^3 \times S^1)$  [MWW23, Corollary 4.2].

Since the proof for the connected sum formula (Proposition 2.2) requires a neck-cutting, we also lose it for  $\mathcal{S}'_0^2$ . Thus, although  $\mathcal{S}'_0^Z$  is a refinement of  $\mathcal{S}_0^Z$ , its computation could be more challenging. We suspect that when  $X$  is simply connected,  $\mathcal{C}_1(X; L)$  is cofinal in  $\mathcal{C}_{\geq 0}(X; L)$  and thus (17) is an isomorphism.

One may also consider versions of skein lasagna modules with various conditions on framings and orientations. For example, if one drops the assumption that surfaces are framed, one can obtain a skein lasagna module for every input TQFT of unframed oriented links in  $S^3$ . The underlying space of skeins in  $X$  rel  $L$ , no matter whether one allows only one input ball or finitely many (or none), is again isomorphic to  $H_2^L(X)$ . The many-balls version of the unframed skein lasagna module for the Khovanov or Lee TQFT seems to agree with its framed counterpart up to renormalizations, in view of the framing-change trick in the proof of Proposition 2.8. In fact, it could be notationally simpler to work with this version throughout the paper, but we follow the existing convention. The structure of the one-ball version of the unframed Khovanov skein lasagna module is less clear.

### 3. VANISHING CRITERION AND EXAMPLES

In this section we prove Theorem 1.4 and give a few examples where the theorem applies. The proof is a straightforward application of Ng's maximal Thurston-Bennequin bound from Khovanov homology [Ng05] and the 2-handlebody formula (13).

**Lemma 3.1** ([Ng05, Theorem 1]). *For an oriented link  $L$ , the Khovanov homology  $Kh^{h,q}(L)$  is supported in the region*

$$TB(L) \leq q - h \leq -TB(-L).$$

*Proof of Theorem 1.4.* It suffices to prove the case  $X = X_n(K)$ ,  $n \geq -TB(-K)$ , since the general case follows from it and Proposition 2.3 applied to  $X_1 = X_n(K)$ ,  $X_2 = X \setminus \text{int}(X_1)$ .

Give  $K$  an orientation, and suppose  $L \subset S^3$  is a framed oriented link disjoint from  $K$  that gives rise to a link  $L \subset \partial X_n(K)$ . Since  $-n \leq TB(-K)$ , we can find a Legendrian representative  $\mathcal{K} \cup \mathcal{L}$  of  $-K \cup -L$  with Thurston-Bennequin number  $tb(\mathcal{K}) = -n$ . Let  $\mathcal{K}(k^+, k^-)$  denote the Legendrian  $(k^+ + k^-)$ -cable of  $\mathcal{K}$  with the orientation on  $k^-$  of the strands reversed. Then the underlying link type of  $\mathcal{K}(k^+, k^-) \cup \mathcal{L}$  is  $-(K(k^+, k^-) \cup L)$  where  $K$  has framing  $n$ . As a framed link with the contact framing, we have

$$tb(\mathcal{K}(k^+, k^-) \cup \mathcal{L}) = w(\mathcal{K}(k^+, k^-) \cup \mathcal{L}) = -w(K(k^+, k^-) \cup L).$$

By Lemma 3.1,  $Kh(-(K(k^+, k^-) \cup L))$  is supported in  $q - h \geq -w(K(k^+, k^-) \cup L)$ . By (8),  $KhR_2(K(k^+, k^-) \cup L)$  is supported in  $q + h \leq w(K(k^+, k^-) \cup L) - w(K(k^+, k^-) \cup L) = 0$ . Now Proposition 2.6 implies  $\mathcal{S}_0^2(X; L; \alpha) = 0$  for any  $\alpha$ , as  $KhR_2(K(r + \alpha^+, r + \alpha^-) \cup L)\{-|\alpha| - 2r\}$  is supported in  $q + h \leq -|\alpha| - 2r \rightarrow -\infty$  as  $r \rightarrow \infty$  yet the map  $\phi$  in (ii) preserves the grading  $q + h$ .  $\square$

**Example 3.2.** The unknot  $K = U$  has maximal  $tb$  number  $-1$ . The  $n$ -trace on  $U$  is  $D(n)$ , the  $D^2$ -bundle over  $S^2$  with euler number  $n$ , which is also a tubular neighborhood of an embedded sphere in a 4-manifold with self-intersection  $n$ . Therefore in this case, Theorem 1.4 says  $\mathcal{S}_0^2(X; L) = 0$  whenever  $X$  contains an embedded sphere with positive self-intersection. This is an adjunction-type obstruction for  $\mathcal{S}_0^2$  to be nontrivial.

For example,  $D(n)$  with  $n > 0$ ,  $\mathbb{C}\mathbb{P}^2$ ,  $S^2 \times S^2$ , or the mirror image of the  $K3$  surface have vanishing Khovanov skein lasagna modules.

**Example 3.3.** The right handed trefoil  $-3_1$  has  $TB(-3_1) = 1$ . Therefore, the  $(\pm 1)$ -trace on  $3_1$  both have vanishing Khovanov skein lasagna modules. The boundary of this 4-manifold is the Poincaré homology sphere  $\Sigma(2, 3, 5)$  for  $-1$ , and the mirror image of the Brieskorn sphere  $\Sigma(2, 3, 7)$  for  $+1$ .

One way to see embedded traces is through a Kirby diagram of 4-manifold  $X$ . If there is an  $n$ -framed 2-handle attached along some knot  $K$  not going over 1-handles, then the 2-handle together with the 4-ball is a copy of  $X_n(K)$  inside  $X$ . Consequently, if we see such a 2-handle with  $n \geq -TB(-K)$ , then  $\mathcal{S}_0^2(X; L) = 0$  for any  $L$ .

*Remark 3.4.* As proved by [MN22, Theorem 1.3] (and will be reproved more generally in Section 6.9),  $\overline{k\mathbb{C}\mathbb{P}^2}$  has nonvanishing skein lasagna module. Therefore, we obtain a slice obstruction from Corollary 1.5, which states that if  $\Sigma$  is a slice disk in  $\overline{k\mathbb{C}\mathbb{P}^2} \setminus \text{int}(B^4)$  for a knot  $K$ , then

$$TB(K) + [\Sigma]^2 < 0.$$

In this special case, however, the obstruction is somewhat weaker than the obstruction coming from Rasmussen  $s$ -invariant from [Ren23, Corollary 1.5] which implies  $s(K) \leq -[\Sigma]^2 - |[\Sigma]|$ . Since  $TB(K) < s(K)$  [Pla06, Proposition 4], this implies  $TB(K) + [\Sigma]^2 + |[\Sigma]| < 0$ .

4. LEE SKEIN LASAGNA MODULES AND LASAGNA  $s$ -INVARIANTS

In this section, we explore the structure of the Lee skein lasagna module for an arbitrary pair  $(X, L)$  of 4-manifold  $X$  and framed oriented link  $L \subset \partial X$ . In particular, we prove the statements in Section 1.2 (except Theorem 1.9) in a more general setup.

Recall from Section 2.4 that the Lee skein lasagna module is defined to be the skein lasagna module with  $KhR_{Lee}$  as the TQFT input. For a pair  $(X, L)$ , its Lee skein lasagna module  $\mathcal{S}_0^{Lee}(X; L)$  is a vector space over  $\mathbb{Q}$  with a homological  $\mathbb{Z}$ -grading, a quantum  $\mathbb{Z}/4$ -grading, a homology class grading by  $H_2^L(X)$ , and an (increasing) quantum filtration (so that 0 has filtration degree  $-\infty$ ). Since the filtration degree of an element can decrease under a filtered map, a nonzero element in the colimit  $\mathcal{S}_0^{Lee}(X; L)$  may also have filtration degree  $-\infty$ . Let  $q: \mathcal{S}_0^{Lee}(X; L) \rightarrow \mathbb{Z} \sqcup \{-\infty\}$  denote the quantum filtration function.

**4.1. Structure of Lee skein lasagna module.** For a pair  $(X, L)$ , let  $L_i$  be the components of  $L$  and  $[L_i] \in H_1(L)$  be the fundamental class of  $L_i$ . Let  $\partial: H_2(X, L) \rightarrow H_1(L)$  be the boundary homomorphism. Define the set of *double classes* in  $X$  rel  $L$  to be

$$H_2^{L, \times 2}(X) := \left\{ (\alpha_+, \alpha_-) \in H_2(X, L)^2 : \partial \alpha_{\pm} = \sum_i \epsilon_{i, \pm} [L_i], \epsilon_{i, \pm} \in \{0, 1\}, \epsilon_{i, +} + \epsilon_{i, -} = 1 \right\}.$$

Thus for any double class  $(\alpha_+, \alpha_-)$  we have  $\alpha_+ + \alpha_- \in H_2^L(X)$ .

The purpose of this section is to prove the following generalization of Theorem 1.6.

**Theorem 4.1.** *The Lee skein lasagna module of  $(X, L)$  is  $\mathcal{S}_0^{Lee}(X; L) \cong \mathbb{Q}^{H_2^{L, \times 2}(X)}$ , with a basis consisting of canonical generators  $x_{\alpha_+, \alpha_-}$ , one for each double class  $(\alpha_+, \alpha_-) \in H_2^{L, \times 2}(X)$ . Moreover,*

- (1)  $x_{\alpha_+, \alpha_-}$  has homological degree  $-2\alpha_+ \cdot \alpha_-$ ;
- (2)  $x_{\alpha_+, \alpha_-}$  has homology class degree  $\alpha := \alpha_+ + \alpha_-$ ;
- (3)  $x_{\alpha_+, \alpha_-} + x_{\alpha_-, \alpha_+}$ ,  $x_{\alpha_+, \alpha_-} - x_{\alpha_-, \alpha_+}$  have quantum  $\mathbb{Z}/4$  degrees  $-\alpha^2 + \#L$ ,  $-\alpha^2 + \#L + 2$ , respectively;
- (4)  $q(x_{\alpha_+, \alpha_-}) = q(x_{\alpha_-, \alpha_+}) = \max(q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+}))$ , which also equals  $\min(q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+})) + 2$  if  $\alpha_+ \neq \alpha_-$  in  $H_2(X, L; \mathbb{Q})$  (in particular if  $L \neq \emptyset$ ).

Here, the bilinear product on  $H_2(X, L)$  is defined using the framing of  $L$ .

When  $X = B^4$ , the bilinear product computes the linking numbers of the boundary links and thus in view of Proposition 2.4, Theorem 4.1 recovers classical structural results on  $KhR_{Lee}(L)$  by Lee [Lee05, Proposition 4.3] and Rasmussen [Ras10, Proposition 2.3, 3.3, Lemma 3.5] [BW08, Section 6.1].

We define a *double skein* in  $X$  rel  $L$  to be a skein  $\Sigma$  together with a partition of its components  $\Sigma = \Sigma_+ \cup \Sigma_-$ . Equivalently, a double skein is a pair  $(\Sigma, \mathfrak{D})$  where  $\Sigma$  is a skein (which is already oriented) and  $\mathfrak{D}$  is an orientation of  $\Sigma$  as an unoriented surface. The equivalence is given by defining  $\Sigma_+$  to be the union of components of  $\Sigma$  on which  $\mathfrak{D}$  matches the given orientation of  $\Sigma$ . A double skein  $\Sigma = \Sigma_+ \cup \Sigma_-$  represents a double class  $([\Sigma_+], [\Sigma_-]) \in H_2^{L, \times 2}(X)$ .

The *canonical Lee lasagna filling* of a double skein  $(\Sigma, \mathfrak{D})$  is the lasagna filling  $x(\Sigma, \mathfrak{D}) = x(\Sigma_+, \Sigma_-) \in KhR_{Lee}(\Sigma)$  defined as follows. Let  $B_i$  be the input balls for  $\Sigma$  and  $K_i = \partial \Sigma|_{\partial B_i}$  be the input links. Assign  $K_i$  the (rescaled) canonical generator  $x_{\mathfrak{D}|_{K_i}} \in KhR_{Lee}(K_i)$ , and put an extra decoration  $\mathbf{a}$  on each component of  $\Sigma_+$  and  $\mathbf{b}$  on each component of  $\Sigma_-$  ( $\mathbf{a}, \mathbf{b}$  are linear combinations of 1 and

the dot decoration; cf. Section 2.3 and Section 2.4 for relevant notations). Define  $x(\Sigma_+, \Sigma_-)$  to be the resulting lasagna filling rescaled by  $2^{-n(\Sigma)}(-1)^{-n(\Sigma_-)}$  where  $n(\cdot)$  is defined as in (11).

Equivalently, the canonical Lee lasagna filling  $x(\Sigma, \mathfrak{D}) = x(\Sigma_+, \Sigma_-)$  can be thought of as obtained by first deleting a local ball on each component of  $\Sigma$ , then assigning  $x_{\mathfrak{D}|_{K_i}}$  to each input link  $K_i$  (which now includes an unknot for each component of  $\Sigma$ ) and rescaling by  $2^{-n(\Sigma^\circ)}(-1)^{-n(\Sigma^\circ)}$ , where  $\Sigma^\circ$  is  $\Sigma$  with local disks deleted. (Strictly speaking this is not a lasagna filling with skein  $\Sigma$  anymore, but on the level of  $\mathcal{S}_0^{Lee}$  they define the same element. The difference will be insignificant for our discussion.)

**Lemma 4.2.** *Let  $\Sigma$  be a skein in  $X$  rel  $L$ . Every element in  $KhR_{Lee}(\Sigma)$  is equivalent in  $\mathcal{S}_0^{Lee}(X; L)$  to a linear combination of canonical Lee lasagna fillings of  $\Sigma$  with various double skein structures.*

*Proof.* Let  $K_i$  be the input links for  $\Sigma$ . Every element in  $KhR_{Lee}(\Sigma)$  (possibly with decorations on surfaces) can be written as a linear combination of ‘‘pure Lee lasagna fillings,’’ which are by definition the ones with a decoration  $x_{\mathfrak{o}_i}$  on each  $K_i$  (for some orientation  $\mathfrak{o}_i$  of  $K_i$  as an unoriented link), and a decoration of  $\mathbf{a}$  or  $\mathbf{b}$  on each component of  $\Sigma$ . Now we claim that such a pure Lee lasagna filling is 0 in  $\mathcal{S}_0^{Lee}(X; L)$  unless it is a (multiple of a) canonical Lee lasagna filling for some double skein structure on  $\Sigma$ .

If a pure Lee lasagna filling does not come from a double skein structure on  $\Sigma$ , then the orientation on  $\Sigma$  defined by the standard orientation on  $\mathbf{a}$ -decorated components and the reverse orientation on the  $\mathbf{b}$ -decorated components is not compatible with  $\mathfrak{o}_i$  for some  $i$ . Take a component  $\Sigma_k$  of  $\Sigma$  and a knot component  $K_{ij} \subset K_i$  where the orientation is incompatible. Think of the  $\mathbf{a}/\mathbf{b}$  decoration on  $\Sigma_k$  to be a multiple of the Lee canonical generator of a local unknot  $U$  on  $\partial\Sigma_k^\circ$ . Then  $U$  and  $K_{ij}$  have incompatible orientations. Choose a path  $\gamma$  in  $\Sigma_k^\circ$  from a point on  $U$  to a point on  $K_{ij}$ . Then deleting a tubular neighborhood of  $\gamma$  induces a map  $K_i \sqcup U \rightarrow K_i$  by connect-summing  $U$  and  $K_{ij} \subset K_i$ , which maps the chosen Lee lasagna filling to zero since the corresponding Lee canonical generators on  $U$  and  $K_i$  are induced by incompatible orientations.  $\square$

If  $[S]: \Sigma \rightarrow \Sigma'$  is a morphism in the category of skeins  $\mathcal{C}(X; L)$ , then a double skein structure  $\mathfrak{D}$  on  $\Sigma$  induces one on  $\Sigma'$  by restricting the orientation, and  $([\Sigma_+], [\Sigma_-]) = ([\Sigma'_+], [\Sigma'_-]) \in H_2^{L, \times 2}(X)$ .

**Lemma 4.3.** *If  $\Sigma = \Sigma_+ \cup \Sigma_-$  is a double skein in  $X$  rel  $L$  and  $[S]: \Sigma \rightarrow \Sigma'$  is a morphism in  $\mathcal{C}(X; L)$  inducing the double skein structure  $\Sigma'_+ \cup \Sigma'_-$  on  $\Sigma'$ , then*

$$KhR_{Lee}([S])(x(\Sigma_+, \Sigma_-)) = x(\Sigma'_+, \Sigma'_-) + \dots$$

where the term  $\dots$  is zero in  $\mathcal{S}_0^{Lee}(X; L)$ . Moreover, there is no  $\dots$  term if every component of  $S$  has a boundary on some input ball of  $\Sigma$ .

*Proof.* This is a consequence of (10). More explicitly, we need to calculate the image of  $\otimes x_{\mathfrak{D}|_{K_i}}$  under the cobordism  $KhR_{Lee}([S]): \otimes KhR_{Lee}(K_i) \rightarrow \otimes KhR_{Lee}(K'_j)$  and keep track of the extra decorations on surfaces. We may assume, by further decomposing  $S$ , that either every component of  $S$  has some boundary in  $\Sigma'$ , or  $S$  is a single cap.

We first deal with the first case which is more complicated. To keep track of the decorations on surfaces, suppose a component  $\Sigma_k \subset \Sigma$  splits into  $n_k > 0$  components in  $\Sigma'$ , then we rewrite its decoration as  $\mathbf{a} = \mathbf{a}^{n_k}$  or  $\mathbf{b} = \mathbf{b}^{n_k}$  and push each copy of  $\mathbf{a}$  or  $\mathbf{b}$  into one corresponding component in  $\Sigma'$ .

By (10),

$$KhR_{Lee}([S])(\otimes x_{\mathfrak{D}|K_i}) = 2^{n(S)}(-1)^{n(S_-)}(\otimes x_{\mathfrak{D}|K'_j}) + \dots$$

where  $S = S_+ \cup S_-$  is induced from  $\Sigma = \Sigma_+ \cup \Sigma_-$ , and  $\dots$  is a linear combination of lasagna fillings incompatible with the double skein structure  $\Sigma'_+ \cup \Sigma'_-$ , thus is zero in  $\mathcal{S}_0^{Lee}(X; L)$  by the proof of Lemma 4.2. If every component of  $S$  has a boundary on some input ball, there is only one compatible orientation  $\mathfrak{D}$  for  $S$  in (10), thus there is no  $\dots$  term.

It now follows that

$$\begin{aligned} & KhR_{Lee}([S])(x(\Sigma_+, \Sigma_-)) \\ &= 2^{-n(\Sigma)}(-1)^{-n(\Sigma_-)} 2^{n(S)}(-1)^{n(S_-)} 2^{n(\Sigma')}(-1)^{n(\Sigma'_-)} x(\Sigma'_+, \Sigma'_-) + \dots \\ &= x(\Sigma'_+, \Sigma'_-) + \dots, \end{aligned}$$

where the additivity of  $n(\cdot)$  ensures  $n(\Sigma) = n(\Sigma' \cup S) = n(\Sigma') + n(S)$  (and similarly for  $n(\Sigma_-)$ ).

Next we consider the case when  $S$  is a cap that annihilates an unknot component in some  $K_i$  (with an identity cobordism in other link components, which we may assume to be empty since it contributes an identity map in a tensor summand). Thus  $\Sigma = S$  is a disk,  $\Sigma' = \emptyset$ , and  $n(\Sigma) = n(S) = n(\Sigma') = 0$ . If  $\Sigma_+ = \Sigma$ , then  $x(\Sigma_+, \Sigma_-) = 2\mathbf{a} \otimes \mathbf{a}$  (one of  $\mathbf{a}$  is the decoration on  $\Sigma$ ) is mapped to  $\epsilon(2\mathbf{a} \cdot \mathbf{a}) = 1 = x(\Sigma'_+, \Sigma'_-)$ . If  $\Sigma_- = \Sigma$ , then  $x(\Sigma_+, \Sigma_-) = -2\mathbf{b} \otimes \mathbf{b}$  is mapped to  $\epsilon(-2\mathbf{b} \cdot \mathbf{b}) = 1 = x(\Sigma'_+, \Sigma'_-)$ .  $\square$

Now we are ready to prove Theorem 4.1.

*Proof of Theorem 4.1.* We define an augmentation map  $\varepsilon: \mathcal{S}_0^{Lee}(X; L) \rightarrow \mathbb{Q}^{H_2^{L, \times 2}(X)}$  by demanding it to map the class represented by a canonical Lee lasagna filling  $x(\Sigma_+, \Sigma_-)$  to  $e_{([\Sigma_+], [\Sigma_-])}$ , the generator of the  $([\Sigma_+], [\Sigma_-])^{\text{th}}$  coordinate of the codomain. Lemma 4.2 and Lemma 4.3 guarantee that  $\varepsilon$  is well-defined.

We show  $\varepsilon$  is an isomorphism. Any pair  $(\alpha_+, \alpha_-) \in H_2^{L, \times 2}(X)$  can be represented by properly embedded immersed surfaces  $\Sigma_{\pm} \subset X$  with  $\partial \Sigma_{\pm}$  forming a partition of  $L \subset X$  as an oriented unframed link. By transversality we may assume all singularities of  $\Sigma = \Sigma_+ \cup \Sigma_-$  are transverse double points. Deleting balls around these singularities makes  $\Sigma$  embedded in  $X$  with some balls deleted, and after deleting one extra local ball on each component of  $\Sigma$  we can put a framing on  $\Sigma$  compatibly with  $L$ . Now the class represented by the Lee canonical lasagna filling  $x(\Sigma_+, \Sigma_-)$  is mapped to  $e_{(\alpha_+, \alpha_-)}$ , proving the surjectivity of  $\varepsilon$ .

We remark that the above proof for surjectivity is a replica of that of (16) with minor changes. We do the same for injectivity but skip the details. It suffices to show any  $x(\Sigma_+, \Sigma_-)$ ,  $x(\Sigma'_+, \Sigma'_-)$  with  $([\Sigma_+], [\Sigma_-]) = ([\Sigma'_+], [\Sigma'_-]) \in H_2^{L, \times 2}(X)$  represent the same element in  $\mathcal{S}_0^{Lee}(X; L)$ . By mimicking the proof of Theorem 2.9, we see that  $\Sigma = \Sigma_+ \cup \Sigma_-$  and  $\Sigma' = \Sigma'_+ \cup \Sigma'_-$  are related by morphisms in  $\mathcal{C}(X; L)$  (or their reverses) compatible with the double skein structures. Namely, we can find a zigzag of morphisms in  $\mathcal{C}(X; L)$

$$\Sigma = \Sigma_0 \leftarrow \Sigma_1 \rightarrow \Sigma_2 \leftarrow \dots \rightarrow \Sigma_{2k} = \Sigma'$$

and double skein structures on each  $\Sigma_i$  compatible with  $\Sigma, \Sigma'$  and the morphisms. Take the corresponding canonical Lee lasagna fillings and apply Lemma 4.3, we obtain a zigzag

$$x(\Sigma_+, \Sigma_-) = x(\Sigma_{0+}, \Sigma_{0-}) \leftarrow x(\Sigma_{1+}, \Sigma_{1-}) \mapsto \dots \mapsto x(\Sigma_{2k+}, \Sigma_{2k-}) = x(\Sigma'_+, \Sigma'_-)$$

up to lasagna fillings that are zero in  $\mathcal{S}_0^{Lee}(X; L)$ . This proves the injectivity of  $\varepsilon$ .

The canonical generator  $x_{\alpha_+, \alpha_-}$  of  $\mathcal{S}_0^{Lee}(X; L)$  can now be defined by  $[x(\Sigma_+, \Sigma_-)]$  for any double skein  $\Sigma = \Sigma_+ \cup \Sigma_-$  representing  $(\alpha_+, \alpha_-)$ . It remains to prove the four addenda to the theorem. (2) is clear since  $[\Sigma] = [\Sigma_+] + [\Sigma_-] = \alpha_+ + \alpha_- = \alpha$ . Denote by  $K_i$ 's the input links of  $\Sigma$ , which also admit decompositions  $K_{i+} \cup K_{i-}$  induced from the decomposition  $\Sigma_+ \cup \Sigma_-$ . The Lee canonical generator  $x_{\mathfrak{D}|K_i}$  has homological degree  $-2lk(K_{i+}, K_{i-})$ . This proves (1) since  $\sum lk(K_{i+}, K_{i-}) = [\Sigma_+] \cdot [\Sigma_-]$ . The class  $x_{\alpha_-, \alpha_+}$  is represented by the reverse double skein  $\Sigma = \Sigma_- \cup \Sigma_+$ , which comes from canonical generators  $x_{\overline{\mathfrak{D}}|K_i}$ , which are conjugates of  $x_{\mathfrak{D}|K_i}$ , meaning that they are equal in quantum  $\mathbb{Z}/4$  degree  $-w(K_i) + \#K_i$  and negatives of each other in degree  $-w(K_i) + \#K_i + 2$ . The decorations  $\mathbf{a}, \mathbf{b}$  (which are equal in degree 0 and negatives of each other in degree 2) also exchange under reversing the double skein structure. Taking into account the extra renormalization factor  $(-1)^{n(\cdot)}$  and the quantum shift  $-\chi(\Sigma)$ , we see that  $x(\Sigma_+, \Sigma_-)$  and  $x(\Sigma_-, \Sigma_+)$  agree in quantum  $\mathbb{Z}/4$  degree

$$\begin{aligned} & \sum (-w(K_i) + \#K_i) - \chi(\Sigma) - 2(n(\Sigma_+) + n(\Sigma_-)) \\ &= (-\alpha^2 + \#\partial_- \Sigma) - \chi(\Sigma) + (\chi(\Sigma) + \#L - \#\partial_- \Sigma) \\ &= -\alpha^2 + \#L, \end{aligned}$$

proving (3).

Finally we prove (4) which is more subtle. If  $q(x_{\alpha_+, \alpha_-}) = q > -\infty$ , we can find a Lee lasagna filling  $(\Sigma, v)$  representing  $x_{\alpha_+, \alpha_-}$  where  $v \in KhR_{Lee}(\Sigma) = \otimes KhR_{Lee}(K_i)\{-\chi(\Sigma)\}$  has  $q(v) = q$  and  $h(v) = h(x_{\alpha_+, \alpha_-}) = -2\alpha_+ \cdot \alpha_-$ . By deleting extra balls if necessary, we may assume every component of  $\Sigma$  has a boundary on some input ball. Define a conjugation action  $\iota$  on  $KhR_{Lee}(\Sigma)$  by 1 on the quantum  $\mathbb{Z}/4$  degree  $-\alpha^2 + \#L$  part and  $-1$  on the other part.

Write  $v = \sum_{\mathfrak{o}} v_{\mathfrak{o}} = \sum_{\mathfrak{o}} \lambda_{\mathfrak{o}} (\otimes_i x_{\mathfrak{o}_i})$  as a linear combination of Lee canonical generators, where  $\mathfrak{o} = (\mathfrak{o}_i)$  runs over orientations of  $K_i$ 's. Each term  $v_{\mathfrak{o}}$  represents  $c(\mathfrak{o})x_{[\Sigma_+(\mathfrak{D})], [\Sigma_-(\mathfrak{D})]}$  for some constant  $c(\mathfrak{o})$  if  $\mathfrak{o}$  comes from a double skein structure  $\mathfrak{D}$  on  $\Sigma$ , and zero otherwise. In the first case,  $[\iota v_{\mathfrak{o}}] = c(\mathfrak{o})x_{[\Sigma_-(\mathfrak{D})], [\Sigma_+(\mathfrak{D})]}$  by (3). In the second case  $[\iota v_{\mathfrak{o}}] = 0$ . Summing up, we have  $[\iota v] = x_{\alpha_-, \alpha_+}$ , thus  $q(x_{\alpha_-, \alpha_+}) \leq q(\iota v) = q(v) = q(x_{\alpha_+, \alpha_-})$ . Now by symmetry  $q(x_{\alpha_+, \alpha_-}) = q(x_{\alpha_-, \alpha_+})$ . By the triangle inequality this also equals  $\max(q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+}))$ .

Finally, suppose  $\alpha_+ \neq \alpha_-$  in  $H_2(X, L; \mathbb{Q})$  and  $q(x_{\alpha_+, \alpha_-}) = q > -\infty$ , we prove  $\min(q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+})) = q - 2$ . Since  $x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+}$  have different  $\mathbb{Z}/4$  grading, we know  $\min(q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+})) \leq q - 2$ . To show the reverse inequality we resort to the following linear algebra lemma.

**Lemma 4.4.** *Let  $W \subset \mathbb{Q}^n$  be a linear subspace. Suppose  $E_1 \sqcup \dots \sqcup E_m$  is a partition of the cube  $E = \{\pm 1\}^n \subset \mathbb{Q}^n$  such that each  $E_i$  is contained in a coset of  $W$ , and  $E_1 = -E_2 \notin W$ . Suppose  $\lambda: E \rightarrow \mathbb{Q}$  satisfies*

$$\sum_{\epsilon \in E_i} \lambda(\epsilon) = \begin{cases} 1, & i = 1 \\ \pm 1, & i = 2 \\ 0, & i > 2. \end{cases}$$

*Then there exist  $a_1, \dots, a_n \in \mathbb{Q}$  such that*

$$\sum_{k=1}^n \sum_{\epsilon \in E_i} \lambda(\epsilon) \epsilon_k a_k = \begin{cases} 1, & i = 1 \\ \mp 1, & i = 2 \\ 0, & i > 2. \end{cases}$$

*Proof.* Pick any  $\ell = (a_1, \dots, a_n) \in W^\perp \subset (\mathbb{Q}^n)^*$  with  $\ell(E_1) = \{1\}$ . □

We now finish the proof of (4). Choose a lasagna filling  $(\Sigma, v)$  representing  $x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+}$  with  $q(v) = q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+})$ ,  $h(v) = -2\alpha_+ \cdot \alpha_-$ , where  $\pm = +$  or  $-$  is the sign that minimizes  $q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+})$ . Assume as before every component of  $\Sigma$  has a boundary on an input ball. Then we can rewrite

$$v = \sum_{\mathfrak{D}} \lambda_{\mathfrak{D}} x(\Sigma_+(\mathfrak{D}), \Sigma_-(\mathfrak{D})) + \dots \quad (18)$$

where  $\mathfrak{D}$  runs over orientations of  $\Sigma$  with  $[\Sigma_+(\mathfrak{D})] \cdot [\Sigma_-(\mathfrak{D})] = \alpha_+ \cdot \alpha_-$ , and  $\dots$  denotes terms that do not come from a double skein structure on  $\Sigma$  (in particular they represent the zero class in  $\mathcal{S}_0^{Lee}(X; L)$ ). Let  $\Sigma_1, \dots, \Sigma_n$  be the components of  $\Sigma$ . Identify an orientation  $\mathfrak{D}$  on  $\Sigma$  as a vector  $\mathfrak{D} \in E = \{\pm 1\}^n \subset \mathbb{Q}^n$ , with  $\mathfrak{D}_i = 1$  if and only if  $\mathfrak{D}$  agrees with the orientation of  $\Sigma$  on  $\Sigma_i$ ; in this way the coefficients  $\lambda_{\mathfrak{D}}$  in (18) define a function  $\lambda: E \rightarrow \mathbb{Q}$ . Partition  $E$  into  $E_1 \sqcup \dots \sqcup E_m$  according to the value of  $[\Sigma_+(\mathfrak{D})] \in H_2(X, L)$ . Then each  $E_i$  is contained in a coset of  $W = \{(k_1, \dots, k_n) \in \mathbb{Q}^n: \sum_i k_i [\Sigma_i] = 0 \in H_2(X, L; \mathbb{Q})\} \subset \mathbb{Q}^n$ . Assume  $E_1, E_2$  are the parts with  $[\Sigma_+(\mathfrak{D})] = \alpha_+, \alpha_-$ , respectively. Then  $E_1, E_2$  satisfy the condition in Lemma 4.4 (they are not contained in  $W$  as  $\alpha_+ \neq \alpha_- \in H_2(X, L; \mathbb{Q})$ ). Also, the condition  $[v] = x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+}$  is exactly the condition on  $\lambda: E \rightarrow \mathbb{Q}$  in Lemma 4.4. Let  $a_1, \dots, a_n$  be given by the conclusion of the lemma applied to our data  $(W, E_1 \sqcup \dots \sqcup E_m, \lambda)$ . Now consider the operator  $L = \sum_{k=1}^n a_k X_k$  on  $KhR_{Lee}(\Sigma)$ , where  $X_k$  is a dot operation on  $\Sigma_k$ . Since  $X\mathbf{a} = \mathbf{a}$  and  $X\mathbf{b} = -\mathbf{b}$ , it follows that

$$[Lv] = \left[ \sum_{k=1}^n \sum_{\mathfrak{D}} \lambda_{\mathfrak{D}} \mathfrak{D}_k a_k \right] = x_{\alpha_+, \alpha_-} \mp x_{\alpha_-, \alpha_+}.$$

Since the operator  $L$  has filtered degree 2, we conclude that

$$q = q(x_{\alpha_+, \alpha_-} \mp x_{\alpha_-, \alpha_+}) \leq q(x_{\alpha_+, \alpha_-} \pm x_{\alpha_-, \alpha_+}) + 2 = \min(q(x_{\alpha_+, \alpha_-}, x_{\alpha_-, \alpha_+})) + 2,$$

as desired.  $\square$

**4.2. Canonical Lee lasagna generators under morphisms.** Recall that the induced map on Lee homology of a link cobordism is determined by its actions on Lee canonical generators, given by (10). In this section we prove a lasagna generalization of this formula. In its most general form, the relevant cobordism map we need to examine is (15).

We recall the notations and explain the setup before stating the generalization to (10). The 4-manifold  $X = X_1 \cup_Y X_2$  is obtained by gluing  $X_1, X_2$  along some common part  $Y, \bar{Y}$  of their boundaries, where  $Y$  is a 3-manifold possibly with boundary. Two framed oriented tangles  $T_1 \subset \partial X_1 \setminus Y$  and  $T_0 \subset Y$  glue to a framed oriented link  $L_1 \subset \partial X_1$ , and  $S \subset X_2$  is a (not necessarily framed) surface with an oriented link boundary  $\partial S = -T_0 \cup T_2 = L_2 \subset \partial X_2$  where  $T_2$  is a tangle in  $\partial X_2 \setminus \bar{Y}$ . Then  $L = T_1 \cup T_2$  is a framed oriented link in  $\partial X$ . The gluing of  $(X_2, S)$  to  $(X_1, L_1)$  along  $(Y, T_0)$  induces the cobordism map

$$\mathcal{S}_0^{Lee}(X_2; S): \mathcal{S}_0^{Lee}(X_1; L_1) \rightarrow \mathcal{S}_0^{Lee}(X; L) \quad (19)$$

as in (15), which has homological degree 0 and quantum filtration degree  $-\chi(S) + \chi(T_0) - [S]^2$ . We recall that by deleting local balls on  $S$  (one for each component) and putting a compatible framing, the punctured surface  $S^\circ$  is a skein in  $X_2$  rel  $L_2$  with a natural lasagna filling by assigning 1 to all boundary unknots, and (19) is defined by putting the class of this lasagna filling in the second tensor summand of the more general gluing map (6).

A class  $\alpha \in H_2^{L_1}(X_1)$  and a class  $\beta \in H_2^{L_2}(X_2)$  glue to a class  $\alpha * \beta \in H_2^L(X)$ . Similarly, a double class  $(\alpha_+, \alpha_-) \in H_2^{L_1, \times 2}(X_1)$  and a double class  $(\beta_+, \beta_-) \in H_2^{L_2, \times 2}(X_2)$  glue to a double class  $(\alpha_+, \alpha_-) * (\beta_+, \beta_-) = (\alpha_+ * \beta_+, \alpha_- * \beta_-) \in H_2^{L, \times 2}(X)$  if they are compatible, i.e. if  $\partial\alpha_+, \partial\beta_+$  (and thus  $\partial\alpha_-, \partial\beta_-$ ) agree on  $T_0$ , and do not glue if they are incompatible.

The punctured surface  $S^\circ$  represents the class  $[S] \in H_2^{L_2}(X_2)$ . Thus the cobordism map (19) decomposes into direct sums of maps  $\mathcal{S}_0^{Lee}(X_1; L_1; \alpha) \rightarrow \mathcal{S}_0^{Lee}(X; L; \alpha * [S])$ .

**Theorem 4.5.** *The cobordism map (19) is determined by*

$$\mathcal{S}_0^{Lee}(X_2; S)(x_{\alpha_+, \alpha_-}) = 2^{m(S, L_1)} \sum_{\mathfrak{D}} (-1)^{m(S^-(\mathfrak{D}), L_{1-})} x_{\alpha_* * [S_+(\mathfrak{D})], \alpha_* * [S_-(\mathfrak{D})]}. \quad (20)$$

Here the sum runs over double skein structures on  $S$  compatible with  $(\alpha_+, \alpha_-)$ ,  $m(S, L_1)$  is an integral constant depending only on the topological type of the input pair, which equals  $n(S)$  if  $S \cap L_1$  is a link, thought of as the negative boundary of  $S$ , and  $L_{1-}$  is the sublink of  $L_1$  with fundamental class  $\partial\alpha_-$ .

*Proof.* This is almost tautological from the construction. Let  $\Sigma = \Sigma_+ \cup \Sigma_-$  be a double skein in  $X_1$  rel  $L_1$  representing  $(\alpha_+, \alpha_-)$ . Then  $x_{\alpha_+, \alpha_-}$  is represented by the canonical Lee lasagna filling  $x(\Sigma_+, \Sigma_-)$  on the skein  $\Sigma$ , defined by decorating each input link  $K_i$  by  $x_{\mathfrak{D}|_{K_i}}$  and each component of  $\Sigma_+, \Sigma_-$  by  $\mathbf{a}, \mathbf{b}$ , respectively, with an overall scaling factor  $2^{-n(\Sigma)} (-1)^{-n(\Sigma_-)}$ . Gluing in  $S$  produces a skein  $\Sigma \cup S^\circ$  in  $X$  rel  $L$ , which has two types of components: the ones that intersect  $\Sigma$  and the ones that do not. The lasagna filling also extends to  $\Sigma \cup S^\circ$ . The components of the first type already have some  $\mathbf{a}$  or  $\mathbf{b}$  decorations from  $\Sigma$ . If there is one such component with both  $\mathbf{a}$  and  $\mathbf{b}$  decorations, then it represents the zero class in  $\mathcal{S}_0^{Lee}(X; L)$ , while there is no orientation on this component compatible with  $(\alpha_+, \alpha_-)$ , making the right hand side of (20) zero as well. Otherwise, every component of the first type has only  $\mathbf{a}$  or  $\mathbf{b}$  decorations that multiply to  $\mathbf{a}$  or  $\mathbf{b}$ , which also determine a unique orientation on the component compatible with  $(\alpha_+, \alpha_-)$ . Each of the components of the second type has two compatible orientations, with a decoration by  $1 = \mathbf{a} + \mathbf{b}$ . The extra unknots around the punctures of  $S^\circ$  can be thought of as assigned  $\mathbf{a}$  or  $\mathbf{b}$  for the components of the first type, and  $\mathbf{a} + \mathbf{b}$  for those of the second type.

Keeping track of the constants, we see  $\mathcal{S}_0^{Lee}(X_2; S)(x_{\alpha_+, \alpha_-})$  is equal to the sum

$$2^{n(\Sigma \cup S) - n(\Sigma)} \sum_{\mathfrak{D}} (-1)^{n(\Sigma_- \cup S^-(\mathfrak{D})) - n(\Sigma_-)} x_{\alpha_* * [S_+(\mathfrak{D})], \alpha_* * [S_-(\mathfrak{D})]}.$$

If  $T_0$  is a link then  $n(\Sigma \cup S) - n(\Sigma) = n(S)$  by the additivity of  $n(\cdot)$ . In general,

$$n(\Sigma \cup S) - n(\Sigma) = -\frac{\chi(S) - \chi(T_0) + \#L - \#L_1}{2} = -\frac{\chi(S) - \chi(\partial S \cap L_1) + \#(\partial S \Delta L_1) - \#L_1}{2}$$

only depends on the topological type of  $(S, L_1)$ . It is an integer since both  $n(\Sigma \cup S)$  and  $n(\Sigma)$  are. Similarly for the exponent on  $-1$ .  $\square$

**4.3. Lasagna  $s$ -invariants and genus bounds.** Let  $(X, L)$  be a pair as before. In view of Theorem 4.1 we make the following definition.

**Definition 4.6.** The *lasagna  $s$ -invariant* of  $(X, L)$  at a double class  $(\alpha_+, \alpha_-) \in H_2^{L, \times 2}(X)$  is

$$s(X; L; \alpha_+, \alpha_-) := q(x_{\alpha_+, \alpha_-}) - 2\alpha_+ \cdot \alpha_- \in \mathbb{Z} \sqcup \{-\infty\}.$$

The *lasagna  $s$ -invariant* of  $(X, L)$  at a class  $\alpha \in H_2^L(X)$  is

$$s(X; L; \alpha) := s(X; L; \alpha, 0) = q(x_{\alpha, 0}) \in \mathbb{Z} \sqcup \{-\infty\}.$$

When  $L = \emptyset$ , we often drop it from the notation of lasagna  $s$ -invariants.

When  $X$  is a 2-handlebody, the isomorphism (14) respects the canonical Lee lasagna generators in the sense that  $x_{\beta_+, \beta_-} \in \mathcal{S}_0^{Lee}(X; (L, \partial\beta); \beta)$  is identified (up to renormalizations) with  $x_{\alpha_+, \alpha_-} \in$



$S_0^{Lee}(X; L; \alpha)$  for the unique pair  $(\alpha_+, \alpha_-) \in H_2^{L, \times 2}(X)$  with  $\alpha_+ - \alpha_- = \beta_+ - \beta_-$ . In particular,

$$s(X; L; \alpha_+, \alpha_-) = s(X; (L, \partial(\alpha_+ - \alpha_-)); \alpha_+ - \alpha_-). \quad (21)$$

Thus, the double class version of lasagna  $s$ -invariants in Definition 4.6 can be recovered from the single class version. When  $X$  is not a 2-handlebody, however, it is not clear that (14) holds in a filtered sense for Lee skein lasagna modules, so we do not have this conclusion.

We prove a few properties of lasagna  $s$ -invariants.

**Theorem 4.7.** *The lasagna  $s$ -invariants satisfy the following properties.*

(0) (Empty manifold)  $s(\emptyset; \emptyset; 0) = 0$ .

(1) (Recover classical  $s$ ) *If  $*$  denotes the unique element in  $H_2^L(B^4)$ , then*

$$s(B^4; L; *) = s_{\text{gl}_2}(L) + 1 = -s(-L) - w(L) + 1,$$

where  $s$  is the classical  $s$ -invariant defined by Rasmussen [Ras10] and Beliakova-Wehrli [BW08].

(2) (Symmetries)  $s(X; L; \alpha_+, \alpha_-) = s(X; L; \alpha_-, \alpha_+) = s(X; L^r; -\alpha_+, -\alpha_-)$ , where  $L^r$  denote the orientation reversal of  $L$ .

(3) (Parity)  $s(X; L; \alpha_+, \alpha_-) \equiv (\alpha_+ + \alpha_-)^2 + \#L \pmod{2}$  (by convention  $-\infty$  has arbitrary parity).

(4) (Connected sums) *If  $(X, L = L_1 \sqcup L_2)$  is a boundary sum, a connected sum, or the disjoint union of  $(X_1, L_1)$  and  $(X_2, L_2)$ , and  $(\alpha_{i,+}, \alpha_{i,-}) \in H_2^{L_i, \times 2}(X_i)$ ,  $i = 1, 2$ , then*

$$s(X; L; \alpha_{1,+} + \alpha_{2,+}, \alpha_{1,-} + \alpha_{2,-}) = s(X_1; L_1; \alpha_{1,+}, \alpha_{1,-}) + s(X_2; L_2; \alpha_{2,+}, \alpha_{2,-}).$$

(5) (Reduced connected sum) *If  $(X, L = L_1 \#_{K_1, K_2} L_2)$  is a (reduced) boundary sum of  $(X_1, L_1)$  and  $(X_2, L_2)$  performed along framed oriented intervals on components  $K_i \subset L_i$ ,  $i = 1, 2$ , and  $(\alpha_{i,+}, \alpha_{i,-}) \in H_2^{L_i, \times 2}(X_i)$  are double classes compatible with the boundary sum, then*

$$s(X; L_1 \#_{K_1, K_2} L_2; \alpha_{1,+} * \alpha_{2,+}, \alpha_{1,-} * \alpha_{2,-}) = s(X_1; L_1; \alpha_{1,+}, \alpha_{1,-}) + s(X_2; L_2; \alpha_{2,+}, \alpha_{2,-}) - 1.$$

(6) (Genus bound) *If  $(X_2, L_2)$  is glued to  $(X_1, L_1)$  along some  $(Y, T_0)$  to obtain  $(X, L)$ , and (unframed)  $S \subset X_2$  is properly embedded with  $\partial S = L_2$ , as in the setup of (19), such that every component of  $S$  has a boundary on  $T_0$ , then for any compatible partition  $S = S_+ \cup S_-$  and double class  $(\alpha_+, \alpha_-) \in H_2^{L_1, \times 2}(X_1)$ ,*

$$s(X; L; \alpha_+ * [S_+], \alpha_- * [S_-]) \leq s(X_1; L_1; \alpha_+, \alpha_-) - \chi(S) + \chi(T_0) - [S]^2.$$

The special case of Theorem 4.7(5) when  $X_1 = X_2 = B^4$ , in view of (1), recovers the connected sum formula

$$s(L_1 \#_{K_1, K_2} L_2) = s(L_1) + s(L_2)$$

for  $s$ -invariant of links in  $S^3$ , a fact that is proved only recently [Man+23, Theorem 7.1].

Since we performed gluing in its most general form, Theorem 4.7(6) has a rather complicated look. We state some special cases as corollaries, roughly in decreasing order of generality. Note that for 2-handlebodies, by (21), the double class versions of the genus bounds are equivalent to the single class versions.

**Corollary 4.8.** *Suppose  $(X_1, L_1 \sqcup L_0)$ ,  $(X_2, -L_0 \sqcup L_2)$  are two pairs that can be glued along some common boundary  $(Y, L_0)$  where  $Y$  is a closed 3-manifold (in particular  $L_1, L_2 \not\subset Y$ ). Suppose (unframed)  $S \subset X_2$  has  $\partial S = -L_0 \sqcup L_2$  and each component of  $S$  has a boundary on  $-L_0$ , then for any compatible partition  $S = S_+ \cup S_-$  and double class  $(\alpha_+, \alpha_-) \in H_2^{L_1 \sqcup L_0, \times 2}(X_1)$ ,*

$$s(X_1 \cup_Y X_2; L_1 \sqcup L_2; \alpha_+ * [S_+], \alpha_- * [S_-]) \leq s(X_1; L_1 \sqcup L_0; \alpha_+, \alpha_-) - \chi(S) - [S]^2.$$

In particular, for any class  $\alpha \in H_2^{L_1 \sqcup L_0}(X_1)$ ,

$$s(X_1 \cup_Y X_2; L_1 \sqcup L_2; \alpha * [S]) \leq s(X_1; L_1 \sqcup L_0; \alpha) - \chi(S) - [S]^2.$$

*Proof.* This is the special case of Theorem 4.7(6) when  $Y$  is closed, so  $T_0 = L_0$  with  $\chi(T_0) = 0$ .  $\square$

*Remark 4.9.* Although Theorem 4.7(6) is more general than Corollary 4.8, it is implied by the latter. This is because we can replace  $(X_2, S)$  by its union with a collar neighborhood of  $(\partial X_1, L_1)$ . Nevertheless, the proof of Theorem 4.7(6) is only notationally more complicated. The same remark applies to Theorem 4.5.

**Corollary 4.10.** *Suppose  $(W, S): (Y_1, L_1) \rightarrow (Y_2, L_2)$  is an unframed cobordism between framed oriented links in closed oriented 3-manifolds such that each component of  $S$  has a boundary on  $L_1$ , and  $X$  is a 4-manifold that bounds  $Y_1$ . Then for any compatible partition  $S = S_+ \cup S_-$  and double class  $(\alpha_+, \alpha_-) \in H_2^{L_1, \times 2}(X)$ ,*

$$s(X \cup_{Y_1} W; L_2; \alpha_+ * [S_+], \alpha_- * [S_-]) \leq s(X; L_1; \alpha_+, \alpha_-) - \chi(S) - [S]^2.$$

*In particular, for any class  $\alpha \in H_2^{L_1}(X)$ ,*

$$s(X \cup_Y W; L_2; \alpha * [S]) \leq s(X; L_1; \alpha) - \chi(S) - [S]^2.$$

*Proof.* This is the special case of Corollary 4.8 when  $\partial X_1 = Y$ .  $\square$

In the special case of Corollary 4.10 when  $S \subset S^3 \times I$  is a link cobordism between links  $L_1, L_2 \subset S^3$ , each component of which has a boundary on  $L_1$ , and when  $X = B^4$ , in view of Proposition 2.4 and Theorem 4.7(1), we have  $s(-L_1) \leq s(-L_2) - \chi(S)$ . Of course we can reverse the orientations to obtain

$$s(L_1) \leq s(L_2) - \chi(S),$$

which recovers the classical genus bound for link cobordisms from  $s$ -invariants [BW08, (7)]<sup>12</sup>.

*Proof of Proposition 1.11.* This is the special case of the single class version of Corollary 4.10 when  $L_1 = L_2 = S = \emptyset$ .  $\square$

**Corollary 4.11.** *Suppose  $(X, L)$  is a pair and  $S_+, S_- \subset X$  are disjoint properly embedded surfaces with  $\partial S_+ \cup \partial S_- = L$ , then*

$$2g(S_+) + 2g(S_-) \geq s(X; L; [S_+], [S_-]) + [S_+]^2 + [S_-]^2 - \#L.$$

*In particular, if  $S \subset X$  is properly embedded with  $\partial S = L$ , then*

$$2g(S) \geq s(X; L; [S]) + [S]^2 - \#L.$$

*Proof.* Delete a local ball on each component of  $S_+$  and  $S_-$ , regard the rest of  $X$  and  $S_{\pm}$  as a cobordism and apply Corollary 4.10, we get

$$s(X; L; [S_+], [S_-]) \leq s((B^4)^{\cup(\#S_+ + \#S_-)}; U^{\cup(\#S_+ + \#S_-)}; *) - \chi(S^\circ) - [S]^2,$$

where  $s((B^4)^{\cup(\#S_+ + \#S_-)}; U^{\cup(\#S_+ + \#S_-)}; *) = (\#S_+ + \#S_-)s(B^4; U; *) = \#S_+ + \#S_-$  by Theorem 4.7(1)(4). The statement follows by simplification.  $\square$

In the special case  $X = B^4$  and  $L = K \subset S^3$  is a knot, the single class version of Corollary 4.11 recovers Rasmussen's slice genus bound for knots [Ras10, Theorem 1].

<sup>12</sup>Note (7) in [BW08] incorrectly missed the condition about components of  $S$  having boundaries on ends.

**Corollary 4.12.** *Suppose  $X$  is a 4-manifold, and  $S_+, S_- \subset X$  are disjoint closed embedded surfaces, then*

$$2g(S_+) + 2g(S_-) \geq s(X; [S_+], [S_-]) + [S_+]^2 + [S_-]^2.$$

*In particular, if  $S \subset X$  is an embedded surface, then*

$$2g(S) \geq s(X; [S]) + [S]^2.$$

*Proof.* This is the special case of Corollary 4.11 when  $L = \emptyset$ . □

The *genus function* of a 4-manifold  $X$ ,  $g(X; \bullet): H_2(X) \rightarrow \mathbb{Z}_{\geq 0}$ , is defined by

$$g(X; \alpha) = \min\{g(\Sigma): \Sigma \subset X \text{ is a closed embedded surface with } [X] = \alpha\}.$$

**Corollary 4.13.** *(Theorem 1.8) The genus function of a 4-manifold  $X$  has a lower bound given by*

$$g(X; \alpha) \geq \frac{s(X; \alpha) + \alpha^2}{2}.$$

*Proof.* This is a reformulation of the second part of Corollary 4.12. □

We return to Theorem 4.7. It is an easy consequence of the hard work that we have done in previous sections.

*Proof of Theorem 4.7.* (0)  $x_{0,0} \in \mathcal{S}_0^{Lee}(\emptyset)$  is represented by the Lee lasagna filling 1 on the empty skein, which has filtration degree 0.

- (1)  $x_{*,0} \in \mathcal{S}_0^{Lee}(B^4; L)$  is represented by the Lee lasagna filling on the skein  $L \times I \subset S^3 \times I = B^4 \setminus \text{int}(\frac{1}{2}B^4)$  with decoration  $x_{o_L}$  on the input link  $L$ . Under the identification  $\mathcal{S}_0^{Lee}(B^4; L) \cong KhR_{Lee}(L)$  given by Proposition 2.4,  $x_{*,0}$  is equal to the Lee canonical generator  $x_{o_L}$ , which has filtration degree  $s_{\mathfrak{gl}_2}(L) + 1 = -s(-L) - w(L) + 1$  by (12).
- (2) The first equality is a consequence of Theorem 4.1(4). Reversing the orientation of all skeins defines an isomorphism  $\mathcal{S}_0^{Lee}(X; L) \cong \mathcal{S}_0^{Lee}(X; L')$  which negates the homology class grading but preserves other gradings and the quantum filtration, and exchanges the canonical Lee lasagna generators. This proves the second equality.
- (3) This is a consequence of Theorem 4.1(3).
- (4) This is a consequence of Proposition 2.2 and the fact that the canonical Lee lasagna generators behave tensorially under the various sum operations (as can be seen on the canonical Lee lasagna filling level).
- (6) This is a consequence of Theorem 4.5. Under the condition on the components of  $S$  and the compatibility assumption, the right hand side of (20) has exactly one term, which has  $S_{\pm}(\mathfrak{D}) = S_{\pm}$ . The statement follows from the fact that  $\mathcal{S}_0^{Lee}(X_2; S)$  has filtered degree  $-\chi(S) + \chi(T_0) - [S]^2$  (cf. the comment after (19)).
- (5) There is a framed saddle cobordism  $(\partial X \times I, S)$  from  $(\partial X, L = L_1 \#_{K_1, K_2} L_2)$  to  $(\partial X, L_1 \sqcup L_2)$ , where  $\partial X$  is bounded by  $X = X_1 \natural X_2$ . Thus Corollary 4.10 (which is a corollary of (6)) and (4) implies

$$s(X; L; \alpha_{1,+} * \alpha_{2,+}, \alpha_{1,-} * \alpha_{2,-}) \geq s(X_1; L_1; \alpha_{1,+}, \alpha_{1,-}) + s(X_2; L_2; \alpha_{2,+}, \alpha_{2,-}) - 1.$$

To prove the reverse inequality, by Theorem 4.1(4), we can choose signs  $\epsilon_i = \pm 1$ ,  $i = 1, 2$ , such that

$$q(x_{\alpha_{i,+}, \alpha_{i,-}} + \epsilon_i x_{\alpha_{i,-}, \alpha_{i,+}}) = s(X_i; L_i; \alpha_{i,+}, \alpha_{i,-}) - 2, \quad i = 1, 2.$$

By Theorem 4.5, the reverse  $S^r$  of  $S$  induces  $\mathcal{S}_0^{Lee}(\partial X \times I; S^r): \mathcal{S}_0^{Lee}(X; L_1 \sqcup L_2) \rightarrow \mathcal{S}_0^{Lee}(X; L)$  that maps  $(x_{\alpha_{1,+}, \alpha_{1,-} + \epsilon_1 x_{\alpha_{1,-}, \alpha_{1,+}}}) \otimes (x_{\alpha_{2,+}, \alpha_{2,-} + \epsilon_2 x_{\alpha_{2,-}, \alpha_{2,+}}})$  to a nonzero multiple of  $x_{\alpha_{1,+} * \alpha_{2,+}, \alpha_{1,-} * \alpha_{2,-} \pm x_{\alpha_{1,-} * \alpha_{2,-}, \alpha_{1,+} * \alpha_{2,+}}}$ . Since  $\mathcal{S}_0^{Lee}(\partial X \times I; S^r)$  has filtered degree 1, it follows that

$$\begin{aligned} & s(X; L; \alpha_{1,+} * \alpha_{2,+}, \alpha_{1,-} * \alpha_{2,-}) \\ & \leq q(x_{\alpha_{1,+} * \alpha_{2,+}, \alpha_{1,-} * \alpha_{2,-} \pm x_{\alpha_{1,-} * \alpha_{2,-}, \alpha_{1,+} * \alpha_{2,+}}) + 2 \\ & \leq q(x_{\alpha_{1,+}, \alpha_{1,-} + \epsilon_1 x_{\alpha_{1,-}, \alpha_{1,+}}}) + q(x_{\alpha_{2,+}, \alpha_{2,-} + \epsilon_2 x_{\alpha_{2,-}, \alpha_{2,+}}}) + 3 \\ & = s(X_1; L_1; \alpha_{1,+}, \alpha_{1,-}) + s(X_2; L_2; \alpha_{2,+}, \alpha_{2,-}) - 1. \end{aligned} \quad \square$$

## 5. NONVANISHING CRITERIA FOR 2-HANDLEBODIES

When  $X$  is a 2-handlebody, the Khovanov or Lee skein lasagna module of a pair  $(X, L)$  has a nice formula (13). In this section, we make use of this 2-handlebody formula to provide some explicit criteria for  $\mathcal{S}_0^2(X; L)$  or  $\mathcal{S}_0^{Lee}(X; L)$  to be nonvanishing, where by convention the latter means the filtration function on  $\mathcal{S}_0^{Lee}(X; L)$  is not identically  $-\infty$  (equivalently, the lasagna  $s$ -invariants of  $(X, L)$  are not all  $-\infty$ ). By our discussions in Section 3 and Section 4, these lead to smooth genus bounds for second homology classes of 4-manifolds. This section is an expansion of Section 1.3 and Section 1.4.

Throughout the section, we assume that  $X$  is the 2-handlebody obtained by attaching 2-handles to  $B^4$  along a framed link  $K = K_1 \cup \dots \cup K_m \subset S^3$ , and  $L \subset \partial X$  is given by a framed oriented link in  $S^3$  disjoint from  $K$ , as in the setup of (13).

**5.1. Rank inequality between Khovanov and Lee skein lasagna modules.** In this section we prove Theorem 1.12, which states that the rank of  $\mathcal{S}_0^2(X; L)$  is bounded below by the dimension of  $gr(\mathcal{S}_0^{Lee}(X; L))$  in every tri-degree  $(h, q, \alpha)$ . In particular, if  $\mathcal{S}_0^{Lee}(X; L)$  is nonvanishing, so is  $\mathcal{S}_0^2(X; L)$ . By Theorem 1.4, this has the following consequence.

**Corollary 5.1.** *If a knot trace  $X_n(K)$  embeds into a 4-manifold  $X$  for some knot  $K$  and framing  $n \geq -TB(-K)$ , then  $\mathcal{S}_0^{Lee}(X; L)$  is vanishing for any  $L \subset \partial X$ . In particular, the lasagna  $s$ -invariants of  $(X, L)$  are all  $-\infty$ .*

*Proof.* This is a consequence of Theorem 1.4 and Theorem 1.12 when  $X = X_n(K)$ . The general case follows from this and Proposition 2.3.  $\square$

Again, by taking the contrapositive and using the trace embedding lemma one obtains the following slice obstruction.

**Corollary 5.2.** *Suppose some class in  $\mathcal{S}_0^{Lee}(X; L)$  has finite filtration degree. If  $K \subset S^3$  is  $n$ -slice in  $X$ , then  $n > TB(K)$ .*  $\square$

We turn to Theorem 1.12. We will only use the 2-handlebody condition in a mild way, namely that the 2-handlebody formula (13) expresses  $\mathcal{S}_0^\bullet(X; L)$  as a filtered colimit instead of an arbitrary colimit (4).

*Proof of Theorem 1.12.* We first rewrite (13) into a filtered colimit. The group  $S(r) := S_{|\alpha_1|+2r_1} \times \dots \times S_{|\alpha_m|+2r_m}$  acts on  $KhR_\bullet(K(r))$ , where  $K(r) := K(\alpha^+ + r, \alpha^- + r) \cup L$ . Then (13) for  $\bullet = Lee$  can be rewritten as

$$\mathcal{S}_0^{Lee}(X; L; \alpha) = \text{colim}_{r \in \mathbb{Z}_{\geq 0}^m} KhR_{Lee}(K(r))^{S(r)} \{-|\alpha| - 2|r|\}, \quad (22)$$

where  $A^{S(r)}$  denotes the subgroup of  $A$  fixed by  $S(r)$ , and the morphisms are given by dotted annular creation maps composed with symmetrization. A similar formula holds for  $\mathcal{S}_0^2(X; L; \mathbb{Q})$ .

Now for a fixed triple  $(h, q, \alpha)$ , pick any linearly independent vectors  $y_1, \dots, y_n \in \mathcal{S}_{0, h, \leq q}^{Lee}(X; L; \alpha)$  whose images in  $gr_q(\mathcal{S}_{0, h}^{Lee}(X; L; \alpha))$  are also linearly independent. Since  $\mathcal{S}_0^{Lee}(X; L; \alpha)$  is expressed as a filtered colimit, we can choose representatives  $v_1, \dots, v_n \in KhR_{Lee}^h(K(r))^{S(r)}\{-|\alpha| - 2|r|\}$  for  $y_1, \dots, y_n$  with  $q(v_i) = q$  ( $q(\cdot)$  denotes the filtration function).

Let  $(CKhR_2(K(r); \mathbb{Q}), d)$  and  $(CKhR_{Lee}(K(r)), d_{Lee})$  be the Khovanov-Rozansky  $\mathfrak{gl}_2$  and Lee cochain complexes for  $K(r)$ , where  $CKhR_{Lee}(K(r)) = CKhR_2(K(r); \mathbb{Q})$  and  $d_{Lee}$  is the sum of  $d$  and another term that decreases the quantum degree. We further lift  $v_1, \dots, v_n$  to the chain level to Lee cocycles  $a_1, \dots, a_n \in CKhR_{Lee}^h(K(r))\{-|\alpha| - 2|r|\}$  with  $a_i = c_i + \epsilon_i$  where  $c_i$  is homogeneous of quantum degree  $q$ , and  $q(\epsilon_i) < q$ . Since  $0 = d_{Lee}a_i = dc_i + \epsilon$  where  $q(\epsilon) < q$ , we see  $c_i$  is a Khovanov cocycle.

We claim that any nontrivial linear combination of  $c_i$ 's represents a nonzero element in  $\mathcal{S}_{0, h, q}^2(X; L; \mathbb{Q})$ , which would imply  $\text{rank}(\mathcal{S}_{0, h, q}^2(X; L)) \geq n$ , proving the theorem. Let  $c = \sum \lambda_i c_i$  where not all  $\lambda_i$  are zero. To prove  $c$  represents a nonzero element, it suffices to check its homology class survives under an arbitrary morphism  $KhR_2(K(r); \mathbb{Q})^{S(r)}\{-|\alpha| - 2|r|\} \rightarrow KhR_2(K(r'); \mathbb{Q})^{S(r')}\{-|\alpha| - 2|r'|\}$  in the filtered system. Since this map is a linear combination of induced maps by link cobordisms, it is induced by a map  $\Phi: CKhR_2(K(r); \mathbb{Q})\{-|\alpha| - 2|r|\} \rightarrow CKhR_2(K(r'); \mathbb{Q})\{-|\alpha| - 2|r'|\}$  on the chain level. Similarly  $KhR_{Lee}(K(r))^{S(r)} \rightarrow KhR_{Lee}(K(r'))^{S(r')}$  is induced by some  $\Phi_{Lee}: CKhR_{Lee}(K(r))\{-|\alpha| - 2|r|\} \rightarrow CKhR_{Lee}(K(r'))\{-|\alpha| - 2|r'|\}$ , where we can assume  $CKhR_{Lee}(K(r')) \cong CKhR_2(K(r'); \mathbb{Q})$  and  $\Phi_{Lee}$  is a sum of  $\Phi$  (which is homogeneous) and some another term that decreases the quantum degree. It follows that  $a = \sum \lambda_i a_i$  satisfies  $\Phi_{Lee}a = \Phi c + \epsilon$  for some  $\epsilon$  with  $q(\epsilon) < q$ . Now, suppose for contrary that  $\Phi c = db$  is a Khovanov coboundary. Then  $\Phi_{Lee}a$  is homologous to  $\Phi_{Lee}a - d_{Lee}b$  which has filtration degree less than  $q$ . Thus the element  $y = \sum_i \lambda_i y_i \in \mathcal{S}_0^{Lee}(X; L)$ , which was represented by  $[a] \in KhR_{Lee}(K(r))^{S(r)}$ , is also represented by  $[\Phi_{Lee}a] \in KhR_{Lee}(K(r'))^{S(r')}$  with filtration level less than  $q$ , implying  $q(y) < q$ , contradicting the choices of  $y_1, \dots, y_n$ .  $\square$

**5.2. Comparison results for Lee skein lasagna modules.** In this section, we prove two comparison results for Lee skein lasagna modules of 2-handlebodies.

Recall that  $(X, L)$  is the pair that arises from  $K \cup L \subset S^3$  after attaching 2-handles along  $K$ . Let  $(X', L')$  be another such pair from  $K' \cup L'$ . Suppose there is a concordance  $C: K \rightarrow K'$ , that is, a genus 0 component-preserving framed link cobordism in  $S^3 \times I$  (thus topologically  $C$  is a disjoint union of annuli). Then the surgery on  $C$  gives a homology cobordism  $W$  between the surgery on  $K$  and the surgery on  $K'$ . In particular,  $W$  (and thus  $C$ ) induces an isomorphism  $\varphi_C: H_*(\partial X) \cong H_*(\partial X')$ . Moreover,  $X' = X \cup_{\partial X} W$ . If  $A \subset W$  is a concordance between  $L$  and  $L'$ , then  $(W, A)$  induces an isomorphism  $H_2(X, L) \cong H_2(X', L')$  that restricts to  $\phi_A: H_2^L(X) \cong H_2^{L'}(X')$ .

**Proposition 5.3.** *Let  $(X, L)$  and  $(X', L')$  be obtained from  $K \cup L, K' \cup L' \subset S^3$  as above, and let  $C: K \rightarrow K'$  be a concordance that induces the homology cobordism  $W: \partial X \rightarrow \partial X'$ .*

- (1) *If  $A \subset W$  is a framed concordance between  $L, L'$ , then  $\mathcal{S}_0^{Lee}(X; L) \cong \mathcal{S}_0^{Lee}(X'; L')$  as tri-graded (by  $\mathbb{Z} \times \mathbb{Z}/4 \times H_2^L(X)$ , where  $H_2^L(X) \cong H_2^{L'}(X')$  via  $\phi_A$ ), quantum filtered vector spaces. Moreover,  $s(X; L; \alpha) = s(X'; L'; \phi_A(\alpha))$  for all  $\alpha \in H_2^L(X)$ .*
- (2) *If  $L$  has components  $L_1, \dots, L_k$ ,  $L'$  has components  $L'_1, \dots, L'_k$ , and  $\varphi_C([L_i]) = [L'_i]$  for all  $i$ , then  $\mathcal{S}_0^{Lee}(X; L)$  is nonvanishing if and only if  $\mathcal{S}_0^{Lee}(X'; L')$  is nonvanishing.*

*Proof.* (1) Gluing  $(W, A)$  to  $(X, L)$  defines a filtered (degree 0) map  $\mathcal{S}_0^{Lee}(W; A): \mathcal{S}_0^{Lee}(X; L) \rightarrow \mathcal{S}_0^{Lee}(X'; L')$ . Turning  $(W, A)$  upside down gives another filtered map  $\mathcal{S}_0^{Lee}(X'; L') \rightarrow \mathcal{S}_0^{Lee}(X; L)$ .

By Theorem 4.5, the two maps interchange the corresponding canonical Lee lasagna generators, thus are inverses to each other. The statement follows.

(2) By symmetry, it suffices to assume the filtration of  $\mathcal{S}_0^{Lee}(X; L)$  is identically  $-\infty$  and prove the same for  $\mathcal{S}_0^{Lee}(X'; L')$ .

Under the given condition, we can find framed immersed cobordisms  $S_i: L_i \rightarrow L'_i$  in  $W$  that intersect each other transversely. After deleting some local balls, we can regard  $S = S_1 \cup \dots \cup S_k$  as a skein in  $W$  rel  $-L \sqcup L'$ . For any double class  $(\alpha_+, \alpha_-) \in H_2^{L, \times 2}(X)$ , there is a unique compatible double skein structure  $S = S_+ \cup S_-$ . Gluing  $(W, S)$  with the canonical Lee lasagna filling  $x(S_+, S_-)$  to  $(X, L)$  maps  $x_{\alpha_+, \alpha_-} \in \mathcal{S}_0^{Lee}(X; L)$  to a nonzero multiple of  $x_{\alpha_+ * [S_+], \alpha_- * [S_-]} \in \mathcal{S}_0^{Lee}(X'; L')$ . Since  $q(x_{\alpha_+, \alpha_-}) = -\infty$ , we see  $q(x_{\alpha_+ * [S_+], \alpha_- * [S_-]}) = -\infty$  as well. Since all canonical Lee lasagna generators of  $(X', L')$  arise this way, we conclude that the filtration function on  $\mathcal{S}_0^{Lee}(X'; L')$  is identically  $-\infty$ .  $\square$

**5.3. Lasagna  $s$ -invariant and  $s$ -invariants of cables.** In this section we show that the lasagna  $s$ -invariants of  $(X, L)$  are determined by the classical  $s$ -invariants of cables of  $K \cup L$ . Recall that as in the setup of (13), the choice  $K \cup L \subset S^3$  gives an identification  $H_2^L(X) \cong \mathbb{Z}^m$ .

**Proposition 5.4.** *For any  $\alpha \in H_2^L(X) \cong \mathbb{Z}^m$ ,*

$$\begin{aligned} s(X; L; \alpha) &= \lim_{r \rightarrow \infty^m} (s_{\mathfrak{gl}_2}(K(\alpha^+ + r, \alpha^- + r) \cup L) - 2|r|) - |\alpha| + 1 \\ &= - \lim_{r \rightarrow \infty^m} (s(-(K(\alpha^+ + r, \alpha^- + r) \cup L)) + 2|r|) - w(K(\alpha^+, \alpha^-) \cup L) - |\alpha| + 1. \end{aligned}$$

*Moreover, the quantity within the limit sign in the first (resp. second) line is nonincreasing (resp. nondecreasing) in  $r$ .*

*Proof.* Since the morphisms in the 2-handlebody formula for  $\bullet = Lee$  are filtered, the quantity within the limit sign in the first line is nonincreasing in  $r$ . In particular, the limit exists (which could be  $-\infty$ ). The  $\mathfrak{gl}_2$   $s$ -invariant relates to the classical  $s$ -invariant via (12). Since  $w(K(\alpha^+ + r, \alpha^- + r) \cup L)$  is independent of  $r$ , the claims on the second equality follow from the first. Thus it remains to prove the first equality.

We describe explicitly the isomorphism (13). Delete a 4-ball  $B^4$  slightly smaller than the 0-handle of  $X$  and put  $K(r) := K(\alpha^+ + r, \alpha^- + r) \cup L$  on its boundary. Cap off the sublink  $K(\alpha^+ + r, \alpha^- + r)$  by some parallel copies of cores of the 2-handles of various orientations corresponding to  $(\alpha^+ + r, \alpha^- + r)$ , and put a cylindrical surface  $L \times I$  connecting the inner link  $L \subset \partial B^4$  and the outer link  $L \subset \partial X$ . This gives a skein in  $(X, L)$  with input link  $K(r)$ . Putting decorations on  $K(r)$  defines  $KhR_{Lee}(K(r)) \rightarrow \mathcal{S}_0^{Lee}(X; L)$ . Taking the direct sum over  $r$  defines a map that descends to the isomorphism (13). It follows that the Lee canonical filling on this skein as a double skein (given by its own orientation) is given up to a scalar by the Lee canonical generator of  $K(r)$  (with its own orientation) decorated on the input ball.

As in the proof of Theorem 1.12, (13) for  $\bullet = Lee$  can be rewritten as (22), where  $S(r) = S_{|\alpha_1| + 2r_1} \times S_{|\alpha_m| + 2r_m}$ . Label the cables of  $K_i$  in  $K(r)$  by  $1, 2, \dots, |\alpha_i| + 2r_i$ . Then the orientations of  $K(r)$  compatible with  $L$  can be identified with tuples  $(X_1, \dots, X_m)$  where each  $X_i$  is a subset of  $[|\alpha_i| + 2r_i] := \{1, 2, \dots, |\alpha_i| + 2r_i\}$ , indicating which strands are given the same orientation as  $K_i$ . Those that give rise to possible orientations of  $K(r)$  compatible with the homology class degree  $\alpha$  correspond to such tuples with  $\#X_i = \alpha_i^+ + r_i$ . Then the canonical Lee lasagna generator  $x_{\alpha, 0} \in \mathcal{S}_0^{Lee}(X; L; \alpha)$

under (22) is represented by

$$x^{inv}(r) := \sum_{\#X_i = \alpha_i^+ + r_i} x_{(X_1, \dots, X_m)} \in KhR_{Lee}(K(r))^{S(r)}$$

with a  $\{-|\alpha| - 2|r|\}$  shift. Define  $s_{\mathfrak{gl}_2}^{inv}(K(r)) := q(x^{inv}(r))$ . Then by (22) we have

$$s(X; L; \alpha) = \lim_{r \rightarrow \infty^m} (s_{\mathfrak{gl}_2}^{inv}(K(r)) - |\alpha| - 2|r|). \quad (23)$$

It remains to show the right hand side of (23) is unchanged if we replace  $s_{\mathfrak{gl}_2}^{inv}(K(r))$  by  $s_{\mathfrak{gl}_2}(K(r)) + 1$ , the latter of which is equal to  $q(x_{(X_1, \dots, X_m)})$  where  $x_{(X_1, \dots, X_m)}$  is any term in the summation that defines  $x^{inv}(r)$ .

Since  $x^{inv}(r)$  is a sum of  $x_{(X_1, \dots, X_m)}$  and some  $S(r)$ -translates of it (which all have the same filtration degree), we have  $q(x^{inv}(r)) \leq q(x_{(X_1, \dots, X_m)})$ , thus

$$\lim_{r \rightarrow \infty^m} (s_{\mathfrak{gl}_2}^{inv}(K(r)) - |\alpha| - 2|r|) \leq \lim_{r \rightarrow \infty^m} (s_{\mathfrak{gl}_2}(K(r)) + 1 - |\alpha| - 2|r|).$$

Now suppose the right hand side is not  $-\infty$ , so the value stabilizes to some limit  $M$  for  $r$  sufficiently large. We claim that for sufficiently large  $r$ , we have  $q(x^{inv}(r)) = q(x_{(X_1, \dots, X_m)}) = M + |\alpha| + 2|r|$ , from which the statement follows.

To this end, we follow the argument in [Ren23, Section 4]. By looking at the  $S(r)$ -action on the canonical generators, we see

$$KhR_{Lee}(K(r)) \cong \bigotimes_{i=1}^m \mathbb{Q} \mathcal{P}([\alpha_i + 2r_i])$$

as an  $S(r)$ -representation, where  $\mathcal{P}$  denotes the power set operation on a set. Each tensor summand can be further decomposed as

$$\begin{aligned} \mathbb{Q} \mathcal{P}([\alpha_i + 2r_i]) &= \bigoplus_{j=0}^{|\alpha_i + 2r_i|} \mathbb{Q} \{X \subset [\alpha_i + 2r_i]; \#X = j\} \\ &= \bigoplus_{j=0}^{|\alpha_i + 2r_i|} \left( \bigoplus_{k=0}^{\min(j, |\alpha_i + 2r_i - j|)} (|\alpha_i + 2r_i - k, k) \right) \end{aligned}$$

into irreducible  $S_{|\alpha_i + 2r_i}$ -representations. Here  $(a, b)$  denote the irreducible  $S_{a+b}$ -representation over  $\mathbb{Q}$  defined by the two-row Young diagram  $(a, b)$ . Since the  $S(r)$ -action on  $KhR_{Lee}(K(r))$  respects the filtration structure, all nonzero elements in each irreducible piece  $V$  in the decomposition of  $S(r)$  have the same filtration degree, which is denoted  $q(V)$ .

Without loss of generality, suppose  $\alpha_i \geq 0$  for all  $i$  (otherwise reverse the orientations of some  $K_i$ 's). Then  $\alpha_i^+ = \alpha_i$ ,  $\alpha_i^- = 0$ . The canonical generator  $x_{(X_1, \dots, X_m)} \in KhR_{Lee}(K(r))$  lies in the  $S(r)$ -subrepresentation (which is optimally small)

$$\bigotimes_{i=1}^m \mathbb{Q} \{X \subset [\alpha_i + 2r_i]; \#X = \alpha_i + r\} = \bigoplus_{k_1=0}^{r_1} \cdots \bigoplus_{k_m=0}^{r_m} \bigotimes_{i=1}^m (\alpha_i + 2r_i - k_i, k_i), \quad (24)$$

while the element  $x^{inv}(r)$  lies in the subrepresentation

$$\bigotimes_{i=1}^m (\alpha_i + 2r_i, 0).$$

Thus  $q(x_{(X_1, \dots, X_m)}) = \max_k (q(\bigotimes_{i=1}^m (\alpha_i + 2r_i - k_i, k_i)))$  (this uses the fact that these representations are pairwise nonisomorphic) and  $q(x^{inv}(r)) = q(\bigotimes_{i=1}^m (\alpha_i + 2r_i, 0))$ .

Suppose for the contrary,  $q(x^{inv}(r)) < q(x_{(X_1, \dots, X_m)})$ . Then there is a summand  $\otimes_{i=1}^m (\alpha_i + 2r_i - k_i, k_i)$  in (24) with some  $k_t \neq 0$  that has  $q(\otimes_{i=1}^m (\alpha_i + 2r_i - k_i, k_i)) = q(x_{(X_1, \dots, X_m)}) = M + |\alpha| + 2|r|$ . The argument in [Ren23, Section 4.3] shows that  $q(\otimes_{i=1}^m (\alpha_i + 2r_i - k_i - \delta_{it}, k_i - \delta_{it})) = q(\otimes_{i=1}^m (\alpha_i + 2r_i - k_i, k_i))$  where  $\otimes_{i=1}^m (\alpha_i + 2r_i - k_i - \delta_{it}, k_i - \delta_{it})$  is the corresponding  $S(r - e_t)$ -irreducible summand of  $KhR_{Lee}(K(r - e_t))$  ( $e_t$  is the  $t^{\text{th}}$  coordinate vector). However, this implies the corresponding canonical generator for  $KhR_{Lee}(K(r - e_t))$  has

$$\begin{aligned} q(x_{(X_1, \dots, X_t \setminus \{*\}, \dots, X_m)}) &\geq q(\otimes_{i=1}^m (\alpha_i + 2r_i - k_i - \delta_{it}, k_i - \delta_{it})) \\ &= q(x_{(X_1, \dots, X_m)}) = M + |\alpha| + 2|r| > M + |\alpha| + 2|r - e_t|. \end{aligned}$$

This contradicts  $r$  being sufficiently large (so that  $\lim(s_{\mathfrak{gl}_2}(K(\cdot)) + 1 - |\alpha| - 2|\cdot|) = M$  already stabilizes at  $r - e_t$ ). The proof is complete.  $\square$

**5.4. Diagrammatic nonvanishing result.** For simplicity, in this section we assume  $L = \emptyset$ . We give a diagrammatic criterion to guarantee the nonvanishingness of  $\mathcal{S}_0^2(X)$  and  $\mathcal{S}_0^{Lee}(X)$ .

As before, assume  $X$  is a 2-handlebody obtained by attaching 2-handles to  $B^4$  along a framed oriented link  $K \subset S^3$  with components  $K_1, \dots, K_m$  and framings  $p_1, \dots, p_m$ . Let  $p = (p_1, \dots, p_m)$  and  $P$  be the diagonal matrix with entries  $p_1, \dots, p_m$ . Choose a link diagram  $D$  of  $K$  as an unframed link. Let  $N$  denote the crossing matrix of  $D$ , where  $N_{ii}$  is the number of crossings of  $K_i$  in diagram  $D$ , and  $N_{ij}$  is a half the number of crossings between  $K_i$  and  $K_j$  in  $D$  for  $i \neq j$ . Let  $w = (w_1, \dots, w_m)$  be the writhe vector of  $K$  in  $D$ , and  $W$  be the diagonal matrix with entries  $w_1, \dots, w_m$ . Finally, for  $f, n \in \mathbb{Z}^m$ , let  $K^f$  denote the knot  $K$  with framing  $f$ , and  $K^f(n)$  denote the  $n$ -cable of  $K$  with framing  $f$ , obtained by replacing  $K_i$  by an  $f_i$ -framed  $|n_i|$ -cable, with orientation reversed if  $n_i < 0$ . Then, any  $K^w(n)$  has a natural diagram induced from  $D$  by taking blackboard-framed cables.

**Theorem 5.5.** *In the notations above, suppose a class  $\alpha \in H_2(X) \cong \mathbb{Z}^m$  satisfies*

- (i)  $\alpha$  maximizes  $h(\alpha') := \alpha'^T (P - W + N) \alpha'$  among all vectors  $\alpha' \in \alpha + 2\mathbb{Z}^m$  (in particular,  $P - W + N$  is nonpositive);
- (ii)  $K^w(\alpha')$  has no negative crossings in the diagram induced from  $D$  for any  $\alpha' \in \alpha + 2\mathbb{Z}^m$  with  $h(\alpha') = h(\alpha)$ .

Then

- (1) For  $\bullet \in \{2, Lee\}$ ,  $\mathcal{S}_{0, > 0}^\bullet(X; \alpha) = 0$ ;
- (2) The dimension of  $\mathcal{S}_{0, 0}^{Lee}(X; \alpha)$  is equal to the number of maximal points of  $h$  on  $\alpha + 2\mathbb{Z}^m$ . The rank of  $\mathcal{S}_{0, 0}^2(X; \alpha)$  is bounded above by this number;
- (3)  $s(X; \alpha) = -s(-K^p(\alpha)) - w(K^p(\alpha)) - |\alpha| + 1$ . In particular,  $X$  has nonvanishing Khovanov and Lee skein lasagna modules.

*Remark 5.6.* It is possible to allow a boundary link  $L \subset \partial X$ , but we refrained from doing so to avoid making the statement too complicated.

**Corollary 5.7.** *In the notations of Theorem 5.5, if  $P - W + N$  is negative definite, then  $s(X; 0) = 0$*

$$\text{and } \mathcal{S}_{0, 0, q}^2(X; 0) \otimes \mathbb{Q} = gr_q(\mathcal{S}_{0, 0}^{Lee}(X; 0)) = \begin{cases} \mathbb{Q}, & q = 0 \\ 0, & q \neq 0. \end{cases}$$

*Proof.* Conditions (i)(ii) in Theorem 5.5 are satisfied for  $\alpha = 0$ , which is the unique maximal point of  $h$ . Thus (3) implies  $s(X; 0) = -s(\emptyset) + 1 = 0$ . By definition of  $s$ , this implies  $\dim gr_0(\mathcal{S}_{0, 0}^{Lee}(X; 0)) \geq 1$ . Now (2) and Theorem 1.12 imply the claims on  $\mathcal{S}_{0, 0}^\bullet(X; 0)$ .  $\square$



*Proof of Theorem 1.14.* For  $X = X_n(K)$ , in the notation of Theorem 5.5 we have  $m = 1$ ,  $P - W + N = n + 2n_-(D)$ . Thus (1) follows from Corollary 5.7.

For (2), take  $D$  to be a positive diagram of  $K$ . Then for  $n \leq 0$ , (i)(ii) in Theorem 5.5 are satisfied for both  $\alpha = 0, 1$ . Thus  $s(X; 0) = -s(\emptyset) + 1 = 0$  and  $s(X; 1) = -s(-K) - n - 1 + 1 = s(K) - n$  by Theorem 5.5(3). For  $n < 0$ ,  $\alpha = \pm 1$  are the unique maximal points of  $h$  on  $1 + 2\mathbb{Z}$ , thus Theorem 5.5(2) implies that the rank of  $\mathcal{S}_{0,0}^2(X; 1)$  is at most 2. Now the claims on  $\mathcal{S}_{0,0}^\bullet(X; 1)$  follows from Theorem 1.6(4) and Theorem 1.12.  $\square$

The rest of the section is devoted to proving Theorem 5.5.

For a link  $L$ , let  $P(L)(t, q)$  be the Poincaré polynomial of  $KhR_2(L)$ . Define

$$\widetilde{KhR}_2(L) := KhR_2(L)[\frac{1}{2}w(L)]\{-\frac{1}{2}w(L)\}$$

to be the renormalized homology of  $L$ , which is independent of the orientation of  $L$  (cf. (9)). Let  $\tilde{P}(L)(t, q)$  be the Poincaré polynomial of  $\widetilde{KhR}_2(L)$  (which is a Laurent polynomial in  $t^{1/2}, q^{1/2}$ ).

For two (Laurent) polynomials  $P, Q$  in variables  $t^{1/2}, q^{1/2}$ , we write  $P \leq Q$  if every coefficient of  $P$  is less than or equal to that of  $Q$ . For two vectors  $a, b \in \mathbb{Z}^m$ , write  $a \leq b$  if  $a_i \leq b_i$  for all  $i$ .

**Lemma 5.8 (Comparison lemma).** *For  $n \in \mathbb{Z}_{\geq 0}^m$  and  $p \in \mathbb{Z}^m$  with  $p \leq w$ , we have*

$$\tilde{P}(K^p(n)) \leq \sum_{\substack{0 \leq n' \leq n \\ 2|n-n'}} R_{n,n',w-p} \tilde{P}(K^w(n')) \quad (25)$$

for some polynomials  $R_{n,n',w-p} \in \mathbb{Z}[t^{\pm 1/2}, q^{\pm 1/2}]$  independent of  $K$  with

$$R_{n,n',w-p}(t, q) = t^{-\sum_i \frac{(w_i - p_i)n_i^2}{2}} q^{\sum_i \frac{(w_i - p_i)(n_i^2 + 2n'_i)}{2}} \prod_{j=1}^m \left( \sum_{r_j=0}^{\frac{n_j - n'_j}{2}} \dim \left( \frac{n_j + n'_j}{2} + r_j, \frac{n_j - n'_j}{2} - r_j \right) q^{2r_j} \right) + \dots$$

where  $\dots$  are terms with lower  $t$ -degrees, and  $(a, b)$  denotes the irreducible  $S_{a+b}$ -representation given by the Young diagram  $(a, b)$ .

Moreover, if (25) is sharp at the highest  $t$ -degree of the left hand side, then for any  $i = 1, \dots, m$  with  $n_i \geq 2$ , the morphism on  $KhR_2$  induced by the dotted annular creation map  $K^p(n - 2e_i) \rightarrow K^p(n)$ <sup>13</sup> is injective in the highest nontrivial homological degree of  $KhR_2(K^p(n))$ .

For reader's convenience, we note that for  $r, n \in \mathbb{Z}_{\geq 0}$ ,  $r \leq n/2$ ,

$$\dim(n - r, r) = \binom{n}{r} - \binom{n}{r-1}.$$

Lemma 5.8 is a consequence of an inductive argument pioneered by Stošić [Sto09]. Although it might have been implicit in the proof of Theorem 1 there, or in more details in [Tag13], for completeness we sketch a proof in Appendix A. For our purpose, we do not really need the quantum shifting information in the lemma.

**Lemma 5.9.** *For  $n \in \mathbb{Z}_{\geq 0}^m$ , the maximal  $t$ -degree of  $\tilde{P}(K^w(n))$  is at most  $\frac{1}{2}n^T Nn$ .*

<sup>13</sup>Really, the dotted annular creation map  $K^p(n - 2e_i) \rightarrow K^p(n - e_i, e_i)$  on  $KhR_2$  composed with the isomorphism that changes the orientation of the reversed  $K_i$  strand.

*Proof.* For any framed link diagram  $D$  of  $L$ , the maximal  $t$ -degree of  $P(L)$  is at most the number of negative crossings of  $D$ , and thus the maximal  $t$ -degree of  $\tilde{P}(L)$  is at most one half the total number of crossings of  $D$ . For  $D$  the standard diagram of  $K^w(n)$ , this number is  $\frac{1}{2}n^T N n$ .  $\square$

*Proof of Theorem 5.5.* Note that adding positive kinks to  $D$  does not affect the conditions and statements, but increases  $w$ . Thus we can assume that  $w \geq p$  and  $D$  has no 0-framed unknot component. We also assume  $\alpha \in \mathbb{Z}_{\geq 0}^m$ , as the general case is only notationally more complicated (and can indeed be deduced from the special case by reversing the orientations of some  $K_i$ 's). Thus  $\alpha^+ = \alpha$ ,  $\alpha^- = 0$ . Write  $N = N_+ + N_-$  to separate contributions coming from positive/negative crossings.

Recall that the 2-handlebody formula for  $X$  and  $\alpha \in \mathbb{Z}_{\geq 0}^m$  (13) can be rewritten as

$$\mathcal{S}_0^\bullet(X; \alpha) = \operatorname{colim}_{r \in \mathbb{Z}_{\geq 0}^m} \operatorname{KhR}_\bullet(K^p(\alpha + r, r))^{S_{\alpha+2r} \{-|\alpha| - 2|r|\}}, \quad (26)$$

where  $S_n = S_{n_1} \times \cdots \times S_{n_m}$  for  $n \in \mathbb{Z}_{\geq 0}^m$ . Here if  $\bullet = 2$  we assume to work over  $\mathbb{Q}$ .

**Step 1:** Isolating the highest  $t$ -degrees in  $\tilde{P}$ .

In this first step we prove that, for any  $n \in \mathbb{Z}_{\geq 0}^m$ ,  $n \equiv \alpha \pmod{2}$ ,

$$\tilde{P}(K^p(n)) = \sum_{\substack{0 \leq n' \leq n \\ 2|n-n' \\ h(n')=h(\alpha)}} R_{n,n',w-p} \tilde{P}(K^w(n')) + \cdots \quad (27)$$

where the highest nontrivial  $t$ -degree in each summand is  $\frac{1}{2}h(\alpha)$ , and  $\cdots$  denotes terms with lower  $t$ -degrees. Moreover, if  $\tilde{P}_{Lee}(K^p(n))(t, q)$  denotes the corresponding renormalized Poincaré polynomial for  $gr(\operatorname{KhR}_{Lee}(K^p(n)))$ , then the highest  $t$ -degree parts of  $\tilde{P}(K^p(n))$  and  $\tilde{P}_{Lee}(K^p(n))$  are equal.

In the comparison lemma 5.8, the highest  $t$ -degree of the  $n'$  term on the right hand side is at most

$$\frac{1}{2}n^{TT}(P - W)n' + \frac{1}{2}n^{TT}Nn' = \frac{1}{2}n^{TT}(P - W + N)n'$$

by Lemma 5.9, which is strictly less than  $h(\alpha)$  if  $h(n') \neq h(\alpha)$  by assumption (i). On the other hand, if  $h(n') = h(\alpha)$ , then by assumption (ii)  $K^w(n')$  has a positive diagram induced by  $D$  with some number  $s(n')$  of connected components (which only depends on the indices of the nonzero coordinates of  $n'$  as  $D$  has no 0-framed unknot component). Then  $\operatorname{KhR}_2^0(K^w(n'))$  (as well as  $\operatorname{KhR}_{Lee}^0(K^w(n'))$ ) has rank  $2^{s(n')}$  [Kho03, Proposition 6.1]. After renormalizing, we see that  $R_{n,n',w-p} \tilde{P}(K^w(n'))$  has maximal  $t$ -degree  $\frac{1}{2}h(\alpha)$ , and in this  $t$ -degree the polynomial has rank (i.e. the evaluation at  $q = 1$ )

$$2^{s(n')} \prod_{j=1}^m \left( \sum_{r_j=0}^{\frac{n_j-n'_j}{2}} \dim \left( \frac{n_j+n'_j}{2} + r_j, \frac{n_j-n'_j}{2} - r_j \right) \right) = 2^{s(n')} \binom{n}{\frac{n-n'}{2}}$$

where for  $a, b \in \mathbb{Z}_{\geq 0}^m$  we write  $\binom{a}{b} = \prod_i \binom{a_i}{b_i}$ . Now it follows from Lemma 5.8 that the highest  $t$ -degree of  $\tilde{P}(K^p(n))$  is at most  $\frac{1}{2}h(\alpha)$ , and in this  $t$ -degree it has rank at most

$$\sum_{\substack{0 \leq n' \leq n \\ 2|n-n' \\ h(n')=h(\alpha)}} 2^{s(n')} \binom{n}{\frac{n-n'}{2}}.$$

On the other hand, the rank of  $\tilde{P}_{Lee}(K^p(n))$  at  $t = \frac{1}{2}h(\alpha)$  is equal to the number of orientations of  $K^p(n)$  with writhe  $h(\alpha)$ . Dividing the count into algebraic cables of  $K^p$ , the rank is again

$$\sum_{\substack{-n \leq n' \leq n \\ 2|n-n'}} \binom{n}{\frac{n-n'}{2}} = \sum_{\substack{-n \leq n' \leq n \\ 2|n-n' \\ h(n')=h(\alpha)}} \binom{n}{\frac{n-n'}{2}} = \sum_{\substack{0 \leq n' \leq n \\ 2|n-n' \\ h(n')=h(\alpha)}} 2^{s(n')} \binom{n}{\frac{n-n'}{2}},$$

$n'^T(P-W+N_+-N_-)n'$                        $h(n')=h(\alpha)$                        $h(n')=h(\alpha)$   
 $=\alpha^T(P-W+N)\alpha$

where the first equality follows from conditions (i) and (ii). The second equality is true because in view of (ii), for  $n' \geq 0$ , changing signs of coordinates of  $n'$  maintains  $h(n')$  if and only if, for each connected component of the diagram of  $K^w(n')$ , we either change all of the corresponding coordinates of  $n'$  or none of them. Since  $\tilde{P}(K^p(n))$  is bounded below by  $\tilde{P}_{Lee}(K^p(n))$  as there is a spectral sequence from  $KhR_2$  to  $KhR_{Lee}$ , we must have the equality (27) as well as its version for  $\tilde{P}_{Lee}(K^p(n))$  as claimed.

**Step 2:** Unrenormalization and proofs of (1) and (2).

For  $n = \alpha + 2r$ , condition (ii) ensures that the writhe of  $K^w(n')$  is same as  $n'^T N n'$ , allowing us to unrenormalize (27) to give

$$P(K^p(\alpha + r, r)) = \sum_{\substack{0 \leq n' \leq n \\ 2|n-n' \\ h(n')=h(\alpha)}} (tq^{-1})^{\frac{1}{2}n'^T(W-P)n'} R_{n,n',w-p} P(K^w(n')) + \dots,$$

where the maximal  $t$ -degree of each term on the right hand side is 0. The same holds for the Lee version. In view of (26), this proves (1).

For (2), by Theorem 4.1(1)(2),  $\mathcal{S}_{0,0}^{Lee}(X; \alpha)$  has a basis consists of  $x_{\alpha_+, \alpha_-}$  with  $\alpha_{\pm} \in H_2(X)$ ,  $\alpha_+ + \alpha_- = \alpha$ ,  $\alpha_+ \cdot \alpha_- = 0$ . Such generators are in one-one-correspondence with  $\alpha' \in \alpha + 2\mathbb{Z}^m$  with  $\alpha'^2 = \alpha^2$ , where the correspondence is given by  $\alpha' := \alpha_+ - \alpha_-$ . The bilinear form on  $H_2(X)$  can be rewritten as  $\beta^2 = \beta^T(P - W + N - 2N_-)\beta$ . Thus by conditions (i) and (ii), we see that such  $\alpha'$  are exactly those with  $h(\alpha') = h(\alpha)$ , proving the statement for  $\mathcal{S}_{0,0}^{Lee}$ .

Write (26) in homological degree 0 for  $\bullet = Lee$  as  $\mathcal{S}_{0,0}^{Lee}(X; \alpha) = \text{colim}_r V_r$  for short. Then by Step 1 and the naturality of the Lee spectral sequence, we know  $\mathcal{S}_{0,0}^2(X; \alpha; \mathbb{Q}) = \text{colim}_r gr(V_r)$ , which equals the subspace of  $gr(\text{colim}_r V_r)$  of finite degree elements together with 0. In particular, the rank of  $\mathcal{S}_{0,0}^2(X; \alpha)$  is bounded above by the dimension of  $\mathcal{S}_{0,0}^{Lee}(X; \alpha)$ .

**Step 3:** Decomposition of  $KhR_{Lee}$  and proof of (3).

As we have already seen once in Step 1 (with  $n = \alpha + 2r$ , and renormalization),  $KhR_{Lee}^0(K^p(\alpha + r, r))$  is generated by canonical generators coming from orientations of  $K^p(\alpha + r, r)$  with writhe  $h(\alpha)$ , and we can divide these orientations according to the their algebraic cable type.

More precisely, for  $-n \leq n' \leq n$  with  $2|n - n'$ ,  $h(n') = h(\alpha)$ , let

$$KhR_{Lee, n'}^0(K^p(\alpha + r, r)) \subset KhR_{Lee}^0(K^p(\alpha + r, r))$$

be the subspace spanned by generators whose orientations make  $K^p(n)$  algebraic  $n'$ -cables of  $K$ . This subspace is an  $S_n$ -subrepresentation.

We will be interested in  $n' = \alpha$ . As in [Ren23, Proposition 4.2], we know

$$KhR_{Lee, \alpha}^0(K^p(\alpha + r, r)) \cong \bigoplus_{0 \leq s \leq r} (n - s, s) \quad (28)$$

as  $S_n$ -representations, where for  $a, b \in \mathbb{Z}_{\geq 0}^m$ ,  $a \geq b$ ,  $(a, b)$  denotes the  $S_{a+b}$ -representation given by the tensor product of the  $S_{a_i+b_i}$ -representations  $(a_i, b_i)$ . Similarly,

$$KhR_{Lee, \alpha}^0(K^p(\alpha + r + e_i, r + e_i)) \cong \bigoplus_{0 \leq s \leq r + e_i} (n + 2e_i - s, s) \quad (29)$$

as  $S_{n+2e_i}$ -representations.

By the argument in [Ren23, Section 4.3], for every  $0 \leq s \leq r$ , the subrepresentation  $(n - s, s)$  in (28) and the subrepresentation  $(n - s + e_i, s + e_i)$  in (29) have the same filtration degree. On the other hand, the dotted annular creation cobordism map restricts to a filtered degree 2 map

$$\phi_i: KhR_{Lee, \alpha}^0(K^p(\alpha + r, r)) \rightarrow KhR_{Lee, \alpha}^0(K^p(\alpha + r + e_i, r + e_i)).$$

By Step 1, the associated graded map of  $\phi_i$  is exactly the corresponding dotted annular creation map on  $KhR_2$ , which is injective by the conclusions of Step 1 and of Lemma 5.8. Thus  $\phi_i$  increases the filtration degree of every nonzero element by exactly 2. Inductively from  $r = 0$  (where  $KhR_{Lee, \alpha}^0(K^p(\alpha)) = (\alpha, 0)$  in filtration degree  $s_{\text{gl}_2}(K^p(\alpha)) + 1$  by definition), we see that the filtration degree of  $(n, 0)$  in (28) is  $s_{\text{gl}_2}(K^p(\alpha)) + 1 + 2|r|$ . Consequently by (13) (cf. (23)) we have

$$s(X; \alpha) = \lim_{r \rightarrow \infty^m} (s_{\text{gl}_2}(K^p(\alpha)) + 1 + 2|r| - |\alpha| - 2|r|) = -s(-K^p(\alpha)) - w(K^p(\alpha)) - |\alpha| + 1 > -\infty.$$

In particular,  $\mathcal{S}_{0,0,s(X;\alpha)}^2(X; \alpha) \neq 0$  by Theorem 1.12.  $\square$

*Remark 5.10.* (1) We expect that in the notation of Theorem 5.5, assuming no component of  $D$  is a 0-framed unknot, then

$$\mathcal{S}_{0,0}^2(X; \alpha) = \bigoplus_{\substack{\alpha' \in \mathbb{Z}_{\geq 0}^m \\ 2|\alpha - \alpha'| \\ h(\alpha') = h(\alpha)}} KhR_2^0(K^w(\alpha')) \left\{ \sum_i \frac{(w_i - p_i)(\alpha_i'^2 + 2\alpha_i')}{2} - \frac{w(K^w(\alpha'))}{2} - |\alpha'| \right\}. \quad (30)$$

In fact, an expanded argument of the proof above shows that the graded rank of the left hand side is bounded above by that of the right hand side. However, we have not been able to work out a proof of (30). We remark that over  $\mathbb{Q}$ , (30) is equivalent to adding to the last claim of Lemma 5.8 the conclusion that the symmetrized dotted annular creation map gives an injection

$$KhR_2^{h_{max}}(K^p(n - 2e_i); \mathbb{Q})^{S_{n-2e_i}} \hookrightarrow KhR_2^{h_{max}}(K^p(n); \mathbb{Q})^{S_n}.$$

where  $h_{max}$  is the maximal nontrivial homological degree of the right hand side.

- (2) In particular, in the setup of Theorem 1.14(1) or (2) when  $n < 0$ , one can easily show the validity of (30). Therefore, the conclusions in Theorem 1.14 for  $\mathcal{S}_{0,0}^2$  can be stated over  $\mathbb{Z}$  instead.

## 6. NONVANISHING EXAMPLES AND APPLICATIONS

In this section we present some explicit calculations and applications for Khovanov and Lee skein lasagna modules and lasagna  $s$ -invariants of some 2-handlebodies (with empty boundary link). Roughly the first half of this section utilizes the tools developed in Section 5, while the second half utilizes results in a paper of the first author [Ren23].

6.1. Knot traces of some small knots.

Knot \ Framing	-4	-3	-2	-1	0	1	2	3	4	5	6
$U$	N	N	N	N	N	V	V	V	V	V	V
$3_1$	N	?	?	V	V	V	V	V	V	V	V
$-3_1$	N	N	N	N	N	?	?	?	?	?V	V
$4_1$	N	N	N	N	N	?V	?V	V	V	V	V

TABLE 1. Vanishing/Nonvanishing of Khovanov skein lasagna modules of some knot traces. An entry N/V means the corresponding knot trace has nonvanishing/vanishing Khovanov skein lasagna module. An entry ?V means it has vanishing Lee skein lasagna module (meaning the filtration function is identically  $-\infty$ ).

**Example 6.1** (Unknot). By Theorem 1.14, for  $n \leq 0$ , the  $D^2$ -bundle  $D(n)$  over  $S^2$  with Euler number  $n$  has  $s(D(n); 0) = 0$ ,  $s(D(n); 1) = -n$ . In particular,  $\mathcal{S}_{0,0}^2(D(n); \alpha) \neq 0$  for  $\alpha = 0, 1$ .

For  $n < 0$ , more explicitly we have  $\mathcal{S}_{0,0}^2(D(n); 0) = \mathbb{Z}$  concentrating in degree 0 and  $\mathcal{S}_{0,0}^2(D(n); 1) = \mathbb{Z}^2$  concentrating in degrees  $n, n - 2$  (cf. Remark 5.10(2)).

On the other hand, for  $n > 0$ , Example 3.2 shows  $\mathcal{S}_0^2(D(n)) = 0$  and consequently all lasagna  $s$ -invariants of  $D(n)$  are  $-\infty$ .

For other examples we only mention the vanishingness/nonvanishingness of the Khovanov skein lasagna modules of the knot traces. For small knots, Corollary 1.16 (which will be proved in Section 6.9) usually provides more effective nonvanishing bounds than Theorem 1.14. We emphasize however that Corollary 1.16 does not provide an explicit  $s$ -invariant in most cases.

**Example 6.2** (Trefoils). The left-handed trefoil, or  $3_1$ , has 3 negative crossings. Therefore 1.14(1) shows  $X_n(3_1)$  has nonvanishing Khovanov skein lasagna module for  $n < -6$ .

We can do better as follows. The knot  $3_1$  can be unknotted by changing a crossing, which is realized by performing a +1 twist on two parallelly oriented strands. This means  $3_1$  is  $(-4)$ -slice in  $\mathbb{C}\mathbb{P}^2$ , thus it is  $n$ -slice in some  $k\mathbb{C}\mathbb{P}^2$  for every  $n \leq -4$ . It follows that  $X_n(3_1)$  embeds in some  $k\overline{\mathbb{C}\mathbb{P}^2}$  for  $n \leq -4$ . By Corollary 1.16,  $X_n(3_1)$  has nonvanishing Khovanov skein lasagna module for  $n \leq -4$ .

On the other hand,  $TB(-3_1) = 1$ , so  $X_n(3_1)$  has vanishing Khovanov skein lasagna module for  $n \geq -1$  by Theorem 1.4. It would be interesting to know whether  $X_n(3_1)$  has nonvanishing Khovanov skein lasagna module for  $n = -3, -2$ .

The right-handed trefoil  $-3_1$  is positive. By either Theorem 1.14 or Corollary 1.16 we know  $X_n(-3_1)$  has nonvanishing Khovanov skein lasagna module for  $n \leq 0$ . On the other hand, since  $TB(3_1) = -6$ ,  $X_n(-3_1)$  has vanishing Khovanov skein lasagna module for  $n \geq 6$ . In this case we have a larger gap of unknown. We remark that Proposition 6.8 will imply that  $X_5(-3_1)$  has vanishing Lee skein lasagna module, as there is a framed concordance in  $\overline{\mathbb{C}\mathbb{P}^2}$  from the 1-framed unknot to the 5-framed  $-3_1$ .

**Example 6.3** (Figure 8). The standard diagram of the figure 8 knot, or  $4_1$ , has 2 negative crossings. Theorem 1.14(1) shows  $X_n(4_1)$  has nonvanishing Khovanov skein lasagna module for  $n < -4$ , while Corollary 1.16 shows so for  $n \leq 0$  as  $4_1$  is 0-slice in  $\mathbb{C}\mathbb{P}^2$ . On the other hand, since  $-4_1 = 4_1$  and  $TB(4_1) = -3$ , we know  $X_n(4_1)$  has vanishing Khovanov skein lasagna module for  $n \geq 3$  by Theorem 1.4.

It would be interesting to know whether  $X_n(4_1)$  has nonvanishing Khovanov skein lasagna module for  $n = 1, 2$ . Again, Proposition 6.8 will imply they both have vanishing Lee skein lasagna modules.

## 6.2. Nonpositive definite plumbings and generalizations.

**Example 6.4** (Nonpositive plumbing tree). Let  $X$  be a nonpositive definite plumbing tree of spheres, i.e. a plumbing of  $D^2$ -bundles over 2-spheres along a tree, whose intersection form is nonpositive definite. Then we can orient the unknots in the underlying Kirby diagram so that all crossings are positive. In particular, in the notation of Theorem 5.5,  $P - W + N = P - W + N_+ - N_-$  is the intersection form of  $X$ , which is nonpositive. Therefore Theorem 5.5(3) implies  $s(X; 0) = 0$  and  $X$  has nonvanishing Khovanov skein lasagna module.

**Example 6.5** (Traces on alternating chainmail links). Let  $G$  be a chainmail graph without loops and with positive edge weights and nonpositive vertex weights, and  $K$  be its associated chainmail link, in the sense of [Ago23, Section 2]. Then  $K$  has a standard positive diagram (as an unframed link), and a nonpositive framing matrix which is equal to  $P - W + N$  for its standard diagram. This means the trace  $X$  on  $K$  has nonvanishing Khovanov skein lasagna module. This generalizes the previous example.

**Example 6.6** (Branched double cover of definite surfaces of alternating links). Suppose  $L \subset S^3$  is a nonsplit alternating link and  $S \subset B^4$  is an unoriented surface bounding  $L$  with negative definite Gordon–Litherland pairing [GL78]. Greene [Gre17] showed every choice of  $S$  comes from one checkerboard surface of an alternating diagram of  $L$  (and conversely one of the checkerboard surface of every alternating diagram is negative definite, which is much easier).

The branched double cover of  $S$  is a 4-manifold  $\Sigma(S)$  bounding the branched double cover  $\Sigma(L)$  of  $L$ . It is also considered by Ozsváth–Szabó [OS05] (denoted  $X_L$  in Section 3 there), and from the description there (see also [Gre13, Section 3.1]) we realize that  $\Sigma(S)$  is a special case of Example 6.5, hence it has nonvanishing Khovanov skein lasagna module. In particular,  $\Sigma(L)$  bounds a 2-handlebody with nonvanishing Khovanov skein lasagna module. Since  $\overline{\Sigma(L)} = \Sigma(-L)$  we see  $\Sigma(L)$  bounds such 2-handlebodies on both sides.

**Example 6.7.** One can further generalize Example 6.5 by connect-summing some components of  $K$  with 0-framed positive knots. The same argument shows the resulting trace has nonvanishing Khovanov skein lasagna module.

**6.3. More exotica from connected sums.** In this section we prove Corollary 1.2. One can show the relevant pairs of manifolds have nonisomorphic Khovanov skein lasagna modules, but we give a simpler argument by showing they have different lasagna  $s$ -invariants.

*Proof of Corollary 1.2.* Write  $W_1 := X_1^{\natural a} \natural X_2^{\natural b}$  and  $W_2 := X_1^{\natural a'} \natural X_2^{\natural b'}$ , where  $a + b = a' + b' = n$  and  $(a, b) \neq (a', b')$ .

Recall from Section 1.5 that  $s(X_1; 1) = 3$ ,  $s(X_2; 1) = 1$ , where 1 is a generator of  $H_2(X_i)$ . Since we did not specify the sign of 1, we equally have  $s(X_1; -1) = 3$ ,  $s(X_2; -1) = 1$ . By the connected sum formula (Theorem 4.7(4)),

$$s(W_1; (\epsilon_1, \dots, \epsilon_n)) = 3a + b$$

for any  $\epsilon_i = \pm 1$ . If there were a diffeomorphism  $W_1 \cong W_2$ , then by considering the intersection form we see it sends each class  $(\epsilon_1, \dots, \epsilon_n)$  to some class  $(\epsilon'_1, \dots, \epsilon'_n)$ ,  $\epsilon'_i = \pm 1$ . However,  $s(W_2; (\epsilon'_1, \dots, \epsilon'_n)) = 3a' + b' \neq 3a + b$ , a contradiction.

By (a Lee version of) [MWW23, Corollary 4.2] we know  $s(S^1 \times S^3; 0) = 0$ . Since  $H_2(S^1 \times S^3) = 0$  the above argument remains valid under connected sums with  $S^1 \times S^3$ . Since  $s(\overline{\mathbb{C}\mathbb{P}^2}; 1) = 1$  by

Example 6.1 for  $n = -1$  and Proposition 2.1, the argument remains valid under further connected sums with  $\overline{\mathbb{C}\mathbb{P}^2}$ .

By a direct calculation (or Theorem 5.5 applied to the 0-framed unknot) we know  $s(S^2 \times D^2; \alpha) = 0$  for all  $\alpha$ . Any diffeomorphism  $k(S^1 \times S^3) \# m \overline{\mathbb{C}\mathbb{P}^2} \# W_1 \natural (S^2 \times D^2)^{\natural c} \cong k(S^1 \times S^3) \# m \overline{\mathbb{C}\mathbb{P}^2} \# W_2 \natural (S^2 \times D^2)^{\natural c}$  sends each class  $(\epsilon_1, \dots, \epsilon_{m+n}, 0^c)$  to some class  $(\epsilon'_1, \dots, \epsilon'_{m+n}, \alpha_1, \dots, \alpha_c)$ . Again, the  $s$ -invariant of the latter class is not equal to the  $s$ -invariant of the former class, a contradiction.

By Example 6.4 we know the negative  $E8$  manifold has  $s(E8; 0) = 0$ . Any diffeomorphism between the above pair with  $d$  further copies of  $E8$  added sends each  $(\epsilon_1, \dots, \epsilon_{m+n}, 0^c, 0^d)$  to  $(\epsilon'_1, \dots, \epsilon'_{m+n}, \alpha_1, \dots, \alpha_c, 0^d)$ , whose  $s$ -invariants still differ, leading to a contradiction.

Finally, since the lasagna  $s$ -invariants of a 4-manifold  $X$  are invariants of the interior of  $X$  (cf. the end of Section 2.1), the interiors of any of the above pairs are not diffeomorphic.  $\square$

#### 6.4. Nonpositive shake genus of positive knots.

We prove Theorem 1.9(1).

*Proof of Theorem 1.9(1).* By Theorem 1.14(2), if  $K'$  is a positive knot and  $n \leq 0$ , then  $s(X_n(K'); 1) = s(K') - n$ . By Proposition 1.13, if  $K$  is a knot concordant to  $K'$ , then  $s(X_n(K); 1) = s(X_n(K'); 1)$ . Thus Equation (1) implies  $g_{sh}^n(K) \geq s(K')/2$ . On the other hand, Rasmussen's classical bound gives  $s(K')/2 \geq g_4(K')$ . Since  $g_4(K') = g_4(K) \geq g_{sh}^n(K)$ , all these quantities are equal.  $\square$

**6.5. Another comparison result for lasagna  $s$ -invariants.** In this section we make use the adjunction inequality for  $s$ -invariants [Ren23, Corollary 1.4] to deduce a comparison result for lasagna  $s$ -invariants in the spirit of Proposition 5.3. Then we prove Theorem 1.9(2). For simplicity, assume the boundary link  $L$  is empty. Instead of working on the 4-manifold level as in Section 5.2 (which is more invariant), we work on the cabled links level directly.

As in the setup of Section 5.2, assume  $X$  is a 2-handlebody obtained by attaching 2-handles to a framed link  $K = K_1 \cup \dots \cup K_m \subset S^3 = \partial B^4$ , and  $X'$  is one obtained by attaching handles to  $K' = K'_1 \cup \dots \cup K'_m$ .

**Proposition 6.8.** *Suppose  $C: K \rightarrow K'$  is a framed concordance in  $\overline{k\mathbb{C}\mathbb{P}^2}$  that restricts to concordances  $C_i: K_i \rightarrow K'_i$ . Then*

$$s(X'; \alpha) \leq s(X; \alpha) - |[C(\alpha)]|$$

for any  $\alpha \in H_2(X) \cong H_2(X') \cong \mathbb{Z}^m$ , where  $[C(\alpha)] = \sum_i \alpha_i [C_i] \in H_2(\overline{k\mathbb{C}\mathbb{P}^2}) \cong \mathbb{Z}^k$ .

*Proof.* The adjunction inequality for the  $s$ -invariant [Ren23, Corollary 1.4] in the  $\mathfrak{gl}_2$  renormalization states that if  $\Sigma: L \rightarrow L'$  is a framed cobordism in  $\overline{k\mathbb{C}\mathbb{P}^2}$ , such that each component of  $\Sigma$  has a boundary on  $L$ , then

$$s_{\mathfrak{gl}_2}(L') \leq s_{\mathfrak{gl}_2}(L) - \chi(\Sigma) - |[\Sigma]|. \quad (31)$$

Now, the  $(\alpha^+ + r, \alpha^- + r)$ -cable of  $C$  gives a concordance between  $K(\alpha^+ + r, \alpha^- + r)$  and  $K'(\alpha^+ + r, \alpha^- + r)$  with homology class  $[C(\alpha)]$ , which by (31) implies

$$s_{\mathfrak{gl}_2}(K'(\alpha^+ + r, \alpha^- + r)) \leq s_{\mathfrak{gl}_2}(K(\alpha^+ + r, \alpha^- + r)) - |[C(\alpha)]|.$$

The statement now follows from Proposition 5.4.  $\square$

Now we prove Theorem 1.9(2).




FIGURE 2. Satellite patterns  $P_{n,m}$  (left) and  $Q_{n,m}$  (right). The boxes denote full twists. The shaded band gives rise to a concordance between  $Q_{n,m}$  and the identity pattern.

*Proof of Theorem 1.9(2).* Putting a framing on  $\Sigma$  makes it a framed cobordism from  $(K, n)$  to  $(K', n - [\Sigma]^2)$ . It follows from Proposition 6.8 that  $s(X_n(K); 1) \geq s(X_{n-[\Sigma]^2}(K'); 1) + |[\Sigma]|$ . If  $n \leq [\Sigma]^2$ , by Theorem 1.14 we know  $s(X_{n-[\Sigma]^2}(K'); 1) = s(K') - n + [\Sigma]^2$ , thus the statement follows.  $\square$

**6.6. Yasui's family of knot trace pairs.** Yasui [Yas15, Figure 10] defined two families of satellite patterns  $P_{n,m}$ ,  $Q_{n,m}$ ,  $n, m \in \mathbb{Z}$  (see Figure 2). He showed that for any  $n, m$  and any knot  $K \subset S^3$ , the two manifolds  $X_n(P_{n,m}(K))$  and  $X_n(Q_{n,m}(K))$ , namely the  $n$ -traces on the two corresponding satellite knots, are homeomorphic. Moreover, he showed that if there exists a Legendrian representative  $\mathcal{K}$  of  $K$  with

$$tb(\mathcal{K}) + |\text{rot}(\mathcal{K})| - 1 = 2g_4(K) - 2, \quad n \leq tb(\mathcal{K}), \quad m \geq 0, \quad (32)$$

then  $X_n(P_{n,m}(K))$  and  $X_n(Q_{n,m}(K))$  are not diffeomorphic, thus form an exotic pair [Yas15, Theorem 4.1]. In fact, under these conditions, the exotica were detected by a discrepancy of shake slice genera:  $g_{sh}^n(P_{n,m}(K)) = g_4(K) + 1 > g_{sh}^n(Q_{n,m}(K))$ . Thus, the interiors of such pairs are also exotic. The case  $K = U$ ,  $n = -1$ ,  $m = 0$  yields the exotic pair  $X_{-1}(-5_2)$  and  $X_{-1}(P(3, -3, -8))$  in Theorem 1.1.

Using lasagna  $s$ -invariants, we present a different family (as stated in Theorem 1.3) of Yasui's knot trace pairs that are exotic. We recall that our hypothesis on  $(K, n, m)$  is that there exists a slice disk  $\Sigma$  of  $K$  in some  $k\mathbb{C}\mathbb{P}^2$  such that

$$s(K) = |[\Sigma]| - [\Sigma]^2, \quad n < -[\Sigma]^2, \quad m \geq 0. \quad (33)$$

Before giving the proof, we compare and discuss the conditions imposed on  $K$  by Yasui (32) and by us (33).



The adjunction inequality for the  $s$ -invariant [Ren23, Corollary 1.5] applied to the slice disk  $\Sigma$  implies that  $s(K) \geq |[\Sigma]| - [\Sigma]^2$ . Therefore the first condition in (33) is that this adjunction inequality attains equality for  $\Sigma$ . Similarly, the first condition in (32) is that the slice-Bennequin inequality attains equality for  $\mathcal{K}$ . These two conditions are satisfied on somewhat different families of knots. For example, as remarked by Yasui, his condition on  $K$  is satisfied for all positive torus knots. On the other hand, our condition on  $K$  is satisfied for all negative torus knots (cf. Lemma 6.10). Our condition also has the advantage of being closed under concordances and connected sums of knots.

*Proof of Theorem 1.3.* As observed by Yasui, adding a band as indicated in Figure 2 and capping off the unknot component give a concordance between  $Q_{n,m}$  and the identity satellite pattern. Therefore,  $Q_{n,m}(K)$  is concordant to  $K$ , thus by Proposition 1.13 and Proposition 5.4 we have

$$\begin{aligned} s(X_n(Q_{n,m}(K)); 1) &= s(X_n(K); 1) \\ &= -\lim_{r \rightarrow \infty} (s(-K(r+1, r)) + 2r) - n - 1 + 1 \leq -s(-K(1, 0)) - n = s(K) - n. \end{aligned} \quad (34)$$

Deleting a local ball in  $B^4$  near  $\Sigma$ , turning the cobordism upside down and reversing the ambient orientation, we obtain a concordance  $-\Sigma^\circ: K \rightarrow U$  in  $k\overline{\mathbb{C}\mathbb{P}^2}$ . Applying the satellite operation  $P_{n,m}$  gives a framed concordance  $C_1 = P_{n,m}(-\Sigma^\circ): (P_{n,m}(K), n) \rightarrow (P_{n+[\Sigma]^2, m}(U), n+[\Sigma]^2)$  in  $k\overline{\mathbb{C}\mathbb{P}^2}$  with  $[C_1] = [-\Sigma^\circ]$  since  $P_{n,m}$  has winding number 1. Since  $n < -[\Sigma]^2$ , there is a framed concordance  $C_2: (P_{n+[\Sigma]^2, m}(U), n+[\Sigma]^2) \rightarrow (P_{-1, m}(U), -1)$  in  $k_1\overline{\mathbb{C}\mathbb{P}^2}$  obtained by adding  $k_1$  positive twists along the meridian of the satellite solid torus, which has  $[C_2] = (1, \dots, 1) \in \mathbb{Z}^{k_1} = H_2(k_1\overline{\mathbb{C}\mathbb{P}^2})$ , where  $k_1 = -1 - n - [\Sigma]^2$ . Similarly, there is a null-homologous framed concordance  $C_3: (P_{-1, m}(U), -1) \rightarrow (P_{-1, 0}(U), -1) = (-5_2, -1)$  in  $k_2\overline{\mathbb{C}\mathbb{P}^2}$  where  $k_2 = m$  (note that the two strands being twisted  $-m$  times in  $P_{n,m}$  are oppositely oriented). Now Proposition 6.8 applied to the concordance  $C_1 \cup C_2 \cup C_3: (P_{n,m}(K), n) \rightarrow (-5_2, -1)$  yields

$$\begin{aligned} s(X_n(P_{n,m}(K)); 1) &\geq s(X_{-1}(-5_2); 1) + |[C_1]| + |[C_2]| + |[C_3]| \\ &= 3 + |[\Sigma]| + k_1 + 0 = |[\Sigma]| - [\Sigma]^2 - n + 2 = s(K) - n + 2. \end{aligned} \quad (35)$$

Together, (34) and (35) imply that  $s(X_n(P_{n,m}(K)); 1) > s(X_n(Q_{n,m}(K)); 1)$ , proving the theorem.  $\square$

*Remark 6.9.* Let  $C \cup K$  be a two-component link in  $S^3$ , where  $C$  is an unknot and  $\ell k(C, K) = 1$ . Regarding  $C$  as a dotted 1-handle and attaching an  $n$ -framed 2-handle along  $P_{n,m}(K)$  (resp.  $Q_{n,m}(K)$ ) yields a Mazur manifold  $Z_P$  (resp.  $Z_Q$ ). The manifolds  $Z_P$  and  $Z_Q$  are homeomorphic, and Hayden-Mark-Piccirillo [HMP21, Theorem 2.7] showed that if  $C \cup K$  satisfies the (very mild) assumptions that

- $K$  is not the meridian of  $C$ ,
- every self homeomorphism of  $\partial Z_P$  preserves the JSJ torus (with orientation) given by the image of the satellite torus of  $P_{n,m}(K)$  in  $\partial Z_P$ ,

then  $Z_P$  being diffeomorphic to  $Z_Q$  implies that  $X_n(P_{n,m}(K))$  is diffeomorphic to  $X_n(Q_{n,m}(K))$ . Therefore, by choosing suitable  $C \cup K$ , Theorem 1.3 also yields various families of exotic Mazur manifolds.

As an example, we present a family of knots  $K$  for which Theorem 1.3 applies.

**Lemma 6.10.** *For every negative torus knot  $-T_{p,q}$ , there exists a slice disk  $\Sigma$  in some  $k\overline{\mathbb{C}\mathbb{P}^2}$  with  $|[\Sigma]| - [\Sigma]^2 = -(p-1)(q-1) = s(-T_{p,q})$ .*

*Proof.* We induct on  $p + q$ . For  $p = q = 1$ ,  $-T_{p,q} = U$  is the unknot and we take  $\Sigma$  to be the trivial slice disk in  $B^4$ .

For the induction step, assume  $p > q \geq 1$ . There is a positive twist along  $q$  strands in  $-T_{p,q}$  that converts it into  $-T_{p-q,q}$ . This gives a concordance  $C: -T_{p-q,q} \rightarrow -T_{p,q}$  in  $\mathbb{C}\mathbb{P}^2$  with  $[C] = q[\mathbb{C}\mathbb{P}^1]$ . By induction hypothesis, there is a slice disk  $\Sigma'$  of  $-T_{p-q,q}$  in  $k'\mathbb{C}\mathbb{P}^2$  with  $||[\Sigma']| - [\Sigma']^2 = -(p-q-1)(q-1)$ . Then  $\Sigma = \Sigma' \cup C$  is a slice disk of  $-T_{p,q}$  in  $(k'+1)\mathbb{C}\mathbb{P}^2$  with  $||[\Sigma]| - [\Sigma]^2 = -(p-1)(q-1)$ .  $\square$

**6.7. Conway knot cables can obstruct its sliceness.** Recently, Piccirillo [Pic20] famously proved that the Conway knot, denoted *Conway*, is not slice. Building on her proof, we show the following.

**Proposition 6.11.** *Let  $Conway(n^+, n^-)$  denote the 0-framed  $(n^+ + n^-)$ -cable of the Conway knot, with the orientation on  $n^-$  of the strands reversed. Then for some  $n > 0$  we have*

$$s(Conway(n+1, n)) > -2n = s(U^{\sqcup(2n+1)}).$$

Here  $U^{\sqcup m}$  is the  $m$ -component unlink.

If *Conway* were slice, then any cable of it is concordant to the corresponding cable of the unknot, implying that they have the same  $s$ -invariant. Therefore, Proposition 6.11 shows that, if people had enough computer power, we could have obstructed the sliceness of the Conway knot by calculating the  $s$ -invariants of its cables. In practice, however, the calculation of  $s(Conway(2, 1))$  seems already out of reach.

*Proof.* Piccirillo [Pic20] found a knot  $K'$  with  $X_0(K') = X_0(Conway)$  and  $s(K') = 2$ . Let  $X$  be the mirror image of this common trace, i.e.  $X = X_0(-Conway) = X_0(-K')$ . By Proposition 5.4, we have

$$s(X; 1) = -\lim_{n \rightarrow \infty} (s(K'(n+1, n)) + 2n) \leq -s(K'(1, 0)) = -s(K') = -2.$$

Again, by Proposition 5.4 we have  $\lim_{n \rightarrow \infty} (s(Conway(n+1, n)) + 2n) = -s(X'; 1) \geq 2$  and hence  $s(Conway(n+1, n)) + 2n \geq 2 > 0$  for all sufficiently large  $n$ .  $\square$

**6.8. Knot traces of torus knots.** The first author [Ren23] calculated the  $s$ -invariants of all torus links with various orientations. In this section, we use this to determine the complete structure of the Lee skein lasagna modules of large negative traces on negative torus links.

For  $d = r + s > 0$  and coprime  $p, q \neq 0$ , let  $T(dp, dq)_{r,s}$  denote the torus link  $T(dp, dq)$  equipped with an orientation where  $r$  of the strands are oriented against the other  $s$  strands, we have

**Lemma 6.12** ([Ren23, Theorem 1.1, Corollary 1.3]). *If  $p, q > 0$ , then*

$$s(T(dp, dq)_{r,s}) = (p|r-s| - 1)(q|r-s| - 1) - 2\min(r, s).$$

*If  $pq < 0$ , then*

$$s(T(dp, dq)_{r,s}) = C(d, p, q, |r-s|)$$

*for some constant  $C$  depending on  $d, p, q, |r-s|$ .*  $\square$

Since  $T(dp, dq)_{r,s}$  is the  $(r, s)$ -cable of  $T(p, q)$  with framing  $pq$ , the first part of Lemma 6.12 and Proposition 5.4 imply that for  $p, q > 0$ ,  $\alpha \in H_2(X_{-pq}(-T(p, q))) \cong \mathbb{Z}$ ,

$$s(X_{-pq}(-T(p, q)); \alpha) = (p+q-1)|\alpha|.$$

More generally, for any  $k \geq pq$ , the adjunction inequality [Ren23, Corollary 1.4] for  $s$ -invariant implies that the  $k$ -framed  $(r, s)$ -cable of  $T(p, q)$ , denoted  $T(p, q)^k(r, s)$ , satisfies

$$\begin{aligned} s(T(p, q)^k(r, s)) &\leq s(T(dp, dq)_{r,s}) + (k - pq)(|r - s|^2 - |r - s|) \\ &= k|r - s|^2 - (k - pq + p + q)|r - s| - 2 \min(r, s) + 1, \end{aligned}$$

which is sharp since

$$\begin{aligned} s(T(p, q)^k(r, s)) &\geq s(T(p, q)^k(|r - s|, 0)) - 2 \min(r, s) \\ &= s(T(p, q)^{pq}(|r - s|, 0)) + (k - pq)(|r - s|^2 - |r - s|) - 2 \min(r, s) \\ &= k|r - s|^2 - (k - pq + p + q)|r - s| - 2 \min(r, s) + 1 \end{aligned}$$

where the first equality is because  $T(p, q)^k(|r - s|, 0)$  can be obtained by adding  $k - pq$  full twists to a positive diagram of  $T(p, q)^{pq}(|r - s|, 0)$  on  $|r - s|$  strands, and for positive link diagrams the  $s$ -invariant equals to the writhe minus twice the number of Seifert circles plus 1.

It follows that for any  $p, q > 0$ ,  $k \geq pq$  and  $\alpha \in H_2(X_{-k}(-T(p, q))) \cong \mathbb{Z}$ ,

$$s(X_{-k}(-T(p, q)); \alpha) = (k - pq + p + q - 1)|\alpha|. \quad (36)$$

Unfortunately, (36) does not yield useful genus bounds via Corollary 4.13.

Theorem 4.1 states that the Lee skein lasagna module of  $X_{-k}(-T(p, q))$  is generated by Lee canonical generators  $x_{\alpha^+, \alpha^-}$ ,  $\alpha^+, \alpha^- \in H_2(X_{-k}(-T(p, q)))$ . The generator  $x_{\alpha^+, \alpha^-}$  has homological degree  $-2\alpha^+ \cdot \alpha^-$  ( $\cdot$  is the intersection form on  $H_2$  instead of the product on  $\mathbb{Z}$ ) and homology class degree  $\alpha^+ + \alpha^-$ . Moreover, (36) and (21) imply it has quantum filtration degree

$$s(X_{-k}(-T(p, q)); \alpha^+ - \alpha^-) + 2\alpha^+ \cdot \alpha^- = (k - pq + p + q - 1)|\alpha^+ - \alpha^-| + 2\alpha^+ \cdot \alpha^-.$$

These data (together with Theorem 4.1(4)) completely determine the structure of  $\mathcal{S}_0^{Lee}(X_{-k}(-T(p, q)))$ . More explicitly, we have the following

**Proposition 6.13.** *If  $p, q > 0$  are coprime and  $k \geq pq$ , then  $s(X_{-k}(-T(p, q)); \alpha) = (k - pq + p + q - 1)|\alpha|$ . Therefore, the associated graded vector space of  $\mathcal{S}_0^{Lee}(X_{-k}(-T(p, q)))$  is given by*

$$gr_*(\mathcal{S}_{0, (\beta^2 - \alpha^2)/2}^{Lee}(X_{-k}(-T(p, q)); \alpha)) = \mathbb{Q}$$

for

$$\beta \equiv \alpha \pmod{2}, \quad * = \begin{cases} \frac{\alpha^2 - \beta^2}{2} + (k - pq + p + q - 1)|\beta| - 1 \pm 1, & \beta \neq 0 \\ \frac{\alpha^2}{2}, & \beta = 0, \end{cases}$$

and 0 in other tri-degrees. □

In the standard diagram, the number of negative crossings of  $-T(p, q)$  is  $pq - p$  or  $pq - q$ . Thus the nonvanishing bound on  $k$  given by Proposition 6.13 is better than that by Theorem 1.14. However, Corollary 1.16 gives an even better bound. For example,  $-T(n, n + 1)$  is  $n^2$ -slice in  $\mathbb{C}\mathbb{P}^2$ , thus  $X_{-k}(-T(n, n + 1))$  has nonvanishing Lee and Khovanov skein lasagna module for  $k \geq n^2$ .

Finally we note that the second part of Lemma 6.12 implies that for  $p, q > 0$ ,  $s(X_{pq}(T(p, q)); \alpha) = -\infty$  for all  $\alpha$ . However, since  $TB(-T(p, q)) = -pq$  [EH01], we already know from Corollary 5.1 that  $s(X_k(T(p, q)); \alpha) = -\infty$  for all  $k \geq pq$  and all  $\alpha$ .

**6.9. Expectations and calculations for  $\overline{\mathbb{C}\mathbb{P}^2}$ .** The 2-handlebody formula (13) expresses the Khovanov skein lasagna module of  $\overline{\mathbb{C}\mathbb{P}^2}$  as a colimit

$$\mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; \alpha; \mathbb{Q}) = \operatorname{colim}_{r \rightarrow \infty} \operatorname{Kh}R_2(-T(|\alpha| + 2r, |\alpha| + 2r)_{\alpha_+, \alpha_-, +r}; \mathbb{Q})^{S_{|\alpha|+2r}} \{-|\alpha| - 2r\} \quad (37)$$

along the symmetrized dotted cobordism maps. Conjecture 6.1 of [Ren23] would imply that these morphisms are injective.

By the argument in [Ren23, Section 4.3], the structure of  $\operatorname{Kh}R_2(T(n, n); \mathbb{Q})$  as an  $S_n$ -representation is determined by the structure of  $\operatorname{Kh}R_2(T(n', n'); \mathbb{Q})^{S_{n'}}$  for  $n' \leq n$ . Using this fact, one can check Conjecture 6.1 (equivalently Conjecture 6.1') of [Ren23] is equivalent to the following complete determination of  $\mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; \mathbb{Q})$  (Note that by (14), all  $\mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; \alpha; \mathbb{Q})$  are determined from those with  $\alpha = 0, 1$ ).

**Conjecture 6.14.** *Let  $K_n(t, q)$  denote the Poincaré polynomial of  $\operatorname{Kh}(T(n, n-1); \mathbb{Q})$  for  $n > 0$ . Then  $\mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; 0; \mathbb{Q})$  has Poincaré polynomial*

$$1 + \sum_{n=1}^{\infty} t^{-2n^2} q^{6n^2-4n+1} K_{2n}(t, q^{-1}) \quad (38)$$

and  $\mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; 1; \mathbb{Q})$  has Poincaré polynomial

$$\sum_{n=1}^{\infty} t^{-2n^2+2n} q^{6n^2-10n+4} K_{2n-1}(t, q^{-1}). \quad (39)$$

In fact,  $K_n(t, q)$  also admits a conjectural recursive formula due to Shumakovitch and Turner (see [Ren23, Conjecture 6.2]). We have verified that this conjecture as well as Conjecture 6.14 conform with the data of  $\operatorname{Kh}(T(n, n))$  for  $n \leq 8$ .

In the rest of the section we present some calculations available for  $\overline{\mathbb{C}\mathbb{P}^2}$ .

Firstly, since  $\overline{\mathbb{C}\mathbb{P}^2}$  is  $X_{-1}(-T(1, 1))$  with a 4-handle attached, Proposition 6.13 and Proposition 2.1 give

$$gr_{2p^2+2p} \mathcal{S}_{0, -2p^2}^{Lee}(\overline{\mathbb{C}\mathbb{P}^2}; 0) = \mathbb{Q}, \quad p \geq 0,$$

which by Theorem 1.12 implies that  $\operatorname{rank}(\mathcal{S}_{0, -2p^2, 2p^2+2p}^2(\overline{\mathbb{C}\mathbb{P}^2}; 0)) \geq 1$ .

On the other hand, one can check that the only contribution to the term  $t^{-2p^2} q^{2p^2+2p}$  in (38) comes from the  $n = p$  term in which  $K_{2p}(t, q^{-1}) = q^{-(4p^2-6p+1)} + q^{-(4p^2-6p+3)} + \dots$  where  $\dots$  are terms with higher  $t$ -degrees (cf. [Kho03, Proposition 6.1], [Ren23, Theorem 2.1(2)]). Thus Conjecture 6.14 predicts  $\mathcal{S}_{0, -2p^2, 2p^2+2p}^2(\overline{\mathbb{C}\mathbb{P}^2}; 0; \mathbb{Q}) = \mathbb{Q}$ .

In fact, we can do better by directly resorting to [Ren23, Theorem 2.1(1)] to see that the generators of  $\operatorname{Kh}^{2(n+p)(n-p), q_{2n, 2n}(2(n+p)(n-p))}(T(2n, 2n)) \cong \mathbb{Z}$  in the notation there after renormalization survive under the colimit of the 2-handlebody formula (37), thus descend to generators of

$$\mathcal{S}_{0, -2p^2, 2p^2+2p}^2(\overline{\mathbb{C}\mathbb{P}^2}; 0) \cong \mathbb{Z}, \quad p \geq 0.^{14}$$

Similarly,

$$\mathcal{S}_{0, -2p^2+2p, 2p^2-1}^2(\overline{\mathbb{C}\mathbb{P}^2}; 1) \cong \mathbb{Z}, \quad p \geq 1,$$

<sup>14</sup>More precisely, since 2 is not invertible in  $\mathbb{Z}$ , Proposition 2.6 only gives the result  $\mathbb{Z}/N$  for some  $N$ . However  $N$  is necessarily 0 since we must have  $\mathbb{Z}/N \otimes \mathbb{Q} = \mathbb{Q}$ .

and a generator comes from generators of  $Kh^{2(n+p-1)(n-p), q_{2n-1, 2n-1}(2(n+p-1)(n-p))}(T(2n-1, 2n-1)) \cong \mathbb{Z}$ . This verifies Conjecture 6.14 in these tri-gradings. Together with Proposition 6.13 for  $p = q = k = 1$ , we have proved Proposition 1.15.

*Proof of Corollary 1.16.* By the connected sum formula for lasagna  $s$ -invariants (Theorem 4.7(4)) and Proposition 1.15,  $s(k\overline{\mathbb{CP}^2}; \alpha) = |\alpha|$ . Thus by Proposition 1.11 we have  $s(X; \alpha) \geq |i_*\alpha|$ . If  $X$  is a 2-handlebody, Proposition 5.4 implies  $s(X; 0) \leq 0$ , proving the claim on the equality. The claim that  $\mathcal{S}_{0,0,q}^2(X; \alpha) \neq 0$  for  $q = s(X; \alpha)$  follows from Theorem 1.12, and for  $q = s(X; \alpha) - 2$ ,  $\alpha \neq 0$  follows moreover from Theorem 4.1(4).  $\square$

When  $X$  is a 2-handlebody, one could also prove Corollary 1.16 using Proposition 6.8.

For the purpose of the next section, we prove the following local finiteness result for  $\mathcal{S}_0^2(\overline{\mathbb{CP}^2}; \mathbb{Q})$ .

**Proposition 6.15.** *When evaluating at  $q = 1$ , the Poincaré polynomial of  $\mathcal{S}_0^2(\overline{\mathbb{CP}^2}; 0; \mathbb{Q})$  is bounded above by (38), and the Poincaré polynomial of  $\mathcal{S}_0^2(\overline{\mathbb{CP}^2}; 1; \mathbb{Q})$  is bounded above by (39). In particular, the total dimension of  $\mathcal{S}_{0, \geq h_0, *}^2(\overline{\mathbb{CP}^2}; \alpha; \mathbb{Q})$  is finite for every  $h_0 \in \mathbb{Z}$ ,  $\alpha \in H_2(\overline{\mathbb{CP}^2}) \cong \mathbb{Z}$ .*

*Proof.* [Ren23, Theorem 2.1(2)] implies that the highest  $t$ -degree of  $t^{-\lfloor n^2/2 \rfloor} K_n$  is at most  $-(n-1)/2$ , which tends to  $-\infty$  as  $n \rightarrow +\infty$ . Thus the last finiteness statement for  $\alpha = 0, 1$  follows from the upper bound statement. The case for general  $\alpha$  follows from (14).

Let  $L_n, M_n$  denote the Poincaré polynomial of  $Kh(T(n, n); \mathbb{Q})$ ,  $Kh(T(n, n); \mathbb{Q})^{S_n}$ , respectively. The argument in [Ren23, Section 4.3] implies

$$L_n = \sum_{i=0}^{\lfloor n/2 \rfloor} \dim(n-i, i) t^{2i(n-i)} q^{6i(n-i)} M_{n-2i}.$$

Define  $M'_n$  inductively by  $M'_0 = M_0$ ,  $M'_1 = M_1$ ,  $M'_n = t^{2n-2} q^{6n-8} M'_{n-2} + q^{n-1} K_n$  for  $n \geq 2$ . Define

$$L'_n = \sum_{i=0}^{\lfloor n/2 \rfloor} \dim(n-i, i) t^{2i(n-i)} q^{6i(n-i)} M'_{n-2i}.$$

It follows from a calculation that Conjecture 6.1' in [Ren23] is equivalent to the assertion that  $M_n = M'_n$ . In any case, it holds true that  $L_n \leq L'_n$ .

By the 2-handlebody formula (37) and taking into account the renormalizations, the Poincaré polynomial of  $\mathcal{S}_0^2(\overline{\mathbb{CP}^2}; 0; \mathbb{Q})$  is bounded above by

$$\liminf_{n \rightarrow \infty} t^{-2n^2} q^{6n^2-2n} M_{2n}(t, q^{-1}), \quad (40)$$

while (38) is equal to

$$\lim_{n \rightarrow \infty} t^{-2n^2} q^{6n^2-2n} M'_{2n}(t, q^{-1}).$$

Here the limits are interpreted coefficientwisely, and the terms in the second limit are nondecreasing.

Now we collapse the quantum grading by setting  $q = 1$ . Assume for contrary that the coefficient of some  $t^m$  in (40) at  $q = 1$  is larger than that in (38). Denote by  $A, B$  these coefficients, respectively,

so  $A > B$ . Then  $\text{coef}_{t^m}(t^{-2n^2} M'_{2n}(t, 1)) \leq B$  for all  $n$  and  $\text{coef}_{t^m}(t^{-2n^2} M_{2n}(t, 1)) \geq A$  for all  $n \geq N$  for some  $N$ . It follows that

$$\begin{aligned} \text{coef}_{t^{2n^2+m}}(L'_{2n}(t, 1)) &= \sum_{i=0}^n \dim(2n-i, i) \text{coef}_{t^m}(t^{-2(n-i)^2} M'_{2n-2i}(t, 1)) \\ &\leq B \sum_{i=0}^n \dim(2n-i, i) = B \binom{2n}{n}. \end{aligned} \quad (41)$$

Similarly, for  $n > N$ ,

$$\begin{aligned} \text{coef}_{t^{2n^2+m}}(L_{2n}(t, 1)) &= \sum_{i=0}^n \dim(2n-i, i) \text{coef}_{t^m}(t^{-2(n-i)^2} M_{2n-2i}(t, 1)) \\ &\geq A \sum_{i=0}^{n-N} \dim(2n-i, i) = A \binom{2n}{n-N}. \end{aligned} \quad (42)$$

Since  $A > B$ , for  $n$  sufficiently large, (42) is larger than (41), contradicting the fact that  $L_{2n} \leq L'_{2n}$ . This proves the statement for  $\mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; 0; \mathbb{Q})$ . The statement for  $\mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; 1; \mathbb{Q})$  is analogous.  $\square$

**6.10.  $\overline{\mathbb{C}\mathbb{P}^2}$ -stably standard homotopy spheres.** We show that the Khovanov skein lasagna module, at least over  $\mathbb{Q}$ , is unable to detect potential exotic 4-spheres that dissolve under some copies of  $\overline{\mathbb{C}\mathbb{P}^2}$ . Note homotopy 4-spheres coming from Gluck twists dissolve under one  $\overline{\mathbb{C}\mathbb{P}^2}$ .

**Proposition 6.16.** *If  $\Sigma$  is a homotopy 4-sphere with  $\Sigma \# k\overline{\mathbb{C}\mathbb{P}^2} = k\overline{\mathbb{C}\mathbb{P}^2}$  for some  $k$ , then  $\mathcal{S}_0^2(\Sigma; \mathbb{Q}) = \mathbb{Q}$ , concentrated in tri-degree  $(0, 0, 0)$ .*

*Proof.* Apply  $\mathcal{S}_0^2$  to  $\Sigma \# k\overline{\mathbb{C}\mathbb{P}^2} = k\overline{\mathbb{C}\mathbb{P}^2}$  and use the connected sum formula (Proposition 2.2), we obtain  $\mathcal{S}_0^2(\Sigma; \mathbb{Q}) \otimes \mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; \mathbb{Q})^{\otimes k} \cong \mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; \mathbb{Q})^{\otimes k}$ . Since  $\mathcal{S}_0^2(\overline{\mathbb{C}\mathbb{P}^2}; \mathbb{Q})$  is nonzero and satisfies a suitable finiteness condition given by Proposition 6.15, we conclude that  $\mathcal{S}_0^2(\Sigma; \mathbb{Q}) = \mathbb{Q}$ .  $\square$

**6.11. Induced map of link cobordisms in  $k\overline{\mathbb{C}\mathbb{P}^2}$ .** Unless otherwise stated, in this section we work over a field  $\mathbb{F}$ , which will be dropped from the notation throughout.

We begin with a general construction. Suppose  $X$  is a closed 4-manifold and  $L_0, L_1 \subset S^3$  are two framed oriented links. A cobordism from  $L_0$  to  $L_1$  in  $X$  is a properly embedded framed oriented surface  $\Sigma \subset X^\circ := X \setminus (\text{int}(B^4) \sqcup \text{int}(B^4))$  with  $\partial\Sigma = -L_0 \sqcup L_1 \subset S^3 \sqcup S^3$ . Thus  $\Sigma$  is a skein in  $X^\circ$  rel  $-L_0 \sqcup L_1$  without input balls, which itself defines a lasagna filling with homological degree 0, quantum degree  $-\chi(\Sigma)$ , and homology class degree  $[\Sigma]$ , which represents a class  $x_\Sigma \in \mathcal{S}_{0,0,-\chi(\Sigma)}^2(X^\circ; -L_0 \sqcup L_1; [\Sigma])$ .

On the other hand, since  $(X^\circ, -L_0 \sqcup L_1) = (X, \emptyset) \# (B^4, -L_0) \# (B^4, L_1)$ , the connected sum formula (Proposition 2.2) and Proposition 2.4 implies that over a field  $\mathbb{F}$ ,

$$\begin{aligned} \mathcal{S}_0^2(X^\circ; -L_0 \sqcup L_1; [\Sigma]) &\cong \mathcal{S}_0^2(X; [\Sigma]) \otimes \mathcal{S}_0^2(B^4; -L_0) \otimes \mathcal{S}_0^2(B^4; L_1) \\ &\cong \mathcal{S}_0^2(X; [\Sigma]) \otimes \text{KhR}_2(-L_0) \otimes \text{KhR}_2(L_1) \\ &\cong \mathcal{S}_0^2(X; [\Sigma]) \otimes \text{Hom}(\text{KhR}_2(L_0), \text{KhR}_2(L_1)). \end{aligned}$$

If  $\theta \in (\mathcal{S}_0^2(X; [\Sigma]))_{h_0, q_0}^*$  is a dual Khovanov skein lasagna class, then we obtain a homomorphism

$$\text{KhR}_2(\Sigma, \theta) := \theta(x_\Sigma): \text{KhR}_2^{*,*}(L_0) \rightarrow \text{KhR}_2^{*+h_0, *+q_0-\chi(\Sigma)}(L_1).$$

In the case when  $X = \overline{\mathbb{CP}^2}$ , there are natural dual lasagna classes in each homological degree. In fact, by Proposition 1.15, for each  $\alpha \in H_2(\overline{\mathbb{CP}^2})$ , we have  $\mathcal{S}_{0,0,|\alpha|}^2(\overline{\mathbb{CP}^2}; \alpha; \mathbb{Z}) = \mathbb{Z}$ . Therefore over a field  $\mathbb{F}$  and up to sign,  $(\mathcal{S}_0^2(\overline{\mathbb{CP}^2}; \alpha))_{0,-|\alpha|}^* \cong \mathbb{F}$  has a canonical dual lasagna generator  $\theta_\alpha$ . Thus for a cobordism  $\Sigma: L_0 \rightarrow L_1$  in  $\overline{\mathbb{CP}^2}$ , there is an induced map

$$KhR_2(\Sigma) := KhR_2(\Sigma, \theta_{[\Sigma]}): KhR_2^{*,*}(L_0) \rightarrow KhR_2^{*,*-\chi(\Sigma)-[|\Sigma|]}(L_1)$$

defined up to sign. Translate back to the more familiar Khovanov homology and reverse the ambient orientation, we see if  $\Sigma: L_0 \rightarrow L_1$  is a cobordism in  $\mathbb{CP}^2$  between oriented links, then there is an induced map  $Kh(\Sigma): Kh(L_0) \rightarrow Kh(L_1)$  defined up to sign, with bidegree  $(0, \chi(\Sigma) - [\Sigma]^2 + [|\Sigma|])$ , at least over a field. This map only depends on the isotopy class of  $\Sigma$  rel boundary.

In fact, we can also give the following more direct construction of  $Kh(\Sigma): Kh(L_0) \rightarrow Kh(L_1)$  over  $\mathbb{Z}$ , which is proposed to us by Hongjian Yang. By transversality, suppose  $\Sigma$  intersects the core  $\mathbb{CP}^1$  of  $\mathbb{CP}^2$  transversely, in some  $p$  points positively and  $q$  points negatively. A tubular neighborhood of  $\mathbb{CP}^1$  has boundary  $S^3$  on which  $\Sigma$  is the torus link  $-T(p+q, p+q)_{p,q} \subset S^3$  (the torus link  $-T(p+q, p+q)$  equipped with an orientation where  $p$  of the strands are oriented against the other  $q$  strands). Tubing this boundary  $S^3$  with the negative boundary of  $(\mathbb{CP}^2)^\circ$  changes  $\Sigma$  into a cobordism  $\Sigma^\circ: L_0 \sqcup (-T(p+q, p+q)_{p,q}) \rightarrow L_1$  in  $S^3 \times I$ , inducing a map

$$Kh(\Sigma^\circ): Kh(L_0) \otimes Kh(-T(p+q, p+q)_{p,q}) \rightarrow Kh(L_1) \quad (43)$$

up to sign with bidegree  $(0, \chi(\Sigma^\circ))$  where  $\chi(\Sigma^\circ) = \chi(\Sigma) - p - q$ . On the other hand, [Ren23, Corollary 2.2] implies that  $Kh^{0, -(p-q)^2 + 2\max(p,q)}(-T(p+q, p+q)_{p,q}; \mathbb{Z}) \cong \mathbb{Z}$ . Plugging a generator of it into (43) defines a map

$$Kh(\Sigma): Kh(L_0) \rightarrow Kh(L_1)$$

over  $\mathbb{Z}$  up to sign with bidegree  $(0, \chi(\Sigma^\circ) - (p-q)^2 + 2\max(p,q)) = (0, \chi(\Sigma) - [\Sigma]^2 + [|\Sigma|])$  as  $[\Sigma] = \pm(p-q)[\mathbb{CP}^1]$ .

We argue briefly that this direct construction agrees with the previous one over a field. We begin with the lasagna construction. Realize  $(\overline{\mathbb{CP}^2})^\circ$  as  $S^3 \times I \# B^4$  with a 2-handle attached to  $\partial B^4$  along a  $(-1)$ -framed unknot and another 4-handle attached. Then a suitable generalization of the 2-handlebody formula (see [MWW23, Theorem 3.2]) gives

$$\begin{aligned} & \mathcal{S}_0^2((\overline{\mathbb{CP}^2})^\circ; -L_0 \sqcup L_1; \alpha) \\ & \cong \left( \bigoplus_r \mathcal{S}_0^2(S^3 \times I \# B^4; -L_0 \sqcup L_1 \sqcup -T(\alpha^+ + r, \alpha^- + r)) \right) \{-|\alpha| - 2r\} \Big/ \sim \end{aligned} \quad (44)$$

for some equivalence relation  $\sim$ , and the element  $x_\Sigma$  for a skein  $\Sigma$  with  $[\Sigma] = \alpha$  is represented by the class of  $\Sigma^\circ$  as a lasagna filling in the right hand side of (44) ( $\Sigma^\circ$  is a skein without input balls in  $S^3 \times I \# B^4$  rel some  $-L_0 \sqcup L_1 \sqcup -T(p+q, p+q)_{p,q}$  obtained by cutting out a tubular neighborhood of a core  $\mathbb{CP}^1 \subset \overline{\mathbb{CP}^2}$ ). The evaluation of  $x_\Sigma$  by the dual lasagna class  $\theta_{[\Sigma]} \in \mathcal{S}_0^2(\overline{\mathbb{CP}^2})^*$  is realized by regarding  $\Sigma^\circ$  as a skein in  $S^3 \times I$  rel  $-L_0 \sqcup L_1$ , where  $-T(p+q, p+q)_{p,q} \subset \partial B^4$  is regarded as an input link on an input ball (so its orientation is reversed) with decoration given by a generator of  $KhR_2^{0, -2\max(p,q)}(T(p+q, p+q)_{p,q}) \cong \mathbb{F}$ . Tubing this input ball to the negative boundary, then Proposition 2.4 shows the equivalence to the direct construction.

In the case when a cobordism  $\Sigma: L_0 \rightarrow L_1$  in  $\mathbb{CP}^2$  is given by a negative full twist along some  $p+q$  strands of  $L$  ( $p$  of which oriented against the other  $q$ ), the induced map  $Kh(\Sigma): Kh(L_0) \rightarrow Kh(L_1)$  can be described by putting a local torus link  $-T(p+q, p+q)_{p,q}$  near  $L_0$ , decorated by a generator

of  $Kh^{0, -(p-q)^2+2\max(p,q)}(-T(p+q, p+q)_{p,q})$ , and performing  $p+q$  saddles between  $-T(p+q, p+q)_{p,q}$  and  $L_0$  to realize the twist.

We note that the same constructions (both the lasagna approach and the direct approach) work for cobordisms in  $k\overline{\mathbb{C}\mathbb{P}^2}$ . For the lasagna approach, this is because the preferred dual lasagna classes  $\theta_\alpha$  of  $\overline{\mathbb{C}\mathbb{P}^2}$  give rise to preferred dual lasagna classes  $\theta_\alpha^{\otimes k}$  of  $k\overline{\mathbb{C}\mathbb{P}^2}$ . For the direct approach, we simply cut out neighborhoods of every core  $\mathbb{C}\mathbb{P}^1$  in  $k\overline{\mathbb{C}\mathbb{P}^2}$ . We summarize as follows.

**Proposition 6.17.** *For an oriented link cobordism  $\Sigma: L_0 \rightarrow L_1$  in  $k\overline{\mathbb{C}\mathbb{P}^2}$ , there is an induced map*

$$Kh(\Sigma): Kh(L_0) \rightarrow Kh(L_1)$$

*defined over any coefficient field up to sign, with bidegree  $(0, \chi(\Sigma) - [\Sigma]^2 + |[\Sigma]|)$ , which only depends on the isotopy class of  $[\Sigma]$  rel boundary. It agrees with the classical induced map for link cobordisms in  $S^3 \times I$  when  $k = 0$ . Moreover, if  $\Sigma': L_1 \rightarrow L_2$  is another oriented link cobordism in some  $k'\overline{\mathbb{C}\mathbb{P}^2}$ , then  $Kh(\Sigma' \circ \Sigma) = Kh(\Sigma') \circ Kh(\Sigma)$ .  $\square$*

The two perspectives compensate each other. The direct construction has the advantage that the induced map is computable from diagrams; however, a priori it is not clear to be independent of the choices of the core  $\mathbb{C}\mathbb{P}^1$ 's and the paths used to tube the extra boundaries to the negative boundary. The lasagna construction is manifestly well-defined and satisfies the functoriality statement in Proposition 6.17; yet it is not directly computable.

It would be interesting to investigate some properties of this construction. For example, consider a link  $L \subset S^3$  and a collection of disjoint 2-disks  $D = \sqcup_{i=1}^k D_i \subset S^3$  intersecting  $L$  transversely, with  $\partial D$  disjoint from  $L$ , such that the algebraic intersection number between each  $D_i$  and  $L$  is zero. Then one might hope that twisting along the various  $D_i$ 's induces maps whose limit recovers the  $\#^k(S^1 \times S^2)$  homology [Roz10; Wil21] of the link  $L' \subset \#^k(S^1 \times S^2)$  obtained by regarding  $L$  as a link in the 0-surgeries along  $\partial D$ , or whose colimit recovers the dual of the  $\#^k(S^1 \times S^2)$  homology of  $-L'$ , the mirror image of  $L'$ .

## REFERENCES

- [Ago23] I. Agol, *Chainmail links and L-spaces*, arXiv preprint arXiv:2306.10918 (2023).
- [Akb77] S. Akbulut, *On 2-dimensional homology classes of 4-manifolds*, in: *Math. Proc. Cambridge Philos. Soc.* Vol. 82, 1, Cambridge University Press, 1977, 99–106.
- [Akb91a] S. Akbulut, *A fake compact contractible 4-manifold*, *J. Differential Geom.* **33** (1991), no. 2, 335–356.
- [Akb91b] S. Akbulut, *An exotic 4-manifold*, *J. Differential Geom.* **33** (1991), no. 2, 357–361.
- [AM97] S. Akbulut and R. Matveyev, *Exotic structures and adjunction inequality*, *Turkish J. Math.* **21** (1997), no. 1, 47–53.
- [Bar02] D. Bar-Natan, *On Khovanov's categorification of the Jones polynomial*, *Algebr. Geom. Topol.* **2** (2002), no. 1, 337–370.
- [Bla10] C. Blanchet, *An oriented model for Khovanov homology*, *J. Knot Theory Ramifications* **19** (2010), no. 02, 291–312.
- [Boy86] S. Boyer, *Simply-connected 4-manifolds with a given boundary*, *Trans. Amer. Math. Soc.* **298** (1986), no. 1, 331–357.
- [BW08] A. Beliakova and S. Wehrli, *Categorification of the colored Jones polynomial and Rasmussen invariant of links*, *Canad. J. Math.* **60** (2008), no. 6, 1240–1266.
- [CS76] S. E. Cappell and J. L. Shaneson, *Some new four-manifolds*, *Ann. of Math. (2)* **104** (1976), no. 1, 61–72.
- [Don90] S. K. Donaldson, *Polynomial invariants for smooth four-manifolds*, *Topology* **29** (1990), no. 3, 257–315.
- [EH01] J. B. Etnyre and K. Honda, *Knots and Contact Geometry I: Torus Knots and the Figure Eight Knot*, *J. Symplectic Geom.* **1** (2001), no. 1, 63–120.
- [Eli90] Y. Eliashberg, *Topological characterization of Stein manifolds of dimension  $> 2$* , *Internat. J. Math.* **1** (1990), no. 01, 29–46.



- [Fre82] M. H. Freedman, *The topology of four-dimensional manifolds*, J. Differential Geom. **17** (1982), no. 3, 357–453.
- [GL78] C. M. Gordon and R. A. Litherland, *On the signature of a link*, Invent. Math. **47** (1978), 53–69.
- [GLW18] J. E. Grigsby, A. M. Licata, and S. M. Wehrli, *Annular Khovanov homology and knotted Schur–Weyl representations*, Compos. Math. **154** (2018), no. 3, 459–502.
- [Gre13] J. E. Greene, *A spanning tree model for the Heegaard Floer homology of a branched double-cover*, J. Topol. **6** (2013), no. 2, 525–567.
- [Gre17] J. Greene, *Alternating links and definite surfaces*, Duke Math. J. **166** (2017), no. 11.
- [HKT94] I. Hambleton, M. Kreck, and P. Teichner, *Nonorientable 4-manifolds with fundamental group of order 2*, Trans. Amer. Math. Soc. **344** (1994), no. 2, 649–665.
- [HMP21] K. Hayden, T. E. Mark, and L. Piccirillo, *Exotic Mazur manifolds and knot trace invariants*, Adv. Math. **391** (2021), 107994.
- [HS21] K. Hayden and I. Sundberg, *Khovanov homology and exotic surfaces in the 4-ball*, arXiv preprint arXiv:2108.04810 (2021).
- [Kho00] M. Khovanov, *A categorification of the Jones polynomial*, Duke Math. J. **101** (2000), no. 3, 359–426.
- [Kho03] M. Khovanov, *Patterns in knot cohomology, I*, Exp. Math. **12** (2003), no. 3, 365–374.
- [KMT12] R. Kirby, P. Melvin, and P. Teichner, *Cohomotopy sets of 4-manifolds*, Proceedings of the Freedman Fest **18** (2012), 161–190.
- [KR08] M. Khovanov and L. Rozansky, *Matrix factorizations and link homology*, Fund. Math. **199** (2008), no. 1, 1–91.
- [Kre06] M. Kreck, *Some closed 4-manifolds with exotic differentiable structure*, in: *Algebraic Topology Aarhus 1982: Proceedings of a conference held in Aarhus, Denmark, August 1–7, 1982*, Springer, 2006, 246–262.
- [Lee05] E. S. Lee, *An endomorphism of the Khovanov invariant*, Adv. Math. **197** (2005), no. 2, 554–586.
- [LM97] P. Lisca and G. Matić, *Tight contact structures and Seiberg–Witten invariants*, Invent. Math. **129** (1997), no. 3, 509–525.
- [Man+23] C. Manolescu, M. Marengon, S. Sarkar, and M. Willis, *A generalization of Rasmussen’s invariant, with applications to surfaces in some four-manifolds*, Duke Math. J. **172** (2023), no. 2, 231–311.
- [MN22] C. Manolescu and I. Neithalath, *Skein lasagna modules for 2-handlebodies*, J. Reine Angew. Math. **2022** (2022), no. 788, 37–76.
- [MWW22] S. Morrison, K. Walker, and P. Wedrich, *Invariants of 4-manifolds from Khovanov–Rozansky link homology*, Geom. Topol. **26** (2022), 3367–3420.
- [MWW23] C. Manolescu, K. Walker, and P. Wedrich, *Skein lasagna modules and handle decompositions*, Adv. Math. **425** (2023), 109071.
- [MWW24] S. Morrison, K. Walker, and P. Wedrich, *Invariants of surfaces in smooth 4-manifolds from link homology*, arXiv preprint arXiv:2401.06600 (2024).
- [Ng05] L. Ng, *A Legendrian Thurston–Bennequin bound from Khovanov homology*, Algebr. Geom. Topol. **5** (2005), no. 4, 1637–1653.
- [OS04] P. Ozsváth and Z. Szabó, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. of Math. (2) (2004), 1027–1158.
- [OS05] P. Ozsváth and Z. Szabó, *On the Heegaard Floer homology of branched double-covers*, Adv. Math. **194** (2005), no. 1, 1–33.
- [Pic19] L. Piccirillo, *Shake genus and slice genus*, Geom. Topol. **23** (2019), no. 5, 2665–2684.
- [Pic20] L. Piccirillo, *The Conway knot is not slice*, Ann. of Math. (2) **191** (2020), no. 2, 581–591.
- [Pla06] O. Plamenevskaya, *Transverse knots and Khovanov homology*, Math. Res. Lett. **13** (2006), no. 4, 571–586.
- [Ras05] J. Rasmussen, *Khovanov’s invariant for closed surfaces*, arXiv preprint math/0502527 (2005).
- [Ras10] J. Rasmussen, *Khovanov homology and the slice genus*, Invent. Math. **182** (2010), no. 2, 419–447.
- [Ren23] Q. Ren, *Lee filtration structure of torus links*, arXiv preprint arXiv:2305.16089 (2023).
- [Roz10] L. Rozansky, *A categorification of the stable  $SU(2)$  Witten–Reshetikhin–Turaev invariant of links in  $S^2 \times S^1$* , arXiv preprint arXiv:1011.1958 (2010).
- [Sto07] M. Stošić, *Homological thickness and stability of torus knots*, Algebr. Geom. Topol. **7** (2007), no. 1, 261–284.
- [Sto09] M. Stošić, *Khovanov homology of torus links*, Topology Appl. **156** (2009), no. 3, 533–541.
- [SW94] N. Seiberg and E. Witten, *Monopoles, duality and chiral symmetry breaking in  $N = 2$  supersymmetric QCD*, Nucl. Phys. B **431** (1994), no. 3, 484–550.
- [SZ24] I. A. Sullivan and M. Zhang, *Kirby belts, categorified projectors, and the skein lasagna module of  $S^2 \times S^2$* , arXiv preprint arXiv:2402.01081v1 (2024).
- [Tag13] K. Tagami, *The maximal degree of the Khovanov homology of a cable link*, Algebr. Geom. Topol. **13** (2013), no. 5, 2845–2896.

- [Tan99] T. Tanaka, *Maximal Bennequin numbers and Kauffman polynomials of positive links*, Proc. Amer. Math. Soc. **127** (1999), no. 11, 3427–3432.
- [Wil21] M. Willis, *Khovanov homology for links in  $\#r(S^2 \times S^1)$* , Michigan Math. J. **70** (2021), no. 4, 675–748.
- [Yas15] K. Yasui, *Corks, exotic 4-manifolds and knot concordance*, arXiv preprint arXiv:1505.02551 (2015).

### APPENDIX A. PROOF OF LEMMA 5.8

We first rewrite Lemma 5.8 in terms of the classical Khovanov homology. For a framed link  $L$ , let  $Q(L)(t, q)$  denote the Poincaré polynomial of  $Kh(L)$ , and  $\tilde{Q}(L)(t, q) := (tq^3)^{-\frac{1}{2}w(L)}Q(L)$ . Then by (8),  $P(L)(t, q) = q^{-w(L)}Q(-L)(t, q^{-1})$  and  $\tilde{P}(L)(t, q) = \tilde{Q}(-L)(t, q^{-1})$ . Thus Lemma 5.8 is equivalent to the following Khovanov version.

**Lemma A.1 (Comparison lemma).** *For  $n \in \mathbb{Z}_{\geq 0}^m$  and  $p \in \mathbb{Z}^m$  with  $p \geq w$ , we have*

$$\tilde{Q}(K^p(n)) \leq \sum_{\substack{0 \leq n' \leq n \\ 2|n-n'}} R_{n, n', p-w} \tilde{Q}(K^w(n')) \quad (45)$$

for some polynomials  $R_{n, n', p-w} \in \mathbb{Z}[t^{\pm 1/2}, q^{\pm 1/2}]$  independent of  $K$  with

$$R_{n, n', p-w}(t, q) = t^{-\sum_i \frac{(p_i - w_i)n_i^2}{2}} q^{-\sum_i \frac{(p_i - w_i)(n_i^2 + 2n'_i)}{2}} \prod_{j=1}^m \left( \sum_{r_j=0}^{\frac{n_j - n'_j}{2}} \dim \left( \frac{n_j + n'_j}{2} + r_j, \frac{n_j - n'_j}{2} - r_j \right) q^{-2r_j} \right) + \dots$$

where  $\dots$  are terms with lower  $t$ -degrees.

Moreover, if (45) is sharp at the highest  $t$ -degree of the left hand side, then for any  $i = 1, \dots, m$  with  $n_i \geq 2$ , the morphism on  $Kh$  induced by the dotted annular creation map  $K^p(n - 2e_i) \rightarrow K^p(n)$  is injective in the highest nontrivial homological degree of  $Kh(K^p(n))$ .

Lemma A.1 can be reduced to an elementary piece. Suppose  $J \cup L \subset S^3$  is a framed oriented link, where  $J$  is a knot and  $L$  is a link. Let  $J^1$  be  $J$  with framing one higher. For  $n \geq 0$ , let  $J(n) \cup L$  denote the link obtained from  $J \cup L$  by replacing  $J$  by an  $n$ -cable of itself, and  $J^1(n) \cup L$  similarly.

**Lemma A.2 (Elementary comparison lemma).**

$$\tilde{Q}(J^1(n) \cup L) \leq \sum_{\substack{0 \leq n' \leq n \\ 2|n-n'}} R_{n, n'} \tilde{Q}(J(n') \cup L) \quad (46)$$

for some polynomials  $R_{n, n'} \in \mathbb{Z}[t^{\pm 1/2}, q^{\pm 1/2}]$  independent of  $J \cup L$  with

$$R_{n, n'}(t, q) = t^{-\frac{n^2}{2}} q^{-\frac{n^2}{2} - n'} \sum_{r=0}^{\frac{n-n'}{2}} \dim \left( \frac{n+n'}{2} + r, \frac{n-n'}{2} - r \right) q^{-2r} + \dots$$

where  $\dots$  are terms with lower  $t$ -degrees.

Moreover, if (46) is sharp at the highest  $t$ -degree of the left hand side, then the morphism on  $Kh$  induced by the dotted annular creation map  $J^1(n-2) \cup L \rightarrow J^1(n) \cup L$  is injective in the highest nontrivial homological degree of  $Kh(J^1(n) \cup L)$ .

It is easy to check that Lemma A.1 is obtained from iteratively applying Lemma A.2 to components of  $K$  until the framings drop to  $w$ . Therefore we only prove Lemma A.2.

Before starting the proof, we introduce an auxiliary family of links defined by Stösić [Sto07; Sto09] and re-exploited by Tagami [Tag13] (our notation aligns with [Ren23, Section 5], which is different from Stösić's or Tagami's).

For  $n, m \geq 0$ ,  $0 \leq i \leq n-1$ , let  $D_{n,m}^i$  denote the braid closure of  $(\sigma_1 \cdots \sigma_{n-1})^m \sigma_1 \cdots \sigma_i \in B_n$ , regarded as an annular link. Thus  $D_{n,m+1}^0 = D_{n,m}^{n-1}$ , and  $D_{n,n}^0 = T_{n,n}$  is the annular torus link  $T(n, n)$ . For  $i > 0$ , in the standard diagram of  $D_{n,m}^i$  defined by the braid word, resolving the last letter  $\sigma_i$  leads to the 0-resolution which is exactly  $D_{n,m}^{i-1}$ , and some 1-resolution, denoted  $E_{n,m}^{i-1}$ .

**Lemma A.3.** *The annular link  $E_{n,m}^{i-1}$  is isotopic to some  $D_{n_0,m_0}^{i_0}$  or its flip (i.e. the braid closure of  $\sigma_{n_0-i_0} \cdots \sigma_{n_0-1} (\sigma_1 \cdots \sigma_{n_0-1})^{m_0}$ , still denoted  $D_{n_0,m_0}^{i_0}$  by an abuse of notation), possibly disjoint union with a local unknot. Moreover,*

- (1)  $0 \leq n_0 < n$ ,  $2|n - n_0$ .
- (2) If  $m < n$ , then  $m_0 < n_0$  or  $n_0 = 0$ .
- (3)  $E_{n,n-1}^{n-2} \simeq D_{n-2,n-3}^{n-3} \sqcup U$  and  $E_{n,n-1}^i \simeq D_{n-2,n-3}^i$  if  $i < n-2$ .
- (4) If  $m < n$ , then the number of negative crossings in the 1-resolution of  $D_{n,m}^i$  oriented via  $D_{n_0,m_0}^{i_0}(\sqcup U)$ , denoted  $\omega_{n,m}^i$ , is less than or equal to  $\frac{n^2 - n_0^2}{2} - 1$ , with equality if and only if  $m = n - 1$ .

*Sketch proof.* The fact that  $E_{n,m}^{i-1}$  is isotopic to some  $D_{n_0,m_0}^{i_0}$  or  $D_{n_0,m_0}^{i_0} \sqcup U$  is checked carefully in [Tag13, Figure 15–20], from which (1) and (2) are also clear. (3) is exactly a half of [Ren23, Lemma 5.1].

For (4), when  $m = n - 1$  there are exactly  $2n - 3 = \frac{n^2 - (n-2)^2}{2} - 1$  negative crossings as claimed in the paragraph after Lemma 5.1 in [Ren23].

When  $m < n - 1$ , we count the contribution as the turnback arc in Figure 15 to Figure 18 of [Tag13] travels along the longitude. Note that the whole isotopy  $E_{n,m}^{i-1} \simeq D_{n_0,m_0}^{i_0}(\sqcup U)$  is decomposed into  $\frac{n-n_0}{2}$  full longitude travels of the turnback arc and an extra final move according to the four types listed in Figure 15 to Figure 18 of [Tag13].

Interpret  $n$  as the number of through strands in the bottom box and  $m$  as the number of full  $\sigma_1 \cdots \sigma_{n-1}$  rotations. Then each full longitude travel decreases  $n$  by 2, decreases  $m$  by 0 or 2 (other than the last longitude travel of Type 4, which will be discussed below). The negative crossing contribution of this full longitude travel is  $m + n - 2 + \epsilon$  if  $m$  is decreased by 2 and  $m + \epsilon$  if  $m$  is not decreased, where  $\epsilon = 0$  or 1 is the contribution outside the bottom box. The last longitude travel has  $\epsilon = 0$ . The negative crossing contribution from the final move is 0 for type 1, 2, and at most  $n - 1$  for type 3, 4 (for these two types, the last longitude travel decreases  $m$  by either 0 or 1, contributing  $m$  negative crossings). In total, the number of negative crossings is at most

$$\sum_{r=0}^{(n-n_0)/2-1} ((m-2r) + n - 2r - 1) \leq \sum_{r=0}^{(n-n_0)/2-1} (2n - 4r - 3) = \frac{n^2 - n_0^2}{2} - \frac{n - n_0}{2} \leq \frac{n^2 - n_0^2}{2} - 1.$$

If the equality holds, we must have  $m = n - 2$  and  $n_0 = n - 2$ . This implies  $i = n - 1$  or  $n - 2$ . One can check  $E_{n,n-2}^{n-2} \simeq D_{n-2,n-4}^{n-3} \simeq E_{n,n-2}^{n-3}$ , and the 1-resolutions of  $D_{n,n-2}^{n-1}$ ,  $D_{n,n-2}^{n-2}$  have  $2n - 4, 2n - 5$  crossings, respectively, which are strictly less than  $\frac{n^2 - (n-2)^2}{2} - 1$ .  $\square$

*Proof of Lemma A.2.* Let  $D_{n,m}^i(J) \cup L \subset S^3$  be the link obtained by replacing  $J$  by its  $D_{n,m}^i$  satellite. For  $i > 0$ , the resolution of  $D_{n,m}^i$  at the last letter  $\sigma_i$  induces a resolution of  $D_{n,m}^i(J) \cup L$  into

$D_{n,m}^{i-1}(J) \cup L$  and  $E_{n,m}^{i-1}(J) \cup L$ , the latter of which is isotopic to some  $D_{n_0,m_0}^{i_0}(J) \cup L$  possibly disjoint union with an unknot.

Fix a framed link diagram of  $J \cup L$ . Then  $D_{n,m}^i(J) \cup L$  have (almost) standard diagrams by taking the blackboard  $n$ -cable of  $J$  and inserting the standard diagrams of  $D_{n,m}^i$  locally. Then, the number of negative crossings of  $D_{n,m}^i(J) \cup L$  is

$$n^2 \cdot n_-(J) + n \cdot n_-(J, L) + n_-(L)$$

where  $n_{\pm}(J), n_{\pm}(L), n_{\pm}(J, L)$  are the numbers of positive/negative crossings of  $J$ , of  $L$ , and between  $J, L$ , respectively. Since the orientation of  $\frac{n-n_0}{2}$  of the strands are reversed in the 1-resolution of  $D_{n,m}^i(J) \cup L$ , the number of negative crossings in the 1-resolution is

$$\begin{aligned} & \left( \left( \frac{n+n_0}{2} \right)^2 + \left( \frac{n-n_0}{2} \right)^2 \right) n_-(J) + 2 \cdot \frac{n+n_0}{2} \cdot \frac{n-n_0}{2} \cdot n_+(J) + \omega_{n,m}^i \\ & + \frac{n+n_0}{2} n_-(J, L) + \frac{n-n_0}{2} n_+(J, L) + n_-(L), \end{aligned}$$

The difference between the numbers of negative crossings of the 1-resolution of  $D_{n,m}^i(J) \cup L$  and  $D_{n,m}^i(J) \cup L$  itself is

$$\omega_{n,m}^i(J, L) := \frac{n^2 - n_0^2}{2} w(J) + (n - n_0) \ell k(J, L) + \omega_{n,m}^i. \quad (47)$$

The skein exact triangle on Khovanov homology corresponding to the resolution is

$$\begin{array}{ccc} Kh(E_{n,m}^{i-1}(J) \cup L)[\omega_{n,m}^i(J, L) + 1]\{3\omega_{n,m}^i(J, L) + 2\} & \longrightarrow & Kh(D_{n,m}^i(J) \cup L) \\ & \swarrow [1] & \searrow \\ & Kh(D_{n,m}^{i-1}(J) \cup L)\{1\}, & \end{array} \quad (48)$$

which implies that

$$Q(D_{n,m}^i(J) \cup L) \leq qQ(D_{n,m}^{i-1}(J) \cup L) + t^{\omega_{n,m}^i(J, L)+1} q^{3\omega_{n,m}^i(J, L)+2} Q(E_{n,m}^{i-1}(J) \cup L). \quad (49)$$

Start with  $J^1(n) \cup L = D_{n,n-1}^{n-1}(J) \cup L$  and apply (49) or the identity  $D_{n,m+1}^0 = D_{n,m}^{n-1}$  iteratively. By Lemma A.3, each  $E_{n,m}^{i-1}$  is some simpler  $D_{n_0,m_0}^{i_0}$  (possibly with an unknot component which contributes a multiplicative factor of  $q + q^{-1}$  to  $Q$ ), and thus we must end with an inequality

$$Q(J^1(n) \cup L) \leq \sum_{\substack{0 \leq n' \leq n \\ 2|n-n'}} S_{n,n',J,L} Q(J(n') \cup L) \quad (50)$$

for some polynomials  $S_{n,n',J,L}$ . These polynomials are determined as follows. Define a graph with vertices  $\{(n, m, i)\}$  with  $(n, m+1, 0)$  identified with  $(n, m, n-1)$ . For  $i > 0$ , construct a directed edge  $(n, m, i) \rightarrow (n, m, i-1)$  with weight  $q$ , and a directed edge  $(n, m, i) \rightarrow (n_0, m_0, i_0)$  with weight  $t^{\omega_{n,m}^i+1} q^{3\omega_{n,m}^i+2}$  if  $E_{n,m}^{i-1} \simeq D_{n_0,m_0}^{i_0}$  and weight  $t^{\omega_{n,m}^i+1} q^{3\omega_{n,m}^i+2} (q + q^{-1})$  if  $E_{n,m}^{i-1} \simeq D_{n_0,m_0}^{i_0} \sqcup U$ . The weight of a directed path from  $(n, m, i)$  to  $(n', m', i')$  is the product of its edge weights, with an extra multiplicative factor  $(tq^3)^{\frac{n^2-n'^2}{2}w(J)+(n-n')\ell k(J,L)}$  to account for the renormalization factors in (47). Then  $S_{n,n',J,L}$  is the sum of the path weights over directed paths from  $(n, n, 0)$  to  $(n', 0, 0)$ .

By Lemma A.3(4), the  $t$ -degree of every path  $(n, n, 0) \rightarrow (n', 0, 0)$  is at most  $\frac{n^2-n'^2}{2}w(J) + (n - n')\ell k(J, L) + \frac{n^2-n'^2}{2}$ , with equality achieved if and only if every edge on it of the form  $(n, m, i) \rightarrow$

$(n_0, m_0, i_0)$  that decreases  $n$  has  $m = n - 1$ . We can also enumerate all paths that achieve equality thanks to Lemma A.3(3). A few path-counting combinatorics then yield (cf. [Ren23, Figure 4])

$$S_{n,n',J,L}(t,q) = (tq^3)^{\frac{n^2-n'^2}{2}w(J)+(n-n')\ell k(J,K)} t^{\frac{n^2-n'^2}{2}} q^{\frac{3n^2-n'^2}{2}-n'} \sum_{r=0}^{\frac{n-n'}{2}} \dim \left( \frac{n+n'}{2} + r, \frac{n-n'}{2} - r \right) q^{-2r} + \dots$$

where  $\dots$  are terms with lower homological degrees. Since  $w(J^1(n) \cup L) = n^2(w(J)+1) + 2n\ell k(J, L) + w(L)$  and  $w(J(n') \cup L) = n'^2w(J) + 2n'\ell k(J, L) + w(L)$ , (50) renormalizes to yield (46), as desired.

Finally, if (46) is sharp in the sense of the condition in the lemma, then the exact triangle (48) at  $(n, m, i) = (n, n - 1, n - 1)$  splits into a short exact sequence in the highest homological degree. In particular

$$Kh(E_{n,n-1}^{n-2}(J) \cup L) \rightarrow Kh(D_{n,n-1}^{n-1}(J) \cup L) \quad (51)$$

is injective in the highest homological degree of the right hand side. Note  $D_{n,n-1}^{n-1}(J) \cup L = J^1(n) \cup L$  and  $E_{n,n-1}^{n-2}(J) \cup L \simeq J^1(n-2) \cup L \sqcup U$ . By [Ren23, Lemma 5.2] we know that (51) is the induced map by the saddle cobordism  $J^1(n-2) \cup L \sqcup U \rightarrow J^1(n) \cup L$ . Since the dotted annular creation cobordism  $Kh(J^1(n-2)) \rightarrow Kh(J^1(n))$  is obtained by restricting the domain of (51) (to the part where  $Kh(U)$  contributes a tensor summand  $X$ ), it is also injective in the highest homological degree.  $\square$

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, BERKELEY, CA 94720, USA

*Email address:* [qiuyu.ren@berkeley.edu](mailto:qiuyu.ren@berkeley.edu)

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843, USA

*Email address:* [mw188@tamu.edu](mailto:mw188@tamu.edu)