

## Software paper for submission to the Journal of Open Research Software

To complete this template, please replace the blue text with your own. The paper has three main sections: (1) Overview; (2) Availability; (3) Reuse potential.

Please submit the completed paper to: editor.jors@ubiquitypress.com

---

### (1) Overview

#### Title

Yamdb: easily accessible thermophysical properties of liquid metals and molten salts

#### Paper Authors

1. Weier, Tom
2. Nash, William
3. Personnettaz, Paolo
4. Weber, Norbert

#### Paper Author Roles and Affiliations

1. Staff scientist, Institute of Fluid Dynamics, Helmholtz-Zentrum Dresden-Rossendorf, Germany
2. PostDoc, Institute of Fluid Dynamics, Helmholtz-Zentrum Dresden-Rossendorf, Germany
3. PostDoc, Centre National d'Etudes Spatiales, Paris, France; Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, IRD, Univ. Gustave Eiffel, ISTERre, 38000 Grenoble, France; Institute of Fluid Dynamics, Helmholtz-Zentrum Dresden-Rossendorf, Germany
4. Staff scientist, Institute of Fluid Dynamics, Helmholtz-Zentrum Dresden-Rossendorf, Germany

#### Abstract

Yamdb (Yet another materials data base) addresses the need to provide thermophysical properties of liquid metals and molten salts in an easily accessible manner. Mathematical relations describing material properties - usually determined by experiment - are taken from the literature. Equations and their coefficients are stored separately. The former can be implemented in any programming language (Python and Go in this case) and the latter are kept in YAML files together with additional information (source, temperature range, composition, accuracy if available, etc).

#### Keywords

material properties; liquid metals; molten salts; YAML; Python; Go

## Introduction

Knowledge of materials properties is a necessary prerequisite for a great number of scientific and engineering tasks, such as designing apparatuses, estimating heat fluxes, ohmic losses and other quantities and is indispensable for many numerical simulations, e.g., in computational fluid dynamics. Correspondingly many material databases exist, typically built to address the needs of a specific field. We concentrate here on liquid metals and fused salts mainly for their application in electrochemical energy storage, i.e., in liquid metal and molten salt batteries [33, 31]. Yamdb was written to offer rapid and easy access to a number of material properties in the liquid state at standard pressure. Typically, the substances useful in this field possess melting temperatures significantly above room temperature.

The scarcity of data on the thermophysical properties of liquid metals compared to, e.g., hydrocarbons is due to the experimental difficulties in obtaining accurate measurements at high temperatures [25]. The same holds true for fused salts. Nevertheless, there are several carefully edited compilations of thermophysical properties of liquid metals [44, 26, 25] available in text. Janz's collection [29, 27] of molten salt properties from the literature published until about 1988 is a seminal work still held in high esteem [53] even if these data are no longer considered Standard Reference Data (SRD) by the National Institute of Standards and Technology (NIST) [30].

The challenges of building an up-to-date electronic molten salts database containing critically evaluated data are vividly described by Gaune-Escard and Fuller [16, 15]. Assessing data quality and accuracy of reporting is a task that should not be underestimated and that entails considerable resource requirements [7]. Even defining a standardized universal data format for thermophysical properties has proved to be a major undertaking [52].

Our aim with Yamdb is much more modest and focused on molten metals and fused salts: we collect correlations, based on experimental findings, describing temperature dependent thermophysical properties from the literature and make them easily accessible from (interactive) programming environments. Value judgments were not attempted. Collecting clean data from the literature is thereby the main work. Though it is seemingly trivial, errors easily creep in or are already present in the original sources [48]. Using multiple sources for identical properties wherever possible helps to discover errors. Since such errors are unavoidable and often difficult to detect, correct numbers (or coefficients) are of greater value and have considerably longer lifetimes than the concrete implementation of equations. Both should therefore be stored separately. To keep this asset, an easily editable, flexible, well documented data format with a permissible license and widespread adoption is needed. Longevity is vital as well but difficult to assess.

We decided here for YAML [3, 21, 22] (YAML Ain't Markup Language™) because it fulfills most of the aforementioned criteria and is a core component of a sufficient number of popular software packages which gives hope for sustained support. In our opinion, YAML strikes a good balance between the complexity of the database format and flexibility.

As mentioned above, over the years a whole series of databases and tools for using thermophysical data have been developed. For a small subset see Tab. 1. The table collects software and databases from different fields. Driven by demand from

Name	Programming	Ref.	License
Cantera	C++/Fortran/Python	[17]	BSD
DDB		[45]	commercial
MSTDB-TP	C++	[56]	copyrighted/license-free
NIST alloy data	web-based	[62]	public domain
NIST Molten Salts Database		[6]	commercial
PYroMat	Python	[37]	GNU GPL v3.0
pyThermoML	Python	[18]	BSD
REFPROP	Fortran	[35]	commercial
Thermo	Python	[2]	MIT
ThermoData Engine	C++	[12]	commercial
ThermoFun	C++/Python	[40]	LGPL
ThermoML	XML/JSON	[50]	open
TPDS-web	web-based	[41, 64]	copyrighted/free of charge

Table 1: Selected software and databases for thermodynamic properties.

the chemical industry, several well established and comprehensive databases with a focus on hydrocarbons exist, e.g., the Dortmund Data Bank (DDB) [49], the ThermoData Engine (TDE) [12] and REFPROP [24] from NIST. They often contain sophisticated modeling capabilities for thermochemical and transport properties especially for mixtures. Cantera [17], PYroMat [37], Thermo [2], and ThermoFun [40] are examples of open source software belonging to this category. The NIST Molten Salts Database [6] is a DOS based program that offers easy access to the collection of correlation equations for the density, dynamic viscosity, electric conductivity, and surface tension of molten salts collected by Janz and co-workers [27]. Recently, NIST made the underlying coefficients and equations electronically available [30]. The Molten Salt Thermal Properties Database (MSTDB) [56] with a thermochemical (MSTDB-TC) and a thermophysical branch (MSTDB-TP) is being actively developed and focuses on the properties of molten salts suitable for the use in molten salt reactors. ThermoML is primarily a data description in Extensible Markup Language (XML) standardized by the International Union of Pure and Applied Chemistry (IUPAC) [13, 14]. NIST provides an archive of material property values stored in their ThermoML Archive [50] with the possibility to access the data in JavaScript Object Notation (JSON) as well. It includes several sets of measured molten salt property values. PyThermoML [18] wraps ThermoML data sets into Python data structures but does not yet - at the time of writing - implement the specification in its entirety. Material properties of molten metals and alloys are available as sets of experimental data from NIST [62] and in a somewhat smaller quantity from the Network Database System for Thermophysical Property Data (TPDS-web) developed by the Japanese National Institute of Advanced Industrial Science and Technology (AIST) [41, 64]. This database also contains data on molten salts.

So far Yamdb has been used in a number of publications [47, 34, 11] on liquid metal and molten salt batteries as well as on interfacial instabilities [60, 43] in liquid metal

```

❶ Na:
❷ Tm: &Tm_Na 370.87
   Tb: 1156.15
   M: 22.98977E-03
❸ density:
❹ IidaGuthrie1988: &density_Na_IidaGuthrie1988
   rho_m: 927
   lambda: .235
❺ equation: Steinberg1974
❻ Ohse1985: # recommended equation
   a:
     - 0.89660679
     - 0.51613430
   default: true
   equation: Shpilrain1985
❼ dynamic_viscosity:
   Hirai1992:
     A: 0.0846
     B: 6.412
     default: true
     equation: Hirai1992
     comment: "p. 401/63 Tab. 1"
   expansion_coefficient:
     IidaGuthrie1988:
❽ <<: *density_Na_IidaGuthrie1988
     default: true

```

Figure 1: Modified snippet from the `metals.yml` database file showing sodium properties. Please see the text for further explanations.

systems with a miscibility gap.

### Implementation and architecture

As mentioned above, the coefficients for the equations expressing the various properties depending on temperature (and concentration) are stored in YAML files. These files contain a nested structure of block mappings [3] so that the symbol of a substance (element or mixture, ‘Na’ in the example in Fig. 1) constitutes the key of the root node ❶. This root node has a number of mandatory child nodes: melting temperature  $T_m$  ❷, boiling temperature  $T_b$ , and molar mass  $M$ . The temperatures are given in K, the molar mass in kg/mol. Thermophysical properties (density ❸, dynamic\_viscosity ❼, and expansion\_coefficient) are sibling nodes of the aforementioned ones and might or might not be present depending on data availability. The property node with the key `density` ❸ has two child nodes `IidaGuthrie1988` ❹ and `Ohse1985` ❻ the keys of which are named according to the references for the original coefficient data: [26, 44]. The child node with the key `equation` ❺ designates the equation used for the density calculations, in this case

$$\rho = \rho_m - \lambda(T - T_m) \quad (1)$$

derived from [54] and [26]<sup>1</sup>.  $\rho$  is the density in kg/m<sup>3</sup> at the temperature  $T$  in K, while  $\rho_m$  is the density at the melting point  $T_m$  and  $\lambda$  the temperature dependence of the liquid density in kg/(m<sup>3</sup> K).

<sup>1</sup>Note that the sign of  $\lambda$  is inverted with respect to the original definition.

If the density equation to be used is not explicitly specified however, it would be calculated according to the data from `Ohse1985` and the equation `Shpilrain1985` because the `Ohse1985` block [6](#) contains the `default` key with the value `true`. A convenient feature of YAML are the ‘anchor’(&)-‘alias’(\*) pairs that allow direct access to the coefficients of the density equation according to `IidaGuthrie1988` [4](#) by the corresponding equation for the expansion coefficient [8](#) without duplicated input. The `default: true` key-value pair is added to the received coefficients. The presence of several keys in the YAML databases are required: `Tm`, `Tb`, `M` for each substance, the necessary parameters for the respective equations, one and only one `default` per property, `equation` per equation record, and `reference` if the equation record’s key cannot be mapped to an entry in the references database. If available, ranges of temperature (`Tmin`, `Tmax`) or fraction in mol% (`xmin`, `xmax`) for which the correlations are valid are included as well as information about the accuracy (`uncertainty`) and a `comment` for remarks and explanations. Further keys can be added as needed. If necessary, additional methods can be implemented that make use of the content referenced by these keys.

As mentioned above, substances are addressed by their chemical symbols. Ordering of these symbols in compounds is alphabetical. In the case of salts, we follow the convention used by Janz [27] and start the formula with the cationic species, e.g., ‘NaCl’ and not ‘ClNa’ as often found in the newer literature, e.g., [20]. Components of mixtures are divided by a dash and ordered alphabetically by compound. The popular mixture of lithium chloride and potassium chloride would therefore be found under ‘KCl-LiCl’.

Since YAML is a human-friendly text format, the database files can be conveniently edited in a number of editors and integrated development environments (IDEs) with YAML support, e.g., Emacs with `yaml-imenu` and `yafolding`, Geany, PyCharm, and Vim.

Python was chosen for the primary implementation because of its well established role as today’s lingua franca of scientific programming and the truly excellent REPL (Read-Eval-Print Loop) IPython. IPython enables interactive, exploratory work [46] and thus easy and rapid access to the available property methods of the dynamical generated substance objects of Yamdb.

There are of course valid reasons to prefer other languages, e.g., escaping frequent deprecation and breaking changes (even in the core language and libraries) [23, 59] or the desire to use less electricity when performing computations [66].

We chose the Go programming language [10] as a case in point for an alternative implementation of a subset of the Yamdb features in the standalone command-line program Goma (Go materials database). Go has a focus on simplicity and long term maintenance and promises compatibility [8] at the source level for the lifetime of the Go 1 specification. In addition, Go offers, among other things, easy portability by supporting a variety of operating systems and processor architectures via native as well as cross compiling. However, the discussion below will mainly focus on the Python implementation Yamdb.

To minimize maintenance effort and increase the longevity of software, it is crucial to reduce dependencies as much as possible and to select components with proven track records [23]. We have tried to heed this advice and aimed for minimal dependencies




Figure 2: Overall program flow when using Yamdb. Please see the text for further explanations.

besides the essential one on a YAML parser. Other than that, only NumPy and an (optional) B $\text{\LaTeX}$  parser are required third party libraries. The remaining external modules are part of the Python Standard library. There is a long-lasting discussion in the Python community [5] on the right place for a YAML parser with voices [65] calling it “Python’s missing battery”. Time will tell, whether YAML will be found worthy of inclusion one day. NumPy is used in the equations for the simple reason that it allows evaluation of thermophysical properties for a vector of temperature values with one function call. NumPy [58, 19] is a fundamental package for scientific computing in Python and therefore not likely to disappear anytime soon. In case of emergency (the release of NumPy 2.0 is nigh), all calls to NumPy should be replaceable by those to the Python Math Library at the expense of loosing the array computing capabilities. To retain the focus and keep the library a manageable size, other functionality like plotting via Matplotlib is deliberately excluded.

The overall program flow is shown in Fig. 2 and followed in the listing below

```

1  import os
2  from importlib import resources as ir
3  from yamdb import yamdb as ydb
4  metals_yaml = os.path.join(ir.files('yamdb'), 'data', 'metals.yml')
5  sdb = ydb.SubstanceDB(metals_yaml)
6  Na_dict = sdb.get_substance('Na')
7  Na = ydb.Properties(Na_dict)
8  T = Na.Tm
9  rho = Na.density(T)
10 print('Na density at %f K is %f kg/m3' % (T, rho))

```

After loading Yamdb first, a database object has to be generated from a supplied YAML file (line 5). The main component of this object is a dictionary (`sdb.materials`; internal use only, not present in the listing) of dictionaries cor-

responding to the structure of the loaded YAML file. The next step on line 6 is the selection of a sub-dictionary `Na_dict` whose key 'Na' in `sdb.materials` corresponds to the symbol of the substance to be investigated. In this case it is the element sodium. From this sub-dictionary, the property object `Na` is dynamically built by a call to `yamdb.Properties`. This is done by assigning values from the sub-dictionary to the three required constant properties melting and boiling point and molar mass (`Tm`, `Tb`, `M`). Furthermore, for the properties that are implemented in the sub-modules of `yamdb.properties` and that are present in the sub-dictionary `Na_dict`, methods are dynamically generated. They select a default equation per property and allow the user to address the non-default equations by the keyword-argument `source`. If one would like to know the density of sodium according to `IidaGuthrie1988` instead of the default `Ohse1985` (see Fig. 1), one would call `rho = Na.density(T, source='IidaGuthrie1988')` instead of the simpler line 9. The property object `Na` now allows one to conveniently access all available property methods by tab-completion at the command line of IPython and recent Python interpreters with access to the readline library. A number of additional methods exist that allow one to extract different kinds of information from the properties object such as listing all available sources for one property (`get_source_list`) or getting the reference key for a certain source (`get_reference`). A more convenient variant of the procedure described above is offered by the function `get_from_metals` in line 2 of the listing below.

```

1 from yamdb import yamdb as ydb
2 Na = ydb.get_from_metals('Na')
3 print('Na density at %f K is %f kg/m3' % (Na.Tm, Na.density(Na.Tm)))

```

It directly accesses the `metals.yml` database included in the `Yamdb` distribution and directly returns the corresponding `Property` object. The first call to `get_from_metals` parses the `YAML` file, and all subsequent calls for other substances use a cached version of the corresponding dictionary.

Mixtures are handled slightly differently because they are not only defined by their components but also by their mole-fractions. Therefore objects of the `Mixture-Properties` class

```

1 from yamdb import yamdb as ydb
2 CaCl2_NaCl = ydb.get_from_salts('CaCl2-NaCl')
3 T = 1020
4 st = CaCl2_NaCl.composition['20-80'].surface_tension(T)
5 print('CaCl2-NaCl 20-80 surface tension at %d K is %4.3f N/m' %
6       (T, st))
7 rho = CaCl2_NaCl.composition['range'].density(1090, 15)

```

```

8 #  $\rho = 1641 \text{ kg/m}^3$ 
9 rho = CaCl2_NaCl.composition['15-85'].density(1090)
10 #  $\rho = 1635 \text{ kg/m}^3$ 

```

are (transparently) generated by the function `get_from_salts` if a hyphen is detected in the substance name. Properties are now methods of substance and composition as shown in line 4 of the listing above. Compositions in the dictionary keys are quoted in mol%, the first value (20) is for the first component ( $\text{CaCl}_2$ ), the second (80) for the second ( $\text{NaCl}$ ) and so forth. Composition keys are kept as short as possible, trailing zeroes are removed and values are rounded to one digit after the decimal point. If correlation equations for a continuous range of compositions are available, the composition dictionary key is ‘range’ as in line 7. The property function (density in this case) requires then the fraction in mol% as a second argument. As can be seen from the comment lines 8 and 10, density values for 1090 K and  $\text{CaCl}_2$ - $\text{NaCl}$  15-85 from the two different correlation functions are quite close. It is essential to check the original sources before using material properties derived from Yamdb in calculations, and publications should cite the original sources. To facilitate these tasks, Yamdb contains a module `yamref` and a  $\text{BIBTEX}$  database with all references `references.bib`.  $\text{BIBTEX}$  was chosen because it can be considered a standard format in scientific publishing, interacts readily with  $\text{LATEX}$  and has a long history without breaking changes. A number of  $\text{BIBTEX}$  parsers are available as third party Python packages. Yamdb currently uses `BibtexParser` [61] that depends on `PyParsing` [38]. At the time of writing, `BibtexParser` undergoes a major version upgrade that entails breaking changes to the application programming interface (API). In future, `Yamref` might switch to an alternative solution for parsing  $\text{BIBTEX}$  files but without changes to the user interface. For now, a simple backup solution is provided in form of a YAML file mirroring the content of the  $\text{BIBTEX}$  file. Access to the YAML file with references is provided by functions implemented in Yamdb’s `yamdb` module. Identity of the information content of both versions ( $\text{BIBTEX}$  and YAML) is covered by an integration test, see section ‘Quality control’. A simple case of looking up the reference for a density value is demonstrated in the listing below:

```

1 from yamdb import yamdb as ydb
2 from yamdb import yamref as yrf
3
4 Ga = ydb.get_from_metals('Ga')
5
6 rho = Ga.density(273.15 + 50)
7
8 yrf.get_from_references(Ga.get_default_source('density'))
9 # 'Iida, Takamichi and Guthrie, Roderick I. L (1988)
10 # The Physical Properties of Liquid Metals.
11 # Clarendon Press. Oxford.'

```



```

9 # alternatively use (references.yml):
10 ydb.get_from_references((Ga.get_default_source('density')))
11 # 'Iida, T., Guthrie, R.I.L., 1988. The physical properties
12 # of liquid metals. Clarendon Press, Oxford.'
```

`get_default_source('density')` returns the source key of the default source (6 in Fig. 1). It is `IidaGuthrie1988` for Ga that gets expanded to the full citation by the call to `yamref.get_from_references`. If different sources from the same reference are present, the sources' keys, e.g., `Ohse1985Rec` and `Ohse1985Exp`, are derived from but do not directly correspond to the reference. In such a case, an intermediate call to `yamdb.get_reference` is necessary.

In contrast to the module `Yamdb`, `Goma` is a single program whose output is controlled by command-line switches. The source code is divided into a library of material property functions and the main program. It basically evaluates the arguments passed to `Goma` and directs them to the proper functions for property evaluation in a `switch` statement. `Goma` depends on the Go Standard library and `go-yaml v.2` [42]. Thanks to the versatile dependency management tools present in the Go-ecosystem, `Goma` includes the `go-yaml v.2` source via

```
> go mod vendor
```

`Goma` calling conventions are described by a help screen (`goma -h` or `goma -help`, command line flags follow Plan9 standards [4]) and a man page. To get the density of Ca at 1130 K enter

```

> goma -S Ca -T 1130 -density
# property  Temp source          value  unit
#           K
# -----  ---  -----
# density   1130 IidaGuthrie2015  1361.7 kg/m³
# density   1130 Shpilrainetal1980 1351.2 kg/m³
# -----  ---  -----
```

at the command line. The available information is returned as a table. Resolving references is not done internally, only the citation keys are provided in the column 'source'. Should the user have access to  $\LaTeX$  and `pandoc` [36], a shell script `gen_report.bash` supplied with `Goma` can be used to generate an overview of all thermophysical properties available for a substance at a given temperature. A (shortened) example of the script's output for Li at 800 K is shown in Fig. 3.

### Quality control

`Yamdb` uses `pytest` for quality control. The single property equations are fully covered by unit tests. They check the correct implementation of the equations by calculating a property value for a predetermined temperature with a defined set of coefficients.

Completeness of the references and instantiation of objects are examined by integration tests. This includes ensuring that all citation keys of `references.bib` are present in `references.yml`.

## Thermophysical properties of Li

property	Temp K	source	value	unit
density	800	Ohse (1985)	483.21	kg/m <sup>3</sup>
dynamic viscosity	800	Hirai (1992)	0.00031787	Pa · s
dynamic viscosity	800	Ohse (1985)	0.00031594	Pa · s
expansion coefficient	800	Takamichi Iida and Guthrie (1988)	0.0002069	1/K
heat capacity	800	Ohse (1985)	4199.8	J/(kg · K)
resistivity	800	Zinkle (1998)	3.9302e-07	Ω · m
sound velocity	800	Blairs (2007)	4326.9	m/s
surface tension	800	Keene (1993)	0.34787	N/m
thermal conductivity	800	Takachimi Iida and Guthrie (2015)	56.301	W/(m · K)

## References

Blairs, S. 2007. “Review of Data for Velocity of Sound in Pure Liquid Metals and Metalloids.” *International Materials Reviews* 52 (6): 321–44.

Figure 3: Script-generated report on the thermophysical properties of Li at 800 K using Goma (shortened). Formatting and bibliography require pandoc and L<sup>A</sup>T<sub>E</sub>X.




Figure 4: Sodium density according to correlation equations from Yamdb (solid lines) compared to experimental data provided via the NIST Alloy data web application [62, 48, 63]. The original data are from Davies (1972) [9], Khairulin et al. (2013) [32], Ruppertsberg and Jost (1989) [51], and Tavo et al. (2009) [55]

Plausibility of the temperature dependency of the properties is tested by comparing (if available) different equations for the same property in one plot. Fig. 4 shows an example for the density values of sodium in the temperature range between melting and boiling point. As can be seen, all correlation equations available from Yamdb collapse fairly well and are in the range of the error margins of the most recent




Figure 5: Surface tension of  $\text{CaCl}_2\text{-NaCl}$  mixtures versus composition for different temperatures adapted from [1]. Correlation expressions (dotted and dashed lines) from Yamdb compared to measured data by Addison, Coldrey [1] (dots and solid lines). Dotted lines result from evaluating the equations by [28] and dashed lines from those of [27]. As expected, the latter are identical to the values by [30].

measurements by Khairulin et al. [32]. Measured values included in Fig. 4 were taken from the NIST alloy data [62]. The `metals.yml` database can be checked by a script `check_metals.py` delivered with Yamdb that produces a large number of plots comparing the temperature dependent thermophysical properties for each metal between different sources.

An example for molten salt mixture properties is given in Fig. 5 adapted from [1]. It shows the surface tension of liquid  $\text{CaCl}_2\text{-NaCl}$  mixtures for different temperatures over the whole range of compositions in the liquid state. In the white areas designated ‘solid’, the mixture is not a liquid. Measured values provided by Addison and Coldrey [1] (dots and solid lines) are compared to equations reported by Janz et al. [28] and Janz [27]. Temperature is encoded by color. It is apparent that the earlier reported equations by Janz et al. [28] (dotted lines) miss the measured values by a difference in surface tension  $\gamma$  corresponding to a temperature difference of about 50 K. The newer equations from Janz [27] (dashed lines) fit much better, especially for the lower and higher temperatures. The values computed with the coefficients from the database by Janz [30] are identical to the ones from Janz [27] (dashed lines).

It should be emphasized that measurements of the thermophysical properties of reactive liquid media at elevated temperatures are far from trivial and very demanding [39]. Certain differences in the reported and calculated values are therefore expected and simply reflect the experimental challenges.

The YAML databases are checked for correct syntax by using yamllint. For the time being, Goma's results for predefined sets of temperature values are simply compared to Yamdb's output.

## (2) Availability

### Operating system

#### Yamdb

An operating system capable of running a supported version of Python is required. Yamdb is tested and developed on ArchLinux (rolling release, 2023-11-03), and has been shown to run on OpenBSD 7.3 and 7.4 (requires installation of python3.11 from the ports tree), Ubuntu 22.04 with python3.11, postmarketOS/Alpine Linux/SXMO, Armbian 23.5, Android 11 via termux, as well as Microsoft Windows 10 with python3.12.

#### Goma

A Go compiler for the operating system and processor architecture is a necessary precondition for compiling Go programs on a system. Alternatively, cross-compilation can provide binaries for non-native architectures and operating systems. Goma is tested and developed on ArchLinux (rolling release, 2023-11-03), and has been shown to run on OpenBSD 7.3 and 7.4, Ubuntu 22.04 with python3.11, postmarketOS/Alpine Linux/SXMO, Armbian 23.5, Android 11 via termux, as well as Microsoft Windows 10.

### Programming language

#### Yamdb

Python 3.11.3

#### Goma

Go 1.20.6

### Additional system requirements

No specific requirements.

### Dependencies

#### Yamdb

- BibTexParser version 1.4.0, <https://github.com/sciunto-org/python-bibtexparser>, BSD license
- NumPy version 1.25.1, <https://numpy.org/>, NumPy license
- PyYAML version 6, <https://github.com/yaml/pyyaml>, MIT license

#### Goma

go-yaml version 2.4.0, <https://github.com/go-yaml/yaml/tree/v2>, Apache License Version 2.0

### List of contributors

Tanja Klopper (Helmholtz-Zentrum Dresden Rossendorf) - Development and testing

William Nash (Helmholtz-Zentrum Dresden Rossendorf) - Microsoft Windows port, testing, and language editing

Hirav Patel (Helmholtz-Zentrum Dresden Rossendorf) - Development and testing

Paolo Personnettaz (Helmholtz-Zentrum Dresden Rossendorf) - Development and testing

Norbert Weber (Helmholtz-Zentrum Dresden Rossendorf) - Material property collection and evaluation

Tom Weier (Helmholtz-Zentrum Dresden Rossendorf) - Design, development and testing, data collection

### Software location:

#### Archive Yamdb

**Name:** RODARE

**Persistent identifier:** DOI: 10.14278/rodare.2549

**Licence:** MIT

**Publisher:** Helmholtz-Zentrum Dresden-Rossendorf

**Version published:** 0.3.0

**Date published:** 05/11/23

#### Archive Goma

**Name:** RODARE

**Persistent identifier:** DOI: 10.14278/rodare.2547

**Licence:** MIT

**Publisher:** Helmholtz-Zentrum Dresden-Rossendorf

**Version published:** 0.1.0

**Date published:** 05/11/23

#### Code repository Yamdb

**Name:** Codebase.Helmholtz

**Persistent identifier:** <https://codebase.helmholtz.cloud/prosa/yamdb>

**Licence:** MIT

**Date published:** 07/11/23

#### Code repository Goma

**Name:** Codebase.Helmholtz

**Persistent identifier:** <https://codebase.helmholtz.cloud/prosa/goma>

**Licence:** MIT

**Date published:** 07/11/23

### Language

English.

### (3) Reuse potential

Yamdb may be used by anybody interested in the thermophysical properties of liquid metals and molten salts. Such data are needed for simulations in application areas like high temperature materials processing, electrochemical engineering, and energy storage. Ease of reuse and extensibility are central goals of Yamdb. It can be incorporated as a library into simulation codes that feature a Python interface. As demonstrated by Goma, the effort to transfer the correlation equations into another programming language is relatively low. Accessing the coefficients from the database files should be trivial in most cases since several programming languages as Crystal and Ruby ship YAML parsers as part of their standard libraries. For most other languages well established third party parsers - often based on the C library libyaml [57] - are available. Direct access to temperature dependent thermophysical properties could thereby be enabled for simulations in the language best suited for the purpose.

Yamdb can be easily extended in a number of ways. In the simplest case just by adding new correlation coefficients for existing substances and already implemented equations. This requires only additions to the YAML databases. New equations are easy to implement as well if existing properties are to be extended. This requires only adding a function with the standard interface

```

1 def Shpilrain1985(Temp, coef=coef):
2     r"""Return density according to Dhse (1985) p.453/454.
3     ... (docstring shortened)
4     """
5     a = coef['a']
6     rho = 0.0
7     tau = Temp/1000.0
8     for i, c in enumerate(a):
9         rho += c * np.power(tau, i)
10    rho *= 1000.0 # apparently needed to convert to kg/m³
11    return rho

```

to the package under `yamdb/properties` containing the functions for the desired property, e.g., `density.py`. To make copying of coefficients from the literature as fail-safe and easy as possible, implementing the equation referred to in a paper is preferred to complex coefficient conversions if nothing similar is already available from the property modules.

If a new property is to be added, a corresponding module named after the property (e.g., `dielectric_constant.py` for the dielectric constants) needs to be added to `yamdb/properties` and the `module_dict` in `yamdb/yamdb.py` has to be extended by an entry for the new module (`'dielectric_constant': None`). In addition, the class `Properties` of `yamdb/yamdb.py` requires a new private wrapper method for the extra property modelled after:

```

1 def _density(self, Temp, *args, source=None):
2     if source is None:
3         source = self.get_default_source('density')

```

```
4     return self._density_func_dict[source](Temp, *args)
```

This can basically be done by replacing `density` with `dielectric_constant` in the listing above.

Adding new coefficients, functions, materials, and properties to Yamdb is explained in detail in Yamdb's documentation.

Support for Yamdb and Goma is available by email to the authors of this paper and by opening issues on Yamdb's or Goma's repository pages, see paragraph 'Software location/code repository'. Documentation and examples are included in the software repositories. Contributing changes back to the main repositories as pull requests or patches is encouraged and highly welcome.

### Acknowledgements

Stimulating discussions with Michael Nimtz and Moritz Streb on the needs for and the implementation of thermophysical materials databases especially for liquid metals and molten salts were instrumental and are highly appreciated. Paolo Peronnetaz worked on the development and testing of this project mainly during its stay at HZDR until February 2022.

### Funding statement

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 963599.

### Competing interests

The authors declare that they have no competing interests.

### References

- [1] C. C. Addison and J. M. Coldrey. Influence of surface reactions on the interface between liquid sodium and molten sodium chloride + calcium chloride mixtures. *Transactions of the Faraday Society*, 56:840, 1960. DOI: [10.1039/tf9605600840](https://doi.org/10.1039/tf9605600840).
- [2] Caleb Bell and Contributors. Thermo: Chemical properties component of chemical engineering design library (ChEDL), 2016-2023. URL <https://github.com/CalebBell/thermo>.
- [3] Oren Ben-Kiki, Clark Evans, and Ingy döt Net. YAML ain't markup language (YAML™) version 1.2, 2021. URL <https://yaml.org/spec/1.2.2/>.
- [4] Matt Butcher and Matt Farina. *Go in Practice*. Manning, Shelter Island, NY, 2016. URL <https://www.manning.com/books/go-in-practice>.
- [5] Brett Cannon, Nathaniel Smith, and Donald Stufft. PEP 518 - Specifying minimum build system requirements for Python projects, 2016. URL <https://peps.python.org/pep-0518/>.

- [6] Malcolm W. Chase and Joan C. Sauerwein. NIST Standard Reference Data Products Catalog 1993. NIST Special Publication NIST SP 782-1993, National Institute of Standards and Technology, Gaithersburg, MD, 1993. DOI: [10.6028/NIST.SP.782-1993](https://doi.org/10.6028/NIST.SP.782-1993).
- [7] Robert D. Chirico, Michael Frenkel, Joseph W. Magee, Vladimir Diky, Chris D. Muzny, Andrei F. Kazakov, Kenneth Kroenlein, Ilmutdin Abdulgatov, Gary R. Hardin, William E. Acree, Joan F. Brenneke, Paul L. Brown, Peter T. Cummings, Theo W. de Loos, Daniel G. Friend, Anthony R. H. Goodwin, Lee D. Hansen, William M. Haynes, Nobuyoshi Koga, Andreas Mandelis, Kenneth N. Marsh, Paul M. Mathias, Clare McCabe, John P. O’Connell, Agilio Pádua, Vicente Rives, Christoph Schick, J. P. Martin Trusler, Sergey Vyazovkin, Ron D. Weir, and Jiangtao Wu. Improvement of quality in publication of experimental thermophysical property data: Challenges, assessment tools, global implementation, and online support. *Journal of Chemical & Engineering Data*, 58(10):2699–2716, 2013. DOI: [10.1021/je400569s](https://doi.org/10.1021/je400569s).
- [8] Russ Cox, Robert Griesemer, Rob Pike, Ian Lance Taylor, and Ken Thompson. The Go programming language and environment. *Communications of the ACM*, 65(5):70–78, 2022. DOI: [10.1145/3488716](https://doi.org/10.1145/3488716).
- [9] H. A. Davies. The density and surface tension of dilute liquid Na-In alloys and comparison with liquid Na-Cd alloys. *Metallurgical Transactions*, 3(11):2917–2921, 1972. DOI: [10.1007/bf02652861](https://doi.org/10.1007/bf02652861).
- [10] Alan A. A. Donovan and Brian W. Kernighan. *The Go Programming Language*. Addison-Wesley, New York, 2015. URL <http://www.gopl.io/>.
- [11] C. Duczek, K. E. Einarsrud, O. E. Godinez-Brizuela, P. Trtik, A. Y. Gelfgat, O. S. Kjos, C. Sommerseth, P. Personnettaz, W. Ding, G. M. Horstmann, S. Landgraf, G. Monrrabal-Marquez, W. Nash, M. Sarma, N. Weber, and T. Weier. Fluid mechanics of Na||Zn liquid metal batteries. *Applied Physics Reviews*, 2023. in preparation.
- [12] Michael Frenkel, Robert D. Chirico, Vladimir Diky, Xinjian Yan, Qian Dong, and Chris Muzny. Thermodata Engine (TDE): Software implementation of the dynamic data evaluation concept. *Journal of Chemical Information and Modeling*, 45(4):816–838, 2005. DOI: [10.1021/ci050067b](https://doi.org/10.1021/ci050067b).
- [13] Michael Frenkel, Robert D. Chirico, Vladimir Diky, Qian Dong, Kenneth N. Marsh, John H. Dymond, William A. Wakeham, Stephen E. Stein, Erich Königsberger, and Anthony R. H. Goodwin. XML-based IUPAC standard for experimental, predicted, and critically evaluated thermodynamic property data storage and capture (ThermoML) (IUPAC recommendations 2006). *Pure and Applied Chemistry*, 78(3):541–612, 2006. DOI: [10.1351/pac200678030541](https://doi.org/10.1351/pac200678030541).
- [14] Michael Frenkel, Vladimir Diky, Robert D. Chirico, Robert N. Goldberg, Heiko Heerklotz, John E. Ladbury, David P. Remeta, John H. Dymond, Anthony R. H. Goodwin, Kenneth N. Marsh, William A. Wakeham, Stephen E. Stein,



- Paul L. Brown, Erich Königsberger, and Peter A. Williams. ThermoML: an XML-based approach for storage and exchange of experimental and critically evaluated thermophysical and thermochemical property data. 5. Speciation and complex equilibria. *Journal of Chemical & Engineering Data*, 56(2):307–316, 2011. DOI: [10.1021/je100999j](https://doi.org/10.1021/je100999j).
- [15] Joan Fuller and Marcelle Gaune-Escard. *The Challenges of Building a Molten Salt Database*, chapter 16, pages 275–294. Green Industrial Applications of Ionic Liquids. Springer Netherlands, 2003. DOI: [10.1007/978-94-010-0127-4\\_16](https://doi.org/10.1007/978-94-010-0127-4_16).
- [16] Marcelle Gaune-Escard and Joan Fuller. Virtual molten salt laboratory: Dream or reality? In S. Seetharaman and Du Sichen, editors, *Proc. 6th Int. Conf. Molten Slags, Fluxes and Salts*, Stockholm, Helsinki, 2000. DOI: [10.1515/HTMP.2001.20.3-4.309](https://doi.org/10.1515/HTMP.2001.20.3-4.309).
- [17] David G Goodwin, Harry K Moffat, Ingmar Schoegl, Raymond L Speth, and Bryan W Weber. Cantera: An object-oriented software toolkit for chemical kinetics, thermodynamics, and transport processes, 2022. DOI: [10.5281/zenodo.6387882](https://doi.org/10.5281/zenodo.6387882).
- [18] Matthias Gültig, Jan P. Range, Benjamin Schmitz, and Jürgen Pleiss. Integration of simulated and experimentally determined thermophysical properties of aqueous mixtures by ThermoML. *Journal of Chemical & Engineering Data*, 67(11):3340–3350, 2022. DOI: [10.1021/acs.jced.2c00391](https://doi.org/10.1021/acs.jced.2c00391).
- [19] Charles R. Harris, K. Jarrod Millman, Stéfan J. van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, Julian Taylor, Sebastian Berg, Nathaniel J. Smith, Robert Kern, Matti Picus, Stephan Hoyer, Marten H. van Kerkwijk, Matthew Brett, Allan Haldane, Jaime Fernández del Río, Mark Wiebe, Pearu Peterson, Pierre Gérard-Marchant, Kevin Sheppard, Tyler Reddy, Warren Weckesser, Hameer Abbasi, Christoph Gohlke, and Travis E. Oliphant. Array programming with NumPy. *Nature*, 585(7825): 357–362, 2020. DOI: [10.1038/s41586-020-2649-2](https://doi.org/10.1038/s41586-020-2649-2).
- [20] W. M. Haynes, David R. Lide, and Thomas J. Bruno, editors. *CRC Handbook of Chemistry and Physics*. CRC Press, Boca Raton, 97 edition, 2016.
- [21] Konrad Hinsén. Caring for your data. *Computing in Science & Engineering*, 14(6):70–74, 2012. DOI: [10.1109/MCSE.2012.108](https://doi.org/10.1109/MCSE.2012.108).
- [22] Konrad Hinsén. Domain-specific languages in scientific computing. *Computing in Science & Engineering*, 20(1):88–92, 2018. DOI: [10.1109/MCSE.2018.011111130](https://doi.org/10.1109/MCSE.2018.011111130).
- [23] Konrad Hinsén. Dealing with software collapse. *Computing in Science & Engineering*, 21(3):104–108, 2019. DOI: [10.1109/MCSE.2019.2900945](https://doi.org/10.1109/MCSE.2019.2900945).
- [24] Marcia L. Huber, Eric W. Lemmon, Ian H. Bell, and Mark O. McLinden. The NIST REFPROP database for highly accurate properties of industrially

- important fluids. *Industrial & Engineering Chemistry Research*, 61(42):15449–15472, 2022. DOI: [10.1021/acs.iecr.2c01427](https://doi.org/10.1021/acs.iecr.2c01427).
- [25] Takachimi Iida and Roderick I. L. Guthrie. *The Thermophysical Properties of Metallic Liquids*. Oxford University Press, Oxford, 2015.
- [26] Takamichi Iida and Roderick I. L. Guthrie. *The Physical Properties of Liquid Metals*. Clarendon Press, Oxford, 1988.
- [27] G. J. Janz. Thermodynamic and transport properties for molten salts: Correlation equations for critically evaluated density, surface tension, electrical conductance, and viscosity data. *Journal of Physical and Chemical Reference Data*, 17(Supplement No. 2), 1988. URL <https://srdata.nist.gov/JPCRD/jpcrdS2Vol17.pdf>.
- [28] G. J. Janz, R. P. T. Tomkins, C. B. Allen, J. R. Downey, G. L. Garner, U. Krebs, and S. K. Singer. Molten salts: Volume 4, part 2, chlorides and mixtures-electrical conductance, density, viscosity, and surface tension data. *Journal of Physical and Chemical Reference Data*, 4(4):871–1178, 1975. DOI: [10.1063/1.555527](https://doi.org/10.1063/1.555527).
- [29] George J. Janz. *Molten Salts Handbook*. Academic Press, New York, 1967.
- [30] George J. Janz. Data from: NIST properties of molten salts database (formerly SRD 27), 1992. DOI: [10.18434/MDS2-2298](https://doi.org/10.18434/MDS2-2298).
- [31] Douglas H. Kelley and Tom Weier. Fluid mechanics of liquid metal batteries. *Applied Mechanics Reviews*, 70(2):020801, 2018. DOI: [10.1115/1.4038699](https://doi.org/10.1115/1.4038699).
- [32] Rashid A. Khairulin, Sergei V. Stankus, and Rasul N. Abdullaev. Density, thermal expansion and binary diffusion coefficients of sodium-lead melts. *High Temperatures - High Pressures*, 42:493–507, 2013.
- [33] H. Kim, D.A. Boysen, J.M. Newhouse, B.L. Spatocco, B. Chung, P.J. Burke, D.J. Bradwell, K. Jiang, A.A. Tomaszowska, K. Wang, W. Wei, L.A. Ortiz, S.A. Barriga, S.M. Poizeau, and D.R. Sadoway. Liquid metal batteries: Past, present, and future. *Chem. Rev.*, 113:2075–2099, 2013. DOI: [10.1021/cr300205k](https://doi.org/10.1021/cr300205k).
- [34] Juhan Lee, Gleidys Monrrabal-Marquez, Martins Sarma, Tobias Lappan, Yvonne Jasmin Hofstetter, Pavel Trtik, Steffen Landgraf, Wenjin Ding, Sumit Kumar, Yana Vaynzof, Norbert Weber, and Tom Weier. Membrane-free alkali metal-iodide battery with a molten salt. *Energy Technology*, page 2300051, 2023. DOI: [10.1002/ente.202300051](https://doi.org/10.1002/ente.202300051).
- [35] E. W. Lemmon, I. H. Bell, M. L. Huber, and M. O. McLinden. NIST Standard Reference Database 23: Reference Fluid Thermodynamic and Transport Properties-REFPROP, version 10.0, National Institute of Standards and Technology, 2018. DOI: [10.18434/T4/1502528](https://doi.org/10.18434/T4/1502528). URL <https://www.nist.gov/srd/refprop>.

- [36] John MacFarlane. Pandoc - a universal document converter, 2023. URL <https://pandoc.org/>.
- [37] Christopher Martin, Joseph Ranalli, and Jacob Moore. PYroMat: A Python package for thermodynamic properties. *Journal of Open Source Software*, 7(79):4757, 2022. DOI: [10.21105/joss.04757](https://doi.org/10.21105/joss.04757).
- [38] Paul McGuire. PyParsing, 2023. URL <https://github.com/pyarsing/pyarsing/>.
- [39] K. C. Mills and P. D. Lee. Plenary lecture: Thermophysical property requirements for modelling of high temperature processes. In *Proceedings of the First International Symposium on Microgravity Research and Applications in Physical Sciences and Biotechnology*, page 555, Sorrento, Italy, 2001. European Space Agency, ESA. URL <https://ui.adsabs.harvard.edu/abs/2001ESASP.454.555M/abstract>.
- [40] George Dan Miron, Allan M. M. Leal, S. V. Dmytrieva, and Dmitrii A. Kulik. Thermofun: A C++/Python library for computing standard thermodynamic properties of substances and reactions across wide ranges of temperatures and pressures. *Journal of Open Source Software*, 8(83):4624, 2023. DOI: [10.21105/joss.04624](https://doi.org/10.21105/joss.04624).
- [41] National Institute of Advanced Industrial Science and Technology. Network database system for thermophysical property data, 2006. URL <https://tpds.db.aist.go.jp/tpds-web/>.
- [42] Gustavo Niemeyer. YAML support for the Go language, 2020. URL <https://github.com/go-yaml/yaml/tree/v2>.
- [43] C. Nore, L. Cappanera, J.-L. Guermond, T. Weier, and W. Herreman. Feasibility of metal pad roll instability experiments at room temperature. *Physical Review Letters*, 126(18):184501, 2021. DOI: [10.1103/PhysRevLett.126.184501](https://doi.org/10.1103/PhysRevLett.126.184501).
- [44] Roland W. Ohse, editor. *Handbook of Thermodynamic and Transport Properties of Alkali Metals*. Number 30 in International Union of Pure and Applied Chemistry Chemical Data Series. Blackwell Scientific Publications, Oxford, London, Edinburgh, Boston, Palo Alto, Melbourne, 1985.
- [45] U. Onken, J. Rarey-Nies, and J. Gmehling. The Dortmund Data Bank: a computerized system for retrieval, correlation, and prediction of thermodynamic properties of mixtures. *International Journal of Thermophysics*, 10(3):739–747, 1989. DOI: [10.1007/BF00507993](https://doi.org/10.1007/BF00507993).
- [46] Fernando Perez, Brian E. Granger, and John D. Hunter. Python: an ecosystem for scientific computing. *Computing in Science & Engineering*, 13(2):13–21, 2011. DOI: [10.1109/MCSE.2010.119](https://doi.org/10.1109/MCSE.2010.119).

- [47] Paolo Personnettaz, Tanja Sophia Klopper, Norbert Weber, and Tom Weier. Layer coupling between solutal and thermal convection in liquid metal batteries. *International Journal of Heat and Mass Transfer*, 188:122555, 2022. DOI: [10.1016/j.ijheatmasstransfer.2022.122555](https://doi.org/10.1016/j.ijheatmasstransfer.2022.122555).
- [48] E. A. Pfeif and K. Kroenlein. Perspective: Data infrastructure for high throughput materials discovery. *APL Materials*, 4(5):053203, 2016. DOI: [10.1063/1.4942634](https://doi.org/10.1063/1.4942634).
- [49] Jürgen Rarey and Jürgen Gmehling. Factual data banks and their application to the synthesis and design of chemical processes and the development and testing of thermophysical property estimation methods. *Pure and Applied Chemistry*, 81(10):1745–1768, 2009. DOI: [10.1351/PAC-CON-08-11-17](https://doi.org/10.1351/PAC-CON-08-11-17).
- [50] Demian Riccardi, Zachary Trautt, Ala Bazyleva, Eugene Paulechka, Vladimir Diky, Joseph W. Magee, Andrei F. Kazakov, Scott A. Townsend, and Chris D. Muzny. Towards improved FAIRness of the ThermoML archive. *Journal of Computational Chemistry*, 43(12):879–887, 2022. DOI: [10.1002/jcc.26842](https://doi.org/10.1002/jcc.26842).
- [51] H. Ruppertsberg and J. Jost. Determination of the heat capacity of liquid alloys according to the  $(\partial p/\partial t)_s$  procedure: Pb/Na. *Thermochimica Acta*, 151:187–195, 1989. DOI: [10.1016/0040-6031\(89\)85348-1](https://doi.org/10.1016/0040-6031(89)85348-1).
- [52] H. G. Semerjian and D. R. Burgess. Data programs at NBS/NIST: 1901-2021. *Journal of Physical and Chemical Reference Data*, 51(1):011501, 2022. DOI: [10.1063/5.0084230](https://doi.org/10.1063/5.0084230).
- [53] R. Serrano-López, J. Fradera, and S. Cuesta-López. Molten salts database for energy applications. *Chemical Engineering and Processing: Process Intensification*, 73:87–102, 2013. DOI: [10.1016/j.cep.2013.07.008](https://doi.org/10.1016/j.cep.2013.07.008).
- [54] D. J. Steinberg. A simple relationship between the temperature dependence of the density of liquid metals and their boiling temperatures. *Metallurgical Transactions*, 5(6):1341–1343, 1974. DOI: [10.1007/BF02646618](https://doi.org/10.1007/BF02646618).
- [55] T. M. Taova, F. M. Mal’surgenova, B. B. Alchagirov, and Kh. L. Khokonov. The density and molar volumes of ternary alloys of cross sections of the sodium-potassium-cesium system technically important temperatures. *High Temperature*, 47(6):815–821, 2009. DOI: [10.1134/s0018151x09060066](https://doi.org/10.1134/s0018151x09060066).
- [56] Nick Termini, Tony Birri, Shane Henderson, and N Dianne Ezell. An overview of the Molten Salt Thermal Properties Database-Thermophysical, Version 2.1.1 (MSTDB-TP v.2.1.1). Technical Report ORNL/TM-2023/2955, Oak Ridge National Laboratory, July 2023. DOI: [10.2172/1988348](https://doi.org/10.2172/1988348).
- [57] The YAML Project. libyaml, 2021. URL <https://github.com/yaml/libyaml>.
- [58] Stéfan van der Walt, S Chris Colbert, and Gaël Varoquaux. The NumPy array: A structure for efficient numerical computation. *Computing in Science & Engineering*, 13(2):22–30, 2011. DOI: [10.1109/MCSE.2011.37](https://doi.org/10.1109/MCSE.2011.37).

- [59] Boris Veytsman. From the president. *TUGboat*, 42(3):223–223, 2021. DOI: [10.47397/tb/42-3/tb132pres](https://doi.org/10.47397/tb/42-3/tb132pres).
- [60] Tom Weier, Ilmārs Grants, Gerrit Maik Horstmann, Steffen Landgraf, Michael Nimtz, Paolo Personnettaz, Frank Stefani, and Norbert Weber. Conductivity influence on interfacial waves in liquid metal batteries and related two-layer systems. *Magnetohydrodynamics*, 56(2-3):237–245, 2020. URL <http://mhd.sal.lv/contents/2020/2/MG.56.2.15.R.html>.
- [61] Michael Weiss. BibtexParser, 2023. URL <https://github.com/sciunto-org/python-bibtexparser>.
- [62] Boris Wilthan. NIST alloy data, National Institute of Standards and Technology, 2019. DOI: [10.18434/M32153](https://doi.org/10.18434/M32153). URL <https://data.nist.gov/od/id/mds2-2153>.
- [63] Boris Wilthan, Erik A. Pfeif, Vladimir V. Diky, Robert D. Chirico, Ursula R. Kattner, and Kenneth Kroenlein. Data resources for thermophysical properties of metals and alloys, Part 1: Structured data capture from the archival literature. *Calphad*, 56:126–138, 2017. DOI: [10.1016/j.calphad.2016.12.004](https://doi.org/10.1016/j.calphad.2016.12.004).
- [64] Kouji Yasuda, Takahiro Ohkubo, Osamu Takeda, Shungo Natsui, and Hidehiro Sekimoto. Activity report on information-gathering of database literatures for molten salts. *Electrochemistry*, 88(4):243–252, 2020. DOI: [10.5796/electrochemistry.20-00058](https://doi.org/10.5796/electrochemistry.20-00058).
- [65] Bartosz Zaczyński. YAML: The missing battery in Python, 2022. URL <https://realpython.com/python-yaml/>.
- [66] Simon Portegies Zwart. The ecological impact of high-performance computing in astrophysics. *Nature Astronomy*, 4(9):819–822, 2020. DOI: [10.1038/s41550-020-1208-y](https://doi.org/10.1038/s41550-020-1208-y).

---

### Copyright Notice

Authors who publish with this journal agree to the following terms:

Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a Creative Commons Attribution License that allows others to share the work with an acknowledgement of the work’s authorship and initial publication in this journal.

Authors are able to enter into separate, additional contractual arrangements for the non-exclusive distribution of the journal’s published version of the work (e.g., post it to an institutional repository or publish it in a book), with an acknowledgement

of its initial publication in this journal.

By submitting this paper you agree to the terms of this Copyright Notice, which will apply to this submission if and when it is published by this journal.