

GENERAL INFINITESIMAL VARIATIONS OF HODGE STRUCTURE OF AMPLE CURVES IN SURFACES

VÍCTOR GONZÁLEZ-ALONSO AND SARA TORELLI

ABSTRACT. Given a smooth projective complex curve inside a smooth projective surface, one can ask how its Hodge structure varies when the curve moves inside the surface. In this paper we develop a general theory to study the infinitesimal version of this question in the case of ample curves. We can then apply the machinery to show that the infinitesimal variation of Hodge structure of a general deformation of an ample curve in $\mathbb{P}^1 \times \mathbb{P}^1$ is an isomorphism.

1. INTRODUCTION

Let $\pi: \mathcal{C} \rightarrow B$ be a family of (smooth projective) complex curves of genus g over a smooth complex manifold B , which we consider as a deformation of a fibre $C = \pi^{-1}(0)$ over a distinguished point $0 \in B$. A tangent vector $v \in T_{B,0}$ induces a first-order infinitesimal deformation of C , whose Kodaira-Spencer class we denote by $\xi_v \in H^1(C, T_C)$. Recall that $H^1(C, T_C)$ is naturally identified with the tangent space of the moduli space (or stack) \mathcal{M}_g at $[C]$, hence we can think of elements in $H^1(C, T_C)$ as tangent directions in \mathcal{M}_g .

The infinitesimal variation of Hodge structure (IVHS) associated with the family $\pi: \mathcal{C} \rightarrow B$ is given by a linear map

$$\theta: T_{B,0} \otimes H^0(C, \Omega_C^1) \rightarrow H^1(C, \mathcal{O}_C)$$

where

$$(1.1) \quad \theta(v \otimes -) : H^0(C, \Omega_C^1) \rightarrow H^1(C, \mathcal{O}_C)$$

is given (up to non-zero scalar) by the product with the Kodaira-Spencer class ξ_v . The kernel of this product map consists of the holomorphic 1-forms on C that extend to the first-order infinitesimal neighbourhood of $C \subseteq \mathcal{C}$ (in the direction v), which can thus be considered as “infinitesimally constant” or invariant in the variation of Hodge structure.

It is known ([LP16, GAT21]) that, for any curve C , the product with a general $\xi \in H^1(C, T_C)$ induces an isomorphism $H^0(C, \Omega_C^1) \rightarrow H^1(C, \mathcal{O}_C)$. Hence a general deformation of C has maximal variation of Hodge structure, since the nearby fibres have no common holomorphic 1-form. We have recently used this fact to prove that for $g \geq 4$, the Torelli map $\mathcal{M}_g \rightarrow \mathcal{A}_g$ (as well as other natural modular maps) is infinitesimally rigid as a morphism of smooth DM-stacks ([CGAT23]).

The above results consider general deformations of C , but in many cases it is useful to restrict the kind of deformations to those preserving certain geometric features of C . In particular, it is natural to consider curves lying in a particular surface S . This set-up has important applications to the study

Date: February 26, 2024.

The project is partially supported by the Alexander von Humboldt Foundation.

of moduli spaces of curves \mathcal{M}_g in low genus, so that a general curve sits inside a given surface. For instance, the general curve of genus 3 is a smooth quartic and therefore properties of the moduli space of genus 3 curves can be given by studying them as plane curves (see for example [GK06]). A very important example is given by curves contained in K3 surfaces, as a general curve of genus $g \leq 9$ or equal to 11 is contained in a K3 surface, and many results about the moduli space \mathcal{M}_g are proved by using related properties (see for example [Laz86], [Muk96], [Voi02], [Voi05]).

The important point for our study is that, as soon as the genus grows, the general curve of genus g is not expected to be sitting inside a prescribed surface. Therefore, it might very well happen that the general deformations *within these restrictions* no longer induce isomorphisms in (1.1). Results in this direction provide geometric information on the tangent bundle of the corresponding sublocus of the moduli space of curves \mathcal{M}_g . Later in the paper we concentrate on the case of curves in $\mathbb{P}^1 \times \mathbb{P}^1$, which is related to the study of the k -gonal locus in \mathcal{M}_g . Indeed, curves admitting a g_k^1 and a g_l^1 , with coprime k, l , can be thought of as curves in $\mathbb{P}^1 \times \mathbb{P}^1$ of bidegree (k, l) .

In this paper we address the case of deformations of an *ample* curve C inside a given fixed surface S . If we denote by $L = \mathcal{O}_S(C)$ the associated (ample) line bundle on S , then we consider only (infinitesimal) *embedded* deformations of C in S within the linear system $|L|$. If $\sigma \in H^0(S, L)$ is a section vanishing precisely on C , such deformations are given by

$$C = \{\sigma + \epsilon\tau = 0\} \subseteq S \times \text{Spec}(\mathbb{C}[\epsilon]/(\epsilon^2)),$$

where $\tau \in H^0(S, L)$ is another section of L and as usual ϵ denotes the parameter of the infinitesimal deformation.

The problem is then to analyze the kernel (or the rank) of the product map $\xi_\tau \cdot: H^0(C, \Omega_C^1) \rightarrow H^1(C, \mathcal{O}_C)$, where $\xi_\tau \in H^1(T_C)$ denotes the associated Kodaira-Spencer class.

Note that there is a natural restriction map $H^0(S, \Omega_S^1) \rightarrow H^0(C, \Omega_C^1)$, which is injective if C is ample (Proposition 2.7). The 1-forms on C arising this way will be trivially preserved in any deformation of C in S . Hence $\ker(\xi_\tau \cdot)$ will always contain a $q(S)$ -dimensional subspace (where $q(S) = h^1(S, \mathcal{O}_S) = h^0(S, \Omega_S^1)$ is the irregularity of the ambient surface).

This motivates the following definition:

Definition 1.1. *Using the above notations, we say:*

- (1) $\tau \in H^0(S, L)$ has maximal (embedded) infinitesimal variation, if the cup-product map with $\xi_\tau \in H^1(C, T_C)$ has the maximal possible rank, i.e.

$$\text{rk}(\xi_\tau \cdot: H^0(C, \Omega_C^1) \rightarrow H^1(C, \mathcal{O}_C)) = g(C) - q(S),$$

- (2) the curve $C \subseteq S$ has maximal (embedded) infinitesimal variation, if there is a $\tau \in H^0(S, L)$ with maximal infinitesimal variation, and
- (3) the (ample) line bundle L has maximal infinitesimal variation if every smooth member $C \in |L|$ has maximal infinitesimal variation.

If only a general $C \in |L|$ has maximal infinitesimal variation, we say that L has generically maximal infinitesimal variation.

Remark 1.2. *The terminology “family of maximal variation” has been used since Viehweg [Vie83] with a different meaning, namely that no subfamily is isotrivial (has mutually birational fibres). For families of smooth curves and using the language of moduli spaces, this means that the map from the base of the family to \mathcal{M}_g has image of dimension $\dim B$. This is the meaning used for example by Y. Dutta and D. Huybrechts in [DH22] regarding linear systems of curves in K3 surfaces. For them, a line bundle L has “maximal variation” if the rational map $|L| \dashrightarrow \mathcal{M}_g$ mapping a smooth member of $|L|$ to its isomorphism class is generically of finite degree, i.e. has image of dimension $\dim |L|$. But it could still happen that all smooth curves in $|L|$ have a non-trivial common Hodge-substructure. Thus our notion of “maximal variation” is much stronger. In order to avoid confusions we have added the word “infinitesimal”, which should remind that we are considering the infinitesimal variations of Hodge structure.*

The question we would like to answer is whether, on any given surface S , sufficiently ample line bundles L have maximal infinitesimal variation. Recall from [Gre85] that a property holds for sufficiently ample line bundles if there is an ample line bundle L_0 such that the property holds for any line bundle L such that $L \otimes L_0^\vee$ is ample (i.e. L is “more ample” than L_0). With this language, our conjecture can be stated as follows:

Conjecture 1.3. *Let S be a projective surface. Then any sufficiently ample line bundle L has maximal infinitesimal variation.*

To our best knowledge, this problem has only been recently addressed in [FP22], where they prove the conjecture for plane curves of degree $d \geq 3$, i.e. for all $L = \mathcal{O}_{\mathbb{P}^2}(d)$ with $d \geq 3$ (thus $L_0 = \mathcal{O}_{\mathbb{P}^2}(2)$).

The main tool in Favale-Pirola’s proof is Macaulay’s theorem on the structure of the Jacobian ring of the plane curve. In this paper we exploit and further develop Green’s generalization of Jacobian ideals and rings for smooth hypersurfaces ([Gre85]) to address the problem for arbitrary projective surfaces, on which some of the nice properties of Jacobian rings of plane curves no longer hold. We can then apply our machinery to prove Conjecture 1.3 for $S = \mathbb{P}^1 \times \mathbb{P}^1$.

Theorem 1.4 (Theorem 4.6). *Let $C \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ be a smooth ample curve. Then a general embedded deformation of C has maximal infinitesimal variation in the sense of Definition 1.1.*

It is natural to ask up to which extent our proof can be extended to more general surfaces. On the one hand, an important step in our proof relies on the second statement of Theorem 4.2, which can be understood as a generalization of the fact that the classical polar map of plane curves is birational onto the image. An equivalent statement for a curve C in a surface S would be that the linear subsystem of $|\mathcal{O}_C(C)|$ given by the image of the natural restriction map

$$H^0(S, T_S) \rightarrow H^0(C, N_{C/S}) = H^0(C, \mathcal{O}_C(C))$$

has no base points and induces a birational map $C \rightarrow \mathbb{P}^r$ (Remark 4.7). This could hold if T_S is positive enough, and it would be interesting to characterize for which surfaces it holds.

On the other hand, the whole strategy of the proof relies on a special property of the 1-forms $\alpha \in H^0(C, \Omega_C^1)$ not arising as restrictions from $H^0(S, \Omega_S^1)$ but lying on the kernel of a general ξ_τ (see

Lemma 3.5), and showing that $\alpha = 0$ is the only possibility. However, this cannot be the case if K_S is positive enough (see Remark 3.6), and somehow limits the applicability of our strategy to surfaces with $p_g(S) \leq 1$, such as rational, K3 or abelian surfaces.

In any case some explicit computations for curves in surfaces $S \subseteq \mathbb{P}^3$ of general type (degrees 5 and 6) make us believe that Conjecture 1.3 actually holds for every surface.

The paper is structured as follows: in Section 2 we develop the general technical framework of generalized Jacobian ideals and rings to study variations of Hodge structures of smooth hypersurfaces, expanding the theory introduced by Green in [Gre85]. In Section 3 we focus in the case of curves on surfaces, proving the crucial Lemma 3.5 and Theorem 3.8, which assumes that the curve does not have maximal infinitesimal variation. Finally in Section 4 we study in detail the case of curves in $\mathbb{P}^1 \times \mathbb{P}^1$, firstly proving that the generalized Jacobian ideals indeed coincide with the naive definition that one could make using the partial derivatives of a bihomogeneous equation, secondly studying the maps induced by the generalized Jacobian ideals in low degrees (analogous to the polar map of a curve in \mathbb{P}^2), and finally proving Theorem 4.4, which contradicts the lower-bound proved in Theorem 3.8 and finishes the proof of Theorem 1.4.

Acknowledgements: We are particularly grateful to Gian Pietro Pirola, for some very insightful conversations we shared during his visit at Leibniz Universität Hannover.

2. GENERALIZED JACOBIAN IDEALS AND RINGS

The main technical tool to prove Theorem 1.4 will be generalized Jacobian ideals and rings, which were introduced by Green in [Gre85] in order to prove the infinitesimal Torelli theorem for sufficiently ample smooth hypersurfaces, as well as to study the local structure of the associated period maps. We will actually need some more precise results than those developed by Green. For the sake of completeness, we will recall the construction of generalized Jacobian ideals.

Since the constructions and first results make sense for (smooth) hypersurfaces on complex manifolds of arbitrary dimension, we start with this more general setting. Only at the end of the section we will restrict ourselves to the case of curves moving on surfaces.

2.1. The bundle Σ_L . Let X be a smooth complex variety of dimension n and L a line bundle on X . In this section we summarize the properties of Σ_L , the bundle of first-order differential operators on sections of L , which is necessary to define the generalized Jacobian ideals and rings. For a more detailed exposition about Σ_L with proofs, we refer to [Ser06].

Suppose (x_1, \dots, x_n) are coordinates on an open subset $U \subseteq X$, and that $L|_U$ is trivialized by a nowhere-vanishing section $e \in H^0(U, L)$, so that any other section $\sigma \in H^0(U, L)$ has the form $\sigma = fe$ for a holomorphic function $f \in \mathcal{O}_X(U)$.

A local section D of Σ_L is then of the form $D = a_0 \cdot 1 + \sum_{i=1}^n a_i D_i$, where $a_0, \dots, a_n \in \mathcal{O}_X(U)$, and D acts on $\sigma = fe$ by differentiating the function f , i.e.

$$(2.1) \quad D(\sigma) = D(fe) := \left(a_0 \cdot f + \sum_{i=1}^n a_i \frac{\partial f}{\partial x_i} \right) e.$$

Imposing that this formula is compatible with the transition functions for L on the overlap of two such open subsets gives the transition functions for Σ_L .

The bundle Σ_L fits into a natural short exact sequence

$$(2.2) \quad 0 \longrightarrow \mathcal{O}_X \longrightarrow \Sigma_L \longrightarrow T_X \longrightarrow 0,$$

where the inclusion $\mathcal{O}_X \hookrightarrow \Sigma_L$ is locally given by $a \mapsto a \cdot 1$ (i.e. by considering holomorphic functions as differential operators of order 0 via multiplication), and the projection $\Sigma_L \rightarrow T_X$ is locally given by $D_i \mapsto \frac{\partial}{\partial x_i}$.

Remark 2.1. *The extension class of (2.2) coincides (up to non-zero scalar multiple) with the first Chern class of L*

$$c_1(L) \in H^1(X, \Omega_X^1) \cong \text{Ext}_{\mathcal{O}_X}^1(\mathcal{O}_X, \Omega_X^1) \cong \text{Ext}_{\mathcal{O}_X}^1(T_X, \mathcal{O}_X).$$

2.2. The differential $d\sigma$ of a section $\sigma \in H^0(L)$ and associated bundles.

Definition 2.2. *Let $\sigma \in H^0(X, L)$ be a global section. The differential of σ is the morphism of vector bundles*

$$d\sigma: \Sigma_L \rightarrow L, \quad D \mapsto d\sigma(D) := D(\sigma)$$

mapping a differential operator $D \in H^0(U, \Sigma_L)$ defined on an open subset $U \subseteq X$ to the derivative of the restriction $\sigma|_U$ along D .

The kernel of $d\sigma$ is denoted by

$$\mathcal{E}_\sigma = \ker d\sigma \subseteq \Sigma_L,$$

The following result summarizes the first properties of $d\sigma$ and \mathcal{E}_σ .

Proposition 2.3. *Let $\sigma \in H^0(X, L)$ be a section of L and denote by $Z = \{\sigma = 0\} \subseteq X$ the vanishing subvariety of σ . Then:*

- (1) *The cokernel of $d\sigma: \Sigma_L \rightarrow L$ is supported on the singular locus of Z . In particular, $d\sigma$ is surjective if Z is smooth.*
- (2) *If Z is smooth there is a commutative diagram with exact rows and columns*

$$(2.3) \quad \begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & \mathcal{E}_\sigma & \xrightarrow{\cong} & T_X(-\log Z) & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathcal{O}_X & \longrightarrow & \Sigma_L & \longrightarrow & T_X \longrightarrow 0 \\ & & \parallel & & \downarrow d\sigma & & \downarrow \widetilde{d\sigma} \\ 0 & \longrightarrow & \mathcal{O}_X & \xrightarrow{\sigma} & L & \longrightarrow & L|_Z \longrightarrow 0 \\ & & & & \downarrow & & \downarrow \\ & & & & 0 & & 0 \end{array}$$

where $\widetilde{d\sigma}: T_X \rightarrow L|_Z$ coincides with the composition $T_X \rightarrow T_{X|Z} \rightarrow N_{Z/X} \cong L|_Z$.

(3) If Z is smooth, then

$$(2.4) \quad \det \mathcal{E}_\sigma \cong K_X^\vee \otimes L^\vee.$$

Proof. (1) Follows directly from the local description (2.1) of $d\sigma(D) = D(\sigma)$ on a trivialization of L , since Z is locally defined by $\{f = 0\}$.

(2) Note first that $d\sigma$ acts on the subsheaf $\mathcal{O}_X \subseteq \Sigma_L$ by multiplying by σ . There is thus a commutative diagram

$$(2.5) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X & \longrightarrow & \Sigma_L & \longrightarrow & T_X \longrightarrow 0 \\ & & \parallel & & \downarrow d\sigma & & \downarrow \widetilde{d\sigma} \\ 0 & \longrightarrow & \mathcal{O}_X & \xrightarrow{\sigma} & L & \longrightarrow & L|_Z \longrightarrow 0 \end{array}$$

where $\widetilde{d\sigma}: T_X \rightarrow L|_Z$ is locally given by differentiating the local equation of σ along a local vector field χ and restricting it to Z , which turns out to be well-defined, independent of the choice of a lifting of χ to Σ_L . Writing the isomorphism $L|_Z \cong N_{Z/X}$ with the normal bundle in local coordinates gives the alternative description of $\widetilde{d\sigma}$ as the composition $T_X \rightarrow T_{X|Z} \rightarrow N_{Z/X} \cong L|_Z$.

With this description it is clear that a vector field lies in $\ker \widetilde{d\sigma}$ if and only if it is tangent to Z along Z . That is, $\ker \widetilde{d\sigma} = T_X(-\log Z) \subseteq T_X$, the dual of the vector bundle $\Omega_X^1(\log Z)$ of meromorphic 1-forms with at most a logarithmic pole along Z .

The diagram (2.3) is obtained by completing (2.5) using the snake lemma and the surjectivity of $d\sigma$.

(3) Since both the central row and the central column are short exact sequences of locally free sheaves, we obtain

$$\det \mathcal{E}_\sigma \cong (\det \Sigma_L) \otimes L^\vee \cong (\det T_X) \otimes L^\vee \cong K_X^\vee \otimes L^\vee.$$

□

Using $d\sigma$ we define the following cohomology class $\delta\sigma \in H^1(X, \mathcal{E}_\sigma \otimes L^\vee)$, which will be very useful for certain formalizations.

Definition 2.4. Let $\sigma \in H^0(L)$ with smooth vanishing locus. We denote by

$$\delta\sigma := \left[0 \rightarrow \mathcal{E}_\sigma \rightarrow \Sigma_L \xrightarrow{d\sigma} L \rightarrow 0 \right] \in \text{Ext}_{\mathcal{O}_X}^1(L, \mathcal{E}_\sigma) = H^1(X, \mathcal{E}_\sigma \otimes L^\vee),$$

the extension class of the central column of (2.5).

Note that $\delta\sigma$ coincides with the image of $1 \in H^0(\mathcal{O}_X)$ under the connecting homomorphism of the same extension tensored by L^\vee :

$$0 \longrightarrow \mathcal{E}_\sigma \otimes L^\vee \longrightarrow \Sigma_L \otimes L^\vee \xrightarrow{d\sigma} \mathcal{O}_X \longrightarrow 0.$$

The Koszul complex of $d\sigma: \Sigma_L \otimes L^\vee \rightarrow \mathcal{O}_X$ has the form

$$0 \longrightarrow \bigwedge^{n+1} \Sigma_L \otimes L^{-n-1} \longrightarrow \bigwedge^n \Sigma_L \otimes L^{-n} \longrightarrow \cdots \longrightarrow \bigwedge^2 \Sigma_L \otimes L^{-2} \longrightarrow \Sigma_L \otimes L^\vee \longrightarrow \mathcal{O}_X \longrightarrow 0$$

Using the isomorphism $\det \Sigma_L \cong \det \mathcal{E}_\sigma \otimes L$ at the first term, the Koszul complex can be split into short exact sequences

$$(2.6) \quad 0 \longrightarrow \bigwedge^r \mathcal{E}_\sigma \otimes L^{-r} \longrightarrow \bigwedge^r \Sigma_L \otimes L^{-r} \longrightarrow \bigwedge^{r-1} \mathcal{E}_\sigma \otimes L^{-r+1} \longrightarrow 0,$$

for $r = 1, \dots, n$, whose extension classes all coincide with $\delta\sigma$ under the canonical identification $\mathrm{Ext}_{\mathcal{O}_X}^1 \left(\bigwedge^{r-1} \mathcal{E}_\sigma \otimes L^{-r+1}, \bigwedge^r \mathcal{E}_\sigma \otimes L^{-r} \right) \cong \mathrm{Ext}_{\mathcal{O}_X}^1 (L, \mathcal{E}_\sigma)$.

2.3. Generalized Jacobian rings. Let now M be another line bundle on X . We still denote with $d\sigma$ the map $\Sigma_L \otimes L^\vee \otimes M \rightarrow M$ induced by tensoring $d\sigma$ with the identity on M .

Definition 2.5. *Following Green, we define*

(1) *the generalized Jacobian ideal (with respect to σ)*

$$J_M := \mathrm{im} \left(d\sigma: H^0(\Sigma_L \otimes M \otimes L^\vee) \longrightarrow H^0(M) \right) \subseteq H^0(M)$$

(2) *and the generalized Jacobian ring $R_M := H^0(M) / J_M$*

on degree M .

Note that $H^0(M)$ is not a ring, so neither $J_M \subseteq H^0(M)$ is an ideal nor R_M is a ring. All these spaces are merely vector spaces. However, the natural multiplication maps $H^0(M) \otimes H^0(N) \rightarrow H^0(M \otimes N)$ map the subspaces $J_M \otimes H^0(N)$ and $H^0(M) \otimes J_N$ into $J_{M \otimes N}$, and this allows also to define a multiplication map $R_M \otimes R_N \rightarrow R_{M \otimes N}$. Loosely speaking, although the following construction only works if $h^1(\mathcal{O}_X) = 0$, we can think of the direct sum $\bigoplus_M H^0(M)$ (over all isomorphism classes of line bundles M) as a commutative ring graded by $\mathrm{Pic}(X)$, and then the subspace $\bigoplus_M J_M$ forms a graded ideal with quotient ring $\bigoplus_M R_M$. The generalized Jacobian ideal J_M and ring R_M are just the summands of degree M .

Example 2.6. *For $S = \mathbb{P}^2$ (with homogeneous coordinates $[x_0 : x_1 : x_2]$) and $L = \mathcal{O}_{\mathbb{P}^2}(n)$ with $n \neq 0$ we have $\Sigma_L \cong \mathcal{O}_{\mathbb{P}^2}(1)^{\oplus 3}$, and the sequence (2.2) is the Euler-sequence with the first map suitably rescaled.*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^2} & \longrightarrow & \mathcal{O}_{\mathbb{P}^2}(1)^{\oplus 3} & \longrightarrow & T_{\mathbb{P}^2} \longrightarrow 0 \\ & & & & 1 \longmapsto & & \frac{1}{n}(x_0, x_1, x_2) \end{array}$$

For any $F \in H^0(\mathbb{P}^2, L) = \mathbb{C}[x_0, x_1, x_2]_n$, the differential $dF: \mathcal{O}_{\mathbb{P}^2}(1)^{\oplus 3} \rightarrow \mathcal{O}_{\mathbb{P}^2}(n)$ is given by the partial derivatives $\left(\frac{\partial F}{\partial x_0}, \frac{\partial F}{\partial x_1}, \frac{\partial F}{\partial x_2} \right)$. The rescaling by $\frac{1}{n}$ ensures that dF restricted to $\mathcal{O}_{\mathbb{P}^2}$ coincides with the F , since the Euler relation gives

$$dF \left(\frac{1}{n}(x_0, x_1, x_2) \right) = \frac{1}{n} \left(x_0 \frac{\partial F}{\partial x_0} + x_1 \frac{\partial F}{\partial x_1} + x_2 \frac{\partial F}{\partial x_2} \right) = F.$$

The generalized Jacobian ideal in “degree” $\mathcal{O}_{\mathbb{P}^2}(m)$ is the image of

$$H^0(\mathcal{O}_{\mathbb{P}^2}(m-n+1))^{\oplus 3} \xrightarrow{dF} H^0(\mathcal{O}_{\mathbb{P}^2}(m)),$$

i.e. it is generated by the partial derivatives of F multiplied by arbitrary polynomials of the correct degree. This means $J_{\mathcal{O}_{\mathbb{P}^2}(m)} = \left(\frac{\partial F}{\partial x_0}, \frac{\partial F}{\partial x_1}, \frac{\partial F}{\partial x_2} \right)_m$ is precisely the piece of degree m of the usual Jacobian ideal $J^F \subseteq \mathbb{C}[x_0, x_1, x_2]$, which motivates the name.

Since for any M and any $r = 1, \dots, n$ the extension class of

$$(2.7) \quad 0 \longrightarrow \bigwedge^r \mathcal{E}_\sigma \otimes L^{-r} \otimes M \longrightarrow \bigwedge^r \Sigma_L \otimes L^{-r} \xrightarrow{d\sigma} \bigwedge^{r-1} \mathcal{E}_\sigma \otimes L^{-r+1} \otimes M \longrightarrow 0$$

is still $\delta\sigma \in H^1(\mathcal{E}_\sigma \otimes L^\vee)$, the coboundary morphisms

$$(2.8) \quad H^r \left(\bigwedge^{r-1} \mathcal{E}_\sigma \otimes L^{-r+1} \otimes M \right) \longrightarrow H^{r+1} \left(\bigwedge^r \mathcal{E}_\sigma \otimes L^{-r} \otimes M \right)$$

are given by cup-product with $\delta\sigma$. In particular we have

$$J_M = \text{im} (d\sigma: H^0(\Sigma_L \otimes M \otimes L^\vee \rightarrow H^0(M))) = \ker (d\sigma: H^0(M) \rightarrow H^1(\mathcal{E}_\sigma \otimes L^\vee \otimes M))$$

and thus cup-product with $\delta\sigma$ induces an inclusion

$$(2.9) \quad R_M = \text{coker} (d\sigma: H^0(\Sigma_L \otimes M \otimes L^\vee \rightarrow H^0(M))) \xrightarrow{\delta\sigma} \\ \cong \ker (H^1(\mathcal{E}_\sigma \otimes L^\vee \otimes M) \rightarrow H^1(\Sigma_L \otimes L^\vee \otimes M)) \subseteq H^1(\mathcal{E}_\sigma \otimes L^\vee \otimes M),$$

which is an equality if for example $H^1(\Sigma_L \otimes L^\vee \otimes M) = 0$.

2.4. Jacobian in degree $K_X \otimes L$. We now focus on the generalized Jacobian ideal and ring on degree $K_X \otimes L$, which turns out to be the piece on which the cup-product with the Kodaira-Spencer class of a general embedded deformation should become an isomorphism.

Proposition 2.7. *Suppose L is ample and $\sigma \in H^0(L)$ is a global section with smooth vanishing divisor $Z := \{\sigma = 0\}$. Then the following hold:*

- (1) $J_{K_X \otimes L} = H^0(K_X) \cdot \sigma = \text{im} \left(H^0(K_X) \xrightarrow{\cdot\sigma} H^0(K_X \otimes L) \right)$.
- (2) The adjunction formula induces a natural inclusion $R_{K_X \otimes L} \hookrightarrow H^0(K_Z)$.
- (3) The natural restriction morphism

$$H^0(\Omega_X^{n-1}) \longrightarrow H^0(\Omega_{X|Z}^{n-1}) \longrightarrow H^0(K_Z),$$

is injective, hence we can identify $H^0(\Omega_X^{n-1})$ with its image in $H^0(K_Z)$.

- (4) Moreover it holds $H^0(K_Z) = H^0(\Omega_X^{n-1}) \oplus R_{K_X \otimes L}$, and in particular $\dim R_{K_X \otimes L} = p_g(Z) - h^{n-1,0}(X)$.

Proof. Twisting the diagram (2.5) with K_X and using the adjunction formula for $Z \subseteq X$ we obtain

$$(2.10) \quad \begin{array}{ccccccc} 0 & \longrightarrow & K_X & \longrightarrow & \Sigma_L \otimes K_X & \longrightarrow & \Omega_X^{n-1} \longrightarrow 0 \\ & & \parallel & & \downarrow d\sigma & & \downarrow \\ 0 & \longrightarrow & K_X & \xrightarrow{\sigma} & K_X \otimes L & \longrightarrow & (K_X \otimes L)|_Z \cong K_Z \longrightarrow 0 \end{array}$$

where the right vertical arrow is the restriction of $(n-1)$ -forms from X to Z .

- (1) The connecting homomorphism $H^0(\Omega_X^{n-1}) \rightarrow H^1(K_X)$ of the long exact sequence in cohomology of the upper row is given by cup product with the corresponding extension class, which is $c_1(L) \in H^1(\Omega_X^1)$. Since L is ample, Lefschetz theorem implies that the connecting homomorphism is an isomorphism. Thus there is an equality $H^0(K_X) = H^0(\Sigma_L \otimes K_X)$ and $J_{K_X \otimes L} = \text{im}(d\sigma: H^0(\Sigma_L \otimes K_X) \rightarrow H^0(K_X \otimes L)) = \text{im}(\cdot\sigma: H^0(K_X) \rightarrow H^0(K_X \otimes L))$.

- (2) Assuming (1) we have

$$\begin{aligned} R_{K_X \otimes L} &= H^0(K_X \otimes L) / J_{K_X \otimes L} = \text{coker}(H^0(K_X) \rightarrow H^0(K_X \otimes L)) = \\ &= \ker(H^0(K_Z) \rightarrow H^1(K_X)) \subseteq H^0(K_Z). \end{aligned}$$

- (3) The first morphism $H^0(\Omega_X^{n-1}) \rightarrow H^0(\Omega_{X|Z}^{n-1})$ is injective, since its kernel is

$$H^0(\Omega_X^{n-1}(-Z)) \cong H^n(\Omega_X^1(Z))^* = 0,$$

where the first isomorphism follows from Serre duality, and the second equality follows from Kodaira-Nakano vanishing theorem.

The second morphism $H^0(\Omega_{X|Z}^{n-1}) \rightarrow H^0(K_Z)$ is also injective, since its kernel is

$$H^0(T_Z \otimes K_{X|Z}) = H^0(T_Z \otimes K_Z \otimes L_{|Z}^\vee) \cong H^{n-1}(\Omega_Z^1 \otimes L_{|Z})^*,$$

which vanishes again by Kodaira-Nakano (on the hypersurface Z) because $L_{|Z}$ is ample.

- (4) From diagram (2.10) we obtain a commutative diagram

$$\begin{array}{ccc} H^0(\Omega_X^{n-1}) & \xrightarrow[\cong]{\cdot(c_1(L))} & H^1(K_X) \\ \downarrow & & \parallel \\ H^0(K_Z) & \xrightarrow{\nu} & H^1(K_X) \end{array}$$

Since the upper map is an isomorphism, we recover the injectivity of the left map and also obtain that $H^0(\Omega_X^{n-1}) \oplus \ker \nu = H^0(K_Z)$. But $\ker \nu = R_{K_X \otimes L}$ by the proof of (2), and the proof is finished. □

Example 2.8. For $X = \mathbb{P}^2$ we have $H^0(K_X) = H^0(\Omega_X^1) = 0$. Hence for any $Z \subseteq X$ defined by an homogeneous polynomial of degree n Proposition 2.7 recovers $J_{K_X \otimes L} = J_{n-3} = 0$ (since the Jacobian ideal is generated by polynomials of degree $n-1$) and $H^0(\Omega_Z^1) \cong R_{n-3} = \mathbb{C}[x_0, x_1, x_2]_{n-3}$.

2.5. Jacobian in degree L . We have seen in Proposition 2.7 that the Jacobian ring $R_{K_X \otimes L}$ of degree $K_X \otimes L$ is a complement in $H^0(K_Z)$ of the set of top-forms on Z obtained by restricting global $(n-1)$ -forms of X . This makes it a good candidate to be the “most variable” part of $H^0(K_Z)$ under embedded deformations of Z in X . We now check that the Jacobian ring R_L of degree $L = \mathcal{O}_X(Z)$ is a good tool to encode the (Kodaira-Spencer classes of) first-order deformations of Z within the linear system $|L|$.

The identification $|L| = \mathbb{P}(H^0(L))$ also gives $T_{[Z]}|L| \cong H^0(L)/\langle\sigma\rangle$, where $Z = \{\sigma = 0\}$. This means that every first order deformation of Z inside the linear system $|L|$ is given by $\{\sigma + \epsilon\tau = 0\}$, where $\tau \in H^0(L)$ is another section and ϵ is the parameter (with $\epsilon^2 = 0$).

We now consider the following two short exact sequences of sheaves:

$$(2.11) \quad 0 \longrightarrow \mathcal{O}_X \xrightarrow{\sigma} L \longrightarrow L|_Z \cong N_{Z/X} \longrightarrow 0$$

and

$$(2.12) \quad 0 \longrightarrow T_Z \longrightarrow T_{X|Z} \longrightarrow N_{Z/X} \longrightarrow 0.$$

From (2.11) we obtain an inclusion $T_{[Z]}|L| = H^0(L)/\langle\sigma\rangle \hookrightarrow H^0(N_{Z/X})$, which composed with the first connecting homomorphism $H^0(N_{Z/X}) \rightarrow H^1(T_Z)$ of (2.12) gives a map

$$(2.13) \quad KS: T_{[Z]}|L| \longrightarrow H^1(T_Z).$$

This is precisely the Kodaira-Spencer map for the family of deformations of Z inside the linear system $|L|$.

Lemma 2.9. *The Kodaira-Spencer map (2.13) factorizes as*

$$(2.14) \quad T_{[Z]}|L| = H^0(L)/\langle\sigma\rangle \longrightarrow R_L = H^0(L)/J_L \longrightarrow H^1(T_Z)$$

Proof. Consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_X & \longrightarrow & \Sigma_L & \longrightarrow & T_X \longrightarrow 0 \\ & & \parallel & & \downarrow d\sigma & & \downarrow \tilde{d}\sigma \\ 0 & \longrightarrow & \mathcal{O}_X & \xrightarrow{\sigma} & L & \longrightarrow & L|_Z \cong N_{Z/X} \longrightarrow 0 \end{array}$$

Note first that $\sigma = d\sigma(1) \in J_L$, hence the first projection $H^0(L)/\langle\sigma\rangle \rightarrow R_L$ in (2.14) is well-defined and has kernel $J_L/\langle\sigma\rangle$.

It remains to show that $J_L/\langle\sigma\rangle \subseteq \ker(KS)$. To this aim, recall that the map $T_X \rightarrow N_{Z/X}$ factorizes as $T_X \rightarrow T_{X|Z} \rightarrow N_{Z/X}$ (Proposition 2.3), and thus

$$\begin{aligned} J_L/\langle\sigma\rangle &= \operatorname{im}(H^0(\Sigma_L) \rightarrow H^0(N_{Z/X})) \subseteq \operatorname{im}(H^0(T_X) \rightarrow H^0(N_{Z/X})) \subseteq \\ &\subseteq \operatorname{im}(H^0(T_{X|Z}) \rightarrow H^0(N_{Z/X})) = \ker\left(H^0(N_{Z/X}) \xrightarrow{KS} H^1(T_Z)\right). \end{aligned}$$

□

Definition 2.10. *By abuse of notation, we will write $KS: H^0(L) \rightarrow H^1(T_Z)$ and $KS: R_L \rightarrow H^1(T_Z)$ for the maps induced by (2.13), and we will call all three maps indistinctly Kodaira-Spencer maps.*

2.6. The top degree. We now consider a third (and last) interesting degree, that plays the role of the socle degree in the case of \mathbb{P}^2 .

We consider the composition of n connecting homomorphisms as in (2.8) (with suitably varying M 's)

$$H^0(M) \longrightarrow H^n \left(\bigwedge^n \mathcal{E}_\sigma \otimes L^{-n} \otimes M \right) = H^n (\det \mathcal{E}_\sigma \otimes L^{-n} \otimes M),$$

which is given by cup-product with the cup-power $(\delta\sigma)^n \in H^n (\bigwedge^n \mathcal{E}_\sigma \otimes L^{-n})$. Composing with the natural isomorphism

$$(2.15) \quad \det \mathcal{E}_\sigma \otimes L^{-n} \cong \det \Sigma_L \otimes L^{-n-1} \cong K_X^\vee \otimes L^{-n-1}$$

we then obtain a map

$$H^0(M) \xrightarrow{(\delta\sigma)^n} H^n (K_X^\vee \otimes L^{-n-1} \otimes M).$$

Since the kernel of the first product with $\delta\sigma$ is J_M , the above map factors through the generalized Jacobian ring as

$$R_M \rightarrow H^n (K_X^\vee \otimes L^{-n-1} \otimes M).$$

In particular, for $M = K_X^2 \otimes L^{n+1}$ we obtain a map

$$(2.16) \quad \text{tr} = \text{tr}_\sigma : R_{K_X^2 \otimes L^{n+1}} \rightarrow H^n (K_X) \cong \mathbb{C}.$$

Definition 2.11. We call the map tr in (2.16) natural trace map with respect to the given section $\sigma \in H^0(L)$.

We now investigate under which conditions the map (2.16) is an isomorphism.

Lemma 2.12. *If L is sufficiently ample, more precisely if $H^1(\Sigma_L \otimes K_X^2 \otimes L^n) = 0$ and*

$$(2.17) \quad H^{r-1} \left(\bigwedge^r \Sigma_L \otimes K_X^2 \otimes L^{n+1-r} \right) = H^r \left(\bigwedge^r \Sigma_L \otimes K_X^2 \otimes L^{n+1-r} \right) = 0$$

for all $r = 2, \dots, n$, then $\text{tr}_\sigma : R_{K_X^2 \otimes L^3} \rightarrow H^n (K_X) \cong \mathbb{C}$ is an isomorphism.

Proof. Set $M = K_X^2 \otimes L^{n+1}$ in the exact sequences (2.7), which become

$$0 \longrightarrow \bigwedge^r \mathcal{E}_\sigma \otimes K_X^2 \otimes L^{n+1-r} \longrightarrow \bigwedge^r \Sigma_L \otimes K_X^2 \otimes L^{n+1-r} \longrightarrow \bigwedge^{r-1} \mathcal{E}_\sigma \otimes K_X^2 \otimes L^{n+2-r} \longrightarrow 0.$$

The condition $H^1(\Sigma_L \otimes K_X^2 \otimes L^n) = 0$ ensures that $R_{K_X^2 \otimes L^{n+1}} \xrightarrow{\delta\sigma} H^1(\mathcal{E}_\sigma \otimes K_X^2 \otimes L^n)$. The remaining conditions ensure that the connecting homomorphisms

$$H^{r-1} \left(\bigwedge^{r-1} \mathcal{E}_\sigma \otimes K_X^2 \otimes L^{n+2-r} \right) \xrightarrow{\delta\sigma} H^r \left(\bigwedge^r \mathcal{E}_\sigma \otimes K_X^2 \otimes L^{n+1-r} \right)$$

are isomorphisms. Thus multiplication with $(\delta\sigma)^n$ followed by the isomorphism (2.15) gives indeed an isomorphism

$$R_{K_X^2 \otimes L^{n+1}} \cong H^n (\det \mathcal{E}_\sigma \otimes K_X^2 \otimes L) \cong H^n (K_X) \cong \mathbb{C}.$$

□

Remark 2.13. *The vanishing conditions in Lemma 2.12 not only involve L but also Σ_L . We haven't deduced any ampleness of Σ_L from the ampleness of L , so it might happen that the exterior powers of Σ_L somehow cancel the ampleness of L and that the conditions (2.17) are rarely satisfied. This is however not the case at all because of the first fundamental exact sequence (2.2), which gives also exact sequences*

$$0 \longrightarrow \bigwedge^{r-1} T_X \longrightarrow \bigwedge^r \Sigma_L \longrightarrow \bigwedge^r T_X \longrightarrow 0$$

and thus tensoring with $K_X^2 \otimes L^{n+1-r}$ we also obtain

$$0 \longrightarrow \Omega_X^{n-r+1} \otimes K_X \otimes L^{n+1-r} \longrightarrow \bigwedge^r \Sigma_L \otimes K_X^2 \otimes L^{n+1-r} \longrightarrow \Omega_X^{n-r} \otimes K_X \otimes L^{n+1-r} \longrightarrow 0.$$

The outer terms of the above exact sequence have no higher cohomology if L is sufficiently ample, and thus the vanishing conditions of Lemma 2.12 are satisfied.

Example 2.14. *For $X = \mathbb{P}^2$ and $Z \subseteq \mathbb{P}^2$ defined by a polynomial of degree n , Lemma 2.12 is precisely the part of and Macaulay's theorem stating that the (one-dimensional) socle of the Jacobian ring is the piece of degree $3n - 6$. Indeed, in this case we have $K_X^2 \otimes L^3 = \mathcal{O}_{\mathbb{P}^2}(3n - 6)$.*

2.7. Natural pairings and duality maps. For any line bundle M , let $M' := K_X^2 \otimes L^{n+1} \otimes M^\vee$ denote the line bundle such that $M \otimes M' \cong K_X^2 \otimes L^{n+1}$. Combining the multiplication map with the natural trace map $\text{tr}_\sigma: R_{K_X^2 \otimes L^{n+1}} \rightarrow H^n(K_X) \cong \mathbb{C}$ we obtain a pairing

$$(2.18) \quad R_M \otimes R_{M'} \longrightarrow R_{K_X^2 \otimes L^{n+1}} \longrightarrow \mathbb{C},$$

and thus also (mutually dual) duality maps

$$(2.19) \quad \lambda_M: R_M \longrightarrow R_{M'}^* \quad \text{and} \quad \lambda_{M'}: R_{M'} \longrightarrow R_M^*.$$

Remark 2.15. *The main content of Macaulay's theorem for curves in \mathbb{P}^2 (or hypersurfaces in \mathbb{P}^n) is precisely that the pairings (2.18) are perfect.*

Of course, if the natural trace is not an isomorphism, there is little hope for the pairing (2.18) to be perfect. But even if $R_{K_X^2 \otimes L^{n+1}} \cong \mathbb{C}$ holds, the pairing will not be perfect in general. Thus we cannot expect the duality maps λ_M to be isomorphisms, but only aim for them to have maximal rank.

We study now the particular cases of duality maps that will be of most interest to us, namely when X is a surface and $M = L$ or $M = K_X \otimes L$.

Proposition 2.16. *If X is a surface and L is an ample line bundle such that $H^1(\Omega_X^1 \otimes L) = 0$, then the duality map $\lambda_{K_X \otimes L^2}$ fits into a natural exact sequence*

$$0 \longrightarrow H^0(\mathcal{O}_X) \xrightarrow{\cdot c_1(L)} H^1(\Omega_X^1) \longrightarrow R_{K_X \otimes L^2} \xrightarrow{\lambda_{K_X \otimes L^2}} R_{K_X \otimes L}^* \longrightarrow 0.$$

Proof. Recall from (2.9) that $\delta\sigma$ induces the isomorphism

$$R_{K_X \otimes L^2} \xrightarrow{\cdot \delta\sigma} \ker(H^1(\mathcal{E}_\sigma \otimes K_X \otimes L) \longrightarrow H^1(\Sigma_L \otimes K_X \otimes L)).$$

Twisting (2.2) with $K_X \otimes L$ we obtain $0 \rightarrow K_X \otimes L \rightarrow \Sigma_L \otimes K_X \otimes L \rightarrow \Omega_X^1 \otimes L \rightarrow 0$. Taking into account the hypothesis $H^1(\Omega_X^1 \otimes L) = 0$ and that $H^1(K_X \otimes L) = 0$ by Kodaira vanishing, the long exact sequence in cohomology gives

$$(2.20) \quad H^1(\Sigma_L \otimes K_X \otimes L) = 0, \quad \text{and thus} \quad R_{K_X \otimes L^2} \cong H^1(\mathcal{E}_\sigma \otimes K_X \otimes L).$$

Again from (2.9) we have an isomorphism

$$R_{K_X \otimes L} \stackrel{\cdot \delta \sigma}{\cong} \ker(H^1(\mathcal{E}_\sigma \otimes K_X) \rightarrow H^1(\Sigma_L \otimes K_X)).$$

Since $H^1(K_X \otimes L) = 0$ because L is ample, we obtain an exact sequence

$$(2.21) \quad 0 \rightarrow R_{K_X \otimes L} \rightarrow H^1(\mathcal{E}_\sigma \otimes K_X) \rightarrow H^1(\Sigma_L \otimes K_X) \rightarrow H^1(K_X \otimes L) = 0.$$

Consider finally the short exact sequence (2.2) twisted by K_X :

$$0 \rightarrow K_X \rightarrow \Sigma_L \otimes K_X \rightarrow \Omega_X^1 \rightarrow 0$$

whose connecting homomorphisms in cohomology are given up to sign by cup-product with $c_1(L)$. In particular we have

$$H^0(\Omega_X^1) \xrightarrow{c_1(L)} H^1(K_X) \xrightarrow{0} H^1(\Sigma_L \otimes K_X) \rightarrow H^1(\Omega_X^1) \xrightarrow{c_1(L)} H^2(K_X) \rightarrow 0,$$

where the first cup-product with $c_1(L)$ is an isomorphism and the second one is surjective because of Lefschetz theorem. Thus in particular we have a short exact sequence

$$0 \rightarrow H^1(\Sigma_L \otimes K_X) \rightarrow H^1(\Omega_X^1) \xrightarrow{c_1(L)} H^2(K_X) \rightarrow 0,$$

which together with (2.21) gives an exact sequence

$$(2.22) \quad 0 \rightarrow R_{K_X \otimes L} \rightarrow H^1(\mathcal{E}_\sigma \otimes K_X) \rightarrow H^1(\Omega_X^1) \xrightarrow{c_1(L)} H^2(K_X) \rightarrow 0.$$

The statement follows after dualizing (2.22) and using Serre duality in $H^2(K_X)^* \cong H^0(\mathcal{O}_X)$ and

$$H^1(\mathcal{E}_\sigma \otimes K_X)^* \cong H^1(\mathcal{E}_\sigma^\vee) \cong H^1(\mathcal{E}_\sigma \otimes K_X \otimes L) \stackrel{(2.20)}{\cong} R_{K_X \otimes L^2},$$

where the isomorphism in the middle follows from

$$\mathcal{E}_\sigma^\vee \cong \det \mathcal{E}_\sigma^\vee \otimes \mathcal{E} \cong (\det \Sigma_L^\vee) \otimes L \otimes \mathcal{E} \cong K_X \otimes L \otimes \mathcal{E}_\sigma.$$

□

Proposition 2.17. *Suppose that X is a surface. If $K_X \otimes L$ is ample and $H^1(\Omega_X^1 \otimes K_X \otimes L) = 0$, the duality map $\lambda_{K_X^2 \otimes L^2}$ fits into a natural exact sequence*

$$H^1(L)^* \xrightarrow{H^1(d\sigma)^*} H^1(\Sigma_L)^* \rightarrow R_{K_X^2 \otimes L^2} \xrightarrow{\lambda_{K_X^2 \otimes L^2}} R_L^* \rightarrow 0.$$

Proof. As in the previous proof, using (2.9) we obtain

$$(2.23) \quad R_{K_X^2 \otimes L^2} \stackrel{\cdot \delta \sigma}{\cong} \ker(H^1(\mathcal{E}_\sigma \otimes K_X^2 \otimes L) \rightarrow H^1(\Sigma_L \otimes K_X^2 \otimes L)).$$

From (2.2), $H^1(\Sigma_L \otimes K_X^2 \otimes L)$ sits into the exact sequence

$$0 = H^1(K_X^2 \otimes L) \rightarrow H^1(\Sigma_L \otimes K_X^2 \otimes L) \rightarrow H^1(T_X \otimes K_X^2 \otimes L) \cong H^1(\Omega_X^1 \otimes K_X \otimes L) = 0,$$

where the first term vanishes because $K_X \otimes L$ is ample, and the last term vanishes by hypothesis. Hence also $H^1(\Sigma_L \otimes K_X^2 \otimes L) = 0$ and (2.23) turns into an isomorphism $R_{K_X^2 \otimes L^2} \xrightarrow{\cdot \delta \sigma} H^1(\mathcal{E}_\sigma \otimes K_X^2 \otimes L)$.

By Serre Duality and the natural isomorphism $\mathcal{E}_\sigma^\vee \cong \mathcal{E}_\sigma \otimes K_X \otimes L$ we also obtain

$$(2.24) \quad R_{K_X^2 \otimes L^2} \cong H^1(\mathcal{E}_\sigma^\vee \otimes K_X) \cong H^1(\mathcal{E}_\sigma)^*.$$

On the other hand, also from (2.9) we can identify $R_L \xrightarrow{\cdot \delta \sigma} \ker(H^1(\mathcal{E}_\sigma) \rightarrow H^1(\Sigma_L))$, hence we have an exact sequence

$$0 \longrightarrow R_L \xrightarrow{\cdot \delta \sigma} H^1(\mathcal{E}_\sigma) \longrightarrow H^1(\Sigma_L) \xrightarrow{H^1(d\sigma)} H^1(L).$$

Dualizing it and using the natural isomorphism (2.24) we obtain the statement. \square

3. IVHS OF CURVES IN SURFACES WITH GENERALIZED JACOBIAN RINGS

In this section we show how the generalized Jacobian rings in the degrees studied above encode almost all necessary information about IVHS of an ample smooth curve Z in a surface X .

In general, an (one-parameter) IVHS of a smooth curve Z is given by a symmetric linear map

$$H^0(K_Z) \rightarrow H^1(\mathcal{O}_Z) \cong H^0(K_Z)^*.$$

If the IVHS comes from first-order infinitesimal deformation of Z , then it is given (up to non-zero scalar) by cup-product with the Kodaira-Spencer class.

Assume from now on that X is a smooth surface and $Z \subseteq X$ is the (smooth) vanishing locus of a section σ of an ample line bundle L . This gives the whole family of generalized Jacobian rings R_M for $M \in \text{Pic}(X)$ with the properties developed in the previous section.

If we consider an embedded first-order deformation given by another section $\tau \in H^0(L)$, the associated IVHS is given by cup-product with $\xi_\tau := KS(\tau) \in H^1(T_Z)$, where $KS: H^0(L) \rightarrow H^1(T_Z)$ is as defined in Definition 2.10 (see Lemma 2.9). In particular, ξ_τ depends only on the class of τ in R_L .

Recall also from Proposition 2.7 that $H^0(K_Z)$ is naturally isomorphic to the direct sum of $R_{K_X \otimes L}$ and $H^0(\Omega_X^1)$. The latter subspace is mapped to zero by any ξ_τ with $[\tau] \in R_L$, since it is preserved in all embedded deformations of Z . Thus the IVHS $\xi_\tau \cdot$ is completely determined by its restriction to $R_{K_X \otimes L}$. Composing this restriction with the dual of the inclusion $\nu: R_{K_X \otimes L} \hookrightarrow H^0(K_Z)$, $\xi_\tau \cdot$ induces a map

$$(3.1) \quad R_{K_X \otimes L} \xrightarrow{\nu} H^0(K_Z) \xrightarrow{\xi_\tau} H^1(\mathcal{O}_Z) \xrightarrow{\nu^*} R_{K_X \otimes L}^*.$$

Since both source and target spaces, as well as ξ_τ , depend on the generalized Jacobian rings, it is natural to expect that the map is also encoded on the generalized Jacobian rings. This is actually true, as we will see in Corollary 3.4, but in order to prove it we need the following technical result.

Proposition 3.1. *Let $\tau \in H^0(L)$ with associated Kodaira-Spencer class $\xi_\tau \in H^1(T_Z)$. Let also $\gamma \in H^0(K_X^2 \otimes L^2)$ and consider its restriction $\gamma|_Z \in H^0(K_X^2 \otimes L|_Z^2) \cong H^0(K_Z^2)$. Then the product*

$$\xi_\tau \cdot \gamma|_Z \in H^1(T_Z \otimes K_Z^2) \cong H^1(K_Z)$$

corresponds to

$$(\delta\sigma)^2 \cdot \tau \cdot \gamma \in H^2(\det(\mathcal{E}_\sigma \otimes L^\vee) \otimes L \otimes (K_X^2 \otimes L^2)) \cong H^2(K_X)$$

under the natural connecting (iso)morphism $H^1(K_Z) \rightarrow H^2(K_X)$.

Proof. Consider first the commutative diagram with exact rows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{E}_\sigma \cong T_X(-\log Z) & \longrightarrow & \Sigma_L & \longrightarrow & L & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & T_Z & \longrightarrow & T_{X|Z} & \longrightarrow & N_{Z/X} \cong L_{|Z} & \longrightarrow & 0 \end{array}$$

where the central vertical arrow is the composition of the surjection $\Sigma_L \twoheadrightarrow T_X$ and the restriction $T_X \rightarrow T_{X|Z}$. The first connecting homomorphisms of both rows fit in the following commutative diagram which summarizes the definition of ξ_τ :

$$(3.2) \quad \begin{array}{ccc} H^0(L) & \longrightarrow & H^1(\mathcal{E}_\sigma) \\ \downarrow & & \downarrow \\ H^0(L_{|Z}) & \longrightarrow & H^1(T_Z) \end{array} \quad \begin{array}{ccc} \tau & \longmapsto & (\delta\sigma) \cdot \tau \\ \downarrow & & \downarrow \\ \tau_{|Z} & \longmapsto & \xi_\tau \end{array}$$

Tensoring with $K_X^2 \otimes L^2$ gives the following diagram

$$(3.3) \quad \begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{E}_\sigma \otimes K_X^2 \otimes L^2 & \longrightarrow & \Sigma_L \otimes K_X^2 \otimes L^2 & \longrightarrow & K_X^2 \otimes L^3 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & T_Z \otimes K_Z^2 \cong K_Z & \longrightarrow & T_{X|Z} \otimes K_Z^2 & \longrightarrow & K_Z^2 \otimes L_{|Z} & \longrightarrow & 0 \end{array}$$

which induces the analogous diagram to (3.2) (in the right square we have multiplied by γ),

$$(3.4) \quad \begin{array}{ccc} H^0(K_X^2 \otimes L^3) & \longrightarrow & H^1(\mathcal{E}_\sigma \otimes K_X^2 \otimes L^2) \\ \downarrow & & \downarrow \\ H^0(K_Z^2 \otimes L_{|Z}) & \longrightarrow & H^1(K_Z) \end{array} \quad \begin{array}{ccc} \tau \cdot \gamma & \longmapsto & (\delta\sigma) \cdot \tau \cdot \gamma \\ \downarrow & & \downarrow \\ \tau_{|Z} \cdot \gamma_{|Z} & \longmapsto & \xi_\tau \cdot \gamma_{|Z} \end{array}$$

Consider now (2.6) with $r = 2 = \text{rk } \mathcal{E}_\sigma$ and tensorize it with $K_X^2 \otimes L^3$ to obtain (using (2.4)):

$$0 \longrightarrow \det \mathcal{E}_\sigma \otimes K_X^2 \otimes L \cong K_X \longrightarrow \bigwedge^2 \Sigma_L \otimes K_X^2 \otimes L \longrightarrow \mathcal{E}_\sigma \otimes K_X^2 \otimes L^2 \longrightarrow 0.$$

From the projection $\Sigma_L \twoheadrightarrow T_X$ we can obtain a map $\bigwedge^2 \Sigma_L \twoheadrightarrow \det T_X = K_X^\vee$, which tensored by $K_X^2 \otimes L$ fits into the following diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K_X & \longrightarrow & \bigwedge^2 \Sigma_L \otimes K_X^2 \otimes L & \longrightarrow & \mathcal{E}_\sigma \otimes K_X^2 \otimes L^2 & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & K_X & \xrightarrow{\sigma} & K_X \otimes L & \longrightarrow & K_Z & \longrightarrow & 0 \end{array}$$

where the rightmost morphism is the same as the left one in (3.3). We have thus commutative diagrams

$$(3.5) \quad \begin{array}{ccc} H^1(\mathcal{E}_\sigma \otimes K_X^2 \otimes L^2) & \longrightarrow & H^2(\mathcal{E}_\sigma \otimes K_X) & & (\delta\sigma) \cdot \tau \cdot \gamma & \longmapsto & (\delta\sigma)^2 \cdot \tau \cdot \gamma \\ \downarrow & & \downarrow & & \downarrow & & \parallel \\ H^1(K_Z) & \longrightarrow & H^2(K_X) & & \xi_\tau \cdot \gamma|_Z & \longmapsto & (\delta\sigma)^2 \cdot \tau \cdot \gamma \end{array}$$

The claim follows combining the lower rows in the right squares of (3.4) and (3.5). \square

In order to write the map (3.1) in terms of the generalized Jacobian rings, we need a bit more of notation. Recall that for any $M \in \text{Pic}(X)$ we have the duality map (2.19) $\lambda_M: R_M \rightarrow R_{M'}^*$, where $M' = M^\vee \otimes K_X^2 \otimes L^3$.

Definition 3.2. For two line bundles M, N on the surface X and a section $\alpha \in H^0(N)$ we define the linear map $\varphi_M(\alpha): R_M \rightarrow R_{(M \otimes N)'}^* = R_{K_X^2 \otimes L^3 \otimes N^\vee \otimes M^\vee}^*$ given as the composition

$$(3.6) \quad \varphi_M(\alpha): R_M \xrightarrow{\cdot\alpha} R_{M \otimes N} \xrightarrow{\lambda_{M \otimes N}} R_{K_X^2 \otimes L^3 \otimes M^\vee \otimes N^\vee}^*$$

of the multiplication by α and the duality map of $M \otimes N$.

Remark 3.3. It follows at once from the definition that

$$(3.7) \quad (\varphi_M(\alpha))^* = \varphi_{K_X^2 \otimes L^3 \otimes M^\vee \otimes N^\vee}(\alpha).$$

With this last piece of notation we can state the announced result:

Corollary 3.4. Let $\tau \in H^0(L)$ represent an embedded first-order deformation of $Z \in |L|$ with Kodaira-Spencer class $\xi_\tau := KS(\tau) \in H^1(T_Z)$. Then the cup-product with ξ_τ fits into the following commutative diagram

$$(3.8) \quad \begin{array}{ccc} H^0(K_Z) & \xrightarrow{\xi_\tau} & H^1(\mathcal{O}_Z) \\ \uparrow \nu & & \downarrow \nu^* \\ R_{K_X \otimes L} & \xrightarrow{\varphi_{K_X \otimes L}(\tau)} & R_{K_X \otimes L}^* \end{array}$$

In particular $\text{rk } \varphi_{K_X \otimes L}(\tau) \leq \text{rk}(\xi_\tau)$.

Proof. Recall from (3.6) that $\varphi_{K_X \otimes L}(\tau)$ is the composition

$$R_{K_X \otimes L} \xrightarrow{\tau} R_{K_X \otimes L^2} \xrightarrow{\lambda_{K_X \otimes L^2}} R_{K_X \otimes L}^*$$

of the multiplication map by τ and the duality map $\lambda_{K_X \otimes L^2}$. Moreover the latter is defined by the natural pairing $R_{K_X \otimes L} \otimes R_{K_X \otimes L^2} \rightarrow R_{K_X^2 \otimes L^3} \xrightarrow{\text{tr}} H^2(K_X) \cong \mathbb{C}$.

Thus the commutativity of (3.8) is equivalent to

$$(3.9) \quad \langle \varphi_{K_X \otimes L}(\tau)(\alpha), \beta \rangle = \langle \nu^*(\nu(\alpha) \cdot \xi_\tau), \beta \rangle$$

for every $\alpha, \beta \in R_{K_X \otimes L}$ (or even their representatives $\alpha, \beta \in H^0(K_X \otimes L)$) where with $\langle -, - \rangle$ we denotes the natural pairing between $R_{K_X \otimes L}$ and its dual $R_{K_X \otimes L}^*$.

On the one hand, by (2.16) and (3.6), the left hand side of (3.9) is identified with the class

$$\mathrm{tr}(\tau\alpha\beta) = (\delta\sigma)^2 \cdot \tau\alpha\beta \in H^2(K_X),$$

which by Proposition 3.1 corresponds to $\xi_\tau \cdot (\alpha\beta)|_Z$ under the isomorphism $H^2(K_X) \cong H^1(K_Z)$.

On the other hand, the right hand side of (3.9) corresponds to the class

$$\nu(\alpha) \cdot \xi_\tau \cdot \nu(\beta) = \xi_\tau \alpha|_Z \beta|_Z \in H^1(K_Z \otimes T_Z \otimes K_Z) = H^1(K_Z).$$

which clearly agrees with $\xi_\tau \cdot (\alpha\beta)|_Z$. \square

We now prove a generalization of [FP22, Lemma 2.4], giving a particular property of sections in the kernel of a general ξ_τ which allows to tackle the problem directly on $R_{K_X \otimes L}$ without need to consider the elements $\tau \in R_L$.

Lemma 3.5. *Let $\tau \in H^0(L)$ be such that $\varphi_{K_X \otimes L}(\tau) : R_{K_X \otimes L} \rightarrow R_{K_X \otimes L}^*$ has maximal rank (among all possible $\tau \in H^0(L)$). Then for every $\alpha \in \ker \varphi_{K_X \otimes L}(\tau)$ it holds $\lambda_{K_X^2 \otimes L^2}(\alpha^2) = 0 \in R_L^*$.*

Proof. The condition $\alpha \in \ker \varphi_{K_X \otimes L}(\tau)$ means that $\lambda_{K_X \otimes L^2}(\tau \cdot \alpha) = 0 \in R_{K_X \otimes L}^*$, i.e.

$$(3.10) \quad \mathrm{tr}(\tau \cdot \alpha \cdot \beta) = 0 \in H^2(K_X)$$

for every $\beta \in R_{K_X \otimes L}$.

Assuming this, we want to prove that $\mathrm{tr}(\alpha^2 \cdot w) = 0$ for every $w \in R_L$.

Let $w \in R_L$ be arbitrary. By semicontinuity, the rank of $\varphi_{K_X \otimes L}(\tau)$ is lower-semicontinuous, i.e. the maximum is attained for τ on a dense open subset of $H^0(L)$. In particular there is a small neighbourhood $U \subseteq \mathbb{C}$ of $0 \in \mathbb{C}$ such that $\varphi_{K_X \otimes L}(\tau + tw)$ is of maximal rank for all $t \in U$. Moreover when t varies, the kernels of $\varphi_{K_X \otimes L}(\tau + tw)$ form a (holomorphic) vector subbundle of the trivial vector bundle with fibre $R_{K_X \otimes L}$ over U . Thus $\alpha \in \ker \varphi_{K_X \otimes L}(\tau)$ can be extended to a (holomorphic) section over U with $\alpha(t) \in \ker \varphi_{K_X \otimes L}(\tau + tw) \subseteq R_{K_X \otimes L}$ for every $t \in U$ and $\alpha(0) = \alpha$. This means

$$\varphi_{K_X \otimes L}(\tau + tw)(\alpha(t)) = 0 \in R_{K_X \otimes L}^* \quad \forall t \in U,$$

or in other words

$$(3.11) \quad \mathrm{tr}((\tau + tw) \cdot \alpha(t) \cdot \beta) = 0 \in H^2(K_X) \cong \mathbb{C} \quad \forall t \in U, \forall \beta \in R_{K_X \otimes L}.$$

Differentiating (3.11) and evaluating at $t = 0$ we obtain

$$\mathrm{tr}(w \cdot \alpha(0) \cdot \beta + \tau \cdot \alpha'(0) \cdot \beta) = 0 \quad \forall \beta \in R_{K_X \otimes L}.$$

In particular for $\beta = \alpha = \alpha(0)$ we have

$$\mathrm{tr}(w \cdot \alpha^2) = -\mathrm{tr}(\tau \cdot \alpha'(0) \cdot \alpha) = 0,$$

where the last equality follows from (3.10) with $\beta = \alpha'(0)$. \square

In view of Lemma 3.5, one way to prove that the general first-order embedded deformation of Z has maximal infinitesimal variation is to prove that the only $\alpha \in R_{K_X \otimes L}$ with $\lambda_{K_X^2 \otimes L^2}(\alpha^2) = 0 \in R_L^*$ is $\alpha = 0$.

Remark 3.6. *This approach can only work if $\dim R_{K_X \otimes L} \leq \dim R_L$, which is not expected if K_X is positive. Indeed, for a given $\tau \in R_L$, the condition $\text{tr}(\tau \cdot \alpha^2) = 0$ defines a quadric $Q_\tau \subseteq \mathbb{P}(R_{K_X \otimes L})$. Thus $\lambda_{K_X^2 \otimes L^2}(\alpha^2) = 0$ is equivalent to α lying on the intersection $Q_{\tau_1} \cap \dots \cap Q_{\tau_m}$ for $\tau_1, \dots, \tau_m \in R_L$ a basis. If $\dim R_L = m \leq \dim \mathbb{P}(R_{K_X \otimes L}) = \dim R_{K_X \otimes L} - 1$ the intersection won't be empty, hence there will be non-zero α 's with $\lambda_{K_X^2 \otimes L^2}(\alpha^2) = 0 \in R_L^*$.*

Thus we only expect this approach to work for curves on surfaces with somehow non-positive canonical bundle, such as del Pezzo surfaces or at most K3 or Abelian surfaces.

It will actually be more convenient to work with the representatives $\alpha \in H^0(K_X \otimes L)$ (which we still denote as α for the sake of simplicity), and thus consider the elements in the kernel of the composition

$$(3.12) \quad \lambda: H^0(K_X^2 \otimes L^2) \longrightarrow R_{K_X^2 \otimes L^2} \xrightarrow{\lambda_{K_X^2 \otimes L^2}} R_L^*$$

Our aim is then to prove that the only $\alpha \in H^0(K_X \otimes L)$ with $\lambda(\alpha^2) = 0$ are those lying on $J_{K_X \otimes L}$. This leads to the following definition.

Definition 3.7. *Let $D \subseteq \mathbb{P}(H^2(K_X^2 \otimes L^2))$ be defined as*

$$D = \{[\alpha\beta] \in \mathbb{P}(H^2(K_X^2 \otimes L^2)) \mid \alpha, \beta \in H^0(K_X \otimes L), \alpha\beta \in \ker \lambda\},$$

where from now on we use square brackets to denote the corresponding point in the projectivization of a vector space.

In other words, D is the intersection of the linear subspace $\mathbb{P}(\ker \lambda)$ and the image of the Segre-multiplication map $\mathbb{P}(H^0(K_X \otimes L)) \times \mathbb{P}(H^0(K_X \otimes L)) \rightarrow \mathbb{P}(H^2(K_X^2 \otimes L^2))$.

Theorem 3.8. *Assume L has not maximal infinitesimal variation at a smooth $Z \in |L|$. Then it holds*

$$\dim D \geq h^0(K_X \otimes L) + p_g(X) - 1.$$

Proof. The proof follows some ideas of [FP22, Proposition 3.4]. Define

$$Y := \{[\alpha] \in \mathbb{P}(H^0(K_X \otimes L)) \mid \alpha^2 \in \ker \lambda\}$$

and the following incidence variety

$$W := \{([\alpha], [\beta]) \in Y \times \mathbb{P}(H^0(K_X \otimes L)) \mid \alpha\beta \in \ker \lambda\}.$$

The projection onto the first factor defines a surjective morphism $\pi: W \rightarrow Y$ ($\beta = \alpha$ gives a preimage for any $[\alpha] \in Y$), and multiplication defines a morphism $\psi: W \rightarrow D$.

Note first that ψ is a finite morphism: the zero locus of any element $[\gamma] \in D \subseteq \mathbb{P}(H^2(K_X^2 \otimes L^2))$ has finitely many decompositions into the sum of two effective divisors, hence in particular γ admits finitely many decompositions (up to multiplication of one factor by $a \in \mathbb{C}^*$) of the form $\gamma = \alpha\beta$ with $\alpha, \beta \in H^0(K_X \otimes L)$. Thus

$$(3.13) \quad \dim W \leq \dim D.$$

We consider now the fibres of π : for $[\alpha] \in Y$ it holds

$$\begin{aligned} \pi^{-1}([\alpha]) &= \{[\alpha]\} \times \{[\beta] \in \mathbb{P}(H^0(K_X \otimes L)) \mid \alpha\beta \in \ker \lambda\} \\ &\cong \mathbb{P}\left(\ker\left(H^0(K_X \otimes L) \xrightarrow{\alpha} H^0(K_X^2 \otimes L^2) \xrightarrow{\lambda} R_L^*\right)\right) \end{aligned}$$

The latter composition factorizes also as $H^0(K_X \otimes L) \rightarrow R_{K_X \otimes L} \xrightarrow{\varphi_{K_X \otimes L}(\alpha)} R_L^*$, where the first map is surjective, and thus

$$\begin{aligned} \dim \pi^{-1}([\alpha]) &= \dim \ker\left(H^0(K_X \otimes L) \rightarrow R_{K_X \otimes L} \xrightarrow{\varphi_{K_X \otimes L}(\alpha)} R_L^*\right) - 1 \\ &= h^0(K_X \otimes L) - \operatorname{rk} \varphi_{K_X \otimes L}(\alpha) - 1 \\ &\stackrel{(3.7)}{=} h^0(K_X \otimes L) - \operatorname{rk} \varphi_L(\alpha) - 1 \end{aligned}$$

Since $\pi: W \rightarrow Y$ is a surjective morphism, it holds

$$(3.14) \quad \dim W = \dim Y + \dim \pi^{-1}([\alpha]) = \dim Y + h^0(K_X \otimes L) - \operatorname{rk} \varphi_L(\alpha) - 1$$

for a general $[\alpha] \in Y$.

It remains to find a lower bound on the dimension of Y . To this aim we introduce one further incidence variety. Denote by λ' the composition

$$\lambda': H^0(K_X \otimes L^2) \rightarrow R_{K_X \otimes L^2} \xrightarrow{\lambda_{K_X \otimes L^2}} R_{K_X \otimes L}^*$$

let $U \subseteq \mathbb{P}(H^0(L))$ be the dense open subset consisting of the $[\tau]$ for which $\varphi_{K_X \otimes L}(\tau)$ has maximal rank, and set

$$I := \{([\tau], [\alpha]) \in U \times \mathbb{P}(H^0(K_X \otimes L)) \mid \tau\alpha \in \ker \lambda'\},$$

which is equipped with two projections $\mu_1: I \rightarrow U$ and $\mu_2: I \rightarrow \mathbb{P}(H^0(K_X \otimes L))$.

The fibre of μ_1 over a given $[\tau] \in U$ is the projectivization of the kernel of the composition

$$H^0(K_X \otimes L) \rightarrow H^0(K_X \otimes L) / J_{K_X \otimes L} = R_{K_X \otimes L} \xrightarrow{\varphi_{K_X \otimes L}(\tau)} R_{K_X \otimes L}^*$$

which obviously contains $J_{K_X \otimes L} = H^0(K_X) \cdot \sigma$ (Proposition 2.7), with equality if and only if $\varphi_{K_X \otimes L}(\tau)$ is an isomorphism. Thus L has maximal infinitesimal variation at $Z \in |L|$ if and only if the general fibre of μ_1 has dimension precisely $\dim \mathbb{P}(J_{K_X \otimes L}) = p_g(X) - 1$. By assumption L has not maximal infinitesimal variation, and hence μ_1 is surjective and its general fibre has dimension at least $p_g(X)$. Therefore it holds

$$(3.15) \quad \dim I \geq \dim U + p_g(X) = \dim \mathbb{P}(H^0(L)) + p_g(X) = h^0(L) + p_g(X) - 1.$$

We now consider the projection μ_2 . By Lemma 3.5, for every $([\tau], [\alpha]) \in I$ it holds $\lambda(\alpha^2) = 0$, i.e. $[\alpha] \in Y$ and thus $\mu_2(I) \subseteq Y$. As for the fibers of μ_2 it holds

$$\mu_2^{-1}([\alpha]) \cong \{[\tau] \in U \mid \varphi_L(\alpha)(\tau) = 0 \in R_{K_X \otimes L}^*\} = U \cap \mathbb{P}\left(\ker\left(H^0(L) \rightarrow R_L \xrightarrow{\varphi_L(\alpha)} R_{K_X \otimes L}^*\right)\right).$$

All in all we obtain

$$(3.16) \quad \dim I = \dim \mu_2(I) + \dim \mu_2^{-1}([\alpha]) \leq \dim Y + \dim \ker \varphi_L(\alpha) + \dim J_L - 1$$

for a general $\alpha \in \mu_2(I)$. Combining (3.15) and (3.16) we obtain

$$\dim Y \geq h^0(L) - \dim J_L - \dim \ker \varphi_L(\alpha) + p_g(X) = \operatorname{rk} \varphi_L(\alpha) + p_g(X),$$

which with (3.13) and (3.14) gives the wanted

$$\dim D \geq \dim W \geq h^0(K_X \otimes L) + p_g(X) - 1.$$

□

4. CURVES IN $\mathbb{P}^1 \times \mathbb{P}^1$

We now show that for $X = \mathbb{P}^1 \times \mathbb{P}^1$ and L sufficiently ample (with a very explicit condition on the bidegree) we can obtain an upper-bound on $\dim D$ contradicting the lower bound of Theorem 3.8, thus proving Theorem 1.4.

Let us then fix $X = \mathbb{P}^1 \times \mathbb{P}^1$. Recall that line bundles on $\mathbb{P}^1 \times \mathbb{P}^1$ are determined by its bidegree. We denote by $\mathcal{O}_X(d, e) = \mathcal{O}_{\mathbb{P}^1}(d) \boxtimes \mathcal{O}_{\mathbb{P}^1}(e)$. More generally, for any sheaf \mathcal{F} of \mathcal{O}_X -modules on X we denote by $\mathcal{F}(d, e) = \mathcal{F} \otimes \mathcal{O}_X(d, e)$.

We also denote by $(x_0 : x_1)$ resp. $(y_0 : y_1)$ homogeneous coordinates on the first resp. second factor. Consider the polynomial ring $S := \mathbb{C}[x_0, x_1, y_0, y_1]$, bigraded so that $\deg(x_i) = (1, 0)$ and $\deg(y_i) = (0, 1)$, and write $S_{d,e}$ for the summand of polynomials of bidegree (d, e) , which correspond with the global sections of $\mathcal{O}_X(d, e)$. Also for any bihomogeneous ideal $I \subseteq S$ we denote by $I_{d,e} = I \cap S_{d,e}$, the summand of elements of bidegree (d, e) .

Let now $L = \mathcal{O}_X(d, e)$ and choose an irreducible non-zero section $F \in H^0(X, L) = S_{d,e}$. Its zero locus $Z := \{F = 0\} \subseteq \mathbb{P}^1 \times \mathbb{P}^1$ is a smooth curve if and only if the four partial derivatives of F have no common zero, i.e. $\left\{ \frac{\partial F}{\partial x_0} = \frac{\partial F}{\partial x_1} = \frac{\partial F}{\partial y_0} = \frac{\partial F}{\partial y_1} = 0 \right\} = \emptyset \subseteq \mathbb{P}^1 \times \mathbb{P}^1$.

To L we can attach the bundle of first-order differential operators Σ_L , and then $F \in H^0(L)$ induces the surjective morphism $dF: \Sigma_L \rightarrow L$. For any $M = \mathcal{O}_X(a, b)$ we can consider the corresponding generalized Jacobian ideal

$$J_{a,b} := J_M = \operatorname{im} \left(H^0(\Sigma_L \otimes L^\vee \otimes M) = H^0(\Sigma_L(a-d, b-e)) \rightarrow H^0(M) \right) \subseteq H^0(M) = S_{a,b}.$$

Lemma 4.1. *The generalized Jacobian ideal coincides with the “usual” Jacobian ideal generated by the partial derivatives of F , i.e.*

$$\begin{aligned} J_{a,b} &= \left(\frac{\partial F}{\partial x_0}, \frac{\partial F}{\partial x_1}, \frac{\partial F}{\partial y_0}, \frac{\partial F}{\partial y_1} \right)_{a,b} \\ &= \left\{ P_0 \frac{\partial F}{\partial x_0} + P_1 \frac{\partial F}{\partial x_1} + Q_0 \frac{\partial F}{\partial y_0} + Q_1 \frac{\partial F}{\partial y_1} \mid P_0, P_1 \in S_{a-d+1, b-e}, Q_0, Q_1 \in S_{a-d, b-e+1} \right\}. \end{aligned}$$

In particular, the pieces of small degree are $J_{d-1, e-1} = 0$,

$$J_{d-1, e} = \left\{ a_0 \frac{\partial F}{\partial x_0} + a_1 \frac{\partial F}{\partial x_1} \mid a_0, a_1 \in \mathbb{C} \right\}, J_{d, e-1} = \left\{ b_0 \frac{\partial F}{\partial y_0} + b_1 \frac{\partial F}{\partial y_1} \mid b_0, b_1 \in \mathbb{C} \right\},$$

and $J_{d,e}$ is generated as a vector space by

$$\left\{ x_0 \frac{\partial F}{\partial x_0}, x_1 \frac{\partial F}{\partial x_0}, x_0 \frac{\partial F}{\partial x_1}, x_1 \frac{\partial F}{\partial x_1}, y_0 \frac{\partial F}{\partial y_0}, y_1 \frac{\partial F}{\partial y_0}, y_0 \frac{\partial F}{\partial y_1}, y_1 \frac{\partial F}{\partial y_1} \right\}.$$

These generators are however not linearly independent, since the Euler relations with respect to each subset of variables give

$$(4.1) \quad \frac{1}{d} \left(x_0 \frac{\partial F}{\partial x_0} + x_1 \frac{\partial F}{\partial x_1} \right) = F = \frac{1}{e} \left(y_0 \frac{\partial F}{\partial y_0} + y_1 \frac{\partial F}{\partial y_1} \right).$$

Proof of Lemma 4.1. We first describe a trivialization of Σ_L . For $i, j = 0, 1$ consider the affine open subset

$$U_{ij} := \{([x_0 : x_1], [y_0 : y_1]) \mid x_i y_j \neq 0\} \xrightarrow{\cong} \mathbb{C}^2 \\ ([x_0 : x_1], [y_0 : y_1]) \mapsto (z_{1-i}, w_{1-j}) := \left(\frac{x_{1-i}}{x_i}, \frac{y_{1-j}}{y_j} \right)$$

On U_{ij} the line bundle L is trivialized by the section $e_{ij} := x_i^d y_j^e$, and any $F(x_0, x_1, y_0, y_1) \in H^0(L)$ can be uniquely written as $F = f_{ij}(z_{1-i}, w_{1-j}) x_i^d y_j^e$ for a polynomial $f_{ij} \in \mathbb{C}[z_{1-i}, w_{1-j}]$. We can obtain f_{ij} explicitly by substituting $x_i = y_j = 1$, $x_{1-i} = z_{1-i}$ and $y_{1-j} = w_{1-j}$.

Moreover Σ_L is trivialized by $\{1, D_z^{ij}, D_w^{ij}\}$, where $D_z^{ij} \mapsto \frac{\partial}{\partial z_{1-i}}$ and $D_w^{ij} \mapsto \frac{\partial}{\partial w_{1-j}}$ under the natural projection $\Sigma_L \rightarrow T_{\mathbb{P}^1 \times \mathbb{P}^1}$. And the morphism $dF: \Sigma_L \rightarrow L$ is given by $dF(1) = f_{ij} e_{ij} = F$,

$$(4.2) \quad dF(D_z^{ij}) = \frac{\partial f_{ij}}{\partial z_{1-i}} e_{ij} = \frac{\partial F}{\partial x_{1-i}} (x_i = 1, y_j = 1) x_i^e y_j^d = x_i \frac{\partial F}{\partial x_{1-i}}$$

and

$$dF(D_w^{ij}) = \frac{\partial f_{ij}}{\partial w_{1-j}} e_{ij} = \frac{\partial F}{\partial y_{1-j}} (x_i = 1, y_j = 1) x_i^e y_j^d = y_j \frac{\partial F}{\partial y_{1-j}}.$$

Since the result of (4.2) is independent of j , we conclude that $D_z^{i0} = D_z^{i1}$ on $U_{i0} \cap U_{i1}$, and analogously $D_w^{0j} = D_w^{1j}$ on $U_{0j} \cap U_{1j}$. We just denote then by D_z^i resp. D_w^j the local section of Σ_L given by D_z^{ij} on U_{ij} resp. D_w^{ij} on U_{ij} .

In order to compare D_z^0 and D_z^1 on $U_{0j} \cap U_{1j}$ we compare their images under dF for an arbitrary $F \in H^0(L)$, which are respectively $x_0 \frac{\partial F}{\partial x_1}$ and $x_1 \frac{\partial F}{\partial x_0}$. Using the Euler relation with respect to the variables x_0, x_1 we obtain

$$(4.3) \quad dF(x_1^2 D_z^0 + x_0^2 D_z^1) = x_1^2 x_0 \frac{\partial F}{\partial x_1} + x_0^2 x_1 \frac{\partial F}{\partial x_0} = x_0 x_1 \left(x_0 \frac{\partial F}{\partial x_0} + x_1 \frac{\partial F}{\partial x_1} \right) = d \cdot x_0 x_1 F = dF(d \cdot x_0 x_1 \cdot 1)$$

Since this should hold for any $F \in H^0(L)$, it must hold

$$(4.4) \quad x_1^2 D_z^0 + x_0^2 D_z^1 = d \cdot x_0 x_1 \cdot 1, \quad \text{i.e.} \quad D_z^1 = d \cdot \frac{x_1}{x_0} \cdot 1 - \left(\frac{x_1}{x_0} \right)^2 D_z^0 = d \cdot z_1 \cdot 1 - z_1^2 D_z^0.$$

Analogously we can prove

$$(4.5) \quad D_w^1 = d \cdot w_1 \cdot 1 - w_1^2 D_w^0.$$

We are now ready to compute a basis of the global sections of $\Sigma_L(a, b)$ for any $a, b \in \mathbb{Z}$.

Note first that the short exact sequence

$$(4.6) \quad 0 \longrightarrow \mathcal{O}_X(a, b) \longrightarrow \Sigma_L(a, b) \longrightarrow T_X(a, b) = \mathcal{O}_X(a+2, b) \oplus \mathcal{O}_X(a, b+2) \longrightarrow 0$$

gives $h^0(\Sigma_L(a, b)) = 0$ unless $a, b \geq -2$. Thus from now on we restrict ourselves to this case.

On U_{ij} , $\Sigma_L(a, b)$ is trivialized by $1 \cdot x_i^a y_j^b$, $D_z^i \otimes x_i^1 y_j^b$ and $D_w^j \otimes x_i^a y_j^b$, and the gluing relations are given by (4.4) and (4.5).

On U_{ij} a global section of $\Sigma_L(a, b)$ is thus described as

$$(A_{ij} \cdot 1 + B_{ij}D_z^i + C_{ij}D_w^j) \otimes x_i^a y_j^b$$

for certain polynomials $A_{ij}, B_{ij}, C_{ij} \in \mathbb{C}[z_{1-j}, w_{1-j}]$. We will now use the relations (4.4) and (4.5) to find relations between these polynomials, as well as upper bounds on their (bi)degrees.

Using (4.3) and $z_0 = \frac{x_0}{x_1} = \frac{1}{z_1}$ we obtain

$$(4.7) \quad A_{0j}(z_1, w_{1-j}) = \left[A_{1j} \left(\frac{1}{z_1}, w_{1-j} \right) + d \cdot z_1 B_{1j} \left(\frac{1}{z_1}, w_{1-j} \right) \right] z_1^a$$

$$(4.8) \quad B_{0j}(z_1, w_{1-j}) = -B_{1j} \left(\frac{1}{z_1}, w_{1-j} \right) z_1^{a+2}$$

$$(4.9) \quad C_{0j}(z_1, w_{1-j}) = C_{1j} \left(\frac{1}{z_1}, w_{1-j} \right) z_1^a$$

From (4.8) and (4.9) it follows $\deg_{z_0} B_{1j} \leq a + 2$ and $\deg_{z_0} C_{1j} \leq a$.

By symmetry, it must also hold

$$\begin{aligned} A_{i0}(z_{1-i}, w_1) &= \left[A_{i1} \left(z_{1-i}, \frac{1}{w_1} \right) + e \cdot w_1 C_{i1} \left(z_{1-i}, \frac{1}{w_1} \right) \right] w_1^b \\ B_{i0} \left(z_{1-i}, \frac{1}{w_1} \right) &= -B_{i1} \left(z_{1-i}, \frac{1}{w_1} \right) w_1^{b+2} \\ C_{i0} \left(z_{1-i}, \frac{1}{w_1} \right) &= C_{i1} \left(z_{1-i}, \frac{1}{w_1} \right) w_1^b \end{aligned}$$

hence in particular $\deg_{w_0} B_{1j} \leq b$ and $\deg_{w_0} C_{1j} \leq b + 2$.

We can try with monomials $B_{11} = z_0^r w_0^s$ and $C_{11} = z_0^n w_0^m$, where $0 \leq r \leq a + 2$, $0 \leq s \leq b$, $0 \leq n \leq a$ and $0 \leq m \leq b + 2$, and check the possibilities for A_{11} .

From (4.7) it follows that $A_{11} \left(\frac{1}{z_1}, w_0 \right) z_1^a + d \cdot z_1^{a+1-r} w_0^s$ must be a polynomial on z_1 and w_0 .

- If $r \leq a + 1$ it is enough that $\deg_{z_0} A_{11} \leq a$, thus we can take any monomial $A_{11}(z_0, w_0) = z_0^t w_0^u$ with $0 \leq t \leq a$.
- If $r = a + 2$, A_{11} must contain the monomial $-d \cdot z_0^{a+1} w_0^s$ in order to cancel the term $d \cdot z_1^{-1} w_0^s$ (and also any monomial as in the case $r \leq a + 1$).

By symmetry:

- If $m \leq b + 1$, A_{11} can contain any monomial $z_0^t w_0^u$ with $0 \leq u \leq b$,
- If $m = b + 2$, then A_{11} must contain $-e \cdot z_0^n w_0^{b+2}$.

Altogether we have the following basis for $H^0(\Sigma_L(a, b))$ (we give their local expressions on U_{11})

- (1) $z_0^t w_0^u \cdot 1 \otimes x_1^a y_1^b$ with $0 \leq t \leq a$ and $0 \leq u \leq b$,
- (2) $z_0^r w_0^s \cdot D_z^1 \otimes x_1^a y_1^b$ with $0 \leq r \leq a + 1$ and $0 \leq s \leq b$,
- (3) $(-d \cdot z_0^{a+1} w_0^s \cdot 1 + z_0^{a+2} w_0^s \cdot D_z^1) \otimes x_1^a y_1^b$ with $0 \leq s \leq b$
- (4) $z_0^n w_0^m \cdot D_w^1 \otimes x_1^a y_1^b$ with $0 \leq n \leq a$ and $0 \leq m \leq b + 1$, and

$$(5) \left(-e \cdot z_0^n w_0^{b+1} \cdot 1 + z_0^n w_0^{b+2} \cdot D_w^1 \right) \otimes x_1^a y_1^b \text{ with } 0 \leq n \leq a.$$

Note that if $a = -2$ or $b = -2$ there are no elements in the above list, hence also $H^0(\Sigma_L(a, b)) = 0$. This is compatible with the short exact sequence (4.6), since in this case we have $H^1(\mathcal{O}_X(a, b)) \neq 0$, and the connecting homomorphism turns out to be an isomorphism.

In order to finish the proof, we just need to compute a set of generators of

$$J_{a,b} = \text{im} \left(dF: H^0(\Sigma_L(a-d, b-e)) \rightarrow H^0(\mathcal{O}_X(a, b)) \right)$$

by applying the differential dF to the elements of the above basis (replacing a by $a-d$ and b by $b-d$). Since $H^0(\Sigma_L(a-d, b-e)) = 0$ unless $a-d, b-e \geq -1$, we can assume $a \geq d-1$ and $b \geq e-1$. Recall that $dF(1) = F$, $dF(D_z^1) = x_1 \frac{\partial F}{\partial x_0}$ and $dF(D_w^1) = y_1 \frac{\partial F}{\partial y_0}$, and thus

$$(1) dF \left(z_0^t w_0^u \cdot 1 \otimes x_1^{a-d} y_1^{b-e} \right) = \frac{x_0^t y_0^u}{x_1^t y_1^u} x_1^{a-d} y_1^{b-e} F = x_0^t x_1^{a-d-t} y_0^u y_1^{b-e-u} F \text{ with } 0 \leq t \leq a-d \text{ and } 0 \leq u \leq b-e, \text{ i.e. we obtain all possible multiples of } F \text{ with bidegree } (a, b);$$

$$(2) dF \left(z_0^r w_0^s \cdot D_z^1 \otimes x_1^{a-d} y_1^{b-e} \right) = x_0^r x_1^{a-d-r+1} y_0^s y_1^{b-e-s} \frac{\partial F}{\partial x_0} \text{ with } 0 \leq r \leq a-d+1 \text{ and } 0 \leq s \leq b-e, \text{ i.e. all multiples of } \frac{\partial F}{\partial x_0} \text{ of bidegree } (a, b);$$

$$(3) dF \left(\left(-d \cdot z_0^{a-d+1} w_0^s \cdot 1 + z_0^{a-d+2} w_0^s \cdot D_z^1 \right) \otimes x_1^{a-d} y_1^{b-e} \right) = -d \cdot x_0^{a-d+1} x_1^{-1} y_0^s y_1^{b-e-s} F + x_0^{a-d+2} x_1^{-2} y_0^s y_1^{b-e-s} x_1 \frac{\partial F}{\partial x_0} = x_0^{a-d+1} x_1^{-1} y_0^s y_1^{b-e-s} \left[-d \cdot F + x_0 \frac{\partial F}{\partial x_0} \right] = x_0^{a-d+1} x_1^{-1} y_0^s y_1^{b-e-s} \left[-x_1 \frac{\partial F}{\partial x_1} \right] = -x_0^{a-d+1} y_0^s y_1^{b-e-s} \frac{\partial F}{\partial x_1} \text{ with } 0 \leq s \leq b-e, \text{ i.e. all multiples of } x_0^{a-d+1} \frac{\partial F}{\partial x_1} \text{ of bidegree } (a, b). \text{ The remaining multiples of } \frac{\partial F}{\partial x_1} \text{ of the same bidegree, admitting powers of } x_1, \text{ can be obtained from (1) and (2) using the Euler relation with respect to } x_0, x_1.$$

$$(4) dF \left(z_0^n w_0^m \cdot D_w^1 \otimes x_1^{a-d} y_1^{b-e} \right) = x_0^n x_1^{a-d-n} y_0^m y_1^{b-e-m+1} \frac{\partial F}{\partial y_0} \text{ with } 0 \leq n \leq a-d \text{ and } 0 \leq m \leq b-e+1, \text{ and}$$

$$(5) dF \left(\left(-e \cdot z_0^n w_0^{b-e+1} \cdot 1 + z_0^n w_0^{b-e+2} \cdot D_w^1 \right) \otimes x_1^{a-d} y_1^{b-e} \right) = -x_0^n x_1^{a-d-n} y_1^{b-e+1} \frac{\partial F}{\partial y_1} \text{ with } 0 \leq n \leq a-d.$$

Thus $J_{a,b}$ contains all products of the partial derivatives of F and monomials (of the right bidegree), which proves the claim. \square

In order to find a contradiction with Theorem 3.8, we need to understand the geometry of the linear system on Z induced by $J_{d,e}$. More strictly speaking, since $F \in J_{d,e}$, the restriction map $J_{d,e} \subseteq H^0(L) \rightarrow H^0(L|_Z) = H^0(\mathcal{O}_Z(d, e))$ is not injective, but its image induces a linear subsystem of $|\mathcal{O}_Z(d, e)|$ that we will still denote $|J_{d,e}|_Z$. It will be helpful to consider also the restrictions of $J_{d-1,e}$ and $J_{d,e-1}$, which by degree reasons inject in $H^0(\mathcal{O}_Z(d-1, e))$ and $H^0(\mathcal{O}_Z(d, e-1))$ respectively.

Theorem 4.2. *Suppose $d, e \geq 2$. Then:*

- (1) *The base divisor of $|J_{d-1,e}|_Z$ coincides with the ramification divisor of the second projection $Z \hookrightarrow X = \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{\pi_2} \mathbb{P}^1$, and after removing the base points it induces the projection of Z onto the first factor.*

(2) The restricted linear system $|J_{d,e}|_Z$ is base point free and the induced morphism $Z \rightarrow |J_{d,e}|_Z^\vee$ is birational into its image.

Proof. (1) Since $\left\{\frac{\partial F}{\partial x_0}, \frac{\partial F}{\partial x_1}\right\}$ is a basis of $J_{d-1,e}$, the base locus of $|J_{d-1,e}|_Z$ consists of the points $p \in X = \mathbb{P}^1 \times \mathbb{P}^1$ such that

$$\frac{\partial F}{\partial x_0}(p) = \frac{\partial F}{\partial x_1}(p) = F(p) = 0,$$

where the third condition $F(p) = 0$ actually follows from the other two because of the Euler identity with respect to x_0, x_1 .

Suppose without loss of generality that $p = ([1 : 0], [1 : 0])$ is a base point. Taking local affine coordinates $(z, w) = \left(\frac{x_1}{x_0}, \frac{y_1}{y_0}\right)$, p corresponds to the origin $(0, 0)$, and the curve Z is locally defined by the equation $f(z, w) := F(1, z, 1, w) = 0$. Being a base point of $J_{d-1,e}$ implies in particular that $\frac{\partial f}{\partial z}(p) = \frac{\partial f}{\partial w}(p) = 0$. Since Z is smooth, it must hold $\frac{\partial f}{\partial w}(p) \neq 0$, and thus the tangent line to Z at p has equation

$$0 = \frac{\partial f}{\partial z}(p)z + \frac{\partial f}{\partial w}(p)w = \frac{\partial f}{\partial w}(p)w$$

or equivalently $w = 0$, which shows that p is a ramification point of the projection onto the second factor (locally given by $(z, w) \mapsto w$).

One can even check that the multiplicity as a base point agrees with the multiplicity in the ramification divisor. Indeed note first that on Z we have $x_0 \frac{\partial F}{\partial x_0} + x_1 \frac{\partial F}{\partial x_1} \equiv 0$, i.e.

$$\frac{\partial F}{\partial x_0|_Z} = -\frac{x_1}{x_0} \frac{\partial F}{\partial x_1|_Z} = -z \frac{\partial F}{\partial x_1|_Z}.$$

Hence the multiplicity of p as a base point of $|J_{d-1,e}|_Z$ coincides with the vanishing order of $\frac{\partial F}{\partial x_1|_Z}$ at p . Using z as a local coordinate for Z around p and expressing $w|_Z$ as a function $w(z)$ we can write (locally around p)

$$\frac{\partial F}{\partial x_1|_Z} = \frac{\partial F}{\partial x_1}(1, z, 1, w(z)) = \frac{\partial f}{\partial z}(z, w(z)).$$

From $f(z, w(z)) \equiv 0$ it follows

$$\frac{\partial f}{\partial z}(z, w(z)) = -\frac{\partial f}{\partial w}(z, w(z))w'(z).$$

Since $\frac{\partial f}{\partial w}(p) \neq 0$, we have

$$\text{ord}_{z=0} \frac{\partial f}{\partial z}(z, w(z)) = \text{ord}_{z=0} w'(z).$$

By the above discussion, the left hand side is the multiplicity of p in the base locus of $|J_{d-1,e}|_Z$, while the right hand side is the order of p in the ramification divisor of the projection onto the second coordinate, as claimed.

The second assertion follows easily from the Euler relation with respect to x_0, x_1 : outside the base locus, the morphism induced by $J_{d-1,e}|_Z$ is given given for example by

$$p \mapsto \left[\frac{\partial F}{\partial x_1}(p) : -\frac{\partial F}{\partial x_0}(p) \right] = [x_0(p) : x_1(p)].$$

(2) The restriction $J_{d,e|Z}$ is generated by the restrictions of $x_i \frac{\partial F}{\partial x_j}$ and $y_i \frac{\partial F}{\partial y_j}$ (Lemma 4.1).

Suppose $p \in Z$ is a base point, so that $x_i(p) \frac{\partial F}{\partial x_j}(p) = y_i(p) \frac{\partial F}{\partial y_j}(p) = 0$ for all i, j .

It holds $x_i(p) \neq 0$ for at least one i , so that $\frac{\partial F}{\partial x_j}(p) = 0$ for both $j = 0, 1$. This implies that p is a ramification point for the second projection $Z \rightarrow \mathbb{P}^1$. Considering the y_i 's one can analogously show that p is also a ramification point for the first projection. This is only possible if Z is not smooth at p , which gives a contradiction.

It remains to show that $J_{d,e|Z}$ induces a birational morphism. Considering the above set of 8 generators, we can describe the induced morphism φ as

$$p \mapsto [\dots : a_{ij} : \dots : b_{ij} : \dots] := \left[\dots : x_i(p) \frac{\partial F}{\partial x_j}(p) : \dots : y_i(p) \frac{\partial F}{\partial y_j}(p) : \dots \right] \in \mathbb{P}^7.$$

Note that, since the generators are not linearly independent (because of the Euler relations (4.1)), hence the image will be contained in the intersection of hyperplanes

$$\left\{ \frac{1}{d}(a_{00} + a_{11}) = \frac{1}{e}(b_{00} + b_{11}) = 0 \right\}.$$

Let's consider now the linear subsystem generated by the $x_i \frac{\partial F}{\partial x_j}$, whose induced rational map $\psi_x: Z \dashrightarrow \mathbb{P}^3$ coincides with the composition of $\varphi: Z \rightarrow \mathbb{P}^7$ with the projection onto the first coordinates $[a_{00} : \dots : a_{11}]$. Furthermore ψ_x can be factorized as

$$\psi_x: Z \dashrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{\Phi} \mathbb{P}^3,$$

where Φ just the Segre embedding, and the first map is given by

$$p \mapsto \left([x_0(p) : x_1(p)], \left[\frac{\partial F}{\partial x_0}(p) : \frac{\partial F}{\partial x_1}(p) \right] \right),$$

i.e. the first map is the product of the first projection and the morphism induced by $J_{d-1,e|Z}$. By the first part of the Theorem, the second factor coincides (up to an automorphism of \mathbb{P}^1) with the first projection.

Thus, with the appropriate choice of coordinates and denoting by Δ the diagonal map, we have

$$\psi_x: Z \xrightarrow{\pi_1} \mathbb{P}^1 \xrightarrow{\Delta} \mathbb{P}^1 \times \mathbb{P}^1 \xrightarrow{\Phi} \mathbb{P}^3$$

i.e. from the projection of φ onto the first four coordinates we recover the first projection of $Z \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$.

Analogously we can recover the second projection π_2 from the composition of φ with the projection onto the last four coordinates $[b_{00} : \dots : b_{11}]$.

Altogether we can recover the original embedding $Z \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^1$ from the morphism φ induced by linear system $|J_{d,e|Z}|$, which implies that φ must be birational. □

We also need to study up to what extent the duality maps $\lambda_{K_X^2 \otimes L^2}$ and $\lambda_{K_X \otimes L^2}$ fail to be isomorphisms.

Lemma 4.3. *Let $L = \mathcal{O}_X(d, e)$. Then the following hold:*

- (1) If $d, e \geq 3$, the duality map $\lambda_{K_X^2 \otimes L^2}: R_{K_X^2 \otimes L^2} = R_{2d-4, 2e-4} \rightarrow R_L^* = R_{d,e}^*$ is an isomorphism. Thus in particular $\ker \lambda = J_{K_X^2 \otimes L^2} = J_{2d-4, 2e-4}$, where λ is as in (3.12).
- (2) If $d = 2$ and $e \geq 2$, then $\lambda_{K_X^2 \otimes L^2}$ is injective.
- (3) If $d, e \geq e$, the duality map $\lambda_{K_X \otimes L^2}: R_{K_X \otimes L^2} = R_{2d-2, 2e-2} \rightarrow R_{K_X \otimes L} = R_{d-2, e-2}^*$ is surjective and has one-dimensional kernel. More precisely, there is a short exact sequence

$$0 \longrightarrow H^0(\mathcal{O}_X) \xrightarrow{\cdot c_1(L)} H^1(\Omega_X^1) \longrightarrow R_{K_X \otimes L^2} \xrightarrow{\lambda_{K_X \otimes L^2}} R_{K_X \otimes L}^* \longrightarrow 0.$$

Proof. (1) The assertion follows from Proposition 2.17 and $H^1(\Sigma_L) = 0$, which in turn follows from the sequence $0 \rightarrow \mathcal{O}_X \rightarrow \Sigma_L \rightarrow T_X \rightarrow 0$.

- (2) In this case we cannot apply Proposition 2.17 because $H^1(\Sigma_L \otimes K_X^2 \otimes L) = H^1(\Sigma_L(-2, e-4)) \neq 0$. Nevertheless we can still try to follow that proof and obtain on the one hand

$$(4.10) \quad R_{K_X^2 \otimes L^2} \cong \ker(H^1(\mathcal{E}_F \otimes K_X^2 \otimes L) \rightarrow H^1(\Sigma_L \otimes K_X^2 \otimes L)).$$

On the other hand for these degrees it still holds $H^1(\Sigma_L) = 0$, which gives

$$(4.11) \quad R_L \cong \ker(H^1(\mathcal{E}_F) \rightarrow H^1(\Sigma_L)) = H^1(\mathcal{E}_F),$$

and thus $R_L^* \cong H^1(\mathcal{E}_F^\vee \otimes K_X) = H^1(\mathcal{E}_F \otimes K_X^2 \otimes L)$. Since both isomorphisms in (4.10) and (4.11) are given by product with the class $\delta F \in \text{Ext}_{\mathcal{O}_X}^1(L, \mathcal{E}_F)$, it turns out that $\lambda_{K_X^2 \otimes L^2}$ is simply the inclusion of the kernel of $H^1(\mathcal{E}_F \otimes K_X^2 \otimes L) \rightarrow H^1(\Sigma_L \otimes K_X^2 \otimes L)$ into $H^1(\mathcal{E}_F \otimes K_X^2 \otimes L)$.

- (3) The assertion follows directly from Proposition 2.16 and the fact that

$$H^1(\Omega_X^1) \cong H^1(\mathcal{O}_X(-2, 0)) \oplus H^1(\mathcal{O}_X(0, -2)) \cong \mathbb{C}^2.$$

□

We are now almost ready to prove the announced upper bound for the dimension of $D \subseteq \mathbb{P}(J_{2d-4, 2e-4})$. To this aim we adapt some ideas from [FP22] to construct a sufficiently high-dimensional linear subspace in $\mathbb{P}(J_{2d-4, 2e-4})$ disjoint from D .

For a fixed $G \in J_L \subseteq S_{d,e}$ with $d, e \geq 3$ consider the following map:

$$\begin{aligned} \mu_G: S_{d-4, e-4} \oplus S_{d-4, e-4} &\longrightarrow S_{2d-4, 2e-4} \\ (A, B) &\longmapsto AF + BG. \end{aligned}$$

Note that, since both $F, G \in J_L$, it holds $\text{im } \mu \subseteq J_{K_X^2 \otimes L^2} = J_{2d-4, 2e-4}$ (for this equality we need $d, e \geq 3$).

Theorem 4.4. *Suppose $d, e \geq 3$. For a general $G \in J_L = J_{d,e}$ (in particular, $G \notin \mathbb{C}F$) it holds:*

- (1) $\ker \mu_G = 0$, and
(2) $D \cap \mathbb{P}(\text{im } \mu_G) = \emptyset$.

Proof. (1) Suppose $\mu_G(A, B) = AF + BG = 0 \in S_{2d-4, 2e-4} = H^0(K_X^2 \otimes L^2)$. Then the restriction to $Z = \{F = 0\}$ is also

$$B|_Z G|_Z \equiv 0 \in H^0\left(\left(K_X^2 \otimes L^2\right)|_Z\right).$$

Since $G|_Z \neq 0$ (because G is not a multiple of F), it must hold $B|_Z \equiv 0$, i.e. B is a multiple of F . But since $B \in S_{d-4, e-4}$ has smaller bidegree than F , it must be $B = 0$. Thus also $FA = 0 \in S_{2d-4, 2e-4}$, and thus $A = 0$ because the map $S_{d-4, e-4} \xrightarrow{\cdot F} S_{2d-4, 2e-4}$ is injective.

(2) Consider the set

$$I = \left\{ (A, B, \alpha, \beta, G) \left| \begin{array}{l} A, B \in S_{d-4, e-4}, \alpha, \beta \in H^0(K_X \otimes L) \setminus \{0\} = S_{d-2, e-2} \setminus \{0\} \\ G \in J_{d, e} \setminus \mathbb{C}F, AF + BG = \alpha\beta \end{array} \right. \right\}$$

which is a quasi-projective subvariety of $S_{d-4, e-4} \times S_{d-4, e-4} \times S_{d-2, e-2} \times S_{d-2, e-2} \times J_{d, e}$.

By definition of D , it holds

$$D \cap \mathbb{P}(\text{im } \mu_G) \neq \emptyset$$

if and only if $G \in \pi(I)$, where $\pi: I \rightarrow J_{d, e} \setminus \mathbb{C}F$ denotes the projection $(A, B, \alpha, \beta, G) \mapsto G$.

We want thus to show that π is not dominant, for which we can replace I by its (open, dense) smooth subset. Assume from now on that π is dominant.

By Theorem 4.2, the restriction of the Jacobian $J_{d, e}$ to Z induces a birational map onto a non-degenerate projective curve $C \subseteq \mathbb{P}^r$ for certain $r \leq \dim J_{d, e} - 2$. The pull-backs of the hyperplane sections of \mathbb{P}^r are by construction the vanishing divisors $(G|_Z)$. Let U denote the subset $\{G \in J_{d, e} \setminus \mathbb{C}F \mid (G|_Z) \text{ is reduced}\}$. Given a fixed $G \in U$ with $(G|_C) = p_1 + p_2 + \dots + p_n$, the fundamental group $\pi_1(U, G)$ induces a monodromy action on $\{p_1, \dots, p_n\}$, i.e. there is a group homomorphism

$$(4.14) \quad \rho: \pi_1(U, G) \rightarrow \text{Sym}(\{p_1, \dots, p_n\}).$$

By the Uniform Position Theorem [ACGH85, Lemma in p. 111] the monodromy group is the full symmetric group, i.e. the group homomorphism (4.14) is surjective.

Replacing U by a non-empty Zariski-open subset the fundamental group does not change, hence we can assume that U is contained in $\pi(I)$ (recall that we are assuming that π is dominant). Replacing I by $\pi^{-1}(U)$ we can assume $\pi: I \rightarrow U$ is a surjective morphism of smooth (quasi-projective) varieties.

Fix $G \in U$ and set $(G|_Z) = p_1 + \dots + p_n$, where $n = \deg \mathcal{O}_Z(d, e) = 2de > 2de - 2d - 2e = 2g(Z) - 2$. By the definition of U , there are A, B, α, β such that $AF + BG = \alpha\beta$. The restriction to $Z = \{F = 0\}$ gives then

$$B|_Z G|_Z = \alpha|_Z \beta|_Z,$$

so $p_1 + \dots + p_n \leq (\alpha|_Z) + (\beta|_Z)$. Up to reordering the indices of the points and switching α and β we can assume $\alpha|_Z(p_i) = 0$ for $i = 1, \dots, m \geq \frac{n}{2}$, and $\alpha|_Z(p_i) \neq 0$ for $i = m + 1, \dots, n$. Moreover since $\alpha|_Z \in H^0(K_Z)$ and thus $\deg(\alpha|_Z) = 2g(Z) - 2 < n$, it holds $m < n$.

Now for every $i = 1, \dots, m$ there is a closed path γ_i on U based at G whose monodromy action permutes p_i and p_{m+1} and leaves all other points fixed. We can lift this path to an open path on I , starting on (A, B, α, β, G) , i.e. we can find a continuous map

$$\tilde{\gamma}_i = \left(\tilde{A}, \tilde{B}, \tilde{\alpha}, \tilde{\beta}, \tilde{G} \right) : [0, 1] \rightarrow I$$

with $\tilde{\gamma}_i(0) = (A, B, \alpha, \beta, G)$ and $\tilde{A}(t)F + \tilde{B}(t)\tilde{G}(t) = \tilde{\alpha}(t)\tilde{\beta}(t)$ for every $t \in [0, 1]$.

Set $\alpha_i := \tilde{\alpha}(1)$, $\beta_i := \tilde{\beta}(1)$, $A_i := \tilde{A}(1)$ and $B_i := \tilde{B}(1)$, which satisfy $A_iF + B_iG = \alpha_i\beta_i$. Thus restricting to Z we obtain

$$p_1 + \dots + p_n = (G|_Z) \leq (B_iG|_Z) = (\alpha_i|_Z) + (\beta_i|_Z).$$

Moreover it holds $\alpha_{i|Z}(p_{m+1}) = \alpha_{i|Z}(p_i) = 0$ and $\alpha_{i|Z}(p_j) = \alpha_{i|Z}(p_j) = 0$ for every $j = 1, \dots, m$, $j \neq i$, while $\alpha_{i|Z}(p_i) = \alpha_{i|Z}(p_{m+1}) \neq 0$.

In this way we obtain $\alpha_0 = \alpha, \alpha_1, \dots, \alpha_m$, whose restrictions to Z are linearly independent holomorphic 1-forms, and hence $m + 1 \leq g(Z)$. But this is a contradiction, since

$$n \leq 2m \leq 2g(Z) - 2 < n.$$

□

As announced, we can use the previous theorem to find an upper bound on $\dim D$.

Corollary 4.5. *If $d, e \geq 3$ it holds*

$$\dim D \leq de - d - e - 4.$$

Proof. Let G be general, so that $D \cap \mathbb{P}(\operatorname{im} \mu_G) = \emptyset \subseteq \mathbb{P}(J_{2d-4, 2e-4})$. Then it must hold $\dim D + \dim \mathbb{P}(\operatorname{im} \mu_G) < \dim \mathbb{P}(J_{2d-4, 2e-4})$, and hence

$$\begin{aligned} \dim D &< \dim J_{2d-4, 2e-4} - \dim \operatorname{im} \mu_G = (\dim S_{2d-4, 2e-4} - \dim R_{2d-4, 2e-4}) - 2 \dim S_{d-4, e-4} = \\ &\stackrel{(*)}{=} (2d-3)(2e-3) - \dim R_{d,e} - 2(d-3)(e-3) = 2de - 9 - (\dim S_{d,e} - \dim J_{d,e}) = \\ &= 2de - 9 - (d+1)(e+1) + \dim J_{d,e} = de - d - e - 10 + \dim J_{d,e} \leq \\ &\leq de - d - e - 10 + h^0(\Sigma_L) = de - d - e - 3. \end{aligned}$$

Note that in $(*)$ we have used $\dim R_{2d-4, 2e-4} = \dim R_{d,e}$, i.e. $\dim R_{K_X^2 \otimes L^2} = \dim R_L$, which follows from $d, e \geq 3$ and Lemma 4.3. □

We are now ready to prove the final result of the paper.

Theorem 4.6. *For any $d, e \geq 1$, $L = \mathcal{O}_X(d, e)$ has maximal infinitesimal variation at every smooth $Z \in |L|$.*

Proof. Let $Z \in |L|$ be a smooth curve defined by a polynomial $F \in H^0(L) = S_{d,e}$. Let's consider first the cases of smaller degrees.

If $d = 1$ or $e = 1$, Z is a rational curve, hence the Hodge structure is trivial and there is nothing to prove.

Suppose now that $d = 2$ and $e \geq 2$. Since $H^0(K_X) = 0$, it holds $R_{K_X \otimes L} = H^0(K_X \otimes L) = S_{0,e-2}$. Moreover it holds $J_{K_X^2 \otimes L^2} = J_{0,2e-4} = 0$ because the Jacobian ideal is generated in degrees $(d-1, e) = (1, e)$ and $(d, e-1) = (2, e-1)$. This means that $R_{K_X^2 \otimes L^2} = H^0(K_X^2 \otimes L^2) = S_{0,2e-4}$.

Suppose that $\alpha \in \ker \varphi_{K_X \otimes L}(\tau)$ for a general $\tau \in H^0(L) = S_{2,e}$, so that $\lambda_{K_X^2 \otimes L^2}(\alpha^2) = 0 \in R_L^*$ by Lemma 3.5. But the injectivity of $\lambda_{K_X^2 \otimes L^2}$ (Lemma 4.3) implies $\alpha^2 = 0 \in S_{0,2e-4}$, and thus $\alpha = 0$.

We can now consider the case $d, e \geq 3$. If L does not have maximal infinitesimal variation at Z , then Theorem 3.8 implies

$$\dim D \geq h^0(K_X \otimes L) + p_g(X) - 1 = h^0(\mathcal{O}_X(d-2, e-2)) - 1 = (d-1)(e-1) - 1 = de - d - e,$$

which contradicts Corollary 4.5. □

Remark 4.7. *Note that a very important point in the proof of Theorem 4.4 is that the generalized Jacobian ideal in degree precisely L (i.e. the image of $\widetilde{d\sigma}: H^0(T_X) \rightarrow H^0(L|_Z)$) induces a birational embedding of Z in a projective space. This property might also hold for more general surfaces, giving some hope that this approach might work in a few more cases.*

REFERENCES

- [ACGH85] Enrico Arbarello, Maurizio Cornalba, Philip Griffiths, and Joe Harris. *Geometry of algebraic curves. Vol. I*, volume 267 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1985.
- [CGAT23] Giulio Codogni, Víctor González-Alonso, and Sara Torelli. Rigidity of modular morphisms via Fujita decomposition, 2023.
- [DH22] Yajnaseni Dutta and Daniel Huybrechts. Maximal variation of curves on $K3$ surfaces. *Tunis. J. Math.*, 4(3):443–464, 2022.
- [FP22] Filippo Francesco Favale and Gian Pietro Pirola. Infinitesimal variation functions for families of smooth varieties. *Milan J. Math.*, 90(1):209–228, 2022.
- [GAT21] Víctor González-Alonso and Sara Torelli. Families of curves with Higgs field of arbitrarily large kernel. *Bull. Lond. Math. Soc.*, 53(2):493–506, 2021.
- [GK06] Martine Girard and David R. Kohel. Classification of genus 3 curves in special strata of the moduli space. In *Algorithmic number theory*, volume 4076 of *Lecture Notes in Comput. Sci.*, pages 346–360. Springer, Berlin, 2006.
- [Gre85] Mark L. Green. The period map for hypersurface sections of high degree of an arbitrary variety. *Compos. Math.*, 55:135–156, 1985.
- [Laz86] Robert Lazarsfeld. Brill-Noether-Petri without degenerations. *J. Differ. Geom.*, 23:299–307, 1986.
- [LP16] Yongnam Lee and Gian Pietro Pirola. On rational maps from the product of two general curves. *Ann. Sc. Norm. Super. Pisa, Cl. Sci. (5)*, 16(4):1139–1152, 2016.
- [Muk96] Shigeru Mukai. Curves and $K3$ surfaces of genus eleven. In *Moduli of vector bundles. Papers of the 35th Taniguchi symposium, Sanda, Japan and a symposium held in Kyoto, Japan, 1994*, pages 189–197. New York, NY: Marcel Dekker, 1996.
- [Ser06] Edoardo Sernesi. *Deformations of algebraic schemes*, volume 334 of *Grundlehren Math. Wiss.*. Berlin: Springer, 2006.
- [Vie83] Eckart Viehweg. Weak positivity and the additivity of the Kodaira dimension for certain fibre spaces. Algebraic varieties and analytic varieties, Proc. Symp., Tokyo 1981, Adv. Stud. Pure Math. 1, 329–353 (1983), 1983.
- [Voi02] Claire Voisin. Green’s generic syzygy conjecture for curves of even genus lying on a $K3$ surface. *J. Eur. Math. Soc. (JEMS)*, 4(4):363–404, 2002.

- [Voi05] Claire Voisin. Green's canonical syzygy conjecture for generic curves of odd genus. *Compos. Math.*, 141(5):1163–1190, 2005.