

Common neighborhood energies and their relations with Zagreb index

Firdous Ee Jannat^a, Rajat Kanti Nath^a, Kinkar Chandra Das^{b,1}

^a*Department of Mathematical Science, Tezpur University,
Napaam -784028, Sonitpur, Assam, India*

E-mail: firdusej@gmail.com, rajatkantinath@yahoo.com

^b*Department of Mathematics, Sungkyunkwan University,
Suwon 16419, Republic of Korea*

E-mail: kinkardas2003@gmail.com

Abstract

In this paper we establish connections between common neighborhood Laplacian and common neighborhood signless Laplacian energies and the first Zagreb index of a graph \mathcal{G} . We introduce the concepts of CNL-hyperenergetic and CNSL-hyperenergetic graphs and showed that \mathcal{G} is neither CNL-hyperenergetic nor CNSL-hyperenergetic if \mathcal{G} is a complete bipartite graph. We obtain certain relations between various energies of a graph. Finally, we conclude the paper with several bounds for common neighborhood Laplacian and signless Laplacian energies of a graph.

MSC: 05C50.

Keywords: Common Neighborhood; Laplacian; Signless Laplacian; Spectrum; Energy; Zagreb index.

1 Introduction

Let \mathcal{G} be a finite simple graph with adjacency matrix $A(\mathcal{G})$ and degree matrix $D(\mathcal{G})$. The spectrum of \mathcal{G} is the set of all the eigenvalues of $A(\mathcal{G})$ with multiplicity. Let $L(\mathcal{G}) := D(\mathcal{G}) - A(\mathcal{G})$ and $Q(\mathcal{G}) := D(\mathcal{G}) + A(\mathcal{G})$ be the Laplacian and signless Laplacian matrices of \mathcal{G} , respectively. More than thousands of research papers are studied on the adjacency, Laplacian and signless Laplacian eigenvalues of graphs, especially the most recent works [2, 23, 26] and the references therein. Then the Laplacian spectrum and the signless Laplacian spectrum of \mathcal{G} are the sets of all the eigenvalues (with multiplicity) of $L(\mathcal{G})$

¹Corresponding author

and $Q(\mathcal{G})$, respectively. Corresponding to these spectra, energy ($E(\mathcal{G})$), Laplacian energy ($LE(\mathcal{G})$) and signless Laplacian energy ($LE^+(\mathcal{G})$) of \mathcal{G} are defined as

$$E(\mathcal{G}) = \sum_{\alpha \in \text{Spec}(\mathcal{G})} |\alpha|, \quad LE(\mathcal{G}) = \sum_{\beta \in \text{L-spec}(\mathcal{G})} \left| \beta - \frac{\text{tr}(D(\mathcal{G}))}{|V(\mathcal{G})|} \right|$$

and

$$LE^+(\mathcal{G}) = \sum_{\gamma \in \text{Q-spec}(\mathcal{G})} \left| \gamma - \frac{\text{tr}(D(\mathcal{G}))}{|V(\mathcal{G})|} \right|,$$

where $\text{Spec}(\mathcal{G})$, $\text{L-spec}(\mathcal{G})$ and $\text{Q-spec}(\mathcal{G})$ are the spectrum, Laplacian spectrum and signless Laplacian spectrum of \mathcal{G} ; $\text{tr}(D(\mathcal{G}))$ is the trace of $D(\mathcal{G})$ and $V(\mathcal{G})$ is the set of vertices of \mathcal{G} .

In 1978, Gutman [17] has introduced the notion of $E(\mathcal{G})$, which has been studied extensively by many mathematicians over the years (see [20] and the references therein). In 2006, Gutman and Zhou [22] have introduced the notion of $LE(\mathcal{G})$; and in 2008, Gutman et al. [19] have introduced the notion of $LE^+(\mathcal{G})$.

Let $N_{\mathcal{G}}(v_i)$ be the neighborhood set of vertex v_i in \mathcal{G} . Also let $d_{\mathcal{G}}(v_i)$ be the degree of the vertex v_i in \mathcal{G} , that is, $d_{\mathcal{G}}(v_i) = |N_{\mathcal{G}}(v_i)|$. Let $m_{\mathcal{G}}(v_i)$ be the average degree of the adjacent vertices of vertex v_i in \mathcal{G} . If v_i is an isolated vertex in \mathcal{G} , then we assume that $m_{\mathcal{G}}(v_i) = 0$. Hence $d_{\mathcal{G}}(v_i) m_{\mathcal{G}}(v_i) = \sum_{v_j: v_i v_j \in E(\mathcal{G})} d_{\mathcal{G}}(v_j)$. The common neighborhood of two vertices v_i and v_j is the set $N_{\mathcal{G}}(v_i) \cap N_{\mathcal{G}}(v_j) = N(v_i, v_j)$, (say), containing all the vertices other than v_i and v_j that are adjacent to both v_i and v_j . The common neighborhood matrix $\text{CN}(\mathcal{G})$ of \mathcal{G} is given by

$$\text{CN}(\mathcal{G})_{i,j} = \begin{cases} |N(v_i, v_j)| & \text{if } i \neq j, \\ 0 & \text{if } i = j, \end{cases}$$

where $\text{CN}(\mathcal{G})_{i,j}$ is the (i, j) -th entry of $\text{CN}(\mathcal{G})$. The common neighborhood spectrum (also known as CN-spectrum) of \mathcal{G} , denoted by $\text{CN-spec}(\mathcal{G})$, is the set of all eigenvalues of $\text{CN}(\mathcal{G})$ with multiplicity. We write $\text{CN-spec}(\mathcal{G}) = \{\mu_1^{a_1}, \mu_2^{a_2}, \dots, \mu_k^{a_k}\}$, where the exponents a_i of μ_i are the multiplicities of μ_i for $i = 1, 2, \dots, k$; and $\mu_1, \mu_2, \dots, \mu_k$ are the distinct eigenvalues of $\text{CN}(\mathcal{G})$. The common neighborhood energy (also known as CN-energy) of

\mathcal{G} , denoted by $E_{CN}(\mathcal{G})$, is defined as

$$E_{CN}(\mathcal{G}) = \sum_{\mu \in \text{CN-spec}(\mathcal{G})} |\mu|.$$

The concepts of CN-spectrum and CN-energy are relatively new and not much explored. A graph \mathcal{G} is called CN-hyperenergetic if $E_{CN}(\mathcal{G}) > E_{CN}(K_{|V(\mathcal{G})|})$, where $K_{|V(\mathcal{G})|}$ is the complete graph on $|V(\mathcal{G})|$ -vertices. The concepts of CN-energy and CN-hyperenergetic graphs were introduced by Alwardi et al. [1] in 2011. It is worth mentioning that the concept of hyperenergetic graph was introduced by Walikar et al. [28] and Gutman [18], independently in 1999. Recall that \mathcal{G} is called hyperenergetic if $E(\mathcal{G}) > E(K_{|V(\mathcal{G})|})$. L-hyperenergetic and Q-hyperenergetic graphs were defined in a similar way and these were introduced in [15]. Very recently, certain mathematical properties on $\text{CN}(\mathcal{G})$ and $E_{CN}(\mathcal{G})$ of \mathcal{G} are studied, see [25] and the references cited therein.

In this paper we introduce the concepts of common neighborhood Laplacian spectrum, common neighborhood signless Laplacian spectrum and their corresponding energies of a graph \mathcal{G} . In Section 2, we establish relations between these energies and the first Zagreb index of a graph. In Section 3, we introduce the concepts of CNL-hyperenergetic and CNSL-hyperenergetic graphs and show that \mathcal{G} is neither CNL-hyperenergetic nor CNSL-hyperenergetic if \mathcal{G} is a complete bipartite graph. In Section 4, we obtain certain relations between various energies of a graph. Finally, in Section 5, we conclude this paper with several bounds for common neighborhood Laplacian and signless Laplacian energies of a graph.

2 Definition and connection with Zagreb index

First we observe that the (i, j) -th entry of $D(\mathcal{G})$ is given by

$$D(\mathcal{G})_{i,j} = \begin{cases} \sum_{k=1}^{|V(\mathcal{G})|} A(\mathcal{G})_{i,k} & \text{if } i = j \text{ and } i = 1, 2, \dots, |V(\mathcal{G})|, \\ 0 & \text{if } i \neq j, \end{cases}$$

where $A(\mathcal{G})_{i,k}$ is the (i, k) -th entry of $A(\mathcal{G})$. Thus $D(\mathcal{G})$ is a diagonal matrix whose diagonal entries are the corresponding row sums of the adjacency matrix of \mathcal{G} . Similarly,

we define common neighborhood row sum matrix (abbreviated as CNRS-matrix) of \mathcal{G} as given below:

$$\text{CNRS}(\mathcal{G})_{i,j} = \begin{cases} \sum_{k=1}^{|\mathcal{V}(\mathcal{G})|} \text{CN}(\mathcal{G})_{i,k} & \text{if } i = j \text{ and } i = 1, 2, \dots, |\mathcal{V}(\mathcal{G})|, \\ 0 & \text{if } i \neq j, \end{cases}$$

where $\text{CNRS}(\mathcal{G})$ is the CNRS-matrix of \mathcal{G} and $\text{CNRS}(\mathcal{G})_{i,j}$ is the (i, j) -th entry of $\text{CNRS}(\mathcal{G})$. The common neighborhood Laplacian matrix and the common neighborhood signless Laplacian matrix (abbreviated as CNL-matrix and CNSL-matrix) of \mathcal{G} , denoted by $\text{CNL}(\mathcal{G})$ and $\text{CNSL}(\mathcal{G})$, respectively, are defined as

$$\text{CNL}(\mathcal{G}) := \text{CNRS}(\mathcal{G}) - \text{CN}(\mathcal{G}) \text{ and } \text{CNSL}(\mathcal{G}) := \text{CNRS}(\mathcal{G}) + \text{CN}(\mathcal{G}).$$

Note that the matrices $\text{CNL}(\mathcal{G})$ and $\text{CNSL}(\mathcal{G})$ are symmetric and positive semidefinite. The set of eigenvalues of $\text{CNL}(\mathcal{G})$ and $\text{CNSL}(\mathcal{G})$ with multiplicities are called common neighborhood Laplacian spectrum and common neighborhood signless Laplacian spectrum (abbreviated as CN-Laplacian spectrum and CN-signless Laplacian spectrum) of \mathcal{G} , respectively. We write $\text{CNL-spec}(\mathcal{G})$ and $\text{CNSL-spec}(\mathcal{G})$ to denote CN-Laplacian spectrum and CN-signless Laplacian spectrum of \mathcal{G} , respectively. By writing $\text{CNL-spec}(\mathcal{G}) = \{\nu_1^{b_1}, \nu_2^{b_2}, \dots, \nu_\ell^{b_\ell}\}$ and $\text{CNSL-spec}(\mathcal{G}) = \{\sigma_1^{c_1}, \sigma_2^{c_2}, \dots, \sigma_m^{c_m}\}$ we mean that $\nu_1, \nu_2, \dots, \nu_\ell$ are the distinct eigenvalues of $\text{CNL}(\mathcal{G})$ with corresponding multiplicities b_1, b_2, \dots, b_ℓ and $\sigma_1, \sigma_2, \dots, \sigma_m$ are the distinct eigenvalues of $\text{CNSL}(\mathcal{G})$ with corresponding multiplicities c_1, c_2, \dots, c_m .

Corresponding to CN-Laplacian spectrum and CN-signless Laplacian spectrum of \mathcal{G} we define common neighborhood Laplacian energy and common neighborhood signless Laplacian energy (abbreviated as CNL-energy and CNSL-energy) of \mathcal{G} . The CNL-energy and CNSL-energy of \mathcal{G} , denoted by $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$, are as defined below:

$$LE_{CN}(\mathcal{G}) := \sum_{\nu \in \text{CNL-spec}(\mathcal{G})} \left| \nu - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|\mathcal{V}(\mathcal{G})|} \right| \quad (1)$$

and

$$LE_{CN}^+(\mathcal{G}) := \sum_{\sigma \in \text{CNSL-spec}(\mathcal{G})} \left| \sigma - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|\mathcal{V}(\mathcal{G})|} \right|. \quad (2)$$

This appends two new entries in the list of energies prepared by Gutman and Furtula [21]. Following the concepts of various hyperenergetic graphs [1, 15, 18, 28], we introduce the concepts of CNL-hyperenergetic and CNSL-hyperenergetic graphs. A graph \mathcal{G} is called CNL-hyperenergetic and CNSL-hyperenergetic if

$$LE_{CN}(\mathcal{G}) > LE_{CN}(K_{|V(\mathcal{G})|}) \text{ and } LE_{CN}^+(\mathcal{G}) > LE_{CN}^+(K_{|V(\mathcal{G})|}),$$

respectively.

The following lemma is useful in computing CN-Laplacian spectrum and CN-signless Laplacian spectrum of a graph having disconnected components.

Lemma 2.1. *If $\mathcal{G} = \mathcal{G}_1 \sqcup \mathcal{G}_2 \sqcup \cdots \sqcup \mathcal{G}_k$ (that is, $\mathcal{G}_1, \mathcal{G}_2, \dots, \mathcal{G}_k$ are the disconnected components of \mathcal{G}), then*

$$\text{CNL-spec}(\mathcal{G}) = \text{CNL-spec}(\mathcal{G}_1) \cup \text{CNL-spec}(\mathcal{G}_2) \cup \cdots \cup \text{CNL-spec}(\mathcal{G}_k)$$

and

$$\text{CNSL-spec}(\mathcal{G}) = \text{CNSL-spec}(\mathcal{G}_1) \cup \text{CNSL-spec}(\mathcal{G}_2) \cup \cdots \cup \text{CNSL-spec}(\mathcal{G}_k)$$

counting multiplicities.

Recall that first Zagreb index $M_1(\mathcal{G})$ is defined as

$$M_1(\mathcal{G}) = \sum_{i=1}^{|V(\mathcal{G})|} d_{\mathcal{G}}(v_i)^2 = \sum_{v_i v_j \in E(\mathcal{G})} (d_{\mathcal{G}}(v_i) + d_{\mathcal{G}}(v_j)).$$

Mathematical properties on the first Zagreb index was reported in [3, 5–10]. By [8], we have

$$M_1(\mathcal{G}) = \sum_{i=1}^{|V(\mathcal{G})|} d_{\mathcal{G}}(v_i)^2 = \sum_{i=1}^{|V(\mathcal{G})|} d_{\mathcal{G}}(v_i) m_{\mathcal{G}}(v_i). \quad (3)$$

The following lemmas are useful in deriving relations between CN-Laplacian energy, CN-signless Laplacian energy and first Zagreb index.

Lemma 2.2. [5] *Let \mathcal{G} be a graph of order $|V(\mathcal{G})|$. Then for each $v_i \in V(\mathcal{G})$,*

$$\sum_{k=1, k \neq i}^{|V(\mathcal{G})|} |N_{\mathcal{G}}(v_i) \cap N_{\mathcal{G}}(v_k)| = \sum_{v_j: v_i v_j \in E(\mathcal{G})} (d_{\mathcal{G}}(v_j) - 1) = d_{\mathcal{G}}(v_i) m_{\mathcal{G}}(v_i) - d_{\mathcal{G}}(v_i),$$

where $d_{\mathcal{G}}(v_i)$ is the degree of the vertex v_i in \mathcal{G} .

Lemma 2.3. *Let \mathcal{G} be a graph with $|e(\mathcal{G})|$ edges and the first Zagreb index $M_1(\mathcal{G})$. Then $tr(\text{CNRS}(\mathcal{G})) = M_1(\mathcal{G}) - 2|e(\mathcal{G})|$.*

Proof. From the definition with Lemma 2.2 and (3), we obtain

$$\begin{aligned}
tr(\text{CNRS}(\mathcal{G})) &= \sum_{i=1}^{|\mathcal{V}(\mathcal{G})|} \text{CNRS}(\mathcal{G})_{i,i} \\
&= \sum_{i=1}^{|\mathcal{V}(\mathcal{G})|} \sum_{j=1, j \neq i}^{|\mathcal{V}(\mathcal{G})|} |N_{\mathcal{G}}(v_i) \cap N_{\mathcal{G}}(v_j)| \\
&= \sum_{i=1}^{|\mathcal{V}(\mathcal{G})|} \left[d_{\mathcal{G}}(v_i) m_{\mathcal{G}}(v_i) - d_{\mathcal{G}}(v_i) \right] \\
&= M_1(\mathcal{G}) - 2|e(\mathcal{G})|.
\end{aligned}$$

This completes the result. □

We conclude this section with the following relations between $LE_{CN}(\mathcal{G})$, $LE_{CN}^+(\mathcal{G})$ and $M_1(\mathcal{G})$ which can be obtained from (1), (2) and Lemma 2.3.

Theorem 2.4. *Let \mathcal{G} be a graph with $|e(\mathcal{G})|$ edges and the first Zagreb index $M_1(\mathcal{G})$. Then*

$$LE_{CN}(\mathcal{G}) = \sum_{\nu \in \text{CNL-spec}(\mathcal{G})} \left| \nu - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|\mathcal{V}(\mathcal{G})|} \right|$$

and

$$LE_{CN}^+(\mathcal{G}) = \sum_{\sigma \in \text{CNL-spec}(\mathcal{G})} \left| \sigma - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|\mathcal{V}(\mathcal{G})|} \right|.$$

3 CN-(signless) Laplacian spectrum and CN-(signless) Laplacian energy

In this section we compute CN-(signless) Laplacian spectrum and CN-(signless) Laplacian energy of some classes of graphs and discuss their properties.

Example 3.1. For $n = 1$, it is clear that $\text{CNL-spec}(K_1) = \{0^1\}$, $\text{CNL-spec}(K_1) = \{0^1\}$ and so $LE_{CN} = 0$, $LE_{CN}^+ = 0$. Therefore, we consider $n \geq 2$. We have

$$\text{CN}(K_n) = (n-2)A(K_n) \quad \text{and} \quad \text{CNRS}(K_n) = \text{diag}[(n-1)(n-2), \dots, (n-1)(n-2)],$$

so $\text{CNL}(K_n) = (n-2)L(K_n)$ and $\text{CNSL}(K_n) = (n-2)Q(K_n)$. Also, $\text{L-spec}(K_n) = \{0^1, n^{n-1}\}$ and $\text{Q-spec}(K_n) = \{(2(n-1))^1, (n-2)^{n-1}\}$. Therefore $\text{CNL-spec}(K_n) = \{0^1, (n(n-2))^{n-1}\}$ and $\text{CNSL-spec}(K_n) = \{(2(n-1)(n-2))^1, ((n-2)^2)^{n-1}\}$. We have $\text{tr}(\text{CNRS}(K_n)) = n(n-1)(n-2)$ and so $\frac{\text{tr}(\text{CNRS}(K_n))}{|V(K_n)|} = (n-1)(n-2)$. Therefore

$$\left| 0 - \frac{\text{tr}(\text{CNRS}(K_n))}{|V(K_n)|} \right| = (n-1)(n-2),$$

$$\left| n(n-2) - \frac{\text{tr}(\text{CNRS}(K_n))}{|V(K_n)|} \right| = n(n-2) - (n-1)(n-2) = (n-2),$$

and

$$\left| 2(n-1)(n-2) - \frac{\text{tr}(\text{CNRS}(K_n))}{|V(K_n)|} \right| = (n-1)(n-2),$$

$$\left| (n-2)^2 - \frac{\text{tr}(\text{CNRS}(K_n))}{|V(K_n)|} \right| = |(n-2)^2 - (n-1)(n-2)| = |-(n-2)| = n-2.$$

Hence, by (1) and (2), we obtain

$$LE_{CN}(K_n) = (n-1)(n-2) + (n-1)(n-2) = 2(n-1)(n-2)$$

and

$$LE_{CN}^+(K_n) = (n-1)(n-2) + (n-1)(n-2) = 2(n-1)(n-2).$$

By [1, Proposition 2.4], it follows that if \mathcal{G}_1 and \mathcal{G}_2 are two disconnected component of \mathcal{G} , then $E_{CN}(\mathcal{G}) = E_{CN}(\mathcal{G}_1) + E_{CN}(\mathcal{G}_2)$. However, $LE_{CN}(\mathcal{G}) \neq LE_{CN}(\mathcal{G}_1) + LE_{CN}(\mathcal{G}_2)$ and $LE_{CN}^+(\mathcal{G}) \neq LE_{CN}^+(\mathcal{G}_1) + LE_{CN}^+(\mathcal{G}_2)$, if $\mathcal{G} = \mathcal{G}_1 \sqcup \mathcal{G}_2$. For example, if $\mathcal{G} = K_4 \sqcup K_6$ then, by Lemma 2.1 with the above result, it follows that

$$\text{CNL-spec}(\mathcal{G}) = \{0^2, 8^3, 24^5\} \quad \text{and} \quad \text{CNSL-spec}(\mathcal{G}) = \{4^3, (12)^1, (16)^5, (40)^1\}.$$

We have

$$\frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} = \frac{\text{tr}(\text{CNRS}(K_4) + \text{tr}(\text{CNRS}(K_6)))}{10} = \frac{24 + 120}{10} = \frac{144}{10}.$$

Therefore, by (1) and (2), we obtain

$$\begin{aligned} LE_{CN}(\mathcal{G}) &= 2 \times \left| 0 - \frac{144}{10} \right| + 3 \times \left| 8 - \frac{144}{10} \right| + 5 \times \left| 24 - \frac{144}{10} \right| \\ &= 2 \times \frac{144}{10} + 3 \times \frac{64}{10} + 5 \times \frac{96}{10} = 96, \end{aligned}$$

but $LE_{CN}(K_4) + LE_{CN}(K_6) = 12 + 40 = 52$, and

$$\begin{aligned} LE_{CN}^+(\mathcal{G}) &= 3 \times \left| 4 - \frac{144}{10} \right| + 1 \times \left| 12 - \frac{144}{10} \right| + 5 \times \left| 16 - \frac{144}{10} \right| + 1 \times \left| 40 - \frac{144}{10} \right| \\ &= 2 \times \frac{104}{10} + \frac{24}{10} + \frac{80}{10} + \frac{256}{10} = \frac{672}{10}, \end{aligned}$$

but $LE_{CN}^+(K_4) + LE_{CN}^+(K_6) = 12 + 40 = 52$.

Example 3.2. We now compute CN-(signless) Laplacian spectrum and CN-(signless) Laplacian energy of the complete bipartite graph $K_{m,n}$ on $(m+n)$ -vertices. For this, let $V(K_{m,n}) = \{v_1, v_2, \dots, v_m, v_{m+1}, v_{m+2}, \dots, v_{m+n}\}$ and $\{v_1, v_2, \dots, v_m\}, \{v_{m+1}, v_{m+2}, \dots, v_{m+n}\}$ be two partitions of $V(K_{m,n})$ such that every vertex in one set is adjacent to every vertex in the other set. We have

$$\text{CN}(K_{m,n}) = \begin{bmatrix} n A(K_m) & 0 \\ 0 & m A(K_n) \end{bmatrix}$$

and

$$\text{CNRS}(K_{m,n}) = \text{diag}[\underbrace{(m-1)n, \dots, (m-1)n}_{m\text{-times}}, \underbrace{(n-1)m, \dots, (n-1)m}_{n\text{-times}}].$$

Thus we have

$$\text{CNL}(K_{m,n}) = \begin{bmatrix} n L(K_m) & 0 \\ 0 & m L(K_n) \end{bmatrix} \text{ and } \text{CNSL}(K_{m,n}) = \begin{bmatrix} n Q(K_m) & 0 \\ 0 & m Q(K_n) \end{bmatrix}.$$

Since $L\text{-spec}(K_m) = \{0^1, m^{m-1}\}$ and $L\text{-spec}(K_n) = \{0^1, n^{n-1}\}$, therefore

$$\text{CNL-spec}(K_{m,n}) = \left\{ (n \times 0)^1, (n \times m)^{m-1}, (m \times 0)^1, (m \times n)^{n-1} \right\} = \left\{ 0^2, (mn)^{m+n-2} \right\}.$$

We have $\frac{\text{tr}(\text{CNRS}(K_{m,n}))}{|V(K_{m,n})|} = \frac{mn(m+n-2)}{m+n}$ and so

$$\left| 0 - \frac{\text{tr}(\text{CNRS}(K_{m,n}))}{|V(K_{m,n})|} \right| = \frac{mn(m+n-2)}{m+n} \text{ and } \left| mn - \frac{\text{tr}(\text{CNRS}(K_{m,n}))}{|V(K_{m,n})|} \right| = \frac{2mn}{m+n}.$$

Hence, by (1), we get

$$LE_{CN}(K_{m,n}) = \frac{2mn(m+n-2)}{m+n} + \frac{2mn(m+n-2)}{m+n} = \frac{4mn(m+n-2)}{m+n}.$$

In particular, for $m = n$, we obtain $\text{CNL-spec}(K_{m,n}) = \{0^2, (m^2)^{2m-2}\}$ and $LE_{CN}(K_{m,n}) = 4m(m-1)$.

Again since $\text{Q-spec}(K_m) = \{(2(m-1))^1, (m-2)^{m-1}\}$ and $\text{Q-spec}(K_n) = \{(2(n-1))^1, (n-2)^{n-1}\}$, therefore,

$$\text{CNSL-spec}(K_{m,n}) = \{(2n(m-1))^1, (n(m-2))^{m-1}, (2m(n-1))^1, (m(n-2))^{n-1}\}.$$

Note that if $m = n = 1$, then $K_{1,1} = K_2$. Hence, $\text{CNSL-spec}(K_{m,n}) = \text{CNSL-spec}(K_2) = \{0^2\}$ and $LE_{CN}^+(K_{1,1}) = LE_{CN}^+(K_2) = 0$. We now assume that $n \geq 2$ or $m \geq 2$. Now,

$$\begin{aligned} \left| 2n(m-1) - \frac{\text{tr}(\text{CNRS}(K_{m,n}))}{|V(K_{m,n})|} \right| &= \left| 2n(m-1) - \frac{mn(m+n-2)}{m+n} \right| \\ &= \begin{cases} \frac{n(n-1)}{n+1} & \text{if } m = 1, \\ \frac{n(m+n)(m-2) + 2mn}{m+n} & \text{if } m \geq 2, \end{cases} \\ \left| n(m-2) - \frac{\text{tr}(\text{CNRS}(K_{m,n}))}{|V(K_{m,n})|} \right| &= \left| n(m-2) - \frac{mn(m+n-2)}{m+n} \right| = \frac{2n^2}{m+n}, \\ \left| 2m(n-1) - \frac{\text{tr}(\text{CNRS}(K_{m,n}))}{|V(K_{m,n})|} \right| &= \left| 2m(n-1) - \frac{mn(m+n-2)}{m+n} \right| \\ &= \begin{cases} \frac{m(m-1)}{m+1} & \text{if } n = 1, \\ \frac{m(m+n)(n-2) + 2mn}{m+n} & \text{if } n \geq 2, \end{cases} \end{aligned}$$

and

$$\left| m(n-2) - \frac{\text{tr}(\text{CNRS}(K_{m,n}))}{|V(K_{m,n})|} \right| = \left| m(n-2) - \frac{mn(m+n-2)}{m+n} \right| = \frac{2m^2}{m+n}.$$

For $m = 1$ and $n \geq 2$, by (2), we have

$$LE_{CN}^+(K_{m,n}) = \frac{n(n-1)}{n+1} + \frac{(n+1)(n-2) + 2n}{n+1} + \frac{2(n-1)}{n+1} = \frac{2(n-1)(n+2)}{n+1}.$$

For $m \geq 2$ and $n = 1$, by (2), we have

$$LE_{CN}^+(K_{m,n}) = \frac{(m+1)(m-2) + 2m}{m+1} + \frac{2(m-1)}{m+1} + \frac{m(m-1)}{m+1} = \frac{2(m-1)(m+2)}{m+1}.$$

For $m, n \geq 2$, by (2), we have

$$\begin{aligned} & LE_{CN}^+(K_{m,n}) \\ &= \frac{n(m+n)(m-2) + 2mn}{m+n} + \frac{2n^2(m-1)}{m+n} + \frac{m(m+n)(n-2) + 2mn}{m+n} + \frac{2m^2(n-1)}{m+n} \\ &= \frac{4(m^2(n-1) + n^2(m-1))}{m+n}. \end{aligned}$$

Hence

$$LE_{CN}^+(K_{m,n}) = \begin{cases} 0 & \text{if } m = 1 \text{ and } n = 1, \\ \frac{2(n-1)(n+2)}{n+1} & \text{if } m = 1 \text{ and } n \geq 2, \\ \frac{2(m-1)(m+2)}{m+1} & \text{if } m \geq 2 \text{ and } n = 1, \\ \frac{4(m^2(n-1) + n^2(m-1))}{m+n} & \text{if } m, n \geq 2. \end{cases}$$

In particular, for $m = n$, we obtain $\text{CNSL-spec}(K_{m,n}) = \left\{ (2m(m-1))^2, (m(m-2))^{2(m-1)} \right\}$ and $LE_{CN}^+(K_{m,n}) = 4m(m-1)$.

Proposition 3.3. *Let $\overline{\mathcal{G}}$ be the complement of a graph \mathcal{G} and $\mathcal{G}_1 \vee \mathcal{G}_2$ be the join of two graphs \mathcal{G}_1 and \mathcal{G}_2 .*

(a) *If $\mathcal{G} = K_{n_1} \vee K_{n_2} \vee \cdots \vee K_{n_k}$, then*

$$\text{CNL-spec}(\mathcal{G}) = \{0^1, ((n_1 + n_2 + \cdots + n_k)(n_1 + n_2 + \cdots + n_k - 2))^{(n_1 + n_2 + \cdots + n_k - 1)}\},$$

$$\text{CNSL-spec}(\mathcal{G}) = \{(2(n_1 + n_2 + \cdots + n_k - 1)(n_1 + n_2 + \cdots + n_k - 2))^1, ((n_1 + n_2 + \cdots + n_k - 2)^2)^{n-1}\}$$

and

$$LE_{CN}(\mathcal{G}) = 2(n_1 + n_2 + \cdots + n_k - 1)(n_1 + n_2 + \cdots + n_k - 2) = LE_{CN}^+(\mathcal{G}).$$

(b) *If $\mathcal{G} = \overline{K}_m \vee \overline{K}_n$, then*

$$\text{CNL-spec}(\mathcal{G}) = \left\{ 0^2, (mn)^{m+n-2} \right\} \text{ and } LE_{CN}(\mathcal{G}) = \frac{4mn(m+n-2)}{m+n},$$

$$\text{CNSL-spec}(\mathcal{G}) = \left\{ (2n(m-1))^1, (n(m-2))^{m-1}, (2m(n-1))^1, (m(n-2))^{n-1} \right\}$$

and

$$LE_{CN}^+(\mathcal{G}) = \begin{cases} \frac{2(n-1)(n+2)}{n+1} & \text{if } m = 1 \text{ and } n \geq 2, \\ \frac{2(m-1)(m+2)}{m+1} & \text{if } m \geq 2 \text{ and } n = 1, \\ \frac{4(m^2(n-1) + n^2(m-1))}{m+n} & \text{if } m, n \geq 2. \end{cases}$$

Proof. The results follow from Examples 3.1 and 3.2 noting that $K_{n_1} \vee K_{n_2} \vee \cdots \vee K_{n_k} = K_{n_1+n_2+\cdots+n_k}$ and $\overline{K_m} \vee \overline{K_n} = K_{m,n}$. \square

Proposition 3.4. *The graph $K_{m,n}$ is not CNL-hyperenergetic.*

Proof. By Examples 3.1 and 3.2, we obtain

$$\begin{aligned} LE_{CN}(K_{m+n}) - LE_{CN}(K_{m,n}) &= 2(m+n-1)(m+n-2) - \frac{4mn(m+n-2)}{m+n} \\ &= \frac{[m(m-1) + n(n-1)](m+n-2)}{m+n} \geq 0 \end{aligned}$$

with equality if and only if $m = n = 1$. Therefore,

$$LE_{CN}(K_{m+n}) \geq LE_{CN}(K_{m,n})$$

with equality if and only if $m = n = 1$. Hence, the result follows. \square

Corollary 3.5. *If S_k denotes the star graph with one internal node and k leaves, then*

$$\text{CNL-spec}(S_k) = \left\{ 0^2, k^{k-1} \right\} \text{ and } LE_{CN}(S_k) = \frac{4k(k-1)}{k+1}.$$

Moreover, S_k is not CNL-hyperenergetic.

Proof. The result follows from Example 3.2 and Proposition 3.4, noting that $S_k = K_{1,k}$. \square

Proposition 3.6. *The graph $K_{m,n}$ is not CNSL-hyperenergetic.*

Proof. If $m = n = 1$ then we have $LE_{CN}^+(K_{1,1}) = LE_{CN}^+(K_2) = 0$. Therefore, we consider the case when m, n are not equal to 1 simultaneously. By Example 3.1, we have

$$LE_{CN}^+(K_{m+n}) = 2(m+n-1)(m+n-2). \quad (4)$$

For $m = 1$ and $n \geq 2$, by (4) and Example 3.2, we have

$$\begin{aligned} LE_{CN}^+(K_{m+n}) - LE_{CN}^+(K_{m,n}) &= 2n(n-1) - \frac{2(n-1)(n+2)}{n+1} \\ &= \frac{2(n-1)(n^2-2)}{n+1} > 0. \end{aligned}$$

For $n = 1$ and $m \geq 2$, by (4) and Example 3.2, we have

$$\begin{aligned} LE_{CN}^+(K_{m+n}) - LE_{CN}^+(K_{m,n}) &= 2m(m-1) - \frac{2(m-1)(m+2)}{m+1} \\ &= \frac{2(m-1)(m^2-2)}{m+1} > 0. \end{aligned}$$

For $m, n \geq 2$, by (4) and Example 3.2, we have

$$\begin{aligned} LE_{CN}^+(K_{m+n}) - LE_{CN}^+(K_{m,n}) &= 2(m+n-1)(m+n-2) - \frac{4(m^2(n-1) + n^2(m-1))}{m+n} \\ &= 2(m^3 + n^3 + m + n) + (m-n)^2 + 2mn(m+n-2) > 0, \end{aligned}$$

since $m+n-2 > 0$. Therefore,

$$LE_{CN}^+(K_{m+n}) \geq LE_{CN}^+(K_{m,n}).$$

Hence, the result follows. \square

Corollary 3.7. *If S_k denotes the star graph with one internal node and k leaves then*

$$\text{CNSL-spec}(S_k) = \left\{ 0^1, (2(k-1))^1, ((k-2))^{k-1} \right\} \text{ and } LE_{CN}^+(S_k) = \frac{2(k-1)(k+2)}{k+1}.$$

Moreover, S_k is not CNSL-hyperenergetic.

Proof. The result follows from Example 3.2 and Proposition 3.6, noting that $S_k = K_{1,k}$. \square

4 Relation between various energies

In this section we derive some relations between E_{CN} , LE_{CN} , LE_{CN}^+ , E , LE and LE^+ of a graph \mathcal{G} . Let $e(\mathcal{G})$ be the set of edges of a graph \mathcal{G} . Let $\text{Spec}(M)$ and $E(M)$ be the

spectrum and energy of a real square symmetric matrix M of size n . Then $\text{Spec}(M) = \{\mu_1, \mu_2, \dots, \mu_n\}$, where $\mu_1, \mu_2, \dots, \mu_n$ are eigenvalues (not necessarily distinct) of M , and

$$E(M) = \sum_{i=1}^n |\mu_i|.$$

The following two lemmas are useful in obtaining a relation between $E_{CN}(\mathcal{G})$ and $E(\mathcal{G})$.

Lemma 4.1. [1, Proposition 2.7] *Let \mathcal{G} be any graph. Then $CN(\mathcal{G}) = A(\mathcal{G})^2 - D(\mathcal{G})$.*

Lemma 4.2. [12, Lemma 2.10] *Let M_1 and M_2 be two real square symmetric matrices of order n and let $M = M_1 + M_2$. Then*

$$E(M) \leq E(M_1) + E(M_2).$$

Theorem 4.3. *Let \mathcal{G} be any graph with $|e(\mathcal{G})|$ edges. Then $E_{CN}(\mathcal{G}) \leq E(\mathcal{G})^2 + 2|e(\mathcal{G})|$.*

Proof. By Lemmas 4.1 and 4.2, we obtain

$$\begin{aligned} E_{CN}(\mathcal{G}) &= E(CN(\mathcal{G})) \\ &= E(A(\mathcal{G})^2 - D(\mathcal{G})) \\ &= E(A(\mathcal{G})^2 + (-D(\mathcal{G}))) \leq E(A(\mathcal{G})^2) + E(-D(\mathcal{G})). \end{aligned} \quad (5)$$

Let $\text{Spec}(\mathcal{G}) = \{\lambda_1, \lambda_2, \dots, \lambda_n\}$, where $n = |V(\mathcal{G})|$. Then $\text{Spec}(A(\mathcal{G})^2) = \{\lambda_1^2, \lambda_2^2, \dots, \lambda_n^2\}$.

Therefore,

$$E(A(\mathcal{G})^2) = \sum_{i=1}^n |\lambda_i^2| \leq \left(\sum_{i=1}^n |\lambda_i| \right)^2 = E(\mathcal{G})^2.$$

Again, let $\text{Spec}(D(\mathcal{G})) = \{d_{\mathcal{G}}(v_1), d_{\mathcal{G}}(v_2), \dots, d_{\mathcal{G}}(v_n)\}$. Then $\text{Spec}(-D(\mathcal{G})) = \{-d_{\mathcal{G}}(v_1), -d_{\mathcal{G}}(v_2), \dots, -d_{\mathcal{G}}(v_n)\}$. Therefore

$$E(-D(\mathcal{G})) = \sum_{i=1}^n |-d_{\mathcal{G}}(v_i)| = \sum_{i=1}^n d_{\mathcal{G}}(v_i) = 2 \times |e(\mathcal{G})| = E(D(\mathcal{G})).$$

Hence, the result follows from (5). □

Corollary 4.4. *Let \mathcal{G} be any graph of order $|V(\mathcal{G})|$ with $|e(\mathcal{G})|$ edges. Then $E_{CN}(\mathcal{G}) \leq 2|e(\mathcal{G})| (|V(\mathcal{G})| + 1)$.*

Proof. It is well-known that $E(\mathcal{G}) \leq \sqrt{2|e(\mathcal{G})||V(\mathcal{G})|}$. Using the above result with Theorem 4.3, we obtain

$$E_{CN}(\mathcal{G}) \leq E(\mathcal{G})^2 + 2|e(\mathcal{G})| \leq 2|e(\mathcal{G})||V(\mathcal{G})| + 2|e(\mathcal{G})|.$$

Hence the result follows. \square

Theorem 4.5. *Let \mathcal{G} be any graph with $|e(\mathcal{G})|$ edges and the first Zagreb index $M_1(\mathcal{G})$. Then $LE_{CN}(\mathcal{G}) \leq E_{CN}(\mathcal{G}) + 2(M_1(\mathcal{G}) - 2|e(\mathcal{G})|)$ and $LE_{CN}^+(\mathcal{G}) \leq E_{CN}(\mathcal{G}) + 2(M_1(\mathcal{G}) - 2|e(\mathcal{G})|)$.*

Proof. By Theorem 2.4, we have

$$\begin{aligned} LE_{CN}(\mathcal{G}) &= \sum_{\nu \in \text{CNL-spec}(\mathcal{G})} \left| \nu - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|} \right| \\ &\leq \sum_{\nu \in \text{CNL-spec}(\mathcal{G})} |\nu| + \sum_{\nu \in \text{CNL-spec}(\mathcal{G})} \left| \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|} \right| \\ &= E(\text{CNL}(\mathcal{G})) + \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|} \sum_{\nu \in \text{CNL-spec}(\mathcal{G})} 1 \\ &= E(\text{CNRS}(\mathcal{G}) - \text{CN}(\mathcal{G})) + M_1(\mathcal{G}) - 2|e(\mathcal{G})|. \end{aligned}$$

Using Lemmas 4.2 & 2.3 and the fact that $E(\text{CN}(\mathcal{G})) = E(-\text{CN}(\mathcal{G}))$ & $E(\text{CNRS}(\mathcal{G})) = \text{tr}(\text{CNRS}(\mathcal{G}))$, we obtain

$$\begin{aligned} LE_{CN}(\mathcal{G}) &\leq E(\text{CNRS}(\mathcal{G})) + E(\text{CN}(\mathcal{G})) + M_1(\mathcal{G}) - 2|e(\mathcal{G})| \\ &= E_{CN}(\mathcal{G}) + 2(M_1(\mathcal{G}) - 2|e(\mathcal{G})|). \end{aligned}$$

Similarly, the bound for $LE_{CN}^+(\mathcal{G})$ follows from Theorem 2.4 and Lemma 4.2. \square

As a consequence of Theorems 4.3 and 4.5, we get the following relations between $E(\mathcal{G})$, $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$.

Corollary 4.6. *Let \mathcal{G} be any graph with $|e(\mathcal{G})|$ edges and the first Zagreb index $M_1(\mathcal{G})$. Then $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$ are bounded above by*

$$E(\mathcal{G})^2 + 2(M_1(\mathcal{G}) - |e(\mathcal{G})|).$$

Remark 4.7. Theorem 4.3 gives relation between $E_{CN}(\mathcal{G})$ and $E(\mathcal{G})$. Theorem 4.5 and Corollary 4.6 give relations between $LE_{CN}(\mathcal{G})$, $LE_{CN}^+(\mathcal{G})$, $E_{CN}(\mathcal{G})$ and $E(\mathcal{G})$. However, using the facts that

$$E(A(\mathcal{G})^2) = \sum_{i=1}^n |\lambda_i^2| = 2|e(\mathcal{G})| \leq E(\mathcal{G})^2$$

and

$$\sum_{\nu \in \text{CNL-spec}(\mathcal{G})} |\nu| = \text{tr}(\text{CNRS}(\mathcal{G})) = M_1(\mathcal{G}) - 2|e(\mathcal{G})| = \sum_{\sigma \in \text{CNSL-spec}(\mathcal{G})} |\sigma|,$$

we get the following better upper bounds for $E_{CN}(\mathcal{G})$, $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$:

$$E_{CN}(\mathcal{G}) \leq 4|e(\mathcal{G})| \leq 2E(\mathcal{G})^2, \quad (6)$$

$$LE_{CN}(\mathcal{G}) \leq 2 \text{tr}(\text{CNRS}(\mathcal{G})) = 2 \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right) \geq LE_{CN}^+(\mathcal{G}). \quad (7)$$

In Section 5, we shall obtain more bounds for $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$.

Recall that the derived graph of \mathcal{G} , denoted by \mathcal{G}^\dagger is the graph with vertex set $V(\mathcal{G})$, in which two vertices are adjacent if and only if their distance in \mathcal{G} is two.

Theorem 4.8. *If \mathcal{G} is a triangle- and quadrangle-free graph, then $LE_{CN}(\mathcal{G}) = LE(\mathcal{G}^\dagger)$ and $LE_{CN}^+(\mathcal{G}) = LE^+(\mathcal{G}^\dagger)$, where \mathcal{G}^\dagger is the derived graph of \mathcal{G} .*

Proof. If \mathcal{G} is a triangle- and quadrangle-free graph, then $CN(\mathcal{G}) = A(\mathcal{G}^\dagger)$. Therefore, $\text{CNRS}(\mathcal{G}) = D(\mathcal{G}^\dagger)$ and so $\text{CNL}(\mathcal{G}) = L(\mathcal{G}^\dagger)$ and $\text{CNSL}(\mathcal{G}) = Q(\mathcal{G}^\dagger)$. Hence, $\text{CNL-spec}(\mathcal{G}) = \text{L-spec}(\mathcal{G}^\dagger)$ and $\text{CNSL-spec}(\mathcal{G}) = \text{Q-spec}(\mathcal{G}^\dagger)$. Since $\text{tr}(\text{CNRS}(\mathcal{G})) = \text{tr}(D(\mathcal{G}^\dagger))$ and $V(\mathcal{G}) = V(\mathcal{G}^\dagger)$, by (1) and (2), we have

$$LE_{CN}(\mathcal{G}) = \sum_{\nu \in \text{L-spec}(\mathcal{G}^\dagger)} \left| \nu - \frac{\text{tr}(D(\mathcal{G}^\dagger))}{|V(\mathcal{G})|} \right| = LE(\mathcal{G}^\dagger)$$

and

$$LE_{CN}^+(\mathcal{G}) = \sum_{\sigma \in \text{Q-spec}(\mathcal{G}^\dagger)} \left| \sigma - \frac{\text{tr}(D(\mathcal{G}^\dagger))}{|V(\mathcal{G})|} \right| = LE^+(\mathcal{G}^\dagger).$$

□

Corollary 4.9. (a) *If T is a tree then $LE_{CN}(T) = LE(T^\dagger)$ and $LE_{CN}^+(T) = LE^+(T^\dagger)$.*

(b) If P_n is the path on n vertices, then $LE_{CN}(P_n) = LE(P_{\lceil \frac{n}{2} \rceil}) + LE(P_{\lfloor \frac{n}{2} \rfloor})$ and $LE_{CN}^+(P_n) = LE^+(P_{\lceil \frac{n}{2} \rceil}) + LE^+(P_{\lfloor \frac{n}{2} \rfloor})$.

(c) Let C_n be the cycle on n vertices.

(i) If n is odd and $n \geq 3$, then $LE_{CN}(C_n) = LE(C_n)$ and $LE_{CN}^+(C_n) = LE^+(C_n)$.

(ii) If n is even and $n > 4$, then $LE_{CN}(C_n) = 2LE(C_{\frac{n}{2}})$ and $LE_{CN}^+(C_n) = 2LE^+(C_{\frac{n}{2}})$. Also, $LE_{CN}(C_4) = 2LE(C_4) = LE_{CN}^+(C_4) = 2LE^+(C_4) = 8$.

Proof. (a) Follows from Theorem 4.8 noting that T is triangle- and quadrangle-free.

(b) Follows from Theorem 4.8 noting that P_n is triangle- and quadrangle-free and

$$P_n^\dagger \cong P_{\lceil \frac{n}{2} \rceil} \cup P_{\lfloor \frac{n}{2} \rfloor}.$$

(c) Let C_n be the cycle on n vertices.

(i) If $n = 3$, then $C_3 = K_3$. Therefore, $LE_{CN}(C_3) = LE_{CN}^+(C_3) = 4 = LE(C_3) = LE^+(C_3)$.

If n is odd and $n > 3$, then C_n is triangle- and quadrangle-free. Also, $(C_n)^\dagger \cong C_n$. Hence, the result follows from Theorem 4.8.

(ii) If n is even and $n > 4$, then C_n is triangle- and quadrangle-free. Also, $(C_n)^\dagger \cong C_{\frac{n}{2}} \cup C_{\frac{n}{2}}$. Therefore, by Theorem 4.8 we get

$$LE_{CN}(C_n) = LE(C_{\frac{n}{2}} \cup C_{\frac{n}{2}}) = 2LE(C_{\frac{n}{2}})$$

and

$$LE_{CN}^+(C_n) = LE^+(C_{\frac{n}{2}} \cup C_{\frac{n}{2}}) = 2LE^+(C_{\frac{n}{2}}).$$

If $n = 4$ then it is easy to see that $CNRS(C_4) = D(C_4)$, which is a 4×4 diagonal matrix such that every element in the diagonal is equal to 2, and $CNL\text{-spec}(C_4) = CNSL\text{-spec}(C_4) = \{0^2, 4^2\}$. Therefore, by (1) and (2), we have

$$LE_{CN}(C_4) = LE_{CN}^+(C_4) = 8.$$

Again, $L\text{-spec}(C_4) = Q\text{-spec}(C_4) = \{0, 2^2, 4\}$ and so $LE(C_4) = LE^+(C_4) = 4$. Thus, $LE_{CN}(C_4) = 2LE(C_4)$ and $LE_{CN}^+(C_4) = 2LE^+(C_4)$. \square

5 More bounds for $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$

In this section we shall obtain several bounds for $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$. Since the matrices $CNL(\mathcal{G})$ and $CNSL(\mathcal{G})$ are positive semidefinite, the elements of $CNL\text{-spec}(\mathcal{G})$ and $CNSL\text{-spec}(\mathcal{G})$ are non-negative. Thus we may write $CNL\text{-spec}(\mathcal{G}) = \{\nu_1, \nu_2, \dots, \nu_{|V(\mathcal{G})|}\}$ and $CNSL\text{-spec}(\mathcal{G}) = \{\sigma_1, \sigma_2, \dots, \sigma_{|V(\mathcal{G})|}\}$, where $\nu_1 \geq \nu_2 \geq \dots \geq \nu_{|V(\mathcal{G})|}$ and $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_{|V(\mathcal{G})|}$. We have

$$\sum_{i=1}^{|V(\mathcal{G})|} \nu_i = \sum_{\nu \in CNL\text{-spec}(\mathcal{G})} \nu = tr(CNRS(\mathcal{G})) = \sum_{\sigma \in CNSL\text{-spec}(\mathcal{G})} \sigma = \sum_{i=1}^{|V(\mathcal{G})|} \sigma_i.$$

Also,

$$\sum_{\nu \in CNL\text{-spec}(\mathcal{G})} \left(\nu - \frac{tr(CNRS(\mathcal{G}))}{|V(\mathcal{G})|} \right) = \sum_{\sigma \in CNSL\text{-spec}(\mathcal{G})} \left(\sigma - \frac{tr(CNRS(\mathcal{G}))}{|V(\mathcal{G})|} \right) = 0.$$

Let α, β ($1 \leq \alpha, \beta \leq |V(\mathcal{G})|$) be the largest integers such that

$$\nu_\alpha \geq \frac{tr(CNRS(\mathcal{G}))}{|V(\mathcal{G})|} = \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|} \text{ and } \sigma_\beta \geq \frac{tr(CNRS(\mathcal{G}))}{|V(\mathcal{G})|} = \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|}. \quad (8)$$

Let $S_\alpha(\mathcal{G}) = \sum_{i=1}^{\alpha} \nu_i$ and $S_\beta^+(\mathcal{G}) = \sum_{i=1}^{\beta} \sigma_i$. Then we have the following useful lemmas.

Lemma 5.1. *For any graph \mathcal{G} , we have*

$$LE_{CN}(\mathcal{G}) = 2S_\alpha(\mathcal{G}) - \frac{2\alpha tr(CNRS(\mathcal{G}))}{|V(\mathcal{G})|} = 2S_\alpha(\mathcal{G}) - \frac{2\alpha (M_1(\mathcal{G}) - 2|e(\mathcal{G})|)}{|V(\mathcal{G})|}$$

and

$$LE_{CN}^+(\mathcal{G}) = 2S_\beta^+(\mathcal{G}) - \frac{2\beta tr(CNRS(\mathcal{G}))}{|V(\mathcal{G})|} = 2S_\beta^+(\mathcal{G}) - \frac{2\beta (M_1(\mathcal{G}) - 2|e(\mathcal{G})|)}{|V(\mathcal{G})|},$$

where $|e(\mathcal{G})|$ is the number of edges and $M_1(\mathcal{G})$ is the first Zagreb index in \mathcal{G} .

Lemma 5.2. *For any graph \mathcal{G} , we have*

$$LE_{CN}(\mathcal{G}) = \max_{1 \leq i \leq |V(\mathcal{G})|} \left\{ 2S_i(\mathcal{G}) - \frac{2i (M_1(\mathcal{G}) - 2|e(\mathcal{G})|)}{|V(\mathcal{G})|} \right\}$$

and

$$LE_{CN}^+(\mathcal{G}) = \max_{1 \leq i \leq |V(\mathcal{G})|} \left\{ 2S_i^+(\mathcal{G}) - \frac{2i \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right)}{|V(\mathcal{G})|} \right\},$$

where $|e(\mathcal{G})|$ is the number of edges and $M_1(\mathcal{G})$ is the first Zagreb index in \mathcal{G} .

Proof. Let k ($1 \leq k \leq |V(\mathcal{G})|$) be any integer. For $k < \alpha$, by (8), we obtain

$$S_\alpha(\mathcal{G}) - S_k(\mathcal{G}) = \sum_{i=k+1}^{\alpha} \nu_i \geq \frac{(\alpha - k) \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right)}{|V(\mathcal{G})|}.$$

For $k > \alpha$, we obtain

$$S_k(\mathcal{G}) - S_\alpha(\mathcal{G}) = \sum_{i=\alpha+1}^k \nu_i < \frac{(k - \alpha) \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right)}{|V(\mathcal{G})|},$$

that is,

$$S_\alpha(\mathcal{G}) - S_k(\mathcal{G}) > \frac{(\alpha - k) \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right)}{|V(\mathcal{G})|}.$$

Moreover, $S_\alpha(\mathcal{G}) = S_k(\mathcal{G})$ for $k = \alpha$. Thus for any value of k ($1 \leq k \leq |V(\mathcal{G})|$), we obtain

$$S_\alpha(\mathcal{G}) - S_k(\mathcal{G}) \geq \frac{(\alpha - k) \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right)}{|V(\mathcal{G})|}$$

and so

$$2S_\alpha(\mathcal{G}) - \frac{2\alpha \operatorname{tr}(\operatorname{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} \geq 2S_k(\mathcal{G}) - \frac{2k \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right)}{|V(\mathcal{G})|}.$$

This gives

$$2S_\alpha(\mathcal{G}) - \frac{2\alpha \operatorname{tr}(\operatorname{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} = \max_{1 \leq i \leq |V(\mathcal{G})|} \left\{ 2S_i(\mathcal{G}) - \frac{2i \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right)}{|V(\mathcal{G})|} \right\}.$$

Similarly, it can be seen that

$$2S_\beta^+(\mathcal{G}) - \frac{2\beta \operatorname{tr}(\operatorname{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} = \max_{1 \leq i \leq |V(\mathcal{G})|} \left\{ 2S_i^+(\mathcal{G}) - \frac{2i \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right)}{|V(\mathcal{G})|} \right\}.$$

Hence, the result follows from Lemma 5.1. \square

Let $(a) := (a_1, a_2, \dots, a_n) \in \mathbb{R}^n$ and $(b) := (b_1, b_2, \dots, b_n) \in \mathbb{R}^n$ be such that $a_1 \geq a_2 \geq \dots \geq a_n$ and $b_1 \geq b_2 \geq \dots \geq b_n$. Then (a) is said to be majorize (b) if

$$\sum_{i=1}^k a_i \geq \sum_{i=1}^k b_i \text{ for } 1 \leq k \leq n-1 \quad \text{and} \quad \sum_{i=1}^n a_i = \sum_{i=1}^n b_i.$$

It is well-known that the spectrum of any symmetric, positive semidefinite matrix majorizes its main diagonal (see [27], [24, p. 218] as noted in [16]). Since $\text{CNL}(\mathcal{G})$ and $\text{CNSL}(\mathcal{G})$ are symmetric and positive semidefinite for any graph \mathcal{G} , we have the following lemma when the elements of $\text{CNL-spec}(\mathcal{G})$, $\text{CNSL-spec}(\mathcal{G})$ and main diagonal elements of $\text{CNRS}(\mathcal{G})$ are arranged in decreasing order.

Lemma 5.3. *For any graph \mathcal{G} , $\text{CNL-spec}(\mathcal{G})$ and $\text{CNSL-spec}(\mathcal{G})$ majorize main diagonal elements of $\text{CNRS}(\mathcal{G})$ when the elements of $\text{CNL-spec}(\mathcal{G})$, $\text{CNSL-spec}(\mathcal{G})$ and main diagonal elements of $\text{CNRS}(\mathcal{G})$ are arranged in decreasing order.*

We write the main diagonal elements of $\text{CNRS}(\mathcal{G})$ as $\text{CNRS}(\mathcal{G})_{i,i}$ for $1 \leq i \leq |V(\mathcal{G})|$, where $\text{CNRS}(\mathcal{G})_{1,1} \geq \text{CNRS}(\mathcal{G})_{2,2} \geq \dots \geq \text{CNRS}(\mathcal{G})_{|V(\mathcal{G})|,|V(\mathcal{G})|}$. Now we give lower bounds for $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$ analogous to the bound given by [11, Theorem 3.1] for $LE(\mathcal{G})$.

Theorem 5.4. *Let \mathcal{G} be a graph with $|e(\mathcal{G})|$ edges and the first Zagreb index $M_1(\mathcal{G})$. Then*

$$LE_{CN}(\mathcal{G}) \geq 2 \left(\Delta(\delta - 1) - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|} \right)$$

and

$$LE_{CN}^+(\mathcal{G}) \geq 2 \left(\Delta(\delta - 1) - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|} \right),$$

where Δ and δ are the maximum degree and the minimum degree in \mathcal{G} , respectively.

Proof. Let v_1 be the maximum degree vertex in \mathcal{G} . Then $d_{\mathcal{G}}(v_1) = \Delta$ and $m_{\mathcal{G}}(v_1) \geq \delta$ as δ is the minimum degree in \mathcal{G} . As a consequence of Lemma 5.3 with Lemma 2.2, we obtain

$$\nu_1 \geq \text{CNRS}(\mathcal{G})_{1,1} = d_{\mathcal{G}}(v_1) m_{\mathcal{G}}(v_1) - d_{\mathcal{G}}(v_1) = \Delta (m_{\mathcal{G}}(v_1) - 1) \geq \Delta(\delta - 1)$$

and

$$\sigma_1 \geq \text{CNRS}(\mathcal{G})_{1,1} = d_{\mathcal{G}}(v_1) m_{\mathcal{G}}(v_1) - d_{\mathcal{G}}(v_1) = \Delta (m_{\mathcal{G}}(v_1) - 1) \geq \Delta(\delta - 1),$$

Using the above result with Lemma 5.2, we obtain

$$\begin{aligned} LE_{CN}(\mathcal{G}) &\geq 2S_1(\mathcal{G}) - \frac{2(M_1(\mathcal{G}) - 2|e(\mathcal{G})|)}{|V(\mathcal{G})|} = 2\nu_1 - \frac{2(M_1(\mathcal{G}) - 2|e(\mathcal{G})|)}{|V(\mathcal{G})|} \\ &\geq 2\left(\Delta(\delta - 1) - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|}\right). \end{aligned}$$

Similarly,

$$LE_{CN}^+(\mathcal{G}) \geq 2\sigma_1 - \frac{2(M_1(\mathcal{G}) - 2|e(\mathcal{G})|)}{|V(\mathcal{G})|} \geq 2\left(\Delta(\delta - 1) - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|}\right).$$

□

Theorem 5.5. *Let \mathcal{G} be a graph with $|e(\mathcal{G})|$ edges and the first Zagreb index $M_1(\mathcal{G})$. Then*

$$LE_{CN}(\mathcal{G}) \geq 2\left(\sum_{i=1}^{\alpha} \text{CNRS}(\mathcal{G})_{i,i} - \frac{\alpha(M_1(\mathcal{G}) - 2|e(\mathcal{G})|)}{|V(\mathcal{G})|}\right)$$

and

$$LE_{CN}^+(\mathcal{G}) \geq 2\left(\sum_{i=1}^{\beta} \text{CNRS}(\mathcal{G})_{i,i} - \frac{\beta(M_1(\mathcal{G}) - 2|e(\mathcal{G})|)}{|V(\mathcal{G})|}\right),$$

where α and β are as given in (8) and $\text{CNRS}(\mathcal{G})_{i,i} = d_{\mathcal{G}}(v_i)(m_{\mathcal{G}}(v_i) - 1)$.

Proof. By Lemma 2.2, we have $\text{CNRS}(\mathcal{G})_{i,i} = d_{\mathcal{G}}(v_i)(m_{\mathcal{G}}(v_i) - 1)$. By Lemma 5.3, we obtain

$$\sum_{i=1}^k \nu_i \geq \sum_{i=1}^k \text{CNRS}(\mathcal{G})_{i,i} \text{ and } \sum_{i=1}^k \sigma_i \geq \sum_{i=1}^k \text{CNRS}(\mathcal{G})_{i,i} \text{ for } 1 \leq k \leq |V(\mathcal{G})|.$$

In particular, we have

$$\sum_{i=1}^{\alpha} \nu_i \geq \sum_{i=1}^{\alpha} \text{CNRS}(\mathcal{G})_{i,i} \text{ and } \sum_{i=1}^{\beta} \sigma_i \geq \sum_{i=1}^{\beta} \text{CNRS}(\mathcal{G})_{i,i}.$$

Therefore,

$$S_{\alpha}(\mathcal{G}) \geq \sum_{i=1}^{\alpha} \text{CNRS}(\mathcal{G})_{i,i} \text{ and } S_{\beta}^+(\mathcal{G}) \geq \sum_{i=1}^{\beta} \text{CNRS}(\mathcal{G})_{i,i}.$$

Hence, the result follows from Lemma 5.1. □

Using Lemma 5.1, we also have the following upper bounds for $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$ analogous to the bound given in [11, Remark 3.8].

Theorem 5.6. *Let \mathcal{G} be a graph of order $|V(\mathcal{G})|$ with $|e(\mathcal{G})|$ edges and the first Zagreb index $M_1(\mathcal{G})$. Then $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$ are bounded above by*

$$2 \left(1 - \frac{1}{|V(\mathcal{G})|} \right) \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right).$$

Proof. We have

$$S_\alpha(\mathcal{G}) \leq S_{|V(\mathcal{G})|}(\mathcal{G}) = \text{tr}(\text{CNRS}(\mathcal{G})) = M_1(\mathcal{G}) - 2|e(\mathcal{G})|$$

and

$$S_\beta^+(\mathcal{G}) \leq S_{|V(\mathcal{G})|}^+(\mathcal{G}) = \text{tr}(\text{CNRS}(\mathcal{G})) = M_1(\mathcal{G}) - 2|e(\mathcal{G})|.$$

Therefore, by Lemma 5.1 with $1 \leq \alpha \leq |V(\mathcal{G})|$, we obtain

$$LE_{CN}(\mathcal{G}) \leq 2 \left(1 - \frac{1}{|V(\mathcal{G})|} \right) \left(M_1(\mathcal{G}) - 2|e(\mathcal{G})| \right).$$

Similarly, we get the bound for $LE_{CN}^+(\mathcal{G})$. □

Note that the bounds obtained in Theorem 5.6 are better than the bounds obtained in (7). We conclude this section with another upper bounds for $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$ analogous to the bound obtained in [13, Theorem 5.5]. The following lemma is useful in this regard.

Lemma 5.7. [13, Lemma 5.1] *Let A be a real symmetric matrix of order n and let d_1, d_2, \dots, d_n be the diagonal entries of the matrix A^2 . Then*

$$E(A) \leq \sum_{i=1}^n \sqrt{d_i}.$$

Theorem 5.8. *Let \mathcal{G} be a graph of order $|V(\mathcal{G})|$ with $|e(\mathcal{G})|$ edges and the first Zagreb index $M_1(\mathcal{G})$. Then*

$$(a) \quad LE_{CN}(\mathcal{G}) \leq \sum_{i=1}^{|V(\mathcal{G})|} \sqrt{\left(\text{CNRS}(\mathcal{G})_{i,i} - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|} \right)^2 + \sum_{k=1, k \neq i}^{|V(\mathcal{G})|} |N_{\mathcal{G}}(v_i) \cap N_{\mathcal{G}}(v_k)|^2},$$

$$(b) \quad LE_{CN}^+(\mathcal{G}) \leq \sum_{i=1}^{|\mathcal{V}(\mathcal{G})|} \sqrt{\left(\text{CNRS}(\mathcal{G})_{i,i} - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|\mathcal{V}(\mathcal{G})|} \right)^2 + \sum_{k=1, k \neq i}^{|\mathcal{V}(\mathcal{G})|} |N_{\mathcal{G}}(v_i) \cap N_{\mathcal{G}}(v_k)|^2},$$

where $\text{CNRS}(\mathcal{G})_{i,i} = d_{\mathcal{G}}(v_i) (m_{\mathcal{G}}(v_i) - 1)$ for $1 \leq i \leq |\mathcal{V}(\mathcal{G})|$ are main diagonal elements of $\text{CNRS}(\mathcal{G})$ such that $\text{CNRS}(\mathcal{G})_{1,1} \geq \text{CNRS}(\mathcal{G})_{2,2} \geq \dots \geq \text{CNRS}(\mathcal{G})_{|\mathcal{V}(\mathcal{G})|,|\mathcal{V}(\mathcal{G})|}$.

Proof. (a) Let $M = \text{CNL}(\mathcal{G}) - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|\mathcal{V}(\mathcal{G})|} I_{|\mathcal{V}(\mathcal{G})|}$, where $I_{|\mathcal{V}(\mathcal{G})|}$ is the identity matrix of size $|\mathcal{V}(\mathcal{G})|$. Then $\text{Spec}(M) = \left\{ \nu_i - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|\mathcal{V}(\mathcal{G})|} : 1 \leq i \leq |\mathcal{V}(\mathcal{G})| \right\}$, where $\text{CNL-spec}(\mathcal{G}) = \{\nu_1, \nu_2, \dots, \nu_{|\mathcal{V}(\mathcal{G})|}\}$. We have

$$\begin{aligned} M^2 &= \left(\text{CNL}(\mathcal{G}) - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|\mathcal{V}(\mathcal{G})|} I_{|\mathcal{V}(\mathcal{G})|} \right)^2 \\ &= (\text{CNL}(\mathcal{G}))^2 - \frac{2 \text{tr}(\text{CNRS}(\mathcal{G}))}{|\mathcal{V}(\mathcal{G})|} \text{CNL}(\mathcal{G}) + \frac{(\text{tr}(\text{CNRS}(\mathcal{G})))^2}{|\mathcal{V}(\mathcal{G})|^2} I_{|\mathcal{V}(\mathcal{G})|}. \end{aligned}$$

Therefore, the i -th diagonal element of M^2 is

$$(M^2)_{i,i} = (\text{CNL}(\mathcal{G}))_{i,i}^2 - \frac{2 \text{tr}(\text{CNRS}(\mathcal{G}))}{|\mathcal{V}(\mathcal{G})|} (\text{CNRS}(\mathcal{G}))_{i,i} + \frac{(\text{tr}(\text{CNRS}(\mathcal{G})))^2}{|\mathcal{V}(\mathcal{G})|^2}.$$

We have

$$\begin{aligned} (\text{CNL}(\mathcal{G}))^2 &= \left(\text{CNRS}(\mathcal{G}) - \text{CN}(\mathcal{G}) \right)^2 \\ &= \text{CNRS}(\mathcal{G})^2 - \text{CNRS}(\mathcal{G}) \text{CN}(\mathcal{G}) - \text{CN}(\mathcal{G}) \text{CNRS}(\mathcal{G}) + \text{CN}(\mathcal{G})^2. \end{aligned}$$

Therefore,

$$(\text{CNL}(\mathcal{G}))_{i,i}^2 = (\text{CNRS}(\mathcal{G}))_{i,i}^2 + (\text{CN}(\mathcal{G}))_{i,i}^2 = (\text{CNRS}(\mathcal{G})_{i,i})^2 + \sum_{k=1, k \neq i}^{|\mathcal{V}(\mathcal{G})|} (\text{CN}(\mathcal{G})_{i,k})^2.$$

Hence,

$$(M^2)_{i,i} = \left(\text{CNRS}(\mathcal{G})_{i,i} - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|\mathcal{V}(\mathcal{G})|} \right)^2 + \sum_{k=1, k \neq i}^{|\mathcal{V}(\mathcal{G})|} (\text{CN}(\mathcal{G})_{i,k})^2.$$

Since $\text{tr}(\text{CNRS}(\mathcal{G})) = M_1(\mathcal{G}) - 2|e(\mathcal{G})|$, by Lemma 5.7, we obtain

$$E(M) \leq \sum_{i=1}^{|\mathcal{V}(\mathcal{G})|} \sqrt{\left(\text{CNRS}(\mathcal{G})_{i,i} - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|\mathcal{V}(\mathcal{G})|} \right)^2 + \sum_{k=1, k \neq i}^{|\mathcal{V}(\mathcal{G})|} |N_{\mathcal{G}}(v_i) \cap N_{\mathcal{G}}(v_k)|^2}.$$

Hence the result follows noting that

$$LE_{CN}(\mathcal{G}) = E(M).$$

(b) Let $N = \text{CNSL}(\mathcal{G}) - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} I_{|V(\mathcal{G})|}$, where $I_{|V(\mathcal{G})|}$ is the identity matrix of size $|V(\mathcal{G})|$. Then $\text{Spec}(N) = \left\{ \sigma_i - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} : 1 \leq i \leq |V(\mathcal{G})| \right\}$, where $\text{CNSL-spec}(\mathcal{G}) = \{\sigma_1, \sigma_2, \dots, \sigma_{|V(\mathcal{G})|}\}$. We have

$$\begin{aligned} N^2 &= \left(\text{CNSL}(\mathcal{G}) - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} I_{|V(\mathcal{G})|} \right)^2 \\ &= (\text{CNSL}(\mathcal{G}))^2 - \frac{2\text{tr}(\text{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} \text{CNSL}(\mathcal{G}) + \frac{(\text{tr}(\text{CNRS}(\mathcal{G})))^2}{|V(\mathcal{G})|^2} I_{|V(\mathcal{G})|}. \end{aligned}$$

Therefore the i -th diagonal element of N^2 is

$$(N^2)_{i,i} = (\text{CNSL}(\mathcal{G}))_{i,i}^2 - \frac{2\text{tr}(\text{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} (\text{CNRS}(\mathcal{G}))_{i,i} + \frac{(\text{tr}(\text{CNRS}(\mathcal{G})))^2}{|V(\mathcal{G})|^2}.$$

We have

$$\begin{aligned} (\text{CNSL}(\mathcal{G}))^2 &= (\text{CNRS}(\mathcal{G}) + \text{CN}(\mathcal{G}))^2 \\ &= \text{CNRS}(\mathcal{G})^2 + \text{CNRS}(\mathcal{G}) \text{CN}(\mathcal{G}) + \text{CN}(\mathcal{G}) \text{CNRS}(\mathcal{G}) + \text{CN}(\mathcal{G})^2. \end{aligned}$$

Therefore,

$$(\text{CNSL}(\mathcal{G}))_{i,i}^2 = (\text{CNRS}(\mathcal{G}))_{i,i}^2 + \sum_{k=1}^{|V(\mathcal{G})|} (\text{CN}(\mathcal{G}))_{i,k}^2.$$

Hence,

$$(N^2)_{i,i} = \left(\text{CNRS}(\mathcal{G})_{i,i} - \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|} \right)^2 + \sum_{k=1}^{|V(\mathcal{G})|} (\text{CN}(\mathcal{G}))_{i,k}^2.$$

Since $\text{tr}(\text{CNRS}(\mathcal{G})) = M_1(\mathcal{G}) - 2|e(\mathcal{G})|$, by Lemma 5.7, we have

$$E(N) \leq \sum_{i=1}^{|V(\mathcal{G})|} \sqrt{\left(\text{CNRS}(\mathcal{G})_{i,i} - \frac{M_1(\mathcal{G}) - 2|e(\mathcal{G})|}{|V(\mathcal{G})|} \right)^2 + \sum_{k=1}^{|V(\mathcal{G})|} |N_{\mathcal{G}}(v_i) \cap N_{\mathcal{G}}(v_k)|^2}.$$

Hence, the result follows noting that

$$LE_{CN}^+(\mathcal{G}) = E(N).$$

□

Theorem 5.9. *If \mathcal{G} is a r -regular graph of order $|V(\mathcal{G})|$, then $LE_{CN}(\mathcal{G})$ and $LE_{CN}^+(\mathcal{G})$ are bounded by*

$$\sum_{i=1}^{|V(\mathcal{G})|} \sqrt{\sum_{k=1}^{|V(\mathcal{G})|} |N_{\mathcal{G}}(v_i) \cap N_{\mathcal{G}}(v_k)|^2}.$$

Proof. Since \mathcal{G} is a regular graph, by Lemma 2.2, we obtain

$$\begin{aligned} \text{tr}(\text{CNRS}(\mathcal{G})) &= \sum_{i=1}^{|V(\mathcal{G})|} \sum_{j=1, j \neq i}^{|V(\mathcal{G})|} |N_{\mathcal{G}}(v_i) \cap N_{\mathcal{G}}(v_j)| \\ &= \sum_{i=1}^{|V(\mathcal{G})|} d_{\mathcal{G}}(v_i) (m_{\mathcal{G}}(v_i) - 1) \\ &= \sum_{i=1}^{|V(\mathcal{G})|} r(r-1) = |V(\mathcal{G})| r(r-1) \end{aligned}$$

and

$$\text{CNRS}(\mathcal{G})_{i,i} = r(r-1) = \frac{\text{tr}(\text{CNRS}(\mathcal{G}))}{|V(\mathcal{G})|}, \quad 1 \leq i \leq |V(\mathcal{G})|.$$

From Theorem 5.8, we get the result. □

Acknowledgement

F. E. Jannat is supported by DST INSPIRE Fellowship (IF200226). K. C. Das is supported by National Research Foundation funded by the Korean government (Grant No. 2021R1F1A1050646).

Competing Interests: The authors declare no conflict of interest.

References

- [1] A. Alwardi, N. D. Soner, I. Gutman, On the common-neighborhood energy of a graph, *Bulletin (Académie Serbe Des Sciences Et Des Arts. Classe Des Sciences Mathématiques Et Naturelles. Sciences Mathématiques)*, **36**, 49–59, 2011.
- [2] X. Chen, Y. Hou, A sharp lower bound on the least signless Laplacian eigenvalue of a graph, *Bulletin Malaysian Math. Sci. Soc.*, **41**, 2011–2018, 2018.

- [3] B. Borovićanin, K. C. Das, B. Furtula, I. Gutman, Bounds for Zagreb indices, *MATCH Commun. Math. Comput. Chem.*, **78**, 17–100, 2017.
- [4] K. C. Das, An improved upper bound for Laplacian graph eigenvalues, *Linear Algebra Appl.*, **368**, 269–278, 2003.
- [5] K. C. Das, Maximizing the sum of the squares of the degrees of a graph, *Discrete Math.*, **285**, 57–66, 2004.
- [6] K. C. Das, Sharp bounds for the sum of the squares of the degrees of a graph, *Kragujevac J. Math.*, **25**, 31–49, 2003.
- [7] K. C. Das, On Comparing Zagreb Indices of Graphs, *MATCH Commun. Math. Comput. Chem.*, **63**, 433–440, 2010.
- [8] K. C. Das, I. Gutman, Some properties of the Second Zagreb Index, *MATCH Commun. Math. Comput. Chem.*, **52**, 103–112, 2004.
- [9] K. C. Das, I. Gutman, B. Horoldagva, Comparison between Zagreb indices and Zagreb coindices, *MATCH Commun. Math. Comput. Chem.*, **68**, 189–198, 2012.
- [10] K. C. Das, I. Gutman, B. Zhou, New upper bounds on Zagreb indices, *J. Math. Chem.*, **46**, 514–521. 2009.
- [11] K. C. Das, S. A. Mojallal, On Laplacian energy of graphs, *Discrete Math.*, **325**, 52-64, 2014.
- [12] K. C. Das, S. A. Mojallal, I. Gutman, On energy of line graphs, *Linear Algebra Appl.*, **499**, 79-89, 2016.
- [13] K. C. Das, S. A. Mojallal, I. Gutman, Relations between degrees, conjugate degrees and graph energies, *Linear Algebra Appl.*, **515**, 24-37, 2017.
- [14] K. C. Das, S. A. Mojallal, I. Gutman, On energy and Laplacian energy of bipartite graphs. *Applied Math. Comput.*, **273**, 759–766, 2016.

- [15] W. N. T. Fafous, R. K. Nath, R. Sharafdini, Various spectra and energies of commuting graphs of finite rings, *Hacet. J. Math. Stat.*, **49**(6), 1915–1925, 2020.
- [16] R. Grone, R. Merris, The Laplacian spectrum of a graph II, *SIAM J. Discrete Math.*, **7**(2), 221–229, 1994.
- [17] I. Gutman, The energy of a graph, *Berichte der Mathematisch-Statistischen Sektion im Forschungszentrum Graz*, **103**, 1–22, 1978.
- [18] I. Gutman, Hyperenergetic molecular graphs, *J. Serb. Chem. Soc.*, **64**, 199–205, 1999.
- [19] I. Gutman, N. M. M. Abreu, C. T. M. Vinagre, A. S. Bonifácio, S. Radenković, Relation between energy and Laplacian energy, *MATCH Commun. Math. Comput. Chem.*, **59**, 343–354, 2008.
- [20] I. Gutman, B. Furtula, Survey of Graph Energies *Mathematics Interdisciplinary Research*, **2**, 85–129, 2017.
- [21] I. Gutman, B. Furtula, Graph energies and their applications, *Bulletin (Académie Serbe Des Sciences Et Des Arts. Classe Des Sciences Mathématiques Et Naturelles. Sciences Mathématiques)*, **44**, 29–45, 2019.
- [22] I. Gutman, B. Zhou, Laplacian energy of a graph, *Linear Algebra Appl.*, **414**, 29–37, 2006.
- [23] M. Liu, G. Zhang, K. C. Das, The maximum number of spanning trees of a graph with given matching number, *Bulletin Malaysian Math. Sci. Soc.*, **44**, 3725–3732, 2021.
- [24] A. W. Marshall, I. Olkin, *Inequalities: Theory of Majorization and Its Applications*, Academic Press, New York, 1979.
- [25] R. K. Nath, W. N. T. Fafous, K. C. Das, Y. Shang, Common neighborhood energy of commuting graphs of finite groups, *Symmetry*, **13** (9), (2021) 1651.

- [26] B. R. Rakshith, K. C. Das, B. J. Manjunatha, Some new families of noncorona graphs with strong anti-reciprocal eigenvalue property, *Bulletin Malaysian Math. Sci. Soc.*, **45**, 2597–2618, 2022. .
- [27] I. Schur, Über eine klasse von Mittelbildungen mit Anwendungen die Determinanten, *Theorie Sitzungsber. Berlin. Math. Gesellschaft*, **22**, 9–20, 1923.
- [28] H. B. Walikar, H. S. Ramane, P. R. Hampiholi, On the energy of a graph, *Graph connections*, Eds. R. Balakrishnan, H. M. Mulder and A. Vijayakumar, Allied publishers, New Delhi, 120–123, 1999.