

Faulhaber's formula, Bernoulli numbers and the equation

$$f(x) + x^k = f(x + 1)$$

Chai Wah Wu

IBM Research

IBM T. J. Watson Research Center, Yorktown Heights, NY, USA*

March 1, 2024

Abstract

In modern usage the Bernoulli numbers and polynomials follow Euler's approach and are defined using generating functions. We consider the functional equation $f(x) + x^k = f(x + 1)$ and show that a solution can be derived from Faulhaber's formula for the sum of powers. We show how these solutions provide a characterization of Bernoulli numbers and related results.

1 Introduction

We consider the functional equation described by

$$f(x) + x^k = f(x + 1) \tag{1}$$

A purpose of this brief note is to show that Faulhaber's formula for the sums of powers (which initiated the study of Bernoulli numbers) can be used to derive a solution to this equation and leads to a characterization of Bernoulli numbers, Bernoulli polynomials and related identities.

2 Faulhaber's formula

Faulhaber's formula (also known as Bernoulli's formula) equates the sum of powers of consecutive integers to a polynomial involving Bernoulli numbers. It was discovered by various people including Takanaga, Faulhaber, and Bernoulli. In particular, for n and p nonnegative integers,

$$F(n, p) = \sum_{i=1}^n i^p = \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^+ n^{p+1-j} \tag{2}$$

where B_j^+ are the Bernoulli numbers of the second kind. There are in general two conventions when defining Bernoulli numbers, with the only difference being the sign of B_1 . In particular, $B_1^+ = \frac{1}{2}$, whereas the Bernoulli numbers of the first kind B_j^- are identical to B_j^+ except for $B_1^- = -\frac{1}{2}$. B_j^- has been used in modern textbooks although recently there are discussions¹ on whether B_j^+ , which was used in the past, should be favored. Computer algebra software such as **Mathematica** currently use B_j^- to denote Bernoulli

*cwwu@us.ibm.com

¹<https://luschny.de/math/zeta/The-Bernoulli-Manifesto.html>, <https://www-cs-faculty.stanford.edu/~knuth/news22.html>

numbers. However, the `bernoulli` function in `Sympy` which until very recently also returns B_j^- has changed from returning B_j^- to B_j^+ starting from version 1.12. Note that the number of terms in the left hand side is n while it is $p + 1$ on the right hand side, meaning that the computation of $F(n, p)$ is more efficient using the right hand side when $n \gg p$.

The Bernoulli numbers are defined in modern usage via the generating functions [1]:

$$\sum_{j=0}^{\infty} \frac{B_j^- x^j}{j!} = \frac{x}{e^x - 1} \quad (3)$$

$$\sum_{j=0}^{\infty} \frac{B_j^+ x^j}{j!} = \frac{x}{1 - e^{-x}} \quad (4)$$

They also satisfy the following recursive definitions [1]:

Lemma 1.

$$\sum_{k=0}^m \binom{m+1}{k} B_k^- = \delta_{m,0} \quad (5)$$

$$\sum_{k=0}^m \binom{m+1}{k} B_k^+ = m + 1 \quad (6)$$

Where $\delta_{m,0}$ is the Kronecker delta function defined as 1 if $m = 0$ and 0 otherwise.

2.1 A variant of Faulhaber's formula

Eq. (2) expresses the sum of powers $F(n, p)$ using Bernoulli numbers of the second kind B_j^+ . The following formula gives a variant of Faulhaber's formula using Bernoulli numbers of the first kind B_j^- [2].

Theorem 1. *If $p > 0$, then*

$$F(n, p) = \sum_{i=1}^n i^p = \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^- (n+1)^{p+1-j} \quad (7)$$

Proof. By Eq. (2),

$$F(n+1, p) = F(n, p) + (n+1)^p = \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^+ (n+1)^{p+1-j}$$

For $j = 1$, the corresponding term is $\frac{1}{2}(n+1)^p$ whereas $\frac{1}{p+1} \binom{p+1}{1} B_1^- (n+1)^p = \frac{1}{2}(n+1)^p$. This means that $F(n+1, p) = (n+1)^p + \frac{1}{p+1} \sum_{j=0}^p \binom{p+1}{j} B_j^- (n+1)^{p+1-j}$ and the conclusion follows. \square

Note that the formula in Eq. (7) is almost identical to the formula in Eq. (2) except for using B_j^- rather than B_j^+ and shifting the base of the powers from n to $n+1$.

3 A solution to the equation $f(x) + x^k = f(x+1)$

Define $f_k(x) = \frac{1}{k+1} \sum_{j=0}^k \binom{k+1}{j} B_j^- x^{k+1-j}$, then the above discussion shows that $f_k(x+1) = f_k(x) + x^k$. Note that $(k+1)f_k(x) = B_{k+1}(x)$, the $k+1$ -th Bernoulli polynomial.² We show that the converse is also true among polynomials in the following sense.

²This implies the following property of Bernoulli polynomials $B_k(x+1) - B_k(x) = kx^{k-1}$, see e.g. [2, 3].

Suppose that p is a minimal degree polynomial that satisfied Eq. (1). Clearly p must have degree larger than or equal to $k - 1$. If p has degree k , then the coefficient of x^k does not match in Eq. (1). Therefore, p must be of degree $k + 1$ or larger. Consider the case where p has degree $k + 1$ written as $p(x) = \sum_{j=0}^{k+1} a_j x^j$. Then Eq. (1) can be written as $\sum_{j=0}^{k+1} a_j x^j + x^k = \sum_{j=0}^{k+1} a_j (x + 1)^j$. Matching the coefficients results in the following $k + 2$ equations $a_j = \sum_{l=0}^{k+1} a_l \binom{l}{j} = \sum_{l=j}^{k+1} a_l \binom{l}{j}$ for $j \neq k$ and $a_k + 1 = \sum_{l=0}^{k+1} a_l \binom{l}{k}$. This last equation can be simplified as: $a_k + 1 = a_k + a_{k+1}(k + 1)$, i.e. $a_{k+1} = \frac{1}{k+1}$. The equation for $j = k + 1$ is the identity $a_{k+1} = a_{k+1}$. The remaining k equations $a_j = \sum_{l=j}^{k+1} a_l \binom{l}{j}$ for $j < k$ uniquely determines the other coefficients a_j . In particular, $a_j = a_j + a_{j+1} \binom{j+1}{j} + \sum_{l=j+2}^{k+1} a_l \binom{l}{j}$, i.e. for $j \leq k$,

$$a_j = -\frac{\sum_{l=j+1}^{k+1} a_l \binom{l}{j-1}}{\binom{j}{j-1}} = -\frac{\sum_{l=j+1}^{k+1} a_l \binom{l}{j-1}}{j}$$

This implies that $a_k = -a_{k+1} \binom{k+1}{k-1} / k = -\frac{1}{2}$.

Let $b_{k+1-j} = (k + 1)a_j / \binom{k+1}{j}$ or equivalently $b_j = (k + 1)a_{k+1-j} / \binom{k+1}{j}$. This implies that $b_0 = 1$ and $b_1 = -\frac{1}{2}$. After straightforward algebraic manipulations we get the following identity regarding b_j 's: $b_j = -\frac{\sum_{w=0}^{j-1} b_w \binom{j+1}{w}}{j+1}$ for $0 < j \leq k + 1$ which is exactly the recursive definition of the Benoulli number B_j^- in Lemma 1. Thus we have shown that

Proposition 1. *The first $k + 2$ Bernoulli numbers B_j^- are defined by the coefficients a_j of the minimal degree polynomial that satisfies the functional equation $f(x) + x^k = f(x + 1)$ via the equation $B_j^- = (k + 1)a_{k+1-j} / \binom{k+1}{j}$.*

Define $g(x) = \frac{1}{k+1} \sum_{j=0}^k \binom{k+1}{j} B_j^+ x^{k+1-j}$, a $(k + 1)^{th}$ -degree polynomial with rational coefficients. Then Eqns 2-7 show that $g(x) + (x + 1)^k = g(x + 1)$. An analogous derivation shows that

Proposition 2. *The first $k + 2$ Bernoulli numbers B_j^+ are defined by the coefficients a_j of the minimal degree polynomial that satisfies the functional equation $f(x) + (x + 1)^k = f(x + 1)$ via the equation $B_j^+ = (k + 1)a_{k+1-j} / \binom{k+1}{j}$.*

Since the Bernoulli polynomials $B_k(x)$ form an Appell sequence [4], i.e., it satisfies the differential equation $\frac{d}{dx} B_k(x) = k B_{k-1}(x)$, it is clear that f_k is an Appell sequence as well. This implies that it satisfies Appell's identity: $f_k(x + y) = \sum_{i=0}^k \binom{k}{i} f_i(x) y^{k-i}$, i.e. $f_k(x) + x^k = f_k(x + 1) = \sum_{i=0}^k \binom{k}{i} f_i(x)$. This simplifies to the equation³ $\sum_{i=0}^{k-1} \binom{k}{i} f_i(x) = x^k$.

References

- [1] Wikipedia, "Bernoulli number — Wikipedia, the free encyclopedia." <http://en.wikipedia.org/w/index.php?title=Bernoulli%20number&oldid=1198154404>, 2024. [Online; accessed 01-February-2024].
- [2] N. D. Larson, *The Bernoulli Numbers: A Brief Primer*. Whitman College, 2019.
- [3] K. Ireland and M. Rosen, *A classical introduction to modern number theory*, vol. 84 of *Graduate texts in mathematics*. Springer, 1982.
- [4] Wikipedia, "Appell sequence — Wikipedia, the free encyclopedia." <http://en.wikipedia.org/w/index.php?title=Appell%20sequence&oldid=1194111356>, 2024. [Online; accessed 01-February-2024].

³This equation can also be derived from the well-known identity $kx^{k-1} = \sum_{i=1}^k \binom{k}{i} B_{k-i}(x)$.

- [5] D. E. Knuth, “Johann Faulhaber and sums of powers,” *Mathematics of Computation*, vol. 61, pp. 277–294, July 1993.
- [6] A. F. Beardon, “Sums of powers of integers,” *The American Mathematical Monthly*, vol. 103, no. 3, pp. 201–213, 1996.
- [7] P. H. N. Luschny, “An introduction to the Bernoulli function.” arXiv:2009.06743, 2021.
- [8] B. C. Kellner, “Faulhaber polynomials and reciprocal Bernoulli polynomials,” *Rocky Mountain J. Math.*, vol. 53, pp. 119–151, 2023.