Research in Astronomy and **A**strophysics

Key words: galaxies: evolution–galaxies: interactions–galaxies: star formation–galaxies: structure– galaxies: bulges–galaxies: photometry

Close Major-merger Pairs at $z = 0$: Star-forming Galaxies with Pseudobulges

Chuan He (0000-0003-1761-5442, hechuan@nao.cas.cn)^{1,2}, Cong Kevin Xu (0000-0002-1588-6700, congxu@nao.cas.cn)¹, Ute Lisenfeld (0000-0002-9471-5423)^{3,4}, Yu Sophia Dai (0000-0002-7928-416X)¹, Taotao Fang (0000-0002-2853-3808)⁵, Jiasheng Huang $(0000-0001-6511-8745)^1$, Wei Wang^{1,6} and Qingzheng Yu (0000-0003-3230-3981)⁵

- ² School of Astronomy and Space Sciences, University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China
- ³ Departamento de Física Teórica y del Cosmos, Universidad de Granada, 18071 Granada, Spain
- 4 Instituto Carlos I de Física Téorica y Computacional, Facultad de Ciencias, 18071 Granada, Spain
- ⁵ Department of Astronomy, Xiamen University, Xiamen, Fujian 361005, People's Republic of China
⁶ CAS Key Laboratory of Ontical Astronomy, National Astronomical Observatories, Chinese
- ⁶ CAS Key Laboratory of Optical Astronomy, National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100101, China

Received 08-Feb-2024; revised 20-Mar-2024; accepted 28-Mar-2024

Abstract We present a study of star-forming galaxies (SFGs) with pseudobulges (bulges with Sérsic index $n < 2$) in a local close major-merger galaxy pair sample (H-KPAIR). With data from new aperture photometries in the optical and near-infrared bands (aperture size of 7 kpc) and from the literature, we find that the mean Age of central stellar populations in Spirals with pseudobulges is consistent with that of disky galaxies and is nearly constant against the bulge-to-total ratio (B/T). Paired Spirals have a slightly lower fraction of pure disk galaxies ($B/T \leq 0.1$) than their counterparts in the control sample. Compared to SFGs with classical bulges, those with pseudobulges have a higher ($> 2 \sigma$) mean of specific star formation rate (sSFR) enhancement ($\text{sSFR}_{\text{enh}} = 0.33 \pm 0.07$ vs sSFR_{enh} = 0.12 ± 0.06) and broader scatter (by \sim 1 dex). The eight SFGs that have the highest SFR_{enh} in the sample all have pseudobulges. A majority (69%) of paired SFGs with strong enhancement (having sSFR more than 5 times the median of the control galaxies) have pseudobulges. The Spitzer data show that the pseudobulges in these galaxies are tightly linked to nuclear/circum-nuclear starbursts. Pseudobulge SFGs in S+S and in S+E pairs have significantly ($> 3\sigma$) different sSFR enhancement, with the means of $\text{sSFR}_{\text{enh}} = 0.45 \pm 0.08$ and -0.04 ± 0.11 , respectively. We find a decrease in the sSFR enhancements with the density of the environment for SFGs with pseudobulges. Since a high fraction (5/11) of pseudobulge SFGs in S+E pairs are in rich groups/clusters (local density $N_{1 \text{Mpc}} \geq 7$), the dense environment might be the cause for their low sSFR_{enh}.

¹ Chinese Academy of Sciences South America Center for Astronomy, National Astronomical Observatories of China, Chinese Academy of Sciences, Beijing 100101, People's Republic of China

1 INTRODUCTION

Broad and intensive studies have shown that galaxy interaction can induce star formation enhancement [\(Toomre & Toomre 1972;](#page-23-0) [Larson & Tinsley 1978](#page-22-0); [Kennicutt](#page-22-1) et al. [1987;](#page-22-1) [Xu & Sulentic 1991](#page-23-1); [Sanders & Mirabel 1996\)](#page-23-2). Both simulations and observations have indicated that the existence and intensity of such enhancement is affected by many factors of the pair such as separation distance [\(Alonso et al. 2004](#page-22-2); [Nikolic et al. 2004\)](#page-22-3), mass ratio [\(Cox et al. 2008\)](#page-22-4), environment [\(Ellison et al. 2010](#page-22-5)), interaction phase [\(Scudder et al. 2012\)](#page-23-3), orbital parameters [\(Patton et al. 2013\)](#page-23-4). Meanwhile, some factors of the paired galaxies such as stellar mass [\(Rodighiero](#page-23-5) et al. [2011](#page-23-5)), redshift [\(Madau & Dickinson](#page-22-6) [2014\)](#page-22-6) and morphology [\(Kennicutt 1998](#page-22-7)) do affect the star formation behavior themselves, too. On the other hand, many quenching mechanisms are reported such as mass [\(Peng et al. 2010\)](#page-23-6), environment [\(Peng et al. 2012\)](#page-23-7), AGN feedback [\(Kormendy & Ho 2013\)](#page-22-8), and bulge-to-total ratios (B/T, [Martig et al.](#page-22-9) [2009;](#page-22-9) [Bluck et al. 2014\)](#page-22-10). The combining effect of many factors especially when studying a big sample with a large range of parameters may bury potential results or even cause statistical bias. This inspires us to delve into the fundamental mechanism of the interaction-induced star formation. For instance, by investigating the kinematic asymmetry utilizing the recent IFS (integral field spectrograph) technique, [Feng et al.](#page-22-11) [\(2020\)](#page-22-11) reveal that the strength of the ongoing tidal effect during certain merging phases is a more basic indicator to understanding the intermittent star formation enhancement in galaxy pairs than projected separations. Also, the SFR level is influenced by both the quantity of gas and its efficiency in forming stars. [Lisenfeld et al. \(2019\)](#page-22-12) reveals that mergers in an earlier stage (i.e. pair) can take a lead in transferring atom gas to molecular gas, which makes the paired sample has a higher SFE when calculating the total gas, however, no significant SFE enhancement is found in their pair sample when calculating the molecular gas. On the other hand, the later stage mergers, who always show their strong strength on star formation activities by appearing as a ULIRG, can pool a large amount of molecular gas (e.g., [Sargent & Scoville 1991;](#page-23-8) [Scoville et al. 1991\)](#page-23-9). By controlling the total gas content, [Li et al.](#page-22-13) [\(2023\)](#page-22-13) found no significant enhancement on SFE in paired galaxies compared to the single galaxies. To conclude, it is still controversial whether and how galaxy interaction can induce star formation enhancement. To acquire solid evidence, one needs to select the sample and their controls carefully, and a detailed classification of their sample as well.

Far-infrared (FIR) observations by Spitzer [\(Xu et al. 2010\)](#page-23-10) and Herschel [\(Cao et al. 2016\)](#page-22-14) on a complete, unbiased close major-merger pairs sample selected from K_s-band [Domingue et al.](#page-22-15) [\(2009\)](#page-22-15) suggest that only SFGs in spiral-spiral (hereafter S+S) pairs have significantly enhanced specific star formation rate ($sSFR = SFR/Mstar$), but not those in Spiral-Elliptical (hereafter S+E) pairs. This situation is also presented in [Park & Choi \(2009](#page-23-11)); [Hwang et al. \(2010](#page-22-16)); [Xu et al.](#page-23-10) [\(2010\)](#page-23-10); [Moon et al.](#page-22-17) [\(2019\)](#page-22-17). [He et al. \(2022\)](#page-22-18) studied the dependence of the interaction-induced specific star formation rate enhancement ($\text{ssFR}_{enh} = \log(\text{sSFR}_{pg}) - \log(\text{sSFR}_{med,ctrl})$ where "pg" stands for paired galaxy and "med,ctrl" for the median of their control galaxies) on the bulge-to-total ratio (B/T). They found a negative dependence of the interaction-induced sSFR enhancement on the bulge-to-total ratio (B/T) and a significant ($> 5\sigma$) enhancement only in paired SFGs with B/T < 0.3 . This is consistent with the results of theoretical simulations which predicted that a massive bulge can stabilize the disk and suppress the SFR during and after close encounters [\(Mihos & Hernquist](#page-22-19) [1996;](#page-22-19) [Cox et al. 2008;](#page-22-4) [Di Matteo et al.](#page-22-20) [2008\)](#page-22-20). However, it appears that SFGs with low B/T ratios, in particular the disky galaxies (B/T \leq 0.1), have very diversified sSFR_{enh} with the value varying in a range of \sim 2.5 orders of magnitude, and even some of such SFGs have sSFR deficit.

In their analysis, [He et al. \(2022\)](#page-22-18) separated pseudobulges (with Sérsic index $n < 2$) from classical bulges (with Sérsic index $n \ge 2$) and assigned $B/T = 0$ to galaxies with pseudobulges. This is because many pseudobulges found in two-component (bulge and disk) models (such as those in [He et al.](#page-22-18) [2022\)](#page-22-18) are misidentified bars, nuclear disks, and nuclear rings [\(Kormendy & Kennicutt 2004\)](#page-22-21) which themselves may be triggered by interaction [\(Chown et al. 2019;](#page-22-22) [Erwin et al. 2021\)](#page-22-23). Meanwhile, the pseudobulges unrelated to nuclear star formation are mostly found in late-type spirals with relatively low B/T ratios [\(Kim et al. 2016\)](#page-22-24), and assigning them to disky galaxies shall not introduce strong bias. However, bars and nuclear rings often have intermittent star formation activity and, when being caught in the "off" phase, may appear to be SFR-quenched [\(Fraser-McKelvie et al. 2020\)](#page-22-25). Also, some massive early-type spirals (such as S0 and Sa galaxies) with low gas content and low SFR may have large bars [\(Herrera-Endoqui et al. 2017](#page-22-26)).

Can the broader scatter of the sSFRenh among disky SFGs be due to the diversity of galaxies with pseudobulges? The major scientific goal of this paper is to answer the above question and to investigate the roles played by pseudobulge SFGs regarding sSFR enhancement in paired galaxies. Different from [He et al. \(2022](#page-22-18)), for galaxies with pseudobulges, we use the original B/T values instead of assigning $B/T = 0$. The SFG sample we use in this paper is introduced in Sect. [2.](#page-2-0) In Sect. [3](#page-2-1) we present a study of the optical color-color diagrams measured in the inner part of the paired galaxies, particularly for galaxies with pseudobulges to separate those with bars, nuclear disks, and nuclear rings from normal pseudobulges of $n < 2$. In Sect. [4](#page-3-0) we present our science analyses and results. A discussion is carried out in Sect. [5.](#page-13-0) We conclude our paper in Sect. [6.](#page-16-0) Throughout this paper, we adopt the Λ-cosmology with $\Omega_{\rm m}=0.3$ and $\Omega_{\Lambda}=0.7$, and $\rm H_{0}=70$ km s⁻¹ Mpc⁻¹.

2 THE SAMPLE

The sample we use in this paper is the same as in [He et al.](#page-22-18) [\(2022\)](#page-22-18), which is the local close majormerger sample H-KPAIR [\(Cao et al. 2016](#page-22-14)). H-KPAIR is a subsample of the KPAIR which is a complete and unbiased K_s -band (Two Micron All Sky Survey, 2MASS) selected pair sample [\(Domingue et al.](#page-22-15) [2009\)](#page-22-15). The H-KPAIR includes 44 Spiral+Spiral (S+S) pairs and 44 Spiral+Elliptical (S+E) pairs, all of which have spectroscopic redshifts in the range of $0.0067 < z < 0.1$, the projected separations in the range of 5 h⁻¹ kpc \leq s(p) \leq 20 h⁻¹ kpc, the radial relative velocity $\delta(\bar{V}z)$ < 500 km s⁻¹ and the K_s -band magnitude differences within 1 mag (corresponding to a mass ratio no greater than 2.5). The strict selection criteria of H-KPAIR gather the most enhanced closed major-merger pairs and makes it an ideal sample to explore the interaction effect to the paired galaxies. All galaxies in H-KPAIR have Herschel imaging observations in the six bands at 70, 110, 160, 250, 350 and 500, and their star formation rates (SFRs) are derived from the Herschel data [\(Cao et al. 2016](#page-22-14)). Seventy pairs have GBT 21cm HI observations [\(Zuo et al. 2018\)](#page-23-12), and 78 spiral galaxies are observed by the IRAM 30m telescope for the CO emissions [\(Lisenfeld et al. 2019](#page-22-12)). The B/T ratios of H-KPAIR galaxies are taken from [He et al. \(2022](#page-22-18)) which are based on 2-D decompositions carried out using the SDSS r -band image. The stellar masses are also taken from [He et al. \(2022](#page-22-18)) who updated the formalism of [\(Cao et al.](#page-22-14) [2016\)](#page-22-14) by including a (g-r) color correction when estimating the mass from the K_s -band luminosity. Throughout this paper, we mainly focus on two subsamples of H-KPAIR: (1) all spiral galaxies Sect. [3](#page-2-1) and [4.1\)](#page-3-1) and (2) all star-forming galaxies (SFGs; Sect. [4.2](#page-7-0) and [4.3\)](#page-12-0). Following [Cao et al. \(2016\)](#page-22-14) and [He et al. \(2022\)](#page-22-18), SFGs are defined by having $\log(\rm sSFR/yr^{-1}) \ge -11.3$. There are 98 SFGs out of 132 Spirals (Table [.1\)](#page-24-0) in the H-KPAIR sample.

3 GALAXY CENTRAL COLORS

We carry out aperture photometry for all galaxies in the H-KPAIR sample using a consistent round aperture in images of the SDSS u, r, i and 2MASS K_s -bands. The u, r, i images are generated using SDSS SAS mosaic tool^{[1](#page-2-2)}, which can stitch together several sky-subtracted, calibrated frames^{[2](#page-2-3)} to form a coherent image over a specified patch of sky using the SWarp [\(Bertin et al. 2002\)](#page-22-27). For the K_s -band, the 2MASS images are resampled into the same pixel scale as the SDSS images of 0.396′′/pix, and the background are subtracted with SWarp. The median Full-Width-at-Half-Maximum (FWHM) of the point spread function (PSF) in the four bands are 1.53″, 1.32″, 1.26″ and 2.9″, respectively.

For each galaxy, we use the same aperture in all four bands which corresponds to a projected diameter of 7 kpc and is centered at its SDSS coordinates. Such a setup is to avoid overlap between apertures of the two member galaxies in a pair, given the minimal pair separation of $5 h^{-1}$ kpc (corresponding approximately to 7 kpc for $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$). Also, the aperture diameters corresponding to

¹ <https://dr12.sdss.org/mosaics/>

² https://data.sdss.org/datamodel/files/BOSS_PHOTOOBJ/frames/RERUN/RUN/CAMCOL/frame.html

7 kpc on our sources range from 4.2′′to 53′′, and most galaxies (141/176) have diameters larger than 2 times 2.9'', which is the worst resolution of the data (i.e. the FWHM = $2.9''$ for the K_s-band). Therefore no aperture correction is needed for any photometry.

The photometric errors of SDSS bands are calculated by combining the Poissonian error of the source flux, the RMS of background, and the calibration error 34 34 . Given the good S/N of our galaxies in the optical bands, errors in these bands are dominated by the calibration error $(1\%$ for the r and i-bands and 2% for the u-band). For the 2MASS K_s -band, in addition to the error sources described above, the coadd noise caused by the resampling and smoothing when generating the Atlas Images is also considered^{[5](#page-3-4)}.

The spiral galaxies in the sample have a median bulge size of $2 \times R_e = 3.6$ kpc (R_e: effective radius of the bulge), and 26% of them have $2 \times R_e \ge 7$ kpc. For the latter galaxies, the colors derived from the 7 kpc aperture photometries provide information on the stellar population within the bulge. For the rest galaxies in the sample, the 7 kpc aperture can include stellar radiations both in the bulge and a small portion of the inner disk, and their relative importance depends on the B/T ratio. As shown in Fig. [1,](#page-4-0) there is a significant correlation between bulge size and B/T (the Spearman's rank correlation coefficient $r_S = 0.55$ and the significance $p_S = 5.6e^{-12}$).

To interpret the colors, predictions of broad-band fluxes are calculated using the Simple Stellar Population (SSP, or instantaneous-burst) model of Bruzual [& Charlot \(2003](#page-22-28)), with the time after the instantaneous-burst τ (i.e. stellar age) and the attenuation A_v as the free parameters. The model uses the Padova (1994) evolution tracks, the Chabrier (2003) IMF with lower and upper mass cutoffs of 0.1–100 M_{\odot} , and the solar metallicity ($Z = 0.02$). The extinctions are calculated following the formalism of [Cardelli et al. \(1989\)](#page-22-29) for the standard diffuse interstellar medium. It is worth noting that the SSP model may be an over-simplification of the real stellar populations in question, and the model predictions on the τ and A_v have large uncertainties. Nevertheless, the model provides a framework to link the colors to the physical properties (e.g. the Age) of the stellar populations which allows us to investigate the difference between galaxies with pseudobulges and classical bulges using color-color diagrams. The results of our aperture photometry and the model fits are listed in Table [.2.](#page-27-0)

4 RESULTS

4.1 Difference between pseudobulges and classical bulges

Two $u-r$ versus i-K_s color-color diagrams are presented in Fig. [2](#page-5-0) and [3,](#page-6-0) with the false color scale representing the B/T ratio and sSFRenh, respectively. The region outlined by the red contour (with a red label "E") in the upper-left corner of the diagram is where the elliptical galaxies (not plotted) are found. Most Spirals located in this region are those with relatively large classical bulges and low SFR_{enh} . The mesh grid in the diagram consists of lines for fixed values of τ and attenuation A_v , respectively, as predicted by the SSP model (see Sect. [3\)](#page-2-1). According to the model, galaxies in the region "E" have old stellar populations of $Age > 1$ Gyr in their central part. For the remaining galaxies, most have their central stellar populations with Ages in the range of [0.1, 1] Gyr and very few with young Ages in the range of [0.01, 0.1] Gyr which are likely associated with nuclear/circum-nuclear starbursts. The range of A_v is between 0 and 3 mag, with most galaxies having 1 mag $\lt A_v \lt 2.5$ mag.

We estimate the Age and A_v for each galaxy via interpolations among the model predictions (in the mesh grid) adjacent to its position in the color-color diagram. Galaxies with $Age > 13$ Gyr are regarded as Age = 13 Gyr and galaxies with $A_v < 0$ are regarded as $A_v = 0$. The plot of A_v versus Age (Fig. [4\)](#page-7-1) shows a strong anti-correlation between the two variables, which is understandable because younger stellar populations are usually associated with stronger star formation activity and higher dust attenuation, and older stellar populations with less star formation activity and dust attenuation. Indeed there are several galaxies with noticeable large classical bulge found to have high Av. We explain this by

³ https://data.sdss.org/datamodel/files/BOSS_PHOTOOBJ/frames/RERUN/RUN/CAMCOL/frame.html

⁴ <https://www.sdss4.org/dr12/algorithms/fluxcal/>

⁵ https://irsa.ipac.caltech.edu/data/2MASS/docs/releases/allsky/doc/sec6_8a.html

Fig. 1: Plot of bulge size $(2 \times R_e)$ versus B/T ratio of all spiral galaxies in H-KPAIR. The horizontal dashed line corresponds to the aperture size of our photometry ($D = 7$ kpc) and a lower cut at $2 \times R_e$ 0.1 kpc (sources with bulge sizes below this line are plotted on it). The filled squares represent galaxies with pseudobulges and the open diamonds galaxies with classical bulges.

two aspects. Firstly, these galaxies have high SFR_{enh} , hence relatively high sSFR, though their control sample shall have low sSFR. The SFR of the sample is derived from Herschel observation [\(Cao et al.](#page-22-14) [2016\)](#page-22-14) and is mainly constrained by the dust temperature, then highly related to the attenuation. Secondly, our $u-r-i-K_s$ color which generated the Av is of the central part of the galaxy instead of the whole galaxy, and old galaxies have dust gathering in the central region (or we can say bulge) is reasonable.

To investigate whether interaction can affect the bulge performance, we select a new control sample (control-sample-2 hereafter). The selection of control-sample-2 is similar to the control sample in [He et al. \(2022,](#page-22-18) , section 4) except for without matching the B/T. The criteria of control-sample-2 go as follows:

- 1. Should be identified as a spiral in Galaxy Zoo [\(Lintott et al. 2008\)](#page-22-30).
- 2. Not in any interacting system, namely no neighbor galaxy in the SDSS database which has projected distance ≤ 100 kpc and observed redshift difference ≤ 1000 km s⁻¹.
- 3. Has reliable B/T ratio, namely having $\chi^2/\nu < 2$ and no bad flag (flag bit 20 = 0) in [Meert et al.](#page-22-31) [\(2015\)](#page-22-31).
- 4. The L_K matches that of the paired galaxy within 0.1 dex
- 5. Match of local density: we adopt a local density indicator $N_{1\text{Mpc}}$, which is the count of galaxies brighter than $M_r = -19.5$ and with redshifts differing less than 1000 km s⁻¹ from that of the target galaxy, in the surrounding sky area of radius $= 1$ Mpc (the count includes the target galaxy itself

Fig. 2: The $u-r-i-K_s$ color-color diagram for Spirals with pseudobulges (filled squares) and classical bulges (open diamonds). The symbol color indicates the B/T ratio in [He et al. \(2022](#page-22-18)). The red contour delineates the region of the distribution of H-KPAIR elliptical galaxies. The mesh grid outlines the time after the instantaneous-burst τ (i.e. stellar age) and attenuation A_v in the model. From bottom to top the dotted lines are τ equal to 0.01, 0.03, 0.04, 0.05, 0.1, 0.3, 1, 13 Gyr, and from left to right the solid lines are A_v of 0, 1, 2, 3 mag.

if it is brighter than $M_r = -19.5$). By means of $N_{1 \text{Mpc}}$, we classify galaxies into 4 environmental categories: field ($N_{1\text{Mpc}} \leq 3$), small group ($4 \leq N_{1\text{Mpc}} \leq 6$), large group ($7 \leq N_{1\text{Mpc}} \leq 10$), and cluster ($N_{1Mpc} > 10$). The control galaxy shall be in the same environmental category as the paired galaxy.

6. Has the closest redshift, among all qualified candidates, to that of the paired galaxy.

In Fig. [5](#page-8-0) we compare the Age and B/T ratio of the paired galaxies which are separated into two subsamples: (1) galaxies with pseudobulges and (2) those with classical bulges. Also plotted are the means of the Age for Spirals in H-KPAIR/control-sample-2 with error bars in the five B/T bins for the two subsamples: (1) $B/T \le 0.1$, (2) $0.1 < B/T \le 0.3$, (3) $0.3 < B/T \le 0.5$, (4) $0.5 < B/T \le 0.7$, (5) $B/T > 0.7$. It shows that, on average, the Age of the central stellar population of galaxies with classical bulges increases with the B/T ratio. This is indeed anticipated because the dominance of the bulge to central colors increases with the B/T ratio (Fig. [1\)](#page-4-0), and in general stellar populations in classical bulges are older than those in disks [\(Driver et al. 2006;](#page-22-32) [Kim et al. 2016\)](#page-22-24). In contrast, for galaxies with pseudobulges, the Age of central stellar populations is rather flat against the B/T ratio, namely the stellar populations have about the same Ages as those in the inner disks of the disky galaxies (with $B/T \sim 0$). This confirms that pseudobulges are different from classical bulges, and the former are

Fig. 3: Same $u-r-i-K_s$ color-color diagram as Figure [2,](#page-5-0) but the color indicates the $\rm sSFR_{enh}$. The data points marked by crosses are SFR undetected in [Cao et al. \(2016\)](#page-22-14)

dominated by the "disk phenomena" such as bars and nuclear disks/rings as assumed in [He et al. \(2022](#page-22-18)). However, some individual galaxies with pseudobulges have central stellar populations with old Ages of > 1 Gyr. Their pseudobulges are unlikely related to any central star formation activity, as indicated by the low star formation enhancement ($sSFR_{\text{enh}} \lesssim 0$). They have relatively low B/T ratios (most with $B/T < 0.3$, all with $B/T < 0.5$) and may be similar to those single late-type galaxies with relatively small pseudobulges found in [Kim et al. \(2016\)](#page-22-24).

It is also interesting to note that a few galaxies with large classical bulges are located in the lower-right quadrant of Fig. [5,](#page-8-0) namely their central stellar populations appear to have relatively young Ages. They may contain starbursts or post-starbursts in the central region which can outshine the old stellar populations and make the bulges "blue". An outstanding example of these galaxies is J13151726+4424255, which has a classical bulge with the $B/T = 0.94$, the Age = 0.11 Gyr, and the $\text{sSFR}_{\text{enh}} = 0.70$. Compared to galaxies in the control sample, paired galaxies with large bulges, particularly those in the forth bin of the B/T ratio ($0.5 < B/T \le 0.7$), have higher Ages. Individual inspections of these galaxies indicate that most of them have elliptical companions (i.e. in SE pairs). Their star formation history might have been affected by the environment associated with the Ellipticals (see the discussion in Section 5).

Fig. [6](#page-9-0) and [7](#page-10-0) show the possibility density function (pdf) of classical bulges and pseudobulges in the five bins for SFGs in H-KPAIR and control-sample-2, respectively. The paired Spirals have lower fraction of pure disk galaxies (bin1, $B/T \leq 0.1$), we attribute this to some paired disky galaxies having nuclear/circum-nuclear starburst and their B/T ratio raising to higher B/T bins. Nevertheless, these

Fig. 4: Plot of the attenuation A_v vs. age of the central stellar population for all spiral galaxies in the H-KPAIR. The color of symbols represents the SFR_{enh} , and the size of symbols represents the B/T ratio. The filled squares and the open diamonds represent pseudobulge galaxies and classical galaxies, respectively. The data points marked by crosses are SFR undetected in [Cao et al.](#page-22-14) [\(2016\)](#page-22-14)

galaxies have a flat profile and is identified as pseudobulge galaxies and are assigned $B/T = 0$ in [He et al. \(2022\)](#page-22-18)

4.2 Paired SFGs with strong enhancement

In Sect. [2](#page-2-0) we defined SFGs by $\log(\rm sSFR/yr^{-1}) \ge -11.3$ (see also [Cao et al. 2016;](#page-22-14) [He et al. 2022](#page-22-18)). Most of the non-SFGs are SFR undetected (marked by crosses in Figure [2,](#page-5-0) [4](#page-7-1) and [5\)](#page-8-0) in [Cao et al. \(2016](#page-22-14)), and they are mostly early-type galaxies with large old classical bulge. Taking these galaxies into the sSFRenh analysis shall not affect our results. From now on we focus on the SFGs subsample.

Fig. [8](#page-11-0) is a plot of SFR_{enh} against B/T ratio, which is similar to Figure 9 of [He et al.](#page-22-18) [\(2022\)](#page-22-18) except for using the original B/T values (instead of $B/T = 0$) for galaxies with pseudobulges, and the 5 bins are divided the same as in Figure [5.](#page-8-0) It shows a broad scatter of $\rm sSFR_{\rm enh}$ for galaxies with pseudobulges, which spans about two orders of magnitude (\sim 1 orders magnitude larger than that for galaxies with classical bulges). This demonstrates that the broad scatter of $sSFR_{\text{enh}}$ for disky galaxies (with $B/T <$ 0.1) found by [He et al. \(2022\)](#page-22-18) is indeed due to galaxies with pseudobulges which are assigned $B/T = 0$ in that paper. The mean SFR_{enh} of pseudobulge SFGs is rather constant against the B/T ratio, in contrast to that of SFGs with classical bulges which decreases with increasing B/T (Fig. [8;](#page-11-0) see also [He et al. 2022\)](#page-22-18). The mean SFR_{enh} of SFGs with pseudobulges ($SFR_{\text{enh}} = 0.33 \pm 0.07$) is higher than that of SFGs with classical bulges ($sSFR_{\text{enh}} = 0.12 \pm 0.06$) at 2σ level.

Fig. 5: Plot of Age vs. B/T ratio for Spirals with pseudobulges (small filled squares) and with classical bulges (small open diamonds). The color of symbols represents SFR_{enh} . The large black squares and large black diamonds represent the means of pseudobulge galaxies and of classical-bulge galaxies in the five B/T ratio bins, respectively. The error bars show the $1-\sigma$ errors of the means. The lines with light and dark shaded areas represent the means and their errors of the two kinds of galaxies in control-sample-2. The data points marked by crosses are SFR undetected in [Cao et](#page-22-14) al. [\(2016\)](#page-22-14)

The eight SFGs with the highest SFR_{enh} all have pseudobulges (Fig. [8\)](#page-11-0). In order to study the galaxies with the strong sSFR enhancement in detail, we concentrate on those having $\text{sSFR}_{\text{enh}} > 0.7$, corresponding to more than 5 times sSFR enhancement compared with the median of the matching controls. With this criterion, there are 13 strongly enhanced SFGs in our sample, nine with pseudobulges and four with classical bulges (see Fig. [8](#page-11-0) and Table [.3\)](#page-31-0). Fig. [9](#page-12-1) shows that their central stellar populations all have Ages younger than 200 Myrs, namely are dominated by stars formed recently. Excluding these galaxies, the rest of our sample have a mean $SFR_{\text{enh}} = 0.11 \pm 0.04$, which is only marginally significant at 2.8 σ level. Statistically, SFGs with strong enhancement represent 13% (=13/98) of SFGs in our sample and 10% (=13/132) of all spiral galaxies in the H-KPAIR sample.

Seven of the 13 galaxies with strong enhancement have Spitzer observations and consequently, the 8 μ m flux densities of both the nuclear emission (aperture size: D = 4 kpc) and the total emis-sion [\(Xu et al. 2010](#page-23-10)). In Fig. [10](#page-13-1) we plot sSFR_{enh} against $\text{C}_{8\mu\text{m}}$ for these galaxies, where $\text{C}_{8\mu\text{m}}$ = $f_{8\mu m, \text{nuclear}}/f_{8\mu m, \text{total}}$ is the nuclear-to-total ratio of the $8\mu m$ emission. $C_{8\mu m}$ is an indicator of the nuclear concentration of star formation in a galaxy. The plot shows a significant linear correlation between sSFR_{enh} and $\text{C}_{8\mu\text{m}}$ (the Pearson's linear correlation coefficient $\text{r}_\text{P} = 0.81$ and the significance $p_P = 0.03$) for these galaxies. Among the seven SFGs in the plot, all the five with pseudobulges have relatively high nuclear concentration ($C_{8\mu m} \geq 0.49$) whereas, for the two SFGs with classical bulges,

Fig. 6: Probability density function of pseudobulge Spirals and classical bulge Spirals in H-KPAIR.

the star formation occurs mostly outside the nuclear region ($C_{8\mu m} < 0.30$). Albeit the small sample size, the plot indicates that the pseudobulges in galaxies with strong enhancement are tightly related to nuclear/circum-nuclear starbursts. Given that most galaxies with strong enhancement have pseudobulges, our result suggests that nuclear/circum-nuclear starburst may be the dominant mode for strong tidally induced star formation in paired galaxies. On the other hand, for a low fraction of paired SFGs (particularly those with classical bulges), the tidally induced star formation may be widely distributed over the entire disk. These results are consistent with the literature which shows that nuclear/circumnuclear starbursts are common among paired galaxies with strong signs of interaction while wider spread star formation enhancement is also observed in some of them [\(Keel et al. 1985;](#page-22-33) [Kennicutt et al. 1987](#page-22-1); [Pan et al. 2019](#page-23-13); [Steffen et al. 2021](#page-23-14)).

Fig. [11](#page-14-0) and Fig. [12](#page-14-1) are optical (u, q, r) color images of the nine pseudobulge galaxies with strong enhancement and four classical-bulge galaxies with strong enhancement, respectively. These images are taken from the database of Dark Energy Spectroscopic Instrument (DESI^{[6](#page-9-1)}) which are deeper and of better angular resolutions than SDSS images and therefore can show the diffuse features more clearly. Four pseudobulge galaxies J07543194+1648214, J09155467+4419510, J09155552+4419580, and J17045097+3449020 (panel b, c, g of Figure [11\)](#page-14-0) and two classical-bulge galaxies J10100079+5440198 and J17045089+3448530 (panel b, d of Figure [12\)](#page-14-1) are found in pairs seemed to be almost coalesced, namely in a late merging stage. This result suggests that a high fraction $(6/13 = 46\%)$ of galaxies with strong enhancement are in late-stage mergers. Here we separate late-stage mergers (pairs with strong interaction signs and very close nuclei) from early-stage mergers (pairs without strong interaction signs

⁶ <https://www.legacysurvey.org/>

Fig. 7: Probability density function of pseudobulge Spirals and classical bulge Spirals in control-sample-2

or the two galaxies being widely separated). The late-stage mergers are likely seen during or after the second close encounter, because it takes relatively long time (more than a few 100 Myrs) for galaxies to develop the interaction features after the first close encounter. In general, only a low fraction of paired galaxies selected in optical and/or near-IR bands (such as H-KPAIR) are late-stage mergers because this stage lasts significantly shorter than the stage before it [\(Di Matteo et al. 2007\)](#page-22-34). Also, for the H-KPAIR sample, the selection criterion of separation ≥ 5 h⁻¹ kpc excludes some late-stage mergers, particularly those which are already coalesced. Hence, the fact that in our sample late-stage mergers represent 46% of SFGs with strong enhancement suggests that the majority of SFGs in late-stage mergers may harbor strong tidally induced star formation whereas only a very low fraction of SFGs in early-stage mergers may do so. For the latter, significant SFR enhancement may be triggered by strong tidal torques only in SFGs that are undergoing or passed the first close encounter (namely pericenter passage in [Feng et al.](#page-22-11) [2020\)](#page-22-11) in low-speed coplanar orbits (namely with well spin–orbit alignment in [Moon et al. \(2021\)](#page-22-35)) as argued by [\(Xu et al. 2021\)](#page-23-15).

The remaining seven galaxies with strong enhancement are in earlier-stage mergers with the two component galaxies clearly separated. Among them, J01183414-0013417, J13153506+6207286, and J16024257+4111501 (panel a, e of Figure [11](#page-14-0) and panel c of Figure [12\)](#page-14-1) are likely in interaction systems with low-inclination orbits and low orbital velocity (all having $\delta v < 70 \text{ km s}^{-1}$ where δv is the measured relative radial velocity of the companion). Particularly J13153506+6207286 (panel e of Figure [11\)](#page-14-0), which is in Arp 238 ($\delta v = 6$ km s⁻¹), has been confirmed to have a coplanar orbit by a simulation of [Holincheck et al. \(2016\)](#page-22-36). [Xu et al. \(2021](#page-23-15)) found a very compact molecular gas concentration in its nucleus which is fueling a strong active nuclear starburst. They argued that the nuclear

Fig. 8: Plot of sSFRenh vs. B/T ratio for SFGs with pseudobulges (small filled squares) and SFGs with classical bulges (small open diamonds). The color of symbols represents the Age of the central stellar population. The large black open squares and large black open diamonds represent the means of pseudobulge galaxies and of classical-bulge galaxies in the five B/T ratio bins, respectively. The error bars show the 1- σ errors of the means. The horizontal solid line and dashed line mark $\rm sSFR_{\rm enh} = 0$ and $sSFR_{\text{enh}} = 0.7$, respectively.

starburst is triggered by a strong tidal torque which has been predicted by simulations for galaxies in low-speed coplanar interactions [\(Barnes & Hernquist 1996;](#page-22-37) [Hopkins et al. 2009](#page-22-38)). This interpretation may also apply to J01183414-0013417 and J16024257+4111501. In contrast, J10332972+4404342, J14005783+4251203, and J14005879+4250427 (panel d, f of Figure [11\)](#page-14-0) may be in high-inclination orbits, and the latter two are members of the same pair (KPAIR J1400+4251) which has a high relative radial velocity ($\delta v = 234 \text{ km s}^{-1}$). These three galaxies all have pseudobulges and might be bar galaxies from the appearance though it is uncertain given the limited resolution of the images. If this is true then their central starbursts may be field by the bars. Finally, J03381222+0110088 (panel a of Figure [12\)](#page-14-1) is the only one that is in an S+E pair among the 13 SFGs with strong enhancement. However, as shown in the optical image, it appears to be interacting with another nearby SFG which is in the same group of the S+E pair. The three galaxies form a close triplet dominated by SFGs. Therefore J03381222+0110088 is not in a de facto S+E pair. It has a very high total gas content with $M_{H2+HI}/M_{star} = 0.56$ and a relatively low star formation efficiency of $SFE = SFR/M_{H2} = 10^{-9.10}$ yr⁻¹, which is below the average SFE of the H-KPAIR sample in [Lisenfeld et al.](#page-22-12) [\(2019\)](#page-22-12). It appears that the high sSFR of J03381222+0110088 is mainly due to its very high gas content which may not be related to any tidal effect.

Fig. 9: Plot of sSFR_{enh} vs. age of the central stellar population SFGs with pseudobulges (filled squares) and SFGs with classical bulges (open diamonds). The color of symbols represents the B/T ratio. The horizontal solid line and dashed line mark $\rm sSFR_{\rm enh} = 0$ and $\rm sSFR_{\rm enh} = 0.7$, respectively.

4.3 Pseudo-bulge galaxies in S+S and S+E pairs

In their study of the B/T dependence of $S\{SFR_{\text{enh}}}$, [He et al. \(2022](#page-22-18)) found that except for in the last bin of $0.5 < B/T \le 1$ where SFGs in S+S and S+E pairs have similar means of sSFR_{enh} (both consistent with $\rm sSFR_{enh} = 0$, in all other B/T bins the former have higher $\rm sSFR_{enh}$ means than the latter. Particularly in the lowest bin of $B/T \le 0.1$, the difference is 0.4 ± 0.1 dex. In [He et al.](#page-22-18) [\(2022\)](#page-22-18) pseudobulge galaxies are assigned with the same $B/T = 0$ and therefore are included in the bin $B/T \le 0.1$. Do pseudobulge SFGs in S+S and in S+E pairs have significantly different sSFR_{enh}?

In Fig. [13](#page-15-0) we compare pseudobulge SFGs in S+S pairs and in S+E pairs. It plots SFR_{enh} against the stellar mass M_{star} , with the color scale indicating the Age and the symbol size the B/T ratio. The plot shows clearly that, compared to pseudobulge SFGs in S+S pairs, those in S+E pairs tend to have older stellar populations and lower sSFR enhancements and none has strong enhancement ($\rm sSFR_{\rm enh} \geq 0.7$). The means of sSFR_{enh} are 0.45 ± 0.08 and -0.04 ± 0.11 for pseudobulge SFGs in S+S pairs and in S+E pairs, respectively, and the difference is above 3σ . Most of the pseudobulge SFGs in S+E pairs (9) of 11) have relatively low stellar mass ($\leq 10^{10.6}$ M_{\odot}).

Fig. [14](#page-16-1) compares pseudobulge SFGs in S+S and S+E in a plot of SFR_{enh} versus the local density indicator $N_{1 \text{Mpc}}$ (defined in [He et al. \(2022\)](#page-22-18) and also introduced in Sect. [4.1\)](#page-3-1). SFGs in highdensity regions with $N_{1Mpc} \geq 7$ have significantly lower sSFR_{enh} than the rest of the sample, and a higher fraction (5/11) of pseudobulge SFGs in S+E, compared to those in S+S, are in this category. These SFGs should be in rich groups/clusters, where star formation may be quenched by galaxy ha-

Fig. 10: Plot of sSFR_{enh} vs. the nuclear-to-total ratio of the 8μ m emission $C_{8\mu m}$.

rassment [\(Moore et al. 1998\)](#page-22-39) and/or ram-pressure stripping [\(Gunn & Gott 1972;](#page-22-40) [Giovanelli & Haynes](#page-22-41) [1983;](#page-22-41) [Gavazzi et al. 2006\)](#page-22-42). Also, these paired SFGs may have very low probability of being in a coplanar orbit because of the frequent disturbance and therefore are less likely to have nuclear starbursts [\(Xu et al. 2021\)](#page-23-15). Detailed inspections of the optical images show that their pseudobulges often correspond to bars or inner rings with low star formation activity (for example J12191866+1201054 which has $sSFR_{\text{enh}} = -0.01$, Fig. [15\)](#page-17-0).

The mean and errors of H_2 gas mass fraction[\(Lisenfeld et al. 2019\)](#page-22-12) and that of the total gas mass derived from FIR dust mass [\(Cao et al. 2016\)](#page-22-14) were plotted in two B/T bins: (1) disk dominant (B/T \leq 0.3), (2) bulge dominant ($B/T > 0.3$) in Figure [16](#page-18-0) and [17,](#page-19-0) as well the SFE calculated from them in Figure [18](#page-20-0) and [19,](#page-21-0) separately. Generally, there is a decreasing sign for both H_2 gas and the total gas on B/T. The pseudobulge SFGs in S+S pairs have the most flat slope, while they have the highest gas fractions in bulge dominant galaxy (bin 2). No obvious trend is found in either SFE versus B/T. When comparing the subsamples, the pseudobulge SFGs have higher SFE_{gas} , and the SFGs in S+S pairs take the lead in both SFE. The most attractive point is that the pseudobulge galaxies in S+E pairs have not as high SFE as the classical bulge galaxies in S+S pairs.

5 DISCUSSION

The significant sSFR enhancement of close major-merger pairs is mainly due to a small population of SFGs with strong enhancement (∼ 15% in our sample; Fig. [8](#page-11-0)). About half of these SFGs are found in late-stage mergers, all of which are in S+S pairs and none in an S+E pair. It is unlikely that any of the late-stage S+S mergers containing strong enhanced SFGs can be misclassified S+E mergers because

Pseudobulge in Paired Galaxies 15

Fig. 11: Dark Energy Spectroscopic Instrument (DESI) images of nine pseudobulge galaxies with strong star formation enhancement. The strongly enhanced galaxies are marked by red plus signs. The white dash on the right-bottom corner of each stamp illustrates the angular scale of 10′′. The pair separation and velocity difference are provided as well.

Fig. 12: DESI images of four classical bulge galaxies with strong star formation enhancement. The red plus signs mark the strongly enhanced galaxies. The blue cross in the image of J0338+0109 marks the E component of the pair. The white dash on the right-bottom corner of each stamp illustrates the angular scale of $10''$. The pair separation and velocity difference are provided as well.

none of the galaxies in these pairs is dominated by a classical bulge (the largest classical bulge is found in J17045088+3448529 shown in panel d of Figure [12](#page-14-1) with a $B/T = 0.48$).

What is the reason for spirals in late-stage S+E mergers to avoid strong starbursts? There are four S+E late-stage mergers in H-KPAIR, which are listed in Table [.4.](#page-31-1) Each of them has the two member galaxies nearly coalesced within a common halo (Fig. [20\)](#page-21-1). They are pairs of massive galaxies with the primaries more massive than $10^{11}/M_{\odot}$. All spiral galaxies in these pairs are rather quiescent in star formation and only one of them (J16354293+2630494; panel d) is included in our SFG sample with an $\text{sSFR} = 10^{-11.26} \text{ yr}^{-1}$. They are uniformly gas-poor with the $\text{M}_{\text{gas}}/\text{M}_{\text{star}}$ ratio less than 4% [\(Cao et al.](#page-22-14) [2016\)](#page-22-14). which is lower than the ratios of 88 percent of SFGs in the H-KPAIR sample, and significantly below the mean of the sample ($M_{\text{gas}}/M_{\text{star}} = 0.13 \pm 0.01$).

Three (J10514450+5101303, J13131470+3910382, and J16354293+2630494) are in pairs containing the most massive galaxies of rich groups or clusters that have more than 12 members [\(Yang et al.](#page-23-16) [2012\)](#page-23-16). For them, the rich-group/cluster environment may be the key to the lack of cold gas and star

Fig. 13: Plot of sSFRenh vs. Mstar for pseudo-bulge SFGs in S+S pairs (pentagram) and in S+E pairs (circle). The symbol color indicates the Age of the central stellar population, and the symbol size shows the B/T ratio. The dashed line corresponds to $\rm sSFR_{\rm enh} = 0.7$.

formation because the ram-pressure stripping and/or evaporation by the hot gas in IGM can remove most of the cold gas in the ISM and tidal tails. J11542299+4932509 is apparently in an isolated pair. Its companion (J11542307+4932456) is a very massive Elliptical with $M_{star} = 10^{11.35} M_{\odot}$. The pair is detected by ROSAT in X-ray with a flux of $f_{0.2-2 \text{ keV}} = 8.22(\pm 1.73) \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$. This may be indicative of the presence of a hot gas halo around the pair which can strip/evaporate the cold gas and be responsible for the low gas content ($M_{\rm gas}/M_{\rm star} = 0.035$) and low sSFR ($< 10^{-11.33}$ yr^{-1}) of J11542299+4932509.

Thus, it appears that the lack of strong enhancement in late-stage S+E mergers may be due to the removal of cold gas by hot gas in the immediate environment [\(Park & Choi 2009;](#page-23-11) [Hwang et al. 2011](#page-22-43)). It should be pointed out that the quiescent spiral galaxies in these late-stage S+E pairs are different from the SFGs studied by [Cao et al.](#page-22-14) [\(2016\)](#page-22-14) and [Lisenfeld et al. \(2019](#page-22-12)). These authors found no significant difference in M_{gas}/M_{star} and in M_{H2+HI}/M_{star} between SFGs in S+S and S+E pairs, and therefore refuted the "cold-gas stripping" hypothesis as far as SFGs in S+E pairs are concerned. A majority of spirals in S+E pairs are SFGs $(64\% = 28/44$ for H-KPAIR), and many of them are gas-rich (e.g. NGC 2936 in Arp 142; [Xu et al. 2021\)](#page-23-15). It will be interesting to find out why they are absent in late-stage S+E mergers.

The interaction-induced star formation enhancement is affected by many factors, some seem to be necessary conditions (e.g., a galaxy should have gas to fuel star formation), but it's hard to say which is a sufficient condition. For instance, the four most enhanced classical bulge SFGs (Figure [12\)](#page-14-1) never reached the ultra enhanced level (say, $sSFR_{\text{enh}} > 1$, corresponding to more than 10 times

Fig. 14: Plot of $sSFR_{\rm enh}$ vs. $N_{1\rm Mpc}$, where $N_{1\rm Mpc}$ is an indicator of local density defined by the number of galaxies brighter than $M_r = -19.7$ mag within 1 Mpc projected distance around the galaxy in question [\(He et al. 2022\)](#page-22-18). filled pentagrams and filled circles represent pseudobulge SFGs in S+S pairs and in S+E pairs, respectively. The dashed line corresponds to $sSFR_{\text{enh}} = 0.7$.

sSFR enhancement compared with the median of the matching controls) as their pseudobulge counterparts. However, as the outstanding cases in classical SFGs, three of them have very low B/T. The other one J17045088+3448529 with $B/T = 0.48$, as well the "blue" bulge dominated galaxy $J13151726+4424255$ whose SFR_{enh} reaches almost to 0.7 as we talked about the Figure [5](#page-8-0) (also see the most upper-right data point in Figure [8\)](#page-11-0), are located in low density environment ($N_{1\text{Mpc}}$ equal to 4 and 3, respectively).

Though KPAIR [\(Domingue et al. 2009](#page-22-15)) was selected from allsky, strict selection criteria make it a relatively small sample. After further refining the classifications such as separating the sample into S+S/S+E pair or pseudo/classical bulge, the subsamples sometimes are too small for a very robust for a robust statistical analysis. Nevertheless, the individual case studies in this paper provide a perspective of the complex mechanism of the interaction-induced star formation enhancement. Future sky surveys with deeper sensitivities and sharper resolutions (e.g. Euclid) will produce larger and better samples for more conclusive studies.

6 CONCLUSION

Here we present a study of SFGs with pseudobulges (bulges with Sérsic index $n < 2$) in a local close major-merger galaxy pair sample (H-KPAIR). The sample is taken from [He et al.](#page-22-18) [\(2022\)](#page-22-18) who did 2-D GALFIT decompositions to study the B/T dependence of the star formation enhancement of paired SFGs.

Fig. 15: DESI images of J1219+1200, an S+E pair. S component, J12191866+1201054 (identified by a red plus sign), is a pseudobulge SFG that contains a bar. The white dash on the right-bottom corner of the figure illustrates the angular scale of 10".

New aperture photometries in the SDSS u, r, i -bands and the 2MASS K_s -band are carried out for the sample to investigate the stellar populations in the central part $(D = 7 \text{ kpc})$ of the galaxies. With the new data and data from the literature, we obtain the following results:

– The mean Age of central stellar populations in spiral galaxies with classical bulges increases with the B/T ratio. On the other hand, the mean Age of central stellar populations in Spirals with pseudobulges is nearly constant against the B/T ratio and is consistent with the mean Age of central stellar populations of disky galaxies (with B/T \sim 0). This confirms that the pseudobulges in the sample are associated mainly with disky phenomena such as bars, nuclear rings, and bright nuclei.

Fig. 16: Plot of means of $\log(M_{H2}/M_{star})$ and errors in two B/T bins for pseudobulge SFGs (square) and classical bulge SFGs (diamond) in S+S (blue) and S+E (red) pair.

- Spiral galaxies have a modestly reduced fraction of pure disk galaxies than in single Spirals, which can be due to the misidentifying as a large pseudobulge of the nuclear/circum-nuclear starburst phenomena.
- Compared with SFGs with classical bulges, those with pseudobulges have a 2 σ higher mean of sSFR enhancement ($\text{SSFR}_{\text{enh}} = 0.33 \pm 0.07$ vs $\text{SSFR}_{\text{enh}} = 0.12 \pm 0.06$) and much broader scatter $(\sim 1$ dex).
- The eight SFGs that have the highest SFR_{enh} in the sample are all pseudobulge galaxies. And a majority (69%) of paired SFGs with strong enhancement (having sSFR more than 5 times the median of the control galaxies) have pseudobulges. The Spitzer data show that the pseudobulges in these galaxies are tightly linked to nuclear starbursts. These results suggest that nuclear starburst may be the dominant mode for strong tidally induced star formation in paired galaxies.
- Pseudobulge SFGs in S+S and in S+E pairs have 3σ significantly different sSFR enhancement: the mean sSFR_{enh} = 0.45 ± 0.08 and -0.04 ± 0.11 for the former and latter, respectively. A high fraction (5/11) of pseudobulge SFGs in S+E are in rich groups/clusters (local density $N_{1\text{Mpc}} \geq 7$), whose star formation enhancement may be hindered by the environment.

In conclusion, we find that paired SFGs with pseudobulges have very diversified central stellar populations and sSFR enhancements. They include galaxies with bars, inner disks and rings, and strong nuclear/circum-nuclear starbursts. The SFGs with strong sSFR enhancement are dominated by pseudobulge galaxies with nuclear/circum-nuclear starbursts. On the other hand, many pseudobulge SFGs,

Fig. 17: Plot of means of $\log(M_{\rm gas}/\rm{Mstar})$ and errors in two B/T bins for pseudobulge SFGs (square) and classical bulge SFGs (diamond) in S+S (blue) and S+E (red) pair.

particularly those in S+E pairs, are barred or ringed galaxies with old central stellar populations and low sSFR enhancements, which are probably caused by environmental effects.

Acknowledgements This work is supported by National Natural Science Foundation of China (NSFC) No. 11873055 and No.11933003, and is sponsored (in part) by the Chinese Academy of Sciences (CAS) through a grant to the CAS South America Center for Astronomy (CASSACA). UL acknowledges support from project PID2020-114414GB-100, financed by MCIN/AEI/10.13039/501100011033 and from the Junta de Andaluciaía (Spain) grant FQM108. TF and QY acknowledge support by National Key R&D Program of China No. 2017YFA0402600, National Natural Science Foundation of China (NSFC) grant Nos. 11890692, 12133008, and 12221003 and China Manned Space Project No. CMS-CSST-2021-A04.

The authors would like to thank Gaoxiang Jin, Cheng Cheng, Chen Cao, Yilun Wang, and Yufu Shen for their helpful discussions.

This publication makes use of data products from the Sloan Digital Sky Survey (SDSS). Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions. SDSS acknowledges support and resources from the Center for High-Performance Computing at the University of Utah. The SDSS website is <www.sdss.org>. SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS Collaboration including the Brazilian Participation Group, the Carnegie Institution for Science, Carnegie Mellon University, Center for Astrophysics — Harvard & Smithsonian (CfA), the Chilean Participation Group, the French Participation Group, Instituto de Astrofísica de

Fig. 18: Plot of means of $\log(\text{SFE}_{H2})$ and errors in two B/T bins for pseudobulge SFGs (square) and classical bulge SFGs (diamond) in S+S (blue) and S+E (red) pair.

Canarias, The Johns Hopkins University, Kavli Institute for the Physics and Mathematics of the Universe (IPMU) / University of Tokyo, the Korean Participation Group, Lawrence Berkeley National Laboratory, Leibniz Institut für Astrophysik Potsdam (AIP), Max-Planck-Institut für Astronomie (MPIA Heidelberg), Max-Planck-Institut für Astrophysik (MPA Garching), Max-Planck-Institut für Extraterrestrische Physik (MPE), National Astronomical Observatories of China, New Mexico State University, New York University, University of Notre Dame, Observatorio Nacional / MCTI, The Ohio State University, Pennsylvania State University, Shanghai Astronomical Observatory, United Kingdom Participation Group, Universidad Nacional Autónoma de México, University of Arizona, University of Colorado Boulder, University of Oxford, University of Portsmouth, University of Utah, University of Virginia, University of Washington, University of Wisconsin, Vanderbilt University, and Yale University.

This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

This publication makes use of image products from the Dark Energy Spectroscopic Instrument (DESI). DESI construction and operations is managed by the Lawrence Berkeley National Laboratory. This research is supported by the U.S. Department of Energy, Office of Science, Office of High-Energy Physics, under Contract No. DE–AC02–05CH11231, and by the National Energy Research Scientific Computing Center, a DOE Office of Science User Facility under the same contract. Additional support

Fig. 19: Plot of means of $\log(SFE_{\rm gas})$ and errors in two B/T bins for pseudobulge SFGs (square) and classical bulge SFGs (diamond) in S+S (blue) and S+E (red) pair.

Fig. 20: DESI images of four late-stage S+E mergers. The red plus signs mark the S components and the blue crosses the E components. The white dash on the right-bottom corner of each stamp illustrates the angular scale of 10′′. The pair separation and velocity difference are provided as well.

for DESI is provided by the U.S. National Science Foundation, Division of Astronomical Sciences under Contract No. AST-0950945 to the NSF's National Optical-Infrared Astronomy Research Laboratory; the Science and Technologies Facilities Council of the United Kingdom; the Gordon and Betty Moore Foundation; the Heising-Simons Foundation; the French Alternative Energies and Atomic Energy Commission (CEA); the National Council of Science and Technology of Mexico (CONACYT); the Ministry of Science and Innovation of Spain, and by the DESI Member Institutions. The DESI collaboration is honored to be permitted to conduct astronomical research on Iolkam Du'ag (Kitt Peak), a mountain with particular significance to the Tohono O'odham Nation.

References

- Alonso, M. S., Tissera, P. B., Coldwell, G., & Lambas, D. G. 2004, MNRAS, 352, 1081 [2](#page-1-0)
- Barnes, J. E., & Hernquist, L. 1996, ApJ, 471, 115 [12](#page-11-1)
- Bertin, E., Mellier, Y., Radovich, M., et al. 2002, in Astronomical Society of the Pacific Conference Series, Vol. 281, Astronomical Data Analysis Software and Systems XI, ed. D. A. Bohlender, D. Durand, & T. H. Handley, 228 [3](#page-2-4)
- Bluck, A. F. L., Mendel, J. T., Ellison, S. L., et al. 2014, MNRAS, 441, 599 [2](#page-1-0)
- Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000 [4](#page-3-5)
- Cao, C., Xu, C. K., Domingue, D., et al. 2016, ApJS, 222, 16 [2,](#page-1-0) [3,](#page-2-4) [5,](#page-4-1) [7,](#page-6-1) [8,](#page-7-2) [9,](#page-8-1) [14,](#page-13-2) [15,](#page-14-2) [16,](#page-15-1) [32](#page-31-2)
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245 [4](#page-3-5)
- Chown, R., Li, C., Athanassoula, E., et al. 2019, MNRAS, 484, 5192 [2](#page-1-0)
- Cox, T. J., Jonsson, P., Somerville, R. S., Primack, J. R., & Dekel, A. 2008, MNRAS, 384, 386 [2](#page-1-0)
- Di Matteo, P., Bournaud, F., Martig, M., et al. 2008, A&A, 492, 31 [2](#page-1-0)
- Di Matteo, P., Combes, F., Melchior, A. L., & Semelin, B. 2007, A&A, 468, 61 [11](#page-10-1)
- Domingue, D. L., Xu, C. K., Jarrett, T. H., & Cheng, Y. 2009, ApJ, 695, 1559 [2,](#page-1-0) [3,](#page-2-4) [17](#page-16-2)
- Driver, S. P., Allen, P. D., Graham, A. W., et al. 2006, MNRAS, 368, 414 [6](#page-5-1)
- Ellison, S. L., Patton, D. R., Simard, L., et al. 2010, MNRAS, 407, 1514 [2](#page-1-0)
- Erwin, P., Seth, A., Debattista, V. P., et al. 2021, MNRAS, 502, 2446 [2](#page-1-0)
- Feng, S., Shen, S.-Y., Yuan, F.-T., Riffel, R. A., & Pan, K. 2020, ApJ, 892, L20 [2,](#page-1-0) [11](#page-10-1)
- Fraser-McKelvie, A., Merrifield, M., Aragón-Salamanca, A., et al. 2020, MNRAS, 499, 1116 [3](#page-2-4)
- Gavazzi, G., Boselli, A., Cortese, L., et al. 2006, A&A, 446, 839 [14](#page-13-2)
- Giovanelli, R., & Haynes, M. P. 1983, AJ, 88, 881 [14](#page-13-2)
- Gunn, J. E., & Gott, J. Richard, I. 1972, ApJ, 176, 1 [14](#page-13-2)
- He, C., Xu, C. K., Domingue, D., Cao, C., & Huang, J.-s. 2022, ApJS, 261, 34 [2,](#page-1-0) [3,](#page-2-4) [5,](#page-4-1) [6,](#page-5-1) [7,](#page-6-1) [8,](#page-7-2) [13,](#page-12-2) [17](#page-16-2)
- Herrera-Endoqui, M., Salo, H., Laurikainen, E., & Knapen, J. H. 2017, A&A, 599, A43 [3](#page-2-4)
- Holincheck, A. J., Wallin, J. F., Borne, K., et al. 2016, MNRAS, 459, 720 [11](#page-10-1)
- Hopkins, P. F., Cox, T. J., Younger, J. D., & Hernquist, L. 2009, ApJ, 691, 1168 [12](#page-11-1)
- Hwang, H. S., Elbaz, D., Lee, J. C., et al. 2010, A&A, 522, A33 [2](#page-1-0)
- Hwang, H. S., Elbaz, D., Dickinson, M., et al. 2011, A&A, 535, A60 [16](#page-15-1)
- Keel, W. C., Kennicutt, R. C., J., Hummel, E., & van der Hulst, J. M. 1985, AJ, 90, 708 [10](#page-9-2)
- Kennicutt, Robert C., J. 1998, ARA&A, 36, 189 [2](#page-1-0)
- Kennicutt, Robert C., J., Keel, W. C., van der Hulst, J. M., Hummel, E., & Roettiger, K. A. 1987, AJ, 93, 1011 [2,](#page-1-0) [10](#page-9-2)
- Kim, K., Oh, S., Jeong, H., et al. 2016, ApJS, 225, 6 [2,](#page-1-0) [6,](#page-5-1) [7](#page-6-1)
- Kormendy, J., & Ho, L. C. 2013, ARA&A, 51, 511 [2](#page-1-0)
- Kormendy, J., & Kennicutt, Robert C., J. 2004, ARA&A, 42, 603 [2](#page-1-0)
- Larson, R. B., & Tinsley, B. M. 1978, ApJ, 219, 46 [2](#page-1-0)
- Li, Y. A., Ho, L. C., & Shangguan, J. 2023, ApJ, 953, 91 [2](#page-1-0)
- Lintott, C. J., Schawinski, K., Slosar, A., et al. 2008, MNRAS, 389, 1179 [5](#page-4-1)
- Lisenfeld, U., Xu, C. K., Gao, Y., et al. 2019, A&A, 627, A107 [2,](#page-1-0) [3,](#page-2-4) [12,](#page-11-1) [14,](#page-13-2) [16,](#page-15-1) [32](#page-31-2)
- Madau, P., & Dickinson, M. 2014, ARA&A, 52, 415 [2](#page-1-0)
- Martig, M., Bournaud, F., Teyssier, R., & Dekel, A. 2009, ApJ, 707, 250 [2](#page-1-0)
- Meert, A., Vikram, V., & Bernardi, M. 2015, MNRAS, 446, 3943 [5](#page-4-1)
- Mihos, J. C., & Hernquist, L. 1996, ApJ, 464, 641 [2](#page-1-0)
- Moon, J.-S., An, S.-H., & Yoon, S.-J. 2019, ApJ, 882, 14 [2](#page-1-0)
- Moon, J.-S., An, S.-H., & Yoon, S.-J. 2021, ApJ, 909, 34 [11](#page-10-1)
- Moore, B., Lake, G., & Katz, N. 1998, ApJ, 495, 139 [14](#page-13-2)
- Nikolic, B., Cullen, H., & Alexander, P. 2004, MNRAS, 355, 874 [2](#page-1-0)
- Pan, H.-A., Lin, L., Hsieh, B.-C., et al. 2019, ApJ, 881, 119 [10](#page-9-2)
- Park, C., & Choi, Y.-Y. 2009, ApJ, 691, 1828 [2,](#page-1-0) [16](#page-15-1)
- Patton, D. R., Torrey, P., Ellison, S. L., Mendel, J. T., & Scudder, J. M. 2013, MNRAS, 433, L59 [2](#page-1-0)
- Peng, Y.-j., Lilly, S. J., Renzini, A., & Carollo, M. 2012, ApJ, 757, 4 [2](#page-1-0)
- Peng, Y.-j., Lilly, S. J., Kovač, K., et al. [2](#page-1-0)010, ApJ, 721, 193 2
- Rodighiero, G., Daddi, E., Baronchelli, I., et al. 2011, ApJ, 739, L40 [2](#page-1-0)
- Sanders, D. B., & Mirabel, I. F. 1996, ARA&A, 34, 749 [2](#page-1-0)
- Sargent, A., & Scoville, N. 1991, ApJ, 366, L1 [2](#page-1-0)
- Scoville, N. Z., Sargent, A. I., Sanders, D. B., & Soifer, B. T. 1991, ApJ, 366, L5 [2](#page-1-0)
- Scudder, J. M., Ellison, S. L., Torrey, P., Patton, D. R., & Mendel, J. T. 2012, MNRAS, 426, 549 [2](#page-1-0)
- Steffen, J. L., Fu, H., Comerford, J. M., et al. 2021, ApJ, 909, 120 [10](#page-9-2)
- Toomre, A., & Toomre, J. 1972, ApJ, 178, 623 [2](#page-1-0)
- Xu, C. K., Lisenfeld, U., Gao, Y., & Renaud, F. 2021, ApJ, 918, 55 [11,](#page-10-1) [14,](#page-13-2) [16](#page-15-1)
- Xu, C. K., Domingue, D., Cheng, Y.-W., et al. 2010, ApJ, 713, 330 [2,](#page-1-0) [9,](#page-8-1) [32](#page-31-2)
- Xu, C., & Sulentic, J. W. 1991, ApJ, 374, 407 [2](#page-1-0)
- Yang, X., Mo, H. J., van den Bosch, F. C., Zhang, Y., & Han, J. 2012, ApJ, 752, 41 [15,](#page-14-2) [32](#page-31-2)
- Zuo, P., Xu, C. K., Yun, M. S., et al. 2018, ApJS, 237, 2 [3,](#page-2-4) [32](#page-31-2)

Table .1: Spiral galaxies in H-KPAIR

name	\boldsymbol{z}	pair type	bulge_type	is_SFG	$r_{\rm mag}$	B/T	$log M_{star}$	$sSFR_{\rm enh}$	$\overline{\mathrm{N}_{1 \mathrm{Mpc}}}$
					(mag)		(M_{\odot})	(dex)	
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
J00202580+0049350	0.0149	SЕ	\overline{p}	Τ	13.66	0.11	10.44	0.34	2
J01183417-0013416	0.0453	SS	p	$\mathbf{1}$	15.60	0.37	10.97	1.34	6
J01183556-0013594	0.0455	SS	$\mathbf c$	$\mathbf{1}$	14.96	0.01	10.57	0.57	6
J02110638-0039191	0.0177	SS	$\mathbf c$	$\mathbf{1}$	14.50	0.27	10.42	0.45	4
J02110832-0039171	0.0181	SS	$\mathbf c$	$\overline{0}$	13.62	0.73	10.63	-1.19	$\overline{4}$
J03381222+0110088	0.0392	SE	$\mathbf c$	$\mathbf{1}$	15.19	0.12	10.67	0.75	10
J07543194+1648214	0.0459	SS	p	$\mathbf{1}$	14.63	0.03	10.95	0.71	3
$J07543221+1648349$	0.0462	SS	p	$\mathbf{1}$	14.53	0.15	11.15	0.62	3
$J08083377+3854534$	0.0402	SE	$\mathbf c$	$\mathbf{1}$	15.45	0.86	10.73	0.11	16
J08233266+2120171	0.0181	SS	p	$\mathbf{1}$	14.42	0.26	10.08	0.39	$\,$ 8 $\,$
$J08233421 + 2120515$	0.0181	SS	$\mathbf c$	$\mathbf{1}$	13.80	0.39	10.35	0.55	8
J08291491+5531227	0.0251	SS	$\, {\bf p}$	$\mathbf{1}$	14.09	0.03	10.56	0.32	3
$J08292083 + 5531081$	0.0252	SS	p	$\mathbf{1}$	14.04	0.11	10.71	0.18	3
J08364482+4722188	0.0526	SS	p	$\mathbf{1}$	15.18	0.19	11.02	-0.46	9
J08364588+4722100	0.0526	SS	\mathbf{c}	$\overline{0}$	14.96	0.87	11.17	-0.67	9
J08381759+3054534	0.0476	SS	$\mathbf c$	$\mathbf{1}$	15.79	0.75	10.77	0.24	3
J08381795+3055011	0.0481	SS	$\mathbf c$	$\mathbf{1}$	15.09	0.73	11.08	-0.37	3
J08390125+3613042	0.0548	SE	$\mathbf c$	$\mathbf{1}$	15.51	0.71	10.94	-0.06	6
J08414959+2642578	0.0848	SE	p	$\mathbf{1}$	15.43	0.46	11.44	-0.54	12
$J09060498 + 5144071$	0.0291	SE	$\mathbf c$	$\mathbf{1}$	14.34	0.14	10.60	0.08	$\overline{4}$
J09123676+3547462	0.0235	SE	$\mathbf c$	$\overline{0}$	15.04	0.37	10.28	-1.15	6
J09134606+4742001	0.0527	SE	$\mathbf c$	1	14.89	0.43	11.05	0.11	23
J09155467+4419510	0.0396	SS	p	$\mathbf{1}$	14.97	0.00	10.76	1.11	$\sqrt{2}$
J09155552+4419580	0.0396	SS	p	$\mathbf{1}$	14.14	0.20	11.11	1.36	$\overline{2}$
$J09264111+0447247$	0.0891	SS	$\mathbf c$	$\mathbf{1}$	16.21	1.00	11.12	-0.52	5
$J09264137+0447260$	0.0907	SS	\mathbf{c}	$\boldsymbol{0}$	16.17	0.70	11.40	-0.82	5
J09374413+0245394	0.0242	SE	$\mathbf c$	$\mathbf{1}$	12.85	0.09	11.24	0.60	\mathfrak{Z}
J10100079+5440198	0.0460	SS	\mathbf{c}	$\mathbf{1}$	14.90	0.00	10.89	0.79	23
J10100212+5440279	0.0463	SS	$\mathbf c$	$\mathbf{1}$	15.67	0.29	10.85	0.29	23
J10155257+0657330	0.0299	SE	p	$\mathbf{1}$	15.09	0.33	10.51	-0.82	3
J10205188+4831096	0.0530	SE	$\mathbf c$	$\mathbf{1}$	16.11	0.34	10.60	0.05	\overline{c}
J10225647+3446564	0.0554	SS	p	$\mathbf{1}$	15.99	0.35	10.67	0.41	\overline{c}
J10225655+3446468	0.0564	SS	$\mathbf c$	$\boldsymbol{0}$	15.03	0.93	11.02	-0.87	$\overline{2}$
J10233658+4220477	0.0456	SS	$\, {\bf p}$	$\mathbf{1}$	15.00	0.14	10.79	0.49	6
J10233684+4221037	0.0454	SS	p	$\mathbf{1}$	15.84	0.63	10.55	0.43	6
J10272950+0114490	0.0236	SE	p	$\mathbf{1}$	14.34	0.52	10.48	0.32	5
J10325316+5306536	0.0640	SE	$\mathbf c$	$\mathbf{1}$	15.49	0.72	11.05	-0.41	19
J10332972+4404342	0.0523	SS	$\, {\bf p}$	$\mathbf{1}$	14.79	0.26	11.12	0.84	4
J10333162+4404212	0.0521	SS	p	$\mathbf{1}$	15.65	0.13	10.95	0.24	4
J10364274+5447356	0.0458	SE	$\mathbf c$	$\boldsymbol{0}$	15.34	0.79	10.81	-1.12	$\sqrt{ }$
J10392338+3904501	0.0435	SE	p	$\overline{0}$	15.26	0.38	10.77	-1.11	\overline{c}
J10435053+0645466	0.0287	SS	$\mathbf c$	$\mathbf{1}$	14.26	0.11	10.55	0.70	5
J10435268+0645256	0.0281	SS	p	1	14.66	0.23	10.48	-0.25	5
J10452478+3910298	0.0268	SE	$\mathbf c$	$\mathbf{1}$	14.44	0.46	10.68	-0.34	4
J10514450+5101303	0.0238	SE	$\mathbf c$	$\overline{0}$	14.06	0.56	10.85	-0.76	8
J10595915+0857357	0.0627	SE	\mathbf{c}	θ	15.63	0.48	10.90	-0.71	9
continue									

26	He et al.

Table 1 continue

Notes: The columns are: (1) galaxy name; (2) redshift; (3) pair type:"S" for Spiral+Spiral pair, "S" for Spiral+Elliptical pair; (4) if the galaxy is an SFG, 1 for true and 0 for false; (5) bulge type: "p" for pseudobulge, "c" for classical-bulge; (6) magnitude of r-band; (7) Bulge-to-Total ratio; (8) stellar mass; (9) specific star formation rate enhancement; (10) number of galaxies brighter than $M_r = -19.7$ mag within a $D = 1$ Mpc circle centered on the galaxy.

Table .2: Aperture photometry and model fits of the H-KPAIR sample

name	type	m_u	e_u	m_r	e_r	m_i	e_i	m_{K_s}	e_{K_s}	Age	A_v
		(mag)	(mag)	(Gyr)	(mag)						
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
J00202580+0049350	ς	16.86	0.02	14.51	0.01	14.09	0.01	13.34	0.02	0.56	1.25
J00202748+0050009	Ε	16.48	0.02	13.87	0.01	13.44	0.01	12.71	0.01	3.21	0.61
J01183417-0013416	${\bf S}$	18.51	0.03	16.17	0.01	15.65	0.01	14.18	0.07	0.05	2.85
J01183556-0013594	${\bf S}$	18.17	0.03	16.55	0.01	16.24	0.01	15.66	0.05	0.09	1.54
J02110638-0039191	${\bf S}$	17.75	0.03	15.25	0.01	14.77	0.01	13.50	0.02	0.10	2.77
J02110832-0039171	${\bf S}$	16.30	0.02	13.91	0.01	13.54	0.01	12.92	0.01	1.61	0.67
J03381222+0110088	${\bf S}$	18.67	0.03	16.39	0.01	15.92	0.01	14.97	0.04	0.17	2.02
J03381299+0109414	E	18.94	0.03	16.09	0.01	15.63	0.01	14.76	0.08	6.31	0.61
J07543194+1648214	${\bf S}$	18.17	0.03	15.93	0.01	15.52	0.01	14.44	0.28	0.09	2.42
J07543221+1648349	${\bf S}$	17.75	0.02	15.64	0.01	15.23	0.01	14.35	0.26	0.13	2.01
J08083377+3854534	${\bf S}$	18.34	0.03	16.01	0.01	15.61	0.01	14.84	0.04	0.46	1.34
J08083563+3854522	E	18.27	0.03	15.49	0.01	15.05	0.01	14.24	0.02	6.07	0.53
J08233266+2120171	${\bf S}$	16.10	0.02	14.73	0.01	14.53	0.01	14.07	0.09	0.07	1.26
J08233421+2120515	${\bf S}$	16.07	0.02	14.47	0.01	14.21	0.01	13.67	0.06	0.09	1.48
J08291491+5531227	${\bf S}$	17.57	0.03	15.37	0.01	14.97	0.01	14.25	0.02	0.34	1.36
J08292083+5531081	$\mathbf S$	17.24	0.02	14.84	0.01	14.44	0.01	13.70	0.01	0.72	1.16
J08364482+4722188	S	18.99	0.03	16.28	0.01	15.83	0.01	14.95	0.03	1.74	1.11
J08364588+4722100	${\bf S}$	18.66	0.03	15.84	0.01	15.41	0.01	14.57	0.02	7.35	0.48
J08381759+3054534	S	18.73	0.03	16.34	0.01	15.85	0.01	14.84	0.02	0.19	2.09
J08381795+3055011	${\bf S}$	19.00	0.03	16.07	0.01	15.52	0.01	14.31	0.01	0.61	2.05
J08385973+3613164	E	18.58	0.03	15.74	0.01	15.30	0.01	14.53	0.03	13.00	0.12
J08390125+3613042	${\bf S}$	19.00	0.03	16.35	0.01	15.89	0.01	15.01	0.05	0.98	1.32
J08414959+2642578	S	19.65	0.03	16.69	0.01	16.16	0.01	15.21	0.03	6.62	0.75
J08415054+2642475	E	20.02	0.04	17.09	0.01	16.63	0.01	16.08	0.05	13.00	0.00
$J09060283 + 5144411$	E	17.61	0.02	14.91	0.01	14.52	0.01	13.74	0.02	4.52	0.56
J09060498+5144071	${\bf S}$	17.99	0.03	15.59	0.01	15.20	0.01	14.49	0.03	0.86	1.05
J09123636+3547180	E	17.15	0.02	14.58	0.01	14.18	0.01	13.56	0.01	7.19	0.12
J09123676+3547462	${\bf S}$	17.65	0.02	15.25	0.01	14.87	0.01	14.34	0.02	4.52	0.13
J09134461+4742165	${\bf E}$	18.57	0.03	15.75	0.01	15.31	0.01	14.56	0.02	13.00	0.02
J09134606+4742001	${\bf S}$	17.54	0.02	15.98	0.01	15.49	0.01	14.43	0.01	0.04	1.86
J09155467+4419510	${\bf S}$	18.19	0.03	15.85	0.01	15.40	0.01	14.22	0.01	0.09	2.60
J09155552+4419580	${\bf S}$	17.58	0.02	15.45	0.01	15.03	0.01	13.93	0.01	0.07	2.39
$J09264111+0447247$	${\bf S}$	20.02	0.04	17.18	0.01	16.73	0.01	15.98	0.15	13.00	0.00
J09264137+0447260	$\mathbf S$	20.03	0.04	17.11	0.01	16.65	0.01	15.69	0.11	4.26	0.93
J09374413+0245394	${\bf S}$	16.77	0.02	14.41	0.01	13.97	0.01	12.83	0.01	0.10	2.57
J09374506+0244504	\overline{E}	16.79	0.02	14.00	0.01	13.57	0.01	12.72	0.01	4.36	0.72
J10100079+5440198	${\bf S}$	18.61	0.03	16.46	0.01	16.01	0.01	14.91	0.01	0.07	2.39
J10100212+5440279	S	18.46	0.03	16.26	0.01	15.83	0.01	14.96	0.01	0.18	1.86
J10155257+0657330	${\bf S}$	17.87	0.02	15.57	0.01	15.20	0.01	14.60	0.15	0.92	0.84
J10155338+0657495	\overline{E}	17.41	0.02	14.79	0.01	14.38	0.01	13.66	0.07	3.77	0.54
J10205188+4831096	${\bf S}$	19.31	0.03	17.00	0.01	16.56	0.01	15.76	0.06	0.36	1.48
J10205369+4831246	E	19.07	0.03	16.18	0.01	15.72	0.01	14.86	0.03	9.75	0.43
J10225647+3446564	${\bf S}$	18.90	0.03	16.70	0.01	16.27	0.01	15.41	0.05	0.18	1.84
J10225655+3446468	${\bf S}$	18.33	0.03	15.79	0.01	15.41	0.01	14.76	0.03	4.26	0.37
J10233658+4220477	${\bf S}$	17.76	0.02	16.02	0.01	15.62	0.01	14.82	0.01	0.06	1.83
J10233684+4221037	S	17.89	0.02	16.28	0.01	15.88	0.01	15.15	0.02	0.06	1.68
continue											

Table2 continue

name	type	m_u	e_u	m_r	e_r	m_i	e_i	m_{K_s}	e_{K_s}	Age	A_v
		(mag)	(mag)	(Gyr)	(mag)						
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
J12191719+1200582	Ε	17.82	0.03	15.17	0.01	14.77	0.01	14.09	0.03	7.98	0.18
J12191866+1201054	${\bf S}$	17.63	0.02	15.69	0.01	15.33	0.01	14.66	0.04	0.17	1.52
J12433887+4405399	${\bf S}$	18.73	0.03	16.07	0.01	15.66	0.01	14.95	0.03	6.43	0.32
J12433936+4406046	E	18.00	0.02	15.31	0.01	14.92	0.01	14.19	0.02	6.70	0.33
J12525011+4645272	${\bf S}$	18.98	0.03	16.48	0.01	16.05	0.01	15.37	0.03	2.10	0.69
J12525212+4645294	${\bf E}$	18.92	0.03	16.11	0.01	15.63	0.01	14.83	0.02	8.83	0.36
J13011662+4803366	${\bf S}$	16.20	0.02	15.00	0.01	14.80	0.01	14.29	0.02	0.05	1.18
J13011835+4803304	S	17.04	0.02	15.56	0.01	15.31	0.01	14.78	0.03	0.07	1.40
J13082737+0422125	S	17.67	0.02	15.98	0.01	15.70	0.01	15.36	0.06	0.28	0.75
J13082964+0422045	S	17.36	0.02	15.32	0.01	14.98	0.01	14.41	0.03	0.37	1.07
J13131429+3910360	${\bf E}$	19.15	0.03	16.26	0.01	15.84	0.01	15.13	0.02	13.00	0.00
J13131470+3910382	S	19.54	0.03	16.78	0.01	16.40	0.01	15.75	0.03	13.00	0.00
J13151386+4424264	S	17.40	0.02	15.46	0.01	15.12	0.01	14.38	0.19	0.13	1.74
J13151726+4424255	S	17.41	0.02	15.02	0.01	14.59	0.01	13.43	0.08	0.11	2.57
J13153076+6207447	S	17.33	0.02	15.56	0.01	15.23	0.01	14.06	0.02	0.04	2.13
J13153506+6207287	S	16.62	0.02	15.13	0.01	14.81	0.01	13.49	0.01	0.02	2.00
J13325525-0301347	S	18.60	0.03	16.37	0.01	15.95	0.01	15.05	0.02	0.17	1.93
J13325655-0301395	S	18.27	0.03	16.19	0.01	15.79	0.01	15.12	0.02	0.27	1.34
J13462001-0325407	S	17.21	0.02	14.62	0.01	14.22	0.01	13.48	0.02	2.57	0.71
J13462215-0325057	E	17.50	0.02	14.82	0.01	14.38	0.01	13.60	0.01	3.34	0.70
J14003661-0254327	S	17.54	0.02	14.91	0.01	14.50	0.01	13.79	0.01	4.94	0.41
J14003796-0254227	S	17.70	0.02	15.13	0.01	14.73	0.01	13.98	0.02	1.85	0.86
J14005783+4251203	S	18.53	0.03	15.94	0.01	15.43	0.01	14.20	0.03	0.16	2.54
J14005879+4250427	S	17.95	0.02	15.78	0.01	15.32	0.01	14.16	0.03	0.06	2.47
J14055079+6542598	S	18.47	0.03	15.98	0.01	15.54	0.01	14.72	0.09	0.66	1.34
J14055334+6542277	${\bf E}$	17.58	0.02	15.07	0.01	14.70	0.01	14.14	0.06	7.91	0.00
$J14062157 + 5043303$	${\bf S}$	14.02	0.02	12.46	0.01	12.20	0.01	11.78	0.01	0.13	1.19
J14064127+5043239	${\bf E}$	15.08	0.02	12.58	0.01	12.16	0.01	11.59	0.01	7.61	0.00
J14070703-0234513	${\bf S}$	19.56	0.04	16.73	0.01	16.24	0.01	15.39	0.04	5.96	0.61
J14070720-0234402	$\mathbf E$	19.35	0.03	16.52	0.01	16.08	0.01	15.26	0.03	9.04	0.39
J14234238+3400324	S	15.37	0.02	13.97	0.01	13.77	0.01	13.36	0.02	0.09	1.22
J14234632+3401012	S	16.35	0.02	14.18	0.01	13.78	0.01	13.01	0.02	0.24	1.56
J14245831-0303597	S	17.81	0.02	15.85	0.01	15.50	0.01	14.90	0.44	0.24	1.26
J14245913-0304012	S	19.41	0.03	16.38	0.01	15.87	0.01	14.92	0.43	12.91	0.48
J14250552+0313590	${\bf E}$	17.84	0.02	15.28	0.01	14.90	0.01	14.07	0.01	0.88	1.27
J14250739+0313560	S	18.70	0.03	16.31	0.01	15.96	0.01	15.10	0.03	0.38	1.57
J14294766+3534275	S	17.10	0.02	14.42	0.01	14.00	0.01	13.25	0.02	5.03	0.49
J14295031+3534122	S	17.73	0.02	15.27	0.01	14.88	0.01	14.20	0.03	1.55	0.79
J14334683+4004512	S	17.29	0.02	14.71	0.01	14.27	0.01	13.37	0.01	0.69	1.45
J14334840+4005392	S	16.33	0.02	14.45	0.01	14.11	0.01	13.21	0.01	0.07	2.02
J14442055+1207429	S	17.28	0.02	15.11	0.01	14.74	0.01	13.96	0.02	0.22	1.62
J14442079+1207552	S	17.59	0.02	14.86	0.01	14.44	0.01	13.66	0.01	5.77	0.47
J15002374+4316559	E	17.60	0.02	14.88	0.01	14.45	0.01	13.62	0.01	3.06	0.81
J15002500+4317131	S	17.81	0.02	15.05	0.01	14.63	0.01	13.83	0.02	6.08	0.50
J15053137+3427534	S	18.98	0.03	16.36	0.01	16.01	0.01	15.47	0.02	13.00	0.00
J15053183+3427526	E	19.10	0.03	16.24	0.01	15.83	0.01	15.12	0.01	13.00	0.00
continue											

Table2 continue

name	type	m_u	e_u	m_r	e_r	m_i	e_i	m_{K_s}	$\mathbf{e}_{\mathbf{K}_s}$	Age	A_v
		(mag)	(mag)	(Gyr)	(mag)						
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)
$J15064391 + 0346364$	S	17.56	0.02	14.90	0.01	14.50	0.01	13.71	0.01	2.88	0.76
J15064579+0346214	${\bf S}$	18.22	0.03	15.60	0.01	15.09	0.01	13.83	0.02	0.15	2.63
J15101587+5810425	$\mathbf S$	17.59	0.02	15.08	0.01	14.70	0.01	14.03	0.10	2.42	0.63
J15101776+5810375	S	19.02	0.03	16.48	0.01	15.95	0.01	14.49	0.15	0.06	3.01
J15144544+0403587	S	18.35	0.03	15.49	0.01	15.03	0.01	14.17	0.01	7.48	0.53
J15144697+0403576	S	18.26	0.03	15.52	0.01	15.12	0.01	14.32	0.01	4.77	0.59
J15233768+3749030	${\bf S}$	17.57	0.02	15.37	0.01	15.04	0.01	14.51	0.06	0.82	0.77
J15233899+3748254	E	17.83	0.03	15.32	0.01	14.93	0.01	14.38	0.06	8.72	0.00
J15264774+5915464	S	18.55	0.03	15.74	0.01	15.33	0.01	14.58	0.03	13.00	0.04
J15264892+5915478	E	18.39	0.03	15.65	0.01	15.25	0.01	14.56	0.03	13.00	0.00
J15281276+4255474	S	16.40	0.02	13.89	0.01	13.42	0.01	12.30	0.01	0.18	2.29
J15281667+4256384	S	16.49	0.02	13.79	0.01	13.35	0.01	12.54	0.02	3.04	0.78
J15523258+4620180	E	18.97	0.03	16.05	0.01	15.63	0.01	14.91	0.02	13.00	0.00
J15523393+4620237	S	19.09	0.03	16.70	0.01	16.25	0.01	15.48	0.03	0.58	1.27
J15562191+4757172	S	17.47	0.02	15.30	0.01	14.93	0.01	14.21	0.07	0.31	1.37
J15562738+4757302	E	17.59	0.02	14.99	0.01	14.60	0.01	13.98	0.03	9.50	0.02
J15583749+3227379	S	19.11	0.03	16.48	0.01	16.05	0.01	15.23	0.03	1.49	1.07
J15583784+3227471	S	18.13	0.02	16.03	0.01	15.60	0.01	14.71	0.28	0.12	2.06
J16024254+4111499	S	16.97	0.02	15.38	0.01	15.09	0.01	14.34	0.10	0.05	1.69
J16024475+4111589	S	17.37	0.02	15.80	0.01	15.52	0.01	14.88	0.16	0.07	1.57
J16080559+2529091	S	18.95	0.03	15.91	0.01	15.37	0.01	14.24	0.02	2.26	1.45
J16080648+2529066	S	19.22	0.03	16.18	0.01	15.53	0.01	13.99	0.01	0.19	3.02
J16082261+2328459	S	19.14	0.03	16.92	0.01	16.53	0.01	15.84	0.05	0.43	1.23
J16082354+2328240	S	18.50	0.03	16.08	0.01	15.66	0.01	14.63	0.02	0.20	2.09
J16145418+3711064	S	18.74	0.03	15.85	0.01	15.41	0.01	14.53	0.02	8.17	0.53
J16145421+3711136	\overline{E}	19.03	0.03	16.17	0.01	15.74	0.01	14.94	0.02	13.00	0.12
J16282497+4110064	S	17.88	0.02	15.25	0.01	14.85	0.01	14.23	0.02	12.99	0.00
J16282756+4109395	S	17.65	0.02	14.91	0.01	14.48	0.01	13.75	0.02	9.85	0.20
J16354293+2630494	S	19.36	0.03	16.44	0.01	15.97	0.01	15.18	0.04	13.00	0.00
J16354366+2630505	E	19.40	0.03	16.43	0.01	15.94	0.01	14.99	0.03	7.35	0.70
J16372583+4650161	S	19.11	0.03	16.19	0.01	15.72	0.01	14.70	0.04	2.52	1.21
J16372754+4650054	S	19.65	0.04	17.09	0.01	16.65	0.01	15.93	0.06	2.64	0.66
J17020320+1900006	E	19.30	0.03	16.17	0.01	15.70	0.01	14.94	0.02	13.00	0.00
J17020378+1859495	S	20.27	0.05	17.57	0.01	17.13	0.01	16.30	0.07	2.54	0.88
J17045089+3448530	${\bf S}$	18.90	0.03	16.73	0.01	16.26	0.01	15.30	0.10	0.11	2.18
J17045097+3449020	S	17.65	0.02	16.00	0.01	15.55	0.01	14.68	0.06	0.05	1.83
J20471908+0019150	S	15.73	0.02	12.82	0.01	12.34	0.01	11.49	0.04	12.15	0.33
J20472428+0018030	E	15.82	0.02	13.09	0.01	12.62	0.01	11.88	0.04	8.22	0.30

Notes: The columns are: (1) galaxy name; (2) galaxy type, "S" for Spiral and "E" for Elliptical; (3), (5), (7), (9) 7 kpc aperture photometry result of u, r, i and K_s -band, respectively, and (4), (6), (8), (10) their errors; (11) stellar age generated from $u-r-i-K_s$ color; (12) attenuation of V-band generated from $u-r-i-K_s$ color

name	\tilde{z}	$\overline{\mathrm{sSFR}_{\text{enh}}}$	pair type	bulge_type	merge_stage	ðν	$log M_{HI+H2}$	$t_{8\mu m, nulear}$	$t_{8\mu m, total}$
		(dex)				(km/s)	(M_{\odot})	(mJy)	(mJy)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
J01183417-0013416	0.0453	.34	SS	D	early	54	10.45	33.67	57.06
$J03381222+0110088$	0.0392	0.75	SE	c	early	432.9	10.45		
$J07543194 + 1648214$	0.0459	0.71	SS		late	104.1	10.09		
$J09155467 + 4419510$	0.0396	1.11	SS		late	2.4			
$J09155552 + 4419580$	0.0396	1.36	SS		late	2.4			
$J10100079 + 5440198$	0.0460	0.79	SS		late	83.7		179.52	187
J10332972+4404342	0.0523	0.84	SS		early	64.2		26.13	49.3
J13153506+6207287	0.0306	1.59	SS		early		10.05	39.86	55.36
J14005783+4251203	0.0327	0.82	SS		early	234.3	10.05	23.52	47.99
J14005879+4250427	0.0335	0.95	SS		early	234.3	9.88		
J16024254+4111499	0.0335	0.77	SS		early	69	10.38		
J17045089+3448530	0.0572	0.80	SS		late	270		22.49	77.56
J17045097+3449020	0.0563	1.04	SS		late	270		4.61	15.88

Table .3: SFGs with strong enhancement

Notes: The columns are: (1) galaxy name; (2) redshift; (3) specific star formation rate enhancement; (4) pair type: "SS" for Spiral+Spiral pair, "SE" for Spiral+Elliptical pair; (5) bulge type: "p" for pseudo-bulge, "c" for classical-bulge; (6) interaction type (detailed description in [Cao et al. 2016](#page-22-14)); (7) difference in radial velocity between two galaxies in a pair; (8) HI gas mass in [Zuo et al. \(2018](#page-23-12)) plus H2 gas mass in [Lisenfeld et al.](#page-22-12) [\(2019](#page-22-12)), where the HI mass is divided by M_{star} ratio for SS pairs, and all assigned to the spiral for SE pairs; (9) Spitzer 8 μ m flux of the galaxy nuclear region [\(Xu et al. 2010](#page-23-10)); (10) Spitzer 8 μ m flux of the total galaxy

Table .4: Galaxies in late-stage S+E mergers

name	\boldsymbol{z}	type	$log M_{star}$	sSFR	$\log M_{\rm gas}$	SDSS_id	GROUP _{id}	richness	rank
			(M_{\odot})		(M_{\odot})				
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
J10514368+5101195	0.0250	Е	11.12	-11.86	8.71	189845	1745	12	
$J10514450 + 5101303$	0.0238	S	10.85	-11.64	< 8.66	189846	1745	12	2
J11542299+4932509	0.0702	S	10.98	-11.35	9.49				
J11542307+4932456	0.0712	E	11.35	-11.72	<9.48	229360	215518		
J13131429+3910360	0.0716	E	11.26	-11.43	<9.36	445348	47	103	
J13131470+3910382	0.0716	S	10.95	-11.12	< 9.48	445349	47	103	2
J16354293+2630494	0.0701	S	11.27	-11.31	<9.52				
J16354366+2630505	0.0713	E	11.27	-11.63	<9.55	306304	177	49	

Notes: The columns are: (1) galaxy name; (2) redshift; (3) galaxy morphology: "S" for Spiral, "E" for Elliptical; (4) stellar mass; (5) specific star formation rate; (6) gas mass from dust mass [\(Cao et al. 2016](#page-22-14)); (7) SDSS-ID; (8) Group-ID; (9) richness (i.e. number of galaxies in the group); (10) rank (1=most massive, 2=other galaxies). Data in column (7)–(10) are from [Yang et al.](#page-23-16) [\(2012](#page-23-16)). Those galaxies with no data in columns (7)–(10) are not included in the catalog of [Yang et al. \(2012](#page-23-16)) for lack of spectroscopic redshift