

Dianoga SIDM: galaxy cluster self-interacting dark matter simulations

Antonio Ragagnin,^{1,2,3*} M. Meneghetti,^{1,4} F. Calura,¹ G. Despali,^{2,1,4} K. Dolag,^{5,6} M. S. Fischer,^{5,7} C. Giocoli,^{1,4} L. Moscardini^{2,1,4}

¹ INAF-Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, Via Piero Gobetti 93/3, I-40129 Bologna, Italy

² Dipartimento di Fisica e Astronomia "Augusto Righi", Alma Mater Studiorum Università di Bologna, via Gobetti 93/2, I-40129 Bologna, Italy

³ IFPU - Institute for Fundamental Physics of the Universe, Via Beirut 2, I-34014 Trieste, Italy

⁴ INFN-Sezione di Bologna, Viale Berti Pichat 6/2, I-40127 Bologna, Italy

⁵ Universitäts-Sternwarte, Fakultät für Physik, Ludwig-Maximilians-Universität München, Scheinerstr. 1, D-81679 München, Germany

⁶ Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, D-85748 Garching, Germany

⁷ Excellence Cluster ORIGINS, Boltzmannstrasse 2, D-85748 Garching, Germany

submitted

ABSTRACT

Context. Self-interacting dark matter (SIDM) can tackle or alleviate small-scale issues within the cosmological standard model Λ CDM, and diverse flavours of SIDM can produce unique astrophysical predictions, resulting in different possible signatures which can be used to test these models with dedicated observations of galaxy clusters.

Aims. This work aims at assessing the impact of DM self-interactions on the properties of galaxy clusters. In particular, the goal is to study the angular dependence of the cross section by testing rare (large angle scattering) and frequent (small angle scattering) SIDM models with velocity-dependent cross sections.

Methods. We re-simulate six galaxy cluster zoom-in initial conditions with a dark matter only run and with a full-physics setup simulations that includes a self-consistent treatment of baryon physics. We test the dark matter only setup and the full physics setup with either collisionless cold dark matter, rare self-interacting dark matter, and frequent self-interacting dark matter models. We then study their matter density profiles as well as their subhalo population.

Results. Our dark matter only SIDM simulations agree with theoretical models, and when baryons are included in simulations, our SIDM models substantially increase the central density of galaxy cluster cores compared to full-physics simulations using collisionless dark matter. SIDM subhalo suppression in full-physics simulations is milder compared to the one found in dark matter only simulations, because of the cuspiest baryonic potential that prevent subhalo disruption. Moreover SIDM with small-angle scattering significantly suppress a larger number of subhaloes compared to large angle scattering SIDM models. Additionally, SIDM models generate a broader range of subhalo concentration values, including a tail of more diffuse subhaloes in the outskirts of galaxy clusters and a population of more compact subhaloes in the cluster cores.

Key words. Galaxies: clusters: general – Cosmology: cosmological parameters – Galaxy: formation – method: numerical – Hydrodynamics

1. Introduction

Galaxy clusters, recognised as the most extensive gravitationally bound systems of galaxies (see Kravtsov & Borgani 2012, for a comprehensive review), play a pivotal role as cosmic laboratories for investigating the large-scale structures within our Universe. The underlying cosmological model impacts their masses (Tinker et al. 2008; Spergel & Steinhardt 2000; Despali et al. 2016; Despali & Vegetti 2017), their lensing signal (Natarajan & Kneib 1997; Moore et al. 1998), the abundance of substructures within them (Tormen et al. 1998; Natarajan & Springel 2004), and their concentration (Giocoli et al. 2012b, 2013; Ragagnin et al. 2021).

The current standard concordance cosmological model Λ CDM assumes the presence of a cosmological constant and a type of dark matter (DM) that is cold and collisionless. Numer-

ical simulations performed within this paradigm are currently showing some tensions with observational data. For instance simulations of galaxy cluster cores (as in Meneghetti et al. 2020) reveal subhaloes that are considerably less compact than their observed counterparts (as illustrated in Bergamini et al. 2019; Granata et al. 2022).

Specifically, the galaxy-galaxy strong lensing (GGSL) signal resulting from simulated substructures of mass $M \approx 10^{11} M_{\odot}$ can be up to a factor of 2 lower than observed values (Meneghetti et al. 2022; Ragagnin et al. 2022; Meneghetti et al. 2023). Concerning the higher mass range of satellites ($M > 10^{11} M_{\odot}$), hydrodynamic simulations can indeed reproduce high mass subhaloes with compactness as high as the one from scaling relations (see the simulations presented in Bahé 2021; Robertson 2021). This can be achieved, for instance, by assuming particularly low AGN efficiencies. However, it is noteworthy that the lensing signal derived from observational data is predominantly influenced by satellites with lower masses (Ragagnin et al.

* e-mail: antonio.ragagnin@unibo.it

2022). In fact the study conducted by Ragagnin et al. (2022) indicates that relying solely on AGN physics does not appear to find a solution to the compactness mismatch without compromising the consistency with observed properties related to stellar mass.

Furthermore, Λ CDM simulations face challenges in reproducing ultra-diffuse galaxies (UDGs, as studied by van Dokkum et al. 2015; Mowla et al. 2017; Greco et al. 2018), whose origin and low concentration parameter may be explained by satellites with cored dark matter profiles (Carleton et al. 2019). Although baryonic processes can also play a crucial role in lowering subhalo compactness (Di Cintio et al. 2019), Λ CDM simulations still fall short in reproducing the UDGs low circular velocity (see Sales et al. 2020, particularly their Fig. 5).

In this paper, we explore the impact of a type of dark matter that is not collisionless. In particular, we study self-interacting DM (SIDM, Spergel & Steinhardt 2000) in the context of hydrodynamical simulations of galaxy clusters. Motivated by particle physics, SIDM models consider interactions between DM elementary particles and a massive mediator through, for instance, a Yukawa potential (Loeb & Weiner 2011). SIDM maintains the theoretical expectations of cold and collisionless DM at large scales while significantly influencing subhalo properties (see, for example, Spergel & Steinhardt 2000; Tulin & Yu 2018; Adhikari et al. 2022; Mastromarino et al. 2023), resulting in objects with a broader range of concentration parameter compared to collisionless CDM.

Notably, DM self-interactions smooth out the density distribution in the central regions of DM subhaloes, leading to profiles that are more cored than their collisionless counterparts lowering the subhalo compactness (see, for instance, Carleton et al. 2019; Nadler et al. 2023; Kong et al. 2022). Moreover SIDM subhaloes that formed with an initially high concentration, will eventually reveal, after the core expansion reaches its minimum, a core-collapse stage that will make them much more compact than their analogues in the Λ CDM model (Outmezguine et al. 2023; Carton Zeng et al. 2023), thus explaining the excess of subhalo compactness observed in galaxy clusters (Yang & Yu 2021).

In this work, we compare two classes of SIDM models. The first is the so-called frequent SIDM (fSIDM) model, which assumes small-angle scattering, in the limit of keeping the momentum transfer cross section fixed but decreasing the scattering angles. The second is the so-called rare SIDM (rSIDM) model, which assumes isotropic scattering. Under these conditions, scattering events are much more frequent in the fSIDM than in the rSIDM model, hence the adjectives frequent and rare. Simulations with a self-consistent treatment of baryonic physics (hereafter FP, as full physics) are necessary to properly study and capture the evolution of galaxy cluster substructures (Kimmig et al. 2023). Therefore in this work we plan to re-simulate our galaxy clusters both with dark matter only (DMO) simulations and FP ones.

The two models can alter the DM distribution in different ways, with fSIDM being more efficient in generating DM-galaxy offsets during mergers because of its small-angle dependency (Fischer et al. 2021). We use the implementation presented in Fischer et al. (2021) and Fischer et al. (2022), a technique that relies on describing the self-interaction between DM particles through an effective drag force, as presented in Kahlhoefer et al. (2014).

The studies of Sagunski et al. (2021), Andrade et al. (2022), Harvey et al. (2019), and Eckert et al. (2022) established a lower limit on the impact of SIDM on cluster scales. Specifically, considering that clusters adhere to an Einasto pro-

Table 1. Regions re-simulated in this work. Different columns from left to right report the name, the virial mass, and the NFW concentration at $z = 0.2$. We present the concentration parameter on the third and fourth columns, computed both in the DMO and FP runs respectively.

region	$M_{\text{vir}}[10^{14}h^{-1}M_{\odot}]$	c_{vir} DMO	c_{vir} FP
D3	4.8	4.0	4.6
D4	2.7	4.6	3.9
D5	1.2	6.0	5.5
D10	11.2	4.1	4.5
D15	11.5	4.8	5.9
D16	12.3	4.6	5.0

Notes. We show mass and concentration of collisionless DM only, because these value are almost identical for the SIDM runs.

file (Einasto 1965), the work of Eckert et al. (2022) shows that the total cross section per unit dark matter particle mass m_{χ} should be constrained to $\sigma/m_{\chi} < 0.19 \text{ g}^{-1} \text{ cm}^2$. This value is notably low when compared to the typical values found in theoretical studies, which hover around $\sigma/m_{\chi} \approx 1 \text{ g}^{-1} \text{ cm}^2$ as used in Mastromarino et al. (2023) (note however that they studied a mass-range that is different than ours).

While this might initially appear as a challenge for SIDM, it is important to note that elementary particle interactions, such as those governed by the Yukawa potential, exhibit cross sections which depend on the relative velocity of the particles. The cross section decreases as a function of the relative velocity, making it applicable for constraining both dwarf spheroidal profiles and group cores (Correa 2021). Consequently, in this study, our focus will be solely on SIDM models with a velocity-dependent cross section.

The paper is structured as follows: we present our suite of zoom-in simulations with collisionless DM and SIDM in Sect. 2; we investigate the overall DM distribution in the cluster cores in Sect. 3; we study the cluster subhalo population in Sect. 4; we draw our conclusions in Sect. 5. Throughout this paper, we use the term "collisionless" to describe the cold and collisionless DM assumed in the standard Λ CDM, distinguishing it from SIDM, which is also a type of cold DM.

2. Numerical Setup

Our simulations are performed using OpenGadget3, a code derived from P-Gadget3, a successor of P-Gadget2 (Springel 2005). The initial conditions mirror those employed in the Di-anoga simulations (as utilised, for instance, in Bonafede et al. 2011; Rasia et al. 2015). These conditions are generated from a parent DMO box with a side length of 1 comoving h^{-1} Gpc and are specifically tailored for conducting hydrodynamic simulations of galaxy clusters. The power spectrum assumed in the parent box corresponds to a Λ CDM model with cosmological parameters $\Omega_m = 0.24$, $\Omega_b = 0.04$, $n_s = 0.96$, $\sigma_8 = 0.8$, and $h = 0.72$.¹ In this work, we re-simulated the initial conditions for six galaxy cluster regions that encompass virial masses in the range $10^{14} - 10^{15} M_{\odot}$ and analyse the results in 15 redshift slices between $z = 0.2 - 0.6$.²

¹ Note that SIDM is a type of collisional, cold DM, therefore the three DM models (collisionless DM, rSIDM, fSIDM) can consistently share the same initial conditions (as opposed to what would happen with warm dark matter).

² Note that the virial mass M_{vir} is defined as the mass within the so-called virial radius R_{vir} , that encloses an average density of

We maintain the same resolution level as presented in [Ragagnin et al. \(2022\)](#). Specifically our simulations employ a gravitational softening of $\epsilon_{\text{DM}} = 3.7 h^{-1} \text{ckpc}$ for DM particles and $\epsilon_{\star} = 2.0 h^{-1} \text{ckpc}$ as the softening parameter for stellar gravitational interactions. The DM particle masses are set to $m_{\text{DM}} = 8.3 \times 10^8 h^{-1} M_{\odot}$.

2.1. Baryon physics

To follow the baryon physics, we simulate the hydrodynamics of gas using an enhanced smoothed particle hydrodynamics (SPH) solver presented in [Beck et al. \(2016\)](#).³ The stellar evolution scheme from [Tornatore et al. \(2007\)](#) which follows 11 chemical elements (H, He, C, N, O, Ne, Mg, Si, S, Ca, Fe) with input cooling tables generated using the CLOUDY photo-ionisation code ([Ferland et al. 1998](#)). For a comprehensive understanding of the modelling of supermassive black holes and energy feedback, readers can refer to [Springel et al. \(2005\)](#); [Fabjan et al. \(2010\)](#) and [Hirschmann et al. \(2014\)](#), where prescriptions for black hole growth and feedback from AGNs are thoroughly described. The identification of haloes and their member galaxies is accomplished using the friends-of-friends halo finder ([Davis et al. 1985](#)) and an improved version of the subhalo finder SUBFIND ([Springel et al. 2001](#)), which accounts for the presence of baryons ([Dolag et al. 2009](#)).

Our feedback scheme is based on the Magneticum subgrid physics model (e.g., [Teklu et al. 2015](#)). The Magneticum AGN model is derived from [Hirschmann et al. \(2014\)](#) and, with a calibration similar to Magneticum suite of simulations, we set a radiative efficiency of $\epsilon_r = 0.2$ to regulate the luminosity of the radiative component and we set the feedback energy per unit time to have a contribution of $\epsilon_f = 0.075$ of the luminosity, as detailed in Eqs. 7-12 in [Steinborn et al. \(2015\)](#).

2.2. Dark matter models

As mentioned in the introduction, in addition to collisionless DM, in this work we will explore the SIDM models implemented in [Fischer et al. \(2023\)](#), which have the advantage of being capable of simulating interactions in the limit of very anisotropic cross sections and of consistently conserving the total energy of the system.

In our SIDM models, DM interactions occur via a light-scalar mediator or a Yukawa potential. An approximation of the scattering cross section, derived using the Born approximation, yields a momentum transfer cross section σ_T expressed as

$$\frac{\sigma_T(v)}{m_{\chi}} = \frac{\sigma_{T0}}{m_{\chi}} \left(1 + \left(\frac{v}{v_c} \right)^2 \right)^{-2}, \quad (1)$$

Here, m_{χ} is the dark matter particle mass, v represents the relative velocity between particles, σ_{T0} denotes a low-velocity plateau normalisation, and v_c represents the knee position before a v^{-4} dependency of the cross section comes into play.

We choose the parameters σ_{T0} and v_c in order for the cross section to impact mainly low mass subhaloes (thus σ must be high for objects with mass $M < 2 \times 10^{10} M_{\odot}$). Moreover we want

³ We used a space-filling curve-aware neighbour search ([Ragagnin et al. 2016](#)).




Fig. 1. SIDM cross section as in Eq. (1). Blue curve reports the cross section with the parameters chosen for this work ($\sigma_{T0} = 40 \text{ g}^{-1} \text{ cm}^2$ and $v_0 = 200 \text{ s}^{-1} \text{ km}$). Vertical shaded area corresponds to the median velocity dispersion for low mass subhaloes ($M < 2 \times 10^{10} M_{\odot}$) in cluster cores, high mass subhaloes ($M > 4 \times 10^{11} M_{\odot}$) in cluster cores, and the typical velocity dispersion for haloes with $M_{\text{vir}} \in [2 - 10] 10^{14} M_{\odot}$.

the cross section to be relatively small for high mass subhaloes (for masses as $M > 4 \times 10^{11} M_{\odot}$, as we are not interested in modifying their properties), and it has to be negligible at cluster scales.

Given these constraints we decided to use the functional form of Eq. (1) with $\sigma_{T0}/m_{\chi} = 40 \text{ cm}^2 \text{ g}^{-1}$ and $v_c = 200 \text{ s}^{-1} \text{ km}$, similar to the constraints from dwarf spheroidal galaxies proposed in [Correa \(2021\)](#). In Fig. 1 present our velocity dependent cross section, together with the relevant mass scales. Here it is easy to appreciate the decreasing importance of the cross section for increasing object masses: small subhaloes ($v \approx 100 \text{ s}^{-1} \text{ km}$) will be greatly impacted by SIDM ($\sigma/m_{\chi} \approx 40 \text{ cm}^2 \text{ g}^{-1}$), while galaxy clusters ($v \approx 1000 \text{ s}^{-1} \text{ km}$) will be only slightly affected by SIDM ($\sigma/m_{\chi} \lesssim 0.1 \text{ cm}^2 \text{ g}^{-1}$).

2.3. Halo Sample

We re-simulate the initial conditions of the regions with DMO and FP simulations and with three dark matter types: collisionless DM, rSIDM (in our case referring to an isotropic cross section), and fSIDM, which means that each galaxy cluster is resimulated six times. We report the main characteristics of our galaxy cluster sample masses and concentration at the lowest redshift available ($z = 0.2$) in Table 1. Here we estimate the dynamical state of haloes by computing the so called concentration parameter (which is known to be a good proxy for it, see e.g., [Ludlow et al. 2012](#)), with objects with higher concentration being mostly early-formed and more relaxed. We compute the concentration parameter by assuming a Navarro–Frank–White (NFW, [Navarro et al. 1997](#)) profile ρ_{NFW} so that

$$\rho_{\text{NFW}}(r) = \frac{\rho_0}{(r/r_s)(1+r/r_s)^2}, \quad (2)$$




Fig. 2. Projected density maps of central region (within $r < 0.15R_{\text{vir}}$) of the D16 region re-simulations at our lowest redshift slice of $z = 0.2$. The upper row shows DMO simulations, while the bottom row refers to FP simulations. Columns from left to right show: collisionless DM simulations, rSIDM simulations, and fSIDM simulations, respectively.

where ρ_0 and r_s are free parameters and the concentration is defined as $c \equiv R_{\text{vir}}/r_s$.⁴ We see that our haloes have concentrations in the range $c \in [2.8, 5.5]$ (we used the predictions from [Ragagnin et al. 2021](#)). Note that region D5 has a higher than average concentration implying that it is a relaxed system ([Ludlow et al. 2012](#)). We report the concentration for both the DMO and FP runs as they are known to differ (see e.g. [Duffy et al. 2010](#)), with DMO concentration being slightly lower than FP concentration. For what concerns the concentration difference between collisionless DM and SIDM, we found no significant changes, as expected SIDM affect the matter distribution mainly on small scales.

Figure 2 shows the projected density maps of the central region (where SIDM has most impact) of the most massive region (D16) at $z = 0.2$ for the corresponding six re-simulations (three DM models with and without baryon physics). Moving from DMO to FP simulations we see much more peaked substructures and that in general SIDM models tend to have substructures that are displaced differently compared to the CDM model, due to the expected different trajectories during mergers (as shown in [Sabarish et al. 2023](#)).

⁴ The fit is performed by minimising the sum of squared log-residuals over 20 logarithmically spaced bins in the range $[0.01, 1]R_{\text{vir}}$.

3. Galaxy cluster matter density profiles

In this section, we will concentrate on the distribution of matter and subhaloes within the cluster cores, specifically within a cluster-centric distance $r < 0.15R_{\text{vir}}$. The reason behind this choice is twofold: on these small scales the effect of SIDM is expected to be largest, as it is a region sensitive to the nature of DM models (see, e.g., [Natarajan et al. 2007](#); [Yang & Yu 2021](#)); moreover, this is the typical field of view of galaxy-galaxy strong lensing analyses from Hubble Frontier Field data (see, e.g., [Bergamini et al. 2019](#); [Granata et al. 2022, 2023](#)).

Figure 3 presents the stacked density profiles within the central regions of our DMO simulations, encompassing redshift slices between $0.2 < z < 0.6$. Given the broad virial mass range across our six regions, we opted to stack all haloes together by rescaling their profiles using the virial radius and virial mass. As expected SIDM simulations yield cored profiles (see e.g., [Kamada & Kim 2020](#)), in particular our SIDM DMO simulations form a core at radii $r < 2 \times 10^{-2}R_{\text{vir}}$. The ratio between collisionless and self-interacting DM (see bottom panel of Fig. 3) becomes significantly different from unity at cluster centric distances $\lesssim 2 \times 10^{-2}R_{\text{vir}}$ (approximately corresponding to a cluster-centric distance of $\approx 100\text{kpc}$). It is worth noting that the profiles are different at distances larger than the fiducial DM softening




Fig. 3. Comparison between stacked central density profile in the DMO runs. We report collisionless DM density profile as blue solid line, rSIDM as green dashed line, and fSIDM as orange dotted line (note that rSIDM and fSIDM lines almost overlap visually). Shaded areas correspond to one standard deviation of the six regions. Upper panel shows the central density, normalised by the average density at R_{vir} for our six regions, while the bottom panel shows the ratio between the profiles of the SIDM runs and the collisionless DM runs. Gray shaded area cores one standard deviation around the fractional fiducial DM softening $2.8 \times \epsilon/R_{\text{vir}}$.

($2.8\epsilon_{\text{DM}}$, indicated by the grey vertical band). Consequently we can conclude that in DMO SIDM simulations, the impact extends across the entire cluster central regions with both SIDM variants (rSIDM and fSIDM) exhibiting $\approx 20\% - 40\%$ lower density compared to collisionless DM.

We present results for the FP simulations in Fig. 4 (top row), where the stacked density profiles of DM, stellar, and total density for our six Dianoga regions are shown. Notably, in SIDM FP simulations, both rare and frequent scenarios tend to exhibit DM central densities that are more pronounced than those observed in the corresponding FP collisionless DM simulations. Notice that Robertson et al. (2019) reported an opposing result, demonstrating that their hydrodynamic SIDM simulations produced cluster cores that, while less cored than the DMO run, still retained a degree of core structure compared to the collisionless run. The reason for this difference remains unclear, however we can speculate that since the baryonic central potential is known to impact the central density evolution (see e.g., Elbert et al. 2018), the different results may be attributed to different central baryon density in our study compared to that in Robertson et al. (2019).

To verify this statement we will conduct the following two analyses: (I) in order to ensure to have our FP collisionless DM profile under control, in Sect. 3.1 we will first verify that the increased DM cuspsiness in collisionless DM runs is due to adiabatic contraction; then (II) in Sect. 3.2 we will compare our SIDM central densities with analytical models of gravothermal

solutions of SIDM haloes in the literature to understand if our increased central density agrees with gravothermal evolution models.

3.1. Adiabatic Contraction

FP simulations (compared to DMO ones) produce a heightened concentration of DM close the cluster centre (as in Despali et al. 2020) due to adiabatic contraction (as found above and for example in Gnedin et al. 2004), where the baryon density in halo cores is high enough to generate a back reaction on the dark matter component and increase its central density significantly compared to DMO runs, see for instance Fig. 5 in Duffy et al. (2010).

In this subsection we will verify that we have under control the results of our collisionless DM simulations, we will use only collisionless DM runs (both FP and DMO), and estimate the contribution to the central baryon density from adiabatic contraction following the theoretical model proposed by Blumenthal et al. (1986). They derive that the halo radius r times mass within it is a constant, namely

$$rM_{\text{dm},i}(< r) = r[M_{\text{dm},f}(< r) + M_{\text{b}}], \quad (3)$$

where $M_{\text{dm},i}$ is the initial dark matter mass, $M_{\text{dm},f}$ is the final dark matter mass and $M_{\text{b},f}$ is the final baryon mass. In our work, we estimate $M_{\text{dm},i}$ as the dark matter mass from DMO runs, $M_{\text{dm},f}$ as the dark matter mass in the FP runs, and $M_{\text{b},f}$ as the baryon mass in the same FP runs. We cross-matched the DMO and FP masses of each halo in all available snapshot and computed the adiabatic contraction factor using the software *contra* (Gnedin et al. 2004).

We report our finding in Fig. 5 (top panel), where we show the stacked DM mass profile of our haloes from collisionless DM simulations from both the DMO and FP runs. The bottom panel of Fig. 5 reports their ratio together with the median prediction from the model by Blumenthal et al. (1986) for each of the cross-matched haloes (the size of the shaded area represents the error on the mean). We can see that the model by Blumenthal et al. (1986) correctly predicts the increased central density of the FP simulations, and we therefore confirmed that the increased DM density found in our collisionless FP (compared to collisionless DMO) simulations is due to adiabatic contraction produced by baryons.

3.2. Gravothermal Solution

We will compare the central density of our haloes with the gravothermal solution for SIDM haloes proposed in Yang et al. (2023). Their model is based on the assumption that evolution of central density and core radius of SIDM NFW haloes can be described by a single parameter β once variables are rescaled as follows: $t_0 = 1/\sqrt{4\pi G\rho_s}$, radii by factor $r_0 = r_s$, densities by a factor $\rho_0 = \rho_s$, and cross sections by a factor $(\sigma/m_\chi)_0 = 1/(\rho_s r_s)$. We denote rescaled quantities with a hat.

We then estimate the core size of our DM haloes of SIDM simulations using the DM profile proposed in their Eq. A5, that is a simil-NFW profile with a possible core radius r_c and a truncation r_{out} , namely

$$\hat{\rho}(\hat{r}) = \frac{\hat{\rho}_c}{1 + (\hat{r}/r_c)^s(1 + \hat{r}/r_{\text{out}})^{3-s}}, \quad (4)$$

where $\hat{\rho}_c$ is the re-scaled central density, and we use $s = 2.19$ as proposed by Yang & Yu (2021).




Fig. 4. Comparison between stacked central density profiles in the FP runs. DM models are represented as in Fig. 3. Columns refer to the results for DM, stellar, and total matter components respectively. Upper panels show DM density profiles, while lower panels show the ratio with respect to the collisionless DM profiles. Gray shaded area represents one standard deviation around the fractional fiducial stellar softening $2.8 \times \epsilon_*/R_{\text{vir}}$.




Fig. 5. Comparison between DM mass profiles of collisionless DM runs. Top panel shows the stacked DM mass profile of DMO runs (black) and FP runs (blue), while bottom panel shows their ratio and the central value predicted by the adiabatic contraction model presented in Blumenthal et al. (1986) as light-blue shaded blue area. The vertical gray shaded area corresponds to one-standard deviation of the DM gravitational softening.

Note that the time rescaling factor t_0 proposed by Yang et al. (2023) does not take into account that FP SIDM simulations have a much shorter evolution time compared to DMO ones (as shown in Zhong et al. 2023). Therefore, for FP simulations we follow the approach of Zhong et al. (2023) and we rescale the time variable by combining Eqs. 15 and 16 in Zhong et al. (2023), namely by a factor of $(\rho_{\text{eff}} \sqrt{|\Phi(0)_{\text{FP}}|}) / (\rho_s \sqrt{|\Phi(0)_{\text{DMO}}|})$, where $\Phi(0)_{\text{DMO}}$ and $\Phi(0)_{\text{FP}}$ are the central gravitational potential, for the DMO and FP counterpart of a halo, and ρ_{eff} is an effective density that captures the contraction effect due to the baryonic




Fig. 6. Rescaled DM central density $\hat{\rho}_c$ as a function of rescaled time $\beta\hat{\sigma}t$, for our FP SIDM (red circles) and DMO SIDM (gray stars) runs. We also overplot the model from (Yang et al. 2023, black solid line). Red shaded area shows the mean value of the FP SIDM points with its one standard deviation, while blue shaded area shows the same for collisionless FP runs.

potential. To this end, for each halo we estimate ρ_{eff} as the central density of baryons and we estimate it using their Eq. 17, namely $\rho_{\text{eff}} \approx \rho_s + \alpha M_b / (2\pi r_s r_h)$, where M_b is the baryon mass of the halo and they set $\alpha = 0.4$. Following their procedure, we compute r_h by fitting an Einasto (Hernquist 1990) profile to the baryonic component (see Appendix A for the fit the details).

We then estimate the cross section acting on the galaxy cluster core by computing the effective cross section σ_{eff} similarly to Yang & Yu (2022), where the average cross-section is weighted with fifth power of the velocity and assuming that the velocities are well described by a Maxwell–Boltzmann distribution (with a characteristic velocity $V_{\text{max}}/\sqrt{3}$, see their Eq. 4.2):




Fig. 7. Comparison between stacked velocity dispersion profiles. Top panel presents the stacked velocity dispersion profiles for our FP simulations galaxy clusters in units of virial velocity $\sqrt{GM_{\text{vir}}/R_{\text{vir}}}$. Bottom panel: ratios with respect to the DMO counter parts. Lines are coloured as in Fig. 4.

$$\sigma_{\text{eff}} = \frac{3}{2} \frac{\langle v^5 \sigma_v(v) \rangle}{\langle v^5 \rangle}, \quad (5)$$

where σ_v is the viscosity cross section given by

$$\sigma_v = 4\pi \int_0^1 \frac{d\sigma}{d\Omega} \sin^2 \theta d \cos \theta. \quad (6)$$

We compute the rescaled time variable $\beta \hat{\sigma} \hat{t}$ as proposed in Yang et al. (2023), where β is the free parameter of their model that should vary around one, therefore we keep it as $\beta = 1$ for simplicity. In Fig. 6 we compare our DM central densities $\hat{\rho}_c$ for both the DMO (gray stars) and FP (red circles) SIDM runs with the prediction of $\hat{\rho}_c(\beta \hat{\sigma} \hat{t})$ from Yang et al. (2023). We can see that DMO $\hat{\rho}_c$ lies in the core-formation phase of Yang et al. (2023) central density evolution, namely in the left part of the plot.

For what concerns the FP simulation, we present mean values (with their one standard deviation) for $\hat{\rho}_c$ in the case of SIDM (red shaded area) and collisionless FP (blue shaded area) simulations. The mean value of $\hat{\rho}_c$ for collisionless FP simulations exceeds the corresponding value in the case of DMO, consistently with expectations from adiabatic contraction. On the other hand, the SIDM FP values of $\hat{\rho}_c$ are higher than the central densities found in collisionless DM FP simulations. This increase is due to the baryonic potential even if the haloes are not in a core-collapse phase. In support to this hypothesis, we illustrate the velocity dispersion profiles of our clusters in Fig. 7, which reveals that the central region of SIDM clusters exhibits an increasing profile, therefore even if the profile is less steep than their collisionless FP counterpart, the FP SIDM clusters are still in their core formation phase (see Appendix A for more details on the density and velocity dispersion time evolution).

4. Subhalo population

In this Section, we analyse the subhalo population in the central regions of our simulated clusters. In particular, we focus first on comparing SIDM subhalo abundances, then we explore whether these values agree with theoretical models on tidal stripping, and finally, we study subhalo compactness as this has potential value for future lensing analyses.

4.1. Subhalo abundance

Our subhalo abundance analyses are specifically focused on subhaloes within a cluster-centric distance of $< 0.15R_{\text{vir}}$ as the central part of the cluster is the one most affected by SIDM properties. Moreover, the central region of clusters is also relevant for strong lensing investigations, as highlighted in Meneghetti et al. (2023). However, more precise comparisons are needed in the future, for instance, considering projection effects and precise computation of the reduced shear to compare simulated and observed data correctly.

In the top panel of Fig. 8 (top panel), we present the cumulative Subhalo Mass Function (SHMF) for our sample at low redshift in the DMO simulations. Notably, there is a pronounced suppression of low-mass subhaloes in these simulations. Satellite suppression is expected in SIDM simulations because of the enhanced self-interactions of satellites with the host (Nadler et al. 2020). In the remaining top panels of Fig. 8, we depict the FP SHMF and dissect it based on the different matter components, including stars and DM. The relative values of the cumulative mass function are presented in the bottom panels of Fig. 8, revealing that SIDM has a smaller impact on the subhalo population in FP simulations compared to DMO simulations. The suppression of satellites in DMO simulations can be substantial, reaching approximately $\approx 50\%$ for rSIDM runs and going as high as $\approx 65\%$ for fSIDM simulations. In FP simulations, the suppression of satellites is more moderate, though still significant, with approximately $\approx 10\%$ for rSIDM and $\approx 35\%$ for fSIDM in terms of total masses. An increased suppression of satellites in fSIDM models compared to rSIDM is expected; see, for instance, Fischer et al. (2022) and Fischer et al. (2023). In general, the smaller satellite suppression produced by SIDM in FP simulations is due to the presence of a stellar component that makes subhaloes more resistant to disruption.

4.2. Tidal stripping

We now investigate the subhalo suppression that we found in SIDM simulations in the light of other theoretical studies. In particular, we want to test if this suppression of substructures is consistent with theoretical works as Peñarrubia et al. (2010), which shows that the suppression details are strongly dependent on the inner log-slope γ of their matter profiles.

In the left panel of Fig. 9, we present the stacked cumulative density profiles of subhaloes in the outskirts of galaxy clusters (outside $r > 0.5R_{\text{vir}}$) for our three DM models in DMO simulations, that we assume to be in an infalling phase. We fit the internal log-slope of the total matter profile and assume it equals to $(3 - \gamma)$ between the softening and 20 kpc. We opted for this radial range because it is large enough to capture the internal log-slope and small enough not to encounter the flattening in the tail of the profile. To show that our radial range is small, we over-plot the mean half mass radius as vertical lines, where we can notice that they are larger than 20 kpc. We can see that the DMO collisionless run has $\gamma \approx 1$, which corresponds to an NFW-like




Fig. 8. Cumulative SHMFs in units of the virial mass for subhaloes with a cluster centric distance $< 0.15R_{\text{vir}}$. The left most panel focus on DMO run with collisionless DM and rare and frequent SIDM; the other panels refer on FP simulations and show the results for different mass components: from left to right, total matter, stellar matter and DM. Colours and line styles are as in Fig. 3. The panels in the bottom row show the ratio between the SHMFs in the SIDM models and the one for the collisionless DM model. Shaded area corresponds to one standard deviation of the mean value.




Fig. 9. Stacked cumulative matter profiles of subhaloes in the outskirts of galaxy clusters (outside $r > 0.5R_{\text{vir}}$), for DMO (left panel) and FP (right panel) simulations for collisionless (blue dotted lines), rSIDM (green dashed line), and fSIDM (orange dotted line) models. Solid lines show the profile fitted in the internal region ($r < 20$ ckpc). Vertical dashed lines correspond to the mean half-mass radius.

core, as expected from theory. The DMO SIDM runs have a significantly lower profile, with fSIDM having a flatter profile than rSIDM, in agreement with the SHMF that we presented in Fig. 8, where fSIDM has a larger suppression than rSIDM.

In the right panel of Fig. 9, we present the same results but for FP simulations. Our γ values correspond to a cusp (as expected, baryons dominate the central part) and both SIDM and collisionless runs have very close values that are in agreement with the much milder SIDM suppression of satellites that we found in the SHMF shown in Fig. 8.

It is possible that the significant subhalo suppression of fSIDM with respect to rSIDM in FP simulations is due to a dif-

ferent effect than tidal stripping; in fact, the two models have similar density profiles (see Fig. 9). Therefore, one can speculate that the different SMHFs could be attributed to the differences in the host potential (Peñarrubia et al. 2010); the fact that the tests performed in Peñarrubia et al. (2010) assume a static host halo; or the effect of dark matter self-interactions between the infalling subhalo and the host.

4.3. Subhaloes compactness

We will now investigate how SIDM impacts the compactness of subhaloes in galaxy cluster cores, as it is the region that is most




Fig. 10. Circular velocity profiles for FP simulations with a cluster centric distance $< 0.15R_{\text{vir}}$. Each column shows a different matter component: total matter on the left, stellar component in the middle and the DM component on the right. Top panel: low mass regime $[1, 2]h^{-1}10^{10}M_{\odot}$, bottom panel: high mass regime $[10, 40]h^{-1}10^{10}M_{\odot}$. Line styles are as in Fig. 3.




Fig. 11. Subhalo compactness and stellar fraction for subhaloes for cluster centric distances $< 0.15R_{\text{vir}}$. The upper panels show the average circular velocity per subhalo mass bin, while the bottom panels refer to the stellar fraction. Each column reports the results for a given re-simulated region (see Table 1). Colour codes and line styles are as in Fig. 3.

affected by SIDM. The work of Peñarrubia et al. (2010) shows that cuspy subhaloes (as the ones we produce in our FP runs) will lose much more mass and decrease circular velocity only slightly (see the upper panel of their Fig. 6). As central substructures are supposed to experience earlier infall, it is expected that they would have undergone substantial dark matter loss while maintaining a circular velocity closely resembling that at the time of infall. The net consequence of this mechanism is an increased circular velocity at fixed subhalo mass.

To verify this, we present the circular velocity profiles in Fig. 10. We observe an increase in the circular velocity of SIDM V_{circ} (up to approximately $\approx 20\% - 30\%$) with respect to the collisionless DM V_{max} . In order to understand the mechanism that increased the compactness of these low-mass subhaloes, we dis-

sect them and analyse the circular velocity profiles of their stellar (central column) and DM component (right column). Here, it is clear that the circular velocity stellar component of the SIDM runs is higher than that of collisionless DM runs. To prove this point better, in Fig. 11, we show the subhalo maximum of circular velocity (upper panels) and stellar fraction (lower panels) versus M_{SH} mean relations for each of our six re-simulated regions

Therefore, we confirm that the increase in the circular velocity of subhaloes in SIDM simulations is associated to an increase in the stellar fraction. In particular, tidal stripping lowers the total mass while the circular velocity decreases only slightly, therefore bringing the data points closer to the observational scaling relation from Bergamini et al. (2019).




Fig. 12. Stellar mass of the BCG as a function of the halo mass M_{200c} . Colour data points are as in Fig. 3. Black stars represent points from Kravtsov et al. (2018).




Fig. 13. Peak of circular velocity for all galaxies with mass cut of $1 < M_{SH} < 2 [h^{-1} 10^{11} M_{\odot}]$ for subhaloes with cluster centric distance in the range $[0.15, 1]R_{vir}$. Colour lines and styles are as in Fig. 3. Shaded vertical bands correspond to the median and one standard deviation on the mean.

It is interesting to note in Fig. 10 that SIDM can produce subhaloes with higher compactness in the low-mass regime of subhaloes. This result reduces the tension between observed and simulated low-mass subhalo compactness Ragagnin et al. (2022) without changing baryon physics.

We now tackle the problem of the BCG stellar masses being overly massive compared to observations. To this end, in Fig. 12 we show the BCG stellar mass vs. the total mass of the halo, where we see that the mass of the BCG is only slightly affected by the inclusion of SIDM models. This is mainly related to the BCG being supposedly accreted by ex-situ material (see, e.g., Montenegro-Taborda et al. 2023). If subhaloes formed their stars in the field, then the subhalo stellar mass would be poorly influenced by SIDM.

4.4. Subhaloes with low circular velocity

In this section, we examine how SIDM models impact the low-concentration tail of subhaloes. To investigate this, we analyse




Fig. 14. Number of objects per mass stellar mass bins (upper panel) and the fraction above Σ_{UDG} defined in Sales et al. (2020) for a given stellar mass bin. Here we consider only objects outside the main halo R_{vir} and inside the main halo FoF group. Colours are organised as in Fig. 3, the shaded area corresponds to the error of the mean in each bin.

the distribution of V_{max} for subhaloes within the mass range of $1 - 2 \cdot 10^{10} h^{-1} M_{\odot}$ and located between $0.15R_{vir}$ and R_{vir} . We exclude subhaloes in the central regions of clusters, as we are aware from Sect. 4.3 that stripping in this region may result in higher compactness. In Fig. 13, we present the PDF of V_{max} for these subhaloes, revealing that SIDM simulations exhibit significantly lower median values of V_{max} . This result is compatible with the increase in circular velocity found in DMO simulations (see Fig. 14 in Fischer et al. 2023), and in this work, we confirm that this effect is still present in hydrodynamic simulations.

We now study if SIDM models are capable of producing an increased fraction of UDGs compared to collisionless DM. We define these objects as in Sales et al. (2020), namely as systems with $M_{*}/R_{eff} < 2.6 \times 10^7 h \text{ kpc}^{-2} M_{\odot}$, where M_{*} is the galaxy stellar mass and R_{eff} is the projected half light radius respectively (defined as in Sales et al. 2020, namely as $4/3$ times the 3D half mass radius). We also use the same resolution cut used in Sales et al. (2020), which imposes a lower limit of 10 DM particles per subhalo. In the upper panel of Fig. 14, we show the fraction of UDGs per mass bin. We notice that SIDM models produce systematically more diffuse objects, as the fraction of SIDM UDGs at $M_{*} \approx 0.3 \times 10^{10} h^{-1} M_{\odot}$ is more than twice as the amount in collisionless DM, in agreement with the fact that SIDM subhaloes in cluster outskirts tend to have lower circular velocities (see e.g., Fig. 6 in Peñarrubia et al. 2010).

To conclude the section, we verify if the increase of UDGs can be due to an overall increased number of substructures. Therefore, in the bottom panel of Fig. 14, we show the total number of galaxies per stellar mass bin, where we can see that SIDM models have systematically significantly fewer subhaloes compared to collisionless DM (stellar masses span 8 log-spaced bins in the range $[0.2, 1] h^{-1} 10^{10} M_{\odot}$).

5. Conclusions

In this study, we conducted a comparative analysis of the impact of rare and frequent velocity-dependent SIDM models within the cores of massive galaxy clusters (with masses around $\approx 10^{14} - 10^{15} M_{\odot}$). Additionally, we examined the influence of incorporating baryons in these simulations. We obtained the following main results:

- DMO simulations of rare and frequent SIDM models form core galaxy cluster profiles in agreement with other theoretical studies. In contrast, SIDM FP simulations show a significantly ($\approx 20\%$) higher central matter density compared to collisionless DM. Therefore we warn that it is not straightforward to observationally constrain SIDM cross-section by just estimating galaxy cluster cores.
- While SIDM DMO simulation strongly suppresses subhaloes in cluster cores compared to collisionless DM, this suppression is dampened in FP simulations; however, it is still significant, corresponding to a factor $\approx 20\%$ for rSIDM and $\approx 35\%$ for fSIDM.
- We found that SIDM produces cluster cores with substructures in the low-mass regime ($< 10^{11} h^{-1} M_{\odot}$) that are more compact and that this increase is due to an enhanced stripping of their DM component.
- SIDM simulations produce a significantly higher fraction of low surface brightness galaxies in cluster outskirts compared to collisionless DM.

In the future, it would be interesting to run and study cluster simulations with a resolution that is high enough to study subhalo core-collapse. Recent studies show that to properly resolve the core-collapse up to redshift $z = 0$ is challenging, as they require a large number of particles (compared to typical cosmological simulations) and small time steps to reproduce theoretical gravothermal evolution models (Yang & Yu 2021; Zhong et al. 2023; Palubski et al. 2024; Mace et al. 2024); to properly resolve the mean free path (see discussion in Fischer et al. 2024); and they also need a relatively high accuracy of gravitational interactions to deal with energy conservation issues of current numerical schemes Fischer et al. (2024).

Acknowledgements. We acknowledge support from the grant PRIN-MIUR 2017 WSCC32. AR acknowledges support by MIUR-DAAD contract number 34843 “The Universe in a Box”. We carried our simulations using the the INAF-Pleiadi project SIDM vs CDM allocated in the Trieste IT framework (Taffoni et al. 2020; Bertocco et al. 2020). We are especially grateful for the support by M. Petkova through the Computational centre for Particle and Astrophysics (C²PAP). LM acknowledges the support from the grants PRIN-MIUR 2022 20227RNLY3 and ASI n.2018-23-HH.0. KD acknowledges support by the COMPLEX project from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme grant agreement ERC-2019-AdG 882679. MSF is support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy – EXC-2094 “Origins” – 390783311. GD acknowledges the funding by the European Union - NextGenerationEU, in the framework of the HPC project – “National Centre for HPC, Big Data and Quantum Computing” (PNRR - M4C2 - I1.4 - CN00000013 – CUP J33C22001170001). We thank all participants of the Darkium SIDM <https://www.darkium.org/> Journal Club for helpful discussions. The authors thank the organisers of the Pollica 2023 SIDM workshop for the interesting discussions. AR and GC thank the support from INAF theory Grant 2022: Illuminating Dark Matter using Weak Lensing by Cluster Satellites, PI: Carlo Giocoli.

References

Adhikari, S., Banerjee, A., Boddy, K. K., et al. 2022, arXiv e-prints, arXiv:2207.10638
 Andrade, K. E., Fuson, J., Gad-Nasr, S., et al. 2022, MNRAS, 510, 54
 Bahé, Y. M. 2021, MNRAS, 505, 1458
 Beck, A. M., Murante, G., Arth, A., et al. 2016, MNRAS, 455, 2110
 Bergamini, P., Rosati, P., Mercurio, A., et al. 2019, A&A, 631, A130
 Bertocco, S., Goz, D., Tornatore, L., et al. 2020, in Astronomical Society of the Pacific Conference Series, Vol. 527, Astronomical Society of the Pacific Conference Series, ed. R. Pizzo, E. R. Deul, J. D. Mol, J. de Plaa, & H. Verkouter, 303
 Blumenthal, G. R., Faber, S. M., Flores, R., & Primack, J. R. 1986, ApJ, 301, 27
 Bonafede, A., Dolag, K., Staszyn, F., Murante, G., & Borgani, S. 2011, MNRAS, 418, 2234
 Bryan, G. L. & Norman, M. L. 1998, ApJ, 495, 80

Carleton, T., Errani, R., Cooper, M., et al. 2019, MNRAS, 485, 382
 Carton Zeng, Z., Peter, A. H. G., Du, X., et al. 2023, arXiv e-prints, arXiv:2310.09910
 Correa, C. A. 2021, MNRAS, 503, 920
 Davis, M., Efstathiou, G., Frenk, C. S., & White, S. D. M. 1985, ApJ, 292, 371
 Despali, G., Giocoli, C., Angulo, R. E., et al. 2016, MNRAS, 456, 2486
 Despali, G., Lovell, M., Vegetti, S., Crain, R. A., & Oppenheimer, B. D. 2020, MNRAS, 491, 1295
 Despali, G. & Vegetti, S. 2017, MNRAS, 469, 1997
 Di Cintio, A., Brook, C. B., Macciò, A. V., Dutton, A. A., & Cardona-Barrero, S. 2019, MNRAS, 486, 2535
 Dolag, K., Borgani, S., Murante, G., & Springel, V. 2009, MNRAS, 399, 497
 Duffy, A. R., Schaye, J., Kay, S. T., et al. 2010, MNRAS, 405, 2161
 Eckert, D., Ettori, S., Robertson, A., et al. 2022, A&A, 666, A41
 Einasto, J. 1965, Trudy Astrofizicheskogo Instituta Alma-Ata, 5, 87
 Eke, V. R., Cole, S., & Frenk, C. S. 1996, MNRAS, 282, 263
 Elbert, O. D., Bullock, J. S., Kaplinghat, M., et al. 2018, ApJ, 853, 109
 Fabjan, D., Borgani, S., Tornatore, L., et al. 2010, MNRAS, 401, 1670
 Ferland, G. J., Korista, K. T., Verner, D. A., et al. 1998, PASP, 110, 761
 Fischer, M. S., Brügggen, M., Schmidt-Hoberg, K., et al. 2021, MNRAS, 505, 851
 Fischer, M. S., Brügggen, M., Schmidt-Hoberg, K., et al. 2022, MNRAS, 516, 1923
 Fischer, M. S., Dolag, K., & Yu, H.-B. 2024, arXiv e-prints, arXiv:2403.00739
 Fischer, M. S., Kasselmann, L., Brügggen, M., et al. 2023, arXiv e-prints, arXiv:2310.07750
 Giocoli, C., Marulli, F., Baldi, M., Moscardini, L., & Metcalf, R. B. 2013, MNRAS, 434, 2982
 Giocoli, C., Tormen, G., & Sheth, R. K. 2012b, MNRAS, 422, 185
 Gnedin, O. Y., Kravtsov, A. V., Klypin, A. A., & Nagai, D. 2004, ApJ, 616, 16
 Granata, G., Bergamini, P., Grillo, C., et al. 2023, A&A, 679, A124
 Granata, G., Mercurio, A., Grillo, C., et al. 2022, A&A, 659, A24
 Greco, J. P., Goulding, A. D., Greene, J. E., et al. 2018, ApJ, 866, 112
 Harvey, D., Robertson, A., Massey, R., & McCarthy, I. G. 2019, MNRAS, 488, 1572
 Hernquist, L. 1990, ApJ, 356, 359
 Hirschmann, M., Dolag, K., Saro, A., et al. 2014, MNRAS, 442, 2304
 Kahlhoefer, F., Schmidt-Hoberg, K., Frandsen, M. T., & Sarkar, S. 2014, MNRAS, 437, 2865
 Kamada, A. & Kim, H. J. 2020, Phys. Rev. D, 102, 043009
 Kimmig, L. C., Remus, R.-S., Dolag, K., & Biffi, V. 2023, ApJ, 949, 92
 Kong, D., Kaplinghat, M., Yu, H.-B., Fraternali, F., & Mancera Piña, P. E. 2022, ApJ, 936, 166
 Kravtsov, A. V. & Borgani, S. 2012, ARA&A, 50, 353
 Kravtsov, A. V., Vikhlinin, A. A., & Meshcheryakov, A. V. 2018, Astronomy Letters, 44, 8
 Loeb, A. & Weiner, N. 2011, Phys. Rev. Lett., 106, 171302
 Ludlow, A. D., Navarro, J. F., Li, M., et al. 2012, MNRAS, 427, 1322
 Mace, C., Carton Zeng, Z., Peter, A. H. G., et al. 2024, arXiv e-prints, arXiv:2402.01604
 Mastromarino, C., Despali, G., Moscardini, L., et al. 2023, MNRAS, 524, 1515
 Meneghetti, M., Cui, W., Rasia, E., et al. 2023, A&A, 678, L2
 Meneghetti, M., Davoli, G., Bergamini, P., et al. 2020, Science, 369, 1347
 Meneghetti, M., Ragagnin, A., Borgani, S., et al. 2022, A&A, 668, A188
 Montenegro-Taborda, D., Rodriguez-Gomez, V., Pillepich, A., et al. 2023, MNRAS, 521, 800
 Moore, B., Governato, F., Quinn, T., Stadel, J., & Lake, G. 1998, ApJ, 499, L5
 Mowla, L., van Dokkum, P., Merritt, A., et al. 2017, ApJ, 851, 27
 Nadler, E. O., Banerjee, A., Adhikari, S., Mao, Y.-Y., & Wechsler, R. H. 2020, ApJ, 896, 112
 Nadler, E. O., Yang, D., & Yu, H.-B. 2023, arXiv e-prints, arXiv:2306.01830
 Natarajan, P., De Lucia, G., & Springel, V. 2007, MNRAS, 376, 180
 Natarajan, P. & Kneib, J.-P. 1997, MNRAS, 287, 833
 Natarajan, P. & Springel, V. 2004, ApJ, 617, L13
 Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490, 493
 Outmezzuine, N. J., Boddy, K. K., Gad-Nasr, S., Kaplinghat, M., & Sagunski, L. 2023, MNRAS, 523, 4786
 Palubski, I., Slone, O., Kaplinghat, M., Lisanti, M., & Jiang, F. 2024, arXiv e-prints, arXiv:2402.12452
 Peñarrubia, J., Benson, A. J., Walker, M. G., et al. 2010, MNRAS, 406, 1290
 Ragagnin, A., Meneghetti, M., Bassini, L., et al. 2022, A&A, 665, A16
 Ragagnin, A., Saro, A., Singh, P., & Dolag, K. 2021, MNRAS, 500, 5056
 Ragagnin, A., Tchipev, N., Bader, M., Dolag, K., & Hammer, N. J. 2016, in Advances in Parallel Computing, Volume 27: Parallel Computing: On the Road to Exascale, Edited by Gerhard R. Joubert, Hugh Leather, Mark Parsons, Frans Peters, Mark Sawyer. IOP Ebook, ISBN: 978-1-61499-621-7, pages 411-420
 Rasia, E., Borgani, S., Murante, G., et al. 2015, ApJ, 813, L17
 Robertson, A. 2021, MNRAS, 504, L7
 Robertson, A., Harvey, D., Massey, R., et al. 2019, MNRAS, 488, 3646

- Sabarish, V. M., Brügggen, M., Schmidt-Hoberg, K., Fischer, M. S., & Kahlhoefer, F. 2023, arXiv e-prints, arXiv:2310.07769
- Sagunski, L., Gad-Nasr, S., Colquhoun, B., Robertson, A., & Tulin, S. 2021, *J. Cosmology Astropart. Phys.*, 2021, 024
- Sales, L. V., Navarro, J. F., Peñafiel, L., et al. 2020, *MNRAS*, 494, 1848
- Spergel, D. N. & Steinhardt, P. J. 2000, *Phys. Rev. Lett.*, 84, 3760
- Springel, V. 2005, *MNRAS*, 364, 1105
- Springel, V., Di Matteo, T., & Hernquist, L. 2005, *MNRAS*, 361, 776
- Springel, V., White, S. D. M., Tormen, G., & Kauffmann, G. 2001, *MNRAS*, 328, 726
- Steinborn, L. K., Dolag, K., Hirschmann, M., Prieto, M. A., & Remus, R.-S. 2015, *MNRAS*, 448, 1504
- Taffoni, G., Becciani, U., Garilli, B., et al. 2020, in *Astronomical Society of the Pacific Conference Series*, Vol. 527, *Astronomical Society of the Pacific Conference Series*, ed. R. Pizzo, E. R. Deul, J. D. Mol, J. de Plaa, & H. Verkouter, 307
- Teklu, A. F., Remus, R.-S., Dolag, K., et al. 2015, *ApJ*, 812, 29
- Tinker, J., Kravtsov, A. V., Klypin, A., et al. 2008, *ApJ*, 688, 709
- Tormen, G., Diaferio, A., & Syer, D. 1998, *MNRAS*, 299, 728
- Tornatore, L., Borgani, S., Dolag, K., & Matteucci, F. 2007, *MNRAS*, 382, 1050
- Tulin, S. & Yu, H.-B. 2018, *Phys. Rep.*, 730, 1
- van Dokkum, P. G., Abraham, R., Merritt, A., et al. 2015, *ApJ*, 798, L45
- Yang, D. & Yu, H.-B. 2021, *Phys. Rev. D*, 104, 103031
- Yang, D. & Yu, H.-B. 2022, *J. Cosmology Astropart. Phys.*, 2022, 077
- Yang, S., Du, X., Zeng, Z. C., et al. 2023, *ApJ*, 946, 47
- Zhong, Y.-M., Yang, D., & Yu, H.-B. 2023, *MNRAS*, 526, 758




Fig. A.1. Density profile of D16 FP galaxy cluster at $z = 0.2$. We show its dark matter density profile (orange solid line); the corresponding NFW fit (dashed blue line) together with the values of r_s and ρ_s (vertical and horizontal dashed blue lines respectively); the corresponding cored-NFW functional form (dashed orange line, see our Eq. 4 or Yang & Yu 2021, Eq. A5), together with the values of ρ_c and r_c (vertical and horizontal dashed orange lines respectively); the profile of baryons (solid purple line); the corresponding Hernquist profile (dotted purple line); and the DM profile of the corresponding DMO simulation (solid green line).

Appendix A: Profile evolution

In this Appendix we provide additional details on the central increase of the DM density and velocity dispersion. First of all, we made sure that we have under control the computation of the central density $\hat{\rho}_c$ and of the central density of the baryons.

In Fig. A.1, we present the DM density profile of the FP run of the D16 galaxy cluster at the lowest redshift slice. There, we can see that the DM has a core and that the core is well fitted by the profile in Eq. 4 (notice how the vertical and horizontal orange dashed lines capture the core density and radius of the orange solid line). We also show that our NFW fits are properly capturing the DM profile (note how the blue dashed lines capture the orange solid line down to a few tens of kpc). Finally, we show that the core of baryons is well captured by a radius of $\approx h^{-1}10\text{ckpc}$ as stated in Sect 3.2.

We now provide more details on the time evolution of the central increase of DM density and circular velocity. The motivation behind this extended study is that in Section 3.2, we found that SIDM profiles on FP clusters have a higher central density compared to collisionless DM profiles of FP clusters. Therefore, we believe it is interesting to show the time evolution of both the velocity dispersion (Figure A.2 left panels) and density profiles (Figure A.2 right panels, for dark matter, stars and total matter, respectively) profile ratio of SIDM FP runs vs. the collisionless FP run values stacked for our six galaxy clusters. We see that at high redshift ($z = 2.3$), both SIDM and collisionless runs of FP clusters match both in terms of velocity dispersion and density profiles; therefore, SIDM and CDM produce similar high-redshift galaxy clusters. As time passes, both the central density and the central velocity dispersion of SIDM runs increase (compared to collisionless FP runs), as well as the relative central velocity dispersion.

Note that the velocity dispersion relative difference has a radial range that starts at ckpc scale (see left column), while the radial range of the relative density starts at $\approx 10\text{ckpc}$ (see right

columns). We use two different radial range in order to better show the deviations of SIDM with respect to CDM. In fact, SIDM velocity dispersion deviates at sub-softening scales (see, e.g., Fig. 7), while density profiles deviate at much larger radii.




Fig. A.2. Relative velocity dispersion and density profiles of FP SIDM runs vs. FP collisionless DM runs. Each row represents a redshift slice. The left panel show the relative dispersion profile, while the right panels report the relative density profiles (of dark matter, stars, and total matter, respectively). Green dashed lines report values for rSIDM, while orange dotted lines report values for fSIDM. Shaded area represent the error on the mean of the 6 regions. Note that the velocity dispersion relative difference has a different radial range compared to the density relative difference (see discussion).