

Construction of quantum codes from (γ, Δ) -cyclic codes

Om Prakash^{*1} · Shikha Patel¹ · Habibul Islam²

Received: date / Accepted: date

Abstract Let \mathbb{F}_q be the finite field of $q = p^m$ elements where p is a prime and m is a positive integer. This paper considers (γ, Δ) -cyclic codes over a class of finite commutative non-chain rings $\mathcal{R}_{q,s} = \mathbb{F}_q[v_1, v_2, \dots, v_s] / \langle v_i - v_i^2, v_i v_j = v_j v_i = 0 \rangle$ where γ is an automorphism of $\mathcal{R}_{q,s}$, Δ is a γ -derivation of $\mathcal{R}_{q,s}$ and $1 \leq i \neq j \leq s$ for a positive integer s . Here, we show that a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$ is the direct sum of (θ, \mathfrak{S}) -cyclic codes of length n over \mathbb{F}_q , where θ is an automorphism of \mathbb{F}_q and \mathfrak{S} is a θ -derivation of \mathbb{F}_q . Further, necessary and sufficient conditions for both (γ, Δ) -cyclic and (θ, \mathfrak{S}) -cyclic codes to contain their Euclidean duals are established. Finally, we obtain many quantum codes by applying the dual containing criterion on the Gray images of these codes. The obtained codes have better parameters than those available in the literature.

Keywords Skew polynomial rings · Skew cyclic codes · (σ, δ) -cyclic codes · Gray map · CSS construction · Quantum codes .

Mathematics Subject Classification (2000) 12Y05 · 16Z05 · 94B05 · 94B35 · 94B15.

1 Introduction

After the pioneering work of Hammons et al. [18] in 1994, codes over finite rings attracted many researchers for better error-correcting codes. Later, several important research has been carried out over finite rings and explored plenty of suitable parameters; we refer [2, 3, 12, 14, 27]. Nevertheless, all of these works have been considered over finite commutative rings. Hence, it is natural to look at these works over the noncommutative ring to obtain codes with better parameters. Towards this, in 2007, Boucher et al. [5] introduced skew cyclic codes as a generalized class of cyclic codes by using a non-trivial automorphism θ on a finite field \mathbb{F}_q . They proved that the noncommutative rings (skew polynomial rings) are worthy alphabets for producing new parameters. In addition, they have provided a few codes with better parameters that are not known earlier over finite commutative rings. Since the factorization of the polynomial $x^n - 1$ plays an important role in the characterization of cyclic codes of length n , more factorization leads to the case of getting many new code parameters. Therefore, skew cyclic codes that generalize cyclic codes in a non-commutative setup attract many researchers. Later, Abualrub et al. [1] and Bhaintwal [4] introduced skew-quasi cyclic codes. From an application point of view, recently, many authors have shown that skew cyclic codes are one of the important resources for producing

¹Department of Mathematics, Indian Institute of Technology Patna, Patna 801 106, India

²School of Computer Science, University of St. Gallen, Switzerland

ORCID: 0000-0002-6512-4229 (Om Prakash)

E-mail: om@iitp.ac.in(*corresponding author), shikha_1821ma05@iitp.ac.in, habibul.pma17@iitp.ac.in

new quantum codes along with classical codes (see [5, 6, 8, 17, 21, 34]).

However, all of the above works have been carried out on skew polynomial rings of automorphism type. Only a few works are available in the literature with both automorphisms and derivations. In [7, 9, 30, 35], the authors generalized the notion of codes over skew polynomial rings with non-trivial automorphism θ and θ -derivation \mathfrak{S} under the usual addition of polynomials and a specific polynomials multiplication involving θ and \mathfrak{S} . For the noncommutative ring $\mathbb{F}_q[x, \theta; \mathfrak{S}]$ where θ is the Frobenius automorphism $a \mapsto a^p$, p is the characteristic of \mathbb{F}_q , the authors [7, 35] defined the inner θ -derivation \mathfrak{S} induced by $\beta \in \mathbb{F}_q^*$ of the form $a \mapsto \beta(\theta(a) - a)$. Further, Boulagouaz and Leroy [9] studied (σ, δ) -codes with σ -derivation induced by the ring element. Recently, Sharma and Bhaintwal [32] have studied skew cyclic codes over $\mathbb{Z}_4 + u\mathbb{Z}_4$, $u^2 = 1$ with both automorphism and inner derivation. In 2021, Ma et al. [26] studied (σ, δ) -skew quasi-cyclic codes over the ring $\mathbb{Z}_4 + u\mathbb{Z}_4$, $u^2 = 1$. Further, in 2021, Patel and Prakash [29] studied (θ, δ_θ) -cyclic codes over the ring $\mathbb{F}_q[u, v]/\langle u^2 - u, v^2 - v, uv - vu \rangle$ via the decomposition method over \mathbb{F}_q . Here, we extend our previous work [29] to a more general structure and propose a fruitful application of (γ, Δ) -cyclic codes in the context of quantum code construction. Per our survey, it is worth mentioning that this is the first article proposing an application of (γ, Δ) -cyclic codes into quantum codes.

Quantum error-correcting codes play a significant role in protecting information against the disturbances such as decoherence occurring in the channel. Shor [33] discovered the first quantum code. After that, Calderbank et al. [11] provided a method to obtain quantum codes from classical codes. This technique became very popular among researchers and is known as the CSS (Calderbank-Shor-Steane) construction. Presently, quantum codes and their implementation from classical codes have gained significant attention. As a consequence, many quantum codes with better parameters have been constructed from different families of linear codes such as cyclic, skew cyclic, skew constacyclic codes, etc., see [2, 3, 12, 14, 19, 25, 27, 28, 31]. But the searching of new methods on different structures are still going on by which one can construct quantum codes efficiently with suitable parameters. Since getting new quantum codes proportionally depend on the abundance of factors of $x^n - 1$, many authors have been exploring quantum codes in the setting of the skew polynomial ring with automorphism where $x^n - 1$ indeed possesses more factorization than the commutative case. However, in this work, we extend all these works in a new direction by considering skew polynomial rings with non-trivial automorphisms and derivations. In this case, we have different derivations for the same Frobenius automorphism having the form $a \mapsto \beta(\theta(a) - a)$ for all $\beta \in \mathbb{F}_q^*$.

The rest of the paper is structured as follows: In Section 2, we present some basic results and notations that will be useful for later sections. In Section 3, we discuss (θ, \mathfrak{S}) -cyclic codes over \mathbb{F}_q and derive a necessary and sufficient condition to contain their duals over \mathbb{F}_q . Further, Section 4 includes the results on (γ, Δ) -cyclic codes over $\mathcal{R}_{q,s}$ and dual-containing property for these codes as well. Section 5 describes the applications of our obtained results by providing many new quantum codes with superior parameters. Finally, Section 6 concludes our work.

2 Preliminaries

In this Section, we provide some preliminary results, definitions and notations which are used throughout this paper. We consider a finite non-chain ring $\mathcal{R}_{q,s} := \mathbb{F}_q[v_1, v_2, \dots, v_s]/\langle v_i - v_i^2, v_i v_j = v_j v_i = 0 \rangle$ where $1 \leq i \neq j \leq s$ and s is a positive integer. This $\mathcal{R}_{q,s}$ is a class of finite commutative ring with unity for different values of q and s . Further, $\mathcal{R}_{q,s}$ can also be represented in the form of $\mathcal{R}_{q,s} = \mathbb{F}_q + v_1\mathbb{F}_q + \dots + v_s\mathbb{F}_q$ with $v_i - v_i^2, v_i v_j = v_j v_i = 0$. Moreover, $\mathcal{R}_{q,s}$ is a non-chain semi-local Frobenius ring having $s + 1$ maximal ideals. For

$s = 2$, there are three maximal ideals $\langle v_1 + v_2 \rangle$, $\langle 1 - v_1 \rangle$ and $\langle 1 - v_2 \rangle$ in $\mathcal{R}_{q,2}$, refer [?]. Consider

$$\zeta_0 = \prod_{i=1}^s (1 - v_i), \quad \text{and} \quad \zeta_j = v_j, \quad 1 \leq j \leq s.$$

It is easy to verify that $\sum_{i=0}^s \zeta_i = 1$ and

$$\zeta_i \zeta_j = \begin{cases} \zeta_i, & \text{if } i = j \\ 0, & \text{if } i \neq j \end{cases}.$$

Therefore, by Chinese Remainder Theorem, $\mathcal{R}_{q,s} = \zeta_0 \mathcal{R}_{q,s} \oplus \zeta_1 \mathcal{R}_{q,s} \oplus \cdots \oplus \zeta_s \mathcal{R}_{q,s} = \zeta_0 \mathbb{F}_q \oplus \zeta_1 \mathbb{F}_q \oplus \cdots \oplus \zeta_s \mathbb{F}_q$. From here we can conclude that any element $t \in \mathcal{R}_{q,s}$ can be uniquely written as $t = \zeta_0 t_0 + \zeta_1 t_1 + \cdots + \zeta_s t_s$, where $t_i \in \mathbb{F}_q$. Also, t is a unit in $\mathcal{R}_{q,s}$ if and only if $t_i \in \mathbb{F}_q^*$ for all i .

Recall that a non-empty subset \mathcal{C} of $\mathcal{R}_{q,s}^n$ is said to be a linear code of length n over $\mathcal{R}_{q,s}$ if it is an $\mathcal{R}_{q,s}$ -submodule of $\mathcal{R}_{q,s}^n$ and the elements of \mathcal{C} are called codewords. The Hamming weight $w_H(c)$ of a codeword $c = (c_0, c_1, \dots, c_{n-1}) \in \mathcal{C}$ is the number of nonzero coordinates in c . The Hamming distance between any two codewords c and c' of \mathcal{C} is defined as $d_H(c, c') = w_H(c - c')$ and the Hamming distance of a linear code \mathcal{C} is defined as $d_H(\mathcal{C}) = \min\{d_H(x, y) \mid x, y \in \mathcal{C}, x \neq y\}$. The Euclidean inner product of c and c' in \mathcal{R}^n is defined by $c \cdot c' = \sum_{i=0}^{n-1} c_i c'_i$ where $c = (c_0, c_1, \dots, c_{n-1})$ and $c' = (c'_0, c'_1, \dots, c'_{n-1})$ are codewords in \mathcal{C} . The dual code of \mathcal{C} is defined by $\mathcal{C}^\perp = \{c \in \mathcal{R}_{q,s}^n \mid c \cdot c' = 0, \text{ for all } c' \in \mathcal{C}\}$. Also, a linear code \mathcal{C} is self-orthogonal if $\mathcal{C} \subseteq \mathcal{C}^\perp$ and self-dual if $\mathcal{C} = \mathcal{C}^\perp$. Further, let $c = (c_0, c_1, \dots, c_{n-1}) \in \mathcal{C} \subseteq \mathbb{F}_q^n$. If \mathcal{C} is an $[n, k, d]$ linear code, then from the Singleton bound, its minimum distance is bounded above by $d \leq n - k + 1$, where d is the minimum distance, k is the dimension, and n is the length of the code. A code achieving the mentioned bound is called maximum-distance-separable (MDS). If the minimum distance of the code is one unit less than the MDS, then the code is called almost MDS. A linear code is said to be optimal if it has the highest possible minimum distance for a given length and dimension.

Definition 1 Let $\mathcal{R}_{q,s}$ be a finite ring and γ be an automorphism of $\mathcal{R}_{q,s}$. Then a map $\Delta : \mathcal{R}_{q,s} \rightarrow \mathcal{R}_{q,s}$ is said to be a γ -derivation of $\mathcal{R}_{q,s}$ if

1. $\Delta(x + y) = \Delta(x) + \Delta(y)$;
2. $\Delta(xy) = \Delta(x)y + \gamma(x)\Delta(y)$

for all $x, y \in \mathcal{R}_{q,s}$.

Let us consider an automorphism $\theta : \mathbb{F}_q \rightarrow \mathbb{F}_q$ defined by $\theta(a) = a^q$, for all $a \in \mathbb{F}_q$ and a θ -derivation $\mathfrak{S} : \mathbb{F}_q \rightarrow \mathbb{F}_q$ defined by $\mathfrak{S}(a) = \beta(\theta(a) - a)$, for all $a \in \mathbb{F}_q$ and $\beta \in \mathbb{F}_q^*$. Now, we extend the above maps over $\mathcal{R}_{q,s}$ and define the skew polynomial ring with both automorphism and derivation over $\mathcal{R}_{q,s}$. Let $\text{Aut}(\mathcal{R}_{q,s})$ be the set of all automorphism of $\mathcal{R}_{q,s}$ and $\gamma \in \text{Aut}(\mathcal{R}_{q,s})$. We consider the set

$$\mathcal{R}_{q,s}[x; \gamma, \Delta] = \{b_l x^l + \cdots + b_1 x + b_0 \mid b_i \in \mathcal{R} \text{ and } l \in \mathbb{N}\},$$

where Δ is a γ -derivation of $\mathcal{R}_{q,s}$. Then $\mathcal{R}_{q,s}[x; \gamma, \Delta]$ is a noncommutative ring unless γ is the identity under the usual addition of polynomials and multiplication is defined with respect to $xb = \gamma(b)x + \Delta(b)$ for $b \in \mathcal{R}_{q,s}$, known as a skew polynomial ring.

Definition 2 An element $f(x) \in \mathcal{R}_{q,s}[x; \gamma, \Delta]$ is said to be a central element of $\mathcal{R}_{q,s}[x; \gamma, \Delta]$ if $f(x)b(x) = b(x)f(x)$, for all $b(x) \in \mathcal{R}_{q,s}[x; \gamma, \Delta]$.

Definition 3 [20,23] A pseudo-linear transformation $T_{\gamma,\Delta} : \mathcal{R}_{q,s}^n \rightarrow \mathcal{R}_{q,s}^n$ is an additive map defined by

$$T_{\gamma,\Delta}(v) = \gamma(v)M + \Delta(v), \quad (1)$$

where $v = (v_1, v_2, \dots, v_n) \in \mathcal{R}_{q,s}^n$, $\gamma(v) = (\gamma(v_1), \gamma(v_2), \dots, \gamma(v_n)) \in \mathcal{R}_{q,s}^n$, M is a matrix of order $n \times n$ over $\mathcal{R}_{q,s}$ and $\Delta(v) = (\Delta(v_1), \Delta(v_2), \dots, \Delta(v_n)) \in \mathcal{R}_{q,s}^n$. If $\Delta = 0$, then T_γ is known as *semi-linear transformation*.

Definition 4 1. A code \mathcal{C} of length n over $\mathcal{R}_{q,s}$ is said to be a (γ, Δ) -linear code if it is a left $\mathcal{R}_{q,s}[x; \gamma, \Delta]$ -submodule of $\frac{\mathcal{R}_{q,s}[x; \gamma, \Delta]}{\langle x^n - 1 \rangle}$. Moreover, if $x^n - 1$ is a central element of $\mathcal{R}_{q,s}[x; \gamma, \Delta]$, then \mathcal{C} is a central (γ, Δ) -linear code.

2. A code \mathcal{C} of length n over $\mathcal{R}_{q,s}$ is said to be a (γ, Δ) -cyclic code if

- \mathcal{C} is a (γ, Δ) -linear code;
- $T_{\gamma,\Delta}(\mathcal{C}) \subseteq \mathcal{C}$, where $T_{\gamma,\Delta}$ is as defined in Equation (1) and M is defined as

$$M = \begin{pmatrix} 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \\ 1 & 0 & \dots & 0 \end{pmatrix}.$$

Remark 1 [15, Exercise 20] Let $\mathcal{R}_{q,s}[x; \gamma, \Delta]$ be a skew polynomial ring, $r \in \mathcal{R}_{q,s}$ and $n \in \mathbb{N}$. Then

$$x^n r = \gamma^n(r)x^n + a_{n-1}x^{n-1} + \dots + a_1x + \Delta^n(r),$$

for some $a_{n-1}, \dots, a_1 \in \mathcal{R}_{q,s}$.

To find generator polynomials of (γ, Δ) -cyclic codes over $\mathcal{R}_{q,s}$, first we derive the right division algorithm in $\mathcal{R}_{q,s}[x; \gamma, \Delta]$.

Theorem 1 (*The Right Division Algorithm*) Let $f(x), g(x) \in \mathcal{R}_{q,s}[x; \gamma, \Delta]$ such that the leading coefficient of $g(x)$ be a unit. Then there exist $q(x), r(x) \in \mathcal{R}_{q,s}[x; \gamma, \Delta]$ such that

$$f(x) = q(x)g(x) + r(x),$$

where $r(x) = 0$ or $\deg r(x) < \deg g(x)$.

Proof If $f(x) = 0$, then result follows by taking $q(x), r(x) = 0$. If $\deg f(x) < \deg g(x)$, then we take $q(x) = 0$ and $r(x) = f(x)$. Further, for $\deg f(x) \geq \deg g(x)$, we prove it by induction on $\deg f(x)$.

It can be easily seen that the result is true for $\deg f(x) = 0$. Now, suppose the result is true for all polynomials of degree less than $\deg f(x)$. Let $f(x) = f_0 + f_1x + \dots + f_sx^s$ and $g(x) = g_0 + g_1x + \dots + g_tx^t$ be two polynomials in $\mathcal{R}_{q,s}[x; \gamma, \Delta]$ such that $f_s \neq 0$ and g_t is a unit. Consider a polynomial

$$h(x) = f(x) - f_s\gamma^{s-t}(g_t^{-1})x^{s-t}g(x).$$

From Remark 1, $h(x)$ can be written as

$$\begin{aligned}
h(x) &= f(x) - f_s \gamma^{s-t} (g_t^{-1}) x^{s-t} g(x) = f(x) - f_s \gamma^{s-t} (g_t^{-1}) x^{s-t} (g_0 + g_1 x + \cdots + g_t x^t) \\
&= f(x) - f_s \gamma^{s-t} (g_t^{-1}) x^{s-t} g_0 - f_s \gamma^{s-t} (g_t^{-1}) x^{s-t} g_1 x - \cdots - f_s \gamma^{s-t} (g_t^{-1}) x^{s-t} g_t x^t \\
&= f(x) - f_s \gamma^{s-t} (g_t^{-1}) (\gamma^{s-t} (g_0) x^{s-t} + a_{s-t-1} x^{s-t-1} + \cdots + a_1 x + \Delta^{s-t} (g_0)) \\
&\quad - \cdots - f_s \gamma^{s-t} (g_t^{-1}) (\gamma^{s-t} (g_t) x^{s-t} + b_{s-t-1} x^{s-t-1} + \cdots + b_1 x + \Delta^{s-t} (g_t)) x^t \\
&= f(x) - f_s \gamma^{s-t} (g_t^{-1}) (\gamma^{s-t} (g_0) x^{s-t} + a_{s-t-1} x^{s-t-1} + \cdots + a_1 x + \Delta^{s-t} (g_0)) \\
&\quad - \cdots - f_s \gamma^{s-t} (g_t^{-1}) \gamma^{s-t} (g_t) x^{s-t} x^t - f_s \gamma^{s-t} (g_t^{-1}) b_{s-t-1} x^{s-t-1} x^t - \cdots - \\
&\quad f_s \gamma^{s-t} (g_t^{-1}) \Delta^{s-t} (g_t) x^t \\
&= f(x) - f_s \gamma^{s-t} (g_t^{-1}) (\gamma^{s-t} (g_0) x^{s-t} + a_{s-t-1} x^{s-t-1} + \cdots + a_1 x + \Delta^{s-t} (g_0)) \\
&\quad - \cdots - f_s x^s - f_s \gamma^{s-t} (g_t^{-1}) b_{s-t-1} x^{s-1} - \cdots - f_s \gamma^{s-t} (g_t^{-1}) \Delta^{s-t} (g_t) x^t \\
&= f_0 + f_1 x + \cdots + f_s x^s - f_s \gamma^{s-t} (g_t^{-1}) (\gamma^{s-t} (g_0) x^{s-t} + a_{s-t-1} x^{s-t-1} + \cdots \\
&\quad + a_1 x + \Delta^{s-t} (g_0)) - \cdots - f_s x^s - f_s \gamma^{s-t} (g_t^{-1}) b_{s-t-1} x^{s-1} - \cdots - f_s \gamma^{s-t} (g_t^{-1}) \\
&\quad \Delta^{s-t} (g_t) x^t \\
&= f_0 + f_1 x + \cdots + f_{s-1} x^{s-1} - f_s \gamma^{s-t} (g_t^{-1}) (\gamma^{s-t} (g_0) x^{s-t} + a_{s-t-1} x^{s-t-1} \\
&\quad + \cdots + a_1 x + \Delta^{s-t} (g_0)) - \cdots - f_s \gamma^{s-t} (g_t^{-1}) b_{s-t-1} x^{s-1} - \cdots - f_s \gamma^{s-t} (g_t^{-1}) \\
&\quad \Delta^{s-t} (g_t) x^t
\end{aligned}$$

where $a_1, a_2, \dots, a_{s-t-1}, b_1, b_2, \dots, b_{s-t-1} \in \mathcal{R}_{q,s}$. Now, we can conclude that $\deg h(x) < \deg f(x)$. Hence, by induction on $\deg h(x)$, there exist $b(x), r(x) \in \mathcal{R}_{q,s}[x; \gamma, \Delta]$ such that

$$h(x) = b(x)g(x) + r(x),$$

where $r(x) = 0$ or $\deg r(x) < \deg g(x)$. Thus,

$$\begin{aligned}
f(x) &= h(x) + f_s \gamma^{s-t} (g_t^{-1}) x^{s-t} g(x) \\
&= b(x)g(x) + r(x) + f_s \gamma^{s-t} (g_t^{-1}) x^{s-t} g(x) \\
&= (b(x) + f_s \gamma^{s-t} (g_t^{-1}) x^{s-t}) g(x) + r(x) \\
&= q(x)g(x) + r(x),
\end{aligned}$$

where $q(x) = b(x) + f_s \gamma^{s-t} (g_t^{-1}) x^{s-t} \in \mathcal{R}_{q,s}[x; \gamma, \Delta]$ and $r(x) = 0$ or $\deg r(x) < \deg g(x)$. This gives the required result.

Similarly, one can define the left division algorithm. In above Theorem 1, if $r(x) = 0$, then $g(x)$ is called a right divisor of $f(x)$ or $f(x)$ is a left multiple of $g(x)$ in $\mathcal{R}_{q,s}[x; \gamma, \Delta]$. Throughout this paper, we consider the right division.

3 (θ, \mathfrak{S}) -cyclic codes over \mathbb{F}_q

In the following, we discuss the main algebraic properties of (θ, \mathfrak{S}) -cyclic codes in $R = \mathbb{F}_q[x; \theta, \mathfrak{S}]$ and provide a necessary and sufficient condition for these codes to contain their Euclidean duals. In [9], Boulagouaz and Leroy introduced the notion of (f, γ, Δ) -cyclic codes. Moreover, a (θ, \mathfrak{S}) -cyclic code \mathcal{C} is the subset of \mathbb{F}_q^n consisting of the coordinates of the elements of $Rg(x)/\langle x^n - 1 \rangle$ in the basis $\{1, x, \dots, x^{n-1}\}$ for some right monic factors $g(x)$ of $x^n - 1$.

Theorem 2 *Let $g(x) = g_0 + g_1 x + \cdots + g_r x^r \in R$ be a monic polynomial.*

1. *A (θ, \mathfrak{S}) -cyclic code of length n corresponding to $Rg(x)/\langle x^n - 1 \rangle$ is a free left \mathbb{F}_q -module of dimension $n - r$ where $r = \deg g(x)$.*

2. If $v = (v_0, v_1, \dots, v_{n-1}) \in \mathcal{C}$, then $T_{\theta, \mathfrak{S}}(v) \in \mathcal{C}$.

3. The rows of the matrix which generates the code \mathcal{C} are given by

$$T_{\theta, \mathfrak{S}}^k(g_0, g_1, \dots, g_r, 0, 0, \dots, 0), \quad \text{for } 0 \leq k \leq n - r - 1.$$

Proof 1. We have $x^n - 1 = h(x)g(x)$ for some monic polynomials $h(x) \in R$. Hence, as left R -module, we have $Rg(x)/\langle x^n - 1 \rangle \cong R/\langle h(x) \rangle$. Since h is monic, $R/\langle h(x) \rangle$ is a free left \mathbb{F}_q -module of rank $\deg h(x) = n - r$.

2. $v = (v_0, v_1, \dots, v_{n-1}) \in \mathcal{C}$ if and only if $v(x) := \sum_{i=0}^{n-1} v_i x^i + \langle x^n - 1 \rangle \in Rg(x)/\langle x^n - 1 \rangle$. Since $xv(x) \in Rg(x)/\langle x^n - 1 \rangle$ and left multiplication by x on $R/\langle x^n - 1 \rangle$ corresponds to the action of $T_{\theta, \mathfrak{S}}$ on \mathbb{F}_q^n , we have $T_{\theta, \mathfrak{S}}(v) \in \mathcal{C}$.

3. We have $T_{\theta, \mathfrak{S}}^k(v_0, v_1, \dots, v_{n-1}) \in \mathcal{C}$ for any $k \geq 0$. On the other hand, it is clear that $g, xg, x^2g, \dots, x^{n-r-1}g$ are left linearly independent over \mathbb{F}_q , all are taken modulo $x^n - 1$ and hence form a basis of $Rg(x)/\langle x^n - 1 \rangle$. In codewords representation, this implies that the vectors $T_{\theta, \mathfrak{S}}^k(g_0, g_1, \dots, g_r, 0, \dots, 0)$ form a left \mathbb{F}_q -basis for \mathcal{C} , $0 \leq k \leq n - r - 1$.

Theorem 3 Let \mathcal{C} be a left R -submodule of $R/\langle x^n - 1 \rangle$. Then \mathcal{C} is a (θ, \mathfrak{S}) -cyclic submodule generated by a monic polynomial of the smallest degree in \mathcal{C} .

Proof Let $g(x) \in \mathcal{C}$ be a monic smallest degree polynomial among nonzero polynomials in \mathcal{C} and $c(x) \in \mathcal{C}$. Then by Theorem 1, there exist unique polynomials $q(x)$ and $r(x)$ in R such that $c(x) = q(x)g(x) + r(x)$ where $r(x) = 0$ or $\deg r(x) < \deg g(x)$. As \mathcal{C} is a left R -submodule, we have $r(x) = c(x) - q(x)g(x) \in \mathcal{C}$. This is a contradiction to the assumption that $g(x)$ is of the smallest degree in \mathcal{C} unless $r(x) = 0$. This implies $c(x) = q(x)g(x)$ and hence \mathcal{C} is a (θ, \mathfrak{S}) -cyclic submodule generated by $g(x)$.

Theorem 4 Let $\mathcal{C} = \langle g(x) \rangle$ be a left R -submodule of $R/\langle x^n - 1 \rangle$, where $g(x)$ is a monic polynomial of smallest degree in \mathcal{C} . Then $g(x)$ is a right divisor of $x^n - 1$.

Proof Consider a monic smallest degree polynomial $g(x)$ in \mathcal{C} . From Theorem 1, there exist polynomials $q(x)$ and $r(x)$ in R such that $x^n - 1 = q(x)g(x) + r(x)$, where $\deg r(x) < \deg g(x)$. Since $g(x)$ and $x^n - 1 = 0$ are in \mathcal{C} , this implies $r(x) = (x^n - 1) - q(x)g(x) \in \mathcal{C}$. But, $g(x)$ is smallest in \mathcal{C} . Therefore, $r(x) = 0$ and hence $g(x)$ is a right divisor of $x^n - 1$.

Let \mathcal{C} be a (θ, \mathfrak{S}) -cyclic code of length n over \mathbb{F}_q generated by the right divisor $g(x)$ of $x^n - 1$, where $g(x) = g_0 + g_1x + \dots + g_r x^r \in R$ and $g_r = 1$. Then from the above discussion, we can conclude that \mathcal{C} is a free left \mathbb{F}_q -module of dimension $k = n - \deg g(x)$. Now, by using [23, Theorem 3.2], the generator matrix of \mathcal{C} is given by

$$G = \begin{pmatrix} g \\ T_{\theta, \mathfrak{S}}(g) \\ \vdots \\ T_{\theta, \mathfrak{S}}^{k-1}(g) \end{pmatrix} \quad (2)$$

where $g = (g_0, g_1, g_2, \dots, g_r)$ is the codeword corresponding to $g(x)$. Moreover, it is well known that $\dim(\mathcal{C}) + \dim(\mathcal{C}^\perp) = n$. Therefore, $\dim(\mathcal{C}^\perp) = n - k = r$. Further, for our convenience, we define a one-to-one correspondence between the algebraic structures and combinatorial structures of (θ, \mathfrak{S}) -cyclic codes as follows:

$$\begin{aligned} \tau : \quad \mathbb{F}_q^n &\longrightarrow \mathbb{F}_q[x; \theta, \mathfrak{S}]/\langle x^n - 1 \rangle \\ (c_0, c_1, c_2, \dots, c_{n-1}) &\longmapsto c_0 + c_1x + c_2x^2 + \dots + c_{n-1}x^{n-1}. \end{aligned}$$

Theorem 5 Let $\mathcal{C} = \langle g(x) \rangle$ be a (θ, \mathfrak{S}) -cyclic code of length n over \mathbb{F}_q , for some right divisor $g(x)$ of $x^n - 1$. Let $x^n - 1 = h(x)g(x) = g(x)h'(x)$ for some monic skew polynomials $g(x), h(x), h'(x) \in R$. Then $c(x) \in \mathbb{F}_q[x; \theta, \mathfrak{S}]/\langle x^n - 1 \rangle$ is contained in \mathcal{C} if and only if $c(x)h'(x) = 0$ in $\mathbb{F}_q[x; \theta, \mathfrak{S}]/\langle x^n - 1 \rangle$.

Proof Let $c(x) \in \mathbb{F}_q[x; \theta, \mathfrak{S}]/\langle x^n - 1 \rangle$ be contained in \mathcal{C} . Then $c(x) = a(x)g(x)$ for some $a(x) \in R$. Now,

$$\begin{aligned} c(x) &= a(x)g(x) \text{ for some } a(x) \in R \\ c(x)h'(x) &= a(x)g(x)h'(x) = a(x)h(x)g(x) \\ &= a(x)(x^n - 1) = 0 \text{ in } \mathbb{F}_q[x; \theta, \mathfrak{S}]/\langle x^n - 1 \rangle. \end{aligned}$$

Conversely, let $c(x)h'(x) = 0$ for some $c(x)$ in $\mathbb{F}_q[x; \theta, \mathfrak{S}]/\langle x^n - 1 \rangle$. Then $c(x)h'(x) = q(x)(x^n - 1)$ for some $q(x) \in \mathbb{F}_q[x; \theta, \mathfrak{S}]/\langle x^n - 1 \rangle$. Also,

$$c(x)h'(x) = q(x)(x^n - 1) = q(x)h(x)g(x) = q(x)g(x)h'(x).$$

This implies that $c(x) = q(x)g(x) \in \langle g(x) \rangle = \mathcal{C}$ as $h'(x)$ is a nonzero polynomial.

Now, with the help of the above-defined correspondence, the following theorem provides the generator matrix of the dual code \mathcal{C}^\perp of (θ, \mathfrak{S}) -cyclic code \mathcal{C} of length n over \mathbb{F}_q .

Theorem 6 Let $\mathcal{C} = \langle g(x) \rangle$ be a (θ, \mathfrak{S}) -cyclic code of length n over \mathbb{F}_q for some right divisor $g(x)$ of $x^n - 1$ and $x^n - 1 = h(x)g(x) = g(x)h'(x)$ for some monic skew polynomials $g(x), h(x), h'(x) \in R$. Then $\deg g(x)$ linearly independent columns of the matrix

$$H = \begin{pmatrix} h' \\ T_{\theta, \mathfrak{S}}(h') \\ \vdots \\ T_{\theta, \mathfrak{S}}^{n-1}(h') \end{pmatrix}$$

form a basis of \mathcal{C}^\perp .

Proof Consider a (θ, \mathfrak{S}) -cyclic code \mathcal{C} of length n over \mathbb{F}_q . Let $\mathcal{C} = \langle g(x) \rangle$ where $g(x)$ is a right divisor of $x^n - 1$, and its leading coefficient is a unit. Then there exists $h(x) = h_0 + h_1x + \cdots + h_kx^k \in \mathbb{F}_q[x; \theta, \mathfrak{S}]/\langle x^n - 1 \rangle$ such that $x^n - 1 = h(x)g(x) = g(x)h'(x)$. Now, for $c(x) = c_0 + c_1x + \cdots + c_{n-1}x^{n-1} \in \mathcal{C}$, we have

$$\tau(c(T_{\theta, \mathfrak{S}})(h')) = c(x)h'(x) = a(x)g(x)h'(x) = a(x)h(x)g(x) = a(x)(x^n - 1) = 0.$$

for some $a(x)$ in $\mathbb{F}_q[x; \theta, \mathfrak{S}]/\langle x^n - 1 \rangle$ and $c(x)h'(x)$ is taken modulo $x^n - 1$. This implies $c(T_{\theta, \mathfrak{S}})(h') = 0$. Thus, $0 = c(T_{\theta, \mathfrak{S}})(h') = c_0 + c_1T_{\theta, \mathfrak{S}}(h') + c_2T_{\theta, \mathfrak{S}}^2(h') + \cdots + c_{n-1}T_{\theta, \mathfrak{S}}^{n-1}(h')$. This shows that $(c_0, c_1, c_2, \dots, c_{n-1}) \cdot H = 0$ for any $c = (c_0, c_1, c_2, \dots, c_{n-1}) \in \mathcal{C}$. Also, $\tau(T_{\theta, \mathfrak{S}}^k(h')) = x^k h'(x)$ for $k = 0, 1, \dots, n - \deg h'(x) - 1 = \deg g(x) - 1$ and hence $\{h', T_{\theta, \mathfrak{S}}(h'), T_{\theta, \mathfrak{S}}^2(h'), \dots, T_{\theta, \mathfrak{S}}^{r-1}(h')\}$ are linearly independent.

We now derive a necessary and sufficient condition for (θ, \mathfrak{S}) -cyclic codes to contain their duals codes over \mathbb{F}_q .

Theorem 7 Let $\mathcal{C} = \langle g(x) \rangle$ be a (θ, \mathfrak{S}) -cyclic code of length n over \mathbb{F}_q , for some right divisor $g(x)$ of $x^n - 1$ and $x^n - 1 = h(x)g(x) = g(x)h'(x)$ for some monic skew polynomials $g(x), h(x), h'(x) \in R$. Then $\mathcal{C}^\perp \subseteq \mathcal{C}$ if and only if $h'(x)h'(x)$ is divisible by $x^n - 1$ from the right.

Proof Let $\mathcal{C} = \langle g(x) \rangle$ be a (θ, \mathfrak{S}) -cyclic code over \mathbb{F}_q such that $\mathcal{C}^\perp \subseteq \mathcal{C}$. Note that $h'(x) \in \mathcal{C}^\perp$ and $\mathcal{C}^\perp \subseteq \mathcal{C} = \langle g(x) \rangle$. Thus, $h'(x) = p(x)g(x)$ for some $p(x) \in R$. Now, multiplying both sides by $h'(x)$ from right, we get

$$h'(x)h'(x) = p(x)g(x)h'(x) = p(x)(x^n - 1).$$

Hence, $h'(x)h'(x)$ is divisible by $x^n - 1$ from the right.

Conversely, let $h'(x)h'(x)$ be divisible by $x^n - 1$ from the right. Then $h'(x)h'(x) = b(x)(x^n - 1)$ for some $b(x) \in R$. Now, consider $a(x) \in \mathcal{C}^\perp = \langle h'(x) \rangle$, then $a(x) = c(x)h'(x)$ for some $c(x) \in R$. Multiplying both sides by $h'(x)$ from right and using $h'(x)h'(x) = b(x)(x^n - 1)$, we get

$$\begin{aligned} a(x)h'(x) &= c(x)h'(x)h'(x) = c(x)b(x)(x^n - 1) \\ &= c(x)b(x)h(x)g(x) = c(x)b(x)g(x)h'(x), \\ \left(a(x) - c(x)b(x)g(x) \right) h'(x) &= 0. \end{aligned}$$

As $h'(x)$ is a nonzero polynomial, we have $a(x) - c(x)b(x)g(x) = 0$, which gives $a(x) = c(x)b(x)g(x)$. Therefore, $a(x) \in \mathcal{C} = \langle g(x) \rangle$. Thus, $\mathcal{C}^\perp \subseteq \mathcal{C}$.

Here, we present an example to show the construction of (θ, \mathfrak{S}) -cyclic codes over \mathbb{F}_q with the help of our derived results.

Example 1 Let $q = 49, n = 14$. In \mathbb{F}_{49} , the Frobenius automorphism $\theta : \mathbb{F}_{49} \rightarrow \mathbb{F}_{49}$ is defined by $\theta(a) = a^7$ whereas the θ -derivation \mathfrak{S} is defined by $\mathfrak{S}(a) = w^2(\theta(a) - a)$ for all $a \in \mathbb{F}_{49}$. Therefore, $R = \mathbb{F}_{49}[x; \theta, \mathfrak{S}]$ is a skew polynomial ring. In $\mathbb{F}_{49}[x; \theta, \mathfrak{S}]$, we have

$$\begin{aligned} x^{14} - 1 &= (w^9x^{12} + 3x^{11} + w^{41}x^{10} + w^{13}x^9 + w^{37}x^8 + w^{47}x^7 + w^{18}x^5 + 6x^4 + w^{38}x^3 \\ &\quad + w^{18}x^2 + w^{28}x + w^{12})(w^{39}x^2 + w^3x + w^{17}) = h(x)g(x) \\ &= (w^{39}x^2 + w^3x + w^{17})(w^9x^{12} + 3x^{11} + w^{41}x^{10} + w^{13}x^9 + w^{37}x^8 + w^{47}x^7 \\ &\quad + w^{33}x^5 + 4x^4 + w^{17}x^3 + w^{37}x^2 + w^{13}x + w^{23}) = g(x)h'(x). \end{aligned}$$

Consider $g(x) = w^{39}x^2 + w^3x + w^{17}$, $h(x) = w^9x^{12} + 3x^{11} + w^{41}x^{10} + w^{13}x^9 + w^{37}x^8 + w^{47}x^7 + w^{18}x^5 + 6x^4 + w^{38}x^3 + w^{18}x^2 + w^{28}x + w^{12}$ and $h'(x) = w^9x^{12} + 3x^{11} + w^{41}x^{10} + w^{13}x^9 + w^{37}x^8 + w^{47}x^7 + w^{33}x^5 + 4x^4 + w^{17}x^3 + w^{37}x^2 + w^{13}x + w^{23}$. Then, by Theorem 6 and Equation 2, \mathcal{C} is a (θ, \mathfrak{S}) -cyclic codes over \mathbb{F}_{49} of length 14 which is generated by $g(x)$. The generator and parity check matrices of \mathcal{C} are given by Equation 2 and Theorem 6 respectively. Since, $h'(x)h'(x)$ is divisible by $x^{14} - 1$ from the right and hence the code \mathcal{C} is also a dual-containing code, i.e., $\mathcal{C}^\perp \subseteq \mathcal{C}$.

4 (γ, Δ) -cyclic codes over $\mathcal{R}_{q,s}$

In this section, our main focus is on the algebraic properties of (γ, Δ) -cyclic codes over $\mathcal{R}_{q,s}$ via decomposition over \mathbb{F}_q . To do so, we consider a linear code \mathcal{C} of length n over $\mathcal{R}_{q,s}$. Now, we define

$$\mathcal{C}_i = \left\{ t_i \in \mathbb{F}_q^n \mid \sum_{i=0}^s \zeta_i t_i \in \mathcal{C}, \text{ for some } t_0, t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_s \in \mathbb{F}_q^n \right\}$$

for $0 \leq i \leq s$. Then \mathcal{C}_i is linear code of length n over \mathbb{F}_q and \mathcal{C} can be decomposed as follows

$$\mathcal{C} = \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i.$$

Further, we consider a map $\gamma : \mathcal{R}_{q,s} \rightarrow \mathcal{R}_{q,s}$ defined by

$$\gamma(r) = \sum_{i=0}^s \zeta_i \theta(r_i)$$

where $r = \sum_{i=0}^s \zeta_i r_i$ and $\theta \in \text{Aut}(\mathbb{F}_q)$ defined by $\theta(r_i) = r_i^{p^t}$ for all $r_i \in \mathbb{F}_q$. Then γ is an automorphism on $\mathcal{R}_{q,s}$. Next, we define a map $\Delta : \mathcal{R}_{q,s} \rightarrow \mathcal{R}_{q,s}$ such that

$$\Delta(r) = (1 + v_1 + v_2 + \cdots + v_r)(\gamma(r) - r)$$

where $r = \sum_{i=0}^r r_i v_i$ and $r_i \in \mathbb{F}_q$.

Theorem 8 *The above defined map Δ is a γ -derivation of $\mathcal{R}_{q,s}$.*

Proof Let $r, t \in \mathcal{R}_{q,s}$, we have

$$\begin{aligned} \Delta(r+t) &= (1 + v_1 + v_2 + \cdots + v_s)(\gamma(r+t) - (r+t)) \\ &= (1 + v_1 + v_2 + \cdots + v_s)(\gamma(r) - r) + (1 + v_1 + v_2 + \cdots + v_s)(\gamma(t) - t) \\ &= \Delta(r) + \Delta(t) \end{aligned}$$

and

$$\begin{aligned} \Delta(rt) &= (1 + v_1 + v_2 + \cdots + v_s)(\gamma(rs) - rt) \\ &= (1 + v_1 + v_2 + \cdots + v_s)(\gamma(r)\gamma(t)) - (1 + v_1 + v_2 + \cdots + v_s)rt \\ &= (1 + v_1 + v_2 + \cdots + v_s)(\gamma(r)\gamma(t)) - (1 + v_1 + v_2 + \cdots + v_s)rt \\ &\quad + (1 + v_1 + v_2 + \cdots + v_s)\gamma(r)t - (1 + v_1 + v_2 + \cdots + v_s)\gamma(r)t \\ &= (1 + v_1 + v_2 + \cdots + v_s)\gamma(r)(\gamma(t) - t) - (1 + v_1 + v_2 + \cdots + v_s)(r - \gamma(r))t \\ &= (1 + v_1 + v_2 + \cdots + v_s)\gamma(r)(\gamma(t) - t) + (1 + v_1 + v_2 + \cdots + v_s)(\gamma(r) - r)t \\ &= \Delta(r)t + \gamma(r)\Delta(t). \end{aligned}$$

Hence, Δ is a γ -derivation of $\mathcal{R}_{q,s}$.

Further, with the help of the defined decomposition of \mathcal{C} , we discuss the algebraic properties of (γ, Δ) -cyclic codes over $\mathcal{R}_{q,s}$.

Theorem 9 *Let $\mathcal{C} = \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$ be a linear code of length n over $\mathcal{R}_{q,s}$ where \mathcal{C}_i is a linear code of length n over \mathbb{F}_q for $i = 0, 1, 2, \dots, s$. Then \mathcal{C} is a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$ if and only if \mathcal{C}_i is a (θ, \mathfrak{S}) -cyclic code of length n over \mathbb{F}_q for $i = 0, 1, 2, \dots, s$.*

Proof Let $\mathcal{C} = \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$ be a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$ and $a^i = (a_0^i, a_1^i, \dots, a_{n-1}^i) \in \mathcal{C}_i$, for $0 \leq i \leq s$. Consider $r_j = \sum_{i=0}^s \zeta_i a_j^i$ for $0 \leq j \leq n-1$. Then $r = (r_0, r_1, \dots, r_{n-1}) \in \mathcal{C}$ and $T_{\gamma, \Delta}(r) \in \mathcal{C}$. Again, we have $\gamma(r_j) = \sum_{i=0}^s \zeta_i \theta(a_j^i)$ and $\Delta(r_j) = \Delta(\sum_{i=0}^s \zeta_i a_j^i) = \Delta(\zeta_0 a_j^0) + \Delta(\zeta_1 a_j^1) + \cdots + \Delta(\zeta_s a_j^s)$ for $0 \leq j \leq n-1$. Also,

$$\begin{aligned} \Delta(\zeta_0 a_j^0) &= \Delta(\zeta_0) a_j^0 + \gamma(\zeta_0) \mathfrak{S}(a_j^0) \\ &= \left((1 + v_1 + \cdots + v_s)(\gamma(\zeta_0) - \zeta_0) \right) a_j^0 + \zeta_0 \mathfrak{S}(a_j^0) \\ &= \zeta_0 \mathfrak{S}(a_j^0). \end{aligned}$$

Similarly, $\Delta(\zeta_i a_j^i) = \zeta_i \mathfrak{S}(a_j^i)$ for $i = 1, 2, \dots, s$ and $0 \leq j \leq n-1$. Hence, $T_{\gamma, \Delta}(r) = \sum_{i=0}^s \zeta_i T_{\theta, \mathfrak{S}}(a^i)$. This implies that $T_{\theta, \mathfrak{S}}(a^i) \in \mathcal{C}_i$ for $i = 0, 1, 2, \dots, s$. Thus, \mathcal{C}_i is a (θ, \mathfrak{S}) -cyclic code of length n over \mathbb{F}_q for $i = 0, 1, 2, \dots, s$.

Conversely, suppose \mathcal{C}_i is a (θ, \mathfrak{S}) -cyclic code of length n over \mathbb{F}_q . Let $r = (r_0, r_1, \dots, r_{n-1}) \in \mathcal{C}$ where $r_j = \sum_{i=0}^s \zeta_i a_j^i$ for $0 \leq j \leq n-1$. Consider, $a^i = (a_0^i, a_1^i, \dots, a_{n-1}^i)$, for $0 \leq i \leq s$. Then $a^i \in \mathcal{C}_i$ and also $T_{\theta, \mathfrak{S}}(a^i) \in \mathcal{C}_i$. Similar to the first part of the proof, we have

$$\gamma(r_j) = \sum_{i=0}^s \zeta_i \theta(a_j^i)$$

and

$$\Delta(r_j) = \Delta\left(\sum_{i=0}^s \zeta_i a_j^i\right) = \sum_{i=0}^s \zeta_i \mathfrak{S}(a_j^i)$$

for $i = 0, 1, 2, \dots, s$ and $0 \leq j \leq n-1$. Then

$$\begin{aligned} T_{\gamma, \Delta}(r) &= \gamma(r)M + \Delta(r) = \left(\gamma(r_{n-1}) + \Delta(r_0), \gamma(r_0) + \Delta(r_1), \gamma(r_1) + \Delta(r_2), \dots, \gamma(r_{n-2}) + \right. \\ &\quad \left. \Delta(r_{n-1})\right) \\ &= \sum_{i=0}^s \zeta_i T_{\theta, \mathfrak{S}}(a^i) \in \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i = \mathcal{C}. \end{aligned}$$

Therefore, \mathcal{C} is a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$.

Theorem 10 Let $\mathcal{C} = \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$ be a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$. Then $\mathcal{C} = \langle \zeta_0 g_0(x), \zeta_1 g_1(x), \dots, \zeta_s g_s(x) \rangle$ and $|\mathcal{C}| = q^{(s+1)n - \sum_{i=0}^s \deg(g_i(x))}$, where $g_i(x)$ is a generator polynomial of \mathcal{C}_i for $i = 0, 1, 2, \dots, s$.

Proof Let $\mathcal{C} = \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$ be a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$. Then, by Theorem 9, \mathcal{C}_i is a (θ, \mathfrak{S}) -cyclic code over \mathbb{F}_q , for $i = 0, 1, 2, \dots, s$. This implies that $\mathcal{C}_i = \langle g_i(x) \rangle \subseteq \mathbb{F}_q[x; \theta, \mathfrak{S}] / \langle x^n - 1 \rangle$ for $i = 0, 1, 2, \dots, s$. Thus,

$$\mathcal{C} = \left\{ r(x) \mid r(x) = \sum_{i=0}^s \zeta_i g_i(x), g_i(x) \in \mathcal{C}_i \right\}.$$

Hence, $\mathcal{C} \subseteq \langle \zeta_0 g_0(x), \zeta_1 g_1(x), \dots, \zeta_s g_s(x) \rangle$.

On the other hand, we consider $\zeta_0 f_0(x)g_0(x) + \zeta_1 f_1(x)g_1(x) + \dots + \zeta_s f_s(x)g_s(x) \in \langle \zeta_0 g_0(x), \zeta_1 g_1(x), \dots, \zeta_s g_s(x) \rangle \subseteq \mathcal{R}_{q,s}[x; \gamma, \Delta] / \langle x^n - 1 \rangle$ where $f_i(x) \in \mathcal{R}_{q,s}[x; \gamma, \Delta] / \langle x^n - 1 \rangle$ for $i = 0, 1, 2, \dots, s$. Then there exists $s_i(x) \in \mathbb{F}_q[x; \theta, \mathfrak{S}] / \langle x^n - 1 \rangle$ such that $\zeta_i f_i(x) = \zeta_i s_i(x)$ for $i = 0, 1, 2, \dots, s$. This implies that $\langle \zeta_0 g_0(x), \zeta_1 g_1(x), \dots, \zeta_s g_s(x) \rangle \subseteq \mathcal{C}$. Thus, $\mathcal{C} = \langle \zeta_0 g_0(x), \zeta_1 g_1(x), \dots, \zeta_s g_s(x) \rangle$.

Moreover, $|\mathcal{C}| = |\mathcal{C}_0| |\mathcal{C}_1| \dots |\mathcal{C}_s| = q^{n - \deg(g_0(x))} q^{n - \deg(g_1(x))} \dots q^{n - \deg(g_s(x))} = q^{(s+1)n - \sum_{i=0}^s \deg(g_i(x))}$.

Theorem 11 Let $\mathcal{C} = \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$ be a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$ and $x^n - 1 = h_i(x)g_i(x) = g_i(x)h'_i(x)$ for some monic skew polynomials $g_i(x), h_i(x), h'_i(x) \in \mathbb{F}_q[x; \theta, \mathfrak{S}]$ for $i = 0, 1, 2, \dots, s$. Then $\mathcal{C}^\perp \subseteq \mathcal{C}$ if and only if $h'_i(x)h_i(x)$ is divisible by $x^n - 1$ from the right.

Proof Let $h'_i(x)h_i(x)$ be divisible by $x^n - 1$ from the right for $i = 0, 1, 2, \dots, s$. Then, by Theorem 7, we have $\mathcal{C}_i^\perp \subseteq \mathcal{C}_i$, $i = 0, 1, 2, \dots, s$. This implies that $\bigoplus_{i=0}^s \zeta_i \mathcal{C}_i^\perp \subseteq \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$.

Hence, $\mathcal{C}^\perp \subseteq \mathcal{C}$.

Conversely, let $\mathcal{C}^\perp \subseteq \mathcal{C}$, then $\bigoplus_{i=0}^s \zeta_i \mathcal{C}_i^\perp \subseteq \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$. Now, considering modulo ζ_i , we get $\mathcal{C}_i^\perp \subseteq \mathcal{C}_i$ for $i = 0, 1, 2, \dots, s$. Thus, $h'_i(x)h_i(x)$ is divisible by $x^n - 1$ on the right for $i = 0, 1, 2, \dots, s$.

The next corollary is a direct consequence of the Theorem 11.

Corollary 1 *Let $\mathcal{C} = \langle g(x) \rangle$ be a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$ and $x^n - 1 = h_i(x)g_i(x) = g_i(x)h'_i(x)$ for some monic skew polynomials $g_i(x), h_i(x), h'_i(x) \in \mathbb{F}_q[x; \theta, \mathfrak{S}]$. Then $\mathcal{C}^\perp \subseteq \mathcal{C}$ if and only if $\mathcal{C}_i^\perp \subseteq \mathcal{C}_i$ for $i = 0, 1, 2, \dots, s$.*

5 Constructions of quantum codes and comparison with the existing codes

The theory of quantum error-correcting codes plays a key role in quantum information theory. For a long time, it has been difficult to provide a satisfactory solution to the problem of protecting information from quantum noises. However, after the introduction of the first quantum error-correcting codes by Shor et al. [33], a stream of great developments has emerged in information theory. Let $H_q(\mathbb{C})$ be a q -dimensional Hilbert vector space. Then the set of n -fold tensor product $H_q^n(\mathbb{C}) = \underbrace{H_q(\mathbb{C}) \otimes H_q(\mathbb{C}) \otimes \dots \otimes H_q(\mathbb{C})}_{n \text{ times}}$ is a q^n -

dimensional Hilbert space. Here, a q^k dimensional subspace of $H_q^n(\mathbb{C})$ is called a quantum code with parameters $[[n, k, d]]_q$ where d is the minimum distance, and k is the dimension of the quantum code. Also, \mathcal{C} is dual-containing if $\mathcal{C}^\perp \subseteq \mathcal{C}$. Moreover, in 1997, the quantum Singleton bound for binary codes was introduced by Knill and Laflamme [22]. In 1998, Calderbank et al. [11] provided the quantum Singleton bound for all codes over finite fields as $k + 2d \leq n + 2$. A quantum code is said to be a quantum MDS code if it attains the Singleton bound. In this section, we study quantum error correcting codes from (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$. In order to construct quantum error-correcting codes, we first derive a necessary and sufficient condition for (γ, Δ) -cyclic codes to be dual containing. Note that a quantum code $[[n, k, d]]_q$ is said to be better than $[[n', k', d']]_q$ if any one of the following or both hold:

1. $d > d'$ when the code rate $\frac{k}{n} = \frac{k'}{n'}$ (Larger distance with same code rate).
2. $\frac{k}{n} > \frac{k'}{n'}$ when the distance $d = d'$ (Larger code rate with same distance).

Now, we define a Gray map and study \mathbb{F}_q -images of (γ, Δ) -cyclic codes. Let $GL_{s+1}(\mathbb{F}_q)$ be the set of all $(s+1) \times (s+1)$ invertible matrices over \mathbb{F}_q . Now, $\varphi : \mathcal{R}_{q,s} \rightarrow \mathbb{F}_q^{s+1}$ define by

$$\varphi(r) = (r_0, r_1, \dots, r_s)G,$$

where $r = \sum_{i=0}^s \zeta_i r_i \in \mathcal{R}_{q,s}$, $G \in GL_{s+1}(\mathbb{F}_q)$ such that $GG^T = kI_{s+1}$, G^T is the transpose matrix of G , $k \in \mathbb{F}_q^*$ and I_{s+1} is the identity matrix of order $s+1$. It is easy to check that φ is a bijection and can be extended over $\mathcal{R}_{q,s}^n$ componentwise. If we define Gray distance for a linear code \mathcal{C} by $d_G(\mathcal{C}) = d_H(\varphi(\mathcal{C}))$, then φ is a linear distance preserving map from $(\mathcal{R}_{q,s}^n, d_G)$ to $(\mathbb{F}_q^{n(s+1)}, d_H)$, where d_H is the Hamming distance in \mathbb{F}_q .

Proposition 1 *The Gray map φ is an \mathbb{F}_q -linear and distance preserving map from $\mathcal{R}_{q,s}^n$ (Gray distance) to $\mathbb{F}_q^{(s+1)n}$ (Hamming distance).*

Proof Let $a = (a_0, a_1, \dots, a_{n-1})$, $b = (b_0, b_1, \dots, b_{n-1}) \in \mathcal{R}_{q,s}^n$, where $a_j = \sum_{i=0}^s \zeta_i a_j^i$, $b_j = \sum_{i=0}^s \zeta_i b_j^i$ for $j = 0, 1, \dots, n-1$ and $a_j^i, b_j^i \in \mathbb{F}_q$. Then

$$\begin{aligned} \varphi(a+b) &= \varphi(a_0 + b_0, a_1 + b_1, \dots, a_{n-1} + b_{n-1}) \\ &= \varphi(\zeta_0(a_0^0 + b_0^0) + \zeta_1(a_0^1 + b_0^1) + \dots + \zeta_s(a_0^s + b_0^s), \dots, \zeta_0(a_{n-1}^0 + b_{n-1}^0) \\ &\quad + \zeta_1(a_{n-1}^1 + b_{n-1}^1) + \dots + \zeta_s(a_{n-1}^s + b_{n-1}^s)) \\ &= [(\zeta_0(a_0^0 + b_0^0), \zeta_1(a_0^1 + b_0^1), \dots, \zeta_s(a_0^s + b_0^s))G, \dots, (\zeta_0(a_{n-1}^0 + b_{n-1}^0), \zeta_1(a_{n-1}^1 + b_{n-1}^1), \dots, \\ &\quad \zeta_s(a_{n-1}^s + b_{n-1}^s))G] \\ &= [(\zeta_0 a_0^0, \zeta_1 a_0^1, \dots, \zeta_s a_0^s)G, \dots, (\zeta_0 a_{n-1}^0, \zeta_1 a_{n-1}^1, \dots, \zeta_s a_{n-1}^s)G] + [(\zeta_0 b_0^0, \zeta_1 b_0^1, \dots, \zeta_s b_0^s)G, \dots, \\ &\quad (\zeta_0 b_{n-1}^0, \zeta_1 b_{n-1}^1, \dots, \zeta_s b_{n-1}^s)G] \\ &= \varphi(a) + \varphi(b). \end{aligned}$$

Now, for any $\lambda \in \mathbb{F}_q$, we have

$$\begin{aligned} \varphi(\lambda a) &= \varphi(\lambda a_0, \lambda a_1, \dots, \lambda a_{n-1}) \\ &= \varphi(\lambda \zeta_0 a_0^0 + \lambda \zeta_1 a_0^1 + \dots + \lambda \zeta_s a_0^s, \dots, \lambda \zeta_0 a_{n-1}^0 + \lambda \zeta_1 a_{n-1}^1 + \dots + \lambda \zeta_s a_{n-1}^s) \\ &= [(\lambda \zeta_0 a_0^0, \lambda \zeta_1 a_0^1, \dots, \lambda \zeta_s a_0^s)G, \dots, (\lambda \zeta_0 a_{n-1}^0, \lambda \zeta_1 a_{n-1}^1, \dots, \lambda \zeta_s a_{n-1}^s)G] \\ &= [\lambda(\zeta_0 a_0^0, \zeta_1 a_0^1, \dots, \zeta_s a_0^s)G, \dots, \lambda(\zeta_0 a_{n-1}^0, \zeta_1 a_{n-1}^1, \dots, \zeta_s a_{n-1}^s)G] \\ &= \lambda[(\zeta_0 a_0^0, \zeta_1 a_0^1, \dots, \zeta_s a_0^s)G, \dots, (\zeta_0 a_{n-1}^0, \zeta_1 a_{n-1}^1, \dots, \zeta_s a_{n-1}^s)G] \\ &= \lambda \varphi(a). \end{aligned}$$

Moreover, $d_G(a, b) = \omega_G(a - b) = \omega_H(\varphi(a - b)) = \omega_H(\varphi(a) - \varphi(b)) = d_H(\varphi(a), \varphi(b))$. Hence, φ is a distance preserving map.

Theorem 12 *If \mathcal{C} is an $[n, k, d_G]$ linear code over $\mathcal{R}_{q,s}$, then $\varphi(\mathcal{C})$ is a $[(s+1)n, k, d_H]$ linear code over \mathbb{F}_q .*

Proof Follows directly from Proposition 1 and the definition of the Gray map.

The Gray map φ preserves the orthogonality as shown in the next result.

Lemma 1 *Let \mathcal{C} be a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$. Then $\varphi(\mathcal{C})^\perp = \varphi(\mathcal{C}^\perp)$. Further, \mathcal{C} is self-dual if and only if $\varphi(\mathcal{C})$ is self-dual.*

Proof Let $c = (c_0, c_1, \dots, c_{n-1}) \in \mathcal{C}$ and $d = (d_0, d_1, \dots, d_{n-1}) \in \mathcal{C}^\perp$ where $c_j = \sum_{i=0}^s \zeta_i c_j^i$, $b_j = \sum_{i=0}^s \zeta_i d_j^i$ for $j = 0, 1, \dots, n-1$ and $c_j^i, d_j^i \in \mathbb{F}_q$. Now, $c \cdot d = \sum_{j=0}^{n-1} c_j d_j = 0$ gives $\sum_{j=0}^{n-1} (c_j^0 d_j^0 + c_j^1 d_j^1 + \dots + c_j^s d_j^s) = 0$. Again,

$$\varphi(c) = [(\zeta_0 c_0^0, \zeta_1 c_0^1, \dots, \zeta_s c_0^s)G, \dots, (\zeta_0 c_{n-1}^0, \zeta_1 c_{n-1}^1, \dots, \zeta_s c_{n-1}^s)G] = (\alpha_0 G, \dots, \alpha_{n-1} G)$$

and

$$\varphi(d) = [(\zeta_0 d_0^0, \zeta_1 d_0^1, \dots, \zeta_s d_0^s)G, \dots, (\zeta_0 d_{n-1}^0, \zeta_1 d_{n-1}^1, \dots, \zeta_s d_{n-1}^s)G] = (\beta_0 G, \dots, \beta_{n-1} G),$$

where $\alpha_j = (c_j^0, c_j^1, \dots, c_j^s)$ and $\beta_j = (d_j^0, d_j^1, \dots, d_j^s)$ for $0 \leq j \leq n-1$ and $GG^T = kI_{s+1}$. Also,

$$\begin{aligned} \varphi(c) \cdot \varphi(d) &= \varphi(c)\varphi(d)^T = \sum_{j=0}^{n-1} \alpha_j GG^T \beta_j^T \\ &= k \sum_{j=0}^{n-1} \alpha_j \beta_j^T \\ &= k \sum_{j=0}^{n-1} (c_j^0 d_j^0 + c_j^1 d_j^1 + \dots + c_j^s d_j^s) = 0. \end{aligned}$$

Since $c \in \mathcal{C}$ and $d \in \mathcal{C}^\perp$ are arbitrary, $\varphi(\mathcal{C}^\perp) \subseteq (\varphi(\mathcal{C}))^\perp$. On the other hand, as φ is a bijective linear map, $|\varphi(\mathcal{C}^\perp)| = |(\varphi(\mathcal{C}))^\perp|$. Therefore, $\varphi(\mathcal{C}^\perp) = (\varphi(\mathcal{C}))^\perp$.

Lemma 2 ([16], Theorem 3) *Let \mathcal{C} be an $[n, k, d]$ linear code over \mathbb{F}_q such that $\mathcal{C}^\perp \subseteq \mathcal{C}$. Then there exists a quantum code with parameters $[[n, 2k - n, d]]_q$.*

Theorem 13 *Let $\mathcal{C} = \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$ be a (γ, Δ) -cyclic code of length n over $\mathcal{R}_{q,s}$. Also, let $\mathcal{C}_i = \langle g_i(x) \rangle$ be a (θ, \mathfrak{S}) -cyclic code over \mathbb{F}_q where $x^n - 1 = h_i(x)g_i(x) = g_i(x)h'_i(x)$ for some monic skew polynomials $g_i(x), h_i(x), h'_i(x) \in \mathbb{F}_q[x; \theta, \mathfrak{S}]$, for $i = 0, 1, \dots, s$. Further, let $h'_i(x)h_i(x)$ be divisible by $x^n - 1$ from the right for $i = 0, 1, \dots, s$. Then there exists a quantum code with parameters $[[s+1)n, 2k - (s+1)n, d_H]]_q$.*

Proof Let $h'_i(x)h_i(x)$ be divisible by $x^n - 1$ from right for $i = 0, 1, \dots, s$. Then from Theorem 11, we have $\mathcal{C}^\perp \subseteq \mathcal{C}$. Also, by Lemma 1, we have $\varphi(\mathcal{C}^\perp) = \varphi(\mathcal{C})^\perp$, and hence $\varphi(\mathcal{C})^\perp \subseteq \varphi(\mathcal{C})$. Thus, $\varphi(\mathcal{C})$ is a dual containing linear code with parameters $[(s+1)n, k, d_H]$ over \mathbb{F}_q . Further, by Lemma 2, there exists a quantum code with parameters $[[s+1)n, 2k - (s+1)n, d_H]]_q$.

Next, with the help of our established results, we construct many new quantum codes possessing better parameters than the existing codes, which are appeared in [13, 36]. In the following examples, $\mathbb{F}_q^* = \langle w \rangle$ denotes the cyclic group of non-zero elements of \mathbb{F}_q generated by $w \in \mathbb{F}_q$. All examples' computations are carried out using the Magma computation system [10].

Example 2 Let $q = 8$, $s = 3$ and $\mathcal{R}_{8,3} = \mathbb{F}_8[v_1, v_2, v_3] / \langle v_1^2 - v_1, v_2^2 - v_2, v_3^2 - v_3, v_1v_2 = v_2v_1 = v_2v_3 = v_3v_2 = v_3v_1 = v_1v_3 = 0 \rangle$, where $\mathbb{F}_8 = \mathbb{F}_2(w)$ and $w^3 + w + 1 = 0$. Let $n = 30$, $\theta : \mathbb{F}_8 \rightarrow \mathbb{F}_8$ be the Frobenius automorphism defined by $\theta(a) = a^2$, and the θ -derivation $\mathfrak{S} : \mathbb{F}_8 \rightarrow \mathbb{F}_8$ is defined by $\mathfrak{S}(a) = w(\theta(a) - a)$ for all $a \in \mathbb{F}_8$. Therefore, $\mathbb{F}_8[x; \theta, \mathfrak{S}]$ is a skew polynomial ring. In $\mathbb{F}_8[x; \theta, \mathfrak{S}]$, we have

$$\begin{aligned}
x^{30} - 1 &= (w^6x^{29} + w^4x^{28} + w^6x^{27} + w^4x^{26} + w^3x^{25} + x^{24} + w^6x^{23} + w^4x^{22} + w^6x^{21} \\
&\quad + w^4x^{20} + w^6x^{19} + w^4x^{18} + w^6x^{17} + w^4x^{16} + w^6x^{15} + w^4x^{14} + w^3x^{13} \\
&\quad + x^{12} + w^6x^{11} + w^4x^{10} + w^3x^9 + x^8 + w^6x^7 + w^4x^6 + w^3x^5 + x^4 + w^6x^3 \\
&\quad + w^4x^2 + w^3x + 1)(w^2x + 1) = h_0(x)g_0(x) \\
&= (w^2x + 1)(w^6x^{29} + w^4x^{28} + w^6x^{27} + w^4x^{26} + w^6x^{25} + w^4x^{24} + w^6x^{23} \\
&\quad + w^4x^{22} + w^6x^{21} + w^4x^{20} + w^6x^{19} + w^4x^{18} + w^6x^{17} + w^4x^{16} + w^6x^{15} \\
&\quad + w^4x^{14} + w^6x^{13} + w^4x^{12} + w^6x^{11} + w^4x^{10} + w^6x^9 + w^4x^8 + w^6x^7 \\
&\quad + w^4x^6 + w^6x^5 + w^4x^4 + w^6x^3 + w^4x^2 + w^6x + w^4) = g_0(x)h'_0(x) \\
x^{30} - 1 &= (w^5x^{28} + w^3x^{27} + w^2x^{26} + w^3x^{25} + w^3x^{24} + w^4x^{23} + w^6x^{22} + w^5x^{21} \\
&\quad + w^6x^{20} + wx^{19} + w^3x^{18} + x^{17} + w^6x^{16} + w^3x^{15} + w^6x^{14} + w^6x^{13} + w^4x^{12} \\
&\quad + w^2x^{11} + w^6x^9 + w^5x^8 + w^6x^6 + w^4x^5 + w^5x^4 + w^4x^2 + w^2x + 1)(wx^2 \\
&\quad + w^4x + w^6) = h_1(x)g_1(x) \\
&= (wx^2 + w^4x + w^6)(w^5x^{28} + w^3x^{27} + w^2x^{26} + w^3x^{25} + x^{24} + w^3x^{22} + w^6x^{21} \\
&\quad + w^4x^{20} + x^{19} + w^5x^{18} + x^{16} + w^6x^{15} + w^5x^{13} + w^3x^{12} + w^2x^{11} + w^3x^{10} \\
&\quad + x^9 + w^3x^7 + w^6x^6 + w^4x^5 + x^4 + w^5x^3 + x + w^6) = g_1(x)h'_1(x) \\
x^{30} - 1 &= (w^6x^{28} + w^6x^{27} + wx^{26} + wx^{24} + x^{23} + w^5x^{22} + x^{20} + w^6x^{19} + w^5x^{17} \\
&\quad + w^3x^{16} + w^2x^{15} + w^3x^{14} + x^{13} + w^4x^{12} + w^6x^9 + wx^8 + w^6x^7 + wx^6
\end{aligned}$$

$$\begin{aligned}
& + wx^5 + w^2x^4 + wx^3 + w^4x^2 + w^2x)(w^4x^2 + w^3x + w) = h_2(x)g_2(x) \\
& = (w^4x^2 + w^3x + w)(w^6x^{28} + w^6x^{27} + wx^{26} + w^2x^{24} + wx^{23} + wx^{22} + w^3x^{21} \\
& \quad + w^2x^{20} + w^2x^{19} + w^5x^{18} + w^5x^{17} + x^{16} + w^2x^{15} + w^6x^{13} + w^6x^{12} + wx^{11} \\
& \quad + w^2x^9 + wx^8 + wx^7 + w^3x^6 + w^2x^5 + w^2x^4 + w^5x^3 + w^5x^2 + x + w^2) \\
& = g_2(x)h'_2(x) \\
x^{30} - 1 & = (x^{28} + w^4x^{27} + w^3x^{26} + w^3x^{25} + wx^{24} + w^4x^{23} + x^{22} + w^4x^{20} + x^{18} + x^{17} \\
& \quad + w^5x^{16} + w^5x^{15} + w^4x^{14} + w^5x^{13} + w^6x^{11} + w^5x^{10} + wx^9 + w^4x^8 + x^7 \\
& \quad + wx^6 + wx^5 + w^3x^3 + w^6x^2 + w^6x + 1)(x^2 + w^2x + w^4) = h_3(x)g_3(x) \\
& = (x^2 + w^2x + w^4)(x^{28} + w^4x^{27} + w^3x^{26} + w^3x^{25} + wx^{24} + w^2x^{23} + w^5x^{22} \\
& \quad + w^2x^{21} + w^4x^{20} + wx^{19} + w^6x^{18} + x^{17} + w^5x^{16} + w^6x^{15} + x^{13} + w^4x^{12} \\
& \quad + w^3x^{11} + w^3x^{10} + wx^9 + w^2x^8 + w^5x^7 + w^2x^6 + w^4x^5 + wx^4 + w^6x^3 \\
& \quad + x^2 + w^5x + w^6) = g_3(x)h'_3(x)
\end{aligned}$$

Now, let $g_0 = w^2x + 1$, $g_1 = wx^2 + w^4x + w^6$, $g_2 = w^4x^2 + w^3x + w$ and $g_3 = x^2 + w^2x + w^4$. Then $\mathcal{C}_i = \langle g_i(x) \rangle$ is a (θ, \mathfrak{S}) -cyclic code of length 30 over \mathbb{F}_8 for $i = 0, 1, 2, 3$. Then by

Theorem 9, $\mathcal{C} = \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$ is a (γ, Δ) -cyclic code of length 30 over $\mathcal{R}_{8,3}$. Let

$$G = \begin{pmatrix} 1 & w & w^3 & 1 \\ w & 1 & 1 & w^3 \\ w^3 & 1 & 1 & w \\ 1 & w^3 & w & 1 \end{pmatrix} \in GL_4(\mathbb{F}_8)$$

such that $GG^T = I_4$. Then $\varphi(\mathcal{C})$ is a $[120, 114, 4]$ linear code over \mathbb{F}_8 . Again,

$$\begin{aligned}
h'_0(x)h'_0(x) & = (w^2x^{28} + x^{27} + x^{26} + w^5x^{25} + w^4x^{24} + w^2x^{22} + x^{21} + x^{20} + w^5x^{19} \\
& \quad + w^2x^{18} + x^{17} + x^{16} + w^5x^{15} + w^2x^{14} + x^{13} + w^3x^{12} + w^4x^{11} + x^{10} \\
& \quad + w^5x^9 + w^4x^8 + w^2x^6 + x^5 + w^3x^4 + w^4x^3 + x^2 + w^5x + w^4)(x^{30} - 1) \\
h'_1(x)h'_1(x) & = (wx^{26} + w^4x^{25} + w^2x^{24} + wx^{23} + wx^{22} + w^3x^{21} + w^5x^{19} + w^4x^{16} \\
& \quad + x^{15} + w^2x^{14} + w^3x^{13} + wx^{12} + w^4x^{11} + w^6x^{10} + w^6x^9 + w^2x^8 + wx^7 \\
& \quad + w^4x^6 + x^4 + x^2 + w^6x + w^6)(x^{30} - 1) \\
h'_2(x)h'_2(x) & = (w^4x^{26} + w^3x^{25} + w^3x^{24} + w^4x^{23} + x^{22} + w^3x^{21} + w^4x^{20} + w^3x^{19} \\
& \quad + w^2x^{18} + wx^{17} + wx^{16} + w^5x^{15} + x^{14} + w^5x^{13} + w^5x^{12} + w^2x^{11} \\
& \quad + w^6x^9 + w^6x^8 + w^2x^7 + w^5x^6 + x^5 + w^4x^4 + w^6x^3 + w^5x^2 + w^5x) \\
& \quad (x^{30} - 1) \\
h'_3(x)h'_3(x) & = (x^{26} + w^2x^{25} + w^5x^{24} + x^{23} + w^2x^{22} + wx^{20} + x^{19} + w^3x^{18} + w^6x^{17} \\
& \quad + wx^{16} + x^{15} + wx^{14} + w^6x^{12} + w^6x^{11} + w^5x^{10} + x^9 + w^3x^8 + x^7 \\
& \quad + x^6 + w^6x^5 + w^6x^3 + w^4x^2 + w^6x + w^6)(x^{30} - 1).
\end{aligned}$$

From above, we see that $h'_i(x)h'_i(x)$ is divisible by $(x^{30} - 1)$ on the right for $i = 0, 1, 2, 3$. Hence, by Theorem 13, there exists a quantum code with parameters $[[120, 108, 4]]_8$ which has the same length and distance but better code rate than the best-known code $[[120, 104, 4]]_8$ given by [13].

Example 3 Let $q = 25$, $s = 3$ and $\mathcal{R}_{25,3} = \mathbb{F}_{25}[v_1, v_2, v_3]/\langle v_1^2 - v_1, v_2^2 - v_2, v_3^2 - v_3, v_1v_2 = v_2v_1 = v_2v_3 = v_3v_2 = v_3v_1 = v_1v_3 = 0 \rangle$. Let $n = 30$, $\theta : \mathbb{F}_{25} \rightarrow \mathbb{F}_{25}$ be the Frobenius

automorphism defined by $\theta(a) = a^5$, and the θ -derivation $\mathfrak{S} : \mathbb{F}_{25} \rightarrow \mathbb{F}_{25}$ is defined by $\mathfrak{S}(a) = w(\theta(a) - a)$ for all $a \in \mathbb{F}_{25}$. Therefore, $\mathbb{F}_{25}[x; \theta, \mathfrak{S}]$ is a skew polynomial ring. In $\mathbb{F}_{25}[x; \theta, \mathfrak{S}]$, we have

$$\begin{aligned}
x^{20} - 1 &= (w^{19}x^{19} + x^18 + w^{20}x^{17} + w^4x^{16} + w^{15}x^{15} + w^{20}x^{14} + x^{13} + w^{19}x^{12} + w^7x^{11} \\
&\quad + w^2x^{10} + w^{10}x^9 + 3x^8 + w^3x^7 + w^{17}x^6 + w^{11}x^5 + 2x^4 + 3x^2 + 4x + 3)(wx \\
&\quad + w^{17}) = h_0(x)g_0(x) \\
&= (wx + w^{17})(w^{19}x^{19} + x^{18} + w^{20}x^{17} + w^4x^{16} + w^{15}x^{15} + wx^{14} + 2x^{13} + w^2x^{12} \\
&\quad + w^{10}x^{11} + w^{21}x^{10} + w^7x^9 + 4x^8 + w^8x^7 + w^{16}x^6 + w^3x^5 + w^{13}x^4 + 3x^3 \\
&\quad + w^{14}x^2 + w^{22}x + w^9) = g_0(x)h'_0(x) \\
x^{20} - 1 &= (w^{14}x^{18} + w^8x^{17} + w^{17}x^{16} + 3x^{15} + 2x^{14} + w^{21}x^{13} + w^8x^{12} + w^{10}x^{11} + wx^{10} \\
&\quad + 4x^9 + w^{19}x^7 + w^{19}x^6 + w^9x^5 + 2x^4 + 4x^3 + x + 2)(w^{10}x^2 + 2x + w^{11}) \\
&= h_1(x)g_1(x) \\
&= (w^{10}x^2 + 2x + w^{11})(w^{14}x^{18} + w^8x^{17} + w^{17}x^{16} + 3x^{15} + 2x^{14} + w^{15}x^{13} \\
&\quad + w^3x^{12} + x^{11} + w^{15}x^{10} + 3x^9 + w^4x^8 + 4x^7 + w^4x^6 + w^{10}x^5 + x^4 + 2x^3 \\
&\quad + w^{11}x^2 + w^{19}x + w^{19}) = g_1(x)h'_1(x) \\
x^{20} - 1 &= (w^{23}x^{19} + w^{19}x^{18} + w^3x^{17} + w^{14}x^{16} + x^{15} + w^4x^{14} + w^{19}x^{13} + w^{10}x^{12} \\
&\quad + w^{13}x^{11} + 2x^{10} + w^2x^9 + w^{13}x^8 + w^7x^7 + w^{21}x^6 + 4x^5 + 2x^4 + 3x^2 + 4x \\
&\quad + 3)(w^5x + 3) = h_2(x)g_2(x) \\
&= (w^5x + 3)(w^{23}x^{19} + w^{19}x^{18} + w^3x^{17} + w^{14}x^{16} + x^{15} + w^5x^{14} + wx^{13} + \\
&\quad w^9x^{12} + w^{20}x^{11} + 2x^{10} + w^{11}x^9 + w^7x^8 + w^{15}x^7 + w^2x^6 + 4x^5 + w^{17}x^4 \\
&\quad + w^{13}x^3 + w^{21}x^2 + w^8x + 3) = g_2(x)h'_2(x) \\
x^{20} - 1 &= (w^{10}x^{18} + 4x^{17} + w^{11}x^{16} + 4x^{15} + w^{16}x^{14} + w^7x^{13} + 3x^{12} + w^8x^{11} \\
&\quad + w^{21}x^{10} + w^{13}x^9 + 3x^8 + 2x^7 + w^{14}x^6 + w^2x^5 + 4x^4 + 4x^3 + x^2 + 4x \\
&\quad + 2)(w^{14}x^2 + w^{19}x + w^{15}) = h_3(x)g_3(x) \\
&= (w^{14}x^2 + w^{19}x + w^{15})(w^{10}x^{18} + 4x^{17} + w^{11}x^{16} + 4x^{15} + w^{16}x^{14} + w^2x^{13} \\
&\quad + x^{12} + 3x^{11} + w^4x^{10} + w^4x^9 + w^7x^8 + 2x^7 + w^{22}x^6 + w^9x^5 + w^{10}x^4 \\
&\quad + x^3 + w^{13}x + w^2) = g_3(x)h'_3(x)
\end{aligned}$$

Now, let $g_0(x) = wx + w^{17}$, $g_1(x) = w^{10}x^2 + 2x + w^{11}$, $g_2(x) = w^5x + 3$ and $g_3(x) = w^{14}x^2 + w^{19}x + w^{15}$. Then $\mathcal{C}_i = \langle g_i(x) \rangle$ is a (θ, \mathfrak{S}) -cyclic code of length 20 over \mathbb{F}_{25} for $i = 0, 1, 2, 3$. Then by Theorem 9, $\mathcal{C} = \bigoplus_{i=0}^3 \zeta_i \mathcal{C}_i$ is a (γ, Δ) -cyclic code of length 20 over $\mathcal{R}_{25,3}$. Let

$$G = \begin{pmatrix} -1 & 1 & 1 & 1 \\ 1 & 1 & 1 & -1 \\ 1 & -1 & 1 & 1 \\ 1 & 1 & -1 & 1 \end{pmatrix} \in GL_4(\mathbb{F}_{25})$$

such that $GG^T = 4I_2$. Then $\varphi(\mathcal{C})$ is a $[80, 74, 4]$ linear code over \mathbb{F}_{25} . Again,

$$\begin{aligned}
h'_0(x)h'_0(x) &= (3x^{18} + w^4x^{17} + w^{19}x^{16} + w^2x^{15} + w^{23}x^{13} + w^8x^{12} + w^9x^{11} + w^{11}x^{10} \\
&\quad + w^{21}x^8 + wx^7 + w^5x^6 + 3x^5 + wx^3 + 2x^2 + 3x + w^{15})(x^{20} - 1) \\
h'_1(x)h'_1(x) &= (w^4x^{16} + w^{13}x^{15} + w^{14}x^{14} + w^2x^{13} + w^{17}x^{12} + w^9x^{11} + w^{20}x^{10} + w^7x^9 \\
&\quad + x^8 + w^{14}x^7 + w^{22}x^6 + w^{21}x^5 + w^8x^4 + 4x^3 + w^{22}x^2 + wx + w^{13})
\end{aligned}$$

Table 1: MDS quantum codes from (θ, \mathfrak{S}) -cyclic codes over \mathbb{F}_q

(n, q)	$\mathfrak{S}(a), a \in \mathbb{F}_q$	$g(x)$	\mathcal{C}	Obtained Codes	Remark
(30, 8)	$w(\theta(a) - a)$	(w^3w^5)	$[30, 29, 2]_8$	$[[30, 28, 2]]_8$	MDS
(30, 8)	$w(\theta(a) - a)$	(ww^4w^6)	$[30, 28, 2]_8$	$[[30, 26, 2]]_8$	Near to MDS
(14, 49)	$w^2(\theta(a) - a)$	$(w^{39}w^3w^{17})$	$[14, 12, 2]_{49}$	$[[14, 10, 2]]_{49}$	Near to MDS
(24, 49)	$w^2(\theta(a) - a)$	(w^41)	$[24, 23, 2]_{49}$	$[[24, 22, 2]]_{49}$	MDS
(24, 49)	$\theta(a) - a$	(w^2w^{17})	$[24, 23, 2]_{49}$	$[[24, 22, 2]]_{49}$	MDS
(20, 25)	$w(\theta(a) - a)$	$(w^{14}w^{19}w^{15})$	$[20, 18, 2]_{25}$	$[[20, 16, 2]]_{25}$	Near to MDS
(24, 25)	$\theta(a) - a$	$(w^3w^4w^{20})$	$[24, 22, 2]_{25}$	$[[24, 20, 2]]_{25}$	Near to MDS
(20, 25)	$w(\theta(a) - a)$	(ww^{17})	$[20, 19, 2]_{25}$	$[[20, 18, 2]]_{25}$	MDS

$$\begin{aligned}
& (x^{20} - 1) \\
h'_2(x)h'_2(x) &= (3x^{18} + w^{20}x^{17} + w^{10}x^{16} + 2x^{15} + w^{19}x^{13} + 4x^{12} + w^{21}x^{11} + 4x^{10} \\
& \quad + w^9x^8 + w^{15}x^7 + 2x^6 + 3x^5 + w^5x^3 + wx^2 + w^{13}x + 1)(x^{20} - 1) \\
h'_3(x)h'_3(x) &= (w^{20}x^{16} + w^{23}x^{15} + 2x^{14} + w^{14}x^{13} + w^{21}x^{12} + w^{13}x^{11} + w^{16}x^{10} \\
& \quad + w^{11}x^9 + w^5x^8 + w^{11}x^7 + w^{11}x^6 + w^{17}x^5 + w^8x^4 + w^8x^3 + w^{11}x^2 \\
& \quad + w^{16}x + w^{20})(x^{20} - 1).
\end{aligned}$$

From above we see that $h'_i(x)h'_i(x)$ is divisible by $x^{20} - 1$ on the right for $i = 0, 1, 2, 3$. Hence, by Theorem 13, there exists a quantum code with parameters $[[80, 68, 4]]_{25}$ which has the same length and distance, but better code rate than the best-known code $[[80, 64, 4]]_{25}$ given by [36].

Let \mathcal{C} be a (θ, \mathfrak{S}) -cyclic code of length n over \mathbb{F}_q where $\mathcal{C} = \langle g(x) \rangle$ and $x^n - 1 = h(x)g(x) = g(x)h'(x)$ for some monic skew polynomials $g(x), h(x), h'(x) \in \mathbb{F}_q[x; \theta, \mathfrak{S}]$. Further, let $h'(x)h'(x)$ be divisible by $x^n - 1$ from the right. Therefore, by Theorem 11, we get the dual containing codes with \mathbb{F}_q -parameters $[n, k, d]_q$ (enlisted in the fourth column of Table 1). Also, by Lemma 2, we construct quantum codes $[[n, k, d]]_q$ (in the fifth column), in which some codes satisfy the equality $n - k + 2 - 2d = 2$ (Near to MDS), and some are

MDS (maximum-distance-separable). Let $\mathcal{C} = \bigoplus_{i=0}^s \zeta_i \mathcal{C}_i$ be a (γ, Δ) -cyclic code of length

n over $\mathcal{R}_{q,s}$ where $\mathcal{C}_i = \langle g_i(x) \rangle$ is a (θ, \mathfrak{S}) -cyclic code of length n over \mathbb{F}_q and $x^n - 1 = h_i(x)g_i(x) = g_i(x)h'_i(x)$ for some monic skew polynomials $g_i(x), h_i(x), h'_i(x) \in \mathbb{F}_q[x; \theta, \mathfrak{S}]$ for $i = 0, 1, 2, \dots, s$. Further, let $h'_i(x)h'_i(x)$ is divisible by $x^n - 1$ from the right for $i = 0, 1, 2, \dots, s$. Therefore, by Theorem 11, we get the dual containing codes with \mathbb{F}_q -parameters $[n, k, d]_q$ (enlisted in the fifth column of Table 2). Also, by Theorem 13, we construct quantum codes $[[n, k, d]]_q$ (in the sixth column), which beat the parameters of best-known codes (in the seventh column) given by the online database [13, 36]. Also, the first and second columns represent s and (n, q) , respectively. Moreover, in third column we present θ -derivations $\mathfrak{S}(a)$ for $a \in \mathbb{F}_q$. Note that in fourth column we give generator polynomials g_i for \mathcal{C}_i ($i = 0, 1, 2, \dots, s$) which is a right factor of $x^n - 1$ in $\mathbb{F}_q[x; \theta, \mathfrak{S}]$. In order to make Table 2 precise, we enlist the coefficients of polynomials in decreasing powers of x . For example, we write w^702w to represent the polynomial $w^7x^3 + 2x + w$.

Table 2: New quantum codes from (γ, Δ) -cyclic codes over $\mathcal{R}_{q,s}$

s	(n, q)	$\mathfrak{S}(a), a \in \mathbb{F}_q$	$[g_0(x), g_1(x), \dots, g_s(x)]$	$\varphi(\mathcal{C})$	Obtained Codes	Existing Codes
2	(48, 9)	$w^2(\theta(a) - a)$	$(w^7 1w^3, w^5 w^2, w^5 12)$	$[144, 139, 3]_9$	$[[144, 134, 3]]_9$	$[[146, 134, 3]]_9$ [13]
3	(36, 9)	$w^2(\theta(a) - a)$	$(w^2 1w^5, 1w^3, w^7 w^2, 2ww^3)$	$[144, 138, 4]_9$	$[[144, 132, 4]]_9$	$[[146, 128, 4]]_9$ [13]
3	(32, 9)	$w^2(\theta(a) - a)$	$(w^2 ww, w^6 w^2 2, 2w^2 ww^3, w^3 1)$	$[128, 120, 4]_9$	$[[128, 112, 4]]_9$	$[[129, 103, 4]]_9$ [13]
2	(42, 9)	$w^2(\theta(a) - a)$	$(ww^6, w^6 ww^3, w^7 w^2)$	$[126, 122, 3]_9$	$[[126, 118, 3]]_9$	$[[130, 118, 3]]_9$ [36]
2	(60, 4)	$w(\theta(a) - a)$	$(www^2, 1w^2 w^2, 11w^2)$	$[180, 174, 3]_4$	$[[180, 168, 3]]_4$	$[[185, 167, 3]]_4$ [36]
3	(20, 25)	$w(\theta(a) - a)$	$(ww^{17}, w^{10} 2w^{11}, w^5 3, w^{14} w^{19} w^{15})$	$[80, 74, 4]_{25}$	$[[80, 68, 4]]_{25}$	$[[80, 64, 4]]_{25}$ [36]
3	(40, 25)	$w(\theta(a) - a)$	$(w^{19} w^{10}, w^{10} w^{14}, w^{11} w^{17}, w^{14} w^4)$	$[120, 116, 3]_{25}$	$[[120, 112, 3]]_{25}$	$[[120, 106, 3]]_{25}$ [36]
3	(30, 8)	$w(\theta(a) - a)$	$(w^2 1, ww^4 w^6, w^4 w^3 w, 1w^2 w^4)$	$[120, 114, 4]_8$	$[[120, 108, 4]]_8$	$[[120, 104, 4]]_8$ [13]
3	(32, 8)	$w(\theta(a) - a)$	$(w^6 w^3 w^4, w^2 w^5 w^5, ww^6, w^6 w^5 w^2)$	$[128, 121, 4]_8$	$[[128, 114, 4]]_8$	$[[128, 112, 4]]_8$ [13]

6 Conclusion

In this paper, we have constructed many quantum codes over a class of finite commutative non-chain rings $\mathcal{R}_{q,s}$, with better parameters than the codes available in recent literature. Particularly, we have obtained (γ, Δ) -cyclic codes using a set of idempotents over $\mathcal{R}_{q,s}$ and established results on their algebraic structure. Towards construction of quantum codes, a necessary and sufficient condition to contain their dual codes has established. Finally, we have obtained many better quantum codes. On the other hand, exploring applications in the quantum computation of these codes is still open as future research work.

Acknowledgements

The first and second authors are thankful to the Department of Science and Technology (DST), Govt. of India, for financial support under CRG/2020/005927, vide Diary No. SERB/F/6780/ 2020-2021 dated 31 December 2020 and Ref No. DST/INSPIRE/03/2016/001445, respectively.

Declarations

Data Availability Statement: The authors declare that [the/all other] data supporting the findings of this study are available within the article. Any clarification may be requested from corresponding author provided it is essential.

Competing interests: The authors declare that there is no conflict of interest regarding the publication of this manuscript.

Use of AI tools declaration The authors declare that they have not used Artificial Intelligence (AI) tools in the creation of this manuscript.

References

1. Abualrub, T., Ghrayeb, A., Aydin, N., Siap, I.: On the construction of skew quasi cyclic codes. *IEEE Trans. Info. Theory* **56**, 2081 – 2090 (2010).
2. Alahmadi, A., Islam, H., Prakash, O., Solé, P., Alkenani, A., Muthana, N., Hijazi, R.: New quantum codes from constacyclic codes over a non-chain ring. *Quantum Inf Process* **20**(60), (2021), <https://doi.org/10.1007/s11128-020-02977-y>.
3. Ashraf, M., Mohammad, G.: Quantum codes over \mathbb{F}_p from cyclic codes over $\mathbb{F}_p[u, v]/\langle u^2 - 1, v^3 - v, uv - vu \rangle$. *Cryptogr. Commun.* **11**(2), 325 – 335 (2019).
4. Bhaintwal, M.: Skew quasi cyclic codes over Galois rings. *Des. Codes Cryptogr.* **62**(1), 85 – 101 (2012).
5. Boucher, D., Geiselmann, W., Ulmer, F.: Skew cyclic codes. *Appl. Algebra Engrg. Comm. Comput.* **18**(4), 379 – 389 (2007).
6. Boucher, D., Solé, P., Ulmer, F.: Skew constacyclic codes over Galois ring. *Adv. Math. Commun.* **2**(3), 273 – 292 (2008).
7. Boucher, D., Ulmer, F.: Linear codes using skew polynomials with automorphisms and derivations. *Des. Codes Cryptogr.* **70**(3), 405 – 431 (2014).
8. Boucher, D., Ulmer, F.: Coding with skew polynomial rings. *J. Symbolic Comput.* **44**(12), 1644 – 1656 (2009).
9. Boulagouaz, M., H., Leroy, A.: (σ, δ) -codes, *Adv. Math. Commun.* **7**(4), 463 – 474 (2013).
10. Bosma, W., Cannon, J.: *Handbook of Magma Functions*. Univ. of Sydney, Sydney (1995).

11. Calderbank, A.R., Rains, E. M., Shor, P. M., Sloane, N. J. A.: Quantum error correction via codes over $GF(4)$. *IEEE Trans. Inform. Theory* **44**, 1369 – 1387 (1998).
12. Dertli, A., Cengellenmis, Y., Eren, S.: On quantum codes obtained from cyclic codes over A_2 . *Int. J. Quantum Inf.* **13**(3), 1550031 (2015).
13. Edel, Y.: Some good quantum twisted codes. <https://www.mathi.uni-heidelberg.de/~yves/Matritzen/QT BCH/QT BCHIndex.html>.
14. Gao, Y., Gao, J., Fu, F. W.: On quantum codes from cyclic codes over the ring $\mathbb{F}_q + v_1\mathbb{F}_q + \dots + v_r\mathbb{F}_q$. *Appl. Algebra Eng. Commun. Comput.* **30**(2), 161 – 174 (2019).
15. Goodearl, K. R., Warfield, R. B.: An introduction to noncommutative Noetherian rings. Cambridge University Press, (1989).
16. Grassl, M., Beth, T.: On optimal quantum codes. *Int. J. Quantum Inf.* **2**, 55-64 (2004).
17. Gursoy, F., Siap, I., Yildiz, B.: Construction of skew cyclic codes over $\mathbb{F}_q + v\mathbb{F}_q$. *Adv. Math. Commun.* **8**(3), 313 – 322 (2014).
18. Hammons, A. R., Kumar, P. V., Calderbank, A. R., Sloane, N. J. A., Sole, P.: The \mathbb{Z}_4 -linearity of Kerdock, Preparata, Goethals, and related codes. *IEEE Trans. Info. Theory* **40**, 301 – 319 (1994).
19. Islam, H., Patel, S., Prakash, O., Solé, P.: A family of constacyclic codes over a class of non-chain rings $A_{q,r}$ and new quantum codes. *J. Appl. Math. Comput.* **68**(4), 2493-2514 (2022).
20. Jacobson, N.: Pseudo-linear transformations. *Ann. Math.* **38**(2), 484 – 507 (1937).
21. Jitman, S., Ling, S., Udomkavanich, P.: Skew constacyclic codes over finite chain rings. *Adv. Math. Commun.*, **6**, 39 – 63 (2012).
22. Knill, E., Laflamme, R.: A theory of quantum error-correcting codes. *Phys. Rev. A* **55**(2) (1997), 900 – 911.
23. Leroy, A.: Pseudo-linear transformation and evaluation in Ore extension. *Bull. Belg. Math. Soc.* **2**, 321 – 345 (1995).
24. Leroy, A.: Noncommutative polynomial maps. *J. Algebra Appl.* **11**(4), 1250076 (2012).
25. Li, J., Gao, J., Fu, F. W., Ma, F.: $\mathbb{F}_q R$ -linear skew constacyclic codes and their application of constructing quantum codes. *Quantum Inf Process.* <https://doi.org/10.1007/s11128-020-02700-x> (2020).
26. Ma, F., Gao, J., Li, J., Fu, F. W.: (σ, δ) -Skew quasi-cyclic codes over the ring $\mathbb{Z}_4 + u\mathbb{Z}_4$. *Cryptogr. Commun.* (2021), <https://doi.org/10.1007/s12095-020-00467-7>.
27. Ma, F., Gao, J., Fu, F. W.: Constacyclic codes over the ring $\mathbb{F}_p + v\mathbb{F}_p + v^2\mathbb{F}_p$ and their applications of constructing new non-binary quantum codes. *Quantum Inf. Process.* **17**(6), 4 (2018).
28. Ozen, M., Ozzaim, T., Ince, H.: Skew quasi cyclic codes over $\mathbb{F}_q + v\mathbb{F}_q$. *J. Algebra Appl.* **18**(4), 1950077, 16 pp (2019).
29. Patel, S., Prakash, O.: (θ, δ_θ) -cyclic codes over $\mathbb{F}_q[u, v]/\langle u^2 - u, v^2 - v, uv - vu \rangle$. *Des. Codes Cryptogr.* **90**(11), 2763-2781 (2022).
30. Patel, S., Prakash, O.: Skew generalized cyclic code over $R[x_1; \sigma_1, \delta_1][x_2; \sigma_2, \delta_2]$. [arXiv:1907.06086](https://arxiv.org/abs/1907.06086) (2019).
31. Prakash, O., Islam, H., Patel, S., Solé, P.: New quantum codes from skew constacyclic codes over a class of non-chain rings $R_{e,q}$. *Internat. J. Theoret. Phys.* **60**(9), 1-19 (2021).
32. Sharma, A., Bhaintwal, M.: A class of skew-cyclic codes over $\mathbb{Z}_4 + u\mathbb{Z}_4$ with derivation. *Adv. Math. Commun.* **12**(4), 1 – 17 (2018).
33. Shor, P. W.: Scheme for reducing decoherence in quantum memory *Phys. Rev. A.* **52**, 2493-2496 (1995).
34. Siap, I., Abualrub, T., Aydin, N., Seneviratne, P.: Skew cyclic codes of arbitrary length. *Int. J. Inf. Coding Theory* **2**(1), 10 – 20 (2011).

35. Tapia Cuitino L.F., Tironi, A. L.: Some properties of skew codes over finite fields. *Des. Codes Cryptogr.* **85**, 359 – 380 (2017).
36. Verma, R. K., Prakash, O., Singh, A., Islam, H.: New quantum codes from skew constacyclic codes. *Adv. Math. Commun.* doi: 10.3934/amc.2021028 (2021).