

Information-Theoretic Generalization Bounds for Deep Neural Networks

Haiyun He, Christina Lee Yu, and Ziv Goldfeld

Abstract

Deep neural networks (DNNs) exhibit an exceptional capacity for generalization in practical applications. This work aims to capture the effect and benefits of depth for supervised learning via information-theoretic generalization bounds. We first derive two hierarchical bounds on the generalization error in terms of the Kullback-Leibler (KL) divergence or the 1-Wasserstein distance between the train and test distributions of the network internal representations. The KL divergence bound shrinks as the layer index increases, while the Wasserstein bound implies the existence of a layer that serves as a generalization funnel, which attains a minimal 1-Wasserstein distance. Analytic expressions for both bounds are derived under the setting of binary Gaussian classification with linear DNNs. To quantify the contraction of the relevant information measures when moving deeper into the network, we analyze the strong data processing inequality (SDPI) coefficient between consecutive layers of three regularized DNN models: Dropout, DropConnect, and Gaussian noise injection. This enables refining our generalization bounds to capture the contraction as a function of the network architecture parameters. Specializing our results to DNNs with a finite parameter space and the Gibbs algorithm reveals that deeper yet narrower network architectures generalize better in those examples, although how broadly this statement applies remains a question.

Index Terms

Deep neural network, generalization error, internal representation, information theory, SDPI

I. INTRODUCTION

Overparameterized deep neural networks (DNNs) have solidified their status as the model of choice for numerous high-dimensional and large-scale learning tasks, primarily due to their remarkable generalization performance. Efforts have been invested into theoretically explaining this phenomenon from various perspectives. This includes approached via norm-based complexity measures [1]–[3], PAC-Bayes bounds [4]–[9], sharpness and flatness of the loss minima [10]–[12], loss landscape [13], implicit regularization induced by the gradient descent algorithms [14]–[16], etc. The reader is referred to the recent survey [17] for a comprehensive literature review. Despite this wealth of research, the precise factors contributing to the generalization capacity of DNNs remain elusive, as indicated in [18], [19]. The goal of this work is to shed new light on the advantages of deep models for learning under the framework of information-theoretic generalization bounds.

The generalization error is the difference between the population risk and the empirical risk on the training data. It measures the extent of overfitting of a trained neural network when the empirical risk is pushed to zero. Information-theoretic generalization bounds have been widely explored in recent years. This line of work was initiated by [20], [21], where a generalization error bound in terms of the mutual information between the input and output of the learning algorithm was derived (see also [22]). These inaugural results inspired various extensions and refinements based on chaining arguments [23], [24], conditioning and processing techniques [25]–[28], as well as other information-theoretic quantities [29]–[32]. However, the aforementioned results were not specialized to the DNN setting and hence did not capture the effect of depth on the generalization bound. Quantifying this effect within such information-theoretic bounds is the main objective of this work.

A. Main Contributions

We present two new hierarchical generalization error bounds for DNNs. The first bound refines the results from [20]–[22] for sub-Gaussian loss functions, bounding the generalization error in terms of the Kullback-Leibler (KL) divergence and mutual information associated with the internal representations of each layer. This bound shrinks as the layer count increases, can adapt to layers of low complexity (e.g., low-dimensional or discrete), and overall highlights the benefits of depth for learning. Our second generalization bound accounts for Lipschitz continuous losses and employs the 1-Wasserstein distance. This bound suggests the existence of a DNN layer that minimizes the generalization upper bound, acting as a *generalization funnel layer*. The bounds are evaluated for the task of binary Gaussian mixture classification and closed-form expressions derived. These expressions demonstrate that the KL divergence bounds shrinks with the ranks of the product of weight matrices, while the

H. He is supported by Cornell CAM Postdoctoral Fellowship. C. Yu is partially supported by NSF grants CNS-1955997, AFOSR grant FA9550-23-1-0301, and an Intel Rising Stars award. Z. Goldfeld is partially supported by NSF grants CCF-2046018, DMS-2210368, and CCF-2308446, and the IBM Academic Award.

This paper was presented in part at the InfoCog workshop at Annual Conference on Neural Information Processing Systems 2023 and submitted in part to International Symposium on Information Theory 2024.

H. He is with Center for Applied Mathematics, C. Yu is with the School of Operations Research and Information Engineering, and Z. Goldfeld is with the School of Electrical and Computer Engineering, Cornell University, Ithaca, NY 14850 USA. (Emails: {hh743, cleeyu, goldfeld}@cornell.edu)

Wasserstein bound (and the resulting generalization funnel layer) depends on the Frobenius norms of the product of weight matrices. We compute the generalization funnel layer on a simple numerical example, illustrating that it can vary between problems and generally depends on the employed training method.

Our hierarchical KL divergence bound qualitatively demonstrates that the generalization bound improves as we delve deeper into the network. To quantify this contraction, we employ the strong data processing inequality (SDPI), which requires some stochasticity in the DNN (otherwise, the SDPI coefficient degenerates to 1). We consider three popular randomized regularization techniques: Dropout [33], DropConnect [34], and Gaussian noise injection [35]–[39]. The SDPI coefficient associated with the stochastic channel induced by each layer is then controlled in terms of the network parameters (depth, width, activation functions, dropout/noise statistics). The desired generalization bound is then obtained by peeling off the DNN layers and aggregating the corresponding contraction coefficients. This results serves to tighten existing generalization bounds for generic supervised learning settings [20]–[22].

Our analysis demonstrates that the product of the contraction coefficients across the layers vanishes as the network depth and dropout probabilities (or noise level) increase, or the layer widths decrease. This highlights the advantage of deep network architectures and stochasticity. We also instantiate our results for the Gibbs algorithm [40], [41], yielding an $O(\frac{1}{n})$ generalization bound that decreases monotonically as the product of the contraction coefficients shrinks. Our bounds and their dependence on the problem parameters are visualized via a simple numerical example of a DNN with a finite parameter space. In this instance, we numerically show a deeper but narrower neural network architecture yields a better generalization performance. Overall, our results provide a information-theoretic perspective for understanding the generalization capabilities of DNNs.

B. Related Works

The generalization performance of deep learning has been extensively studied over the past decades, although a comprehensive and satisfactory account is still obscure. Motivated by empirical evidence that overparameterized DNNs tend to generalize better [18], [42], [43], numerous attempts to pin this observation down theoretically were made. Some control the generalization error using norm-based complexity measures [1]–[3], such as the Rademacher complexity with norm constraints [1], the Frobenius/spectral/ ℓ_1 -norm of weight matrices [2], or the Fisher-Rao norm [3]. A comparisons between several norm-based complexity measures can be found in [2], [3]. Other works employ PAC-Bayes bounds [6], [7] to understand DNNs in terms of compressibility and robustness [4], [5], [8], [9]. The relationship between generalization and the sharpness/flatness of the loss minima or the landscape of loss was studied in [10]–[13]. The seminal works [18], [44] analyzed the finite-sample expressivity of DNNs and proved that any two-layer NN with ReLU activation can generalize, given sufficiently many parameters. A numerically-tight DNN generalization bound based on validation and training datasets were derived in [19]. They also suggested that factors like explicit regularization, minimum norm solution, low complexity, stability, flat minima, etc., many not be necessary for achieving good generalization (see also [18]). The effect of implicit regularization induced by the gradient descent algorithms on generalization was explored in [14]–[16], [45].

The information-theoretic study of generalization was initiated by [20]–[22], where the generalization error was controlled by the mutual information between the input and output of the learning algorithm. These results capture the intuition that a learning algorithm generalizes when it extracts relevant information from the training data while disregarding irrelevant information (cf. [35], [46]–[49] for related work on the information bottleneck principle for deep learning). Inspired by this novel perspective, many followup works set to derive tighter bounds, e.g., using chaining arguments [23] or conditional mutual information [25]–[27]. Information-theoretic bounds that employ other information/discrepancy measures were also explored, encompassing f -divergences, α -Rényi divergences, the generalized Jensen-Shannon divergence, the Wasserstein distance, and more [29]–[32], [50]. Some works have attempted to obtain information-theoretic generalization bounds by considering specific optimization algorithms. Generalization of the stochastic gradient Langevin dynamics (SGLD) was studied in [26], [51], [52], capturing the effect of training iterations, step size, and noise level. By relating the dynamics in SGLD to stochastic gradient descent (SGD), generalization bounds for SGD were also derived [53], [54]. However, these previous works did not relate the generalization bounds to the architecture of DNNs. Our goal is to fill this gap by quantifying the effects of the network depth, layer width, activation functions, and injected stochasticity on generalization.

C. Paper Outline

The rest of our paper is organized as follows. In Section II, we define the notations and formulate the supervised learning problem under a feedforward DNN model. In Section III, we present the hierarchical generalization bounds based on the KL divergence and the Wasserstein distance, respectively. We then specialize our bounds to the case of binary Gaussian classification and derive the analytical expressions. In Section IV, we quantify the contraction of the relevant information measures in the hierarchical KL divergence bounds as the layer index grows. Subsequently, we derive tighter generalization bounds and visualize them using an instance of DNNs with a finite parameter space. Furthermore, we obtain the analytical expressions of the bounds for the Gibbs algorithm. Section V concludes our discussion and outlines avenues for future work. The proofs of our results are provided in the appendices.




Fig. 1: L -layer feedforward network.

II. PRELIMINARIES AND PROBLEM FORMULATION

A. Notation

The class of Borel probability measures on $\mathcal{X} \subseteq \mathbb{R}^d$ is denoted by $\mathcal{P}(\mathcal{X})$. A random variable $X \sim P_X \in \mathcal{P}(\mathcal{X})$ is called σ -sub-Gaussian, if $\mathbb{E}[\exp(\lambda(X - \mathbb{E}[X]))] \leq \lambda^2 \sigma^2 / 2$ for any $\lambda \in \mathbb{R}$. For two measures $\mu, \nu \in \mathcal{P}(\mathcal{X})$, $\mu \ll \nu$ means that $\nu(E) = 0$ implies $\mu(E) = 0$ for all measurable set E . The f -divergence between $\mu, \nu \in \mathcal{P}(\mathcal{X})$ ($\mu \ll \nu$) is defined by $D_f(\mu \parallel \nu) := \int f(d\mu/d\nu) d\nu$, where $f : (0, +\infty) \rightarrow \mathbb{R}$ is convex and $f(1) = 0$. The Kullback-Leibler (KL) divergence is defined by taking $f(u) = u \log u$. The Hellinger (H^2) distance is defined by taking $f(u) = (1 - \sqrt{u})^2$. The total variation (TV) distance is defined by taking $f(u) = \frac{1}{2}|u - 1|$. The mutual information between $(X, Y) \sim P_{X,Y} \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ is defined as $I(X; Y) := D_{\text{KL}}(P_{X,Y} \parallel P_X \otimes P_Y)$. The Shannon entropy of a discrete random variable $X \sim P_X \in \mathcal{P}(\mathcal{X})$ is $H(X) = H(P_X) = \log(|\mathcal{X}|) - D_{\text{KL}}(P_X \parallel \text{Unif}(\mathcal{X}))$. For $p \in [1, \infty)$ and a pair of probability measures μ, ν on a metric space $(\mathcal{X}, \|\cdot\|)$ with $\|\cdot\|$ being the Euclidean distance, the p -Wasserstein distance between them is defined as $W_p(\mu, \nu) := (\inf_{\pi \in \Pi(\mu, \nu)} \mathbb{E}_{(x, x') \sim \pi} [\|x - x'\|^p])^{1/p}$, where $\Pi(\mu, \nu)$ is the set of couplings on \mathcal{X}^2 with marginal distributions μ and ν . For a d -dimensional vector X and integers $1 \leq i < j \leq d$, we use the shorthands $X_i^j := (X_i, \dots, X_j)$ and $[j] := \{1, 2, \dots, j\}$. For a function $f : \mathbb{R}^d \rightarrow \mathbb{R}^d$, where $f = (f_1, \dots, f_d)$, we define $\|f\|_\infty := \sup_{x \in \mathbb{R}^d} \max_{i=1, \dots, d} |f_i(x)|$. For a vector v , define $\|v\| := \sqrt{v^\top v}$ as the Euclidean norm. For a matrix \mathbf{A} , define $\|\mathbf{A}\|_{\text{op}} := \sup\{\|\mathbf{A}v\| \mid \|v\| = 1\}$ as the operator norm and $\|\mathbf{A}\|_{\text{F}} := \sqrt{\text{tr}(\mathbf{A}\mathbf{A}^*)}$ as the Frobenius norm.

B. Supervised Learning Problem

Consider a data space $\mathcal{X} \subseteq \mathbb{R}^{d_0}$ and label set $\mathcal{Y} = [K] \subseteq \mathbb{N}$. Fix a data distribution $P_{X,Y} \in \mathcal{P}(\mathcal{X} \times \mathcal{Y})$ and let $(X, Y) \sim P_{X,Y}$ be a nominal data feature-label pair. The training dataset $D_n = \{(X_i, Y_i)\}_{i=1}^n$ comprises independently and identically distributed (i.i.d.) copies of (X, Y) ; note that $P_{D_n} = P_{X,Y}^{\otimes n}$. We consider a feedforward DNN model with L layers for predicting the label Y from the test sample X via $\hat{Y} := g_{\mathbf{w}_L} \circ g_{\mathbf{w}_{L-1}} \circ \dots \circ g_{\mathbf{w}_1}(X)$, where $g_{\mathbf{w}_l}(t) = \phi_l(\mathbf{w}_l t)$, $l \in [L]$, for a weight matrix $\mathbf{w}_l \in \mathbb{R}^{d_l \times d_{l-1}}$ and an activation function $\phi_l : \mathbb{R} \rightarrow \mathbb{R}$ (acting on vectors element-wise). Denote all the network parameters by $\mathbf{w} = (\mathbf{w}_1, \dots, \mathbf{w}_L)$ and the parameter space by $\mathcal{W} \subseteq \mathbb{R}^{d_1 \times d_0} \times \dots \times \mathbb{R}^{d_L \times d_{L-1}}$. We denote the internal representation of the l^{th} layer by $T_l := g_{\mathbf{w}_l} \circ \dots \circ g_{\mathbf{w}_1}(X)$, $l \in [L]$, noting that $T_0 = X$. When the input to the network is X_i (rather than X), we add a subscript i to the internal representation notation, writing $T_{l,i}$ instead of T_l . See Figure 1 for an illustration. The setup can be generalized to regression problems by setting $\mathcal{Y} \subseteq \mathbb{R}$. Furthermore, our arguments extend to the case when the training dataset D_n comprises dependent but identically distributed data samples, e.g., ones generated from a Markov chain Monte Carlo method.

Let $\ell : \mathcal{W} \times \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R}_+$ be the loss function. Given any $\mathbf{w} \in \mathcal{W}$, the *population risk* and the *empirical risk* are respectively defined as

$$\mathcal{L}_{\text{P}}(\mathbf{w}, P_{X,Y}) := \mathbb{E}[\ell(\mathbf{w}, X, Y)], \quad ; \quad \mathcal{L}_{\text{E}}(\mathbf{w}, D_n) := \frac{1}{n} \sum_{i=1}^n \ell(\mathbf{w}, X_i, Y_i),$$

where the loss function ℓ penalizes the discrepancy between the true label Y and the DNN prediction $\hat{Y} = g_{\mathbf{w}_L} \circ \dots \circ g_{\mathbf{w}_1}(X)$, i.e., $\ell(\mathbf{w}, x, y) = \tilde{\ell}(g_{\mathbf{w}_L} \circ \dots \circ g_{\mathbf{w}_1}(x), y)$. A learning algorithm trained with D_n can be characterized by a stochastic mapping $P_{\mathbf{W}|D_n}$. Given any $(P_{\mathbf{W}|D_n}, P_{X,Y})$, the *expected generalization error* is defined as the expected gap between the population empirical risks:

$$\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y}) := \mathbb{E}[\mathcal{L}_{\text{P}}(\mathbf{W}, P_{X,Y}) - \mathcal{L}_{\text{E}}(\mathbf{W}, D_n)], \quad (1)$$

where the expectation is w.r.t. $P_{(X,Y), D_n, \mathbf{W}} = P_{X,Y}^{\otimes(n+1)} P_{\mathbf{W}|D_n}$.

III. HIERARCHICAL GENERALIZATION BOUND

Existing results such as [20]–[22] bound the generalization error from (1) in terms of the mutual information terms $I(D_n; \mathbf{W})$ or $\sum_{i=1}^n I(X_i, Y_i; \mathbf{W})$, which only depend on the raw input dataset and the algorithm. We next establish two improved generalization bounds, whose hierarchical structure captures the effect of the internal representations T_l . The first bound shrinks

as one moves deeper into the network, providing new evidence for the benefits of deep models for learning. The second bound is minimized by one of the network layers, suggesting the existence of a ‘funnel’ layer that governs generalization.

A. KL Divergence Bound

We present the following generalization bound for the above described setting.

Theorem 1 (Hierarchical generalization bound). *Suppose that the loss function $\ell(\mathbf{w}, X, Y)$ is σ -sub-Gaussian under $P_{X,Y}$, for all $\mathbf{w} \in \mathcal{W}$. We have*

$$|\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y})| \leq \text{UB}(L) \leq \text{UB}(L-1) \leq \dots \leq \text{UB}(0), \quad (2)$$

where $\text{UB}(l) := \frac{\sigma\sqrt{2}}{n} \sum_{i=1}^n \sqrt{l(T_{l,i}, Y_i; \mathbf{W}_{l+1}^L | \mathbf{W}_1^L) + \text{D}_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_{l,i}, Y_i | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L})}$, $l = 0, \dots, L$.

Theorem 1 is derived by first establishing the $\text{UB}(L)$ upper bound via the Donsker-Varadhan variational representation of the KL divergence and the sub-Gaussianity of the loss function. We then invoke the data processing inequality (DPI) to successively peel off the layers to arrive at the remaining bounds. See Appendix A for a detailed proof. While the $\text{UB}(L)$ forms the tightest bound, the state hierarchy highlights the benefit of depth for learning and lend well for comparison to existing results. Indeed, observing that $\text{UB}(0) = \sqrt{2\sigma^2} n^{-1} \sum_{i=1}^n \sqrt{l(X_i, Y_i; \mathbf{W})}$, we see that our bound is tighter than the one from [22].

Theorem 1 shows that the model generalizes when both $l(T_{l,i}, Y_i; \mathbf{W}_{l+1}^L | \mathbf{W}_1^L)$ and $\text{D}_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_{l,i}, Y_i | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L})$ are small, for some layer $l = 0, \dots, L$. This happens when the parameters of subsequent layers are not overly dependent on the l^{th} input internal representation, and when the learned posterior of this internal representation highly matches the prior.

Remark 1 (Special cases). *To gain further intuition, we present two special cases under which the bounds simplify:*

1) One-to-one mapping: *If $g_{\mathbf{W}_l}$ is one-to-one for all $l \in [L]$, the DPI holds with equality:*

$$\begin{aligned} \text{D}_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_{l,i}, Y_i | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}) &= \text{D}_{\text{KL}}(P_{X_i, Y_i | \mathbf{W}_1^L} \| P_{X_i, Y_i | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}) = l(X_i, Y_i; \mathbf{W}_1^L), \\ l(T_{l,i}, Y_i; \mathbf{W}_{l+1}^L | \mathbf{W}_1^L) &= l(X_i, Y_i; \mathbf{W}_{l+1}^L | \mathbf{W}_1^L). \end{aligned}$$

Thus, the upper bounds are equal: $\text{UB}(L) = \text{UB}(L-1) = \dots = \text{UB}(0)$, which implies that the representation at each layer has the same effect on the generalization error.

2) Discrete latent space: *The generalization bound simplifies when T_l can only take finitely many values (e.g., the discrete latent layer in the VQ-VAE [55]). Assuming that $t_l(\mathbf{w}_1^L) := \min_{t \in \mathcal{T}_l, y \in \mathcal{Y}} P_{T_{l,i}, Y_i | \mathbf{W}_1^L}(t, y | \mathbf{w}_1^L) \in (0, |\mathcal{T}_l \times \mathcal{Y}|^{-1})$ and $\bar{t}_l := \sup_{\mathbf{w}_1^L} t_l(\mathbf{w}_1^L)$, we have*

$$\text{UB}(l) \leq \sqrt{2\sigma^2 \log(K^2 / \bar{t}_l)}.$$

As \bar{t}_l grows, we see that $P_{T_{l,i}, Y_i | \mathbf{W}_1^L}$ tends to the uniform distribution on $\mathcal{T}_l \times \mathcal{Y}$ and its entropy/variance increases. This, in turn, shrinks the generalization error, which is consistent with the intuition that stochasticity leads to better generalization. The proof is provided in Appendix B.

B. Wasserstein Distance Bound

Akin to Theorem 1, we present a generalization error bound based on the Wasserstein distance. Unlike the KL divergence, Wasserstein distances do not generally follow the DPI, and hence the presented bound does not adhere to a descending hierarchical structure. Instead, it shows that there exists a layer that minimizes the Wasserstein generalization bound.

Theorem 2 (Min Wasserstein generalization bound). *Suppose that the loss function $\tilde{\ell} : \mathcal{Y} \times \mathcal{Y} \rightarrow \mathbb{R}_{\geq 0}$ is ρ_0 -Lipschitz and the activation function $\phi_l : \mathbb{R} \rightarrow \mathbb{R}$ is ρ_l -Lipschitz, for each $l = 1, \dots, L$. We have*

$$\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y}) \leq \min_{l=0, \dots, L} \frac{\rho_0}{n} \sum_{i=1}^n \mathbb{E} \left[\left(1 \vee \prod_{j=l+1}^L \rho_j \|\mathbf{W}_j\|_{\text{op}} \right) W_1(P_{T_{l,i}, Y_i | \mathbf{W}}(\cdot | \mathbf{W}), P_{T_{l,i}, Y_i | \mathbf{W}}(\cdot | \mathbf{W})) \right].$$

The derivation of the bound relies on Kantorovich–Rubinstein duality, which ties W_1 to the difference of expectations defining the generalization error. See Appendix C for the proof details. As the Wasserstein distance is monotonically increasing in the order (i.e., $W_p \leq W_q$ whenever $p \leq q$), the 1-Wasserstein distance provides the sharpest bound. Compared to [32, Theorem 2] and [56, Theorem 1], we make a weaker assumption on the loss function and take Lipschitz continuity of activation functions into account, which holds for the commonly used loss and activation functions. Compared to the KL divergence bound from Theorem 1, which degenerates when the considered distributions are supported on different domains, the Wasserstein distance is robust to mismatched supports and the corresponding bound is meaningful even in that setting. Theorem 2 suggests that the generalization bound is controlled by a certain layer that achieves the smallest weighted 1-Wasserstein distance between




Fig. 2: Illustration of binary Gaussian mixture data with $d_0 = 2$ and the Bayes optimal linear classifier.

the distributions of the training and test internal representations. This layer serves as a funnel that determines the overall generalization performance; thus, we call it *generalization funnel layer*. It suggests that within a DNN, there exists a specific layer that exerts a stronger impact on generalization compared to others.

Remark 2 (Comparison with KL divergence bound). *Assume that the loss function (bounded within $[0, A] \subset \mathbb{R}_{\geq 0}$) and the activation functions in the DNN model satisfy the Lipschitz continuity conditions in Theorem 2. Under this assumption, the loss function $\ell(\mathbf{w}, X, Y)$, where $(X, Y) \sim P_{X, Y}$, is $\frac{A}{2}$ -subGaussian for all \mathbf{w} . When $\rho_0 K^2 \leq A$, the generalization bound given in Theorem 2 is tighter than $\text{UB}(L)$ in Theorem 1. A proof of this claim is provided in Appendix D, and utilizes [57, Theorem 4], Pinsker's and Bretagnolle-Huber inequalities.*

C. Case Study: Binary Gaussian Mixture Classification

To better understand the generalization bounds from Theorems 1 and 2 and assess their dependence on depth, we consider the following binary Gaussian mixture example and evaluate the bounds analytically.

Classification problem setting. Consider the binary classification problem illustrated in Fig. 2, where the input data distribution is a binary Gaussian mixture: $P_Y = \text{Unif}\{-1, +1\}$ and $P_{X|Y=y} = \mathcal{N}(y\mu_0, \sigma_0^2 \mathbf{I}_{d_0})$ for $y \in \{\pm 1\}$, where $\mu_0 \in \mathbb{R}^d$ and $\sigma_0 > 0$. The goal is to classify the binary label Y given the feature X . Notice that under this setting, the Bayes optimal classifier is $Y^* = \tanh(\mu_0^\top X)$.

Model and algorithm. Consider a classifier that is realized by a linear L -layer neural network composed with a hyperbolic tangent nonlinearity, i.e., $\hat{Y}(\mathbf{w}) = \tanh(\mathbf{w}_{\otimes L}^\top X)$, where $\mathbf{w}_{\otimes l} := \mathbf{w}_l \mathbf{w}_{l-1} \cdots \mathbf{w}_1$. To train the model to approach the Bayes optimal classifier $\tanh(\mu_0^\top X)$, we consider a set of algorithms $P_{\mathbf{W}|D_n}$ under which the output network parameters satisfy $\mathbf{W}_{\otimes L}^\top = \frac{1}{n} \sum_{i=1}^n Y_i X_i$, and set the prediction to $\hat{Y} = \hat{Y}(\mathbf{W})$. The rationale behind this choice of algorithm comes from observing that $Y_i X_i \sim \mathcal{N}(\mu_0, \sigma_0^2 \mathbf{I}_{d_0})$ are i.i.d. for $i = 1, \dots, n$, and by the strong law of large numbers we have $\mathbf{W}_{\otimes L}^\top \rightarrow \mu_0$ almost surely, as $n \rightarrow \infty$. Performance is measured using the quadratic loss function $\ell(\mathbf{w}, X, Y) = (Y - \tanh(\mathbf{w}_{\otimes L}^\top X))^2$, which is bounded inside $[0, 4]$ and is thus 2-sub-Gaussian under $P_{X, Y}$, for all \mathbf{w} .

Analysis. We move to evaluate the generalization bounds in Theorems 1 and 2 by computing the prior and posterior distributions and the divergences between them. Proofs of subsequent claims are all deferred to Appendix E.

Lemma 3 (Prior and posterior of (X_i, Y_i)). *For any $i = 1, \dots, n$ and $y \in \{\pm 1\}$, the prior distribution of $X_i|Y_i = y$ is given by $P_{X_i|Y_i=y} = \frac{1}{2} \mathcal{N}(y\mu_0, \sigma_0^2 \mathbf{I}_{d_0})$, while its posterior distribution given the model $\mathbf{W}_{\otimes L}$ is $P_{X_i|Y_i=y, \mathbf{W}_{\otimes L}} = \frac{1}{2} \mathcal{N}(y \mathbf{W}_{\otimes L}^\top, \frac{(n-1)\sigma_0^2}{n} \mathbf{I}_{d_0})$. Furthermore, we have $P_{Y_i|\mathbf{w}, \mathbf{W}_{\otimes L}} = P_{Y_i|\mathbf{W}_{\otimes L}} = \text{Unif}\{-1, +1\}$ and $P_{X_i, Y_i|\mathbf{w}, \mathbf{W}_{\otimes L}} = P_{X_i, Y_i|\mathbf{W}_{\otimes L}}$.*

Given the above expressions for the involved distributions, we evaluate the KL divergence generalization bound from Theorem 1 as follows.

Proposition 4 (KL divergence bound evaluation). *Under the binary Gaussian classification setting, we have*

$$|\text{gen}(P_{\mathbf{W}|D_n}, P_{X, Y})| \leq \widetilde{\text{UB}}(L) \leq \widetilde{\text{UB}}(L-1) \leq \cdots \leq \widetilde{\text{UB}}(0),$$

where $\widetilde{\text{UB}}(l) := 2\sqrt{\mathbb{E}[r_l](\log \frac{n}{n-1} - \frac{1}{n}) + \frac{d_0}{n}}$, $r_0 = d_0$, and $r_l = \text{rank}(\mathbf{W}_{\otimes l})$, for $l \in [L]$.

As a sanity check, observe that $\widetilde{\text{UB}}_n(l)$ converges to 0 as $n \rightarrow \infty$, for all $l = 0, 1, \dots, L$, as expected. Recalling that $\text{rank}(\mathbf{AB}) \leq \text{rank}(\mathbf{A}) \wedge \text{rank}(\mathbf{B})$, we see that $r_L \leq r_{L-1} \leq \cdots \leq r_1 \leq r_0 = d_0$. Consequently, the contraction from

TABLE I: The generalization funnel layer index l^* for differently generated model \mathbf{W} when $L = 10$ in Example 1.

Generating methods $\prod_{j=1}^{l'} C_j = 0.2 \ \frac{1}{n} \sum_{i=1}^n Y_i X_i\ $	$l' = 3$	$l' = 5$	$l' = 7$
Generalization funnel layer l^*	3	5	7

$\widetilde{\text{UB}}_n(l-1)$ to $\widetilde{\text{UB}}_n(l)$ is evident and quantified by the gap between the ranks of $\mathbf{W}_{\otimes(l-1)}$ and $\mathbf{W}_{\otimes l}$, namely, $r_{l-1} - r_l$. Note that in our example, $\text{rank}(\mathbf{W}_L) = 1$ and thus $\text{rank}(\mathbf{W}_{\otimes L}) = 1$, independent of the depth L , which means that the tightest bound, $\widetilde{\text{UB}}(L)$, does not change with L . Nevertheless, the intermediate bounds $\widetilde{\text{UB}}(l)$, for $l \in [L-1]$, generally shrink as L grows, depicting the trajectory of generalization performance of the internal layers. Extending the above example beyond the classification setting to representation learning, where the output representation dimension d_L varies according to the network structure, would enable observing a similar effect for $\widetilde{\text{UB}}(L)$ as well. Our focus on the binary classification case is motivated by its analytic tractability, and we leave further extensions for future work.

We proceed to evaluate the Wasserstein generalization bound under the considered setting.

Proposition 5 (Wasserstein distance based bound evaluation). *Under the binary Gaussian classification setting and from Theorem 2, we have*

$$\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y}) \leq \frac{4\sqrt{2}\sigma_0(\sqrt{d_0} + (\sqrt{n} - \sqrt{n-1}))}{\sqrt{n}} \min_{l=0,\dots,L} \mathbb{E} \left[\left(1 \vee \prod_{j=l+1}^L \|\mathbf{W}_j\|_{\text{op}}^2 \right) \|\mathbf{W}_{\otimes l}\|_{\text{F}}^2 \right]^{\frac{1}{2}},$$

where $\mathbf{W}_{\otimes l} = \mathbf{W}_l \mathbf{W}_{l-1} \cdots \mathbf{W}_1$ for $l \in [L]$, and $\mathbf{W}_{\otimes 0} = \mathbf{I}_{d_0}$.

This upper bound is computed using the 2-Wasserstein distance between two Gaussian distributions based on the fact that $\mathbf{W}_1 \leq \mathbf{W}_2$. Note that this upper bound also vanishes as $n \rightarrow \infty$. In this case, the *generalization funnel* layer that yields the tightest upper bound depends on the Frobenius norm of the product of network weight matrices up to the current layer $\|\mathbf{W}_{\otimes l}\|_{\text{F}}$ and the product of subsequent layers' operator norms. We notice that $\|\mathbf{W}_{\otimes l}\|_{\text{F}} = \sqrt{\text{tr}(\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^{\text{T}})}$ not only depends on $\text{rank}(\mathbf{W}_{\otimes l})$ but also on the singular values of $\mathbf{W}_{\otimes l}$. Thus, the generalization funnel layer is not necessarily the last one. In the following example, by considering a simple neural network model with different training methods, we empirically show that the generalization funnel layer depends on the training method. It implies that there indeed exists a network layer that plays a more important role in controlling the generalization performance of the learning algorithm.

Example 1 (Numerical evaluation of Proposition 5). *Consider a DNN with $L = 10$ layers, each of width 2, i.e., $d_0 = d_1 = \cdots = d_{L-1} = 2$. Let the training dataset D_n contain $n = 100$ samples that are i.i.d. from the aforementioned binary Gaussian mixture with $\mu_0 = (0.5 \ 0)^{\text{T}}$ and $\sigma_0 = 1$. We consider a learning algorithm $P_{\mathbf{W}|D_n}$ that proceeds as follows: randomly generate (2×2) rotation matrices $\mathbf{W}_1, \dots, \mathbf{W}_{L-1}$ and the vector \mathbf{W}_L such that $\mathbf{W}_{\otimes L}^{\text{T}} = \frac{1}{n} \sum_{i=1}^n Y_i X_i$. Specifically, let $\mathbf{W}_l = C_l \begin{pmatrix} \cos \theta_l & \sin \theta_l \\ -\sin \theta_l & \cos \theta_l \end{pmatrix}$, for $l = 1, \dots, L-1$, be the rotation matrix, and set the last layer parameter vector as $\mathbf{W}_L = (0, C_L)$. The rotation angles θ_l and scaling factors C_l are then calibrated based on D_n to yield the desired distribution, i.e., $\mathbf{W}_{\otimes L} = \frac{1}{n} \sum_{i=1}^n Y_i X_i$. To that end, we let $\theta_l = \frac{1}{L-1} \arccos \langle \mathbf{W}_L, \frac{1}{n} \sum_{i=1}^n Y_i X_i \rangle$ for $l = 1, \dots, L-1$, while randomly generate $\{C_l\}_{l=1}^L$ such that $\prod_{l=1}^L C_l = \|\frac{1}{n} \sum_{i=1}^n Y_i X_i\|$.*

Under this algorithm, we have that $\mathbf{W}_{\otimes l}$ is full-rank, $\|\mathbf{W}_{\otimes l}\|_{\text{F}} = \sqrt{2} \prod_{j=1}^l C_j$ for $l = 1, \dots, L-1$, $\|\mathbf{W}_{\otimes L}\|_{\text{F}} = \prod_{l=1}^L C_l = \|\frac{1}{n} \sum_{i=1}^n Y_i X_i\|$ and $\|\mathbf{W}_l\|_{\text{op}} = C_l$ for $l = 1, \dots, L$. Given any training dataset D_n , the generalization funnel layer is determined by the way we generate $\{C_l\}_{l=1}^L$. We first let $\prod_{j=l+1}^L C_j \leq 1$ for all $l = 0, 1, \dots, L-1$. If we pick any $1 \leq l' \leq L-1$ and let $\prod_{j=1}^{l'} C_j$ be sufficiently small (at least smaller than $\sqrt{d_0} \wedge \|\frac{1}{n} \sum_{i=1}^n Y_i X_i\|$), then the generalization funnel layer is equal to l' . For numerical illustration, we generate L numbers from the uniform distribution $\text{Unif}\{[0, 1]\}$ and scale them to be C_l 's such that $\prod_{l=1}^L C_l = \|\frac{1}{n} \sum_{i=1}^n Y_i X_i\|$ and $\prod_{j=1}^{l'} C_j = 0.2 \|\frac{1}{n} \sum_{i=1}^n Y_i X_i\|$, for $l' \in \{3, 5, 7\}$. We compute the generalization funnel layer index as the minimizer of the sample mean from 10^4 output network parameters \mathbf{W} (trained on 100 datasets D_n): $l^ = \arg \min_{l=0,1,\dots,L} \text{SampleMean}((1 \vee \prod_{j=l+1}^L \|\mathbf{W}_j\|^2) \|\mathbf{W}_{\otimes l}\|_{\text{F}}^2) = \arg \min_{l=0,1,\dots,L} \text{SampleMean}(\|\mathbf{W}_{\otimes l}\|_{\text{F}}^2)$. As shown in Table I, the generalization funnel layer varies according to the parameter generating methods.*

IV. TIGHTER GENERALIZATION BOUND VIA CONTRACTION

Inspired by Theorem 1, we next aim to capture the contraction of the information measures in our bound as the layer count grows. To that end, we use the *strong* DPI (SDPI) [58]–[60], for which some preliminaries are needed.

A. Strong Data Processing Inequality

Given $P_X, Q_X \in \mathcal{P}(\mathcal{X})$ and a transition kernel (channel) $P_{Y|X}$, write $P_Y = P_{Y|X} \circ P_X$ and $Q_Y = P_{Y|X} \circ Q_X$ for the marginal distributions at the output of the channel when we feed it with P_X or Q_X , respectively. Assuming $P_X \ll Q_X$ and




Fig. 3: Examples of DNNs with stochasticity. (a) l^{th} layer with Dropout probability δ_l . (b) l^{th} layer with DropConnect probabilities $\{\delta_{l,i,j}\}_{j=1}^{d_{l-1}}$. (c) Noisy DNN with injected isotropic Gaussian noise to the l^{th} layer.

that Q_X is not a point mass, the SDPI coefficient for $P_{Y|X}$ under the f -divergence D_f is

$$\eta_f(P_{Y|X}) := \sup_{P_X, Q_X} \frac{D_f(P_{Y|X} \circ P_X \| P_{Y|X} \circ Q_X)}{D_f(P_X \| Q_X)} \in [0, 1].$$

We write $\eta_{\text{KL}}(P_{Y|X})$ and $\eta_{\text{TV}}(P_{Y|X})$ for the coefficients under the KL divergence and the total variation distance, respectively. Proposition II.4.10 in [61] shows that for any f -divergence, we have

$$\eta_f(P_{Y|X}) \leq \eta_{\text{TV}}(P_{Y|X}) = \sup_{x, x' \in \mathcal{X}} \|P_{Y|X=x} - P_{Y|X=x'}\|_{\text{TV}},$$

where $\eta_{\text{TV}}(P_{Y|X})$ is also known as *Dobrushin's coefficient* [59]. It can be shown that if $Y = g(X)$ for some deterministic function $g : \mathcal{X} \rightarrow \mathcal{Y}$, then $\eta_f(P_{g(X)|X}) = \eta_{\text{TV}}(P_{g(X)|X}) = 1$ (cf., e.g., Proposition II.4.12 from [61]).

According to the above, if all the feature maps $g_{\mathbf{w}_l}$, for $l = 1, \dots, L$, in the DNN are deterministic, the contraction coefficients we are looking for degenerate to 1, landing us back at the bound from Theorem 1. To arrive at a nontrivial contraction coefficient, we consider the following common regularization techniques in machine learning: Dropout [33], DropConnect [34], and Gaussian noise injection [35], [37]–[39]. These methods have been used to introduce stochasticity to neural networks to enhance generalization and robustness.

Before presenting the subsequent results, we find it succinct to define the notations: for any vector V , let $V(i)$ be i^{th} element of V ; for any matrix \mathbf{M} , let $\mathbf{M}(i, j)$ be the $(i, j)^{\text{th}}$ element of \mathbf{M} .

B. Generalization Bound for DNNs with Dropout

The Dropout [33] technique has been shown to be a simple way to prevent neural networks from overfitting. We hereby analyse the contraction behaviour in the generalization bounds caused by Dropout. Consider the l^{th} layer of the neural network that applies a Dropout probability $\delta_l \in (0, 1)$, $l = 0, \dots, L - 1$, as shown in Figure 3a. Let $E_l \sim \text{Bern}(1 - \delta_l)^{\otimes d_l}$ be an i.i.d. sequence of d_l Bernoulli variables. Then the activation output of the $(l + 1)^{\text{th}}$ layer is given by

$$T_{l+1} = \phi_{l+1}(\mathbf{W}_{l+1}(T_l \odot E_l)) =: \phi_{l+1}(\mathbf{W}_{l+1}\tilde{T}_l), \quad (3)$$

where \odot denotes the elementwise product operation. When $T_l(i)$ is not equal to 0, with probability δ_l , $T_l(i)$ is deactivated as 0 before being passed to the next layer; with probability $1 - \delta_l$, $T_l(i)$ retains its value. When $T_l(i) = 0$, it always remains unchanged. The channel $P_{\tilde{T}_l|T_l}$ can then be regarded as a composition of d_l parallel independent generalized Z-channels [62]. Thus, the stochastic channel $P_{T_{l+1}|T_l, \mathbf{w}} = P_{T_{l+1}|\tilde{T}_l, \mathbf{w}} \circ P_{\tilde{T}_l|T_l}$ yields a non-trivial SDPI coefficient.

Lemma 6 (Dropout SDPI coefficient). *The SDPI coefficient of the Dropout channel $P_{\tilde{T}_l|T_l}$ with Dropout probability $\delta_l \in (0, 1)$ is $\eta_{\text{KL}}(P_{\tilde{T}_l|T_l}) = 1 - \delta_l^{d_l}$, for $l = 0, \dots, L$.*

Lemma 6, which is proven in Appendix F, implies that whenever Dropout is applied to a certain layer, it leads to a strict contraction in the generalization bound in Theorem 1 since the coefficient $\eta_{\text{KL}}(P_{\tilde{T}_l|T_l})$ is strictly less than 1. Consequently, the layer mappings in a DNN with Dropout gives rise to the following result.

Theorem 7 (DNN with Dropout generalization bound). *Consider the DNN model with Dropout probability $\delta_l \in (0, 1)$ at the l^{th} layer (cf. (3)), $l = 0, \dots, L - 1$. If the loss function $\ell(\mathbf{w}, X, Y)$ is σ -sub-Gaussian under $P_{X,Y}$, for all $\mathbf{w} \in \mathcal{W}$, we have*

$$|\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y})| \leq \frac{\sigma\sqrt{2}}{n} \sum_{i=1}^n \sqrt{\prod_{l=0}^{L-1} (1 - \delta_l^{d_l}) \mathcal{I}(X_i; \mathbf{W}|Y_i) + \mathcal{I}(Y_i; \mathbf{W})}.$$

The proof of Theorem 7 initiates at the bound $\text{UB}(L)$ from Theorem 1 and first factors out the terms that depend on the label Y from the KL divergence. This yields the summand $\mathcal{I}(Y_i; \mathbf{W})$. This step is necessary since the label is not processed by the DNN with Dropout, and the corresponding SDPI coefficient (without factoring Y_i out) would degenerate to 1. For the

remaining term, we invoke the SDPI L times collecting the coefficients and invoking Lemma 6 to arrive at the desired bound. The details are provided in Appendix G.

We make the following observations regarding the bound from Theorem 7. The coefficient product $\prod_{l=0}^{L-1} (1 - \delta_l^{d_l})$ exhibits a decrement from 1 to 0 with: (i) increasing Dropout probabilities $\{\delta_l\}_{l=0}^{L-1}$; (ii) elevated network depth L ; and (iii) shrinking layer widths $\{d_l\}_{l=0}^{L-1}$. In addition, using Taylor's expansion, we have $\log(1 - \delta_l^{d_l}) \approx -\delta_l^{d_l}$ and $\delta_l^{d_l} = (1 - (1 - \delta_l))^{d_l} \approx 1 - d_l(1 - \delta_l)$. Then the product of coefficient can be approximated as

$$\prod_{l=0}^{L-1} (1 - \delta_l^{d_l}) = \exp\left(\sum_{l=0}^{L-1} \log(1 - \delta_l^{d_l})\right) \approx \exp\left(-\sum_{l=0}^{L-1} \delta_l^{d_l}\right) \approx \exp\left(-\sum_{l=0}^{L-1} (1 - d_l(1 - \delta_l))\right),$$

highlighting an exponential decay. Since the label follows a categorical distribution of parameter K , we have $I(Y_i; \mathbf{W}) \leq \log K$. The smaller the number of distinct labels K , the better the generalization bound. The behavior of $I(X_i; \mathbf{W}|Y_i)$, on the other hand, is harder to pin down as it depends on the data distribution and the learning algorithm at hand. In the following statement, we investigate the behavior of $I(X_i, Y_i; \mathbf{W})$ w.r.t. the input layer Dropout probability δ_0 . At the end of Section IV, we investigate the generalization bound w.r.t. the layer dimension and network depth with an instance of DNN with a finite parameter space.

Lemma 8 (Upper bound on $I(X_i, Y_i; \mathbf{W})$ in DNNs with Dropout). *For a DNN model with Dropout probability $\delta_0 \in (0, 1)$ at the input layer, we have*

$$I(X_i, Y_i; \mathbf{W}) \leq \text{MIUB}_i(\delta_0) := \sum_{k=0}^{d_0} \delta_0^{d_0-k} (1 - \delta_0)^k \binom{d_0}{k} I_k^{(i)}, \quad \forall i = 1, \dots, n,$$

where $I_k^{(i)} = \max_{\mathcal{J}(k) \subseteq [d_i], |\mathcal{J}(k)|=k} I((X_i(j))_{j \in \mathcal{J}(k)}, Y_i; \mathbf{W})$, for $k = 0, \dots, d_0$. The upper bound $\text{MIUB}_i(\delta_0)$ monotonically shrinks to 0 as δ_0 increases from 0 to 1.

The proof of Lemma 8, fully provided in Appendix H, hinges on the application of the chain rule of mutual information. Lemma 8 suggests that the generalization bound (c.f. UB(0) in Theorem 1) shrinks even when Dropout is only applied to the input data, commonly employed as a data augmentation technique. Since $I(X_i; \mathbf{W}|Y_i), I(Y_i; \mathbf{W}) \leq I(X_i, Y_i; \mathbf{W})$, Theorem 7 and Lemma 8 show that the generalization bound of a DNN with Dropout diminishes as the Dropout probabilities increase. This aligns with the intuition that introducing stochasticity, as facilitated by Dropout mechanisms, enhances the model's capacity for generalization.

C. Generalization Bound for DNNs with DropConnect

Compared to Dropout which effectively deactivates some columns of the network weight matrices, DropConnect [34] operates in a more precise way by randomly deactivating each element of the weight matrices. Consider the l^{th} layer of the neural network adopts a DropConnect probability $\delta_{l,i,j} \in (0, 1)$ at the $(i, j)^{\text{th}}$ entry of the weight matrix \mathbf{W}_l , $l = 1, \dots, L$, as shown in Figure 3b. Let $\mathbf{E}_l \sim \prod_{i=1}^{d_l} \prod_{j=1}^{d_{l-1}} \text{Bern}(1 - \delta_{l,i,j})$ be a $d_l \times d_{l-1}$ binary matrix with i.i.d. Bernoulli elements. Then the activation output of the l^{th} layer is given by

$$T_l = \phi_l((\mathbf{W}_l \odot \mathbf{E}_l)T_{l-1}) =: \left[\phi_l\left(\sum_{j=1}^{d_{l-1}} \mathbf{W}_l(1, j)\tilde{T}_{l-1}(1, j)\right), \dots, \phi_l\left(\sum_{j=1}^{d_{l-1}} \mathbf{W}_l(d_l, j)\tilde{T}_{l-1}(d_l, j)\right) \right]^T, \quad (4)$$

where \odot denotes the elementwise product operation, \tilde{T}_{l-1} is a $d_l \times d_{l-1}$ matrix, and $\tilde{T}_{l-1}(i, j) := \mathbf{E}_l(i, j)T_{l-1}(j)$. When $\mathbf{W}_l(i, j)T_{l-1}(j)$ is not equal to 0, with probability $\delta_{l,i,j}$, $\mathbf{W}_l(i, j)T_{l-1}(j)$ is deactivated as 0 before being passed to the i^{th} neuron of the next layer; with probability $1 - \delta_{l,i,j}$, $\mathbf{W}_l(i, j)T_{l-1}(j)$ keeps its value. When $\mathbf{W}_l(i, j)T_{l-1}(j) = 0$, it always remains unchanged. In a similar manner to Section IV-B, the channel $P_{\tilde{T}_{l-1}|T_{l-1}}$ can then be regarded as a composition of $d_l d_{l-1}$ parallel independent generalized Z-channels. Thus, the stochastic channel $P_{T_l|T_{l-1}, \mathbf{W}} = P_{T_l|\tilde{T}_{l-1}, \mathbf{W}} \otimes P_{\tilde{T}_{l-1}|T_{l-1}}$ also yields a non-trivial SDPI coefficient.

Lemma 9 (DropConnect SDPI coefficient bound). *The SDPI coefficient of the DropConnect channel $P_{\tilde{T}_{l-1}|T_{l-1}}$ with DropConnect probabilities $\delta_{l,i,j} \in (0, 1)$, $i = 1, \dots, d_l, j = 1, \dots, d_{l-1}$, satisfies $\eta_{\text{KL}}(P_{\tilde{T}_{l-1}|T_{l-1}}) \leq 1 - \prod_{i=1}^{d_l} \prod_{j=1}^{d_{l-1}} \delta_{l,i,j}$, for $l = 1, \dots, L$.*

Lemma 9, which is proven in Appendix I, implies that the application of DropConnect at a certain layer also leads to a strict contraction in the generalization bound in Theorem 1 since the coefficient $\eta_{\text{KL}}(P_{\tilde{T}_{l-1}|T_{l-1}}) < 1$. Compared to the coefficient shown in Lemma 6, this coefficient decreases with an increase in individual probability $\delta_{l,i,j}$. This observation highlights the advantageous flexibility of DropConnect in improving generalization performance. Consequently, the layer mappings in a DNN with DropConnect gives rise to the following non-trivial result.

Theorem 10 (DNN with DropConnect generalization bound). *Consider the DNN model with DropConnect probabilities $\delta_{l,i,j} \in (0,1)$, $i = 1, \dots, d_l, j = 1, \dots, d_{l-1}$ at the l^{th} layer (cf. (4)), $l = 1, \dots, L$. If the loss function $\ell(\mathbf{w}, X, Y)$ is σ -sub-Gaussian under $P_{X,Y}$, for all $\mathbf{w} \in \mathcal{W}$, we have*

$$|\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y})| \leq \frac{\sigma\sqrt{2}}{n} \sum_{i=1}^n \sqrt{\prod_{l=1}^L \left(1 - \prod_{i=1}^{d_l} \prod_{j=1}^{d_{l-1}} \delta_{l,i,j}\right)} \mathsf{I}(X_i; \mathbf{W}|Y_i) + \mathsf{I}(Y_i; \mathbf{W}).$$

The proof of Theorem 10 can be derived following the same procedures in that of Theorem 7 and is thus omitted. We observe that the product of coefficients also decrease from 1 to 0 as the DropConnect probabilities increase, the layer dimensions shrink, or the number of layers increases. An exponential decay similar to that discussed after Theorem 7 also occurs in this case. By combining DropConnect with Dropout at the input layer X_i , a contraction is enforced upon the mutual information terms $(\mathsf{I}(X_i; \mathbf{W}|Y_i), \mathsf{I}(Y_i; \mathbf{W}))$, resulting in a generalization bound that decays with the increase of dropout probabilities.

D. Generalization Bound for Noisy DNNs

Injecting Gaussian noise to neuron outputs is another form of DNN regularization that has been explored in various research studies [35], [37]–[39]. The idea is to add random Gaussian noise to the input, the network parameters or the internal representation during training. This can help prevent overfitting and improve the model’s generalization by introducing a level of uncertainty and robustness [37]. In this section, to arrive at nontrivial SDPI contraction coefficient we consider a noisy DNN model where the feature map at each layer is perturbed by isotropic Gaussian noise, i.e.,

$$\tilde{T}_l = g_{\mathbf{W}_l}(\tilde{T}_{l-1}) + \epsilon_l Z_l, \quad l = 1, \dots, L, \quad (5)$$

where $\tilde{T}_0 = X$, $Z_l \sim N(0, \mathbf{I}_{d_l})$ is independent of the input and $\epsilon_l \in \mathbb{R}_{>0}$ is a constant. The illustration is shown in Figure 3c. Such noisy DNNs were explored in [35] and were shown to serve as good approximations of classical (deterministic) networks.

To analyze generalization error under the noisy DNN model, we present the following lemma that bounds the SDPI coefficient for the aforementioned channel.

Lemma 11 (Noisy DNN SDPI coefficient bound). *Let $X \sim P_X \in \mathcal{P}(\mathbb{R}^{d_x})$ and consider a bounded function $g : \mathbb{R}^{d_x} \rightarrow \mathbb{R}^{d_y}$. Set $Y = g(X) + \epsilon N$, where $\epsilon > 0$ and $N \sim N(0, \mathbf{I}_{d_y})$ is independent of X . The SDPI coefficient of the induced channel $P_{Y|X}$ satisfies*

$$\eta_f(P_{Y|X}) \leq \eta_{\text{TV}}(P_{Y|X}) \leq 1 - 2\mathsf{Q}\left(\frac{\sqrt{2d_y}\|g\|_\infty}{2\epsilon}\right),$$

where $\mathsf{Q}(x) := \int_x^\infty \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt$ is the Gaussian complimentary cumulative distribution function.

Lemma 11, which is proven in Appendix J, implies that whenever $\sqrt{d_y}\|g\|_\infty \epsilon^{-1} > 0$, we have $\eta_{\text{KL}}(P_{Y|X}) < 1$. Consequently, the layer mappings in a noisy DNN with bounded activations present non-trivial contraction, which gives rise to the following result.

Theorem 12 (Noisy DNN generalization bound). *Consider the noisy DNN model from (5) with bounded activation functions $\phi_l, l = 1, \dots, L$. If the loss function $\ell(\mathbf{w}, X, Y)$ is σ -sub-Gaussian under $P_{X,Y}$, for all $\mathbf{w} \in \mathcal{W}$, we have*

$$|\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y})| \leq \frac{\sigma\sqrt{2}}{n} \sum_{i=1}^n \sqrt{\prod_{l=1}^L \left(1 - 2\mathsf{Q}\left(\frac{\sqrt{2d_l}\|\phi_l\|_\infty}{2\epsilon_l}\right)\right)} \mathsf{I}(X_i; \mathbf{W}|Y_i) + \mathsf{I}(Y_i; \mathbf{W}).$$

The proof of Theorem 12 is provided in Appendix K, which follows the similar procedures as in the proof of Theorem 7. The term $\mathsf{I}(Y_i; \mathbf{W})$ is factored out since the label is not processed by the noisy DNN and upper bounded by $\log K$. The SDPI is invoked L times to arrive at the product of coefficient for the remaining term. We also note that $\|\phi_l\|_\infty$ is typically small, e.g., $\|\phi_l\|_\infty = 1$ if $\phi_l \in \{\text{sigmoid}, \text{softmax}, \text{tanh}\}$. From Theorem 12, we observe that for fixed noise parameters $\epsilon_1, \dots, \epsilon_L$, the coefficient product decreases from 1 to 0 as the layer dimensions $\{d_l\}_{l=1}^L$ shrink and the number of layers L grows. If $\min_{l \in [L]} (1 - 2\mathsf{Q}(\frac{\sqrt{2d_l}\|\phi_l\|_\infty}{2\epsilon_l})) \leq q \in (0, 1)$ (i.e., $\max_{l \in [L]} d_l \leq (\frac{2\epsilon_l \mathsf{Q}^{-1}(\frac{1-q}{2})}{\|\phi_l\|_\infty})^2$), the coefficient is less than $\exp(-L \log \frac{1}{q})$, which decays exponentially in L . Furthermore, by increasing the noise level ϵ_l ’s, the coefficient product shrinks as well. The observations are consistent with those of Theorems 7 and 10.

To analyze the behavior of $\mathsf{I}(X_i; \mathbf{W}|Y_i)$ and generalization bounds in the aforementioned theorems concerning layer dimensions and network depth, we examine the following tractable instance of DNNs.

Example 2 (Numerical evaluation of tightened generalization bounds). *Consider the DNN parameter space is constrained to be finite, e.g., $\mathcal{W} = [B]^{d_1 \times d_0} \times \dots \times [B]^{d_L \times d_{L-1}}$ for some $B \in \mathbb{N}$. We can upper bound the mutual information by $\mathsf{I}(X_i; \mathbf{W}|Y_i) \leq \mathsf{H}(\mathbf{W}) \leq \sum_{l=1}^L d_l d_{l-1} \log B$, which increases as d_l and L grow. We consider the following two cases of increasing the depth of an L -layer neural net with discrete parameters as above.*




Fig. 4: Examples of tightened generalization bounds for two types of NNs with stochasticity by adding an extra hidden layer.




Fig. 5: Examples of tightened generalization bounds for two types of NNs with stochasticity by dividing one hidden layer into two separate hidden layers.

1) Add an extra hidden layer with dimension d_{l^*} . We compare a 2-layer neural net with layer dimensions $\{d_0 = 10, d_1 = 20, d_2 = 2\}$ to a 3-layer neural net with layer dimensions $\{d_0 = 10, d_1 = d_{l^*}, d_2 = 20, d_3 = 2\}$. Figures 4a and 4b plot the said mutual information bounds (ignoring $\sigma\sqrt{2}$ without loss of generality) from Theorems 7 and 12, respectively. The tightened bounds with the contraction coefficient behave similarly for both the NN with Dropout and NN with Gaussian noise. We observe that if d_{l^*} is sufficiently small, the generalization bound shrinks as a result of the added layer.

2) Divide the dimension of one hidden layer into two separate hidden layers. We compare a 2-layer neural net with layer dimensions $\{d_0 = 10, d_1 = 30, d_2 = 2\}$ to a 3-layer neural net with layer dimensions $\{d_0 = 10, d_1 = d, d_2 = 30 - d, d_3 = 2\}$. By varying $d \in [1, 29]$, Figures 5a and 5b plot the said mutual information bounds (ignoring $\sigma\sqrt{2}$ without loss of generality) from Theorems 7 and 12, respectively. The tightened bounds with the contraction coefficient still behave similarly for both the NN with Dropout and NN with Gaussian noise. We observe that if one of the splitted layer dimensions, i.e., d or $30 - d$ in this example, is sufficiently small, the generalization bound shrinks as a result of the layer division.

This example suggests that a deep but narrower network may generalize better. We observe that when d_{l^*} or d is equal to 1, the generalization bound is minimized. This phenomenon is due to the fact that we are taking a worst-case upper bound on $H(\mathbf{W})$. Thus, to draw more compelling conclusions, it is necessary to conduct thorough analyses and experiments for general algorithms/architectures. Next, we prove that the implications from this example can be generalized to the Gibbs algorithm.

E. Applications to Gibbs Algorithm

We further characterize the tightened generalization bounds in Theorems 7, 10 and 12, for DNNs with Dropout, DropConnect and injected Gaussian noise, under the Gibbs algorithm. The Gibbs algorithm is a tractable and idealized model for learning algorithms with various regularization approaches, e.g., stochastic optimization methods or relative entropy regularization [40].

This allows us to investigate the concrete impacts of the DNN structure parameters on the mutual information terms in the generalization bounds.

Given any training dataset $D_n = s$, consider the $(\alpha, \pi(\mathbf{W}), \mathcal{L}_E(\mathbf{W}, s))$ -Gibbs algorithm [41]

$$P_{\mathbf{W}|D_n}^\alpha(\mathbf{W}|s) := \frac{\pi(\mathbf{W}) \exp(-\alpha \mathcal{L}_E(\mathbf{W}, s))}{\Lambda_\alpha(s)},$$

where $\alpha > 0$ denotes the inverse temperature, $\pi \in \mathcal{P}(\mathcal{W})$ is an arbitrary prior distribution on \mathbf{W} and $\Lambda_\alpha(s) = \mathbb{E}[\exp(-\alpha \mathcal{L}_E(\mathbf{W}, s))]$ is the partition function. The $(\alpha, \pi(\mathbf{W}), \mathcal{L}_E(\mathbf{W}, s))$ -Gibbs algorithm can be viewed as a randomized version of empirical risk minimization (ERM). As the inverse temperature $\alpha \rightarrow \infty$, then network parameters generated by the Gibbs algorithm converges to those from the standard ERM. In [40], it is proven that under some conditions on the loss function, the posterior $P_{\mathbf{W}|D_n}^\alpha$ induced by Stochastic Gradient Langevin Dynamics (SGLD) algorithm [63] is close to the Gibbs posterior $P_{\mathbf{W}|D_n}^\alpha$ in 2-Wasserstein distance for sufficiently large iterations.

When DNNs with non-trivial SDPI contraction coefficients (c.f. Lemmas 6, 9 and 11) are trained with the Gibbs algorithm using a loss function bounded within $[0, 1]$, we obtain the distribution-free bound that decreases as the number of network layers increases.

Proposition 13 (Tightened generalization bound for Gibbs algorithm). *Consider that the DNN model from Eq. (3) with Dropout probability $\delta_l \in (0, 1)$, Eq. (4) with DropConnect probability $\delta_{l,i,j} \in (0, 1)$, or Eq. (5) with bounded activation functions ϕ_l , $l = 1, \dots, L$, is trained by the $(\alpha, \pi(\mathbf{W}), \mathcal{L}_E(\mathbf{W}, D_n))$ -Gibbs algorithm with any loss function $\ell : \mathcal{W} \times \mathcal{X} \times \mathcal{Y} \rightarrow [0, 1]$. The SDPI contraction coefficients for these DNN models are denoted by $\tilde{\eta}_{\text{KL}}(l) := 1 - \delta_{l-1}^{d_l-1}$, $\tilde{\eta}_{\text{KL}}(l) := 1 - \prod_{i=1}^{d_l} \prod_{j=1}^{d_{l-1}} \delta_{l,i,j}$ and $\tilde{\eta}_{\text{KL}}(l) := 1 - 2\mathcal{Q}(\frac{\sqrt{2d_l} \|\phi_l\|_\infty}{2\epsilon_l})$, respectively. There exists a constant $\gamma \in [0, 1]$ such that*

$$|\text{gen}(P_{\mathbf{W}|D_n}^\alpha, P_{X,Y})| \leq \frac{\alpha}{4n} \sqrt{\gamma \prod_{l=1}^L \tilde{\eta}_{\text{KL}}(l) + 1 - \gamma}.$$

Proposition 13, which is obtained by combining [64, Theorem 2] and Theorems 7, 10 and 12, is tighter than [64, Theorem 2]. Compared to the exact characterization of generalization error in [41, Theorem 1], which did not explicitly show the dependence on the neural net structure, Proposition 13 shows that the bound is of order $\mathcal{O}(\frac{1}{n})$ and shrinks with increasing L and decreasing d_l due to the SDPI coefficient $\tilde{\eta}_{\text{KL}}(l) \in (0, 1)$. As per the analyses presented in Sections IV-B–IV-D, the generalization bound for the Gibbs algorithm approximately exhibits an exponential decay with increasing network depth L and decreasing layer dimension d_l . This observation suggests the benefit of deeper network depth and narrower layer dimensions when the DNN model is trained using regularization techniques, such as Dropout, DropConnect and noise injection, in conjunction with the Gibbs algorithm. Some other studies have also shown the generalization capability of deep and narrow neural networks [18], [19], [43], [65], [66].

Proof of Proposition 13. For any loss function $\ell : \mathcal{W} \times \mathcal{X} \times \mathcal{Y} \rightarrow [0, 1]$, $\ell(\mathbf{w}, X, Y)$ is $\frac{1}{2}$ -sub-Gaussian under $P_{X,Y}$ for any $\mathbf{w} \in \mathcal{W}$. Let $D_n^{-i} = D_n \setminus (X_i, Y_i)$ for any $i \in [n]$. The Gibbs algorithm $P_{\mathbf{W}|D_n}^\alpha$ with a loss function $\ell \in [0, 1]$ is $(\frac{\alpha^2}{8n^2}, P_{X,Y})$ -stable in mutual information (cf. [64, Definition 1, Theorem 4]), i.e., $\frac{1}{n} \sum_{i=1}^n I(\mathbf{W}; X_i, Y_i | D_n^{-i}) \leq \frac{\alpha^2}{8n^2}$. Since (X_i, Y_i) is independent of D_n^{-i} , we have $\frac{1}{n} \sum_{i=1}^n I(\mathbf{W}; X_i, Y_i) \leq \frac{1}{n} \sum_{i=1}^n I(\mathbf{W}; X_i, Y_i | D_n^{-i}) \leq \frac{\alpha^2}{8n^2}$. There exists an independent constant $\gamma \in [0, 1]$ such that $\frac{1}{n} \sum_{i=1}^n I(\mathbf{W}; X_i | Y_i) \leq \frac{\gamma \alpha^2}{8n^2}$ and $\frac{1}{n} \sum_{i=1}^n I(\mathbf{W}; Y_i) \leq \frac{(1-\gamma)\alpha^2}{8n^2}$. By further applying the Jensen's inequality to Theorem 7, 10 or 12, we obtain the bound. \square

V. CONCLUDING REMARKS

This work set to quantify the generalization error of DNNs within the framework of information theory. We derived two hierarchical generalization bounds that capture the effect of depth through the internal representations of the corresponding layers. The two bounds compare the distributions of internal representations of the training and test data under (i) the KL divergence, and (ii) the 1-Wasserstein distance. The KL divergence bound diminishes as the layer index increases, indicating the advantage of deep network architectures. The Wasserstein bound is minimized by the so-called *generalization funnel layer*, providing new a insight that certain layers play a more prominent role than others in governing generalization performance. We instantiated these results to a binary Gaussian mixture classification task with linear DNNs. Simple analytic expressions for the two generalization bounds we obtained, with the KL divergence reducing to depend on (and shrinks with) the rank of the product of weight matrices, while the Wasserstein bound simplified to depend on the Frobenius norm of the weight matrix product. The latter further implied that the generalization funnel layer of a given model varies with different training methods.

Next, we set to quantify the contraction of the KL divergence bound as a function of depth. To that end, we analyze the SDPI coefficient between consequent layers in three regularized DNN models: Dropout, DropConnect, and independent Gaussian noise injection (stochasticity is necessary to avoid a degenerate SDPI coefficient). This enables refining our KL divergence generalization bound to capture the contraction through the product of SDPI coefficient associated with different layers, which

diminishes from 1 to 0 as the depth L increases or the per-layer width d_l shrinks. It was further demonstrated that the rate of decay is approximately exponential and that the SDPI coefficient decays with an increase of network stochasticity, such as higher Dropout/DropConnect probabilities and higher injected noise level. Numerical evaluation of our bounds are provided for a simple DNN model, where the parameter space is finite. The simulations suggest that, in this example, deeper yet narrower network architectures generalize better. While the extent to which this statement applies generally is unclear, we provide a proof that it is true when learning is performed via the Gibbs algorithm. Overall, our results provide an information-theoretic perspective on generalization of deep models, encompassing quantitative hierarchical bounds to insights into architectures that may generalize better.

Several appealing future research directions extend from this work. First, we can generalize our analyses to other neural network architectures such as convolution layer, pooling layer, residual connection, attention layer and so on. It would be interesting to investigate the effects of the properties of different layer types on generalization performance. Second, tightening SDPI coefficients would be an interesting direction. If we have some prior knowledge about the input distributions, can we improve the SDPI coefficient from the current worst-case one? It would also be insightful to leverage non-linear SDPI [60] to obtain contraction curve for deterministic neural networks and a wider range of activation functions. Third, the behaviour of the mutual information term is still worth investigating. Even though the SDPI coefficient product shrinks with deeper and narrower neural network, the mutual information $I(X_i, Y_i; \mathbf{W})$ may increase or decrease according to various DNN architectures and training algorithms. We can set out from the stability perspective. By showing the dependence of the stability parameters on DNN architecture, we can make similar analysis to that of the Gibbs algorithm. We can also extend the assumption of finite parameter space to the setting in the two-stage algorithm [20, Section 4.2], where the parameter space is countable w.r.t. the training dataset in the first stage. The mutual information will be upper bounded by a VC dimension, which can be further quantified by DNN architecture. Finally, since generalization error tracks the gap between test and training losses, for non-empirical-risk-minimization algorithms, we need to consider excess risk to better understand the performance of the algorithms. It would be interesting to extend our established tools to bounding excess risk. Through these analyses, we will have a clearer view on the impact of DNN architectures on generalization capability. Moving forward, it would be valuable to conduct thorough analyses and experiments for general algorithms and network structures to draw more compelling conclusions on the impact of network depth and width and find out the generalization funnel layer.

REFERENCES

- [1] N. Golowich, A. Rakhlin, and O. Shamir, “Size-independent sample complexity of neural networks,” in *Conference On Learning Theory*. PMLR, 2018, pp. 297–299.
- [2] B. Neyshabur, S. Bhojanapalli, D. McAllester, and N. Srebro, “Exploring generalization in deep learning,” in *Advances in Neural Information Processing Systems*, vol. 30, 2017, pp. 5949–5958.
- [3] T. Liang, T. Poggio, A. Rakhlin, and J. Stokes, “Fisher-Rao metric, geometry, and complexity of neural networks,” in *The 22nd International Conference on Artificial Intelligence and Statistics*. PMLR, 2019, pp. 888–896.
- [4] S. Arora, R. Ge, B. Neyshabur, and Y. Zhang, “Stronger generalization bounds for deep nets via a compression approach,” in *International Conference on Machine Learning*. PMLR, 2018, pp. 254–263.
- [5] G. K. Dziugaite and D. M. Roy, “Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data,” in *Proceedings of the 33rd Conference on Uncertainty in Artificial Intelligence, UAI 2017*. AUAI Press, 2017.
- [6] D. A. McAllester, “Some PAC-Bayesian theorems,” in *Proceedings of the 11th Annual Conference on Computational Learning Theory*, 1998, pp. 230–234.
- [7] —, “PAC-Bayesian model averaging,” in *Proceedings of the 12th Annual Conference on Computational Learning Theory*, 1999, pp. 164–170.
- [8] B. Neyshabur, S. Bhojanapalli, and N. Srebro, “A PAC-Bayesian approach to spectrally-normalized margin bounds for neural networks,” in *International Conference on Learning Representations*, 2018.
- [9] W. Zhou, V. Veitch, M. Austern, R. P. Adams, and P. Orbanz, “Non-vacuous generalization bounds at the imagenet scale: a PAC-Bayesian compression approach,” in *International Conference on Learning Representations*, 2018.
- [10] S. Hochreiter and J. Schmidhuber, “Flat minima,” *Neural Computation*, vol. 9, no. 1, pp. 1–42, 1997.
- [11] L. Dinh, R. Pascanu, S. Bengio, and Y. Bengio, “Sharp minima can generalize for deep nets,” in *International Conference on Machine Learning*. PMLR, 2017, pp. 1019–1028.
- [12] N. S. Keskar, D. Mudigere, J. Nocedal, M. Smelyanskiy, and P. T. P. Tang, “On large-batch training for deep learning: Generalization gap and sharp minima,” in *International Conference on Learning Representations*, 2017.
- [13] L. Wu, Z. Zhu, and W. E, “Towards understanding generalization of deep learning: Perspective of loss landscapes,” *ICML 2017 Workshop on Principled Approaches to Deep Learning, Sydney, Australia*, 2017.
- [14] D. Soudry, E. Hoffer, M. S. Nacson, S. Gunasekar, and N. Srebro, “The implicit bias of gradient descent on separable data,” *The Journal of Machine Learning Research*, vol. 19, no. 1, pp. 2822–2878, 2018.
- [15] S. L. Smith and Q. V. Le, “A Bayesian perspective on generalization and stochastic gradient descent,” in *International Conference on Learning Representations*, 2018.
- [16] S. Chatterjee and P. Zielinski, “On the generalization mystery in deep learning,” *arXiv preprint arXiv:2203.10036*, 2022.
- [17] D. Jakobovitz, R. Giryes, and M. R. Rodrigues, “Generalization error in deep learning,” in *Compressed Sensing and Its Applications: Third International MATHEON Conference 2017*. Springer, 2019, pp. 153–193.
- [18] C. Zhang, S. Bengio, M. Hardt, B. Recht, and O. Vinyals, “Understanding deep learning (still) requires rethinking generalization,” *Communications of the ACM*, vol. 64, no. 3, pp. 107–115, 2021.
- [19] K. Kawaguchi, L. P. Kaelbling, and Y. Bengio, “Generalization in deep learning,” in *Mathematical Aspects of Deep Learning*. Cambridge University Press, 2022.
- [20] A. Xu and M. Raginsky, “Information-theoretic analysis of generalization capability of learning algorithms,” *Advances in Neural Information Processing Systems*, vol. 30, 2017.
- [21] D. Russo and J. Zou, “How much does your data exploration overfit? controlling bias via information usage,” *IEEE Transactions on Information Theory*, vol. 66, no. 1, pp. 302–323, 2019.

- [22] Y. Bu, S. Zou, and V. V. Veeravalli, "Tightening mutual information-based bounds on generalization error," *IEEE Journal on Selected Areas in Information Theory*, vol. 1, no. 1, pp. 121–130, 2020.
- [23] A. Asadi, E. Abbe, and S. Verdú, "Chaining mutual information and tightening generalization bounds," in *Advances in Neural Information Processing Systems*, 2018, pp. 7234–7243.
- [24] E. Clerico, A. Shidani, G. Deligiannidis, and A. Doucet, "Chained generalisation bounds," in *Conference on Learning Theory*. PMLR, 2022, pp. 4212–4257.
- [25] H. Hafez-Kolahi, Z. Golgooni, S. Kasaei, and M. Soleymani, "Conditioning and processing: Techniques to improve information-theoretic generalization bounds," *Advances in Neural Information Processing Systems*, vol. 33, 2020.
- [26] M. Haghifam, J. Negrea, A. Khisti, D. M. Roy, and G. K. Dziugaite, "Sharpened generalization bounds based on conditional mutual information and an application to noisy, iterative algorithms," *Advances in Neural Information Processing Systems*, 2020.
- [27] T. Steinke and L. Zakyntinou, "Reasoning about generalization via conditional mutual information," in *Conference on Learning Theory*. PMLR, 2020, pp. 3437–3452.
- [28] H. Harutyunyan, M. Raginsky, G. Ver Steeg, and A. Galstyan, "Information-theoretic generalization bounds for black-box learning algorithms," *Advances in Neural Information Processing Systems*, vol. 34, pp. 24670–24682, 2021.
- [29] A. R. Esposito, M. Gastpar, and I. Issa, "Generalization error bounds via Rényi-, f-divergences and maximal leakage," *IEEE Transactions on Information Theory*, vol. 67, no. 8, pp. 4986–5004, 2021.
- [30] G. Aminian, S. Masiha, L. Toni, and M. R. Rodrigues, "Learning algorithm generalization error bounds via auxiliary distributions," *arXiv preprint arXiv:2210.00483*, 2022.
- [31] G. Aminian, L. Toni, and M. R. Rodrigues, "Information-theoretic bounds on the moments of the generalization error of learning algorithms," in *IEEE International Symposium on Information Theory (ISIT)*, 2021.
- [32] H. Wang, M. Diaz, J. C. S. Santos Filho, and F. P. Calmon, "An information-theoretic view of generalization via Wasserstein distance," in *2019 IEEE International Symposium on Information Theory (ISIT)*. IEEE, 2019, pp. 577–581.
- [33] N. Srivastava, G. Hinton, A. Krizhevsky, I. Sutskever, and R. Salakhutdinov, "Dropout: A simple way to prevent neural networks from overfitting," *Journal of Machine Learning Research*, vol. 15, no. 56, pp. 1929–1958, 2014.
- [34] L. Wan, M. Zeiler, S. Zhang, Y. Le Cun, and R. Fergus, "Regularization of neural networks using dropconnect," in *International Conference on Machine Learning*. PMLR, 2013, pp. 1058–1066.
- [35] Z. Goldfeld, E. Van Den Berg, K. Greenewald, I. Melnyk, N. Nguyen, B. Kingsbury, and Y. Polyanskiy, "Estimating information flow in deep neural networks," in *International Conference on Machine Learning*. PMLR, 2019, pp. 2299–2308.
- [36] Z. Goldfeld, K. Greenewald, J. Niles-Weed, and Y. Polyanskiy, "Convergence of smoothed empirical measures with applications to entropy estimation," *IEEE Transactions on Information Theory*, vol. 66, no. 7, pp. 4368–4391, 2020.
- [37] I. Gitman, H. Lang, P. Zhang, and L. Xiao, "Understanding the role of momentum in stochastic gradient methods," *Advances in Neural Information Processing Systems*, vol. 32, 2019.
- [38] A. Neelakantan, L. Vilnis, Q. V. Le, I. Sutskever, L. Kaiser, K. Kurach, and J. Martens, "Adding gradient noise improves learning for very deep networks," *arXiv preprint arXiv:1511.06807*, 2015.
- [39] C. M. Bishop, "Training with noise is equivalent to tikhonov regularization," *Neural Computation*, vol. 7, no. 1, pp. 108–116, 1995.
- [40] M. Raginsky, A. Rakhlin, and M. Telgarsky, "Non-convex learning via stochastic gradient Langevin dynamics: a nonasymptotic analysis," in *Conference on Learning Theory*. PMLR, 2017, pp. 1674–1703.
- [41] G. Aminian, Y. Bu, L. Toni, M. Rodrigues, and G. Wornell, "An exact characterization of the generalization error for the Gibbs algorithm," *Advances in Neural Information Processing Systems*, vol. 34, pp. 8106–8118, 2021.
- [42] B. Neyshabur, R. Tomioka, and N. Srebro, "In search of the real inductive bias: On the role of implicit regularization in deep learning," in *3rd International Conference on Learning Representations, ICLR 2015, San Diego, CA, USA, May 7-9, 2015, Workshop Track Proceedings*, 2015.
- [43] K. He, X. Zhang, S. Ren, and J. Sun, "Deep residual learning for image recognition," in *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, 2016, pp. 770–778.
- [44] C. Zhang, S. Bengio, M. Hardt, B. Recht, and O. Vinyals, "Understanding deep learning requires rethinking generalization," in *International Conference on Learning Representations*, 2017.
- [45] G. Vardi, "On the implicit bias in deep-learning algorithms," *Communications of the ACM*, vol. 66, no. 6, pp. 86–93, 2023.
- [46] R. Shwartz-Ziv and N. Tishby, "Opening the black box of deep neural networks via information," 2017, arXiv:1703.00810 [cs.LG].
- [47] A. M. Saxe, Y. Bansal, J. Dapello, M. Advani, A. Kolchinsky, B. D. Tracey, and D. D. Cox, "On the information bottleneck theory of deep learning," in *International Conference on Learning Representations*, 2018.
- [48] R. A. Amjad and B. C. Geiger, "Learning representations for neural network-based classification using the information bottleneck principle," *IEEE Transactions on Pattern Analysis and Machine Intelligence*, vol. 42, no. 9, pp. 2225–2239, 2019.
- [49] Z. Goldfeld and Y. Polyanskiy, "The information bottleneck problem and its applications in machine learning," *IEEE Journal on Selected Areas in Information Theory*, vol. 1, no. 1, pp. 19–38, 2020.
- [50] A. T. Lopez and V. Jog, "Generalization error bounds using Wasserstein distances," in *2018 IEEE Information Theory Workshop (ITW)*. IEEE, 2018, pp. 1–5.
- [51] A. Pensa, V. Jog, and P.-L. Loh, "Generalization error bounds for noisy, iterative algorithms," in *2018 IEEE International Symposium on Information Theory (ISIT)*. IEEE, 2018, pp. 546–550.
- [52] H. Wang, R. Gao, and F. P. Calmon, "Generalization bounds for noisy iterative algorithms using properties of additive noise channels," *Journal of machine learning research*, vol. 24, pp. 26:1–26:43, 2023.
- [53] G. Neu, G. K. Dziugaite, M. Haghifam, and D. M. Roy, "Information-theoretic generalization bounds for stochastic gradient descent," in *Conference on Learning Theory*. PMLR, 2021, pp. 3526–3545.
- [54] Z. Wang and Y. Mao, "On the generalization of models trained with SGD: Information-theoretic bounds and implications," in *International Conference on Learning Representations*, 2022.
- [55] A. van den Oord, O. Vinyals, and K. Kavukcuoglu, "Neural discrete representation learning," *Advances in Neural Information Processing Systems*, vol. 30, pp. 6306–6315, 2017.
- [56] B. Rodríguez Gálvez, G. Bassi, R. Thobaben, and M. Skoglund, "Tighter expected generalization error bounds via wasserstein distance," *Advances in Neural Information Processing Systems*, vol. 34, pp. 19109–19121, 2021.
- [57] A. L. Gibbs and F. E. Su, "On choosing and bounding probability metrics," *International Statistical Review*, vol. 70, no. 3, pp. 419–435, 2002.
- [58] J. E. Cohen, Y. Iwasa, G. Rautu, M. Beth Ruskai, E. Seneta, and G. Zbaganu, "Relative entropy under mappings by stochastic matrices," *Linear Algebra and its Applications*, vol. 179, pp. 211–235, 1993.
- [59] R. L. Dobrushin, "Central limit theorem for nonstationary markov chains. i," *Theory of Probability & Its Applications*, vol. 1, no. 1, pp. 65–80, 1956.
- [60] Y. Polyanskiy and Y. Wu, "Strong data-processing inequalities for channels and Bayesian networks," in *Convexity and Concentration*. Springer, 2017, pp. 211–249.
- [61] J. Cohen, J. H. Kempermann, and G. Zbaganu, *Comparisons of stochastic matrices with applications in information theory, statistics, economics and population*. Springer Science & Business Media, 1998.
- [62] T. M. Cover and J. A. Thomas, *Elements of Information Theory (Wiley Series in Telecommunications and Signal Processing)*. USA: Wiley-Interscience, 2006.

- [63] M. Welling and Y. W. Teh, “Bayesian learning via stochastic gradient langevin dynamics,” in *Proceedings of the 28th international conference on machine learning (ICML-11)*, 2011, pp. 681–688.
- [64] M. Raginsky, A. Rakhlin, M. Tsao, Y. Wu, and A. Xu, “Information-theoretic analysis of stability and bias of learning algorithms,” in *2016 IEEE Information Theory Workshop (ITW)*. IEEE, 2016, pp. 26–30.
- [65] P. Kidger and T. Lyons, “Universal approximation with deep narrow networks,” in *Conference on learning theory*. PMLR, 2020, pp. 2306–2327.
- [66] J. Lee, J. Y. Choi, E. K. Ryu, and A. No, “Neural tangent kernel analysis of deep narrow neural networks,” in *International Conference on Machine Learning*. PMLR, 2022, pp. 12 282–12 351.
- [67] C. Villani, *Topics in optimal transportation*. American Mathematical Soc., 2021, vol. 58.
- [68] O. Ordentlich and Y. Polyanskiy, “Strong data processing constant is achieved by binary inputs,” *IEEE Transactions on Information Theory*, vol. 68, no. 3, pp. 1480–1481, 2021.
- [69] Y. Polyanskiy and Y. Wu, “Dissipation of information in channels with input constraints,” *IEEE Transactions on Information Theory*, vol. 62, no. 1, pp. 35–55, 2015.
- [70] S. Barsov and V. V. Ul’yanov, “Estimates of the proximity of Gaussian measures,” in *Soviet Mathematics–Doklady*, vol. 34, 1987, pp. 462–466.

APPENDIX

A. Proof of Theorem 1

Let us rewrite the risks and generalization error under the DNN setup. Let $(X, Y) \sim P_{X,Y}$ be a pair of test data sample. At each layer l , the internal representation T_l of a test data feature X is conditionally independent of W_{l+1}^L given W_1^l . For any $\mathbf{W} \in \mathcal{W}$, let the loss function be rewritten as $\ell(\mathbf{W}, X, Y) = \ell(g_{\mathbf{W}_L} \circ g_{\mathbf{W}_{L-1}} \circ \dots \circ g_{\mathbf{W}_1}(X), Y)$. The expected population risk over all possible \mathbf{W} is given by

$$\begin{aligned} \mathbb{E}_{\mathbf{W}}[\mathcal{L}_{\mathcal{P}}(\mathbf{W}, P_{X,Y})] &= \mathbb{E}[\ell(g_{\mathbf{W}_L} \circ g_{\mathbf{W}_{L-1}} \circ \dots \circ g_{\mathbf{W}_1}(X), Y)] \\ &= \mathbb{E}[\ell(g_{\mathbf{W}_L} \circ g_{\mathbf{W}_{L-1}} \circ \dots \circ g_{\mathbf{W}_{l+1}}(T_l), Y)] \\ &= \mathbb{E}[\mathbb{E}[\ell(g_{\mathbf{W}_L} \circ g_{\mathbf{W}_{L-1}} \circ \dots \circ g_{\mathbf{W}_{l+1}}(T_l), Y) | \mathbf{W}_1^l]] \end{aligned}$$

where $l \in [L]$ and given \mathbf{W}_1^l , (T_l, Y) are independent of \mathbf{W}_{l+1}^L .

Denote the overall feature mapping function as $f_{\mathbf{W}} \triangleq g_{\mathbf{W}_L} \circ g_{\mathbf{W}_{L-1}} \circ \dots \circ g_{\mathbf{W}_1}$. Similarly, for any $l \in [L]$, the expected empirical risk can also be rewritten as

$$\begin{aligned} \mathbb{E}[\mathcal{L}_{\mathcal{E}}(\mathbf{W}, D_n)] &= \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n \ell(f_{\mathbf{W}}(X_i), Y_i)\right] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\ell(f_{\mathbf{W}}(X_i), Y_i)] \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\ell(g_{\mathbf{W}_L} \circ g_{\mathbf{W}_{L-1}} \circ \dots \circ g_{\mathbf{W}_{l+1}}(T_{l,i}), Y_i)] \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\mathbb{E}[\ell(g_{\mathbf{W}_L} \circ g_{\mathbf{W}_{L-1}} \circ \dots \circ g_{\mathbf{W}_{l+1}}(T_{l,i}), Y_i) | \mathbf{W}_1^l]]. \end{aligned}$$

For notational simplicity, let $g_{\mathbf{W}_k^j} := g_{\mathbf{W}_k} \circ g_{\mathbf{W}_{k-1}} \circ \dots \circ g_{\mathbf{W}_j}$ for any $k < j$ and $k, j \in \mathbb{N}$. Then the expected generalization error can be rewritten as

$$\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}\left[\mathbb{E}[\ell(g_{\mathbf{W}_{l+1}^L}(T_l), Y) | \mathbf{W}_1^l] - \mathbb{E}[\ell(g_{\mathbf{W}_{l+1}^L}(T_{l,i}), Y_i) | \mathbf{W}_1^l]\right]. \quad (6)$$

If the loss function $\ell(\mathbf{w}, X, Y)$ is σ -sub-Gaussian under $P_{X,Y}$ for all $\mathbf{w} \in \mathcal{W}$, we also have for any $l \in [0 : L]$, $\ell(g_{\mathbf{W}_{l+1}^L}(T_l), Y)$ is σ -sub-Gaussian under $P_{T_l, Y | \mathbf{W}=\mathbf{w}}$ for all $\mathbf{w} \in \mathcal{W}$. From Donsker-Varadhan representation, we have for any $\lambda \in \mathbb{R}$,

$$\begin{aligned} &D_{\text{KL}}(P_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^l} \| P_{T_l, Y | \mathbf{W}_1^l} \otimes P_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^l}) \\ &\geq \mathbb{E}_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^l}[\lambda \ell(g_{\mathbf{W}_{l+1}^L}(T_{l,i}), Y_i)] - \log \mathbb{E}_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^l} \mathbb{E}_{T_l, Y | \mathbf{W}_1^l}[\exp(\lambda \ell(g_{\mathbf{W}_{l+1}^L}(T_l), Y))] \\ &\geq \lambda (\mathbb{E}_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^l}[\lambda \ell(g_{\mathbf{W}_{l+1}^L}(T_{l,i}), Y_i)] - \mathbb{E}_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^l} \mathbb{E}_{T_l, Y | \mathbf{W}_1^l}[\ell(g_{\mathbf{W}_{l+1}^L}(T_l), Y)]) - \frac{\lambda^2 \sigma^2}{2}. \end{aligned}$$

We can decompose $D_{\text{KL}}(P_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^l} \| P_{T_l, Y | \mathbf{W}_1^l} \otimes P_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^l} | P_{\mathbf{W}_1^l})$ as follows

$$\begin{aligned} &D_{\text{KL}}(P_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^l} \| P_{T_l, Y | \mathbf{W}_1^l} \otimes P_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^l} | P_{\mathbf{W}_1^l}) \\ &= D_{\text{KL}}(P_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^l} \| P_{T_{l,i}, Y_i | \mathbf{W}_1^l} \otimes P_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^l} | P_{\mathbf{W}_1^l}) + D_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^l} \| P_{T_l, Y | \mathbf{W}_1^l} | P_{\mathbf{W}_1^l}) \\ &= I(T_{l,i}, Y_i; \mathbf{W}_{l+1}^L | \mathbf{W}_1^l) + D_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^l} \| P_{T_l, Y | \mathbf{W}_1^l} | P_{\mathbf{W}_1^l}). \end{aligned} \quad (7)$$

Thus, we have

$$\begin{aligned}
& I(T_{l,i}, Y_i; \mathbf{W}_{l+1}^L | \mathbf{W}_1^L) + \text{D}_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_l, Y | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}) \\
&= \text{D}_{\text{KL}}(P_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_l, Y | \mathbf{W}_1^L} \otimes P_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}) \\
&\geq \lambda \mathbb{E}_{\mathbf{W}_1^L} [\mathbb{E}_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^L} [\ell(g_{\mathbf{W}_{l+1}^L}(T_{l,i}), Y_i)] - \mathbb{E}_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^L} \mathbb{E}_{T_l, Y | \mathbf{W}_1^L} [\ell(g_{\mathbf{W}_{l+1}^L}(T_l), Y)]]] - \frac{\lambda^2 \sigma^2}{2}.
\end{aligned}$$

By optimizing the RHS over $\lambda > 0$ and $\lambda \leq 0$, respectively, we finally obtain

$$\begin{aligned}
& \left| \mathbb{E}_{\mathbf{W}_1^L} \mathbb{E}_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^L} [\ell(g_{\mathbf{W}_{l+1}^L}(T_{l,i}), Y_i)] - \mathbb{E}_{\mathbf{W}_1^L} \mathbb{E}_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^L} \mathbb{E}_{T_l, Y | \mathbf{W}_1^L} [\ell(g_{\mathbf{W}_{l+1}^L}(T_l), Y)] \right| \\
&\leq \sqrt{2\sigma^2 (I(T_{l,i}, Y_i; \mathbf{W}_{l+1}^L | \mathbf{W}_1^L) + \text{D}_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_l, Y | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}))},
\end{aligned}$$

which holds for all $l \in [L]$. Conditioned on \mathbf{W}_l , $T_{l,i}$ and T_l are generated by the same process from $T_{l-1,i}$ and T_{l-1} , respectively. By the data-processing inequality, the KL divergence in (7) can be bounded as follows:

$$\begin{aligned}
& \text{D}_{\text{KL}}(P_{\mathbf{W}_{l+1}^L, T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_l, Y | \mathbf{W}_1^L} \otimes P_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}) \\
&\leq \text{D}_{\text{KL}}(P_{\mathbf{W}_{l+1}^L, T_{l-1,i}, Y_i | \mathbf{W}_1^L} \| P_{T_{l-1}, Y | \mathbf{W}_1^L} \otimes P_{\mathbf{W}_{l+1}^L | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}) \\
&= \text{D}_{\text{KL}}(P_{\mathbf{W}_l^L, T_{l-1,i}, Y_i | \mathbf{W}_1^{l-1}} \| P_{T_{l-1}, Y | \mathbf{W}_1^{l-1}} \otimes P_{\mathbf{W}_l^L | \mathbf{W}_1^{l-1}} | P_{\mathbf{W}_1^{l-1}}) \\
&\quad \vdots \\
&\leq \text{D}_{\text{KL}}(P_{X_i, Y_i, \mathbf{W}_1^L} \| P_{X, Y} \otimes P_{\mathbf{W}_1^L}) \\
&= I(X_i, Y_i; \mathbf{W}).
\end{aligned}$$

Therefore, the expected generalization error in (6) can be upper bounded as follows:

$$\begin{aligned}
& |\text{gen}(P_{\mathbf{W} | D_n}, P_{X, Y})| \\
&\leq \frac{1}{n} \sum_{i=1}^n \sqrt{2\sigma^2 \text{D}_{\text{KL}}(P_{T_{L,i}, Y_i | \mathbf{W}} \| P_{T_L, Y | \mathbf{W}} | P_{\mathbf{W}})} \\
&\leq \frac{1}{n} \sum_{i=1}^n \sqrt{2\sigma^2 (I(T_{L-1,i}, Y_i; \mathbf{W}_L | \mathbf{W}_1^{L-1}) + \text{D}_{\text{KL}}(P_{T_{L-1,i}, Y_i | \mathbf{W}_1^{L-1}} \| P_{T_{L-1}, Y | \mathbf{W}_1^{L-1}} | P_{\mathbf{W}_1^{L-1}}))} \\
&\quad \vdots \\
&\leq \frac{1}{n} \sum_{i=1}^n \sqrt{2\sigma^2 (I(T_{l,i}, Y_i; \mathbf{W}_l^L | \mathbf{W}_1^L) + \text{D}_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_l, Y | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}))} \\
&\quad \vdots \\
&\leq \frac{1}{n} \sum_{i=1}^n \sqrt{2\sigma^2 (I(T_{1,i}, Y_i; \mathbf{W}_1^L | \mathbf{W}_1) + \text{D}_{\text{KL}}(P_{T_{1,i}, Y_i | \mathbf{W}_1} \| P_{T_1, Y | \mathbf{W}_1} | P_{\mathbf{W}_1}))} \\
&\leq \frac{1}{n} \sum_{i=1}^n \sqrt{2\sigma^2 I(X_i, Y_i; \mathbf{W})}.
\end{aligned}$$

B. Proof for Remark 1

We prove the simplified generalization bound for the discrete latent space case. The information-theoretic quantities in $\text{UB}(l)$ can be upper bounded as follows:

$$\begin{aligned}
& I(T_{l,i}, Y_i; \mathbf{W}_{l+1}^L | \mathbf{W}_1^L) + \text{D}_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_l, Y | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}) \\
&= I(T_{l,i}; \mathbf{W}_{l+1}^L | \mathbf{W}_1^L) + I(Y_i; \mathbf{W}_{l+1}^L | T_{l,i}, \mathbf{W}_1^L) + \text{D}_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{W}_1^L} \| P_{T_l, Y | \mathbf{W}_1^L} | P_{\mathbf{W}_1^L}) \\
&= H(T_{l,i} | \mathbf{W}_1^L) - H(T_{l,i} | \mathbf{W}_1^L, \mathbf{W}_{l+1}^L) + H(Y_i | T_{l,i}, \mathbf{W}_1^L) - H(Y_i | T_{l,i}, \mathbf{W}_1^L, \mathbf{W}_{l+1}^L) \\
&\quad - H(T_{l,i}, Y_i | \mathbf{W}_1^L) - \mathbb{E}_{P_{T_{l,i}, Y_i, \mathbf{W}_1^L}} [\log P_{T_l, Y | \mathbf{W}_1^L}] \\
&= H(T_{l,i} | \mathbf{W}_1^L) - H(T_{l,i} | \mathbf{W}_1^L, \mathbf{W}_{l+1}^L) + H(Y_i | T_{l,i}, \mathbf{W}_1^L) - H(Y_i | T_{l,i}, \mathbf{W}_1^L, \mathbf{W}_{l+1}^L) \\
&\quad - H(T_{l,i} | \mathbf{W}_1^L) + H(Y_i | T_{l,i}, \mathbf{W}_1^L) - \mathbb{E}_{P_{T_{l,i}, Y_i, \mathbf{W}_1^L}} [\log P_{T_l, Y | \mathbf{W}_1^L}] \\
&\stackrel{(a)}{\leq} 2H(Y_i | T_{l,i}, \mathbf{W}_1^L) - \mathbb{E}_{P_{T_{l,i}, Y_i, \mathbf{W}_1^L}} [\log P_{T_l, Y | \mathbf{W}_1^L}]
\end{aligned}$$

$$\begin{aligned}
&\leq 2 \log |\mathcal{Y}| - \mathbb{E}_{P_{T_l, i, Y_i, \mathbf{w}_1^l}} [\log P_{T_l, Y} | \mathbf{w}_1^l] \\
&= 2 \log K - \mathbb{E}_{P_{T_l, i, Y_i, \mathbf{w}_1^l}} [\log P_{T_l, Y} | \mathbf{w}_1^l],
\end{aligned}$$

where (a) follows since $H(T_{l,i} | \mathbf{W}_1^l, \mathbf{W}_{l+1}^L) \geq 0$. Assuming $t_l(\mathbf{w}_1^l) := \min_{t \in \mathcal{T}_l, y \in \mathcal{Y}} P_{T_l, Y} | \mathbf{w}_1^l(t, y | \mathbf{w}_1^l) \in (0, |\mathcal{T}_l \times \mathcal{Y}|^{-1})$ and $\bar{t}_l := \sup_{\mathbf{w}_1^l} t_l(\mathbf{w}_1^l)$, then

$$\begin{aligned}
\mathbb{E}_{P_{T_l, i, Y_i, \mathbf{w}_1^l}} [\log P_{T_l, Y} | \mathbf{w}_1^l] &\geq \log \bar{t}_l, \quad \text{and} \\
I(T_{l,i}, Y_i; \mathbf{W}_{l+1}^L | \mathbf{W}_1^l) + D_{\text{KL}}(P_{T_l, i, Y_i} | \mathbf{w}_1^l \| P_{T_l, Y} | \mathbf{w}_1^l | P_{\mathbf{w}_1^l}) &\leq \log \frac{K^2}{\bar{t}_l}.
\end{aligned}$$

C. Proof of Theorem 2

Recall the Kantorovich-Rubinstein duality [67]: for any two probability measures $P, Q \in \mathcal{P}(\mathcal{X})$, $W_1(P, Q) = \sup_{f \in \text{Lip}_1(\mathcal{X})} \mathbb{E}_P[f] - \mathbb{E}_Q[f]$, where $\text{Lip}_k(\mathcal{X}) = \{f \in \{f : \mathcal{X} \rightarrow \mathbb{R}\} : |f(x) - f(y)| \leq k \|x - y\|, \forall x, y \in \mathcal{X}\}$, for any $k \in \mathbb{R}_{\geq 0}$

Since $\tilde{\ell}(g_{\mathbf{w}_L} \circ \dots \circ g_{\mathbf{w}_1}(X), Y)$ is ρ_0 -Lipschitz in $(g_{\mathbf{w}_L} \circ \dots \circ g_{\mathbf{w}_1}(X), Y)$ and $\phi_l(\cdot)$ is ρ_l -Lipschitz, we have for any \mathbf{w} ,

$$\begin{aligned}
|\ell(\mathbf{w}, x, y) - \ell(\mathbf{w}, x', y')| &= |\tilde{\ell}(g_{\mathbf{w}_L} \circ \dots \circ g_{\mathbf{w}_1}(x), y) - \tilde{\ell}(g_{\mathbf{w}_L} \circ \dots \circ g_{\mathbf{w}_1}(x'), y')| \\
&\leq \rho_0 \|(g_{\mathbf{w}_L}(g_{\mathbf{w}_1^{L-1}}(x)), y) - (g_{\mathbf{w}_L}(g_{\mathbf{w}_1^{L-1}}(x')), y')\| \\
&= \rho_0 \sqrt{(\phi_L(\mathbf{w}_L g_{\mathbf{w}_1^{L-1}}(x)) - \phi_L(\mathbf{w}_L g_{\mathbf{w}_1^{L-1}}(x')))^2 + (y - y')^2} \\
&\leq \rho_0 \sqrt{(\rho_L \|\mathbf{w}_L\| \|g_{\mathbf{w}_1^{L-1}}(x) - g_{\mathbf{w}_1^{L-1}}(x')\|)^2 + (y - y')^2} \\
&= \rho_0 \sqrt{\rho_L^2 \|\mathbf{w}_L\|^2 \|\phi_{L-1}(\mathbf{w}_{L-1} g_{\mathbf{w}_1^{L-2}}(x)) - \phi_{L-1}(\mathbf{w}_{L-1} g_{\mathbf{w}_1^{L-2}}(x'))\|^2 + (y - y')^2} \\
&\leq \rho_0 \sqrt{\rho_L^2 \rho_{L-1}^2 \|\mathbf{w}_L\|^2 \|\mathbf{w}_{L-1}\|^2 \|g_{\mathbf{w}_1^{L-2}}(x) - g_{\mathbf{w}_1^{L-2}}(x')\|^2 + (y - y')^2} \\
&\vdots \\
&\leq \rho_0 \sqrt{\prod_{j=1}^L \rho_j^2 \|\mathbf{w}_j\|^2 \|x - x'\|^2 + (y - y')^2} \\
&\leq \bar{\rho}_0(\mathbf{w}) \sqrt{\|x - x'\|^2 + (y - y')^2}
\end{aligned}$$

where $\bar{\rho}_0(\mathbf{w}) := \rho_0(1 \vee \prod_{j=1}^L \rho_j \|\mathbf{w}_j\|)$. It means $\ell(\mathbf{W}, X, Y)$ is $\bar{\rho}_0(\mathbf{W})$ -Lipschitz in (X, Y) for any \mathbf{W} . Then we have

$$\begin{aligned}
\text{gen}(P_{\mathbf{W}|D_n}, P_{X, Y}) &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\ell(\mathbf{W}, X, Y) - \ell(\mathbf{W}, X_i, Y_i)] \\
&\leq \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\bar{\rho}_0(\mathbf{W}) W_1(P_{X_i, Y_i} | \mathbf{w}, P_{X, Y} | \mathbf{w})].
\end{aligned}$$

For $l = 1, \dots, L$, similarly, we have $\tilde{\ell}(g_{\mathbf{w}_l} \circ \dots \circ g_{\mathbf{w}_{l+1}}(T_l), Y)$ is $\rho_0(1 \vee \prod_{j=l+1}^L \rho_j \|\mathbf{W}_j\|)$ -Lipschitz in (T_l, Y) for all \mathbf{W} .

Let $\bar{\rho}_l(\mathbf{W}) = \rho_0(1 \vee \prod_{j=l+1}^L \rho_j \|\mathbf{W}_j\|)$. Then from the definition in (6), we have

$$\text{gen}(P_{\mathbf{W}|D_n}, P_{X, Y}) \leq \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\bar{\rho}_l(\mathbf{W}) W_1(P_{T_{l,i}, Y_i} | \mathbf{w}, P_{T_l, Y} | \mathbf{w})].$$

The proof is completed by taking the minimum over $l = 0, \dots, L$.

D. Proof for Remark 2

Let $\text{diam}(\mathcal{X}) := \sup\{\|x - y\| : x, y \in \mathcal{X}\}$. From [57, Theorem 4], Pinsker's and Bretagnolle-Huber inequalities, for any two probability distributions $\mu, \nu \in \mathcal{P}(\mathcal{X})$, we have

$$W_1(\mu, \nu) \leq \text{diam}(\mathcal{X}) D_{\text{TV}}(\mu, \nu) \leq \text{diam}(\mathcal{X}) \sqrt{\left(\frac{1}{2} D_{\text{KL}}(\mu \| \nu) \wedge (1 - \exp(-D_{\text{KL}}(\mu \| \nu)))\right)}.$$

From Theorem 1 and [52], the generalization error can be bounded as follows:

$$\left| \text{gen}(P_{\mathbf{W}|D_n}, P_{X, Y}) \right| \leq \frac{A}{n} \sum_{i=1}^n D_{\text{TV}}(P_{T_{L,i}, Y_i} | \mathbf{w}, P_{T_L, Y} | \mathbf{w} | P_{\mathbf{W}}) \leq \text{UB}(L),$$

where $\text{UB}(L) := \frac{\sqrt{2}A}{2n} \sum_{i=1}^n \sqrt{D_{\text{KL}}(P_{T_L,i,Y_i|\mathbf{W}} \| P_{T_L,Y|\mathbf{W}} | P_{\mathbf{W}})}$. It can be observed that the total variation distance based bound is tighter under this condition.

Let $l^* := \min_{l=0,\dots,L} \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\bar{\rho}_l(\mathbf{W}) \mathbf{W}_1(P_{T_{l,i,Y_i|\mathbf{W}}}, P_{T_{l,Y|\mathbf{W}}})]$.

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\bar{\rho}_{l^*}(\mathbf{W}) \mathbf{W}_1(P_{T_{l^*,i,Y_i|\mathbf{W}}}, P_{T_{l^*,Y|\mathbf{W}}})] &\leq \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\bar{\rho}_L(\mathbf{W}) \mathbf{W}_1(P_{T_{L,i,Y_i|\mathbf{W}}}, P_{T_{L,Y|\mathbf{W}}})] \\ &= \frac{\rho_0}{n} \sum_{i=1}^n \mathbf{W}_1(P_{T_{L,i,Y_i|\mathbf{W}}}, P_{T_{L,Y|\mathbf{W}}} | P_{\mathbf{W}}) \\ &\leq \frac{\rho_0 \text{diam}(\mathcal{T}_L \times \mathcal{Y})}{n} \sum_{i=1}^n D_{\text{TV}}(P_{T_{L,i,Y_i|\mathbf{W}}}, P_{T_{L,Y|\mathbf{W}}} | P_{\mathbf{W}}). \end{aligned}$$

Under our supervised classification setting, $\text{diam}(\mathcal{T}_L \times \mathcal{Y}) = K^2$. Thus, when $\rho_0 K^2 \leq A$, 1-Wasserstein distance based bound is even tighter than the one based on the TV-distance.

E. Proofs for the case study of binary Gaussian mixture

To simplify some of the notation ahead, the distribution of a Gaussian random variable X with mean μ and variance σ^2 is denoted by $\mathcal{N}_X(\mu, \sigma^2)$. The subscript stressing the random variable at play is convenient for telling apart the various Gaussian distributions to follow; when there is no ambiguity, the subscript is dropped.

Under the binary Gaussian mixture classification setting, we first know that the prior of $\mathbf{W}_{\otimes L}^\top$ is $P_{\mathbf{W}_{\otimes L}^\top} = \mathcal{N}_{\mathbf{W}_{\otimes L}^\top}(\mu_0, \frac{\sigma_0^2}{n} \mathbf{I}_{d_0})$. Given any pair of training data sample (X_i, Y_i) , we have

$$\mathbf{W}_{\otimes L}^\top | (X_i, Y_i) = \frac{1}{n} Y_i X_i + \frac{1}{n} \sum_{j \neq i} Y_j X_j \sim \mathcal{N}_{\mathbf{W}_{\otimes L}^\top}(\mu_{\mathbf{W}_{\otimes L} | i}, \Sigma_{\mathbf{W}_{\otimes L} | i}),$$

where $\mu_{\mathbf{W}_{\otimes L} | i} = \frac{1}{n} Y_i X_i + \frac{n-1}{n} \mu_0$ and $\Sigma_{\mathbf{W}_{\otimes L} | i} = \frac{n-1}{n^2} \sigma_0^2 \mathbf{I}_{d_0}$. Then the posterior distribution of (X_i, Y_i) given $\mathbf{W}_{\otimes L}$ is given by

$$\begin{aligned} P_{X_i, Y_i | \mathbf{W}_{\otimes L}} &= \frac{P_{\mathbf{W}_{\otimes L}^\top | X_i, Y_i} P_{X_i, Y_i}}{P_{\mathbf{W}_{\otimes L}}} \\ &= \frac{\mathcal{N}_{\mathbf{W}_{\otimes L}^\top}(\mu_{\mathbf{W}_{\otimes L} | i}, \Sigma_{\mathbf{W}_{\otimes L} | i})}{\mathcal{N}_{\mathbf{W}_{\otimes L}^\top}(\mu_0, \frac{\sigma_0^2}{n} \mathbf{I}_{d_0})} \times \frac{1}{2} \mathcal{N}_{X_i}(Y_i \mu_0, \sigma_0^2 \mathbf{I}_{d_0}) \\ &= \frac{1}{2} \mathcal{N}_{X_i}(Y_i \mu_0, \sigma_0^2 \mathbf{I}_{d_0}) \times C_i \mathcal{N}_{\mathbf{W}_{\otimes L}^\top} \left(Y_i X_i, \frac{(n-1)\sigma_0^2}{n} \mathbf{I}_{d_0} \right) \\ &= \frac{1}{2} \mathcal{N}_{Y_i X_i} \left(\mathbf{W}_{\otimes L}^\top, \frac{(n-1)\sigma_0^2}{n} \mathbf{I}_{d_0} \right), \end{aligned}$$

where $C_i = n^{d_0} \sqrt{\left(\frac{2\pi\sigma_0^2}{n}\right)^{d_0}} \exp\left\{-\frac{1}{2\sigma_0^2} (Y_i X_i - \mu_0)^\top (Y_i X_i - \mu_0)\right\}$. By integrating $P_{X_i, Y_i | \mathbf{W}_{\otimes L}}$ over X_i , we obtain $P_{Y_i=1 | \mathbf{W}_{\otimes L}} = \frac{1}{2} = P_{Y_i=-1 | \mathbf{W}_{\otimes L}} = \frac{1}{2}$. We can also conclude that $P_{X_i, Y_i | \mathbf{W}, \mathbf{W}_{\otimes L}} = P_{X_i, Y_i | \mathbf{W}_{\otimes L}}$ and $P_{Y_i | \mathbf{W}, \mathbf{W}_{\otimes L}} = P_{Y_i | \mathbf{W}_{\otimes L}} = \text{Unif}\{-1, +1\}$.

Next, we compute the divergences between the prior and posterior.

Proof of Proposition 4. For any $l \in [L]$, conditioned on $(Y_i, \mathbf{W}, \mathbf{W}_{\otimes L})$, the distribution of $T_{l,i} = \mathbf{W}_{\otimes l} X_i$ is Gaussian with the mean and covariance

$$\mathbb{E}[T_{l,i} | Y_i, \mathbf{W}, \mathbf{W}_{\otimes L}] = Y_i \mathbf{W}_{\otimes l} \mathbf{W}_{\otimes L}^\top, \quad \text{Cov}[T_{l,i} | Y_i, \mathbf{W}, \mathbf{W}_{\otimes L}] = \frac{(n-1)\sigma_0^2}{n} \mathbf{W}_{\otimes l} \mathbf{W}_{\otimes L}^\top.$$

Similarly for a test data sample, when conditioned on $(Y, \mathbf{W}, \mathbf{W}_{\otimes L})$, the distribution of $T_l = \mathbf{W}_{\otimes l} X$ is Gaussian with mean and covariance

$$\mathbb{E}[T_l | Y, \mathbf{W}, \mathbf{W}_{\otimes L}] = \mathbb{E}[T_l | Y, \mathbf{W}] = Y_l \mathbf{W}_{\otimes l} \mu_0, \quad \text{Cov}[T_l | Y_l, \mathbf{W}, \mathbf{W}_{\otimes L}] = \sigma_0^2 \mathbf{W}_{\otimes l} \mathbf{W}_{\otimes L}^\top.$$

Note that when $\mathbf{W}_{\otimes l}$ is not a full-rank matrix, the covariance matrices $\text{Cov}[T_{l,i} | Y_i, \mathbf{W}, \mathbf{W}_{\otimes L}]$ and $\text{Cov}[T_l | Y_l, \mathbf{W}, \mathbf{W}_{\otimes L}]$ are singular. Thus, the posteriors $P_{T_{l,i} | Y_i, \mathbf{W}, \mathbf{W}_{\otimes L}}$ and $P_{T_l | Y_l, \mathbf{W}, \mathbf{W}_{\otimes L}}$ are the push-forwards of two d_l -dimensional nonsingular Gaussian distributions to the lower-dimensional space. In fact, the dimension can be further proven to be $\text{rank}(\mathbf{W}_{\otimes l})$ via eigendecomposition.

Take the test data sample $(X, Y = 1)$ for example in the followings. Let $X = \mu_0 + Z$ and the l^{th} representation $T_l = \mathbf{W}_{\otimes l} X = \mathbf{W}_{\otimes l} \mu_0 + \mathbf{W}_{\otimes l} Z$, where $Z \sim \mathcal{N}(0, \sigma_0^2 \mathbf{I}_{d_0})$ and $\mathbf{W}_{\otimes l} Z \sim \mathcal{N}(0, \sigma_0^2 \mathbf{W}_{\otimes l} \mathbf{W}_{\otimes L}^\top)$ is a singular Gaussian distribution.

Since $\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top$ is a $(d_l \times d_l)$ positive semi-definite and symmetric matrix, the eigendecomposition (or singular value decomposition (SVD)) is given by

$$\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^\top = (\mathbf{U} \mathbf{\Lambda}^{\frac{1}{2}})(\mathbf{U} \mathbf{\Lambda}^{\frac{1}{2}})^\top,$$

where \mathbf{U} is a $(d_l \times d_l)$ orthogonal matrix, $\mathbf{\Lambda}$ is a $(d_l \times d_l)$ diagonal matrix whose entries are the eigenvalues of $\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top$. Without loss of generality, we assume that the diagonal values (i.e., eigenvalues) in $\mathbf{\Lambda}$ are in descending order. Note that the SVD of $\mathbf{W}_{\otimes l}$ is $\mathbf{W}_{\otimes l} = \mathbf{U} \mathbf{S} \mathbf{V}^\top$ where \mathbf{S} is a $(d_l \times d_0)$ diagonal matrix, $\mathbf{S} \mathbf{S}^\top = \mathbf{\Lambda}$, \mathbf{V} is a $(d_0 \times d_0)$ orthogonal matrix. Construct the following two d_l -dimensional vectors:

$$Z_0 := (Z, \underbrace{0, \dots, 0}_{(d_l - d_0) \text{ entries}})^\top, \quad \tilde{Z} := (\underbrace{\tilde{Z}_1, \dots, \tilde{Z}_{\text{rank}(\mathbf{W}_{\otimes l})}}_{\sim \mathcal{N}(0, \sigma_0^2 \mathbf{I}_{\text{rank}(\mathbf{W}_{\otimes l})})}, \underbrace{0, \dots, 0}_{(d_l - \text{rank}(\mathbf{W}_{\otimes l}))^\top \text{ entries}}).$$

Since the last $(d_l - \text{rank}(\mathbf{W}_{\otimes l}))$ columns of $\mathbf{U} \mathbf{\Lambda}^{\frac{1}{2}}$ are all zero, the following equalities hold:

$$\mathbf{W}_{\otimes l} Z \stackrel{d}{=} \mathbf{U} \mathbf{\Lambda}^{\frac{1}{2}} Z_0 \stackrel{d}{=} \mathbf{U} \mathbf{\Lambda}^{\frac{1}{2}} \tilde{Z}, \quad \text{and} \quad T_l \stackrel{d}{=} \mathbf{W}_{\otimes l} \mu_0 + \mathbf{U} \mathbf{\Lambda}^{\frac{1}{2}} \tilde{Z}.$$

Thus, only the a subset of $\text{rank}(\mathbf{W}_{\otimes l})$ covariates of T_l are effective random variables and the Gaussian PDF of T_l is on the $\text{rank}(\mathbf{W}_{\otimes l})$ -dimensional space. Let $r_l = \text{rank}(\mathbf{W}_{\otimes l})$. We can define a restriction of Lebesgue measure to the $\text{rank}(\mathbf{W}_{\otimes l})$ -dimensional affine subspace of \mathbb{R}^{r_l} where the Gaussian distribution is supported. With respect to this measure the distribution of T_l given $(\mathbf{W}, \mathbf{W}_{\otimes l}, Y = 1)$ has the density of the following motif:

$$p_{\mathbf{W}, \mathbf{W}_{\otimes l}, Y=1}(T_l) = \frac{\exp\left(-\frac{1}{2\sigma_0^2} (T_l - \mathbf{W}_{\otimes l} \mu_0)^\top (\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top)^\dagger (T_l - \mathbf{W}_{\otimes l} \mu_0)\right)}{\sqrt{(2\pi)^{r_l} \det^*(\sigma_0^2 \mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top)}} \quad (8)$$

where $(\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top)^\dagger = \mathbf{U} \mathbf{\Lambda}^\dagger \mathbf{U}^\top$ is the generalized inverse of $\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top$, $\mathbf{\Lambda}^\dagger$ is the pseudo-inverse of $\mathbf{\Lambda}$, and \det^* is the pseudo-determinant. In a similar manner, the density of $T_{l,i}$ can be obtained by replacing the mean $\mathbf{W}_{\otimes l} \mu_0$ with $\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes L}^\top$ and σ_0^2 with $\frac{(n-1)\sigma_0^2}{n}$.

Recall that the KL divergence between any two Gaussian distributions $P = \mathcal{N}(\mu_p, \mathbf{\Sigma}_p), Q = \mathcal{N}(\mu_q, \mathbf{\Sigma}_q) \in \mathcal{P}(\mathbb{R}^k)$ for some $k \in \mathbb{N}$ is given by

$$D_{\text{KL}}(P||Q) = \frac{1}{2} \left(\log \frac{\det^*(\mathbf{\Sigma}_q)}{\det^*(\mathbf{\Sigma}_p)} - k + (\mu_p - \mu_q)^\top \mathbf{\Sigma}_q^\dagger (\mu_p - \mu_q) + \text{tr}(\mathbf{\Sigma}_q^\dagger \mathbf{\Sigma}_p) \right).$$

For $l = 1, \dots, L$, the UB(l) in Theorem 1 can be written as

$$\begin{aligned} \text{UB}(l) &= \frac{\sqrt{2}\sigma}{n} \sum_{i=1}^n \sqrt{D_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{w}} || P_{T_l, Y | \mathbf{w}} | P_{\mathbf{w}})} \\ &\leq \frac{\sqrt{2}\sigma}{n} \sum_{i=1}^n \sqrt{D_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{w}, \mathbf{w}_{\otimes L}} || P_{T_l, Y | \mathbf{w}, \mathbf{w}_{\otimes L}} | P_{\mathbf{w}, \mathbf{w}_{\otimes L}})} =: \widetilde{\text{UB}}(l), \end{aligned}$$

where the hierarchical structure $\widetilde{\text{UB}}(L) \leq \widetilde{\text{UB}}(L-1) \leq \dots \leq \widetilde{\text{UB}}(0)$ still holds.

From the probability density function (PDF) of T_l and $T_{l,i}$ (c.f. (8)), the KL divergence term in the upper bound $\widetilde{\text{UB}}(l)$ can be rewritten as: for $l = 1, \dots, L$,

$$\begin{aligned} &D_{\text{KL}}(P_{T_{l,i}, Y_i | \mathbf{w}, \mathbf{w}_{\otimes L}} || P_{T_l, Y | \mathbf{w}, \mathbf{w}_{\otimes L}} | P_{\mathbf{w}, \mathbf{w}_{\otimes L}}) \\ &= \frac{1}{2} D_{\text{KL}}(P_{T_{l,i}, Y_i=1 | \mathbf{w}, \mathbf{w}_{\otimes L}} || P_{T_l, Y=1 | \mathbf{w}, \mathbf{w}_{\otimes L}} | P_{\mathbf{w}, \mathbf{w}_{\otimes L}}) + \frac{1}{2} D_{\text{KL}}(P_{T_{l,i}, Y_i=-1 | \mathbf{w}, \mathbf{w}_{\otimes L}} || P_{T_l, Y=-1 | \mathbf{w}, \mathbf{w}_{\otimes L}} | P_{\mathbf{w}, \mathbf{w}_{\otimes L}}) \\ &= \frac{1}{2} \mathbb{E} \left[\mathbb{E} \left[r_l \left(\log \frac{n}{n-1} - 1 + \frac{n-1}{n} \right) + \frac{1}{\sigma_0^2} (\mu_0 - \mathbf{W}_{\otimes L}^\top)^\top (\mathbf{W}_{\otimes l})^\top (\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top)^\dagger \mathbf{W}_{\otimes l} (\mu_0 - \mathbf{W}_{\otimes L}^\top) \middle| \mathbf{W}, \mathbf{W}_{\otimes L} \right] \right] \\ &= \frac{1}{2} \mathbb{E} \left[\mathbb{E} \left[r_l \left(\log \frac{n}{n-1} - \frac{1}{n} \right) + \frac{1}{\sigma_0^2} (\mu_0 - \mathbf{W}_{\otimes L}^\top)^\top \mathbf{V} \mathbf{S}^\top \mathbf{U}^\top \mathbf{U} \mathbf{\Lambda}^\dagger \mathbf{U} \mathbf{S} \mathbf{V}^\top (\mu_0 - \mathbf{W}_{\otimes L}^\top) \middle| \mathbf{W}, \mathbf{W}_{\otimes L} \right] \right] \\ &= \frac{1}{2} \mathbb{E} \left[\mathbb{E} \left[r_l \left(\log \frac{n}{n-1} - \frac{1}{n} \right) + \frac{1}{\sigma_0^2} \|\mu_0 - \mathbf{W}_{\otimes L}^\top\|^2 \middle| \mathbf{W}, \mathbf{W}_{\otimes L} \right] \right] \\ &= \mathbb{E} \left[\frac{r_l}{2} \left(\log \frac{n}{n-1} - \frac{1}{n} \right) + \frac{1}{2\sigma_0^2} \mathbb{E}[\|\mu_0 - \mathbf{W}_{\otimes L}^\top\|^2] \right] \end{aligned}$$

$$= \frac{\mathbb{E}[r_l]}{2} \left(\log \frac{n}{n-1} - \frac{1}{n} \right) + \frac{d_0}{2n},$$

where the last equality follows since $\frac{\sqrt{n}}{\sigma_0}(\mu_0 - \mathbf{W}_{\otimes L}^\top) \sim \mathcal{N}(0, \mathbf{I}_{d_0})$, $\frac{n}{\sigma_0^2} \|\mu_0 - \mathbf{W}_{\otimes L}^\top\|^2 \sim \chi_{d_0}^2$ and

$$\frac{1}{2\sigma_0^2} \mathbb{E}[\|\mu_0 - \mathbf{W}_{\otimes L}^\top\|^2] = \frac{d_0}{2n}.$$

The KL divergence term in the upper bound $\widetilde{\text{UB}}(0)$ can be rewritten as

$$\begin{aligned} & \text{D}_{\text{KL}}(P_{X_i, Y_i | \mathbf{W}, \mathbf{W}_{\otimes L}} \| P_{X, Y} | P_{\mathbf{W}, \mathbf{W}_{\otimes L}}) \\ &= \mathbb{E} \left[\mathbb{E} \left[\text{D}_{\text{KL}} \left(\frac{1}{2} \mathcal{N}_{X_i} \left(-\mathbf{W}_{\otimes L}^\top, \frac{(n-1)\sigma_0^2}{n} \mathbf{I}_{d_0} \right) \left\| \frac{1}{2} \mathcal{N}_X(-\mu_0, \sigma_0^2 \mathbf{I}_{d_0}) \right\| \right) \right. \right. \\ & \quad \left. \left. + \text{D}_{\text{KL}} \left(\frac{1}{2} \mathcal{N}_{X_i} \left(\mathbf{W}_{\otimes L}^\top, \frac{(n-1)\sigma_0^2}{n} \mathbf{I}_{d_0} \right) \left\| \frac{1}{2} \mathcal{N}_X(\mu_0, \sigma_0^2 \mathbf{I}_{d_0}) \right\| \right) \middle| \mathbf{W}, \mathbf{W}_{\otimes L} \right] \right] \\ &= \frac{1}{2} \mathbb{E} \left[\mathbb{E} \left[d_0 \left(\log \frac{n}{n-1} - 1 + \frac{n-1}{n} \right) + \frac{1}{\sigma_0^2} \|\mu_0 - \mathbf{W}_{\otimes L}^\top\|^2 \middle| \mathbf{W}, \mathbf{W}_{\otimes L} \right] \right] \\ &= \frac{d_0}{2} \left(\log \frac{n}{n-1} - \frac{1}{n} \right) + \frac{1}{2\sigma_0^2} \mathbb{E}[\|\mu_0 - \mathbf{W}_{\otimes L}^\top\|^2] \\ &= \frac{d_0}{2} \left(\log \frac{n}{n-1} - \frac{1}{n} \right) + \frac{d_0}{2n}. \end{aligned}$$

Note that $r_l = \text{rank}(\mathbf{W}_{\otimes l}) \leq \min\{\text{rank}(\mathbf{W}_l), \text{rank}(\mathbf{W}_{\otimes(l-1)})\} \leq \text{rank}(\mathbf{W}_{\otimes(l-1)}) = r_{l-1}$, for any $l = 2, \dots, L$, and $r_1 \leq \min\{d_0, d_1\}$. \square

Proof of Proposition 5. Since the closed form of W_1 between two Gaussian distributions is not known but known for W_2 and $W_1(\cdot, \cdot) \leq W_2(\cdot, \cdot)$, we consider analysing $W_2(P_{T_{l,i}, Y_i | \mathbf{W}}, P_{T_{l,i}, Y} | P_{\mathbf{W}})$ as a surrogate of the upper bound in Theorem 2. Following the proof of Theorem 2, we can obtain

$$\begin{aligned} \text{gen}(P_{\mathbf{W} | D_n}, P_{X, Y}) &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\ell(\mathbf{W}, X, Y) - \ell(\mathbf{W}, X_i, Y_i)] \\ &= \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\mathbb{E}[\ell(\mathbf{W}, X, Y) - \ell(\mathbf{W}, X_i, Y_i) | \mathbf{W}, \mathbf{W}_{\otimes L}, Y_i]] \\ &\stackrel{(a)}{\leq} \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\bar{\rho}_0(\mathbf{W}) W_1(P_{X_i | Y_i, \mathbf{w}, \mathbf{W}_{\otimes L}}, P_{X | Y, \mathbf{w}, \mathbf{W}_{\otimes L}})] \\ &\leq \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\bar{\rho}_0(\mathbf{W}) W_2(P_{X_i | Y_i, \mathbf{w}, \mathbf{W}_{\otimes L}}, P_{X | Y, \mathbf{w}, \mathbf{W}_{\otimes L}})], \end{aligned}$$

where (a) follows since $P_{Y_i | \mathbf{W}, \mathbf{W}_{\otimes L}} = P_Y = \text{Unif}\{-1, +1\}$. Similarly, we also have for all $l = 1, \dots, L$.

$$\text{gen}(P_{\mathbf{W} | D_n}, P_{X, Y}) \leq \frac{1}{n} \sum_{i=1}^n \mathbb{E}[\bar{\rho}_l(\mathbf{W}) W_2(P_{T_{l,i} | Y_i, \mathbf{w}, \mathbf{W}_{\otimes L}}, P_{T_{l,i} | Y, \mathbf{w}, \mathbf{W}_{\otimes L}})].$$

By plugging the $P_{T_{l,i} | Y_i, \mathbf{w}, \mathbf{W}_{\otimes L}}$ and $P_{T_{l,i} | Y, \mathbf{w}, \mathbf{W}_{\otimes L}}$ (c.f. (8)) into the upper bound, we have

$$\begin{aligned} & W_2(P_{T_{l,i} | Y_i, \mathbf{w}, \mathbf{W}_{\otimes L}}, P_{T_{l,i} | Y, \mathbf{w}, \mathbf{W}_{\otimes L}}) \\ &= \left(\|\mathbf{W}_{\otimes l}(\mathbf{W}_{\otimes L}^\top - \mu_0)\|^2 + \text{tr} \left(\frac{(n-1)\sigma_0^2}{n} (\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top) + \sigma_0^2 (\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top) \right. \right. \\ & \quad \left. \left. - 2 \left(\frac{(n-1)\sigma_0^4}{n} (\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top)^{\frac{1}{2}} (\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top) (\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top)^{\frac{1}{2}} \right) \right)^{\frac{1}{2}} \\ &= \sqrt{\|\mathbf{W}_{\otimes l}(\mathbf{W}_{\otimes L}^\top - \mu_0)\|^2 + \left(\frac{\sqrt{n-1}}{\sqrt{n}} - 1 \right)^2 \sigma_0^2 \text{tr}(\mathbf{W}_{\otimes l} \mathbf{W}_{\otimes l}^\top)} \\ &= \sqrt{\|\mathbf{W}_{\otimes l}(\mathbf{W}_{\otimes L}^\top - \mu_0)\|^2 + \frac{(\sqrt{n-1} - \sqrt{n})^2 \sigma_0^2}{n} \|\mathbf{W}_{\otimes l}\|_{\mathbb{F}}^2}. \end{aligned}$$

Similarly, we have

$$W_2(P_{X_i|Y_i, \mathbf{w}, \mathbf{w}_{\otimes L}}, P_{X|Y, \mathbf{w}, \mathbf{w}_{\otimes L}}) = \sqrt{\|(\mathbf{W}_{\otimes L}^\top - \mu_0)\|^2 + \frac{d_0(\sqrt{n-1} - \sqrt{n})^2 \sigma_0^2}{n}}.$$

Since the activation functions are all 1-Lipschitz, i.e., $\rho_l = 1$ for all $l = 1, \dots, L$ and the loss function $\tilde{\ell}$ is $4\sqrt{2}$ -Lipschitz, we have $\bar{\rho}_l(\mathbf{W}) = 4\sqrt{2}(1 \vee \prod_{j=l+1}^L s_{\max}(\mathbf{W}_j))$ for $l = 0, 1, \dots, L$.

For notational simplicity, let $\mathbf{W}_{\otimes 0} = \mathbf{W}_0 = \mathbf{I}_{d_0}$ and $r_0 = d_0$. Here we use $\|\cdot\|_{\mathbb{F}}$ of a vector to equivalently denote its Euclidean norm, with a slight abuse of notations. Then the generalization error is upper bounded by

$$\begin{aligned} & \text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y}) \\ & \leq \min \left\{ \min_{l=1, \dots, L} \mathbb{E} \left[\bar{\rho}_l(\mathbf{W}) \sqrt{\| \mathbf{W}_{\otimes l} (\mathbf{W}_{\otimes L}^\top - \mu_0) \|^2 + \frac{(\sqrt{n-1} - \sqrt{n})^2 \sigma_0^2}{n}} \| \mathbf{W}_{\otimes l} \|_{\mathbb{F}}^2 \right], \right. \\ & \quad \left. \mathbb{E} \left[\bar{\rho}_0(\mathbf{W}) \sqrt{\| (\mathbf{W}_{\otimes L}^\top - \mu_0) \|^2 + \frac{d_0(\sqrt{n-1} - \sqrt{n})^2 \sigma_0^2}{n}} \right] \right\} \\ & \stackrel{(a)}{\leq} \min \left\{ \min_{l=1, \dots, L} \mathbb{E} \left[\bar{\rho}_l(\mathbf{W}) \left(\| \mathbf{W}_{\otimes l} (\mathbf{W}_{\otimes L}^\top - \mu_0) \| + \frac{(\sqrt{n} - \sqrt{n-1}) \sigma_0}{\sqrt{n}} \| \mathbf{W}_{\otimes l} \|_{\mathbb{F}} \right) \right], \right. \\ & \quad \left. \mathbb{E} \left[\bar{\rho}_0(\mathbf{W}) \left(\| (\mathbf{W}_{\otimes L}^\top - \mu_0) \| + \frac{\sqrt{d_0}(\sqrt{n} - \sqrt{n-1}) \sigma_0}{\sqrt{n}} \right) \right] \right\} \\ & \stackrel{(b)}{\leq} \min \left\{ \min_{l=1, \dots, L} \mathbb{E} \left[\bar{\rho}_l(\mathbf{W}) \| \mathbf{W}_{\otimes l} \|_{\mathbb{F}} \left(\| (\mathbf{W}_{\otimes L}^\top - \mu_0) \| + \frac{(\sqrt{n} - \sqrt{n-1}) \sigma_0}{\sqrt{n}} \right) \right], \right. \\ & \quad \left. \mathbb{E} \left[\bar{\rho}_0(\mathbf{W}) \sqrt{d_0} \left(\| (\mathbf{W}_{\otimes L}^\top - \mu_0) \| + \frac{(\sqrt{n} - \sqrt{n-1}) \sigma_0}{\sqrt{n}} \right) \right] \right\} \\ & = \min_{l=0, \dots, L} \mathbb{E} \left[\bar{\rho}_l(\mathbf{W}) \| \mathbf{W}_{\otimes l} \|_{\mathbb{F}} \left(\| \mathbf{W}_{\otimes L}^\top - \mu_0 \| + \frac{(\sqrt{n} - \sqrt{n-1}) \sigma_0}{\sqrt{n}} \right) \right] \\ & \stackrel{(c)}{\leq} \min_{l=0, \dots, L} \mathbb{E} \left[\bar{\rho}_l(\mathbf{W})^2 \| \mathbf{W}_{\otimes l} \|_{\mathbb{F}}^2 \right]^{\frac{1}{2}} \mathbb{E} \left[\left(\| \mathbf{W}_{\otimes L}^\top - \mu_0 \| + \frac{(\sqrt{n} - \sqrt{n-1}) \sigma_0}{\sqrt{n}} \right)^2 \right]^{\frac{1}{2}} \\ & \stackrel{(d)}{\leq} \min_{l=0, \dots, L} \mathbb{E} \left[\bar{\rho}_l(\mathbf{W})^2 \| \mathbf{W}_{\otimes l} \|_{\mathbb{F}}^2 \right]^{\frac{1}{2}} \left(\frac{\sqrt{d_0} \sigma_0}{\sqrt{n}} + \frac{(\sqrt{n} - \sqrt{n-1}) \sigma_0}{\sqrt{n}} \right) \\ & = \frac{4\sqrt{2} \sigma_0 (\sqrt{d_0} + (\sqrt{n} - \sqrt{n-1}))}{\sqrt{n}} \min_{l=0, \dots, L} \mathbb{E} \left[\left(1 \vee \prod_{j=l+1}^L \| \mathbf{W}_j \| \right)^2 \| \mathbf{W}_{\otimes l} \|_{\mathbb{F}}^2 \right]^{\frac{1}{2}}, \end{aligned}$$

where (a) follows since $\sqrt{a^2 + b^2} \leq |a| + |b|$, (b) follows from the Cauchy-Schwarz inequality, (c) follows from the Hölder's inequality, and (d) follows from Minkowski's inequality and $\frac{n}{\sigma_0^2} \| \mathbf{W}_{\otimes L}^\top - \mu_0 \|^2 \sim \chi_{d_0}^2$. \square

Proof of Example 1. Since \mathbf{W}_l is (2×2) rotation matrix multiplied by a scalar factor C_l and $W_L = (0, C_L)$ is a row vector, $\| \mathbf{W}_l \| = C_l$ for $l = 1, \dots, L$. We have $\bar{\rho}_l(\mathbf{W}) = 4\sqrt{2}(1 \vee \prod_{j=l+1}^L \| \mathbf{W}_j \|) = 4\sqrt{2}(1 \vee \prod_{j=l+1}^L C_j)$ for $l = 0, 1, \dots, L$. \square

F. Proof of Lemma 6

Recall the definition of Dropout technique in (3). Fix any $l \in \{0, \dots, L\}$. Given \mathbf{W} , the Markov chain $T_l \rightarrow \tilde{T}_l \rightarrow T_{l+1}$ holds and we have the following SDPI:

$$\begin{aligned} & D_{\text{KL}}(P_{T_{l+1}, i|Y_i, \mathbf{w}} \| P_{T_{l+1}|Y, \mathbf{w}} | P_{Y_i, \mathbf{w}}) \\ & \leq \mathbb{E} \left[\mathbb{E}[\eta_{\text{KL}}(P_{T_{l+1}|\tilde{T}_l, \mathbf{w}}) D_{\text{KL}}(P_{\tilde{T}_l, i|Y_i, \mathbf{w}} \| P_{\tilde{T}_l|Y, \mathbf{w}}) | Y_i, \mathbf{W}] \right] \\ & \leq \eta_{\text{KL}}(P_{\tilde{T}_l|T_l}) \mathbb{E} \left[\mathbb{E}[\eta_{\text{KL}}(P_{T_{l+1}|\tilde{T}_l, \mathbf{w}}) D_{\text{KL}}(P_{T_{l+1}, i|Y_i, \mathbf{w}} \| P_{T_l|Y, \mathbf{w}}) | Y_i, \mathbf{W}] \right] \\ & \stackrel{(a)}{=} \eta_{\text{KL}}(P_{\tilde{T}_l|T_l}) D_{\text{KL}}(P_{T_{l+1}, i|Y_i, \mathbf{w}} \| P_{T_l|Y, \mathbf{w}} | P_{Y_i, \mathbf{w}}) \end{aligned}$$

where (a) follows from $\eta_{\text{KL}}(P_{T_{l+1}|\tilde{T}_l, \mathbf{w}}) = 1$ since $P_{T_{l+1}|\tilde{T}_l, \mathbf{w}}$ is a deterministic mapping. We observe that $P_{\tilde{T}_l|T_l}$ is composed of d_l parallel identical channels and thus, let $P_{\tilde{T}_l|T_l} := P_{\tilde{T}_l|T}^{\otimes d_l}$, where $P_{\tilde{T}_l|T}$ denotes the channel for $\tilde{T} = T \cdot E$ with $E \sim \text{Bern}(1 - \delta_l)$. Let $\tilde{\mathcal{T}}, \mathcal{T} \subset \mathbb{R}$ be the alphabets for \tilde{T}, T , respectively. In the following, to avoid any confusion, we denote $P_{\tilde{T}_l|T_l}^{\delta_l}$ as $P_{\tilde{T}_l|T}$ with cross probability $1 - \delta_l$.

a) *Upper bound for $\eta_{\text{KL}}(P_{\tilde{T}|T}^{\delta_l})$* : By applying [60, Theorem 5] d_l times, we have

$$\eta_{\text{KL}}(P_{\tilde{T}|T}^{\delta_l}) \leq 1 - (1 - \eta_{\text{KL}}(P_{\tilde{T}|T}))^{d_l}. \quad (9)$$

If $T \neq 0$, the channel $P_{\tilde{T}|T}^{\delta_l}$ is equivalent to the erasure channel with an erasure probability δ_l . From [68], which shows that the SDPI coefficient is achieved by binary inputs, we have

$$\eta_{\text{KL}}(P_{\tilde{T}|T}^{\delta_l}) = \sup_{t, t' \in \mathcal{T}} \sup_{P, Q \in \mathcal{P}(\{t, t'\})} \frac{D_{\text{KL}}(P_{\tilde{T}|T}^{\delta_l} \circ P \| P_{\tilde{T}|T}^{\delta_l} \circ Q)}{D_{\text{KL}}(P \| Q)} = \eta_{\text{KL}}(\text{BEC}(\delta_l)),$$

If $tt' \neq 0$, the channel $P_{\tilde{T}|T}^{\delta_l} : \{t, t'\} \rightarrow \{t, t', 0\}$ is equivalent to the binary the erasure channel with an erasure probability δ_l , denoted as $\text{BEC}(\delta_l)$. If $tt' = 0$, the channel $P_{\tilde{T}|T}^{\delta_l} : \{t, t'\} \rightarrow \{t, t'\}$ is equivalent to the Z-channel with a cross probability δ_l , denoted as $\text{ZC}(\delta_l)$. Thus, the SDPI coefficient is equal to

$$\eta_{\text{KL}}(P_{\tilde{T}|T}^{\delta_l}) = \eta_{\text{KL}}(\text{BEC}(\delta_l)) \vee \eta_{\text{KL}}(\text{ZC}(\delta_l)).$$

Given the $\text{BEC}(\delta_l)$, for any two binary input distributions $\text{Bern}(\alpha)$ and $\text{Bern}(\beta)$, we have

$$D_{\text{KL}}(\text{BEC}(\delta_l) \circ \text{Bern}(\alpha) \| \text{BEC}(\delta_l) \circ \text{Bern}(\beta)) = (1 - \delta_l) D_{\text{KL}}(\text{Bern}(\alpha) \| \text{Bern}(\beta)), \forall \alpha, \beta \in [0, 1].$$

Thus, $\eta_{\text{KL}}(\text{BEC}(\delta_l)) = 1 - \delta_l$.

The SDPI coefficient for $\text{ZC}(\delta_l)$ is upper bounded by the Dobrushin's coefficient:

$$\eta_{\text{KL}}(\text{ZC}(\delta_l)) \leq \eta_{\text{TV}}(\text{ZC}(\delta_l)) = \frac{1}{2}(|1 - \delta_l| + |0 - (1 - \delta_l)|) = 1 - \delta_l.$$

On the other hand, the lower bound for $\eta_{\text{KL}}(\text{ZC}(\delta_l))$ is derived as follows. Rewrite $\eta_{\text{KL}}(\text{ZC}(\delta_l))$ in terms of mutual informations:

$$\eta_{\text{KL}}(\text{ZC}(\delta_l)) = \sup_{P_{U, X}: U \rightarrow T \rightarrow \tilde{T}} \frac{I(U; \tilde{T})}{I(U; T)}$$

where we assume $T, \tilde{T} \in \{0, 1\}$ without loss of generality. Let $U \sim \text{Bern}(\frac{1}{2})$, $P_{T|U=0} = \text{Bern}(\alpha)$ and $P_{T|U=1} = \text{Bern}(1 - \alpha)$ for some $\alpha \in [0, 1]$. We have $I(U; T) = H(U) - H(U|T) = 1 - h_b(\alpha)$, where $h_b(\alpha) = H(\text{Bern}(\alpha))$ and

$$I(U; \tilde{T}) = 1 - \frac{1 - \delta_l}{2} h_b(\alpha) - \frac{1 + \delta_l}{2} h_b\left(\frac{\alpha + (1 - \alpha)\delta_l}{1 + \delta_l}\right).$$

When $\alpha \rightarrow \frac{1}{2}$, by applying the L'Hôpital's rule, we have

$$\eta_{\text{KL}}(\text{ZC}(\delta_l)) \geq \frac{I(U; T)}{I(U; \tilde{T})} = \frac{1 - \frac{1 - \delta_l}{2} h_b(\alpha) - \frac{1 + \delta_l}{2} h_b\left(\frac{\alpha + (1 - \alpha)\delta_l}{1 + \delta_l}\right)}{1 - h_b(\alpha)} \rightarrow 1 - \delta_l.$$

Therefore, $\eta_{\text{KL}}(\text{ZC}(\delta_l)) = 1 - \delta_l$ and $\eta_{\text{KL}}(P_{\tilde{T}|T}) = \max\{\eta_{\text{KL}}(\text{BEC}(\delta_l)), \eta_{\text{KL}}(\text{ZC}(\delta_l))\} = 1 - \delta_l$. By plugging $\eta_{\text{KL}}(P_{\tilde{T}|T})$ back to (9), we get

$$\eta_{\text{KL}}(P_{\tilde{T}|T}^{\delta_l}) \leq 1 - \delta_l^{d_l}. \quad (10)$$

b) *Lower bound for $\eta_{\text{KL}}(P_{\tilde{T}|T}^{\delta_l})$* : Let $\mathcal{T}_l \subset \mathbb{R}^{d_l}$ be the alphabet of T_l and $\mathcal{T}_l^{-0} \subset (\mathbb{R} \setminus \{0\})^{d_l}$. From [68], we have

$$\eta_{\text{KL}}(P_{\tilde{T}|T}^{\delta_l}) \geq \frac{1}{2} \sup_{t, t' \in \mathcal{T}_l} D_{\text{H}^2}(P_{\tilde{T}_l|T_l=t}^{\delta_l}, P_{\tilde{T}_l|T_l=t'}^{\delta_l}) \geq \frac{1}{2} \sup_{t, t' \in \mathcal{T}_l^{-0}} D_{\text{H}^2}(P_{\tilde{T}_l|T_l=t}^{\delta_l}, P_{\tilde{T}_l|T_l=t'}^{\delta_l})$$

where D_{H^2} denotes the Hellinger distance. Given any $t \in \mathcal{T}_l^{-0}$, the output \tilde{T}_l follows the binomial distribution $\text{Binom}(d_l, \delta_l)$ on the alphabet $\tilde{\mathcal{T}}_l(t)$, where $\{\mathbf{0}\} \subset \tilde{\mathcal{T}}_l(t)$. It is straightforward that $\sup_{t, t' \in \mathcal{T}_l^{-0}} D_{\text{H}^2}(P_{\tilde{T}_l|T_l=t}^{\delta_l}, P_{\tilde{T}_l|T_l=t'}^{\delta_l}) = \sup_{t, t' \in \mathcal{T}_l^{-0}: \text{Hamming}(t, t')=d_l} D_{\text{H}^2}(P_{\tilde{T}_l|T_l=t}^{\delta_l}, P_{\tilde{T}_l|T_l=t'}^{\delta_l})$. Note that for any $t, t' \in \mathcal{T}_l^{-0}$ such that $\text{Hamming}(t, t') = d_l$, we have $(\tilde{\mathcal{T}}_l(t) \setminus \{\mathbf{0}\}) \cap (\tilde{\mathcal{T}}_l(t') \setminus \{\mathbf{0}\}) = \emptyset$. Then the corresponding Hellinger distance is given by

$$\begin{aligned} D_{\text{H}^2}(P_{\tilde{T}_l|T_l=t}^{\delta_l}, P_{\tilde{T}_l|T_l=t'}^{\delta_l}) &= \sum_{u \in \tilde{\mathcal{T}}_l(t) \cup \tilde{\mathcal{T}}_l(t')} \left(\sqrt{P_{\tilde{T}_l|T_l=t}^{\delta_l}(u)} - \sqrt{P_{\tilde{T}_l|T_l=t'}^{\delta_l}(u)} \right)^2 \\ &= \left(\sqrt{P_{\tilde{T}_l|T_l=t}^{\delta_l}(\mathbf{0})} - \sqrt{P_{\tilde{T}_l|T_l=t'}^{\delta_l}(\mathbf{0})} \right)^2 + \sum_{u \in \tilde{\mathcal{T}}_l(t) \setminus \{\mathbf{0}\}} P_{\tilde{T}_l|T_l=t}^{\delta_l}(u) + \sum_{u \in \tilde{\mathcal{T}}_l(t') \setminus \{\mathbf{0}\}} P_{\tilde{T}_l|T_l=t'}^{\delta_l}(u) \end{aligned}$$

$$\begin{aligned}
&\stackrel{(a)}{=} 2 \sum_{k=0}^{d_l-1} \binom{d_l}{k} \delta_l^k (1-\delta_l)^{n-k} \\
&= 2(1-\delta_l^{d_l}),
\end{aligned}$$

where (a) follows since $P_{\tilde{T}_l|T_l=t}^{\delta_l}(\mathbf{0}) = P_{\tilde{T}_l|T_l=t'}^{\delta_l}(\mathbf{0}) = \delta_l^{d_l}$. The lower bound for $\eta_{\text{KL}}(P_{\tilde{T}_l|T_l}^{\delta_l})$ is then given by

$$\eta_{\text{KL}}(P_{\tilde{T}_l|T_l}^{\delta_l}) \geq 1 - \delta_l^{d_l}. \quad (11)$$

By combining (10) and (11), we conclude that $\eta_{\text{KL}}(P_{\tilde{T}_l|T_l}^{\delta_l}) = 1 - \delta_l^{d_l}$.

G. Proof of Theorem 7

From the proof of Lemma 6 in Appendix F, we have

$$\text{D}_{\text{KL}}(P_{T_{l+1},i|Y_i,\mathbf{W}} \| P_{T_{l+1}|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}}) \leq (1 - \delta_l^{d_l}) \text{D}_{\text{KL}}(P_{T_{l,i}|Y_i,\mathbf{W}} \| P_{T_l|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}}), \quad l = 0, \dots, L.$$

Recall the upper bound (2) in Theorem 1:

$$\begin{aligned}
|\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y})| &\leq \frac{1}{n} \sum_{i=1}^n \sqrt{2\sigma^2 \text{D}_{\text{KL}}(P_{T_{L,i}|Y_i,\mathbf{W}} \| P_{T_L|Y,\mathbf{W}} | P_{\mathbf{W}})} \\
&= \frac{1}{n} \sum_{i=1}^n \sqrt{2\sigma^2 (\text{D}_{\text{KL}}(P_{T_{L,i}|Y_i,\mathbf{W}} \| P_{T_L|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}}) + I(Y_i; \mathbf{W}))}, \quad (12)
\end{aligned}$$

where (12) follows since for any $l \in [L]$,

$$\begin{aligned}
\text{D}_{\text{KL}}(P_{T_{L,i}|Y_i,\mathbf{W}} \| P_{T_L|Y,\mathbf{W}} | P_{\mathbf{W}}) &= \text{D}_{\text{KL}}(P_{T_{L,i}|Y_i,\mathbf{W}} P_{Y_i|\mathbf{W}} \| P_{T_L|Y,\mathbf{W}} P_{Y|\mathbf{W}} | P_{\mathbf{W}}) \\
&= \text{D}_{\text{KL}}(P_{T_{L,i}|Y_i,\mathbf{W}} \| P_{T_L|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}}) + \text{D}_{\text{KL}}(P_{Y_i|\mathbf{W}} \| P_Y | P_{\mathbf{W}}) \\
&= \text{D}_{\text{KL}}(P_{T_{L,i}|Y_i,\mathbf{W}} \| P_{T_L|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}}) + I(Y_i; \mathbf{W}).
\end{aligned}$$

It can be observed that the data-processing inequality is only applied to $\text{D}_{\text{KL}}(P_{T_{L,i}|Y_i,\mathbf{W}} \| P_{T_L|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}})$ since Y_i is not processed. Furthermore, since $Y_i \in \mathcal{Y} = [K]$ is a discrete random variable, we have $I(Y_i; \mathbf{W}) \leq H(Y_i) \leq \log K$.

By induction, if we have L layers

$$\begin{aligned}
\text{D}_{\text{KL}}(P_{T_{L,i}|Y_i,\mathbf{W}} \| P_{T_L|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}}) &\leq (1 - \delta_{L-1}^{d_{L-1}}) \text{D}_{\text{KL}}(P_{T_{L-1,i}|Y_i,\mathbf{W}} \| P_{T_{L-1}|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}}) \\
&\leq \prod_{l=L-2}^{L-1} (1 - \delta_l^{d_l}) \text{D}_{\text{KL}}(P_{T_{L-2,i}|Y_i,\mathbf{W}} \| P_{T_{L-2}|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}}) \\
&\vdots \\
&\leq \prod_{l=0}^{L-1} (1 - \delta_l^{d_l}) \text{D}_{\text{KL}}(P_{T_{0,i}|Y_i,\mathbf{W}} \| P_{T_0|Y,\mathbf{W}} | P_{Y_i,\mathbf{W}}) = \prod_{l=1}^L (1 - \delta_l^{d_l}) I(X_i; \mathbf{W} | Y_i). \quad (13)
\end{aligned}$$

The proof is completed by plugging (13) into (12).

H. Proof of Lemma 8

In this proof, we show that the mutual information terms decay as the Dropout probability δ_0 increases. Let $\mathcal{J}(k) \subseteq [d]$ be a set of coordinate indices, $|\mathcal{J}(k)| = k \in [d]$, and $X_i(\mathcal{J}(k)) = \{X_i(j)\}_{j \in \mathcal{J}(k)}$ be a subset of coordinates of X_i . Some coordinates of the input X_i are randomly deactivated as 0 before being passed into the neural network. Consider an independent binary mask vector $E_t \sim \text{Bern}(1 - \delta_0)^{\otimes d_0}$ for the input X_i . It can be observed that $X_i \odot E_t$ is a sufficient statistic of X_i for \mathbf{W} . We decompose the mutual information term as

$$\begin{aligned}
I(X_i, Y_i; \mathbf{W}) &= I(X_i \odot E_t, Y_i; \mathbf{W}) \\
&\stackrel{(a)}{=} \sum_{e \in \{0,1\}^{d_0}} P_{E_t}(e) I(X_i \odot e, Y_i; \mathbf{W})
\end{aligned}$$

where (a) follows since E_t is independent of (X_i, Y_i) . Let $e_{\mathcal{J}(k)} \in \{0,1\}^{d_0}$ denote the binary vector with non-zeros elements at indices $\mathcal{J}(k)$, for $k = 0, \dots, d_0$. Then $P_{E_t}(e_{\mathcal{J}(k)}) = \delta_0^{d_0-k} (1 - \delta_0)^k$.

Let $I_k^{(i)} = \max_{\mathcal{J}(k) \subseteq [d], |\mathcal{J}(k)|=k} I(X_i(\mathcal{J}(k)), Y_i; \mathbf{W})$, for $k = 0, \dots, d_0$. We have $0 = I_0^{(i)} \leq I_1^{(i)} \leq \dots \leq I_{d_0}^{(i)} = I(X_i, Y_i; \mathbf{W})$. Thus, we have

$$\begin{aligned}
& \sum_{e \in \{0,1\}^{d_0}} P_{E_t}(e) I(X_i \odot e, Y_i; \mathbf{W}) \\
&= \sum_{k=0}^{d_0} \sum_{\mathcal{J}(k) \subseteq [d]} \delta_0^{d_0-k} (1-\delta_0)^k I(X_i(\mathcal{J}(k)), Y_i; \mathbf{W}) \\
&= \sum_{k=0}^{d_0} \delta_0^{d_0-k} (1-\delta_0)^k \sum_{\mathcal{J}(k) \subseteq [d]} I(X_i(\mathcal{J}(k)), Y_i; \mathbf{W}) \\
&\leq \sum_{k=0}^{d_0} \delta_0^{d_0-k} (1-\delta_0)^k \binom{d_0}{k} I_k^{(i)} \\
&= \sum_{k \leq \lfloor \frac{d_0}{2} \rfloor} \delta_0^{d_0-k} (1-\delta_0)^k \binom{d_0}{k} I_k^{(i)} + \sum_{k > \lfloor \frac{d_0}{2} \rfloor} \delta_0^{d_0-k} (1-\delta_0)^k \binom{d_0}{k} I_k^{(i)}. \tag{14}
\end{aligned}$$

As δ_0 increases, the increase in the first term of (14) is less than the decrease in the second term of (14) since $I_k^{(i)}$ is increasing in k . Thus, (14) monotonically decreases as δ_0 increases. Note that when δ_0 increases to 1, the upper bound (14) shrinks to 0.

Consider a special case when $I_k^{(i)}$ is proportional to k , i.e., $I_k^{(i)} = \alpha k$ for some constant $\alpha \in \mathbb{R}_{>0}$. Then (14) can be rewritten as

$$\sum_{e \in \{0,1\}^{d_0}} P_{E_t}(e) I(X_i \odot e, Y_i; \mathbf{W}) \leq \sum_{k=0}^{d_0} \binom{d_0}{k} \delta_0^{d_0-k} (1-\delta_0)^k \alpha k = \alpha d_0 (1-\delta_0)$$

which decays linearly in δ_0 .

I. Proof of Lemma 9

From (4), for $i = 1, \dots, d_l$, the i^{th} element of T_l is given by

$$T_l(i) = \phi_l \left(\sum_{j=1}^{d_{l-1}} \mathbf{W}_l(i, j) \mathbf{E}_l(i, j) T_{l-1}(j) \right) =: \phi_l \left(\sum_{j=1}^{d_{l-1}} \mathbf{W}_l(i, j) \tilde{T}_{l-1}(i, j) \right),$$

where \tilde{T}_{l-1} is a $d_l \times d_{l-1}$ matrix. Thus, given the network weights \mathbf{W} , at any layer l , we have d_l Markov chains: for $i = 1, \dots, d_l$, $T_{l-1} \rightarrow \tilde{T}_{l-1}(i, \cdot) \rightarrow T_l(i)$, where $\tilde{T}_{l-1}(i, \cdot)$ denotes the i^{th} row of \tilde{T}_{l-1} . We observe that for any $i \in [d_l]$, $P_{T_l(i) | \tilde{T}_{l-1}(i, \cdot), \mathbf{W}}$ is a deterministic mapping, while $P_{\tilde{T}_{l-1}(i, \cdot) | T_{l-1}}$ is composed of d_{l-1} parallel independent Z-channels, i.e., $P_{\tilde{T}_{l-1}(i, \cdot) | T_{l-1}} = \prod_{j=1}^{d_{l-1}} P_{\tilde{T} | T}^{\delta_{l-1, i, j}}$, with $P_{\tilde{T} | T}^{\delta_{l-1, i, j}}$ defined in Appendix F. Since each row of \tilde{T}_{l-1} are independent with each other given T_{l-1} , $P_{\tilde{T}_{l-1} | T_{l-1}} = \prod_{i=1}^{d_l} P_{\tilde{T}_{l-1}(i, \cdot) | T_{l-1}} = \prod_{i=1}^{d_l} \prod_{j=1}^{d_{l-1}} P_{\tilde{T} | T}^{\delta_{l-1, i, j}}$.

The the following SDPI holds: for any training data sample index $i \in [n]$,

$$\begin{aligned}
& D_{\text{KL}}(P_{T_{l+1, i} | Y_i, \mathbf{W}} \| P_{T_{l+1} | Y, \mathbf{W}} | P_{Y_i, \mathbf{W}}) \\
&\leq \mathbb{E} [\mathbb{E} [\eta_{\text{KL}}(P_{T_{l+1} | \tilde{T}_l, \mathbf{W}}) D_{\text{KL}}(P_{\tilde{T}_l, i | Y_i, \mathbf{W}} \| P_{\tilde{T}_l | Y, \mathbf{W}}) | Y_i, \mathbf{W}]] \\
&\leq \mathbb{E} [\mathbb{E} [\eta_{\text{KL}}(P_{T_{l+1} | \tilde{T}_l, \mathbf{W}}) \eta_{\text{KL}}(P_{\tilde{T}_l | T_l}) D_{\text{KL}}(P_{T_l, i | Y_i, \mathbf{W}} \| P_{T_l | Y, \mathbf{W}}) | Y_i, \mathbf{W}]] \\
&\stackrel{(a)}{=} \eta_{\text{KL}} \left(\prod_{j=1}^{d_{l+1}} \prod_{k=1}^{d_l} P_{\tilde{T} | T}^{\delta_{l, j, k}} \right) D_{\text{KL}}(P_{T_l, i | Y_i, \mathbf{W}} \| P_{T_l | Y, \mathbf{W}} | P_{Y_i, \mathbf{W}}) \\
&\stackrel{(b)}{\leq} \left(1 - \prod_{j=1}^{d_{l+1}} \left(1 - \eta_{\text{KL}} \left(\prod_{k=1}^{d_l} P_{\tilde{T} | T}^{\delta_{l, j, k}} \right) \right) \right) D_{\text{KL}}(P_{T_l, i | Y_i, \mathbf{W}} \| P_{T_l | Y, \mathbf{W}} | P_{Y_i, \mathbf{W}})
\end{aligned}$$

where (a) follows from $\eta_{\text{KL}}(P_{T_{l+1} | \tilde{T}_l, \mathbf{W}}) = 1$ since $P_{T_{l+1} | \tilde{T}_l, \mathbf{W}}$ is a deterministic mapping, (b) follows from [60, Theorem 5]. By applying [60, Theorem 5] again, we have

$$\eta_{\text{KL}} \left(\prod_{k=1}^{d_l} P_{\tilde{T} | T}^{\delta_{l, j, k}} \right) \leq 1 - \prod_{k=1}^{d_l} \delta_{l, j, k}, \quad \text{and} \quad 1 - \prod_{j=1}^{d_{l+1}} \left(1 - \eta_{\text{KL}} \left(\prod_{k=1}^{d_l} P_{\tilde{T} | T}^{\delta_{l, j, k}} \right) \right) \leq 1 - \prod_{j=1}^{d_{l+1}} \prod_{k=1}^{d_l} \delta_{l, j, k}.$$

J. Proof of Lemma 11

When $Y = g(X) + \epsilon N$, where N is an independent noise and $\epsilon > 0$ is a constant controlling the signal-to-noise ratio (SNR), then $P_{Y|X} = P_{g(X) + \epsilon N}$. The Dobrushin's coefficient is given by

$$\eta_{\text{TV}}(P_{Y|X}) = \sup_{x, x' \in \mathbb{R}^{d_x}} \|P_{g(x) + \epsilon N} - P_{g(x') + \epsilon N}\|_{\text{TV}}.$$

Let N be a Gaussian noise generated from $\mathcal{N}(\mathbf{0}, \mathbf{I}_{d_y})$. Denote the vector function $g(\cdot)$ by $g(\cdot) = (g_1(\cdot), \dots, g_{d_y}(\cdot))$. Following the proof in [69] and the total variation distance between two Gaussians with the same covariance matrix in [70, Theorem 1], we have for any $x, x' \in \mathcal{X}$,

$$\begin{aligned} D_{\text{TV}}(P_{g(x) + \epsilon N}, P_{g(x') + \epsilon N}) &= \|\mathcal{N}(g(x), \epsilon^2 \mathbf{I}_{d_y}) - \mathcal{N}(g(x'), \epsilon^2 \mathbf{I}_{d_y})\|_{\text{TV}} \\ &= 1 - 2\text{Q}\left(\frac{\|g(x) - g(x')\|}{2\epsilon}\right) \\ &= 1 - 2\text{Q}\left(\frac{\sqrt{\sum_{i=1}^{d_y} (g_i(x) - g_i(x'))^2}}{2\epsilon}\right) \\ &\leq 1 - 2\text{Q}\left(\frac{\sqrt{d_y}(\|g(x)\|_{\infty}^2 + \|g(x')\|_{\infty}^2)}{2\epsilon}\right) \\ &\leq 1 - 2\text{Q}\left(\frac{\sqrt{2d_y}\|g\|_{\infty}}{2\epsilon}\right) \end{aligned}$$

where $\text{Q}(x) = \int_x^{\infty} \frac{1}{\sqrt{2\pi}} e^{-t^2/2} dt$ is the Gaussian complimentary CDF. Finally, we have

$$\eta_f(P_{Y|X}) \leq \eta_{\text{TV}}(P_{Y|X}) \leq 1 - 2\text{Q}\left(\frac{\sqrt{2d_y}\|g\|_{\infty}}{2\epsilon}\right).$$

K. Proof of Theorem 12

Among the commonly used activation functions, the following functions and their gradients are bounded: for any $u \in \mathbb{R}$,

- sigmoid function: $\text{sigmoid}(u) = \frac{1}{1+e^{-u}} \in [0, 1]$, $\text{sigmoid}'(u) = \frac{e^{-u}}{(1+e^{-u})^2} \in [0, 1]$.
- softmax function: $\text{softmax}(u)_i = \frac{e^{u_i}}{\sum_j e^{u_j}} \in [0, 1]$, $\text{softmax}'(u)_i = \text{softmax}(u)_i(1 - \text{softmax}(u)_i) \in [0, 1]$.
- tanh function: $\tanh u = \frac{e^u - e^{-u}}{e^u + e^{-u}} \in [-1, 1]$, $\tanh'(u) = 1 - \tanh^2 u \in [0, 1]$.

As shown in (5), that is, $\tilde{T}_l = g_{\mathbf{W}_l}(\tilde{T}_{l-1}) + \epsilon_l Z_l = \phi_l(\mathbf{W}_l \tilde{T}_{l-1}) + \epsilon_l Z_l$ and from Lemma 11, the Dobrushin's coefficient at the l^{th} layer is upper bounded by

$$\eta_{\text{TV}}(P_{\tilde{T}_l|\tilde{T}_{l-1}, \mathbf{W}}) \leq 1 - 2\text{Q}\left(\frac{\sqrt{2d_l}\|g_{\mathbf{W}_l}\|_{\infty}}{2\epsilon_l}\right) = 1 - 2\text{Q}\left(\frac{\sqrt{2d_l}\|\phi_l\|_{\infty}}{2\epsilon_l}\right),$$

where $\|\phi_l\|_{\infty} = 1$ for $\phi_l \in \{\text{sigmoid}, \text{softmax}, \text{tanh}\}$.

In the following proof, at no risk of confusion, we let $T_l = \tilde{T}_l$ and $T_{l,i} = \tilde{T}_{l,i}$, for simplicity. Conditioned on \mathbf{W} , $P_{T_{l-1,i}|Y_i, \mathbf{W}} \neq P_{T_{l-1}|Y, \mathbf{W}}$ but $P_{T_{l,i}|T_{l-1,i}, \mathbf{W}} = P_{T_l|T_{l-1}, \mathbf{W}}$. Thus, we have

$$\begin{aligned} D_{\text{KL}}(P_{T_{l,i}|Y_i, \mathbf{W}} \| P_{T_l|Y, \mathbf{W}} | P_{Y_i, \mathbf{W}}) \\ &= D_{\text{KL}}(P_{T_{l,i}|T_{l-1,i}, \mathbf{W}} \circ P_{T_{l-1,i}|Y_i, \mathbf{W}} \| P_{T_l|T_{l-1}, \mathbf{W}} \circ P_{T_{l-1}|Y, \mathbf{W}} | P_{Y_i, \mathbf{W}}) \\ &\leq \left(1 - 2\text{Q}\left(\frac{\sqrt{2d_l}\|\phi_l\|_{\infty}}{2\epsilon_l}\right)\right) D_{\text{KL}}(P_{T_{l-1,i}|Y_i, \mathbf{W}} \| P_{T_{l-1}|Y, \mathbf{W}} | P_{Y_i, \mathbf{W}}). \end{aligned}$$

By induction, if we have L layers

$$\begin{aligned} D_{\text{KL}}(P_{T_{L,i}|Y_i, \mathbf{W}} \| P_{T_L|Y, \mathbf{W}} | P_{Y_i, \mathbf{W}}) &\leq \left(1 - 2\text{Q}\left(\frac{\sqrt{2d_L}\|\phi_L\|_{\infty}}{2\epsilon_L}\right)\right) D_{\text{KL}}(P_{T_{L-1,i}|Y_i, \mathbf{W}} \| P_{T_{L-1}|Y, \mathbf{W}} | P_{Y_i, \mathbf{W}}) \\ &\vdots \\ &\leq \prod_{l=1}^L \left(1 - 2\text{Q}\left(\frac{\sqrt{2d_l}\|\phi_l\|_{\infty}}{2\epsilon_l}\right)\right) D_{\text{KL}}(P_{T_{0,i}|Y_i, \mathbf{W}} \| P_{T_0|Y, \mathbf{W}} | P_{Y_i, \mathbf{W}}) \\ &= \prod_{l=1}^L \left(1 - 2\text{Q}\left(\frac{\sqrt{2d_l}\|\phi_l\|_{\infty}}{2\epsilon_l}\right)\right) I(X_i; \mathbf{W} | Y_i). \end{aligned} \tag{15}$$

By combining (15) with (12), the expected generalization error is upper bounded by

$$|\text{gen}(P_{\mathbf{W}|D_n}, P_{X,Y})| \leq \frac{\sqrt{2\sigma^2}}{n} \sum_{i=1}^n \sqrt{\prod_{l=1}^L \left(1 - 2\text{Q}\left(\frac{\sqrt{2d_l}\|\phi_l\|_\infty}{2\epsilon_l}\right)\right)} \mathbb{I}(X_i; \mathbf{W}|Y_i) + \mathbb{I}(Y_i; \mathbf{W}).$$

The proof is thus completed.