

OGLE-2018-BLG-0971, MOA-2023-BLG-065, and OGLE-2023-BLG-0136: Microlensing events with prominent orbital effects

Cheongho Han¹, Andrzej Udalski², Ian A. Bond³, Chung-Uk Lee⁴, Andrew Gould^{5,6}

(Leading authors)

Michael D. Albrow⁷, Sun-Ju Chung⁴, Kyu-Ha Hwang⁴, Youn Kil Jung⁴, Hyoun-Woo Kim⁴, Yoon-Hyun Ryu⁴,
Yossi Shvartzvald⁸, In-Gu Shin⁹, Jennifer C. Yee⁹, Hongjing Yang¹⁰, Weicheng Zang^{9,10}, Sang-Mok Cha^{4,11},
Doeon Kim¹, Dong-Jin Kim⁴, Seung-Lee Kim⁴, Dong-Joo Lee⁴, Yongseok Lee^{4,11}, Byeong-Gon Park⁴,

Richard W. Pogge⁶

(The KMTNet Collaboration)

Przemek Mróz², Michał K. Szymański², Jan Skowron², Radosław Poleski², Igor Soszyński², Paweł Pietrukowicz²,
Szymon Kozłowski², Krzysztof A. Rybicki^{2,8}, Patryk Iwanek², Krzysztof Ulaczyk¹², Marcin Wrona²,

Mariusz Gromadzki², Mateusz J. Mróz²

(The OGLE Collaboration)

Fumio Abe¹³, Richard Barry¹⁴, David P. Bennett^{14,15}, Aparna Bhattacharya^{13,14}, Hirosame Fujii¹³, Akihiko Fukui^{16,17},
Ryusei Hamada¹⁸, Yuki Hirao¹⁸, Stela Ishitani Silva^{15,19}, Yoshitaka Itow¹³, Rintaro Kirikawa¹⁸, Naoki Koshimoto²⁰,
Yutaka Matsubara¹³, Shota Miyazaki¹⁸, Yasushi Muraki¹³, Greg Olmschenk¹⁴, Clément Ranc²¹,

Nicholas J. Rattenbury²², Yuki Satoh¹⁸, Takahiro Sumi¹⁸, Daisuke Suzuki¹⁸, Mio Tomoyoshi¹⁸, Paul J. Tristram²³,
Aikaterini Vandenbroucke^{14,15}, Hibiki Yama¹⁸, and Kansuke Yamashita¹⁸

(The MOA Collaboration)

(Affiliations can be found after the references)

Received ; accepted

ABSTRACT

Aims. We undertake a project to reexamine microlensing data gathered from high-cadence surveys. The aim of the project is to reinvestigate lensing events with light curves exhibiting intricate anomaly features associated with caustics, yet lacking prior proposed models to explain these features.

Methods. Through detailed reanalyses considering higher-order effects, we identify that accounting for orbital motions of lenses is vital in accurately explaining the anomaly features observed in the light curves of the lensing events OGLE-2018-BLG-0971, MOA-2023-BLG-065, and OGLE-2023-BLG-0136.

Results. We estimate the masses and distances to the lenses by conducting Bayesian analyses using the lensing parameters of the newly found lensing solutions. From these analyses, we identify that the lenses of the events OGLE-2018-BLG-0971 and MOA-2023-BLG-065 are binaries composed of M dwarfs, while the lens of OGLE-2023-BLG-0136 is likely to be a binary composed of an early K-dwarf primary and a late M-dwarf companion. For all lensing events, the probability of the lens residing in the bulge is considerably higher than that of it being located in the disk.

Key words. Gravitational lensing: micro

1. Introduction

In general, light curves of microlensing events are modeled by assuming a rectilinear relative motion between the lens and the source. However, deviations from this assumption arise due to accelerations affecting the motion of either the observer, lens, or source. For instance, an observer experiences acceleration due to the Earth's orbital motion around the Sun, known as microlens-parallax effects (Gould 1992, 2000). Additionally, if a source of a lensing event is part of a binary system in which two stars orbit a common barycenter, the source's motion also undergoes acceleration (Han & Gould 1997; Rahvar & Dominik 2009). Similarly, the lens's orbital motion induces acceleration, causing deviations

from a rectilinear relative motion between the lens and source, known as lens-orbital effects.

There have been instances of lensing events in which accounting for lens orbital motions was crucial for accurately interpreting observed lensing light curves. MACHO 97-BLG-41 (Alcock et al. 2000) notably marked the first binary-lens single-source (2L1S) system displaying significant deviations from the assumption of a static binary configuration. Initially, these deviations were attributed to the presence of a third body of the lens, specifically a circumbinary planet (Bennett et al. 1999). However, Albrow et al. (2000), based on the analysis using an independent data set, later proposed a solution involving an orbiting binary lens. The controversy was definitively resolved by Jung et al. (2018), who, through a direct comparison of the

Table 1. Coordinates of events.

Event	(RA, Dec) _{J2000}	(<i>l</i> , <i>b</i>)	Other ID references
OGLE-2018-BLG-0971	(17:59:01.63, -28:13:42.20)	(2°:1082, -2°:1727)	MOA-2018-BLG-173, KMT-2018-BLG-2336
MOA-2023-BLG-065	(18:00:35.68, -29:13:23.95)	(1°:4140, -2°:9645)	KMT-2023-BLG-2430
OGLE-2023-BLG-0136	(18:09:16.97, -30:36:47.00)	(1°:1132, -5°:2851)	KMT-2023-BLG-2849

two models using a combined data set, found that the orbiting binary-lens interpretation is preferred over the circumbinary planet model.

OGLE-2006-BLG-109 was the second instance in which lens orbital effects played a crucial role in providing accurate explanations for the lensing light curve (Gaudi et al. 2008; Bennett et al. 2010). The light curve of this event exhibited a complex anomaly pattern, comprising multiple distinctive features. A comprehensive understanding of these anomalies was achieved only after accounting for combined higher-order effects resulting from the parallactic motion of Earth and the orbital motion of the lens. From the analysis considering these higher-order effects, the lens was proven to be the first double planet system discovered with the gravitational microlensing method.

OGLE-2005-BLG-018 marked the third instance in which the significance of lens orbital effects was established. The light curve of the event displayed multiple anomaly features, comprising two neighboring strong anomalies and a comparatively weak anomaly positioned apart from the stronger ones. Although a model based on a static binary lens configuration could approximately explain the two strong anomalies, accurately describing the separate weak anomaly was challenging. As a result, this event had not been previously addressed until Shin et al. (2011) revisited it and demonstrated that accounting for the lens orbital motion was essential for accurately describing all the anomaly features in the lensing light curve.

Prompted by the work of Shin et al. (2011), Park et al. (2013) revisited microlensing data available until then and conducted thorough analyses of binary-lens events. They focused on cases in which static binary models fell short in accurately describing observed light curves. Through these analyses, they revealed that the substantial residuals of the light curves of two lensing events OGLE-2006-BLG-277 and OGLE-2012-BLG-0031 from static 2L1S models were predominantly attributed to the influence of lens orbital motion.

OGLE-2009-BLG-020 was identified as a binary-lens event where orbital motion was initially predicted through light curve analysis (Skowron et al. 2011). Subsequent 3.5 years of radial velocity monitoring confirmed an orbit consistent with the predictions derived from the microlensing light curve analysis (Yee et al. 2016).

In the case OGLE-2013-BLG-0723, the lensing light curve was initially interpreted by a triple-lens model, in which the lens consisted of a Venus-mass planet and binary brown-dwarf host (Udalski et al. 2015a). Later, Han et al. (2016a) reexamined the event, incorporating lens-orbital effects, and proposed a revised interpretation involving two-body lens instead of the previously suggested three-body lens solution. This updated model provided a notably better fit to the observed light curve.

Gaia16aye, detected toward the northern Galactic disk field, was a binary microlensing event and stood among the earliest instances detected from the alerts issued by the Gaia space mission. The brightening of the source induced by lensing endured for more than two years, and thus it was crucial to include the orbital motion of the lens for the precise description of the lensing light curve (Wyrzykowski et al. 2020).

The light curve of the lensing event KMT-2021-BLG-0322 exhibited multiple sets of caustic-crossing features. The overall features of the light curve were approximately described by a 2L1S model, but the model left substantial residuals. From the reanalysis, Han et al. (2021) found that the residuals could be explained either by considering a non-rectilinear lens-source motion caused by the combination of microlens-parallax and lens-orbital effects or by adding an additional low-mass companion to the binary lens, and hence three lens components (3L1S system). The degeneracy between the higher-order 2L1S model and the 3L1S model was very severe, making it difficult to single out a correct solution based on the photometric data. This degeneracy was known before for two previous events (MACHO-97-BLG-41 and OGLE-2013-BLG-0723), which led to the false detections of planets in binary systems, and thus the identification of the degeneracy for the event illustrated that this degeneracy could be common.

In this study, we present comprehensive analyses on three 2L1S lensing events: OGLE-2018-BLG-0971, MOA-2023-BLG-065, and OGLE-2023-BLG-0136. These events bear resemblances, displaying anomalies in their light curves with intricate and complex features that prove challenging to interpret using static binary-lens models. We demonstrate the importance of considering orbital motions of lenses in precisely describing the observed anomaly features in the lensing light curves.

2. Event selections and data

We undertook a project involving the reexamination of microlensing data obtained from three ongoing high-cadence microlensing surveys: the Korea Microlensing Telescope Network (KMTNet: Kim et al. 2016), the Optical Gravitational Lensing Experiment (OGLE: Udalski et al. 2015a), and the Microlensing Observations in Astrophysics survey (MOA: Bond et al. 2001). In this project, we directed our attention to lensing events displaying intricate anomaly features related to caustics, yet lacking prior proposed models to explain these features. We commenced our investigation by analyzing the KMTNet data spanning from the 2016 season to the 2023 season, specifically focusing on identifying anomalous lensing events characterized by conspicuous caustic-crossing features. Subsequently, we examined the presence of lensing models corresponding to these events, filtering out candidate events for which either no models were proposed or presented models failed to accurately delineate the anomalous features. Finally, in the concluding step, we verified the availability of additional data obtained from the OGLE and MOA surveys for the candidate events. Through this process, we identified three lensing events for which the consideration of orbital motions of lenses played crucial role in accurately describing the observed anomaly features in the lensing light curves, including OGLE-2018-BLG-0971, MOA-2023-BLG-065, and OGLE-2023-BLG-0136. In Table 1, we present the equatorial and Galactic coordinates of the events. All these events were captured and observed by multiple surveys. We provide ID references assigned by the respective surveys, using the

ID references from the initial discovery surveys in subsequent discussions.

Observations of the events by the individual surveys were conducted using the telescopes operated by the respective survey groups. The KMTNet group utilizes three identical telescopes, each featuring a 1.6-meter aperture and equipped with a camera capable of capturing a field spanning 4 square degrees. For continuous coverage of lensing events, the KMTNet telescopes are strategically distributed throughout three countries in the Southern Hemisphere: at the Siding Spring Observatory in Australia (KMTA), the Cerro Tololo Interamerican Observatory in Chile (KMTC), and the South African Astronomical Observatory in South Africa (KMTS). The MOA survey employs a telescope with a 1.8 m aperture located at the Mt. John Observatory in New Zealand. The camera mounted on the MOA telescope has the capacity to capture a 2.2 square degree area of the sky in a single shot. The OGLE survey operates the 1.3-meter Warsaw telescope situated at the Las Campanas Observatory in Chile. The camera mounted on the OGLE telescope provides a field of view that spans 1.4 square degrees. The primary observations conducted by the KMTNet and OGLE surveys were done in the I band, whereas observations by the MOA survey were conducted in the custom MOA- R band. In all surveys, a portion of images was acquired in the V band for the color measurements of source stars.

The data of the events were processed using photometry pipelines that are customized to the individual survey groups: KMTNet employed the Albrow et al. (2009) pipeline, OGLE utilized the Udalski (2003) pipeline, and MOA employed the Bond et al. (2001) pipeline. For the use of the optimal data, the KMTNet data set was refined through a re-reduction process using the code developed by Yang et al. (2023). For each data set, error bars estimated from the photometry pipelines were recalibrated not only to ensure consistency of the errorbars with the scatter of data but also to set the χ^2 value per degree of freedom (d.o.f.) for each data to unity. This normalization process was done in accordance with the procedure outlined by Yee et al. (2012).

3. Light curve modeling

The light curves of all the analyzed events show anomalous features that are characteristic of caustics. Caustics arise when a lens system comprises multiple masses, and they represent the source positions at which the magnification of a point source diverges to infinity. Consequently, the presence of caustic-related features in the light curves implies the involvement of lenses composed of multiple masses in producing these events.

Taking the caustic-related features into account, our analysis begins by modeling the light curves within a static 2L1S framework. This framework operates under the assumption of rectilinear relative motion between the lens and the source. Within this static binary-lens model, a lensing light curve is defined by seven basic parameters. The first three of these parameters characterizes the source's approach to the lens, denoted as (t_0, u_0, t_E) . These parameters represent the time of the closest lens-source approach, the separation at that instant (impact parameter), and the event time scale, respectively. Here u_0 is scaled to the angular Einstein radius θ_E , and t_E is defined as the duration for the source to traverse the Einstein radius. The additional set of three parameters (s, q, α) characterizes the binary lens configuration. These parameters represent the projected separation (scaled to θ_E), the mass ratio between the binary lens components (M_1 and M_2), and the angle (source trajectory angle) formed between the

lens-source proper motion vector $\boldsymbol{\mu}$ and the axis defined by M_1 and M_2 . The last parameter ρ , which is defined as the ratio of the angular source radius θ_* to the Einstein radius, that is, $\rho = \theta_*/\theta_E$, quantifies the deformation of lensing light curves during caustic crossings due to finite-source effects. In the 2L1S modeling, we begin by exploring the binary parameters s and q using a grid approach, employing multiple initial values of α . Subsequently, we determine the remaining parameters through a downhill method based on the Markov Chain Monte Carlo (MCMC) technique. The lensing solutions are further refined by enabling variation in all parameters.

As shown in the following section, interpreting the light curves of the events examined in this study solely through static 2L1S models poses a significant challenge. In such instances, we undertake additional modeling to account for higher-order effects which induces deviations in the relative lens-source motion from a rectilinear path. Our modeling approach explores the influences of both lens orbital motion and microlens-parallax effects. To integrate these higher-order effects into the modeling, additional parameters beyond the fundamental set need to be included. The additional parameters introduced for modeling with microlens-parallax effects encompass $(\pi_{E,N}, \pi_{E,E})$, representing the north and east components of the microlens-parallax vector, $\boldsymbol{\pi}_E$, respectively. The microlens-parallax vector is defined as

$$\boldsymbol{\pi}_E = \left(\frac{\pi_{\text{rel}}}{\theta_E} \right) \left(\frac{\boldsymbol{\mu}}{\mu} \right), \quad (1)$$

where $\pi_{\text{rel}} = au(1/D_L - 1/D_S)$ denotes the relative lens-source parallax, and (D_L, D_S) represent the distances to the lens and source, respectively. Under the first-order approximation of small changes in the positions of the lenses during lensing magnifications, the lens-orbital effects can be characterized by two parameters: $(ds/dt, d\alpha/dt)$. These parameters represent the rates of change in the binary separation and the source trajectory angle, respectively. In the parallax+orbit modeling, we enforced a condition where the projected kinetic-to-potential energy ratio was required to remain below unity. This condition ensured that the planet remained gravitationally bound to its host. The energy ratio was computed from the higher-order lensing parameters by

$$\left(\frac{\text{KE}}{\text{PE}} \right)_{\perp} = \frac{(a_{\perp}/\text{au})^3}{8\pi^2(M/M_{\odot})} \left[\left(\frac{1}{s} \frac{ds}{dt} \right)^2 + \left(\frac{d\alpha}{dt} \right)^2 \right]. \quad (2)$$

Here a_{\perp} denotes the physical separation between the binary lens components. Fully describing the Keplerian orbital motion necessitates the inclusion of additional parameters.¹ However, determining these extra parameters poses a challenge due to the limitations of gravitational lensing, which is insensitive to the motion of the lens along the line of sight together with the partial light curve coverage spanning only a minor fraction of the rotation period (Albrow et al. 2000).

Another higher-order effect that causes accelerations of the relative lens-source motion is the orbital motion of the source, known as the "xallarap effect" (Griest & Hu 1992; Han & Gould 1997; Poindexter et al. 2005; Zhu et al. 2017; Satoh et al. 2023). While the lens is confirmed to be binary, there is no prior justification to assume a binary source, although this possibility cannot be entirely dismissed. Therefore, we refrain from testing the xallarap effect as long as the observed anomalies can be explained by lens-orbital effects.

¹ To access a comprehensive description of the orbital lensing parameters, refer to the summary provided in the Appendix of Skowron et al. (2011).




Fig. 1. Light curve of the lensing event OGLE-2018-BLG-0971. The solid and dotted curves drawn over data points represent the model curves obtained from the 2L1S analyses with and without considering the lens orbital motion, respectively. The arrows with labels t_1 , t_2 , and t_3 denote the times of the major anomaly features.

On occasion, light curves of 2L1S events affected by higher-order effects can imitate the patterns seen in 3L1S event light curves. This confusion typically arises when a distinct anomaly feature in a lensing light curve is isolated from the primary ones, as seen in previous events such as MACHO-97-BLG-41, OGLE-2013-BLG-0723, and KMT-2021-BLG-0322. For the analyzed events in this work, we find that such degeneracies do not exist. In the following subsections, we offer detailed analyses of each individual event.

3.1. OGLE-2018-BLG-0971

The OGLE group first identified the lensing event OGLE-2018-BLG-0971 on 4 June, 2018. Four days later, the MOA group verified the event, and the KMTNet group later retrieved it from a post-season data examination. The MOA and KMTNet groups designated the event as MOA-2018-BLG-173 and KMT-2018-BLG-2336, respectively. Figure 1 shows the light curve of the event. The source of the event was situated within the overlapping region of the three KMTNet fields – BLG02, BLG03, and BLG43. To differentiate between the individual data sets, we designate labels corresponding to the respective fields. The light curve manifests multiple anomaly features centered at $\text{HJD} \equiv \text{HJD} - 2400000 \sim 58277$ (t_1), ~ 58280 (t_2), and ~ 58283 (t_3). The symmetric pattern with respect to t_1 suggests that the anomaly around this epoch likely stems from the source’s approach to a caustic cusp, while the U-shaped pattern spanning t_2 to t_3 indicates that these epochs correspond to the times of caustic entrance and exit. Apart from these anomaly features, there is an additional subtle anomaly feature appearing just after the caustic exit. Figure 2 offers a detailed view of this specific region.

After modeling the light curve using a static 2L1S framework, we identified a solution that broadly captures the features of the anomaly. In Table 2, we list the lensing parameters of the static solution. However, it is found that the static model exhibits subtle residuals, particularly in the vicinity of t_3 , as illustrated in the magnified view presented in Figure 2.

In light of deviations from the static model, we checked the feasibility of explaining the deviation with higher-order ef-




Fig. 2. Enlarged view of the OGLE-2018-BLG-0971 light curve in the region around t_2 and t_3 marked in Fig. 2.




Fig. 3. Scatter plots of points in the MCMC chain on the $\pi_{E,E} - \pi_{E,N}$ planes for the four events analyzed in this paper. Points are color-coded to represent those with $< 1\sigma$ (red), $< 2\sigma$ (yellow), $< 3\sigma$ (green), and $< 4\sigma$ (cyan).

fects. The event duration, estimated at $t_E \sim 7.1$ days from the static model, is short. Consequently, we initially considered the lens-orbital effect in the modeling. Through this approach, we derived a solution that accounts for all anomaly features. The model curve of this solution is shown as a solid line in Figure 1, offering a comprehensive view, and in Figure 2, providing an enlarged view around t_3 . The fit significantly improved with the inclusion of lens orbital motion, by $\Delta\chi^2 = 3949.4$ compared to the static model. Subsequent examination of microlens-parallax effects through additional modeling revealed a minimal fit improvement, by $\Delta\chi^2 = 0.9$, indicating the predominant influence of lens-orbital effects. In Figure 3, we plot the scatter plot of points in the MCMC chain on the $\pi_{E,E} - \pi_{E,N}$ plane. In Table 2, we provide the lensing parameters for both the orbit-only and orbit+parallax solutions. The parameters defining the binary lens are $(s, q) \sim (1.01, 0.87)$, indicating that the event was

Table 2. Lensing parameters of OGLE-2018-BLG-0971

Parameter	Static	Orbit	Orbit + Parallax
χ^2	13921.7	9973.2	9972.3
t_0 (HJD')	58278.5148 ± 0.0085	58279.0241 ± 0.0098	58279.0245 ± 0.0097
u_0	0.2771 ± 0.0006	0.2956 ± 0.0005	0.2955 ± 0.0008
t_E (days)	7.143 ± 0.012	7.126 ± 0.009	7.128 ± 0.011
s	0.9552 ± 0.0006	1.0070 ± 0.0009	1.0074 ± 0.0010
q	0.7161 ± 0.0062	0.8669 ± 0.0064	0.8653 ± 0.0066
α (rad)	2.1609 ± 0.0024	2.2964 ± 0.0027	2.2968 ± 0.0025
ρ (10^{-2})	1.078 ± 0.017	1.218 ± 0.011	1.221 ± 0.011
$\pi_{E,N}$	–	–	0.07 ± 0.27
$\pi_{E,E}$	–	–	-0.016 ± 0.091
ds/dt (yr^{-1})	–	-1.722 ± 0.048	-1.720 ± 0.094
$d\alpha/dt$ (rad yr^{-1})	–	0.931 ± 0.026	0.939 ± 0.054




Fig. 4. Lens system configuration for OGLE-2018-BLG-0971. The arrowed line represents the source trajectory, and the closed figures composed of concave curves represent caustics. The caustic shape and location evolve over time due to the orbital motion of the lens. The sets of caustics drawn in black, green, and red correspond to the three anomaly epochs t_1 , t_2 , and t_3 marked in Fig. 1. The right panels display still frames capturing the source's approach to the caustic at these three epochs. The orange circle on the source trajectory appearing in each right panel is sized to indicate the angular dimension of the source relative to the size of the caustic.

generated a binary system composed of roughly equal masses with a separation close to the Einstein radius of the lens system. Although the time scale of the event, $t_E \sim 7.1$ days, represents only a small fraction of the Earth's orbital period, we conducted separate modeling specifically to account for the microlens-parallax effect. From this analysis, it was observed that not only did the fit perform worse compared to the orbit-only model by $\Delta\chi^2 = 42.6$, but also the derived parallax parameters ($\pi_{E,E}, \pi_{E,N}$) $\sim (35.80, 32.50)$ appeared absurdly large for a typical Galactic lensing event.

The configuration of the lens system for the lensing event OGLE-2018-BLG-0971 is shown in Figure 4. This configuration reveals that the lens system forms a single set of resonant caustics featuring six cusps – two along the binary axis and four




Fig. 5. Lensing light curve of MOA-2023-BLG-065. The left and right insets show the zoom-in views of the regions around the caustic spikes at t_1 and t_3 .


positioned away from the axis. The source initially approached the left on-axis cusp around t_1 , entered the caustic near t_2 , and exited the caustic at around t_3 . These caustic approach and crossings gave rise to anomalies at the corresponding epochs. The primary deviation of the static 2L1S model, particularly around t_3 , stems from its inability to account for the variation in the caustic caused by the orbital motion of the binary lens.

3.2. MOA-2023-BLG-065

Figure 5 shows the lensing light curve of the MOA-2023-BLG-065 event. The source flux magnification induced by lensing was initially identified on 17 March, 2023 (HJD' = 60021) through the survey conducted by the MOA group, subsequently confirmed by the KMTNet group. The ID reference designated by the KMTNet survey is KMT-2023-BLG-2430. The light curve displays a complex pattern comprising multiple anomaly features. Notably, two spike features at $t_1 \sim 60020$ and $t_3 \sim 60035$ appear to be a pair of caustic-crossing spikes. Furthermore, the symmetry observed between the ascending and descending segments of the anomaly feature centered at $t_4 \sim 60043$ implies that it originated from the source approach to a cusp of the caustic. While the magnification between caustic spikes typically follows

Table 3. Lensing parameters of MOA-2023-BLG-065


Parameter	Static	Parallax	Orbit	Orbit + Parallax
χ^2	2399.0	1876.6	1882.7	1872.6
t_0 (HJD')	60030.308 ± 0.024	60030.140 ± 0.045	60029.951 ± 0.079	60030.356 ± 0.090
u_0	-0.3893 ± 0.0053	-0.426 ± 0.009	-0.4539 ± 0.0079	-0.4378 ± 0.0083
t_E (days)	30.74 ± 0.51	37.27 ± 0.59	39.03 ± 0.54	37.81 ± 0.75
s	1.327 ± 0.011	1.452 ± 0.008	1.471 ± 0.007	1.467 ± 0.010
q	1.089 ± 0.013	0.877 ± 0.041	0.939 ± 0.023	0.887 ± 0.022
α (rad)	4.7757 ± 0.0001	4.7668 ± 0.0015	4.7705 ± 0.0050	4.7477 ± 0.0062
ρ (10^{-3})	2.018 ± 0.039	1.724 ± 0.041	1.763 ± 0.033	1.739 ± 0.037
$\pi_{E,N}$	–	0.069 ± 0.119	–	-0.20 ± 0.26
$\pi_{E,E}$	–	-0.157 ± 0.045	–	-0.214 ± 0.054
ds/dt (yr^{-1})	–	–	0.015 ± 0.071	-0.296 ± 0.082
$d\alpha/dt$ (rad yr^{-1})	–	–	-0.531 ± 0.159	0.217 ± 0.506


Fig. 6. Enlarged view of the MOA-2023-BLG-065 light curve in the region around t_4 marked in Fig. 5.

a U-shape pattern, the region between the caustic spikes at t_1 and t_3 deviates significantly from the U-shape, displaying a distinctive rise and fall in the region centered at $t_2 \sim 60023$. This deviation is indicative of a source asymptotically approaching a fold of the caustic.

From the 2LIS analysis of the lensing light curve under a static binary frame, we found that the model falls short of precisely describing the data, even though it approximately outlines the anomaly features. In Figure 5, the static 2LIS model is represented by a dotted curve. This static model exhibits a notably inadequate fit especially in the region of the anomaly centered at t_4 , as highlighted in the enlarged view presented in Figure 6. The complete lensing parameters for the static 2LIS solution are detailed in Table 3.

Although the static solution cannot adequately describe the anomaly feature at around t_4 , it does exhibit an anomaly around the time of that anomaly. This suggests the possibility that it can be described with a slight deformation of the source trajectory caused by higher-order effects. In light of this possibility, we conducted three additional sets of modeling: the first two models separately incorporated microlens-parallax and lens-orbital effects, while the third model encompassed both effects simultaneously. In Table 3, we present the lensing parameters for the


Fig. 7. Lens system configuration for MOA-2023-BLG-065.

three models. Comparing the static and higher-order solution demonstrates a notable enhancement in fit, with a $\Delta\chi^2 = 526.4$ compared to the static solution. In Figure 3, we present the scatter plot of points in the MCMC chain on the $\pi_{E,E}-\pi_{E,N}$ plane. It shows that the east component of the microlens-parallax vector, $\pi_{E,E}$, is constrained, although the uncertainty of the north component, $\pi_{E,N}$, is large. Additionally, the normalized source radius $\rho = (1.739 \pm 0.037) \times 10^{-3}$ was measured from the deformation of the light curve by finite source effects during the epochs around t_2 and t_3 . The magnified views of these specific regions are shown in the two insets of Figure 5.

Figure 7 illustrates the lens-system configuration of MOA-2023-BLG-065. It shows that the binary lens, characterized by parameters $(s, q) \sim (1.5, 0.9)$, creates a resonant caustic, and the source traversed the left side of this caustic. Initially, the source passed the upper fold, creating the first caustic spike at t_1 . Subsequently, it approached the upper left fold asymptotically, resulting in rising and falling features around t_2 . The source then exited the caustic, generating the second caustic spike at t_3 , and proceeded to pass the tip of the lower left cusp, causing the final anomaly feature at around t_4 . The distortion of the caustic due to the orbital motion of the lens led the light curve to deviate from




Fig. 8. Lensing light curve of OGLE-2023-BLG-0136. The insets provides closer look at the regions surrounding t_2 and t_3 .

the anticipated behavior according to the static model. This discrepancy was particularly noticeable in the part of the light curve during the final cusp approach, enabling the detection of the lens orbital effect.

3.3. OGLE-2023-BLG-0136

The lensing event OGLE-2023-BLG-0136 was first found during its early phase by the OGLE group on 1 April, 2023 (HJD' = 60036). Subsequently, the KMTNet survey validated the event, designating it as KMT-2023-BLG-2849. Figure 8 illustrates the lensing light curve of the event. The curve exhibits a complex pattern of anomalies, comprised of multiple distinct features: a caustic-crossing feature around $t_1 = 59997$ and two additional features centered at around $t_2 = 60089$ and $t_3 = 60103$. The first feature had limited coverage because it was observed solely by the KMTc telescope, while the other KMTNet telescopes and the OGLE telescope did not start observations of the 2023 season at the time of the anomaly. However, the subsequent two anomaly features were extensively observed by the combined data from both the OGLE and KMTNet surveys.

We initially modeled the lensing light curve under a static 2L1S framework. The model curve of the static solution is depicted by a dotted curve in Figure 8, and the lensing parameters of the solution are listed in Table 4. Upon examination of the fit, it is observed that this model roughly captures the anomaly features around t_2 and t_3 , yet it lacks accuracy in explaining the feature at t_1 . Figure 9 offers a closer look at the model fit around t_1 . While the static model does not precisely capture the first caustic-crossing anomaly, it does display a weak bump feature which seems to result from a caustic approach. This suggests that the source might pass over the caustic, potentially influenced by a slight shift in the caustic position due to the orbital motion of the binary lens. Considering this, we move forward with additional modeling, incorporating the effects of the lens orbital motion.

The solid curves in Figures 8 and 9 represent the model derived from the orbital solution. The lensing parameters of the solution are listed in Table 4. From the inspection of the fit, it is found that the orbital solution accurately accounts for the anomaly around t_1 , improving the fit by $\Delta\chi^2 = 2367.4$ com-




Fig. 9. Enlarged view of the OGLE-2023-BLG-0136 light curve in the region around t_1 marked in Fig. 8.




Fig. 10. Configuration of the lens system for OGLE-2023-BLG-0136.

pared to the static solution. The estimated event time scale, $t_E \sim 60$ days, is moderately long, and thus we further examined whether incorporating the microlens-parallax effect could enhance the fit. From the model derived with the consideration both higher-order effects, we found a marginal enhancement in the fit, $\Delta\chi^2 = 4.5$, compared to the orbital solution. This suggests that the dominant higher-order effect is attributed to the lens orbital motion. Details of the lensing parameters for the parallax and orbit+parallax solutions are outlined in Table 4 and the scatter plot on the $\pi_{E,E} - \pi_{E,N}$ plane is shown in Figure 3. The estimated binary lens parameters are $(s, q) \sim (0.71, 0.30)$. The normalized source radius, $\rho = (2.005 \pm 0.029) \times 10^{-3}$, was precisely measured from the deformed light curve during the anomaly at around t_2 . The enlarged view of this region is shown in the inset of Figure 8.

Table 4. Lensing parameters of OGLE-2023-BLG-0136

Parameter	Static	Parallax	Orbit	Orbit + Parallax
χ^2	3180.0	2773.6	812.6	809.4
t_0 (HJD')	60030.726 ± 0.093	60029.065 ± 0.132	60030.276 ± 0.107	60030.08 ± 0.11
u_0	0.2685 ± 0.0007	0.2290 ± 0.0008	0.2808 ± 0.0012	0.2833 ± 0.0017
t_E (days)	66.70 ± 0.13	86.00 ± 0.2999499	61.25 ± 0.13	59.35 ± 0.48
s	0.66848 ± 0.00042	0.68593 ± 0.00034	0.71297 ± 0.00035	0.71101 ± 0.00085
q	0.3117 ± 0.0015	0.2607 ± 0.0025	0.2851 ± 0.0027	0.2978 ± 0.0034
α (rad)	1.4503 ± 0.0009	1.3623 ± 0.0019	1.4268 ± 0.0016	1.42832 ± 0.0027
ρ (10^{-3})	1.965 ± 0.029	1.535 ± 0.03	1.992 ± 0.029	2.005 ± 0.029
$\pi_{E,N}$	–	0.016 ± 0.003	–	0.080 ± 0.021
$\pi_{E,E}$	–	-0.220 ± 0.001	–	0.030 ± 0.010
ds/dt (yr^{-1})	–	–	-0.4403 ± 0.0037	-0.4410 ± 0.0039
$d\alpha/dt$ (rad yr^{-1})	–	–	-0.237 ± 0.007	-0.408 ± 0.050

In Figure 10, we illustrate the lens system configuration for OGLE-2023-BLG-0136. At the time of the first anomaly, the caustic exhibited a resonant form in which the central caustic and peripheral caustics were interconnected by narrow bridges. The first anomaly feature at around t_1 was generated when the source traversed the lower bridge of this resonant caustic. As the source proceeded, the binary separation decreased, leading to the detachment of the peripheral caustics from the central caustic. The second and third anomaly features arose from the successive passages of the source over the lower and right cusps of the upper peripheral caustic. The static model effectively captured the second and third anomaly features due to their close temporal proximity, which minimized the deformation of the caustic caused by the lens orbital motion. However, the time gap between these features and the first anomaly feature exceeded 100 days, resulting in substantial alterations to the caustic shape and position. As a result, the static model fell short in accurately representing the observed light curve.

It is important to note that event time scales can significantly vary between static and non-static models. For instance, in the case of MOA-2023-BLG-065, the time scale shifts from ~ 30.7 days for the static solution to ~ 37.8 days for the higher-order solution. Similarly, for OGLE-2023-BLG-0136, the time scale changes from ~ 66.7 days for the static solution to ~ 59.4 days for the higher-order solution. The time scale serves as a fundamental observable for constraining the physical parameters of the lens. Therefore, accounting for higher-order effects in modeling is essential for accurately determining these parameters.

4. Source stars and Einstein radii

In this section, we specify the source stars and determine the angular Einstein radii of the events. The source stars are specified based on their colors and magnitudes, accounting for corrections due to reddening and extinction. In this process, we first conducted photometry of the I and V -band data using the pyDIA code (Albrow 2017), and then estimated the instrumental source color and magnitude, $(V - I)_S$, by regressing the data of the individual passbands with respect to the model. Calibration of the source color and magnitude follows the method outlined by Yoo et al. (2004), leveraging the centroid of the red giant clump (RGC) in the color-magnitude diagram (CMD) for this purpose. The RGC centroid can be used for a reference because its de-reddened color and magnitude, represented as $(V - I)_{\text{RGC},0}$, are established from studies conducted by Bensby et al. (2013) and Nataf et al. (2013). In the calibration process, we first po-




Fig. 11. Positions of source stars with respect to the centroids of red giant clump (RGC) in the instrumental color-magnitude diagrams. For MOA-2023-BLG-065 and OGLE-2023-BLG-0136, the positions of the blends are additionally marked.

sitioned the source in the CMD constructed using the pyDIA code, measured the offsets of the source in color and magnitude, $\Delta(V - I, I)$, from the RGC centroid, and then estimated the de-reddened source color and magnitude as:

$$(V - I, I)_{S,0} = (V - I, I)_{\text{RGC},0} + \Delta(V - I, I). \quad (3)$$

In Figure 11, we indicate the locations of source stars for the individual events relative to the RGC centroids on the instrumental CMDs constructed using the KMTC data sets. The values estimated for $(V - I)_S$, $(V - I)_{\text{RGC}}$, $(V - I)_{\text{RGC},0}$, and $(V - I)_{S,0}$ through the described procedure are compiled in Table 5. Based on the derived colors and magnitudes, we determine that the source star of OGLE-2018-BLG-0971 is a K-type subgiant. Additionally, the sources of MOA-2023-BLG-065 and OGLE-2023-BLG-0136 are G-type main-sequence stars.

The angular Einstein radius of each event was estimated from the relation

$$\theta_E = \frac{\theta_*}{\rho}, \quad (4)$$

where the angular source radius θ_* is deduced from the color and magnitude, and the normalized source radius ρ is measured from

Table 5. Source parameters, Einstein radii, and relative proper motions

Parameter	OGLE-2018-BLG-0971	MOA-2023-BLG-065	OGLE-2023-BLG-0136
$(V - I)_S$	2.446 ± 0.021	1.730 ± 0.011	1.488 ± 0.035
I_S	18.876 ± 0.002	20.490 ± 0.002	19.373 ± 0.005
$(V - I, I)_{\text{RGC}}$	(2.645, 116.173)	(2.131, 15.788)	(1.817, 15.452)
$(V - I, I)_{\text{RGC},0}$	(1.060, 14.371)	(1.060, 14.386)	(1.060, 14.393)
$(V - I)_{S,0}$	$0.861 \pm 0.045,$	$0.659 \pm 0.041,$	$0.731 \pm 0.053,$
I_0	17.073 ± 0.020	19.089 ± 0.020	18.315 ± 0.021
Type	K1IV	G0V	G4V
θ_* (μas)	1.433 ± 0.119	0.451 ± 0.037	0.699 ± 0.061
θ_E (mas)	0.117 ± 0.010	0.256 ± 0.021	0.351 ± 0.031
μ (mas/yr)	6.01 ± 0.50	2.42 ± 0.20	2.13 ± 0.19

the light curve analysis. To derive the source radius, we initially transformed the measured $V - I$ color into $V - K$ color using the [Bessell & Brett \(1988\)](#) relation. Subsequently, we determined θ_* using the relationship provided by [Kervella et al. \(2004\)](#) between $(V - K, V)$ and θ_* . The estimated angular radii of source stars for the individual lensing events are listed in Table 5, along with the corresponding angular Einstein radii calculated using the relationship described in Eq. (3). Also listed in the table are the relative lens-source proper motion estimated by the relation

$$\mu = \frac{\theta_E}{t_E}. \quad (5)$$

For MOA-2023-BLG-065 and OGLE-2023-BLG-0136, we were able to constrain the blended light. In Figure 11, we mark the positions of blend in the CMDs. To assess the likelihood of the lens being the primary source of the blended flux, we measured the centroids of the source image during lensing magnification and at the baseline. In the case of MOA-2023-BLG-065, the measured astrometric offset between the source positions is $\delta\theta = 0.46 \pm 0.08$ arcsec, and this excludes the possibility that the blended light comes mainly from the lens. For OGLE-2023-BLG-0136, the significant astrometric uncertainty prevents drawing a meaningful conclusion regarding the origin of the blended light.

5. Physical lens parameters

The physical parameters of a lens are constrained by the lensing observables of t_E , θ_E , and π_E . When all these parameters are simultaneously measured, the mass and distance to the lens are uniquely determined by

$$M = \frac{\theta_E}{\kappa\pi_E}; \quad D_L = \frac{\text{au}}{\pi_E\theta_E + \pi_S}. \quad (6)$$

Here $\kappa = 4G/(c^2\text{au})$ and $\pi_S = \text{au}/D_S$ represents the parallax of the source lying at a distance D_S ([Gould 2000](#)). For all analyzed events, the observables of t_E and θ_E were precisely measured, but the constraint on the microlens parallax was relatively weak because of its subtle effects. Due to these incomplete measurements of the lensing observables, we determine the physical lens parameters through Bayesian analyses of the individual events. This approach integrates constraints from measured lensing observables with priors derived from the physical and dynamic distributions, as well as the mass function of lens objects within the Galaxy.

The Bayesian analysis began by generating a large number of synthetic events via Monte Carlo simulation. Within this simulation, the physical parameters of the lens mass were deduced

from a model mass function, while the distances to the lens and source, along with their relative proper motion, were derived from a Galaxy model. Our approach incorporated the mass function model suggested by [Jung et al. \(2018\)](#) and utilized the Galaxy model introduced by [Jung et al. \(2021\)](#). For each synthetic event defined by physical parameters ($M_i, D_{L,i}, D_{S,i}, \mu_i$), we calculated the values of the corresponding lensing observables using the relations

$$t_{E,i} = \frac{\theta_{E,i}}{\mu_i}; \quad \theta_{E,i} = \sqrt{\kappa M_i \pi_{\text{rel},i}} \quad (7)$$

for the event time scale and Einstein radius, respectively, and using the relation in Eq. (1) for the microlens parallax. Then, the posteriors of M and D_L were obtained by assigning a weight to each event of $w_i = \exp(-\chi_i^2/2)$, where χ_i^2 value is computed by

$$\chi_i^2 = \frac{\Delta t_{E,i}^2}{\sigma^2(t_E)} + \frac{\Delta \theta_{E,i}^2}{\sigma^2(\theta_E)} + \sum_{j=1}^2 \sum_{k=1}^2 b_{j,k} (\pi_{E,j,i} - \pi_{E,i})(\pi_{E,k,i} - \pi_{E,i}). \quad (8)$$

Here, $\Delta t_{E,i} = t_{E,i} - t_E$, $\Delta \theta_{E,i} = \theta_{E,i} - \theta_E$, (t_E, θ_E) stand for the measured values of the observables, $[\sigma(t_E), \sigma(\theta_E)]$ indicate their corresponding uncertainties, and $b_{j,k}$ represents the inverse covariance matrix of the microlens-parallax vector π_E , $(\pi_{E,1}, \pi_{E,2})_i = (\pi_{E,N}, \pi_{E,E})_i$ are the parallax parameters of each simulated event, and $(\pi_{E,N}, \pi_{E,E})$ represent the parallax parameters measured from modeling. [Han et al. \(2016b\)](#) showed that parallax measurements can be important even when there is little χ^2 improvement, because they can constrain π_E to be small. We, therefore, consider the constraints given by the measured parallax parameters in the Bayesian analyses.

In Figures 12 and 13, we present the Bayesian posteriors of the primary lens masses and distances to the lens systems. In Table 6, we summarize the estimated physical parameters for the individual lensing events. Among the parameters, M_1 and M_2 denote the masses of the primary and companion of the lens, and a_\perp denotes the projected physical separation between M_1 and N_2 . We present the median value derived from the Bayesian posterior distribution as a representative value for each physical parameter, with uncertainties estimated within the 16% to 84% range of the distribution. Also listed in the table are the relative probabilities for the lens being in the Galactic disk, p_{disk} , and in the bulge, p_{bulge} . According to the estimated masses, the lenses of the events OGLE-2018-BLG-0971 and MOA-2023-BLG-065 are binaries composed of M dwarfs. On the other hand, the lens of OGLE-2023-BLG-0136 is likely to be a binary composed of an early K-dwarf primary and a late M-dwarf companion. Across all the analyzed events, p_{bulge} is substantially higher than p_{disk} , suggesting a higher likelihood of the all lenses being located in the bulge rather than the disk. The probabilities p_{disk} and

Table 6. Physical lens parameters

Parameter	OGLE-2018-BLG-0971	MOA-2023-BLG-065	OGLE-2023-BLG-0136
$M_1 (M_\odot)$	$0.215^{+0.231}_{-0.126}$	$0.30^{+0.27}_{-0.15}$	$0.88^{+0.23}_{-0.18}$
$M_2 (M_\odot)$	$0.186^{+0.200}_{-0.109}$	$0.28^{+0.25}_{-0.14}$	$0.259^{+0.068}_{-0.052}$
D_L (kpc)	$7.50^{+0.98}_{-1.01}$	$7.62^{+0.89}_{-0.84}$	$7.52^{+0.92}_{-1.13}$
a_\perp (au)	$1.39^{+0.18}_{-0.19}$	$3.10^{+0.36}_{-0.34}$	$1.92^{+0.23}_{-0.29}$
p_{disk}	26%	9%	21%
p_{bulge}	74%	91%	79%




Fig. 12. Bayesian posterior distributions of the primary lens mass (M_1) for the lensing events. In each panel, the event rate contributions from the disk and bulge lens populations are depicted by the blue and red curves, respectively, while black curve represents the sum of the contributions from two lens populations.

p_{bulge} were found by analyzing the proportion of artificial lensing events where the lenses came from either the disk or bulge distributions within the Galaxy model utilized in the Monte Carlo simulation process.

6. Summary

We have analyzed microlensing data collected from high-cadence surveys to reevaluate lensing events that lacked proposed interpretations for their intricate anomaly features. Through detailed reanalyses considering higher-order effects, we identified that accounting for the orbital motions of lenses was vital in accurately explaining the anomaly features observed in the lensing light curves of OGLE-2018-BLG-0971, MOA-2023-BLG-065, and OGLE-2023-BLG-0136.

By conducting Bayesian analyses based on the lensing parameters from the newly found solutions and together with the constraints derived from lensing observables, we estimated masses and distances to the lenses. These analyses revealed that the lenses for events OGLE-2018-BLG-0971 and MOA-2023-BLG-065 are binary systems consisting of M dwarfs. Addition-




Fig. 13. Bayesian posterior distributions of the distances to the lens (D_L). The notations correspond to those used in Fig. 12.

ally, for OGLE-2023-BLG-0136, the lens is likely to be a binary system comprising an early K-dwarf primary and a late M-dwarf companion. Notably, across all observed lensing events, the likelihood of the lens being in the bulge significantly outweighs its likelihood of being in the disk.

Acknowledgements. Work by C.H. was supported by the grants of National Research Foundation of Korea (2019R1A2C2085965). J.C.Y. and I.-G.S. acknowledge support from U.S. NSF Grant No. AST-2108414. Y.S. acknowledges support from BSF Grant No. 2020740. This research has made use of the KMTNet system operated by the Korea Astronomy and Space Science Institute (KASI) at three host sites of CTIO in Chile, SAAO in South Africa, and SSO in Australia. Data transfer from the host site to KASI was supported by the Korea Research Environment Open Network (KREONET). This research was supported by KASI under the R&D program (project No. 2023-1-832-03) supervised by the Ministry of Science and ICT. W.Z. and H.Y. acknowledge support by the National Natural Science Foundation of China (Grant No. 12133005). W. Zang acknowledges the support from the Harvard-Smithsonian Center for Astrophysics through the CfA Fellowship. The MOA project is supported by JSPS KAKENHI Grant Number JP24253004, JP26247023, JP16H06287 and JP22H00153.

References

- Albrow, M. D., Beaulieu, J.-P., Caldwell, J. A. R., et al. 2000, *ApJ*, 534, 894
- Albrow, M., Horne, K., Bramich, D. M., et al. 2009, *MNRAS*, 397, 2099
- Albrow, M. 2017, MichaelDAIbrow/pyDIA: Initial Release on Github, Version v1.0.0, Zenodo, doi:10.5281/zenodo.268049

- Alcock, C., Allsman, R. A., Alves, D., et al. 2000, *ApJ*, 541, 270
- Bennett, D. P., Rhie, S. H., Becker, A. C., et al. 1999, *Nature*, 402, 57
- Bennett, D. P., Rhie, S. H., Nikolaev, S., et al. 2010, *ApJ*, 713, 837
- Bensby, T., Yee, J. C., Feltzing, S., et al. 2013, *A&A*, 549, A147
- Bessell, M. S., & Brett, J. M. 1988, *PASP*, 100, 1134
- Bond, I. A., Abe, F., Dodd, R. J., et al. 2001, *MNRAS*, 327, 868
- Gaudi, B. S., Bennett, D. P., Udalski, A., et al. 2008, *Science*, 319, 927
- Gould, A. 1992, *ApJ*, 392, 442
- Gould, A. 2000, *ApJ*, 542, 785
- Griest, K., & Hu, W. 1992, *ApJ*, 397, 362
- Han, C., & Gould, A. 1997, *ApJ*, 480, 196
- Han, C., Bennett, D. P., Udalski, A., et al. 2016a, *ApJ*, 825, 8
- Han, C., Udalski, A., Gould, A., et al. 2016b, *ApJ*, 828, 53
- Han, C., Gould, A., Hirao, Y., et al. 2021, *A&A*, 655, A24
- Jung, Y. K., Han, C., Gould, A., et al. 2013, *ApJ*, 768, L7
- Jung, Y. K., Udalski, A., Gould, A., et al. 2018, *AJ*, 155, 219
- Jung, Y. K., Han, C., Udalski, A., et al. 2021, *AJ*, 161, 293
- Kervella, P., Thévenin, F., Di Folco, E., & Ségransan, D. 2004, *A&A*, 426, 29
- Kim, S.-L., Lee, C.-U., Park, B.-G., et al. 2016, *JKAS*, 49, 37
- Nataf, D. M., Gould, A., Fouqué, P., et al. 2013, *ApJ*, 769, 88
- Park, H., Udalski, A., Han, C., et al. 2013, *ApJ*, 778, 134
- Poindexter, S., Afonso, C., Bennett, D. P., et al. 2005, *ApJ*, 633, 914
- Rahvar, S., & Dominik, M. 2009, *MNRAS*, 392, 1193
- Satoh, Y. K., Koshimoto, N., Bennett, D. P., et al. 2023, *AJ*, 166, 116
- Shin, I.-G., Udalski, A., Han, C., et al. 2011, *ApJ*, 735, 85
- Skowron, J., Udalski, A., Gould, A., et al. 2011, *ApJ*, 738, 87
- Smith, M. C., Mao, S., & Paczyński, B. 2003, *MNRAS*, 339, 925
- Udalski, A. 2003, *Acta Astron.*, 53, 291
- Udalski, A., Jung, Y. K., Han, C., et al. 2015a, *ApJ*, 812, 47
- Udalski, A., Szymański, M. K., Szymański, G., et al. 2015, *Acta Astron.*, 65, 1
- Wyrzykowski, Ł., Mróz, P., Rybicki, K. A., et al. 2020, *A&A*, 633, A98
- Yang, H., Yee, J. C., Hwang, K.-H., et al. 2023, *MNRAS*, submitted
- Yee, J. C., Shvartzvald, Y., Gal-Yam, A., et al. 2012, *ApJ*, 755, 102
- Yee, J. C., Johnson, J. A., Skowron, J., et al. 2016, *ApJ*, 821, 121
- Yoo, J., DePoy, D. L., Gal-Yam, A. et al. 2004, *ApJ*, 603, 139
- Zhu, W., Udalski, A., Novati, S. C., et al. 2017, *AJ*, 154, 210
- ¹⁹ Department of Physics, The Catholic University of America, Washington, DC 20064, USA
- ²⁰ Department of Astronomy, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
- ²¹ Sorbonne Université, CNRS, UMR 7095, Institut d'Astrophysique de Paris, 98 bis bd Arago, 75014 Paris, France
- ²² Department of Physics, University of Auckland, Private Bag 92019, Auckland, New Zealand
- ²³ University of Canterbury Mt. John Observatory, P.O. Box 56, Lake Tekapo 8770, New Zealand

-
- ¹ Department of Physics, Chungbuk National University, Cheongju 28644, Republic of Korea
- ² Astronomical Observatory, University of Warsaw, Al. Ujazdowskie 4, 00-478 Warszawa, Poland
- ³ Institute of Natural and Mathematical Science, Massey University, Auckland 0745, New Zealand
- ⁴ Korea Astronomy and Space Science Institute, Daejeon 34055, Republic of Korea
- ⁵ Max Planck Institute for Astronomy, Königstuhl 17, D-69117 Heidelberg, Germany
- ⁶ Department of Astronomy, The Ohio State University, 140 W. 18th Ave., Columbus, OH 43210, USA
- ⁷ University of Canterbury, Department of Physics and Astronomy, Private Bag 4800, Christchurch 8020, New Zealand
- ⁸ Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot 76100, Israel
- ⁹ Center for Astrophysics | Harvard & Smithsonian 60 Garden St., Cambridge, MA 02138, USA
- ¹⁰ Department of Astronomy and Tsinghua Centre for Astrophysics, Tsinghua University, Beijing 100084, China
- ¹¹ School of Space Research, Kyung Hee University, Yongin, Kyeonggi 17104, Republic of Korea
- ¹² Department of Physics, University of Warwick, Gibbet Hill Road, Coventry, CV4 7AL, UK
- ¹³ Institute for Space-Earth Environmental Research, Nagoya University, Nagoya 464-8601, Japan
- ¹⁴ Code 667, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA
- ¹⁵ Department of Astronomy, University of Maryland, College Park, MD 20742, USA
- ¹⁶ Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
- ¹⁷ Instituto de Astrofísica de Canarias, Vía Láctea s/n, E-38205 La Laguna, Tenerife, Spain
- ¹⁸ Department of Earth and Space Science, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan