

# CONTROL OF THE SCHRÖDINGER EQUATION IN $\mathbb{R}^3$ : THE CRITICAL CASE

P. BRAZ E SILVA, R. DE A. CAPISTRANO-FILHO, J. D. DO N. CARVALHO,  
AND D. DOS SANTOS FERREIRA

ABSTRACT. This article deals with the  $H^1$ -level local null controllability for the defocusing critical nonlinear Schrödinger equation in  $\mathbb{R}^3$ . Firstly, we show the problem under consideration to be well-posed using Strichartz estimates. Moreover, through the Hilbert uniqueness method, we prove the linear Schrödinger equation to be controllable. Finally, we use a perturbation argument and show local controllability for the critical nonlinear Schrödinger equation.

## 1. INTRODUCTION

The study of the energy-critical case of the nonlinear Schrödinger equation (C-NLS) in four dimensions, namely,

$$(1.1) \quad \begin{cases} i\partial_t u + \Delta u - |u|^4 u = 0, & \text{on } \mathbb{R}^3 \times [0, +\infty), \\ u(x, 0) = u_0(x) \in H^1(\mathbb{R}^3), \end{cases}$$

is motivated by several deep mathematical and physical reasons. In the context of partial differential equations, the term “energy-critical” refers to the fact that the associated energy of the equation is invariant with respect to its natural scaling. For the usual NLS (with  $|u|^2 u$  instead of  $|u|^4 u$  in (1.1)), this scaling symmetry is crucial because it allows for the potential development of self-similar blow-up solutions, which are central to understanding singularity formation. In four dimensions, the equation (1.1) is energy-critical, meaning the nonlinearity is precisely balanced with the dispersive nature of the equation. This balance leads to delicate analytical challenges, such as the need for sophisticated tools to study global well-posedness, scattering, and the behavior of solutions.

Understanding whether solutions scatter (i.e., behave like linear solutions at infinity) is a central question for large data. The system (1.1) in four dimensions is a key case where one can study the global dynamics of solutions, including whether all solutions with sub-critical energy scatter or if some can lead to blow-up. The four-dimensional case is often studied using concentration compactness methods, which are crucial for understanding the dynamics near critical points. These methods have broader implications for other dispersive equations as well.

In summary, the energy-critical system (1.1) in four dimensions is a rich and challenging problem with deep connections to pure and applied mathematics, making it a significant and highly motivated area of study. For details we strongly encourage the reader to see [7, 12, 34, 20, 31, 13, 6] and the references therein.

---

*Date:* November 4, 2024.

2020 *Mathematics Subject Classification.* 93B05, 93B07, 35Q55, 49K40.

*Key words and phrases.* NLS system, Critical case, Internal controllability, HUM.

\*Corresponding author: ddsf@math.cnrs.fr.

Braz e Silva was partially supported by CAPES/MATH-AMSUD #8881.520205/2020-01, CAPES/PRINT #88887.311962/2018-00, CAPES/COFECUB #88887.879175/2023-00, CNPq #421573/2023-6, #305233/2021-1, #308758/2018-8, #432387/2018-8. Capistrano-Filho was partially supported by CAPES/MATH-AMSUD #8881.520205/2020-01, CAPES/PRINT #88887.311962/2018-00, CAPES/COFECUB #88887.879175/2023-00, CNPq #421573/2023-6, #307808/2021-1 and #401003/2022-1 and Propesqi (UFPE). Carvalho was partially supported by CNPq, CAPES-MATHAMSUD #8881.520205/2020-01 and FACEPE BFD-0014-1.01/23.

**1.1. Addressed issues and review of the literature.** In this article, we will consider the  $H^1(\mathbb{R}^3)$  local controllability for the defocusing critical nonlinear Schrödinger equation (C-NLS)

$$(1.2) \quad \begin{cases} i\partial_t u + \Delta u - |u|^4 u = f(x, t), & \text{on } \mathbb{R}^3 \times [0, +\infty), \\ u(x, 0) = u_0(x) \in H^1(\mathbb{R}^3), \end{cases}$$

where  $u = u(x, t)$  is a complex-valued function of two variables  $x \in \mathbb{R}^3$  and  $t \in \mathbb{R}$ , the subscripts denote the corresponding partial derivatives, while the function  $f(x, t)$  is a control input. We are mainly concerned with the following null control problem for system (1.2).

**Control problem:** Let  $T > 0$  be given. For any given  $u_0 \in H^1(\mathbb{R}^3)$ , can one find a control  $f(x, t)$  such that system (1.2) admits a solution  $u \in C([0, T]; H^1(\mathbb{R}^3))$  satisfying  $u(x, T) = 0$  in  $\mathbb{R}^3$ ?

Control properties of Schrödinger equations have received much attention in the last decades. For example, regarding control issues, one may see [14, 22, 23, 24] and the references therein. As for Carleman estimates and applications to inverse problems, we cite [2, 4, 5, 18, 21, 33] and the references therein. An excellent review of the contributions up to 2003 is in [36].

Let us detail some recent results. The results due to Illner *et al.* [9, 10] considered internal controllability of the nonlinear Schrödinger equation posed on a finite interval  $(-\pi, \pi)$

$$(1.3) \quad \begin{cases} i\partial_t v + \partial_{xx} v + \lambda|v|^2 v = f(x, t), & x \in (-\pi, \pi), \\ v(-\pi, t) = v(\pi, t), \quad \partial_x v(-\pi, t) = \partial_x v(\pi, t), \end{cases}$$

where the forcing function  $f = f(x, t)$ , supported in a sub-interval of  $(-\pi, \pi)$ , is a control input. They showed that system (1.3) is locally exactly controllable in the space

$$H_p^1(-\pi, \pi) := \{v \in H^1(-\pi, \pi) : v(-\pi) = v(\pi)\}.$$

Later, Lange and Teismann [15] considered the internal control of the same nonlinear Schrödinger equation in (1.3) posed on a finite interval but with the homogeneous Dirichlet boundary conditions

$$v(-\pi, t) = v(\pi, t) = 0,$$

and showed that this is locally exactly controllable in the space  $H_0^1(0, \pi)$  around a special ground state of the equation (see also [16] for internal controllability of the nonlinear Schrödinger equation posed on a finite interval).

In [27], Rosier and Zhang considered the nonlinear Schrödinger equation

$$(1.4) \quad i\partial_t u + \Delta u + \lambda|u|^2 u = 0$$

posed on a bounded domain  $K$  in  $\mathbb{R}^n$  with both Dirichlet boundary conditions and Neumann boundary conditions. They showed that if either

$$s > \frac{n}{2}$$

or

$$0 \leq s < \frac{n}{2}, \quad \text{with } 1 \leq n < 2 + 2s$$

or

$$s = 0, 1 \quad \text{with } n = 2,$$

then both systems, with Dirichlet and Neumann conditions, when the control inputs are acting on the whole boundary of  $K$ , are locally exactly controllable in the classical Sobolev space  $H^s(K)$  around any smooth solution of the Schrödinger equation.

In [28], the authors extend the results of Rosier and Zhang [26]. More precisely, they assume that the spatial variable belongs to the rectangle

$$\mathcal{R} = (0, l_1) \times \cdots \times (0, l_n)$$

and investigate the control properties of the semi-linear Schrödinger equation

$$i\partial_t u + \Delta u + \lambda|u|^\alpha u = ia(x)h(x, t), \quad x \in \mathbb{T}^n \quad t \in (0, T),$$

where  $\lambda \in \mathbb{R}$  and  $\alpha \in 2\mathbb{N}^*$  by combining new linear controllability results in the spaces  $H^s(\mathcal{R})$  with Bourgain analysis. In this case, the geometric control condition is not required (see [28] for more details). It is important to note that these cases are studied in the subcritical case.

Finally, considering a  $2d$ -compact Riemann manifold  $M$  without boundary, Dehman *et al.* [8] studied internal control and stabilization of nonlinear Schrödinger equations

$$i\partial_t w + \Delta w - |w|^2 w = f(x, t), \quad x \in M.$$

They showed, in particular, that the system is semi-globally exactly controllable and semi-globally exponentially stabilizable in the space  $H^1(M)$  assuming that both the geometric control condition and a unique continuation condition are satisfied (see [8] for more details). The work [3] extended this one: The authors studied global controllability and stabilization properties for the fractional Schrödinger equation on  $d$ -dimensional compact Riemannian manifolds without boundary  $(M, g)$ . They used microlocal analysis to show the propagation of regularity which, together with the geometric control condition (GCC) and a unique continuation property, allowed them to prove global control results.

We conclude by mentioning a recent work. In a very interesting work [32], Taüfer demonstrated that the controllability of the linear Schrödinger equation in  $\mathbb{R}^d$  holds for any time  $T > 0$  with internal control supported on nonempty, periodic, open sets. This result, in particular, showed that the controllability of the linear Schrödinger equation in full space extends to a strictly larger class of control supports than that of the wave equation.

**1.2. Main results.** As discussed above, most of the available results at the moment are for the classical Schrödinger equation (1.4) in different domains and considering the control inputs acting on the whole boundary. However, in the critical case, namely, system (1.2), the internal control problem remains open.

Motivated by the ideas contained in [27], let us present the first answer to the control problem stated at the beginning of the section. To do this, consider the control system

$$(1.5) \quad \begin{cases} i\partial_t u + \Delta u - |u|^4 u = \varphi(x)h(x, t), & \text{on } \mathbb{R}^3 \times [0, +\infty), \\ u(x, 0) = u_0(x) \in H^1(\mathbb{R}^3), \end{cases}$$

where the function  $\varphi \in C^\infty(\mathbb{R}^3, [0, 1])$  and satisfies

$$(1.6) \quad \varphi(x) = \begin{cases} 0, & \text{if } |x| \leq R, \\ 1, & \text{if } |x| \geq R + 1, \end{cases}$$

for some  $R > 0$  large enough. Our result below gives a first answer in this direction.

**Theorem 1.1** (Local controllability). *Let  $T > 0$  be given. There exists  $\delta > 0$  such that for any  $u_0 \in H^1(\mathbb{R}^3)$  satisfying*

$$\|u_0\|_{H^1} \leq \delta,$$

*one can find  $h(x, t) \in C(\mathbb{R}; H^1(\mathbb{R}^3))$  such that system (1.5) admits a solution  $u \in C([0, T]; H^1(\mathbb{R}^3))$  satisfying  $u(T) = 0$ .*

In our arguments, Strichartz-type inequalities play a fundamental role in giving the well-posedness of the system (1.5). In addition, as a first step, we have used a unique continuation ensured by a Carleman estimate proved by [21] (see also [17]).

**Theorem 1.2.** *For every initial data  $u_0 \in H^1(\mathbb{R}^3)$  and every  $T > 0$ , there exists a control  $h(x, t) \in C(\mathbb{R}; H^1(\mathbb{R}^3))$  with support in  $\mathbb{R} \times (\mathbb{R}^3 \setminus B_R(0))$ ,  $R > 0$ , such that the unique solution of the linear system associated to (1.5) satisfies  $u(T, \cdot) = 0$ .*

With Theorem 1.2 in hand, a perturbation argument ensures that we can get a local controllability result for the critical Schrödinger equation (1.5), so giving Theorem 1.1.

**1.3. Structure of this work.** We finish our introduction by giving an outline of this work. It is divided as follows:

- In Section 2, we give auxiliary results that are important in establishing our control result. Precisely, we present a review of the Cauchy problem for the Schrödinger equation.
- In Section 3, we present the proof of our main results. Firstly, we show the local null controllability around the null trajectories for the linear system associated with (1.5), proving Theorem 1.2. Then, through a perturbation argument, we show Theorem 1.1.
- Finally, we discuss some future perspectives in Section 4.

## 2. A REVIEW OF THE CAUCHY PROBLEM

**2.1. Smoothing.** For the sake of completeness, we discuss the smoothing properties of the linear Schrödinger equation,

$$(2.1) \quad \begin{cases} i\partial_t u + \Delta u = 0, & \text{on } \mathbb{R}^3 \times \mathbb{R}, \\ u(x, 0) = \psi(x), & x \in \mathbb{R}^3, \end{cases}$$

which will play an important role in establishing the controllability for the defocusing critical nonlinear Schrödinger equation (1.5). To this end, for  $j \in \{1, 2, 3\}$ , let  $P_j$  be the differential operator on  $\mathbb{R}^4$  defined by

$$(2.2) \quad P_j v(t, x) = (x_j + 2it\partial_j)v(t, x).$$

For a multi-index  $\alpha$ , define the differential operator  $P_\alpha$  on  $\mathbb{R}^4$  by

$$P_\alpha = \prod_{j=1}^3 P_j^{\alpha_j}.$$

Additionally, for  $x \in \mathbb{R}^3$ , consider

$$x^\alpha = \prod_{j=1}^3 x_j^{\alpha_j}.$$

For any smooth function  $u(t, x)$ , one has

$$P_j u(t, x) = 2ite^{i\frac{|x|^2}{4t}} \frac{\partial}{\partial x_j} (e^{-i\frac{|x|^2}{4t}} u(t, x)).$$

Indeed, note that

$$\begin{aligned} 2ite^{i\frac{|x|^2}{4t}} \frac{\partial}{\partial x_j} (e^{-i\frac{|x|^2}{4t}} u) &= -2ite^{i\frac{|x|^2}{4t}} \frac{2ix_j}{4t} e^{-i\frac{|x|^2}{4t}} u(t, x) + 2ite^{i\frac{|x|^2}{4t}} e^{-i\frac{|x|^2}{4t}} \frac{\partial}{\partial x_j} u(t, x) \\ &= x_j u(t, x) + 2it \frac{\partial}{\partial x_j} u(t, x). \end{aligned}$$

Hence,

$$P_\alpha u(t, x) = (2it)^{|\alpha|} e^{i\frac{|x|^2}{4t}} D^\alpha (e^{-i\frac{|x|^2}{4t}} u(t, x)).$$

On the other hand, we easily obtain

$$[P_j, i\partial_t + \Delta] = 0.$$

Thus, considering  $u \in C(\mathbb{R}, H^1(\mathbb{R}^3))$  to be any solution of the linear Schrödinger equation (2.1), one has that  $P_j u$  and  $P_\alpha u$  is also a solution.

Now, with the previous analysis in hand and taking into account the relation (2.2), we present the next result, which gives a local smoothing property for the linear Schrödinger equation (2.1).

**Proposition 2.1.** *Let  $\alpha$  be a multi-index and  $T > 0$  be given. Let  $\psi \in H^1(\mathbb{R}^3)$  be such that  $x^\alpha \psi \in H^1(\mathbb{R}^3)$ . Then, the corresponding solution  $u$  of the IVP (2.1) satisfies*

$$P_\alpha u \in C(\mathbb{R}; H^1(\mathbb{R}^3))$$

and there exists a constant  $C$  depending only on  $T$  and  $\alpha$  such that

$$\|P_\alpha u\|_{H^1(\mathbb{R}^3)} \leq C \|x^\alpha \psi\|_{H^1(\mathbb{R}^3)}$$

holds for any  $t \in [-T, T]$ . In particular, if  $\psi \in H^1(\mathbb{R}^3)$  has compact support, then  $u$  is infinitely smooth everywhere except at  $t = 0$ .

*Proof.* A standard density argument assures that it is sufficient to prove the result for  $\psi \in \mathcal{S}(\mathbb{R}^3)$ . To this end, assume, first, that  $|\alpha| = 1$ , so that  $P_\alpha = P_j$  for some  $j \in \{1, 2, 3\}$ . Now, note that

$$\|u(t)\|_{H^1(\mathbb{R}^3)} = \|\psi\|_{H^1(\mathbb{R}^3)},$$

for any  $t \in [-T, T]$ . Let  $w^j(t, x) = P_j u(t, x)$ . Applying the operator  $P_j$  to (2.1) yields

$$\begin{cases} i\partial_t w^j + \Delta w^j = 0, \\ w^j(0, x) = x_j \psi, \end{cases}$$

due to the fact that  $P_j u(0, x) = x_j u(0, x)$ . Since

$$w^j(t) = e^{it\Delta}(x_j \psi),$$

we get that

$$\|w^j(t)\|_{H^1(\mathbb{R}^3)} \leq C \|x_j \psi\|_{H^1(\mathbb{R}^3)}.$$

The general case ( $|\alpha| > 1$ ) follows by induction.  $\square$

**2.2. Local existence.** Here, we are interested in studying the  $H^1$  critical, defocusing, Cauchy problem for the nonlinear Schrödinger equation (C-NLS)

$$(2.3) \quad \begin{cases} i\partial_t u + \Delta u - |u|^4 u = g, & \text{on } [0, T] \times \mathbb{R}^3, \\ u(0) = u_0 \in H^1(\mathbb{R}^3), \end{cases}$$

where  $g(t, x) = g \in L_{loc}^\infty(\mathbb{R}, H^1(\mathbb{R}^3))$  and  $T > 0$ . Let us first present some definitions used throughout the paper.

*Definition 1.* A pair  $(q, r)$  is called  $L^2$ -admissible if  $r \in [2, 6]$  and  $q$  satisfies

$$\frac{2}{q} + \frac{3}{r} = \frac{3}{2}.$$

Such a pair is called  $H^1$ -admissible if  $r \in [6, +\infty)$  and  $q$  satisfies

$$\frac{2}{q} + \frac{3}{r} = \frac{1}{2}.$$

With these definitions in hand, we present two results that are paramount to prove that the Cauchy problem (2.3) is well-posed. The first one is *Strichartz estimates* and the second one is a standard Sobolev embedding. These results can be found in [7, 11].

**Lemma 2.2.** *Let  $(q, r)$  be a  $L^2$ -admissible pair. We have*

$$(2.4) \quad \|e^{it\Delta} h\|_{L_t^q L_x^r} \leq c \|h\|_{L_x^2},$$

$$(2.5) \quad \left\| \int_0^T e^{i(t-\tau)\Delta} g \, d\tau \right\|_{L_t^q L_x^r} + \left\| \int_0^t e^{i(t-\tau)\Delta} g \, d\tau \right\|_{L_t^q L_x^r} \leq c \|g\|_{L_t^{m'} L_x^{n'}},$$

and

$$\left\| \int_0^T e^{it\Delta} g(\tau) \, d\tau \right\|_{L_x^2} \leq C \|g\|_{L_t^{q'} L_x^{r'}}.$$

where  $(q, r)$ ,  $(m, n)$  are any  $L^2$ -admissible pair.

**Lemma 2.3.** For  $v \in L^{10}([0, T]; L^{10}(\mathbb{R}^3))$  such that  $\nabla v \in L^{10}([0, T]; L^{\frac{10}{13}}(\mathbb{R}^3))$ , we have

$$\|v\|_{L_t^{10} L_x^{10}} \leq C \|\nabla v\|_{L_t^{10} L_x^{\frac{30}{13}}}.$$

For an interval  $I$ , define the norms  $S(I)$ ,  $W(I)$  and  $Z(I)$  by

$$\|u\|_{S(I)} = \|u\|_{L^{10}(I; L^{10}(\mathbb{R}^3))}, \quad \|u\|_{Z(I)} = \|u\|_{L^{10}(I; L^{\frac{30}{13}}(\mathbb{R}^3))} \quad \text{and} \quad \|u\|_{W(I)} = \|u\|_{L^{\frac{10}{3}}(I; L^{\frac{10}{3}}(\mathbb{R}^3))}.$$

The following theorem gives us the solution to problem (2.3).

**Theorem 2.4.** Let  $u_0 \in H^1(\mathbb{R}^3)$ . If  $\|u_0\|_{H^1}$  is small enough, then there exists an interval  $I$  and an unique solution  $u(t, x)$  of problem (2.3) in  $I \times \mathbb{R}^3$ , with  $u \in C(I; H^1(\mathbb{R}^3))$ , satisfying

$$\|\nabla u\|_{W(I)} < \infty, \quad \|u\|_{S(I)} < \infty \quad \text{and} \quad \|\nabla u\|_{Z(I)} < \infty.$$

*Proof.* We follow the ideas from [12]. Assume, without loss of generality, that  $t_0 = 0$ . Observe that the Cauchy problem (2.3) is equivalent to the integral equation (Duhamel's formula)

$$u(t) = e^{it\Delta} u_0 - \int_0^t e^{i(t-\tau)\Delta} [|u|^4 u + g] d\tau.$$

Define

$$\| \|u\| \| = \sup_{t \in I} \|u(t)\|_{L^2} + \sup_{t \in I} \|\nabla u(t)\|_{L^2} + \|u\|_{S(I)} + \|\nabla u\|_{W(I)} + \|\nabla u\|_{Z(I)}.$$

For  $R > 0$  to be conveniently chosen later on, consider the set

$$B_R := \left\{ u(t, x) \text{ on } I \times \mathbb{R}^3 : \| \|u\| \| \leq R \right\}.$$

We want to show that the operator  $\Phi_{u_0} : B_R \rightarrow B_R$  defined by

$$\Phi_{u_0}(u) = e^{it\Delta} u_0 - \int_0^t e^{i(t-\tau)\Delta} [|u|^4 u + g] d\tau$$

has a fixed point for  $R$  small enough. To this end, first, observe that

$$\begin{aligned} \|\Phi_{u_0}(u)\|_{L_x^2} &\leq \|e^{it\Delta} u_0\|_{L_x^2} + \left\| \int_0^t \nabla e^{i(t-\tau)\Delta} [|u|^4 u + g] d\tau \right\|_{L_x^2} \\ &\leq \|u_0\|_{L^2} + C \| |u|^4 u \|_{L_t^1 L_x^2} + \|g\|_{L_t^1 L_x^2} \\ &\leq C \|u_0\|_{H^1} + C \|u\|_{S(I)}^5 + C_I \|g\|_{L_t^\infty H_x^1} \end{aligned}$$

and

$$\begin{aligned} \|\nabla \Phi_{u_0}(u)\|_{L_x^2} &\leq \|\nabla e^{it\Delta} u_0\|_{L_x^2} + \left\| \int_0^t \nabla e^{i(t-\tau)\Delta} [|u|^4 u + g] d\tau \right\|_{L_x^2} \\ &\leq \|\nabla u_0\|_{L^2} + C \|\nabla |u|^4 u\|_{L_t^{\frac{10}{7}} L_x^{\frac{10}{7}}} + \|\nabla g\|_{L_t^1 L_x^2} \\ &\leq C \|u_0\|_{H^1} + C \|u\|_{S(I)}^4 \|\nabla u\|_{W(I)} + C_I \|g\|_{L_t^\infty H_x^1}. \end{aligned}$$

Choosing the length of  $I$  small enough such that  $C_I \|g\|_{L_t^\infty H_x^1} \leq C \|u_0\|_{H^1}$ , we have

$$\|\nabla \Phi_{u_0}(u)\|_{L_x^2} \leq 2C \|u_0\|_{H^1} + CR^5.$$

Secondly, notice that

$$\begin{aligned} \|\nabla \Phi_{u_0}(u)\|_{W(I)} &\leq \|\nabla e^{it\Delta} u_0\|_{W(I)} + \left\| \int_0^t \nabla e^{i(t-\tau)\Delta} [|u|^4 u + g] d\tau \right\|_{W(I)} \\ &\leq \|\nabla u_0\|_{L^2} + C \|\nabla |u|^4 u\|_{L_t^{\frac{10}{7}} L_x^{\frac{10}{7}}} + \|\nabla g\|_{L_t^1 L_x^2}. \end{aligned}$$

Due to Hölder's inequality with  $p = \frac{7}{4}$  and  $q = \frac{7}{3}$ , we get

$$\|\nabla |u|^4 u\|_{L_t^{\frac{10}{7}} L_x^{\frac{10}{7}}} \leq C \|u\|_{S(I)}^4 \|\nabla u\|_{W(I)}.$$

Thus

$$\begin{aligned}\|\nabla\Phi_{u_0}(u)\|_{W(I)} &\leq C\left(\|\nabla u_0\|_{L^2} + \|u\|_{S(I)}^4\|\nabla u\|_{W(I)} + \|\nabla g\|_{L_t^1 L_x^2}\right) \\ &\leq C\|u_0\|_{H^1} + CR^5 + C_I\|g\|_{L_t^\infty H_x^1}.\end{aligned}$$

Choosing the length of  $I$  small enough such that  $C_I\|g\|_{L_t^\infty H_x^1} \leq C\|u_0\|_{H^1}$ , one gets

$$\|\nabla\Phi_{u_0}(u)\|_{W(I)} \leq 2C\|u_0\|_{H^1} + CR^5.$$

On other hand, using estimate (2.4) with  $q = 10$  and  $r = 30/13$ , due to the inequality (2.5) with the same  $q, r$  and  $m' = 2$  and  $n' = \frac{6}{5}$ , Hölder's inequality gives

$$\begin{aligned}\|\nabla\Phi_{u_0}(u)\|_{Z(I)} &\leq \|\nabla e^{it\Delta}u_0\|_{Z(I)} + \left\| \int_0^t \nabla e^{i(t-\tau)\Delta}[|u|^4u + g] d\tau \right\|_{Z(I)} \\ &\leq \|\nabla u_0\|_{L^2} + C\|\nabla|u|^4u\|_{L_t^2 L_x^{\frac{6}{5}}} + C\|\nabla g\|_{L_t^1 L_x^2} \\ &\leq \|\nabla u_0\|_{L^2} + C\|\nabla u\|_{Z(I)}\|u\|_{S(I)}^4 + C\|\nabla g\|_{L_t^1 L_x^2} \\ &\leq C\|u_0\|_{H^1} + CR^5 + C_I\|g\|_{L_t^\infty H_x^1} \\ &\leq 2C\|u_0\|_{H^1} + CR^5\end{aligned}$$

since  $C_I\|g\|_{L_t^\infty H_x^1} \leq C\|u_0\|_{H^1}$ . Finally,

$$\begin{aligned}\|\Phi_{u_0}(u)\|_{S(I)} &\leq \|\nabla\Phi_{u_0}(u)\|_{Z(I)} \\ &\leq \|\nabla e^{it\Delta}u_0\|_{Z(I)} + \left\| \int_0^t \nabla e^{i(t-\tau)\Delta}[|u|^4u + g] d\tau \right\|_{Z(I)} \\ &\leq \|\nabla u_0\|_{L^2} + C\|\nabla u\|_{Z(I)}\|u\|_{S(I)}^4 + C\|\nabla g\|_{L_t^1 L_x^2} \\ &\leq C\|u_0\|_{H^1} + CR^5 + C_I\|g\|_{L_t^\infty H_x^1} \\ &\leq 2C\|u_0\|_{H^1} + CR^5,\end{aligned}$$

since  $C_I\|g\|_{L_t^\infty H_x^1} \leq C\|u_0\|_{H^1}$ . Summing up, we get

$$\|\Phi_{u_0}(u)\| \leq 2C\|u_0\|_{H^1} + CR^5 \leq R$$

as long as  $\|u_0\|_{H^1} \leq \frac{R}{2C} - \frac{R^5}{2}$ , where  $R < (1/C)^{1/4}$ .

Next, denoting  $f(u) = |u|^4u$ , we get

$$\begin{aligned}\|\Phi_{u_0}(u) - \Phi_{u_0}(v)\|_{L_x^2} &\leq C\|f(u) - f(v)\|_{L_t^1 L_x^2} \\ &\leq C\|u - v\|_{S(I)}(\|u\|_{S(I)}^4 + \|v\|_{S(I)}^4).\end{aligned}$$

Moreover, we also have

$$\begin{aligned}\|\nabla\Phi_{u_0}(u) - \nabla\Phi_{u_0}(v)\|_{L_x^2} &\leq C\|\nabla f(u) - \nabla f(v)\|_{L_t^{\frac{10}{7}} L_x^{\frac{10}{7}}} \\ &\leq C\left(\left\| |u|^4|\nabla u - \nabla v| \right\|_{L_t^{\frac{10}{7}} L_x^{\frac{10}{7}}} + \left\| |u - v||u|^3|\nabla v| \right\|_{L_t^{\frac{10}{7}} L_x^{\frac{10}{7}}} \right. \\ &\quad \left. + \left\| |u - v||v|^3|\nabla v| \right\|_{L_t^{\frac{10}{7}} L_x^{\frac{10}{7}}} \right) \\ &\leq C\left(\|u\|_{S(I)}^4\|\nabla u - \nabla v\|_{W(I)} + \|u - v\|_{S(I)}\|\nabla v\|_{W(I)}\|u\|_{S(I)}^3 + \right. \\ &\quad \left. + \|u - v\|_{S(I)}\|\nabla v\|_{W(I)}\|v\|_{S(I)}^3\right) \\ &\leq CR^4\|u - v\|_{S(I)} + CR^4\|\nabla u - \nabla v\|_{W(I)},\end{aligned}$$

and

$$\begin{aligned}
\|\nabla\Phi_{u_0}(u) - \nabla\Phi_{u_0}(v)\|_{W(I)} &\leq C\|\nabla f(u) - \nabla f(v)\|_{L_t^{\frac{10}{7}}L_x^{\frac{10}{7}}} \\
&\leq C\left(\| |u|^4|\nabla u - \nabla v| \|_{L_t^{\frac{10}{7}}L_x^{\frac{10}{7}}} + \| |u-v||u|^3|\nabla v| \|_{L_t^{\frac{10}{7}}L_x^{\frac{10}{7}}} \right. \\
&\quad \left. + \| |u-v||v|^3|\nabla v| \|_{L_t^{\frac{10}{7}}L_x^{\frac{10}{7}}} \right) \\
&\leq C\left(\|u\|_{S(I)}^4\|\nabla u - \nabla v\|_{W(I)} + \|u-v\|_{S(I)}\|\nabla v\|_{W(I)}\|u\|_{S(I)}^3 + \right. \\
&\quad \left. + \|u-v\|_{S(I)}\|\nabla v\|_{W(I)}\|v\|_{S(I)}^3\right) \\
&\leq CR^4\|u-v\|_{S(I)} + CR^4\|\nabla u - \nabla v\|_{W(I)}.
\end{aligned}$$

Following the same reasoning as before,

$$\begin{aligned}
\|\nabla\Phi_{u_0}(u) - \nabla\Phi_{u_0}(v)\|_{Z(I)} &\leq C\left(\|u\|_{S(I)}^4\|\nabla u - \nabla v\|_{Z(I)} + \|u-v\|_{S(I)}\|\nabla v\|_{Z(I)}\|u\|_{S(I)}^3 + \right. \\
&\quad \left. + \|u-v\|_{S(I)}\|\nabla v\|_{Z(I)}\|v\|_{S(I)}^3\right) \\
&\leq CR^4\|\nabla u - \nabla v\|_{Z(I)} + CR^4\|u-v\|_{S(I)}.
\end{aligned}$$

Moreover, by Sobolev embedding, we have

$$\|\Phi_{u_0}(u) - \Phi_{u_0}(v)\|_{S(I)} \leq \|\nabla\Phi_{u_0}(u) - \nabla\Phi_{u_0}(v)\|_{Z(I)} \leq CR^4\|\nabla u - \nabla v\|_{Z(I)} + CR^4\|u-v\|_{S(I)}.$$

Summing up yields

$$\begin{aligned}
\|\Phi_{u_0}(u) - \Phi_{u_0}(v)\| &\leq CR^4\|\nabla u - \nabla v\|_{Z(I)} + CR^4\|u-v\|_{S(I)} + CR^4\|\nabla u - \nabla v\|_{W(I)} \\
&\leq CR^4\|\nabla u - \nabla v\|_{Z(I)} + CR^4\|u-v\|_{S(I)} + CR^4\|\nabla u - \nabla v\|_{W(I)} \\
&\quad + CR^4\sup_{t \in I}\|\nabla u(t) - \nabla v(t)\|_{L_x^2} \\
&\leq CR^4\|u-v\|
\end{aligned}$$

Thus, if  $R > 0$  is such that  $CR^4 < 1$ , then  $\Phi_{u_0}$  is a contraction in  $B_R$  and, therefore, has a unique fixed point, i.e., problem (2.3) has a local solution defined on a maximal interval  $[0, T]$ .  $\square$

### 3. CONTROLLABILITY FOR THE CRITICAL NONLINEAR SCHRÖDINGER EQUATION

In this section, we prove Theorem 1.1 using a duality strategy which reduces the controllability problem (1.5) to prove an observability inequality, the so-called ‘‘Hilbert Uniqueness Method’’ [19], for the solutions of the linear system

$$(3.1) \quad \begin{cases} i\partial_t u + \Delta u = \varphi(x)h(x, t), & \text{on } (0, T) \times \mathbb{R}^3, \\ u(0) = u_0, \end{cases}$$

where  $\varphi = \varphi(x)$  is defined by (1.6).

**3.1. Linear Schrödinger equation: Null controllability.** In this section, we will show Theorem 1.2. Note that the null controllability of system (3.1) follows from the observability inequality, namely,

$$(3.2) \quad \|v_0\|_{H^{-1}}^2 \leq C \int_0^T \|\varphi v(t)\|_{H^{-1}}^2 dt,$$

where  $v(x, t)$  is a solution to the adjoint system associated to (3.1)

$$(3.3) \quad \begin{cases} i\partial_t v + \Delta v = 0 \text{ on } \mathbb{R} \times \mathbb{R}^3, \\ v(0) = v_0 \in H^{-1}(\mathbb{R}^3). \end{cases}$$

The observability inequality (3.2) is given by the following result.

**Proposition 3.1.** *Let  $\varphi$  be a  $C^\infty$  real function on  $\mathbb{R}^3$  as in (1.6). Then, for every  $T > 0$ , there exists a constant  $C = C(T) > 0$  such that inequality (3.2) holds for every solution  $v(t, x)$  of system (3.3).*

*Proof.* We will split the proof into several steps.

**First step:**  $H^1$ -observability.

**Lemma 3.2.** *Consider the system*

$$(3.4) \quad \begin{cases} i\partial_t w + \Delta w = 0, \text{ on } (0, T) \times \mathbb{R}^3, \\ w(0) = w_0 \in H^1(\mathbb{R}^3). \end{cases}$$

There exists a constant  $C > 0$  such that for each  $w_0 \in H^1(\mathbb{R}^3)$ , the solution  $w(t)$  of (3.4) satisfies

$$(3.5) \quad \|w_0\|_{H^1(\mathbb{R}^3)}^2 \leq C \int_0^T \|\varphi w(t)\|_{H^1(\mathbb{R}^3)}^2 dt.$$

*Proof.* Let  $q \in C_0^\infty(\mathbb{R}^3; \mathbb{R}^3)$  such that

$$q(x) = \begin{cases} x, & \text{if } |x| \leq R + 2, \\ 0, & \text{if } |x| \geq R + 3. \end{cases}$$

Multiplying the equation in (3.4) by  $q \cdot \nabla \bar{w} + \frac{1}{2} \bar{w}(\operatorname{div}_x q)$ , taking the real part and integrating by parts gives

$$(3.6) \quad \begin{aligned} \frac{1}{2} \operatorname{Im} \int_{\mathbb{R}^3} (wq \cdot \nabla \bar{w}) dx \Big|_0^T + \frac{1}{2} \operatorname{Re} \int_0^T \int_{\mathbb{R}^3} w \nabla(\operatorname{div}_x q) \cdot \nabla \bar{w} dx dt \\ + \operatorname{Re} \int_0^T \int_{\mathbb{R}^3} \sum_{j,k=1}^3 \left( \frac{\partial q_k}{\partial x_j} \frac{\partial \bar{w}}{\partial x_k} \frac{\partial w}{\partial x_j} \right) dx dt = 0, \end{aligned}$$

where we have used the fact that the function  $q(x)$  has compact support. Notice that there is conservation of energy in  $H^1(\mathbb{R}^3)$

$$(3.7) \quad \|w(t)\|_{H^1(\mathbb{R}^3)}^2 = \|w(0)\|_{H^1(\mathbb{R}^3)}^2 = \|w_0\|_{H^1(\mathbb{R}^3)}^2, \quad t \in [0, T].$$

It follows from (3.6) that

$$\begin{aligned} \int_0^T \int_{B_{R+2}(0)} |\nabla w|^2 dx dt &\leq C_\varepsilon \left( \int_0^T \int_{B_{R+3}(0) \setminus B_{R+2}(0)} |\nabla w|^2 dx dt + \|w_0\|_{L^2(\mathbb{R}^3)}^2 \right) \\ &\quad + \varepsilon \int_0^T \|w(t)\|_{H^1(\mathbb{R}^3)}^2 dt, \end{aligned}$$

for any  $\varepsilon > 0$  and some constant  $C_\varepsilon > 0$ . We also have

$$\|w(t)\|_{H^1(\mathbb{R}^3)}^2 \leq C_R \left( \int_{B_{R+2}(0)} |\nabla w|^2 dx + \|\varphi w(t)\|_{H^1(\mathbb{R}^3)}^2 \right)$$

Indeed, observe that

$$\begin{aligned} \|\nabla w(t)\|_{L^2(\mathbb{R}^3)}^2 &= \|\nabla w(t)\|_{L^2(B_{R+1}(0))}^2 + \|\nabla w(t)\|_{L^2(\mathbb{R}^3 \setminus B_{R+1}(0))}^2 \\ &\leq \|\nabla w(t)\|_{L^2(B_{R+2}(0))}^2 + \|\varphi w(t)\|_{H^1(\mathbb{R}^3)}^2 \end{aligned}$$

and that

$$\begin{aligned} \|w(t)\|_{L^2(\mathbb{R}^3)}^2 &= \|w(t)\|_{L^2(B_{R+1}(0))}^2 + \|w(t)\|_{L^2(\mathbb{R}^3 \setminus B_{R+1}(0))}^2 \\ &\leq |B_{R+1}(0)|^{\frac{2}{3}} \|w(t)\|_{L^6(B_{R+1}(0))}^2 + \|\varphi w(t)\|_{L^2(\mathbb{R}^3 \setminus B_{R+1}(0))}^2 \\ &\leq |B_{R+1}(0)|^{\frac{2}{3}} \|w(t)\|_{L^6(\mathbb{R}^3)}^2 + \|\varphi w(t)\|_{H^1(\mathbb{R}^3)}^2 \end{aligned}$$

and using the Gagliardo-Nirenberg-Sobolev inequality

$$\|w(t)\|_{L^6(\mathbb{R}^3)} \leq C_1 \|\nabla w(t)\|_{L^2(\mathbb{R}^3)}$$

shows the claim. Moreover, we also obtain

$$(3.8) \quad \|w_0\|_{H^1(\mathbb{R}^3)}^2 = \frac{1}{T} \int_0^T \|w(t)\|_{H^1(\mathbb{R}^3)}^2 dt \leq C \left( \int_0^T \|\varphi w(t)\|_{H^1(\mathbb{R}^3)}^2 dt + \|w_0\|_{L^2(\mathbb{R}^3)}^2 \right).$$

Indeed we have

$$\begin{aligned} \int_0^T \|w(t)\|_{H^1(\mathbb{R}^3)}^2 dt &\leq C_R \left( \int_0^T \|\nabla w(t)\|_{L^2(B_{R+2}(0))}^2 dt + \int_0^T \|\varphi w(t)\|_{H^1(\mathbb{R}^3)}^2 dt \right) \\ &\leq C_\varepsilon C_R \left( \int_0^T \int_{B_{R+3}(0) \setminus B_{R+2}(0)} |\nabla w|^2 dx dt + \|w_0\|_{L^2(\mathbb{R}^3)}^2 \right) \\ &\quad + C_R \int_0^T \|\varphi w(t)\|_{H^1(\mathbb{R}^3)}^2 dt + C_R \varepsilon \int_0^T \|w(t)\|_{H^1(\mathbb{R}^3)}^2 dt \\ &\leq C_\varepsilon C_R \|w_0\|_{L^2(\mathbb{R}^3)}^2 + (C_\varepsilon + 1) C_R \int_0^T \|\varphi w(t)\|_{H^1(\mathbb{R}^3)}^2 dt \\ &\quad + C_R \varepsilon \int_0^T \|w(t)\|_{H^1(\mathbb{R}^3)}^2 dt \end{aligned}$$

and it suffices to choose  $\varepsilon$  small enough to absorb the second right-hand side term in the left hand-side term. So, it remains to show

$$(3.9) \quad \|w_0\|_{L^2(\mathbb{R}^3)}^2 \leq c \int_0^T \|\varphi w(t)\|_{H^1(\mathbb{R}^3)}^2 dt$$

to achieve Lemma 3.2. To this end, let us argue by contradiction, that is, suppose that (3.9) does not hold. If this is the case, there exists a sequence  $\{w_{n,0}\}$  in  $H^1(\mathbb{R}^3)$  such that the corresponding sequence of solutions  $\{w_n\}$  of (3.4) satisfies

$$(3.10) \quad 1 = \|w_{n,0}(t)\|_{L^2(\mathbb{R}^3)}^2 \geq n \int_0^T \|\varphi w_n(t)\|_{H^1(\mathbb{R}^3)}^2 dt, \quad n = 1, 2, \dots$$

Due to inequalities (3.8) and (3.10), we get

$$\|w_{n,0}\|_{H^1(\mathbb{R}^3)}^2 = \frac{1}{T} \int_0^T \|w_n(t)\|_{H^1(\mathbb{R}^3)}^2 dt \leq C \left( \int_0^T \|\varphi w_n(t)\|_{H^1(\mathbb{R}^3)}^2 dt + \|w_{n,0}\|_{L^2(\mathbb{R}^3)}^2 \right) \leq 2C,$$

hence the sequence  $\{w_n(0) = w_{n,0}\}_{n \in \mathbb{N}}$  is bounded in  $H^1(\mathbb{R}^3)$ . Extracting a subsequence and still denoting it by  $\{w_{n,0}\}$ , we may assume that

$$w_{n,0} \rightharpoonup w_0 \text{ weakly in } H^1(\mathbb{R}^3)$$

and

$$w_n \rightharpoonup w \text{ weakly in } L^2((0, T); H^1(\mathbb{R}^3)),$$

where  $w \in C([0, T]; H^1(\mathbb{R}^3))$  is a solution of (3.4). By inequality (3.10),

$$\varphi w_n \rightarrow 0 \text{ in } L^2((0, T); H^1(\mathbb{R}^3)) \text{ strongly.}$$

Since  $\varphi w_n \rightharpoonup 0$  in  $L^2((0, T); H^1(\mathbb{R}^3))$  weakly, we conclude that  $\varphi w \equiv 0$  on  $(0, T) \times \mathbb{R}^3$ . Therefore,

$$w \equiv 0, \quad |x| > R + 1, \quad t \in (0, T).$$

According to Proposition 2.1, one has  $w \in C^\infty(\mathbb{R}^3 \times (0, T))$ . Now, we are in a position to use the unique continuation property for the Schrödinger equation showed in [21] to conclude that

$$w \equiv 0 \text{ on } \mathbb{R}^3 \times (0, T).$$

Since  $\varphi w_n \rightarrow 0$  strongly in  $L^2((0, T); H^1(\mathbb{R}^3))$ , we get

$$(3.11) \quad w_n \rightarrow 0 \text{ strongly in } L^2((0, T); H^1(\mathbb{R}^3 \setminus B_{R+1}(0))).$$

On the other hand taking into account (3.8) we have

$$\frac{1}{T} \int_0^T \|w_n(t)\|_{H^1(B_{R+1}(0))}^2 dt \leq C \left( \int_0^T \|\varphi w_n(t)\|_{H^1(\mathbb{R}^3)}^2 dt + \|w_{n,0}\|_{L^2(\mathbb{R}^3)}^2 \right) \leq 2C,$$

and using the equation (3.8),

$$\begin{aligned} \int_0^T \|\partial_t w_n(t)\|_{H^{-1}(B_{R+1}(0))}^2 dt &= \int_0^T \|\Delta w_n(t)\|_{H^{-1}(B_{R+1}(0))}^2 dt \\ &\leq C \int_0^T \|w_n(t)\|_{H^1(B_{R+1}(0))}^2 dt \end{aligned}$$

as well as

$$\int_0^T \|\nabla w_n(t)\|_{H^{-1}(B_{R+1}(0))}^2 dt \leq C \int_0^T \|w_n(t)\|_{H^1(\mathbb{R}^3)}^2 dt.$$

Therefore,

$$w_n \text{ is bounded in } L^2((0, T); H^1(B_{R+1}(0))) \cap H^1((0, T); H^{-1}(B_{R+1}(0))).$$

Due to Aubin's lemma (see [29]) and the convergence (3.11), we conclude that for a subsequence, still denoted by  $\{w_n\}$ ,

$$w_n \rightarrow w = 0 \text{ strongly in } L^2((0, T); L^2(\mathbb{R}^3))$$

which contradicts (3.10). So, the estimate (3.5) follows from (3.8) and (3.9) showing the lemma.  $\square$

**Second step: Weak observability inequality.**

We prove a bound which is weaker than the observability inequality (3.2).

**Lemma 3.3.** *Let  $v$  be the solution of system (3.3) with  $v_0 \in H^{-1}(\mathbb{R}^3)$ . Then,*

$$(3.12) \quad \|v_0\|_{H^{-1}}^2 \leq C \left( \int_0^T \|\varphi v(t)\|_{H^{-1}}^2 dt + \|(1 - \varphi(x/2))v_0\|_{H^{-2}}^2 \right).$$

*Proof.* Again, let us argue by contradiction. If inequality (3.12) is not verified, there exists a sequence  $\{v_n\}$  of solutions to problem (3.3) in  $C([0, T]; H^{-1}(\mathbb{R}^3))$  such that

$$(3.13) \quad 1 = \|v_n(0)\|_{H^{-1}}^2 \geq n \left( \int_0^T \|\varphi v_n(t)\|_{H^{-1}}^2 dt + \|(1 - \varphi(x/2))v_n(0)\|_{H^{-2}}^2 \right).$$

Up to a subsequence, we may assume that

$$v_n \rightharpoonup v \text{ in } L^\infty((0, T); H^{-1}(\mathbb{R}^3)) \text{ weak}^*$$

and

$$(3.14) \quad v_n(0) \rightharpoonup v(0) \text{ in } H^{-1}(\mathbb{R}^3) \text{ weak},$$

where  $v \in C([0, T]; H^{-1}(\mathbb{R}^3))$  is a solution of problem (3.3). By inequality (3.13),

$$\varphi v_n \rightarrow 0 \text{ (strongly) in } L^2((0, T); H^{-1}(\mathbb{R}^3)).$$

Since

$$\varphi v_n \rightharpoonup \varphi v \text{ in } L^\infty((0, T); H^{-1}(\mathbb{R}^3)) \text{ weak}^*,$$

we conclude that  $\varphi v \equiv 0$ . Therefore,  $v(t, x) = 0$  for  $|x| > R + 1$  and  $t \in (0, T)$ . So, using the unique continuation property as in Step 1, we get that  $v \equiv 0$ . In particular,  $v(0) = 0$ .

Now, we claim that

$$(3.15) \quad \|\varphi(x/2)v_n(0)\|_{H^{-2}}^2 \leq C \int_0^T \|\varphi v_n(t)\|_{H^{-1}}^2 dt.$$

To prove (3.15), introduce the function  $\tilde{v}_n(x, t) = \varphi(x/2)v_n(x, t)$  which satisfies

$$i\partial_t \tilde{v}_n + \Delta \tilde{v}_n = f_n$$

where  $f_n = [\Delta \varphi(x/2)]v_n + 2\nabla \varphi(x/2)\nabla v_n$ . Then, the fact that  $\text{supp}[\varphi(x/2)] \subset \{\varphi = 1\}$  yields

$$\begin{aligned} \|\tilde{v}_n(0)\|_{H^{-2}(\mathbb{R}^3)}^2 &\leq c \left( \int_0^T \|\tilde{v}_n(t)\|_{H^{-2}(\mathbb{R}^3)}^2 dt + \int_0^T \|f_n(t)\|_{H^{-2}(\mathbb{R}^3)}^2 dt \right) \\ &\leq c \int_0^T \|\varphi v_n(t)\|_{H^{-1}(\mathbb{R}^3)}^2 dt, \end{aligned}$$

giving (3.15). Now, using (3.13), one has

$$\begin{aligned} \|v_n(0)\|_{H^{-2}}^2 &\leq 2 \left( \|\varphi(x/2)v_n(0)\|_{H^{-2}}^2 + \|(1 - \varphi(x/2))v_n(0)\|_{H^{-2}}^2 \right) \\ &\leq c \int_0^T \|\varphi v_n(t)\|_{H^{-1}}^2 dt + 2\|(1 - \varphi(x/2))v_n(0)\|_{H^{-2}}^2 \rightarrow 0, \end{aligned}$$

that is,

$$(3.16) \quad v_n(0) \rightarrow 0 \text{ strongly in } H^{-2}(\mathbb{R}^3).$$

Let  $w_n = (1 - \Delta)^{-1}v_n$ . Then  $w_n \in C([0, T]; H^1(\mathbb{R}^3))$  is a solution of the equation (3.3). By the convergences (3.14) and (3.16), we can ensure that

$$w_n(0) \rightharpoonup 0 \text{ in } H^1(\mathbb{R}^3) \text{ weakly}$$

and

$$(3.17) \quad w_n \rightarrow 0 \text{ in } C([0, T]; L^2(\mathbb{R}^3)) \text{ strongly.}$$

Now, split  $\varphi w_n$  as

$$\varphi w_n = (1 - \Delta)^{-1}(\varphi v_n) - (1 - \Delta)^{-1}[\varphi, (1 - \Delta)]w_n.$$

Observe that the operator  $[\varphi, (1 - \Delta)]$  maps  $L^2(\mathbb{R}^3)$  continuously into  $H^{-1}(\mathbb{R}^3)$ . So, due to the convergence (3.17), we get that

$$(3.18) \quad (1 - \Delta)^{-1}[\varphi, (1 - \Delta)]w_n \rightarrow 0 \text{ in } C([0, T]; H^1(\mathbb{R}^3)).$$

On the other hand, by (3.13),

$$(3.19) \quad (1 - \Delta)^{-1}(\varphi v_n) \rightarrow 0 \text{ in } L^2((0, T); H^1(\mathbb{R}^3)).$$

Therefore, by the convergences (3.18) and (3.19) above, it follows that

$$\varphi w_n \rightarrow 0 \text{ in } L^2((0, T); H^1(\mathbb{R}^3)).$$

Since  $w_n$  satisfies (3.3), using Lemma 3.2, more precisely, the observability inequality (3.5), we conclude that

$$w_n(0) \rightarrow 0 \text{ in } H^1(\mathbb{R}^3) \text{ strongly,}$$

and so

$$v_n(0) \rightarrow 0 \text{ in } H^{-1}(\mathbb{R}^3) \text{ strongly,}$$

which is a contradiction to the fact that  $\|v_n(0)\|_{H^{-1}}^2 = 1$ , for all  $n$ . This finishes the proof.  $\square$

**Third step:** *Proof of the observability inequality (3.2).*

If (3.2) is false, then there exists a sequence  $\{v_n\}$  of solutions to (3.3) in  $C([0, T]; H^{-1}(\mathbb{R}^3))$  such that

$$(3.20) \quad 1 = \|v_n(0)\|_{H^{-1}}^2 \geq n \int_0^T \|\varphi v_n(t)\|_{H^{-1}}^2 dt, \quad \forall n \geq 0.$$

Extracting a subsequence, still denoted by the same indexes, we have that

$$v_n \rightharpoonup v \text{ in } L^\infty((0, T); H^{-1}(\mathbb{R}^3)) \text{ weak}^*$$

and

$$v_n(0) \rightharpoonup v(0) \text{ in } H^{-1}(\mathbb{R}^3) \text{ weak},$$

for some solution  $v \in C([0, T]; H^{-1}(\mathbb{R}^3))$  of the system (3.3). Note that

$$\varphi v_n \rightharpoonup \varphi v \text{ in } L^\infty(0, T; H^{-1}(\mathbb{R}^3)) \text{ weak}^*$$

and this, combined with (3.20) ( $\varphi v_n \rightarrow 0$  in  $L^2((0, T); H^{-1}(\mathbb{R}^3))$ ), yields  $\varphi v \equiv 0$  and, hence,  $v \equiv 0$  for  $|x| > R + 1$ ,  $t \in (0, T)$ . So, by the unique continuation property as in Step 2, we deduce that  $v \equiv 0$  on  $\mathbb{R}^3 \times (0, T)$ . On the other hand, the sequence  $(1 - \varphi(x/2))v_n(0)$  is bounded in  $H^{-1}(\mathbb{R}^3)$  and has compact support contained in  $B_{2R+2}(0)$ . Therefore, extracting a subsequence, we may assume that it converges strongly in  $H^{-2}(\mathbb{R}^3)$ . Moreover, its limit is necessarily 0 since

$$(1 - \varphi(x/2))v_n(0) \rightarrow 0 \text{ in } H^{-2}(\mathbb{R}^3).$$

Using inequality (3.12), we conclude that  $\|v_n(0)\|_{H^{-1}} \rightarrow 0$ , which contradicts (3.20). This proves the desired observability inequality (3.3) and finishes the proof of Proposition 3.1.  $\square$

We are now in a position to prove Theorem 1.2.

*Proof of Theorem 1.2.* We prove Theorem 1.2 using Hilbert's uniqueness method. First, note that since the Schrödinger equation (3.1) is backward well-posed, we may assume that  $u(T) = 0$  without loss of generality.

Now, consider the two systems

$$(3.21) \quad \begin{cases} i\partial_t u + \Delta u = \varphi(x)h(x, t) \text{ on } [0, T] \times \mathbb{R}^3, \\ u(T) = 0, \end{cases}$$

with  $\varphi(x)$  satisfying (1.6), and

$$\begin{cases} i\partial_t v + \Delta v = 0 \text{ on } [0, T] \times \mathbb{R}^3, \\ v(0) = v_0 \in H^{-1}(\mathbb{R}^3). \end{cases}$$

Multiplying the equation of the first system by  $\bar{v}$  and integrating by parts, we obtain

$$i \int_{\mathbb{R}^3} [\bar{v}(T)u(T) - \bar{v}_0 u(0)] dx = \int_0^T \int_{\mathbb{R}^3} \varphi(x)h(x, t)\overline{v(x, t)} dx dt.$$

Hence, taking  $L^2(\mathbb{R}^3)$  as pivot space, one has

$$(3.22) \quad \langle v_0, -iu_0 \rangle = \int_0^T \langle \varphi(x)v, h(t) \rangle dt,$$

where  $\langle \cdot, \cdot \rangle$  denotes the duality pairing between  $H^{-1}(\mathbb{R}^3)$  and  $H^1(\mathbb{R}^3)$ . Now, consider the isomorphism between  $H^1$  and  $H^{-1}$  defined as follows:

$$\Lambda : H^1(\mathbb{R}^3) \rightarrow H^{-1}(\mathbb{R}^3)$$

giving by  $\Lambda v = (v, \cdot)_{H^1}$ . For any  $v_0 \in H^{-1}(\mathbb{R}^3)$ , let  $h(t) = \Lambda^{-1}(\varphi v(t))$  ( $h \in C([0, T]; H^1(\mathbb{R}^3))$ ) and let  $u$  be the corresponding solution of system (3.21). Finally, set  $\Gamma(v_0) = -iu(\cdot, 0)$ . Then, we have

$$\langle v_0, \Gamma(v_0) \rangle = \int_0^T \|\varphi v(t)\|_{H^{-1}(\mathbb{R}^3)}^2 dt \geq c \|v_0\|_{H^{-1}(\mathbb{R}^3)}^2,$$

in view of the observability inequality (3.2) and identity (3.22). Since the operator  $\Gamma$  is continuous and coercive, it follows from the Lax-Milgram theorem that  $\Gamma$  defines an isomorphism, and this concludes the proof of Theorem 1.2.  $\square$

**3.2. Nonlinear system: Proof of Theorem 1.1.** The proof is based on a perturbation argument due to E. Zuazua [35]. To use it, consider the following two Schrödinger systems with initial data in  $H^{-1}$  and null initial data, namely

$$\begin{cases} i\partial_t\Phi + \Delta\Phi = 0 \text{ on } [0, T] \times \mathbb{R}^3, \\ \Phi(0) = \Phi_0 \in H^{-1}(\mathbb{R}^3), \end{cases}$$

and

$$(3.23) \quad \begin{cases} i\partial_t u + \Delta u - |u|^4 u = A\Phi \text{ on } [0, T] \times \mathbb{R}^3, \\ u(T) = 0, \end{cases}$$

respectively, where  $A$  is defined by

$$A\Phi := \Lambda^{-1}(\varphi(x)\Phi),$$

with  $\Lambda$  as in Theorem 1.2.

Now, define the operator

$$\begin{aligned} \mathcal{L} : H^{-1}(\mathbb{R}^3) &\rightarrow H^1(\mathbb{R}^3) \\ \Phi_0 &\mapsto \mathcal{L}\Phi_0 = u_0 = u(0). \end{aligned}$$

The goal is then to show that  $\mathcal{L}$  is onto in a small neighborhood of the origin of  $H^1(\mathbb{R}^3)$ . To this end, split  $u$  as  $u = v + \Psi$ , where  $\Psi$  is a solution of

$$\begin{cases} i\partial_t\Psi + \Delta\Psi = A\Phi \text{ on } [0, T] \times \mathbb{R}^3, \\ \Psi(T) = 0, \end{cases}$$

and  $v$  is a solution of

$$(3.24) \quad \begin{cases} i\partial_t v + \Delta v = |u|^4 u \text{ on } [0, T] \times \mathbb{R}^3, \\ v(T) = 0. \end{cases}$$

Clearly  $u, v$  and  $\Psi$  belong to  $C([0, T], H^1(\mathbb{R}^3)) \cap L^{10}([0, T]; L^{10}(\mathbb{R}^3))$  and  $u(0) = v(0) + \Psi(0)$ . We may write

$$\mathcal{L}\Phi_0 = \mathcal{J}\Phi_0 + \Gamma\Phi_0,$$

where  $\mathcal{J}\Phi_0 = v_0$ . Observe that  $\mathcal{L}\Phi_0 = u_0$ , or equivalently,  $\Phi_0 = -\Gamma^{-1}\mathcal{J}\Phi_0 + \Gamma^{-1}u_0$ .

Now, define the operator

$$\begin{aligned} \mathcal{B} : H^{-1}(\mathbb{R}^3) &\rightarrow H^{-1}(\mathbb{R}^3) \\ \Phi_0 &\rightarrow \mathcal{B}\Phi_0 = \Gamma^{-1}\mathcal{J}\Phi_0 + \Gamma^{-1}u_0, \end{aligned}$$

where we are taking into account that  $\Gamma$  is the linear control isomorphism between  $H^{-1}$  and  $H^1$ , due to Theorem 1.2.

Now, the goal is to prove that  $\mathcal{B}$  has a fixed point near the origin of  $H^{-1}(\mathbb{R}^3)$ . More precisely, let us prove that if  $\|u_0\|_{H^1}$  is small enough, then  $\mathcal{B}$  is a contraction on a small ball  $B_R$  of  $H^{-1}(\mathbb{R}^3)$ . We may assume  $T < 1$  and we will denote by  $C > 0$  any constant that may have its numerical value changed line by line. Since  $\Gamma$  is an isomorphism, we have

$$(3.25) \quad \begin{aligned} \|\mathcal{B}\Phi_0\|_{H^{-1}} &\leq \|\Gamma^{-1}\mathcal{J}\Phi_0\|_{H^{-1}} + \|\Gamma^{-1}u_0\|_{H^{-1}} \\ &\leq C(\|\mathcal{J}\Phi_0\|_{H^1} + \|u_0\|_{H^1}) \\ &\leq C(\|v(0)\|_{H^1} + \|u_0\|_{H^1}). \end{aligned}$$

**Claim 1:** There exists  $C > 0$  such that

$$(3.26) \quad \|v(0)\|_{H^1} \leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^5.$$

Indeed, note that due to the classical energy estimate for system (3.24), Strichartz estimates (see Lemma 2.2) and a Sobolev embedding (see Lemma 2.3), we have

$$\begin{aligned} \|v(0)\|_{L^2} &\leq \|v(T)\|_{L^2} + \left\| \int_0^t e^{i(t-\tau)\Delta} |u|^4 u \, d\tau \right\|_{L_x^2} \\ &\leq C \|u^5\|_{L_t^1 L_x^2} \\ &\leq C \|u\|_{L_t^{10} L_x^{10}}^5 \\ &\leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^5 \end{aligned}$$

and

$$\begin{aligned} \|\nabla v(0)\|_{L^2} &\leq \|\nabla v(T)\|_{L^2} + \left\| \int_0^t e^{i(t-\tau)\Delta} \nabla |u|^4 u \, d\tau \right\|_{L_x^2} \\ &\leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}} \|u\|_{L_t^{10} L_x^{10}}^4 \\ &\leq \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^5. \end{aligned}$$

Thus,

$$\|v(0)\|_{L^2}^2 \leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^{10}$$

and

$$\|\nabla v(0)\|_{L^2}^2 \leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^{10}.$$

Summing up, we have (3.26), thus showing Claim 1.

**Claim 2:** There exists  $C > 0$  such that

$$(3.27) \quad \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}} \leq C \|\Phi_0\|_{H^{-1}}.$$

In fact, applying Lemma 2.2 to system (3.23), one gets

$$\begin{aligned} \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}} &\leq \|\nabla u(T)\|_{L^2} + C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}} \|u\|_{L_t^{10} L_x^{10}}^4 + C \|\nabla A\Phi\|_{L_t^1 L_x^2} \\ &\leq C \left( \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^5 + \|A\Phi\|_{L_t^2 H_x^1} \right). \end{aligned}$$

Note that, using the fact that  $\Lambda$  is an isomorphism, we get

$$\|A\Phi\|_{H^1} = \|\Lambda^{-1}(\varphi\Phi)\|_{H^1} \leq C \|\varphi\Phi\|_{H^{-1}}$$

or, equivalently,

$$\|A\Phi\|_{L_t^2 H_x^1} \leq \left( \int_0^T \|\varphi\Phi\|_{H^{-1}}^2 \, dt \right)^{\frac{1}{2}}.$$

Then, the duality (3.22) yields

$$\begin{aligned} \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}} &\leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^5 + C \left( \int_0^T \|\varphi\Phi\|_{H^{-1}}^2 \, dt \right)^{\frac{1}{2}} \\ &\leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^5 + C \left( \langle \Gamma\Phi_0, \Phi_0 \rangle \right)^{\frac{1}{2}} \\ &\leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^5 + C \left( \|\Gamma\Phi_0\|_{H^1} \|\Phi_0\|_{H^{-1}} \right)^{\frac{1}{2}} \\ &\leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^5 + C \left( \|\Phi_0\|_{H^{-1}}^2 \right)^{\frac{1}{2}} \\ &\leq C \|\nabla u\|_{L_t^{10} L_x^{\frac{30}{13}}}^5 + C \|\Phi_0\|_{H^{-1}}. \end{aligned}$$

Using a bootstrap argument, taking  $\|\Phi_0\|_{H^{-1}} \leq R$  with  $R$  small enough, we get (3.27), showing Claim 2.

Note that putting together inequalities (3.26) and (3.27) into (3.25), we conclude

$$\|\mathcal{B}\Phi_0\|_{H^{-1}} \leq C\left(\|v(0)\|_{H^1} + \|u_0\|_{H^1}\right) \leq C\left(\|\Phi_0\|_{H^{-1}}^5 + \|u_0\|_{H^1}\right).$$

Then, choosing  $R$  small enough so that  $CR^5 \leq R/2$  and

$$\|u_0\|_{H^1} \leq \frac{R}{2C},$$

we get

$$\|\mathcal{B}\Phi_0\|_{H^{-1}} \leq R$$

and, therefore,  $\mathcal{B}$  reproduces the closed ball of radius  $R$  of  $H^{-1}(\mathbb{R})$ .

Finally, we prove that  $\mathcal{B}$  is a contraction map. To do this, let us study the systems

$$\begin{cases} i\partial_t(u_1 - u_2) + \Delta(u_1 - u_2) - |u_1|^4u_1 + |u_2|^4u_2 = A(\Phi^1 - \Phi^2), \\ (u_1 - u_2)(T) = 0, \end{cases}$$

and

$$(3.28) \quad \begin{cases} i\partial_t(v_1 - v_2) + \Delta(v_1 - v_2) = |u_1|^4u_1 - |u_2|^4u_2, \\ (v_1 - v_2)(T) = 0. \end{cases}$$

As before, we also have

$$(3.29) \quad \|\mathcal{B}\Phi_0^1 - \mathcal{B}\Phi_0^2\|_{H^{-1}} \leq C\|v_1(0) - v_2(0)\|_{H^1}.$$

We now estimate  $v_1(0) - v_2(0)$  in the  $H^1$ -norm. First, applying Lemma 2.3 yields that

$$\begin{aligned} \|v_1(0) - v_2(0)\|_{L^2} &\leq \left\| \int_0^t e^{i(t-\tau)\Delta} (|u_1|^4u_1 - |u_2|^4u_2) d\tau \right\|_{L_x^2} \\ &\leq \| |u_1|^4u_1 - |u_2|^4u_2 \|_{L_t^1 L_x^2} \\ &\leq C \|u_1 - u_2\|_{L_t^5 L_x^{10}} \left( \|u_1\|_{L_t^5 L_x^{10}}^4 + \|u_2\|_{L_t^5 L_x^{10}}^4 \right) \\ &\leq C \|u_1 - u_2\|_{L_t^{10} L_x^{10}} \left( \|u_1\|_{L_t^{10} L_x^{10}}^4 + \|u_2\|_{L_t^{10} L_x^{10}}^4 \right) \\ &\leq C \|\nabla u_1 - \nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}} \left( \|\nabla u_1\|_{L_t^{10} L_x^{\frac{30}{13}}}^4 + \|\nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}}^4 \right) \\ &\leq CR^4 \|\nabla u_1 - \nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}} \end{aligned}$$

since we have

$$\|\nabla u_j\|_{L_t^{10} L_x^{\frac{30}{13}}} \leq C\|\Phi_0\|_{H^{-1}} \leq CR$$

and then applying Strichartz estimates (see Lemma 2.2)

$$\begin{aligned}
\|\nabla v_1(0) - \nabla v_2(0)\|_{L^2} &\leq \left\| \int_0^t \nabla e^{i(t-\tau)\Delta} (|u_1|^4 u_1 - |u_2|^4 u_2) d\tau \right\|_{L_t^2 L_x^2} \\
&\leq \|\nabla(|u_1|^4 u_1 - |u_2|^4 u_2)\|_{L_t^2 L_x^{\frac{6}{5}}} \\
&\leq C \|\nabla(u_1 - u_2)\|_{L_t^{10} L_x^{\frac{30}{13}}} \|u_1\|_{L_t^{10} L_x^{10}}^4 \\
&\leq C \|\nabla(u_1 - u_2)\|_{L_t^{10} L_x^{\frac{30}{13}}} \|\nabla u_1\|_{L_t^{10} L_x^{\frac{30}{13}}}^4 \\
&\quad + C \|\nabla u_1 - \nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}} \left( \|\nabla u_1\|_{L_t^{10} L_x^{\frac{30}{13}}}^3 \|\nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}} \right. \\
&\quad \left. + \|\nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}}^3 \|\nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}} \right) \\
&\leq CR^4 \|\nabla(u_1 - u_2)\|_{L_t^{10} L_x^{\frac{30}{13}}} + CR^4 \|\nabla(u_1 - u_2)\|_{L_t^{10} L_x^{\frac{30}{13}}} \\
&\leq CR^4 \|\nabla(u_1 - u_2)\|_{L_t^{10} L_x^{\frac{30}{13}}}.
\end{aligned}$$

Thus,

$$\|v_1(0) - v_2(0)\|_{L^2}^2 \leq C^2 R^8 \|\nabla u_1 - \nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}}^2$$

and

$$\|\nabla v_1(0) - \nabla v_2(0)\|_{L^2}^2 \leq C^2 R^8 \|\nabla u_1 - \nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}}^2.$$

These bounds together give us the  $H^1$ -estimate

$$\|v_1(0) - v_2(0)\|_{H^1} \leq CR^4 \|\nabla u_1 - \nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}}.$$

Now, let us bound the right-hand side of this inequality. To this end, first notice that

$$\begin{aligned}
\|\nabla(u_1 - u_2)\|_{L_t^{10} L_x^{\frac{30}{13}}} &\leq \|\nabla(|u_1|^4 u_1 - |u_2|^4 u_2)\|_{L_t^2 L_x^{\frac{6}{5}}} + \|\nabla A(\Phi^1 - \Phi^2)\|_{L_t^1 L_x^2} \\
&\leq CR^4 \|\nabla u_1 - \nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}} + C \|A(\Phi^1 - \Phi^2)\|_{L_t^2 H_x^1} \\
&\leq CR^4 \|\nabla u_1 - \nabla u_2\|_{L_t^{10} L_x^{\frac{30}{13}}} + C \|\Phi_0^1 - \Phi_0^2\|_{H^{-1}}.
\end{aligned}$$

So, choosing  $R > 0$  small enough, we get

$$\|\nabla(u_1 - u_2)\|_{L_t^{10} L_x^{\frac{30}{13}}} \leq C \|\Phi_0^1 - \Phi_0^2\|_{H^{-1}}.$$

Therefore,

$$\begin{aligned}
(3.30) \quad \|v_1(0) - v_2(0)\|_{H^1} &= \left( \|v_1(0) - v_2(0)\|_{L^2}^2 + \|\nabla v_1(0) - \nabla v_2(0)\|_{L^2}^2 \right)^{\frac{1}{2}} \\
&\leq CR^4 \|\Phi_0^1 - \Phi_0^2\|_{H^{-1}}.
\end{aligned}$$

Finally, we get by inequalities (3.29) and (3.30) that

$$\|\mathcal{B}\Phi_0^1 - \mathcal{B}\Phi_0^2\|_{H^{-1}} \leq C \|v_1(0) - v_2(0)\|_{H^1} \leq CR^4 \|\Phi_0^1 - \Phi_0^2\|_{H^{-1}},$$

concluding that  $\mathcal{B}$  is a contraction on a small ball  $B_R$  of  $H^{-1}$ . This completes the proof of Theorem 1.1.  $\square$

## 4. COMMENTS AND OPEN ISSUES

In this work, we have studied the local null controllability of the critical C-NLS in dimension 3. Considering the defocusing critical nonlinear Schrödinger equation (1.5), we showed this system to be controllable in  $H^1(\mathbb{R}^3)$ -level, that is, it satisfies  $u(T) = 0$  for  $u_0 \in H^1(\mathbb{R}^3)$  satisfying  $\|u_0\|_{H^1} \leq \delta$ , and  $h(x, t) \in C([0, T]; H^1(\mathbb{R}^3))$ . Concerning our main results, Theorems 1.1 and 1.2, the following remarks are worth mentioning.

- The main tools to achieve these results were:
  - *Strichartz estimates* for the Schrödinger operator [7, 11], which give the well-posedness theory for system (1.5);
  - *A unique continuation property* for the linear system associated with (1.5), which follows by the Carleman estimate showed in [21] (see also [17]);
  - *A perturbation argument*, as presented in [35], which allows one to extend the result for the nonlinear system (1.5).
- For the  $3d$ -case, Laurent [17] showed large time global internal controllability for the nonlinear Schrödinger equation on some compact manifolds of dimension 3. The results therein were obtained for the nonlinearity  $|u|^2u$  instead of  $|u|^4u$ . In this sense, our result completes the local control result given by the author for the critical case (1.5) in  $\mathbb{R}^3$ .
- Since we are working on the whole space  $\mathbb{R}^3$  and we have control taking the form  $\varphi(x)h(x, t)$  in the system (1.5) (remember the definition of  $\varphi$  in (1.6)), the geometric control condition (see, for instance, [1, 25]) is easily satisfied. This corroborates with the recent work of Taüfer [32], that is, our work expresses that we do not need a strong GCC to achieve the control properties for the Schrödinger equation in  $\mathbb{R}^3$ .

Thus, our work gives the first step to understanding control problems for the C-NLS in dimension 3, and, therefore, some important open issues appear. Let us present them below.

- (A) **Stabilization problem:** Can one find a feedback control law  $f = Ku$  so that the resulting closed-loop system

$$i\partial_t u + \Delta u - |u|^4 u = Ku, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^3,$$

is asymptotically stable at an equilibrium point as  $t \rightarrow +\infty$ ?

- (B) **Global control problem:** If the answer to the previous question is positive, another natural issue is to obtain a global control result for the system (1.5), i.e., a control result for large data. This would be a consequence of the global stabilization result together with the local control result shown in Theorem 1.1.
- (C) **Geometric control condition (GCC) or thick-set conditions:** Our work considered the control acting on  $B(0, R)^c$ . From the perspective of the observation region of the linear Schrödinger equation, an interesting open problem appears: Are there more general sets satisfying the (GCC) such that the observability inequality is verified?

**Acknowledgment.** This work is part of Carvalho's Ph.D. thesis at the Department of Mathematics of the Universidade Federal de Pernambuco. It was mostly done while the third author was visiting Université de Lorraine. The third author thanks the host institution for its warm hospitality.

## REFERENCES

- [1] C. Bardos, G. Lebeau, and J. Rauch, Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary, *SIAM J. Control Optim.* 305 (1992) 1024–1065.
- [2] L. Baudouin and J.-P. Puel, *Uniqueness and stability in an inverse problem for the Schrödinger equation*, *Inverse Problems* 18 (2001) 1537–1554.
- [3] R. A. Capistrano-Filho and A. Pampu, *The fractional Schrödinger equation on compact manifolds: Global controllability results*, *Mathematische Zeitschrift*, 301 (2022) 3817–3848.
- [4] L. Cardoulis and P. Gaitan, *Simultaneous identification of diffusion coefficient and the potential for the Schrödinger operator with only one observation*, *Inverse Problems* 26 (2010).

- [5] L. Cardoulis, M. Cristofol, and P. Gaitan, *Inverse problem for the Schrödinger operator in an unbounded strip*, J. Inverse Ill-Posed Probl. 16 (2008) 127–146.
- [6] T. Cazenave and F. B. Weissler, *The Cauchy problem for the critical nonlinear Schrödinger equation in  $H^s$* , Nonlinear Analysis: Theory, Methods & Applications, 14(10), (1990) 807–836.
- [7] T. Cazenave, *Semilinear Schrödinger Equations*, Amer. Math. Soc., Providence, RI (2003).
- [8] B. Dehman, P. Gérard, and G. Lebeau, *Stabilization and control for the nonlinear Schrödinger equation on a compact surface*, Mathematische Zeitschrift, 254 (2006) 729–749.
- [9] R. Illner, H. Lange, and H. Teismann, *A note on the exact internal control of nonlinear Schrödinger*, in: CRM Proc. Lecture Notes, 33 (2003) 127–137.
- [10] R. Illner, H. Lange, and H. Teismann, *Limitations on the control of Schrödinger equations*, ESAIM Control Optim. Calc. Var. 12 (2006) 615–635.
- [11] M. Keel and T. Tao, *Endpoint Strichartz estimates*, Am. J. Math. 120 (1998) 955–980.
- [12] C. E. Kenig and F. Merle, *Global well-posedness, scattering, and blow-up for the energy-critical, focusing, nonlinear Schrödinger equation in the radial case*, Invent. Math. 166 (2006) 645–675.
- [13] K.C. E. Kenig and F. Merle, *Nondispersive radial solutions to energy supercritical non-linear wave equations, with applications*, American Journal of Mathematics, 130(3), (2008) 633–665.
- [14] V. Komornik and P. Loreti, *Fourier Series in Control Theory Springer-Verlag*, (2005).
- [15] H. Lange and H. Teismann, *Controllability of the nonlinear Schrödinger equation in the vicinity of the ground state*, Math. Meth. Appl. Sci. 30, (2007) 1483–1505.
- [16] C. Laurent, *Global controllability and stabilization for the nonlinear Schrödinger equation on an interval*, ESAIM Control Optimisation and Calculus of Variations, 2 (2010) 356–379.
- [17] C. Laurent, *Global controllability and stabilization for the nonlinear Schrödinger equation on some compact manifolds of dimension 3*, SIAM Journal on Mathematical Analysis, 42, (2010) 785–832.
- [18] I. Lasiecka, R. Triggiani, and X. Zhang, *Carleman estimates at the  $H^1(\Omega)$ - and  $L^2(\Omega)$ -level for nonconservative Schrödinger equations with unobserved Neumann B.C.*, Arch. Inequal. Appl. 2, (2004) 215–338.
- [19] J.-L. Lions, *Exact controllability, stabilization and perturbations for distributed systems*, SIAM Rev. 30, (1988) 1–68.
- [20] F. Merle and P. Raphaël, *Sharp upper bound on the blow-up rate for the critical nonlinear Schrödinger equation*, Geometric and Functional Analysis, 13(3), (2003) 591–642.
- [21] A. Mercado, A. Osses, and L. Rosier, *Inverse problems for the Schrödinger equation via Carleman inequalities with degenerate weights*, Inverse Problems 24, (2008) 1–18.
- [22] L. Miller, *How violent are fast controls for Schrödinger and plate vibrations?*, Arch. Rational Mech. Anal. 172, (2004) 429–456.
- [23] K.-D. Phung, *Observability and controllability for Schrödinger equations*, SIAM J. Control Optim. 40, (2001) 211–230.
- [24] K. Ramdani, T. Takahashi, G. Tenenbaum, and M. Tucsnak, *A spectral approach for the exact observability of infinite-dimensional systems with skew-adjoint generator*, J. Funct. Anal. 226, (2005) 193–229.
- [25] J. Rauch and M. Taylor, *Exponential decay of solutions to hyperbolic equations in bounded domains*, Indiana Univ. Math. J. 24(1), (1975) 79–86.
- [26] L. Rosier and B.-Y. Zhang, *Local exact controllability and stabilizability of the nonlinear Schrödinger equation on a bounded domain*, SIAM J. Control Optim. 48, (2009) 972–992.
- [27] L. Rosier and B.-Y. Zhang, *Exact boundary controllability of the nonlinear Schrödinger equation*, J. Differential Equations 246, (2009) 4129–4153.
- [28] L. Rosier and B.-Y. Zhang, *Control and Stabilization of the Nonlinear Schrödinger Equation on Rectangles*, Mathematical Models, and Methods in Applied Sciences, 12, (2010) 2293–2347.
- [29] J. Simon, *Compact sets in the space  $L^p(0, T; B)$* , Ann. Mat. Pura Appl. (4) CXLVI (1987) 65–97.
- [30] C. Sulem and P. L. Sulem, *The Nonlinear Schrödinger Equation: Self-Focusing and Wave Collapse*, Applied Mathematical Sciences, Vol. 139, Springer (1999).
- [31] T. Tao, *Nonlinear Dispersive Equations: Local and Global Analysis*. CBMS Regional Conference Series in Mathematics, American Mathematical Society (2009).
- [32] M. Täufer, *Controllability of the Schrödinger equation on unbounded domains without geometric control condition*, ESAIM Control Optim. Calc. Var., 29 (59), (2023) 1-11.
- [33] G. Yuan and M. Yamamoto, *Carleman estimates for the Schrödinger equation and applications to an inverse problem and an observability inequality*, Chinese Annals of Mathematics 31(4), (2010) 555–578.
- [34] M. Visan, *The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions*, Duke Mathematical Journal, 138(2), (2007) 281–374.
- [35] E. Zuazua, *Exact controllability for the semilinear wave equation*, J. Math. Pures Appl. 69, 33–55 (1990)
- [36] E. Zuazua, *Remarks on the controllability of the Schrödinger equation*, Quantum Control: Mathematical and Numerical Challenges, 193–211, CRM Proc. Lecture Notes, Vol. 33 (Amer. Math. Soc., 2003).

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE PERNAMBUCO, S/N CIDADE UNIVERSITÁRIA,  
50740-545, RECIFE (PE), BRAZIL

*Email address:* `pablo.braz@ufpe.br`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE PERNAMBUCO, S/N CIDADE UNIVERSITÁRIA,  
50740-545, RECIFE (PE), BRAZIL

*Email address:* `roberto.capistranofilho@ufpe.br`

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE PERNAMBUCO, S/N CIDADE UNIVERSITÁRIA,  
50740-545, RECIFE (PE), BRAZIL

*Email address:* `jackellyny.dassy@ufpe.br`

INSTITUT ÉLIE CARTAN DE LORRAINE, UMR CNRS 7502, ÉQUIPE SPHINX, INRIA, UNIVERSITÉ DE LORRAINE,  
F-54506 VANDOEUVRE-LES-NANCY CEDEX, FRANCE.

*Email address:* `dds@math.cnrs.fr`