A CLASSIFICATION OF CONSTANT GAUSSIAN CURVATURE SURFACES IN THE THREE-DIMENSIONAL HYPERBOLIC SPACE

JUN-ICHI INOGUCHI AND SHIMPEI KOBAYASHI

ABSTRACT. Weakly complete constant Gaussian curvature $-1 < K < 0$ surfaces will be classified in terms of holomorphic quadratic differentials. For this purpose, we will first establish a loop group method for constant Gaussian curvature surfaces in \mathbb{H}^3 with $K > -1$ but $K \neq 0$ by using harmonicities of Lagrangian and Legendrian Gauss maps. Then we will show that a spectral parameter deformation of the Lagrangian harmonic Gauss map gives a harmonic map into \mathbb{H}^2 for $-1 < K < 0$ or \mathbb{S}^2 for $K > 0$, respectively.

INTRODUCTION

As is well known, Hilbert proved the nonexistence of isometric immersions of the hyperbolic plane \mathbb{H}^2 into the Euclidean 3-space \mathbb{E}^3 . Analogous results hold for surfaces in the 3-sphere \mathbb{S}^3 and the hyperbolic 3-space \mathbb{H}^3 as follows, see for example Spivak [\[14\]](#page-22-0):

Fact.

- (1) There is no complete surface in \mathbb{S}^3 with constant Gaussian curvature with $K < 1$ but $K \neq 0$, and the only complete immersions of constant Gaussian curvature $K \geq 1$ in S ³ are totally umbilic round spheres. In particular, the only complete surface of $K = 1$ are great spheres.
- (2) There is no complete surface in \mathbb{H}^3 with constant Gaussian curvature $K < -1$, and the only complete immersions of constant Gaussian curvature $K \geq 0$ in \mathbb{H}^3 are totally umbilic spheres or equidistant surfaces from geodesics (the only $K = 0$).

On the one hand, it is known that there are complete surfaces of $K = 0$ in \mathbb{S}^3 [\[5\]](#page-22-1) or $K = -1$ in \mathbb{H}^3 [\[11\]](#page-22-2), respectively. When Spivak wrote the book [\[14\]](#page-22-0), the only known examples of complete surfaces of constant Gaussian curvature in \mathbb{H}^3 satisfying $-1 < K < 0$ are equidistant surfaces and surfaces of revolution. Then Rosenberg and Spruck [\[13\]](#page-22-3) proved that given a smooth curve Γ in the ideal boundary of \mathbb{H}^3 and a constant $k \in (-1,0)$, there exists a embedded complete surface M with Gaussian curvature $K = k$ whose boundary at infinity is Γ. Other outstanding results about complete surfaces solving a Plateau problem at infinity were also given by Labourie [\[8\]](#page-22-4).

Date: April 15, 2024.

²⁰²⁰ Mathematics Subject Classification. Primary 53C42, 58D10, Secondary 53C43, 53A10.

Key words and phrases. Constant Gaussian curvature; hyperbolic 3-space; loop groups.

The first named author is partially supported by Kakenhi 24540063, 15K04834, 19K03461, 23K03081.

The second named author is partially supported by Kakenhi 26400059, 18K03265, 22K03265.

2 J. INOGUCHI AND S.-P. KOBAYASHI

In our previous paper with Brander [\[1\]](#page-22-5), we have given a loop group method for constructing constant Gaussian curvature surfaces in \mathbb{S}^3 with $K < 1$ but $K \neq 0$. The key clue of [\[1\]](#page-22-5) is that every constant Gaussian curvature surfaces is uniquely determined by a Lorentzian harmonic map into the unit 2-sphere \mathbb{S}^2 . The Lorentzian harmonic map is a normal Gauss map of the surface. On the other hand in [\[3\]](#page-22-6), we have established a unified loop group method for constant mean curvature surfaces in \mathbb{H}^3 . The loop group method of [\[3\]](#page-22-6) is based on the characterization of constant mean curvature condition in terms of Gauss map which takes values in the unit tangent sphere bundle $U\mathbb{H}^3$ of the hyperbolic 3-space \mathbb{H}^3 .

The first purpose of the present paper is to establish a loop group method for constant Gaussian curvature surfaces in \mathbb{H}^3 with $K > -1$ but $K \neq 0$. Contrary to the S³-case, we do not have \mathbb{H}^2 -valued harmonic maps naturally associated to constant Gaussian curvature surfaces in \mathbb{H}^3 . Thus it is not easy to modify [\[1\]](#page-22-5) to the \mathbb{H}^3 -case. To overcome this difficulty, we will first establish a Ruh-Vilms type characterization for Legendrian and Lagrangian Gauss maps of constant Gaussian curvature surfaces in \mathbb{H}^3 :

Theorem A. Let $f : M \to \mathbb{H}^3$ be a surface of Gaussian curvature $K > -1$ with unit normal n. Let $F = (f, n) : M \to \text{U} \mathbb{H}^3$ and $L = f \wedge n : M \to \text{Geo}(\mathbb{H}^3)$ be the Legendrian and Lagraingian Gauss maps of f, respectively. Then the following statements are mutually equivalent:

- (1) The Gaussian curvature of f is constant.
- (2) The Legendrian Gauss map F is harmonic.
- (3) The Lagrangian Gauss map L is harmonic.
- (4) The Klotz differential is holomorphic.

Note that $\text{Geo}(\mathbb{H}^3)$ is the space of all oriented geodesics of \mathbb{H}^3 . The precise definitions of the Klotz differential, Legendrian and Lagrangian Gauss maps will be given in Section [1,](#page-3-0) and the proof of Theorem [A](#page-1-0) will be given in Section [2.2.](#page-10-0) From Theorem [A](#page-1-0) and the structure equations of a constant Gaussian curvature surface f , there is a family of constant Gaussian curvature surfaces f^{λ} , the so-called *associated family*, parameterized by the spectral parameter $\lambda \in \mathbb{S}^1 \subset \mathbb{C}$ such that $f^{\lambda}|_{\lambda=1} = f$.

The second purpose of the present paper is that we characterize constant Gaussian curvature surface with $K > -1$ but $K \neq 0$ in terms of other simple harmonic maps by using a loop group method. For this purpose, we will consider a deformation of the spectral parameter off the unit circle to the spectral parameter family of Lagrangian Gauss maps. Then it will be turned out that every constant Gaussian curvature surfaces with K > −1 but $K \neq 0$ are recovered from the Lax equation of harmonic maps into \mathbb{H}^2 and S 2 , respectively:

Theorem B. Let $f : M \to \mathbb{H}^3$ be a constant Gaussian curvature surface with $K > -1$ but $K \neq 0$, and let f^{λ} and n^{λ} the corresponding associated family and a family of unit normals of f^{λ} . Moreover, let $L^{\lambda} = f^{\lambda} \wedge n^{\lambda}$ be a family of Lagrangian Gauss harmonic maps. Then the following statements hold:

(1) Assume $-1 < K < 0$ and take a constant $\lambda_0 \in \mathbb{C} \setminus \overline{\mathbb{D}}$ with $\overline{\mathbb{D}} = {\lambda \in \mathbb{C} \mid |\lambda| \leq 1}$ such that

$$
|\lambda_0| = \exp\left(\operatorname{arcosh}\sqrt{-1/K}\right)
$$

holds. Then the map $L^{\lambda}|_{\lambda=\lambda_0}$ is a harmonic local diffeomorphism into \mathbb{H}^2 up to a conjugation.

(2) Assume $K > 0$ and take a constant $\lambda_0 \in \mathbb{C} \setminus \overline{\mathbb{D}}$ such that

$$
|\lambda_0| = \exp\left(\operatorname{arsinh}\sqrt{1/K}\right)
$$

holds. Then the map $L^{\lambda}|_{\lambda=\lambda_0}$ is a harmonic local diffeomorphism into \mathbb{S}^2 up to a conjugation.

Moreover, if the surface f^{λ} is not totally umbilic, the harmonic Lagrangian Gauss map $L^{\lambda}|_{\lambda=\lambda_0}$ is non-conformal.

The proof will be given in Section [2.4.](#page-13-0)

The final and principal purpose of the present paper is a classification of constant Gaussian curvature $-1 < K < 0$ surfaces. By Theorems [A](#page-1-0) and [B,](#page-1-1) we obtain the harmonic local diffeomorphism $L^{\lambda}|_{\lambda=\lambda_0}$ into \mathbb{H}^2 corresponding to f. It is known [\[15\]](#page-22-7) that all complete ∂ -energy harmonic maps from the unit disc D or the complex plane C into \mathbb{H}^2 bijectively correspond to holomorphic quadratic differentials. In our case the holomorphic quadratic differentials are exactly the Klotz differentials in Theorem [A,](#page-1-0) however, the induced metric of the surface f is not related to the ∂ -energy of the harmonic map. We thus introduce an another natural metric for a surface in \mathbb{H}^3 in terms of the unit tangent sphere bundle U \mathbb{H}^3 : The so-called weak metric is defined by

$$
dt^{2} = \langle df, df \rangle + \frac{1}{1 + K} \langle dn, dn \rangle.
$$

Then we will show that equivalence of completeness of the weak metrics and the ∂-energy of a harmonic map, see Lemma [3.1.](#page-17-0) Our main result of the paper is a classification of weakly complete constant Gaussian curvature $-1 < K < 0$ surfaces in \mathbb{H}^3 in terms of holomorphic quadratic differentials.

Theorem C. Let K_K be a set of equivalent classes (by rigid motions on \mathbb{H}^3) of weakly complete constant Gaussian curvature $-1 < K < 0$ surfaces with conformal type the unit disk $\mathbb{D} := \{z \in \mathbb{C} \mid |z| \leq 1\}$ or the complex plane \mathbb{C} . Moreover, let \mathcal{QD} be a set of equivalent classes (by Möbius transformations on $\mathbb D$ or $\mathbb C$) of holomorphic quadratic differentials on $\mathbb D$ or $\mathbb C$, which are not constant on $\mathbb C$. Then there exists a bijective correspondence between \mathcal{K}_K and \mathcal{QD} .

The proof will be given in Section [3.1.](#page-17-1) Finally in Corollary [3.2](#page-19-0) we will show that complete constant Gaussian curvature $-1 < K < 0$ surfaces will be obtained by bounded holomorphic quadratic differentials.

The paper is organized as follows: In Section [1,](#page-3-0) we will recall fundamental results/notion about surfaces in \mathbb{H}^3 . In Section [2,](#page-7-0) we will discuss harmonicities of Lagrangian and Legendrian Gauss maps for surfaces with Gaussian curvature $K > -1$ and establish a loop group

4 J. INOGUCHI AND S.-P. KOBAYASHI

method for constant Gaussian curvature surfaces. In Section [3,](#page-16-0) we will classify weakly complete constant Gaussian curvature $-1 < K < 0$ surfaces in terms of holomorphic quadratic differentials.

1. preliminaries

In this section, we recall fundamental results/notion about surfaces with $K > -1$ in \mathbb{H}^3 . We will introduce conformal coordinates with respect to the second fundamental form of such surfaces. Moreover, the unit tangent sphere bundle $U\mathbb{H}^3$ and the space of all oriented geodesics $Geo(\mathbb{H}^3)$ will be recalled, and the Lagrangian and Legendrian Gauss maps will be introduced.

1.1. Surfaces in the hyperbolic 3-space. Let us denote by $\mathbb{E}^{1,3}$ the *Minkowski* 4-space, that is, $\mathbb{E}^{1,3} = (\mathbb{R}^4(x_0, x_1, x_2, x_3), -dx_0^2 + dx_1^2 + dx_2^2 + dx_3^2)$. Then the *hyperbolic* 3-space \mathbb{H}^3 is defined by an unit central hyperquadric:

$$
\mathbb{H}^3 = \{ (x_0, x_1, x_2, x_3) \in \mathbb{E}^{1,3} \mid -x_0^2 + x_1^2 + x_2^2 + x_3^2 = -1, \quad x_0 > 0 \}.
$$

It is known that \mathbb{H}^3 is a simply connected Riemannian 3-manifold of constant curvature -1 . Denote by $\text{SO}_{1,3}^+$ the identity component of the Lorentz group $\text{O}_{1,3}$. Then \mathbb{H}^3 is represented by $\mathbb{H}^3 = SO_{1,3}^+/SO_3$.

Let $f: M \to \mathbb{H}^3$ be a surface with unit normal n and let us take a local orthonormal frame field $\{e_1, e_2, e_3 = n\}$ of f. Moreover take a dual coframe field $\{\omega^1, \omega^2, \omega^3\}$ of $\{e_1, e_2, e_3 = n\}$. Then this coframe field satisfies

$$
\mathrm{d}\omega^{\alpha}=\omega^{\beta}\wedge\omega_{\beta}^{\alpha},
$$

where $1 \leq \alpha, \beta \leq 3$, see for example [\[16\]](#page-22-8). Note that we have used the Einstein summation convention. Moreover, the structure equation is

(1.1)
$$
d\omega_{\alpha}^{\ \beta} = \omega_{\alpha}^{\ \gamma} \wedge \omega_{\gamma}^{\ \beta} + \Omega_{\alpha}^{\ \beta},
$$

where $1 \leq \alpha, \beta, \gamma \leq 3$ and $\Omega_{\alpha}^{\ \beta}$ is the curvature 2-form of \mathbb{H}^3 defined by

$$
\Omega_{\alpha}^{\ \beta} = \omega^{\alpha} \wedge \omega^{\beta}.
$$

On the surface (M, f) , we have $\omega^i \wedge {\omega_i}^3 = 0$, $(1 \le i \le 2)$. Cartan's lemma implies that ω_i^3 is expressed as $\omega_i^3 = b_{ij} \omega^j$, $b_{ij} = b_{ji}$, $(1 \le i, j \le 2)$. One can check that the first and second fundamental forms $I = \langle df, df \rangle$ and $I = -\langle df, dn \rangle$ are respectively represented as

(1.2)
$$
\mathbf{I} = \omega^1 \otimes \omega^1 + \omega^2 \otimes \omega^2, \quad \mathbf{I} = b_{ij} \omega^i \otimes \omega^j,
$$

where $1 \leq i, j \leq 2$. Then, by [\(1.1\)](#page-3-1) and [\(1.2\)](#page-3-2), the Gauss curvature K and the mean curvature H can be computed as

(1.3)
$$
K = -1 + \det(b_{ij}), \quad H = \frac{1}{2}(b_{11} + b_{22}).
$$

The third fundamental form $\mathbb{I} = \langle dn, dn \rangle$ can be computed as

(1.4)
$$
\mathbb{I} = \omega_1^3 \otimes \omega_1^3 + \omega_2^3 \otimes \omega_2^3,
$$

and (1.2) , (1.3) and (1.4) imply a relation

$$
\mathbb{II} = 2H\mathbb{I} - (K+1)\mathbb{I}.
$$

From now on we assume that the Gaussian curvature K satisfies $K > -1$. Then from (1.2) and (1.3) , it is easy to see that the second fundamental form II derived from n is a Riemannian metric on M. We equip a conformal structure $[\mathbb{I}]$ determined by \mathbb{I} . This conformal structure is called the second conformal structure of M with respect to the orientation determined by n.

Remark 1.1. If the Gaussian curvature K satisfies $K < -1$, then the second fundamental form II defines a Lorentzian metric on M and hence induce a Lorentz conformal structure $[\mathbb{I}]$ on M .

Using a positive function σ , let us represent the Gaussian curvature $K > -1$ by

$$
K = -1 + \sigma^2
$$

and take a conformal coordinate $z = x + yi$ with respect to the second conformal structure [II]. Then one can check that the quadratic differential

$$
Q\,dz^2 = \langle \partial f, \partial f \rangle \,\mathrm{d}z^2
$$

is well defined on the Riemann surface $(M, [\mathbb{I}])$, see for example [\[6\]](#page-22-9). Here we use the following notation for operators ∂ and ∂ :

$$
\partial = \frac{1}{2} \left(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \right), \quad \bar{\partial} = \frac{1}{2} \left(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \right).
$$

This quadratic differential $Q dz^2$ will be called the Klotz differential of (M, f) . We will prove later that the constancy of the Gaussian curvature $K > -1$ is characterized by holomorphicity of the Klotz differential, see Theorem [A.](#page-1-0)

From the above argument, the first and second fundamental forms are given by

$$
\mathbf{I} = Q \, \mathrm{d}z^2 + 2\ell \, \mathrm{d}z \mathrm{d}\bar{z} + \bar{Q} \, \mathrm{d}\bar{z}^2, \quad \mathbf{I} = 2m \, \mathrm{d}z \mathrm{d}\bar{z},
$$

where $\ell > 0$ and $m > 0$ are some positive functions. From these expressions, it is easy to see that $p \in M$ is an umbilic point if and only if $Q(p) = 0$. Then from [\(1.3\)](#page-3-3) and the Gaussian curvature $K = -1 + \sigma^2$, the equation $\sigma^{-2}m^2 = \ell^2 - |Q|^2$ holds. Let us introduce a real function u by $\ell + \sigma^{-1}m = e^u$. Then from the equation $\ell - \sigma^{-1}m = |Q|^2 e^{-u}$, we have

$$
\ell = \frac{1}{2}(e^u + |Q|^2 e^{-u}), \quad m = \frac{\sigma}{2}(e^u - |Q|^2 e^{-u}).
$$

Note that since $m > 0$, we have a condition $e^{2u} > |Q|^2$. Therefore the first and second fundamental forms are rephrased as

(1.6)
$$
I = Q dz^{2} + (e^{u} + |Q|^{2} e^{-u}) dz d\bar{z} + \bar{Q} d\bar{z}^{2}, \quad I = \sigma (e^{u} - |Q|^{2} e^{-u}) dz d\bar{z}.
$$

Note that determinant of the coefficient matrix of I and the mean curvature H can be computed respectively as $(e^u - |Q|^2 e^{-u})^2$ and

(1.7)
$$
H = \frac{\sigma}{2} \frac{e^{2u} + |Q|^2}{e^{2u} - |Q|^2}.
$$

Using the relation [\(1.5\)](#page-4-0), we compute the third fundamental form $\mathbb{I} = \langle d\mathbf{n}, d\mathbf{n} \rangle$ as

(1.8)
$$
\mathbb{I} = \sigma^2 \left(-Q \, dz^2 + (e^u + |Q|^2 e^{-u}) \, dz \, d\bar{z} - \bar{Q} \, d\bar{z}^2 \right).
$$

1.2. The 2×2 -matrix model of \mathbb{H}^3 . Let $\{e_0, e_1, e_2, e_3\}$ be the standard basis of $\mathbb{E}^{1,3}$. We identify e_0, e_1, e_2 and e_3 with the following 2×2 matrices:

$$
(1.9) \qquad \boldsymbol{e}_0 = \left(\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array}\right), \quad \boldsymbol{e}_1 = \left(\begin{array}{cc} 1 & 0 \\ 0 & -1 \end{array}\right), \ \boldsymbol{e}_2 = \left(\begin{array}{cc} 0 & -i \\ i & 0 \end{array}\right) \text{ and } \boldsymbol{e}_3 = \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right).
$$

Then the Minkowski 4-space $\mathbb{E}^{1,3}$ is identified with the space of all complex Hermitian 2×2 matrices Her₂C, that is, $\mathbb{E}^{1,3} \cong \text{Her}_2\mathbb{C} = \left\{\xi_0 e_0 + \xi_1 e_1 + \xi_2 e_2 + \xi_3 e_3 \mid \xi_0, \xi_1, \xi_2, \xi_3 \in \mathbb{R} \right\}.$ It is easy to see that for $\xi \in \text{Her}_2\mathbb{C}$, $-\det \xi = -\xi_0^2 + \xi_1^2 + \xi_2^2 + \xi_3^2$. Thus the Lorentzian metric of $\text{Her}_2\mathbb{C}$ is described as

$$
\langle \xi, \eta \rangle = -\frac{1}{2} \text{tr}(\xi \mathbf{e}_2{}^t \eta \mathbf{e}_2).
$$

In particular, we have $\langle \xi, \xi \rangle = - \det \xi$ for $\xi \in \text{Her}_2\mathbb{C}$. Note that $\{e_0, e_1, e_2, e_3\}$ is an orthonormal basis of $\mathbb{E}^{1,3}$. Then we have the identification: $\mathbb{H}^3 = \{ \xi \in \text{Her}_2\mathbb{C} \mid \det \xi =$ 1, $\text{tr } \xi > 0$. Moreover, the special linear group $\text{SL}_2\mathbb{C}$ acts isometrically and transitively on the hyperbolic 3-space via the action: $SL_2\mathbb{C} \times \mathbb{H}^3 \to \mathbb{H}^3$ by $(g,\xi) \mapsto g \xi g^*$. Here the superscript * denotes the Hermitian conjugate, that is, $g^* = {}^t\bar{g}$ for a matrix g. The isotropy subgroup of this action at $e_0 = (1, 0, 0, 0) \approx$ id is the special unitary group SU₂. Hence \mathbb{H}^3 is represented by $\mathbb{H}^3 = SL_2\mathbb{C}/SU_2$ as a *Riemannian symmetric space*. The natural projection $\pi : SL_2\mathbb{C} \to \mathbb{H}^3$ is explicitly given by $\pi(g) = gg^*, g \in SL_2\mathbb{C}$. In other words, \mathbb{H}^3 is represented as

$$
\mathbb{H}^3 = \{ gg^* \mid g \in \mathrm{SL}_2\mathbb{C} \}.
$$

Remark 1.2. In this context, the simple Lie group $SL_2\mathbb{C}$ is regarded as a simple real Lie group and as a double covering of the special Lorentz group $\text{SO}_{1,3}^+$. The real Lie algebra $\mathfrak{sl}_2\mathbb{C}$ is spanned by the basis $\mathcal{E} = \{ie_1, ie_2, ie_3, e_1, e_2, e_3\}.$ With respect to the scalar product

$$
\langle X, Y \rangle_K = \frac{1}{2} \operatorname{tr}(XY), \quad (X, Y \in \mathfrak{sl}_2\mathbb{C}),
$$

 $\mathcal E$ has signature $(-, -, -, +, +, +)$ and induces a bi-invariant semi-Riemannian metric on SL₂C. The Lie algebras \mathfrak{u}_1 and \mathfrak{su}_2 are expressed as $\mathfrak{u}_1 = \mathbb{R}(ie_1)$ and $\mathfrak{su}_2 = \mathbb{R}(ie_1) \oplus$ $\mathbb{R}(ie_2) \oplus \mathbb{R}(ie_3)$, respectively. Thus \mathfrak{su}_2 is negative definite linear subspace of $\mathfrak{sl}_2\mathbb{C}$.

The tangent space $\mathfrak{m} = T_{\mathcal{C}_0} \mathbb{H}^3$ is given by

$$
\mathfrak{m}=\mathfrak{sl}_2\mathbb{C}\cap\mathrm{Her}_2\mathbb{C}=\mathbb{R}e_1\oplus\mathbb{R}e_2\oplus\mathbb{R}e_3.
$$

One can see that the restriction of the scalar product $\langle \cdot, \cdot \rangle_K$ to m coincides with the Lorentz scalar product on $\mathfrak{m} \subset \text{Her}_2\mathbb{C}$. The induced Riemannian metric on $\mathbb{H}^3 = SL_2\mathbb{C}/SU_2$ from $\langle \cdot, \cdot \rangle_K$ is of constant curvature -1 .

1.3. Unit tangent sphere bundle. The unit tangent sphere bundle $U\mathbb{H}^3$ of \mathbb{H}^3 is represented as,

$$
\mathrm{U}\mathbb{H}^3=\{(\boldsymbol{x},\boldsymbol{v})\in\mathrm{Her}_2\mathbb{C}\times\mathrm{Her}_2\mathbb{C}\,\,|\,\,\det\boldsymbol{x}=1,\mathrm{tr}\,\boldsymbol{x}>0,\,\det\boldsymbol{v}=-1,\,\langle\boldsymbol{x},\boldsymbol{v}\rangle=0\},
$$

see [\[3\]](#page-22-6) in detail. The special liner group $SL_2\mathbb{C}$ acts isometrically and transitively on $U\mathbb{H}^3$ via the action $g \cdot (\bm{x}, \bm{v}) = (g\bm{x}g^*, g\bm{v}g^*)$. The isotropy subgroup of SL₂C at (e_0, e_1) is

$$
H = \left\{ \mathrm{diag}(e^{i\theta}, e^{-i\theta}) \mid \theta \in \mathbb{R} \right\} = \mathrm{U}_1.
$$

Thus the unit tangent sphere bundle $U\mathbb{H}^3$ is represented by $SL_2\mathbb{C}/U_1$ as a homogeneous manifold. The Lie algebra h of H is $h = \{diag(ia_2, -ia_2) | a_2 \in \mathbb{R}\}$. The tangent space $\mathfrak{p} := T_{(\boldsymbol{e}_0, \boldsymbol{e}_1)} \cup \mathbb{H}^3$ is given by

(1.10)
$$
\mathfrak{p} = \left\{ \left(\begin{array}{cc} a_1 & b_1 + ib_2 \\ c_1 + ic_2 & -a_1 \end{array} \right) \middle| a_1, b_1, b_2, c_1, c_2 \in \mathbb{R} \right\}.
$$

Note that

$$
\mathfrak{p}=\{\mathbb{R} e_1\oplus\mathbb{R} e_2\oplus\mathbb{R} e_3\}\oplus\{\mathbb{R}(ie_2)\oplus\mathbb{R}(ie_3)\}=\mathfrak{m}\oplus\{\mathbb{R}(ie_2)\oplus\mathbb{R}(ie_3)\}.
$$

The scalar product $\langle \cdot, \cdot \rangle_K$ induces a $SL_2\mathbb{C}$ -invariant semi-Riemannian metric on UH³. The resulting homogeneous semi-Riemannian space $G/H = SL_2\mathbb{C}/\mathrm{U}(1)$ is naturally reductive. The unit tangent sphere bundle admits two kinds of CR-structures J_1 and J_2 . The resulting CR-manifold $(\mathrm{U}\mathbb{H}^3, J_1)$ is the *twistor* CR-manifold of \mathbb{H}^3 in the sense of LeBrun [\[9\]](#page-22-10).

1.4. Natural projections and two Gauss maps. We have several natural projections from the unit tangent sphere bundle U \mathbb{H}^3 . To define the projections, we introduce notion of the space $\text{Geo}(\mathbb{H}^3)$ of all oriented geodesics in \mathbb{H}^3 which is identified with the Grassmannian manifold $\text{Gr}_{1,1}(\mathbb{E}^{1,2})$ of all oriented timelike planes in $\mathbb{E}^{1,3}$. The unit tangent sphere bundle U \mathbb{H}^3 is a principal line bundle over $\text{Gr}_{1,1}(\mathbb{E}^{1,2})$. The natural projection $\pi_0: \text{U}\mathbb{H}^3 \to \text{Geo}(\mathbb{H}^3)$ is given explicitly by $\pi_0(\boldsymbol{x}, \boldsymbol{v}) = \boldsymbol{x} \wedge \boldsymbol{v}$.

The space $\text{Gr}_{1,1}(\mathbb{E}^{1,2})$ is a homogeneous space of $G = \text{SL}_2\mathbb{C}$. The isotropy subgroup K at $e_0 \wedge e_1$ is

$$
K = GL_1 \mathbb{C} = \{ \text{diag}(a, a^{-1}) \mid a \in \mathbb{C}^\times \} \cong GL_1 \mathbb{C}.
$$

The Lie algebra \mathfrak{k} of K is $\mathfrak{k} = \{ \text{diag}(ia_1, -a_1) \mid a_1 \in \mathbb{C} \}$. The tangent space of $\text{Gr}_{1,1}(\mathbb{E}^{1,2})$ at $e_0 \wedge e_1$ is identified with

(1.11)
$$
\mathfrak{q} = \left\{ \left(\begin{array}{cc} 0 & b_1 + ib_2 \\ c_1 + ic_2 & 0 \end{array} \right) \middle| b_1, b_2, c_1, c_2 \in \mathbb{R} \right\}.
$$

One can see that $G/K = SL_2\mathbb{C}/GL_1\mathbb{C}$ is a semi-Riemannian symmetric space with respect to the metric induced from $\langle \cdot, \cdot \rangle$. For more detail, see [\[3\]](#page-22-6).

Next for $(x, v) \in \text{U} \mathbb{H}^3$, it is easy to see that v takes values in the *de Sitter 3-space*

$$
\mathbb{S}^{1,2} = \{ (x_0, x_1, x_2, x_3) \in \mathbb{E}^{1,3} \mid -x_0^2 + x_1^2 + x_2^2 + x_3^2 = 1 \},\
$$

which is a simply connected Lorentzian 3-manifold of constant sectional curvature 1. Thus there are two natural projections $\pi_+ : \text{U} \mathbb{H}^3 \to \mathbb{H}^3$ and $\pi_- : \text{U} \mathbb{H}^3 \to \mathbb{S}^{1,2}$, $\pi_+(\mathbf{x}, \mathbf{v}) = \mathbf{x}$ and $\pi_-(x, v) = v$, respectively. Finally there is also a natural projection from $\pi_* : SL_2\mathbb{C} \to \mathbb{U} \mathbb{H}^3$

by $\pi_*(g) = (gg^*, ge_1g^*)$. Note that the projection $\pi : SL_2\mathbb{C} \to \mathbb{H}^3 = SL_2\mathbb{C}/SU_2$ is given by the composition $\pi = \pi_+ \circ \pi_*$. see [\[3\]](#page-22-6).

Then we have a natural projection $\pi_0 : \text{U} \mathbb{H}^3 \to \text{Geo}(\mathbb{H}^3)$ by $\pi_0(\boldsymbol{x}, \boldsymbol{v}) = \boldsymbol{x} \wedge \boldsymbol{v}$. Next for $(x, v) \in \text{U} \mathbb{H}^3$, it is easy to see that v takes values in the de Sitter 3-space

$$
\mathbb{S}^{1,2} = \{ (x_0, x_1, x_2, x_3) \in \mathbb{E}^{1,3} \mid -x_0^2 + x_1^2 + x_2^2 + x_3^2 = 1 \},\
$$

which is a simply connected Lorentzian 3-manifold of constant sectional curvature 1. Thus there are two natural projections $\pi_+ : \text{U} \mathbb{H}^3 \to \mathbb{H}^3$ and $\pi_- : \text{U} \mathbb{H}^3 \to \mathbb{S}^{1,2}$, $\pi_+(\mathbf{x}, \mathbf{v}) = \mathbf{x}$ and $\pi_-(x, v) = v$, respectively. Finally there is also a natural projection from $\pi_* : SL_2(\mathbb{C} \to \mathbb{U} \mathbb{H}^3)$ by $\pi_*(g) = (gg^*, ge_1g^*)$. Note that the projection $\pi : SL_2\mathbb{C} \to \mathbb{H}^3 = SL_2\mathbb{C}/SU_2$ is given by the composition $\pi = \pi_+ \circ \pi_*$.

Using these projections, we will recall two Gauss maps for a surface in \mathbb{H}^3 , which are important in this paper. Note that several natural Gauss maps were already defined in see [\[3,](#page-22-6) Appendix of the archive version].

Definition 1.1. Let $f : M \to \mathbb{H}^3$ be a surface with unit normal n , and let $\text{Gr}_{1,1}(\mathbb{E}^{1,3}) =$ Geo(\mathbb{H}^3). The Legendrian Gauss map F and the Lagraingian Gauss map L of f are respective defined by

$$
F = (f, n) : M \to \mathbf{U} \mathbb{H}^3 \quad \text{and} \quad L = f \wedge n : M \to \mathbf{Gr}_{1,1}(\mathbb{E}^{1,3}).
$$

Remark 1.3. Let us denote by η and Ω the canonical contact form of UH³ and the canonical symplectic form of $\text{Gr}_{1,1}(\mathbb{E}^{1,3})$. Then one can see that F is Legendre, that is, $F^*\eta = 0$ and L is Lagrangian, that is, $L^*\Omega = 0$, respectively, see [\[3\]](#page-22-6). In [3], using the conformal structure induced from the first fundamental form, we characterized constant mean curvature surfaces by harmonicity of the Legendrian Gauss map.

2. Harmonic Gauss maps

In this section, we characterize constant Gaussian curvature $K > -1$ surfaces in terms of harmonicity of Legendrian and Lagrangian Gauss maps, respectively. These harmonic Gauss maps naturally induce one parameter family, the so-called spectral parameter family, of constant Gaussian curvature $K > -1$ surfaces. Finally a deformation of the spectral parameter induces harmonic maps into \mathbb{S}^2 in case of $K > 0$ or \mathbb{H}^2 in case of $-1 < K < 0$, respectively.

2.1. Lax representation. Let $f : M \to \mathbb{H}^3$ be a surface of Gaussian curvature $K =$ $-1 + \sigma^2$ with some positive function σ . Let $D \subset M$ be a simply connected domain and $z = x + yi$ on D a conformal coordinate on D with respect to [II]. We define two matrices as

(2.1)
$$
\hat{\mathbf{e}}_2 = -\frac{1}{2} (\mathbf{e}_2 - i\mathbf{e}_3) = \begin{pmatrix} 0 & i \\ 0 & 0 \end{pmatrix}, \quad \hat{\mathbf{e}}_3 = -\frac{1}{2} (\mathbf{e}_2 + i\mathbf{e}_3) = \begin{pmatrix} 0 & 0 \\ -i & 0 \end{pmatrix}.
$$

Note that $\hat{\mathbf{e}}_3^* = \hat{\mathbf{e}}_2$ and $\langle \hat{\mathbf{e}}_i, \hat{\mathbf{e}}_i \rangle = 0$ $(i = 2, 3)$ and $\langle \hat{\mathbf{e}}_2, \hat{\mathbf{e}}_3 \rangle = 1/2$, and thus $\{\hat{\mathbf{e}}_2, \hat{\mathbf{e}}_3\}$ forms null basis. Let us consider a change of basis ${e, e^*}$ defined by

$$
(e, e^*) = (\hat{e}_2, \hat{e}_3) \begin{pmatrix} e^{u/2} & \bar{Q}e^{-u/2} \\ Qe^{-u/2} & e^{u/2} \end{pmatrix}.
$$

Then it is easy to see that $\langle e, e \rangle = Q$, $\langle e^*, e^* \rangle = \overline{Q}$ and $\langle e, e^* \rangle = (e^u + |Q|^2 e^{-u})/2$. Since ${e_0, e_1, e, e^*}$ is a basis in $\mathbb{E}^{1,3}$, there exists a map $\Psi : D \subset M \to SL_2\mathbb{C}$ satisfying the following equations:

$$
f = \Psi \mathbf{e}_0 \Psi^*, \ n = \Psi \mathbf{e}_1 \Psi^*, \ \partial f = \Psi \mathbf{e} \Psi^* \text{ and } \bar{\partial} f = \Psi \mathbf{e}^* \Psi^*.
$$

It is clear that Ψ is a moving frame along the surface f. The moving frame Ψ will be called the *coordinate frame*. We obtain the following Lax representation for f .

Proposition 2.1. The coordinate frame Ψ satisfies the Lax equation:

(2.2)
$$
\partial \Psi = \Psi U, \quad \bar{\partial} \Psi = \Psi V,
$$

where U and V are given by

(2.3)
$$
U = \left(\frac{1}{4}\partial u + p\right)e_1 + \frac{1+\sigma}{2}\left(e^{u/2}\hat{e}_2 + Qe^{-u/2}\hat{e}_3\right),
$$

(2.4)
$$
V = -\left(\frac{1}{4}\bar{\partial}u + \bar{p}\right)\mathbf{e}_1 + \frac{1-\sigma}{2}\left(\bar{Q}e^{-u/2}\hat{\mathbf{e}}_2 + e^{u/2}\hat{\mathbf{e}}_3\right),
$$

and e_1 , \hat{e}_2 and \hat{e}_3 are defined in [\(1.9\)](#page-5-0) and [\(2.1\)](#page-7-1), and

$$
p = \frac{-\bar{\partial}Q + e^{-u}Q\partial\bar{Q}}{2(e^u - |Q|^2e^{-u})}.
$$

The compatibility conditions, that is, the Gauss-Codazzi equations for a surface f , are given as follows:

(2.5)
$$
\bar{\partial}\partial u + \frac{K}{2}(e^u - |Q|^2 e^{-u}) + 2(\partial \bar{p} + \bar{\partial}p) = 0,
$$

(2.6)
$$
2\bar{\partial}(Q\sigma) + (e^u + |Q|^2 e^{-u})\partial \sigma = 0.
$$

Conversely, let u be a real smooth function and Q a complex valued smooth function and $K = -1 + \sigma^2$ with some positive function σ be solutions of the Gauss-Codazzi equations (2.5) and (2.6) such that $e^{2u} - |Q|^2 > 0$ holds, and let Ψ be a solution of the Lax equation [\(2.2\)](#page-8-2). Define respective maps

$$
f = \Psi \mathbf{e}_0 \Psi^*
$$
 and $n = \Psi \mathbf{e}_1 \Psi^*$.

Then f is a surface in \mathbb{H}^3 with unit normal n and the Gaussian curvature of f is $K =$ $-1 + \sigma^2 > -1.$

Proof. Differentiating $f = \Psi e_0 \Psi^*$ by z, we have $\partial f = \Psi (U + V^*) \Psi^*$. Comparing it with the formula $\partial f = \Psi(e^{u/2}\hat{\boldsymbol{e}}_2 + Qe^{-u/2}\hat{\boldsymbol{e}}_3) \Psi^*$, we obtain

$$
(2.7) \t v_{11} = -\overline{u_{11}}, \ v_{22} = -\overline{u_{22}}, \ u_{12} + \overline{v_{21}} = ie^{u/2}, \ u_{21} + \overline{v_{12}} = -iQe^{-u/2},
$$

where we set $U = (u_{ij})_{1 \leq i,j \leq 2}$ and $V = (v_{ij})_{1 \leq i,j \leq 2}$. Note that since Ψ takes values in $SL_2\mathbb{C}$, we have $u_{11} + u_{22} = 0 = v_{11} + v_{22}$. Since $\langle \partial f, \partial n \rangle = 0$ and $\partial n = \Psi (U \mathbf{e}_1 + \mathbf{e}_1 V^*) \Psi^*$, we have

$$
\text{tr}\{(e^{u/2}\hat{\boldsymbol{e}}_2 + Qe^{-u/2}\hat{\boldsymbol{e}}_3)\cdot \boldsymbol{e}_2{}^t(U\boldsymbol{e}_1 + \boldsymbol{e}_1 V^*)\boldsymbol{e}_2\} = 0,
$$

thus $e^{u/2}(u_{21} - \overline{v_{12}}) + Qe^{-u/2}(u_{12} - \overline{v_{21}}) = 0$. Using [\(2.7\)](#page-8-3), this is rephrased as

(2.8)
$$
u_{21} + Qe^{-u}u_{12} = 0.
$$

Next by the second fundamental form II in [\(1.2\)](#page-3-2) we have $-\langle \bar{\partial} f, \partial n \rangle = \sigma (e^u - |Q|^2 e^{-u})/2$. A direct computation similar to the above shows that this is equivalent with $\sigma(e^u |Q|^2 e^{-u}$)/2 = $i(\bar{Q}e^{-u/2}u_{21} + e^{u/2}u_{12}) - (e^u - |Q|^2 e^{-u})/2$. Comparing this with [\(2.8\)](#page-9-0) and (2.7) , we obtain

$$
u_{12} = \frac{i}{2}(1+\sigma)e^{u/2}, \quad u_{21} = -\frac{i}{2}(1+\sigma)Qe^{-u/2},
$$

$$
v_{12} = \frac{i}{2}(1-\sigma)\bar{Q}e^{-u/2}, \quad v_{21} = -\frac{i}{2}(1-\sigma)e^{u/2}.
$$

To obtain the diagonal part of U and V, we use the compatibility condition $\bar{\partial}\partial f = \partial\bar{\partial}f$. A direct computation together with the relation in [\(2.7\)](#page-8-3) shows that this is equivalent to

$$
-2\overline{u}_{11} + \frac{\overline{\partial}u}{2} = 2\overline{Q}e^{-u}u_{11} + e^{-u}\left(\overline{Q}_z - \frac{\partial u}{2}\overline{Q}\right),
$$

$$
-2Q\overline{u}_{11} - \overline{\partial}Q + \frac{\overline{\partial}u}{2}Q = \left(2u_{11} - \frac{\partial u}{2}\right)e^u.
$$

From these equations, we obtain

(2.9)
$$
u_{11} = -\overline{v_{11}} = -\frac{\partial u}{4} + p, \quad p = \frac{-\overline{\partial}Q + e^{-u}Q\partial\overline{Q}}{2(e^u - |Q|^2e^{-u})}
$$

Now the compatibility conditions, that is $\bar{\partial}\partial\Psi = \partial\bar{\partial}\Psi$, are given by a straightforward computation: First note that $\bar{\partial}\partial\Psi = \partial\bar{\partial}\Psi$ is equivalent to the zero curvature equation $\mathcal{L} := \partial U - \partial V + [V, U] = 0$. Then (1, 1)- and (2, 2)-entries of \mathcal{L} give the equation [\(2.5\)](#page-8-0), that is, the Gauss equation. Then $(1, 2)$ - and $(2, 1)$ -entries of $\mathcal{L} = 0$ can be computed as

.

(2.10)
$$
i(\bar{\partial}X - 2\bar{p}X)e^{u/2} - i(\bar{Q}\partial Y + Y\partial \bar{Q} + 2Y\bar{Q}p)e^{-u/2} = 0,
$$

(2.11)
$$
i(-Q\bar{\partial}X - X\bar{\partial}Q - 2XQ\bar{p})e^{-u/2} - i(-\partial Y + 2Yp)e^{u/2} = 0,
$$

where $X = (1 + \sigma)/2$ and $Y = (1 - \sigma)/2$. From these equations and the expression p in [\(2.9\)](#page-9-1), we have the equation [\(2.6\)](#page-8-1), that is, the Codazzi equation.

The converse statement is almost clear by a direct computation for $f = \Psi e_0 \Psi^*$ with $n = \Psi e_1 \Psi^*$. The first fundamental form of f is non-degenerate by the assumption, that is, f gives a surface. Then *n* is clearly the unit normal of f. 2.2. Harmonicities of Gauss maps. We now characterize a constant Gaussian curvature surface $K > -1$ in \mathbb{H}^3 as follows in terms of various Gauss maps as in Theorem [A.](#page-1-0)

Proof of Theorem [A](#page-1-0). (1) \Rightarrow (4): This is clear from the Codazzi equation, that is, the equation [\(2.6\)](#page-8-1). (4) \Rightarrow (1): Conversely assume that $Q dz^2$ is holomorphic. Thus note that zeros of Q are isolated. On a region such that $Q \neq 0$, we consider the Codazzi equation [\(2.6\)](#page-8-1) and its conjugation with multiplying $-(e^u + |Q|^2 e^{-u})/2$. Adding it to the Codazzi equation multiplying \overline{Q} , we obtain $-\overline{(\overline{\partial}\sigma)}(e^u - |Q|^2 e^{-u})^2/2 = 0$. Since $\langle \overline{\partial}f, \partial n \rangle =$ $\sigma(e^u - |Q|^2 e^{-u})/2 > 0$, this equation holds if and only if σ is constant, thus the surface is a constant Gaussian curvature surface with $K > -1$. Then we use the argument of continuity and it implies that the surface is a constant Gaussian curvature surface $K > -1$ on M.

We now show that (1) \Leftrightarrow (2). We use the harmonicity criterion [\(B.1\)](#page-21-0) for $F : M \rightarrow$ $G/H = SL_2\mathbb{C}/U_1$ with reductive decomposition $\mathfrak{sl}_2\mathbb{C} = \mathfrak{u}_1 \oplus \mathfrak{p}$, where \mathfrak{p} is given by [\(1.10\)](#page-6-0). From Proposition [2.1,](#page-8-4) it is easy to see that $\alpha = \Psi^{-1} d\Psi = \alpha_{\mathfrak{h}} + \alpha'_{\mathfrak{p}} + \alpha''_{\mathfrak{p}}$ is given by

$$
\alpha_{\mathfrak{h}} = \left(\frac{\partial u}{4} + p\right) e_1 \mathrm{d}z - \left(\frac{\bar{\partial}u}{4} + \bar{p}\right) e_1 \mathrm{d}\bar{z}, \ \alpha_{\mathfrak{p}}' = X e \mathrm{d}z \ \text{and} \ \alpha_{\mathfrak{p}}'' = Y e^* \mathrm{d}\bar{z},
$$

where **e** is defined in [\(2.1\)](#page-7-1) and $X = (1+\sigma)/2$ and $Y = (1-\sigma)/2$. By a direct computation, we have $[\alpha'_{\mathfrak{p}} \wedge \alpha''_{\mathfrak{p}}] = XY(e^u - |Q|^2 e^{-u}) e_1 dz \wedge d\bar{z}$. From this it is clear that $[\alpha'_{\mathfrak{p}} \wedge \alpha''_{\mathfrak{p}}]_{\mathfrak{p}} = 0$. Next we compute $d(*\alpha_p) + [\alpha \wedge *\alpha_p]$. It is easy to see that the $(1, 1)$ - and $(2, 2)$ -entries are zero. Then the $(1, 2)$ - and $(2, 1)$ -entries can be computed as

$$
\left\{ (2\bar{p}X - \bar{\partial}X)e^{u/2} - (2p\bar{Q}Y + \partial(Y\bar{Q}))e^{-u/2} \right\} dz \wedge d\bar{z},
$$

$$
\left\{ (\bar{\partial}(XQ) + 2\bar{p}QX)e^{-u/2} + (\partial Y - 2pY)e^{u/2} \right\} dz \wedge d\bar{z}.
$$

Then using the equations [\(2.10\)](#page-9-2) and [\(2.11\)](#page-9-3), $d(*\alpha_{p}) + [\alpha \wedge *\alpha_{p}] = 0$ is equivalent with

$$
\bar{\partial}X - 2\bar{p}X = \partial Y + 2pY = 0.
$$

These equations hold if and only if $p = 0$ and this happens if and only if $\overline{\partial}Q = 0$, that is, the Klotz differential is holomorphic. Since (1) is equivalent with (4), the claim follows.

Finally we show (1) \Leftrightarrow (3). Let us consider the harmonicity of $L : M \rightarrow G/K =$ $SL_2\mathbb{C}/GL_1\mathbb{C}$ equipped with the reductive decomposition $\mathfrak{sl}_2\mathbb{C} = \mathfrak{k} \oplus \mathfrak{q}$, where \mathfrak{q} is given by [\(1.11\)](#page-6-1). Since Geo(\mathbb{H}^3) is a symmetric space, the first condition in [\(B.1\)](#page-21-0), that is, $[\alpha'_{\mathsf{q}} \wedge \alpha''_{\mathsf{q}}]_{\mathsf{q}} =$ 0 is vacuous. Then the harmonicity of L is exactly the same as the case (2) and the claim follows. This completes the proof.

Corollary 2.1. Let U^{λ} and V^{λ} be a \mathbb{S}^1 -family of matrices parameterized by $\lambda \in \mathbb{S}^1$:

(2.12)
$$
U^{\lambda} = \left(\frac{1}{4}\partial u + p\right)\mathbf{e}_1 + \frac{1+\sigma}{2}\lambda^{-1}\left(e^{u/2}\hat{\mathbf{e}}_2 + Qe^{-u/2}\hat{\mathbf{e}}_3\right),
$$

(2.13)
$$
V^{\lambda} = -\left(\frac{1}{4}\bar{\partial}u + \bar{p}\right)\mathbf{e}_1 + \frac{1-\sigma}{2}\lambda\left(\bar{Q}e^{-u/2}\hat{\mathbf{e}}_2 + e^{u/2}\hat{\mathbf{e}}_3\right).
$$

Moreover, set a \mathbb{S}^1 -family of Maure-Cartan forms as

$$
\alpha^{\lambda} = U^{\lambda} \mathrm{d} z + V^{\lambda} \mathrm{d} \bar{z}.
$$

Then any statement in Theorem [A](#page-1-0) is equivalent with the following statement:

(5) $d + \alpha^{\lambda}$ is a family of flat connections on $M \times SL_2\mathbb{C}$.

Proof. The family of Maurer-Cartan form α^{λ} can be decomposed as

$$
\alpha^{\lambda} = \lambda^{-1} \alpha'_{\mathfrak{p}} + \alpha_{\mathfrak{k}} + \lambda \alpha''_{\mathfrak{p}},
$$

according to the decomposition of $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ of the symmetric space $\text{Geo}(\mathbb{H}^3) = G/K$ and the complex structure of M. Here note that $\lambda^{-1} \alpha'_{\mathfrak{p}} = \mathrm{Off}(U^{\lambda})dz$ and $\lambda \alpha''_{\mathfrak{p}} = \mathrm{Off}(V^{\lambda})d\bar{z}$, where $\mathrm{Off}(X)$ denotes the off-diagonal part of X. It is well known that, see for example Theorem [B.1](#page-21-1) in Appendix [B,](#page-21-2) $d + \alpha^{\lambda}$ is flat if and only if the corresponding Lagrangian Gauss map $L: M \to \text{Geo}(\mathbb{H}^3)$ is harmonic.

Remark 2.1.

- (1) If the surface is a flat surface $K = 0$, then the harmonic Legendrian and Lagrangian Gauss maps become conformal. In fact a direct computation shows that $\langle \partial F, \partial F \rangle =$ $\langle \partial L, \partial L \rangle = 0$. Moreover under the identification, Geo(\mathbb{H}^3) = $\mathbb{S}^2 \times \mathbb{S}^2 \setminus \triangle$, the Lagrangian Gauss map L is the pair of hyperbolic Gauss maps. For flat surfaces, Gálvez et al [\[4\]](#page-22-11) showed that both hyperbolic Gauss maps are conformal.
- (2) Here $SL_2\mathbb{C}$ is the double cover of $SO_{1,3}^+$ and we consider it as a real Lie group. Thus $\alpha'_{\mathfrak{p}}$ - and $\alpha''_{\mathfrak{p}}$ -parts of the Maurer-Cartan form in the proof of Theorem [A](#page-1-0) should be in the complexification of $\mathfrak{sl}_2\mathbb{C}$, that is, it is given by $\mathfrak{sl}_2\mathbb{C}\times\mathfrak{sl}_2\mathbb{C}$, see for example [\[3,](#page-22-6) Section 7.1].
- (3) The family of flat connections $d + \alpha^{\lambda}$ induces a family of maps $\{\Psi^{\lambda}\}_{\lambda \in \mathbb{S}^1}$ such that $(\Psi^{\lambda})^{-1} d\Psi^{\lambda} = \alpha^{\lambda}$, the so-called extended frame, see Section [2.3.](#page-11-0)

2.3. Spectral parameter. We now show that the parameter $\lambda \in \mathbb{S}^1$ in Corollary [2.1](#page-10-1) naturally appears in the Gauss-Codazzi equations of a constant Gaussian curvature $K >$ -1 surface. From (2.5) and (2.6) , they are given as

(2.14)
$$
\bar{\partial}\partial u + \frac{K}{2}(e^u - |Q|^2 e^{-u}) = 0 \text{ and } \bar{\partial}Q = 0.
$$

In particular, note that the function $p = (-\bar{\partial}Q + e^{-u}Q\partial\bar{Q})(2e^u - 2|Q|^2e^{-u})^{-1}$ vanishes by holomorphicity of Q.

Remark 2.2. Since the Klotz differential $Q dz^2$ is holomorphic, on a (simply connected) region free of umbilics, we may reparameterize z so that $Q = 1$. With respect to such a coordinate z, the Gauss-Codazzi equation has the form

$$
\bar{\partial}\partial u + K\sinh u = 0, \quad (K > -1).
$$

This equation appears as a harmonic map equation for $\mathbb{D} \subset \mathbb{C} \to \mathbb{H}^2$ in case of $-1 < K < 0$ or \mathbb{S}^2 in case of $K > 0$, respectively, see [\[7\]](#page-22-12).

It is easy to see that the Gauss-Codazzi equations [\(2.14\)](#page-11-1) are invariant under the deformation $Q \mapsto \lambda^{-2}Q$ for $\lambda \in \mathbb{S}^1$. Then from fundamental theory of surface, there is a \mathbb{S}^1 -family of constant Gaussian curvature $K > -1$ surfaces in \mathbb{H}^3 . Let $\widetilde{\Psi}$ be the corresponding family of moving frames, that is,

(2.15)
$$
\widetilde{\Psi}^{-1} d\widetilde{\Psi} = U dz + V d\overline{z},
$$

where U and V and defined in [\(2.3\)](#page-8-5) and [\(2.4\)](#page-8-6) with $p = 0$, σ constant and the holomorphic Q and anti-holomorphic \overline{Q} are replaced by $\lambda^{-2}Q$ and $\lambda^2\overline{Q}$, respectively. Consider a gauge transformation by right:

$$
\widetilde{\Psi} \longmapsto \Psi^{\lambda} = \widetilde{\Psi} \operatorname{diag}(\lambda^{1/2}, \lambda^{-1/2}).
$$

Then a straightforward computation shows that $U^{\lambda} = (\Psi^{\lambda})^{-1} \partial \Psi^{\lambda}$ and $V^{\lambda} = (\Psi^{\lambda})^{-1} \bar{\partial} \Psi^{\lambda}$ are given in [\(2.12\)](#page-10-2) and [\(2.13\)](#page-10-3) with $p = \bar{p} = 0$, respectively. In this way the parameter $\lambda \in \mathbb{S}^1$ appears in the Gauss-Codazzi equations of a constant Gaussian curvature surface.

Definition 2.1. The solution Ψ^{λ} of

(2.16)
$$
(\Psi^{\lambda})^{-1} d\Psi^{\lambda} = U^{\lambda} dz + V^{\lambda} d\bar{z} \text{ with } \Psi^{\lambda}|_{z=z_{*}} = id
$$

is called the *extended frame* of a constant Gaussian curvature $K > -1$ surface f and the Lagrangian harmonic Gauss map $L: D \to \text{Geo}(\mathbb{H}^3)$, where z_* is some base point $z_* \in D$. Moreover, the parameter $\lambda \in \mathbb{S}^1$ in the extended frame Ψ^{λ} will be called the *spectral* parameter.

One can easily check that $U(\lambda) = U^{\lambda}$ and $V(\lambda) = V^{\lambda}$ satisfy the following conditions:

$$
\mathrm{Ad}(\mathbf{e}_1)U(\lambda) = U(-\lambda), \quad \mathrm{Ad}(\mathbf{e}_1)V(\lambda) = V(-\lambda).
$$

Thus the extended frame Ψ^{λ} takes values in the twisted loop group $\Lambda SL_2\mathbb{C}_\tau$, see Appendix [A](#page-20-0) for the definition. From the extended frame Ψ^{λ} , it is easy to obtain a \mathbb{S}^{1} -family of constant Gaussian curvature $K > -1$ surfaces.

Proposition 2.2. Let Ψ^{λ} be the extended frame of some constant Gaussian curvature $K > -1$ surface f and define respective maps f^{λ} and n^{λ} by

$$
f^{\lambda} = \Psi^{\lambda} e_0 (\Psi^{\lambda})^*
$$
 and $n^{\lambda} = \Psi^{\lambda} e_1 (\Psi^{\lambda})^*$.

Then, for each $\lambda \in \mathbb{S}^1$, f^{λ} is a constant Gaussian curvature $K > -1$ surface with unit normal n^{λ} and the following first and second fundamental forms:

$$
I^{\lambda} = \lambda^{-2}Q dz^{2} + (e^{u} + |Q|^{2}e^{-u}) dz d\bar{z} + \lambda^{2} \bar{Q} d\bar{z}^{2}, \quad I I^{\lambda} = \sigma(e^{u} - |Q|^{2}e^{-u}) dz d\bar{z}.
$$

In particular $f^{\lambda}|_{\lambda=1}$ and the original constant Gaussian curvature surface f are the same surface up to a rigid motion. Moreover, the maps

$$
F^{\lambda} = (f^{\lambda}, n^{\lambda}) : M \to \text{UIII}^3
$$
 and $L^{\lambda} = f^{\lambda} \wedge n^{\lambda} : M \to \text{Gr}_{1,1}(\mathbb{E}^{1,3})$

are the Legendrian harmonic map and the Lagrangian harmonic map of f^{λ} , respectively.

Proof. Since $\lambda \in \mathbb{S}^1$, we have

$$
f^{\lambda} = \Psi^{\lambda} \mathbf{e}_0 (\Psi^{\lambda})^* = \widetilde{\Psi} \mathbf{e}_0 \widetilde{\Psi}^*,
$$

where $\tilde{\Psi}$ is defined in [\(2.15\)](#page-12-0). Now the claim follows directly from Proposition [2.1.](#page-8-4) \Box

Definition [2.2](#page-12-1). A family of surfaces $\{f^{\lambda}\}_{\lambda \in \mathbb{S}^1}$ defined in Proposition 2.2 will be called the associated family of the constant Gaussian curvature $K > -1$ surface $f(= f^{\lambda}|_{\lambda=1}).$

2.4. A spectral parameter deformation of the harmonic Lagrangian Gauss maps. From Remark [2.2,](#page-11-2) for a constant Gaussian curvature surfaces with $K > -1$ but $K \neq 0$ in \mathbb{H}^3 , it is expected that there are harmonic maps into $\mathbb{H}^2 = SU_{1,1}/U_1$ or $\mathbb{S}^2 = SU_2/U_1$ which are obtained from the constant Gaussian curvature surface. Thus it is also expected that a family of frames taking values in the loop group of $SU_{1,1}$ or SU_2 , that is, $\Lambda SU_{1,1}$ or $\Lambda SU_{2\tau}$, see [A](#page-20-0)ppendix A for the definitions. Unfortunately, the extended frame Ψ^{λ} of a constant Gaussian curvature $K > -1$ surface does not take values in $\Lambda SU_{1,1}$ nor ΛSU_{2r} for any $\lambda \in \mathbb{S}^1$. Thus we consider a deformation of the spectral parameter:

$$
\lambda \in \mathbb{S}^1 \to \lambda \in \mathbb{C}^\times \left(:= \{ c \in \mathbb{C} \mid c \neq 0 \} \right).
$$

Then the extended frame Ψ^{λ} still takes values of $\Lambda SL_2\mathbb{C}_{\tau}$. In Theorem [B](#page-1-1) stated in Introduction, we claim that there exist two special absolute values $|\lambda_0| \neq 1$ of the spectral parameter such that the Lagrangian Gauss maps take values \mathbb{S}^2 or \mathbb{H}^2 at λ_0 . We now prove Theorem [B:](#page-1-1)

Proof of Theorem [B](#page-1-1). We first rephrase the conditions of λ_0 .

 $-1 < K < 0$ and by using a positive $\sigma > 0$ such that $K = -1 + \sigma^2$ holds: For $-1 < K < 0$, that is, $0 < \sigma < 1$, the condition $|\lambda_0| = \exp \left(\operatorname{arcosh} \sqrt{-1/K} \right)$ is equivalent to

(2.17)
$$
|\lambda_0| = \sqrt{\frac{1+\sigma}{1-\sigma}}.
$$

For $K > 0$, that is, $\sigma > 1$, the condition $|\lambda_0| = \exp\left(\arcsin\sqrt{1/K}\right)$ is equivalent to

(2.18)
$$
|\lambda_0| = \sqrt{\frac{1+\sigma}{-1+\sigma}}
$$

Let us represent the Lagrangian Gauss map by

$$
L^{\lambda} = i\Psi^{\lambda} \mathbf{e}_1 (\Psi^{\lambda})^{-1}.
$$

.

It is easy to see that $L^{\lambda}|_{\lambda \in \mathbb{C}^{\times}}$ still takes values in the symmetric space $SL_2\mathbb{C}/GL_1\mathbb{C}$. Then $L^{\lambda}|_{\lambda=\lambda_0\in\mathbb{C}^{\times}}$ takes values in \mathbb{S}^2 (resp. \mathbb{H}^2), up to a conjugation by a constant factor in $SL_2\mathbb{C}$ if and only if $\Psi^{\lambda}|_{\lambda=\lambda_0\in\mathbb{C}^{\times}}$ takes values in SU_2 (resp. $SU_{1,1}$), up to a conjugation by the constant factor in SL₂C. Let $U(\lambda) = U^{\lambda}$ and $V(\lambda) = V^{\lambda}$ be the matrices defined in [\(2.12\)](#page-10-2) and [\(2.13\)](#page-10-3), respectively. It is clear that $\Psi^{\lambda}|_{\lambda=\lambda_0\in\mathbb{C}^{\times}}$ takes values in SU₂ (resp. SU_{1,1}), up to a conjugation by a constant factor in $SL_2\mathbb{C}$, if and only if

$$
-{}^{t}\overline{U(\lambda_0)}=V(\lambda_0), \quad (\text{resp. } -\boldsymbol{e}_1 {}^{t}\overline{U(\lambda_0)}\boldsymbol{e}_1=V(\lambda_0))
$$

holds. A straightforward computation shows that this is equivalent with the equation (2.17) (resp. (2.18)).

In case of \mathbb{S}^2 , λ_0 in [\(2.17\)](#page-13-1) exists if and only the surface has constant Gauss curvature $K = -1 + \sigma^2 > 0$. In case of \mathbb{H}^2 , λ_0 in [\(2.18\)](#page-13-2) exists if and only if the surface has constant Gauss curvature $-1 < K = -1 + \sigma^2 < 0$. Moreover λ_0 is unique up to phase. Since harmonicity of the Lagrangian Gauss map does not depend on the values of $\lambda \in \mathbb{C}^{\times}$, that is, for any $\lambda \in \mathbb{C}^{\times}$, the Lagrangian Gauss map L^{λ} is a harmonic map, thus $L^{\lambda}|_{\lambda=\lambda_0}$ is a harmonic map into \mathbb{S}^2 or \mathbb{H}^2 .

Consider the inner product $\langle dL^{\lambda}, dL^{\lambda} \rangle$. Here \langle , \rangle denotes the Killing metric on \mathfrak{su}_2 or $\mathfrak{su}_{1,1}$:

$$
\langle A, B \rangle_{\mathfrak{su}_2} = -\frac{1}{2} \operatorname{tr} AB \quad \text{or} \quad \langle A, B \rangle_{\mathfrak{su}_{1,1}} = \frac{1}{2} \operatorname{tr} AB,
$$

see also Remark [1.2.](#page-5-1) Then using the form of U^{λ} and V^{λ} , we compute

$$
\partial L^{\lambda} = 2i\lambda^{-1} X \Psi^{\lambda} (-e^{u/2} \hat{\mathbf{e}}_2 + Q e^{-u/2} \hat{\mathbf{e}}_3) (\Psi^{\lambda})^{-1}
$$

$$
\bar{\partial} L^{\lambda} = 2i\lambda Y \Psi^{\lambda} (-\bar{Q}e^{-u/2} \hat{\mathbf{e}}_2 + e^{u/2} \hat{\mathbf{e}}_3) (\Psi^{\lambda})^{-1},
$$

where $X = (1 + \sigma)/2$, $Y = (1 - \sigma)/2$ and \hat{e}_2 , \hat{e}_3 are defined in [\(2.10\)](#page-9-2), [\(2.11\)](#page-9-3) and [\(2.1\)](#page-7-1), respectively. Then in case of \mathbb{H}^2 , that is, in case of $-1 < K < 0$, using the condition in (2.17) , we compute

$$
\langle \partial L^\lambda, \partial L^\lambda\rangle_{\mathfrak{su}_{1,1}}\big|_{\lambda=\lambda_0}=-Ke^{-2i\theta}Q,\quad \langle \partial L^\lambda, \bar{\partial} L^\lambda\rangle_{\mathfrak{su}_{1,1}}\big|_{\lambda=\lambda_0}=-K\frac{e^u+|Q|^2e^{-u}}{2},
$$

and $\langle \bar{\partial} L^{\lambda}, \bar{\partial} L^{\lambda} \rangle_{\mathfrak{su}_{1,1}}|_{\lambda=\lambda_0} = -Ke^{2i\theta} \bar{Q}$. Similary, in case of \mathbb{S}^2 , that is, in case of $K > 0$, using the condition in [\(2.18\)](#page-13-2), we compute

$$
\langle \partial L^\lambda, \partial L^\lambda\rangle_{\mathfrak{su}_2}\big|_{\lambda=\lambda_0}=-Ke^{-2i\theta}Q,\quad \langle \partial L^\lambda, \bar{\partial} L^\lambda\rangle_{\mathfrak{su}_2}\big|_{\lambda=\lambda_0}=K\frac{e^u+|Q|^2e^{-u}}{2},
$$

and $\langle \bar{\partial} L^{\lambda}, \bar{\partial} L^{\lambda} \rangle_{\mathfrak{su}_2}\Big|_{\lambda=\lambda_0} = -Ke^{2i\theta} \bar{Q}$. From these equations it is easy to see that $L^{\lambda}|_{\lambda=\lambda_0}$ is a local diffeomorphism. Moreover if f is a non-totally umbilic constant Gaussian curvature surface with $K > -1$ but $K \neq 0$, then $L^{\lambda}|_{\lambda = \lambda_0}$ is a non-conformal harmonic map. \square

Remark 2.3.

- (1) For a flat $(K = 0, \text{ that is, } \sigma = 1)$ surface in \mathbb{H}^3 , the Lagrangian Gauss map does not take \mathbb{S}^2 nor \mathbb{H}^2 , however, the extended frame Ψ^{λ} is in fact holomorphic with respect to z by a diagonal gauge $D = \text{diag}(e^{u/4}, e^{-u/4})$, that is, $(\Psi^{\lambda}D)^{-1}\overline{\partial}(\Psi^{\lambda}D) = 0$, see [\(2.13\)](#page-10-3). Moreover, the Weierstrass type formula is known for flat surfaces in \mathbb{H}^3 , [\[4\]](#page-22-11). Thus from now on, we will not consider flat surfaces.
- (2) The absolute value $|\lambda_0|$ in Theorem [B](#page-1-1) takes values in $(1,\infty)$. Instead one can choose

$$
|\lambda_0| = \exp\left(-\operatorname{arcosh}\sqrt{-1/K}\right), \quad \text{or} \quad |\lambda_0| = \exp\left(-\operatorname{arsinh}\sqrt{1/K}\right),
$$

in case of $-1 < K < 0$ or in case of $K > 0$, respectively. Then $|\lambda_0|$ takes values $(0, 1)$, and $L^{\lambda}|_{\lambda=\lambda_0}$ is a harmonic local diffeomorphism into \mathbb{H}^2 or a harmonic map into S 2 , respectively.

Conversely let \hat{L} be a non-conformal harmonic map into $G/K = \mathbb{S}^2$ or \mathbb{H}^2 . Then there exists a family of maps $\hat{\Psi}^{\lambda}$ taking values in $\Lambda SU_{2\tau}$ or $\Lambda SU_{1,1\tau}$ such that

$$
(\hat{\Psi}^{\lambda})^{-1} \mathrm{d} \hat{\Psi}^{\lambda} = \hat{\alpha}^{\lambda} = \lambda^{-1} \hat{\alpha}_{\mathfrak{p}}' + \hat{\alpha}_{\mathfrak{k}} + \lambda \hat{\alpha}_{\mathfrak{p}}'' \quad \text{with} \quad \hat{\Psi}^{\lambda}|_{z_{*}} = \mathrm{id}
$$

holds, that is, $\hat{\Psi}^{\lambda}$ is the extended frame of \hat{L} . Using $\hat{\Psi}^{\lambda}$, we can give a family of constant Gaussian curvature surface in the following theorem.

Theorem 2.1. Let \hat{L} be a non-conformal harmonic map into \mathbb{S}^2 or \mathbb{H}^2 and $\hat{\Psi}^{\lambda}$ the corresponding extended frame. Define respective maps

$$
\hat{f}^{\lambda} = \hat{\Psi}^{\lambda} \mathbf{e}_0 (\hat{\Psi}^{\lambda})^* \quad and \quad \hat{n}^{\lambda} = \hat{\Psi}^{\lambda} \mathbf{e}_1 (\hat{\Psi}^{\lambda})^*.
$$

Then the following statements hold:

(1) In case of \mathbb{H}^2 : For any $\lambda \in \mathbb{C}^\times$ such that $0 < |\lambda_1|^{\pm 1} < 1$, $\hat{f}^{\lambda}|_{\lambda = \lambda_1}$ is a surface in \mathbb{H}^3 with constant Gaussian curvature

$$
-1 < K = -\left(\frac{2|\lambda_1|}{|\lambda_1|^2 + 1}\right)^2 < 0.
$$

(2) In case of \mathbb{S}^2 : For any $\lambda \in \mathbb{C}^\times$ such that $0 < |\lambda_1|^{\pm 1} < 1$, $\hat{f}^{\lambda}|_{\lambda = \lambda_1}$ is a surface in \mathbb{H}^3 with constant Gaussian curvature

$$
K = \left(\frac{2|\lambda_1|}{|\lambda_1|^2 - 1}\right)^2 > 0.
$$

Proof. The Maurer-Cartan form $\hat{\alpha}^{\lambda} = (\hat{\Psi}^{\lambda})^{-1} d \hat{\Psi}^{\lambda}$ for the extended frame $\hat{\Psi}^{\lambda}$ of a nonconformal harmonic map \overline{L} has the following form:

$$
\hat{\alpha}^{\lambda} = \lambda^{-1} \hat{\alpha}_{\mathfrak{p}}' + \hat{\alpha}_{\mathfrak{k}} + \lambda \hat{\alpha}_{\mathfrak{p}}'',
$$

where

$$
\alpha_{\mathfrak{k}} = \frac{1}{4} \left(\partial \hat{u} \, \mathrm{d}z - \bar{\partial} \hat{u} \, \mathrm{d}\bar{z} \right) \mathbf{e}_1, \quad \alpha'_{\mathfrak{p}} = \frac{1}{2} \left(e^{\hat{u}/2} \hat{\mathbf{e}}_2 + \hat{Q} e^{-\hat{u}/2} \hat{\mathbf{e}}_3 \right) \mathrm{d}z,
$$

and $\alpha''_{\mathfrak{p}} = \pm^t \overline{\alpha'_{\mathfrak{p}}}$. Here, the plus sign is chosen in case of \mathbb{H}^2 and the minus sign is chosen in case of \mathbb{S}^2 . Moreover, \hat{Q} d z^2 is a holomorphic quadratic differential of the harmonic map \hat{L} defined by $\hat{Q} = \langle \partial \hat{L}, \partial \hat{L} \rangle$ and the $e^{\hat{u}(z)} \rho^{-2}(z)$, where $\rho^2(z) = 4/(1 - |z|^2)^2$, is the ∂ -energy of \hat{L} . Note that relation of the coefficient of \hat{e}_2 and the coefficient of e_1 is given by the Maurer-Cartan equation $(d\hat{\alpha}^{\lambda} + \frac{1}{2})$ $\frac{1}{2} [\hat{\alpha}^{\lambda} \wedge \hat{\alpha}^{\lambda}]$ $|_{\lambda=1} = 0$ and the non-conformality of the harmonic map L .

In case of \mathbb{H}^2 : Choose $0 < |\lambda_1|^{\pm 1} < 1$, and define $e^{u/2}$ and Q by

$$
e^{u/2} = \frac{1 + |\lambda_1|^2}{2|\lambda_1|} e^{\hat{u}/2}
$$
 and $Q = \left(\frac{1 + |\lambda_1|^2}{2|\lambda_1|}\right)^2 \hat{Q}.$

It is easy to see that $\partial \hat{u} = \partial u$ and $\bar{\partial} \hat{u} = \bar{\partial} u$. Then the Maurer-Cartan form $\hat{\alpha}^{\lambda}$ at $\lambda_1 = |\lambda_1|e^{it}, \ (t \in \mathbb{R})$

(2.19)
$$
\hat{\alpha}^{\lambda}|_{\lambda=\lambda_1} = \frac{1}{4} \left(\partial u \, dz - \bar{\partial} u \, d\bar{z} \right) \mathbf{e}_1 + \frac{1+\sigma}{2} e^{-it} \left(e^{u/2} \hat{\mathbf{e}}_2 + Q e^{-u/2} \hat{\mathbf{e}}_3 \right) dz + \frac{1-\sigma}{2} e^{it} \left(\bar{Q} e^{-u/2} \hat{\mathbf{e}}_2 + e^{u/2} \hat{\mathbf{e}}_3 \right) d\bar{z}
$$

with $|\sigma| = |(1 - |\lambda_1|^2)(1 + |\lambda_1|^2)^{-1}| < 1$, that is, $\hat{\Psi}^{\lambda}|_{\lambda = \lambda_1}$ is the extended frame of some constant Gaussian curvature $-1 < K = -1 + \sigma^2 < 0$ surface in \mathbb{H}^3 . Therefore by using Proposition [2.2,](#page-12-1) the map $\hat{f}^{\lambda}|_{\lambda_1}$ gives a surface in \mathbb{H}^3 with constant Gaussian curvature

$$
-1 < K = -1 + \sigma^2 = -\left(\frac{2|\lambda_1|}{|\lambda_1|^2 + 1}\right)^2 < 0.
$$

In case of \mathbb{S}^2 : Choose $0 < |\lambda_1|^{\pm} < 1$, and define $e^{u/2}$ and Q by

$$
e^{u/2} = \frac{1 - |\lambda_1|^2}{2|\lambda_1|} e^{\hat{u}/2}
$$
 and $Q = \left(\frac{1 - |\lambda_1|^2}{2|\lambda_1|}\right)^2 \hat{Q}.$

It is easy to see that $\partial \hat{u} = \partial u$ and $\bar{\partial} \hat{u} = \bar{\partial} u$. Then the Maurer-Cartan form $\hat{\alpha}^{\lambda}$ at $\lambda_1 = |\lambda_1|e^{it}$, $(t \in \mathbb{R})$ becomes [\(2.19\)](#page-16-1) with $|\sigma| = |(1 + |\lambda_1|^2)(1 - |\lambda_1|^2)^{-1}| > 1$, that is, $\hat{\Psi}^{\lambda}|_{\lambda=\lambda_1}$ is the extended frame of some constant Gaussian curvature $K = -1 + \sigma^{-2} > 0$ surface in \mathbb{H}^3 . Therefore by using Proposition [2.2,](#page-12-1) the map $\hat{f}^{\lambda}|_{\lambda_1}$ gives a surface in \mathbb{H}^3 with constant Gaussian curvature

$$
K = -1 + \sigma^2 = \left(\frac{2|\lambda_1|}{|\lambda_1|^2 - 1}\right)^2 > 0.
$$

This completes the proof. \Box

3. Classification of weakly complete constant Gaussian curvature **SURFACES**

It is known that the ∂ -energy density of a harmonic map into \mathbb{H}^2 is $e^u \rho^{-2}$, where $\rho^2 = 4/(1 - |z|^2)^2$ and the metric e^u dzd \overline{z} is a natural metric for the harmonic map, [\[17\]](#page-22-13). Unfortunately, this metric is not directly related to the metric of the constant Gaussian curvature $K > -1$ surface. Thus we consider an another natural metric which is induced from sum of the first and third fundamental forms, the so-called weak metric, that is the pull back of the Sasaki metric on the unit tangent sphere bundle $U\mathbb{H}^3$, and we show that the equivalence of completeness of the weak metric and the metric $e^u dz d\bar{z}$. Then we give a classification of constant Gaussian curvature $-1 < K < 0$ surfaces with the weak metric such that it is complete, the so-called weakly complete metric. Finally, we show the existence of complete and equivariant constant Gaussian curvature $-1 < K < 0$ surfaces, respectively.

3.1. Weakly complete metrics and a classification of constant Gaussian curvature surfaces. Let $f : M \to \mathbb{H}^3$ be a surface with the unit normal $n : M \to \mathbb{S}^{1,2}$. First we define a weakly complete metric of the surface f with $K > -1$.

Definition 3.1. For a surface f with $K > -1$, a metric

(3.1)
$$
\mathrm{d}t^2 = \langle \mathrm{d}f, \mathrm{d}f \rangle + \frac{1}{1+K} \langle \mathrm{d}n, \mathrm{d}n \rangle
$$

is called the *weak metric* and if the weak metric dt^2 is complete, it is called the *weakly* complete metric and the surface f will be called the weakly complete surface.

Remark 3.1.

- (1) The notion of weakly completeness was originally defined for a constant negative Gaussian curvature surfaces in \mathbb{R}^3 , see [\[10\]](#page-22-14).
- (2) For $K = 0$, the weak metric is a pull back of the *Sasaki metric* of the unit tangent sphere bundle $\text{U}\mathbb{H}^3$, $\mathrm{d}t^2 = \langle \mathrm{d}f, \mathrm{d}f \rangle + \langle \mathrm{d}n, \mathrm{d}n \rangle$. Note that the natural metric of the Legendrian immersion $F = (f, n)$ is given by $\langle dF, dF \rangle = \langle df, df \rangle - \langle dn, dn \rangle$, which is indefinite, [\[3\]](#page-22-6).

Lemma 3.1. Let $f : M \to \mathbb{H}^3$ be a constant Gaussian curvature $K > -1$ surface. Moreover let $ds^2 = e^u dz d\bar{z}$ be the metric defined by the function e^u given in [\(1.6\)](#page-4-1). Then the surface f is a weakly complete surface if and only if the metric ds^2 is complete.

Proof. Let $dt^2 = I + \frac{1}{1+K} \mathbb{I}$ be the weak metric of the surface f, where $I = \langle df, df \rangle$ and $\mathbb{I} = \langle d\mathbf{n}, d\mathbf{n} \rangle$ are the first and third fundamental forms, respectively. Then it is easy to see that, by (1.6) and (1.8) , dt^2 can be computed as

$$
\mathrm{d}t^2 = 2(e^u + |Q|^2 e^{-u}) \, \mathrm{d}z \mathrm{d}\bar{z}.
$$

Moreover, since f is a constant Gaussian curvature, if necessary by choosing the unit normal as $-n$, we have $e^{2u} > |Q|^2$, see [\(1.6\)](#page-4-1). If e^u dzd \overline{z} is complete, then it is easy to see that dt^2 is complete, that is, the surface is a weakly complete surface. Conversely if dt^2 is complete, then $\overline{}$ $\mathcal{C}_{0}^{(n)}$ $\sqrt{2(e^u+|Q|^2e^{-u})}$ dz| needs to diverge along any divergent curve C on M, that is, either

$$
\sqrt{2} \int_C e^{u/2} |\mathrm{d}z| \quad \text{or} \quad \sqrt{2} \int_C e^{-u/2} |Q| |\mathrm{d}z|
$$

needs to diverge. The latter case also implies that $\mathcal{C}_{0}^{(n)}$ $e^{u/2}$ |dz| diverges for any divergent curve C on M, since $e^u > |Q|$. This completes the proof.

As stated Theorem [C](#page-2-0) in Introduction, we classify weakly complete constant negative Gaussian curvature $-1 < K < 0$ surfaces.

Proof of Theorem [C](#page-2-0). From Theorem [A,](#page-1-0) for a constant negative Gaussian curvature surface $-1 < K < 0$ in \mathbb{H}^3 , the Klotz differential is holomorphic. Since the equivalent class of the surface, that is a rigid motion, does not change the Klotz differential, thus we can define a map g from \mathcal{K}_K into \mathcal{QD} .

Let us consider the Gauss-Codazzi equations

(3.2)
$$
\bar{\partial}\partial u + \frac{K}{2}(e^u - |Q|^2 e^{-u}) = 0, \quad \bar{\partial}Q = 0
$$

for a constant Gaussian curvature $-1 < K < 0$ surface f. Note that the second equation is nothing but holomorphicity equation of Q. It is well known that, using a conformal change of coordinates $z \to \sqrt{-K}z$, the first equation is the elliptic PDE for harmonic maps from $\mathbb D$ or $\mathbb C$ into $\mathbb H^2$, see [\[17,](#page-22-13) [18\]](#page-22-15). It is also well known that for a given quadratic differential $Q dz^2$ on $\mathbb D$ or $\mathbb C$ which is not identically zero on $\mathbb C$, there exists a unique solution u of the first equation in [\(3.2\)](#page-18-0) such that e^u dzd \bar{z} is complete. Moreover, it has been shown in [\[15,](#page-22-7) Theorem 3.2 that if the holomorphic quadratic differential $Q dz^2$ is not constant on \mathbb{C} , then the corresponding harmonic map is global diffeomorphism and satisfies the inequality

$$
e^{2u} > |Q|^2,
$$

see [\[15,](#page-22-7) Lemma 1.3 (2)]. Therefore by fundamental theorem of surface theory there exists a unique constant Gaussian curvature $-1 < K = -1 + \sigma^2 < 0$ surface f, up to rigid motion, such that the holomorphic Klotz differential is $Q dz^2$ and $I^{(1,1)} + \sigma^{-1}I^{(1,1)} = 2e^u dz d\bar{z}$, where the superscript $(1, 1)$ denotes the $(1, 1)$ -part of the fundamental forms. Note that since $\det \tilde{I} = (e^u - |Q|^2 e^{-u})^2 > 0$, where \tilde{I} is the coefficient matrix of the first fundamental form I, f never degenerate. Moreover, from Lemma [3.1,](#page-17-0) the resulting surface f is a weakly complete surface. Note that since the equivalent class of the holomorphic quadratic differential is given by a Möbius transformation on $\mathbb D$ or $\mathbb C$, it corresponds to change of conformal parameters of the corresponding constant Gaussian curvature surface. Therefore, $g: \mathcal{K}_K \to \mathcal{QD}$ is bijective. This completes the proof.

Remark 3.2.

- (1) In general, weakly complete constant Gaussian curvature $-1 < K < 0$ surfaces would have non-trivial topology. The most simplest example is an equidistant cylinder which have the non-zero constant Klotz differential. We will discuss further examples in a forthcoming paper.
- (2) If the holomorphic quadratic differential is non-zero constant on \mathbb{C} , then the corresponding harmonic map is not diffeomorphism, and in fact $e^{2u} = |Q|^2 = c \neq 0$ holds and the corresponding surface becomes degenerate, that is, a geodesic in \mathbb{H}^3 .
- (3) Theorem [C](#page-2-0) does not hold for weakly complete constant Gaussian curvature $K > 0$ surfaces in \mathbb{H}^3 , since there are infinitely many such surfaces for a given holomorphic quadratic differential $Q \, dz^2$. Moreover, the resulting surface could be degenerate at some point, that is, $e^{2u} = |Q|^2$ occurs.

3.2. Complete constant Gaussian curvature surfaces. As corollaries of Theorem [C](#page-2-0) we show the existence of complete and equivariant constant Gaussian curvature $-1 < K <$ 0 surfaces in \mathbb{H}^3 , respectively.

Corollary 3.1. Let $Q dz^2$ be a bounded holomorphic quadratic differential on the unit disk D with respect to the Poincare metric $ds_p^2 = \rho^2(z) dz d\overline{z}$ with $\rho^2(z) = 4(1 - |z|^2)^{-2}$, that is,

$$
\sup_{z \in \mathbb{D}} |Q(z)| \rho^{-2}(z) < \infty.
$$

Then there exists a unique complete constant Gaussian curvature $-1 < K < 0$ surface in \mathbb{H}^3 whose Klotz differential is $Q \, dz^2$.

Proof. We need to prove completeness of the constant Gaussian curvature surface f with the bounded Klotz differential $Q dz^2$. The first fundamental form I of the surface f constructed in the proof of Theorem [C](#page-2-0) is given as

$$
I = Q dz2 + (eu + |Q|2 e-u) dz d\overline{z} + \overline{Q} d\overline{z}2.
$$

Then it can be rephrased as

$$
\mathcal{I} = \frac{1}{|K|} e^u \left| d\hat{z} + \mu d\bar{\hat{z}} \right|^2, \quad \mu = \frac{\bar{Q}}{e^u}, \quad \hat{z} = \sqrt{-K}z.
$$

Note that the coordinates \hat{z} normalizes the Gauss equation in [\(2.14\)](#page-11-1) as the standard harmonic map setting, and for simplicity, we write \hat{z} by z. Since $Q dz^2$ is bounded, by [\[15,](#page-22-7) Theorem 2.1, there exists a quasi-conformal diffeomorphism ζ on $\mathbb D$ which satisfies the Beltrami equation $\partial \zeta = \mu \partial \zeta$. Here μ can be rephrased as

$$
\mu = \frac{\bar{Q}}{e^u} = \frac{\bar{Q}\rho^{-2}}{e^u \rho^{-2}}, \quad \rho^2(z) = \frac{4}{(1 - |z|^2)^2},
$$

and $e^u \rho^{-2}$ is the ∂-energy of the deformed Lagrangian Gauss map $L^{\lambda}|_{\lambda=\lambda_0}$ into \mathbb{H}^2 . More-over by [\[15,](#page-22-7) Lemma 2.4] the solution ζ is a harmonic diffeomorphism onto $(\mathbb{D}, \rho^2 |d\zeta|^2)$ and the ∂ -energy of ζ is again $e^u \rho^{-2}$. Therefore the first fundamental I can be rephrased in the coordinates ζ as

(3.3)
$$
I = \frac{1}{|K|} e^u |dz + \mu dz|^2 = \frac{1}{|K|} \frac{e^u}{|\partial \zeta|^2} |d\zeta|^2 = \frac{1}{|K|} \frac{e^u \rho^{-2}}{|\partial \zeta|^2} \rho^2 |d\zeta|^2 = \frac{1}{|K|} \rho^2 |d\zeta|^2.
$$

Thus I defines a complete metric. This completes the proof.

From Gauss-Bonnet theorem, it is easy to see that there does not exist a constant Gaussian curvature $-1 < K < 0$ surface in \mathbb{H}^3 of genus zero. Therefore in the following corollary, we omitt the genus zero case.

Corollary 3.2. Let $Q \, dz^2$ be a holomorphic quadratic differential defined on a Riemann surface $M = \widetilde{M}/\Gamma$, where \widetilde{M} is the unit disk $\mathbb D$ or $\mathbb C$ and Γ is a Fuchsian group, respectively. Moreover assume that in case of $\widetilde{M} = \mathbb{C}$, $Q \, dz^2$ is not constant. Then there exists a unique weakly complete constant Gaussian curvature $-1 < K < 0$ surface $f : \widetilde{M} \to \mathbb{H}^3$ satisfying the following equivariant property: There exist a representation ρ from Γ to the isometry group Iso \mathbb{H}^3 such that for any $\gamma \in \Gamma$ the following relation holds:

$$
f\circ\gamma=\rho(\gamma)\circ f.
$$

Proof. From Theorem [C,](#page-2-0) for a given holomorphic quadratic differential $Q dz^2$ on D or \mathbb{C} , there exists a unique weakly complete constant Gaussian curvature $-1 < K = -1 + \sigma^2 < 0$ surface f from D or C into \mathbb{H}^3 such that the Klotz differential of f is Q d z^2 . Moreover since $Q dz^2$ is invariant with respect to Γ, that is, for any $\gamma \in \Gamma$, $Q \circ \gamma = Q$ holds and the uniqueness of the surface f, there exists a representation $\rho : \Gamma \to \text{Iso } \mathbb{H}^3$ such that for any $\gamma \in \Gamma$

$$
f \circ \gamma = \rho(\gamma) \circ f
$$

holds. This completes the proof. \Box

Remark 3.3.

- (1) In [\[8,](#page-22-4) Thoerem D], a similar statement to Corollary [3.2](#page-19-0) has been proved using completely different idea, the so-called "asymptotic Plateau problem", that is, the existence of a constant Gaussian curvature $-1 < K < 0$ surface which are asymptotically to a given curve γ in the ideal boundary of \mathbb{H}^3 .
- (2) To show the existence of weakly complete constant Gaussian curvature $-1 < K < 0$ surfaces f which have nontrivial topology, we need

$$
\rho(\gamma) = \mathrm{id}
$$

for some $\gamma \in \Gamma$. It is not easy to see this condition holds for a given holomorphic quadratic differential.

Appendix A. Loop groups

Let $SL_2\mathbb{C}$ be the special linear group of degree 2 and its Lie algebra $\mathfrak{sl}_2\mathbb{C}$. The twisted loop group of $SL_2\mathbb{C}$ is a set of smooth maps

$$
\Lambda SL_2 \mathbb{C}_\tau = \{ g : \mathbb{S}^1 \to SL_2 \mathbb{C} \mid \tau g(\lambda) = g(-\lambda) \},
$$

where τ is defined by $\tau g = \text{Ad}(e_1)g$. It is known that $\Lambda SL_2\mathbb{C}_\tau$ is a complex Banach Lie group with respect to a suitable topology, see [\[12\]](#page-22-16). The Lie algebra of $\Lambda SL_2\mathbb{C}_7$ is

$$
\Lambda \mathfrak{sl}_2 \mathbb{C}_\tau = \{ g : \mathbb{S}^1 \to \mathfrak{sl}_2 \mathbb{C} \mid \tau g(\lambda) = g(-\lambda) \},
$$

and it is called the *twisted loop algebra* of $\text{ASL}_2\mathbb{C}_\tau$. We then define two subgroups $\text{ASU}_{2\tau}$ and $\Lambda SU_{1.1\tau}$ as

$$
\begin{aligned} \Lambda \text{SU}_{2\tau} &= \{ g \in \Lambda \text{SL}_2\mathbb{C}_\tau \mid {}^t \overline{g(1/\bar{\lambda})}^{-1} = g(\lambda) \}, \\ \Lambda \text{SU}_{1,1\tau} &= \{ g \in \Lambda \text{SL}_2\mathbb{C}_\tau \mid \mathbf{e}_1 {}^t \overline{g(1/\bar{\lambda})}^{-1} \mathbf{e}_1 = g(\lambda) \}, \end{aligned}
$$

and they are called the *twisted loop groups* of SU_2 and $SU_{1,1}$, respectively. It is easy to see that these subgroups are real forms of the complex Banach Lie group $\Lambda SL_2\mathbb{C}_\tau$, that is, they are given by fixed point sets of anti-linear involutions on $\text{ASL}_2\mathbb{C}_\tau$.

22 J. INOGUCHI AND S.-P. KOBAYASHI

Appendix B. Harmonic maps into symmetric spaces

B.1. We recall that harmonicity from a Riemann surface M into a homogeneous semi-Riemannian space $N = G/H$. For more detail, see for examples [\[2,](#page-22-17) [3\]](#page-22-6). Let $\varphi : M \to G/H$. be a smooth map of a simply connected Riemann surface into a reductive homogeneous semi-Riemannian space $N = G/H$ equipped with a reductive decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$. Then there exits a smooth map $\Phi : M \to G$ such that $\Phi H = \varphi$. Such a map Φ is called a frame of φ . The Lie algebra g-valued 1-form $\alpha = \Phi^{-1} d\Phi$ is called the *Maurer-Cartan form* of Φ. The Maurer-Cartan form α defines a flat connection $d + \alpha$ on the principal bundle $M \times G$. The flatness is expressed as the following zero curvature equation:

$$
d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0.
$$

We decompose the Maurer-Cartan form α by

$$
\alpha = \alpha_{\mathfrak{k}} + \alpha_{\mathfrak{p}}, \quad \alpha_{\mathfrak{p}} = \alpha_{\mathfrak{p}}' + \alpha_{\mathfrak{p}}''
$$

according to the reductive decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ and \prime and $\prime\prime$ denotes $(1, 0)$ - and $(0, 1)$ parts on M, respectively.

The harmonicity of a map φ can be characterized by the Maure-Cartan form α as

(B.1)
$$
[\alpha'_{\mathfrak{p}} \wedge \alpha''_{\mathfrak{p}}]_{\mathfrak{p}} = 0 = d(*\alpha_{\mathfrak{p}}) + [\alpha \wedge * \alpha_{\mathfrak{p}}],
$$

where the subscript $\mathfrak h$ and $\mathfrak p$ denote the $\mathfrak h$ - and $\mathfrak p$ -part, respectively. Moreover $*$ denotes the Hodge star operator, for example $*dz = -i dz$ and $*d\bar{z} = i d\bar{z}$. In particular if the target homogeneous space G/H is a semi-Riemannian symmetric space, then the first condition is vacuous.

B.2. It is known that harmonic maps from a Riemann surface into a semi-Riemannian symmetric space can be characterized by loop groups, see for an example [\[2\]](#page-22-17). In this paper we consider two particular symmetric spaces, that is, the 2-sphere \mathbb{S}^2 and the hyperbolic 2-space \mathbb{H}^2 . Thus we briefly explain how harmonic maps into these symmetric spaces are characterized.

Let M be a simply connected Riemann surface and G/K be a Riemannian symmetric space $\mathbb{S}^2 = \mathrm{SU}_2/\mathrm{U}_1$ or $\mathbb{H}^2 = \mathrm{SU}_{1,1}/\mathrm{U}_1$. Moreover, let $\varphi : M \to G/K$ be a smooth map and Φ a framing of φ . Consider the Maurer-Cartan form $\alpha = \Phi^{-1} d\Phi = \alpha_{\mathfrak{k}} + \alpha'_{\mathfrak{p}} + \alpha''_{\mathfrak{p}}$ and define a family of Maurer-Cartan forms α^{λ} as

$$
\alpha^{\lambda} = \lambda^{-1} \alpha'_{\mathfrak{p}} + \alpha_{\mathfrak{k}} + \lambda \alpha''_{\mathfrak{p}}, \quad \lambda \in \mathbb{C}^{\times}.
$$

From the construction, it is clear to see that $d + \alpha^{\lambda}|_{\lambda=1}$ gives the original flat connection $d + \alpha$. Then harmonicity of the map φ is characterized as follows.

Theorem B.1 ([\[2\]](#page-22-17)). A smooth map $\varphi : M \to G/K$ of a simply connected Riemann surface into a semi-Riemannian symmetric space G/K is harmonic if and only if $d + \alpha^{\lambda}$ is flat connection for all λ , that is, the partial differential equation

$$
d\alpha^{\lambda} + \frac{1}{2} [\alpha^{\lambda} \wedge \alpha^{\lambda}] = 0
$$

is satisfied for all $\lambda \in \mathbb{C}^{\times}$.

REFERENCES

- [1] D. Brander, J. Inoguchi, S.-P. Kobayashi, Constant Gaussian curvature surfaces in the 3-sphere via loop groups, Pacific J. Math. 269(2014), no. 2, 281–303.
- [2] F. E. Burstall, F. Pedit, Harmonic maps via Adler-Kostant-Symes theory, in Harmonic maps and integrable systems, 221–272, Aspects Math., E23, Friedr. Vieweg, Braunschweig, 1994.
- [3] J. F. Dorfmeister, J. Inoguchi, S.-P. Kobayashi, Constant mean curvature surfaces in hyperbolic 3 space via loop group, J. Reine Angew. Math. $686(2014)$, 1–36, [arXiv:1108.1641](http://arxiv.org/abs/1108.1641) [math DG].
- [4] J. A. Gálvez, A. Martinez, F. Milan, Flat surfaces in the hyperbolic 3-space, *Math. Ann.* $316(2000)$, 419–435.
- [5] Y. Kitagawa, Periodicity of the asymptotic curves on flat tori in S^3 . J. Math. Soc. Japan 40(1988), no. 3, 457–476.
- [6] T. Klotz, Some uses of the second conformal structure on strictly convex surfaces, Proc. Amer. Math. Soc. **14**(1963), 793–799.
- [7] S.-P. Kobayashi, Real forms of complex surfaces of constant mean curvature. Trans. Amer. Math. Soc. **363**(2011), no. 4, 1765–1788.
- [8] F. Labourie, Un lemme de Morse pour les surfaces convexes, Invent. Math. 141(2000), 239–297.
- [9] C. R. LeBrun, Twistor CR manifolds and three-dimensional conformal geometry, Trans. Amer. Math. Soc. 284(1984), 601–616
- [10] M. Melko, I. Sterling, Application of soliton theory to the construction of pseudospherical surfaces in \mathbb{R}^3 , Ann. Global Anal. Geom. 11(1993), no. 1, 65–107.
- [11] K. Nomizu, Isometric immersions of the hyperbolic plane into the hyperbolic space, Math. Ann. 205(1973), 181–192.
- [12] A. Pressley and G. Segal, Loop groups, Oxford Mathematical Monographs. Oxford University Press, New York, 1986.
- [13] H. Rosenberg, H. Spruck, On the existence of convex hypersurfaces of constant Gauss curvature in hyperbolic space, *J. Differential Geom.* **40**(1994), 379–409.
- [14] M. Spivak, A comprehensive Introduction to Differential Geometry IV, 2nd ed., Publish or Perish, Wilmington, Del., 1979.
- [15] L-F. Tam, T. Y-H Wan, Harmonic diffeomorphisms into Cartan-Hadamard surfaces with prescribed Hopf differentials. Comm. Anal. Geom., 2(1994), no. 4, 593–625.
- [16] K. Tenenblat, Transformations of manifolds and applications to differential equations, Longman, 1998.
- [17] T. Y-H Wan, Constant mean curvature surface, harmonic maps, and universal Teichmüller space. J. Differential Geom. **35**(1992), no. 3, 643–657.
- [18] T. Y-H Wan, T. K-K. Au, Parabolic constant mean curvature spacelike surfaces. Proc. Amer. Math. Soc. 120(1994), no. 2, 559–564.

Department of Mathematics, Hokkaido University, Sapporo, 060-0810, Japan Email address: inoguchi@math.sci.hokudai.ac.jp

Department of Mathematics, Hokkaido University, Sapporo, 060-0810, Japan Email address: shimpei@math.sci.hokudai.ac.jp