

Open Cluster Dynamics under the Influence of Outflow-Ambient Interactions

MUXIN LIU ^{1,2} LILE WANG ^{2,1} XIAOTING FU ³ AND LUIS C. HO ^{2,1}

¹*Department of Astronomy, School of Physics, Peking University, Beijing 100871, China*

²*Kavli Institute for Astronomy and Astrophysics, Peking University, Beijing 100871, China*

³*Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210023, China*

ABSTRACT

Outflowing stars impinging upon ambient gas experience accelerations due to the gravitational feedback from the interaction morphology between the outflow and the ambient gas. Such “negative dynamical friction” (NDF), in contrast to the conventional “dynamical friction” (DF), is studied for its impact on the dynamics of open clusters (OCs) immersed in a uniform ambient gas. We modify the N -body integration code REBOUND with both NDF and DF implemented according to the outflow conditions of each star in a consistently constructed OC. The evolution of stars is also involved in determining the gas-star interactions throughout their stellar lives. Compared to DF-only and gas-free models with identical initial conditions, the NDF-affected cluster is puffier and evaporates faster, as indicated by various diagnostics, including lower velocity dispersions and larger half-mass and half-light radii. Neutron stars with fast winds are expelled from the cluster due to their intensive NDF effect, even without the “kicks” by asymmetric supernovae. Exploration of parameter space confirms that the NDF effect is generally enhanced with higher ambient gas densities, in qualitative agreement with the expression of acceleration. Outflow-ambient interactions should be considered for the proper interpretation of the stellar dynamics evolution in clusters.

Keywords: Open star clusters (1160), Stellar dynamics (1596), Stellar winds (1636), Stellar mass loss (1613), Dynamical friction (422), Neutron stars (1108), N-body simulations (1083)

1. INTRODUCTION

Open clusters (OCs) offer valuable insights into the formation and evolution of stars. Comprised of approximately 10^2 to 10^4 stars that are gravitationally bound to each other (Binney & Tremaine 2008), OCs comprise well-defined single stellar populations. They form from a common molecular cloud collapse (Krumholz et al. 2019), with the same metallicity and age. As a result, OCs serve as valuable tracers of star formation, chemical evolution, kinematics, and gravitational dynamics (see, e.g., Friel 1995; Krumholz et al. 2019; Cantat-Gaudin 2022; Fu et al. 2022; Magrini et al. 2023). For instance, they can be utilized to determine the metallicity of stellar populations in different regions of the Milky Way, as their distances can be constrained by their color-magnitude diagrams (see, e.g., Sandage 1953). More-

over, the kinematic properties of OCs can be used to trace the rotation curve of the Milky Way (see, e.g., Tarricq et al. 2021). Thanks to the contributions of missions like Gaia (Gaia Collaboration et al. 2016) in providing stellar astrometry information, including precise position, parallax, and proper motion, significant advancements have been made (e.g., Gaia Collaboration et al. 2018, 2021, 2023). The member stars of OCs and the properties of several thousand clusters have been determined based on Gaia data (see, e.g., Cantat-Gaudin 2022; Hunt & Reffert 2023). Some OCs show tidal structures and indicate interactions with their environments (see, e.g., Pang et al. 2022; Tarricq et al. 2022). These advancements enable the dynamical characteristics study of OCs through observations and facilitate comparison with theoretical models.

Most Galactic OCs travel in the Milky Way thin disk on near-circular orbits that pass through the Galactic mid-plane several times in one orbital revolution (see, e.g., Fu et al. 2022). The impact of the detailed interactions between the cluster member stars and the gas

clouds in the Galactic plane remains unclear. Do the interactions change the morphology of the star clusters and lead to the loss of member stars? Do the interactions differ among different kinds of stars? These are still open questions.

The ambient gas density likely will affect the dynamical characteristics of an OC as it travels through a gas cloud. A massive object moving through a giant gas cloud experiences dynamic friction (DF) (Chandrasekhar 1943; Ostriker 1999; Edgar 2004). According to the standard Bondi-Hoyle-Lyttleton accretion model (Hoyle & Lyttleton 1939; Bondi & Hoyle 1944; Edgar 2004), it is understood that due to the long-range nature of gravity, the gas outside the Bondi radius is also influenced by gravity, resulting in the formation of an overdense tail downstream of the massive object. This overdense region exerts a pulling force on the object in the opposite direction of its motion, causing it to decelerate. This force is known as DF force. Works such as Kim & Kim (2007) and Baruteau et al. (2011) have shown that gas DF can lead to the hardening of binaries, and Tagawa et al. (2020) have pointed out that binaries can form through single-single interactions by dissipating kinetic energy in a gaseous medium. Other works, such as Tanaka et al. (2002), have shown that DF force can dampen the velocity dispersion of black holes (BHs) and stars, and Just et al. (2012) have demonstrated that this dissipative force acting on stars in the disk can result in an increased mass flow toward the supermassive BH and an asymmetry in the phase space distribution due to a rotating accretion disk.

However, the situation changes if stellar objects launch powerful winds. Gruzinov et al. (2020) demonstrated that when the wind speed is sufficiently high, the extent of the underdense region becomes substantial, resulting in an overall gravitational force from the gas that aligns with the object’s motion. Consequently, the DF becomes negative in this scenario, causing the object to accelerate. This phenomenon is known as negative dynamic friction (NDF). In other related works, Li et al. (2020) employed hydrodynamic simulations to explore the changes in accretion rate and the strength of NDF in the presence of outflows from compact objects. Additionally, Wang & Li (2022) conducted a study on NDF in the context of a binary system using global 3D hydrodynamic simulations. This work uses the N -body integrator REBOUND (Rein & Liu 2012) to study the NDF impact on the dynamics of OCs immersed in a uniform ambient gas. Thanks to the support of REBOUND in allowing the incorporation of additional physics, both NDF and DF are implemented according to the outflow conditions of each star in a consistently constructed OC.

This paper is structured as follows. Section 2 provides comprehensive descriptions of the gas-star interactions incorporated, the stellar evolution model, and the setup of the fiducial model. Section 3 analyzes and compares the impacts of different types of gas-star interactions—namely, “DF”, “NDF”, and “None”—on the dynamics and stellar evolution of the fiducial model OC, both with and without bulk motion. Section 4 explores scenarios with various ambient gas densities. Discussion and a summary are given in Section 5.

2. METHODS

This work simulates the dynamics of gas-coupled OCs using N -body simulations by adding modules to REBOUND. For accuracy, we adopt the IAS15 integrator (Rein & Spiegel 2015), which is a 15th order scheme with adaptive step size control. In order to model the clusters properly, stellar particles are created with proper mass distributions and evolved with stellar evolution models that include compact objects epochs along with the dynamics.

2.1. Gas-star Interactions

The interaction between gas and stars can generate either friction or anti-friction, depending on the strength of the stellar outflows. For stars that have no outflows, the analytic approximations for DF are adopted from Ostriker (1999):

$$a_{\text{DF}} = \frac{4\pi G^2 \rho}{v_*^2} \times \begin{cases} \ln \left[\Lambda \left(1 - \frac{1}{\mathcal{M}^2} \right)^{1/2} \right], & \mathcal{M} > 1 \\ \frac{1}{2} \ln \left(\frac{1 + \mathcal{M}}{1 - \mathcal{M}} \right) - \mathcal{M}, & \mathcal{M} < 1, \end{cases} \quad (1)$$

where v_* is the stellar velocity, ρ is the ambient gas density, $\mathcal{M} \equiv v_*/c_s$ is the Mach number in the gas, and $\Lambda \equiv b_{\text{max}}/b_{\text{min}}$ is the Coulomb factor. In this work, we assume that the a_{DF} direction is always aligned with the relative motion direction, because the timescales of hydrodynamic phenomena τ_{hyd} are typically much shorter than the stellar orbital dynamic timescales τ_{dyn} ,

$$\begin{aligned} \tau_{\text{hyd}} &\sim 0.02 \text{ yr} \times \left(\frac{l_{\text{hyd}}}{R_{\odot}} \right) \left(\frac{v_*}{1 \text{ km s}^{-1}} \right)^{-1}, \\ \tau_{\text{dyn}} &\sim 10^5 \text{ yr} \times \left(\frac{l_{\text{dyn}}}{0.1 \text{ pc}} \right) \left(\frac{v_*}{1 \text{ km s}^{-1}} \right)^{-1}, \end{aligned} \quad (2)$$

in which l_{hyd} and l_{dyn} are the typical spatial scale of the hydrodynamic processes and the stellar dynamics, respectively. The overdense structures causing DF is typically a few times the stellar size in absence of outflows (see e.g., Ostriker 1999), and the typical orbital scales of stars in OCs are no less than $\sim 10^{-1}$ pc (see e.g.,

Nilakshi et al. 2002) (unless in close encounter events, which are typically rapid and dwarf DF effects by the ambient gas). Therefore, it is almost always eligible to assume that the star-gas interaction structures settle at the steady or quasi-steady states, and the direction of DF stays along the stellar motion relative to the gas. We use the Bondi radius of each star, $b_{\min} \equiv 2GM/c_s^2$, for the minimum impact parameter, and we approximate b_{\max} with the size upper limit of a giant molecular cloud (~ 100 pc in this work; e.g., Solomon et al. 1987; Miville-Deschênes et al. 2017; Sun et al. 2018).

Whenever a star has outflows, the gravitational feedback coming from the interactions between such outflows and the ambient gas will likely accelerate the star. We adopt the analytic approximation described in Gruzinov et al. (2020) and Li et al. (2020),

$$a_{\text{NDF}} = \pi G \rho \int_0^\pi d\theta \cos\theta \sin\theta R_s \times \left\{ \frac{3}{2} \left[1 + \frac{2u(1 - \cos\theta)}{R_s^2 \sin^2\theta / R_0^2} \right]^2 - 2 \left[1 + \frac{u^2}{R_s^2 / R_0^2} \right] \right\}, \quad (3)$$

where $R_0 \equiv [\dot{m}_w v_w / (4\pi \rho v_*^2)]^{1/2}$ is the standoff distance (where the total pressure of the incoming medium equals to that of the outflow), \dot{m}_w is the outflow wind mass-loss rate, v_w is the wind radial velocity, $R_s = R_0 [3(1 - \theta \cot\theta) / \sin^2\theta]^{1/2}$ is the contact discontinuity location at the polar angle θ (where the gas over-densities are located), and $u \equiv v_* / v_w$. In general, for typical OCs, v_* spans $0.5 - 5$ km s $^{-1}$ (e.g., Tarricq et al. 2021; Soubiran et al. 2018; Mathieu 1984), which is significantly lower than the v_w of almost all stellar outflows. Even asymptotic giant branch (AGB) stars, which are known for their slow outflows, still have $v_w \gtrsim 30$ km s $^{-1}$ (Habing & Olofsson 2004; Ramstedt et al. 2008). Therefore, one can take the $u \rightarrow 0$ limit of Eq. (3) and adopt a simple analytic approximation, $a_{\text{NDF}} \approx 8.18 G \rho R_0$ (see also Li et al. 2020).

Similar to the DF case, the NDF direction stays along the direction of star to gas relative motion. While the stellar dynamic timescales are estimated similar to Eq. (2), the estimation of hydrodynamic timescales approximately reads,

$$\tau_{\text{hyd}} \sim 5 \text{ yr} \times \left(\frac{l_{\text{hyd}}}{10^2 \text{ AU}} \right) \left(\frac{v_w}{10^2 \text{ km s}^{-1}} \right)^{-1}, \quad (4)$$

where the spatial scale l_{hyd} should stand for the distance from the star to the bow shock, and v_w is the stellar wind velocity. As one can easily observe, the inequality $\tau_{\text{hyd}} \ll \tau_{\text{dyn}}$ still holds in the NDF cases.

There are multiple types of stars in an OC, and their interactions with the ambient gas are different. Unless

otherwise noted, we use the anti-friction recipes for main sequence stars, red giant branch (RGB) stars, and neutron stars (NSs), while we assume that white dwarfs (WDs) and BHs have *no* outflows and thus obey a_{DF} in Eq. (1). Admittedly, WDs and BHs in binaries can accrete and have consequent activities (e.g., disk winds, decretions, and jets), yet detailed discussions on those phenomena only add to the complications and obscure the effects concerned, and are beyond the scope of the current paper. For main sequence stars and giants, the wind properties are calibrated for stars with solar metallicity. The mass-loss rates are approximated by Reimers formula (e.g., Kudritzki & Reimers 1978; see also Choi et al. 2016),

$$\dot{m}_w \approx 4 \times 10^{-13} M_\odot \text{ yr}^{-1} \times \left(\frac{L}{L_\odot} \right) \left(\frac{R}{R_\odot} \right) \left(\frac{M}{M_\odot} \right)^{-1} \\ \propto 4 \times 10^{-13} M_\odot \text{ yr}^{-1} \times \left(\frac{M}{M_\odot} \right)^{3.3}, \quad (5)$$

where we use the approximate scaling laws $L \propto M^{3.5}$ (Kuiper 1938) and $R \propto M^{0.8}$ (Demircan & Kahraman 1991). Such mass-loss rates are on the high end of typical values, yet stars in an OC are generally young and tend to have stronger stellar winds. The stellar wind velocity has significant variations and uncertainties from star to star. We adopt a simplified power-law fitting for main sequence stars,

$$v_w \approx 400 \text{ km s}^{-1} \times \left(\frac{M}{M_\odot} \right)^{0.6}, \quad (6)$$

which yields observed values of $v_w = 400$ km s $^{-1}$ for solar-mass stars (e.g., Brooks et al. 2015), and $v_w \approx 2000$ km s $^{-1}$ for type O8 stars (e.g., Bernabeu et al. 1989). The uncertainties of wind properties are even greater for RGB stars; for simplicity, we assume all of them have $v_{w,\text{RGB}} \approx 30$ km s $^{-1}$ and $\dot{m}_{w,\text{RGB}} \approx 3 \times 10^{-9} M_\odot \text{ yr}^{-1}$ (Habing & Olofsson 2004; Ramstedt et al. 2008). NSs can have significant outflows due to complex acceleration mechanisms (such as winds originating from young and intensely hot neutron stars are propelled by the absorption of high-energy neutrinos by photons and neutrons near the stellar surface (Salpeter & Shapiro 1981; Duncan et al. 1986)), and we assume $v_w \approx 3 \times 10^4$ km s $^{-1} = 0.1c$ (c for the speed of light), and $\dot{m}_w \approx 3 \times 10^{-9} M_\odot \text{ yr}^{-1}$ (e.g., Kato 1983; Paczynski 1990; Mor et al. 2023).

2.2. Stellar Evolution Properties

Stars in an OC evolve for hundreds of millions of years before the cluster evaporates (Portegies Zwart et al.

2010; Krumholz et al. 2019; Krause et al. 2020). Given that the main sequence lifetime of an $8 M_{\odot}$ star with solar metallicity is about 35 Myr (see stellar model PARSEC v1.2s¹; Bressan et al. 2012; Tang et al. 2014; Chen et al. 2015), before the cluster evaporates all massive stars will have evolved off the main sequence. Very massive stars ($M \gtrsim 25 M_{\odot}$) are not considered in this work. We assume that each star has finished its main sequence phase and entered its RGB stage at the time we consider significant gas-star interactions to occur. The evolution time of the RGB phase is approximated by a piecewise function in line with the solar metallicity and solar composition stellar models ($Z=0.017$, $Y=0.279$) of PARSEC v1.2s:

$$t_{\text{RGB}} = \begin{cases} 10^5 \text{ yr} , & M > 15 M_{\odot} \\ 3 \times 10^8 \times \left(\frac{M}{M_{\odot}}\right)^{-4} \text{ yr} , & M \leq 15 M_{\odot} . \end{cases} \quad (7)$$

After the RGB stage, we assume that stars of $M < 8 M_{\odot}$ will undergo their AGB phase and end up as WDs. Because the AGB stage experiences massive outflows that lose $\sim 50\%$ of the stellar mass within a relatively short period ($\Delta t \lesssim 10^5$ yr), the impact of AGB outflows due to anti-friction can be accounted as a pulse momentum injection, by integrating $dv/dt = a_{\text{NDF}}$ assuming $u \ll 1$ and $\dot{m}_w \approx \Delta M/\Delta t$. The increments regarding the velocity is then better estimated by the change in v^2 ,

$$\begin{aligned} v \, dv &= a_{\text{NDF}} v \, dt ; \\ \Delta v^2 &\approx 2 \int_0^{\Delta t} a_{\text{NDF}} v \, dt \\ &\approx 3 \times 10^{-2} \text{ km}^2 \text{ s}^{-2} \times \left(\frac{\rho}{30 m_p \text{ cm}^{-3}}\right)^{1/2} \\ &\quad \times \left(\frac{\Delta M}{M_{\odot}}\right)^{1/2} \left(\frac{v_w}{30 \text{ km s}^{-1}}\right)^{1/2} \left(\frac{\Delta t}{10^5 \text{ yr}}\right)^{1/2} , \end{aligned} \quad (8)$$

where m_p is the proton mass, and ΔM is the total mass-loss during the period. We assume that the consequent WD has $M_{\text{WD}} = \min\{0.5M, M_{\odot}\}$ (here M is for the progenitor RGB star mass), and ΔM is given by the decrease of mass before becoming a WD. For stars with $M > 8 M_{\odot}$, we estimate their Δv^2 similarly, assuming that the expansion period of the supernova ejecta is $\Delta t = 10^4$ yr, $v_w = 0.1c$, and ΔM is also deduced from $M_{\text{NS}} = \min\{0.1M, 1.5M_{\odot}\}$ and the progenitor mass M . Note that we consider this pulse momentum injection by AGB outflow in all types of gas-star interactions and intentionally ignore the “kicks” in NS (WD) momenta

induced by the asymmetries of supernovae (mass-loss in WD progenitor), so as not to obscure the NDF by such kicks (while NDF only adds to the effects of the kicks). Other epochs of stellar evolution are ignored as they have negligible impact on the stellar cluster dynamics.

2.3. Setup of Fiducial and Other Models

The fiducial model studies an OC immersed in the cold neutral medium (CNM) in the Galaxy, which has $\rho \approx 30 m_p \text{ cm}^{-3}$ and $T \approx 100$ K (Draine 2011). Considering the generality of the studied OC, the initial sample is generated in two steps. First, a rough sample is obtained using a two-phase cold collapse method. Second, based on the parameters of this sample (e.g., the number of stars, concentration, half-mass radii, etc.), a more realistic sample is generated using MCLUSTER. MCLUSTER is an open-source tool that can set initial conditions for N -body simulations or generate artificial star clusters for direct study (Küpper et al. 2011).

The first phase of the two-phase cold collapse method generates a 2000-star ensemble using the broken power-law initial mass function over the stellar mass range $0.2 < (M/M_{\odot}) < 25$. (Kroupa et al. 1993; Kroupa 2002; Reid et al. 2002)

$$\xi(M) \propto \begin{cases} M^{-2.35} , & M > 0.5 M_{\odot} \\ M^{-1.3} , & M < 0.5 M_{\odot} . \end{cases} \quad (9)$$

At the beginning of this phase, all stars have zero velocity and are distributed uniformly within an $r < 1$ pc sphere. In the second phase, this initial ensemble of stars is evolved using REBOUND *without* any stellar evolution processes or interactions with gas through 100 Myr. During this process, the OC undergoes sufficient relaxation, reaching virial equilibrium (i.e., the ratio of total kinetic energy E_k to total potential energy E_p is ~ -0.5).

In the next step, the parameters of the sample generated by the two-phase method were used as inputs, and a final sample was generated using MCLUSTER. The final sample consists of 1633 stars, with a total mass of $\Sigma M = 1249.21 M_{\odot}$, a concentration $c \sim 2.5$ (King (1966) model), and a half-mass radius $r_{1/2} \sim 3$ pc (see Figures 1–2). The consequent star ensemble is then adopted as the initial condition of the fiducial simulation, which is then evolved for $\Delta t_{\text{evo}} = 200$ Myr within the $L_{\text{box}} = 500$ pc bounding box. An important assumption is that star formation in the cluster is a single-age event, and once the cluster is formed, all the gas is expelled.

Different models exploring physical parameters (especially those about the ambient gas properties) use the

¹ <https://people.sissa.it/~sbressan/parsec.html>




Figure 1. The initial condition of the fiducial model includes logarithmic density histograms of distance (to the cluster center) and velocity in the upper and right panels, respectively. Mass bins are indicated by colors in the scattered plot. The distances and velocities histograms are calculated with respect to the centers of the space and velocity distributions, instead of the centers of mass. These centers are estimated by taking the median value along each dimension in configuration and velocity spaces, and we have verified that rotating the coordinate system does not affect the locations of centers significantly. The velocity dispersion of this sample is $\sim 0.96 \text{ km s}^{-1}$.

same initial condition of the fiducial model. Simulations for each model, including the fiducial model specific to the CNM, are conducted with three types of gas-star interactions, indicated as “NDF”, “DF” and “None” as the suffixes of the models, to emphasize the modes of the gas-star interactions. Note that the stars that do not launch winds (e.g., WDs) always experience DF effect even in the model marked as “NDF”.

3. FIDUCIAL MODEL RESULTS

The impacts of the gas-star interactions on the model OC immersed in the fiducial uniform CNM gas ($\rho_0 = 30 m_p \text{ cm}^{-3}$, $T = 100 \text{ K}$; CNMF hereafter for this case) are illustrated by the histograms in Figures 3. Thanks to the NDF acceleration, the stars generally reside on the side with higher mechanical energy in the CNMF-NDF run, compared to its CNMF-DF and CNMF-None counterparts. The distribution of distances to the cluster center (“distances” hereafter) tilts to the more distant side, while the velocity tends to be smaller as the stars




Figure 2. The initial condition of the fiducial model includes the mass distribution (top panel) and logarithmic density histograms of mass (bottom panel). The red and purple dashed lines in the top panel indicate half-mass and half-light radius, respectively.

are statistically farther from the global gravitational potential minimum.

3.1. Cluster Sizes and Velocity Dispersions

Statistics directly related to observables are presented in the left column of Figure 4. All three models exhibit the same trend of dynamical evaporation, indicated by their increasing half-mass and half-light radii (in each time interval without star deaths), and decreasing velocity dispersions. The half-mass radii for CNMF-NDF and CNMF-DF are both larger than the gas-free CNMF-None model, yet the underlying reasons should be different if one inspects the comparison in velocity dispersions. In self-gravitating systems, acceleration forces pushing the stars forward (such as NDF) tend to relocate them to shallower places in the potential well, and eventually reduces the magnitudes of velocity dispersions. Deceleration forces, in contrast, should put the stars closer to the cluster center first. However, this will lead to more frequent close encounters, which scatters stars into larger distances and also results in increased cluster sizes eventually.

The half-light radii exhibit a “cliff” structure when the most massive stars die (the middle left panel of Figure 4). Particularly, a pronounced cliff occurs at $t_{\text{evo}} \sim 8 \text{ Myr}$, corresponding to the death of the most




Figure 3. The impact of NDF and DF on the distance (upper panel) and velocity (lower panel) distributions of OC in the CNMF. Note that both panels display the differences compared to the None case in their distribution functions.

massive star ($\sim 22.3 M_{\odot}$). The differences in half-light radii among the three models are much smaller compared to those in half-mass radii due to the evaporation of massive stars from the cluster. This is because half-light radii are weighted more heavily towards the more massive main sequence stars, given the mass-luminosity scaling relation $L \propto M^{3.5}$. For low-mass stars, the differences in their luminosities are not significant, and their stellar winds are not as strong, resulting in the NDF effects being not prominent. Moreover, the half-light radius of the DF model cannot match the NDF model since gravitational scatterings tend to eject less massive stars due to the conservation of momentum and mechanical energy.

The binaries formation within OC have drawn the concern in some recent research works (e.g., Li & Mao 2018). Figure 5 illustrates the fraction of bounded pairs $F_b \equiv N_{\text{pair,b}}/N_{\text{pair,tot}}$, where $N_{\text{pair,tot}} = N(N-1)/2$ is the total number of pairs in the system, and $N_{\text{pair,b}}$ is for the bounded pairs that have negative total energy in their center-of-mass frame. The three runs for the

CNMF model does not exhibit appreciable differences in F_b , as the number of stars in this work is insufficient to make the binaries important in the fiducial runs.

3.2. Mass Segregated Clusters

As star clusters dynamically evolve, they exhibit an increasing degree of mass segregation, with a general tendency to continue progressing towards even greater mass segregation (e.g., Chandrasekhar 1943; Chandrasekhar & von Neumann 1942). In this work, the fiducial sample used was obtained without specifying the degree of mass segregation. Thanks to the help of MCLUSTER, it is possible to generate samples with a specified degree of mass segregation. In this section, another comparable sample corresponding to completely segregated conditions are generated (see Figures 6). Similar to the fiducial sample, this completely segregated samples are in virial equilibrium with 1633 stars, half-mass radii of about 3 pc, a cluster metallicity of $Z=0.017$, and a density profile following the King (1966) model with $c \sim 2.5$.

As shown in the right column of Figure 4, unlike the results obtained with the unsegregated sample, the degeneracy in the half-light radius between the NDF and the other two models is completely resolved. After 200 Myr evolution, both the half-mass radii and half-light radii for the NDF model are $\sim 2\times$ as large as those in other gas-star interaction models. Combining the two columns in Figure 4, these evolution tracks imply that the increase in cluster size for the NDF model is primarily driven by more massive stars, whose faster and stronger stellar winds push them to larger distances through NDF effects.

In the DF case, the half-light radius shows a "cliff" structure, which is related to the positional changes of the most massive star. Around $t_{\text{evo}} \sim 100$ Myr, some of the relatively massive stars are located closer to the cluster's center, while others are farther away. As the most massive stars (those closer to the center) lose mass and die, the remaining most massive stars are located farther from the cluster's center. These more distant massive stars contribute more to the half-light radius, causing it to rise. The half-light radius decreases only when these massive stars at the periphery either die or escape from the cluster, making the most massive stars once again those closer to the center.

3.3. Bulk Motion of Clusters

The gas-star interactions—both NDF and DF—are the results of stellar outflows and motion. Although the CNMF model runs assume no bulk motion of the whole cluster, the realistic are usually more complicated. The centers of mass of an OC and a gas cloud can move




Figure 4. The half-mass radii (upper panel), half-light radii (middle panel), and velocity dispersions (lower panel) of the OC in the CNMF model during its 200 Myr evolution period. The left column shows the results using the unsegregated (fiducial) sample, and the right column shows the results using the completely segregated sample. The half-light radii are calculated using the $L \propto M^{3.5}$ scaling relation for the main sequence and RGB stars, and there are “cliffs” at $t_{\text{evo}} \sim 8$ Myr when the most luminous stars end their main sequence and RGB lives. Note that the statistics are calculated in three dimensions, not projected along the line of sight. When calculating the half-mass and half-light radii, the center of mass and “center of light” is determined using stars within a 500 pc range centered around the median space coordinate. Velocity dispersion is calculated based on bounded main sequence stars with negative mechanical energy in the OC’s center-of-mass frame.

on their own tracks when evolving in galaxies, and the relative bulk motion between the two objects should be expected. We hence introduce models CNM3 and CNM5 to study such bulk motions at center of mass speeds $v_c = 3 \text{ km s}^{-1}$ and 5 km s^{-1} , respectively (see also Table 1).

Due to the morphologies of the stellar wind-ambient interaction regions, the magnitudes of NDF acceleration decreases with faster stellar velocities (Eq. 3). This relation qualitatively yields the phenomena in Figure 7, where the increase in mechanical energy caused by NDF is slightly suppressed with higher v_c . Regarding the in-

crease in half-mass and half-light radii, a similar suppression is observed in Figure 8. Concerning velocity dispersion, the NDF suppression results in stars settling in deeper regions of the potential well, leading to larger velocity dispersion than the CNMF. While these patterns can be qualitatively anticipated for higher v_c velocities, detailed analyses should be based on cluster-specific calculations rather than extrapolations.

3.4. Stellar Evolution and Compact Objects

Outflowing stars are expected to be pushed further away from the cluster center once NDF is in action. For typical main sequence stars immersed in the CNM, how-




Figure 5. The evolution of bounded pairs fraction of OC for all runs in the CNMF model.




Figure 6. The distance distribution for different mass bins in the unsegregated (fiducial) and completely segregated samples.

ever, such effects are generally obscured by the overall trend of OC dynamical evaporation. When an outflowing star travels at a small initial speed in a uniform medium, the asymptotic relation near $t \rightarrow \infty$ between the travel distance l and the time t roughly reads, using




Figure 7. The impact of NDF on the distance (upper panel) and velocity (lower panel) distributions of OC with different v_c . Note that both panels display the differences compared to CNMF-NDF in their distribution functions.

the approximate expression for Eq. (3),

$$l \sim 10 \text{ pc} \times \left(\frac{\rho}{30 m_p \text{ cm}^{-3}} \right)^{1/2} \left(\frac{v_w}{400 \text{ km s}^{-1}} \right)^{1/2} \times \left(\frac{\dot{m}_w}{10^{-13} M_\odot \text{ yr}^{-1}} \right)^{1/2} \left(\frac{t}{100 \text{ Myr}} \right)^{3/2} \propto t^{3/2}. \quad (10)$$

In the meantime, the velocity scales as $v \propto t^{1/2}$ asymptotically. These estimations also give the *lower* limit of time required to drive a star out of the cluster even *without* the gravitational potential well. When the potential well is present, direct exclusion of a solar-mass main sequence star only via NDF is not possible.

For objects with intensive outflows, however, the accelerations are much more significant dynamically. Although $a_{\text{NDF}} \propto v_w^{1/2}$ increases sub-linearly with v_w , the NS with $v_w = 0.1c$ can still escape from the stellar ensemble well before the evaporation of OCs. Figure 9 illustrates the efficient expulsion of NS (upper panel) by NDF (solid lines), compared to the DF (dotted lines)




Figure 8. The evolution of differences in half-mass radii (upper panel), half-light radii (middle panel), and velocity dispersions (lower panel) across different scenarios (CNMF, CNM3 and CNM5). The differences are obtained by subtracting the values of the gas-free case (“None”) from the NDF case).

and gas-free (dashed lines) cases where the ordinary “evaporation” process of OC and the increment of star’s velocity by AGB-outflow-induced pulse present. NS are driven out to $\gtrsim 10^2$ pc from the center of the cluster within only ~ 50 Myr, shorter than the lifetime of a typical OC. After escaping from the OC potential, the $l \propto t^{3/2}$ scaling and Eq. (10) holds semi-quantitatively for each of the NS at large distances ($\gtrsim 10$ pc). Assuming that the gaseous disk half-thickness is ~ 0.5 kpc in the Galaxy, such expulsion process will eventually take ~ 100 Myr to remove an NS from the disk. Such quick expulsions will likely lead to the scarcity of NS in OCs, and probably in the whole gaseous galactic disk. Figure 10 illustrates a reduced expulsion of NSs attributed to the attenuation of the NDF effect by the bulk motion. The NS expulsion is ubiquitous for gas-immersed OCs unless the cluster bulk motion is too fast relative to the gas. Previous works often attribute the NS exclusion from clusters to the “kicks” during asymmetric




Figure 9. The evolution trajectories of stars evolving into compact objects (NS in the upper panel, and WD in the lower panel), showing the distances r to the cluster center, in CNMF with distinct stars represented by different colors. The upper and lower panels show the tracks of main sequence stars transitioning into NS and WDs under different gas-star interactions. Thick lines represent trajectories during the main sequence phase, while thin lines indicate trajectories after they evolve into NS or WD. These trajectories only depict stars exiting the main sequence within the evolution time. Note that all NSs are expelled from OC under the influence of NDF.

supernovae explosions (e.g., Fragione & Banerjee 2020). With the NDF effects, however, NS exclusion can still take place without these “kicks”. Note that even in the case of intensive outflows from stars, the winds of the stars cannot combine to drive a wind from the OC. This is because the standoff distance of the contact discontinuity,

$$R_0 \approx 0.004 \text{ pc} \times \left(\frac{v_w}{400 \text{ km s}^{-1}} \right)^{1/2} \left(\frac{v_*}{1 \text{ km s}^{-1}} \right)^{-1} \times \left(\frac{\dot{m}_w}{4 \times 10^{-13} M_\odot \text{ yr}^{-1}} \right)^{1/2} \left(\frac{\rho}{30 m_p \text{ cm}^{-3}} \right)^{-1/2}, \quad (11)$$




Figure 10. The evolutionary trajectories of main sequence stars transitioning into NSs are depicted in the scenarios of CNM3 (upper panel) and CNM5 (lower panel). In both CNM3 and CNM5, some NSs remain within the OC but tend to move outward. This phenomenon is attributed to the bulk motion attenuating the NDF effect.

is much smaller than the distance between member stars (e.g., the minimum distance in the initial OC of the fiducial case is ~ 0.01 pc).

In contrast to NS, WDs do not exhibit outflows in our model (which applies to stand-alone WDs, yet accreting WDs are likely to produce accretion disk winds or even jets). The expulsion phenomenon primarily results from pulses induced by AGB outflows rather than NDF effects (Figure 9, lower panel). Only a fraction of WDs are expelled from the cluster due to these pulses, and the remaining WDs continue to reside within the cluster.

However, the deficiency of WDs in young OCs has been observed (e.g., Kalirai et al. 2001). Fellhauer et al. (2003) investigates that the paucity of WDs could result from the isotropic recoil speed attained from nonspherically symmetric mass loss during post-main-sequence evolution. In order to consider this deficiency, following the approach of Fellhauer et al. (2003), the transition pulse of a star into a WD is set with a direction that uniformly distributes randomly in space and a magnitude




Figure 11. The evolutionary trajectories of main sequence stars transitioning into WDs. Note that in this scenario, the pulse is set with a direction that uniformly distributes randomly in space and a magnitude of 2 km s^{-1} .

of 2 km s^{-1} . The result is demonstrated in Figure 11. Under this setup, almost all WDs are expelled from the OC. In the scenario of AGB-outflow-induced pulse, it is assumed that AGB outflows are isotropic, while nonspherically symmetric mass loss leads to stronger pulses, explaining the difference in WD expulsion phenomena.

In the current OC model, where binary systems are very rare (Figure 5), the assumption that WDs have no outflows should hold reasonably well. In future works studying more stars and significantly higher occurrence rates of binaries, the fate of WDs with outflows in binaries should be revisited with proper modeling of their accretion feedbacks and the non-isotropic outflow patterns (e.g., Li et al. 2020).

3.5. Cluster in Galactic Potential

As a star departs from its cluster, its path is computed considering the Galaxy's external potential. Incorporating the Galactic potential in models of OC offers a more realistic scenario. Following Allen & Santillan (1991), the Galactic potential ϕ_{Gal} is composed of three parts: the central part ϕ_1 , the disc ϕ_2 , and the halo ϕ_3 , i.e., $\phi_{\text{Gal}} = \phi_1 + \phi_2 + \phi_3$. These three parts of the potential can be expressed in cylindrical coordinates (R, z) as follows:

$$\begin{aligned} \phi_1 &= -\frac{M_1}{(R^2 + z^2 + b_1^2)^{1/2}}, \\ \phi_2 &= -\frac{M_2}{\left\{R^2 + \left[a_2 + (z^2 + b_2^2)^{1/2}\right]^2\right\}^{1/2}}, \\ \phi_3 &= -\frac{M(r)}{r} - \frac{M_3}{1.02a_3} \\ &\quad \times \left[-\frac{1.02}{1 + (r/a_3)^{1.02}} + \ln(1 + (r/a_3)^{1.02}) \right]_r^{r_d}, \end{aligned} \quad (12)$$




Figure 12. Similar to Figure 4, but the results shown are the NDF scenario considered the Galaxy’s potential.

where

$$\begin{aligned}
 r &= (R^2 + z^2)^{1/2}, \\
 M(r) &= -\frac{M_3(r/a_3)^{2.02}}{1 + (r/a_3)^{1.02}}.
 \end{aligned}
 \tag{13}$$

The parameters for the above equations are $M_1 = 1.41 \times 10^{10} M_\odot$, $b_1 = 387$ pc, $M_2 = 8.56 \times 10^{10} M_\odot$, $a_2 = 532$ pc, $b_2 = 250$ pc, $M_3 = 10.7 \times 10^{10} M_\odot$, $a_3 = 1.2 \times 10^4$ pc and $r_d = 10^5$ pc. The OC are positioned on circular orbits at a speed of 220 km s^{-1} , specifically $(v_R, v_\phi, v_z) = (0, 220, 0) \text{ km s}^{-1}$, at a distance of 8.5 kpc from the Galactic center.

The impacts of the gas-star interactions in this scenario are illustrated in Figure 12. The overall trend is consistent with Figure 4, where both the half-mass and half-light radii increase over time during the time interval without star deaths, showing an increase in cluster size. Also, due to the absence of massive stars and the effect of the tidal field, the differences of the half-light radii among the three gas-star interactions are more indistinguishable.

The velocity dispersion among the three models decreases during the first ~ 50 Myr and then tends to stabilize around ~ 0.37 . This results from the combined gravitational effects of the external tidal field and the interactions among the stars within the cluster. At ~ 50 Myr, the cluster deforms under the influence of the tidal field and gradually forms a tidal tail structure. [Dinbier & Kroupa \(2020\)](#) studied the dynamical evolution of the tidal tails of OC in detail and provided a semi-analytical model. They found that the velocity dispersion of tails formed by gas expulsion exhibits an oscillation-like pattern over time. They concluded that such oscillations depend solely on the local galactic frequencies, i.e., the orbital frequency and the epicyclic frequency, and do not depend on any cluster properties (e.g., the initial velocity dispersion or mass).

It must be acknowledged that the tidal field is one of the significant effects of the evolution of internal dynamical features within OC. However, this work focuses on the effect of gas-star interactions. While incorporating the tidal field would render the scenario more realistic, it also complicates the physical mechanisms, making it challenging to analyze the effects of different mechanisms clearly. Therefore, the tidal field will not be considered in subsequent scenarios, meaning the external Galaxy’s potential will be excluded. In future work, the influence of both tidal fields and gas interactions on the dynamical characteristics of stars or clusters should be included.

4. MODELS EXPLORING AMBIENT GAS PROPERTIES

The galaxy disk contains multiple phases of the interstellar medium, whereas the CNM phase that we have explored in §3 only occupies a relatively small volume fraction (typically a few per cent, see e.g., [Ferrière 2001](#); [Draine 2011](#)). We therefore conduct various simulations to study the interactions between OCs and other types of gases, including the warm neutral medium (WNM), diffuse molecular regions (MD), dense and intermediate molecular clouds, and even the mid-plane of active galactic nucleus (AGN) disks ([Draine 2011](#); [Cantiello et al.](#)

Table 1. Physical parameters for N -body simulations for open clusters in different scenarios.

Name	Description [†]
CNMF	Fiducial Model (Cold Neutral Media; Section 2.3)
CNM3	$v_c = 3 \text{ km s}^{-1}$
CNM5	$v_c = 5 \text{ km s}^{-1}$
WNM0	$[\rho] = 0, T = 5000 \text{ K}$ (Warm Neutral Media)
WNM1	$[\rho] = 1, T = 1000 \text{ K}$ (Warm Neutral Media)
MD	$[\rho] = 3, T = 30 \text{ K}$ (Diffuse Molecular Regions)
M4*	$[\rho] = 4, T = 30 \text{ K}$ (Molecular Clouds)
M5*	$[\rho] = 5, T = 30 \text{ K}$ (Molecular Clouds)
M7*	$[\rho] = 7, T = 30 \text{ K}$ (Molecular Clouds)
AGND*	$[\rho] = 9, T = 30 \text{ K}$ (AGN Disk Gases)

NOTE—[†]: Only the properties different from the fiducial model are described; $[\rho] \equiv \log_{10}[\rho_0/(0.3 m_p \text{ cm}^{-3})]$
 *: Evolution time $\Delta t_{\text{evo}} = 20 \text{ Myr}$.

2021). Properties of these gases are summarized in Table 1.

4.1. Distributions in the Configuration and Velocity Spaces

In a virialized self-gravitating system, stars staying at larger distances from the cluster center generally move at slower velocities. When the NDF and DF accelerations are sufficiently low and gradual, their impact on the cluster configurations is mostly adiabatic, and the system remains virialized during the evolution. When the ambient gas density is raised to $3 \times 10^8 m_p \text{ cm}^{-3}$ in the model AGND, however, the NDF acceleration for a solar-mass star moving at 1 km s^{-1} becomes $a_{\text{NDF}} \sim 10^{-9} \text{ cm s}^{-2}$, which is $\sim 10\times$ the gravitational acceleration by a $10^3 M_\odot$ cluster at 10 pc. A cluster evolving under such strong NDF or DF accelerations no longer remains adiabatic; the dispersal takes place directly within the first $\sim 3 \text{ Myr}$ resulting in a quickly expanding spatial size and higher velocity dispersions (Figure 13). This phenomenon also appears in the models M7 and M5 that stand for typical molecular clouds, while the models for diffuse molecular regions (model MD) and more diffuse gas (WNM0, WNM1) are qualitatively similar to the CNMF case.

Another effect that emerges in various ambient models is the stratification of stars. The acceleration by NDF on stars tend to increase their energy, which is similar to some “buoyancy” in self-gravitating OCs. As more massive stars launch more powerful stellar winds (Section 2.2), they tend to move quicker to the “surfaces” of OCs under NDF, and vice versa. We notice that such “buoyancy” will not make the young OCs “inversely seg-




Figure 13. Similar to Figure 8, but for various ambient gas densities without v_c , showing only the first 20 Myr. Note that the cut-off at around 17 Myr in the velocity dispersion of AGND exists, as no stars have negative mechanical energy in the OC’s center-of-mass frame.

regated” initially, as the timescales required to fully “inversely segregate” the cluster ($\sim 10^7 - 10^8 \text{ yr}$) (see e.g. Figure 14) is longer than the time period of star formation ($\lesssim 10^7 \text{ yr}$) (Palla & Stahler 1999; Palla et al. 2005).

Since the stellar luminosity scales as $L \propto M^{3.5}$, the half-light radii are considerably greater than half-mass radii in NDF-affected clusters. For those models that have the most dense ambients (M5, M7, AGND), the half-light radii are $\gtrsim 3\times$ greater than the half-light radii after only $\sim 10 \text{ Myr}$, indicating an obvious radial strat-




Figure 14. The distance distribution for different mass bins in the M5-NDF scenario after 5 Myr (upper panel) and 20 Myr (lower panel) evolution. Due to their stronger winds, massive stars are subject to greater NDF than low-mass stars, resulting in a reduced mass segregation.

ification in terms of stellar types. When attempting to infer the age of an observed OC by the observations in astrometry and dynamics, one should take care of the possible encounters with dense clouds, whose effect in puffing up the OCs is similar to the intrinsic evaporation of clusters. Statistics on the radial distributions of mass and light in such clusters, therefore, may be helpful in reducing this type of parameter degeneracies.

4.2. Puff-up and Dispersal of Open Clusters in Gases

According to Eq. (3), the magnitude of NDF acceleration depends much more sensitively on ρ_0 than on T . Figure 15 reveals the general trend that the NDF effects increase with denser ambient gas, raising the total mechanical energy and easing the evaporation of cluster stars. The scaling relation in Eq. (3) is sub-linear with respect to ρ_0 ($\sim \rho_0^{1/2}$). Assuming a relatively invariant velocity distribution over the same period of time, one can infer the scaling relation $dE/dt \sim \rho_0^{1/2}$, and subsequently $\Delta|1/r| \sim \rho_0^{1/2}$. This pattern is qualitatively seen in Figure 13, yet we note that such simple scaling




Figure 15. Similar to Figure 7, but the results shown are the difference in their distribution functions between the labeled models and the MD (diffuse molecular regions) model (see also Table 1) for denser ambient gas without v_c , influenced by NDF.

no longer applies quantitatively when there are significant increases in velocity dispersions.

Models M5 and M7 show that OCs in relatively dense ambients will quickly puff up or even disperse within ~ 20 Myr, a timescale that is comparable or even shorter than the crossing time of an OC over a dense molecular cloud. If one adopts the typical sizes ~ 20 pc of dense molecular clouds ($\rho_0 \gtrsim 10^4 m_p \text{ cm}^{-3}$) (Cernicharo 1991; Bergin & Tafalla 2007) with an OC crossing time ~ 40 Myr (assuming bulk velocity $\sim 0.5 \text{ km s}^{-1}$), Model M5 indicates that a cluster should puff up to $r_{1/2} \gtrsim 10$ pc after this crossing. A similar process will also happen for the cluster-gas encounter with a denser molecular cloud described by Model M7, even if the lifetime limits of a few Myr for these dense clouds are considered.

AGN disks are also considered as places where stars can form, which have lifetimes on the order of $10^0 - 10^3$ Myr (Marconi et al. 2004; Martini & Weinberg 2001). If a $10^3 M_\odot$ OC forms and stays inside the AGN disk for $\gtrsim 5$ Myr, this OC should no longer be considered as a cluster anymore, even if the destruction by orbital motion shears is not taken into account. In other




Figure 16. Similar to Figure 10, but for the scenarios of WNM0 (upper panel) and WNM1 (lower panel).




Figure 17. Similar to Figure 5, but for the AGND scenario with an evolution time of 20 Myr.

words, the likelihood that one can find OCs formed in AGN disks should not be significant.

4.3. Compact Objects and Binaries

As elaborated in Sections 3.3 and 4.1, higher stellar velocities and lower ambient gas densities suppress

the NDF acceleration. In various models, the expulsion of NS via NDF behaves accordingly. Figure 16 illustrates that, even for Model WNM1 whose ambient density is merely $3 m_p \text{ cm}^{-3}$, such expulsion mechanism is still working, although the time required is considerably longer. This effect is, nevertheless, still more perceptible than the CNM5 case (see Figure 10). For Model WNM0, the expulsion mechanism weakened compared to WNM1, but all NSs still tend to move away from the OC.

The fraction of bounded pairs corresponding to various models does not show any considerable changes compared to the fiducial CNMF model, except for the AGND model with DF only (Figure 17). The number of bounded pairs increased drastically by $\sim 10^1$ times in the first ~ 20 Myr of evolution. Strong deceleration caused by DF shrinks the separations between stars and reduces stellar velocities, facilitating the formation of binaries. Several studies have explored this phenomenon. Rafikov (2013, 2016) proposed that gas-assisted inspirals efficiently merge supermassive BH binaries within a Hubble time. The introducing NDF would nonetheless “flip” this scenario, widening the binary separations. Wang & Li (2022) demonstrated in simulations of AGB star-outflowing pulsars that a dense and slow outflow exerts a positive torque on the binary, causing $\gtrsim 10\%$ orbit expansion. Extrapolating these mechanisms to the OCs immersed in dense gas can cause a severe inhibition of binary occurrence rates.

5. DISCUSSION AND SUMMARY

In this paper we study the OC evolution immersed in galactic gases. Overall, the introduction of NDF makes the OC puffer and eases the evaporation of cluster stars, resulting in reduced velocity dispersions and increased half-mass and half-light radii. This effect scales sub-linearly with the ambient density and is mostly irrelevant to the ambient temperature as the ram pressures on both sides dominate the gas-star interactions. NS with powerful winds are expelled from the cluster due to the intense NDF effect, while WD stars undergo conventional evaporation due to the absence of outflow.

5.1. Impacts of the NDF Effects on Cluster Evolution

While gas DF has garnered most of the attention, NDF has been relatively overlooked. Opposite to DF, NDF accelerates stars along their motion direction. Such difference in the effects between NDF versus DF can “flip” the signs of multiple physical mechanisms. For instance, one may expect that the OC evaporation is suppressed or even inhibited by DF when encountered with gas, while the actual scenario could be right the opposite due to NDF. In the densest gases including dark

molecular clouds, AGN disks, and AGN tori (whose average column density is as high as $\sim 10^{24} \text{ cm}^{-2}$; see, e.g., [Zhao et al. 2021](#)), dispersal of OCs can be almost instant compared to their dynamical lifetimes as clusters. Interpretations about observations on cluster kinetics should take the possibility of NDF into account.

Observation missions like Gaia ([Gaia Collaboration et al. 2016](#)) provide abundant astrometric data for stars in our galaxy, facilitating the study of OCs’ dynamical evolution (e.g., [Maurya et al. 2023](#)). Because of the radial stratification in stellar masses (Section 4.1), one can prospectively tell the effects of gas-star interactions by measuring the half-mass and half-light radii, as well as the velocity dispersions. The combination of high precision photometry with astrometry can further help the researchers to find out the radial distribution functions of different stellar types. These measurements can be a direct characterization of gas-star interaction history that are potentially important for identifying an OC’s “invasion” into dense gases during its evolution history.

5.2. Neutron Star Depletion

Due to substantial outflows, NSs gradually drift away from their associated OC under the influence of NDF in all simulations. Conversely, WD stars influenced by NDF experience a pulse momentum injection from the main sequence to the AGB stage and subsequently to the WD stage. However, upon transitioning into the WD phase, WDs decelerate due to DF as they lack outflows. This mechanism ensures that part of WDs remain confined within the OC and experience conventional evaporation. If the kick received by a WD is strong enough (such as through an asymmetric mass loss process ([Fragione & Banerjee 2020](#))), it is more likely to be expelled from the OC (see in Figure 11).

It is commonly assumed that kicks by asymmetric supernovae drive NS to high velocities relative to the cluster centers, and eventually expel them from OCs. [Lai \(2004\)](#) reviewed various physical mechanisms leading to kicks, and [Bortolas et al. \(2017\)](#) suggested that NSs are generally scattered away from SgrA* due to such kicks. [Contenta et al. \(2015\)](#) indicated that the natal kick of NSs can alter the cluster’s lifetime by almost a factor of ~ 4 . Admittedly, the kicks serve as an effective expulsion mechanism of NS. However, kicks only entail a one-time momentum injection, whose intensity and direction of momentum injection is stochastic and can well result in the NS’s deceleration. The NDF effects are, in contrast, a much more steady and robust mechanism that almost ensures the expulsion without gravitational scatterings. Such effects can further lead to the reduction of NS binaries and mergers within OCs, which is poten-

tially relevant to the spatial distribution of NS-related mergers and gravitational wave events.

5.3. Future Works

Due to the limitations in physical modeling and computation, this work does have some caveats and issues that should be addressed in future works. For example, simplistic isotropic stellar wind models are adopted for each type of star, which is suitable only for a fraction of NS. Many NS—especially those with accretion disks—may launch directed or bipolar outflows, which could affect the intensity and direction of the NDF forces ([Li et al. 2020](#)). The WD model in this study lacks wind and is susceptible to DF only. While stand-alone WD stars may not have outflows, accretion from their companions may cause disk winds and jets. This study ignores the time lag in forming stars within the OC and does not consider the interaction between star and star-forming gas. However, the expulsion of residual star-forming gas does not occur instantaneously after the completion of the star formation process in realistic scenario. Observations reveal that star formation efficiencies within OCs vary widely, ranging from several percent to 30 percent for dense clumps within molecular clouds ([Lada & Lada 2003](#); [Higuchi et al. 2009](#)), and from 0.1 percent to a few percent for their associated giant molecular clouds ([Evans et al. 2009](#); [Murray 2011](#)). This implies that newly formed or forming stars (such as protostars and pre-MS stars, which have outflows) would interact with star-forming gas, resulting in the influence of NDF on the initial phase space distribution of stars within young OC. Moreover, the stellar winds from these stars may blow unprocessed gas out of the cluster, inhibiting the ongoing star formation process. Refined models in the future should take the complexities of stellar outflows and the star formation process into account.

Although our study considers the highly dense scenario of an AGN disk (AGND scenario), we have omitted multiple physical conditions relevant to AGN in order to isolate and emphasize the impact by NDF. Incorporating the shear effect is crucial for understanding stellar formation and evolution within AGN disks. However, this topic is specific to disks and beyond the scope of this paper, which should be addressed in a separate study. Future studies on AGN disks will require additional considerations, such as orbital motion and vertical stratification of gas. The limited thickness of AGN disks differs substantially from the assumed uniform gas ambient in this paper, especially considering the vertical density gradients. Furthermore, isolated stars would experience the DF effects in the highly-dense environment

of a galaxy's circumnuclear regions. Future works shall also explore these possibilities of NDF-affected stars.

In addition, our studies focus on relatively small and sparse OCs with $\sim 10^3$ stars. In high-density environments, particularly in AGN disks, super star clusters are often present instead of OCs, whose star formation efficiencies are typically high. Investigating larger ensembles of stars such as globular clusters can impose additional challenges. Globular clusters have $\sim 10^6$ or even more stars, while each of the stars might still be dynamically important. Accurate treatments of frequent close encounters are beyond the capacity of ordinary orbital integrators with tree-based or particle-mesh based gravity solvers, and special algorithms are required for sufficient accuracy. During close encounters, shock structures caused by gas-star interactions on both sides may collide, resulting in highly complex gas morphologies and interaction patterns. Understanding how such complex physical scenarios affect the dynamical evolution of stars within globular clusters could be explored once proper algorithms are prepared for complicated compu-

tations. Even without proper treatments of frequent close encounters in more massive clusters, the dynamical evolution of compact objects (NS, WD, and BH) are still qualitatively feasible within the current framework for globular clusters and super star clusters. Detailed discussions are nonetheless beyond the scope of the current paper focusing on OCs, and a subsequent paper is being composed specifically addressing this issue.

- 1 L. C. Ho was supported by the National Science Founda-
- 2 tion of China (11991052, 12233001), the National
- 3 Key R&D Program of China (2022YFF0503401), and
- 4 the China Manned Space Project (CMS-CSST-2021-
- 5 A04, CMS-CSST-2021-A06). X. Fu thanks the support
- 6 of the National Natural Science Foundation of China
- 7 (NSFC) No. 12203100 and the China Manned Space
- 8 Project with NO. CMS-CSST-2021-A08. M. Liu and L.
- 9 Wang appreciates the computational resources provided
- 10 by the Kavli Institute of Astronomy and Astrophysics at
- 11 Peking University. We also thank our colleagues Renyue
- 12 Cen, Kohei Inayoshi, Fangzhou Jiang, and Meng Sun for
- 13 helpful discussions and suggestions.

REFERENCES

- Allen, C., & Santillan, A. 1991, *RMxAA*, 22, 255
- Baruteau, C., Cuadra, J., & Lin, D. N. C. 2011, *ApJ*, 726, 28, doi: [10.1088/0004-637X/726/1/28](https://doi.org/10.1088/0004-637X/726/1/28)
- Bergin, E. A., & Tafalla, M. 2007, *ARA&A*, 45, 339, doi: [10.1146/annurev.astro.45.071206.100404](https://doi.org/10.1146/annurev.astro.45.071206.100404)
- Bernabeu, G., Magazzu, A., & Stalio, R. 1989, *A&A*, 226, 215
- Binney, J., & Tremaine, S. 2008, *Galactic Dynamics: Second Edition* (Princeton University Press)
- Bondi, H., & Hoyle, F. 1944, *MNRAS*, 104, 273, doi: [10.1093/mnras/104.5.273](https://doi.org/10.1093/mnras/104.5.273)
- Bortolas, E., Mapelli, M., & Spera, M. 2017, *MNRAS*, 469, 1510, doi: [10.1093/mnras/stx930](https://doi.org/10.1093/mnras/stx930)
- Bressan, A., Marigo, P., Girardi, L., et al. 2012, *MNRAS*, 427, 127, doi: [10.1111/j.1365-2966.2012.21948.x](https://doi.org/10.1111/j.1365-2966.2012.21948.x)
- Brooks, D. H., Ugarte-Urra, I., & Warren, H. P. 2015, *Nature Communications*, 6, 5947, doi: [10.1038/ncomms6947](https://doi.org/10.1038/ncomms6947)
- Cantat-Gaudin, T. 2022, *Universe*, 8, 111, doi: [10.3390/universe8020111](https://doi.org/10.3390/universe8020111)
- Cantiello, M., Jermyn, A. S., & Lin, D. N. C. 2021, *ApJ*, 910, 94, doi: [10.3847/1538-4357/abdf4f](https://doi.org/10.3847/1538-4357/abdf4f)
- Cernicharo, J. 1991, in *NATO Advanced Study Institute (ASI) Series C, Vol. 342, The Physics of Star Formation and Early Stellar Evolution*, ed. C. J. Lada & N. D. Kylafis, 287, doi: [10.1007/978-94-011-3642-6_8](https://doi.org/10.1007/978-94-011-3642-6_8)
- Chandrasekhar, S. 1943, *ApJ*, 97, 255, doi: [10.1086/144517](https://doi.org/10.1086/144517)
- Chandrasekhar, S., & von Neumann, J. 1942, *ApJ*, 95, 489, doi: [10.1086/144420](https://doi.org/10.1086/144420)
- Chen, Y., Bressan, A., Girardi, L., et al. 2015, *MNRAS*, 452, 1068, doi: [10.1093/mnras/stv1281](https://doi.org/10.1093/mnras/stv1281)
- Choi, J., Dotter, A., Conroy, C., et al. 2016, *ApJ*, 823, 102, doi: [10.3847/0004-637X/823/2/102](https://doi.org/10.3847/0004-637X/823/2/102)
- Contenta, F., Varri, A. L., & Heggie, D. C. 2015, *MNRAS*, 449, L100, doi: [10.1093/mnrasl/slv026](https://doi.org/10.1093/mnrasl/slv026)
- Demircan, O., & Kahraman, G. 1991, *Ap&SS*, 181, 313, doi: [10.1007/BF00639097](https://doi.org/10.1007/BF00639097)
- Dinnbier, F., & Kroupa, P. 2020, *A&A*, 640, A84, doi: [10.1051/0004-6361/201936570](https://doi.org/10.1051/0004-6361/201936570)
- Draine, B. T. 2011, *Physics of the Interstellar and Intergalactic Medium* (Princeton University Press)
- Duncan, R. C., Shapiro, S. L., & Wasserman, I. 1986, *ApJ*, 309, 141, doi: [10.1086/164587](https://doi.org/10.1086/164587)
- Edgar, R. 2004, *NewAR*, 48, 843, doi: [10.1016/j.newar.2004.06.001](https://doi.org/10.1016/j.newar.2004.06.001)
- Evans, Neal J., I., Dunham, M. M., Jørgensen, J. K., et al. 2009, *ApJS*, 181, 321, doi: [10.1088/0067-0049/181/2/321](https://doi.org/10.1088/0067-0049/181/2/321)

- Fellhauer, M., Lin, D. N. C., Bolte, M., Aarseth, S. J., & Williams, K. A. 2003, *ApJL*, 595, L53, doi: [10.1086/379005](https://doi.org/10.1086/379005)
- Ferrière, K. M. 2001, *Reviews of Modern Physics*, 73, 1031, doi: [10.1103/RevModPhys.73.1031](https://doi.org/10.1103/RevModPhys.73.1031)
- Fragione, G., & Banerjee, S. 2020, *ApJL*, 901, L16, doi: [10.3847/2041-8213/abb671](https://doi.org/10.3847/2041-8213/abb671)
- Friel, E. D. 1995, *ARA&A*, 33, 381, doi: [10.1146/annurev.aa.33.090195.002121](https://doi.org/10.1146/annurev.aa.33.090195.002121)
- Fu, X., Bragaglia, A., Liu, C., et al. 2022, *A&A*, 668, A4, doi: [10.1051/0004-6361/202243590](https://doi.org/10.1051/0004-6361/202243590)
- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, 595, A1, doi: [10.1051/0004-6361/201629272](https://doi.org/10.1051/0004-6361/201629272)
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1, doi: [10.1051/0004-6361/201833051](https://doi.org/10.1051/0004-6361/201833051)
- . 2021, *A&A*, 649, A1, doi: [10.1051/0004-6361/202039657](https://doi.org/10.1051/0004-6361/202039657)
- Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2023, *A&A*, 674, A1, doi: [10.1051/0004-6361/202243940](https://doi.org/10.1051/0004-6361/202243940)
- Gruzinov, A., Levin, Y., & Matzner, C. D. 2020, *MNRAS*, 492, 2755, doi: [10.1093/mnras/staa013](https://doi.org/10.1093/mnras/staa013)
- Habing, H. J., & Olofsson, H. 2004, *Asymptotic Giant Branch Stars* (Springer-Verlag New York), doi: [10.1007/978-1-4757-3876-6](https://doi.org/10.1007/978-1-4757-3876-6)
- Higuchi, A. E., Kurono, Y., Saito, M., & Kawabe, R. 2009, *ApJ*, 705, 468, doi: [10.1088/0004-637X/705/1/468](https://doi.org/10.1088/0004-637X/705/1/468)
- Hoyle, F., & Lyttleton, R. A. 1939, *Proceedings of the Cambridge Philosophical Society*, 35, 405, doi: [10.1017/S0305004100021150](https://doi.org/10.1017/S0305004100021150)
- Hunt, E. L., & Reffert, S. 2023, *A&A*, 673, A114, doi: [10.1051/0004-6361/202346285](https://doi.org/10.1051/0004-6361/202346285)
- Just, A., Yurin, D., Makukov, M., et al. 2012, *ApJ*, 758, 51, doi: [10.1088/0004-637X/758/1/51](https://doi.org/10.1088/0004-637X/758/1/51)
- Kalirai, J. S., Ventura, P., Richer, H. B., et al. 2001, *AJ*, 122, 3239, doi: [10.1086/324463](https://doi.org/10.1086/324463)
- Kato, M. 1983, *PASJ*, 35, 33
- Kim, H., & Kim, W.-T. 2007, *ApJ*, 665, 432, doi: [10.1086/519302](https://doi.org/10.1086/519302)
- King, I. R. 1966, *AJ*, 71, 64, doi: [10.1086/109857](https://doi.org/10.1086/109857)
- Krause, M. G. H., Offner, S. S. R., Charbonnel, C., et al. 2020, *SSRv*, 216, 64, doi: [10.1007/s11214-020-00689-4](https://doi.org/10.1007/s11214-020-00689-4)
- Kroupa, P. 2002, *Science*, 295, 82, doi: [10.1126/science.1067524](https://doi.org/10.1126/science.1067524)
- Kroupa, P., Tout, C. A., & Gilmore, G. 1993, *MNRAS*, 262, 545, doi: [10.1093/mnras/262.3.545](https://doi.org/10.1093/mnras/262.3.545)
- Krumholz, M. R., McKee, C. F., & Bland-Hawthorn, J. 2019, *ARA&A*, 57, 227, doi: [10.1146/annurev-astro-091918-104430](https://doi.org/10.1146/annurev-astro-091918-104430)
- Kudritzki, R. P., & Reimers, D. 1978, *A&A*, 70, 227
- Kuiper, G. P. 1938, *ApJ*, 88, 472, doi: [10.1086/143999](https://doi.org/10.1086/143999)
- Küpper, A. H. W., Maschberger, T., Kroupa, P., & Baumgardt, H. 2011, *MNRAS*, 417, 2300, doi: [10.1111/j.1365-2966.2011.19412.x](https://doi.org/10.1111/j.1365-2966.2011.19412.x)
- Lada, C. J., & Lada, E. A. 2003, *ARA&A*, 41, 57, doi: [10.1146/annurev.astro.41.011802.094844](https://doi.org/10.1146/annurev.astro.41.011802.094844)
- Lai, D. 2004, in *Cosmic explosions in three dimensions*, ed. P. Höflich, P. Kumar, & J. C. Wheeler, 276, doi: [10.48550/arXiv.astro-ph/0312542](https://doi.org/10.48550/arXiv.astro-ph/0312542)
- Li, X., Chang, P., Levin, Y., Matzner, C. D., & Armitage, P. J. 2020, *MNRAS*, 494, 2327, doi: [10.1093/mnras/staa900](https://doi.org/10.1093/mnras/staa900)
- Li, Z., & Mao, C. 2018, *ApJ*, 859, 36, doi: [10.3847/1538-4357/aabc09](https://doi.org/10.3847/1538-4357/aabc09)
- Magrini, L., Viscasillas Vázquez, C., Spina, L., et al. 2023, *A&A*, 669, A119, doi: [10.1051/0004-6361/202244957](https://doi.org/10.1051/0004-6361/202244957)
- Marconi, A., Risaliti, G., Gilli, R., et al. 2004, *MNRAS*, 351, 169, doi: [10.1111/j.1365-2966.2004.07765.x](https://doi.org/10.1111/j.1365-2966.2004.07765.x)
- Martini, P., & Weinberg, D. H. 2001, *ApJ*, 547, 12, doi: [10.1086/318331](https://doi.org/10.1086/318331)
- Mathieu, R. D. 1984, *ApJ*, 284, 643, doi: [10.1086/162447](https://doi.org/10.1086/162447)
- Maurya, J., Joshi, Y. C., Samal, M. R., Rawat, V., & Gour, A. S. 2023, *Journal of Astrophysics and Astronomy*, 44, 71, doi: [10.1007/s12036-023-09959-3](https://doi.org/10.1007/s12036-023-09959-3)
- Miville-Deschênes, M.-A., Murray, N., & Lee, E. J. 2017, *ApJ*, 834, 57, doi: [10.3847/1538-4357/834/1/57](https://doi.org/10.3847/1538-4357/834/1/57)
- Mor, R., Livne, E., & Piran, T. 2023, *MNRAS*, 518, 623, doi: [10.1093/mnras/stac2775](https://doi.org/10.1093/mnras/stac2775)
- Murray, N. 2011, *ApJ*, 729, 133, doi: [10.1088/0004-637X/729/2/133](https://doi.org/10.1088/0004-637X/729/2/133)
- Nilakshi, Sagar, R., Pandey, A. K., & Mohan, V. 2002, *A&A*, 383, 153, doi: [10.1051/0004-6361:20011719](https://doi.org/10.1051/0004-6361:20011719)
- Ostriker, E. C. 1999, *ApJ*, 513, 252, doi: [10.1086/306858](https://doi.org/10.1086/306858)
- Paczynski, B. 1990, *ApJ*, 363, 218, doi: [10.1086/169332](https://doi.org/10.1086/169332)
- Palla, F., Randich, S., Flaccomio, E., & Pallavicini, R. 2005, *ApJL*, 626, L49, doi: [10.1086/431668](https://doi.org/10.1086/431668)
- Palla, F., & Stahler, S. W. 1999, *ApJ*, 525, 772, doi: [10.1086/307928](https://doi.org/10.1086/307928)
- Pang, X., Tang, S.-Y., Li, Y., et al. 2022, *ApJ*, 931, 156, doi: [10.3847/1538-4357/ac674e](https://doi.org/10.3847/1538-4357/ac674e)
- Portegies Zwart, S. F., McMillan, S. L. W., & Gieles, M. 2010, *ARA&A*, 48, 431, doi: [10.1146/annurev-astro-081309-130834](https://doi.org/10.1146/annurev-astro-081309-130834)
- Rafikov, R. R. 2013, *ApJ*, 774, 144, doi: [10.1088/0004-637X/774/2/144](https://doi.org/10.1088/0004-637X/774/2/144)
- . 2016, *ApJ*, 827, 111, doi: [10.3847/0004-637X/827/2/111](https://doi.org/10.3847/0004-637X/827/2/111)
- Ramstedt, S., Schöier, F. L., Olofsson, H., & Lundgren, A. A. 2008, *A&A*, 487, 645, doi: [10.1051/0004-6361:20078876](https://doi.org/10.1051/0004-6361:20078876)
- Reid, I. N., Gizis, J. E., & Hawley, S. L. 2002, *AJ*, 124, 2721, doi: [10.1086/343777](https://doi.org/10.1086/343777)

- Rein, H., & Liu, S. F. 2012, *A&A*, 537, A128, doi: [10.1051/0004-6361/201118085](https://doi.org/10.1051/0004-6361/201118085)
- Rein, H., & Spiegel, D. S. 2015, *MNRAS*, 446, 1424, doi: [10.1093/mnras/stu2164](https://doi.org/10.1093/mnras/stu2164)
- Salpeter, E. E., & Shapiro, S. L. 1981, *ApJ*, 251, 311, doi: [10.1086/159466](https://doi.org/10.1086/159466)
- Sandage, A. R. 1953, *AJ*, 58, 61, doi: [10.1086/106822](https://doi.org/10.1086/106822)
- Solomon, P. M., Rivolo, A. R., Barrett, J., & Yahil, A. 1987, *ApJ*, 319, 730, doi: [10.1086/165493](https://doi.org/10.1086/165493)
- Soubiran, C., Cantat-Gaudin, T., Romero-Gómez, M., et al. 2018, *A&A*, 619, A155, doi: [10.1051/0004-6361/201834020](https://doi.org/10.1051/0004-6361/201834020)
- Sun, J., Leroy, A. K., Schruba, A., et al. 2018, *ApJ*, 860, 172, doi: [10.3847/1538-4357/aac326](https://doi.org/10.3847/1538-4357/aac326)
- Tagawa, H., Haiman, Z., & Kocsis, B. 2020, *ApJ*, 898, 25, doi: [10.3847/1538-4357/ab9b8c](https://doi.org/10.3847/1538-4357/ab9b8c)
- Tanaka, H., Takeuchi, T., & Ward, W. R. 2002, *ApJ*, 565, 1257, doi: [10.1086/324713](https://doi.org/10.1086/324713)
- Tang, J., Bressan, A., Rosenfield, P., et al. 2014, *MNRAS*, 445, 4287, doi: [10.1093/mnras/stu2029](https://doi.org/10.1093/mnras/stu2029)
- Tarricq, Y., Soubiran, C., Casamiquela, L., et al. 2022, *A&A*, 659, A59, doi: [10.1051/0004-6361/202142186](https://doi.org/10.1051/0004-6361/202142186)
- . 2021, *A&A*, 647, A19, doi: [10.1051/0004-6361/202039388](https://doi.org/10.1051/0004-6361/202039388)
- Wang, L., & Li, X. 2022, *ApJ*, 932, 108, doi: [10.3847/1538-4357/ac6ce6](https://doi.org/10.3847/1538-4357/ac6ce6)
- Zhao, X., Marchesi, S., Ajello, M., et al. 2021, *A&A*, 650, A57, doi: [10.1051/0004-6361/202140297](https://doi.org/10.1051/0004-6361/202140297)