

COMPLEMENTARY ASYMPTOTIC ANALYSIS FOR A MINIMAL RANDOM WALK.

CRISTIAN F. COLETTI, MANUEL GONZÁLEZ-NAVARRETE, AND VÍCTOR HUGO VÁZQUEZ GUEVARA

ABSTRACT. We discuss a complementary asymptotic analysis of the so called minimal random walk. More precisely, we present a version of the almost sure central limit theorem as well as a generalization of the recently proposed quadratic strong laws. In addition, alternative demonstrations of the functional limit theorems will be supplied based on a Pólya urn scheme instead of a martingale approach.

Keywords: Minimal Random Walk, Almost Sure Central Limit Theorem, Pólya urns.

1. INTRODUCTION

In this paper we complement the asymptotic analysis developed in [4] and [5] for the minimal random walk (MRW) proposed by Harbola, Kumar and Lindenberg [8] in 2014. More precisely, we provide the almost sure central limit theorem (ASCLT) for the diffusive and critical regimes as well as the convergence to even moments of a Gaussian random variable, that generalizes the quadratic strong laws previously obtained in [4]. Additionally, we achieve an alternative demonstration for the functional central limit theorems (FCLT) through its relation with an urn of Pólya type in the spirit of [1] which was originally proved in [4]. It must be mentioned that most of our technical display will be based; to some extent, on some side properties provided by [4], situation that will be properly treated in Appendix A.

The rest of the paper is organized as follows: Section 2 presents to the MRW as well as its basic conditional properties. Section 3 provides versions of the ASCLT and convergence to even moments of Gaussian distribution in the diffusive and critical regimes. Section 4 explores the relation of the MRW and a Pólya urn scheme with balls of two colors to provide alternative proofs of the versions of the FCLT obtained in [4]. Four appendices are considered: Appendix A summarizes some properties taken from [4] and provides the conditional moments of the MRW. Appendices B and C deal with the demonstrations corresponding to results of Section 3. Last appendix is devoted to the proof of the FCLT's of Section 4.

2. THE MINIMAL RANDOM WALK

In this section we present the main topic of this paper: The MRW, whose movement and position at time $n \geq 1$ are denoted by X_n and S_n respectively behaves as follow:

- At time $n = 0$ a particle following the MRW is at the origin; i.e., $S_0 = 0$.
- At time $n = 1$, the particle decides to move one unit right with probability $s \in (0, 1)$ or remain at the origin with probability $1 - s$. Then $S_1 = X_1$ has the Bernoulli distribution with parameter s .
- At any time $n \geq 2$, the particle chooses uniformly at random some point in the past; let us say $1 \leq k(n) \leq n - 1$. Then, its following movement; X_n , will be conditioned by the observed value of $X_{k(n)}$ as follows:

On the one hand, if $X_{k(n)} = 0$ then X_n posses a Bernoulli conditional distribution with parameter $q \in (0, 1]$ (It should be underlined that the case $q = 0$ will not be treated in this paper). While, on the other hand, if $X_{k(n)} = 1$ then, the conditional distribution of X_n is also Bernoulli but with parameter $p \in [0, 1]$, where p does not necessarily is equal to q .

- The position of the particle at time $n \geq 1$ satisfies that

$$(1) \quad S_n = S_{n-1} + X_n,$$

with $S_0 = 0$ and $X_n \in \{0, 1\}$.

In order to consider the martingale approach of [4] for the MRW, let (\mathcal{F}_n) be the increasing sequence of σ -algebras $\mathcal{F}_n = \sigma(X_1, \dots, X_n)$; i.e. \mathcal{F}_n represents the knowledge up to time n .

From [8], we know that for $x = 0, 1$, $\mathbb{P}(X_{n+1} = x | \mathcal{F}_n) = 1 - x + (2x - 1)(q + \alpha \frac{S_n}{n})$, then it may be found that the conditional expected movement and position of the particle at any time are; respectively, given by

$$(2) \quad \mathbb{E}[X_{n+1} | \mathcal{F}_n] = q + \alpha \frac{S_n}{n} \quad \text{a.s.}$$

and

$$(3) \quad \mathbb{E}[S_{n+1} | \mathcal{F}_n] = q + \gamma_n S_n \quad \text{a.s.}$$

where, $\alpha := p - q$ and $\gamma_n = \frac{n+\alpha}{n}$.

Let us now consider the discrete time scalar martingale (M_n) , given for $n \geq 0$, by

$$(4) \quad M_n = a_n S_n - q A_n,$$

where $a_1 = 1$ and, for $n \geq 2$

$$(5) \quad a_n = \prod_{k=1}^{n-1} \gamma_k^{-1} = \frac{\Gamma(n)\Gamma(\alpha+1)}{\Gamma(n+\alpha)},$$

where Γ stands for the Euler gamma function. In addition, $A_0 = 0$ and, for $n \geq 1$,

$$(6) \quad A_n = \sum_{k=1}^n a_k.$$

The study of asymptotic properties of the scalar MRW is usually done in 3 regimes:

- (1) If $\alpha < 1/2$ then we are in the diffusive regime.
- (2) If $\alpha = 1/2$ then the regime is the critical one, and
- (3) If $\alpha > 1/2$ then the regime is called superdiffusive.

Provided results in following sections will be; basically of the almost sure central limit type, convergence to even moments of Gaussian distribution and functional central limit theorems.

3. DEEPER RESULTS VIA THE MARTINGALE APPROACH

In the paper [5], it has been proved versions of the central limit theorem and the law of iterated logarithm in the case $\alpha \leq 1/2$, which were complemented by [4], including law of large numbers, functional central limit theorems, an asymptotic analysis of the center of mass and a better understanding of the superdiffusive case $\alpha > 1/2$ by providing the almost sure and in L^2 convergence as well as the Gaussian fluctuations around its limit, in addition the first two moments of this

random variable were explicitly calculated. Furthermore, in [9] it has been considered the case $q = 0$ where a central limit theorem and a law of iterated logarithm were obtained.

In this section, we will display proper versions of the almost sure central limit theorem and of a generalization of the recently proposed quadratic strong laws under the condition $\alpha \leq 1/2$. For this, in all the sequel we will make use of the asymptotic variance

$$(7) \quad \sigma^2 := \frac{q(1-p)}{(1-\alpha)^2}$$

3.1. The diffusive regime. We initiate the asymptotic analysis by dealing with the almost sure central limit theorem (ASCLT) for the diffusive regime ($\alpha < 1/2$).

Theorem 3.1. *If $\alpha < 1/2$ then, we have the following almost sure convergence of measures*

$$(8) \quad \frac{1}{\log n} \sum_{k=1}^n \frac{1}{k} \delta_{\sqrt{k} \left(\frac{S_k}{k} - \frac{q}{1-\alpha} \right)} \implies N \left(0, \frac{\sigma^2}{1-2\alpha} \right)$$

where σ^2 is given in (7).

In a complementary way, we provide the corresponding convergence of moments, which generalizes the quadratic strong law found in [4].

Theorem 3.2. *If $\alpha < 1/2$, then we have; for $r \geq 1$, the following almost sure convergence*

$$(9) \quad \lim_{n \rightarrow \infty} \frac{1}{\log n} \sum_{k=1}^n k^{r-1} \left(\frac{S_k}{k} - \frac{q}{1-\alpha} \right)^{2r} = \frac{\sigma^{2r} (2r)!}{2^r r! (1-2\alpha)^r}$$

3.2. The critical regime. We will investigate now, analogue results to those presented in previous subsection for the critical regime, starting with the ASCLT.

Theorem 3.3. *If $\alpha = 1/2$, then we have that almost surely*

$$(10) \quad \frac{1}{\log \log n} \sum_{k=2}^n \frac{1}{k \log k} \delta_{\sqrt{\frac{k}{\log k}} \left(\frac{S_k}{k} - 2q \right)} \implies N(0, 4q(1-p))$$

Next theorem establishes the convergence to even moments of Gaussian distribution

Theorem 3.4. *If $\alpha = 1/2$, then the following almost sure convergence holds*

$$(11) \quad \lim_{n \rightarrow \infty} \frac{1}{\log \log n} \sum_{k=2}^n \left(\frac{1}{\log k} \right)^{r+1} k^{r-1} \left(\frac{S_k}{k} - 2q \right)^{2r} = \frac{(4q(1-p))^r (2r)!}{2^r r!}$$

where $r \geq 1$.

4. RELATION OF THE MRW WITH PÓLYA URNS

In this section, we will perform a counterpart display of the ideas exposed in [1] for the elephant random walk. Hence, in order to establish a relationship between the MRW and an urn of Pólya type, let us consider

- An urn with balls of two colors: red and blue.
- At $n = 1$ we put in a blue one with probability (w.p.) s .
- At $n > 1$ we choose uniformly at random one ball, observe its color, put it back and if it is red, then we put another red w.p. $1 - q$.
- If it is blue then we put another blue w.p. p .

Hence, S_n is the number of blue balls after n extractions which evolves as the MRW. Furthermore, we obtain the so-called mean replacement matrix for the earlier urn associated with the MRW:

$$(12) \quad A := \begin{pmatrix} 1-q & 1-p \\ q & p \end{pmatrix}$$

whose eigenvalues are $\lambda_1 := 1$ and $\lambda_2 := \alpha$ with $\lambda_2 \leq \lambda_1$.

In succeeding subsection, the theory for generalized Pólya urns established in [7] will be employed in order to find alternative demonstrations for the functional central limit theorems in both the diffusive and critical regimes as well as the almost sure convergence in the superdiffusive regime. In all the sequel $D([0, \infty[)$ denotes to the Skorokhod space of right-continuous functions with left-hand limits.

4.1. Limit theorems via the Pólya urns scheme. Subsequent results were obtained in [4] via the martingale approach considered in sections 2 and 3, nevertheless in Appendix D the link between the MRW and the previously urn scheme will be employed in order to prove them.

Theorem 4.1. *If $\alpha < 1/2$ then, we have the distributional convergence in $D([0, \infty[)$,*

$$\left(\sqrt{n} \left(\frac{S_{\lfloor nt \rfloor}}{\lfloor nt \rfloor} - \frac{q}{1-a} \right), t \geq 0 \right) \Longrightarrow (W_t, t \geq 0)$$

where $(W_t, t \geq 0)$ is a real-valued centered Gaussian process starting at the origin with covariance given, for all $0 < s \leq t$, by

$$\mathbb{E}[W_s W_t] = \frac{\sigma^2}{(1-2a)t} \left(\frac{t}{s} \right)^a.$$

In particular, we have the asymptotic normality

$$\sqrt{n} \left(\frac{S_n}{n} - \frac{q}{1-a} \right) \xrightarrow{\mathcal{L}} N \left(0, \frac{\sigma^2}{1-2a} \right)$$

Theorem 4.2. *If $\alpha = 1/2$ then, we have the distributional convergence in $D([0, \infty[)$,*

$$\left(\sqrt{\frac{n^t}{\log n}} \left(\frac{S_{\lfloor n^t \rfloor}}{\lfloor n^t \rfloor} - 2q \right), t \geq 0 \right) \Longrightarrow (2\sqrt{q(1-p)} B_t, t \geq 0)$$

where $(B_t, t \geq 0)$ is a standard Brownian motion.

In particular, we have the central limit theorem

$$\sqrt{\frac{n}{\log n}} \left(\frac{S_n}{n} - 2q \right) \xrightarrow{\mathcal{L}} N(0, 4q(1-p))$$

Theorem 4.3. *If $\alpha > 1/2$ then, we have the almost sure convergence,*

$$n^{1-a} \left(\frac{S_n}{n} - \frac{q}{1-a} \right) \Longrightarrow L$$

where L is a non-degenerated random variable and whose two first moments are calculated in Theorem 2.6 of [4].

ACKNOWLEDGMENTS

Cristian F. Coletti was supported by grant #2017/10555-0 São Paulo Research Foundation (FAPESP). Manuel González-Navarrete was partially supported by Fondecyt Iniciación 11200500.

APPENDIX A
INITIAL ANALYSIS

In this Appendix, we will state some useful facts on the MRW in order to achieve Theorems 3.1 - 3.4:

For each $n \geq 1$ and $k \geq 1$, the definition of the MRW ensures that, $X_n \leq 1$, then $0 \leq S_n \leq n$ and $X_n = X_n^k$. Moreover,

$$\mathbb{E} [X_{n+1}^k | \mathcal{F}_n] = q + \alpha \frac{S_n}{n} \quad \text{a.s.}$$

and

$$\mathbb{E} [S_{n+1}^{2k} | \mathcal{F}_n] = S_n^{2k} + \left[q + \alpha \frac{S_n}{n} \right] \sum_{j=1}^{2k} \binom{2k}{j} S_n^{2k-j}.$$

In addition, the martingale (M_n) may be rewritten as $M_n = \sum_{k=1}^n a_k \varepsilon_k$, where $\varepsilon_{n+1} = X_{n+1} - (q + \alpha \frac{S_n}{n})$. It also holds that

$$\mathbb{E} [\varepsilon_{n+1}^k | \mathcal{F}_n] = \sum_{j=0}^{k-2} \binom{k}{j} \left(q + \alpha \frac{S_n}{n} \right)^{j+1} (-1)^j + (-1)^{k-1} \left(q + \alpha \frac{S_n}{n} \right)^k (k-1) < \infty.$$

Therefore, Theorems 2.1 and 2.3 of [4] guide us to

$$(A.1) \quad \lim_{n \rightarrow \infty} \mathbb{E} [\varepsilon_{n+1}^2 | \mathcal{F}_n] = \sigma^2 \quad \text{a.s.},$$

finally, a very useful consequence of Lemma B.1 of [3] which will be repeatedly used is

$$(A.2) \quad \left| \frac{A_n}{na_n} - \frac{1}{1-\alpha} \right| \sim \frac{1}{n^{1-\alpha}}.$$

In addition, by continuing the notation of [4], let us consider the following entities for each $n \geq 1$:

$$v_n := \sum_{k=1}^n a_k^2.$$

and the explosion coefficient associated with martingale (M_n)

$$f_n := a_n^2 / v_n$$

.

APPENDIX B
ASCLT FOR THE DIFFUSIVE REGIME

Proof of theorem 3.1 and 3.2: For this regime, it has been previously obtained [4] that, as $n \rightarrow \infty$:

$$(B.1) \quad \frac{v_n}{n^{1-2\alpha}} \rightarrow \ell := \frac{\Gamma^2(\alpha+1)}{1-2\alpha}$$

and

$$(B.2) \quad f_n \sim \frac{1-2\alpha}{n} \rightarrow 0.$$

Additionally, (A.2) implies the following useful relation:

$$(B.3) \quad \frac{M_k}{\sqrt{v_{k-1}}} \sim \sqrt{\frac{1-2\alpha}{k}} \left(S_k - k \frac{q}{1-\alpha} \right)$$

In order to utilize Theorem A.1 of [12] for proving Theorem 3.1, we note from last statements that for $\eta > 0$

$$(B.4) \quad \sum_{k=1}^{\infty} \frac{1}{v_k} \mathbb{E} [|\Delta M_k|^2 \mathbb{I}_{|\Delta M_k| \geq \eta \sqrt{v_k}} | \mathcal{F}_{k-1}] \leq \frac{1}{\eta^2} \sum_{k=1}^{\infty} \frac{1}{v_k^2} \mathbb{E} [|\Delta M_k|^4 | \mathcal{F}_{k-1}]$$

$$(B.5) \quad \leq \sup_{k \geq 1} \mathbb{E} [\varepsilon_k^4 | \mathcal{F}_{k-1}] \frac{1}{\eta^2} \sum_{k=1}^{\infty} \frac{a_k^4}{v_k^2} \sim \frac{\sup_{k \geq 1} \mathbb{E} [\varepsilon_k^4 | \mathcal{F}_{k-1}]}{\eta^2} \sum_{k=1}^{\infty} \frac{(1-2\alpha)^2}{k^2} < \infty.$$

On the same fashion, it may be found that

$$\sum_{k=1}^{\infty} \frac{1}{v_k^2} \mathbb{E} [|\Delta M_k|^4 \mathbb{I}_{|\Delta M_k| \leq \sqrt{v_k}} | \mathcal{F}_{k-1}] \leq \sum_{k=1}^{\infty} \frac{a_k^4}{v_k^2} \mathbb{E} [\varepsilon_k^4 | \mathcal{F}_{k-1}] < \infty.$$

Hence, Theorem A.1 of [12] let us conclude that

$$(B.6) \quad \frac{1}{\log v_n} \sum_{k=1}^n f_k \delta_{M_k / \sqrt{v_{k-1}}} \Rightarrow G \quad \text{a.s.}$$

where G is the Gaussian distribution with mean zero and variance σ^2 . Finally, convergence (B.1), (B.2) and relation (B.3) imply

$$(B.7) \quad \frac{1}{(1-2\alpha) \log n} \sum_{k=1}^n \frac{(1-2\alpha)}{k} \delta_{\sqrt{1-2\alpha} \left(\frac{S_k}{k} - \frac{q}{1-\alpha} \right) \sqrt{k}} \Rightarrow G \quad \text{a.s.}$$

which conduces us to convergence (8).

For obtaining Theorem 3.2, we may observe from (A.1) and (A.2), together with Theorem 3 of [2] (or Theorem A.2 of [12]) that; for $r \geq 1$:

$$(B.8) \quad \frac{1}{\log n} \sum_{k=1}^n f_k \left(\frac{M_k^2}{v_{k-1}} \right)^r \rightarrow \frac{\sigma^{2r} (2r)!}{2^r r!},$$

however, definition of M_n and f_n , convergences (B.1) and (B.2) lead us to

$$(B.9) \quad \frac{1}{\log n} \sum_{k=1}^n \frac{1}{k^{r+1}} \left(S_k - q \frac{A_k}{a_k} \right)^{2r} \rightarrow \frac{\sigma^{2r} (2r)! (1-2\alpha)}{2^r r! (1-2\alpha)^{r+1}},$$

that implies

$$(B.10) \quad \frac{1}{\log n} \sum_{k=1}^n k^{r-1} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^{2r} \rightarrow \frac{\sigma^{2r} (2r)!}{2^r r! (1-2\alpha)^r}.$$

Nevertheless, we have that

$$(B.11) \quad \frac{1}{\log n} \sum_{k=1}^n k^{r-1} \left(\frac{S_k}{k} - \frac{q}{1-\alpha} \right)^{2r} = \frac{1}{\log n} \sum_{k=1}^n k^{r-1} \sum_{i=0}^{2r} \binom{2r}{i} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^i \left(q \frac{A_k}{ka_k} - \frac{q}{1-\alpha} \right)^{2r-i}.$$

However, it may be noticed from (A.2); for $0 \leq i < 2r$, that

$$\begin{aligned}
& \frac{1}{\log n} \sum_{k=1}^n k^{r-1} \binom{2r}{i} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^i \left(q \frac{A_k}{ka_k} - \frac{q}{1-\alpha} \right)^{2r-i} \\
& \leq \frac{C}{\log n} \sqrt{\sum_{k=1}^n k^{i-1} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^{2i}} \sqrt{\sum_{k=1}^n k^{2r-i-1} \left(q \frac{A_k}{ka_k} - \frac{q}{1-\alpha} \right)^{4r-2i}} \\
& \sim \frac{C}{\log n} \sqrt{\sum_{k=1}^n k^{i-1} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^{2i}} \sqrt{\sum_{k=1}^n k^{2r-i-1} \left(\frac{1}{k^{2(1-\alpha)}} \right)^{2r-i}} \\
& \hspace{20em} \rightarrow 0,
\end{aligned}$$

since $2(1-\alpha)(2r-1) - (2r-i-1) > 1$. Hence, we infer that

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \frac{1}{\log n} \sum_{k=1}^n k^{r-1} \left(\frac{S_k}{k} - \frac{q}{1-\alpha} \right)^{2r} \\
& = \lim_{n \rightarrow \infty} \frac{1}{\log n} \sum_{k=1}^n k^{r-1} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^{2r} \\
& \hspace{15em} = \frac{\sigma^{2r}(2r)!}{2^r r! (1-2\alpha)^r}.
\end{aligned}$$

APPENDIX C

ASCLT FOR THE CRITICAL REGIME

Proof of theorems 3.3 and 3.4: On this regime, it is useful to highlight that [4]

$$(C.1) \quad \frac{v_n}{\log n} \rightarrow \frac{\pi}{4},$$

which leads us to ensure that

$$(C.2) \quad f_n \sim \frac{1}{n \log n} \rightarrow 0,$$

and also that $\sum_{k=1}^{\infty} \frac{a_k^2}{v_n^2} < \infty$. Therefore, from analogue arguments to those exposed in Appendix B we conclude that hypothesis of the ASCLT expressed in Theorem A.1 of [12] are fulfilled. Hence, from such ASCLT, relation (A.2) and convergence (C.1) we infer (10).

In order to demonstrate Theorem 3.4, we may note; from similar argumentation than in Appendix B, that (A.1) and (A.2), together with Theorem 3 of [2] and convergences (C.1) and (C.2); for $r \geq 1$, that

$$(C.3) \quad \frac{1}{\log \log n} \sum_{k=2}^n k^{r-1} \left(\frac{1}{\log k} \right)^{r+1} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^{2r} \rightarrow \frac{\sigma^{2r}(2r)!}{2^r r!}.$$

However, we may observe that

$$\begin{aligned}
\text{(C.4)} \quad & \frac{1}{\log \log n} \sum_{k=2}^n k^{r-1} \left(\frac{1}{\log k} \right)^{r+1} \left(\frac{S_k}{k} - 2q \right)^{2r} \\
&= \frac{1}{\log \log n} \sum_{k=2}^n k^{r-1} \left(\frac{1}{\log k} \right)^{r+1} \sum_{i=0}^{2r} \binom{2r}{i} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^i \left(q \frac{A_k}{ka_k} - 2q \right)^{2r-i}.
\end{aligned}$$

For $0 \leq i < 2r$, we observe that Cauchy-Schwarz inequality, relation (A.2) and convergence (C.3) imply that

$$\frac{1}{\log \log n} \sum_{k=2}^n k^{r-1} \left(\frac{1}{\log k} \right)^{r+1} \binom{2r}{i} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^i \left(q \frac{A_k}{ka_k} - 2q \right)^{2r-i} \rightarrow 0$$

because

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \left(\frac{1}{\log k} \right)^{2r-i+1} \frac{1}{k} < \infty$$

since $0 \leq i < 2r$ and $r \geq 1$.

All this leads us to find out that

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \frac{1}{\log \log n} \sum_{k=2}^n k^{r-1} \left(\frac{1}{\log k} \right)^{r+1} \left(\frac{S_k}{k} - 2q \right)^{2r} \\
&= \lim_{n \rightarrow \infty} \frac{1}{\log \log n} \sum_{k=2}^n k^{r-1} \left(\frac{1}{\log k} \right)^{r+1} \left(\frac{S_k}{k} - q \frac{A_k}{ka_k} \right)^{2r} \\
&= \frac{\sigma^{2r} (2r)!}{2^r r!}.
\end{aligned}$$

APPENDIX D

THE FUNCTIONAL CENTRAL LIMIT THEOREM BASED ON A PÓLYA URN SCHEME

In the sequel, we will employ the notation of [7] in an attempt to facilitate consultation of the theory contained therein and be able to expand the details of this Appendix; if it is necessary. Hence, at this setting, it is not hard to see that conditions (A1) – (A6) (of that paper) hold for the MRW and his eigenvalues. Additionally, we find out that right and left eigenvectors given by

$$v_1 = \frac{q}{1-\alpha} \left(\frac{1-p}{q}, 1 \right)^t, \quad v_2 = \frac{1}{2} (-1, 1)^t$$

and

$$u_1 = (1, 1), \quad u_2 = \frac{2(p-1)}{\alpha-1} \left(\frac{q}{p-1}, 1 \right);$$

respectively, are such that $u_1 \cdot v_1 = u_2 \cdot v_2 = 1$. Finally, we denote by $W_n = (P_n, S_n)$ to the composition of the urn at stage $n \geq 1$ where P_n states for the number of red balls and S_n is the number of blue ones.

Proof of theorem 4.1 (the diffusive regime):

For this regime we will make use of Theorem 3.31 i) of [7] that asserts that

$$\text{(D.1)} \quad (n^{-1/2} (W_{\lfloor nt \rfloor} - nt v_1), t \geq 0) \implies (V(t), t \geq 0).$$

where $(V(t), t \geq 0)$ is a centered Gaussian vector process such that $V(0) = (0, 0)$ and with covariance structured given for $0 < s \leq t$ by (see Remark 5.7 of [7])

$$(D.2) \quad \mathbb{E} [V(s)V^t(t)] = s\Sigma_I e^{\log(t/s)A^t},$$

where Σ_I was defined in equation (2.15) of [7] and given for the MRW by

$$(D.3) \quad \Sigma_I = \frac{q(1-p)}{(1-2\alpha)(1-\alpha)^2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}.$$

Also, we have that

$$\begin{pmatrix} t \\ s \end{pmatrix}^{A^t} = P^{-1} \begin{pmatrix} t \\ s \end{pmatrix}^\alpha I_2 P$$

where I_2 is identity matrix of order 2 and

$$(D.4) \quad P = I_2 - v_1 u_1^t = \frac{1}{1-\alpha} \begin{pmatrix} q & p-1 \\ -q & 1-p \end{pmatrix}.$$

Hence, (D.2) implies

$$(D.5) \quad \mathbb{E} [V(s)V^t(t)] = s \begin{pmatrix} t \\ s \end{pmatrix}^\alpha \frac{q(1-p)}{(1-2\alpha)(1-\alpha)^2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix},$$

On the other hand, we observe that $S_n = (0, 1)W_n^t$, which together with the continuous mapping theorem leads us to the conclusion of Theorem 4.1.

Proof of theorem 4.2 (the critical regime):

On this regime, Theorem 3.31 ii) of [7] conduces us to

$$\left(\sqrt{\frac{1}{n^t \log n}} (W_{[n^t]} - n^t v_1), t \geq 0 \right) \Longrightarrow (V(t), t \geq 0),$$

where $(V(t), t \geq 0)$ is a centered Gaussian vector process such that $V(0) = (0, 0)$ and with covariance structured given for $0 < s \leq t$ by (see equation (3.27) of [7])

$$(D.6) \quad \mathbb{E} [V(s)V^t(t)] = 4sq(1-p) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}.$$

By considering the same applied transformation as in the diffusive regime, the continuous mapping theorem implies the conclusion of Theorem 4.2.

Proof of Theorem 4.3 (the superdiffusive regime): This theorem follows essentially from Theorem 3.24 of [7]. More precisely, since at this regime the lowest eigenvalue is greater than the half of the largest one (that is $\alpha > 1/2$), we have that, almost surely

$$n^{-\alpha} (W_n - n v_1) \rightarrow \hat{W},$$

where \hat{W} is a random vector with values in $\{w \in \mathbb{R}^2 / w = \lambda(-1, 1) \text{ for some } \lambda \in \mathbb{R} - \{0\}\}$, therefore (again) the continuous mapping theorem conduces us to conclusion of Theorem 4.3.

REFERENCES

- [1] E. Baur and J. Bertoin. Elephant Random Walks and their connection to Pólya-type urns. *Phys. Rev. E* 94, 052134 (2016).
- [2] B. Bercu. On the convergence of moments in the almost sure central limit theorem for martingales with statistical applications. *Stochastic Process. Appl.* 111, 1, pp. 157–173 (2004).
- [3] B. Bercu. A martingale approach for the elephant random walk, *J. Phys. A: Math. Theor.* 51 015201 (2018).
- [4] B. Bercu and V.H. Vázquez Guevara. Further results on the minimal random walk, *J. Phys. A: Math. Theor.* 55 415001 (2022).
- [5] C.F. Coletti, R. J. Gava and L. R. de Lima. Limit theorems for a minimal random walk model, *J. Stat. Mech.* 083206 (2019).
- [6] M. Duflo. *Random Iterative Models*, Springer Verlag, Berlin (1997).
- [7] S. Janson. Functional limit theorems for multitype branching processes and generalized Pólya urns. *Stoch. Proc. Appl.* 110(2), 177–245 (2004).
- [8] U. Harbola, N. Kumar and K. Lindenberg. Memory-induced anomalous dynamics in a minimal random walk model. *Phys. Rev. E* 90, 022136 (2014).
- [9] T. Miyazaki and M. Takei. Limit Theorems for the ‘Laziest’ Minimal Random Walk Model of Elephant Type. *J Stat Phys* 181, 587–602 (2020).
- [10] G. M. Schütz and S. Trimper. Elephants can always remember: Exact long-range memory effects in a non-Markovian random walk. *Phys. Rev. E* 70, 045101 (2004).
- [11] W.F. Stout. Almost sure convergence, *Probability and Mathematical Statistics*, Vol. 24, Academic Press, New York-London (1974).
- [12] V. H. Vázquez Guevara. On the almost sure central limit theorem for the Elephant Random Walk. *J.Phys. A: Math. Theor.* 52 475201 (2019).

Email address: cristian.coletti@ufabc.edu.br

Email address: manuel.gonzaleznavarrete@ufrontera.cl

Email address: vvazquez@fcfm.buap.mx

UNIVERSIDADE FEDERAL DO ABC, CENTER OF MATHEMATICS, COMPUTATION AND COGNITION, SAO PAULO, BRAZIL.

UNIVERSIDAD DE LA FRONTERA, DEPARTAMENTO DE MATEMÁTICA Y ESTADÍSTICA, TEMUCO, CHILE.

BENEMÉRITA UNIVERSIDAD AUTÓNOMA DE PUEBLA, FACULTAD DE CIENCIAS FÍSICO, PUEBLA, MÉXICO.