

Fluctuation of a bilayer composed by surfactants

Shunta Kikuchi and Hiroshi Watanabe*

*Faculty of Science and Technology, Department of Applied Physics and
Physico-Informatics, Keio University, Yokohama 223-8522, Japan*

We investigated the properties of lipid bilayer fluctuations using molecular dynamics. We modeled the surfactant as a diatomic molecule, constructed a bilayer in the solvent, and observed the Fourier spectrum of its fluctuations. The results showed that q^4 behavior was dominant at the high temperature, whereas a crossover from q^4 to q^2 was observed at the low temperature. This behavior is complementary to the results for monolayer membranes, where q^2 behavior was dominant when the interfacial tension was high, whereas a crossover from q^2 to q^4 was observed when the interfacial tension was lowered [S. Kikuchi and H. Watanabe, *J. Chem. Phys.*, **158** 12 (2023)]. These results suggest that the restoring force is dominated by elasticity at high temperatures and by interfacial tension at low temperatures. We also observed radial distribution functions to investigate the structure of surfactants in lipid bilayers. The structure of the surfactant depends on temperature in the direction horizontal to the interface, while it does not depend on temperature in the direction perpendicular to the interface. This result suggests that lipid bilayers increase fluctuations while maintaining the membrane structure in high temperature.

1. Introduction

The lipid bilayer consists of two layers of lipids, with the hydrophilic heads facing outward and the hydrophobic tails facing inward.^{1,2)} Since the lipid bilayer is a fundamental structure forming the basis of the cell membrane, understanding lipid bilayers is essential both in biology and drug delivery.³⁾ However, describing lipid bilayers using the continuum representation is challenging since the atoms' behavior significantly affects the membrane's properties. Recently, developments of computational power allow us to perform all-atom simulations of membranes by molecular dynamics (MD) simulations.^{4,5)} The numerical studies investigated the generation process of the vesicles,^{6,7)} the relation between the fluidity and

*hwatanabe@appi.keio.ac.jp

the molecular orientation,^{8,9)} and the drug permeation through lipid bilayers.¹⁰⁾

One of the fundamental properties of the membrane is its elasticity. For example, when the cell is deformed, the elasticity of the bilayer maintains its shape.¹¹⁾ In the ventricular assist devices, the local pressure gradient causes hemolysis with damage in the red blood cells.^{12,13)} In addition to the deformation, the lipid bilayers are affected by the blood flow and the thermal fluctuations.^{14,15)} In the previous models of lipid bilayers with realistic parameters, the elasticity was dominant for the restoring force. However, in the case of the monolayer, the main restoring force of the membrane was found to originate from the interfacial tension with finite interfacial tension, whereas the crossover from the interfacial tension to the elasticity was observed as the interfacial tension decreased.¹⁶⁾ These results implied that the parameters at the interface affected the fluctuations, and therefore, it is expected that crossovers similar to those observed in monolayers can be observed in bilayers. On the other hand, there can be differences in behavior derived from the structure of monolayers and bilayers, but how they differ needs to be clarified. In this paper, we aim to investigate the similarities and differences in the behavior of monolayers and bilayers by molecular dynamics.

The rest of the paper is organized as follows. In section 2, we describe the method. The results are described in Sec. 3. Section 4 is devoted to the summary and discussion.

2. Method

We adopted the Lennard-Jones (LJ) and the Weeks-Chalender-Andresen (WCA) potential as interactions of lipid molecules. The LJ potential is described by

$$\phi(r) = \begin{cases} 4\varepsilon \left\{ \left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right\} & (r < r_c), \\ 0 & (r \geq r_c), \end{cases} \quad (1)$$

where σ is the atomic diameter, ε is the well depth and r is the distance between the atoms. The cutoff length is denoted by r_c . In the following, we adopt the reduced unit such that the σ , ε , and k_B are unity. We set $r_c = 3.0$ for the LJ potential and $r_c = 2^{1/6}$ for the WCA potential. The LJ interaction represents the attraction between atoms while the WCA represents the repulsion.

In this paper, the solvent molecules, the hydrophilic and hydrophobic molecules of the lipid molecules were represented as one atom, respectively. The lipid molecule is modeled as a diatomic molecule with a hydrophobic atom and a hydrophilic atom bonded together. In order to investigate the fluctuations of the lipid bilayer, we prepare the bilayer separating the solvent as shown in Fig. 1 (a). The lipid molecule are placed at the liquid-liquid interface and act as the amphiphilic molecules. The z axis is the normal to the interface. As shown in

Fig. 1 (b), the solvent, the hydrophili, and the hydrophobic atoms are referred to as A, B, and C atoms, respectively. We set the interaction between the solvent atoms and the hydrophilic atoms (A-B) as the LJ potential, the interaction between A-C as the WCA potential, and the interaction between B-C as the WCA potential with the well depth $\varepsilon = 1.05$. The interaction between the same atoms is set as the LJ potential. We adopted the harmonic potential as the bond interaction in the surfactants which is

$$V(r) = K(r - l_0)^2 \quad (2)$$

where l_0 is the bond length and K is the spring constant. The bond length is set to $l_0 = 1.50$ and the spring constant is set to $K = 200$ throughout this study. Note that the interactions between atoms of the same surfactant molecule are not considered.




Fig. 1. (Color online) (a) The schematic view of the simulation box. The bilayer membrane separates the solvent. The interface is normal to the z -axis. The solvent, hydrophilic, and hydrophobic atoms are referred to as A, B, and C atoms, respectively. (b) The typical snapshot of the simulations. The interface fluctuates.

The typical snapshot is shown in Fig. 1 (b). We used Visual Molecular Dynamics (VMD) for the visualization.¹⁷⁾ The total number of the atoms was set to $N = 1\,600\,000$. The simulations were performed using the Large-scale Atomic/Molecular Massively Parallel Simulator (LAMMPS).¹⁸⁾ The temperature and pressure in the simulation box are maintained by the Nosé-Hoover thermostat and the Andersen barostat.^{19–21)} The pressure was set to $P = 0.20$ and the temperature range $0.65 \leq T \leq 0.80$ was studied. The velocity Verlet algorithm is used for the time integration with a time step of 0.005.²²⁾ Simulations were performed up to 3 000 000 steps.

After the system had equilibrated, we calculated the spectra of membrane fluctuations and the radial density function (RDF) $g(r)$ of atoms. The area near the interface ($80 < z < 120$) of the simulation box was divided into $G \times G$ sections in the $x-y$ direction. The center of gravity of the lipids in each rectangle was defined as the interface position $h(x, y)$, where $G = 30$ in

this study. Then we obtain the spectrum $h(q)$ by the Fourier transformation of $h(x, y)$, where q is the wavenumber. The Fourier transform of the height $h(x, y)$ is given by

$$h(q_x, q_y) = \frac{1}{G} \sum_{x=0}^{G-1} \sum_{y=0}^{G-1} h(x, y) \exp\{-i2\pi(q_x x + q_y y)/G\}. \quad (3)$$

After calculating the $h(q_x, q_y)$, we obtain the $h(q)$ by averaging $h(q_x, q_y)$ for the angular distribution, where $q = \sqrt{q_x^2 + q_y^2}$. The spectra of fluctuations are denoted by⁴⁾

$$|h(q)|^2 = \frac{k_B T}{\gamma q^2 + \kappa q^4}. \quad (4)$$

where γ is the interface tension, κ is the elasticity of the interface, $k_B = 1$ is the Boltzmann constant, respectively. Equation (4) means that the q^2 component of the membrane fluctuation originates from the interfacial tension and the q^4 components from the elasticity.

3. Results

The snapshots of the simulations are shown in Fig. 2. The membrane was flat in low temperature ($T = 0.70$), while the membrane fluctuated significantly in high temperature ($T = 0.80$). Therefore, it is expected that the fluctuation properties of the interface depend on temperature.




Fig. 2. (Color online) The snapshots of the simulations in (a) low temperature ($T = 0.70$) and (b) high temperature ($T = 0.80$). Only surfactants are shown for the visibility. While the membrane of the surfactants is almost flat in low temperature, it is fluctuating in high temperature.

The density profile of the solvent atoms is shown in Fig. 3. At the high temperatures, sol-

vent densities were finite near the interface, and solvent appeared to seep into the membrane. However, this behavior originated from membrane fluctuations and not the seeping of solvent molecules, as will be shown later.




Fig. 3. The solvents' density profile. The density is almost zero at low temperature ($T = 0.70$). Therefore, the membrane completely separates the interface. The density of the solvents at the interface increases as the temperature increases.

In the previous work on the fluctuations of the monolayer membrane, the restoring force originated from the interfacial tension when the interfacial tension was finite, whereas the elasticity became the dominant force when the interfacial tension was virtually zero.¹⁶⁾ This investigation suggests that the fluctuation's nature also changes with the membranes' interfacial tension. To study the relation between the surface tension and the fluctuation of the membrane, we investigated the temperature dependence of the interface tension, which is shown in Fig. 4. At the low temperature ($T = 0.70$), when the membrane was almost flat, the interfacial tension was finite, while at the high temperature ($T = 0.80$), when the membrane was highly fluctuating, the interfacial tension was virtually zero.

In the case of the monolayer membrane, the crossover from q^2 to q^4 was observed as the interface tension decreased, which reflects that the restoring force was changed from the interfacial tension to the elasticity. Therefore, we investigated the fluctuation spectra of the membrane to identify its restoring force. The temperature dependence of the fluctuation spectra is shown in Fig. 5. While the crossover from q^2 to q^4 was observed at the low temperature, only q^4 behaviors were observed at high temperatures. To identify the power exponents of the




Fig. 4. The temperature dependence of the interfacial tension. The interfacial tension decreases as the temperature increases and becomes virtually zero at $T = 0.80$.

spectra, $q^2|h(q)|$ and $q^4|h(q)|$ are plotted as functions of q in Fig. 6.²³⁾ The q^2 behavior exists at low temperatures, but the q^4 behavior becomes dominant at high temperatures. These results indicate that q^2 behavior was observed when the interfacial tension was finite, and only q^4 behaviors were observed when the interfacial tension was virtually zero. The change in the fluctuation can be originated from the change in the crystalline structure of the membrane. For example, the membrane can be stiff at a low temperature since the surfactants are aligned in an alternating crystal structure. Still, the membrane becomes soft since the crystalline structure melts as temperature increases. To investigate the crystal structure, we calculated the RDF of the membrane. The RDF between solvents (A-A) is shown in Fig. 7 (a), and the RDF between solvent and hydrophobic atoms (A-C) is shown in Fig. 7 (b). The temperature dependence was observed in the RDF between A-A. This result is consistent with the general solvent. On the other hand, the RDF between A-C was almost zero, regardless of the temperature. This behavior indicates that the solvent atoms did not penetrate the membrane, and the interaction between the solvent and the hydrophobic atoms did not change the interfacial tension. Figure 8 (a) shows the RDF between the hydrophobic atoms (B-B), and Fig. 7 (a) shows the RDF between B-C. The apparent temperature dependence was observed in the RDFs between the same types of atoms. These results imply that the horizontal fluidity in the membrane increases as the temperature increases. On the other hand, the temperature dependence was not observed in the RDF between the hydrophilic atoms and the hydrophobic

atoms. This result shows that the structure of the membrane did not change with temperature. Consequently, the membrane increases its fluctuation, keeping its microscopic structure as the temperature increases.




Fig. 5. The temperature dependence of the spectra $|h(q)|^2$. The Decimal logarithm is taken for both axes. (a) The low temperature ($T = 0.70$). q^4 was only observed. (b) The high temperature ($T = 0.75, 0.80$). The crossover from q^2 to q^4 was observed.




Fig. 6. Graphs of (a) $q^2|h(q)|^2$ and (b) $q^4|h(q)|^2$. The Decimal logarithm is taken for both axes. At the low temperature ($T = 0.70$), $q^2|h(q)|^2$ was constant in the small wavenumber region. This behavior indicates that q^2 is dominant. With the high temperature ($T = 0.80$), $q^4|h(q)|^2$ was constant in the small wavenumber region. This behavior indicates that q^4 is dominant.

4. Summary and Discussion

In this paper, we investigated the fluctuations of bilayer membranes using MD simulations. The power spectra of the membrane fluctuations changed with temperature. Since the membrane separated the solvents, the interfacial tension was slightly affected, and q^4 behavior was only observed with low temperatures. However, when the temperature was low




Fig. 7. Graphs of the RDF $g(r)$. (a) The RDF between the solvent atoms (A-A). The temperature dependence was seen. (b) The RDF between the solvent atoms and the hydrophobic atoms (A-C). The RDF was almost zero, which indicates that the solvent did not penetrate the membrane.








Fig. 8. Graphs of the RDF $g(r)$. (a) The RDF between the hydrophilic atoms (B-B). There exists the temperature dependence. (b) The RDF between the hydrophilic atoms and the hydrophobic atoms (B-C). The RDFs exhibit virtually no temperature dependence.

enough to see the effect of the interfacial tension, the crossover from q^4 to q^2 behaviors was observed.

Here, we discuss the relation between the monolayer and the bilayer membranes. Table I summarizes the relation between the fluctuations and the interfacial tension, combined with the results of the monolayers.¹⁶⁾ In the previous study, q^2 behavior was observed, and the interfacial tension was dominant for the fluctuations of monolayer when the interfacial tension was finite, whereas the crossover from q^2 to q^4 was observed, which indicates that the elasticity was dominant when the interfacial tension was almost zero. In this paper, increasing the temperature in the bilayer membrane corresponds to reducing the interfacial tension, which is complementary to the results for the monolayer.

We also observed RDFs to investigate the relationship between membrane fluctuation and microscopic atomic interactions. We found that the interaction between the same types

Table I. Table of fluctuations in monolayer and bilayer membranes. In the monolayer membrane, q^2 behavior was only observed in the region where the interfacial tension was finite, and the crossover from q^2 to q^4 was observed when the interfacial tension was almost zero. In the bilayer membrane, in contrast, the crossover from q^2 to q^4 in the region where the interfacial tension was finite, while q^4 behavior was only observed when the interfacial tension was almost zero.

	q^2 only	Crossover q^2 to q^4	q^4 only
Monolayer	 $\gamma \neq 0$	 $\gamma = 0$	
Bilayer		 $\gamma \neq 0$	 $\gamma = 0$

of atoms depends on the temperature. In contrast, the interaction between different surfactant atoms did not depend on temperature. These results imply that the lipid bilayers are anisotropic in their interactions, and the anisotropy alters the structure of the membrane.

The effect of the asymmetry on the fluctuations is the subject of future work. Although we modeled lipid bilayers as symmetric membranes, the *real* lipid bilayers are composed of various lipids, and asymmetry facilitates deformation in specific directions.²⁴⁾ Understanding their behavior will lead to further insights into lipid bilayers.

Acknowledgment

The authors would like to thank H. Noguchi and H. Nakano for fruitful discussions. This research was supported by JSPS KAKENHI, Grant No. JP21K11923. The computation was partly carried out using the facilities of the Supercomputer Center, Institute for Solid State Physics (ISSP), University of Tokyo.

References

- 1) J. F. Nagle and S. Tristram-Nagle: *Biochimica et Biophysica Acta (BBA)-Reviews on Biomembranes* **1469** (2000) 159.
- 2) M. Bloom, E. Evans, and O. G. Mouritsen: *Quarterly Reviews of Biophysics* **24** (1991) 293.
- 3) C. Peetla, A. Stine, and V. Labhassetwar: *Molecular Pharmaceutics* **6** (2009) 1264.
- 4) R. Goetz and R. Lipowsky: *The Journal of Chemical Physics* **108** (1998) 7397.
- 5) R. Goetz, G. Gompper, and R. Lipowsky: *Physical Review Letters* **82** (1999) 221.
- 6) S. J. Marrink and A. E. Mark: *Journal of the American Chemical Society* **125** (2003) 15233.
- 7) W. Shinoda, R. DeVane, and M. L. Klein: *The Journal of Physical Chemistry B* **114** (2010) 6836.
- 8) P. S. Coppock and J. T. Kindt: *Langmuir* **25** (2009) 352.
- 9) R. M. Venable, Y. Zhang, B. J. Hardy, and R. W. Pastor: *Science* **262** (1993) 223.
- 10) M. Orsi and J. W. Essex: *Soft Matter* **6** (2010) 3797.
- 11) G. Fabrikant, S. Lata, J. D. Riches, J. A. Briggs, W. Weissenhorn, and M. M. Kozlov: *PLoS Computational Biology* **5** (2009) e1000575.
- 12) A. Viallat and M. Abkarian: *International Journal of Laboratory Hematology* **36** (2014) 237.
- 13) H. Yu, S. Engel, G. Janiga, and D. Thévenin: *Artificial Organs* **41** (2017) 603.
- 14) K. De Haas, C. Blom, D. Van den Ende, M. H. Duits, and J. Mellema: *Physical Review E* **56** (1997) 7132.
- 15) A. Zgorski, R. W. Pastor, and E. Lyman: *Journal of Chemical Theory and Computation* **15** (2019) 6471.
- 16) S. Kikuchi and H. Watanabe: *The Journal of Chemical Physics* **158** (2023).
- 17) W. Humphrey, A. Dalke, and K. Schulten: *Journal of Molecular Graphics* **14** (1996) 33.
- 18) S. Plimpton: *Journal of Computational Physics* **117** (1995) 1.
- 19) S. Nosé: *Molecular Physics* **52** (1984) 255.
- 20) W. G. Hoover: *Physical Review A* **31** (1985) 1695.
- 21) H. C. Andersen: *The Journal of Chemical Physics* **72** (1980) 2384.

- 22) L. Verlet: *Physical Review* **159** (1967) 98.
- 23) M. C. Watson, E. G. Brandt, P. M. Welch, and F. L. Brown: *Physical Review Letters* **109** (2012) 028102.
- 24) J. A. Op den Kamp: *Annual Review of Biochemistry* **48** (1979) 47.