DOMAINS OF EXISTENCE OF SLICE REGULAR FUNCTIONS IN ONE QUATERNIONIC VARIABLE

XINYUAN DOU, MING JIN, GUANGBIN REN, AND IRENE SABADINI

ABSTRACT. Recently, we introduced domains of slice regularity in the space $\mathbb H$ of quaternions and also proved that domains of slice regularity satisfy a symmetry with respect to paths, called 2-path-symmetry. In this paper, we give a full characterization by showing that all 2-path-symmetric slice-open sets are domains of slice regularity. In fact, we will prove a counterpart of the Cartan-Thullen theorem for slice regular functions, namely that a slice-open set is a domain of existence for some slice regular function if and only if it is a domain of slice regularity, if and only if it is slice-regularly convex, if and only if it is 2-path-symmetric. As a tool, we also prove an interpolation theorem of independent interest.

1. Introduction

The objective of this paper is to present a version of the Cartan-Thullen theorem adapted to the case of slice regular functions in one quaternionic variable. The Cartan-Thullen theorem [6] is a fundamental result in the theory of several complex variables, providing a powerful characterization of domains of holomorphy in complex analysis. This theorem establishes an equivalence between holomorphically convex domains, domains of holomorphy and the domains of existence of holomorphic functions

In the context of one complex variable, all open sets are domains of holomorphy. However, in several complex variables, Hartogs' phenomenon, see [34] or e.g. [35], demonstrates that not all open sets are necessarily domains of holomorphy and so the study of domains of holomorphy plays a crucial role in this context.

This paper aims to provide a topological equivalent characterization of the domains of existence of slice regular functions in one quaternionic variable on non-axially symmetric slice-open sets with respect to slice topology. In fact, we prove a Cartan-Thullen theorem and an interpolation theorem for such functions. The study is not a straightforward generalization of the case of one complex variable, although slice regular functions are one of the possible counterparts of holomorphic functions in the case of one quaternionic variable, and indeed we appeal more to techniques from the case of several complex variables.

²⁰²⁰ Mathematics Subject Classification. Primary: 30G35; Secondary: 32A30, 32D05.

Key words and phrases. Domains of holomorphy; quaternions; slice regular functions; representation formula; slice topology.

This work was supported by the China Postdoctoral Science Foundation (BX20220327) and the NNSF of China (12171448). The fourth author, member of GNSAGA, INdAM, is partially supported by PRIN 2022 Real and Complex Manifolds: Geometry and Holomorphic Dynamics. The authors express their gratitude to Xiangyu Zhou, Xieping Wang, and Zhuo Liu for engaging in insightful discussions on the topic of several complex variables.

Slice regular functions were introduced by Gentili and Struppa [30] in case of one quaternionic variable, elaborating an idea of Cullen [17] with the aim of defining a class of quaternionic "holomorphic" functions including polynomials and power series. Roughly speaking, slice regular functions are functions that are holomorphic with respect to imaginary unit I when restricted to the complex plane $\mathbb{C}_I := \mathbb{R} + \mathbb{R}I$. Slice regular functions were then extended to the case of functions defined on Clifford algebras [13], octonions [31], real alternative *-algebras [33] and, more in general, to even-dimensional Euclidean spaces [21] and also to vector-valued functions, see [4]. This class of functions is particularly successful [3,11,14,25,37], especially in view of the applications in operator theory [2,4,7–9,12,15,24].

For over a decade, the research on slice regular functions mainly focused on axially symmetric slice domains, where the representation formula shows that the values of a slice regular function are determined by the restriction to a complex plane (or restrictions to two different complex semi-planes). Interested readers can refer to the books [4,15,16,29] and the references therein to learn more about slice regular functions.

Slice analysis on non-axially symmetric domains is systematically studied after the introduction of slice topology, see [19,20] although it is also studied in [27,28], but with a substantially different approach still relying on the Euclidean topology. In [19,20] notions such as the path-representation formula and path-slice functions, are introduced to deal with non-axially symmetric domains.

To further develop the global theory of slice analysis, it is necessary to understand the properties of domains of slice regularity, an analogue of domains of holomorphy to this setting. Proposition 9.7 in [20] shows that all the domains of holomorphy satisfy a certain topology condition, called 2-path-symmetry. In this paper we will prove the converse, i.e. all 2-path-symmetric open sets in slice topology are domains of holomorphy. We will also show that a open set in slice topology is a domain of holomorphy if and only if it is the domain of existence of some slice regular function, if and only if it satisfies certain convexity, called slice regular convexity. These equivalent conditions form the Cartan-Tullen theorem for slice regular functions.

As we already pointed out, the proof of Cartan-Thullen theorem for slice regular functions has some similarities to that one given in complex analysis in several variables, for example the use of an interpolation theorem, but also relevant differences and it can be summarized as follows. We firstly prove that for any 2path-symmetric slice-open set Ω , there is $I \in \mathbb{S}$ such that the cardinality of $J \in \mathbb{S}$ such that $\Psi_J^I(\Omega_J) \neq \Omega_I$, where $\Psi_J^I: x + yJ \mapsto x + yI$, is countable, see Proposition 3.8. This allows us to construct a Riemann domain $\check{\Omega}$ over \mathbb{C} , see Theorem 5.10. Then we use some holomorphic maps on certain subsets $\tilde{\Omega}^{[I]}$, $I \in \mathbb{S}$ of the Riemann domain $\hat{\Omega}$, to construct some suitable slice regular functions, see Proposition 7.9. By the properties of these holomorphic functions, we deduce that the corresponding slice regular functions have the good properties in Lemma 8.5. This result implies that there is a slice regular function such that its absolute value is smaller than $\epsilon > 0$ on an invariant subset K of Ω (an analogue of compact set K with $K = \hat{K}_{\Omega}$ in complex analysis) and f=1 at a given point $q\in\Omega\backslash K$. Using these "good" slice regular functions, we prove an interpolation theorem, see Theorem 8.8 useful in the proof part of the Cartan-Thullen theorem in this framework, see Section 9.

It is important to point out that to prove the full statement of the Cartan-Thullen theorem, we need some notions that are specific to the quaternionic case, and deeply differ from those in the complex analysis case.

We provide here a brief account of some of these differences to show that the results in this paper are far from being trivial or easy. We consider a continuous function r^+ on $\tilde{\Omega}$ (see Proposition 5.14) corresponding to the σ -distance in Ω (see (6.16)). The function r^+ plays a similar role as the distance between the compact set and the boundary of the domain of holomorphy in complex analysis (see the proof of Proposition 8.7 and Proposition 9.1). Since a Riemann domain $\tilde{\Omega}$ is axially symmetric, we define conjugation on $\tilde{\Omega}$ (see (6.8)); note that the values of r^+ on conjugated points coincide (see Proposition 6.4). We then introduce stem discrete sets (see Definition 7.1) which, in the proof of the interpolation theorem, play a role similar to discrete points in complex analysis. Note that stem discrete points may not be discrete in Ω , and their corresponding set may also not be discrete in $\check{\Omega}$. They are discrete in the sense of σ -distance, that is, the corresponding set $\psi_{\Omega}^{\mathbb{S}}(A \cap \Omega^r)$ in $\check{\Omega}$ of intersection of a stem discrete set A and $\Omega^r=\{q\in\Omega:r\leq\sigma(q,\mathbb{H}\backslash\Omega)\}$ is discrete for each r > 0. An Ω -invariant set (see Definition 7.2) plays a role similar to a compact set K with $\hat{K}_{\Omega} = K$. Moreover, a constant C_x depending on the point $x \in \Omega$ appears in the definition of slice regular convex hull \hat{K}_{Ω} of K (see Definition 2.7), but this constant does not have a counterpart in complex analysis.

The structure of the paper is as follows. In section 2 we introduce the basic notions useful to state and prove the main results of this paper. In section 3 we give some properties of 2-path-symmetric st-domains, among which the crucial property of countability of distinct slices. In section 4 we introduce slice-piecewise paths to describe path-connectedness of st-domains. In section 5 we introduce a notion of 2-path-symmetric Riemann domain associated with a 2-path-symmetric st-domain, via a suitably defined equivalence relation. In Section 6 we study the correspondence between 2-path-symmetric st-domains of $\mathbb H$ and the associated 2-path-symmetric Riemann domains over $\mathbb C$ by proving a number of properties. In Section 7 we introduce stem discrete sets which serve as prescribed points for interpolation theorem in the case of slice analysis. In Section 8, we prove a couple of extension results of functions and an interpolation theorem. Finally, in Section 9 we summarize what we prepared in the preceding sections and we prove the Cartan-Thullen theorem in this framework. At the end, we added a list of symbols an notions to help the reader to navigate the paper.

2. Main results

In this section, we state our main results. To this end, we recall some definitions and results from [20]. Denote by

$$\mathbb{H} := \mathbb{R} + i\mathbb{R} + j\mathbb{R} + k\mathbb{R}$$

the quaternion number system where

$$i^2 = j^2 = k^2 = ijk = -1.$$

The set of imaginary units of quaternions \mathbb{H} is denoted by

$$\mathbb{S} := \{ I \in \mathbb{H} : I^2 = -1 \},$$

and let

$$\mathbb{S}^2_* := \{ (I, J) \in \mathbb{S}^2 : I \neq J \}.$$

Below, we will explore the extension of slice regularity along paths, akin to what is done in the holomorphic setting. If the slice domain Ω in $\mathbb H$ does not intersect the real axis, then, by the definition of the slice topology, the theory mirrors that of the holomorphic setting. However, if Ω intersects the real axis, we encounter various challenges and, consequently, below we consider continuous paths starting from points lying on the real axis.

Let $U \subset \mathbb{C}$ and set

$$\mathscr{P}_0(U) := \{ \delta : [0,1] \to U : \delta \text{ is continuous and } \delta(0) \in \mathbb{R} \}.$$

Definition 2.1. $\Omega \subset \mathbb{H}$ is called 2-path-symmetric if

$$\gamma^K \subset \Omega$$
.

for any $K \in \mathbb{S}$, $\gamma \in \mathscr{P}_0(\mathbb{C})$ and $(I,J) \in \mathbb{S}^2_*$ with $\gamma^I, \gamma^J \subset \Omega$, where $\gamma^I := \Psi^I_i \circ \gamma$,

$$\Psi_i^I: \quad \mathbb{C} \quad \longrightarrow \quad \mathbb{C}_I,$$

$$x+yi \quad \longmapsto \quad x+yI.$$

and $\gamma^K \subset \Omega$ is short for $\gamma^K([0,1]) \subseteq \Omega$.

The slice topology on \mathbb{H} is defined by

$$\tau_s(\mathbb{H}) := \{ U \subset \mathbb{H} : U_I \in \tau(\mathbb{C}_I), \forall \ I \in \mathbb{S} \},\$$

where $\tau(\mathbb{C}_I)$ the Euclidean topology on the complex plane $\mathbb{C}_I := \mathbb{R} + \mathbb{R}I$, and $U_I := U \cap \mathbb{C}_I$.

Conventions: We call $U \subset \mathbb{H}$ is slice-connected, if U is connected in $\tau_s(\mathbb{H})$; we call $U \subset \mathbb{H}$ is slice-open, if U is open in $\tau_s(\mathbb{H})$ and so on. A domain in $\tau_s(\mathbb{H})$ will be called st-domain since the terminology 'slice domain' is used in the literature for a different concept. Note also that the symbol \subset denotes the inclusion of sets, allowing equality.

Let $z \in \mathbb{C}$, $U \subset \mathbb{C}$, $I \in \mathbb{S}$ and $\mathbb{S}' \subset \mathbb{S}$. Denote

$$z^I := \Psi_i^I(z)$$
, and $U^{\mathbb{S}'} := \{ w^J \in \mathbb{H} : w \in U, \text{ and } J \in \mathbb{S}' \}$.

Moreover, we write $z^{\mathbb{S}'} := \{z\}^{\mathbb{S}'}$ and $U^I := U^{\{I\}}$ for short.

Example 2.2. Let $I, J \in \mathbb{S}$ with $I \neq \pm J$. Set

$$U_{a,b} := \{ \lambda e^{i\theta} + 3i \in \mathbb{C} : 2 < \lambda < 4, \ a < \theta < b \}.$$

Then

$$E_1 := \left(U_{-\frac{\pi}{2},\frac{3\pi}{4}}\right)^I \bigcup \left(U_{\frac{\pi}{4},\frac{3\pi}{2}}\right)^J \bigcup \left[U_{0,3\pi} \cap \operatorname{Conj}_{\mathbb{C}}\left(U_{0,3\pi}\right)\right]^{\mathbb{S}}$$

is a 2-path-symmetric st-domain, where $\operatorname{Conj}_{\mathbb{C}}: \mathbb{C} \to \mathbb{C}, \ z \mapsto \overline{z}$. Note that $U_{0,3\pi} \cap \operatorname{Conj}_{\mathbb{C}}(U_{0,3\pi})$ is the connected component of $U_{-\frac{\pi}{2},\frac{3\pi}{4}} \cap U_{\frac{\pi}{4},\frac{3\pi}{2}}$ which intersects the real axis. The intersection of E_1 with the complex planes \mathbb{C}_I , \mathbb{C}_J , \mathbb{C}_K , $K \neq \pm I, \pm J$ is shown in Figure 1.

Example 2.3. Let U be an open set in \mathbb{R} with infinite connected components. Then there are $a_1, a_2, ..., b_1, b_2, ... \in \mathbb{R}$ such that

$$U = \bigcup_{\ell=1}^{+\infty} (a_{\ell}, b_{\ell}) \quad \text{with} \quad (a_{i}, b_{i}) \cap (a_{j}, b_{j}) = \varnothing, \quad \forall i \neq j.$$




Figure 1.

Let $I_1, I_2, ... \in \mathbb{S}$ with $I_i \neq \pm I_j$ for each $i \neq j$. Then

$$E_2 := \bigcup_{\ell=1}^{+\infty} \left[(V_\ell)^{\mathbb{S}} \cup (W_\ell)^{I_\ell} \right]$$

is a 2-path-symmetric st-domain, where

$$\begin{cases} V_{\ell} := \left\{ x + yi \in \mathbb{C} : \ a_{\ell} < x < b_{\ell}, \ |y| < 1 \right\}, \\ W_{\ell} := W_{\ell}^{1} \cup W_{\ell}^{2} \cup W_{\ell}^{3}, \\ W_{\ell}^{1} := \left\{ x + yi \in \mathbb{C} : \left(x - \frac{a_{\ell} + b_{\ell}}{2} \right) \left(x - \frac{a_{\ell+1} + b_{\ell+1}}{2} \right) < 0, \ 2 < y < 3 \right\}, \\ W_{\ell}^{2} := \left\{ x + yi \in \mathbb{C} : \frac{a_{\ell} + b_{\ell}}{2} < x < b_{\ell}, \ 0 < y < 3 \right\}, \\ W_{\ell}^{3} := \left\{ x + yi \in \mathbb{C} : a_{\ell+1} < x < \frac{a_{\ell+1} + b_{\ell+1}}{2}, \ 0 < y < 3 \right\}. \end{cases}$$

See Figure 2.




FIGURE 2.

Definition 2.4. Let $I \in \mathbb{S}$ and $U \in \tau(\mathbb{C}_I)$. A function $f : U \to \mathbb{H}$ is called (I-)holomorphic, if f is real differentiable on $U \cap \mathbb{C}_I$ and

$$\frac{1}{2} \left(\frac{\partial}{\partial x} + I \frac{\partial}{\partial y} \right) f(x + yI) = 0, \qquad \forall \ x + yI \in U.$$

Denote by $\mathcal{O}_I(U)$ the class of I-holomorphic functions defined on U.

Definition 2.5. Let $\Omega \in \tau_s(\mathbb{H})$. A function $f: \Omega \to \mathbb{H}$ is called **slice regular**, if for each $I \in \mathbb{S}$, $f_I := f|_{\Omega_I}$ is I-holomorphic. Denote by $\mathcal{SR}(\Omega)$ the class of slice regular functions defined on Ω .

Definition 2.6. Let $\Omega \in \tau_s(\mathbb{H})$ and $\mathcal{F} \subset \mathcal{SR}(\Omega)$. Ω is called an \mathcal{F} -domain of existence if there are no $\Omega_1, \Omega_2 \in \tau_s(\mathbb{H})$ with the following properties:

- (i) $\varnothing \neq \Omega_1 \subset \Omega_2 \cap \Omega$.
- (ii) Ω_2 is τ_s -connected and $\Omega_2 \nsubseteq \Omega$.
- (iii) For each $f \in \mathcal{F}$, there is $\widetilde{f} \in \mathcal{SR}(\Omega_2)$ such that $f = \widetilde{f}$ in Ω_1 .

If there are slice-open sets Ω , Ω_1 , Ω satisfying (i)-(iii), then we call $(\Omega, \Omega_1, \Omega_2)$ an \mathcal{F} -triple. Moreover, we call $(\Omega, \Omega_1, \Omega_2)$ special, if Ω_1 is a slice-connected component of $\Omega \cap \Omega_2$.

 Ω is called a **domain of slice regularity** if it is an $\mathcal{SR}(\Omega)$ -domain of existence and Ω is called the **domain of existence** of $f \in \mathcal{SR}(\Omega)$ if Ω is $\{f\}$ -domain of existence

We now extend holomorphic convexity to the theory of slice regular functions:

Definition 2.7. Let $\Omega \in \tau_s(\mathbb{H})$ and $K \subset \Omega$. We call

$$\widehat{K}_{\Omega} := \left\{ x \in \Omega : \exists \ C_x > 0, \ s.t. \ |f(x)| \le C_x \sup_{y \in K} |f(y)|, \ \forall \ f \in \mathcal{SR}(\Omega) \right\},$$

the slice regularly convex hull of K.

For any $\Omega \subset \mathbb{H}$, denote

(2.1)
$$\Omega_{2ps} := \left\{ z \in \mathbb{C} : \exists \ \gamma \in \mathscr{P}_0(\mathbb{C}) \text{ and } (I, J) \in \mathbb{S}^2_*, \right. \\ s.t. \ \gamma^I, \gamma^J \subset \Omega \text{ and } \gamma(1) = z \right\},$$

and

$$\mathbb{S}_{\mathcal{O}}^{2ps} := \{ I \in \mathbb{S} : \Omega_I \neq (\Omega_{2ps})^I \}.$$

Definition 2.8. $K \subset \mathbb{H}$ is called **finite-non-symmetric compact**, if K_I is compact in $\tau(\mathbb{C}_I)$ and $|\mathbb{S}_K^{2ps}| < +\infty$.

Definition 2.9. $\Omega \in \tau_s(\mathbb{H})$ is called **slice regularly convex**, if

$$(\widehat{K}_{\Omega})_I \subseteq \Omega_I$$
,

for each finite-non-symmetric compact set $K \subset \Omega$ and $I \in \mathbb{S}$.

The main result is as follows:

Theorem 2.10. (Cartan-Thullen) Let $\Omega \in \tau_s(\mathbb{H})$. Then the following statements are equivalent:

- (i) Ω is the domain of existence of some slice regular function.
- (ii) Ω is a domain of slice regularity.
- (iii) Ω is slice regularly convex.
- (iv) Ω is 2-path-symmetric.

3. 2-PATH-SYMMETRIC ST-DOMAINS

In this section, we prove various properties of 2-path-symmetric st-domains. In particular, next Proposition 3.8 shows that 2-path-symmetric st-domains have at most countable distinct slices, where distinct is meant in the sense of some symmetry. To begin with, we define a binary relation which is an equivalence relation on 2-path-symmetric open sets.

Definition 3.1. Define a binary relation \approx_{Ω} on $\Omega \subset \mathbb{H}$: Let $p, q \in \Omega$. Then $p \approx_{\Omega} q$ if and only if there is $\gamma \in \mathscr{P}_0(\mathbb{C})$ and $I, J \in \mathbb{S}$ such that

(3.1)
$$\gamma^{I}, \gamma^{J} \subset \Omega, \qquad \gamma^{I}(1) = p \quad and \quad \gamma^{J}(1) = q.$$

Let us set

$$[p]_{\approx_{\Omega}} := \{ w \in \Omega : w \approx_{\Omega} p \}.$$

Proposition 3.2. Let $\Omega \subset \mathbb{H}$. Then the following statements holds:

- (i) Let $p, z^L \in \Omega$, with $z \in \mathbb{C}$, $L \in \mathbb{S}$ and $p \approx_{\Omega} z^L$. Then $p = z^K$ for some $K \in \mathbb{S}$.
- (ii) If Ω is 2-path-symmetric, then \approx_{Ω} is an equivalence relation on Ω .
- (iii) Let $z^I, z^J \in \Omega$ with $z^I \approx_{\Omega} z^J$. Then there is $\gamma \in \mathscr{P}_0(\mathbb{C})$ such that

$$\gamma(1) = z$$
 and $\gamma^I, \gamma^J \subset \Omega$.

(iv) Let $z^{I_i} \in \Omega$ with $z^{I_i} \in [z^{I_1}]_{\approx_{\Omega}}$, i = 1, ..., k. If Ω is 2-path-symmetric, then there is $\gamma \in \mathscr{P}_0(\mathbb{C})$ such that

$$\gamma(1)=z, \qquad and \qquad \gamma^{I_{\imath}}\subset \Omega, \qquad \imath=1,...,k.$$

(v) Let $z^I \in \Omega$ with $|[z^I]_{\approx_{\Omega}}| > 1$. If Ω is 2-path-symmetric, then $[z^I]_{\approx_{\Omega}} = z^{\mathbb{S}}$.

Proof. (i) Since $p \approx_{\Omega} z^L$, there is $\gamma \in \mathscr{P}_0(\mathbb{C})$ and $I, J \in \mathbb{S}$ such that (3.1) holds. Then $\gamma(1) = \operatorname{Conj}_{\mathbb{C}}^{\lambda}(z)$ for some $\lambda \in \{0,1\}$, where $\operatorname{Conj}_{\mathbb{C}}: \mathbb{C} \to \mathbb{C}, z \mapsto \overline{z}$ and $\operatorname{Conj}_{\mathbb{C}}^{\lambda} := (\operatorname{Conj}_{\mathbb{C}})^{\lambda}$ (i.e. we set $\operatorname{Conj}_{\mathbb{C}}^0 = \operatorname{Identity}$ and $\operatorname{Conj}_{\mathbb{C}}^1 = \operatorname{Conj}_{\mathbb{C}}$). By (3.1),

$$q = \gamma^J(1) = \left[\operatorname{Conj}_{\mathbb{C}}^{\lambda}(z)\right]^J = z^{(-1)^{\lambda}J} = z^K,$$

where $K := (-1)^{\lambda} L \in \mathbb{S}$.

(ii) The reflexivity and symmetry of \approx_{Ω} hold directly by definition. We only need to prove the transitivity. Let $u, v = x + yL, w \in \Omega$ with $u \approx_{\Omega} v$ and $v \approx_{\Omega} w$. By hypothesis, there is $\gamma \in \mathscr{P}_0(\mathbb{C})$ and $I, J \in \mathbb{S}$ such that

$$\gamma^I, \gamma^J \subset \Omega, \qquad \gamma^I(1) = u, \qquad \text{and} \qquad \gamma^J = v.$$

According to (i) and since $v \approx_{\Omega} w$, we have that $w = x + yK = \gamma^K(1)$ for some $K \in \mathbb{S}$. It implies that $u \approx_{\Omega} w$.

(iii) If $z \in \mathbb{R}$, then (iii) holds by taking $\gamma \equiv z$. Otherwise, $z \notin \mathbb{R}$. By definition, there is $\beta \in \mathscr{P}_0(\mathbb{C})$ and $I', J' \in \mathbb{S}$ such that $\beta^{I'}, \beta^{J'} \subset \Omega$,

$$\beta^{I'}(1) = z^I$$
, and $\beta^{J'}(1) = z^J$.

It follows that $\beta(1) = \operatorname{Conj}_{\mathbb{C}}^{\lambda}(z)$ for some $\lambda \in \{0, 1\}$. Then

$$z^{I} = \beta^{I'}(1) = \left[\operatorname{Conj}_{\mathbb{C}}^{\lambda}(z)\right]^{I'} = z^{(-1)^{\lambda}I'}.$$

It implies that $I' = (-1)^{\lambda} I$ and $J' = (-1)^{\lambda} I$. Take $\gamma := \operatorname{Conj}_{\mathbb{C}}^{\lambda}(\beta)$. Then

$$\gamma(1) = \left[\operatorname{Conj}_{\mathbb{C}}^{\lambda}(\beta)\right](1) = \operatorname{Conj}_{\mathbb{C}}^{\lambda}(\beta(1)) = \operatorname{Conj}_{\mathbb{C}}^{\lambda}\left(\operatorname{Conj}_{\mathbb{C}}^{\lambda}(z)\right) = z,$$

and

$$\gamma^{I} = \left[\operatorname{Conj}_{\mathbb{C}}^{\lambda}(\beta)\right]^{I} = \beta^{(-1)^{\lambda}I} = \beta^{I'} \subset \Omega.$$

Similarly, $\gamma^J \subset \Omega$. It implies that (iii) holds.

(iv) If $I_1 = \cdots I_k$, then (iv) holds by taking $\gamma \equiv z$. Otherwise, without loss of generality, we suppose $I_1 \neq I_2$. By (iii), there is $\gamma \in \mathscr{P}_0(\mathbb{C})$ such that $\gamma(1) = z$ and

 $\gamma^{I_1}, \gamma^{I_2} \subset \Omega$. Since Ω is 2-path-symmetric, $\gamma^J \subset \Omega$ for each $J \in \mathbb{S} \supset \{I_3, ..., I_k\}$. It implies that (iv) holds.

(v) It follows from (i) that $[z^I]_{\approx_0} \subset z^{\mathbb{S}}$. We only need to prove that

$$(3.2) z^{\mathbb{S}} \subset [z^I]_{\approx_{\Omega}}.$$

Since $|[z^I]_{\approx_{\Omega}}| > 1$, there is $w^L \approx_{\Omega} q$ for some $w \in \mathbb{C}$ and $L \in \mathbb{S}$ with $w^L \neq q$. By (i), there is $J \in \mathbb{S} \setminus \{I\}$ such that $w^L = z^J$. According to (iv), there is $\gamma \in \mathscr{P}_0(\mathbb{C})$ such that $\gamma(1) = z$ and $\gamma^I, \gamma^J \subset \Omega$. Since Ω is 2-path-symmetric, we have $\gamma^K \subset \Omega$ for each $K \in \mathbb{S}$. It implies that $z^K = \gamma^K(1) \in [z^I]_{\approx_{\Omega}}$ for each $K \in \mathbb{S}$. Therefore (3.2) holds.

Remark 3.3. Let $\Omega \subset \mathbb{H}$. It is easy to check by Proposition 3.2 (iii) that

(3.3)
$$\Omega_{2ps} = \left\{ z \in \mathbb{C} : z^I, z^J \in \Omega \text{ with } z^I \approx_{\Omega} z^J, \text{ for some } (I, J) \in \mathbb{S}^2_* \right\}.$$

Note also that, by definition,

$$(3.4) \Omega_{\mathbb{R}} := \Omega \cap \mathbb{R} \subset \Omega_{2ns}.$$

Moreover, if $\Omega_{\mathbb{R}} = \emptyset$, then $\Omega_{2ps} = \emptyset$.

Denote by

$$\mathcal{L}_{z}^{w}: [0,1] \longrightarrow \mathbb{C},$$

$$t \longmapsto tw + (1-t)z.$$

the segment from $z \in \mathbb{C}$ to $w \in \mathbb{C}$.

Proposition 3.4. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric. Then $\Omega_{2ps} \in \tau(\mathbb{C})$.

Proof. Let $z \in \Omega_{2ps}$. By definition, there is a $\gamma \in \mathscr{P}_0(\mathbb{C})$ and $(I, J) \in \mathbb{S}^2_*$ such that $\gamma^I, \gamma^J \subset \Omega$ and $\gamma(1) = z$. It implies that $z^I \in \Omega_I$ and $z^J \in \Omega_J$. Therefore there is r > 0 such that

$$[B_{\mathbb{C}}(z,r)]^{I}, [B_{\mathbb{C}}(z,r)]^{J} \subset \Omega,$$

where $[B_{\mathbb{C}}(z,r)]^I$ is the disc in the plane \mathbb{C}_I with center at z^I and with radius r > 0. For each $w \in B_{\mathbb{C}}(z,r)$, it is easy to check that

$$\gamma \circ \mathcal{L}_{z}^{w} \in \mathscr{P}_{0}(\mathbb{C}), \quad (\gamma \circ \mathcal{L}_{z}^{w})^{I} \subset \Omega_{I}, \quad (\gamma \circ \mathcal{L}_{z}^{w})^{J} \subset \Omega_{J} \quad \text{and} \quad \gamma \circ \mathcal{L}_{z}^{w}(1) = w.$$

It implies by definition that $w \in \Omega_{2ps}$. Then $B_{\mathbb{C}}(z,r) \subset \Omega_{2ps}$, and z is an interior point of Ω_{2ps} . Therefore Ω_{2ps} is open.

Proposition 3.5. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric. Then

(3.5)
$$\partial(\Omega_{2ps}) \bigcap \Psi_I^i(\Omega_I) \bigcap \Psi_J^i(\Omega_J) = \varnothing, \qquad \forall \ (I,J) \in \mathbb{S}^2_*,$$

where
$$\Psi_I^i := (\Psi_I^I)^{-1}: \ \Omega_I \to \Omega, \ \Psi_I^i(x+Iy) = x+iy.$$

Proof. If $\Omega_{\mathbb{R}} = \emptyset$, then $\Omega_{2ps} = \emptyset$, so is $\partial(\Omega_{2ps})$. It implies that (3.5) holds. Otherwise $\Omega_{\mathbb{R}} \neq \emptyset$. Suppose that (3.5) does not hold. Then there is a point

$$z \in \partial(\Omega_{2ps}) \bigcap \Psi_I^i(\Omega_I) \bigcap \Psi_J^i(\Omega_J).$$

Since $\Omega \in \tau_s(\mathbb{H})$, by definition Ω_I is open in \mathbb{C}_I and then $\Psi_I^i(\Omega_I)$ is open in \mathbb{C} . Thence there is r > 0 such that the open ball $\mathbb{B} := B_{\mathbb{C}}(z, r)$ in \mathbb{C} is contained in $\Psi_I^i(\Omega_I) \cap \Psi_J^i(\Omega_J)$.

It follows from $z \in \mathbb{B} \cap \partial(\Omega_{2ps})$ that $\mathbb{B} \cap \Omega_{2ps} \neq \emptyset$. Let $u \in \mathbb{B} \cap \Omega_{2ps}$. By (2.1), there is $\alpha \in \mathscr{P}_0(\mathbb{C})$, such that $\alpha^I, \alpha^J \subset \Omega$ and $\alpha(1) = u$. Then by

$$(\mathcal{L}_{u}^{z})^{K} \subset \mathbb{B}^{K} \subset (\Psi_{K}^{i}(\Omega))^{K} = \Omega_{K}, \qquad K = I, J,$$

we have

$$(\alpha \circ \mathcal{L}_{u}^{z})^{K} = \alpha^{K} \circ (\mathcal{L}_{u}^{z})^{K} \subset \Omega_{K}, \qquad K = I, J.$$

It implies that

$$z = (\alpha \circ \mathcal{L}_{u}^{z})(1) \in \Omega_{2ps}$$

which contradicts $z \in \partial(\Omega_{2ps})$.

Proposition 3.6. Let $\Omega \subset \mathbb{H}$ be an st-domain with $\Omega_{\mathbb{R}} := \Omega \cap \mathbb{R} \neq \emptyset$. Then each connected component of Ω_I in topological space \mathbb{C}_I , $I \in \mathbb{S}$, intersects \mathbb{R} .

Proof. Proof by contradiction. If U is a connected component of Ω_I for some $I \in \mathbb{S}$ with $U_{\mathbb{R}} \neq \emptyset$, then it is easy to check that U and $\Omega \setminus U$ are two disjoint nonempty slice-open sets. It implies that Ω is not slice-connected, a contradiction.

Remark 3.7. Let $\Omega \subset \mathbb{H}$ be an st-domain with $\Omega_{\mathbb{R}} = \emptyset$. According to [20, Proposition 3.8], $\Omega \subset \mathbb{C}_I$ for some $I \in \mathbb{S}$. It follows by connectedness that Ω is a domain in the topological space \mathbb{C}_I .

Proposition 3.8. Let $\Omega \subset \mathbb{H}$ be a non-empty 2-path-symmetric st-domain. Then $\mathbb{S}^{2ps}_{\Omega}$ is countable.

Proof. (i) Suppose that $\Omega_{\mathbb{R}} = \emptyset$. By Remark 3.7, Ω is a domain \mathbb{C}_I for some $I \in \mathbb{S}$. Therefore $\Omega_J = \emptyset$ for each $J \neq \pm I$. Recalling that, by (2.2), $\mathbb{S}^{2ps}_{\Omega} = \{I \in \mathbb{S} : \Omega_I \neq (\Omega_{2ps})^I\}$, it follows from $\Omega_{2ps} = \emptyset$ that $\mathbb{S}^{2ps}_{\Omega} = \{\pm I\}$ is countable.

(ii) Suppose that $\Omega_{\mathbb{R}} \neq \emptyset$. Let \mathcal{B} be a countable topological bases of \mathbb{C} . Let

(ii) Suppose that $\Omega_{\mathbb{R}} \neq \emptyset$. Let \mathcal{B} be a countable topological bases of \mathbb{C} . Let $I \in \mathbb{S}^{2ps}_{\Omega}$. By definition $\Omega_{I} \neq (\Omega_{2ps})^{I}$. Let U be a connected component of Ω_{I} in \mathbb{C}_{I} with $U \nsubseteq (\Omega_{2ps})^{I}$. By Proposition 3.6, $U_{\mathbb{R}} \neq \emptyset$. It follows by $U_{\mathbb{R}} \subset \Omega_{\mathbb{R}} \subset \Omega_{2ps}$ that $U \cap (\Omega_{2ps})^{I} \supset U_{\mathbb{R}} \neq \emptyset$.

If $U \cap \partial_I \left[(\Omega_{2ps})^I \right] = \emptyset$, then U is the union of two disjoint nonempty open set $U \setminus (\Omega_{2ps})^I$ and $U \cap (\Omega_{2ps})^I$, where $\partial_I \left[(\Omega_{2ps})^I \right]$ is the boundary of $(\Omega_{2ps})^I$ in \mathbb{C}_I . So U is not connected, a contradiction.

Otherwise $U \cap \partial_I [(\Omega_{2ps})^I] \neq \emptyset$. Let $z_I \in U \cap \partial_I [(\Omega_{2ps})^I]$ and $V_{(I)} \in \mathcal{B}$ with $z_I \in (V_{(I)})^I \subset \Omega_I$. Consider the map

$$\varphi: \quad \mathbb{S}^{2ps}_{\Omega} \quad \longrightarrow \quad \mathcal{B},$$

$$J \quad \longmapsto \quad V_{(J)}.$$

We will prove that φ is injective by contradiction: If $\varphi(K) = \varphi(L)$ for some $(K, L) \in \mathbb{S}^2_*$, then by $V_{(K)} = \varphi(K) = \varphi(L) = V_{(L)}$ we have

$$\Psi_K^i(z_K) \in \partial(\Omega_{2ps}) \cap V_{(K)} \cap V_{(L)} \subset \partial(\Omega_{2ps}) \cap \Psi_K^i(\Omega_K) \cap \Psi_L^i(\Omega_L).$$

It implies that (3.5) does not hold, a contradiction. Therefore φ is injective and the cardinality $|\mathbb{S}_{\mathcal{O}}^{2ps}| < |\mathcal{B}|$. So $\mathbb{S}_{\mathcal{O}}^{2ps}$ is countable.

Proposition 3.9. Let $\Omega \subset \mathbb{H}$ be 2-path-symmetric and $U \subset \mathbb{C}$ be path-connected with $U \cap \Omega_{2ps} \neq \emptyset$. If $U^I \subset \Omega_I$ and $U^J \subset \Omega_J$ for some $I \neq J$, then $U \subset \Omega_{2ps}$.

Proof. Let $z \in U \cap \Omega_{2ps}$. Then there are $\gamma \in \mathscr{P}_0(\mathbb{C})$ and $(I', J') \in \mathbb{S}^2_*$, s.t. $\gamma^{I'}, \gamma^{J'} \subset \Omega$ and $\gamma(1) = z$. Since Ω is 2-path-symmetric, $\gamma^K \subset \Omega$ for each $K \in \mathbb{S}$. Take $w \in U$, then there is a path β in U from z to w. Thus

$$\beta^L \subset U^L \subset \Omega_L, \quad \forall \ L \in \{I, J\}.$$

Hence $(\gamma\beta)^I$, $(\gamma\beta)^J \subset \Omega$ and $(\gamma\beta)(1) = w$. It implies $w \in \Omega_{2ps}$ and so $U \subset \Omega_{2ps}$. \square

Let $\Omega \subset \mathbb{H}$ and set

$$\widetilde{\Omega} := \bigcup_{x+yI \in \Omega} x + y \mathbb{S}.$$

 Ω is called *axially symmetric* if $\Omega = \widetilde{\Omega}$. We point out that $\widetilde{\Omega}$ is said to be the *axially symmetric completion* of Ω according to the terminology in [10].

In the sequel, it is useful to introduce the notation

$$\Omega^{2ps} := \{ x + yI : x + yi \in \Omega_{2ps}, \ I \in \mathbb{S} \}$$

to denote the axially symmetric completion of $\Omega_{2ps},$ i.e. $\Omega^{2ps}=\widetilde{\Omega^{2ps}}.$

Remark 3.10. Let $\Omega \subset \mathbb{H}$ is 2-path-symmetric. Then the following statements follow from the definitions:

- (i) Ω^{2ps} is a subset of Ω .
- (ii) Let $I \in \mathbb{S}$. Then $x + yI \in \Omega^{2ps}$ if and only if $x + yJ \in \Omega^{2ps}$ for each $J \in \mathbb{S}$.
- (iii) If $\Omega \in \tau_s(\mathbb{H})$, then $\Omega^{2ps} \in \tau_s(\mathbb{H})$.
- (iv) If $\Omega^{2ps} \in \tau_s(\mathbb{H})$, then for each $f \in \mathcal{SR}(\Omega^{2ps})$,

$$f(x+yI) = (1,I) \begin{pmatrix} 1 & J \\ 1 & K \end{pmatrix} \begin{pmatrix} f(x+yJ) \\ f(x+yK) \end{pmatrix}, \qquad \forall \ x+yI \in \Omega^{2ps}.$$

For all $J,K \in \mathbb{S}$ and $x+yI \in \Omega^{2ps}$. It implies that $f:(\Omega^{2ps},\tau) \to (\mathbb{H},\tau)$ is continuous.

(v)
$$\Omega_{2ps} = \operatorname{Conj}_{\mathbb{C}}(\Omega_{2ps}) := \{ z \in \mathbb{C} : \overline{z} \in \Omega_{2ps} \}.$$

4. SLICE-PIECEWISE PATH

In this brief section, we introduce the concept of slice-piecewise paths. These paths can describe the path-connectedness of st-domains in a finer way. In particular, we show that any two points in an st-domain can be connected by a slice-piecewise path.

Definition 4.1. Let $n \in \mathbb{N} \setminus \{0\}$, $I_i \in \mathbb{S}$,

$$I_i \in \mathbb{S}, \qquad \gamma_i \in \mathscr{P}(\mathbb{C}) := \{ \delta : [0,1] \to \mathbb{C} : \delta \text{ is continuous} \}, \qquad i = 1, ..., n.$$

We call $((\gamma_1)^{I_1}, ..., (\gamma_n)^{I_n})$, an (n-)slice-piecewise path from $[(\gamma_1)^{I_1}]$ (0) to $[(\gamma_n)^{I_n}]$ (1) if

$$\gamma_i(1) = \gamma_{i+1}(0), \qquad i = 1, ..., n-1.$$

Moreover, we call an (n-)slice-piecewise path $\beta = (\beta_1, ..., \beta_n)$ is in $\Omega \subset \mathbb{H}$, if $\beta_i \subset \Omega$, i = 1, ..., n.

Proposition 4.2. Let $\Omega \subset \mathbb{H}$ be an st-domain, and $p, q \in \Omega$. Then there is a slice-piecewise path β in Ω from p to q.

Proof. According to [20, Proposition 3.10], (\mathbb{H}, τ_s) is connected and locally pathconnected, so is (Ω, τ_s) . Then there is a path α in Ω from p to q. Let $t \in [0, 1]$. As a result of [20, Proposition 3.5], there is an st-domain $U_t \subset \Omega$ containing $\alpha(t)$, such that U_t is real-connected (that is $(U_t)_{\mathbb{R}}$ being connected in \mathbb{R}). Then the connected component W_t of $\alpha^{-1}(U_t)$ containing t is open in [0, 1]. Since [0, 1] is compact, it follows that there is a minimal $m \in \mathbb{N}^+$, and $t_i \in [0, 1]$, i = 1, 2, ..., m, such that

$$[0,1] \subset \bigcup_{i=1}^{m} W_{t_i}$$
 and $0 < t_0 < t_1 < \dots < t_m < 1$.

In fact, $\bigcup_{i=1}^m W_{t_i}$ covers each point in [0,1] at most twice. Otherwise, there is a point $q \in W_a \cap W_b \cap W_c$ in [0,1], and $a,b,c \in \{t_1,t_2,...,t_m\}$ are different from each other. Then

$$W := W_a \cup W_b \cup W_c$$

is an open interval in [0,1]. Without loss of generality, we can suppose that

$$\inf(W_a) = \inf(W)$$
 and $\sup(W_b) = \sup(W)$.

Then $W = W_a \cup W_b$. So we can remove W_c , and m is not minimal.

Therefore there is a bijection

$$L: \{1, 2, ..., m\} \rightarrow \{1, 2, ..., m\}$$

(reordering $\{W_{t_i}\}_{i=1}^m$ by the order of $\sup(W_{t_i})$), and

$$s_i \in [0, 1], \qquad i = 1, 2, ..., m + 1$$

with

$$0 = s_1 < s_2 < \ldots < s_{m+1} = 1 \qquad \text{and} \qquad [s_{\jmath}, s_{\jmath+1}] \subset W_{t_{L(\jmath)}}, \quad \jmath = 1, 2, \ldots, m.$$

Then

$$\alpha([s_i, s_{i+1}]) \subset U_{t_{L(i)}}, \quad i = 1, 2, ..., m.$$

Thanks to [20, Corollary 3.9], there are two paths β_{2i-1} and β_{2i} in U such that $\beta_{2i-1}\beta_{2i}$ be a path from $\alpha(s_i)$ to $\alpha(s_{i+1})$ for each $i \in \{1, 2, ..., m\}$, and β_j , j = 1, ..., 2m is a path on a slice \mathbb{C}_{I_j} for some $I_j \in \mathbb{S}$. Then $\prod_{i=1}^{2m}\beta_i := \beta_1 \cdots \beta_{2m}$ is a path from p to q, i.e. $(\beta_1, ..., \beta_{2m})$ is a slice-piecewise path from p to q.

5. 2-PATH-SYMMETRIC RIEMANN DOMAIN

In this section, we introduce a suitable notion of 2-path-symmetric Riemann domain. To this end, we first define an equivalence relation \sim_{Ω} on $\bigsqcup_{I \in \mathbb{S}} \Omega_I \subset \mathbb{H} \times \mathbb{S}$. Then, for any 2-path-symmetric st-domain $\Omega \subset \mathbb{H}$ we define a new topological space $\check{\Omega}$, see (5.5), and a map $\pi_{\check{\Omega}}$ which turns out to be a local homomorphism, see Proposition 5.7. In Theorem 5.10 we prove that $(\check{\Omega}, \pi_{\check{\Omega}})$ is a Riemann domain over \mathbb{C} when Ω is a 2-path-symmetric domain and this is the reason to refer to $\check{\Omega}$ as to the 2-path-symmetric Riemann domain associated with Ω .

Let $\Omega \subset \mathbb{H}$ and denote by

$$\bigsqcup_{I\in\mathbb{S}}\Omega_I:=\bigcup_{I\in\mathbb{S}}\left(\Omega_I\times\{I\}\right)=\left\{(u,I)\in\mathbb{H}\times\mathbb{S}:u\in\Omega_I\right\},$$

the disjoint union of $\{\Omega_I\}_{I\in\mathbb{S}}$ with the disjoint union topology

$$\tau\left(\bigsqcup_{I\in\mathbb{S}}\Omega_I\right):=\left\{U\subset\bigsqcup_{I\in\mathbb{S}}\Omega_I:U\cap\left(\Omega_I\times\{I\}\right)\in\tau\left(\Omega_I\times\{I\}\right),\;\forall\;I\in\mathbb{S}\right\}.$$

Let $u \in \bigsqcup_{I \in \mathbb{S}} \Omega_I$. Denote by (z_u, I_u) , the unique pair in $\mathbb{C} \times \mathbb{S}$ such that

$$(5.1) u = (z_u^{I_u}, I_u),$$

where $z_u^{I_u} := (z_u)^{I_u}$.

Define a binary relation \sim_{Ω} : Let $u, v \in \bigsqcup_{I \in \mathbb{S}} \Omega_I$. Then $u \sim_{\Omega} v$ if and only if the following statements hold:

(5.2)
$$(i) \quad z_u = z_v.$$

$$(ii) \quad I_u = I_v \text{ when } z_u \notin \Omega_{2ps}.$$

It is easy to check that if $u \sim_{\Omega} v$ and $u \neq v$, then

(5.3)
$$I_u \neq I_v$$
, and $z_u = z_v \in \Omega_{2ps}$.

According to (5.2),

$$(5.4) z_a^{I_a} \approx_{\Omega} z_b^{I_b}, \forall \ a,b \in \bigsqcup_{I \in \mathbb{S}} \Omega_I \text{ with } a \sim_{\Omega} b.$$

Proposition 5.1. Let $\Omega \subset \mathbb{H}$. Then the binary relation \sim_{Ω} is an equivalence relation.

Proof. The reflexivity of \sim_{Ω} is directly by definition.

Proof of transitivity: Let $u \sim_{\Omega} v$ and $v \sim_{\Omega} w$, we need to prove $u \sim_{\Omega} w$. If two of u, v and w coincide, then the transitivity holds. Otherwise, u, v and w are distinct to each other. $u \sim_{\Omega} v$ implies by (5.3) that $z_u = z_v \in \Omega_{2ps}$. Similarly, $z_v = z_w \in \Omega_{2ps}$. In summary,

$$z_u = z_v = z_w \in \Omega_{2ps}$$
.

It is clear by (5.2) that $u \sim w$.

Proof of symmetry: Let $u \sim_{\Omega} v$. If u = v, then $v \sim_{\Omega} u$ by reflexivity. Otherwise $u \neq v$. According to (5.3), $z_u = z = z_v \in \Omega_{2ps}$. Hence $v \sim_{\Omega} u$.

Let X be a set and let P(X) be the power set of X, namely the collection of subsets of X:

$$P(X) := \{ U \subset X \}.$$

Let $\Omega \in \tau_s(\mathbb{H})$ be a 2-path-symmetric. We define the topological space

$$(5.5) \qquad \qquad \breve{\Omega} := \bigsqcup_{I \in \mathbb{S}} \Omega_I / \sim_{\Omega}$$

with the final topology

induced by inclusions $\left\{i_{\scriptscriptstyle\Omega}^I\right\}_{I\in\mathbb{S}}$ defined by

$$i_{\Omega}^{I}: \quad \Omega_{I} \times \{I\} \quad \longrightarrow \quad \check{\Omega} \subset P\left(\bigsqcup_{I \in \mathbb{S}} \Omega_{I}\right),$$

$$u \qquad \longmapsto \quad \left[u\right]_{\sim_{\Omega}} := \left\{v \in \bigsqcup_{I \in \mathbb{S}} \Omega_{I} : v \sim_{\Omega} u\right\}.$$

It is easy to check that the definition implies that i_{Ω}^{I} is continuous and

$$(5.7) (i_{\Omega})^{-1}\left([u]_{\sim_{\Omega}}\right) = \begin{cases} \left\{\left(z_{u}^{J}, J\right) : J \in \mathbb{S}\right\}, & z_{u} \in \Omega_{2ps}, \\ \left\{u = \left(z_{u}^{I_{u}}, I_{u}\right)\right\}, & \text{otherwise,} \end{cases}$$

where $i_{\Omega}: u \mapsto [u]_{\sim_{\Omega}}$ is the canonical projection, moreover

$$(5.8) [(w^I, I)]_{\sim_{\Omega}} = [(w^J, J)]_{\sim_{\Omega}}, \forall w \in \Omega_{2ps} \text{ and } I, J \in \mathbb{S},$$

Hence for each $U \subset \Omega_{2ps}$ and $I, J \in \mathbb{S}$,

$$(5.9) \qquad [U^I \times \{I\}]_{\sim_{\Omega}} = \bigcup_{\xi \in U} [(\xi^I, I)]_{\sim_{\Omega}} = \bigcup_{\xi \in U} [(\xi^J, J)]_{\sim_{\Omega}} = [U^J \times \{J\}]_{\sim_{\Omega}},$$

where

$$[\boldsymbol{U}^I \times \{I\}]_{\sim_{\Omega}} := \left\{ [\boldsymbol{v}]_{\sim_{\Omega}} : \boldsymbol{v} \in \boldsymbol{U}^I \times \{I\} \right\}.$$

Proposition 5.2. Let $\Omega \subset \mathbb{H}$ be a 2-path-symmetric set, $I \in \mathbb{S}$ and $U \subset \Omega_I$. Then

$$(5.10) \qquad (i_{\Omega}^{J})^{-1} \left(\left[U \times \{I\} \right]_{\sim_{\Omega}} \right) = \begin{cases} \left(\Psi_{I}^{J}(U) \cap \Omega^{2ps} \right) \times \{J\}, & J \neq I, \\ U \times \{I\}, & J = I, \end{cases}$$

where $\Psi_I^J := \Psi_i^J \circ \Psi_I^i$.

Proof. "Proof of \subset " Let

$$u \in (i_{\Omega}^{J})^{-1} ([U \times \{I\}]_{\sim_{\Omega}}) \subset (i_{\Omega}^{J})^{-1} (\check{\Omega}) = \Omega_{J} \times \{J\}.$$

Then there is $v \in U \times \{I\}$ such that $u \sim_{\Omega} v$. By (5.2), $z_u = z_v$.

If I = J, then

$$u = \left(z_u^J, J\right) = \left(z_v^I, I\right) = v \in U \times \{I\}.$$

Otherwise, $I \neq J$. By (5.2), $z_u^J \in \Omega^{2ps}$. It follows from $z_v^I \in U$ that $z_u^J = z_v^J \in \Psi_I^J(U)$. In summary,

$$u=\left(z_u^J,J\right)\in \left(\Psi_I^J(U)\cap\Omega^{2ps}\right)\times\{J\}.$$

"Proof of \supseteq " The case of J=I: Let $u\in U\times\{I\}$. Since $i_{\Omega}^{I}(u)=[u]_{\sim_{\Omega}}\in [U\times\{I\}]_{\sim_{\Omega}}$, we have $u\in \left(i_{\Omega}^{I}\right)^{-1}\left(\left[U\times\{I\}\right]_{\sim_{\Omega}}\right)$.

The case of $J \neq I$: Let $u \in (\Psi_I^J(U) \cap \Omega^{2ps}) \times \{J\}$. Then $z_u^J \in \Omega^{2ps}$. It implies that $u \sim_{\Omega} (z_u^I, I) =: v$ and $v \in U \times \{I\}$. Hence

$$i_{\scriptscriptstyle\Omega}^J(u) = [u]_{\sim_\Omega} = [v]_{\sim_\Omega} \in \left[U \times \{I\}\right]_{\sim_\Omega},$$

i.e.
$$u \in (i_{\Omega}^{J})^{-1}([U \times \{I\}]_{\sim_{\Omega}}).$$

Let $\Omega \subset \mathbb{H}$ be a 2-path-symmetric set, $\mu \in \check{\Omega}$ and $u, v \in (i_{\Omega})^{-1}(\mu)$. It follows from $u \sim_{\Omega} v$ that $z_u = z_v$. So

$$(5.11) z_{\mu} := z_{\nu}$$

do not depend on the choice of $u \in (i_{\Omega})^{-1}(\mu)$. Denote

$$(5.12) \mathbb{S}_{\mu} := \left\{ I \in \mathbb{S} : z_{\mu}^{I} \in \Omega \text{ with } \left[(z_{\mu}^{I}, I) \right]_{\sim_{\Omega}} = \mu \right\}.$$

It follows from (5.7) that

(5.13)
$$\mathbb{S}_{\mu} = \begin{cases} \mathbb{S}, & z_{\mu} \in \Omega_{2ps}, \\ \{I_{\mu}\}, & \text{otherwise}, \end{cases}, \quad \forall \; \mu \in \check{\Omega},$$

where we denote by I_{μ} the unique element in \mathbb{S}_{μ} .

Proposition 5.3. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, and $\mu \in \check{\Omega}$. If $\mu =$ $[(w^J,J)]_{\sim_{\Omega}}$ for some $w \in \mathbb{C}$ and $J \in \mathbb{S}$, then $w = z_{\mu}$.

Proposition 5.4. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $I \in \mathbb{S}$ and $U \in \tau(\Omega_I)$. Then

$$(5.14) \qquad \qquad \left(i_{\Omega}^{J}\right)^{-1}\left([U\times\{I\}]_{\sim_{\Omega}}\right)\in\tau\left(\Omega_{J}\times\{J\}\right), \qquad \forall\ J\in\mathbb{S},$$

and

$$[U \times \{I\}]_{\sim_{\Omega}} \in \tau(\breve{\Omega}).$$

Moreover,

$$i_{\Omega}^{I}: \Omega_{I} \times \{I\} \longrightarrow [\Omega_{I} \times \{I\}]_{\sim_{\Omega}},$$

$$u \longmapsto i_{\Omega}^{I}(u).$$

is a homeomorphism.

Proof. According to Proposition 3.4, $\Omega_{2ps} \in \tau(\mathbb{C})$. It implies by definition that $\Omega^{2ps} \cap \mathbb{C}_J = (\Omega_{2ps})^J$ is open in \mathbb{C}_J . Then

$$\Psi_I^J(U) \cap \Omega^{2ps} = \Psi_I^J(U) \cap \mathbb{C}_J \cap \Omega^{2ps} = \Psi_I^J(U) \cap (\mathbb{C}_J \cap \Omega^{2ps}) \in \tau(\mathbb{C}_J).$$

It follows from (5.10) that (5.14) holds. And (5.15) holds by (5.14) and (5.6).

Let $v, u \in \Omega_I \times \{I\}$ with $\iota_{\Omega}^I(u) = \iota_{\Omega}^I(v) =: \mu$. Then $u, v \in (\iota_{\Omega}^I)^{-1}(\mu)$. By (5.11), $z_u = z_\mu = z_v$. Therefore

$$u = (z_u^I, I) = (z_v^I, I) = v.$$

It implies that i_{Ω}^{I} is injective. And by definition i_{Ω}^{I} is surjective. Let $V \in \tau(\Omega_{I} \times \{I\})$. Then $V = W \times \{I\}$ for some $W \in \tau(\Omega_{I})$. By (5.15),

$$\left[\left(\imath_{\Omega}^{I}\right)^{-1}\right]^{-1}(V)=\imath_{\Omega}^{I}(W\times\{I\})=i_{\Omega}^{I}(W\times\{I\})=[W\times\{I\}]_{\sim_{\Omega}}\in\tau(\check{\Omega})$$

Hence $(i_{\Omega}^{I})^{-1}$ is continuous. Since i_{Ω}^{I} is continuous, so is i_{Ω}^{I} . In summary, i_{Ω}^{I} is a homeomorphism.

Let $z \in \mathbb{C}$, $I \in \mathbb{S}$ and recall that $B_I(z^I, r) := [B_{\mathbb{C}}(z, r)]^I$ denotes the ball with center z^I and radius $r \geq 0$ in \mathbb{C}_I , where $B_{\mathbb{C}}(z,r)$ denotes the open ball with center z and radius r in \mathbb{C} . Note that $B_{\mathbb{C}}(z,0) = \emptyset$.

Let $\mu \in \tilde{\Omega}$. Denote

(5.16)
$$r_{\mu}^{I} := \begin{cases} \delta_{I}(z_{\mu}^{I}, \mathbb{C}_{I} \setminus \Omega_{I}), & I \in \mathbb{S}_{\mu}, \\ 0, & \text{otherwise,} \end{cases}$$

where δ_I is the distance function on \mathbb{C}_I . Moreover, we set

$$(5.17) r_{\mu}^{+} := \sup_{I \in \mathbb{S}} r_{\mu}^{I}, r_{\mu}^{-} := \inf_{I \in \mathbb{S}} r_{\mu}^{I},$$

$$(5.18) \mathbb{S}_{\mu}^{+} := \{ I \in \mathbb{S} : r_{\mu}^{I} = r_{\mu}^{+} \}, \text{and} \mathbb{S}_{\mu}^{-} := \{ I \in \mathbb{S} : r_{\mu}^{I} = r_{\mu}^{-} \}.$$

Proposition 5.5. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, and $\mu \in \check{\Omega}$. Then the following statements holds:

$$(i)$$
 $|\mathbb{S} \setminus \mathbb{S}_{\mu}^{-}| \leq 1$.

(5.19)
$$\mathbb{S}_{\mu}^{+} = \begin{cases} \mathbb{S} \backslash \mathbb{S}_{\mu}^{-}, & |\mathbb{S}_{\mu}^{+}| = 1, \\ \mathbb{S} = \mathbb{S}_{\mu} = \mathbb{S}_{\mu}^{-}, & otherwise. \end{cases}$$

Moreover

$$(5.20) \hspace{1cm} r_{\mu}^{\lambda} = r_{\mu}^{I}, \hspace{1cm} \forall \hspace{1mm} I \in \mathbb{S}_{\mu}^{\lambda}, \hspace{1cm} \lambda \in \{+, -\}.$$

(iii) Let
$$r \in (0, r_{\mu}^+]$$
. Then

$$(5.21) B_{\check{\Omega}}(\mu, r) := \left[\left(B_{\mathbb{C}}(z_{\mu}, r) \right)^{I} \times \left\{ I \right\} \right]_{\mathcal{C}_{0}},$$

does not depend on the choice of $I \in \mathbb{S}_{\mu}$ with $B_I(z_{\mu}^I, r) \subset \Omega$.

(iv)
$$\beta_{\check{\Omega}} := \{B_{\check{\Omega}}(\nu, r) : \nu \in \check{\Omega}, \ 0 < r \le r_{\nu}^+\} \text{ is a basis for the topology } \tau(\check{\Omega}).$$

Proof. (i) Proof by contradiction. If $|\mathbb{S}\backslash\mathbb{S}^-| \geq 2$, then there is $(I,J) \in \mathbb{S}^2_*$ such that

$$r_{\mu}^{I} \ge r_{\mu}^{J} > r_{\mu}^{-} \ge 0.$$

By (5.16), $I, J \in \mathbb{S}_{\mu}$. It follows from (5.13) that $z_{\mu} \in \Omega_{2ps}$ and $\mathbb{S}_{\mu} = \mathbb{S}$. Let $U := B_{\mathbb{C}}(z_{\mu}, r_{\mu}^{J})$. It is easy to check by (5.16) that $U^{I} \subset \Omega_{I}$ and $U^{J} \subset \Omega_{J}$. According to Proposition 3.9, $U \subset \Omega_{2ps}$. Hence

$$U^K \subset \Omega_K, \quad \forall K \in \mathbb{S} = \mathbb{S}_{\mu}.$$

Again by (5.13), $r_{\mu}^{K} \geq r_{\mu}^{J}$. Therefore $r_{\mu}^{-} = \inf_{L \in \mathbb{S}} r_{\mu}^{L} \geq r_{\mu}^{J}$, a contradiction. (ii) Suppose that $|\mathbb{S} \setminus \mathbb{S}_{\mu}^{-}| = 0$. Then $\mathbb{S} = \mathbb{S}_{\mu}^{-}$. By (5.18),

$$(5.22) r_{\mu}^{I} = r_{\mu}^{-}, \forall I \in \mathbb{S}.$$

It implies from (5.17) that $r_{\mu}^+ = r_{\mu}^-$ and $\mathbb{S}_{\mu}^+ = \mathbb{S}_{\mu}^-$. According to (5.16) and (5.22), $\mathbb{S}_{\mu} = \mathbb{S}$. In summary,

$$\mathbb{S}_{\mu}^{+} = \mathbb{S}_{\mu}^{-} = \mathbb{S}_{\mu} = \mathbb{S}$$

and then (5.19) holds.

Suppose that $|\mathbb{S}\backslash\mathbb{S}_{\mu}^-|=1$. Then $\mathbb{S}\backslash\mathbb{S}_{\mu}^-=\{I\}$ for some $I\in\mathbb{S}$. By definition, $r_{\mu}^{+} \neq r_{\mu}^{-}$. It is easy to check that $\mathbb{S}_{\mu}^{+} = \{I\} = \mathbb{S} \setminus \mathbb{S}_{\mu}^{-}$. Therefore (5.19) holds by (i). (iii) We only need to prove (iii) holds when $|\mathbb{S}_{\mu}| > 1$. Let $I, J \in \mathbb{S}_{\mu}$ with $I \neq J$, $B_I(z_\mu^J,r)\subset\Omega$ and $B_J(z_\mu^J,r)\subset\Omega$. By similar method in (i), $B_{\mathbb{C}}(z_\mu,r)\subset\Omega_{2ps}$. By (5.9),

$$\left[(B_{\mathbb{C}}(z_{\mu}, r))^J \times \{J\} \right]_{\sim_{\Omega}} = \left[(B_{\mathbb{C}}(z_{\mu}, r))^I \times \{I\} \right]_{\sim_{\Omega}}.$$

It clear that (iii) holds.

(iv) According to (5.15),

(5.23)
$$B_{\check{\Omega}}(\nu, r) \in \tau(\check{\Omega}), \quad \forall \nu \in \check{\Omega} \text{ and } r \in (0, r_{\nu}^{+}].$$

Let $U \in \tau(\check{\Omega})$ and $\xi \in U$. We will find $V \in \beta_{\check{\Omega}}$ such that $\xi \in V \subset U$: Fix $J \in \mathbb{S}_{\xi}$. Then by (5.6),

$$(z_{\xi}^{J}, J) \in (i_{\Omega}^{J})^{-1}(U) \in \tau(\Omega_{J} \times \{J\}).$$

It implies that there is r > 0 such that

$$[B_{\mathbb{C}}(z_{\xi},r)]^{J} \times \{J\} = B_{J}(z_{\xi}^{J},r) \times \{J\} \subset (i_{\Omega}^{J})^{-1}(U) \subset \Omega_{J} \times \{J\}.$$

Hence

(5.25)
$$\xi \in \left[\left[B_{\mathbb{C}}(z_{\xi}, r) \right]^{J} \times \{J\} \right]_{\sim_{\Omega}} \subset U.$$

By (5.24), $[B_{\mathbb{C}}(z_{\xi},r)]^J \subset \Omega$. It implies that $r \leq r_{\xi}^J \leq r_{\xi}^+$. Then

$$\left[\left[B_{\mathbb{C}}(z_{\xi},r)\right]^{J}\times\{J\}\right]_{\sim_{\Omega}}=B_{\breve{\Omega}}(\xi,r)\in\beta_{\breve{\Omega}}.$$

According (5.23) and (5.25), $\beta_{\check{\Omega}}$ is a basis for $\tau(\check{\Omega})$.

Proposition 5.6. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $\mu, \nu \in \check{\Omega}$ with $z_{\mu} = z_{\nu} \in \overline{\Omega_{2ps}}$. Then $\mu = \nu$.

Proof. If $z_{\mu} \in \Omega_{2ps}$, then choose $I \in \mathbb{S}_{\mu}$. By definition, $\mu = [(z_{\mu}^{I}, I)]_{\sim_{\Omega}}$. According to (5.7),

$$\begin{split} \left(i_{\Omega}\right)^{-1}\left(\mu\right) &= \left(i_{\Omega}\right)^{-1}\left([(z_{\mu}^{I},I)]_{\sim_{\Omega}}\right) = \left\{\left(z_{\mu}^{J},J\right): J \in \mathbb{S}\right\} \\ &= \left\{\left(z_{\nu}^{J},J\right): J \in \mathbb{S}\right\} = \left(i_{\Omega}\right)^{-1}\left(\nu\right). \end{split}$$

It implies that $\mu = \nu$.

Otherwise $z_{\mu} \in \overline{\Omega_{2ps}} \backslash \Omega_{2ps} = \partial(\Omega_{2ps})$. Let $I \in \mathbb{S}_{\mu}$ and $J \in \mathbb{S}_{\nu}$. Then $z_{\mu}^{I} \in \Omega_{I}$, $z_{\nu}^{J} \in \Omega_{J}$ and

$$z_{\mu} = z_{\nu} \in \partial(\Omega_{2ps}) \bigcap \Psi_I^i(\Omega_I) \bigcap \Psi_J^i(\Omega_J).$$

It follows by (3.5) that I = J. Then

$$\mu=[(z_\mu^I,I)]_{\sim_\Omega}=[(z_\nu^I,I)]_{\sim_\Omega}=\nu.$$

Let $\Omega \subset \mathbb{H}$ be a 2-path-symmetric st-domain. According to (5.11), we define the map

$$\pi_{\check{\Omega}}: \ \check{\Omega} \longrightarrow \mathbb{C},$$

$$\mu \longmapsto z_{\mu}.$$

Proposition 5.7. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $I \in \mathbb{S}$ and $U \in \tau(\Omega_I)$. Then

is a homeomorphism. Therefore, $\pi_{\breve{\Omega}}$ is a local homeomorphism.

Proof. (i) It is easy to check by (5.7) that the preimage

$$(5.27) \qquad \left(\pi_{\breve{\Omega}}|_{[U\times\{I\}]_{\sim_{\Omega}}}\right)^{-1}(z) = \left\{\left[\left(z^{I},I\right)\right]_{\sim_{\Omega}}\right\}, \qquad \forall \ z\in\Psi_{I}^{i}(U),$$

is a singleton. It follows that $\pi_{\breve{\Omega}}|_{[U\times\{I\}]_{\sim_{\Omega}}}$ is bijective, moreover $\pi_{\Omega}|_{[U\times\{I\}]_{\sim_{\Omega}}}$ is continuous. In fact, let $V\in\tau\left(\Psi_{I}^{i}(U)\right)$. It implies that

$$V^{I} \in \tau (U) \subset \tau (\Omega_{I})$$
.

It follows from (5.27) and (5.15) that

$$\left(\pi_{\breve{\Omega}}|_{[U\times\{I\}]_{\sim_{\Omega}}}\right)^{-1}(V) = \left[\Psi_{i}^{I}(V)\times\{I\}\right]_{\sim_{\Omega}} \in \tau(\breve{\Omega}).$$

Since $\left[\Psi_i^I(V) \times \{I\}\right]_{\sim_{\Omega}} \subset \left[U \times \{I\}\right]_{\sim_{\Omega}}$, we have

$$\left(\pi_{\breve{\Omega}}|_{[U\times\{I\}]_{\sim_{\Omega}}}\right)^{-1}(V)\in\tau\left(\left[U\times\{I\}\right]_{\sim_{\Omega}}\right).$$

It implies that $\pi_{\breve{\Omega}}|_{[U \times \{I\}]_{\sim_{\Omega}}}$ is continuous.

Also $\left(\pi_{\breve{\Omega}}|_{[U\times\{I\}]_{\sim\Omega}}\right)^{-1}$ is continuous: let $W\in\tau\left([U\times\{I\}]_{\sim\Omega}\right)$. Then there is $D\subset U$ such that $W=[D\times\{I\}]_{\sim\Omega}$. By (5.10) and (5.6),

$$D \times \{I\} = \left(i_{\Omega}^{I}\right)^{-1} \left(\left[D \times \{I\}\right]_{\sim_{\Omega}}\right) \in \tau\left(\Omega_{I} \times \{I\}\right).$$

Since $D \subset U$, we have $D \in \tau(U)$ and $\Psi_I^i(D) \in \tau(\Psi_I^i(U))$. Therefore by (5.26)

$$\left(\left(\pi_{\breve{\Omega}}|_{[U\times\{I\}]_{\sim_{\Omega}}}\right)^{-1}\right)^{-1}(W) = \pi_{\breve{\Omega}}|_{[U\times\{I\}]_{\sim_{\Omega}}}\left([D\times\{I\}]_{\sim_{\Omega}}\right) = \Psi_{I}^{i}(D) \in \tau\left(\Psi_{I}^{i}(U)\right).$$

It implies that $(\pi_{\tilde{\Omega}}|_{[U\times\{I\}]_{\sim_{\Omega}}})^{-1}$ is continuous. In summary, $\pi_{\tilde{\Omega}}|_{[U\times\{I\}]_{\sim_{\Omega}}}$ is a homeomorphism.

(ii) According to (5.15), for each $\mu \in \check{\Omega}$, the set $[\Omega_{I_{\mu}} \times \{I_{\mu}\}]_{\sim_{\Omega}}$ is an open neighborhood of μ . It follows from (i) that π_{Ω} is a local homeomorphism.

Proposition 5.8. Let $\Omega \subset \mathbb{H}$ be a 2-path-symmetric st-domain. Then $(\check{\Omega}, \tau)$ is Hausdorff.

Proof. Let $\mu, \nu \in \check{\Omega}$ with $\mu \neq \nu$. According to Proposition 5.5 (iii), there is r' > 0 such that $\mu \in B_{\check{\Omega}}(\mu, r') \in \tau(\check{\Omega})$, and $\nu \in B_{\check{\Omega}}(\nu, r') \in \tau(\check{\Omega})$. If there is $r \in (0, r')$ such that

$$(5.28) B_{\breve{O}}(\mu, r) \cap B_{\breve{O}}(\mu, r) = \varnothing,$$

then $\tau(\check{\Omega})$ is Hausdorff.

(i) If $z_{\mu} \neq z_{\nu}$, then set $r = \min \left\{ \frac{1}{2} |z_{\mu} - z_{\nu}|, r' \right\}$. By (5.10),

$$(i_{\Omega}^{I})^{-1}\left(B_{\check{\Omega}}(\xi,r)\right)\subset B_{I}(z_{\xi}^{I},r)\times\{I\},\qquad \xi=\mu,\nu,\ \forall\ I\in\mathbb{S}.$$

Hence

$$\left(\left(i_{\Omega}^{I}\right)^{-1}\left(B_{\widecheck{\Omega}}(\mu,r)\right)\right)\bigcap\left(\left(i_{\Omega}^{I}\right)^{-1}\left(B_{\widecheck{\Omega}}(\nu,r)\right)\right) \qquad \forall \ I\in\mathbb{S}.$$

$$\subset\left(B_{I}(z_{\mu}^{I},r)\times\left\{I\right\}\right)\cap\left(B_{I}(z_{\nu}^{I},r)\times\left\{I\right\}\right)=\varnothing,$$

It implies that

$$(5.29) \qquad \left(\left(i_{\Omega} \right)^{-1} \left(B_{\check{\Omega}}(\mu, r) \right) \right) \bigcap \left(\left(i_{\Omega} \right)^{-1} \left(B_{\check{\Omega}}(\nu, r) \right) \right) = \emptyset$$

Then (5.28) holds.

(ii) Otherwise, $z_{\mu} = z_{\nu}$. By Proposition 5.6, $z_{\mu} \notin \overline{\Omega_{2ps}}$. Then there is $r \in (0, r')$ such that $B_{\mathbb{C}}(z_{\mu}, r) \cap \Omega_{2ps} = \emptyset$. It follows from (5.10) that

$$(5.30) (i_{\Omega})^{-1} \left(B_{\check{\Omega}}(\xi, r) \right) = \bigcup_{I \in \mathbb{S}_{\varepsilon}} \left(B_{I}(z_{\xi}^{I}, r) \right), \xi = \mu, \nu.$$

Since $\mu \neq \nu$, we have $\mathbb{S}_{\mu} \cap \mathbb{S}_{\nu} = \emptyset$. It follows from (5.30) that (5.29) holds, and then (5.28) holds.

Proposition 5.9. Let $\Omega \subset \mathbb{H}$ be a 2-path-symmetric st-domain, and $\mu, \nu \in \check{\Omega}$. Then there is a slice-piecewise path $\beta = (\beta_1, ..., \beta_m)$ from $p \in \Omega$ to $q \in \Omega$, and $I_1, ..., I_m \in \mathbb{S}$ such that

$$\mu = [(p, I_1)]_{\sim_{\Omega}}, \quad \nu = [(q, I_m)]_{\sim_{\Omega}}, \quad and \quad \beta_i \subset \Omega_{I_i}.$$

Proof. By definition, there is $p, q \in \Omega$ and $J_1, J_m \in \mathbb{S}$ such that

$$\mu = [(p, J_1)]_{\sim_{\Omega}}$$
 and $\nu = [(q, J_m)]_{\sim_{\Omega}}$.

According to Proposition 4.2, there is a slice-piecewise path $\beta = (\beta_1, ..., \beta_m)$ in Ω from p to q. Then

$$\beta_1 \subset \mathbb{C}_{K_1}, \qquad \beta_m \subset \mathbb{C}_{K_m} \quad \text{and} \quad \beta_i \subset \mathbb{C}_{I_i}, \qquad i = 2, ..., m-1,$$

for some $K_1, K_m, I_i \in \mathbb{S}$.

We will choose $I_1 \in \{J_1, K_1\}$ such that $\mu = [(p, I_1)]_{\sim_{\Omega}}$ and $\beta_1 \subset \Omega_{I_1}$: If $K_1 = \pm J_1$, then let $I_1 = J_1$. It implies that $\beta_1 \subset \mathbb{C}_{K_1} = \mathbb{C}_{J_1} = \mathbb{C}_{I_1}$. Otherwise $K_1 \neq \pm J_1$. Let $I_1 = K_1$. It follows that $p \in \mathbb{C}_{J_1} \cap \mathbb{C}_{K_1} = \mathbb{R}$. By definition, $p \in \Omega^{2ps}$. Then

$$\mu = [(p, J_1)]_{\sim \Omega} = [(p, K_1)]_{\sim \Omega} = [(p, I_1)]_{\sim \Omega}.$$

Similarly, there is $I_m \in \{J_m, K_m\}$ such that $\nu = [(q, I_m)]_{\Omega}$ and $\beta_m \subset \Omega_{I_m}$.

Recall that (see e.g. [22, Page 88]) a Riemann domain over \mathbb{C} is a pair (G, π) with the following properties:

- (i) G is a connected Hausdorff space.
- (ii) $\pi: G \to \mathbb{C}$ is a local homeomorphism.

Theorem 5.10. Let $\Omega \subset \mathbb{H}$ be a 2-path-symmetric st-domain. Then $(\check{\Omega}, \pi_{\check{\Omega}})$ is a Riemann domain over \mathbb{C} .

Proof. Let $\mu, \nu \in \check{\Omega}$. By Proposition 5.9, there is a slice-piecewise path $\beta = (\beta_1, ..., \beta_m)$ from $p \in \Omega$ to $q \in \Omega$, and $I_1, ..., I_m \in \mathbb{S}$ such that

$$\mu = [(p, I_1)]_{\sim_{\Omega}}, \qquad \nu = [(q, I_m)]_{\sim_{\Omega}} \quad \text{and} \quad \beta_i \subset \Omega_{I_i}.$$

Note that $i_{\Omega}^{I_i} \circ \beta_i'$ is a path in $\tilde{\Omega}$, where

$$\beta_i': [0,1] \longrightarrow \Omega_{I_i} \times \{I_i\},$$

$$t \longmapsto (\beta(t), I_i).$$

It implies that

$$\prod_{i=1}^{m} \left(i_{\Omega}^{I_{i}} \circ \beta_{i}' \right) = \left(i_{\Omega}^{I_{1}} \circ \beta_{1}' \right) \circ \cdots \circ \left(i_{\Omega}^{I_{m}} \circ \beta_{m}' \right)$$

is a path in $\check{\Omega}$ from $\mu = [(p, I_1)]_{\sim_{\Omega}}$ to $\nu = [(q, I_m)]_{\sim_{\Omega}}$. Hence $\check{\Omega}$ is path-connected and then connected. According to Propositions 5.7 and 5.8, $\check{\Omega}$ is Hausdorff and $\pi_{\check{\Omega}}$ is a local homeomorphism. In summary, $(\check{\Omega}, \pi_{\check{\Omega}})$ is a Riemann domain over \mathbb{C} . \square

In the sequel we shall refer to $\check{\Omega}$ as the 2-path-symmetric Riemann domain associated with Ω .




FIGURE 3.

Example 5.11. Let $E_1, U_{a,b}, I, J$ be defined as in Example 2.2. Then

$$\breve{E}_1 \cong \left[\left(U'_{-\frac{\pi}{2}, \frac{3\pi}{4}} \times \{1\} \right) \bigcup \left(U'_{\frac{\pi}{4}, \frac{3\pi}{2}} \times \{2\} \right) \right] / \sim,$$

with the structure-preserving map

$$\phi\left(\left[(z^K,K)\right]_{\sim_{\Omega}}\right) = \begin{cases} [(z,1)]_{\sim}, & K = \pm I, \\ [(z,2)]_{\sim}, & otherwise, \end{cases} \quad \forall \ z^K \in E_1,$$

where

$$U'_{a,b} := U_{a,b} \cap \operatorname{Conj}_{\mathbb{C}}(U_{a,b})$$

and $(z,n) \sim (w,m)$ with $(z,n) \neq (w,m)$ if and only if $n \neq m$ and $z = w \in U_{0,3\pi} \cap \operatorname{Conj}_{\mathbb{C}}(U_{0,3\pi})$.

Example 5.12. Let E_2 , V_{ℓ} , W_{ℓ} , I_{ℓ} be defined as in Example 2.3. Then

$$reve{E_2} \cong igcup_{\ell=1}^{+\infty} \left[V_\ell \cup W_\ell \cup \operatorname{Conj}_\mathbb{C}(W_\ell) \right] imes \left\{ \ell \right\} \middle/ \sim$$

with the structure-preserving map

$$\phi\left([(z^K, K)]_{\sim_{\Omega}}\right) = [(z, \ell)]_{\sim}, \qquad \forall \ z^K \in (V_{\ell})^{\mathbb{S}} \cup (W_{\ell})^{I_{\ell}} \cup (V_{\ell+1})^{\mathbb{S}},$$

where $(z,n) \sim (w,m)$ with $(z,n) \neq (w,m)$ if and only if |n-m| = 1 and $z = w \in V_{\max(n,m)}$. See Figure 3 where the broken lines denote different branches.

Remark 5.13. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric. Then $\check{\Omega}$ is the disjoint union of some Riemann domains over \mathbb{C} , i.e.

$$\breve{\Omega} = \bigcup_{U \in \mathcal{SCC}(\Omega)} \breve{U},$$

where $SCC(\Omega)$ is the family of slice-connected components of Ω .

Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $\mu \in \check{\Omega}$ and $r \in (0, r_{\mu}^+]$. By Proposition 5.5 (iii),

$$\pi_{\breve{\Omega}}\left[B_{\breve{\Omega}}(\mu,r)\right] = \pi_{\breve{\Omega}}\left[\left[\left(B_{\mathbb{C}}\left(z_{\mu},r\right)\right)^{I} \times \left\{I\right\}\right]_{\sim_{\Omega}}\right] = B_{\mathbb{C}}\left(z_{\mu},r\right).$$

Let $r_1, r_2 \in (0, r)$ with $r_1 + r_2 \le r$, and let $\nu \in B_{\breve{o}}(\mu, r_1)$. Then

$$z_{\nu} = \pi_{\breve{\Omega}}(\nu) \in \pi_{\breve{\Omega}} \left[B_{\breve{\Omega}}(\mu, r_1) \right] = B_{\mathbb{C}} \left(z_{\mu}, r_1 \right),$$

and

$$(5.31) B_{\tilde{\Omega}}(\nu, r_2) = \left[(B_{\mathbb{C}}(z_{\nu}, r_2))^I \times \{I\} \right]_{\sim_{\Omega}}$$

$$\subset \left[(B_{\mathbb{C}}(z_{\mu}, r))^I \times \{I\} \right]_{\sim_{\Omega}} = B_{\tilde{\Omega}}(\mu, r).$$

Proposition 5.14. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric. The map

$$r^+: \check{\Omega} \longrightarrow (0, +\infty) =: \mathbb{R}^+,$$

$$\mu \longmapsto r_{\mu}^+,$$

is continuous.

Proof. Let $U \in \tau(\mathbb{R}^+)$ and $\mu \in (r^+)^{-1}(U)$. Then there is $t_1 \in (0, r_\mu^+)$ such that $B_{\mathbb{R}}(r_\mu^+, t_1) \subset U$. Let $\nu \in B_{\breve{\Omega}}(\mu, t_1/2)$. By (5.31),

$$B_{\breve{\Omega}}(\nu, r_{\mu}^+ - t_1/2) \subset B_{\breve{\Omega}}(\mu, r_{\mu}^+),$$

and $\nu \in (r^+)^{-1}(U)$. Hence μ is a interior point in $(r^+)^{-1}(U)$, and $(r^+)^{-1}(U)$ is open. It is clear that r^+ is continuous.

Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, and $U \subset \check{\Omega}$. Denote

$$r^+(U) := r_U^+ := \inf_{\mu \in U} r_\mu^+.$$

If U is compact, then by (5.20).

$$r_U^+ = r_\nu^+ > 0$$
, for some $\nu \in U$.

6. Some properties of 2-path-symmetric Riemann domains

In this section, we carefully study a relation between the points in Ω and the points in the associated 2-path-symmetric Riemann domain $\check{\Omega}$. This relation is useful to construct slice regular functions that cannot be extended slice regularly to a larger domain, in the sense of Riemann domains. Moreover we prove a number of technical results which are crucial in the proof of the interpolation result in Section 8 and in the proof of the Cartan-Thullen theorem.

Let (Ω, π) be a Riemann domain over \mathbb{C} and $U \subset \Omega$. Denote by

$$\delta_{\Omega}(U, \partial\Omega) := \sup \{r > 0 : \exists B'_{\Omega}(z, r), \forall z \in U\},$$

the boundary distance of U on Ω , where $B'_{\Omega}(z,r)$ is the unique subset (if it exists) of Ω such that $z \in B'_{\Omega}(z,r)$ and $\pi: B'_{\Omega}(z,r) \to B_{\mathbb{C}}(\pi(z),r)$ is a homeomorphism. Let

$$U_{\Omega}^{\wedge} := \widehat{U}_{\Omega} := \left\{ z \in \Omega : |f(z)| \le \sup_{w \in U} |f(w)|, \ f \in \mathscr{O}(\Omega) \right\}$$

be the holomorphically convex hull, where $\mathcal{O}(\Omega)$ is the class of holomorphic functions on Ω . Note that, by definition

$$(6.1) U \subset \widehat{U}_{\Omega}.$$

Recall each (1-dimensional) Riemann domain over \mathbb{C} is a domain of holomorphy. By [36, Lemma 7.5.5] we have

$$\delta_{\Omega}(U,\partial\Omega) = \delta_{\Omega}(\widehat{U}_{\Omega},\partial\Omega).$$

By Theorem 5.10 we know that if $\Omega \subset \mathbb{H}$ is a 2-path-symmetric st-domain then $(\check{\Omega}, \pi_{\Omega})$ is a Riemann domain over \mathbb{C} . It follows from

$$\pi_{\check{\Omega}}\left[B_{\check{\Omega}}(\mu,r)\right] = B_{\mathbb{C}}(z_{\mu},r) = \pi_{\check{\Omega}}\left[B_{\check{\Omega}}'(\mu,r)\right], \quad \forall \ \mu \in \check{\Omega} \quad \text{and} \quad r \leq r_{\mu}^{+},$$

that

$$B_{\check{\Omega}}(\mu,r) = B'_{\check{\Omega}}(\mu,r), \qquad \forall \ \mu \in \check{\Omega} \quad \text{and} \quad r \le r_{\mu}^+,$$

where $B_{\check{\alpha}}(\mu, r)$ is defined by (5.21).

Note that sometimes, for graphical reasons, it is convenient to write U_{Ω}^{\wedge} instead of \widehat{U}_{Ω} , like in the result below:

Proposition 6.1. Let Ω be a Riemann domain over \mathbb{C} , $K \subset \Omega$ be compact with $\hat{K}_{\Omega} = K$, $U \in \tau(\Omega)$ and r > 0. Then $K \cap U^r$ is compact in U with

$$(6.2) (K \cap U^r)_U^{\wedge} = K \cap U^r,$$

where

$$U^r := \{ z \in U : \delta_U(z, \partial U) \ge r \}.$$

Proof. Since

$$\delta_U((K \cap U^r)_U^{\wedge}, \partial U) = \delta_U(K \cap U^r, \partial U) \ge \delta_U(U^r, \partial U) = r,$$

we have

$$(6.3) (K \cap U^r)_U^{\wedge} \subset U^r.$$

Let $z \in U \backslash K$. Then there is $f \in \mathscr{O}(\Omega)$ such that

$$|f(z)| \ge \sup_{w \in K} |f(w)| \ge \sup_{w \in K \cap U^r} |f(w)|.$$

It implies that $z \notin (K \cap U^r)_U^{\wedge}$. Therefore $(K \cap U^r)_U^{\wedge} \subset K$, and then (6.2) holds by (6.3) and (6.1).

The σ -ball with center $q \in \mathbb{H}$ and radius s > 0 is introduced in [26] as follows:

$$\Sigma(q, s) := \{ q \in \mathbb{H} : \sigma(p, q) < s \}$$

and is induced by the so-called σ -distance

$$\sigma(p,q) := \begin{cases} |q-p|, & \exists \ I \in \mathbb{S}, \ s.t. \ p,q \in \mathbb{C}_I, \\ \sqrt{(Re(q-p))^2 + |Im(q)|^2 + |Im(p)|^2}, & \text{otherwise.} \end{cases}$$

Let $p \in \mathbb{H}$ and $U, V \subset \mathbb{H}$. Set

$$\sigma_V^p := \sigma(p,V) := \inf_{q \in V} \left(\sigma(p,q) \right), \qquad \text{and} \qquad \sigma_V^U := \sigma(U,V) := \inf_{p \in U, q \in V} \left(\sigma(p,q) \right).$$

Proposition 6.2. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $I \in \mathbb{S}$, $p \in \Omega_I$ and $U \subset \Omega_I$. Then

(6.4)
$$\sigma(p, \mathbb{H} \backslash \Omega) = \delta_I(p, \mathbb{C}_I \backslash \Omega_I),$$

and

(6.5)
$$\sigma(U, \mathbb{H} \backslash \Omega) = \delta_I(U, \mathbb{C}_I \backslash \Omega_I).$$

Proof. $\sigma^p_{\mathbb{H}\setminus\Omega} \leq \delta_I(p,\mathbb{C}_I\setminus\Omega_I)$ holds directly by $B_I(p,r)\subset\Sigma(p,r)$ for each r>0. Conversely, suppose $B_I(p,r)\subset\Omega$ for some fixed r>0. If $B_I(p,r)\cap\mathbb{R}=\varnothing$, then

$$\Sigma(p,r) = B_I(p,r) \subset \Omega.$$

Since the choice of r > 0 with $B_I(p, r) \subset \Omega$ is arbitrary,

(6.6)
$$\sigma(p, \mathbb{H} \backslash \Omega) \ge \delta_I(p, \mathbb{C}_I \backslash \Omega_I)$$

and (6.4) holds. Otherwise $B_I(p,r) \cap \mathbb{R} \neq \emptyset$. Then for each $x+yJ \in \Sigma(p,r)$, it is each to check that

$$\mathcal{L}_x^{x+yi} \in \mathscr{P}_0(\mathbb{C}) \quad \text{and} \quad (\mathcal{L}_x^{x+yi})^I, (\mathcal{L}_x^{x+yi})^{-I} \subset \Omega.$$

Since Ω is 2-path-symmetric, $(\mathcal{L}_x^{x+yi})^J \subset \Omega$. Therefore

$$x + yJ = (\mathcal{L}_x^{x+yi})^J(1) \in \Omega.$$

Hence $\Sigma(p,r) \subset \Omega$. Since the choice of r > 0 with $B_I(p,r) \subset \Omega$ is arbitrary, (6.6) holds. Therefore (6.4) holds. Then

$$\sigma(U, \mathbb{H} \backslash \Omega) = \inf_{q \in U} \sigma(q, \mathbb{H} \backslash \Omega) = \inf_{q \in U} \delta_I(q, \mathbb{C}_I \backslash \Omega_I) = \delta_I(U, \mathbb{C}_I \backslash \Omega_I),$$

i.e.
$$(6.5)$$
 holds.

Proposition 6.3. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $\mu \in \check{\Omega}$ and $I, J \in \mathbb{S}_{\mu}$. Then (6.7) $\left[\left(\overline{z_{\mu}}^{-I}, -I\right)\right]_{\Omega \cap \sigma} = \left[\left(\overline{z_{\mu}}^{-J}, -J\right)\right]_{\Omega \cap \sigma}.$

Proof. If $z_{\mu} \in \Omega_{2ps}$, then by Remark 3.10 (v) we have $\overline{z_{\mu}} \in \Omega_{2ps}$. It implies from (5.7) that $(\overline{z_{\mu}}^{-I}, -I) \sim_{\Omega} (\overline{z_{\mu}}^{-J}, -J)$. Hence (6.7) holds. Otherwise, $z_{\mu} \notin \Omega_{2ps}$. By (5.13), we have $\mathbb{S}_{\mu} = \{I_{\mu}\}$ and then $I = J = I_{\mu}$. It follows from $\overline{z_{\mu}}^{-I} = z_{\mu}^{I} \in \Omega$ that (6.7) holds.

For each $\mu \in \Omega$, we choose a fixed $J_{\mu} \in \mathbb{S}_{\mu}$. Define

(6.8)
$$\begin{array}{ccc}
\operatorname{Conj}_{\breve{\Omega}} : & \breve{\Omega} & \longrightarrow & \breve{\Omega}, \\
\mu & \longmapsto & \overline{\mu} := \left[\left(\overline{z_{\mu}}^{-J_{\mu}}, -J_{\mu} \right) \right]_{\sim_{\Omega}} = \left[\left(z_{\mu}^{J_{\mu}}, -J_{\mu} \right) \right]_{\sim_{\Omega}}.
\end{array}$$

According to Proposition 6.3, $\operatorname{Conj}_{\check{\Omega}}$ does not depend on the choice of $J_{\mu} \in \mathbb{S}_{\mu}$. It is easy to check that

(6.9)
$$z_{\overline{\mu}} = \overline{z_{\mu}}, \quad \text{and} \quad \mathbb{S}_{\overline{\mu}} = -\mathbb{S}_{\mu}.$$

If $z_{\mu} = \Omega_{2ps}$, then by (5.2),

$$\overline{\mu} = [(\overline{z_{\mu}}^I, I)]_{\sim_{\Omega}}, \quad \forall I \in \mathbb{S}_{\mu} = \mathbb{S}.$$

Proposition 6.4. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $\mu \in \Omega$, and $U \subset \check{\Omega}$. Then

$$r_{\mu}^{+} = r_{\overline{\mu}}^{+}, \quad and \quad r_{U}^{+} = r_{\text{Conjo}(U)}^{+}.$$

Proof. Let $I \in \mathbb{S}_{\mu}^+$. By (5.20), $r_{\mu}^+ = r_{\mu}^I > 0$. Then $-I \in -\mathbb{S}_{\mu} = \mathbb{S}_{\overline{\mu}}$, and

$$B_{-I}\left(z_{\overline{\mu}}^{-I}, r_{\mu}^{+}\right) = B_{-I}\left(\overline{z_{\mu}}^{-I}, r_{\mu}^{+}\right) = B_{I}(z_{\mu}^{I}, r_{\mu}^{I}) \subset \Omega.$$

It implies that $r_{\overline{\mu}}^+ \ge r_{\overline{\mu}}^{-I} \ge r_{\mu}^+$. Similarly, $r_{\mu}^+ = r_{\overline{\overline{\mu}}}^+ \ge r_{\overline{\mu}}^+$. Therefore $r_{\mu}^+ = r_{\overline{\mu}}^+$. Then

$$r_U^+ = \inf_{\nu \in U} r_\nu^+ = \inf_{\nu \in U} r_{\overline{\nu}}^+ = \inf_{\overline{\nu} \in \operatorname{Conj}_{\widetilde{\Omega}}(U)} r_{\overline{\nu}}^+ = r_{\operatorname{Conj}_{\widetilde{\Omega}}(U)}^+.$$

Proposition 6.5. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric. Then $\operatorname{Conj}_{\check{\Omega}} : \check{\Omega} \to \check{\Omega}$ is a homeomorphism.

Proof. Conj_{$\check{\Omega}$} is bijective directly by Conj²_{$\check{\Omega}$} = id_{$\check{\Omega}$}, where id_{$\check{\Omega}$} is the identity map on $\check{\Omega}$. Let $U \in \tau(\check{\Omega})$ and $\mu \in \operatorname{Conj}_{\check{\Omega}}^{-1}(U)$. It implies that

$$\overline{\mu}=\operatorname{Conj}_{\breve{\Omega}}(\mu)=\operatorname{Conj}_{\breve{\Omega}}^{-1}(\mu)\in\operatorname{Conj}_{\breve{\Omega}}^{-1}\left[\operatorname{Conj}_{\breve{\Omega}}^{-1}(U)\right]=U,$$

and there is $r \in (0, r_{\mu}^+)$ such that $B_{\breve{\Omega}}(\overline{\mu}, r) \subset U$. Let $J_{\mu} \in \mathbb{S}_{\mu}$. Then

$$-J_{\mu} \in -\mathbb{S}_{\mu} = \mathbb{S}_{\overline{\mu}}$$

and

$$B_{\breve{\Omega}}(\mu,r) = \left[B_{J_{\mu}} \left(z_{\mu}^{J_{\mu}}, r \right) \times \{J_{\mu}\} \right]_{\sim_{\Omega}} = \operatorname{Conj}_{\breve{\Omega}}^{2} \left(\left[B_{J_{\mu}} \left(z_{\mu}^{J_{\mu}}, r \right) \times \{J_{\mu}\} \right]_{\sim_{\Omega}} \right)$$

$$= \operatorname{Conj}_{\breve{\Omega}} \left(\left[B_{J_{\mu}} \left(\overline{z_{\mu}}^{-J_{\mu}}, r \right) \times \{-J_{\mu}\} \right]_{\sim_{\Omega}} \right)$$

$$= \operatorname{Conj}_{\breve{\Omega}} \left(\left[B_{-J_{\mu}} \left(z_{\overline{\mu}}^{-J_{\mu}}, r \right) \times \{-J_{\mu}\} \right]_{\sim_{\Omega}} \right) = \operatorname{Conj}_{\breve{\Omega}} \left(B_{\breve{\Omega}}(\overline{\mu}, r) \right)$$

$$\subset \operatorname{Conj}_{\breve{\Omega}}(U) = \operatorname{Conj}_{\breve{\Omega}}^{-1}(U).$$

Therefore, μ is an interior point in $\operatorname{Conj}_{\check{\Omega}}^{-1}(U)$, and $\operatorname{Conj}_{\check{\Omega}}^{-1}(U)$ is open. It is clear that $\operatorname{Conj}_{\check{\Omega}}$ is continuous, so is $\operatorname{Conj}_{\check{\Omega}}^{-1} = \operatorname{Conj}_{\check{\Omega}}$. In summary, $\operatorname{Conj}_{\check{\Omega}}$ is a homeomorphism.

Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric and $t \geq 0$ and set

$$(6.10) \qquad \Omega^t := \{ q \in \Omega : t \le \sigma(q, \mathbb{H} \setminus \Omega) \}, \qquad \Omega^{t^\circ} := \{ q \in \Omega : t < \sigma(q, \mathbb{H} \setminus \Omega) \}.$$

$$\breve{\Omega}_+^t := \{ \mu \in \breve{\Omega} : t \leq r_\mu^+ \}, \qquad \text{and} \qquad \breve{\Omega}_+^{t^\circ} := \{ \mu \in \breve{\Omega} : t < r_\mu^+ \}.$$

Proposition 6.6. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $I \in S$, $w \in \Omega_I$ with $\mathbb{S}^+_{[(w,I)]_{\sim_{\Omega}}} = \mathbb{S}$. Then

(6.11)
$$\Omega^r \cap [w]_{\approx_{\Omega}} = \varnothing, \quad \forall \ r > \sigma(w, \mathbb{H} \setminus \Omega).$$

Proof. Let $u \in [w]_{\sim \Omega}$. Then $u \approx_{\Omega} w = z^I$, where $z := \Psi_I^i(w)$. According to Proposition 3.2 (i), $u = z^K$ for some $K \in \mathbb{S}$. By (5.19), $\mathbb{S}_{\mu} = \mathbb{S}_{\mu}^+ = \mathbb{S}$, where $\mu := [(w,I)]_{\sim_{\Omega}} = [(z^I,I)]_{\sim_{\Omega}}$. It follows from $z_{\mu} = z$, (5.16), (5.20) and (6.4) that

$$\sigma(u, \mathbb{H} \setminus \Omega) = \delta_K \left(z^K, \mathbb{C}_K \setminus \Omega_K \right) = r_\mu^K$$

= $r_\mu^I = r_\mu^I = \delta_I \left(z^I, \mathbb{C}_I \setminus \Omega_I \right) = \sigma(w, \mathbb{H} \setminus \Omega) < r$.

According to (6.10), $u \notin \Omega^r$. Then (6.11) holds.

Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $\mathbb{S}' \subset \mathbb{S}$, $t \geq 0$ and $\rho \in \{t, t^{\circ}\}$. Denote

$$\psi_{\Omega}^{\mathbb{S}'}: P(\Omega) \longrightarrow P(\check{\Omega}),$$

$$U \longmapsto \left[\bigcup_{I \in \mathbb{S}'} U_I \times \{I\}\right]_{\sim_{\Omega}},$$

$$\varphi_{\Omega}^{\rho}: P(\check{\Omega}) \longrightarrow P(\Omega),$$

$$(6.12) \qquad V \longmapsto \left\{q \in \Omega^{\rho}: \psi_{\Omega}^{\mathbb{S}}(\{q\}) \cap V \neq \varnothing\right\},$$

$$\phi_{\Omega}^{\rho}: P(\Omega) \longrightarrow P(\Omega),$$

$$U \longmapsto \left[U \cup \left(U \cap \Omega^{2ps}\right)^{\sim}\right] \cap \Omega^{\rho} = [U]_{\approx_{\Omega}}^{\cup} \cap \Omega^{\rho},$$

$$\phi_{\check{\Omega}}^{\rho}: P(\check{\Omega}) \longrightarrow P(\check{\Omega}),$$

$$V \longmapsto \left[V \cup \operatorname{Conj}_{\check{\Omega}}(V)\right] \cap \check{\Omega}_{+}^{\rho},$$

where

$$[U]^{\cup}_{\approx_{\Omega}}:=\bigcup_{V\in [U]_{\approx_{\Omega}}}V=\{q\in\Omega:q\in[p]_{\approx_{\Omega}},\text{ for some }p\in U\},$$

and $(U \cap \Omega^{2ps})^{\sim}$ denotes the axially symmetric completion of $U \cap \Omega^{2ps}$. By definition, ϕ_{Ω}^{ρ} and ϕ_{Ω}^{ρ} are projection, i.e. $(\phi_{\Omega}^{\rho})^2 = \phi_{\Omega}^{\rho}$. Let $x \in \Omega_{\mathbb{R}}$. By (3.4), $x \in \Omega_{\mathbb{R}} \subset \Omega_{2ps}$. According to (5.2),

$$(x^I, I) \sim_{\Omega} (x^J, J), \quad \forall I, J \in \mathbb{S}.$$

Thence for each $J \in \mathbb{S}$,

$$\begin{split} \psi_{\Omega}^{\mathbb{S}}(\{x\}) &= \left[\bigcup_{I \in \mathbb{S}} \{x\}_I \times \{I\}\right]_{\sim_{\Omega}} = \left[\bigcup_{I \in \mathbb{S}} \{(x,I)\}\right]_{\sim_{\Omega}} \\ &= \bigcup_{I \in \mathbb{S}} \left\{\left[(x^I,I)\right]_{\sim_{\Omega}}\right\} = \left[(x^J,J)\right]_{\sim_{\Omega}}. \end{split}$$

It implies that

$$(6.13) \qquad \quad \psi_{\Omega}^{\mathbb{S}}(\{z^I\}) = \left\{ \left[(z^I, I) \right]_{\sim_{\Omega}} \right\} \bigcup \left\{ \left[(\overline{z}^{-I}, -I) \right]_{\sim_{\Omega}} \right\}, \qquad \forall \ z^I \in \Omega.$$

Let $z \in \Omega_{2ps}$ and $I, J \in \mathbb{S}$. Then it follows from

$$[(z^I,I)]_{\sim_{\Omega}} = [(z^J,J)]_{\sim_{\Omega}}, \qquad \text{and} \qquad [(\overline{z}^{-I},-I)]_{\sim_{\Omega}} = [(z^{-J},-J)]_{\sim_{\Omega}}$$

that

(6.14)
$$\psi_{\Omega}^{\mathbb{S}}(\{z^I\}) = \psi_{\Omega}^{\mathbb{S}}(\{z^J\}).$$

Proposition 6.7. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, t > 0, and $\rho \in \{t, t^{\circ}\}$. Then

$$\mu := [(w^I, I)]_{\sim_{\Omega}} \in \breve{\Omega}_+^{\rho}, \quad \forall \ w^I \in \Omega^{\rho}.$$

Moreover,

$$\psi_{\Omega}^{\mathbb{S}}(\Omega^{\rho}) \subset \breve{\Omega}_{+}^{\rho}.$$

Proof. (i) Let $w^I \in \Omega^{\rho}$. By (5.1) and (5.11), $z_{\mu} = z_{(w^I,I)} = w$. There is

(6.15)
$$\varepsilon \in \begin{cases} \{0\}, & \rho = t, \\ (0, +\infty), & \rho = t^{\circ}, \end{cases}$$

such that

$$B_I(z_\mu^I, t + \varepsilon) = B_I(w^I, t + \varepsilon) \subset \Omega_I.$$

According to (5.17) and (5.20), $r_{\mu}^{+} \geq r_{\mu}^{I} \geq t + \varepsilon$. It implies that $\mu \in \check{\Omega}_{+}^{\rho}$. (ii) Let $\xi^{J} \in \Omega^{\rho}$. By (6.13)

$$\psi_{\scriptscriptstyle \Omega}^{\mathbb{S}}(\{\xi^J\}) = \bigg\{ \big[\big(\xi^J,J\big)\big]_{\sim_{\Omega}} \bigg\} \bigcup \bigg\{ \big[\big(\overline{\xi}^{-J},-J\big)\big]_{\sim_{\Omega}} \bigg\}, \qquad \forall \; z^J \in \Omega.$$

It implies by $\overline{\xi}^{-J} = \xi^J \in \Omega^{\rho}$ and (i) that

$$[(\xi^J,J)]_{\sim_{\Omega}},[(\overline{\xi}^{-J},J)]_{\sim_{\Omega}}\in \check{\Omega}^{\rho}_{+}.$$

Therefore

$$\psi_{\scriptscriptstyle \Omega}^{\mathbb{S}}(\Omega^{\rho}) = \bigcup_{\xi^J \in \Omega^{\rho}} \psi_{\scriptscriptstyle \Omega}^{\mathbb{S}}\left(\{\xi^J\}\right) \subset \check{\Omega}_+^{\rho}.$$

The technical result in the next lemma will be useful in the next section.

Lemma 6.8. Let $\Omega \subset \tau_s(\mathbb{H})$ be 2-path-symmetric, t > 0 and $\rho \in \{t, t^{\circ}\}$. Then

(6.16)
$$\begin{cases} \varphi_{\Omega}^{\rho} \circ \psi_{\Omega}^{\mathbb{S}} = \phi_{\Omega}^{\rho}, \\ \psi_{\Omega}^{\mathbb{S}} \circ \varphi_{\Omega}^{\rho} = \phi_{\Omega}^{\rho}. \end{cases}$$

Proof. Let $U \in P(\Omega)$ and $V \in P(\check{\Omega})$.

(i) Suppose $p \in \varphi_{\check{\Omega}}^{\rho} \circ \psi_{\Omega}^{\mathbb{S}}(U)$. Then

$$(6.17) \qquad \varphi_{\Omega}^{\rho} \circ \psi_{\Omega}^{\mathbb{S}}(U) = \varphi_{\Omega}^{\rho} \left(\left[\bigcup_{I \in \mathbb{S}} U_{I} \times \{I\} \right]_{\sim_{\Omega}} \right)$$

$$= \left\{ q \in \Omega^{\rho} : \psi_{\Omega}^{\mathbb{S}}(\{q\}) \cap \left[\bigcup_{I \in \mathbb{S}} U_{I} \times \{I\} \right]_{\sim_{\Omega}} \neq \varnothing \right\}.$$

implies that $p \in \Omega^{\rho}$ and

$$\left[\bigcup_{I\in\mathbb{S}}\{p\}_I\times\{I\}\right]_{\sim_{\Omega}}\cap\left[\bigcup_{I\in\mathbb{S}}U_I\times\{I\}\right]_{\sim_{\Omega}}=\psi_{\scriptscriptstyle\Omega}^{\mathbb{S}}(\{p\})\cap\left[\bigcup_{I\in\mathbb{S}}U_I\times\{I\}\right]_{\sim_{\Omega}}\neq\varnothing,$$

i.e. there is $w \in U$ and $J, K \in \mathbb{S}_{\mu}$ such that

$$\mu:=\left[(p,J)\right]_{\sim_{\Omega}}=\left[(w,K)\right]_{\sim_{\Omega}}\in\left[\bigcup_{I\in\mathbb{S}}\left\{p\right\}_{I}\times\left\{I\right\}\right]_{\sim_{\Omega}}\bigcap\left[\bigcup_{I\in\mathbb{S}}U_{I}\times\left\{I\right\}\right]_{\sim_{\Omega}}.$$

It follows from (5.4) and $(p, J) \sim_{\Omega} (w, K)$ that

$$p = z_{(p,J)}^{I_{(p,J)}} \approx_{\Omega} z_{(w,k)}^{I_{(w,k)}} = w.$$

It implies that $p \in [w]_{\approx_{\Omega}} \subset [U]_{\approx_{\Omega}}$. Hence $p \in [U]_{\approx_{\Omega}} \cap \Omega^{\rho} = \varphi^{\rho}_{\Omega}(U)$. Then

(6.18)
$$\varphi^{\rho}_{\bullet} \circ \psi^{\mathbb{S}}_{\Omega}(U) \subset \varphi^{\rho}_{\Omega}(U).$$

(ii) Suppose $p \in \phi_{\Omega}(U) = [U]_{\approx_{\Omega}} \cap \Omega^{p}$. Then there is $z^{K} \in U$ with $p \approx_{\Omega} z^{K}$. By Proposition 3.2 (i), there is $J \in \mathbb{S}$ such that $p = z^{J}$. According to (3.3), J = K or $z \in \Omega_{2ps}$. Then by (5.2), $(p, J) = (z^{J}, J) \sim_{\Omega} (z^{K}, K)$. Therefore

$$\left[(p,J)\right]_{\sim_{\Omega}} = \left[(z^K,K)\right]_{\sim_{\Omega}} \in \left[U_K \times \{K\}\right]_{\sim_{\Omega}} \subset \left[\bigcup_{I \in \mathbb{S}} U_I \times \{I\}\right]_{\sim_{\Omega}}.$$

Hence

$$[(p,J)]_{\sim_{\Omega}} \in \left[\bigcup_{I \in \mathbb{S}} \left\{p\right\}_{I} \times \left\{I\right\}\right]_{\sim_{\Omega}} \bigcap \left[\bigcup_{I \in \mathbb{S}} U_{I} \times \left\{I\right\}\right]_{\sim_{\Omega}} \neq \varnothing.$$

By (6.17), $p \in \varphi_{\Omega}^{\rho} \circ \psi_{\Omega}^{\mathbb{S}}(U)$. Then $\phi_{\Omega}(U) \subset \varphi_{\Omega}^{\rho} \circ \psi_{\Omega}^{\mathbb{S}}(U)$. According to (6.18), (6.19) $\phi_{\Omega} = \varphi_{\Omega}^{\rho} \circ \psi_{\Omega}^{\mathbb{S}}.$

(iii) Suppose that

$$\mu \in \psi_{\circ}^{\mathbb{S}} \circ \varphi_{\circ}^{\rho}(V) = \psi_{\circ}^{\mathbb{S}} \left\{ q \in \Omega^{\rho} : \psi_{\circ}^{\mathbb{S}}(\{q\}) \cap V \neq \varnothing \right\}.$$

By (6.13) and Proposition 6.7, there is $z^I \in \Omega^{\rho}$ with $\psi_0^{\mathbb{S}}(\{z^I\}) \cap V \neq \emptyset$ such that

$$\mu \in \psi_{\Omega}^{\mathbb{S}}(\{z^{I}\}) = \left\{ \left[(z^{I}, I) \right]_{\sim_{\Omega}} \right\} \bigcup \left\{ \left[(\overline{z}^{-I}, -I) \right]_{\sim_{\Omega}} \right\}$$
$$\subset (V \cup \operatorname{Conj}_{\breve{\Omega}}(V)) \cap \breve{\Omega}_{+}^{\rho} = \phi_{\breve{\Omega}}^{\rho}(U).$$

It implies that

(6.20)
$$\psi_{\Omega}^{\mathbb{S}} \circ \varphi_{\check{\Omega}}^{\rho}(V) \subset \phi_{\check{\Omega}}^{\rho}(U).$$

(iv) Suppose that $\mu \in \varphi_{\Omega}^{\rho}(V) = (V \cap \operatorname{Conj}_{\check{o}}(V)) \cap \Omega^{\rho}$. Then

$$\nu := \operatorname{Conj}_{0}^{\lambda}(\mu) \in V, \quad \text{for some} \quad \lambda \in \{0, 1\}$$

It follows from $\mu \in \check{\Omega}_{+}^{\rho}$ that $r_{\mu}^{+} \geq t + \varepsilon$ for some ε satisfying (6.15). Let $I \in \mathbb{S}_{\mu}^{+}$. Then $r_{\mu}^{+} = r_{\mu}^{I}$ and $B_{I}\left(z_{\mu}^{I}, r_{\mu}^{I}\right) \subset \Omega$. It implies that

$$(6.21) \quad z_{\nu}^{J} = z_{\operatorname{Conj}_{\Omega}^{\lambda}(\mu)}^{J} = \operatorname{Conj}_{\mathbb{C}}^{\lambda}(z_{\mu})^{(-1)^{\lambda}I} = z_{\mu}^{I} \in \Omega^{\rho}, \quad \text{where} \quad J = (-1)^{\lambda}I.$$

Hence

$$\nu = \left[\left(\operatorname{Conj}_{\check{\Omega}}^{\lambda}(z_{\mu})^{(-1)^{\lambda}I}, (-1)^{\lambda}I \right) \right]_{\sim_{\Omega}} = \left[\left(z_{\nu}^{J}, J \right) \right]_{\sim_{\Omega}} \in \left[\bigcup_{I \in \mathbb{S}} \left\{ z_{\nu}^{J} \right\}_{I} \times \left\{ I \right\} \right]_{\sim_{\Omega}}.$$

Therefore

$$\nu \in \left[\bigcup_{I \in \mathbb{S}} \left\{ z_{\nu}^{J} \right\} \times \left\{ I \right\} \right]_{\sim_{\Omega}} \cap V \neq \varnothing.$$

According to (6.21) and Proposition 6.7,

$$\mu = \left[\left(z_{\mu}^{J}, J \right) \right]_{\sim_{\Omega}} \in \check{\Omega}_{+}^{\rho}.$$

It implies that $z_{\nu}^{J} \in \varphi_{\circ}^{\rho}(V)$. By (6.21), $z_{\mu}^{I} = z_{\nu}^{J}$. Then

$$\mu = \left[\left(z_{\mu}^{I}, I \right) \right]_{\sim_{\Omega}} = \left[\left(z_{\nu}^{J}, I \right) \right]_{\sim_{\Omega}} \in \bigcup_{L \in \mathbb{S}, \ z_{\nu}^{J} \in \mathbb{C}_{L}} \left[\left(z_{\nu}^{J}, L \right) \right]_{\sim_{\Omega}}$$

$$= \left[\bigcup_{L \in \mathbb{S}} \left\{z_{\nu}^{J}\right\}_{L} \times \left\{L\right\}\right]_{\sim \alpha} = \psi_{\Omega}^{\mathbb{S}}\left(\left\{z_{\nu}^{J}\right\}\right) \subset \psi_{\Omega}^{\mathbb{S}}\left(\varphi_{\Omega}^{\rho}(V)\right).$$

It implies that $\phi^{\rho}_{\breve{\Omega}}(U) \subset \psi^{\mathbb{S}}_{\Omega} \circ \varphi^{\rho}_{\breve{\Omega}}(V)$. By (6.20), $\psi^{\mathbb{S}}_{\Omega} \circ \varphi^{\rho}_{\breve{\Omega}} = \phi^{\rho}_{\breve{\Omega}}$. It follows from (6.19) that (6.16) holds.

7. Stem discrete sets

In this section, we give a concept of stem discrete sets which applies to 2-path-symmetric open sets. The stem discrete sets play same roles as discrete sets in interpolation theorem in complex analysis. When these sets have the property of invariance in Definition 7.2, we can deduce further consequences that we describe in various results in this section and that will be used in Section 8. In particular, in Proposition 7.9, we show how to construct a slice regular function defined on such a 2-path-symmetric open set using a formula that mimics a representation formula.

Definition 7.1. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric. $A \subset \Omega$ is called $(\check{\Omega}$ -)stem discrete, if for each r > 0, $\psi_{\Omega}^{\mathbb{S}}(A \cap \Omega^r)$ has no accumulation point in $\check{\Omega}$. Moreover, we call A is $(\check{\Omega}$ -)stem-finite if $|\psi_{\Omega}^{\mathbb{S}}(A)| < +\infty$.

Definition 7.2. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric. $K \subset \check{\Omega}$ is called $(\check{\Omega}$ -)invariant, if K is compact and

$$K = K_{\check{\alpha}}^{\wedge} = \operatorname{Conj}_{\check{\alpha}}(K).$$

Moreover, $A \subset \Omega$ is called (Ω -)invariant, if $\psi_{\Omega}^{\mathbb{S}}(A)$ is invariant and $\phi_{\Omega}^{r}(A) = A$ for some r > 0.

Proposition 7.3. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, and $K \subset \Omega$ be invariant. Then

(7.1)
$$\phi_{\Omega}^{\sigma(K,\mathbb{H}\setminus\Omega)}(K) = K.$$

Proof. It follows from $K \subset \Omega^{\sigma(K, \mathbb{H} \setminus \Omega)}$ that

$$\phi_{\Omega}^{\sigma(K,\mathbb{H}\backslash\Omega)}(K) = [K]_{\approx_{\Omega}}^{\cup} \bigcap \Omega^{\sigma(K,\mathbb{H}\backslash\Omega)} \supset K.$$

Since K is invariant,

$$K = \phi_{\Omega}^r(K) = [K]_{\approx_{\Omega}}^{\cup} \bigcap \Omega^r, \quad \text{for some} \quad r > 0.$$

Hence

(7.3)
$$\sigma(K, \mathbb{H}\backslash\Omega) = \sigma\left([K]_{\approx_{\Omega}}^{\cup} \bigcap \Omega^{r}, \mathbb{H}\backslash\Omega\right) \geq \sigma\left(\Omega^{r}, \mathbb{H}\backslash\Omega\right) = r.$$

Therefore, $\Omega^{\sigma(K,\mathbb{H}\setminus\Omega)}\subset\Omega^r$. It implies that

$$(7.4) K = [K]_{\approx_{\Omega}}^{\cup} \bigcap \Omega^{r} \supset [K]_{\approx_{\Omega}}^{\cup} \bigcap \Omega^{\sigma(K, \mathbb{H} \setminus \Omega)} = \phi_{\Omega}^{\sigma(K, \mathbb{H} \setminus \Omega)}(K).$$

Then (7.1) holds by (7.2) and (7.4).

Proposition 7.4. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $K \subset \Omega$ be invariant, $p \in \Omega$ with $|[p]_{\approx_{\Omega}} \cap K| \geq 2$. Then $[p]_{\approx_{\Omega}} \subset K$.

Proof. Let $q \in [p]_{\approx_{\Omega}}$ and $z^I, \xi \in [p]_{\approx_{\Omega}} \cap K$ with $z^I \neq \xi$. By Proposition 3.2 (i),

$$q=z^J, \qquad \text{and} \qquad \xi=z^L, \qquad \text{for some} \qquad J,L\in\mathbb{S} \text{ with } L\neq I.$$

According to (3.3), $z \in \Omega_{2ps}$. Let $\mu := [(z^I, I)]_{\sim_{\Omega}}$. It follows from (5.13) that $\mathbb{S}_{\mu} = \mathbb{S}$. By (5.16) and (6.4),

$$\sigma(z^{\mathcal{I}}, \mathbb{H} \backslash \Omega) = \delta_{\mathcal{I}}(z^{\mathcal{I}}, \mathbb{C}_{\mathcal{I}} \backslash \Omega_{\mathcal{I}}) = r_{u}^{\mathcal{I}}, \qquad \mathcal{I} = I, J, L.$$

It implies by $z^{\mathcal{I}} \in K$, $\mathcal{J} = I, L$, that

$$r_{\mu}^{\mathcal{J}} = \sigma(z^{\mathcal{J}}, \mathbb{H} \backslash \Omega) \ge \sigma(K, \mathbb{H} \backslash \Omega).$$

By (5.19),

$$\sigma(q,\mathbb{H}\backslash\Omega)=\sigma(z^J,\mathbb{H}\backslash\Omega)=r_\mu^J\geq r_\mu^-=\min\{r_\mu^I,r_\mu^L\}\geq\sigma(K,\mathbb{H}\backslash\Omega).$$

Then $q \in \Omega^{\sigma(K, \mathbb{H} \setminus \Omega)}$. By (7.1),

$$q \in [p]_{\approx_{\Omega}} \cap \Omega^{\sigma(K, \mathbb{H} \backslash \Omega)} \subset [K]^{\cup}_{\approx_{\Omega}} \cap \Omega^{\sigma(K, \mathbb{H} \backslash \Omega)} = \phi^{\sigma(K, \mathbb{H} \backslash \Omega)}_{\Omega}(K).$$

It implies that $[p]_{\approx_{\Omega}} \subset K$.

Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $U \in \tau(\check{\Omega}), g: U \to \mathbb{C}$ and $\mu \in \check{\Omega}$. Define

$$B^{\mu}_{\mathbb{C}}:=B_{\mathbb{C}}(z_{\mu},r_{\mu}^{+}), \qquad B^{\mu}_{\S}:=B_{\S}(\mu,r_{\mu}^{+}),$$

and

(7.5)
$$g_{\mu}: \pi_{\breve{\Omega}}(U \cap B_{\breve{\Omega}}^{\mu}) \longrightarrow \mathbb{C},$$
$$z \longmapsto g \circ \left(\pi_{\breve{\Omega}}|_{B_{\breve{\Omega}}^{\mu}}\right)^{-1}(z).$$

If g is holomorphic, then by definition $g_{\mu} = g \circ \left(\pi_{\breve{\Omega}}|_{B_{\breve{\Omega}}^{\mu}}\right)^{-1}$ is also holomorphic. It follows from chain rule, $\frac{\partial}{\partial \overline{z}}g = 0$ and $\frac{\partial}{\partial z}\operatorname{Conj}_{\mathbb{C}} = 0$ that

(7.6)
$$\frac{\partial}{\partial z} \left(g_{\mu} \circ \operatorname{Conj}_{\mathbb{C}}|_{B_{\mathbb{C}}^{\mu}} \right) = \left[\left(\frac{\partial}{\partial z} g_{\mu} \right) \circ \operatorname{Conj}_{\mathbb{C}}|_{B_{\mathbb{C}}^{\mu}} \right] \cdot \left(\frac{\partial}{\partial z} \operatorname{Conj}_{\mathbb{C}}|_{B_{\mathbb{C}}^{\mu}} \right) \\
+ \left[\left(\frac{\partial}{\partial \overline{z}} g_{\mu} \right) \circ \operatorname{Conj}_{\mathbb{C}}|_{B_{\mathbb{C}}^{\mu}} \right] \cdot \left(\frac{\partial}{\partial z} \overline{\operatorname{Conj}_{\mathbb{C}}|_{B_{\mathbb{C}}^{\mu}}} \right) = 0.$$

Let $\nu \in B^{\mu}_{\check{\Omega}} \cap U$. Then for each $\xi \in B^{\mu}_{\check{\Omega}} \cap B^{\nu}_{\check{\Omega}} \cap U$,

$$\left(\pi_{\breve{\Omega}}|_{B^{\mu}_{\breve{\Omega}}}\right)^{-1}(z_{\xi}) = \xi = \left(\pi_{\breve{\Omega}}|_{B^{\nu}_{\breve{\Omega}}}\right)^{-1}(z_{\xi}).$$

It follows that

$$g_{\nu}(z_{\xi}) = g \circ \left(\pi_{\breve{\Omega}}|_{B_{\breve{\Omega}}^{\nu}}\right)^{-1}(z_{\xi}) = g(\xi) = g \circ \left(\pi_{\breve{\Omega}}|_{B_{\breve{\Omega}}^{\mu}}\right)^{-1}(z_{\xi}) = g_{\mu}(z_{\xi}).$$

Hence

$$(7.7) g_{\nu} = g_{\mu}, \text{near} z_{\nu}.$$

Let $I \in \mathbb{S}_{\mu}^+$. By Proposition 6.4, $r_{\mu}^+ = r_{\overline{\mu}}^+$. Then

(7.8)
$$\operatorname{Conj}_{\check{\Omega}}(B_{\check{\Omega}}^{\mu}) = \operatorname{Conj}_{\check{\Omega}}\left(\left[B_{\mathbb{C}}(z_{\mu}, r_{\mu}^{+})^{I} \times \{I\}\right]_{\sim_{\Omega}}\right) \\ = \left[B_{\mathbb{C}}(\overline{z_{\mu}}, r_{\overline{\mu}}^{+})^{-I} \times \{-I\}\right]_{\sim_{\Omega}} = B_{\check{\Omega}}^{\overline{\mu}}.$$

If $\overline{\mu} \in U$, then $\overline{\nu} \in \operatorname{Conj}_{\breve{\Omega}}(B^{\mu}_{\breve{\Omega}}) \cap U = B^{\overline{\mu}}_{\breve{\Omega}} \cap U$ and

(7.9)
$$g_{\overline{\nu}} = g_{\overline{\mu}}, \quad \text{near} \quad z_{\overline{\nu}} = \overline{z_{\nu}}.$$

Remark 7.5. Let Ω be 2-path-symmetric, $I \in \mathbb{S}$, $z^J \in \Omega$ and $\mu := [(z^J, J)]_{\sim_{\Omega}}$ with $J \neq \pm I$. Then

$$\mu, \overline{\mu} \in [\Omega_J \times \{\pm J\}]_{\sim_{\Omega}} \subset \psi_{\Omega}^{\mathbb{S} \setminus \{-I\}}(\Omega).$$

Let $I, J \in \mathbb{S}$. By calculation,

$$\begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix} \begin{pmatrix} 1 & \\ & -1 \end{pmatrix} = \begin{pmatrix} 1 & -I \\ 1 & I \end{pmatrix} = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix} \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}.$$

Then check the matrices below

$$(7.10) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix} = (1, J) \begin{pmatrix} 1 & 1 \\ & -1 \end{pmatrix}^{-1} \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1}$$

$$= (1, J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} 1 & 1 \\ 1 & \end{pmatrix}^{-1}$$

$$= (1, J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} 1 & 1 \\ 1 & \end{pmatrix}.$$

Similarly, it follows from

$$\begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix} \begin{pmatrix} & -1 \\ 1 & \end{pmatrix} = \begin{pmatrix} I & -1 \\ -I & -1 \end{pmatrix} = \begin{pmatrix} I & \\ & -I \end{pmatrix} \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}$$

that

$$\begin{split} J(1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} = & (1,J) \begin{pmatrix} & -1 \\ 1 & \end{pmatrix} \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \\ = & (1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} I \\ & -I \end{pmatrix}. \end{split}$$

It implies that

(7.11)
$$\overline{\partial_J}(1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} = (1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \overline{\partial_I} & \\ & \overline{\partial_{-I}} \end{pmatrix},$$

where

$$\overline{\partial_L} := \frac{\partial}{\partial \overline{z}^L} := \frac{1}{2} \left(\frac{\partial}{\partial x} + L \frac{\partial}{\partial y} \right), \qquad L \in \mathbb{S} \cup \{i\}.$$

It is easy to check that

(7.12)
$$\begin{cases} \frac{\partial}{\partial \overline{z}^I} \Psi_i^I = \Psi_i^I \frac{\partial}{\partial \overline{z}^i} = \Psi_i^I \frac{\partial}{\partial \overline{z}}, \\ \frac{\partial}{\partial \overline{z}^{-I}} \Psi_i^I = \Psi_i^I \frac{\partial}{\partial \overline{z}^{-i}} = \Psi_i^I \frac{\partial}{\partial z}. \end{cases}$$

Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $U \subset \Omega$ and $r \geq 0$. Denote

$$\check{\Omega}^{\scriptscriptstyle{[I]}} := \psi_{\scriptscriptstyle \Omega}^{\scriptscriptstyle{\mathbb{S}\backslash\{-I\}}}(\Omega), \qquad \qquad U_{\scriptscriptstyle \check{\Omega}}^{\scriptscriptstyle{[I]}} := \psi_{\scriptscriptstyle \Omega}^{\scriptscriptstyle{\mathbb{S}\backslash\{-I\}}}(U) = \left[\bigcup_{J \neq -I} U_J \times \{J\}\right]_{\scriptscriptstyle{\mathbb{Q}\backslash\Omega}},$$

and

and

$$(7.13) U_{\breve{\Omega}}^{r,I} := \psi_{\Omega}^{s}(U) \bigcap \left(\breve{\Omega}^{[I]} \right)^{r}.$$

Lemma 7.6. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $K \subset \Omega$ be invariant and $p \in \Omega$ with $[p]_{\approx_{\Omega}} \nsubseteq K$ and $|[p]_{\approx_{\Omega}}| > 1$. Then

(7.14)
$$[p]_{\approx_{\Omega}} \cap K = \{w^{-J}\}, \quad \text{for some} \quad w \in \mathbb{C} \quad \text{and} \quad J \in \mathbb{S}.$$

$$(7.15) [(w^J, J)]_{\sim_{\Omega}} \notin K_{\breve{o}}^{[J]}.$$

Proof. (7.14) holds directly by Proposition 7.4. According to Proposition 3.2 (v), $[p]_{\approx_{\Omega}} = w^{\mathbb{S}}$. Suppose that (7.15) does not hold, i.e.

$$\mu := [(w^J, J)]_{\sim_{\Omega}} \in K_{\breve{\Omega}}^{[J]} = \left[\bigcup_{\hbar \neq -J} K_{\hbar} \times \{\hbar\}\right]_{\sim_{\Omega}}.$$

It implies that there is $v \in \mathbb{C}$ and $L \in \mathbb{S} \setminus \{-J\}$ such that $v^L \in K_L$ and $[(v^L, L)]_{\sim_{\Omega}} = \mu$. By Proposition 5.3, $v = z_{\mu} = w$. Hence

$$w^L = v^L \in K_L \subset K,$$

a contradiction to (7.14). Therefore, (7.15) holds.

Proposition 7.7. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $K \subset \Omega$ be Ω -invariant and $r \geq 0$. Then

(7.16)
$$K_{\breve{\Omega}}^{r,I} = \left(K_{\breve{\Omega}}^{r,I}\right)_{\breve{\Omega}^{[I]}}^{\wedge}, \quad \forall \ I \in \mathbb{S}.$$

Proof. Since K is Ω-invariant, $\psi_{\Omega}^{\mathbb{S}}(K) = [\psi_{\Omega}^{\mathbb{S}}(K)]_{\check{\Omega}}^{\wedge}$. It follows from $\check{\Omega}^{[I]} \subset \check{\Omega}$ and $K_{\check{\alpha}}^{r,I} \subset \psi_{\Omega}^{\mathbb{S}}(K)$ that

$$\left(K_{\breve{\Omega}}^{r,I}\right)_{\breve{\Omega}^{[I]}}^{\wedge} \subset \left(K_{\breve{\Omega}}^{r,I}\right)_{\breve{\Omega}}^{\wedge} \subset \left[\psi_{\Omega}^{\mathbb{S}}(K)\right]_{\breve{\Omega}}^{\wedge} = \psi_{\Omega}^{\mathbb{S}}(K)$$

According to (7.5.6) in [36, Lemma 7.5.5],

$$\delta_{\breve{\Omega}^{[I]}}\left(\left[K_{\breve{\Omega}}^{r,I}\right]_{\breve{\Omega}^{[I]}}^{\wedge},\partial\breve{\Omega}^{[I]}\right)=\delta_{\breve{\Omega}^{[I]}}\left(K_{\breve{\Omega}}^{r,I},\partial\breve{\Omega}^{[I]}\right)\geq\delta_{\breve{\Omega}^{[I]}}\left(\left(\breve{\Omega}^{[I]}\right)^{r},\partial\breve{\Omega}^{[I]}\right)=r,$$

i.e. $\left[K_{\breve{\Omega}}^{r,I}\right]_{\breve{\Omega}^{[I]}}^{\wedge} \subset \left(\breve{\Omega}^{[I]}\right)^{r}$. It implies that

$$\left(K_{\breve{\Omega}}^{r,I}\right)_{\breve{\Omega}^{[I]}}^{\land} \subset \psi_{\Omega}^{\mathbb{S}}(K) \bigcap \left(\breve{\Omega}^{[I]}\right)^{r} = K_{\breve{\Omega}}^{r,I}.$$

By definition, $K_{\breve{\Omega}}^{r,I} \subset \left(K_{\breve{\Omega}}^{r,I}\right)_{\breve{\Omega}^{[I]}}^{\wedge}$. Therefore (7.16) holds.

Proposition 7.8. Let $\Omega \in \tau_s(\Omega)$ be 2-path-symmetric, $I \in \mathbb{S}$ and $U \subset \Omega$. Then

$$U_{\breve{\Omega}}^{[I]} \subset U_{\breve{\Omega}}^{\sigma_{\mathbb{H}\backslash\Omega}^{U},I}.$$

Proof. Let $\mu \in U^{[I]}_{\check{\Omega}} = \bigcup_{L \in \mathbb{S} \setminus \{-I\}} [U_L \times \{L\}]_{\sim_{\Omega}}$. Then $\mu = [(z^J_{\mu}, J)]_{\sim_{\Omega}}$ for some $\mu \in \mathbb{S}_{\mu} \setminus \{-I\}$. It implies that

$$B_J(z^J, \sigma^{\scriptscriptstyle U}_{\scriptscriptstyle \mathbb{H}\setminus\Omega})\subset \Sigma(z^J, \sigma^{\scriptscriptstyle U}_{\scriptscriptstyle \mathbb{H}\setminus\Omega})\subset\Omega.$$

Therefore

$$B_{\breve{\Omega}}(\mu,\sigma^{\scriptscriptstyle U}_{\scriptscriptstyle \mathbb{H}\backslash\Omega}) = [B_J(z^J,\sigma^{\scriptscriptstyle U}_{\scriptscriptstyle \mathbb{H}\backslash\Omega})\times\{J\}]_{\sim_\Omega} \subset \bigcup_{L\in\mathbb{S}\backslash\{-I\}} [U_L\times\{L\}]_{\sim_\Omega} = U^{\scriptscriptstyle [I]}_\Omega.$$

Then $\mu \in (U^{[I]}_{\check{\Omega}})^{\sigma^U_{\mathbb{H} \backslash \Omega}}$ and $U^{[I]}_{\check{\Omega}} \subset (U^{[I]}_{\check{\Omega}})^{\sigma^U_{\mathbb{H} \backslash \Omega}}$. Therefore

$$U_{\check{\Omega}}^{[I]} \subset \psi_{\Omega}^{\mathfrak{s}}(U) \bigcap \left(\check{\Omega}^{[I]} \right)^{\sigma_{\mathbb{H} \backslash \Omega}^{U}} = U_{\check{\Omega}}^{\sigma_{\mathbb{H} \backslash \Omega}^{U}, I}.$$

Proposition 7.9. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $I \in \mathbb{S}$, $g : \check{\Omega}^{[I]} \to \mathbb{C}$ be holomorphic. Define

$$g^I_{\Omega}: \Omega \longrightarrow \mathbb{H},$$

(7.17)
$$z^{J} \longmapsto \begin{cases} \Psi_{i}^{I} \circ g_{\mu}(z) = \Psi_{i}^{I} \circ g(\mu), & J = I, \\ h(z, J), & J \neq \pm I, \end{cases}$$

 $\label{eq:condition} \textit{for each } z^J \in \Omega \textit{ with } z \in \mathbb{C} \textit{ and } J \in \mathbb{S} \backslash \{-I\}, \textit{ where } \mu = [(z^J, J)]_{\sim_{\Omega}} \textit{ and }$

$$h(z,J) = (1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \Psi_i^I \circ g_{\mu}(z) \\ \Psi_i^I \circ g_{\overline{\mu}}(\overline{z}) \end{pmatrix}.$$

Then g_{Ω}^{I} is well-defined and slice regular. Moreover,

(7.18)
$$\varphi_{\breve{\Omega}}^{0}\left(\mathcal{Z}(g)\cap\operatorname{Conj}_{\breve{\Omega}}\circ\mathcal{Z}(g)\right)\subset\mathcal{Z}(g_{\Omega}^{I}).$$

Proof. (i) Let $z^J = w^L$ for some $z, w \in \mathbb{S}$ and $J, L \in \mathbb{S} \setminus \{-I\}$ with $(z, J) \neq (w, L)$. If $z \in \mathbb{R}$, then by (3.4), $z = z^J \in \Omega_{\mathbb{R}} \subset \Omega_{2ps}$. It follows that $w^L = z^J \in \mathbb{R}$ that $w = w^L = z^J = z$. According to (5.2),

$$\mu = [(z^J,J)]_{\sim_\Omega} = [(z^{-J},-J)]_{\sim_\Omega} = [(z^J,-J)]_{\sim_\Omega} = \overline{\mu}.$$

Similarly, $\nu = \overline{\nu} = \mu$, where $\nu = [(w^L, L)]_{\sim_{\Omega}}$. Hence if $L \neq I$, then

$$h(w,L) = (1,L) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \Psi_i^I \circ g_{\nu}(w) \\ \Psi_i^I \circ g_{\overline{\nu}}(\overline{w}) \end{pmatrix}$$
$$= \frac{1}{2} (1,L) \begin{pmatrix} 1 & 1 \\ -I & I \end{pmatrix} \begin{pmatrix} \Psi_i^I \circ g_{\mu}(z) \\ \Psi_i^I \circ g_{\mu}(z) \end{pmatrix} = \Psi_i^I \circ g_{\mu}(z).$$

Similarly, if $J \neq I$, then $h(z,J) = \Psi_i^I \circ g_\mu(z)$. It is easy to check that g_Ω^I is well-defined in this case.

Otherwise, $z \notin \mathbb{R}$ and it is immediate that $w = \overline{z}$ and $L = -J \neq \pm I$. Then

$$\nu = [(w^L, L)]_{\sim_{\Omega}} = [(\overline{z}^{-J}, -J)]_{\sim_{\Omega}} = \overline{\mu}.$$

By (7.10),

$$\begin{split} h(w,L) = & (1,L) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \Psi_i^I \circ g_\nu(w) \\ \Psi_i^I \circ g_{\overline{\nu}}(\overline{w}) \end{pmatrix} \\ = & (1,-J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \Psi_i^I \circ g_{\overline{\mu}}(\overline{z}) \\ \Psi_i^I \circ g_\mu(z) \end{pmatrix} \\ = & (1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \begin{pmatrix} \Psi_i^I \circ g_{\overline{\mu}}(\overline{z}) \\ \Psi_i^I \circ g_\mu(z) \end{pmatrix} \\ = & (1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \Psi_i^I \circ g_\mu(z) \\ \Psi_i^I \circ g_{\overline{\mu}}(\overline{z}) \end{pmatrix} = h(z,J). \end{split}$$

In summary, h(w,L)=h(z,J) for each $w^L=z^J$ with $L,J\in\mathbb{S}\backslash\{-I\}$. Hence g^I_Ω is well-defined.

(ii) Let
$$w^J \in \Omega$$
, $\nu = [(w^J, J)]_{\sim_{\Omega}}$ and $z^J \in B_J(w^J, r_{\nu}^J)$. Then

$$\mu := [(z^J,J)]_{\sim_{\Omega}} \in \left[B_{\mathbb{C}}\left(w,r_{\nu}^J\right)^J \times \{J\}\right]_{\sim_{\Omega}} \subset B_{\check{\Omega}}\left(\nu,r_{\nu}^J\right) = B_{\check{\Omega}}\left(\nu,r_{\nu}^+\right) = B_{\check{\Omega}}^{\nu}.$$

If J = I, then by (7.7) and (7.12) that

$$\frac{\partial}{\partial \overline{z}^J}g_{\Omega}^I(z^J) = \frac{\partial}{\partial \overline{z}^I}\Psi_i^I \circ g_{\mu}(z) = \frac{\partial}{\partial \overline{z}}g_{\nu}(z) = 0.$$

If J = -I, then

$$[(\overline{z}^I,I)]_{\sim_{\Omega}} = [(\overline{z}^{-J},-J)]_{\sim_{\Omega}} = [(z^J,-J)]_{\sim_{\Omega}} = \overline{\mu}.$$

It follows from (7.9), (7.12) and (7.6) that

$$\frac{\partial}{\partial \overline{z}^J}g_{\Omega}^I(z^J) = \frac{\partial}{\partial \overline{z}^{-I}}g_{\Omega}^I(\overline{z}^I) = \frac{\partial}{\partial \overline{z}^{-I}}\Psi_i^I \circ g_{\overline{\mu}}(\overline{z}) = \Psi_i^I \frac{\partial}{\partial z}g_{\overline{\nu}} \circ \mathrm{Conj}_{\mathbb{C}}(z) = 0.$$

Otherwise $J \neq \pm I$. By the same method as above two cases and according to (7.11),

$$\begin{split} \frac{\partial}{\partial \overline{z}^{J}}g_{\Omega}^{I}(z^{J}) &= \frac{\partial}{\partial \overline{z}^{J}}(1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \Psi_{i}^{I} \circ g_{\mu}(z) \\ \Psi_{i}^{I} \circ g_{\overline{\mu}}(\overline{z}) \end{pmatrix} \\ &= (1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \frac{\partial}{\partial \overline{z}^{I}} & \\ & \frac{\partial}{\partial \overline{z}^{-I}} \end{pmatrix} \begin{pmatrix} \Psi_{i}^{I} \circ g_{\nu}(z) \\ \Psi_{i}^{I} \circ g_{\overline{\nu}}(\overline{z}) \end{pmatrix} \\ &= (1,J) \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \Psi_{i}^{I} & \\ & \Psi_{i}^{I} \end{pmatrix} \begin{pmatrix} \frac{\partial}{\partial \overline{z}} g_{\nu}(z) \\ \frac{\partial}{\partial z} g_{\overline{\nu}} \circ \operatorname{Conj}_{\mathbb{C}}(z) \end{pmatrix} = 0. \end{split}$$

Since the choice of $z^J \in B_J\left(w^J, r_\nu^J\right)$ is arbitrary, $g_\Omega^I|_{B_J\left(w^J, r_\nu^J\right)}$ is holomorphic. It follows that for each $J \in \mathbb{S}$, $g_\Omega^I|_{\Omega_J}$ is holomorphic at each point $w^J \in \Omega_J$, and then holomorphic on Ω_J . Therefore, g_Ω^I is slice regular.

(iii) Let $z^J \in \varphi_{\tilde{\Omega}}^0(\mathcal{Z}(g) \cap \operatorname{Conj}_{\tilde{\Omega}} \circ \mathcal{Z}(g))$ with $J \neq -I$, and $\mu := [(z^J, J)]_{\sim_{\Omega}}$. It follows from (6.13) that there is $\lambda \in \{0, 1\}$ such that

$$\operatorname{Conj}_{\tilde{\Omega}}^{\lambda}(\mu) = [(z^J, (-1)^{\lambda}J)]_{\sim_{\Omega}} \in \psi_{\Omega}^{\mathbb{S}}(\{z^J\}) \cap \mathcal{Z}(g) \cap \operatorname{Conj}_{\tilde{\Omega}} \circ \mathcal{Z}(g) \neq \varnothing.$$

Then for each $\kappa \in \{0, 1\}$

$$\operatorname{Conj}_{\breve{\Omega}}^{\kappa}(\mu) = \operatorname{Conj}_{\breve{\Omega}}^{\lambda+\kappa} \left[\operatorname{Conj}_{\breve{\Omega}}^{\lambda}(\mu) \right]$$
$$\in \operatorname{Conj}_{\breve{\Omega}}^{\lambda+\kappa} \left[\mathcal{Z}(g) \cap \operatorname{Conj}_{\breve{\Omega}} \circ \mathcal{Z}(g) \right] = \mathcal{Z}(g) \cap \operatorname{Conj}_{\breve{\Omega}} \circ \mathcal{Z}(g).$$

Therefore $\mu, \overline{\mu} \in \mathcal{Z}(g)$. According to (7.5),

$$g_{\mu}(z) = g \circ \left(\pi|_{B^{\mu}_{\tilde{\Omega}}}\right)^{-1}(z) = g(\mu) = 0.$$

Similarly $g_{\overline{\mu}}(\overline{z}) = 0$. It is easy to check by (7.17) that $g_{\Omega}^{I}(z^{J}) = 0$. Hence $z^{J} \in \mathcal{Z}(g_{\Omega})$. It implies that (7.18) holds.

We recall some facts in slice analysis. Let $u, v \in \mathbb{H}$ and suppose that $u\overline{v} \in \mathbb{C}_I$ for some $I \in \mathbb{C}$. Then for any $J \in \mathbb{S}$,

$$\min_{L=\pm I} |u+Lv| \leq |u+Jv| \leq \max_{L=\pm I} |u+Lv| \,.$$

It implies that for each $w \in (\mathbb{C}_I)^{2 \times 1}$ and $J \in \mathbb{S}$,

(7.19)
$$|(1,J)w| \le \max_{L=\pm I} |(1,L)w|.$$

Proposition 7.10. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $U \subset \Omega$ with $r := \sigma(U, \mathbb{H} \setminus \Omega) > 0$, $I \in \mathbb{S}$ and $g : \check{\Omega}^{[I]} \to \mathbb{C}$. Then

(7.20)
$$\sup_{U} |g_{\Omega}^{I}| \leq \sup_{U_{\varkappa}^{[I]}} |g|.$$

Proof. Let
$$z^J \in U$$
 with $J \in \mathbb{S} \setminus \{-I\}$, and $\mu := \{z^J\}$. If $J = I$ then
$$|g^I_{\Omega}(z^I)| = |\Psi^I_i \circ g_{\mu}(z)| = |g_{\mu}(z)| = |g(\mu)| \leq \sup_{U^{[I]}_{\Omega}} |g|.$$

Otherwise, $J \in \mathbb{S} \setminus \{\pm I\}$. Then

$$w := \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix}^{-1} \begin{pmatrix} \Psi_i^I \circ g_{\mu}(z) \\ \Psi_i^I \circ g_{\overline{\mu}}(\overline{z}) \end{pmatrix} \in (\mathbb{C}_I)^{2 \times 1},$$

and

$$\begin{pmatrix} (1,I)w \\ (1,-I)w \end{pmatrix} = \begin{pmatrix} 1 & I \\ 1 & -I \end{pmatrix} w = \begin{pmatrix} \Psi_i^I \circ g_\mu(z) \\ \Psi_i^I \circ g_{\overline{\mu}}(\overline{z}) \end{pmatrix} = \begin{pmatrix} \Psi_i^I \circ g(\mu) \\ \Psi_i^I \circ g(\overline{\mu}) \end{pmatrix}.$$

By $z^J \in U$, we have

$$\mu = [(z^J, J)]_{\sim_{\Omega}} = [U_J \times \{J\}]_{\sim_{\Omega}} \subset \psi_{\Omega}^{\mathbb{S} \setminus \{I\}}(U) = U_{\alpha}^{[I]}$$

and

$$\overline{\mu} = [(z^J, -J)]_{\sim_{\Omega}} = [U_{-J} \times \{-J\}]_{\sim_{\Omega}} \subset \psi_{\Omega}^{\mathfrak{S} \setminus \{I\}}(U) = U_{\breve{\Omega}}^{[I]}.$$

According to (7.19),

$$\begin{split} |g_{\scriptscriptstyle \Omega}^{\scriptscriptstyle I}(z^I)| = &|(1,J)w| \leq \max_{L=\pm I} |(1,L)w| = \max_{\nu=\mu,\overline{\mu}} |\Psi_i^I \circ g(\nu)| \\ = \max_{\nu=\mu,\overline{\mu}} |g(\nu)| \leq \sup_{U_{\scriptscriptstyle \Omega}^{\scriptscriptstyle [I]}} |g|. \end{split}$$

Therefore

$$|g_{\scriptscriptstyle \Omega}^{\scriptscriptstyle I}(q)| \leq \sup_{U_{\scriptscriptstyle O}^{\scriptscriptstyle [I]}} |g|, \qquad \forall \ q \in U.$$

and (7.20) holds.

Proposition 7.11. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $p \in \Omega$, and $U \subset \Omega$. Then (7.21) $r^+(\psi_{\Omega}^{\mathbb{S}}(\{p\})) \geq \sigma(p, \mathbb{H} \setminus \Omega)$,

and

$$r^+ \left(\psi_{\Omega}^{\mathbb{S}}(U) \right) \ge \sigma(U, \mathbb{H} \backslash \Omega).$$

Proof. Let $w^I = p$ and $\mu = [(w^I, I)]_{\sim_{\Omega}}$. By (6.4),

$$[B_{\mathbb{C}}(z_{\mu}, \sigma^{p}_{\mathbb{H}\setminus\Omega})]^{I} = B_{I}(p, \sigma^{p}_{\mathbb{H}\setminus\Omega}) = B_{I}(p, \delta_{I}(p, \mathbb{C}_{I}\setminus\Omega_{I})) \subset \Omega.$$

It follows from Proposition 6.4 that $\sigma^p_{\mathbb{H}\setminus\Omega}\leq r^I_\mu\leq r^I_\mu=r^+_{\overline{\mu}}$. Then (7.21) holds by (6.13). It is clear that

$$r^{+} \left(\psi_{\Omega}^{\mathbb{S}}(U) \right) = r^{+} \left(\bigcup_{p \in U} \psi_{\Omega}^{\mathbb{S}}(\{p\}) \right) = \inf_{p \in U} r^{+} \left(\psi_{\Omega}^{\mathbb{S}}(\{p\}) \right)$$
$$\geq \inf_{p \in U} \sigma(p, \mathbb{H} \backslash \Omega) = \sigma(U, \mathbb{H} \backslash \Omega).$$

Proposition 7.12. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, and $\mu \in \check{\Omega}$. Then $\mu \notin \psi_{\Omega}^{\mathbb{S} \setminus \mathbb{S}_{\mu}}(\Omega)$.

Proof. If $\mathbb{S} = \mathbb{S}_{\mu}$, then $\mu \notin \emptyset = \psi_{\Omega}^{\mathbb{S} \setminus \mathbb{S}_{\mu}}(\Omega)$ and Proposition 7.12 holds. Otherwise, $\mathbb{S} \neq \mathbb{S}_{\mu}$. Suppose that $\mu \in [\Omega_I \times \{I\}]_{\sim_{\Omega}}$ for some $I \in \mathbb{S} \setminus \mathbb{S}_{\mu}$. Let $J \in \mathbb{S}_{\mu}$. According to (5.13), $z_{\mu} \notin \Omega_{2ps}$. Hence there is $w^I \in \Omega_I$ such that

$$[(z_{\mu}^{J}, J)]_{\sim_{\Omega}} = \mu = [(w^{I}, I)]_{\sim_{\Omega}}, \quad \text{i.e.} \quad (z_{\mu}^{J}, J) \sim_{\Omega} (w^{I}, I).$$

It follows from (5.2) that J = I, a contradiction. Thence $\mu \notin [\Omega_I \times \{I\}]_{\sim_{\Omega}}$ and

$$\mu \notin \bigcup_{L \in \mathbb{S} \setminus \mathbb{S}_{\mu}} [\Omega_L \times \{L\}]_{\sim_{\Omega}} = \psi_{\Omega}^{\mathbb{S} \setminus \mathbb{S}_{\mu}}(\Omega).$$

Proposition 7.13. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $K \subset \Omega$ be invariant, $z^I \in \Omega \backslash K$, and $\mu = [(z^I, I)]_{\sim_{\Omega}}$. Then

$$\begin{cases} \overline{\mu} \in \widecheck{\Omega}^{[J]} \backslash \left(\widecheck{\Omega}^{[J]}\right)^{\sigma_{\mathbb{H} \backslash \Omega}^{K}}, & [z^{I}]_{\approx_{\Omega}} \cap K = \{z^{J}\} \text{ for some } J \in \mathbb{S}, \\ \mu, \overline{\mu} \notin \psi_{\Omega}^{\mathbb{S}}(K), & [z^{I}]_{\approx_{\Omega}} \cap K = \varnothing, & \text{otherwise.} \end{cases}$$

Proof. (i) Suppose $[z^I]_{\approx_0} \cap K = \{z^J\}$ for some $J \in \mathbb{S}$. Then

$$r_{\mu}^{+} \geq r_{\mu}^{J} = \sigma(z^{J}, \mathbb{H} \backslash \Omega) \geq \sigma(K, \mathbb{H} \backslash \Omega).$$

If
$$\sigma(z^I, \mathbb{H}\backslash\Omega) \ge \sigma(K, \mathbb{H}\backslash\Omega)$$
, then $z^I \in \Omega^{\sigma(z^I, \mathbb{H}\backslash\Omega)} \subset \Omega^{\sigma(K, \mathbb{H}\backslash\Omega)}$. By (6.14), $\psi_{\Omega}^{\mathbb{S}}(z^I) \cap \psi_{\Omega}^{\mathbb{S}}(K) = \psi_{\Omega}^{\mathbb{S}}(z^J) \cap \psi_{\Omega}^{\mathbb{S}}(K) = \psi_{\Omega}^{\mathbb{S}}(z^J) \ne \emptyset$,

(6.16) and (7.1) that

$$z^I \in \varphi^{\sigma(K, \mathbb{H} \backslash \Omega)}_{\breve{\Omega}} \left(\psi^{\mathbb{S}}_{\Omega}(K) \right) = \phi^{\sigma(K, \mathbb{H} \backslash \Omega)}_{\Omega}(K) = K,$$

a contradiction. Therefore $\sigma(z^I, \mathbb{H}\backslash\Omega) < \sigma(K, \mathbb{H}\backslash\Omega)$. Let

$$w^I \in \left[B_{\mathbb{C}}(z, \sigma^K_{\mathbb{H} \setminus \Omega})\right]^I \setminus \Omega.$$

Then

$$\nu := \left[(w^J,J) \right]_{\sim_{\Omega}} \in \left[\left[B_{\mathbb{C}}(z,\sigma^K_{\mathrm{H}\backslash\Omega}) \right]^J \times \{J\} \right]_{\sim_{\Omega}} = B_{\breve{\Omega}}(\mu,\sigma^K_{\mathrm{H}\backslash\Omega}).$$

It implies by $I \notin \mathbb{S}_{\nu}$, (5.13) and (6.9) that $\mathbb{S}_{\overline{\nu}} = -\mathbb{S}_{\nu} = -\{J\} = \{-J\}$. According to Proposition 7.12, $\overline{\nu} \notin \psi_{\Omega}^{\mathbb{S} \setminus \{-J\}}(\Omega)$. It follows from

$$\overline{\nu} = \operatorname{Conj}_{\breve{\Omega}}(\nu) \in \operatorname{Conj}_{\breve{\Omega}}\left[B_{\breve{\Omega}}(\mu, \sigma_{{\mathbb{H}} \backslash \Omega}^K)\right] = B_{\breve{\Omega}}(\overline{\mu}, \sigma_{{\mathbb{H}} \backslash \Omega}^K)$$

that

$$B_{\breve{\Omega}}(\overline{\mu}, \sigma_{\mathbb{H}\backslash\Omega}^K) \not\subset \psi_{\Omega}^{\mathbb{S}\backslash\{-J\}}(\Omega) = \breve{\Omega}^{[J]}.$$

Thence $\overline{\mu} \notin (\check{\Omega}^{[J]})^{\sigma_{\mathbb{H}\backslash\Omega}^K}$. According (3.3) and $z^I \approx_{\Omega} z^J$, we have $z \in \Omega_{2ps}$ and $\overline{z}^J = z^{-J} \in \Omega$. According to (5.8),

$$\overline{\mu} = [(\overline{z}^{-J}, -J)]_{\sim_{\Omega}} = [(\overline{z}^{J}, J)]_{\sim_{\Omega}} \in [\Omega_{J} \times \{J\}]_{\sim_{\Omega}} \subset \psi_{\Omega}^{\mathbb{S} \backslash \{-J\}}(\Omega) = \widecheck{\Omega}^{[J]}.$$

In summary, $\overline{\mu} \in \check{\Omega}^{[J]} \setminus (\check{\Omega}^{[J]})^{\sigma_{\mathbb{H} \setminus \Omega}^K}$.

(ii) Otherwise,
$$\left| [z^I]_{\approx_{\Omega}} \cap K \right| \neq 1$$
. If $\left| [z^I]_{\approx_{\Omega}} \cap K \right| \geq 2$, then by Proposition 7.4, $z^I \in [z^I]_{\approx_{\Omega}} \subset K$,

a contradiction. Thence $[z^I]_{\approx_{\Omega}} \cap K = \varnothing$. If $\{\mu, \overline{\mu}\} \cap \psi_{\Omega}^{\mathbb{S}}(K) \neq \varnothing$. Then there is $w^J \in K$ and $\lambda, \kappa \in \{0,1\}$ such that $\operatorname{Conj}_{\Omega}^{\lambda}(\mu) = [(w^J, (-1)^{\kappa}J)]_{\sim_{\Omega}}$. Then

$$[(z^I, I)]_{\sim_{\Omega}} = \mu = \operatorname{Conj}_{\check{\Omega}}^{\lambda}[(w^J, (-1)^{\kappa}J)]_{\sim_{\Omega}} = [(w^J, (-1)^{\kappa+\lambda}J)]_{\sim_{\Omega}} = [(w'^{J'}, J')]_{\sim_{\Omega}},$$

where $w' = \operatorname{Conj}_{\mathbb{C}}^{\kappa+\lambda}(w)$ and $J' = (-1)^{\kappa+\lambda}J$. By (5.4), $z^I \approx_{\Omega} {w'}^{J'}$. It implies that

$$w^J = {w'}^{J'} \in [z^I]_{\approx_{\Omega}} \cap K,$$

a contradiction. Thence $\{\mu, \overline{\mu}\} \cap \psi_0^{\mathbb{S}}(K) = \emptyset$, i.e. $\mu, \overline{\mu} \notin \psi_0^{\mathbb{S}}(K)$.

8. Interpolation Theorem

In this section, we will prove an interpolation theorem for slice regular functions, which is crucial for proving the Cartan-Thullen Theorem 2.10 but also of independent interest. We begin by defining when an open set Ω is 2-path-symmetric and when a function is 2-path-slice, so that we can prove how to obtain a unique extension of 2-path-slice function to such a set Ω , see Proposition 8.3. Another extension result, this time to obtain a slice regular function, is proved in Proposition 8.6; this result will be needed to prove the Interpolation Theorem 8.8 for slice regular function defined on 2-path-symmetric slice-open sets.

Definition 8.1. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $U \subset \Omega$ and let

$$U_{\Omega,2} := \{ p \in U : |[p]_{\approx_{\Omega}} \cap U| \ge 2 \}.$$

We call

(8.1)
$$U_{\Omega,\bar{2}} := \left(\bigcup_{p \in U_{\Omega,2}} [p]_{\approx_{\Omega}}\right) \bigcup U$$

the Ω -2-path-symmetric completion of U. We say that U is Ω -2-path-symmetric, if $U = U_{\Omega,\tilde{\omega}}$.

Definition 8.2. Let $U \subset \Omega \subset \mathbb{H}$. $f: U \to \mathbb{H}$ is called Ω -path-slice, if for each $\gamma \in \mathscr{P}_0(\mathbb{C})$, $I, J, K \in \mathbb{S}$ with $\gamma^I, \gamma^J, \gamma^K \subset \Omega$, $J \neq K$ and $\gamma^I(1), \gamma^J(1), \gamma^K(1) \in U$, we have

$$(8.2) f \circ \gamma^I(1) = (1, I) \begin{pmatrix} 1 & J \\ 1 & K \end{pmatrix}^{-1} \begin{pmatrix} f \circ \gamma^J(1) \\ f \circ \gamma^K(1) \end{pmatrix}.$$

Proposition 8.3. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $U \subset \Omega$, and $f: U \to \mathbb{H}$ be Ω -path-slice. Then there is a unique Ω -path-slice function $g:=f_{\Omega,\tilde{2}}:U_{\Omega,\tilde{2}}\to \mathbb{H}$ such that $g|_U=f$.

Proof. (i) Denote

$$\mathcal{U}:=\{[p]_{\approx_\Omega}\cap U\in P(\Omega): p\in U\}.$$

By Proposition 3.2 (i), for each $\mu \in \mathcal{U} \subset P(\Omega)$, there is $z_{\mu} \in \mathbb{C}$ such that

$$\mu = \left\{ \left(z_{\mu} \right)^{I} \in U : I \in \mathbb{S} \right\}.$$

If $|\mu| \geq 2$, then $z_{\mu} \notin \mathbb{R}$ and we choose $I_{\mu}, J_{\mu} \in \mathbb{S}$ with $z_{\mu}^{I_{\mu}}, z_{\mu}^{J_{\mu}} \in \mu$ and $I_{\mu} \neq J_{\mu}$. For each $q \in [z_{\mu}^{I_{\mu}}]_{\approx_{\Omega}}$ with $|\mu| \geq 2$, denote

$$z_q:=z_\mu, \qquad I_q:=I_\mu, \qquad J_q:=J_\mu$$

and L_q be the unique imaginary unit such that $q=z_q^{I_q}$. Define

$$f_{\Omega,\tilde{2}}: U_{\Omega,\tilde{2}} \longrightarrow \mathbb{H},$$

$$q \longmapsto \begin{cases} f(q), & q \in U, \\ h(q), & \text{otherwise,} \end{cases}$$

where

(8.3)
$$h(q) := (1, L_q) \begin{pmatrix} 1 & I_q \\ 1 & J_q \end{pmatrix}^{-1} \begin{pmatrix} f(z_q^{I_q}) \\ f(z_q^{J_q}) \end{pmatrix}.$$

Now we show that

$$(8.4) \quad f_{\Omega,\tilde{z}}(q) = (1, L_q) \begin{pmatrix} 1 & I_q \\ 1 & J_q \end{pmatrix}^{-1} \begin{pmatrix} f(z_q^{I_q}) \\ f(z_q^{J_q}) \end{pmatrix}, \quad \forall \ q \in U_{\Omega,\tilde{z}} \text{ with } |[q]_{\approx_{\Omega}} \cap U| \ge 2.$$

Let $q \in U_{\Omega,\tilde{2}}$ with $|[q]_{\approx_{\Omega}} \cap U| \geq 2$. If $q \notin U$, then (8.4) holds by (8.3). Otherwise, $q \in U$. It follows from Proposition 3.2 (iv) and $z_q^{I_q}, z_q^{J_q} \in [q]_{\approx_{\Omega}}$ that there is $\beta \in \mathscr{P}_0(\mathbb{C})$ such that

(8.5)
$$\beta^{L_q}, \beta^{I_q}, \beta^{J_q} \subset \Omega, \quad \text{and} \quad \beta(1) = z_q.$$

Since f is Ω -path-slice,

$$f_{\Omega,\tilde{2}}(q) = f(q) = f(\beta^{L_q}(1)) = (1, L_q) \begin{pmatrix} 1 & I_q \\ 1 & J_q \end{pmatrix}^{-1} \begin{pmatrix} f(\beta^{I_q}(1)) \\ f(\beta^{J_q}(1)) \end{pmatrix}$$
$$= (1, L_q) \begin{pmatrix} 1 & I_q \\ 1 & J_q \end{pmatrix}^{-1} \begin{pmatrix} f(z_q^{I_q}) \\ f(z_q^{J_q}) \end{pmatrix}.$$

Thence (8.4) holds.

Let $\gamma \in \mathscr{P}_0(\mathbb{C})$, $I, J, K \in \mathbb{S}$ with $\gamma^I, \gamma^J, \gamma^K \subset \Omega$, $J \neq K$ and $\gamma^I(1), \gamma^J(1), \gamma^K(1) \in U_{\Omega, \tilde{2}}$. If $w := \gamma(1) \notin \mathbb{R}$, then $|\mu| \geq |\{w^J, w^K\}| \geq 2$, where $\mu := [w^I]_{\approx_{\Omega}} \cap U$. Then there is $\lambda \in \{0, 1\}$ such that

$$z_{\mu} = \operatorname{Conj}_{\mathbb{C}}^{\lambda}(w), \quad \text{and} \quad z_{\mu}^{(-1)^{\lambda}\Gamma} = w^{\Gamma}, \quad \Gamma = I, J, K.$$

It follows from (8.4) that

$$(1,I) \begin{pmatrix} 1 & J \\ 1 & K \end{pmatrix}^{-1} \begin{pmatrix} f_{\Omega,\bar{z}}(\gamma^{J}(1)) \\ f_{\Omega,\bar{z}}(\gamma^{K}(1)) \end{pmatrix} = (1,I) \begin{pmatrix} 1 & J \\ 1 & K \end{pmatrix}^{-1} \begin{pmatrix} f_{\Omega,\bar{z}}(w^{J}) \\ f_{\Omega,\bar{z}}(w^{K}) \end{pmatrix}$$

$$= (1,I) \begin{pmatrix} 1 & J \\ 1 & K \end{pmatrix}^{-1} \begin{pmatrix} 1 & J \\ 1 & K \end{pmatrix} \begin{pmatrix} 1 & (-1)^{\lambda} \end{pmatrix} \begin{pmatrix} 1 & I_{\mu} \\ 1 & J_{\mu} \end{pmatrix}^{-1} \begin{pmatrix} f(z_{\mu}^{I_{\mu}}) \\ f(z_{\mu}^{J_{\mu}}) \end{pmatrix}$$

$$= (1,I) \begin{pmatrix} 1 & (-1)^{\lambda} \end{pmatrix} \begin{pmatrix} 1 & I_{\mu} \\ 1 & J_{\mu} \end{pmatrix}^{-1} \begin{pmatrix} f(z_{\mu}^{I_{\mu}}) \\ f(z_{\mu}^{J_{\mu}}) \end{pmatrix}$$

$$= (1,(-1)^{\lambda}I) \begin{pmatrix} 1 & I_{\mu} \\ 1 & J_{\mu} \end{pmatrix}^{-1} \begin{pmatrix} f(z_{\mu}^{I_{\mu}}) \\ f(z_{\mu}^{J_{\mu}}) \end{pmatrix} = f_{\Omega,\bar{z}}(w^{I}) = f_{\Omega,\bar{z}}(\gamma^{I}(1)).$$

Otherwise, $w \in \mathbb{R}$. Then $\gamma^{\Gamma}(1) = w$, $\Gamma = I, J, K$. It implies that

$$(8.7) \qquad (1,I) \begin{pmatrix} 1 & J \\ 1 & K \end{pmatrix}^{-1} \begin{pmatrix} f_{\Omega,\tilde{z}}(\gamma^{J}(1)) \\ f_{\Omega,\tilde{z}}(\gamma^{K}(1)) \end{pmatrix} = (1,I) \begin{pmatrix} 1 & J \\ 1 & K \end{pmatrix}^{-1} \begin{pmatrix} 1 & J \\ 1 & K \end{pmatrix} \begin{pmatrix} f_{\Omega,\tilde{z}}(w) \\ 0 \end{pmatrix} = (1,I) \begin{pmatrix} f_{\Omega,\tilde{z}}(w) \\ 0 \end{pmatrix} = f_{\Omega,\tilde{z}}(\gamma^{I}(1)).$$

According (8.6) and (8.7), $f_{\Omega,\tilde{2}}$ is Ω -path-slice.

(ii) Suppose that $h: U_{\Omega,\tilde{z}} \to \mathbb{H}$ is Ω -path-slice with $h|_U = f$. Let $q \in U_{\Omega,\tilde{z}} \setminus U$. It is easy to check that $|[q]_{\approx_{\Omega}} \cap U| \geq 2$. According to Proposition 3.2 (iv), there is $\beta \in \mathscr{P}_0(\mathbb{C})$ such that (8.5) holds. By (8.4),

$$\begin{split} h(q) = & h(\beta^{L_q}(1)) = (1, L_q) \begin{pmatrix} 1 & I_q \\ 1 & J_q \end{pmatrix}^{-1} \begin{pmatrix} h(\beta^{I_q}(1)) \\ h(\beta^{J_q}(1)) \end{pmatrix} \\ = & (1, L_q) \begin{pmatrix} 1 & I_q \\ 1 & J_q \end{pmatrix}^{-1} \begin{pmatrix} f(z_q^{I_q}) \\ f(z_q^{J_q}) \end{pmatrix} = f_{\Omega, \tilde{2}}(q). \end{split}$$

It implies from $h|_U = f = f_{\Omega,\tilde{2}}|_U$ that $h = f_{\Omega,\tilde{2}}$.

Recalling the notion of analytic set, see e.g. [36, Definition 2.3.1], we prove the next result:

Lemma 8.4. Let (Ω, π) be a Riemann domain over \mathbb{C} , $K \subset \Omega$ be compact with $K_{\Omega}^{\wedge} = K$, A be an analytic set with $A \neq \Omega$, $z \in A \setminus K$ and $\varepsilon > 0$. Then there is $f \in \mathcal{O}(\Omega)$ such that

(8.8)
$$f(z) = 1, \quad f|_{A \setminus \{z\}} \equiv 0, \quad and \quad \sup_{K} |f| < \varepsilon.$$

Proof. Since A is an analytic set with $A \neq \Omega$, z is an isolated point of A. Therefore the function

$$g: A \longrightarrow \mathbb{C},$$

$$\xi \longmapsto \begin{cases} 1, & \xi = z, \\ 0, & \text{otherwise,} \end{cases}$$

is holomorphic. Since every non-compact Riemann surface is a Stein manifold, the Riemann domain Ω over $\mathbb C$ is a Stein manifold. By interpolation theorem (e.g. [36, Theorem 4.5.11]), there is $\widetilde{g} \in \mathscr{O}(\Omega)$ such that $\widetilde{g}|_A = g$. Since $z \notin K = K_{\Omega}^{\wedge}$, there is $h \in \mathscr{O}(\Omega)$ such that $|h(z)| > \sup_K |h|$. Then there is $h \in \mathbb{N}$ such that

$$\sup_{K} \left| \left(\frac{h}{h(z)} \right)^{\lambda} \right| < \frac{\varepsilon}{\max_{K} |\widetilde{g}| + 1}.$$

It is easy to check that $f := \widetilde{g} \cdot \left(\frac{h}{h(z)}\right)^{\lambda} \in \mathscr{O}(\Omega)$ satisfies (8.8).

Lemma 8.5. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $K \subset \Omega$ be invariant, $\varepsilon > 0$, $A \subset \Omega$ be stem-finite, and $g: A \to \mathbb{H}$ be an Ω -path-slice function with $A \setminus [p]_{\approx_{\Omega}}, A_{\Omega,\tilde{2}} \cap K \subset \mathcal{Z}(g_{\Omega,\tilde{2}})$ for some $p \in A$. Then there is $f \in \mathcal{SR}(\Omega)$ such that

(8.9)
$$f|_{A} = g, \quad and \quad \sup_{K} |f| < \varepsilon.$$

Proof. If $A \cap [p]_{\approx_{\Omega}} \subset K$, then $g \equiv 0$ and (8.9) holds for $f \equiv 0$. Otherwise, let

$$(8.10) z^I \in A \cap [p]_{\approx_{\Omega}} \backslash K, \text{for some} z \in \mathbb{C} \text{and} I \in \mathbb{S}.$$

Denote $\mu := [(z^I, I)]_{\sim_{\Omega}}$. By (7.3), $r := \sigma(K, \mathbb{H} \setminus \Omega) > 0$.

(i) If $|[p]_{\approx_{\Omega}}| = 1$, then $[p]_{\approx_{\Omega}} = \{z^I\}$. Since A is stem-finite,

$$|A_{\bullet}^{[I]}| = |\psi_{\bullet}^{\mathbb{S}\setminus\{-I\}}(A)| \le |\psi_{\bullet}^{\mathbb{S}}(A)| < \infty.$$

Thence $A_{\check{\Omega}}^{[I]}$ is an analytic set in the Riemann domain $\check{\Omega}^{[I]}$. Proposition 7.13 implies that $\mu \notin K_{\check{\Omega}}^{r,I} \subset \psi_{\Omega}^{\mathbb{S}}(K)$. According to Proposition 7.7, $K_{\check{\Omega}}^{r,I} = \left(K_{\check{\Omega}}^{r,I}\right)_{\check{\Omega}^{[I]}}^{\wedge}$. According to Lemma 8.4, there is $h_1 \in \mathscr{O}(\check{\Omega}^{[I]})$ such that

$$(8.11) h_1(\mu) = 1, h_1|_{A_{\Omega}^{[I]} \setminus \{\mu\}} \equiv 0, \text{and} \sup_{K_{\widetilde{O}}^{r,I}} |h_1| < \frac{\varepsilon}{|g(z^I)| + 1}.$$

Let

$$(8.12) f := (h_1)^I_{\Omega} \cdot g(z^I).$$

Then

$$f(z^{I}) = (h_{1})_{\Omega}^{I}(z^{I}) \cdot g(z^{I}) = \Psi_{i}^{I}((h_{1})_{\mu}(z)) \cdot g(z^{I})$$

= $\Psi_{i}^{I}(h_{1}(\mu)) \cdot g(z^{I}) = \Psi_{i}^{I}(1) \cdot g(z^{I}) = g(z^{I}).$

If $A = \{z^I\}$, then $f|_A = g$. Otherwise, there is $w^L \in A \setminus [p]_{\approx_{\Omega}} = A \setminus \{z^I\} \subset \mathcal{Z}(h_1)$ with $L \in \mathbb{S} \setminus \{-I\}$, and $\nu := [(w^L, L)]_{\sim_{\Omega}}$. According to (5.2), $\nu \notin \{\mu, \overline{\mu}\}$. It follows from

$$\nu = [(w^L, L)]_{\sim_{\Omega}} \in [A \times \{L\}]_{\sim_{\Omega}} \subset \psi_{\Omega}^{\text{sh}-1}(A) = A_{\tilde{\Omega}}^{[I]}$$

that $\nu \in A_{\breve{o}}^{[I]} \setminus \{\mu\}$. If J = I, then

$$f(w^{J}) = (h_{1})_{\Omega}^{I}(w^{J}) \cdot g(z^{I}) = \Psi_{i}^{I}((h_{1})_{\nu}(w)) \cdot g(z^{I})$$

= $\Psi_{i}^{I}(h_{1}(\nu)) \cdot g(z^{I}) = \Psi_{i}^{I}(0) \cdot g(z^{I}) = 0 = g_{\Omega,\tilde{2}}(w^{J}) = g(w^{J}).$

Otherwise, $J \neq I$. Then

$$\overline{\nu} = [(w^L, -L)]_{\sim_{\Omega}} \subset [U_{-L} \times \{-L\}]_{\sim_{\Omega}} \subset \psi_{\Omega}^{\mathbb{S} \setminus \{-I\}}(A) = A_{\mathbb{O}}^{[I]},$$

implies that $\overline{\nu}\in A_{\breve{\Omega}}^{{\scriptscriptstyle [I]}}\backslash\{\mu\}$ and

$$\nu \in \operatorname{Conj}_{\breve{\Omega}}(A^{[I]}_{\breve{\Omega}} \setminus \{\mu\}) \subset \operatorname{Conj}_{\breve{\Omega}}(\mathcal{Z}(h_1)).$$

By (7.18),

$$w^J \in \varphi_{\breve{\Delta}}^0\left(\{\nu\}\right) \subset \varphi_{\breve{\Delta}}^0\left(\mathcal{Z}(h_1) \cap \operatorname{Conj}_{\breve{\Delta}} \circ \mathcal{Z}(h_1)\right) \subset \mathcal{Z}((h_1)_{\Omega}^I).$$

It is clear that

$$f(w^J) = (h_1)_{\Omega}^I(w^J) \cdot g(z^I) = 0 \cdot g(z^I) = 0 = g(w^J).$$

In summary,

$$(8.13) f|_{A\setminus [p]_{\approx_{\Omega}}} \equiv 0 \equiv g|_{A\setminus [p]_{\approx_{\Omega}}} and f|_{A} = g.$$

By Proposition 7.8, $K^{[I]}_{\check{\Omega}}\subset K^{r,I}_{\check{\Omega}}.$ It follows from Proposition 7.10 and (8.11) that

$$\sup_{K} |(h_1)_{\Omega}^I| \leq \sup_{K_{\tilde{\Omega}}^{[I]}} |h_1| \leq \sup_{K_{\tilde{\Omega}}^{r,I}} |h_1| < \frac{\varepsilon}{|g(z^I)|+1}.$$

According to (8.12), $\sup_K |f| < \varepsilon$. Hence (8.9) holds.

(ii) If $|[p]_{\approx_{\Omega}}| > 1$ and $[p]_{\approx_{\Omega}} \cap K \neq \emptyset$, then by Lemma 7.6,

$$[p]_{\approx_{\Omega}} \cap K = \{z^{-J}\}, \quad \text{for some} \quad J \in \mathbb{S},$$

and

$$\mu = [(z^I, I)]_{\sim_{\Omega}} = [(z^J, J)]_{\sim_{\Omega}} \notin K_{\Diamond}^{[J]}.$$

Since $|[p]_{\approx_{\Omega}}| > 1$, $z \notin \mathbb{R}$ and $\mu \neq \overline{\mu}$. According to (7.16) and Lemma 8.4, there is $h_2 \in \mathscr{O}(\check{\Omega}^{[J]})$ such that

$$(8.14) \quad h_2(\mu) = 1, \qquad h_2|_{A^{[J]}_{\tilde{\Omega}} \cup \{\overline{\mu}\} \setminus \{\mu\}} \equiv 0, \qquad \text{and} \qquad \sup_{K^{r,J}_{\tilde{C}}} |h_2| < \frac{\varepsilon}{|g(z^I)| + 1}.$$

Define

(8.15)
$$f := (h_2)_{\Omega}^J \cdot (1, J) \begin{pmatrix} 1 & I \\ 1 & -J \end{pmatrix}^{-1} \begin{pmatrix} g(z^I) \\ 0 \end{pmatrix}.$$

By (7.17) and (8.14),

$$\begin{cases} (h_2)_{\Omega}^J(z^J) = h_2(\mu) = 1, \\ (h_2)_{\Omega}^J(z^{-J}) = (h_2)_{\Omega}^J(\overline{z}^J) = h_2([(\overline{z}^J, J)]_{\sim_{\Omega}}) = h_2(\overline{\mu}) = 0. \end{cases}$$

It implies by (8.15) that

$$f(z^L) = (1, L) \begin{pmatrix} 1 & I \\ 1 & -J \end{pmatrix}^{-1} \begin{pmatrix} g(z^I) \\ 0 \end{pmatrix}, \qquad L = \pm J,$$

and then

$$\begin{split} f(z^I) = & (1,I) \begin{pmatrix} 1 & J \\ 1 & -J \end{pmatrix}^{-1} \begin{pmatrix} f(z^J) \\ f(z^{-J}) \end{pmatrix} \\ = & (1,I) \begin{pmatrix} 1 & J \\ 1 & -J \end{pmatrix}^{-1} \begin{pmatrix} 1 & J \\ 1 & -J \end{pmatrix} \begin{pmatrix} 1 & I \\ 1 & -J \end{pmatrix}^{-1} \begin{pmatrix} g(z^I) \\ 0 \end{pmatrix} = g(z^I). \end{split}$$

If $|[p]_{\approx_{\Omega}} \cap A| = 1$, then by (8.10), $[p]_{\approx_{\Omega}} \cap A = \{z^I\}$. It implies that

$$(8.16) f|_{[p]_{\approx_{\Omega}} \cap A} = g|_{[p]_{\approx_{\Omega}} \cap A}.$$

Otherwise, $|[p]_{\approx_{\Omega}} \cap A| > 1$. Then $p \in A_{\Omega,2}$ and $z^{-J} \in A_{\Omega,\bar{2}} \cap K \subset \mathcal{Z}(g_{\Omega,\bar{2}})$. It implies that

$$\begin{cases} f(z^I) = g(z^I) = g_{\Omega,\tilde{z}}(z^I), \\ f(z^J) = 0 = g(z^J). \end{cases}$$

Since f and $g_{\Omega,\tilde{2}}$ are Ω -path-slice, $f = g_{\Omega,\tilde{2}}$ on $[p]_{\approx_{\Omega}}$. Hence (8.16) holds. By similar method in (i), (8.13) holds and $\sup_K |f| < \varepsilon$. Hence (8.9) holds.

(iii) If $|[p]_{\approx_{\Omega}}| > 1$, $[p] \cap K = \emptyset$ and $|[p]_{\approx_{\Omega}} \cap A| = 1$, then by (8.10), $[p]_{\approx_{\Omega}} \cap A = \{z^I\}$. By similar method in (i), there is $h_1 \in \mathscr{O}(\check{\Omega}^{[I]})$ satisfying (8.11), and $f := (h_1)_{\Omega}^I \cdot g(z^I)$ satisfies (8.9).

(iv) If
$$|[p]_{\approx_{\Omega}}| > 1$$
, $[p] \cap K = \emptyset$ and $|[p]_{\approx_{\Omega}} \cap A| > 1$, then by Proposition 7.13 $\mu, \overline{\mu} \notin K_{\times}^{r,I} \cup K_{\times}^{r,-I} \subset \psi_{\Omega}^{\mathbb{S}}(K)$.

By similar method in (i), there is $h_3 \in \mathcal{O}(\check{\Omega}^{[J]})$ and $h_4 \in \mathcal{O}(\check{\Omega}^{[-J]})$ such that

$$\begin{split} h_3(\mu) &= h_4(\overline{\mu}) = 1, \qquad h_3|_{A^{[I]}_\Omega \setminus \{\mu\}} \equiv 0, \qquad h_4|_{A^{[-I]}_\Omega \setminus \{\overline{\mu}\}} \equiv 0 \\ \sup_{K^{r,I}_\Omega} |h_3| &< \frac{\varepsilon}{2|g(z^I)|+1}, \qquad \text{and} \qquad \sup_{K^{r,-I}_{\widetilde{\Omega}}} |h_4| &< \frac{\varepsilon}{2|g_{\Omega,\widetilde{2}}(z^{-I})|+1}. \end{split}$$

Define $f_3 := 2 \cdot (h_3)^I_{\Omega} \cdot g(z^I)$, $f_4 := 2 \cdot (h_4)^{-I}_{\Omega} \cdot g_{\Omega,\tilde{z}}(z^{-I})$ and $f := \frac{f_3 + f_4}{2}$. By similar method in (i),

$$(8.17) f'|_{A \setminus [p]_{\approx_{\Omega}}} = g|_{A \setminus [p]_{\approx_{\Omega}}} \equiv 0, \text{and} \sup_{K} |f'| < \epsilon, f' = f_3, f_4, f.$$

By similar method in (ii), $f_3(z^I) = g(z^I) = 2 \cdot g_{\Omega,\tilde{z}}(z^I)$, $f_4(z^{-I}) = 2 \cdot g_{\Omega,\tilde{z}}(z^{-I})$, and $f_3(z^{-I}) = f_4(z^I) = 0$. Thence

$$\begin{cases} f(z^I) = \frac{f_3(z^I) + f_4(z^I)}{2} = g_{\Omega,\bar{2}}(z^I), \\ f(z^{-I}) = \frac{f_3(z^I) + f_4(z^I)}{2} = g_{\Omega,\bar{2}}(z^{-I}). \end{cases}$$

By similar method in (ii), $f|_{[p]_{\approx_{\Omega}}} = g_{\Omega,\tilde{z}}|_{[p]_{\approx_{\Omega}}}$ and $f|_{A\cap[p]_{\approx_{\Omega}}} = g|_{A\cap[p]_{\approx_{\Omega}}}$. Hence (8.9) hold by (8.17).

Proposition 8.6. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $K \subset \Omega$ be invariant, $\varepsilon > 0$, $A \subset \Omega$ be stem-finite, and $g: A \to \mathbb{H}$ be an Ω -path-slice function with $g_{\Omega,\tilde{z}}|_{A_{\Omega,\tilde{z}}\cap K} = 0$. Then there is $f \in \mathcal{SR}(\Omega)$ such that (8.9) holds.

Proof. Since A is Ω -stem-finite, $\psi_{\Omega}^{\mathbb{S}}(A)$ is finite. Then there is $z_1,...,z_m \in \mathbb{C}$ and $I_1,...,I_m \in \mathbb{S}$ and such that

$$\psi_{\Omega}^{\mathbb{S}}(A) = \bigcup_{\ell=1}^{m} \left\{ \left[\left(z_{\ell}^{I_{\ell}}, I_{\ell} \right) \right]_{\sim_{\Omega}} \right\}.$$

Let $u^L \in A$. Then $[(u^L, L)]_{\sim_{\Omega}} \in \psi_{\Omega}^{\mathbb{S}}(A)$. It implies that there is $\ell \in \{1, ..., m\}$ such that $[(u^L, L)]_{\sim_{\Omega}} = \left[\left(z_{\ell}^{I_{\ell}}, I_{\ell}\right)\right]_{\sim_{\Omega}}$, i.e. $(u^L, L) \sim_{\Omega} \left(z_{\ell}^{I_{\ell}}, I_{\ell}\right)$. According to (5.4), $u^L \approx_{\Omega} z_{\ell}^{I_{\ell}}$, i.e. $u^L \in \left[z_{\ell}^{I_{\ell}}\right]_{\approx_{\Omega}}$. It implies that

$$A \subset \bigcup_{i=1}^m \left\{ \left[z_i^{I_i} \right]_{\approx_{\Omega}} \right\}.$$

Let $v_1, ..., v_{m'} \in \{1, ..., m\}$ such that

$$\bigcup_{i=1}^{m'} \left\{ \left[z_{v_i}^{I_{v_i}} \right]_{\approx_{\Omega}} \right\} = \bigcup_{i=1}^{m} \left\{ \left[z_i^{I_i} \right]_{\approx_{\Omega}} \right\} \supset A,$$

and

$$\left[z_{v_i}^{I_{v_i}}\right]_{\approx_{\Omega}} \cap \left[z_{v_j}^{I_{v_j}}\right]_{\approx_{\Omega}} = \varnothing, \qquad i \neq j.$$

Let $i \in \{1, ..., m'\}$. By Lemma 8.5, there is $f_{\ell} \in \mathcal{SR}(\Omega)$ such that

$$f_i|_A = g_i,$$
 and $\sup_K |f_i| < \frac{\varepsilon}{m'},$

where $g_i: A \to \mathbb{H}$ defined by

$$g_i(q) := \begin{cases} g(q), & q \in \left[z_{v_i}^{I_{v_i}}\right]_{\approx_{\Omega}}, \\ 0, & \text{otherwise.} \end{cases}$$

It is easy to check that $f := \sum_{j=1}^{m'} f_j$ satisfying (8.9).

Proposition 8.7. Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, then there are Ω -invariant subsets K_{ℓ} , $\ell \in \mathbb{N}$ of Ω such that

$$(8.18) (K_{\ell})_I \subset (K_{\ell+1})_I^{\circ}, \bigcup_{\ell=0}^{+\infty} (K_{\ell})_I^{\circ} = \Omega_I, \forall I \in \mathbb{S},$$

where $(K_{\ell})_{I}^{\circ}$ denote the set of interior points of $(K_{\ell})_{I}$.

Proof. According to [18, Remark 6.8], there are compact sets $K'_{\ell} \subset \check{\Omega}$, $\ell \in \mathbb{N}$ such that

$$K'_{\ell} \subset (K'_{\ell+1})^{\circ}, \qquad (K'_{\ell})^{\wedge}_{\tilde{\Omega}} = K'_{\ell}, \qquad \text{and} \qquad \bigcup_{\ell=0}^{+\infty} (K'_{\ell})^{\circ} = \check{\Omega}.$$

By similar method of proof of [18, Remark 6.8], there are $\check{\Omega}$ -invariant subsets K''_{ℓ} , $\ell \in \mathbb{N}$ of $\check{\Omega}$ such that

(8.19)
$$K''_{\ell} \subset (K''_{\ell+1})^{\circ}, \quad \text{and} \quad \bigcup_{\ell=0}^{+\infty} (K''_{\ell})^{\circ} = \widecheck{\Omega}:$$

We may define K''_{ℓ} inductively by

$$K_0'' := \left[K_0' \cap \operatorname{Conj}(K_0') \right]_{\check{\Omega}}^{\wedge}$$

 $i_{\ell} > \ell$ being the smallest nature number such that $K''_{\ell} \subset (K'_{i_{\ell}})^{\circ}$, and

(8.20)
$$K''_{\ell+1} := \left[K'_{i_{\ell}} \cap \operatorname{Conj}_{\check{\Omega}}(K'_{i_{\ell}}) \right]_{\check{\Omega}}^{\wedge}.$$

It is easy to check that K''_{ℓ} is compact and

$$K''_{\ell} = (K''_{\ell})^{\wedge}_{\check{\Omega}} = \operatorname{Conj}_{\check{\Omega}}(K''_{\ell}).$$

Therefore, K_{ℓ} is $\check{\Omega}$ -invariant. According to (8.20), $K''_{\ell} \subset (K''_{i_{\ell}})^{\circ} \subset (K''_{\ell+1})^{\circ}$ and

(8.21)
$$\bigcup_{\ell=0}^{+\infty} (K''_{\ell})^{\circ} \supset \bigcup_{\ell=0}^{+\infty} (K'_{i_{\ell}})^{\circ} \supset \bigcup_{\ell=0}^{+\infty} (K'_{\ell})^{\circ} = \widecheck{\Omega}.$$

Therefore, (8.19) holds.

(i) Let

$$K_{\ell} := \varphi_{\breve{\Delta}}^{r_{\ell}}(K_{\ell}''), \quad \text{where} \quad r_{\ell} := \frac{r^{+}(K_{\ell}'')}{\ell + 1}.$$

Since K''_{ℓ} is $\check{\Omega}$ -invariant, $K''_{\ell} = \operatorname{Conj}_{\check{\Omega}}(K''_{\ell})$. It follows from (6.16) and

$$K''_{\ell} \subset \Omega^{r^+(K''_{\ell})}_+ \subset \Omega^{r_{\ell}}_+.$$

that

$$\begin{split} \psi_{\Omega}^{\mathbb{S}}(K_{\ell}) = & \psi_{\Omega}^{\mathbb{S}} \circ \varphi_{\check{\Omega}}^{r_{\ell}}(K_{\ell}'') = \varphi_{\check{\Omega}}^{r_{\ell}}(K_{\ell}'') \\ = \left[K_{\ell}'' \cup \operatorname{Conj}_{\check{S}}(K_{\ell}'') \right] \cap \Omega_{+}^{r_{\ell}} = K_{\ell}'' \cap \Omega_{+}^{r_{\ell}} = K_{\ell}''. \end{split}$$

According to (6.16),

$$\phi^{r_\ell}_{\scriptscriptstyle\Omega}(K_\ell) = \varphi^{r_\ell}_{\scriptscriptstyle\tilde\Omega} \circ \psi^{\mathbb{S}}_{\scriptscriptstyle\Omega}(K_\ell) = \varphi^{r_\ell}_{\scriptscriptstyle\tilde\Omega}(K_\ell'') = K_\ell.$$

It implies that K_{ℓ} is Ω -invariant.

(ii) Let $I \in \mathbb{S}$ and $q \in (K_{\ell})_I$. Then $q = z^I$, for some $z \in \mathbb{C}$. Then

$$\mu := [(z^I, I)]_{\sim_{\Omega}} \in [(K_{\ell})_I \times \{I\}]_{\sim_{\Omega}} \subset \psi_{\Omega}^{\mathbb{S}}(K_{\ell}) = K_{\ell}'' \subset (K_{\ell+1}'')^{\circ}.$$

Since $z^I \in K_\ell = \varphi_{\breve{o}}^{r_\ell}(K''_\ell) \subset \Omega^{r_\ell}$,

$$r_{\mu}^{I} = \delta_{I}(z_{I}, \mathbb{C}_{I} \setminus \Omega_{I}) = \sigma(z_{I}, \mathbb{H} \setminus \Omega) \ge r_{\ell}$$

$$= \frac{r^{+}(K_{\ell}^{"})}{\ell+1} > \frac{r^{+}(K_{\ell}^{"})}{\ell+2} \ge \frac{r^{+}(K_{\ell+1}^{"})}{(\ell+1)+1} = r_{\ell+1}.$$

Therefore, there is $r \in (0, r_{\mu}^I - r_{\ell+1})$ such that $B_{\check{\Omega}}(\mu, r) \subset (K_{\ell+1}'')^{\circ}$. Then for each $w^I \in B_I(z^I, r)$,

$$[(w^I, I)]_{\sim_{\Omega}} \subset [B_I(z^I, r) \times \{I\}]_{\sim_{\Omega}} = B_{\check{\Omega}}(\mu, r) \subset (K''_{\ell+1})^{\circ} \subset K''_{\ell+1}.$$

It implies that $[(w^I, I)]_{\sim_{\Omega}} \in \psi_{\Omega}^{\mathbb{S}}(\{w^I\}) \cap (K_{\ell+1}'') \neq \emptyset$ and

$$\sigma(w^I, \mathbb{H} \backslash \Omega) \geq \sigma(z^I, \mathbb{H} \backslash \Omega) - \sigma(z^I, w^I) > r_\mu^I - r > r_{\ell+1}.$$

Hence $w^I \in \Omega^{r_{\ell+1}}$. By (6.12), $w^I \in \varphi_{\Omega}^{r_{\ell+1}}(K_{\ell+1}'') = K_{\ell+1}$. It follows that $B_I(z^I, r) \subset (K_{\ell+1})_I$ and $z^I \in (K_{\ell+1})_I^\circ$. Therefore, $(K_\ell)_I \subset (K_{\ell+1})_I^\circ$.

(iii) Let $z^I \in \Omega_I$. Then there is $\ell \in \mathbb{N}$ such that $\sigma(z^I, \mathbb{H} \setminus \Omega) > \frac{r^+(K_0'')}{\ell+1}$. Then

$$\sigma(z^I, \mathbb{H} \setminus \Omega) > \frac{r^+(K_0'')}{\ell+1} \ge \frac{r^+(K_m'')}{m+1} = r_m, \quad \forall m > \ell.$$

According to (8.21), there is $i > \ell$ such that $[(z^I, I)]_{\sim_{\Omega}} \in K_i''$. By similar method in (ii), $z^I \in K_i \subset (K_{i+1})^{\circ}$. Then $z^I \in (K_{i+1})^{\circ}$ and then $z^I \in (K_{i+1})_I^{\circ}$, for each $z^I \in \Omega_I$. It implies that $\bigcup_{\ell=0}^{+\infty} (K_{\ell})_I^{\circ} = \Omega_I$. Therefore, Proposition 8.7 holds by (i)-(iii).

The Weierstrass factorization theorem for entire slice regular function is studied in [32], see also [16]. Some interpolation theorems for slice regular functions defined on axially symmetric domains are treated in [1,5,23]. Now we give an interpolation theorem for slice regular function defined on 2-path-symmetric slice-open sets.

Theorem 8.8. (Interpolation Theorem) Let $\Omega \in \tau_s(\mathbb{H})$ be 2-path-symmetric, $A \subset \Omega$ be stem discrete, and $g: A \to \mathbb{H}$ be an Ω -path-slice function. Then there is $f \in \mathcal{SR}(\Omega)$ such that $f|_A = g$.

Proof. According to Proposition 8.7, there are Ω-invariant subsets K_{ℓ} , $\ell \in \mathbb{N}$ of Ω such that (8.18) holds. Define

$$A_{\ell} := (A \cap K_{\ell+1}) \backslash K_{\ell},$$

and $g_{\ell}: \bigcup_{i=1}^{\ell} A_i \to \mathbb{H}$ by

$$g_{\ell}(q) = \begin{cases} g(q) - \sum_{i=1}^{\ell-1} f_i(q), & q \in A_{\ell}, \\ 0, & \text{otherwise} \end{cases}$$

Here, by Proposition 8.6, we can construct a function $f_{\ell} \in \mathcal{SR}(\Omega)$ such that

$$|f_{\ell}|_{\bigcup_{i=1}^{\ell} A_i} = g_{\ell}, \quad \text{and} \quad \sup_{K_{\ell-1}} |f_{\ell}| < \frac{\varepsilon}{2^{\ell}}.$$

Let $f := \sum_{i=1}^{+\infty} f_i$. Since $\sum_{i=1}^{+\infty} f_i$ converges uniformly on $(K_j)_I$, $j \in \mathbb{N} \setminus \{0\}$, we have f_I is I-holomorphic. It implies that f is slice regular.

Let $a \in A$. Then $a \in A_i$ for some $i \in \mathbb{N}$. Then

$$f(a) = \sum_{j=1}^{+\infty} f_j(a) = \sum_{j=1}^{i-1} f_j(a) + f_i(a) + \sum_{j=i+1}^{+\infty} f_j(a)$$
$$= \sum_{j=1}^{i-1} f_j(a) + \left[g(a) - \sum_{j=1}^{i-1} f_j(a) \right] + 0 = g(a).$$

It implies that $f|_A = g$. Hence Theorem 8.8 holds.

9. Proof of Cartan-Thullen Theorem

In this section, we will prove the counterpart of the Cartan-Thullen theorem in the theory of slice regular functions. As in the complex case, we need a notion of domain of existence, see Definition 9.2. The preparation results provided in the previous sections together with the description of slice regularly convex domains in the next proposition allow to prove the main Theorem 2.10.

Proposition 9.1. A set $\Omega \in \tau_s(\mathbb{H})$ is slice regularly convex, if and only if Ω is 2-path-symmetric.

Proof. Let Ω be a slice regularly convex slice-open set and let us suppose, by absurd, that Ω is not 2-path-symmetric. Then there exists $\gamma \in \mathscr{P}_0(\mathbb{C})$ and $I_1, I_2, J \in \mathbb{S}$, $I_1 \neq I_2$, such that $\gamma^{I_1}, \gamma^{I_2} \subset \Omega$ but $\gamma^J \nsubseteq \Omega$. Since $\gamma^J(0) = \gamma^{I_1}(0) \in \Omega$, there is $t \in (0,1]$ such that $\gamma^J([0,t]) \subset \Omega$ and $\gamma^J([0,t]) \nsubseteq \Omega$. Let

$$K := \gamma^{I_1}([0,1]) \cup \gamma^{I_2}([0,1])$$

be a finite-non-symmetric compact subset. Then $\gamma^{I_1}, \gamma^{I_2} \subset K$. By the representation formula,

$$\sup_{\gamma^{J}((0,t))} |f| \le |c_1| \sup_{\gamma^{I_1}((0,t))} |f| + |c_2| \sup_{\gamma^{I_2}((0,t))} |f|, \qquad \forall \ f \in \mathcal{SR}(\Omega).$$

where

$$(c_1, c_2) = (1, J) \begin{pmatrix} 1 & I_1 \\ 1 & I_2 \end{pmatrix}^{-1}.$$

It implies that $\gamma^J([0,t)) \subset \widehat{K}_{\Omega} \cap \Omega_J = (\widehat{K}_{\Omega})_J$ and $(\widehat{K}_{\Omega})_J$ is not relatively compact in Ω_J . Thus Ω is not slice regularly convex, contradicting the hypothesis.

To prove the converse, suppose that Ω is 2-path-symmetric and let $K \subset \Omega$ be a finite-non-symmetric compact. According to (2.1), $K_{2ps} \subset \Omega_{2ps}$. By Remark 3.10 (i), $K^{2ps} \subset \Omega^{2ps}$. Let $I, J \in \mathbb{S} \backslash \mathbb{S}_K^{2ps}$ and $\mu \in [K_I \times \{I\}]_{\sim_{\Omega}}$. According to (2.2), $z_{\mu}^I \in K_I = (K_{2ps})^I \subset (\Omega_{2ps})^I$. Therefore $z_{\mu} \in K_{2ps} \subset \Omega_{2ps}$, and $z_{\mu}^J \in (K_{2ps})^J = K_J$. By (5.6), $(z_{\mu}^I, I) \sim_{\Omega} (z_{\mu}^J, J)$ and

$$[(z_\mu^I,I)]_{\sim_\Omega} = [(z_\mu^J,J)]_{\sim_\Omega} \subset [K_J \times \{J\}]_{\sim_\Omega}.$$

It implies that $[K_I \times \{I\}]_{\sim_{\Omega}} \subset [K_J \times \{J\}]_{\sim_{\Omega}}$. Similarly, $[K_J \times \{J\}]_{\sim_{\Omega}} \subset [K_I \times \{I\}]_{\sim_{\Omega}}$. Hence

$$(9.1) [K_I \times \{I\}]_{\sim_{\Omega}} = [K_J \times \{J\}]_{\sim_{\Omega}}, \forall I, J \in \mathbb{S} \setminus \mathbb{S}_K^{2ps}.$$

Let $L \in \mathbb{S}$. Since K_L is compact in $\tau(\mathbb{C}_L)$, $K_L \times L$ is compact in $\tau(\mathbb{C}_L) \times \{L\}$. According to Proposition 5.4, ι_{Ω}^I is a homeomorphism. Therefore $[K_I \times \{I\}]_{\sim_{\Omega}} =$ $\iota_{\Omega}^{I}(K_{L} \times L)$ is compact in $[\Omega_{I} \times \{I\}]_{\sim_{\Omega}}$ and also compact in $\check{\Omega} \supset [\Omega_{I} \times \{I\}]_{\sim_{\Omega}}$. Let $J_{0} \in \mathbb{S} \backslash \mathbb{S}_{K}^{2ps}$. Since \mathbb{S}_{K}^{2ps} is finite and (9.1),

$$\psi_{\Omega}^{\mathbb{S}}(K) = \bigcup_{I \in \mathbb{S}} [K_I \times \{I\}]_{\sim_{\Omega}}$$

$$= \left(\bigcup_{I \in \mathbb{S}_K^{2ps}} [K_I \times \{I\}]_{\sim_{\Omega}}\right) \bigcup \left(\bigcup_{I \in \mathbb{S} \setminus \mathbb{S}_K^{2ps}} [K_I \times \{I\}]_{\sim_{\Omega}}\right)$$

$$= \left(\bigcup_{I \in \mathbb{S}_K^{2ps}} [K_I \times \{I\}]_{\sim_{\Omega}}\right) \bigcup [K_{J_0} \times \{J_0\}]_{\sim_{\Omega}}$$

is compact in $\check{\Omega}$. It follows from $\operatorname{Conj}_{\check{\Omega}}([K_I \times \{I\}]_{\sim_{\Omega}}) = [K_{-I} \times \{-I\}]_{\sim_{\Omega}}$ that $\operatorname{Conj}_{\check{\Omega}}\left[\psi_{\Omega}^{\mathbb{S}}(K)\right] = \psi_{\Omega}^{\mathbb{S}}(K)$. By similar method in the proof of Proposition 8.7, $K' := \left[\psi_{\Omega}^{\mathbb{S}}(K)\right]_{\check{\Omega}}^{\wedge}$ is $\check{\Omega}$ -invariant and $K'' := \varphi_{\check{\Omega}}^{r^+(K')}(K')$ is Ω -invariant with $K' = \psi_{\Omega}^{\mathbb{S}}(K'')$. Next we show that $(K'')_I$ is compact. Let $I \in \mathbb{S}$ and $\{w_\ell^I\}_{\ell \in \mathbb{N}}$ be a Cauchy sequence in K_I'' with

$$w^I := \lim_{\ell \to +\infty} w^I_{\ell}.$$

Since $K'' = \varphi_{\check{\Omega}}^{r^+(K')}(K') \subset \Omega^{r^+(K')}, \ w^I \in \Omega^{r^+(K')} \subset \Omega_I$. Then $\{(w_\ell^I, I)\}_{\ell \in \mathbb{N}}$ is a Cauchy sequence in $K_I'' \times \{I\}$. By Proposition 5.4, $\{[(w_\ell^I, I)]_{\sim_{\Omega}} = \iota_{\Omega}^I((w_\ell^I, I))\}$ is a Cauchy sequence in $\check{\Omega}$ and

$$\lim_{\ell \to +\infty} [(w_\ell^I, I)]_{\sim_\Omega} = i_\Omega^I \left(\lim_{\ell \to +\infty} (w_\ell^I, I) \right) = i_\Omega^I ((w^I, I)) = [(w^I, I)]_{\sim_\Omega}.$$

Since $[(w_\ell^I,I)]_{\sim_{\Omega}} \in \psi_{\Omega}^{\mathbb{S}}(\{w_\ell^I\}) \subset \psi_{\Omega}^{\mathbb{S}}(K'') = K'$ and K' is compact, $[(w^I,I)]_{\sim_{\Omega}} \in K'$. It implies that

$$[(w^I,I)]_{\sim_{\Omega}} \in \psi_{\Omega}^{\mathbb{S}}(\{w^I\}) \cap K' \neq \varnothing.$$

By (6.12), $w^I \in \varphi^{r^+(K')}_{\S}(K') = K''$. It implies that $w^I \in (K'')_I$ and $(K'')_I$ is compact.

Let $p \in \Omega \backslash K''$. If $p \in \widehat{K}_{\Omega}$, then there is $C_p > 0$ such that

(9.2)
$$|f(p)| \le C_p \sup_{q \in K} |f(q)|, \qquad \forall \ f \in \mathcal{SR}(\Omega).$$

However, by Proposition 8.6, there is $h \in \mathcal{SR}(\Omega)$ such that

$$h(p) = 1,$$
 and $\sup_{q \in K} |h(q)| \le \sup_{q \in K''} |h(q)| < \frac{1}{C_p + 1},$

a contradiction to (9.2). Hence $\widehat{K}_{\Omega} \subset K''$ and then $(\widehat{K}_{\Omega})_I \subset (K'')_I \in \Omega_I$, for each $I \in \mathbb{S}$. It implies that K is slice regularly convex.

Definition 9.2. Let $I \in \mathbb{S}$, $U \in \tau(\mathbb{C}_I)$ and $\mathcal{F} \subset \mathcal{O}_I(U)$. U is called an \mathcal{F} -domain of existence if there are no $U_1, U_2 \in \tau(\mathbb{C}_I)$ with the following properties:

- (i) $\varnothing \neq U_1 \subset U_2 \cap U$.
- (ii) U_2 is connected in $\tau(\mathbb{C}_I)$ and $U_2 \nsubseteq U$.
- (iii) For each $f \in \mathcal{F}$, there is $\widetilde{f} \in \mathcal{SR}(U_2)$ such that $f = \widetilde{f}$ in U_1 .

If there are sets $U, U_1, U_2 \in \tau(\mathbb{C}_I)$ satisfying (i)-(iii), then we call (U, U_1, U_2) a \mathcal{F} -triple. Moreover, we say that (U, U_1, U_2) is **special**, if U_1 is a connected component of $U \cap U_2$ in \mathbb{C}_I .

We now have all the needed terminology and arguments to prove the main result of this paper, namely Theorem 2.10, that is repeated here for the reader's convenience:

Theorem 9.3. (Cartan-Thullen) Let $\Omega \in \tau_s(\mathbb{H})$. Then the following statements are equivalent:

- (i) Ω is the domain of existence of some slice regular function.
- (ii) Ω is a domain of slice regularity.
- (iii) Ω is slice regularly convex.
- (iv) Ω is 2-path-symmetric.

Proof. (iii)⇔(iv) follows directly by Proposition 9.1.

- (i)⇒(ii) follows by definition.
- (ii) \Rightarrow (iv) Directly by [20, Proposition 9.7].
- (iv) \Rightarrow (i) Let Ω be 2-path-symmetric. We only need to prove that every slice-connected component of Ω is a domain of existence of some slice regular function; without loss of generality, we can assume that Ω is slice-connected.

Let $J_0 \in \mathbb{S} \setminus \mathbb{S}_{\Omega}^{2ps}$ and consider

$$A_{[I]} := \{ q \in \Omega_I : q = x + yI, \text{ for some } x, y \in \mathbb{Q} \}, \quad \forall I \in \mathbb{S}$$

and

$$A := \bigcup_{I \in \mathbb{S}_{\mathcal{O}}^{2ps} \cup \{J_0\}} A_{[I]} \times \{I\}.$$

By Proposition 3.8, $\mathbb{S}_{\Omega}^{2ps}$ is countable, and so is A. Let $\vartheta_A:A\to\mathbb{N}$ be a bijection and $(a_i,I_i):=\vartheta_A^{-1}(i)$.

Let

$$(9.3) b_i \in B_{I_i}(a_i, \sigma(a, \mathbb{H} \setminus \Omega))$$

with $\sigma(b_i, \mathbb{H} \setminus \Omega) < \frac{1}{i+1}$ and $[b_i]_{\approx_{\Omega}} \neq [b_j]_{\approx_{\Omega}}$ for j < i. Set

(9.4)
$$B_{i} := \begin{cases} [b_{i}]_{\approx_{\Omega}}, & \mathbb{S}^{+}_{[(b_{i},I_{i})]_{\sim_{\Omega}}} = \mathbb{S}, \\ \{b_{i}\}, & \text{otherwise}, \end{cases}$$

and

$$B:=\bigcup_{i\in\mathbb{N}}B_i.$$

According to Proposition 6.6, $B_{\ell} \cap \Omega^{\frac{1}{i+1}} = \emptyset$, for all $\ell \geq i$. It implies that

$$B\cap\Omega^{\frac{1}{\imath+1}}\subset\bigcup_{\ell=0,1,\ldots,\imath-1}[b_\ell]_{\approx_\Omega}.$$

It follows from (6.4), (6.5) and Proposition 3.2 (i) that

$$\psi_{\scriptscriptstyle \Omega}^{\mathbb{S}}\left(B\cap\Omega^{\frac{1}{\mathfrak{r}+1}}\right)\subset\bigcup_{\ell=0,1,...,\mathfrak{r}-1}\bigg\{[(b_{\ell},I_{\ell})]_{\sim_{\Omega}},[(b_{\ell},-I_{\ell})]_{\sim_{\Omega}}\bigg\}.$$

Then $\left|\psi_{\Omega}^{\mathbb{S}}\left(B\cap\Omega^{\frac{1}{i+1}}\right)\right|\leq 2i$, i.e. the cardinality of $\psi_{\Omega}^{\mathbb{S}}\left(B\cap\Omega^{\frac{1}{i+1}}\right)$ is finite and has no accumulation point in $\tilde{\Omega}$. It implies that B is stem-discrete. Define $g: B \to \mathbb{H}$ by

$$g(q) = i, \qquad q \in [b_i]_{\approx_{\Omega}}.$$

By Interpolation Theorem 8.8, there is $f \in \mathcal{SR}(\Omega)$ such that $f|_B = g$.

By the fact in complex analysis, for each $I\in\mathbb{S}^{2ps}_{\Omega},\ f_I$ can not extend to a holomorphic function near any point of $\partial_I(\Omega_I)$, i.e.

(9.5)
$$\Omega_I$$
 is the domain of existence of f_I , for each $I \in \mathbb{S}^{2ps}_{\Omega}$.

Suppose that there is $J \in \mathbb{S} \backslash \mathbb{S}_{\Omega}^{2ps}$ such that Ω_J is not the domain of existence of f_J . Then there is a special $\{f_J\}$ -triple (Ω_J, U_1, U_2) and a J-holomorphic functions

h: $U_2 \to \mathbb{H}$ such that $h|_{U_1} = f_J|_{U_1}$. Let $z^J \in \partial_J(\Omega_J) \cap \partial_J(U_1) \cap U_2$. If $z^I \in \Omega$ for some $I \in \mathbb{S}$, then it follows from $z^J \notin \Omega$ that $z \notin \Omega_{2ps}$. Therefore $I \in \mathbb{S}_2^{2ps}$ and $z^{-I} \in \partial_I(\Omega_I)$. By representation formula with similar method in the proof of [20, Proposition 9.7], f_I can extend to a I-holomorphic function near $z^{-I} \in \partial_I(\Omega_I)$, a contradiction to (9.5). Otherwise, $z^I \notin \Omega$ for all $I \in \mathbb{S}$. Then there is r > 0 such that $B_J(z^J, 2r) \subset U_2$.

Denote

$$W := \Psi^{i}_{J}(U_{1} \cap B_{J}(z^{J}, r)).$$

Let $w \in W$ and $L \in \mathbb{S}$. Since $|w^L - z^L| < r$ and $z^L \in \partial_L(\Omega_L)$, we have $\sigma(w^L, \mathbb{H} \setminus \Omega) < r$ r. It follows from

$$B_L(w^L, \sigma(w^L, \mathbb{H} \setminus \Omega)) \subset B_L(z^J, 2r) \cap \Omega_L \subset \Psi_J^L(U_2) \cap \Omega_L = \Psi_J^L(U_2 \cap \Omega),$$

the connectedness of $B_L(w^L, \sigma(w^L, \mathbb{H} \backslash \Omega))$ and (Ω_J, U_1, U_2) being special that

(9.6)
$$B_L(w^L, \sigma(w^L, \mathbb{H} \setminus \Omega)) \subset \Psi_J^L(U_1).$$

It implies from $U_1 \subset \Omega_J \subset \Omega^{2ps} \subset \Omega$ that

$$\Psi_J^I(U_1) \subset \Omega$$
, and $\sigma(w^L, \mathbb{H} \setminus \Omega) = \sigma(w^L, \Psi_J^L(U_1)) = \sigma(W^J, U_1)$.

Therefore,

(9.7)
$$\mathbb{S}^{+}_{[(w^{L},L)]_{\infty}} = \mathbb{S}, \quad \forall w \in W \text{ and } L \in \mathbb{S}.$$

Denote

$$W_* := \Psi_J^i \left(U_1 \cap B_J \left(z^J, \frac{r}{2} \right) \right)$$

Let $w_1 \in W_*$ and $L_1 \in \mathbb{S}$. By $\sigma(w_1^{L_1}, \mathbb{H} \setminus \Omega) \leq |w_1^{L_1} - z^{L_1}| < \frac{r}{2}$, we have

$$B_{L_1}\left(w_1^{L_1}, \sigma\left(w_1^{L_1}, \mathbb{H}\backslash\Omega\right)\right) \subset B_{L_1}(z^{L_1}, r).$$

According to (9.6),

(9.8)
$$B_{L_1}\left(w_1^{L_1}, \sigma\left(w_1^{L_1}, \mathbb{H}\backslash\Omega\right)\right) \subset W^{\mathbb{S}}.$$

Let $J_1 \in \mathbb{S} \setminus \mathbb{S}_{\Omega}^{2ps}$. Then there is $\{c_\ell\}_{\ell \in \mathbb{N}}$ with $c_\ell \in A_{[J_1]} \cap W_*^{J_1}$ such that

$$\lim_{\ell \to +\infty} c_{\ell} = z^{J_1}.$$

Since $c_{\ell} \in \Omega_{J_1} \subset \Omega^{2ps}$,

$$\Psi_{J_1}^{J_0}(c_\ell) \in A_{[J_0]} \subset A.$$

Then there is $\{v_\ell\}_{\ell\in\mathbb{N}}$ with $v_\ell\in\mathbb{N}$ such that $a_{v_\ell}=\Psi_{J_1}^{J_0}(c_\ell)$. It follows from

$$a_{v_{\ell}} = \Psi_{J_1}^{J_0}(c_{\ell}) \in W_*^{\mathbb{S}},$$

(9.3) and (9.8) that

$$b_{v_{\ell}} \in B\left(a_{v_{\ell}}, \sigma\left(a_{v_{\ell}}, \mathbb{H} \backslash \Omega\right)\right) \subset W^{\mathbb{S}}.$$

According to (9.7) and (9.4), $B_{v_{\ell}} = [b_{v_{\ell}}]_{\approx_{\Omega}}$. Since $b_{v_{\ell}} \in \Omega_{J_0} \subset \Omega^{2ps}$,

$$d_{v_{\ell}} := \Psi_{J_0}^{J_1}(b_{v_{\ell}}) \in B_{v_{\ell}}.$$

Then

$$\lim_{i \to +\infty} d_{v_i} = \lim_{i \to +\infty} \left(\Psi^{J_0}_{J_1}(b_{v_i}) \right) = \Psi^{J_0}_{J_1} \lim_{i \to +\infty} b_{v_i} = \Psi^{J_0}_{J_1} \lim_{i \to +\infty} a_{v_i} = \lim_{i \to +\infty} c_{v_i} = z^{J_1}.$$

By definition,

$$h(z^{J}) = \lim_{i \to +\infty} h(\Psi_{I}^{J}(d_{v_{i}})) = \lim_{i \to +\infty} f(\Psi_{I}^{J}(d_{v_{i}}))$$
$$= \lim_{i \to +\infty} g(\Psi_{I}^{J}(d_{v_{i}})) = \lim_{i \to +\infty} v_{i} = +\infty,$$

a contradiction. Therefore Ω_J is the domain of existence of f_J .

In summary, Ω_I is the domain of existence of f_I for each $I \in \mathbb{S}$. Following the method in the proof of [20, Proposition 9.4], Ω is the domain of existence of f. \square

Conflict of Interest. The authors declare that they have no competing interests regarding the publication of this paper.

Availability of data. There are no data associated with the research in this paper.

References

- [1] Daniel Alpay, Vladimir Bolotnikov, Fabrizio Colombo, and Irene Sabadini, Self-mappings of the quaternionic unit ball: multiplier properties, the Schwarz-Pick inequality, and the Nevanlinna-Pick interpolation problem, Indiana Univ. Math. J. **64** (2015), no. 1, 151–180. MR3320522
- [2] Daniel Alpay, Fabrizio Colombo, Jonathan Gantner, and Irene Sabadini, A new resolvent equation for the S-functional calculus, J. Geom. Anal. 25 (2015), no. 3, 1939–1968. MR3358079
- [3] Daniel Alpay, Fabrizio Colombo, and Irene Sabadini, Pontryagin-de Branges-Rovnyak spaces of slice hyperholomorphic functions, J. Anal. Math. 121 (2013), 87–125. MR3127378
- [4] ______, Slice hyperholomorphic Schur analysis, Operator Theory: Advances and Applications, vol. 256, Birkhäuser/Springer, Cham, 2016. MR3585855
- [5] Vladimir Bolotnikov, On the intersection of left and right ideals in the ring of quaternion polynomials, J. Algebra Appl. 22 (2023), no. 12, Paper No. 2350249, 23. MR4663899
- [6] Henri Cartan and Peter Thullen, Zur Theorie der Singularitäten der Funktionen mehrerer komplexen Veränderlichen, Math. Ann. 106 (1932), no. 1, 617–647. MR1512777
- [7] Fabrizio Colombo and Jonathan Gantner, Quaternionic closed operators, fractional powers and fractional diffusion processes, Operator Theory: Advances and Applications, vol. 274, Birkhäuser/Springer, Cham, 2019. MR3967697
- [8] Fabrizio Colombo, Jonathan Gantner, and David P. Kimsey, Spectral theory on the S-spectrum for quaternionic operators, Operator Theory: Advances and Applications, vol. 270, Birkhäuser/Springer, Cham, 2018. MR3887616
- [9] Fabrizio Colombo, Jonathan Gantner, David P. Kimsey, and Irene Sabadini, Universality property of the S-functional calculus, noncommuting matrix variables and Clifford operators, Adv. Math. 410 (2022), Paper No. 108719, 39. MR4496722
- [10] Fabrizio Colombo, Graziano Gentili, Irene Sabadini, and Daniele Struppa, Extension results for slice regular functions of a quaternionic variable, Adv. Math. 222 (2009), no. 5, 1793– 1808. MR2555912

- [11] Fabrizio Colombo, Roman Lávivcka, Irene Sabadini, and Vladimír Souvcek, The Radon transform between monogenic and generalized slice monogenic functions, Math. Ann. 363 (2015), no. 3-4, 733-752. MR3412341
- [12] Fabrizio Colombo and Irene Sabadini, The quaternionic evolution operator, Adv. Math. 227 (2011), no. 5, 1772–1805. MR2803786
- [13] Fabrizio Colombo, Irene Sabadini, and Daniele C. Struppa, Slice monogenic functions, Israel J. Math. 171 (2009), 385–403. MR2520116
- [14] _____, Duality theorems for slice hyperholomorphic functions, J. Reine Angew. Math. **645** (2010), 85–105. MR2673423
- [15] ______, Noncommutative functional calculus, Progress in Mathematics, vol. 289, Birkhäuser/Springer Basel AG, Basel, 2011. Theory and applications of slice hyperholomorphic functions. MR2752913
- [16] ______, Entire slice regular functions, SpringerBriefs in Mathematics, Springer, Cham, 2016. MR3585395
- [17] Charles. G. Cullen, An integral theorem for analytic intrinsic functions on quaternions, Duke Math. J. 32 (1965), 139–148. MR0173012
- [18] Jean-Pierre Demailly, Complex analytic and differential geometry, 2012.
- [19] Xinyuan Dou, Ming Jin, Guangbin Ren, and Irene Sabadini, A new approach to slice analysis via slice topology, Adv. Appl. Clifford Algebr. 31 (2021), no. 5, Paper No. 67, 36. MR4309797
- [20] Xinyuan Dou, Guangbin Ren, and Irene Sabadini, Extension theorem and representation formula in non-axially-symmetric domains for slice regular functions, J. Eur. Math. Soc. (JEMS) 25 (2023), no. 9, 3665–3694. MR4634680
- [21] ______, A representation formula for slice regular functions over slice-cones in several variables, Ann. Mat. Pura Appl. (4) 202 (2023), no. 5, 2421–2446. MR4634271
- [22] Klaus Fritzsche and Hans Grauert, From holomorphic functions to complex manifolds, Graduate Texts in Mathematics, vol. 213, Springer-Verlag, New York, 2002. MR1893803
- [23] Sorin G. Gal and Irene Sabadini, Walsh equiconvergence theorems in the quaternionic setting, Complex Var. Elliptic Equ. 59 (2014), no. 12, 1589–1607. MR3254690
- [24] Jonathan Gantner, Operator theory on one-sided quaternion linear spaces: intrinsic S-functional calculus and spectral operators, Mem. Amer. Math. Soc. 267 (2020), no. 1297, iii+101. MR4195727
- [25] Graziano Gentili, Simon Salamon, and Caterina Stoppato, Twistor transforms of quaternionic functions and orthogonal complex structures, J. Eur. Math. Soc. (JEMS) 16 (2014), no. 11, 2323–2353. MR3283400
- [26] Graziano Gentili and Caterina Stoppato, Power series and analyticity over the quaternions, Math. Ann. 352 (2012), no. 1, 113–131. MR2885578
- [27] ______, Geometric function theory over quaternionic slice domains, J. Math. Anal. Appl. 495 (2021), no. 2, Paper No. 124780, 38. MR4182982
- [28] _____, A local representation formula for quaternionic slice regular functions, Proc. Amer. Math. Soc. 149 (2021), no. 5, 2025–2034. MR4232195
- [29] Graziano Gentili, Caterina Stoppato, and Daniele C. Struppa, Regular functions of a quaternionic variable, Springer Monographs in Mathematics, Springer, Heidelberg, 2013. MR3013643
- [30] Graziano Gentili and Daniele C. Struppa, A new theory of regular functions of a quaternionic variable, Adv. Math. 216 (2007), no. 1, 279–301. MR2353257
- [31] ______, Regular functions on the space of Cayley numbers, Rocky Mountain J. Math. 40 (2010), no. 1, 225–241. MR2607115
- [32] Graziano Gentili and Irene Vignozzi, The Weierstrass factorization theorem for slice regular functions over the quaternions, Ann. Global Anal. Geom. 40 (2011), no. 4, 435–466. MR2836832
- [33] Riccardo Ghiloni and Alessandro Perotti, Slice regular functions on real alternative algebras, Adv. Math. 226 (2011), no. 2, 1662–1691. MR2737796
- [34] Fritz Hartogs, Einige folgerungen aus der cauchyschen integralformel bei funktionen mehrerer veränderlichen, Sitzungsberichte, vol. 1906,6, München, 1907.
- [35] Steven G. Krantz, Function theory of several complex variables, AMS Chelsea Publishing, Providence, RI, 2001. Reprint of the 1992 edition. MR1846625
- [36] Junjiro Noguchi, Analytic function theory of several variables, Springer, Singapore, 2016. Elements of Oka's coherence. MR3526579

[37] Caterina Stoppato, A new series expansion for slice regular functions, Adv. Math. 231 (2012), no. 3-4, 1401–1416. MR2964609

INDEX

2-path-symmetric, 4	$\psi_{\Omega}^{\mathbb{S}'},\varphi_{\Omega}^{\rho},\phi_{\Omega}^{\rho},\phi_{\Sigma}^{\rho},24$
$B_{\mathbb{C}}^{\mu}, B_{\mathbb{Q}}^{\mu}, 28$	τ_{Ω}^{Ω} , τ_{Ω}^{Ω} , τ_{Ω}^{Ω} , τ_{Ω}^{Ω} , τ_{Ω}^{Ω}
$B_{\check{\mathbf{O}}}(\mu,r), 15$	
P(X), 12	$egin{array}{l} au_s,\ 4\ \widehat{K}_\Omega,\ 6 \end{array}$
$U^{\sim}, 24$	K_{Ω} , σ
U^r , 21	$\widetilde{\Omega}$, 10
$U_{\Omega}^{\gamma}, \widehat{U}_{\Omega}, 20$	$f_{\Omega, \widetilde{2}}, 35$
	$g_{\mu}, 29$
$U_{\alpha}^{[I]}, U_{\alpha}^{r,I}, 29$	$g_{\Omega}^{I},31$
$U_{\Omega,2},U_{\Omega,ar{2}},35$	$i_{\Omega}^{I},12$
$[U]_{\approx_{\Omega}}^{\cup}, 24$	$r^+, 20$
Ω -2-path-symmetric, 35	$r^+(U), \ r_U^+, \ 20$
Ω -path-slice, 35	$r_{\mu}^{+},r_{\mu}^{-},r_{\mu}^{I},14$
$\Omega_{+}^{t}, \Omega_{+}^{t^{\circ}}, \check{\Omega}_{+}^{t}, \check{\Omega}_{+}^{t^{\circ}}, 23$	z^{I} , $z^{\mathbb{S}'}$, U^{I} , $U^{\mathbb{S}'}$, 4
$\Omega^{2ps}, 10$	$\operatorname{Conj}_{reve{\omega}}, \overline{\mu}, 22$
$\Omega_{\mathbb{R}},8$	(I-)holomorphic, 5
$\Omega_{2ps}, 6$	$(\tilde{\Omega}$ -)invariant, $(\Omega$ -)invariant, 27
$\Psi_{\underline{I}}^{J},13$	$(\check{\Omega})$ stem discrete, $(\check{\Omega})$ stem-finite, 27
$\Psi_i^I,~4$	(12-)stem discrete, (12-)stem-inite, 21
\approx_{Ω} , 7	Axially symmetric, 10
$\breve{\Omega},12$	Axially symmetric completion, 10
$reve{\Omega}^{{\scriptscriptstyle [I]}},29$	iminify symmotric completion, 10
$\delta_{\Omega}, 20$	Domain of existence, 6
$\gamma^I,4$	Domain of slice regularity, 6
$\mathbb{S}, 3$	3,4,4,4,7,7
\mathbb{S}^2_* , 3	Finite-non-symmetric compact, 6
\mathbb{S}^2_* , 3 $\mathbb{S}^{2ps}_{\Omega}$, 6	•
$\mathbb{S}_{\mu}^{::}$, 13	Slice regular, 5
$\mathbb{S}_{\mu}^{+}, \mathbb{S}_{\mu}^{-}, 14$	Slice regularly convex, 6
\mathcal{F} -domain of slice existence, 5, 44	Slice regularly convex hull, 6
$\mathcal{L}_z^w, 8$	Slice topology, 4
$\mathcal{SR}(\Omega)$, 5	Slice-connected, slice-open,
$\mathscr{O}(\Omega),20$	st-domain, 4
$\mathcal{O}_{I}(U), 5$	Slice-triple, 6, 45
$\mathscr{P}_0(U), 4$	Special slice-triple, 6, 45
- \ / /	- , ,

 $Email\ address, \ {\tt Xinyuan\ Dou:\ douxinyuan@ustc.edu.cn}$

Department of Mathematics, University of Science and Technology of China, Hefei 230026, China

INSTITUTE OF MATHEMATICS, AMSS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA

Email address, Ming Jin: mjin@must.edu.mo

Faculty of Innovation Engineering, Macau University of Science and Technology, Macau, China

 ${\it Email \ address}, \ {\it Guangbin \ Ren:} \ {\tt rengb@ustc.edu.cn}$

INDEX 51

Department of Mathematics, University of Science and Technology of China, Hefei 230026, China

Email address, Irene Sabadini: irene.sabadini@polimi.it

Dipartimento di Matematica, Politecnico di Milano, Via Bonardi, 9, 20133 Milano, Italy