

# First Order Linear Proportional Difference Equation with Integration Factor the $(s, t)$ -Pantograph Function

Ronald Orozco López

May 21, 2024

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Preliminaries</b>	<b>3</b>
2.1	Notation for $q$ -calculus . . . . .	3
2.2	Generalized Fibonacci polynomials . . . . .	4
2.3	Calculus on generalized Fibonacci polynomials . . . . .	4
2.4	Deformed $(s, t)$ -exponential function . . . . .	6
2.5	The $(s, t)$ -pantograph function . . . . .	7
<b>3</b>	<b>Deformed <math>(s, t)</math>-chain rule</b>	<b>9</b>
<b>4</b>	<b>First order functional-difference <math>(s, t)</math>-equation</b>	<b>12</b>
4.1	An equation general . . . . .	12
4.1.1	Equation $\mathbf{D}_{s,t}y(x) + \alpha(x) \frac{\exp_{s,t}[uA(x),u]}{\exp_{s,t}[A(\varphi_{s,t}x),u]}y(\varphi'_{s,t}x) = \beta(x)$ . . . . .	13
4.1.2	Solution of the equation $\mathbf{D}_{s,t}y + \alpha(x)y(\varphi'_{s,t}x) = \beta(x)$ . . . . .	14
4.1.3	Equation involving partial theta function . . . . .	16
4.2	Equation $\mathbf{D}_{s,t}y = \alpha(ay(x) + by(ux)) + \beta(x)$ . . . . .	16
4.3	Operator method . . . . .	19
4.4	Deformed $(s, t)$ -Bernoulli Equation . . . . .	20

## Abstract

In this paper we find solutions to first-order linear proportional difference equations via the  $(s, t)$ -integration factor method. The  $(s, t)$ -integration factor involves the  $(s, t)$ -Pantograph function, which is a generalization of the partial Theta function. Other equations are solved including the  $(s, t)$ -analogue of the Bernoulli equation.

2020 *Mathematics Subject Classification*: Primary 39B22. Secondary 11B39, 34K06.

*Keywords*: Proportional equation, generalized Fibonacci polynomials, partial Theta function, Bernoulli equation.

# 1 Introduction

Orozco [8] studied the existence of equations in differences with proportional delay of the form

$$y(ax) - y(bx) = (a - b)xf(x, y(x), y(ux)), \quad y(\eta) = \xi, \quad (1)$$

where  $a, b, u \in \mathbb{R}$ . In this paper, we will be mainly interested in solving equations in proportional differences of the form

$$y(ax) - y(bx) = (a - b)x(\beta(x) - \alpha(x)y(ux)), \quad (2)$$

or equivalently

$$\mathbf{D}_{a,b}y(x) + \alpha(x)y(ux) = \beta(x),$$

by means of the integration factor method involving the  $(s, t)$ -Pantograph function

$$\mathbf{E}_{s,t}(a, b; x, u) = \sum_{n=0}^{\infty} (a \oplus b)_{1,u}^{(n)} \frac{x^n}{\{n\}_{s,t}!}$$

which is a solution of the equation

$$\mathbf{D}_{s,t}y(x) = ay(x) + by(ux). \quad (3)$$

The special case  $s = 2$  and  $t = -1$  is studied in [5, 6]. The function  $\mathbf{E}_{s,t}(a, b; x, u)$  generalises the  $(s, t)$ -exponential and the partial Theta function  $\Theta_0(x, y)$  of Ramanujan. Some solutions will then be expressed in terms of partial Theta functions. Since the integration factor  $I = \exp(\int_{\eta}^x \alpha(t)dt)$  for ordinary differential equations involves the composition of functions, then it will be necessary to introduce a type of composition analogous to the  $q$ -composition of Gessel [9] and Johnson [10].

In addition, we shall give a power series solution to the equation

$$y(ax) - y(bx) = (a - b)x(\alpha cy(x) + dy(ux)) + \beta(x),$$

and we will provide a solution via operators for the equation

$$y(ax) - y(bx) = (a - b)x(c\beta y(x) + \gamma y(ux) + \delta \mathbf{E}_{s,t}(c, d; \alpha x, u)).$$

Finally, we will consider the deformed  $(s, t)$ -Bernoulli equation

$$D_{\varphi^{n-1}, \varphi'^{(n-1)}}y(x) + \alpha(x) \frac{(D_q \mathbf{E}_{s,t})(a, b; x, u) \square A(x)}{\mathbf{E}_{s,t}[a, b; A(\varphi_{s,t}), u]} y(\varphi_{s,t}^{n-1}x) = \beta(x) \prod_{j=0}^{\infty} \frac{y(q^j \varphi_{s,t}^{n-1}x)}{y(q^{n+j} \varphi_{s,t}^{n-1}x)},$$

where

$$D_{\varphi^{n-1}, \varphi'^{(n-1)}}f(x) = \frac{f(\varphi^{n-1}x) - f(\varphi'^{(n-1)})}{(\varphi^{n-1} - \varphi'^{(n-1)})x}$$

is a  $(\varphi^{n-1}, \varphi'^{(n-1)})$ -derivative.

We divide this paper as follows. Section 1 is a short introduction. Section 2 deals with the results and formulas of the  $q$ -calculus and the calculus on generalized Fibonacci polynomials needed to develop this paper. Section 3 introduces the  $(s, t)$ -composition, the deformed  $(s, t)$ -chain rule, and its properties are given. The final section discusses the main objective of this paper, which is to solve first-order linear proportional difference equations.

## 2 Preliminaries

### 2.1 Notation for $q$ -calculus

The  $q$ -shifted factorial is

$$(a; q)_n = \begin{cases} 1, & n = 0; \\ \prod_{k=0}^{n-1} (1 - aq^k), & n \geq 1. \end{cases} \quad (4)$$

Also, define

$$(a; q)_\infty = \lim_{n \rightarrow \infty} (a; q)_n = \prod_{k=0}^{\infty} (1 - aq^k). \quad (5)$$

For a complex number  $a$  and  $q \neq 1$  the  $q$ -number  $[a]_q$  is defined by

$$[a]_q = \frac{1 - q^a}{1 - q}. \quad (6)$$

In terms of  $q$ -numbers the  $q$ -number factorial  $[n]_q!$  is defined for a nonnegative integer  $n$  by  $[n]_q! = \prod_{k=1}^n [k]_q$ . Another expression for  $[n]_q!$  is

$$[n]_q! = \prod_{k=0}^{n-1} \frac{1 - q^{k+1}}{1 - q} = \frac{(q; q)_n}{(1 - q)^n}. \quad (7)$$

The  ${}_r\phi_s$  basic hypergeometric series is define by

$${}_r\phi_s \left( \begin{matrix} a_1, a_2, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, z \right) = \sum_{n=0}^{\infty} \frac{(a_1; q)_n (a_2; q)_n \cdots (a_r; q)_n}{(q; q)_n (b_1; q)_n \cdots (b_s; q)_n} \left[ (-1)^n q^{\binom{n}{2}} \right]^{1+s-r} z^n. \quad (8)$$

The  $q$ -binomial theorem states that

$${}_1\phi_0(a; q, z) = \frac{(az; q)_\infty}{(z; q)_\infty} = \sum_{n=0}^{\infty} \frac{(a; q)_n}{(q; q)_n} z^n. \quad (9)$$

The  $q$ -exponential  $e_q(z)$  is defined by

$$\begin{aligned} e_q(z) &= \sum_{n=0}^{\infty} \frac{z^n}{[n]_q!} = \sum_{n=0}^{\infty} \frac{((1 - q)z)^n}{(q; q)_n} \\ &= {}_1\phi_0 \left( \begin{matrix} 0 \\ - \end{matrix}; q, -(1 - q)z \right) = \frac{1}{((1 - q)z; q)_\infty}. \end{aligned} \quad (10)$$

Another  $q$ -analogue of the classical exponential function is

$$\begin{aligned} E_q(z) &= \sum_{n=0}^{\infty} q^{\binom{n}{2}} \frac{z^n}{[n]_q!} = \sum_{n=0}^{\infty} q^{\binom{n}{2}} \frac{((1 - q)z)^n}{(q; q)_n} \\ &= {}_1\phi_1 \left( \begin{matrix} 0 \\ 0 \end{matrix}; q, -(1 - q)z \right) = (-(1 - q)z; q)_\infty \end{aligned} \quad (11)$$

## 2.2 Generalized Fibonacci polynomials

The generalized Fibonacci polynomials depending on the variables  $s, t$  are defined by

$$\begin{aligned}\{0\}_{s,t} &= 0, \\ \{1\}_{s,t} &= 1, \\ \{n+2\}_{s,t} &= s\{n+1\}_{s,t} + t\{n\}_{s,t}.\end{aligned}$$

As special cases, we obtain the sequences of Pell, Jacobsthal, Chebysheff of the second kind, Mersenne, and Repunits, among others. When  $s = P$  and  $t = -Q$ , with  $P$  and  $Q$  integers, we obtain the  $(P, -Q)$ -Lucas differential calculus and if  $Q = -1$ , we obtain the  $P$ -Fibonacci differential calculus. The  $(s, t)$ -Fibonacci constant is the ratio toward which adjacent  $(s, t)$ -Fibonacci polynomials tend. This is the only positive root of  $x^2 - sx - t = 0$ . We will let  $\varphi_{s,t}$  denote this constant, where

$$\varphi_{s,t} = \frac{s + \sqrt{s^2 + 4t}}{2}$$

and

$$\varphi'_{s,t} = s - \varphi_{s,t} = -\frac{t}{\varphi_{s,t}} = \frac{s - \sqrt{s^2 + 4t}}{2}.$$

The  $(s, t)$ -Fibonomial coefficients are define by

$$\binom{n}{k}_{s,t} = \frac{\{n\}_{s,t}!}{\{k\}_{s,t}! \{n-k\}_{s,t}!},$$

where  $\{n\}_{s,t}! = \{1\}_{s,t} \{2\}_{s,t} \cdots \{n\}_{s,t}$  is the  $(s, t)$ -factorial or generalized fibotorial. From Eq.(7),

$$\{n\}_{s,t}! = \varphi_{s,t}^{\binom{n}{2}} \frac{(q; q)_n}{(1-q)^n}.$$

## 2.3 Calculus on generalized Fibonacci polynomials

For all  $u \in \mathbb{C}$  and for all  $s, t \in \mathbb{R}$  such that  $s \neq 0, t \neq 0$ , define the  $(s, t)$ -derivative  $\mathbf{D}_{s,t}$  of a function  $f(x)$  as

$$(\mathbf{D}_{s,t}f)(x) = \frac{f(\varphi_{s,t}x) - f(\varphi'_{s,t}x)}{(\varphi_{s,t} - \varphi'_{s,t})x},$$

for all  $x \neq 0$  and  $(\mathbf{D}_{s,t}f)(0) = f'(0)$ , provided  $f'(0)$  exists. The equivalence that exists between the  $q$ -derivative and the  $(s, t)$ -derivative is given by the following identity:

$$(\mathbf{D}_{s,t}f)(x) = (D_qf)(\varphi_{s,t}x),$$

where

$$D_qf(x) = \frac{f(x) - f(qx)}{(1-q)x}$$

and  $q = \varphi'_{s,t}/\varphi_{s,t}$ . We will say that a function  $f(x)$  is  $q$ -periodic if  $f(y) = f(qy)$ , with  $y = \varphi_{s,t}x$ . Let  $\mathbb{P}_{s,t}$  denote the set of  $q$ -periodic functions. The  $q$ -periodic functions form

the kernel of the operator  $\mathbf{D}_{s,t}$ . Set  $s \neq 0$  and  $t < 0$ . From the condition of  $q$ -periodicity of  $f(x)$  it follows that  $f(y) = f(qy)$ , with  $q > 0$ . Then

$$\begin{aligned} f(q^y) &= f(q^{y+1}) \\ G(y) &= G(y+1) \end{aligned}$$

where  $G$  is an arbitrary periodic function with period one and  $f(x) = G(\log_q(x))$ ,  $x > 0$ . If  $t > 0$ , then  $q < 0$  and  $f(x) = G(\log(x)/(\log(-q) + i\pi))$ , so that  $x \in \mathbb{C}/\{0\}$  and  $f(x) \in \mathbb{P}_{s,t}$ .

Let  $f$  be an arbitrary function. In [7] the following  $(p, q)$ -integrals of  $f$  are defined

$$\begin{aligned} \int_0^a f(x) d_{p,q}x &= (p-q)a \sum_{k=0}^{\infty} \frac{q^k}{p^{k+1}} f\left(\frac{q^k}{p^{k+1}}a\right) \quad \text{if } \left|\frac{p}{q}\right| > 1, \\ \int_0^a f(x) d_{p,q}x &= (q-p)a \sum_{k=0}^{\infty} \frac{p^k}{q^{k+1}} f\left(\frac{p^k}{q^{k+1}}a\right) \quad \text{if } \left|\frac{p}{q}\right| < 1 \end{aligned}$$

$$\int_a^b f(x) d_{p,q}x = \int_0^b f(x) d_{p,q}x - \int_0^a f(x) d_{p,q}x$$

For  $\alpha \in \mathbb{R}$ , the set  $\alpha I \subset \mathbb{R}$  is defined by  $\alpha I = \{\alpha x : x \in I\}$ . Set  $|q|$  such that  $0 < |q| < 1$ . We will say that  $I$  is a  $(q, \varphi)$ -geometric set if it contains all geometric sequences  $\{xq^n/\varphi_{s,t}\}$ ,  $n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$ ,  $x \in I$ . For a pair of real numbers  $a$  and  $b$ , we define

$$[a, b]_q = \left\{ a \frac{q^n}{\varphi_{s,t}} : n \in \mathbb{N}_0 \right\} \cup \left\{ b \frac{q^n}{\varphi_{s,t}} : n \in \mathbb{N}_0 \right\}$$

which we will call a  $(q, \varphi)$ -interval with extreme points  $a$  and  $b$ . For real numbers  $a$  and  $b$ , the following property holds:

$$\text{If } a, b \in I \implies [a, b]_q \subset I.$$

Consider the function  $f : [a, b]_q \rightarrow \mathbb{K}$ , where  $\mathbb{K}$  is either  $\mathbb{R}$  or  $\mathbb{C}$ . The  $(s, t)$ -integral in  $[a, b]_q$  of  $f$  is

$$\int_a^b f(x) d_{s,t}x = (1-q) \sum_{n=0}^{\infty} \left[ bf\left(b \frac{q^n}{\varphi_{s,t}}\right) - af\left(a \frac{q^n}{\varphi_{s,t}}\right) \right] q^n = \int_a^b f(x/\varphi_{s,t}) d_qx. \quad (12)$$

We will call  $f$  a function  $(s, t)$ -integrable if the series in Eq.(12) is convergent.

For any real number  $p \geq 1$  and  $a, b \in I$ , we will denote by  $\mathcal{L}_{s,t}^p[a, b]$  the set of functions  $f : [a, b]_q \rightarrow \mathbb{K}$  such that  $|f|^p$  is  $(s, t)$ -integrable in  $[a, b]_q$ , i.e.,

$$\mathcal{L}_{s,t}^p[a, b] = \left\{ f : [a, b]_q \rightarrow \mathbb{K} : \int_{[a,b]_q} |f|^p d_{s,t} < \infty \right\}.$$

We also set

$$\mathcal{L}_{s,t}^\infty[a, b] = \left\{ f : [a, b]_q \rightarrow \mathbb{K} : \sup_{n \in \mathbb{N}_0} \left\{ \left| f\left(\frac{aq^n}{\varphi_{s,t}}\right) \right|, \left| f\left(\frac{bq^n}{\varphi_{s,t}}\right) \right| \right\} < \infty \right\}$$

**The fundamental theorem of  $(s, t)$ -calculus.** Let  $0 < |q| < 1$ , and  $I = [a, b]_q$  a  $(q, \varphi)$ -interval. Fix  $a, b \in I$ , and let  $f : I \rightarrow \mathbb{K}$  be a function such that  $\mathbf{D}_{s,t}f \in \mathcal{L}_{s,t}^1[a, b]$ . Then: if  $f$  is continuous at 0, then

$$\int_a^b (\mathbf{D}_{s,t}f)(x) d_{s,t}x = f(b) - f(a). \quad (13)$$

**The  $(s, t)$ -integration by parts formula.** Assume  $0 < |q| < 1$ , and let  $I = [a, b]_q$  be a  $(q, \varphi)$ -interval. Fix  $a, b \in I$ , and let  $f : I \rightarrow \mathbb{K}$  be a function such that  $\mathbf{D}_{s,t}f \in \mathcal{L}_{s,t}^1[a, b]$ . Then the equality

$$\int_I (\mathbf{D}_{s,t}f)(x) g(\varphi'_{s,t}x) d_{s,t}x = \left[ f \cdot g \right]_a^b - \int_I f(\varphi_{s,t}x) (\mathbf{D}_{s,t}g)(x) d_{s,t}x \quad (14)$$

holds, provided  $f, g \in \mathcal{L}_{s,t}^1[a, b]$ ,  $(\mathbf{D}_{s,t}f)(x)$  and  $(\mathbf{D}_{s,t}g)(x)$  are bounded for all  $x \in I$ , and the limits exists. If, in addition,  $f$  and  $g$  are continuous at 0, then

## 2.4 Deformed $(s, t)$ -exponential function

Set  $s \neq 0, t \neq 0, s, t \in \mathbb{R}$ . For all  $u \in \mathbb{C}$ , the deformed  $(s, t)$ -exponential function

$$\exp_{s,t}(z, u) = \begin{cases} \sum_{n=0}^{\infty} u \binom{n}{2} \frac{z^n}{\{n\}_{s,t}!} & \text{if } u \neq 0; \\ 1 + z & \text{if } u = 0, \end{cases}$$

is solution of the equation  $\mathbf{D}_{s,t}y = y(ux)$ . The  $(s, t)$ -antiderivative of  $\exp_{s,t}(x, u)$  is

$$\int \exp_{s,t}(x, u) d_{s,t}x = u \exp_{s,t}(x/u, u). \quad (15)$$

Also, we define

$$\begin{aligned} \text{Exp}_{s,t}(x) &= \exp_{s,t}(x, \varphi_{s,t}) \\ \text{Exp}'_{s,t}(x) &= \exp_{s,t}(x, \varphi'_{s,t}) \end{aligned}$$

and

$$\exp_{s,t}((\alpha \oplus \beta)_{\varphi, \varphi'}x) = \sum_{n=0}^{\infty} (\alpha \oplus \beta)_{\varphi, \varphi'}^n \frac{x^n}{\{n\}_{s,t}!} = \text{Exp}_{s,t}(x) \text{Exp}'_{s,t}(x) \quad (16)$$

where

$$(\alpha \oplus \beta)_{\varphi, \varphi'}^n = \begin{cases} \prod_{k=0}^{n-1} (\alpha \varphi_{s,t}^k + \beta \varphi'_{s,t}{}^k), & \text{if } n \geq 1; \\ 1, & \text{if } n = 0. \end{cases} \quad (17)$$

**Proposition 1.** For all  $\alpha, \beta \in \mathbb{C}$ . Then

$$1. \mathbf{D}_{s,t} \exp_{s,t}((\alpha \oplus \beta)_{\varphi, \varphi'}x) = \alpha \exp_{s,t}((\alpha \oplus \beta)_{\varphi, \varphi'}\varphi_{s,t}x) + \beta \exp_{s,t}((\alpha \oplus \beta)_{\varphi, \varphi'}\varphi'_{s,t}x).$$

$$2. \int \exp_{s,t}((\alpha \oplus \beta)_{\varphi, \varphi'}x) d_{s,t}x = \frac{\varphi_{s,t}\varphi'_{s,t}}{\alpha\varphi'_{s,t} + \beta\varphi_{s,t}} \exp_{s,t}(((\alpha/\varphi_{s,t}) \oplus (\beta/\varphi'_{s,t}))_{\varphi, \varphi'}x)$$

*Proof.* From Eq.(17),

$$\begin{aligned}
\mathbf{D}_{s,t} \exp_{s,t}((\alpha \oplus \beta)_{\varphi,\varphi'} x) &= \mathbf{D}_{s,t} \left( \sum_{n=0}^{\infty} (\alpha \oplus \beta)_{\varphi,\varphi'}^n \frac{x^n}{\{n\}_{s,t}!} \right) \\
&= \sum_{n=0}^{\infty} (\alpha \oplus \beta)_{\varphi,\varphi'}^{n+1} \frac{x^n}{\{n\}_{s,t}!} \\
&= \alpha \sum_{n=0}^{\infty} (\alpha \oplus \beta)_{\varphi,\varphi'}^n \frac{(\varphi_{s,t} x)^n}{\{n\}_{s,t}!} + \beta \sum_{n=0}^{\infty} (\alpha \oplus \beta)_{\varphi,\varphi'}^n \frac{(\varphi'_{s,t} x)^n}{\{n\}_{s,t}!} \\
&= \alpha \exp_{s,t}((\alpha \oplus \beta)_{\varphi,\varphi'} \varphi_{s,t} x) + \beta \exp_{s,t}((\alpha \oplus \beta)_{\varphi,\varphi'} \varphi'_{s,t} x).
\end{aligned}$$

From  $(s, t)$ -integration by parts formula Eq.(14),

$$\begin{aligned}
\int \exp_{s,t}((\alpha \oplus \beta)_{\varphi,\varphi'} x) d_{s,t} x &= \int \text{Exp}_{s,t}(\alpha x) \text{Exp}'_{s,t}(\beta x) d_{s,t} x \\
&= \frac{\varphi_{s,t}}{\alpha} \int \text{Exp}'_{s,t}(\beta(\varphi'_{s,t} x)/\varphi_{s,t}) \mathbf{D}_{s,t} \text{Exp}_{s,t}(\alpha x/\varphi_{s,t}) d_{s,t} x \\
&= \frac{\varphi_{s,t}}{\alpha} \left[ \text{Exp}_{s,t}(\beta x/\varphi'_{s,t}) \text{Exp}_{s,t}(\alpha x/\varphi_{s,t}) \right. \\
&\quad \left. - \frac{\beta}{\varphi'_{s,t}} \int \text{Exp}_{s,t}(\alpha x) \text{Exp}'_{s,t}(\beta x) d_{s,t} x \right].
\end{aligned}$$

Then

$$\left( 1 + \frac{\varphi_{s,t} \beta}{\varphi'_{s,t} \alpha} \right) \int \text{Exp}_{s,t}(\alpha x) \text{Exp}'_{s,t}(\beta x) d_{s,t} x = \frac{\varphi_{s,t}}{\alpha} \text{Exp}'_{s,t}(\beta x/\varphi'_{s,t}) \text{Exp}_{s,t}(\alpha x/\varphi_{s,t})$$

and

$$\begin{aligned}
\int \text{Exp}_{s,t}(\alpha x) \text{Exp}'_{s,t}(\beta x) d_{s,t} x &= \frac{\varphi_{s,t} \varphi'_{s,t}}{\alpha \varphi'_{s,t} + \beta \varphi_{s,t}} \text{Exp}_{s,t}(\alpha x/\varphi_{s,t}) \text{Exp}'_{s,t}(\beta x/\varphi'_{s,t}) \\
&= \frac{\varphi_{s,t} \varphi'_{s,t}}{\alpha \varphi'_{s,t} + \beta \varphi_{s,t}} \exp_{s,t}(((\alpha/\varphi_{s,t}) \oplus (\beta/\varphi'_{s,t}))_{\varphi,\varphi'} x).
\end{aligned}$$

The proof is completed. □

## 2.5 The $(s, t)$ -pantograph function

The  $(1, u)$ -deformed  $(s, t)$ -pantograph function is defined by

$$\mathbf{E}_{s,t}(a, b; z, u) = \sum_{n=0}^{\infty} (a \oplus b)_{1,u}^n \frac{z^n}{\{n\}_{s,t}!}, \tag{18}$$

and this is solution of the equation  $\mathbf{D}_{s,t} y = ay(x) + by(ux)$ , where

$$(a \oplus b)_{1,u}^n = \begin{cases} \prod_{k=0}^{n-1} (a + bu^k), & \text{if } n \neq 0; \\ 1, & \text{if } n = 0. \end{cases} \tag{19}$$

For  $a \neq 0$ , it is very easy to notice that  $(a \oplus b)_{1,u}^n = a^{\binom{n}{2}} (b/a; u)_n$ . Then

$$\mathbf{E}_{s,t}(a, b; x, u) = \sum_{n=0}^{\infty} \left( \frac{a}{\varphi_{s,t}} \right)^{\binom{n}{2}} \frac{(b/a; u)_n}{(q; q)_n} ((1-q)x)^n \tag{20}$$

is a  $q$ -series. Some special values of the function  $E_{s,t}(a, b, u; x)$  are

$$\begin{aligned}
E_{s,t}(a, -a; x, u) &= 0, \\
E_{s,t}(a, 0; x, -) &= \exp_{s,t}(ax), \\
E_{s,t}(0, a; x, u) &= \exp_{s,t}(ax, u), \\
E_{s,t}(a, b; x, 1) &= \exp_{s,t}((a+b)x), \\
E_{s,t}(ac, bc; x, u) &= E_{s,t}(a, b; cx, u), \\
E_{s,t}(\varphi_{s,t}, -b; -x, q) &= {}_1\phi_0 \left( \begin{matrix} b/\varphi_{s,t} \\ - \end{matrix}; q, (1-q)x \right) = \frac{((b/\varphi_{s,t})(1-q)x; q)_\infty}{((1-q)x; q)_\infty}, \\
E_{s,t}(1, -q; x, q) &= \Theta_0((1-q)x, \varphi_{s,t}^{-1}),
\end{aligned}$$

where  $q = \varphi'_{s,t}/\varphi_{s,t}$  and

$$\Theta_0(x, y) = \sum_{n=0}^{\infty} y^{\binom{n}{2}} x^n \quad (21)$$

is the Partial Theta function [?]. From here, is defined the function

$$\psi(q) = \Theta_0(q, q) = \sum_{n=0}^{\infty} q^{\binom{n+1}{2}} = \frac{(q^2; q^2)_\infty}{(q; q^2)_\infty}.$$

Also,

$$E_{s,t}(\varphi'_{s,t}, -b; -x, q) = \sum_{n=0}^{\infty} (-1)^n q^{\binom{n}{2}} \frac{(b/\varphi'_{s,t}; q)_n}{(q; q)_n} (1-q)^n x^n, \quad (22)$$

and

$$E_{s,t}(\varphi'_{s,t}, -\varphi'_{s,t}q; -x, q) = \sum_{n=0}^{\infty} (-1)^n q^{\binom{n}{2}} (1-q)^n x^n, \quad (23)$$

**Theorem 1.** *Set  $a, b, u \in \mathbb{C}$  such that  $a \neq 0$  and  $|b/au| < 1$ . Then*

$$\int E_{s,t}(a, b; x, u) d_{s,t}x = \frac{1}{a} \sum_{k=0}^{\infty} (-1)^k \left( \frac{b}{au} \right)^k E_{s,t}(a, b; u^k x, u). \quad (24)$$

*Proof.* From Eq.(18)

$$\begin{aligned}
\int E_{s,t}(a, b; x, u) d_{s,t}x &= \sum_{n=0}^{\infty} (a \oplus b)_{1,u}^n \frac{x^{n+1}}{\{n+1\}_{s,t}!} \\
&= \sum_{n=1}^{\infty} \frac{(a \oplus b)_{1,u}^n}{a + bu^{n-1}} \frac{x^n}{\{n\}_{s,t}!} \\
&= \frac{1}{a} \sum_{n=1}^{\infty} (a \oplus b)_{1,u}^n \frac{x^n}{\{n\}_{s,t}!} \sum_{k=0}^{\infty} (-1)^k \left( \frac{b}{au} \right)^k u^{kn} \\
&= \frac{1}{a} \sum_{k=0}^{\infty} (-1)^k \left( \frac{b}{au} \right)^k \sum_{n=1}^{\infty} (a \oplus b)_{1,u}^n \frac{(u^k x)^n}{\{n\}_{s,t}!} \\
&= \frac{1}{a} \sum_{k=0}^{\infty} (-1)^k \left( \frac{b}{au} \right)^k (E_{s,t}(a, b; u^k x, u) - 1) \\
&= \frac{1}{a} \sum_{k=0}^{\infty} (-1)^k \left( \frac{b}{au} \right)^k E_{s,t}(a, b; u^k x, u) - \frac{u}{au + b}.
\end{aligned}$$



As  $\mathbf{D}_{s,t}\left(\frac{u}{au+b}\right) = 0$ , then  $\frac{1}{a} \sum_{k=0}^{\infty} (-1)^k \left(\frac{b}{au}\right)^k \mathbf{E}_{s,t}(a, b; u^k x, u)$  is the  $(s, t)$ -antiderivative of  $\mathbf{E}_{s,t}(a, b; x, u)$ .  $\square$

**Corollary 1.** *If  $|q| < 1$ , then*

$$\int \Theta_0((1-q)x, \varphi_{s,t}^{-1}) d_{s,t}x = \sum_{k=0}^{\infty} (\Theta_0((1-q)q^k x, \varphi_{s,t}^{-1}) - 1). \quad (25)$$

### 3 Deformed $(s, t)$ -chain rule

This section is based on Gessel's paper [9], slightly modified by Johnson [10], on  $q$ -function composition and  $q$ -chain rule. We will define the deformed  $(s, t)$ -composition of functions and give the  $(s, t)$ -chain rule.

Consider a function  $f(x)$  of the form

$$f(x) = \sum_{n=1}^{\infty} f_n \frac{x^n}{[n]_q!}.$$

Then for  $k \geq 0$ , the  $k$ th symbolic power  $f^{[k]}(x)$  of  $f$  is defined by  $f^{[0]}(x) = 1$  and for  $k \geq 1$  define the  $k$ th symbolic power of  $f$  inductively by

$$D_q f^{[k]}(x) = [k]_q f^{[k-1]}(x) D_q f(x), \quad \text{with } f^{[k]}(0) = 0.$$

When  $k = 1$ ,  $\mathbf{D}_q f^{[1]}(x) = \mathbf{D}_q f(x)$ . Then  $f^{[1]}(x) = f(x)$  because their derivatives coincide and because both are zero at  $x = 0$ . Note also that  $\mathbf{D}_q x^{[k]} = [k]_q x^{[k-1]}$  and by induction it is shown that  $x^{[k]} = x^k$ , because  $x^{[1]} = x$ . If  $r$  does not depend on  $x$ , then

$$(t f(x))^{[k]} = t^k f^{[k]}(x). \quad (26)$$

For exchange  $D_q$  by  $\mathbf{D}_{s,t}$  in the definition of Gessel of  $q$ -chain rule, we obtain the  $(s, t)$ -chain rule of  $f^{[n]}(x)$

$$\mathbf{D}_{s,t} f^{[n]}(x) = \{n\}_{s,t} f^{[n-1]}(x) \mathbf{D}_{s,t} f(x), \quad \text{with } f^{[k]}(0) = 0 \quad (27)$$

when  $f(x)$  is defined by

$$f(x) = \sum_{n=1}^{\infty} f_n \frac{x^n}{\{n\}_{s,t}!}.$$

Next, we will define the deformed  $(s, t)$ -composition of functions.

**Definition 1.** *Suppose a function  $g_{s,t}(x, u)$  defined by*

$$g_{s,t}(x, u) = \sum_{n=0}^{\infty} u^{\binom{n}{2}} g_n \frac{x^n}{\{n\}_{s,t}!}$$

*Then the  $u$ -deformed  $(s, t)$ -composition  $g_{s,t}[f, u]$  is defined to be*

$$g_{s,t}(x, u) \square f(x) = g_{s,t}[f, u] = \sum_{n=0}^{\infty} u^{\binom{n}{2}} g_n \frac{f^{[n]}(x)}{\{n\}_{s,t}!}. \quad (28)$$

More generally,

**Definition 2.** Suppose a function  $g_{s,t}(a, b; x, u)$  defined by

$$g_{s,t}(a, b; x, u) = \sum_{n=0}^{\infty} (a \oplus b)_{1,u}^n g_n \frac{x^n}{\{n\}_{s,t}!}$$

Then the  $(1, u)$ -deformed  $(s, t)$ -composition  $g_{s,t}[a, b; f, u]$  is defined to be

$$g_{s,t}(a, b; x, u) \square f(x) = g_{s,t}[a, b; f(x), u] = \sum_{n=0}^{\infty} (a \oplus b)_{1,u}^n g_n \frac{f^{[n]}(x)}{\{n\}_{s,t}!}. \quad (29)$$

When  $a = 0$  and  $b = 1$ , then

$$g_{s,t}[0, 1; f(x), u] = g_{s,t}[f(x), u].$$

Then it is easy to obtain the  $(s, t)$ -chain rule for  $(1, u)$ -deformed functions.

**Proposition 2.** Set  $a, b \in \mathbb{R}$  such that  $a \neq -b$ . Then

$$\begin{aligned} \mathbf{D}_{s,t}(g_{s,t}[a, b; f(x), u]) &= \left( a(\mathbf{D}_{s,t}g_{s,t})[a, b; f(x), u] \right. \\ &\quad \left. + b(\mathbf{D}_{s,t}g_{s,t})[a, b; uf(x), u] \right) (\mathbf{D}_{s,t}f)(x). \end{aligned} \quad (30)$$

*Proof.* From Eq.(27), we have that

$$\begin{aligned} \mathbf{D}_{s,t}(g_{s,t}[a, b; f(x), u]) &= \mathbf{D}_{s,t} \left( \sum_{n=0}^{\infty} (a \oplus b)_{1,u}^n g_n \frac{f^{[n]}(x)}{\{n\}_{s,t}!} \right) \\ &= \sum_{n=1}^{\infty} (a \oplus b)_{1,u}^n g_n \frac{\{n\}_{s,t} f^{[n-1]}(x)}{\{n\}_{s,t}!} \mathbf{D}_{s,t}f(x) \\ &= \sum_{n=0}^{\infty} (a \oplus b)_{1,u}^{n+1} g_{n+1} \frac{f^{[n]}(x)}{\{n\}_{s,t}!} \mathbf{D}_{s,t}f(x) \\ &= \left( a(\mathbf{D}_{s,t}g_{s,t})[a, b; f(x), u] \right. \\ &\quad \left. + b(\mathbf{D}_{s,t}g_{s,t})[a, b; uf(x), u] \right) \mathbf{D}_{s,t}f(x) \end{aligned}$$

and the proof is reached. □

Immediately, we have the following Corollary.

**Corollary 2.**  $\mathbf{D}_{s,t}g_{s,t}[f(x), u] = (\mathbf{D}_{s,t}g_{s,t})[uf(x), u] \mathbf{D}_{s,t}f(x)$ .

*Proof.* Set  $a = 0$  and  $b = 1$  in the above Proposition. □

**Definition 3.** For  $n \in \mathbb{N}$ , define

$$\int_{f(r)}^{g(r)} x^n d_{s,t}x \equiv \left( \int x^n d_{s,t}x \right) \Big|_{f(r)}^{g(r)} = \frac{x^{n+1} \square g(r) - x^{n+1} \square f(r)}{\{n+1\}_{s,t}}. \quad (31)$$

We formally state that for all  $n, m \in \mathbb{N}$ ,

1.

$$\int_{f(r)}^{f(g(r))} (\alpha x^n + \beta x^m) d_{s,t}x = \alpha \int_{f(r)}^{f(g(r))} x^n d_{s,t}x + \beta \int_{f(r)}^{f(g(r))} x^m d_{s,t}x.$$

2.

$$\int_{f(r)}^{f(g(r))} \sum_{n=0}^{\infty} a_n x^n d_{s,t}x = \sum_{n=0}^{\infty} a_n \int_{f(r)}^{f(g(r))} x^n d_{s,t}x.$$

**Theorem 2 (Substitution formula).** For all  $n \in \mathbb{N}$ ,

1.

$$\int_a^b f^{[n]}(x) \mathbf{D}_{s,t}f(x) d_{s,t}x = \int_{f(a)}^{f(b)} x^n d_{s,t}x. \quad (32)$$

2.

$$\int_a^b f[g(x), u] \mathbf{D}_{s,t}g(x) d_{s,t}x = \int_{g(a)}^{g(b)} f(x, u) d_{s,t}x. \quad (33)$$

*Proof.* From Definition 3

$$\begin{aligned} \int_a^b f^{[n]}(x) \mathbf{D}_{s,t}f(x) d_{s,t}x &= \frac{1}{\{n+1\}_{s,t}} \int_a^b \mathbf{D}_{s,t}f^{[n+1]}(x) d_{s,t}x \\ &= \frac{f^{[n+1]}(b) - f^{[n+1]}(a)}{\{n+1\}_{s,t}} = \int_{f(a)}^{f(b)} x^n d_{s,t}x. \end{aligned}$$

By using Eq.(32), we have formally

$$\begin{aligned} \int_a^b f[g(x), u] \mathbf{D}_{s,t}g(x) d_{s,t}x &= \int_a^b \left( \sum_{m=0}^{\infty} u^{(m)} f_m \frac{g^{[m]}(x)}{\{m\}_{s,t}!} \mathbf{D}_{s,t}g(x) \right) d_{s,t}x \\ &= \sum_{m=0}^{\infty} \frac{u^{(m)}}{\{m\}_{s,t}!} f_m \int_a^b g^{[m]}(x) \mathbf{D}_{s,t}g(x) d_{s,t}x \\ &= \sum_{m=0}^{\infty} \frac{u^{(m)}}{\{m\}_{s,t}!} f_m \int_{g(a)}^{g(b)} x^m d_{s,t}x \\ &= \int_{g(a)}^{g(b)} \sum_{m=0}^{\infty} u^{(m)} f_m \frac{x^m}{\{m\}_{s,t}!} d_{s,t}x \\ &= \int_{g(a)}^{g(b)} f(x, u) d_{s,t}x. \end{aligned}$$

□

From Eq.(28) we have the following composition

$$\exp_{s,t}[f(x), u] = \sum_{n=0}^{\infty} u^{(n)} \frac{f^{[n]}(x)}{\{n\}_{s,t}!}.$$

Also, denote

$$\begin{aligned} \text{Exp}_{s,t}[f(x)] &= \exp_{s,t}[f(x), \varphi_{s,t}], \\ \text{Exp}'_{s,t}[f(x)] &= \exp_{s,t}[f(x), \varphi'_{s,t}] \end{aligned}$$

Then,  $\text{Exp}_{s,t}[x] = \text{Exp}_{s,t}(x)$  and  $\text{Exp}'_{s,t}[x] = \text{Exp}'_{s,t}(x)$ . Also

$$\mathbf{D}_{s,t} \text{Exp}_{s,t}[f(x)] = \text{Exp}_{s,t}[\varphi_{s,t}f(x)]\mathbf{D}_{s,t}f(x), \quad (34)$$

$$\mathbf{D}_{s,t} \text{Exp}'_{s,t}[f(x)] = \text{Exp}'_{s,t}[\varphi'_{s,t}f(x)]\mathbf{D}_{s,t}f(x) \quad (35)$$

and in general

$$\mathbf{D}_{s,t} \exp_{s,t}[f(x), u] = \exp_{s,t}[uf(x), u]\mathbf{D}_{s,t}f(x),$$

$$\mathbf{D}_{s,t} \mathbf{E}_{s,t}[a, b; f(x), u] = \left( a\mathbf{E}_{s,t}[a, b; g(x), u] + b\mathbf{E}_{s,t}[a, b; ug(x), u] \right) \mathbf{D}_{s,t}f(x).$$

**Example 1.** *On the one hand,*

$$\begin{aligned} \{n\}_{s,t} \int_0^x r^{n-1} \exp_{s,t}[ur^n, u] d_{s,t}r &= \int_0^x \mathbf{D}_{s,t} \exp_{s,t}[r^n, u] d_{s,t}r \\ &= \exp_{s,t}[x^n, u] - 1 \end{aligned}$$

*and on the other hand,*

$$\begin{aligned} \{n\}_{s,t} \int_0^x r^{n-1} \exp_{s,t}[ur^n, u] d_{s,t}r &= \int_0^{x^n} \exp_{s,t}(ur, u) d_{s,t} \\ &= \exp_{s,t}[x^n, u] - 1. \end{aligned}$$

## 4 First order functional-difference $(s, t)$ -equation

### 4.1 An equation general

**Theorem 3.** *Set  $a \neq -b$  and  $q = \varphi'_{s,t}/\varphi_{s,t}$ . Assume  $0 < |q| < 1$  and suppose  $\alpha$  and  $\beta$  are continuous functions on an interval  $[\eta, x]$ ,  $\eta > 0$ . Set  $A(x) = \int_{\eta}^x \alpha(r) d_q r$ . Then the function  $y(x)$  given by*

$$y(x) = \frac{1}{\mathbf{E}_{s,t}[a, b; A(x), u]} \left( \int_{\eta}^x \beta(r) \mathbf{E}_{s,t}[a, b; A(\varphi_{s,t}r), u] d_{s,t}r + G(\log_q(x)) \right) \quad (36)$$

*is a solution to the equation*

$$\mathbf{D}_{s,t}y + \alpha(x) \frac{(\mathbf{D}_{s,t} \mathbf{E}_{s,t})(a, b; x, u) \square A(x)}{\mathbf{E}_{s,t}[a, b; A(\varphi_{s,t}x), u]} y(\varphi'_{s,t}x) = \beta(x), \quad (37)$$

*where*

$$(\mathbf{D}_{s,t} \mathbf{E}_{s,t})(a, b; x, u) \square A(x) = a\mathbf{E}_{s,t}[a, b; A(x), u] + b\mathbf{E}_{s,t}[a, b; uA(x), u],$$

*with initial value problem  $y(\eta) = G(\log_q(\eta))$  for some  $G(\log_q(x)) \in \mathbb{P}_{s,t}$ . If  $\eta \geq 0$ , then the function  $y(x)$  given by*

$$y(x) = \frac{1}{\mathbf{E}_{s,t}[a, b; A(x), u]} \left( \int_{\eta}^x \beta(r) \mathbf{E}_{s,t}[a, b; A(\varphi_{s,t}r), u] d_{s,t}r \right) \quad (38)$$

*is a solution to the Eq.(37) with initial value problem  $y(\eta) = \xi$ , for  $\eta \in \mathbb{R}$ .*

*Proof.* By multiplying Eq.(37) by  $E_{s,t}[a, b; A(\varphi_{s,t}x), u]$ , we obtain

$$\begin{aligned} E_{s,t}[a, b; A(\varphi_{s,t}x), u] \mathbf{D}_{s,t}y + \alpha(x) (aE_{s,t}[a, b; A(x), u] + bE_{s,t}[a, b; uA(x), u]) y(\varphi'_{s,t}x) \\ = \beta(x)E_{s,t}[a, b; A(\varphi_{s,t}x), u]. \end{aligned}$$

As

$$\begin{aligned} \mathbf{D}_{s,t} (E_{s,t}[a, b; A(x), u]y) (x) = E_{s,t}[a, b; A(\varphi_{s,t}x), u] \mathbf{D}_{s,t}y(x) \\ + \alpha(x) (aE_{s,t}[a, b; A(x), u] + bE_{s,t}[a, b; uA(x), u]) y(\varphi'_{s,t}x), \end{aligned}$$

then

$$\mathbf{D}_{s,t} (E_{s,t}[a, b; A(x), u]y) = \beta(x)E_{s,t}[a, b; A(\varphi_{s,t}x), u].$$

From Theorem 13

$$\begin{aligned} \int_{\eta}^x \mathbf{D}_{s,t} (E_{s,t}[a, b; A(r), u]y) d_{s,t}r = E_{s,t}[a, b; A(x), u]y(x) - G(\log_q(\eta)) \\ = \int_{\eta}^x \beta(r)E_{s,t}[a, b; A(\varphi_{s,t}r), u]d_{s,t}r \end{aligned}$$

and from this, it follows that

$$y(x) = \frac{G(\log_q(\eta))}{E_{s,t}[a, b; A(x), u]} + \frac{1}{E_{s,t}[a, b; A(x), u]} \int_{\eta}^x \beta(r)E_{s,t}[a, b; A(\varphi_{s,t}r), u]d_{s,t}r$$

is solution of Eq.(37) if  $\eta > 0$ . By changing  $G(\log_q(x))$  by  $G(\log_q(\eta))$  above, then we obtain a more general solution. If  $\eta \geq 0$ , then the solution of Eq.(38) is

$$y(x) = \frac{\xi}{E_{s,t}[a, b; A(x), u]} + \frac{1}{E_{s,t}[a, b; A(x), u]} \int_{\eta}^x \beta(r)E_{s,t}[a, b; A(\varphi_{s,t}r), u]d_{s,t}r$$

and the proof is reached.  $\square$

By interchanging  $\varphi_{s,t}$  and  $\varphi'_{s,t}$  in the above theorem, we obtain the solution to the equation

$$\mathbf{D}_{s,t}y + \alpha(x) \frac{(\mathbf{D}_{s,t}E_{s,t})(a, b; x, u) \square A(x)}{E_{s,t}[a, b; A(\varphi'_{s,t}x), u]} y(\varphi_{s,t}x) = \beta(x). \quad (39)$$

#### 4.1.1 Equation $\mathbf{D}_{s,t}y(x) + \alpha(x) \frac{\exp_{s,t}[uA(x), u]}{\exp_{s,t}[A(\varphi_{s,t}x), u]} y(\varphi'_{s,t}x) = \beta(x)$

**Corollary 3.** Set  $q = \varphi'_{s,t}/\varphi_{s,t}$ . Assume  $0 < |q| < 1$  and suppose  $\alpha$  and  $\beta$  are continuous functions on an interval  $[\eta, x]$ ,  $\eta > 0$ . Set  $A(x) = \int_{\eta}^x \alpha(r)d_qr$ . Then the function  $y(x)$  given by

$$y(x) = \frac{1}{\exp_{s,t}[A(x), u]} \left( \int_{\eta}^x \beta(r) \exp_{s,t}[A(\varphi_{s,t}r), u]d_{s,t}r + G(\log_q(x)) \right) \quad (40)$$

is a solution to the initial value problem

$$\mathbf{D}_{s,t}y + \alpha(x) \frac{\exp_{s,t}[uA(x), u]}{\exp_{s,t}[A(\varphi_{s,t}x), u]} y(\varphi'_{s,t}x) = \beta(x), \quad y(\eta) = G(\log_q(\eta)) \quad (41)$$

for some  $G(\log_q(x)) \in \mathbb{P}_{s,t}$ . If  $\eta \geq 0$ , then the function  $y(x)$  given by

$$y(x) = \frac{1}{\exp_{s,t}[A(x), u]} \left( \int_{\eta}^x \beta(r) \exp_{s,t}[A(\varphi_{s,t}r), u] d_{s,t}r + \xi \right), \quad (42)$$

is a solution to the initial value problem

$$\mathbf{D}_{s,t}y + \alpha(x) \frac{\exp_{s,t}[uA(x), u]}{\exp_{s,t}[A(\varphi_{s,t}x), u]} y(\varphi'_{s,t}x) = \beta(x), \quad y(\eta) = \xi \quad (43)$$

for  $\eta \in \mathbb{R}$ .

**Example 2.** Let us now take into account the equation

$$\mathbf{D}_{s,t}y(x) + x^{n-1} \frac{\exp_{s,t}[ux^n, u]}{\exp_{s,t}[\varphi_{s,t}^n x^n, u]} y(\varphi'_{s,t}x) = x^{n-1}, \quad y(0) = \xi$$

where  $n \neq 0$ . From Eq.(40),

$$\begin{aligned} y(x) &= \frac{1}{\exp_{s,t}[x^n, u]} \left( \int_0^x r^{n-1} \exp_{s,t}[ur^n, u] d_{s,t}r + \xi \right) \\ &= \frac{1}{\exp_{s,t}[x^n, u]} \left( \frac{\exp_{s,t}[x^n, u] - 1}{\{n\}_{s,t}} + \xi \right) \\ &= \frac{\exp_{s,t}[x^n, u] - 1}{\{n\}_{s,t} \exp_{s,t}[x^n, u]} + \frac{\xi}{\exp_{s,t}[x^n, u]}. \end{aligned}$$

#### 4.1.2 Solution of the equation $\mathbf{D}_{s,t}y + \alpha(x)y(\varphi'_{s,t}x) = \beta(x)$

**Corollary 4.** Set  $q = \varphi'_{s,t}/\varphi_{s,t}$ . Assume  $0 < |q| < 1$  and suppose  $\alpha$  and  $\beta$  are continuous functions on an interval  $[\eta, x]$ ,  $\eta > 0$ . Set  $A(x) = \int_{\eta}^x \alpha(r) d_q r$ . Then the function  $y(x)$  given by

$$y(x) = \frac{1}{\text{Exp}_{s,t}[A(x)]} \left( \int_{\eta}^x \beta(r) \text{Exp}_{s,t}[A(\varphi_{s,t}r)] d_{s,t}r + G(\log_q(x)) \right) \quad (44)$$

is a solution to the initial value problem

$$\mathbf{D}_{s,t}y + \alpha(x)y(\varphi'_{s,t}x) = \beta(x), \quad y(\eta) = G(\log_q(\eta)) \quad (45)$$

for some  $G(\log_q(x)) \in \mathbb{P}_{s,t}$ . If  $\eta \geq 0$ , then the function  $y(x)$  given by

$$y(x) = \frac{1}{\text{Exp}_{s,t}[A(x)]} \left( \int_{\eta}^x \beta(r) \text{Exp}_{s,t}[A(\varphi_{s,t}r)] d_{s,t}r + \xi \right), \quad (46)$$

is a solution to the initial value problem

$$\mathbf{D}_{s,t}y + \alpha(x)y(\varphi'_{s,t}x) = \beta(x), \quad y(\eta) = \xi \quad (47)$$

for  $\eta \in \mathbb{R}$ .

**Example 3.** Let us consider the equation

$$\mathbf{D}_{s,t}y(x) - y(\varphi'_{s,t}x) = x^m$$

where  $m \geq 0$ . From Eq.(44)

$$\begin{aligned} y(x) &= \frac{1}{\text{Exp}_{s,t}[-x]} \left( \int r^m \text{Exp}_{s,t}[-\varphi_{s,t}r] d_{s,t}r + G(\log_q(x)) \right) \\ &= \frac{1}{\text{Exp}_{s,t}(-x)} \left( \int r^m \text{Exp}_{s,t}(-\varphi_{s,t}r) d_{s,t}r + G(\log_q(x)) \right) \end{aligned}$$

and from Eq.(14)

$$\begin{aligned} \int x^m \text{Exp}_{s,t}(-\varphi_{s,t}x) d_{s,t}x &= -\varphi'_{s,t}{}^{(-m)} \int (\varphi'_{s,t}x)^m \mathbf{D}_{s,t} \text{Exp}_{s,t}(-x) d_{s,t}x \\ &= -\varphi'_{s,t}{}^{(-m)} x^m \text{Exp}_{s,t}(-x) \\ &\quad + \varphi'_{s,t}{}^{(-m)} \{m\}_{s,t} \int x^{m-1} \text{Exp}_{s,t}(-\varphi_{s,t}x) d_{s,t}x. \end{aligned}$$

Then

$$\int x^m \text{Exp}_{s,t}(-\varphi_{s,t}x) d_{s,t}x = - \left( \sum_{k=0}^m \varphi'_{s,t}{}^{(-m(k+1)+\binom{k+1}{2})} \frac{\{m\}_{s,t}!}{\{m-k\}_{s,t}!} x^{m-k} \right) \text{Exp}_{s,t}(-x)$$

and

$$y(x) = G(\log_q(x)) \text{Exp}'_{s,t}(x) - \sum_{k=0}^m \varphi'_{s,t}{}^{(-m(k+1)+\binom{k+1}{2})} \frac{\{m\}_{s,t}!}{\{m-k\}_{s,t}!} x^{m-k}. \quad (48)$$

**Example 4.** Let us consider the equation

$$\mathbf{D}_{s,t}y(x) + y(\varphi_{s,t}x) = \alpha \text{Exp}'_{s,t}(\beta x),$$

where  $\beta \neq -\varphi'_{s,t}$ . From Eq.(44)

$$\begin{aligned} y(x) &= \frac{1}{\text{Exp}_{s,t}(x)} \left( \alpha \int \text{Exp}'_{s,t}(\beta x) \text{Exp}_{s,t}(\varphi_{s,t}x) d_{s,t}x + G(\log_q(x)) \right) \\ &= \frac{1}{\text{Exp}_{s,t}(x)} \left( \alpha \int \exp_{s,t}((\varphi_{s,t} \oplus \beta)_{\varphi, \varphi'}x) d_{s,t}x + G(\log_q(x)) \right) \\ &= \frac{\alpha \varphi'_{s,t}}{\varphi'_{s,t} + \beta} \text{Exp}'_{s,t}(\beta x / \varphi'_{s,t}) + \frac{G(\log_q(x))}{\text{Exp}_{s,t}(x)}. \end{aligned}$$

**Example 5.** Let us consider the equation

$$\mathbf{D}_{s,t}y(x) + y(\varphi_{s,t}x) = \alpha x^m \text{Exp}'_{s,t}(-\varphi_{s,t}x)$$

From Eq.(44),

$$\begin{aligned} y(x) &= \frac{1}{\text{Exp}_{s,t}(x)} \left( \alpha \int x^m \text{Exp}'_{s,t}(-\varphi_{s,t}x) \text{Exp}_{s,t}(\varphi_{s,t}x) d_{s,t}x + G(\log_q(x)) \right) \\ &= \frac{1}{\text{Exp}_{s,t}(x)} \left( \alpha \int x^m d_{s,t}x + G(\log_q(x)) \right) \\ &= \frac{\alpha}{\{m+1\}_{s,t}} \frac{x^{m+1}}{\text{Exp}_{s,t}(x)} + \frac{G(\log_q(x))}{\text{Exp}_{s,t}(x)} \end{aligned}$$

### 4.1.3 Equation involving partial theta function

**Corollary 5.** Set  $q = \varphi'_{s,t}/\varphi_{s,t}$ . Assume  $0 < |q| < 1$  and suppose  $\alpha$  and  $\beta$  are continuous functions on an interval  $[\eta, x]$ ,  $\eta > 0$ . Set  $A(x) = \int_{\eta}^x \alpha(r) d_q r$ . Then the function  $y(x)$  given by

$$y(x) = \frac{1}{\Theta_0[(1-q)A(x), \varphi_{s,t}^{-1}]} \left( \int_{\eta}^x \beta(r) \Theta_0[(1-q)A(\varphi_{s,t} r), \varphi_{s,t}^{-1}] d_{s,t} r + G(\log_q(x)) \right) \quad (49)$$

is a solution to the initial value problem

$$\mathbf{D}_{s,t} y(x) + \alpha(\varphi_{s,t} x) \frac{(D_q \Theta_0((1-q)x, \varphi_{s,t}^{-1})) \square A(\varphi_{s,t} x)}{\Theta_0[(1-q)A(\varphi_{s,t} x), \varphi_{s,t}^{-1}]} y(\varphi'_{s,t} x) = \beta(x), \quad (50)$$

$y(\eta) = G(\log_q(\eta))$ , for some  $G(\log_q(x)) \in \mathbb{P}_{s,t}$ . If  $\eta \geq 0$ , then the function  $y(x)$  given by

$$y(x) = \frac{1}{\Theta_0[(1-q)A(x), \varphi_{s,t}^{-1}]} \left( \int_{\eta}^x \beta(r) \Theta_0[(1-q)A(\varphi_{s,t} r), \varphi_{s,t}^{-1}] d_{s,t} r + \xi \right) \quad (51)$$

is a solution to the initial value problem

$$\mathbf{D}_{s,t} y + \alpha(\varphi_{s,t} x) \frac{(D_q \Theta_0((1-q)x, \varphi_{s,t}^{-1})) \square A(\varphi_{s,t} x)}{\Theta_0[(1-q)A(\varphi_{s,t} x), \varphi_{s,t}^{-1}]} y(\varphi'_{s,t} x) = \beta(x), \quad y(\eta) = \xi \quad (52)$$

for  $\eta \in \mathbb{R}$ .

**Example 6.** The initial value problem

$$\mathbf{D}_{s,t} y(x) + \frac{\mathbf{D}_{s,t} \Theta_0((1-q)x, \varphi_{s,t}^{-1})}{\Theta_0((1-q)\varphi_{s,t} x, \varphi_{s,t}^{-1})} y(\varphi'_{s,t} x) = 1, \quad y(0) = \xi$$

has solution

$$\begin{aligned} y(x) &= \frac{1}{\Theta_0((1-q)x, \varphi_{s,t}^{-1})} \int_0^x \Theta_0[(1-q)\varphi_{s,t} r, \varphi_{s,t}^{-1}] d_{s,t} r + \frac{\xi}{\Theta_0((1-q)x, \varphi_{s,t}^{-1})} \\ &= \frac{1}{\varphi_{s,t} \Theta_0((1-q)x, \varphi_{s,t}^{-1})} \sum_{k=0}^{\infty} (\Theta_0((1-q)q^k \varphi_{s,t} x, \varphi_{s,t}^{-1}) - 1) + \frac{\xi}{\Theta_0((1-q)x, \varphi_{s,t}^{-1})}. \end{aligned}$$

If  $x = \frac{\varphi_{s,t}^{-1}}{1-q}$ , then

$$y \left( \frac{\varphi_{s,t}^{-1}}{1-q} \right) = \frac{(\varphi_{s,t}^{-1}; \varphi_{s,t}^{-2})_{\infty}}{(\varphi_{s,t}^{-2}; \varphi_{s,t}^{-2})_{\infty}} \sum_{k=0}^{\infty} (\Theta_0(q^k, \varphi_{s,t}^{-1}) - 1) + \frac{(\varphi_{s,t}^{-1}; \varphi_{s,t}^{-2})_{\infty}}{(\varphi_{s,t}^{-2}; \varphi_{s,t}^{-2})_{\infty}} \xi.$$

## 4.2 Equation $\mathbf{D}_{s,t} y = \alpha(ay(x) + by(ux)) + \beta(x)$

**Theorem 4.** Consider the following equation

$$\mathbf{D}_{s,t} y(x) = \alpha(ay(x) + by(ux)) + \beta(x), \quad y(0) = a_0, \quad (53)$$

where  $\alpha$  is a constant, and  $\beta(x)$  is an analytic function with expansion

$$\beta(x) = \sum_{n=0}^{\infty} b_n x^n.$$



Then its solution is given by

$$y(x) = a_0 E_{s,t}(a, b; \alpha x, u) + \sum_{n=1}^{\infty} \left( \sum_{k=0}^{n-1} \frac{\{k\}_{s,t}! \alpha^{n-k-1}}{(a \oplus b)_{1,u}^{k+1}} b_k \right) (a \oplus b)_{1,u}^n \frac{x^n}{\{n\}_{s,t}!}. \quad (54)$$

*Proof.* Assuming that the power series expansion of  $y(x)$  is

$$y(x) = \sum_{n=0}^{\infty} a_n x^n,$$

and substituting it into the above equation and setting all coefficients of each term  $x^n$  to be zeros, we obtain the formulas of  $a_n$  as follows

$$a_{n+1} = \frac{a + bu^n}{\{n+1\}_{s,t}} \alpha a_n + \frac{b_n}{\{n+1\}_{s,t}}.$$

For  $n \geq 1$ , we have

$$a_n = \frac{(a \oplus b)_{1,u}^n}{\{n\}_{s,t}!} \alpha^n a_0 + \frac{(a \oplus b)_{1,u}^n}{\{n\}_{s,t}!} \sum_{k=0}^{n-1} \frac{\{k\}_{s,t}! \alpha^{n-1-k}}{(a \oplus b)_{1,u}^{k+1}} b_k.$$

So the solution can be represented by

$$\begin{aligned} y(x) &= a_0 \sum_{n=0}^{\infty} \frac{(a \oplus b)_{1,u}^n}{\{n\}_{s,t}!} \alpha^n x^n + \sum_{n=1}^{\infty} \frac{(a \oplus b)_{1,u}^n}{\{n\}_{s,t}!} \left( \sum_{k=0}^{n-1} \frac{\{k\}_{s,t}! \alpha^{n-k-1}}{(a \oplus b)_{1,u}^{k+1}} b_k \right) x^n \\ &= a_0 E_{s,t}(a, b; \alpha x, u) + \sum_{n=1}^{\infty} \frac{(a \oplus b)_{1,u}^n}{\{n\}_{s,t}!} \left( \sum_{k=0}^{n-1} \frac{\{k\}_{s,t}! \alpha^{n-k-1}}{(a \oplus b)_{1,u}^{k+1}} b_k \right) x^n. \end{aligned}$$

The proof is completed.  $\square$

**Corollary 6.** Consider the following equation

$$\mathbf{D}_{s,t} y(x) = \alpha y(x) + \beta(x), \quad y(0) = a_0, \quad (55)$$

where  $\alpha$  is a constant, and  $\beta(x)$  is an analytic function with expansion

$$\beta(x) = \sum_{n=0}^{\infty} b_n x^n.$$

Then its solution is given by

$$y(x) = a_0 \exp_{s,t}(\alpha x, u) + \sum_{n=1}^{\infty} \left( \sum_{k=0}^{n-1} \frac{\{k\}_{s,t}! \alpha^{n-1-k}}{u^{\binom{k+1}{2}}} b_k \right) u^{\binom{n}{2}} \frac{x^n}{\{n\}_{s,t}!}. \quad (56)$$

**Corollary 7.** Set  $q = \varphi'_{s,t} / \varphi_{s,t}$ . Consider the following equation

$$\mathbf{D}_{s,t} y(x) = y(x) - qy(qx) + \beta(x), \quad y(0) = a_0, \quad (57)$$

where  $\alpha$  is a constant, and  $\beta(x)$  is an analytic function with expansion

$$\beta(x) = \sum_{n=0}^{\infty} b_n x^n.$$

Then its solution is given by

$$y(x) = a_0 \Theta_0((1-q)x, \varphi_{s,t}^{-1}) + \sum_{n=1}^{\infty} \left( \sum_{k=0}^{n-1} \frac{\varphi_{s,t}^{\binom{k}{2}}}{(1-q)^k (1-q^{k+1})} b_k \right) (1-q)^n \varphi_{s,t}^{-\binom{n}{2}} x^n. \quad (58)$$

**Example 7.** The solution of

$$\mathbf{D}_{s,t}y(x) = \alpha(ay(x) + by(ux)) + \delta + \epsilon x, \quad y(0) = a_0$$

is

$$y(x) = \left( c + \frac{\delta}{\alpha(a \oplus b)_{1,u}^1} + \frac{\epsilon}{\alpha^2(a \oplus b)_{1,u}^2} \right) \mathbf{E}_{s,t}(a, b; \alpha x, u) - \frac{\delta}{\alpha(a \oplus b)_{1,u}^1} - \frac{\epsilon}{\alpha^2(a \oplus b)_{1,u}^2}$$

where  $c = a_0$ .

**Example 8.** The solution of

$$\mathbf{D}_{s,t}y(x) = \alpha(ay(x) + by(ux)) + x^m, \quad y(0) = a_0$$

is

$$y(x) = c\mathbf{E}_{s,t}(a, b; \alpha x, u) + \frac{\{m\}_{s,t}!}{\alpha^{m+1}(a \oplus b)_{1,u}^{m+1}} \left( \mathbf{E}_{s,t}(a, b; \alpha x, u) - \sum_{n=0}^m \frac{(a \oplus b)_{1,u}^n}{\{n\}_{s,t}!} (\alpha x)^n \right)$$

where  $c = a_0$ .

**Theorem 5.** The solution of

$$\mathbf{D}_{s,t}y(x) = \varphi'_{s,t}(ay(x) + by(ux)) + \beta(a\mathbf{E}_{s,t}(a, b, u; \varphi_{s,t}x) + b\mathbf{E}_{s,t}(a, b, u; \varphi_{s,t}ux))$$

is given by

$$y(x) = c\mathbf{E}_{s,t}(a, b; \varphi'_{s,t}x, u) + \beta x(a\mathbf{E}_{s,t}(a, b; x, u) + b\mathbf{E}_{s,t}(a, b; ux, u)),$$

where  $c = a_0$ .

*Proof.* The power series expansion of  $\beta(a\mathbf{E}_{s,t}(a, b; \varphi_{s,t}x, u) + b\mathbf{E}_{s,t}(a, b; \varphi_{s,t}ux, u))$  is

$$\beta(a\mathbf{E}_{s,t}(a, b; \varphi_{s,t}x, u) + b\mathbf{E}_{s,t}(a, b; \varphi_{s,t}ux, u)) = \beta \sum_{k=0}^{\infty} (a \oplus b)_{1,u}^{k+1} \varphi_{s,t}^k \frac{x^k}{\{k\}_{s,t}!}$$

which gives

$$b_k = \beta \frac{(a \oplus b)_{1,u}^{k+1}}{\{k\}_{s,t}!} \varphi_{s,t}^k.$$

Substituting it into the Eq.(54) yields

$$\begin{aligned} y(x) &= a_0\mathbf{E}_{s,t}(a, b; \varphi'_{s,t}x, u) + \beta \sum_{n=1}^{\infty} \left( \sum_{k=0}^{n-1} \frac{\{k\}_{s,t}! \varphi_{s,t}'^{(n-1-k)}}{(a \oplus b)_{1,u}^{k+1}} \frac{(a \oplus b)_{1,u}^{k+1}}{\{k\}_{s,t}!} \varphi_{s,t}^k \right) (a \oplus b)_{1,u}^n \frac{x^n}{\{n\}_{s,t}!} \\ &= a_0\mathbf{E}_{s,t}(a, b; \varphi'_{s,t}x, u) + \beta \sum_{n=1}^{\infty} \left( \sum_{k=0}^{n-1} \varphi_{s,t}^k \varphi_{s,t}'^{(n-1-k)} \right) (a \oplus b)_{1,u}^n \frac{x^n}{\{n\}_{s,t}!} \\ &= a_0\mathbf{E}_{s,t}(a, b; \varphi'_{s,t}x, u) + \beta \sum_{n=1}^{\infty} \{n\}_{s,t} (a \oplus b)_{1,u}^n \frac{x^n}{\{n\}_{s,t}!} \\ &= a_0\mathbf{E}_{s,t}(a, b; \varphi'_{s,t}x, u) + \beta \sum_{n=1}^{\infty} (a \oplus b)_{1,u}^n \frac{x^n}{\{n-1\}_{s,t}!} \\ &= a_0\mathbf{E}_{s,t}(a, b; \varphi'_{s,t}x, u) + \beta x \sum_{n=0}^{\infty} (a \oplus b)_{1,u}^{n+1} \frac{x^n}{\{n\}_{s,t}!} \\ &= a_0\mathbf{E}_{s,t}(a, b; \varphi'_{s,t}x, u) + \beta x(a\mathbf{E}_{s,t}(a, b; x, u) + b\mathbf{E}_{s,t}(a, b; ux, u)). \end{aligned}$$

The proof is completed.  $\square$

**Corollary 8.** *The solution of*

$$\mathbf{D}_{s,t}y(x) = \varphi'_{s,t}y(ux) + \beta \exp_{s,t}(\varphi_{s,t}ux, u)$$

is given by

$$y(x) = c \exp_{s,t}(\varphi'_{s,t}x, u) + \beta x \exp_{s,t}(ux, u),$$

where  $c = a_0$ .

**Corollary 9.** *The solution of*

$$\mathbf{D}_{s,t}y(x) = \varphi'_{s,t}(y(x) - qy(qx)) + \beta(\Theta_0((1-q)\varphi_{s,t}x, \varphi_{s,t}^{-1}) - q\Theta_0((1-q)\varphi_{s,t}qx, \varphi_{s,t}^{-1}))$$

is given by

$$y(x) = c\Theta_0((1-q)\varphi'_{s,t}x, \varphi_{s,t}^{-1}) + \beta x(\Theta_0((1-q)x, \varphi_{s,t}^{-1}) - q\Theta_0((1-q)qx, \varphi_{s,t}^{-1})),$$

where  $c = a_0$ .

### 4.3 Operator method

We use an operator method to deal with a linear proportional difference equation. Two basic operator are  $(s, t)$ -derivative operator  $\mathbf{D}_{s,t}$  and the scale  $T_u$  defined by

$$T_u y(x) = y(ux).$$

Two basic properties of these two operators are

$$\begin{aligned} \mathbf{D}_{s,t}T_u &= uT_u\mathbf{D}_{s,t}, \\ T_uT_v &= T_{uv}. \end{aligned}$$

**Theorem 6.** *For the first-order equation*

$$\mathbf{D}_{s,t}y(x) = a\beta y(x) + \gamma T_u y(x) + \delta \mathbf{E}_{s,t}(a, b; \alpha x, u) \quad (59)$$

the general solution is given by

$$y(x) = c\mathbf{E}_{s,t}(a\beta, \gamma; x, u) + \frac{u\delta}{b\alpha - u\gamma}\mathbf{E}_{s,t}(a, b; \alpha x/u, u) \quad (60)$$

*Proof.* Formally, we have

$$y(x) = \delta(\mathbf{D}_{s,t} - a\beta 1 - \gamma T_u)^{-1}\mathbf{E}_{s,t}(a, b; \alpha x, u).$$

To compute the right side of the above equation, we use the following formula

$$(\mathbf{D}_{s,t} - a\beta 1 - \gamma T_u)\mathbf{E}_{s,t}(a, b; \beta x, u) = (b\beta - \gamma)\mathbf{E}_{s,t}(a, b; u\beta x, u).$$

So, taking  $\beta = \frac{\alpha}{u}$  gives

$$(\mathbf{D}_{s,t} - a\frac{\alpha}{u}1 - \gamma T_u)^{-1}\mathbf{E}_{s,t}(a, b; \alpha x, u) = \frac{1}{\frac{b\alpha}{u} - \gamma}\mathbf{E}_{s,t}(a, b; \alpha x/u, u).$$

Therefore, a special solution is given by

$$y^*(x) = \frac{u\delta}{b\alpha - u\gamma}\mathbf{E}_{s,t}(a, b; \alpha x/u, u)$$

and the general solution is

$$y(x) = c\mathbf{E}_{s,t}(a\beta, \gamma; x, u) + \frac{u\delta}{b\alpha - u\gamma}\mathbf{E}_{s,t}(a, b; \alpha x/u, u).$$

□

#### 4.4 Deformed $(s, t)$ -Bernoulli Equation

In this section we will focus on solving  $(s, t)$ -equations in the form

$$D_{\varphi^{n-1}, \varphi^{(n-1)}} y(x) + \alpha(x) \frac{(D_q E_{s,t})(a, b; x, u) \square A(x)}{E_{s,t}[a, b; A(\varphi_{s,t} x), u]} y(\varphi_{s,t}^{n-1} x) = \beta(x) \prod_{j=0}^{\infty} \frac{y(q^j \varphi_{s,t}^{n-1} x)}{y(q^{n+j} \varphi_{s,t}^{n-1} x)} \quad (61)$$

and

$$D_{\varphi^{n-1}, \varphi^{(n-1)}} y(x) + \alpha(x) \frac{(D_q E_{s,t})(a, b; x, u) \square A(x)}{E_{s,t}[a, b; A(\varphi'_{s,t} x), u]} y(\varphi'_{s,t}{}^{(n-1)} x) = \beta(x) \prod_{j=0}^{\infty} \frac{y(q^j \varphi_{s,t}^{n-1} x)}{y(q^{n+j} \varphi_{s,t}^{n-1} x)} \quad (62)$$

where  $\alpha(x)$  and  $\beta(x)$  are continuous functions,  $n$  is any real number, and where

$$D_{\varphi^{n-1}, \varphi^{(n-1)}} f(x) = \frac{f(\varphi_{s,t}^{n-1} x) - f(\varphi'_{s,t}{}^{(n-1)} x)}{(\varphi_{s,t}^{n-1} - \varphi'_{s,t}{}^{(n-1)})x}$$

is a  $(p, q)$ -derivative. Note that  $n \neq 0$  and  $n \neq 1$ . Also, note that when  $s = 2$  and  $t = -1$ , those  $(s, t)$ -difference equations Eqs.(61,62) lead to the usual Bernoulli equations.

To solve  $(s, t)$ -difference equation Eq.(61), it is first divided by  $\prod_{j=0}^{\infty} \frac{y(q^j \varphi_{s,t}^{n-1} x)}{y(q^{n+j} \varphi_{s,t}^{n-1} x)}$ ,

$$D_{\varphi^{n-1}, \varphi^{(n-1)}} y(x) \prod_{j=0}^{\infty} \frac{y(q^{n+j} \varphi_{s,t}^{n-1} x)}{y(q^j \varphi_{s,t}^{n-1} x)} + \alpha(x) \prod_{j=0}^{\infty} \frac{y(q^{n+j} \varphi_{s,t}^{n-1} x)}{y(q^{j+1} \varphi_{s,t}^{n-1} x)} = \beta(x).$$

Substitution of  $z(x) = \prod_{j=0}^{\infty} \frac{y(q^{n-1+j} \varphi_{s,t}^{n-2} x)}{y(q^j \varphi_{s,t}^{n-2} x)}$  is used to convert the above equation into a  $(s, t)$ -difference equation in terms of  $z(x)$ . The  $(s, t)$ -derivative of  $z(x)$  is as follows

$$\begin{aligned} \mathbf{D}_{s,t} z(x) &= \mathbf{D}_{s,t} \left( \prod_{j=0}^{\infty} \frac{y(q^{n-1+j} \varphi_{s,t}^{n-2} x)}{y(q^j \varphi_{s,t}^{n-2} x)} \right) \\ &= \frac{1}{(\varphi_{s,t} - \varphi'_{s,t})x} \left( \prod_{j=0}^{\infty} \frac{y(q^{n-1+j} \varphi_{s,t}^{n-1} x)}{y(q^j \varphi_{s,t}^{n-1} x)} - \prod_{j=0}^{\infty} \frac{y(q^{n-1+j} \varphi_{s,t}^{n-2} \varphi'_{s,t} x)}{y(q^j \varphi_{s,t}^{n-2} \varphi'_{s,t} x)} \right) \\ &= -\frac{\varphi_{s,t}^{n-1} - \varphi'_{s,t}{}^{(n-1)}}{(\varphi_{s,t} - \varphi'_{s,t})} \frac{y(\varphi_{s,t}^{n-1} x) - y(\varphi'_{s,t}{}^{(n-1)} x)}{(\varphi_{s,t}^{n-1} - \varphi'_{s,t}{}^{(n-1)})x} \prod_{j=0}^{\infty} \frac{y(q^{n+j} \varphi_{s,t}^{n-1} x)}{y(q^j \varphi_{s,t}^{n-1} x)} \\ &= -\{n-1\}_{s,t} D_{\varphi^{n-1}, \varphi^{(n-1)}} y(x) \prod_{j=0}^{\infty} \frac{y(q^{n+j} \varphi_{s,t}^{n-1} x)}{y(q^j \varphi_{s,t}^{n-1} x)}. \end{aligned}$$

Then the  $(s, t)$ -difference equation equivalent is

$$-\frac{1}{\{n-1\}_{s,t}} \mathbf{D}_{s,t} z(x) + \alpha(x) \frac{(D_q E_{s,t})(a, b; x, u) \square A(x)}{E_{s,t}[a, b; A(\varphi_{s,t} x), u]} z(\varphi'_{s,t} x) = \beta(x). \quad (63)$$

The solution  $z(x)$  is used to find the required solution  $y(x)$ . Let us rewrite it as

$$y(x) = \begin{cases} y(q^{n-1}x) \frac{z(qx/\varphi_{s,t}^{n-2})}{z(x/\varphi_{s,t}^{n-2})}, & \text{if } n \neq 2; \\ \frac{1}{z(x)}, & \text{if } n = 2. \end{cases}$$

The previous expression can be rewrite for  $y(x)$  as follows:

1. For  $0 < |q| < 1$ ,  $n > 1$  and  $n \neq 2$ :

$$y(x) = y(0) \prod_{i=0}^{\infty} \frac{z(q^{i(n-1)+1}x/\varphi_{s,t}^{n-2})}{z(q^{i(n-1)}x/\varphi_{s,t}^{n-2})}.$$

2. For  $|q| > 1$  and  $n < 1$ :

$$y(x) = y(0) \prod_{i=0}^{\infty} \frac{z(q^{i(n-1)+1}x/\varphi_{s,t}^{n-2})}{z(q^{i(n-1)}x/\varphi_{s,t}^{n-2})}.$$

3. For  $n = 2$ ,

$$y(x) = \frac{1}{z(x)}.$$

4. Otherwise

$$y(x) = y(\infty) \prod_{i=0}^{\infty} \frac{z(q^{i(n-1)+1}x/\varphi_{s,t}^{n-2})}{z(q^{i(n-1)}x/\varphi_{s,t}^{n-2})}.$$

In a similar way, the  $(s, t)$ -difference equation in Eq.(62) is converted to the following  $(s, t)$ -difference equation in term of  $z(x)$

$$-\frac{1}{\{n-1\}_{s,t}} \mathbf{D}_{s,t}z(x) + \alpha(x) \frac{(D_q E_{s,t})(a, b; x, u) \square A(x)}{E_{s,t}[a, b; A(\varphi'_{s,t}x), u]} z(\varphi_{s,t}x) = \beta(x) \quad (64)$$

and the required solution is found to be as the previous one. Finally, the Eqs.(63,64) are solved for  $z(x)$  as previously carried out.

**Example 9.** *The equation*

$$D_{\varphi, \varphi'} y(x) + y(\varphi_{s,t}x) = x^2 y(\varphi_{s,t}x) y(\varphi'_{s,t}x)$$

is a  $(s, t)$ -Bernoulli difference equation for  $n = 2$ . Let  $z(x) = \frac{1}{y(x)}$ . Now the  $(s, t)$ -difference equation is transformed on  $z(x)$ , to

$$\mathbf{D}_{s,t}z(x) - z(\varphi'_{s,t}x) = -x^m.$$

From Example 3, the solution is

$$z(x) = G(\log_q(x)) \text{Exp}'_{s,t}(x) + \sum_{k=0}^m \varphi_{s,t}'^{(-m(k+1)+\binom{k+1}{2})} \frac{\{m\}_{s,t}!}{\{m-k\}_{s,t}!} x^{m-k}.$$

Finally, we define the following  $u$ -deformed  $(s, t)$ -Bernoulli difference equations

$$D_{\varphi^{n-1}, \varphi^{(n-1)}} y(x) + \alpha y(\varphi_{s,t}^{n-1} ux) = \beta(x) \prod_{j=0}^{\infty} \frac{y(q^j \varphi_{s,t}^{n-1} ux)}{y(q^{n+j} \varphi_{s,t}^{n-1} ux)} \quad (65)$$

and

$$D_{\varphi^{n-1}, \varphi^{(n-1)}} y(x) + \alpha y(\varphi'_{s,t}{}^{(n-1)} ux) = \beta(x) \prod_{j=0}^{\infty} \frac{y(q^j \varphi_{s,t}^{n-1} ux)}{y(q^{n+j} \varphi_{s,t}^{n-1} ux)} \quad (66)$$

with initial value problem  $y(0) = a_0$ . With the substitution

$$z(x) = \prod_{j=0}^{\infty} \frac{y(q^{n-1+j} \varphi_{s,t}^{n-2} ux)}{y(q^j \varphi_{s,t}^{n-2} ux)}$$

the Eqs.(65,66) are transformed into the equations

$$-\frac{1}{\{n-1\}_{s,t}} \mathbf{D}_{s,t} z(x) + \alpha z(\varphi'_{s,t} ux) = \beta(x)$$

and

$$-\frac{1}{\{n-1\}_{s,t}} \mathbf{D}_{s,t} z(x) + \alpha z(\varphi_{s,t} ux) = \beta(x)$$

which can be solved by using the Corollary 6.

## References

- [1] V. Kac, C. Pokman, *Quantum Calculus*, Universitext, Springer-Verlag, 2002
- [2] M. Ward, A calculus of sequences, *Am. J. Math.* **58**(1936) 255–266.
- [3] A.F. Horadam, Basic Properties of a Certain Generalized Sequence of Numbers, *The Fibonacci Quart.*, **3:3** (1965), 161–176.
- [4] R. Ranjan, *Sources in the Development of Mathematics: Series and Products from the fifteenth to the Twenty-first Century*, Cambridge University Press.
- [5] A. Iserles, On the generalized pantograph functional-differential equation. *European J. Appl. Math.* **4**, (1993), 1–38
- [6] A. Iserles, Y. Liu, On functional-differential equation with proportional delays, *J. Math. Anal. Appl.* **207**, 73–95 (1997)
- [7] P. Njionou Sadjang, On the fundamental Theorem of  $(p, q)$ -Calculus and Some  $(p, q)$ -Taylor Formulas. *Results Math* **73**, 39 (2018) <https://doi.org/10.1007/s00025-018-0783-z>
- [8] R. Orozco, Existence of Solutions of Functional-Difference Equations with Proportional Delay on Deformed Generalized Fibonacci Polynomials via Successive Approximation and Bell Polynomials (under revision), arXiv:2211.04450 (2022)
- [9] I. M. Gessel, A  $q$ -analog of the exponential formula. *Discrete Math.* **40** (1982), 69–80
- [10] W. P. Johnson, A  $q$ -analogue of Faá di Bruno's formula. *J. Comb. Theory, A.* **76** (1996), 305–314