

# Energy Window Augmented Plane Waves (EWAPW)

Garry Goldstein

*garrygoldsteinwinnipeg@gmail.com*

In this work we present a new basis set for electronic structure calculations of crystalline solids using Density Functional theory (DFT) methods. In this construction we take advantage of the fact that most DFT calculations use a convergence loop in order to obtain the eigenstates of a final Khon Sham (KS) Hamiltonian matrix whose eigenstates also give the appropriate electron density needed to obtain the KS potential needed for that KS Hamiltonian matrix. Here we propose that for the basis of each step of the iteration we use the previous eigenstate basis in the interstitial region but augmented inside the MT sphere with the solution to the spherically averaged KS Hamiltonian for the energy window of that eigenstate. To reduce the number of times the KS potential needs to be solved inside the MT spheres it is advantageous to use energy windows and solve the KS Hamiltonian inside the MT region only once per window (at some energy inside the window) so that the KS Hamiltonian needs only be solved a small number of times per iteration, for practical applications on the order of 10 to 100 windows. This method combines the energy dependence of methods such as Projected Augmented Wave functions (PAW) with the ability of the basis set to adjust to the solid state (rather than atomic) environment of basis sets such as Linearized Augmented Plane Waves (LAPW).

## I. INTRODUCTION

The usefulness of a basis set for a Density Functional Theory (DFT) calculation is mainly determined by its linearization error (that is energy errors for states inside the valence band), its ability to handle semi-core states, that is the situation when there is more than one relevant band (valence band) just at or below the Fermi energy, the effectiveness with which the basis set converges to the exact eigenstates of the KS Hamiltonian when the cutoff  $\mathbf{K}_{max}$  is raised, the efficiently one can write down the Khon Sham (KS) Hamiltonian in that basis amongst other things [1–3]. There are now several known competitors in the literature for which basis set is optimal. These can be divided into mostly plane wave basis sets and mostly Muffin Tin (MT) basis set [2]. One approach, which is mostly MT, is the Linear Muffin Tin Orbitals (LMTO) basis set [2, 4–8]. In this basis set the main goal is to work with as small a basis set as possible but still obtain good results for eigenstates and eigen-energies of the KS Hamiltonian, but only near the Fermi surface (valence band). For these basis sets one looks for solutions to the spherically averaged KS problem inside a muffin tin (MT) sphere of an atom, in some angular momentum channel  $Y_{lm}$ , and matches it with outgoing spherical Bessel function solutions multiplied by spherical Harmonics,  $Y_{lm}$ . The key condition enforced in the LMTO basis set is that the Bessel function “tails” from all MT spheres cancel and inside the MT spheres as such only the “head” functions corresponding to solutions of the spherically averaged KS Hamiltonian remain inside the MT spheres. This produces a very small and efficient basis set, which can be made even more efficient in the Atomic Sphere Approximation (ASA) where the spheres are space filling and the Bessel functions, often now taken just at zero energy, are not explicitly part of the basis wave functions and are used as an intermediate step to ensure smoothness. This eliminates the need for

having plane wave wave functions (or Bessel functions) in the interstitial thereby simplifying the situation. The LMTO basis is most efficient but not necessarily simplest or most accurate as it is only good near the middle of the valence band and uses very large MT spheres where spherical symmetry is poor. The Korringa Khon Rostoker (KKR) basis set [2, 9, 10] is related to the LMTO basis set [11] however it uses a variable energy to compute the eigenstate of the KS potential in the MT sphere with the energy being determined by tail cancellation inside the MT sphere. This allows one to obtain eigenstates at all energies not just near the Fermi surface (valence band). This method is highly inefficient numerically, unfortunately, as it requires one to solve the KS Hamiltonian at a large number of eigen-energies which slows down computational times by an order of magnitude or more. This method within the ASA approximation also eliminates the need for all plane wave (or Bessel function) basis elements.

The competitor method to Muffin Tin wave functions is the plane wave method, which strives for high accuracy at the cost of greater computational times [1, 2, 4]. Within the plane wave method, on the other hand, the plane wave basis set is very simple but is extremely inefficient at obtaining good eigenstates of the KS Hamiltonian for reasonably small wave vector cutoffs. Indeed near the nuclei, where the electron density is high and highly concentrated (oscillating), the KS potential has high wave vector components so that the eigenstates of the KS Hamiltonian have high wave vector components as well. This means that impractically large numbers of plane waves are needed to accurately describe the environment near atomic nuclei, making the situation generically numerically impossible. There are two main approaches to cure this problem: pseudopotentials and augmentation, both of which we now describe in turn.

The simplest pseudopotentials are norm conserving pseudopotentials [2, 3]. The idea behind norm conserv-

ing pseudopotentials is that to obtain the eigen-energies (but not the eigenstates) of the KS Hamiltonian, if one assumes a spherically symmetric Atomic Sphere around the nucleus, one does not need the exact nuclear eigen-wavefunction for each angular momentum channel, but any wavefunction, with the same scattering phase shift at the atomic sphere radius, would do just as well [2]. As such one can replace the atomic sphere around a nucleus with a collection of scattering phase shifts in every angular momentum channel and for every energy. However only the middle of the valence band is of interest for many DFT calculations so the scattering phase shift there matters the most. As such for practical calculations associated with modern code one replaces the atomic KS potential with a collection of pseudopotentials (potentials with explicit angular momentum projection) in each angular momentum channel, which have the scattering phase shifts the same as the one in the middle of the valence band for the atomic problem in that angular momentum channel. Furthermore since one need only match a single scattering phase shift the pseudopotential may be quite smooth and as such well described by a small number of wavevectors. By also adding the norm conserving condition one ensures that the first derivative of the scattering phase shift, with respect to energy, matches with the one from the atomic calculations and that the Madelung energies are roughly correct (in that charge is conserved inside the atomic sphere, however higher multipoles do not match as the wavefunctions does not match the atomic wavefunctions at all radii so the Madelung energy is not exact but good for small spheres). This is an efficient method to study electronic structures.

There are many ways to improve upon norm conserving pseudopotentials. Ultrasoft pseudopotentials allow one to study the scattering phase shift at a variety of energies where the scattering phase shift of an eigenstate, with a specific energy from the set of eigenstates chosen to make the ultrasoft potential, matches the scattering phase shift of the atomic potential at that energy and as such by extension for a variety of energies [1-3, 12]. Furthermore norm conservation is no longer enforced but Madelung energies are preserved (at least the first multipole) by introducing a fictitious non-identity overlap matrix between the smooth pseudopotential eigenwavefunctions which leads to the eigenstates having the same overlaps inside the atomic sphere as if they were norm conserving. Furthermore, since we have scattering phase shifts at different eigenenergies the second advantage of norm conservation (the correctness of the scattering phase shift near the middle of the valence band due to matching of derivatives of the scattering phase shift between the pseudopotential and the actual potential) is no longer essential. The relaxation of the norm conserving condition allows for smoother pseudopotentials at various energies and makes the cutoff  $\mathbf{K}_{max}$  smaller (softer). Indeed if all derivatives of the scattering phase shift with respect to energy are matched with the atomic potential and as such the pseudopotential matches the atomic

potential at all orders in perturbation theory and then the pseudopotential must be the exact potential of the atom and we have gained nothing - this is we might as well expand the exact KS problem with plane waves. As such this relaxation of norm conservation leads to much smoother potentials and lower cutoffs.

Another method to improve upon norm conserving pseudopotentials is the Projected Augmented Waves (PAW) method [2, 3, 13]. In PAW there is a further advantage in that the atomic eigenstates with various scattering shifts and norms are linearly transformed into smooth wavefunctions within some atomic spheres (no phase shift matching is needed). These smooth wavefunctions have a good expansion in terms of plane waves (with good convergence often as low as 30 Rydberg). Whereby the transform between the atomic wavefunctions and the smooth wavefunctions can be used to setup a all electron KS problem which represents the exact many body problem but with a plane wave basis (this is similar to but more effective than the Phillips-Kleinman pseudopotential approach). In all pseudopotential methods there are no changes to the scattering pseudopotentials or atomic wavefunctions (PAW) as a function of the all electron KS problem, which is a marked disadvantage as the atomic environment, even for small atomic spheres, is not a perfect match for the solid state environment.

In augmentation methods we overcome the difficulty of having a fixed atomic sphere and adjust the basis to the exact KS problem. In particular, in Linearized Augmented Plane Waves (LAPW) plane waves are augmented with the solutions to the KS Hamiltonian at the linearization energy (typically the middle of the valence band)  $\psi(r)$  and its derivative with respect to energy  $\dot{\psi}(r) = \frac{\partial}{\partial E}\psi(r)$  by matching the plane wave wavefunction and its derivative with respect to the radius at the MT sphere radius [1, 2, 5, 14, 15]. This allows for efficient description of the valence band. This method can be further improved as in Higher Derivative Localized Orbitals (HDLO) where  $\ddot{\psi}(r) = \frac{\partial^2}{\partial E^2}\psi(r)$  is used inside the MT sphere and one more derivative is matched which reduces linearization errors. However once again the more derivatives are matched the more the radial wavefunction inside the MT sphere look like Bessel functions the higher cutoff, harder, the basis is [16, 17]. If there are several relevant bands one can use LO or lo basis set [1, 18, 19] (localized orbitals at that band energy) or multi-window options (in extreme cases where it is of interest to have multiple copies of basis sets one at each linearization energy [1]). While LAPW and other augmented methods do have direct connections between the augmentation wavefunctions (their norms, multipole moments and phase shifts) with the exact KS problem they are limited in that they are closer to norm conserving pseudopotentials rather than ultrasoft and PAW in that there is usually one linearization energy much like there is one scattering phase shift for norm conserving pseudopotentials and limited adjustment for multiple bands of interest. In Slater's APW [20] method the wavefunc-

tion inside the MT sphere depends on the energy of the eigenstate but it has the same problem as KKR, the KS Hamiltonian inside the MT sphere has to be solved at a very large number of energies leading to an increase by more than an order of magnitude of the computer resources needed.

In this work we overcome these difficulties and combine the advantages of PAW (the ability to adjust to different energies) with LAPW (the ability to adjust to the solid state environment) without resorting to a large number of solutions of the KS problem inside the MT sphere but instead to a solution of the problem in energy windows (on the order of ten to one hundred windows for practical applications). Our new method is based on using a convergence loop to adjust the form of the Energy Window Augmented Plane Wave (EWAPW) with each iteration. We use the eigenstate basis from the previous EWAPW calculation within the interstitial region as the basis for the next EWAPW calculation except the augmented pieces inside the MT spheres are the eigenstates of the spherically averaged to KS Hamiltonian from the window relevant to the previous eigenenergy. More precisely we keep the plane wave piece with all the expansion coefficients:

$$\chi_E(\mathbf{k}, \mathbf{r}) = \frac{1}{\sqrt{V}} \sum_{\mathbf{K}} A_E^{\mathbf{K}}(\mathbf{k}) \exp(i(\mathbf{k} + \mathbf{K}) \cdot \mathbf{r}) \quad (1)$$

Here  $\mathbf{r}$  is inside the interstitial and  $\chi_E(\mathbf{k}, \mathbf{r})$  is the eigenstate from the previous iteration and  $A_E^{\mathbf{K}}(\mathbf{k})$  are its expansion coefficients in terms of plane waves inside the interstitial region. Here  $V$  is the volume of a unit cell, and match it to the spherically averaged solutions to the KS Hamiltonian within each MT sphere  $\mu$ :

$$\left[ -\frac{d^2}{dr^2} + \frac{l(l+1)}{r^2} + \bar{J}_{KS}(r) \right] r\psi_{l\mu}^{E_W}(r) = E_W r\psi_{l\mu}^{E_W}(r) \quad (2)$$

and  $\bar{J}_{KS}(r)$  is the spherically average Khon Sham (KS) potential. We also choose the coefficients  $A_{lm}^{\mu E_W, \mathbf{k}}$  such that the wave function is continuous as  $S_\mu$ , whereby we must have that:

$$A_{lm}^{\mu E_W, \mathbf{k}} = \sum_{\mathbf{K}} A_E^{\mathbf{K}}(\mathbf{k}) \frac{\frac{1}{\sqrt{V}} 4\pi i^l J_l(|\mathbf{k} + \mathbf{K}| S_\mu) Y_{lm}^*(\widehat{\mathbf{k} + \mathbf{K}})}{\psi_{l\mu}^{E_W}(S_\mu)} \times \exp(i(\mathbf{k} + \mathbf{K}) \cdot \mathbf{r}_\mu) \quad (3)$$

Whereby the basis wavefunctions for the next iteration are given by:

$$\begin{aligned} \check{\chi}^{\mathbf{k}, E}(\mathbf{r}; \mathbf{r}_{\mu,1}, \dots, \mathbf{r}_{\mu,M}) &= \\ &= \frac{1}{\sqrt{V}} \sum_{\mathbf{K}} A_E^{\mathbf{K}}(\mathbf{k}) \exp(i(\mathbf{k} + \mathbf{K}) \cdot \mathbf{r}) \prod_{\mu} \Theta(|\mathbf{r} - \mathbf{r}_\mu| - S_\mu) \\ &+ \sum_{\mu=1}^M \sum_{l,m} Y_{lm}(\widehat{\mathbf{r} - \mathbf{r}_\mu}) \left[ A_{lm}^{\mu E_W, \mathbf{k}} \psi_{l\mu}^{E_W}(\mathbf{r} - \mathbf{r}_\mu) \right] \times \\ &\times \Theta(S_\mu - |\mathbf{r} - \mathbf{r}_\mu|) \end{aligned} \quad (4)$$

Here  $E_W$  is in the middle of an energy window (we divide the entire energy range of the calculation into a number of windows, on the order of 10 to 100 for practical calculations, and use those energies for linearizations) which contains  $E$ . Here  $M$  is the total number of atoms per unit cell and  $\Theta$  is the heavy-side function. We will now use  $\check{\chi}_{\mathbf{K}}^{\mathbf{k}, E}(\mathbf{r}; \mathbf{r}_{\mu,1}, \dots, \mathbf{r}_{\mu,M})$  as the basis for the next convergence iteration and so forth until convergence to the solution of the KS problem. The key formula needed for Eq. (3) is given by [21]:

$$\begin{aligned} &\frac{1}{\sqrt{V}} \exp(i(\mathbf{k} + \mathbf{K}) \cdot \mathbf{r}) \\ &= \exp(i(\mathbf{k} + \mathbf{K}) \cdot \mathbf{r}_\mu) \frac{1}{\sqrt{V}} 4\pi \times \\ &\times \sum_{l,m} Y_{lm}^*(\widehat{\mathbf{k} + \mathbf{K}}) Y_{lm}(\widehat{\mathbf{r} - \mathbf{r}_\mu}) i^l J_l(|\mathbf{k} + \mathbf{K}| |\mathbf{r} - \mathbf{r}_\mu|) \end{aligned} \quad (5)$$

## II. MAIN CONSTRUCTION

We begin with the eigenstate equation:

$$\sum_{\mathbf{K}'} [\bar{H}_{KS}^{\mathbf{k}}]_{E, E'} A_{n, E'}^{\mathbf{k}} = \epsilon_n^{\mathbf{k}} \sum_{\mathbf{K}'} [\bar{O}^{\mathbf{k}}]_{E, E'} A_{n, E'}^{\mathbf{k}} \quad (6)$$

Where  $[\bar{H}_{KS}^{\mathbf{k}}]_{E, E'}$  and  $[\bar{O}^{\mathbf{k}}]_{E, E'}$  are obtained in Section III. Now we have that  $\chi_E(\mathbf{k}, \mathbf{r})$  is given by Eq. (1) as such we have that

$$\chi^{\mathbf{k}, E} \equiv \chi^{\mathbf{k}, \epsilon_n^{\mathbf{k}}} \frac{1}{\sqrt{V}} \sum_{\mathbf{K}, E} A_{n, E}^{\mathbf{k}} A_E^{\mathbf{K}}(\mathbf{k}) \exp(i(\mathbf{k} + \mathbf{K}) \cdot \mathbf{r}) \quad (7)$$

As such we have that:

$$A_{E=\epsilon_n^{\mathbf{k}}}^{\mathbf{K}}(\mathbf{k}) = \sum_E A_{n, E}^{\mathbf{k}} A_E^{\mathbf{K}}(\mathbf{k}) \quad (8)$$

Whereby we use Eqs. (1) through (4) to obtain the next basis set.

## III. EWFAPW

Here we would like to do full potential calculations for the EWAPW basis or EWFAPW. We break everything into simpler pieces which are easier to code and reuses many pieces of existing codes [22–28]. Because the wavefunction and its derivative are nearly continuous near the MT sphere (see Section IV D) it is appropriate to use FLAPW appropriate to LAPW basis set, however corrections due to discontinuity of derivatives and the wavefunction (due to angular momentum truncation) are possible [21].

### A. Overlap

The overlap is a sum of the interstitial part and the MT part which we compute in turn.

#### 1. MT Part

We introduce

$$[O]_{\varepsilon_1, \varepsilon_2}^{l, \mu} = \int_0^{S_\mu} r^2 dr \psi_{\varepsilon_1}^l(r) \psi_{\varepsilon_2}^l(r) \quad (9)$$

This means that

$$[O^{\mathbf{k}}]_{E, E'}^{MT, \mu} = \sum_{l, m} A_{lm}^{\mu E_W, \mathbf{k}} A_{l-m}^{\mu E'_W, \mathbf{k}} [O]_{E_W, E'_W}^{l, \mu} \quad (10)$$

#### 2. Interstitial

We write

$$[O^{\mathbf{k}}]_{\mathbf{K}, \mathbf{K}'}^I = \Theta_{\mathbf{K}-\mathbf{K}'} \quad (11)$$

where

$$\Theta_{\mathbf{K}} = \delta_{\mathbf{K}, 0} - \sum_{\mu} \exp(-i\mathbf{K} \cdot \mathbf{r}_\mu) \frac{(4\pi S_\mu)^3}{V} \cdot \frac{J_1(|\mathbf{K}| S_\mu)}{|\mathbf{K}| S_\mu} \quad (12)$$

is the Fourier transform of

$$\Theta(\mathbf{r}) \equiv \prod_{\mu} \Theta(|\mathbf{r} - \mathbf{r}_\mu| - S_\mu) \quad (13)$$

whereby

$$[O^{\mathbf{k}}]_{E, E'}^I = \sum_{\mathbf{K}, \mathbf{K}'} [O^{\mathbf{k}}]_{\mathbf{K}, \mathbf{K}'}^I A_E^{\mathbf{K}*}(\mathbf{k}) A_{E'}^{\mathbf{K}'}(\mathbf{k}) \quad (14)$$

### B. KS Hamiltonian

The KS Hamiltonian is a sum of the MT part and the interstitial part which we compute in turn. We write:

$$J(\mathbf{r}) = \begin{cases} \sum_{\mathbf{K}} J_I^{\mathbf{K}} \exp(i\mathbf{K} \cdot \mathbf{r}) & \text{Interstitial} \\ \sum_{l, m} J_{l, m}(|\mathbf{r}|) Y_{l, m}(\hat{\mathbf{r}}) & \text{MT} \end{cases} \quad (15)$$

#### 1. MT Part

The diagonal piece is well known [2] so we focus on the non-diagonal pieces. We write:

$$\begin{aligned} & [-\bar{\Delta}^{\mathbf{k}} + \bar{J}^{\mathbf{k}}]_{E, E'}^{MT} \\ &= \sum_{\mu} \sum_{l, m} \sum_{l', m'} A_{lm}^{\mu E_W, \mathbf{k}*} \cdot t_{lm, l'm'}^{\mu, E, E'} \cdot A_{l'm'}^{\mu E'_W, \mathbf{k}} \end{aligned} \quad (16)$$

Where there no sum over repeated indices and

$$t_{lm, l'm'}^{\mu, E, E'} = \sum_{l''} I_{l'l''}^{\mu\alpha\beta} G_{l''l'm''}^{m'mm''} \quad (17)$$

and:

$$\begin{aligned} G_{l, l', l''}^{m, m', m''} &= \int Y_{l, m}^* Y_{l', m'} Y_{l'', m''} d\Omega, \\ I_{l, l', l''}^{\mu E E'} &= \int \psi_{E_W}^{l*}(r) J_{l''}^{\mu}(r) \psi_{E'_W}^{l'}(r) r^2 dr \end{aligned} \quad (18)$$

#### 2. Interstitial

Now in the interstitial we have that

$$[-\bar{\Delta}^{\mathbf{k}} + \bar{J}^{\mathbf{k}}]_{\mathbf{K}, \mathbf{K}'}^I = [J_I \Theta]_{\mathbf{K}-\mathbf{K}'} + \frac{1}{2m} [\mathbf{k} + \mathbf{K}']^2 \Theta_{\mathbf{K}-\mathbf{K}'} \quad (19)$$

Where

$$[AB]_{\mathbf{K}, \mathbf{K}'} = \sum_{\mathbf{K}''} [A]_{\mathbf{K}, \mathbf{K}''} [B]_{\mathbf{K}'', \mathbf{K}'} \quad (20)$$

whereby

$$[-\bar{\Delta}^{\mathbf{k}} + \bar{J}^{\mathbf{k}}]_{E, E'}^I = \sum_{\mathbf{K}, \mathbf{K}'} [-\bar{\Delta}^{\mathbf{k}} + \bar{J}^{\mathbf{k}}]_{\mathbf{K}, \mathbf{K}'}^I A_E^{\mathbf{K}*}(\mathbf{k}) A_{E'}^{\mathbf{K}'}(\mathbf{k}) \quad (21)$$

## IV. COMMENTS

There are many minor technical issues that need to be further handled here we describe the solutions to many of them.

### A. Initialization

We can initialize within the LAPW basis set with only one linearization energy or using the initial basis set for the interstitial region  $\chi(\mathbf{k} + \mathbf{K}) = \frac{1}{\sqrt{V}} \exp(i(\mathbf{k} + \mathbf{K}) \cdot \mathbf{r})$  with energy

$$E(\chi(\mathbf{k} + \mathbf{K})) = \frac{1}{2m} (\mathbf{k} + \mathbf{K})^2 + J_{KS}(\mathbf{Q} = 0) \quad (22)$$

where we use the augmentation energy  $E_W$  for the window relevant to  $E(\chi(\mathbf{k} + \mathbf{K}))$ , see Eqs. (3) (4), with

$$A_E^{\mathbf{K}'}(\mathbf{k}) = \delta(\mathbf{K} - \mathbf{K}'). \quad (23)$$

### B. Partial Diagonalization [1]

In many situations it is advantageous to only look for the lowest few bands and not diagonalize the entire KS

Hamiltonian matrix. In which case since we are interested in only the lowest bands, say only  $\mathcal{M}$  of them we now choose new “eigenkets” with energy given by Eq. (22) and expansion given by Eq. (23) where we order the bands according to the value of  $E(\chi(\mathbf{k} + \mathbf{K}))$  and do not use the first  $\mathcal{M}$  ones and add them as a basis to the  $\mathcal{M}$  diagonalized basis elements.

### C. Expansion of basis set during convergence iteration [29]

In many case it is worthwhile to use small basis sets at the beginning of the convergence loop and larger basis sets later. In which case many basis elements of the for  $\chi(\mathbf{k} + \mathbf{K})$  with  $\mathbf{K}$  large, in which case we can use the same procedure as in the initialization with augmentation being determined by Eqs. (22) and (23). This allows for basis expansion during the solution of the KS problem.

### D. EWAPW vs. EWLAPW and multiradius options [29]

We notice that the KS Hamiltonian has no singularities on the MT sphere radius. As such the wavefunction when expressed within the plane wave basis set and the basis set adapted to the MT sphere should be the same near the MT radius. Therefore once the two wavefunctions match on the surface of the sphere once convergence is well established so do their derivatives and higher derivatives, as such there is limited advantage to EWLAPW or using a variable multi radius basis to match additional derivatives.

## V. CONCLUSIONS

In this work we studied various methods to solve the KS problem once the form of the KS potential (Lo-

cal Density Approximation (LDA), Generalized Gradient Approximation GGA metaGGA etc.) has been fixed. We presented three historical candidates for optimal solutions. Indeed LMTO is optimized to get sufficiently accurate results with minimal basis sets and as such minimal computer work. PAW and other pseudopotentials as well as augmented methods which are optimized to solve the KS problem accurately but generically require significantly higher resources than LMTO like methods. Between PAW and augmentation methods, it is reasonable to disagree which is more accurate. On the one hand PAW and ultrasoft pseudopotentials incorporate the atomic potential at various energies on the other hand augmentation methods incorporate the solid state crystal potential at the linearization energy which can be significantly different than the atomic one. As a rule of thumb, if the system is highly correlated then studying the crystal potential at the linearization energy is better than studying the potential at various energies so LAPW is more accurate than PAW, while for highly itinerant systems PAW is likely better than LAPW. This suggests the use LDA+U (or LDA+DMFT) with PAW for many systems [30]. Here instead we combine the features of both PAW and LAPW by allowing us to study the energy dependence of the crystal KS Hamiltonian at a variety of linearization energies within the new EWAPW basis set, furthermore we showed that for practical implementations using EWFAPW it is possible to heavily reuse current FLAPW code leading to efficient implementations. In the future it would be of interest to go further and study LDA+U (or LDA+DMFT) with our new EWAPW basis as well as to study Pulay forces, pressure and stress for all methods. This could have real life applications to many real materials.

- 
- [1] D. J. Singh and D. Nordstrom, *Planewaves, pseudopotentials, and the LAPW method* (Springer, New York, 2006).
  - [2] R. M. Martin, *Electronic Structures Basic Theory and Practical Methods* (Cambridge University Press, Cambridge, 2020).
  - [3] D. Marx and J. Hutter, *Ab Initio Molecular Dynamics Basic Theory and Advanced Methods* (Cambridge University Press, Cambridge, 2009).
  - [4] J. M. Wills, M. Alouani, P. Anderson, A. Dellin, O. Eriksson, and O. Grechnev, *Full-Potential Electronic Structure Method Energy and Force Calculations with Density Functional Theory and Dynamical Mean Field Theory* (Springer, New York, 2010).
  - [5] O. K. Andersen, Phys. Rev. B **12**, 3060 (1975).
  - [6] H. L. Skriver, *The LMTO method Muffin-Tin Orbitals and Electronic Structure* (Springer, New York, 1984).
  - [7] O. K. Andersen, and O. Jepsen, Phys. Rev. Lett. **53**, 2571 (1984).
  - [8] O. K. Andersen, T. S.-Dasgupta, and S. Ezhof, Bull. Mat. Sci. **26**, 19 (2003).
  - [9] W. Kohn, N. Rostoker, Phys. Rev. **94**, 1111 (1954).
  - [10] J. Korringa, Physica **13**, 392 (1947).
  - [11] G. Goldstein, arXiv 2403.12846.
  - [12] D. Vanderbilt, Phys. Rev. B **41**, 7892 (1990).
  - [13] P. E. Blochl, Phys. Rev. B **50**, 17953 (1994).
  - [14] J. M. Soler, and A. R. Williams, Phys. Rev. B **40**, 1560 (1989).
  - [15] J. M. Soler and A. R. Williams, Phys. Rev. B **42**, 9728 (1990).

- [16] G. Michalícek, *Extending the precision and efficiency of all-electron full-potential linearized augmented plane-wave density functional theory* (Aachen University, 2014, thesis).
- [17] G. Michalícek, M. Betzinger, C. Freidrich and S. Blugel, *Comp. Phys. Comm.* **184**, 2670 (2013).
- [18] D. J. Singh, *Phys. Rev. B* **43**, 6388 (1991).
- [19] E. Sjostedt, L. Nordstrom, and D. J. Singh, *Sol. Sta. Comm.* **114**, 15 (2000).
- [20] J. C. Slater, *Phys. Rev.* **51**, 846 (1937).
- [21] T. L. Loucks, *Augmented Plane Wave Method* (W. A. Benjamin Inc., New York, 1967).
- [22] D. R. Hamann, *Phys. Rev. Lett.* **42**, **662** (1979).
- [23] E. Wimmer, H. Krakauer, M. Weinert and A. J. Freeman, *Phys. Rev. B* **24**, 864 (1981).
- [24] H. J. F. Jansen and A. J. Freeman, *Phys. Rev. B* **30**, 561 (1984).
- [25] S.-H. Wei, and H. Krakauer, *Phys. Rev. Lett.* **55**, 1200 (1985).
- [26] S.-H. Wei, H. Krakauer and M. Weinert, *Phys. Rev. B* **32**, 7792 (1985).
- [27] L. F. Mattheis, and D. R. Hamann, *Phys. Rev. B* **33**, 823 (1986).
- [28] P. Blaha, K. Schwarz, P. Soratin, and S. B. Trickey, *Comp. Phys. Comm.* **59**, 399 (1990).
- [29] G. Goldstein, arXiv 2403.15954.
- [30] G. Kotliar, S. Y. Savrasov, K. Haule, V. S. Oudovenko, O. Parcollet, C. A. Marianetti, *Rev. Mod. Phys.* **78**, 865 (2006).