The influence of the Sun and Moon on the observation of very high energy gamma-ray sources using EAS arrays

Tao Wen^{1,2}, Songzhan Chen^{2,3} and Benzhong Dai¹

- ¹ School of Physics and Astronomy, Yunnan University, 650091 Kunming, Yunnan, China; *bzhdai@ynu.edu.cn, wentao@ihep.ac.cn*
- ² Key Laboratory of Particle Astrophyics, Institute of High Energy Physics, Chinese Academy of Sciences, 100049 Beijing, China; *chensz@ihep.ac.cn*
- ³ Tianfu Cosmic Ray Research Center, 610000 Chengdu, Sichuan, China

Received 20xx month day; accepted 20xx month day

Abstract With great advance of ground-based extensive air shower array, such as LHAASO and HAWC, many very high energy (VHE) gamma-ray sources have been discovered and are been monitored regardless of the day and the night. Hence, the Sun and Moon would have some compact on the observation of gamma-ray sources, which have not been taken into account in previous analysis. In this paper, the influence of the Sun and Moon on the observation of very high energy gamma-ray sources when they are near the line of sight of the Sun or Moon is estimated. The tracks of all the known VHE sources are scanned and several VHE sources are found to be very close to the line of sight of the Sun or Moon during some period. The absorption of very high energy gamma-ray by sunlight is estimated with detailed method and some usefully conclusions are achieved. The main influence is the block of the Sun and Moon on gamma-ray and their shadow on the cosmic ray background. The influence is investigated considering the detector angular resolution and some strategy on data analysis are proposed to avoid the underestimation of the gamma-ray emission.

Key words: $\gamma\gamma$ absorption — SUN light — Cosmic Microwave Background

1 INTRODUCTION

Very high energy (VHE, E>0.1 TeV) gamma-rays, located at the highest energy band of the cosmic electromagnetic radiation, are a powerful probe for astrophysics and fundamental physics under extreme conditions. Thanks to the advancements in ground-based gamma-ray detectors, our knowledge about the VHE gamma-ray universe has made impressive progress over the past two decades. The successful operation of the second generation Imaging Atmospheric Cherenkov Telescopes (IACTs), such as H.E.S.S. (Jung 2001), MAGIC (Baixeras 2003), and VERITAS (Weekes et al. 2002), has significantly increased the number of detected VHE gamma-ray sources from about 10 to over 200 since 2004. Due to the limitation of this technique which adopt optical detectors, the observation can only be implemented during the clear and moonless night. Therefore, the influence of the Sun and Moon and their light on the very high energy gamma-ray sources are ignored for a long time.

Recently, the sensitivity of another ground-based detector technique EAS arrays has greatly improved thanks to the successful operation of the new generation arrays, such as HAWC and LHAASO. The HAWC (DeYoung & HAWC Collaboration 2012; Springer 2016) and LHAASO (He & the

LHAASO Collaboration 2018) observatories has detected 65 and 90 VHE sources(Albert et al. 2020; Cao et al. 2023), respectively. Since the EAS arrays adopt particle detectors, they have a large field of view which cover 1/7 of the whole sky at each moment, and they are usually operated with nearly full duty cycle regardless of the day and the night. In such a situation, some of the gamma-ray sources may be very close to the line of sight of the Sun or Moon during observation which would lead to the underestimation of the sources flux. Therefore, the influence of the Sun and the Moon on the very high energy gamma-ray sources near the line of sight should be carefully studied and some strategy is needed to guide the data analysis to avoid the underestimation of the gamma-ray emission.

When a gamma-ray sources is very close to the line of sight of the Sun or the Moon, the measurement of the gamma-ray emission may be affected due to four factors. Firstly, when a source is blocked by the Sun or the Moon, the gamma-rays from the source will not reach the earth. Secondly, when a source is close to the line of sight of the Sun, the gamma-rays from the source will suffer the $\gamma - \gamma$ absorption by the sunlight during their propagation to the earth. The quantitative absorption should depend on the energy of the gamma-rays and also depend on the space angle between source and the Sun. The extinction of the VHE gamma-ray by sunlight has been noticed in a brief paper (Loeb 2022) when studying the TeV gamma-ray background, however, no detailed results are presented and the quantitative impact on gamma-rays from a specific source is not estimated. Thirdly, the Sun and Moon will produce a shadow on the cosmic ray background, which will lead to a deficit source that cancel the gamma-ray signals. These shadows have been well measured by EAS arrays with high significance and were adopted to calibrate detectors and perform physics measurements (Ambrosio et al. 1998, 2003; Aielli et al. 2011; Bartoli et al. 2011, 2017; Aartsen et al. 2021). Fourthly, the Sun could also produce gamma-ray emission due to interaction of cosmic ray with the solar atmosphere. The solar gamma-rays have been well detected and studied using Fermi-LAT data at GeV band (Abdo et al. 2011; Ng et al. 2016; Linden et al. 2018; Tang et al. 2018). The emission at VHE band have been studied by several literature (Albert et al. 2018; Bartoli et al. 2019; Li et al. 2024) and a positive signal was also reported by HAWC collaboration recently (Albert et al. 2023).

In this paper, we will study the influence of the Sun and Moon on the very high energy gamma-ray sources near the line of sight. Especially, the absorption of very high energy gamma-ray by sunlight will quantitative calculated for the first time. Then, we will explore some strategy on data analysis to avoid affection of the Sun and Moon on the measurement of the gamma-ray emission from the sources. The paper is organized as follows: the known VHE gamma-ray sources that may be very close to the Sun and Moon are explored in Section 2. The absorption of very high energy gamma-ray by sunlight is calculated in Section 3 including method and results. Some strategy to avoid the affection of the Sun and Moon are presented in Section 4. Section 5 is the summary.

2 VHE GAMMA-RAY SOURCES THAT COULD BE CLOSE TO THE LINE OF SIGHT OF THE SUN AND MOON

Due to the Earth's orbit around the Sun, the position of the Sun in the celestial coordinates varies annually. Figure 1 shows the track of the Sun in the in celestial coordinates. The declination of the Sun can vary from -23.6° to 23.6° . The track is about same every year. Due to the Moon's orbit around the Earth, the position of the Moon in the celestial coordinates varies with a duration of 29.53 days. Besides this monthly variation, the varying range of the declination also vary with a duration of 18.61 year. The maximum (minimum) declination can change from 18.2° (-18.2°) to 28.7° (-28.7°). The track of the Moon in the celestial coordinates during the 18.61 year is also shows in Figure 1. It is worth to note that both the Sun and Moon are extended source in the sky with a radius about 0.26° .

Up to now, about 300 VHE gamma-ray sources have been detected according to TeVCat¹. All these sources are also presented in the Figure 1. According to the Figure 1, it is clear that some sources can be blocked by the Sun or Moon and some sources would be very close to the Sun or Moon during some period. Table 1 list the 20 known VHE sources that with the closest space angle less than 2° from

¹ http://tevcat.uchicago.edu/



Fig. 1: The track of Sun and Moon in the Equatorial Coordinates. The Points are TeV Sources listed in TeVCat, where the red points are Extragal Sources, the purple points are Galactic Sources, the orange points represent the UNID Sources. The light cyan region is the field of view of LHAASO. The orange line indicates the trajectory of Sun in one year. The skyblue belt is the trajectory of moon in 18.6 years.

the Sun or Moon. The right ascension (R.A.) and declination (Dec) of the sources and their closest space angle to Sun and Moon are presented in the table 1. These sources include both Galactic and extra-galactic sources. Obviously, these sources should be with careful treatment when measure their gamma-ray emission using the EAS array data. The impact of the Sun and the Moon will be studied later.

3 THE ABSORPTION OF VHE GAMMA-RAYS BY SUNLIGHT

It is known that that VHE gamma-rays emitted from distant astronomical sources would be absorbed by low energy background photons through photon-photon interaction. These absorption leads to opacity of the universe to the VHE gamma-rays. The absorption of gamma-rays at a specific energy E is mainly due to the low energy photons with wavelength around $\lambda \sim 1.5(E/1\text{TeV}) \,\mu\text{m}$. The gamma-ray opacity above 100 TeV is mainly due to the absorption f Cosmic Microwave Background (CMB) (Hauser & Dwek 2001). The gamma-ray opacity bellow 100 TeV is mainly due to the absorption of Extragalactic background light (EBL) in the intergalactic space (Finke et al. 2010; Saldana-Lopez et al. 2021). The interstellar radiation fields (ISRF) within our Galaxy would also absorb the gamma-rays bellow 100 TeV with a visible fraction for some cases (Moskalenko et al. 2006). Before a cosmic VHE gamma-ray photon reaches the Earth, it should also pass through the sunlight. Therefore, the gamma-ray should suffer the $\gamma - \gamma$ absorption by the sunlight. The density of sunlight dependent on the distance to the Sun r, which is in inverse proportion to R_{Ω}^2 . The absorption should dependent on the space angle between

Source name	$\mathbf{R}a(^{\circ})$	$Dec(^{\circ})$	closest angle to $Sun(^{\circ})$	closest angle to Moon ($^{\circ}$)
Bouree hume	1(4()	Dee()	closest angle to buil ()	closest angle to Moon ()
HAWC J0543+233	85.78	23.40	0.313	0.325
W 28	270.43	-23.33	0.315	0.375
3C 279	194.05	-5.79	0.513	0.428
HESS J1800-240B	270.11	-24.04	0.601	0.573
RBS 0413	49.95	18.76	0.603	0.410
HESS J1800-240A	270.49	-23.96	0.634	0.301
HESS J1800-240C	269.71	-24.05	0.709	0.436
IC 443	94.21	22.50	0.983	0.354
Crab Pulsar	83.63	22.01	1.307	0.558
Crab	83.63	22.01	1.308	0.560
Terzan 5	266.95	-24.81	1.413	0.563
HESS J1804-216	271.13	-21.70	1.735	0.426
SNR G004.8+6.2	263.35	-21.57	1.737	0.077
2HWC J1309-054	197.31	-5.49	1.765	0.278
Kepler's SNR	262.67	-21.49	1.790	0.478
VER J0521+211	80.44	21.21	1.928	0.241
1ES 0647+250	102.69	25.05	2.157	0.559
OJ 287	133.70	20.10	2.600	0.443
GRB 180720B	0.53	-2.94	2.911	0.442
HESS J1808-204	272.16	-20.43	2.997	0.265

Table 1: The VHE sources with the closest space angle less than 2° from the Sun or Moon

a gamma-ray source and the Sun. In the following, the Crab Nebula will be taken as an example to estimate the absorption by sunlight.

3.1 Estimation method

To quantity estimate the absorption by sunlight, the radiation spectrum from Sun is approximated as a black-body spectrum with a temperature of 5800K. The radiation power is in direct proportion to the surface area of the Sun. The luminosity at Earth is roughly consistent with the measurement achieved by American Society for Testing and Materials (ASTM) (American Society for Testing and Materials 2000), the ASTM E-490 solar spectral irradiance is based on data from satellites, space shuttle missions, high-altitude aircraft, rocket soundings, ground-based solar telescopes, and modeled spectral irradiance, more details can be found at website ², as shown in the Figure 2. The peak luminosity is around 0.5 μ m and the total emission power is 1366.1 W/m^2 at the Earth. The density of sunlight photon $n_{pn}(\nu, r)$ dependent on the photon frequency ν and the distance to the Sun r, which can be written in the form of:

$$n_{ph}(\nu, r) = \frac{2\pi\nu^2 R_{\odot}^2}{c^3 r^2} \frac{1}{e^{h\nu/KT} - 1}$$
(1)

where c is the speed of light, R_{\odot} =6.963×10¹⁰ cm is the radius of the Sun, T is the temperature, K is the Boltzmann constant.

The cross section of the γ - γ absorption presented in (Dermer & Menon 2010) is adopted here:

$$\sigma_{\gamma\gamma}(s) = \frac{1}{2}\pi r_e^2 (1 - \beta_{cm}^2) \left[(3 - \beta_{cm}^4) ln \left(\frac{1 + \beta_{cm}}{1 - \beta_{cm}} \right) - 2\beta_{cm} (2 - \beta_{cm}^2) \right]$$
(2)

where r_e is the classical electron radius, $\beta_{cm} = \sqrt{1 - s^{-1}}$, $s = \frac{1}{2}[E\epsilon(1 - cos(\theta))]$ is the center-ofmomentum frame Lorentz factor of the produced electron and position, E is the energy of the gammaray, ϵ is the energy of the low energy photon, θ is the space angle between the direction of the low energy photon and that of the gamma-ray.

² https://www.pveducation.org/pvcdrom/appendices/standard-solar-spectra



Fig. 2: The luminosity of sunlight as the function of photon wavelength. The black line is the expected luminosity at Earth from a black-body spectrum with a temperature of 5800K at the surface of the Sun. The red line indicates the measurement achieved by American Society for Testing and Materials (ASTM).



Fig. 3: A schematic diagram shows the interaction between the gamma-rays from Crab Nebula with sunlight and the parameters used in the equations.



Fig. 4: The survival fractions the gamma-ray after the absorption of the sunlight. Different lines indicate the results using different space angle between gamma-ray source and the Sun.

With the density of sunlight photon and the cross section of the γ - γ , the optical depth of the gammaray from the Crab Nebula to Earth due to the absorption of the sunlight can be estimated using the function:

$$\tau_{\gamma\gamma,\text{SunLight}} = \frac{1}{2} \int dx \int_{1/\epsilon}^{\infty} (1 - \cos(\theta)) n_{ph}(\epsilon, r) \sigma_{\gamma\gamma}(s) d\epsilon \tag{3}$$

Where dx is integrated along the line from the Crab Nebula to the Earth as illustrated in the schematic diagram shown in Figure 3. In this figure, R_o =100 AU, and the β is the space angle between the Sun and the Crab Nebula. In the calculation, We divide the integration interval into two segments, i.e., with r less than R_o and r larger than R_o . When $r < R_o$, the angle between the direction of the sun-light photon and the direction the gamma-ray change rapidly. When the $r > R_o$, the collision angle will be considered as consistent value $\phi = 180^{\circ}$.

3.2 Estimation result

Using the equation (3), we calculate the optical depth (τ) of gamma-ray at different energies for different space angle between the Sun and the Crab Nebula. The survival fractions the gamma-ray $e^{-\tau}$ are shown in the Figure 4. The absorption is clearly dependent on the gamma-ray energy and the space angle between gamma-ray and the Sun. According to the Figure 4, the absorption of gamma-ray happen at a wide energy range from about 0.1 TeV to 100 TeV with maximum absorption approximately at 0.89 TeV. When the space angle between Crab Nebula and Sun is 0.25°, which is at the boundary of the Sun, the maximum absorbed percentage is 1.2%. It is worth to note that we have ignored the absorption due to the interaction between the gamma-ray and the solar atmosphere, which should be heavy when the



Fig. 5: Upper: The space angle between the Sun and Crab Nebula as function of the time in one year. Down: The absorption of the sunlight on gamma-rays with different energies, i.e., 0.2, 0.4, 0.89, 5, and 25 TeV, as function of the time in one year.

space angle is less than 0.25° . When the space angle is 1.29° , the maximum absorption ratio is 0.25%. The maximum absorption is less than 0.05% when the space angle is larger than 5° .

Due to the Earth's orbit around the Sun, the position of the Sun in the celestial coordinates varies annually. Hence the space angle between the Sun and Crab Nebula varies annually as shown in Figure 5, with maximum angle larger than 150° and minimum around 1.29°. We also estimation of the absorption of the sunlight for gamma-rays with different energies from the Crab Nebula as the function the time as shown in Figure 5. Since the minimum space angle between Crab Nebula and Sun is 1.29° occurred in the day of 3th July, the maximum absorption is less than 0.25%.

3.3 Cross check about the calculation procedure

When gamma-rays from the Crab Nebula travel towards the Earth, they are not only absorbed by lowenergy sunlight photons, but also by cosmic microwave background (CMB) photons. The absorption process has been calculated in (Cao et al. 2021). In order to validate the reliability of the calculation



Fig. 6: The survival fractions the gamma-ray after the absorption by the sunlight and the CMB photons. Different lines indicate the results using different space angle between gamma-ray source and the Sun.The yellow region represents the absorption of sunlight, while the sky-blue region mainly corresponds to the absorption of CMB.

procedure presented in this paper, we have used the same computational procedures for sunlight photons to calculate the absorption of gamma-rays from the direction of the Crab Nebula by the CMB. In our calculations, the CMB radiation spectrum adopts a 2.7K blackbody spectrum, and the CMB density is isotropic and position-independent, in contrast to the density of sunlight photons. The density of CMB is:

$$n_{ph,cmb}(\nu,t) = \frac{8\pi\nu^2}{c^3} \frac{1}{\exp\left(\frac{h\nu}{KT}\right) - 1}$$
(4)

For the Crab, we don't need to consider the evolution of the universe, the the optical depth of CMB is:

$$\tau_{\gamma\gamma,cmb} = \frac{1}{2} \int dx \int_{-1}^{1} (1-\mu)d\mu \int_{1/\epsilon}^{\infty} n_{cmb}(\epsilon)\sigma_{\gamma\gamma}(s)d\epsilon$$
(5)

where $\mu = 1 - \cos(\theta)$. The total survival ratio of gamma-rays as a function of energy is $e^{-\tau} = e^{-(\tau_{cmb} + \tau_{Sunlight})}$, as shown in Figure 6. The absorption of CMB on gamma-rays is primarily observed at energies above 200 TeV, with the maximum absorption occurring at around 2 PeV, and an absorption probability of 25%. This result aligns with the findings in reference (Cao et al. 2021), thus providing basic validation of the correctness of the calculations in this paper.

4 THE AFFECTION OF THE SUN AND MOON ON GAMMA-RAY SOURCE OBSERVATION

Based on the calculations from the previous section, it is evident that the absorption of gamma-rays by sunlight is extremely minimal and can be disregarded during most observation periods. The Moon primarily reflects sunlight and has a much lower brightness compared to the Sun, resulting in an even smaller absorption of gamma-rays by moonlight, which can be directly ignored. However, when the angle between the direction of the gamma-ray source and the directions of the Sun and Moon is less than 0.25° , the gamma-rays will be completely blocked by the Sun and Moon, and cannot be ignored during this time period. Therefore, it is necessary to exclude this data during the analysis.

Additionally, when the angle between the direction of the gamma-ray source and the directions of the Sun and Moon is small, the Sun and Moon directly block the cosmic ray background, creating a missing negative source. This shadow can counteract the gamma-ray signal, and its impact depends on the ratio of the number of gamma-ray signals to the background. For previous arrays like ARGO-YBJ, which lacked the ability to distinguish between gamma-ray and cosmic ray background, the gamma-ray signals only accounted for a small portion of the backgrounds. For example, in observations of the brightest gamma-ray source, the Crab Nebula, the gamma-ray signals were only about 1% of the backgrounds (Bartoli et al. 2015), leading to a significant counteracting effect from the shadow on the gamma-ray signals.

For the LHAASO array, due to its excellent ability to distinguish between gamma-rays and cosmic ray backgrounds, the proportion of gamma-rays to backgrounds has significantly increased. For example, in observations of the Crab Nebula, the ratio of gamma-ray signals to backgrounds for the LHAASO-WCDA array is 20% at 1 TeV, 80% at 6 TeV(Aharonian et al. 2021a), and for the LHAASO-KM2A array, the corresponding ratios are 12% at 10 TeV, 260% at 25 TeV, and 4100% at 100 TeV(Aharonian et al. 2021b). However, considering the current detection sensitivity of LHAASO is at 1% of the Crab flux level, the impact at energies bellow 100 TeV cannot be ignored.

Fig.7 shows the counteracting effect of Sun or Moon shadow on the gamma-ray signals at different space angles. In this example, we assume the detector angular resolution to be 0.5° and the number of deficit shadow events to be three times that of the gamma-ray signals. This situation is similar to the observation of LHAASO-KM2A on a gamma-ray source with a flux around 10% Crab unit at 10 TeV. It is clear that the gamma-ray signals will be completely counteracted by the shadow when they are very close the direction of Sun or Moon. The counteracting effect decreases as the increasing of the space angle. To ensure safety, we recommend removing data when the space angle between the gamma-ray source and the directions of the Sun and Moon is less than 3 times the angular resolution plus 0.25° . For example, the observation time of the EAS array for Crab is approximately 7 hours per day, and only during a few days the space angle between Crab and Sun or Moon is less than 2° in one year. The data from this period of observation can be excluded for data analysis, thus eliminating its impact. Since the angle between the source and the directions of the Sun and Moon varies periodically, this portion of the data accounts for a very small percentage, generally less than 1% of the total observation time in one year, and therefore does not significantly affect the observation of the source.

Table 1 lists known gamma-ray sources with a minimum angle less than 2° between the source and the Sun or Moon. When measuring their radiation using EAS arrays such as LHAASO and HAWC, special attention should be paid to excluding data when the space angle between the source and the Sun or Moon is small.

5 SUMMARY

With significant advancements in ground-based extensive air shower arrays, numerous very high-energy (VHE) gamma-ray sources have been discovered and are being continuously monitored by LHAASO and HAWC, regardless of day or night. With investigating the celestial coordinates of the Sun and Moon, we identified 20 sources that can come close to the line of sight of the Sun or Moon during some certain periods. We have conducted the first estimation of sunlight absorption on very high-energy



Fig. 7: The impact of the shadow caused by the Sun or Moon on the observation of gamma-ray sources at different space angles, i.e., 0.25°, 2*PSF+0.25°, 3*PSF+0.25°, and 4*PSF+0.25°. The detector angular resolution (denoted as PSF) is assumed to be 0.5°. The number of deficit shadow events is assumed to be three times that of the gamma-ray signals from the source.

gamma-rays. The absorption primarily affects gamma-rays with energies ranging from 0.1 TeV to 100 TeV, with maximum absorption occurring around 0.9 TeV. The absorption is contingent on the spatial angle between the Sun and the sources; however, the overall absorption is minimal, with a maximum of less than 1.2%. The primary influence is the obstruction of the Sun and Moon on gamma-rays, and their shadow on the cosmic ray background. The impact is also contingent on the detector's angular resolution. To mitigate its effects, we recommend excluding data when the angle between the gamma-ray source and the directions of the Sun and Moon is less than a certain value taking into account the experiment angular resolution.

Acknowledgements This work is supported by the National Science Foundation of China under Grant Nos.12393854, 12022502, 12263007.

References

Aartsen M. G., Abbasi R., Ackermann M. et al. 2021, Phys. Rev. D, 103, 042005 2 Abdo A. A., Ackermann M., Ajello, M. et al. 2011, ApJ, 734, 116 2 Aharonian, F., An, Q., Axikegu, et al. 2021a, Chinese Physics C, 45, 085002 9 Aharonian, F., An, Q., Axikegu, et al. 2021b, Chinese Physics C, 45, 025002 9 Aielli, G., Bacci, C., Bartoli, B., et al. 2011, ApJ, 729, 113 2 Albert, A., Alfaro, R., Alvarez, C., et al. 2018, Phys. Rev. D, 98, 123011 2

- Albert, A., Alfaro, R., Alvarez, C. et al. 2020, ApJ, 905, 76 2
- Albert A., Alfaro R., Alvarez C. et al. 2023, Phys. Rev. Lett., 131, 051201 2
- Ambrosio, M., Antolini, R., Aramo, C., et al. 1998, Phys. Rev. D, 59, 012003 2
- Ambrosio, M., Antolini, R., Baldini, A., et al. 2003, Astroparticle Physics, 20, 145 2
- American Society for Testing and Materials. 2000, Standard Solar Constant and Zero Air Mass Solar Spectral Irradiance Tables, Tech. Rep. E490, ASTM International 4
- Baixeras, C. 2003, Nuclear Physics B Proceedings Supplements, 114, 247 1
- Bartoli, B., Bernardini, P., Bi, X. J., et al. 2011, Phys. Rev. D, 84, 022003 2
- Bartoli, B., Bernardini, P., Bi, X. J. et al. 2015, ApJ, 798, 119 9
- Bartoli, B., Bernardini, P., Bi, X., et al. 2017, Astroparticle Physics, 90, 20 2
- Bartoli, B., Bernardini, P., Bi, et al. 2019, ApJ, 872, 143 2
- Cao, Z., Aharonian, F. A., An, Q., et al. 2021, Nature, 594, 33 7, 8
- Cao, Z., Aharonian, F., An, Q., et al. 2023, arXiv e-prints, arXiv:2305.17030 2
- Dermer, C. D., & Menon, G. 2010, High Energy Radiation from Black Holes. Gamma Rays, Cosmic Rays, and Neutrinos 4
- DeYoung, T., & HAWC Collaboration. 2012, Nuclear Instruments and Methods in Physics Research A, 692, 72 1
- Finke, J. D., Razzaque, S., & Dermer, C. D. 2010, ApJ, 712, 238 3
- Hauser, M. G., & Dwek, E. 2001, ARA&A, 39, 249 3
- He, H., & the LHAASO Collaboration, F. 2018, Radiation Detection Technology and Methods, 2, 7 1
- Jung, I. 2001, H.E.S.S. The High Energy Stereoscopic System, ed. M. M. Shapiro, T. Stanev, & J. P. Wefel (Dordrecht: Springer Netherlands), 345–348 1
- Li, Z., Ng, K. C. Y., Chen, S., Nan, Y., & He, H. 2024, Chinese Physics C, 48, 045101 2
- Linden, T., Zhou, B., Beacom, J. F., et al. 2018, Phys. Rev. Lett., 121, 131103 2
- Loeb, A. 2022, Research Notes of the American Astronomical Society, 6, 148 2
- Moskalenko, I. V., Porter, T. A., & Strong, A. W. 2006, ApJ, 640, L155 3
- Ng, K. C. Y., Beacom, J. F., Peter, A. H. G., & Rott, C. 2016, Phys. Rev. D, 94, 023004 2
- Saldana-Lopez, A., Domínguez, A., Pérez-González, P. G., et al. 2021, MNRAS, 507, 5144 3
- Springer, R. 2016, Nuclear and Particle Physics Proceedings, 279-281, 87 1
- Tang, Q.-W., Ng, K. C. Y., Linden, T., et al. 2018, Phys. Rev. D, 98, 063019 2
- Weekes, T. C., Badran, H., Biller, S. D., et al. 2002, Astroparticle 1