

# Quantum Many-Body Scarring in a Non-Abelian Lattice Gauge Theory

Giuseppe Calajó<sup>1,\*</sup>,<sup>1</sup> Giovanni Cataldi<sup>1,2,3,\*</sup>,<sup>1,2,3</sup> Marco Rigobello<sup>1,2,3</sup>,<sup>1,2,3</sup> Darvin Wanisch<sup>1,2,3</sup>,<sup>1,2,3</sup> Giuseppe Magnifico<sup>4,5</sup>,<sup>4,5</sup> Pietro Silvi<sup>1,2,3</sup>,<sup>1,2,3</sup> Simone Montangero<sup>1,2,3</sup>,<sup>1,2,3</sup> and Jad C. Halimeh<sup>6,7,8,9</sup>

<sup>1</sup>*Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Padova, I-35131 Padova, Italy.*

<sup>2</sup>*Dipartimento di Fisica e Astronomia “G. Galilei”, Università di Padova, I-35131 Padova, Italy.*

<sup>3</sup>*Padua Quantum Technologies Research Center, Università degli Studi di Padova*

<sup>4</sup>*Dipartimento di Fisica, Università di Bari, I-70126 Bari, Italy.*

<sup>5</sup>*Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Bari, I-70125 Bari, Italy.*

<sup>6</sup>*Department of Physics and Arnold Sommerfeld Center for Theoretical Physics (ASC), Ludwig Maximilian University of Munich, 80333 Munich, Germany*

<sup>7</sup>*Max Planck Institute of Quantum Optics, 85748 Garching, Germany*

<sup>8</sup>*Munich Center for Quantum Science and Technology (MCQST), 80799 Munich, Germany*

<sup>9</sup>*Dahlem Center for Complex Quantum Systems, Free University of Berlin, 14195 Berlin, Germany*

(Dated: May 27, 2024)

Quantum many-body scarring (QMBS) is an intriguing mechanism of ergodicity breaking that has recently spurred significant attention. Particularly prominent in Abelian lattice gauge theories (LGTs), an open question is whether QMBS nontrivially arises in non-Abelian LGTs. Here, we present evidence of robust QMBS in a non-Abelian SU(2) LGT with dynamical matter. Starting in product states that require little experimental overhead, we show that prominent QMBS arises for certain quenches, facilitated through meson and baryon-antibaryon excitations, highlighting its non-Abelian nature. The uncovered scarred dynamics manifests as long-lived coherent oscillations in experimentally accessible local observables as well as prominent revivals in the state fidelity. Our findings bring QMBS to the realm of non-Abelian LGTs, highlighting the intimate connection between scarring and gauge symmetry, and are amenable for observation in a recently proposed trapped-ion qudit quantum computer.

**Introduction.**—The nature of equilibration in an isolated interacting quantum many-body system is a fundamental question in physics [1]. Whereas open quantum systems are expected to thermalize due to exchanging energy with a bath [2], the conditions under which an isolated quantum many-body system thermalizes are stipulated within the Eigenstate Thermalization Hypothesis (ETH) [3–6]. An *ergodic* Hamiltonian is expected to lead to *quantum thermalization* for generic initial states, where the information of the initial state is locally erased [7].

In recent years, a new paradigm of ergodicity breaking has emerged that violates the ETH. In certain ergodic Hamiltonians, there is a polynomial (in system size) number of special non-thermal *scar* eigenstates, that are roughly equally spaced in energy over the whole spectrum [8–10], and that exhibit anomalously low bipartite entanglement entropy [11]. Upon initializing the system in an initial state with a high overlap with these scar eigenstates, the subsequent quench dynamics give rise to *quantum many-body scarring* (QMBS), which manifests as prominent oscillations in local observables and persistent revivals in the (local and global) Loschmidt fidelity that last longer than all relevant timescales, thereby circumventing expected thermalization [12–18]. QMBS has received a lot of attention since its initial discovery in

[12], and has since been the subject of many quantum simulation experiments [19–23].

After the discovery of QMBS, it has been shown that the effective model quantum-simulated in [12] is a spin-1/2 U(1) lattice gauge theory (LGT), where the electric field is represented by a spin-1/2  $z$ -operator [24]. The robustness of QMBS also depends on the stability of the underlying gauge symmetry in this model [25]. QMBS also arises in spin- $S$  U(1) LGTs [26, 27], in LGTs with other Abelian gauge groups [28–30], and in 2 + 1D LGTs [31–36]. However, all of these works involve Abelian gauge groups except the recent Ref. [33], which considers a pure (without dynamical matter) SU(2) LGT. Looking at the bipartite entanglement entropy, they identify a few scar-eigenstate candidates in a certain parameter regime. However, no experimentally feasible scarred initial states have been proposed, and the type of scarring exhibited vanishes beyond the crudest truncation of the electric field basis.

Given the intimate connection between QMBS and LGTs, and the current large effort to quantum simulate the latter [37–47], it is important to investigate the origin of QMBS in connection to gauge symmetry by exploring its possible occurrence in non-Abelian LGTs with dynamical matter. In particular, to faithfully investigate a high-energy context of QMBS, including dynamical matter is crucial, since nature hosts dynamical matter fields. In this Letter, we provide exact diagonalization (ED) and matrix

\* These authors contributed equally to this work.

product state (MPS) results that showcase robust QMBS dynamics in a non-Abelian SU(2) LGT with dynamical matter starting in simple initial product states. We show how the QMBS dynamics involve state transfer through meson and baryon–anti-baryon bare states, highlighting the non-Abelian nature of this scarred dynamics.

**Model.**—As a prototypical model to probe non-ergodic dynamics in non-Abelian LGTs, we consider a 1 + 1D matter-coupled hardcore-gluon [48–53] SU(2) LGT. The dynamics is governed by the Kogut-Susskind Hamiltonian, on an  $N$ -site lattice with spacing  $a_0$  ( $c, \hbar = 1$ ) [54]:

$$\hat{H}_0 = \frac{1}{2a_0} \sum_{n=1}^N \sum_{a,b \in \{r,g\}} \left[ -i\hat{\psi}_{na}^\dagger \hat{U}_{n,n+1}^{ab} \hat{\psi}_{n+1,b} + \text{H.c.} \right] + m_0 \sum_{n,a} (-1)^n \hat{\psi}_{na}^\dagger \hat{\psi}_{na} + \frac{a_0 g_0^2}{2} \sum_n \hat{E}_{n,n+1}^2. \quad (1)$$

The model describes a matter quark field of mass  $m_0$ , living on lattice sites  $n$ , coupled, with strength  $g_0$ , to a truncated SU(2) gauge field, defined on lattice links — see Fig. 1. The quark field is a staggered fermion doublet  $\hat{\psi}_{na}$ , obeying anticommutation relations  $\{\hat{\psi}_{na}, \hat{\psi}_{n'b}^\dagger\} = \delta_{n,n'} \delta_{a,b}$  and  $\{\hat{\psi}_{na}, \hat{\psi}_{n'b}\} = 0$  [55]. A basis for matter sites is  $\{|0\rangle, |r\rangle = \hat{\psi}_r^\dagger |0\rangle, |g\rangle = \hat{\psi}_g^\dagger |0\rangle, |2\rangle = \hat{\psi}_r^\dagger \hat{\psi}_g^\dagger |0\rangle\}$ . Gauge link states can be expanded in the basis  $|j, m_L, m_R\rangle$ , where  $m_L, m_R \in \{-j, \dots, +j\}$  index spin- $j$  states, and the link energy density operator is diagonal and coincides with the quadratic Casimir,  $\hat{E}^2 |j, m_L, m_R\rangle = j(j+1) |j, m_L, m_R\rangle$  [56]. Adopting a hardcore-gluon truncation, we restrict  $j$  to  $\{0, 1/2\}$ , namely keep only basis states reachable from the singlet  $|00\rangle$  via at most a single application of the parallel transporter  $\hat{U}^{ab}$ :  $\{|00\rangle, |rr\rangle, |gg\rangle, |gr\rangle, |rg\rangle\}$ , where  $r, g$  are short-hands for  $\pm 1/2$  and  $j$  was left implicit.

Non-Abelian Gauss law mandates that physical states are local gauge singlets,  $\hat{G}_n^\nu |\Psi_{\text{phys}}\rangle \equiv 0$ ,  $\forall n, \nu \in \{x, y, z\}$ , where  $\hat{G}_n^\nu$  are the generators of local rotations at site  $n$ . In numerical simulations, we enforce this constraint using a dressed site approach, which yields a defermionized qudit model with a 6-dimensional local basis [51]. Its derivation is detailed in the Supplemental Material [48]. In the qudit model, it is convenient to remove the explicit dependence on  $a_0$  by rescaling the energy, mass, and coupling in terms of the dimensionless quantities  $(\hat{H}, m, g^2) = 4\sqrt{2}a_0(\hat{H}_0, m_0, 3a_0g_0^2/16)$ . Values reported in the Figures refer to these rescaled quantities.

**Non-Abelian scarred dynamics.**—To investigate the presence of non-ergodic behavior in this model, we consider the Schrödinger time evolution of different initial states  $|\Psi(t=0)\rangle$ , across various parameter regimes. The time evolution is performed on a chain of  $N = 30$  sites via MPS methods. The details of the simulation are reported in [48]. As initial configuration, we first consider the *polarized bare vacuum* (PV), which consists of the matter sites in the bare vacuum configuration, while the gauge

sites in the excited electric field state,

$$|\Psi_{\text{PV}}\rangle = |\dots 2020 \dots\rangle_{\text{m}} \left| \dots \frac{rg - gr}{\sqrt{2}} \dots \frac{rg - gr}{\sqrt{2}} \dots \right\rangle_{\text{g}}, \quad (2)$$

where  $|\cdot\rangle_{\text{m}}$  and  $|\cdot\rangle_{\text{g}}$  refer to the matter and gauge subsystems, respectively. Here  $(m_R)_{n-1,n}$  and  $(m_L)_{n,n+1}$  around each bulk site  $n$  are valence-bonded in a singlet. In the qudit formulation of the model, this state becomes a product state [48].

We identify two regimes avoiding thermalization: the large-mass one,  $m \gg (g^2, 1)$ , and the large-coupling one,  $g^2 \gg (m, 1)$ . For each regime, we compute the return fidelity between the initial state and the evolved state,  $\mathcal{F}(t) = |\langle \Psi(t) | \Psi(0) \rangle|^2$ , shown in Fig. 1. After an initial complete system relaxation, we observe persistent revivals, signaling a clear deviation from the expected ergodic behavior. The revivals occur approximately periodically, with the period indicated in the figure for both the considered regimes.

The observed dynamics can be understood as a process of persistent particle pair creation out of an initial false vacuum [57–59]. To see this, we define the site occupancy operator,

$$\hat{Q}_n = (-1)^n \sum_a \left[ \hat{\psi}_{na}^\dagger \hat{\psi}_{na} - \frac{1 - (-1)^n}{2} \right], \quad (3)$$

which counts the number of quarks (on even sites) or antiquarks (on odd sites). Then, we compute the average occupancy  $\rho_q(t)$  of each  $\hat{Q}_n$  eigenvalue (single site quark occupancy,  $q \in \{0, 1, 2\}$ ), projecting on the corresponding eigensubspace of  $\hat{Q}_n$  and averaging over all sites  $n$ :

$$\rho_q(t) = \frac{1}{N} \sum_n \langle \Psi(t) | \delta_{\hat{Q}_n, q} | \Psi(t) \rangle. \quad (4)$$

They oscillate around average values that do not coincide with the ones predicted by thermal ensembles, confirming deviations from ergodic behavior, as detailed in [48]. Large values of either mass or coupling confine the system dynamics close to the initial state, creating few particle pair excitations in the system. In the large-mass regime, most of the excitations are meson-like (quark and antiquark adjacent pairs with an excited shared gauge link). In the large-coupling regime, the formation of pairs of baryon- (quark pair) and antibaryon- (antiquark pair) like excitations becomes likely as well, as shown by the enhanced double occupancy with respect to the large-mass regime. Finally, the last row of Fig. 1 shows the evolution of the bipartite entanglement entropy defined as  $S = -\text{Tr}[\hat{\rho}_A \log \hat{\rho}_A]$ , where  $A$  and  $B$  indicate the two halves of the chain and  $\hat{\rho}_A = \text{Tr}_B[\hat{\rho}_{AB}]$  is the density operator reduced to the  $B$  subsystem. In both cases, we observe a slow growth of the entanglement entropy after an initial fast increase up to the first fidelity revival peak. The oscillations on top of the growth are driven by the




FIG. 1. Many-body scarring dynamics of the polarized bare vacuum (left) and the bare vacuum (right) initial states. *Top*: cartoons depicting the classical configurations that give the leading contribution to the dynamics; with circles and ellipses denoting matter sites and gauge links respectively. *Bottom*: return fidelity, average quark occupancy, and bipartite entanglement entropy as a function of time, for different mass and coupling regimes, reported in Figure. MPS simulation with  $N = 30$  sites in open boundary conditions (OBC), maximum bond dimension  $D_{\max} = 350$ , and truncation tolerance  $\text{tol} = 10^{-7}$  [48].

successive fidelity revivals, which indeed have the same period.

In addition to the polarized bare vacuum initial state, we also observe similar scarred-like dynamics for the *bare vacuum* (V) state: the ground state of the Hamiltonian in Eq. (1) without the hopping term, made of alternated doubly occupied and empty sites, and trivial links,

$$|\Psi_V(t=0)\rangle = |\dots 2020 \dots\rangle_m |\dots 0000 \dots\rangle_g. \quad (5)$$

The resulting non-ergodic dynamics is also shown in Fig. 1. Compared to the polarized vacuum case, the oscillation's period in the fidelity revivals and occupancy is smaller. It can be explained by the many-body spectrum as discussed in the following section. The large coupling regime is characterized by a faster entanglement growth, consistently with a corresponding lower return fidelity. Although in the regimes considered this product state is energetically close to the true ground state of the model, the observed dynamics is driven by nontrivial physics. This is evident from the significant oscillations observed in the average occupancy, which lead the system significantly away from its initial configuration.

For both initial states, non-ergodic dynamics persists also at longer times and remains robust with increasing

system size, as we discuss in [48]. Importantly, this behavior is not observed for other initial state configurations or different coupling-mass regimes [48], suggesting that the non-ergodicity originates directly from many-body scars in the spectrum. In the following, we demonstrate that is indeed the case by analyzing the full many-body spectrum of the model.

**Tower of scar states.**—To determine whether the observed non-ergodic dynamics originates from quantum many-body scars, we perform ED up to  $N = 10$  sites and, for each eigenstate in the many-body spectrum  $\{|\Phi_s\rangle\}$ , we compute the overlap  $\mathcal{O}_{PV/V} = |\langle \Psi_{PV/V} | \Phi_s \rangle|^2$  and the bipartite entanglement entropy  $S$  with the two considered initial states (polarized, PV, and unpolarized bare vacua, V). These quantities are plotted in Fig. 2, for both regimes where we observe revivals in the return fidelity. For each candidate QMBS regime, we find a ‘tower of scar states’ (highlighted by red circles) characterized by a high overlap with the initial product states and entanglement low enough (compared to the rest of the many-body spectrum) to give rise to scarring behavior. In each case, the energy gap between the scar states in the tower,  $\Delta E$  (reported in Fig. 2), approximately matches the frequency of the revivals observed in the return fidelity from Fig. 1,




FIG. 2. Spectrum analysis for a lattice chain of  $N=10$  sites in OBC, in the two parameter regimes of Fig. 1. Red circles highlight the scar states. *First row*: the overlap of the many-body spectrum with the polarized bare vacuum (left) and the bare vacuum (right); only states with  $\mathcal{O}_{(P)V} > 10^{-9}$  are included. *Second row*: bipartite entanglement entropy of each eigenstate.

$\Delta E \approx 2\pi/T$ . The correspondence between spectral and dynamical features leads us to conclude that QMBS is indeed the underlying mechanism behind the non-ergodic dynamics shown in Fig. 1.

**Ergodic vs non-ergodic dynamics.** — To confirm that the observed non-ergodic dynamics is indeed due to scarring, we should observe thermal relaxation for other initial states at the same energy. To verify this, we focus on the case of the polarized vacuum state and evolve the *microcanonical* ensemble (ME) state constructed as a uniform coherent superposition of all eigenstates lying within a small energy window around the quench energy  $E_{PV} = \langle \Psi_{PV} | \hat{H} | \Psi_{PV} \rangle$ :

$$|\Psi_{PV}^{\text{ME}}\rangle = \frac{1}{\sqrt{N_{E_{PV}, \delta E}}} \sum_{s, |E_s - E_{PV}| < \delta E} |\Phi_s\rangle, \quad (6)$$

where  $\delta E = \sqrt{\langle \Psi_{PV} | \hat{H}^2 | \Psi_{PV} \rangle - E_{PV}^2}$  defines the energy shell  $[E_{PV} - \delta E, E_{PV} + \delta E]$ , and  $N_{E_{PV}, \delta E}$  is the number of eigenstates in this energy shell [7]. Figure 3 compares the evolution of this state to that of  $|\Psi_{PV}\rangle$ , in the identified scarred regimes. As the plot illustrates, the *microcanonical* state exhibits ergodic behavior: return fidelities relax toward zero (up to minor oscillations attributable to the finite lattice size) and the single occupancy remains close to its microcanonical ensemble average  $\langle \rho_1 \rangle_{\text{ME}}$  [7, 48]. In contrast, for the polarized vacuum,  $\rho_1$  widely oscillates around its long-time average  $\langle \rho_1 \rangle_{\text{DE}}$ , whose value is predicted by the diagonal ensemble (DE) and largely deviates from the thermal result (see [48]). We thus conclude that the observed non-ergodic dynamics strictly depends on the initial conditions and escapes an otherwise thermal behavior through the occurrence of scar states in the many-body spectrum.

**Summary and outlook.** — In this letter, we presented strong numerical evidence for the occurrence of quantum




FIG. 3. Dynamics of the *microcanonical* state in Eq. (6) for the same parameter regimes of Fig. 1. Gray curves reproduce the polarized bare vacuum dynamics, for comparison. The blue horizontal lines reproduce the thermal  $\rho_1$  occupancy, obtained via microcanonical ensemble (ME). The black ones show the diagonal ensemble (DE) results for the long-time average of  $\rho_1$  on  $|\Psi_{PV}(t)\rangle$ , highlighting its lack of thermalization [48]. ED results for  $N=10$  sites in PBC.

many-body scars in a non-Abelian  $SU(2)$  lattice gauge theory with dynamical matter. In particular, starting from simple experimentally friendly vacuum product states, scarred dynamics manifests as persistent oscillations in local observables and periodic revivals in the state fidelity. This dynamics arises from a state transfer through meson and baryon-antibaryon bare states, which allow the system to escape otherwise expected thermalization. Even though the results were obtained for a non-Abelian lattice gauge theory model with the gauge-link truncated at the lowest nontrivial irreducible representations, in the



Supplemental Material [48] we provide evidence that the observed scar dynamics persists even at higher gauge-link truncations, similar to what has recently been observed for Abelian models [27], hinting at the possible survival of quantum many-body scarring in the Kogut-Susskind limit of the SU(2) model under consideration. These findings demonstrate that the strong connection between QMBS and LGTs persists even for non-Abelian gauge groups, raising the intriguing question of whether such behavior represents an intrinsic feature of more fundamental theories such as quantum chromodynamics.

Our work opens the door to exciting future directions: In the SM [48], we present numerical evidence of scar eigenstates with very low bipartite entanglement entropy throughout the entire spectrum even in regimes where both  $m$  and  $g$  are quite small. It would be interesting to find simple product states showing large overlap with these eigenstates. Additionally, it would be interesting to map the scarring phase diagram for this model as has been done for the PXP model [60], and to see how driving protocols can enhance non-Abelian scarring [61]. Another venue to pursue is exploring the robustness of the non-Abelian scarring uncovered in this work in two spatial dimensions, as it has been recently done for Abelian gauge theories [35, 62]. This can shed further light on the nature of this non-Abelian scarring and its possible occurrence in the quantum-field-theory limit. Furthermore, scarring is known to give rise to complex periodicities in the corresponding dynamical quantum phase transitions of Abelian LGTs [63, 64], and it would be interesting to see if this carries over to the non-Abelian case.

Finally, we stress that, as addressed in Ref. [51], the qudit formulation of the proposed model is well-suited for conducting a digital quantum simulation on a recently demonstrated trapped-ion qudit quantum processor [65]. This opens up the intriguing possibility to experimentally observe this exotic non-ergodic dynamics in the near future.

**Acknowledgments.**—The authors thank Anthony Ciavarella, Jean-Yves Desaules, and Paolo Stornati for valuable discussions. They acknowledge financial support from: the European Union via QuantERA2017 project QuantHEP, via QuantERA2021 project T-NiSQ, via the Quantum Technology Flagship project PASQuanS2, and via the NextGenerationEU project CN00000013 - Italian Research Center on HPC, Big Data and Quantum Computing (ICSC); the Italian Ministry of University and Research (MUR) via PRIN2022-PNRR project TANQU, via Progetti Dipartimenti di Eccellenza project Frontiere Quantistiche (FQ) and project Quantum Sensing and Modelling for One-Health (QuaSiModO); the Max Planck Society and the Emmy Noether Programme of the German Research Foundation (DFG) via grant no. HA 8206/1-1; the WCRI-Quantum Computing and Simulation Center (QCSC) of Padova University; the University of Bari through the 2023-UNBACLE-0244025 grant the Istituto

Nazionale di Fisica Nucleare (INFN) via project NPQCD and project QUANTUM. The authors also acknowledge computational resources by Cloud Veneto and the ITensors Library for the MPS calculations [66, 67] as well as the ed-igt Library for Exact Diagonalization [68].

- 
- [1] P. Reimann, *Phys. Rev. Lett.* **101**, 190403 (2008).
  - [2] I. Reichental, A. Klemmner, Y. Kafri, and D. Podolsky, *Phys. Rev. B* **97**, 134301 (2018).
  - [3] J. M. Deutsch, *Phys. Rev. A* **43**, 2046 (1991).
  - [4] M. Srednicki, *Phys. Rev. E* **50**, 888 (1994).
  - [5] L. D’Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, *Advances in Physics* **65**, 239 (2016).
  - [6] J. M. Deutsch, *Reports on Progress in Physics* **81**, 082001 (2018).
  - [7] M. Rigol, V. Dunjko, and M. Olshanii, *Nature* **452**, 854 (2008).
  - [8] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papić, *Nature Physics* **14**, 745 (2018).
  - [9] S. Moudgalya, N. Regnault, and B. A. Bernevig, *Phys. Rev. B* **98**, 235156 (2018).
  - [10] M. Schecter and T. Iadecola, *Phys. Rev. Lett.* **123**, 147201 (2019).
  - [11] C.-J. Lin and O. I. Motrunich, *Phys. Rev. Lett.* **122**, 173401 (2019).
  - [12] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, *Nature* **551**, 579 (2017).
  - [13] S. Moudgalya, S. Rachel, B. A. Bernevig, and N. Regnault, *Phys. Rev. B* **98**, 235155 (2018).
  - [14] H. Zhao, J. Vovrosh, F. Mintert, and J. Knolle, *Phys. Rev. Lett.* **124**, 160604 (2020).
  - [15] P. N. Jepsen, Y. K. E. Lee, H. Lin, I. Dimitrova, Y. Margalit, W. W. Ho, and W. Ketterle, *Nature Physics* **18**, 899 (2022).
  - [16] M. Serbyn, D. A. Abanin, and Z. Papić, *Nature Physics* **17**, 675 (2021).
  - [17] S. Moudgalya, B. A. Bernevig, and N. Regnault, *Reports on Progress in Physics* **85**, 086501 (2022).
  - [18] A. Chandran, T. Iadecola, V. Khemani, and R. Moessner, *Annual Review of Condensed Matter Physics* **14**, 443 (2023).
  - [19] D. Bluvstein, A. Omran, H. Levine, A. Keesling, G. Semeghini, S. Ebadi, T. T. Wang, A. A. Michailidis, N. Maskara, W. W. Ho, S. Choi, M. Serbyn, M. Greiner, V. Vuletić, and M. D. Lukin, *Science* **371**, 1355 (2021).
  - [20] D. Bluvstein, H. Levine, G. Semeghini, T. T. Wang, S. Ebadi, M. Kalinowski, A. Keesling, N. Maskara, H. Pichler, M. Greiner, V. Vuletić, and M. D. Lukin, *Nature* **604**, 451 (2022).
  - [21] G.-X. Su, H. Sun, A. Hudomal, J.-Y. Desaules, Z.-Y. Zhou, B. Yang, J. C. Halimeh, Z.-S. Yuan, Z. Papić, and J.-W. Pan, *Phys. Rev. Res.* **5**, 023010 (2023).
  - [22] P. Zhang, H. Dong, Y. Gao, L. Zhao, J. Hao, J.-Y. Desaules, Q. Guo, J. Chen, J. Deng, B. Liu, W. Ren, Y. Yao, X. Zhang, S. Xu, K. Wang, F. Jin, X. Zhu, B. Zhang, H. Li, C. Song, Z. Wang, F. Liu, Z. Papić, L. Ying, H. Wang, and Y.-C. Lai, *Nature Physics* **19**, 120 (2023).
  - [23] H. Dong, J.-Y. Desaules, Y. Gao, N. Wang, Z. Guo,

- J. Chen, Y. Zou, F. Jin, X. Zhu, P. Zhang, H. Li, Z. Wang, Q. Guo, J. Zhang, L. Ying, and Z. Papić, *Science Advances* **9**, eadj3822 (2023).
- [24] F. M. Surace, P. P. Mazza, G. Giudici, A. Leroise, A. Gambassi, and M. Dalmonte, *Phys. Rev. X* **10**, 021041 (2020).
- [25] J. C. Halimeh, L. Barbiero, P. Hauke, F. Grusdt, and A. Bohrdt, *Quantum* **7**, 1004 (2023).
- [26] J.-Y. Desaulles, D. Banerjee, A. Hudomal, Z. Papić, A. Sen, and J. C. Halimeh, *Phys. Rev. B* **107**, L201105 (2023).
- [27] J.-Y. Desaulles, A. Hudomal, D. Banerjee, A. Sen, Z. Papić, and J. C. Halimeh, *Phys. Rev. B* **107**, 205112 (2023).
- [28] T. Iadecola and M. Schechter, *Phys. Rev. B* **101**, 024306 (2020).
- [29] A. S. Aramthottil, U. Bhattacharya, D. González-Cuadra, M. Lewenstein, L. Barbiero, and J. Zakrzewski, *Phys. Rev. B* **106**, L041101 (2022).
- [30] J.-Y. Desaulles, T. Iadecola, and J. C. Halimeh, arXiv eprints 10.48550/arxiv.2404.11645 (2024), arXiv:2404.11645 [cond-mat.quant-gas].
- [31] D. Banerjee and A. Sen, *Phys. Rev. Lett.* **126**, 220601 (2021).
- [32] S. Biswas, D. Banerjee, and A. Sen, *SciPost Phys.* **12**, 148 (2022).
- [33] L. Ebner, A. Schäfer, C. Seidl, B. Müller, and X. Yao, arXiv eprints 10.48550/arxiv.2401.15184 (2024), arXiv:2401.15184 [hep-lat].
- [34] I. Sau, P. Stornati, D. Banerjee, and A. Sen, *Phys. Rev. D* **109**, 034519 (2024).
- [35] J. Osborne, I. P. McCulloch, and J. C. Halimeh, arXiv eprints 10.48550/arxiv.2403.08858 (2024), arXiv:2403.08858 [cond-mat.quant-gas].
- [36] T. Budde, M. K. Marinković, and J. C. P. Barros, arXiv eprints 10.48550/arxiv.2403.08892 (2024), arXiv:2403.08892 [hep-lat].
- [37] M. Dalmonte and S. Montangero, *Contemporary Physics* **57**, 388 (2016).
- [38] M. C. Bañuls, R. Blatt, J. Catani, A. Celi, J. I. Cirac, M. Dalmonte, L. Fallani, K. Jansen, M. Lewenstein, S. Montangero, C. A. Muschik, B. Reznik, E. Rico, L. Tagliacozzo, K. Van Acoleyen, F. Verstraete, U.-J. Wiese, M. Wingate, J. Zakrzewski, and P. Zoller, *The European Physical Journal D* **74**, 165 (2020).
- [39] E. Zohar, J. I. Cirac, and B. Reznik, *Reports on Progress in Physics* **79**, 014401 (2015).
- [40] Y. Alexeev, D. Bacon, K. R. Brown, R. Calderbank, L. D. Carr, F. T. Chong, B. DeMarco, D. Englund, E. Farhi, B. Fefferman, A. V. Gorshkov, A. Houck, J. Kim, S. Kimmel, M. Lange, S. Lloyd, M. D. Lukin, D. Maslov, P. Maunz, C. Monroe, J. Preskill, M. Roetteler, M. J. Savage, and J. Thompson, *PRX Quantum* **2**, 017001 (2021).
- [41] M. Aidelsburger, L. Barbiero, A. Bermudez, T. Chanda, A. Dauphin, D. González-Cuadra, P. R. Grzybowski, S. Hands, F. Jendrzejewski, J. Jünemann, G. Juzeliūnas, V. Kasper, A. Piga, S.-J. Ran, M. Rizzi, G. Sierra, L. Tagliacozzo, E. Tirrito, T. V. Zache, J. Zakrzewski, E. Zohar, and M. Lewenstein, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **380**, 20210064 (2022).
- [42] E. Zohar, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **380**, 20210069 (2022).
- [43] N. Klco, A. Roggero, and M. J. Savage, *Reports on Progress in Physics* **85**, 064301 (2022).
- [44] C. W. Bauer, Z. Davoudi, A. B. Balantekin, T. Bhattacharya, M. Carena, W. A. de Jong, P. Draper, A. El-Khadra, N. Gemelke, M. Hanada, D. Kharzeev, H. Lamm, Y.-Y. Li, J. Liu, M. Lukin, Y. Meurice, C. Monroe, B. Nachman, G. Pagano, J. Preskill, E. Rinaldi, A. Roggero, D. I. Santiago, M. J. Savage, I. Siddiqi, G. Siopsis, D. Van Zanten, N. Wiebe, Y. Yamauchi, K. Yeter-Aydeniz, and S. Zorzetti, *PRX Quantum* **4**, 027001 (2023).
- [45] A. D. Meglio, K. Jansen, I. Tavernelli, C. Alexandrou, S. Arunachalam, C. W. Bauer, K. Borras, S. Carrazza, A. Crippa, V. Croft, R. de Putter, A. Delgado, V. Dunjko, D. J. Egger, E. Fernandez-Combarro, E. Fuchs, L. Funcke, D. Gonzalez-Cuadra, M. Grossi, J. C. Halimeh, Z. Holmes, S. Kuhn, D. Lacroix, R. Lewis, D. Lucchesi, M. L. Martinez, F. Meloni, A. Mezzacapo, S. Montangero, L. Nagano, V. Radescu, E. R. Ortega, A. Roggero, J. Schuhmacher, J. Seixas, P. Silvi, P. Spentzouris, F. Tacchino, K. Temme, K. Terashi, J. Tura, C. Tuysuz, S. Vallecorsa, U.-J. Wiese, S. Yoo, and J. Zhang, arXiv eprints 10.48550/arxiv.2307.03236 (2023), arXiv:2307.03236 [quant-ph].
- [46] J. C. Halimeh, M. Aidelsburger, F. Grusdt, P. Hauke, and B. Yang, arXiv eprints 10.48550/arxiv.2310.12201 (2023), arXiv:2310.12201 [cond-mat.quant-gas].
- [47] Y. Cheng and H. Zhai, arXiv eprints 10.48550/arxiv.2401.07708 (2024), arXiv:2401.07708 [cond-mat.quant-gas].
- [48] See Supplemental Material for details about the derivation of the model, the characterization of the ergodic regime, and the long-time dynamics.
- [49] G. Cataldi, G. Magnifico, P. Silvi, and S. Montangero, arXiv eprints 10.48550/arxiv.2307.09396 (2024), arXiv:2307.09396 [hep-lat].
- [50] M. Rigobello, G. Magnifico, P. Silvi, and S. Montangero, arXiv eprints 10.48550/arxiv.2308.04488 (2023), arXiv:2308.04488 [hep-lat].
- [51] G. Calajò, G. Magnifico, C. Edmunds, M. Ringbauer, S. Montangero, and P. Silvi, arXiv eprints 10.48550/arxiv.2402.07987 (2024), arXiv:2402.07987 [quant-ph].
- [52] P. Silvi, Y. Sauer, F. Tschirsich, and S. Montangero, *Phys. Rev. D* **100**, 074512 (2019).
- [53] E. Zohar and J. I. Cirac, *Phys. Rev. D* **99**, 114511 (2019).
- [54] J. Kogut and L. Susskind, *Phys. Rev. D* **11**, 395 (1975).
- [55] L. Susskind, *Phys. Rev. D* **16**, 3031 (1977).
- [56] E. Zohar and M. Burrello, *Physical Review D* **91**, 054506 (2015).
- [57] D. Banerjee, M. Dalmonte, M. Müller, E. Rico, P. Stebler, U.-J. Wiese, and P. Zoller, *Physical Review Letters* **109**, 10.1103/physrevlett.109.175302 (2012).
- [58] S. Kühn, J. I. Cirac, and M.-C. Bañuls, *Phys. Rev. A* **90**, 042305 (2014).
- [59] G. Magnifico, M. Dalmonte, P. Facchi, S. Pascazio, F. V. Pepe, and E. Ercolessi, *Quantum* **4**, 281 (2020).
- [60] A. Daniel, A. Hallam, J.-Y. Desaulles, A. Hudomal, G.-X. Su, J. C. Halimeh, and Z. Papić, *Phys. Rev. B* **107**, 235108 (2023).
- [61] A. Hudomal, J.-Y. Desaulles, B. Mukherjee, G.-X. Su, J. C. Halimeh, and Z. Papić, *Phys. Rev. B* **106**, 104302 (2022).
- [62] T. Budde, M. K. Marinković, and J. C. P. Barros, arXiv eprints 10.48550/arxiv.2403.08892 (2024), arXiv:2403.08892 [hep-lat].
- [63] M. Van Damme, T. V. Zache, D. Banerjee, P. Hauke, and J. C. Halimeh, *Phys. Rev. B* **106**, 245110 (2022).

- [64] M. Van Damme, J.-Y. Desaulles, Z. Papić, and J. C. Halimeh, *Phys. Rev. Res.* **5**, 033090 (2023).
- [65] M. Ringbauer, M. Meth, L. Postler, R. Stricker, R. Blatt, P. Schindler, and T. Monz, *Nature Physics* **18**, 1053 (2022).
- [66] M. Fishman, S. R. White, and E. M. Stoudenmire, *SciPost Phys. Codebases* , 4 (2022).
- [67] M. Fishman, S. R. White, and E. M. Stoudenmire, *SciPost Phys. Codebases* , 4 (2022).
- [68] G. Cataldi, *ed-igt*, Code for Exact Diagonalization of Quantum Many-Body Hamiltonians and Lattice Gauge Theories. (2024).
- [69] P. Silvi, E. Rico, T. Calarco, and S. Montangero, *New Journal of Physics* **16**, 103015 (2014).
- [70] M. Fannes, B. Nachtergaele, and R. F. Werner, *Communications in Mathematical Physics* **144**, 443 (1992).
- [71] A. Klumper, A. Schadschneider, and J. Zittartz, *Journal of Physics A: Mathematical and General* **24**, L955 (1991).
- [72] A. Klumper, A. Schadschneider, and J. Zittartz, *Europhysics Letters* **24**, 293 (1993).
- [73] G. Vidal, *Physical review letters* **91**, 147902 (2003).
- [74] R. Orús, *Annals of Physics* **349**, 117 (2014).
- [75] J. Eisert, M. Cramer, and M. B. Plenio, *Rev. Mod. Phys.* **82**, 277 (2010).
- [76] F. Verstraete, V. Murg, and J. I. Cirac, *Advances in physics* **57**, 143 (2008).
- [77] U. Schollwöck, *Annals of physics* **326**, 96 (2011).
- [78] G. Vidal, *Phys. Rev. Lett.* **93**, 040502 (2004).
- [79] J. Haegeman, J. I. Cirac, T. J. Osborne, I. Pizorn, H. Verschelde, and F. Verstraete, *Phys. Rev. Lett.* **107**, 070601 (2011).
- [80] S. Paeckel, T. Köhler, A. Swoboda, S. R. Manmana, U. Schollwöck, and C. Hubig, *Annals of Physics* **411**, 167998 (2019).
- [81] A. W. Sandvik, *AIP Conference Proceedings* **1297**, 135 (2010).

# Supplemental Online Material for “Quantum Many-Body Scarring in a Non-Abelian Lattice Gauge Theory”

In this Supplemental Material, we provide details on the model derivation and on the numerical techniques employed in this study. Additionally, we show: persistence of non-ergodic dynamics at higher gauge truncations, evidence of ergodic dynamics for different parameter regimes than the ones considered in the main text, time evolution for different system sizes, and longer time.

## HARDCORE-GLUON QUDIT MODEL

In order to map the Kogut-Susskind  $SU(2)$  lattice Yang-Mills Hamiltonian of Eq. (1) to the qudit model used in numerical simulations, we: (i) perform a hardcore-gluon truncation of the gauge field; (ii) build the gauge-singlet local dressed basis; and (iii) gauge-defermionize. The procedure is conveniently formulated by decomposing matter and gauge local Hilbert spaces in  $SU(2)$  irreducible representations (irreps). This preliminary step is already inbuilt in the local bases introduced in the main text: the irrep basis  $|j, m_L, m_R\rangle$  for gauge links, and the Fock basis  $\{|q\rangle\} = \{|0\rangle, |r\rangle, |g\rangle = \hat{\psi}_{r,g}^\dagger|0\rangle, |2\rangle = \hat{\psi}_r^\dagger\hat{\psi}_g^\dagger|0\rangle\}$  for matter sites. Matter basis states are associated with the following spin labels  $(j, m)$ :

$$|0\rangle \leftrightarrow (0, 0) \quad |r\rangle, |g\rangle \leftrightarrow (0, \pm 1/2) \quad |2\rangle \leftrightarrow (0, 0). \quad (\text{S1})$$

Irrep decomposition specifies how local gauge rotations act on a site  $n$  and its neighboring links. Infinitesimal rotations are generated by  $\hat{\mathbf{G}}_n = \hat{\mathbf{R}}_{n-1,n} + \hat{\mathbf{Q}}_n + \hat{\mathbf{L}}_{n,n+1}$ , where  $\hat{Q}_n^\nu = \sum_{ab} \hat{\psi}_{na}^\dagger S_{ab}^{(\frac{1}{2})^\nu} \hat{\psi}_{nb}$  rotates the quark field at  $n$ , while  $\hat{\mathbf{R}}_{n-1,n}$  and  $\hat{\mathbf{L}}_{n,n+1}$  account for the transformation of the gauge links at its left and its right respectively [39]:

$$\langle j' m'_L m'_R | \hat{L}^\nu | j m_L m_R \rangle = \delta_{j,j'} S_{m'_L, m_L}^{(j)^\nu} \delta_{m'_R, m_R}, \quad (\text{S2a})$$

$$\langle j' m'_L m'_R | \hat{R}^\nu | j m_L m_R \rangle = \delta_{j,j'} \delta_{m'_L, m_L} S_{m'_R, m_R}^{(j)^\nu}; \quad (\text{S2b})$$

$S^{(j)^\nu}$  being the spin- $j$   $\mathfrak{su}(2)$  matrices, with  $\nu \in \{x, y, z\}$ . The chromoelectric energy operator is just their square  $\hat{E}^2 = \hat{\mathbf{R}}^2 = \hat{\mathbf{L}}^2$ , i.e. the quadratic Casimir [39]:

$$\hat{E}^2 |j m_L m_R\rangle = j(j+1) |j m_L m_R\rangle. \quad (\text{S3})$$

In principle, the gauge field could occupy arbitrarily high spin shells; however, in the hardcore-gluon model, we restrict to  $j \in \{0, 1/2\}$ , corresponding to a cutoff  $\|\hat{E}^2\| \leq 3/4$  on the chromoelectric field operator spectrum ( $\|\cdot\|$  denotes the matrix norm). Recovering the untruncated physics is not crucial for the present study. Nevertheless, we stress that hardcore gluons give a good approximation of the untruncated low energy physics in the strong coupling limit  $g_0 \gg 1$ , where the gauge field energy term dominates the Hamiltonian.

To enforce the physical state condition, i.e., non-Abelian Gauss law,  $\hat{\mathbf{G}}_n |\Psi_{\text{phys}}\rangle \equiv 0$ , we observe that  $\hat{\mathbf{L}}$

and  $\hat{\mathbf{R}}$  from Eq. (S2) act nontrivially only on the  $m_L$  and  $m_R$  index respectively. This suggests [69] to factorize each gauge link in a pair of new rishon degrees of freedom, living at its edges. Upon fusing each site with its adjacent rishons, we forge a composite site where Gauss law becomes an internal constraint [50]. We label the basis states of each rishon as  $|0\rangle = |j=0, m=0\rangle$  and  $|r\rangle, |g\rangle = |j=\frac{1}{2}, m=\pm\frac{1}{2}\rangle$ . The left and right rishons on each link are then constrained to be in the same spin shell  $j$  by restricting to the even parity sector of a local  $\mathbb{Z}_2$  symmetry, whose action reads

$$\hat{P} = +|0\rangle\langle 0| - (|r\rangle\langle r| + |g\rangle\langle g|). \quad (\text{S4})$$

In this way, the original physical link states are selected. Solving Gauss law in the composite site amounts to finding the local singlets via Clebsh-Gordan decomposition, which provides the following 6-dimensional local dressed basis,  $\{|\alpha\rangle = |m_R^\alpha(n-1, n), q^\alpha(n), m_L^\alpha(n, n+1)\}_{\alpha=1}^6$  [51]:

$$\begin{aligned} |1\rangle &= |0, 0, 0\rangle, & |2\rangle &= \frac{|r, 0, g\rangle - |g, 0, r\rangle}{\sqrt{2}}, \\ |3\rangle &= \frac{|g, r, 0\rangle - |r, g, 0\rangle}{\sqrt{2}}, & |4\rangle &= \frac{|0, r, g\rangle - |0, g, r\rangle}{\sqrt{2}}, \\ |5\rangle &= |0, 2, 0\rangle, & |6\rangle &= \frac{|r, 2, g\rangle - |g, 2, r\rangle}{\sqrt{2}}. \end{aligned} \quad (\text{S5})$$

In the irrep basis for gauge link, the parallel transporter is given in terms of Clebsh-Gordan coefficients [56]:

$$\langle j' m'_L m'_R | \hat{U}^{ab} | j m_L m_R \rangle = \sqrt{\frac{2j+1}{2j'+1}} \frac{C_{\frac{1}{2}, a; j', m'_L}^{j, m_L}}{C_{\frac{1}{2}, b; j, m_R}^{j', m'_R}}.$$

we realize it in terms of rishon operators as

$$\hat{U}_{n,n+1}^{ab} \rightarrow \frac{1}{\sqrt{2}} \hat{\zeta}_{n,n+1}^{(L)a} (\hat{\zeta}_{n,n+1}^{(R)b})^\dagger, \quad (\text{S6})$$

with

$$\hat{\zeta}^r = |0\rangle\langle r| + |g\rangle\langle 0|, \quad \hat{\zeta}^g = |0\rangle\langle g| - |r\rangle\langle 0|. \quad (\text{S7})$$

Observe that the right-hand side of Eq. (S6) preserves the link parity, as desired. Conversely, a single rishon operator always inverts the local parity:  $\{\hat{P}, \hat{\zeta}^a\} = 0$ . The existence of a  $\mathbb{Z}_2$ -grading in the rishon spaces allows to gauge-defermionize the model [49]: we take  $\hat{\zeta}^a$  to be fermionic and introduce the bosonic operators,

$$\hat{Q}_n^{(L,R)} = \sum_a (\hat{\zeta}_{n,n+1}^{(L,R)a})^\dagger \hat{\psi}_{n,a}, \quad \hat{M} = \sum_a \hat{\psi}_{n,a}^\dagger \hat{\psi}_{n,a}. \quad (\text{S8})$$



For potential quantum simulation implementations, it is convenient to rewrite these solely in terms of Hermitian operators, introducing [51]

$$\begin{aligned}\hat{A}^{(1)} &= \hat{Q}^{(L)} + \hat{Q}^{(L)\dagger}, & \hat{B}^{(1)} &= \hat{Q}^{(R)} + \hat{Q}^{(R)\dagger}, \\ \hat{A}^{(2)} &= i[\hat{Q}^{(L)} - \hat{Q}^{(L)\dagger}], & \hat{B}^{(2)} &= i[\hat{Q}^{(R)} - \hat{Q}^{(R)\dagger}].\end{aligned}\quad (\text{S9})$$

Finally, we map Eq. (1) to the following 6-dimensional qudit Hamiltonian [51]:

$$\hat{H} = \sum_{j \in \{1,2\}} \sum_n \hat{A}_n^{(j)} \hat{B}_{n+1}^{(j)} + m(-1)^n \hat{M}_n + g^2 \sum_n \hat{C}_n, \quad (\text{S10})$$

where we rescaled energy to absorb the hopping prefactor,  $\hat{H} = 4\sqrt{2}a_0\hat{H}_0$ , and defined the dimensionless couplings  $m = 4\sqrt{2}a_0m_0$  and  $g^2 = \frac{3\sqrt{2}}{4}a_0g_0^2$ . The qubit operators appearing in Eq. (S10) are reported below [51]:

$$\hat{Q}^{(L)} = \sqrt{2}|1\rangle\langle 4| + |2\rangle\langle 3| + |3\rangle\langle 6| + \sqrt{2}|4\rangle\langle 5|, \quad (\text{S11a})$$

$$\hat{Q}^{(R)} = \sqrt{2}|1\rangle\langle 3| + |2\rangle\langle 4| + \sqrt{2}|3\rangle\langle 5| + |4\rangle\langle 6|, \quad (\text{S11b})$$

$$\hat{M} = |3\rangle\langle 3| + |4\rangle\langle 4| + 2|5\rangle\langle 5| + 2|6\rangle\langle 6|, \quad (\text{S11c})$$

$$\hat{C} = 2|2\rangle\langle 2| + |3\rangle\langle 3| + |4\rangle\langle 4| + 2|6\rangle\langle 6|. \quad (\text{S11d})$$

Finally, in the dressed qudit basis given in Eq. (S5) the initial states considered in the main text read respectively

$$|\Psi_{\text{PV}}\rangle = |6\rangle|2\rangle\dots|6\rangle|2\rangle, \quad (\text{S12})$$

for the polarized bare vacuum and

$$|\Psi_{\text{V}}\rangle = |5\rangle|1\rangle\dots|5\rangle|1\rangle, \quad (\text{S13})$$

for the bare vacuum.

### Scars at higher gauge link truncation

The rishon decomposition of  $\hat{U}^{ab}$  proposed in Eq. (S6) can be generalized to arbitrary truncation of the maximum allowed spin shell  $j_{\text{max}}$ , although at a (manageable) added cost. Starting from a given spin shell  $j$ , we have to separately account for the action when both rishons are increased to shell  $j + \frac{1}{2}$ , and both are decreased to shell  $j - \frac{1}{2}$ . We can then decompose  $\hat{U}^{ab}$  as follows [49]:

$$\hat{U}_{\mathbf{j},\mathbf{j}+\boldsymbol{\mu}}^{ab} = \hat{\zeta}_{A,\mathbf{j},\boldsymbol{\mu}}^a \hat{\zeta}_{B,\mathbf{j}+\boldsymbol{\mu},-\boldsymbol{\mu}}^{b\dagger} + \hat{\zeta}_{B,\mathbf{j},\boldsymbol{\mu}}^a \hat{\zeta}_{A,\mathbf{j}+\boldsymbol{\mu},-\boldsymbol{\mu}}^{b\dagger}, \quad (\text{S14})$$

where the two  $\zeta$ -rishon species, A and B, act respectively as raising and lowering the spin shell of the SU(2) gauge irreducible representation. Interestingly, they are related to each other as:

$$\hat{\zeta}_A^a = i\sigma_{a,b}^y \hat{\zeta}_B^{b\dagger}, \quad \hat{\zeta}_A^{a\dagger} = i\sigma_{a,b}^y \hat{\zeta}_B^b. \quad (\text{S15})$$

We can then rewrite Eq. (S14) just in terms of one species, e.g. B. Dropping the index, i.e.  $\hat{\zeta}_B^a = \hat{\zeta}^a$ , it holds:

$$\hat{U}_{\mathbf{j},\mathbf{j}+\boldsymbol{\mu}}^{ab} = i\sigma_{a,c}^y \hat{\zeta}_{\mathbf{j},\boldsymbol{\mu}}^{c\dagger} \hat{\zeta}_{\mathbf{j}+\boldsymbol{\mu},-\boldsymbol{\mu}}^{b\dagger} + i\sigma_{b,c}^y \hat{\zeta}_{\mathbf{j},\boldsymbol{\mu}}^a \hat{\zeta}_{\mathbf{j}+\boldsymbol{\mu},-\boldsymbol{\mu}}^c. \quad (\text{S16})$$

For a chosen truncation  $j_{\text{max}}$  of the SU(2) irreducible representation,  $\zeta$ -rishons are defined as follows [49]:

$$\hat{\zeta}^{g(r)} = \left[ \sum_{j=0}^{j_{\text{max}}-\frac{1}{2}} \sum_{m=-j}^j \chi(j, m, g(r)) |j, m\rangle \langle j + \frac{1}{2}, m \mp \frac{1}{2}| \right]_F, \quad (\text{S17})$$

where the function  $\chi(j, m, \alpha)$  reads

$$\chi(j, m, g(r)) = \sqrt{\frac{j \mp m + 1}{(2j+1)(2j+2)}}. \quad (\text{S18})$$

For instance, truncating the gauge link to the shell  $j = 1$ , we obtain a local Hilbert qudit basis of 10 states. It is interesting to highlight that the scarring dynamics observed for the minimal truncation in Fig. 1, is not a feature of the truncation itself. Indeed, a similar behavior is displayed in the next  $j = 1$  gauge truncation, as shown in Fig. S1, where we return the fidelity and the entanglement entropy for the same two cases discussed in the main text. Analogously with what has been observed in abelian LGTs, these results remarkably suggest that the scars dynamics of non-abelian theories also persist at higher gauge truncations.

## ERGODIC VS NON-ERGODIC BEHAVIOR

In this section, we provide further evidence that the non-ergodic dynamics discussed in the main text are indeed attributable to the occurrence of many-body scars. Specifically, we first demonstrate how, for other parameter regimes an ergodic behavior is observed as predicted by thermal ensembles. Secondly, we illustrate how scars-induced revivals persist at larger system sizes and over longer timescales.

### Thermal ensembles

The long-time average of a generic observable  $\hat{O}$  resulting from the unitary dynamics of a given initial state  $|\Psi(0)\rangle = \sum_s C_s |\Phi_s\rangle$  expressed in the eigenbasis of the Hamiltonian (1) is well described by the expectation value predicted by the *diagonal* ensemble [7]:

$$\begin{aligned}\bar{O} &\equiv \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \langle \Psi(t) | \hat{O} | \Psi(t) \rangle \\ &= \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T dt \sum_{s,p} C_s^* C_p \langle \Phi_s | e^{it\hat{H}} \hat{O} e^{-it\hat{H}} | \Phi_p \rangle \\ &= \sum_{s,p} C_s^* C_p \langle \Phi_s | \hat{O} | \Phi_p \rangle \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T e^{i(E_s - E_p)t} dt \\ &= \sum_s |C_s|^2 \langle \Phi_s | \hat{O} | \Phi_s \rangle = \text{Tr}\{\hat{O} \hat{\rho}_{\text{DE}}\} \equiv \langle \hat{O} \rangle_{\text{DE}},\end{aligned}\quad (\text{S19})$$




FIG. S1. *Scarring at higher gauge truncation.* Many body scarring dynamics for the polarized bare vacuum and the bare vacuum for the truncated SU(2) YM LGT at  $j_{\max} = 1$  (pink line) in comparison with the corresponding one at  $j_{\max} = 1/2$  (cyan line). Each column reports the fidelity and bipartite entanglement entropy as a function of time for the same two cases ( $m, g$ ) considered in Fig. 1. The simulations are obtained via ED for  $N = 10$  sites.

where  $\hat{\rho}_{\text{DE}} = \sum_s |C_s\rangle\langle C_s| |\Phi_s\rangle\langle\Phi_s|$  is the diagonal ensemble. Then, if the expectation value of the observable  $\langle\hat{O}(t)\rangle$  approaches long-time average  $\langle\hat{O}\rangle_{\text{DE}}$  over long times, the system *equilibrates*. Rather, *thermalization* is achieved when the steady state coincides with the prediction of the microcanonical ensemble whose thermal average reads

$$\langle\hat{O}\rangle_{\text{ME}} = \text{Tr}\{\hat{O}\hat{\rho}_{\text{ME}}\}, \quad (\text{S20})$$

where the microcanonical ensemble is defined as a superposition within the energy shell  $[E_q - \delta E, E_q + \delta E]$ , with the quench energy  $E_q = \langle\Psi(0)|\hat{H}|\Psi(0)\rangle$ , containing  $N_{E,\delta E}$  eigenstates:

$$\hat{\rho}_{\text{ME}} = \frac{1}{N_{E,\delta E}} \sum_{|E_s - E_q| < \delta E} |\Phi_s\rangle\langle\Phi_s|. \quad (\text{S21})$$

### Ergodic dynamics

We have shown that the two considered initial product states polarized bare vacuum (PV) and bare vacuum (V), escape thermalization for the two regimes ( $m = 5, g^2 = 1$ ) and ( $m = 1, g^2 = 5$ ). To confirm the scarring nature of the observed non-ergodic dynamics we investigate the absence of scars states in the many-body spectrum for other parameter regimes. In Fig. S2, we look at the overlap of the full many-body spectrum with the two considered initial states PV and V and the corresponding bipartite entanglement entropy for different values of  $g^2 = m$ .

We do not observe a clear emergence of a tower of states (rather than in Fig. 2), and thus, we expect ergodic dynamics for the two considered states. This is also confirmed by the time evolution of the two initial states for




FIG. S2. *Many-body spectrum.* Spectrum analysis for a lattice chain of  $N = 10$  sites with open boundary conditions in a few paradigmatic regimes (corresponding to different columns) where we observe ergodic dynamics. The first and second rows show the overlap of the many-body spectrum with the bare vacuum and polarized bare vacuum state, respectively. Only states with  $\mathcal{O}_{\text{PV}} > 10^{-9}$  are included in the plot. In the third row, we display the bipartite entanglement entropy of all eigenstates in the energy spectrum as a function of their respective eigenvalues.

$g^2 = m = 1$  displayed in Fig. S3. In this case, the fidelity does not exhibit persistent revivals, and the entanglement entropy shows rapid growth compared to the previously observed scarred dynamics. Moreover, as expected for thermalization, the occupancy relaxes to a steady value compatible with the microcanonical ensemble averages.




FIG. S3. *Ergodic dynamics.* First to third row: return fidelity, average quark occupancy, and bipartite entanglement entropy as a function of time for the two considered initial states, polarized bare vacuum (PV), and bare vacuum (V) at  $m=g^2=1$ . The horizontal lines in the occupancy panels indicate the thermal expectation value of the single particle occupancy density obtained through ED of a lattice of  $N = 10$  with PBS for the microcanonical ensemble. The simulation was performed with MPS for  $N = 20$  sites by setting open boundary conditions, maximum bond dimension  $D_{\max} = 350$ , and truncation tolerance  $\text{tol} = 10^{-7}$ .

We leave any further investigations on scarring dynamics for other candidate initial states to future work. To this extent, it could be interesting to investigate the  $g^2 = m = 0.1$  case, whose many-body spectrum contains extremely low entanglement entropy states.

### Scaling with the system size

Many-body scars represent a polynomial (in number) subset of special non-thermal eigenstates within the full many-body spectrum. This implies that the overlap with the initially considered product states decreases as the system size increases. To investigate this behavior, in Fig. S4(a), we plot the return fidelity between the evolved state and two initial states discussed in the main text: the polarized bare vacuum (PV) and the bare vacuum (V). Exploiting the slow entanglement growth in the many-body scars regime, we considered system sizes ranging from  $N = 10$  to  $N = 50$ . Although the fidelity peaks decrease with increasing  $N$  as expected, pronounced revivals are still visible even for large chains with  $N = 50$  sites. The peaks become sharper, with the fidelity minima approach-




FIG. S4. *Finite-size scaling.* Return fidelity during the evolution of the polarized bare vacuum (a) and bare vacuum (b) initial states, for  $m = 5$ ,  $g^2 = 1$ , open boundary conditions, and various systems sizes  $N$ . The side plots (c) and (d) show the finite-size scaling of the highest fidelity peak, highlighted by a blue line in (a) and (b). Notice the log  $y$ -scale. The pink line is an exponential fit  $\mathcal{F}_N = f^N$ , and the extrapolated peak fidelity per site  $f$  is reported in the Figure. Simulations performed with maximum MPS bond dimension  $D_{\max} = 350$ , and truncation tolerance  $\text{tol} = 10^{-7}$ .

ing zero, accordingly with the expected revivals induced by scars. The maximum amplitude of these peaks follows an exponential scaling in the system size,  $\mathcal{F}_N = f^N$ , with the extrapolated peak fidelity per site being  $f = 0.97$  and  $f = 0.98$  for the PV and V states, respectively. This estimate further demonstrates the robustness of the observed scar dynamics even at large system sizes.

### Long-time dynamics

To illustrate how the observed non-ergodic dynamics persist over long times, we conducted simulations for an extended duration. In Fig. S5, we present the same dynamics presented in Fig. 1 of the main text, now up to  $t = 30$ , nearly ten times longer than previously shown. At this extended duration, we continue to observe persistent revivals and oscillations in both the return fidelity and occupancy. Moreover, in addition to the rapid oscillations with a period  $T < 1$  discussed in the main text, we observe a beating with frequency  $T \sim 1$  attributed to the non-uniform distribution of scars in energy spacing. Finally, we still observe a slow growth in entanglement at such extended times, consistent with dynamics constrained to the many-body scars subset of the spectrum.




FIG. S5. *Long time dynamics.* First to third row: return fidelity, average quark occupancy, and bipartite entanglement entropy as a function of time for the two considered initial states, polarized bare vacuum, and bare vacuum. We considered different values of mass and coupling as indicated in the Figure. The simulation was performed with MPS for  $N = 30$  sites by setting open boundary conditions, maximum bond dimension  $D_{\max} = 350$ , and truncation tolerance  $\text{tol} = 10^{-7}$ .

## NUMERICAL METHODS

### Matrix product state simulation

To solve the dynamics described by the model Eq. (S10) we employ a numerical simulation based on Matrix product states (MPS).

Consider a quantum many-body system defined on a one-dimensional lattice with  $N$  sites. A generic pure quantum state  $|\psi\rangle$  of this system lives in the tensor product  $\mathcal{H} = \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_N$  of  $N$  local Hilbert spaces  $\mathcal{H}_j$ , each one being of finite dimension  $d$ . This state  $|\psi\rangle$  can be expanded as  $|\psi\rangle = \sum_{i_1, \dots, i_N} \psi_{i_1, i_2, \dots, i_N} |i_1, i_2, \dots, i_N\rangle$ , where  $\{|i\rangle_j\}$  is the canonical basis of the site  $j$ . Providing an exact description of this state entails knowing all the  $d^N$  complex coefficients  $\psi_{i_1, i_2, \dots, i_N}$ . From a computational point of view, this becomes infeasible for large values of  $N$  due to the exponential increase of the memory and resources you would need.

The main idea of the MPS representation consists in reshaping the generic quantum state  $|\psi\rangle$  of the one-dimensional system into the form [70–72]:

$$|\psi\rangle = \sum_{i_1, \dots, i_N} \text{Tr} \left[ A_1^{(i_1)} A_2^{(i_2)} \dots A_N^{(i_N)} \right] |i_1, i_2, \dots, i_N\rangle, \quad (\text{S22})$$

where, for each specific set of physical indices  $\{i_1, i_2, \dots, i_N\}$ ,

the trace of the product of the  $A_j^{(i_j)}$  matrices gives back the state coefficient  $\psi_{i_1, i_2, \dots, i_N}$ . Each matrix  $A_j^{(i_j)}$  has dimension  $D_{j-1} \times D_j$  and, when working with open-boundary conditions, finite-edge constraints are used, i.e.  $D_1 = 1$  and  $D_N = 1$ . In principle, every state of a one-dimensional quantum many-body system can be written exactly in the MPS form, with bond dimensions  $D_j$  growing exponentially with the lattice sites  $N$  [73]. However, when dealing with low-energy states of local Hamiltonians, and their real-time dynamics for small-to-intermediate time, an approximate but efficient representation at fixed maximum bond dimensions is usually very accurate in capturing the essential features of the systems [74, 75]. In this case, the number of parameters in the MPS representation grows only polynomially with the system's size, drastically reducing the overall computational complexity.

Bond dimension  $D_j$  are crucial parameters since they allow interpolating between separable states (all  $D_j = 1$ ) and strongly entangled states ( $D_j \gg 1$ ). Thus, they estimate the quantum correlations and entanglement present in the MPS representation. This implies that dynamical problems characterized by a slow entanglement entropy growth, as in the many-body scars dynamics, can be efficiently and accurately described by MPS ansatz with small or intermediate bond dimensions [76, 77].

To compute the time evolution of the system, we made



use of state-of-the-art algorithms for MPS dynamical simulations: time-evolving block decimation (TEBD) algorithm [78] or the time-dependent variational principle (TDVP) [79] preferring the first to capture scars dynamics, with a precise time step resolution, and the second to capture the more computationally demanding ergodic dynamics. For a detailed description of the methods, please see [80]). With both methods during the time evolution, we impose a fixed tolerance,  $\text{tol}$ , for the truncation in the employed Schmidt decomposition. The maximum bond dimension employed in the simulation to ensure such precision during the time evolution is upper bounded by a maximum bond dimension  $D_{\text{max}}$ . These quantities define the accuracy of the simulation and are reported in the Figure's caption.

### Exact diagonalization

To obtain the full spectrum  $\{|\Phi_s\rangle\}$  of the model given in Eq. (1) and extract the overlap and the entanglement

behaviors displayed in Fig. 2, we use an Exact Diagonalization (ED) code [68] that exploits all the main symmetries of the model.

For instance, on each link, we can resolve the  $\mathbb{Z}_2$  symmetry obtained from the gauge link decomposition into a pair of rishons, and constrain the same electric Casimir on both sides. Throughout all the simulations, we also select the zero baryon number density sector,  $\hat{N}_b = \frac{1}{2} \sum_n (\hat{M}_n - 1) = 0$ , corresponding to the half-filling case.

Ultimately, in the case of periodic boundary conditions (PBC), we also exploit translational invariance [81] and restrict simulations within momentum sector  $k = 0$ , which states such as the bare vacuum in Eq. (5) and polarized vacuum Eq. (2) belong to. Notice that, due to the staggered mass term in Eq. (1), the model is invariant under translations of two sites.