Training-Conditional Coverage Bounds under Covariate Shift

Mehrdad Pournaderi and Yu Xiang Department of Electrical and Computer Engineering University of Utah {m.pournaderi, yu.xiang}@utah.edu

Abstract

Training-conditional coverage guarantees in conformal prediction concern the concentration of the error distribution, conditional on the training data, below some nominal level. The conformal prediction methodology has recently been generalized to the covariate shift setting, namely, the covariate distribution changes between the training and test data. In this paper, we study the training-conditional coverage properties of a range of conformal prediction methods under covariate shift via a weighted version of the Dvoretzky–Kiefer–Wolfowitz (DKW) inequality tailored for distribution change. The result for the split conformal method is almost assumption-free, while the results for the full conformal and jackknife+ methods rely on strong assumptions including the uniform stability of the training algorithm.

1 Introduction

Conformal prediction is a framework for constructing *distribution-free* predictive confidence regions as long as the training and test data are exchangeable [Vovk et al.](#page-6-0) [\[2005\]](#page-6-0) (also see [Shafer and Vovk \[2008\]](#page-6-1), [Vovk et al. \[2009\]](#page-6-2), [Vovk \[2012\]](#page-6-3)). Specifically, let $((X_1, Y_1), (X_2, Y_2), \ldots, (X_n, Y_n), (X_{test}, Y_{test}))$ denote a tuple of exchangeable data points, consisting of a training sequence of n samples $\mathbf{D}_n := ((X_1, Y_1), \dots, (X_n, Y_n)) \in (\mathcal{X} \times \mathcal{Y})^n$ and one test sample $(X_{test}, Y_{test}) \in \mathcal{X} \times \mathcal{Y}$. The conformal prediction provides a coverage of Y_{test} in the sense of

$$
\mathbb{P}(Y_{\text{test}} \in \hat{C}_{\alpha}(X_{\text{test}})) \ge 1 - \alpha,\tag{1}
$$

where \hat{C}_{α} : $\mathcal{X} \to 2^{\mathcal{Y}}$ is a data-dependent map. This type of guarantee is referred to as *marginal* coverage, as it is averaged over all the training and test data.

One natural direction to stronger results is to devise *conditional* coverage guarantee

$$
\mathbb{P}(Y_{\text{test}} \in \hat{\mathcal{C}}_{\alpha}(X_{\text{test}})|X_{\text{test}}) \geq 1 - \alpha.
$$

However, it has been shown in [Vovk \[2012\]](#page-6-3), [Foygel Barber et al. \[2021\]](#page-6-4), [Lei and Wasserman](#page-7-0) [\[2014\]](#page-7-0) that it is *impossible* to obtain (non-trivial) distribution-free prediction regions $\mathcal{C}(x)$ in the finite-sample regime; relaxed versions of this type of guarantee have been extensively studied (see [Jung et al. \[2022](#page-7-1)], [Gibbs et al.](#page-7-2) [\[2023\]](#page-7-2), [Vovk et al.](#page-7-3) [\[2003\]](#page-7-3) and references therein). As an alternative approach, several results (e.g., [Vovk](#page-6-3) [\[2012\]](#page-6-3), [Bian and Barber \[2023\]](#page-7-4)) have been reported on the *training-conditional* guarantee by conditioning on D_n , which is also more appealing than the marginal guarantee as can be seen below. Define the following miscoverage rate as a function of the training data,

$$
P_e(\mathbf{D}_n) := \mathbb{P}(Y_{\text{test}} \notin \hat{\mathcal{C}}(X_{\text{test}})|\mathbf{D}_n).
$$

Note that the marginal coverage in [\(1\)](#page-0-0) is equivalent to $\mathbb{E}[P_e(\mathbf{D}_n)] \leq \alpha$. The training-conditional guarantees are of the following form, for some small $\delta > 0$,

$$
\mathbb{P}(P_e(\mathbf{D}_n) \ge \alpha) \le \delta
$$

or its asymptotic variants. Roughly speaking, this guarantee means that the $(1 - \alpha)$ -level coverage lower bounds hold for a *generic* dataset.

For the *K*-fold CV+ with m samples in each fold, the conditional coverage bound

$$
\mathbb{P}\left(P_e(\mathbf{D}_n) \ge 2\alpha + \sqrt{2\log(K/\delta)/m}\right) \le \delta \tag{2}
$$

is established in [Bian and Barber \[2023](#page-7-4)]. They have also shown that distribution-free trainingconditional guarantees for full conformal and *jackknife+* methods are impossible without further assumptions; in particular, they conjectured that a certain form of algorithmic stability is needed for full conformal and jackknife+. Recently, [Liang and Barber \[2023](#page-7-5)] proposed conditional coverage bounds for jackknife+ and full conformal prediction sets under the assumption that the training algorithm is symmetric and satisfies certain stability conditions (see Section [4](#page-5-0) for more details). In this line of research, samples are assumed to be i.i.d., which is not only exchangeable but also ergodic and admits some nice concentration properties. However, this assumption can be violated in the application. In particular, the input data distribution during deployment can differ from the distribution observed during training. This phenomenon is called *distribution shift* and it is a crucial problem in trustworthy machine learning (see Section [2.3\)](#page-3-0). In this regard, split and full conformal prediction methods as the central methods for distribution-free uncertainty quantification of black-box models have recently been extended to handle a popular type of distribution shift called *covariate shift* [Tibshirani et al. \[2019\]](#page-7-6). In the covariate shift setting, the distribution of covariates in the test data differs from the one observed in the training data, but, the conditional distribution of the response given the features remains the same across the training and test populations. A similar extension has been made for the jackknife+ method [Prinster et al.](#page-7-7) [\[2022\]](#page-7-7).

In this paper, we study the training-conditional coverage properties of the conformal prediction methods under covariate shift. We present concentration bounds of the training-conditional errors for weighted jackknife+ (JAW) [Prinster et al. \[2022\]](#page-7-7), full, and split conformal methods. In terms of distributions, we assume the existence of dQ/dP and consider two regimes: (1) $\frac{dQ}{dP}\right|_{\infty} < \infty$ and (2) $\|dQ/dP\|_{P,2} < \infty$, where Q and P denote the distribution of the test and training data, respectively. These two settings have also been considered in [Ma et al.](#page-7-8) [\[2023\]](#page-7-8) to study the optimality of the reproducing kernel Hilbert space (RKHS) regressions under the covariate shift setting, where it is assumed that the true regression function (i.e., $\mathbb{E}(Y | X = x)$) belongs to the RKHS. Regarding the training algorithm, no assumption is made for the split conformal method. However, full conformal and jackknife+ methods are analyzed under the assumption of *uniform stability* as explained below. Let $\hat{\mu}_{\textbf{D}_n}$ denote the predictor function estimated using the training data \textbf{D}_n . A training algorithm is called uniformly stable if,

$$
\|\hat{\mu}_{\mathbf{D}_n} - \hat{\mu}_{\mathbf{D}'_n}\|_{\infty} \leq \beta \tag{3}
$$

with $\beta = O(1/n)$ for any two datasets (D_n, D'_n) differing in one (training) data sam-ple [Bousquet and Elisseeff \[2002\]](#page-7-9). This is a stronger notion of algorithmic stability than the (m, n) stability assumed in [Liang and Barber](#page-7-5) [\[2023\]](#page-7-5). Nevertheless, uniform stability is satisfied by a class of regression models known as RKHS regression [Bousquet and Elisseeff \[2002\]](#page-7-9), i.e., regularized empirical risk minimization over an RKHS [Paulsen and Raghupathi \[2016\]](#page-7-10), [Schölkopf and Smola](#page-7-11) [\[2002\]](#page-7-11). Examples of reproducing kernel Hilbert spaces are certain Sobolev spaces of smooth functions [Wahba \[1990](#page-7-12)].

2 Background and Related Work

2.1 Full conformal and split conformal

Define the shorthand $D_n \cup (x, y) := ((X_1, Y_1), (X_2, Y_2), \dots, (X_n, Y_n), (x, y))$ and let $\hat{\mu}_{(x,y)} :=$ $T(D_n \cup (x, y))$ denote the regression function obtained by running the training algorithm T on $D_n \cup$ (x, y) . Define the score function $s(x', y'; \mu) := f(\mu(x'), y')$ via some arbitrary (measurable) cost function $f: \mathcal{Y} \times \mathcal{Y} \to \mathbb{R}$ and predictor function $\mu: \mathcal{X} \to \mathcal{Y}$. For instance, $s(x', y'; \mu) = |y' - \mu(x')|$ when $f(y, y') = |y - y'|$. Define the (training) multiset $\mathcal{D}_n := \{(X_i, Y_i) \in \mathcal{X} \times \mathcal{Y} : i \in [n]\}$ where $[n] := \{1, 2, ..., n\}$, and let

$$
\mathcal{S}(D;\mu) := \{ s(x',y';\mu) : (x',y') \in D \}, \quad D \in \cup_{n \geq 1} (\mathcal{X} \times \mathcal{Y})^n.
$$

Observe that if T is symmetric, i.e., $T(D)$ is invariant under permutations of the elements of the training data D, then the elements of $\mathcal{S}(\mathcal{D}_n \cup (X_{\text{test}}, Y_{\text{test}}); \hat{\mu}_{(X_{\text{test}}, Y_{\text{test}})})$ are exchangeable. Therefore,

$$
\mathbb{P}\left(s(X_{\text{test}}, Y_{\text{test}}; \hat{\mu}_{(X_{\text{test}}, Y_{\text{test}})}) \leq \hat{F}_{\mathcal{S}(\mathcal{D}_n \cup (X_{\text{test}}, Y_{\text{test}}); \hat{\mu}_{(X_{\text{test}}, Y_{\text{test}})})}(1-\alpha)\right) \geq 1-\alpha,
$$

where $\hat{F}_S^{-1}(1-\alpha)$ denotes the *empirical* quantile function with respect to S. Thus,

$$
\mathbb{P}(Y_{\text{test}} \in \hat{C}_{\alpha}^{\text{full}}(X_{\text{test}})) \ge 1 - \alpha,
$$

where the following confidence region is referred to as *full conformal* in the literature

$$
\hat{C}_{\alpha}^{\text{full}}(x) = \{ y : s(x, y; \hat{\mu}_{(x, y)}) \le \hat{F}_{\mathcal{S}(\mathcal{D}_{n} \cup (x, y); \hat{\mu}_{(x, y)})}^{-1}(1 - \alpha) \} \subseteq \{ y : s(x, y; \hat{\mu}_{(x, y)}) \le \hat{F}_{\mathcal{S}(\mathcal{D}_{n}; \hat{\mu}_{(x, y)}) \cup \{\infty\}}^{-1}(1 - \alpha) \},
$$
\n(4)

It is well-known that this approach can be computationally intensive when $\mathcal{Y} = \mathbb{R}$ since to find out whether $y \in \hat{C}_{\alpha}^{\text{full}}(x)$ one needs to train the model with the dataset including (x, y) . One simple way to alleviate this issue is to split the data into training and calibration datasets, namely $\mathcal{D}_n = \mathcal{D}^{\text{train}} \cup \mathcal{D}^{\text{cal}}$. Without loss of generality (when splitting), we assume $\mathbf{D}^{\text{train}} =$ $((X_1,Y_1),\ldots,(X_{n_0},Y_{n_0}))$ and $\mathbf{D}^{\text{cal}} = ((X_{n_0+1},Y_{n_0+1}),\ldots,(X_n,Y_n)).$ In the split conformal method, one first finds the regression predictor $\hat{\mu} = T(\hat{\mathbf{D}}^{\text{train}})$ and treats $\hat{\mu}$ as fixed. Note that the elements of $\mathcal{S}(\mathcal{D}^{\text{cal}} \cup (X_{\text{test}}, Y_{\text{test}}); \hat{\mu})$ are exchangeable. Hence, we get

$$
\mathbb{P}\left(s(X_{\text{test}}, Y_{\text{test}}; \hat{\mu}) \leq \hat{F}_{\mathcal{S}(\mathcal{D}^{\text{cal}} \cup (X_{\text{test}}, Y_{\text{test}}); \hat{\mu})}(1-\alpha)\right) \geq 1-\alpha.
$$

Hence, we have $\mathbb{P}(Y_{\text{test}} \in \hat{C}_{\alpha}^{\text{split}}(X_{\text{test}})) \geq 1 - \alpha$ for

$$
\hat{C}_{\alpha}^{\text{split}}(x) = \left\{ y : s(x, y; \hat{\mu}) \le \hat{F}_{\mathcal{S}(\mathcal{D}^{\text{cal}}; \hat{\mu}) \cup \{\infty\}}^{-1}(1 - \alpha) \right\}
$$
\n
$$
\supseteq \left\{ y : s(x, y; \hat{\mu}) \le \hat{F}_{\mathcal{S}(\mathcal{D}^{\text{cal}} \cup (x, y); \hat{\mu})}^{-1}(1 - \alpha) \right\}.
$$
\n(5)

2.2 Jackknife+

Although the split-conformal approach resolves the computational efficiency problem of the full conformal method, it is somewhat inefficient in using the data and may not be useful in situations where the number of samples is limited. A heuristic alternative has long been known in the literature, namely, jackknife or leave-one-out cross-validation that can provide a compromise between the full conformal and split conformal methods. In particular,

$$
\hat{C}_{\alpha}^{\mathbf{J}}(x) = \{ y : s(x, y; \hat{\mu}) \leq \hat{F}_{\mathcal{S}^{\mathbf{J}}}^{-1} (1 - \alpha) \},
$$

where $\hat{\mu} = T(\mathbf{D}_n)$ and $\mathcal{S}^J := \{ s(X_i, Y_i; \hat{\mu}^{-i}) : 1 \leq i \leq n \}$ and $\hat{\mu}^{-i} :=$ $T(((X_1,Y_1), \ldots, (X_{i-1}, Y_{i-1}), (X_{i+1}, Y_{i+1}), \ldots, (X_n, Y_n)))$. Despite its effectiveness, no general finite-sample guarantees are known for jackknife. Recently, [Barber et al. \[2021\]](#page-7-13) proposed jackknife+, a modified version of the jackknife for $\mathcal{Y} = \mathbb{R}$ and $f(y, y') = |y - y'|$, and established $(1 - 2\alpha)$ coverage lower bound for it. Let

$$
\mathcal{S}^{-}(x) = \{\hat{\mu}^{-i}(x) - |Y_i - \hat{\mu}^{-i}(X_i)| : i \in [n]\} \cup \{-\infty\},
$$

$$
\mathcal{S}^{+}(x) = \{\hat{\mu}^{-i}(x) + |Y_i - \hat{\mu}^{-i}(X_i)| : i \in [n]\} \cup \{\infty\}.
$$

The jackknife+ prediction interval is defined as

$$
\hat{C}_{\alpha}^{\mathbf{J}\mathbf{+}}(x) = [\hat{F}_{\mathcal{S}^{-}(x)}^{-1}(\alpha), \ \hat{F}_{\mathcal{S}^{+}(x)}^{-1}(1-\alpha)].
$$

In the same paper, an ϵ -inflated version of the jackknife+

$$
\hat{C}_{\alpha}^{J+,\epsilon}(x) = [\hat{F}_{\mathcal{S}^{-}(x)}^{-1}(\alpha) - \epsilon, \ \hat{F}_{\mathcal{S}^{+}(x)}^{-1}(1-\alpha) + \epsilon]. \tag{6}
$$

is proposed which has $1 - \alpha - 4\sqrt{\nu}$ coverage lower bound (instead of $1 - 2\alpha$), if the training procedure satisfies

$$
\max_{i\in [n]} \mathbb{P}(|\hat{\mu}(X_{\text{test}}) - \hat{\mu}^{-i}(X_{\text{test}})| > \epsilon) < \nu.
$$

Also, the jackknife+ has been generalized to CV+ for K-fold cross-validation, and $(1 - 2\alpha \sqrt{\frac{2}{|\mathcal{D}^{\text{train}}|}}$ coverage lower bound is established.

2.3 Conformal prediction under distribution shift

Covariate shift concerns the setting when the covariate distribution changes between training and test data, while the conditionals remain the same. Specifically, we have

$$
Z_i := (X_i, Y_i) \stackrel{\text{i.i.d.}}{\sim} P_X \times P_{Y|X} =: P \quad \text{training data};
$$

$$
Z_{\text{test}} := (X_{\text{test}}, Y_{\text{test}}) \sim Q_X \times P_{Y|X} =: Q \quad \text{test data}.
$$

This notion has been extensively studied in machine learning (e.g., see [Sugiyama et al.](#page-7-14) [\[2007\]](#page-7-14), [Quinonero-Candela et al.](#page-7-15) [\[2008\]](#page-7-15), [Sugiyama and Kawanabe](#page-7-16) [\[2012\]](#page-7-16), [Wen et al. \[2014\]](#page-7-17), [Reddi et al.](#page-7-18) [\[2015\]](#page-7-18), [Chen et al. \[2016](#page-7-19)] and the references therein).

To handle the covariate shift, a weighted version of conformal prediction was first proposed in the seminal work by Tibshirani et al. [Tibshirani et al. \[2019\]](#page-7-6). The key assumption is that the ratio of test to training covariate, dQ/dP , needs to be known; we follow the same assumption in this work. This reweighting scheme has been extended to various settings in conformal prediction [Podkopaev and Ramdas](#page-7-20) [\[2021\]](#page-7-20), [Lei and Candès \[2021](#page-7-21)], [Fannjianga et al.](#page-8-0), [Guan \[2023\]](#page-8-1). The general domain adaption problem has also been an active area from a causal perspective (e.g., [Zhang et al. \[2013a\]](#page-8-2), [Peters et al. \[2016](#page-8-3)], [Gong et al.](#page-8-4) [\[2016\]](#page-8-4), [Chen and Bühlmann](#page-8-5) [\[2021\]](#page-8-5), [Du and Xiang](#page-8-6) [\[2023\]](#page-8-6)).

The weighted procedures require computing a weight associated with each sample,

$$
w_i = \frac{\frac{dQ}{dP}(Z_i)}{\frac{dQ}{dP}(Z_{\text{test}}) + \sum_{i \in \mathcal{I}} \frac{dQ}{dP}(Z_i)}, \ i \in \mathcal{I} \quad \text{and} \quad w_{\text{test}} = \frac{\frac{dQ}{dP}(Z_{\text{test}})}{\frac{dQ}{dP}(Z_{\text{test}}) + \sum_{i \in \mathcal{I}} \frac{dQ}{dP}(Z_i)}
$$

where $\mathcal{I} = [n]$ for full conformal and jackknife+, and $\mathcal{I} \subset [n]$ for the split conformal method which corresponds to the calibration (or hold-out) dataset. In particular, to generalize the full and split conformal methods to distribution shift setting, according to [Tibshirani et al. \[2019\]](#page-7-6) one needs to replace the empirical CDF used in construction of [\(4\)](#page-2-0) and [\(5\)](#page-2-1)

$$
\hat{F}_{\mathcal{S}}(t) = \frac{1}{n+1} \sum_{s \in \mathcal{S}} \mathbf{1} \{ s \le t \}, \quad \mathcal{S} = \{ s(Z_i) : i \in \mathcal{I} \} \cup \{ \infty \}
$$

by the weighted version

$$
F_{Q,n}(t) = w_{\text{test}} \mathbf{1}\{t = \infty\} + \sum_{i \in \mathcal{I}} w_i \mathbf{1}\{s(Z_i) \le t\}.
$$

Similarly, to extend the jackknife+ method, [Prinster et al.](#page-7-7) [\[2022\]](#page-7-7) proposes replacing $\hat{F}_{\mathcal{S}^-}$ and $\hat{F}_{\mathcal{S}^+}$ by their corresponding weighted versions where w_{test} is associated to $-\infty$ and ∞ , respectively.

3 Conditional Coverage Guarantees

3.1 Split conformal

The split conformal training-conditional coverage guarantee under the i.i.d. setting has been studied in [Vovk \[2012\]](#page-6-3), [Bian and Barber \[2023\]](#page-7-4). The proof in this setting relies on the Dvoretzky–Kiefer–Wolfowitz (DKW) inequality. The following theorem concerns training-conditional coverage guarantee under covariate shift, where the extension is made by developing a weighted version of the DKW inequality. Let $\mathbf{D}_n = (Z_1, \ldots, Z_n) \in (\mathcal{X} \times \mathcal{Y})^n$ denote an *n*-tuple of data points containing both the training and calibration data sets. Define the training conditional probability of error

$$
P_e^{\text{split}}(\mathbf{D}_n) = \mathbb{P}\left(Y_{\text{test}} \notin \hat{C}_{\alpha}^{\text{split}}(X_{\text{test}}) \middle| \mathbf{D}_n\right).
$$
\n(7)

Theorem 1 Let $m < n$ denote the size of the calibration data set. Assume $Q \ll P$ and $dQ/dP \le$ $B < \infty$ *. Then,*

$$
\mathbb{P}\left(P_e^{split}(\mathbf{D}_n) > \alpha + \left(\sqrt{2B\log(4/\delta)} + 3C\right)\sqrt{\frac{B}{m}}\right) \le \delta.
$$

for all $\delta > 0$ *, where* $C > 0$ *is a universal constant. The probability is taken with respect to P since each entry in* \mathbf{D}_n *follows* P *.*

Remark 1 *Relaxing the assumption* $dQ/dP \leq B$ *to* $\|dQ/dP\|_{P,2} \leq B$ *, we get*

$$
\mathbb{P}\left(P_e^{split}(\mathbf{D}_n) > \alpha + \frac{2B(3C+1)}{\delta\sqrt{m}}\right) \le \delta, \qquad 0 < \delta \le 1.
$$
 (8)

For simulations, we refer the readers to the experiments on Airfoil data [Tibshirani et al.](#page-7-6) [\[2019\]](#page-7-6), which shows the empirical distribution of $P_e^{\text{split}}(\mathbf{D}_n)$. Note that the theoretical guarantees in [Tibshirani et al. \[2019\]](#page-7-6) only concern the marginal coverage, e.g., $\mathbb{E}[P_e^{\text{split}}(\mathbf{D}_n)] = \mathbb{P}(Y_{\text{test}} \notin$ $\hat{C}_{\alpha}^{\text{split}}(X_{\text{test}})$) for split conformal.

3.2 Full conformal and jackknife+

Let $\mu_{\beta} : \mathcal{X} \to \mathbb{R}$ denote a predictor function parameterized by $\beta \in \mathbb{R}^p$. By a slight abuse of notation, let the map $T: \bigcup_{n\geq 1} (\mathcal{X} \times \mathcal{Y})^n \to \mathbb{R}^p$ denote a training algorithm for estimating β , hence, $\hat{\beta}_n = T(\mathbf{D}_n)$ In this case, we have $\hat{\mu}_{\mathbf{D}_n} = \mu_{\hat{\beta}_n}$.

Assumption 1 (Uniform stability) *For all* $i \in [n]$ *, we have*

$$
\sup_{z_1,\ldots,z_n} \|\mu_{T(z_1,\ldots,z_{i-1},z_{i+1}\ldots,z_n)} - \mu_{T(z_1,\ldots,z_i,\ldots,z_n)}\|_{\infty} \le \frac{c_n}{2}.
$$

In the case of the ridge regression [Hoerl and Kennard](#page-8-7) [\[1970\]](#page-8-7) with $\mathcal{Y} = [-I, I]$ and $\mathcal{X} = \{x :$ $||x||_2 \le b$, this assumption holds with $c_n = 16 b^2 I^2/(\lambda n)$, where λ denotes the regularization parameter. See [Bousquet and Elisseeff \[2002\]](#page-7-9) for the general result on the uniform stability of RKHS regression.

Assumption 2 *The model is bi-Lipschitz (Lipeomorphism) in parameters, i.e.,*

$$
\kappa_1 \|\beta - \beta'\|_{\infty} \le \|\mu_{\beta} - \mu_{\beta'}\|_{\infty} \le \kappa_2 \|\beta - \beta'\|_{\infty},
$$

with $\kappa_1 > 0$ *and* $\kappa_2 < \infty$ *.*

Remark 2 *It is worth noting that if the parameter space* $\Theta \subseteq \mathbb{R}^p$ *is compact,* $\Phi: U \to L^{\infty}(\mathcal{X})$ *given by* $\beta \mapsto \mu_\beta$ *is continuously differentiable for some open* $U \supseteq \Theta$ *, then* $\kappa_2 < \infty$ *. Moreover, the inverse function theorem (for Banach spaces), gives the sufficient condition under which the inverse is continuously differentiable over* $\Phi(U)$ *and hence* $\kappa_1 > 0$ *.*

In the case of linear regression with $\mathcal{X} = \{x : ||x||_2 \leq b\}$, one can verify that Assumption [2](#page-4-0) holds with $\kappa_1 = b$ and $\kappa_2 = \sqrt{p}b$. Let $\overline{\beta}_n = \mathbb{E} \hat{\beta}_n$, $\hat{\beta}_{-i} = T(Z_1, \dots, Z_{i-1}, Z_{i+1}, \dots, Z_n)$ with $Z_i = (X_i, Y_i)$ and $\overline{\beta}_{-i} = \mathbb{E} \hat{\beta}_{-i}$. Define

$$
F^{(n-1)}(t) := \mathbb{P}_{Z_1 \sim P}\left(\left|Y_1 - \mu_{\overline{\beta}_{-1}}(X_1)\right| \le t\right),
$$

$$
F_Q^{(n-1)}(t) := \mathbb{P}_{Z_1 \sim Q}\left(\left|Y_1 - \mu_{\overline{\beta}_{-1}}(X_1)\right| \le t\right),
$$

Assumption 3 (Bounded density) $F^{\prime (n)} < L_n$ and $F^{\prime (n)}_Q < L_{Q,n}$.

We introduce the following shorthand:

$$
A(n, p, \epsilon) := 2 \kappa_2 c_{n-1} \left(\frac{1}{\kappa_1} + \sqrt{\frac{n}{2 \kappa_1^2} \log \frac{2p}{\epsilon}} \right).
$$

Theorem 2 (Jackknife+ under exchangeability) *Under Assumptions* $1-\lambda$, for all $\epsilon, \delta > 0$, it *holds that*

$$
\mathbb{P}\left(P_e^{J*}(\mathbf{D}_n) > \alpha + \sqrt{\frac{\log(2/\delta)}{2n}} + L_{n-1}A(n, p, \epsilon)\right) \leq \epsilon + \delta.
$$

Theorem 3 (Jackknife+ under covariate shift) Assume $Q \ll P$ and $dQ/dP \leq B$. Under As*sumptions* $1-3$ *, for all* $\epsilon, \delta > 0$ *, it holds that*

$$
\mathbb{P}\left(P_e^{J*}(\mathbf{D}_n) > \alpha + \left(\sqrt{2B\log 4/\delta} + 3C\right)\sqrt{\frac{B}{n}} + L_{Q,n-1}A(n, p, \epsilon)\right) \le \epsilon + \delta,
$$

where C is a universal constant and $A(n, p, \epsilon)$ *is the same as in Theorem* [2.](#page-4-3)

Using the same arguments as in the proof of this theorem, one can get a coverage bound for the CV+ as well. Unlike [\(2\)](#page-1-0) which is meaningful only if the number of samples in each fold m is large, the bound we present in the following corollary is suitable for cases where $m/n \to 0$.

Corollary [1](#page-4-1) (CV+) *Under Assumptions 1[—3,](#page-4-2) for all* $\epsilon, \delta > 0$ *, it holds that*

$$
\mathbb{P}\left(P_e^{CV+}(\mathbf{D}_n) > \alpha + \sqrt{\frac{\log(2/\delta)}{2n}} + 2m L_{n-m} \kappa_2 c_{n-m} \left(\frac{1}{\kappa_1} + \sqrt{\frac{n}{2\kappa_1^2} \log \frac{2p}{\epsilon}}\right)\right) \le \epsilon + \delta.
$$

The following theorem concerns the training-conditional guarantees for the full conformal prediction. We again introduce a shorthand:

$$
E(n, p, \epsilon) := c_{n+1} + \sqrt{2n \log \frac{2p}{\epsilon}} \frac{\kappa_2 c_n}{\kappa_1}.
$$

Theorem 4 (Full conformal under exchangeability) *Under Assumptions [1—](#page-4-1)[3,](#page-4-2) for all* $\epsilon, \delta > 0$, *it holds that*

$$
\mathbb{P}\left(P_e^{\text{full}}(\mathbf{D}_n) > \alpha + \sqrt{\frac{\log(2/\delta)}{2n}} + L_n E(n, p, \epsilon)\right) \leq \epsilon + \delta.
$$

Theorem 5 (Full conformal under covariate shift) *Assume* $Q \ll P$ *and* $dQ/dP \leq B$ *. Under Assumptions [1](#page-4-1)[—3,](#page-4-2) for all* $\epsilon, \delta > 0$ *, it holds that*

$$
\mathbb{P}\left(P_e^{\text{full}}(\mathbf{D}_n) > \alpha + \left(\sqrt{2B\log(4/\delta)} + 3C\right)\sqrt{\frac{B}{n}} + L_{Q,n}E(n, p, \epsilon)\right) \le \epsilon + \delta.
$$

where C is a universal constant and $E(n, p, \epsilon)$ *is the same as in Theorem [4.](#page-5-1)*

We note that similar to Remark [1,](#page-4-4) one can relax assumption $dQ/dP \leq B$ to $||dQ/dP||_{P,2} \leq B$ and in this case the bounds for the jackknife+ and full conformal methods hold with the slow rate $O(1/(\delta\sqrt{n}))$ instead of $O(\log{(1/\delta)}/\sqrt{n})$.

4 Discussion

We have presented training-conditional coverage guarantees for conformal prediction methods under two covariate shift regimes: (1) $\|dQ/dP\|_{\infty} < \infty$ and (2) $\|dQ/dP\|_{P,2} < \infty$. The split conformal method has been analyzed for the black-box training algorithm while the full conformal and jackknife+ have been analyzed under three assumptions. Although Assumptions [1](#page-4-1) and [2](#page-4-0) have been verified only for the ridge regression in this paper, we conjecture that they are satisfied by a truncated version of the general RKHS regressions which we leave for future research. Truncated and sketched versions of the RKHS models have been extensively studied in the previous literature from the computational efficiency perspective (see [Amini](#page-8-8) [\[2021\]](#page-8-8), [Williams and Seeger](#page-8-9) [\[2000\]](#page-8-9), [Zhang et al. \[2013b\]](#page-8-10), [Alaoui and Mahoney \[2015\]](#page-8-11), [Cortes et al. \[2010\]](#page-8-12) and references therein). The results in this paper quantify the training sample size (or calibration sample size for split conformal) needed for the coverage under covariate shift.

The (m, n) -stability parameters were recently introduced in [Liang and Barber](#page-7-5) [\[2023\]](#page-7-5) and used to compute training-conditional coverage bounds for inflated full conformal and jackknife+ prediction

intervals under exchangeability. Unlike uniform stability which is a distribution-free property of a training process, (m, n) -stability depends on both the training algorithm and the distributions of the data as follows

$$
\psi_{m,n}^{\text{out}} = \mathbb{E}_{\mathbf{D}_{n+m}} |\hat{\mu}_{\mathbf{D}_n}(X_{\text{test}}) - \hat{\mu}_{\mathbf{D}_{n+m}}(X_{\text{test}})|,
$$
\n(9)

$$
\psi_{m,n}^{\text{in}} = \mathbb{E}_{\mathbf{D}_{n+m}} |\hat{\mu}_{\mathbf{D}_n}(X_1) - \hat{\mu}_{\mathbf{D}_{n+m}}(X_1)|,\tag{10}
$$

where $\hat{\mu}_{\mathbf{D}_n} = T(\mathbf{D}_n)$ and $X_{\text{test}} \perp \!\!\! \perp \mathbf{D}_{n+m}$ with $\mathbf{D}_{n+m} = ((X_1, Y_1), ..., (X_{n+m}, Y_{n+m}))$. Although weaker than uniform stability, these parameters are yet not well-understood in a practical sense. To elaborate on the difference between this approach and uniform stability, we evaluate these bounds for the ridge regression under exchangeability.

Assume $\mathcal{X} = \{x : ||x|| \leq b\}$ and $\mathcal{Y} = [-I, I]$. As stated in the previous section, this regression model satisfies $c_n = 16 b^2 I^2/(\lambda n)$, $\kappa_1 = b$ and $\kappa_2 = \sqrt{p} b$. Hence, we get the following bound for both full conformal and jackknife+ methods,

$$
\mathbb{P}\left(P_e^{\text{full}}(\mathbf{D}_n) > \alpha + O\big(n^{-1/2}\big(\sqrt{\log(1/\delta)} + \sqrt{p\log(2p/\epsilon)}\big)\big)\right) \le \epsilon + \delta.
$$

On the other hand, the following bound is proposed for the γ-inflated jackknife+ in [Liang and Barber](#page-7-5) [\[2023\]](#page-7-5),

$$
\mathbb{P}\left(P_e^{\mathbf{J}+\gamma}(\mathbf{D}_n) > \alpha + 3\sqrt{\frac{\log(1/\delta)}{\min(m,n)}} + 2\sqrt[3]{\frac{\psi_{m,n-1}^{\text{out}}}{\gamma}}\right) \le 3\delta + \sqrt[3]{\frac{\psi_{m,n-1}^{\text{out}}}{\gamma}},\tag{11}
$$

for all $m \ge 1$. We get $\psi_{m,n}^{\text{out}} = O(mc_n)$ since $\psi_{1,n}^{\text{out}} \le c_{n+1}/2$ by definition [\(9\)](#page-6-5) and Assumption [1,](#page-4-1) and $\psi_{m,n}^{\text{out}} \leq \sum_{k=n}^{n+m-1} \psi_{1,k}^{\text{out}}$ holds according to in [Liang and Barber \[2023](#page-7-5), Lemma 5.2]. Substituting for $\psi_{m,n-1}^{\text{out}}$ in bound [\(11\)](#page-6-6), we obtain

$$
\mathbb{P}\left(P_e^{J+\gamma}(\mathbf{D}_n) > \alpha + O\left(\sqrt{\frac{\log(1/\delta)}{\min(m,n)}} + \sqrt[3]{\frac{m\,c_{n-1}}{\gamma}}\right)\right) \le 3\delta + O\left(\sqrt[3]{\frac{mc_{n-1}}{\gamma}}\right). \tag{12}
$$

Letting $m^{-1/2} = (m/n)^{1/3}$ to balance the two terms $\sqrt{\frac{\log(1/\delta)}{\min(m,n)}}$ and $\sqrt[3]{mc_{n-1}/\gamma}$, we get $m =$ $n^{2/5}$. By plugging $m = n^{2/5}$ into [\(12\)](#page-6-7), we have

$$
\mathbb{P}\left(P_e^{1+\gamma}(\mathbf{D}_n) > \alpha + O\big(n^{-1/5}\big(\sqrt{\log(1/\delta)} + \gamma^{-1/3}\big)\big)\right) \le 3\delta + O\left(n^{-1/5}\gamma^{-1/3}\right). \tag{13}
$$

This bound, although dimension-free (i.e., does not depend on p), is very slow with respect to the sample size. In [Liang and Barber](#page-7-5) [\[2023\]](#page-7-5), the same bound as [\(11\)](#page-6-6) is established for γ -inflated full conformal method except with $\psi_{m-1,n+1}^{\text{in}}$ instead of $\psi_{m,n-1}^{\text{out}}$. Hence, the same bound as [\(13\)](#page-6-8) can be obtained for the γ -inflated full conformal method via $\psi^{\text{in}}_{m,n} = O(mc_n)$.

References

- Vladimir Vovk, Alexander Gammerman, and Glenn Shafer. *Algorithmic learning in a random world*, volume 29. Springer, 2005.
- Glenn Shafer and Vladimir Vovk. A tutorial on conformal prediction. *Journal of Machine Learning Research*, 9(3), 2008.
- Vladimir Vovk, Ilia Nouretdinov, and Alex Gammerman. On-line predictive linear regression. *The Annals of Statistics*, pages 1566–1590, 2009.
- Vladimir Vovk. Conditional validity of inductive conformal predictors. In *Asian conference on machine learning*, pages 475–490. PMLR, 2012.
- Rina Foygel Barber, Emmanuel J Candes, Aaditya Ramdas, and Ryan J Tibshirani. The limits of distribution-free conditional predictive inference. *Information and Inference: A Journal of the IMA*, 10(2):455–482, 2021.
- Jing Lei and Larry Wasserman. Distribution-free prediction bands for non-parametric regression. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 76(1):71–96, 2014.
- Christopher Jung, Georgy Noarov, Ramya Ramalingam, and Aaron Roth. Batch multivalid conformal prediction. *arXiv preprint arXiv:2209.15145*, 2022.
- Isaac Gibbs, John J Cherian, and Emmanuel J Candès. Conformal prediction with conditional guarantees. *arXiv preprint arXiv:2305.12616*, 2023.
- Vladimir Vovk, David Lindsay, Ilia Nouretdinov, and Alex Gammerman. Mondrian confidence machine. *Technical Report*, 2003.
- Michael Bian and Rina Foygel Barber. Training-conditional coverage for distribution-free predictive inference. *Electronic Journal of Statistics*, 17(2):2044–2066, 2023.
- Ruiting Liang and Rina Foygel Barber. Algorithmic stability implies training-conditional coverage for distribution-free prediction methods. *arXiv preprint arXiv:2311.04295*, 2023.
- Ryan J Tibshirani, Rina Foygel Barber, Emmanuel Candes, and Aaditya Ramdas. Conformal prediction under covariate shift. *Advances in neural information processing systems*, 32, 2019.
- Drew Prinster, Anqi Liu, and Suchi Saria. Jaws: Auditing predictive uncertainty under covariate shift. *Advances in Neural Information Processing Systems*, 35:35907–35920, 2022.
- Cong Ma, Reese Pathak, and Martin J Wainwright. Optimally tackling covariate shift in RKHSbased nonparametric regression. *The Annals of Statistics*, 51(2):738–761, 2023.
- Olivier Bousquet and André Elisseeff. Stability and generalization. *The Journal of Machine Learning Research*, 2:499–526, 2002.
- Vern I Paulsen and Mrinal Raghupathi. *An introduction to the theory of reproducing kernel Hilbert spaces*, volume 152. Cambridge university press, 2016.
- Bernhard Schölkopf and Alexander J Smola. *Learning with kernels: support vector machines, regularization, optimization, and beyond*. MIT press, 2002.
- Grace Wahba. *Spline models for observational data*. SIAM, 1990.
- Rina Foygel Barber, Emmanuel J Candes, Aaditya Ramdas, and Ryan J Tibshirani. Predictive inference with the jackknife+. 2021.
- Masashi Sugiyama, Matthias Krauledat, and Klaus-Robert Müller. Covariate shift adaptation by importance weighted cross validation. *Journal of Machine Learning Research*, 8(5), 2007.
- Joaquin Quinonero-Candela, Masashi Sugiyama, Anton Schwaighofer, and Neil D Lawrence. *Dataset shift in machine learning*. MIT Press, 2008.
- Masashi Sugiyama and Motoaki Kawanabe. *Machine learning in non-stationary environments: Introduction to covariate shift adaptation*. 2012.
- Junfeng Wen, Chun-Nam Yu, and Russell Greiner. Robust learning under uncertain test distributions: Relating covariate shift to model misspecification. In *International Conference on Machine Learning*, pages 631–639. PMLR, 2014.
- Sashank Reddi, Barnabas Poczos, and Alex Smola. Doubly robust covariate shift correction. In *Proceedings of the AAAI conference on artificial intelligence*, volume 29, 2015.
- Xiangli Chen, Mathew Monfort, Anqi Liu, and Brian D Ziebart. Robust covariate shift regression. In *Artificial Intelligence and Statistics*, pages 1270–1279. PMLR, 2016.
- Aleksandr Podkopaev and Aaditya Ramdas. Distribution-free uncertainty quantification for classification under label shift. In *Uncertainty in artificial intelligence*, pages 844–853. PMLR, 2021.
- Lihua Lei and Emmanuel J Candès. Conformal inference of counterfactuals and individual treatment effects. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 83(5):911–938, 2021.
- Clara Fannjianga, Stephen Batesa, Anastasios Angelopoulosa, Jennifer Listgartena, and Michael I Jordana. Conformal prediction for the design problem.
- Leying Guan. Localized conformal prediction: A generalized inference framework for conformal prediction. *Biometrika*, 110(1):33–50, 2023.
- Kun Zhang, Bernhard Schölkopf, Krikamol Muandet, and Zhikun Wang. Domain adaptation under target and conditional shift. In *International Conference on Machine Learning*, pages 819–827. PMLR, 2013a.
- Jonas Peters, Peter Bühlmann, and Nicolai Meinshausen. Causal inference by using invariant prediction: identification and confidence intervals. *Journal of the Royal Statistical Society. Series B (Statistical Methodology)*, pages 947–1012, 2016.
- Mingming Gong, Kun Zhang, Tongliang Liu, Dacheng Tao, Clark Glymour, and Bernhard Schölkopf. Domain adaptation with conditional transferable components. In *International Conference on Machine Learning*, pages 2839–2848. PMLR, 2016.
- Yuansi Chen and Peter Bühlmann. Domain adaptation under structural causal models. *Journal of Machine Learning Research*, 22:1–80, 2021.
- Kang Du and Yu Xiang. Learning invariant representations under general interventions on the response. *IEEE Journal on Selected Areas in Information Theory*, 2023.
- Arthur E Hoerl and Robert W Kennard. Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics*, 12(1):55–67, 1970.
- Arash A Amini. Spectrally-truncated kernel ridge regression and its free lunch. *Electronic Journal of Statistics*, 15(2):3743–3761, 2021.
- Christopher Williams and Matthias Seeger. Using the nyström method to speed up kernel machines. *Advances in neural information processing systems*, 13, 2000.
- Yuchen Zhang, John Duchi, and Martin Wainwright. Divide and conquer kernel ridge regression. In *Conference on learning theory*, pages 592–617. PMLR, 2013b.
- Ahmed Alaoui and Michael W Mahoney. Fast randomized kernel ridge regression with statistical guarantees. *Advances in neural information processing systems*, 28, 2015.
- Corinna Cortes, Mehryar Mohri, and Ameet Talwalkar. On the impact of kernel approximation on learning accuracy. In *Proceedings of the thirteenth international conference on artificial intelligence and statistics*, pages 113–120. JMLR Workshop and Conference Proceedings, 2010.
- Aad W Van Der Vaart and Jon A Wellner. *Weak convergence and empirical processes: with applications to statistics*. Springer, 1996.
- Colin McDiarmid et al. On the method of bounded differences. *Surveys in Combinatorics*, 141(1): 148–188, 1989.
- JE Yukich. Laws of large numbers for classes of functions. *Journal of multivariate analysis*, 17(3): 245–260, 1985.
- John Lafferty, Han Liu, and Larry Wasserman. Concentration of measure. *On-line. Available: https://www.stat.cmu.edu/ larry/=sml/Concentration.pdf*, 2008.
- Aryeh Dvoretzky, Jack Kiefer, and Jacob Wolfowitz. Asymptotic minimax character of the sample distribution function and of the classical multinomial estimator. *The Annals of Mathematical Statistics*, pages 642–669, 1956.

A Technical Lemmas: Weighted DKW Inequalities

Let $F_Q(x) = \mathbb{P}_{Z \sim Q}(s(Z) \leq x)$ where $s(\cdot)$ denotes some fixed score function. Define,

$$
\hat{F}_{Q,n}(x) := \sum_{i \in [n]} \hat{w}_i \mathbf{1} \{ s(Z_i) \le x \}, \qquad \hat{w}_i = \frac{\frac{dQ}{dP}(Z_i)}{\sum_{i \in [n]} \frac{dQ}{dP}(Z_i)}.
$$
\n(14)

.

where $Z_i \stackrel{\text{i.i.d.}}{\sim} P$.

Lemma 1 (Bounded likelihood ratio) Assume $Q \ll P$ and $dQ/dP \leq B$. Then, for all $\delta > 0$

$$
\mathbb{P}\left(\sup_{x\in\mathbb{R}}\left|\hat{F}_{Q,n}(x)-F_Q(x)\right|>\delta+3\,C\sqrt{\frac{B}{n}}\right)\leq 4e^{-n\delta^2/(2B^2)}.
$$

where C > 0 *is some universal constant.*

Proof: By substituting the formula for \hat{w}_i in the definition of $\hat{F}_{Q,n}$, we get

$$
\hat{F}_{Q,n} = \frac{\frac{1}{n} \sum_{i \in [n]} \frac{dQ}{dP}(Z_i) \mathbf{1}\{s(Z_i) \le x\}}{\frac{1}{n} \sum_{i \in [n]} \frac{dQ}{dP}(Z_i)}
$$

Hence,

$$
\sup_{x \in \mathbb{R}} |\hat{F}_{Q,n}(x) - F_Q(x)| \le \sup_{x \in \mathbb{R}} \left| \frac{1}{n} \sum_{i \in [n]} \frac{dQ}{dP}(Z_i) \mathbf{1}\{s(Z_i) \le x\} - F_Q(x) \right| + \left| \frac{1}{\frac{1}{n} \sum_{i \in [n]} \frac{dQ}{dP}(Z_i)} - 1 \right| \sup_{x \in \mathbb{R}} \left| \frac{1}{n} \sum_{i \in [n]} \frac{dQ}{dP}(Z_i) \mathbf{1}\{s(Z_i) \le x\} \right|
$$

$$
= \sup_{x \in \mathbb{R}} \left| \frac{1}{n} \sum_{i \in [n]} \frac{dQ}{dP}(Z_i) \mathbf{1}\{s(Z_i) \le x\} - F_Q(x) \right| + \left| 1 - \frac{1}{n} \sum_{i \in [n]} \frac{dQ}{dP}(Z_i) \right| \tag{15}
$$

Regarding the first term we have,

$$
\sup_{x \in \mathbb{R}} \left| \frac{1}{n} \sum_{i \in [n]} \frac{dQ}{dP}(Z_i) \mathbf{1}\{s(Z_i) \le x\} - F_Q(x) \right| = \sup_{x \in \mathbb{R}} \left| \int \frac{dQ}{dP}(z) \mathbf{1}\{s(z) \le x\} \, d(P_n - P) \right|
$$

$$
= \sup_{x \in \mathbb{R}} \left| \int f_x \, d(P_n - P) \right|
$$

where $f_x \in L^1(\mathcal{X} \times \mathcal{Y}, P)$ is defined as $f_x(z) := \frac{dQ}{dP}(z) \mathbf{1}\{s(z) \leq x\}$, and $P_n(A) =$ $\frac{1}{n} \sum_{i \in [n]} \mathbf{1} \{ Z_i \in A \}$ is the empirical measure.

We note,

$$
1 \leq ||dQ/dP||_{P,2}^2 = \int (dQ/dP)^2 \, dP \leq ||dQ/dP||_{\infty} \int dQ/dP \, dP \leq B.
$$

according to the Holder's inequality. Hence, $||dQ/dP||_{P,2} \le \sqrt{B}$ and b[yVan Der Vaart and Wellner](#page-8-13) [\[1996,](#page-8-13) Theorem 2.14.2] we have

$$
\mathbb{E}\left(\sup_{x\in\mathbb{R}}\left|\int f_x\,d(P_n-P)\right|\right)\leq C\sqrt{\frac{B}{n}}\int_0^1\sqrt{1+\log N_{[]}(\epsilon\|dQ/dP\|_{P,2},\mathcal{F},L^2(P))}\,d\epsilon,\quad(16)
$$

with some universal constant $C > 0$, and $N_{[]} (\epsilon ||dQ/dP||_{P,2}, \mathcal{F}, L^2(P))$ denotes the bracketting number for function class $\mathcal{F} = \{f_x : x \in \mathbb{R}\}.$ It remains to compute an upper bound for $N_{[]}$ $(\epsilon, \mathcal{F}, L^2(P))$. Let $t_0 = -\infty$ and define,

$$
t_{i+1} := \sup \left\{ t \in \mathbb{R} : \int f^2(z) \mathbf{1} \{ s(z) \in (t_i, t] \} \, dP < \epsilon^2 \right\},\tag{17}
$$

where $f = dQ/dP$. We note that there exists $\ell < \infty$ such that $t_{\ell} = \infty$. This is true since $t_{j+1} < \infty$ implies

$$
\int f^2(z)\mathbf{1}\{s(z)\in(t_i,t_{i+1}]\}\,dP\geq\epsilon^2,\quad 0\leq i\leq j,
$$

according to [\(17\)](#page-9-0) and dominated convergence theorem. Hence,

$$
(j+1)\epsilon^2 \le \sum_{i=0}^j \int f^2(z)\mathbf{1}\{s(z) \in (t_i, t_{i+1}]\} \, dP \le \int f^2 \, dP = \|dQ/dP\|_{P,2}^2,\tag{18}
$$

and therefore $j + 1 \leq (\|dQ/dP\|_{P,2}/\epsilon)^2$. Let $k = \min{\{\ell \in \mathbb{Z} : t_{\ell} = \infty\}}$. Now define ϵ -brackets as follows

$$
b_i = \{g(z) : f(z)\mathbf{1}\{s(z) \le t_i\} \le g(z) \le f(z)\mathbf{1}\{s(z) < t_{i+1}\}\}
$$

where $0 \le i \le k-1$. Clearly, $\mathcal{F} \subset \bigcup_{i=0}^k b_i$ with $b_k = \{f\}$. Brackets b_0, \ldots, b_{k-1} have size

$$
\left(\int f^2(z)\mathbf{1}\{s(z)\in(t_i,t_{i+1})\} dP\right)^{1/2} \leq \epsilon, \quad 0 \leq i \leq k-1
$$

according to [\(17\)](#page-9-0) and bracket b_k has size 0. Hence, we have $k+1$ brackets in total with size smaller than ϵ . By [\(18\)](#page-10-0), we have $k-1 \leq (\|dQ/dP\|_{P,2}/\epsilon)^2$ which implies that $N_{[]}(\epsilon, \mathcal{F}, L^2(P)) \leq k+1 \leq$ $2 + (||dQ/dP||_{P,2}/\epsilon)^2$. Plugging this into [\(18\)](#page-10-0), we get

$$
N_{[]} (\epsilon ||dQ/dP||_{P,2}, \mathcal{F}, L^2(P)) \leq 2 + \frac{1}{\epsilon^2}.
$$

Therefore,

$$
\mathbb{E}\left(\sup_{x\in\mathbb{R}}\left|\int f_x\,d(P_n-P)\right|\right)\leq 3\,C\sqrt{\frac{B}{n}}=C'\sqrt{\frac{B}{n}},\tag{19}
$$

since

$$
\int_0^1 \sqrt{1 + \log(2 + 1/\epsilon^2)} \le 1 + \left(\int_0^1 \log(2 + 1/\epsilon^2)\right)^{1/2} \le 1 + \left(2\int_0^1 (\epsilon^2 - \log \epsilon)\right)^{1/2} \le 3.
$$

Now let P'_n denote the empirical measure obtained after replacing Z_i by some arbitrary Z'_i . In this case, we observe that

$$
\left| \sup_{x \in \mathbb{R}} \left| \int f_x d(P_n - P) \right| - \sup_{x \in \mathbb{R}} \left| \int f_x d(P_n' - P) \right| \right| \leq \sup_{x \in \mathbb{R}} \left| \int f_x d(P_n - P_n') \right|
$$

$$
\leq \frac{1}{n} \sup_{x \in \mathbb{R}} |f_x(Z_i) - f_x(Z_i')| \leq \frac{B}{n}.
$$

Therefore, by McDiarmid's inequality [McDiarmid et al.](#page-8-14) [\[1989\]](#page-8-14) we get

$$
\mathbb{P}\left(\left|\sup_{x\in\mathbb{R}}\left|\int f_x d(P_n-P)\right|-\mathbb{E}\left(\sup_{x\in\mathbb{R}}\left|\int f_x d(P_n-P)\right|\right)\right|>\delta\right)\leq 2e^{-2n\delta^2/B^2}.
$$

Combining with [\(19\)](#page-10-1), we obtain,

$$
\mathbb{P}\left(\sup_{x\in\mathbb{R}}\left|\int f_x\,d(P_n-P)\right|>\delta+C'\sqrt{\frac{B}{n}}\right)\leq 2e^{-2n\delta^2/B^2}.
$$

Back to [\(15\)](#page-9-1), for the second term Hoeffding's inequality implies

$$
\mathbb{P}\left(\left|1-\frac{1}{n}\sum_{i\in[n]}\frac{dQ}{dP}(Z_i)\right|>\delta\right)\leq 2e^{-2n\delta^2/B^2}.\tag{20}
$$

Combining the bounds for the two terms, we obtain,

$$
\mathbb{P}\left(\sup_{x\in\mathbb{R}}\left|\hat{F}_{Q,n}(x)-F_Q(x)\right|>\delta+C'\sqrt{\frac{B}{n}}\right)\leq 4e^{-n\delta^2/(2B^2)}.
$$

The following version relaxes the assumption $dQ/dP \leq B$ to $||dQ/dP||_{P,2} \leq K$ at the cost of slower rates.

Lemma 2 (Alternative version: bounded second moment) *Assume* $Q \ll P$ *and* $||dQ/dP||_{P,2} \le$ *K. Then, for all* $\delta > 0$

$$
\mathbb{P}\left(\sup_{x\in\mathbb{R}}\left|\hat{F}_{Q,n}(x)-F_Q(x)\right|>\delta\right)\leq \frac{6\,CK}{\delta\sqrt{n}}+\frac{4(K^2-1)}{n\delta^2}.
$$

where C > 0 *is some universal constant.*

Proof: By the same argument that led to [\(19\)](#page-10-1) in the proof of Lemma [1,](#page-9-2) we get

$$
\mathbb{E}\left(\sup_{x\in\mathbb{R}}\left|\int f_x\,d(P_n-P)\right|\right)\leq\frac{3\,CK}{\sqrt{n}}=\frac{C'K}{\sqrt{n}}.
$$

Combining this with Markov's inequality, we obtain

$$
\mathbb{P}\left(\sup_{x\in\mathbb{R}}\left|\int f_x\,d(P_n-P)\right|>\delta\right)\leq\frac{C'K}{\delta\sqrt{n}}.\tag{21}
$$

This bounds the first term of [\(15\)](#page-9-1). Regarding the second term, Chebyshev's inequality gives

$$
\mathbb{P}\left(\left|1-\frac{1}{n}\sum_{i\in[n]}\frac{dQ}{dP}(Z_i)\right|>\delta\right)\leq \frac{\text{Var}\left(\frac{1}{n}\sum_{i\in[n]}\frac{dQ}{dP}(Z_i)\right)}{\delta^2}\leq \frac{K^2-1}{n\delta^2}.\tag{22}
$$

The result follows from combining [\(21\)](#page-11-0) and [\(22\)](#page-11-1).

In the following, we present yet another version by dropping the dependency on the constant C.

Lemma 3 (Alternative version) Assume $Q \ll P$ and $dQ/dP \leq B$. Then,

$$
\mathbb{P}\left(\sup_{x\in\mathbb{R}}|\hat{F}_{Q,n}(x)-F_Q(x)|>\delta\right) \le (72/\delta)e^{-n\delta^2/(4B)} + 2e^{-n\delta^2/(2B^2)}
$$

for all $\delta > 0$ *.*

Proof: To prove this lemma, we compute the upper bound for the first term of [\(15\)](#page-9-1) differently. According to [Yukich](#page-8-15) [\[1985,](#page-8-15) Theorem 2.1] and [Lafferty et al. \[2008,](#page-8-16) Theorem 7.86] we get

$$
\mathbb{P}\left(\sup_{x\in\mathbb{R}}\left|\int f_x\,d(P_n-P)\right|>\epsilon\right)\leq 4N_{[]}(\epsilon/8,\mathcal{F},L^1(P))\,e^{-n\epsilon^2/B}\tag{23}
$$

for $\epsilon \le 2/3$, where $N_{\parallel}(\epsilon/8, \mathcal{F}, L^1(P))$ denotes the bracketting number for function class $\mathcal{F} =$ ${f_x : x \in \mathbb{R}}$. Similar to the proof of Lemma [1,](#page-9-2) it can be shown that $N_{[]}(\epsilon, \mathcal{F}, L^1(P)) \leq 2 + 1/\epsilon$. Therefore,

$$
\mathbb{P}\left(\sup_{x\in\mathbb{R}}\left|\int f_x\,d(P_n-P)\right|>\epsilon\right)\leq 8(1+4/\epsilon)\,e^{-n\epsilon^2/B}\leq (36/\epsilon)e^{-n\epsilon^2/B}.
$$

Combining with [\(20\)](#page-10-2) for the second term of [\(15\)](#page-9-1), we obtain

$$
\mathbb{P}\left(\sup_{x\in\mathbb{R}}|\hat{F}_{Q,n}(x)-F_Q(x)|>\delta\right) \le (72/\delta)e^{-n\delta^2/(4B)} + 2e^{-n\delta^2/(2B^2)}
$$

$$
\le (74/\delta)\exp\left(-\frac{n\delta^2}{2(B+1)^2}\right).
$$

B Proof of Theorem [1](#page-3-1)

Recall that

$$
F_{Q,n}(x) = w_{\text{test}} \mathbf{1}\{t = \infty\} + \sum_{i \in \mathcal{I}} w_i \mathbf{1}\{s(Z_i) \le x\}, \qquad w_i = \frac{\frac{dQ}{dP}(Z_i)}{\frac{dQ}{dP}(Z_{\text{test}}) + \sum_{i \in \mathcal{I}} \frac{dQ}{dP}(Z_i)}, \tag{24}
$$

where $Z_i \stackrel{\text{i.i.d.}}{\sim} P$, $Z_{\text{test}} \sim Q$ and $|\mathcal{I}| = m$. We note

$$
P_e(\mathcal{D}_n) = \mathbb{P}\left(Y_{\text{test}} \notin \hat{C}_{\alpha}^{\text{split}}(X_{\text{test}}) \middle| \mathcal{D}_n\right) = \mathbb{P}\left(s(X_{\text{test}}, Y_{\text{test}}; \hat{\mu}) > F_{Q,m}^{-1}(1-\alpha) \middle| \mathcal{D}_n\right)
$$

$$
\leq \mathbb{P}\left(s(X_{\text{test}}, Y_{\text{test}}; \hat{\mu}) > \hat{F}_{Q,m}^{-1}(1-\alpha) \middle| \mathcal{D}_n\right)
$$

$$
= 1 - F_Q(\hat{F}_{Q,m}^{-1}(1-\alpha)),
$$

where $F_Q(x) = \mathbb{P}_{Z\sim Q}(s(Z) \leq x | \mathcal{D}_n)$ and $\hat{F}_{Q,m}^{-1}$ is defined according to [\(14\)](#page-9-3) with index set \mathcal{I} instead of $[n]$. Using the weighted DKW inequality from Lemma [1](#page-9-2) we get

$$
F_Q(\hat{F}_{Q,m}^{-1}(1-\alpha)) \ge \hat{F}_{Q,m}(\hat{F}_{Q,m}^{-1}(1-\alpha)) - \delta - C' \sqrt{\frac{B}{m}} \ge 1 - \alpha - \delta - C' \sqrt{\frac{B}{m}}, \text{ when.}
$$

Hence, we have

$$
\mathbb{P}\left(P_e(\mathcal{D}_n) > \alpha + \delta + C'\sqrt{\frac{B}{m}}\right) \le 4e^{-m\delta^2/(2B^2)}
$$

for all $\delta > 0$, or equivalently,

$$
\mathbb{P}\left(P_e(\mathcal{D}_n) > \alpha + \sqrt{\frac{2B^2}{m} \log \frac{4}{\delta}} + C'\sqrt{\frac{B}{m}}\right) = \mathbb{P}\left(P_e(\mathcal{D}_n) > \alpha + \left(\sqrt{2B\log 4/\delta} + C'\right)\sqrt{\frac{B}{m}}\right) \le \delta.
$$

This completes the proof of Theorem [1.](#page-3-1)

B.1 Proof of Remark [1](#page-4-4)

To see[\(8\)](#page-4-5), we note that by using Lemma [2](#page-11-2) instead of Lemma [1](#page-9-2) we get

$$
\mathbb{P}\left(P_e(\mathcal{D}_n) > \alpha + \delta\right) \le \frac{6\,CK}{\delta\sqrt{m}} + \frac{4(K^2 - 1)}{m\delta^2}
$$

for all $\delta > 0$. Hence,

$$
\mathbb{P}\left(P_e(\mathcal{D}_n) > \alpha + 2\left(\frac{3CK + \sqrt{\delta(K^2 - 1)}}{\delta\sqrt{m}}\right)\right) \le \delta
$$

for all $\delta > 0$, which implies

$$
\mathbb{P}\left(P_e(\mathcal{D}_n) > \alpha + \frac{2K(3C+1)}{\delta\sqrt{m}}\right) \le \delta, \qquad 0 < \delta \le 1.
$$

C Proof for Jackknife+

Lemma 4 *If Assumption [1](#page-4-1) and [2](#page-4-0) hold, then*

$$
\mathbb{P}\left(\left\|\hat{\beta}_n - \mathbb{E}\,\hat{\beta}_n\right\|_{\infty} \geq \epsilon\right) \leq 2p \exp\left(-\frac{2\kappa_1^2 \epsilon^2}{n c_n^2}\right).
$$

Proof: Assumption [1](#page-4-1) and [2](#page-4-0) imply that

$$
\sup_{z_1,\ldots,z_n,z'_i}||T(z_1,\ldots,z_i\ldots,z_n)-T(z_1,\ldots,z'_i,\ldots,z_n)||_{\infty}\leq \frac{c_n}{\kappa_1}.
$$

By McDiarmid's inequality [McDiarmid et al. \[1989](#page-8-14)] we get

$$
\mathbb{P}\left(\|\hat{\beta}_n - \mathbb{E}\,\hat{\beta}_n\|_{\infty} \geq \epsilon\right) = \mathbb{P}\left(\left\|T(Z_1, \ldots, Z_n) - \mathbb{E}\,T(Z_1, \ldots, Z_n)\right\|_{\infty} \geq \epsilon\right) \leq 2p \exp\left(-\frac{2\kappa_1^2 \epsilon^2}{nc_n^2}\right)
$$

for independent Z_i and all $\epsilon > 0$.

dependent Z_i and all $\epsilon > 0$.

Lemma 5 *Under Assumptions [1](#page-4-1) and [2](#page-4-0) we have*

$$
\mathbb{P}\left(\left.\max_{i}\left\|\mu_{\hat{\beta}_{-i}}-\mu_{\overline{\beta}_{-1}}\right\|_{\infty}\geq\epsilon\right)\leq 2p\exp\left(-\frac{2\kappa_1^2}{n}\left(\frac{\epsilon}{\kappa_2c_{n-1}}-\frac{1}{\kappa_1}\right)^2\right).
$$

Proof: From Assumption [1](#page-4-1) and [2,](#page-4-0) it follows that

$$
\max_{i,j} \|\hat{\beta}_{-i} - \hat{\beta}_{-j}\|_{\infty} \le \frac{c_{n-1}}{\kappa_1}.
$$
 (25)

Also, according to [\(4\)](#page-12-0), we have $\|\hat{\beta}_{-1} - \overline{\beta}_{-1}\|_{\infty} < \epsilon$ with probability at least 1 − $2p \exp(-2\kappa_1^2 \epsilon^2/(n c_{n-1}^2))$. We note that,

$$
\mathbb{P}\left(\max_{i} \left\|\mu_{\hat{\beta}_{-i}} - \mu_{\overline{\beta}_{-1}}\right\|_{\infty} \geq \epsilon\right) \stackrel{(*)}{\leq} \mathbb{P}\left(\kappa_{2} \max_{i} \left\|\hat{\beta}_{-i} - \overline{\beta}_{-1}\right\|_{\infty} \geq \epsilon\right)
$$

$$
\stackrel{(**)}{\leq} \mathbb{P}\left(\kappa_{2} \left(\frac{c_{n-1}}{\kappa_{1}} + \left\|\hat{\beta}_{-1} - \overline{\beta}_{-1}\right\|_{\infty}\right) \geq \epsilon\right)
$$

$$
\leq 2p \exp\left(-\frac{2\kappa_{1}^{2}}{n}\left(\frac{\epsilon}{\kappa_{2}c_{n-1}} - \frac{1}{\kappa_{1}}\right)^{2}\right).
$$

where (*) and (**) hold according to Assumption [2](#page-4-0) and [\(25\)](#page-13-0), respectively. \blacksquare

C.1 Proof of Theorem [2](#page-4-3)

We note,

$$
\hat{\mathcal{C}}_{\alpha}^{\text{J+}}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{-i}}(X_{i}) \right| \geq \left| y - \mu_{\hat{\beta}_{-i}}(X_{\text{test}}) \right| \right\} > \alpha \right\}
$$
\n
$$
\supseteq \left\{ y \in \mathbb{R} : \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\overline{\beta}_{-i}}(X_{i}) \right| - \left| \mu_{\hat{\beta}_{-i}}(X_{i}) - \mu_{\overline{\beta}_{-1}}(X_{i}) \right| \geq \right\}
$$
\n
$$
\left| y - \mu_{\overline{\beta}_{-1}}(X_{\text{test}}) \right| + \left| \mu_{\hat{\beta}_{-i}}(X_{\text{test}}) - \mu_{\overline{\beta}_{-1}}(X_{\text{test}}) \right| \right\} > \alpha \right\},
$$

where the first relation holds according to [Bian and Barber \[2023\]](#page-7-4). This first step can also be recovered by letting $w_i = 1/(n + 1)$ and $\hat{w}_i = 1/n$ in the first four steps of C.2. Assuming $\max_i \|\mu_{\hat{\beta}_{-i}} - \mu_{\overline{\beta}_{-1}}\|_{\infty} < \epsilon$, we obtain

$$
\hat{C}_{\alpha}^{\mathcal{J}+}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\overline{\beta}_{-1}}(X_{i}) \right| \geq \left| y - \mu_{\overline{\beta}_{-1}}(X_{\text{test}}) \right| + 2\epsilon \right\} > \alpha \right\}
$$

$$
\supseteq \left\{ y \in \mathbb{R} : 1 - \hat{F}^{(n-1)} \left(\left| y - \mu_{\overline{\beta}_{-1}}(X_{\text{test}}) \right| + 2\epsilon \right) > \alpha \right\}.
$$

Assuming $\left\| \hat{F}^{(n-1)} - F^{(n-1)} \right\|_{\infty} < \delta$, we obtain

$$
\hat{\mathcal{C}}_{\alpha}^{\mathcal{J}+}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : 1 - F^{(n-1)}\left(\left| y - \mu_{\overline{\beta}_{-1}}(X_{\text{test}}) \right| + 2\epsilon \right) > \alpha + \delta \right\}
$$

$$
\supseteq \left\{ y \in \mathbb{R} : 1 - F^{(n-1)}\left(\left| y - \mu_{\overline{\beta}_{-1}}(X_{\text{test}}) \right| \right) > \alpha + \delta + 2\epsilon L_{n-1} \right\}
$$

Therefore,

$$
P_e(\mathbf{D}_n) = \mathbb{P}(Y_{n+1} \notin \hat{\mathcal{C}}_{\alpha}^{J+}(X_{\text{test}})|\mathbf{D}_n) \le \mathbb{P}\left(1 - F^{(n-1)}\left(\left|Y_{n+1} - \mu_{\overline{\beta}_{-1}}(X_{\text{test}})\right|\right) \le \alpha + \delta + 2\epsilon L_{n-1}\right)
$$

$$
= \alpha + \delta + 2\epsilon L_{n-1}
$$

 $\text{for } \mathbf{D}_n \in \mathcal{A} \ \cap \ \mathcal{B} \ \ \text{where } \ \mathcal{A} \ := \ \ \left\{ D : \max_i \| \mu_{\hat{\beta}_{-i}} - \mu_{\overline{\beta}_{-1}} \|_\infty < \epsilon \right\} \ \ \text{and } \ \ \mathcal{B} \quad :=$ $\left\{ D : \left\| \hat{F}^{(n-1)} - F^{(n-1)} \right\|_{\infty} < \delta \right\}$. From Lemma [5,](#page-13-1) we know

$$
\mathbb{P}(\mathbf{D}_n \notin \mathcal{A}) \le 2p \exp\left(-\frac{2\kappa_1^2}{n}\left(\frac{\epsilon}{\kappa_2 c_{n-1}} - \frac{1}{\kappa_1}\right)^2\right).
$$

Also, according to Dvoretzky–Kiefer–Wolfowitz inequality [Dvoretzky et al. \[1956\]](#page-8-17), we have $\mathbb{P}(\mathbf{D}_n \notin \mathcal{B}) \leq 2e^{-2n\delta^2}$. Thus,

$$
\mathbb{P}(P_e(\mathbf{D}_n) > \alpha + \delta + \epsilon) \le \mathbb{P}\left((\mathcal{A} \cap \mathcal{B})^c\right) \le 2e^{-2n\delta^2} + 2p\exp\left(-\frac{2\kappa_1^2}{n}\left(\frac{\epsilon}{2L_{n-1}\kappa_2c_{n-1}} - \frac{1}{\kappa_1}\right)^2\right),
$$

or equivalently,

$$
\mathbb{P}\left(P_e(\mathbf{D}_n) > \alpha + \sqrt{\frac{\log(2/\delta)}{2n}} + 2L_{n-1}\kappa_2 c_{n-1}\left(\frac{1}{\kappa_1} + \sqrt{\frac{n}{2\kappa_1^2}\log\frac{2p}{\epsilon}}\right)\right) \leq \epsilon + \delta. \quad \blacksquare
$$

C.2 Proof of Theorem [3](#page-5-2)

We note,

$$
\hat{C}_{\alpha}^{1+}(X_{\text{test}}) = \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} w_{i} 1 \left\{ \mu_{\hat{\beta}_{-i}}(X_{\text{test}}) + \left| Y_{i} - \mu_{\hat{\beta}_{-i}}(X_{i}) \right| < y \right\} < 1 - \alpha \right\} \bigcap
$$
\n
$$
\left\{ y \in \mathbb{R} : \sum_{i=1}^{n} w_{i} 1 \left\{ \mu_{\hat{\beta}_{-i}}(X_{\text{test}}) - \left| Y_{i} - \mu_{\hat{\beta}_{-i}}(X_{i}) \right| > y \right\} < 1 - \alpha \right\}
$$
\n
$$
\supseteq \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} w_{i} 1 \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{-i}}(X_{i}) \right| < \left| y - \mu_{\hat{\beta}_{-i}}(X_{\text{test}}) \right| \right\} < 1 - \alpha \right\} \bigcap
$$
\n
$$
\left\{ y \in \mathbb{R} : \sum_{i=1}^{n} w_{i} 1 \left\{ \left| \mu_{\hat{\beta}_{-i}}(X_{\text{test}}) - y \right| > \left| Y_{i} - \mu_{\hat{\beta}_{-i}}(X_{i}) \right| \right\} < 1 - \alpha \right\}
$$
\n
$$
= \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} w_{i} 1 \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{-i}}(X_{i}) \right| < \left| y - \mu_{\hat{\beta}_{-i}}(X_{\text{test}}) \right| \right\} < 1 - \alpha \right\}
$$
\n
$$
\supseteq \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} \hat{w}_{i} 1 \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{-i}}(X_{i}) \right| < \left| y - \mu_{\hat{\beta}_{-i}}(X_{\text{test}}) \right| \right\} < 1 - \alpha \right\}
$$
\n
$$
= \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} \hat{w}_{i}
$$

where the first and last relations hold by the definition of $\hat{C}_{\alpha}^{J+}(X_{\text{test}})$ and triangle inequality, respectively. Assuming $\max_i ||\mu_{\hat{\beta}_{-i}} - \mu_{\overline{\beta}_{-1}}||_{\infty} < \epsilon$, we obtain

$$
\hat{C}_{\alpha}^{\mathbf{J}+}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} \hat{w}_i \mathbf{1} \left\{ \left| Y_i - \mu_{\overline{\beta}_{-1}}(X_i) \right| \ge \left| y - \mu_{\overline{\beta}_{-1}}(X_{\text{test}}) \right| + 2\epsilon \right\} > \alpha \right\}
$$

$$
\supseteq\left\{y\in\mathbb{R}:\sum_{i=1}^{n}\hat{w}_{i}\mathbf{1}\left\{\left|Y_{i}-\mu_{\overline{\beta}_{-1}}(X_{i})\right|\geq\left|y-\mu_{\overline{\beta}_{-1}}(X_{\text{test}})\right|+2\epsilon\right\}>\alpha\right\}
$$
\n
$$
\supseteq\left\{y\in\mathbb{R}:1-\hat{F}_{Q}^{(n-1)}\left(\left|y-\mu_{\overline{\beta}_{-1}}(X_{\text{test}})\right|+2\epsilon\right)>\alpha\right\}.
$$

where $\hat{F}_Q^{(n-1)}(t) := \sum_{i=1}^n \hat{w}_i \mathbf{1} \{ \left| Y_i - \mu_{\overline{\beta}_{-1}}(X_i) \right| \le t \}$ and, \hat{w}_i and w_i are defined in [\(14\)](#page-9-3) and [\(24\)](#page-11-3), respectively. Define,

$$
F_Q^{(n-1)}(t) := \mathbb{P}_{Z_1 \sim Q} \left(\left| Y_1 - \mu_{\overline{\beta}_{-1}}(X_1) \right| \le t \right),
$$

Assuming $\left\| \hat{F}_Q^{(n-1)} - F_Q^{(n-1)} \right\|$ $\left\| \mathcal{Q}^{(n-1)} \right\|_{\infty} < \delta$, we obtain $\sqrt{ }$

$$
\hat{C}_{\alpha}^{\mathbf{J}+}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : 1 - F_Q^{(n-1)}\left(\left| y - \mu_{\overline{\beta}_{-1}}(X_{\text{test}}) \right| + 2\epsilon \right) > \alpha + \delta \right\}
$$

$$
\supseteq \left\{ y \in \mathbb{R} : 1 - F_Q^{(n-1)}\left(\left| y - \mu_{\overline{\beta}_{-1}}(X_{\text{test}}) \right| \right) > \alpha + \delta + 2\epsilon L_{Q, n-1} \right\}
$$

Therefore,

$$
P_e(\mathbf{D}_n) = \mathbb{P}(Y_{n+1} \notin \hat{\mathcal{C}}_\alpha^{\mathrm{J}+}(X_{\mathrm{test}})|\mathbf{D}_n) \le \mathbb{P}\left(1 - F_Q^{(n-1)}\left(\left|Y_{n+1} - \mu_{\overline{\beta}_{-1}}(X_{\mathrm{test}})\right|\right) \le \alpha + \delta + 2\epsilon L_{Q,n-1}\right)
$$

$$
= \alpha + \delta + 2\epsilon L_{Q,n-1}
$$

for $\mathbf{D}_n \in \mathcal{A} \cap \mathcal{B}(\delta)$ where $\mathcal{A} := \left\{ D : \max_i ||\mu_{\hat{\beta}_{-i}} - \mu_{\overline{\beta}_{-1}}||_{\infty} < \epsilon \right\}$ and $\mathcal{B}(\delta) :=$ $\left\{D : \left\|\hat{F}_Q^{(n-1)} - F_Q^{(n-1)}\right\| \right\}$ $\left\| \mathcal{Q} \right\|_{\infty} \leq \delta$. From Lemma [5,](#page-13-1) we know

$$
\mathbb{P}(\mathbf{D}_n \notin \mathcal{A}) \le 2p \exp\left(-\frac{2\kappa_1^2}{n} \left(\frac{\epsilon}{\kappa_2 c_{n-1}} - \frac{1}{\kappa_1}\right)^2\right)
$$

Also, according to weighted DKW inequality from Lemma [1,](#page-9-2) we have

$$
\mathbb{P}\left(\mathbf{D}_n \notin \mathcal{B}\left(\delta + 3C\sqrt{\frac{B}{n}}\right)\right) \le 4e^{-n\delta^2/(2B^2)}
$$

Thus,

$$
\begin{split} \mathbb{P}\left(P_e(\mathbf{D}_n) > \alpha + \delta + \epsilon + 3\,C\sqrt{\frac{B}{n}}\right) &\leq 4e^{-n\delta^2/(2B^2)} \\ &+ 2p\exp\left(-\frac{2\kappa_1^2}{n}\left(\frac{\epsilon}{2L_{Q,n-1}\kappa_2 c_{n-1}} - \frac{1}{\kappa_1}\right)^2\right), \end{split}
$$

or equivalently,

$$
\mathbb{P}\left(P_e(\mathbf{D}_n) > \alpha + \left(\sqrt{2B\log\frac{4}{\delta}} + 3C\right)\sqrt{\frac{B}{n}} + 2L_{Q,n-1}\kappa_2 c_{n-1} \left(\frac{1}{\kappa_1} + \sqrt{\frac{n}{2\kappa_1^2}\log\frac{2p}{\epsilon}}\right)\right) \le \epsilon + \delta.
$$

D Proof for full conformal

Lemma 6 *Under Assumptions [1](#page-4-1) and [2,](#page-4-0) we have*

$$
\mathbb{P}\left(\left\|\mu_{\hat{\beta}_n}-\mu_{\overline{\beta}_n}\right\|_{\infty}\geq \epsilon\right)\leq 2p\exp\left(-\frac{2\kappa_1^2\epsilon^2}{n\kappa_2^2c_n^2}\right).
$$

Proof: According to Lemma [4,](#page-12-0) we have $\|\hat{\beta}_n - \overline{\beta}_n\|_{\infty} < \epsilon$ with probability at least 1 – $2p \exp \left(-\frac{2\kappa_1^2 \epsilon^2}{n c_n^2}\right)$ $\frac{\kappa_1^2 \epsilon^2}{n c_n^2}$). It follows from Assumption [2](#page-4-0) that,

$$
\mathbb{P}\left(\left\|\mu_{\hat{\beta}_n} - \mu_{\overline{\beta}_n}\right\|_{\infty} \geq \epsilon\right) \leq \mathbb{P}\left(\kappa_2 \left\|\hat{\beta}_n - \overline{\beta}_n\right\|_{\infty} \geq \epsilon\right) \leq 2p \exp\left(-\frac{2\kappa_1^2 \epsilon^2}{n\kappa_2^2 c_n^2}\right).
$$

We need the following notation for the proof of Theorem [4:](#page-5-1)

$$
\hat{\beta}_{X_{\text{test}},y} := T(((X_1, Y_1), \dots, (X_n, Y_n), (X_{\text{test}}, y))).
$$

D.1 Proof of Theorem [4](#page-5-1)

We note,

$$
\hat{\mathcal{C}}_{\alpha}^{\text{full}}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{i}) \right| \geq \left| y - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{\text{test}}) \right| \right\} > \alpha \right\}
$$
\n
$$
\supseteq \left\{ y \in \mathbb{R} : \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{n}}(X_{i}) \right| - \left| \mu_{\hat{\beta}_{n}}(X_{i}) - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{i}) \right| \geq \right.
$$
\n
$$
\left| y - \mu_{\hat{\beta}_{n}}(X_{\text{test}}) \right| + \left| \mu_{\hat{\beta}_{n}}(X_{\text{test}}) - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{\text{test}}) \right| \right\} > \alpha \right\}
$$
\n
$$
\supseteq \left\{ y \in \mathbb{R} : \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{n}}(X_{i}) \right| \geq \left| y - \mu_{\hat{\beta}_{n}}(X_{\text{test}}) \right| + c_{n+1} \right\} > \alpha \right\},
$$

where the first and last relations hold according to the definition of $\hat{C}_{\alpha}^{\text{full}}(X_{\text{test}})$ and Assumption [1.](#page-4-1) Assuming $\|\mu_{\hat{\beta}_n} - \mu_{\overline{\beta}_n}\|_{\infty} < \epsilon$, we obtain

$$
\hat{\mathcal{C}}_{\alpha}^{\text{full}}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : \frac{1}{n} \sum_{i=1}^{n} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\overline{\beta}_{n}}(X_{i}) \right| \ge \left| y - \mu_{\overline{\beta}_{n}}(X_{\text{test}}) \right| + c_{n+1} + 2\epsilon \right\} > \alpha \right\}
$$

$$
\supseteq \left\{ y \in \mathbb{R} : 1 - \hat{F}^{(n)} \left(\left| y - \mu_{\overline{\beta}_{n}}(X_{\text{test}}) \right| + c_{n+1} + 2\epsilon \right) > \alpha \right\}.
$$

Assuming $\left\| \hat{F}^{(n)} - F^{(n)} \right\|_{\infty} < \delta$, we obtain

$$
\hat{C}_{\alpha}^{\text{full}}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : 1 - F^{(n)} \left(\left| y - \mu_{\overline{\beta}_n}(X_{\text{test}}) \right| + c_{n+1} + 2\epsilon \right) > \alpha + \delta \right\}
$$

$$
\supseteq \left\{ y \in \mathbb{R} : 1 - F^{(n)} \left(\left| y - \mu_{\overline{\beta}_n}(X_{\text{test}}) \right| \right) > \alpha + \delta + (2\epsilon + c_{n+1}) L_n \right\}.
$$

Therefore,

$$
P_e(\mathbf{D}_n) = \mathbb{P}(Y_{n+1} \notin \hat{\mathcal{C}}_{\alpha}^{\text{full}}(X_{\text{test}})|\mathbf{D}_n)
$$

\n
$$
\leq \mathbb{P}\left(1 - F^{(n)}\left(\left|Y_{n+1} - \mu_{\overline{\beta}_n}(X_{\text{test}})\right|\right) \leq \alpha + \delta + (2\epsilon + c_{n+1})L_n\right)
$$

\n
$$
= \alpha + \delta + (2\epsilon + c_{n+1})L_n
$$

for $\mathbf{D}_n \in \mathcal{A} \cap \mathcal{B}$ where $\mathcal{A} := \left\{ D : ||\mu_{\hat{\beta}_n} - \mu_{\overline{\beta}_n}||_{\infty} < \epsilon \right\}$ and $\mathcal{B} := \left\{ D : \left\| \hat{F}^{(n)} - F^{(n)} \right\|_{\infty} < \delta \right\}$. From Lemma [5,](#page-13-1) we know $\mathbb{P}(\mathbf{D}_n \notin \mathcal{A}) \leq 2p \exp \left(-\frac{2\kappa_1^2 \epsilon^2}{n \kappa_2^2 c_n^2}\right)$ $n \kappa_2^2 c_n^2$. Also, according to Dvoretzky–Kiefer–Wolfowitz inequality, we have $\mathbb{P}(\mathbf{D}_n \notin \mathcal{B}) \leq 2e^{-2n\delta^2}$. Thus,

$$
\mathbb{P}(P_e(\mathbf{D}_n) > \alpha + \delta + \epsilon) \le \mathbb{P}\left((\mathcal{A} \cap \mathcal{B})^c\right) \le 2e^{-2n\delta^2} + 2p\exp\left(-\left(\frac{\kappa_1(\epsilon/L_n - c_{n+1})}{\sqrt{2n}\kappa_2c_n}\right)^2\right),
$$

or equivalently,

$$
\mathbb{P}\left(P_e(\mathbf{D}_n) > \alpha + \sqrt{\frac{\log(2/\delta)}{2n}} + L_n \left(c_{n+1} + \sqrt{2n\log\frac{2p}{\epsilon}} \frac{\kappa_2 c_n}{\kappa_1}\right)\right) \le \epsilon + \delta. \quad \blacksquare
$$

D.2 Proof of Theorem [5](#page-5-3)

We note,

$$
\hat{\mathcal{C}}_{\alpha}^{\text{full}}(X_{\text{test}}) = \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} w_{i} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{i}) \right| < \left| y - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{\text{test}}) \right| \right\} < 1 - \alpha \right\}
$$
\n
$$
\supseteq \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} \hat{w}_{i} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{i}) \right| < \left| y - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{\text{test}}) \right| \right\} < 1 - \alpha \right\}
$$
\n
$$
= \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} \hat{w}_{i} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{i}) \right| \ge \left| y - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{\text{test}}) \right| \right\} > \alpha \right\}
$$
\n
$$
\supseteq \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} \hat{w}_{i} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{n}}(X_{i}) \right| - \left| \mu_{\hat{\beta}_{n}}(X_{i}) - \mu_{\hat{\beta}_{X_{\text{test}},y}}(X_{i}) \right| \right\} > \alpha \right\}
$$
\n
$$
\supseteq \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} \hat{w}_{i} \mathbf{1} \left\{ \left| Y_{i} - \mu_{\hat{\beta}_{n}}(X_{i}) \right| \ge \left| y - \mu_{\hat{\beta}_{n}}(X_{\text{test}}) \right| + c_{n+1} \right\} > \alpha \right\},
$$

where the first and last relations hold according to the definition of $\hat{C}_{\alpha}^{\text{full}}(X_{\text{test}})$ (under covariate shift) and Assumption [1.](#page-4-1) Assuming $\|\mu_{\hat{\beta}_n} - \mu_{\overline{\beta}_n}\|_{\infty} < \epsilon$, we obtain

$$
\hat{\mathcal{C}}_{\alpha}^{\text{full}}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : \sum_{i=1}^{n} \hat{w}_i \mathbf{1} \left\{ \left| Y_i - \mu_{\overline{\beta}_n}(X_i) \right| \ge \left| y - \mu_{\overline{\beta}_n}(X_{\text{test}}) \right| + c_{n+1} + 2\epsilon \right\} > \alpha \right\}
$$

$$
\supseteq \left\{ y \in \mathbb{R} : 1 - \hat{F}_{Q}^{(n)} \left(\left| y - \mu_{\overline{\beta}_n}(X_{\text{test}}) \right| + c_{n+1} + 2\epsilon \right) > \alpha \right\}.
$$

Assuming $\left\| \hat{F}_Q^{(n)} - F_Q^{(n)} \right\|$ $\left\| \mathcal{Q}^{(n)} \right\|_{\infty} < \delta$, we obtain

$$
\hat{C}_{\alpha}^{\text{full}}(X_{\text{test}}) \supseteq \left\{ y \in \mathbb{R} : 1 - F_Q^{(n)} \left(\left| y - \mu_{\overline{\beta}_n}(X_{\text{test}}) \right| + c_{n+1} + 2\epsilon \right) > \alpha + \delta \right\}
$$

$$
\supseteq \left\{ y \in \mathbb{R} : 1 - F_Q^{(n)} \left(\left| y - \mu_{\overline{\beta}_n}(X_{\text{test}}) \right| \right) > \alpha + \delta + (2\epsilon + c_{n+1}) L_{Q,n} \right\}.
$$

Therefore,

$$
P_e(\mathbf{D}_n) = \mathbb{P}(Y_{n+1} \notin \hat{C}_{\alpha}^{\text{full}}(X_{\text{test}})|\mathbf{D}_n)
$$

\n
$$
\leq \mathbb{P}\left(1 - F_Q^{(n)}\left(\left|Y_{n+1} - \mu_{\overline{\beta}_n}(X_{\text{test}})\right|\right) \leq \alpha + \delta + (2\epsilon + c_{n+1})L_{Q,n}\right)
$$

\n
$$
= \alpha + \delta + (2\epsilon + c_{n+1})L_{Q,n}
$$

for $\mathbf{D}_n \in \mathcal{A} \cap \mathcal{B}(\delta)$ where $\mathcal{A} := \left\{ D : ||\mu_{\hat{\beta}_n} - \mu_{\overline{\beta}_n}||_{\infty} < \epsilon \right\}$ and $\mathcal{B}(\delta) :=$ $\left\{ D : \left\| \hat{F}_Q^{(n)} - F_Q^{(n)} \right\| \right.$ $\mathbb{E}[Q_{Q}^{(n)}] \leq \delta$. From Lemma [5,](#page-13-1) we know $\mathbb{P}(\mathbf{D}_n \notin \mathcal{A}) \leq 2p \exp \left(-\frac{2\kappa_1^2 \epsilon^2}{n\kappa_2^2 c_n^2}\right)$ $n \kappa_2^2 c_n^2$. Also, according to weighted DKW inequality from Lemma [1,](#page-9-2) we have

$$
\mathbb{P}\left(\mathbf{D}_n \notin \mathcal{B}\left(\delta + 3C\sqrt{\frac{B}{n}}\right)\right) \le 4e^{-n\delta^2/(2B^2)}
$$

Thus,

$$
\mathbb{P}\left(P_e(\mathbf{D}_n) > \alpha + \delta + 3C\sqrt{\frac{B}{n}} + \epsilon\right) \le 4e^{-n\delta^2/(2B^2)} + 2p\exp\left(-\left(\frac{\kappa_1(\epsilon/L_{Q,n} - c_{n+1})}{\sqrt{2n}\kappa_2c_n}\right)^2\right),
$$

or equivalently,

$$
\mathbb{P}\left(P_e(\mathbf{D}_n) > \alpha + \left(\sqrt{2B\log 4/\delta} + 3C\right)\sqrt{\frac{B}{n}} + L_{Q,n}\left(c_{n+1} + \sqrt{2n\log\frac{2p}{\epsilon}}\frac{\kappa_2 c_n}{\kappa_1}\right)\right) \le \epsilon + \delta.
$$