

# GENERATOR POLYNOMIALS OF CYCLIC EXPURGATED OR EXTENDED GOPPA CODES

XUE JIA, FENGWEI LI, HUAN SUN, AND QIN YUE

ABSTRACT. Classical Goppa codes are a well-known class of codes with applications in code-based cryptography, which are a special case of alternant codes. Many papers are devoted to the search for Goppa codes with a cyclic extension or with a cyclic parity-check subcode. Let  $\mathbb{F}_q$  be a finite field with  $q = 2^l$  elements, where  $l$  is a positive integer. In this paper, we determine all the generator polynomials of cyclic expurgated or extended Goppa codes under some prescribed permutations induced by the projective general linear automorphism  $A \in PGL_2(\mathbb{F}_q)$ . Moreover, we provide some examples to support our findings.

## 1. INTRODUCTION

In coding theory, cyclic codes [11, 16, 34] are an interesting type of linear codes and have wide applications in communication and storage systems due to their efficient encoding and decoding algorithms. Let  $\mathbb{F}_q$  be a finite field with  $q$  elements, where  $q = p^s$ ,  $p$  is a prime, and  $s$  is a positive integer. An  $[n, k, d]$  linear code  $\mathcal{C}$  is a  $k$ -dimensional subspace of  $\mathbb{F}_q^n$  with minimum distance  $d$ . It is called cyclic if  $(c_0, c_1, \dots, c_{n-1}) \in \mathcal{C}$  implies  $(c_{n-1}, c_0, c_1, \dots, c_{n-2}) \in \mathcal{C}$ . By identifying the codeword  $(c_0, c_1, \dots, c_{n-1}) \in \mathbb{F}_q^n$  with the polynomial

$$c_0 + c_1x + c_2x^2 + \dots + c_{n-1}x^{n-1} \in \mathbb{F}_q[x]/(x^n - 1),$$

a code  $\mathcal{C}$  of length  $n$  over  $\mathbb{F}_q$  corresponds to a subset of  $\mathbb{F}_q[x]/(x^n - 1)$ . Then  $\mathcal{C}$  is cyclic if and only if the corresponding subset is an ideal of  $\mathbb{F}_q[x]/(x^n - 1)$ . Note that every ideal of  $\mathbb{F}_q[x]/(x^n - 1)$  is principal. Hence  $\mathcal{C} = \langle u(x) \rangle$ , where  $u(x)$  is a monic divisor of  $x^n - 1$  over  $\mathbb{F}_q$ . Then  $u(x)$  is called the generator polynomial and  $h(x) = (x^n - 1)/u(x)$  is called the parity-check polynomial of the cyclic code  $\mathcal{C}$ . A cyclic code is totally determined by its generator polynomial, so it is significant to investigate the generator polynomial of a cyclic code.

---

2010 *Mathematics Subject Classification.* 94B05.

*Key words and phrases.* Cyclic codes, Expurgated Goppa codes, Extended Goppa codes, Generator polynomials.

The paper is supported by the National Natural Science Foundation of China ( No. 62172219 and No. 12171420), the Natural Science Foundation of Shandong Province under Grant ZR2021MA046, the Natural Science Foundation of Jiangsu Province under Grant BK20200268, the Innovation Program for Quantum Science and Technology under Grant 2021ZD0302902.

(\*Corresponding authours: Fengwei Li and Qin Yue).

Goppa codes [14] are subfield subcodes of generalized Reed-Solomon (GRS) codes and a special case of alternant codes [16, 21, 34]. In recent decades, Goppa codes have received extensive attention in the field of coding and cryptography. In 1978 [22], McEliece introduced the first code-based cryptosystem, in which he used a random binary irreducible Goppa code as the private key. Now the principle of the McEliece system still plays an important role in the design of code-based cryptography. In particular, the Classic McEliece originated from McEliece is one of the four candidate algorithms for the fourth round of the National Institute of Standards and Technology (NIST) post-quantum cryptography competition, and is considered to be very promising to be standardized by NIST [23]. In terms of structural cryptanalysis, there are several effective attacks on families associated with random irreducible Goppa codes [1, 8, 13, 26, 32]. Moreover, there exist some key-recovery approaches that work on unstructured Goppa codes, but whose complexity is exponential at least in the more general framework [9, 17]. It can be seen that the key using Goppa codes is sufficiently secure. Recall that there are two general constructions in coding theory that allow to define new codes from a given code, namely expurgated and extended codes, which satisfy certain linear constraints and are designed to increase code redundancy. In summary, it is very meaningful to study the structures and properties of these algebraic codes.

It is well known that some of Goppa codes have a cyclic extension (cf. [21], Research Problem 12.3). Many papers are devoted to the search for such families of Goppa codes [7], [12], [27–31]. In addition, cyclic Goppa codes, cyclic expurgated and extended Goppa codes were also investigated in [2–6] and [19]. Specifically, in [5], Berger proved that the parity-check subcodes of Goppa codes and the extended Goppa codes are both alternant codes and gave new families of Goppa codes with a cyclic extension, and some families of non-cyclic Goppa codes with a cyclic parity-check subcode. Berger presented the construction of Goppa codes invariant under some prescribed permutation [4]. Moreover, Berger constructed some other families of Goppa codes with a cyclic extension and gave a necessary and sufficient condition for a Goppa code to have a cyclic extension [3]. However, it is not yet known the generator polynomials of these cyclic codes. Based on the above results, we shall determine all the generator polynomials of cyclic expurgated or extended Goppa codes under some prescribed permutations induced by the projective general linear automorphism  $A \in PGL_2(\mathbb{F}_q)$ .

This paper is organized as follows. In section 2, we recall some definitions of Goppa codes, expurgated and extended Goppa codes. Besides, we recall the method to construct cyclic expurgated or extended Goppa codes. In section 3, we show generator polynomials of cyclic expurgated or extended Goppa codes and provide some examples to support our findings. In section 4, we conclude the paper.

## 2. PRELIMINARIES

In this paper, we always assume that  $q = 2^l$ ,  $\mathbb{F}_q$  denotes a finite field with  $2^l$  elements,  $\overline{\mathbb{F}}_q = \mathbb{F}_q \cup \{\infty\}$  is a set of coordinates for the projective line, and  $\mathbb{F}_q[x]$  is the polynomial ring over  $\mathbb{F}_q$ . We shall review some basic notations and necessary preliminaries.

## 2.1. Goppa Codes, Expurgated and Extended Goppa codes.

In this subsection, we shall recall the definitions of Goppa codes, expurgated and extended Goppa codes, which can be found in [4, 5, 16, 21, 25, 34].

**Definition 2.1.** Let  $\mathcal{L} = (\alpha_1, \dots, \alpha_n)$  be an  $n$ -tuple of distinct elements of  $\mathbb{F}_q$  and  $g(x) = \sum_{i=0}^r g_i x^i \in \mathbb{F}_q[x]$  a polynomial of degree  $r (< n)$  such that  $g(\alpha_i) \neq 0$  for  $i = 1, \dots, n$ . The Goppa code  $\Gamma(\mathcal{L}, g)$  with the support  $\mathcal{L}$  and the Goppa polynomial  $g(x)$  is defined as

$$\Gamma(\mathcal{L}, g) = \left\{ \mathbf{c} = (c_1, c_2, \dots, c_n) \in \mathbb{F}_2^n : \sum_{i=1}^n \frac{c_i}{x - \alpha_i} \equiv 0 \pmod{g(x)} \right\}.$$

The expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g)$  of  $\Gamma(\mathcal{L}, g)$  is defined as

$$\tilde{\Gamma}(\mathcal{L}, g) = \left\{ \mathbf{c} = (c_1, c_2, \dots, c_n) \in \Gamma(\mathcal{L}, g) : \sum_{i=1}^n c_i = 0 \right\}.$$

The extended Goppa code  $\overline{\Gamma}(\mathcal{L}, g)$  is defined as

$$\overline{\Gamma}(\mathcal{L}, g) = \left\{ \mathbf{c} = (c_1, \dots, c_n, c_{n+1}) \in \mathbb{F}_2^{n+1} : (c_1, c_2, \dots, c_n) \in \Gamma(\mathcal{L}, g), \sum_{i=1}^{n+1} c_i = 0 \right\}.$$

Moreover, the parity-check matrices of  $\Gamma(\mathcal{L}, g)$ ,  $\tilde{\Gamma}(\mathcal{L}, g)$ , and  $\overline{\Gamma}(\mathcal{L}, g)$  are given by

$$H = \begin{pmatrix} g(\alpha_1)^{-1} & g(\alpha_2)^{-1} & \cdots & g(\alpha_n)^{-1} \\ \alpha_1 g(\alpha_1)^{-1} & \alpha_2 g(\alpha_2)^{-1} & \cdots & \alpha_n g(\alpha_n)^{-1} \\ \vdots & \vdots & & \vdots \\ \alpha_1^{r-1} g(\alpha_1)^{-1} & \alpha_2^{r-1} g(\alpha_2)^{-1} & \cdots & \alpha_n^{r-1} g(\alpha_n)^{-1} \end{pmatrix},$$

$$\tilde{H} = \begin{pmatrix} g(\alpha_1)^{-1} & g(\alpha_2)^{-1} & \cdots & g(\alpha_n)^{-1} \\ \alpha_1 g(\alpha_1)^{-1} & \alpha_2 g(\alpha_2)^{-1} & \cdots & \alpha_n g(\alpha_n)^{-1} \\ \vdots & \vdots & & \vdots \\ \alpha_1^r g(\alpha_1)^{-1} & \alpha_2^r g(\alpha_2)^{-1} & \cdots & \alpha_n^r g(\alpha_n)^{-1} \end{pmatrix}, \quad (2.1)$$

$$\overline{H} = \begin{pmatrix} g(\alpha_1)^{-1} & g(\alpha_2)^{-1} & \cdots & g(\alpha_n)^{-1} & 0 \\ \alpha_1 g(\alpha_1)^{-1} & \alpha_2 g(\alpha_2)^{-1} & \cdots & \alpha_n g(\alpha_n)^{-1} & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ \alpha_1^{r-1} g(\alpha_1)^{-1} & \alpha_2^{r-1} g(\alpha_2)^{-1} & \cdots & \alpha_n^{r-1} g(\alpha_n)^{-1} & 0 \\ \alpha_1^r g(\alpha_1)^{-1} & \alpha_2^r g(\alpha_2)^{-1} & \cdots & \alpha_n^r g(\alpha_n)^{-1} & g(\infty)^{-1} \end{pmatrix}, \quad (2.2)$$

where  $g(\infty) = g_r$ . If  $g(x)$  is irreducible over  $\mathbb{F}_q$ , then  $\Gamma(\mathcal{L}, g)$ ,  $\tilde{\Gamma}(\mathcal{L}, g)$ , and  $\bar{\Gamma}(\mathcal{L}, g)$  are called irreducible.

**Lemma 2.2.** (*[21]*) *Let  $\Gamma(\mathcal{L}, g)$  be a binary Goppa code in Definition 2.1. Suppose that the Goppa polynomial  $g(x) \in \mathbb{F}_q[x]$  has no multiple zeros, then  $\Gamma(\mathcal{L}, g) = \Gamma(\mathcal{L}, g^2)$ .*

## 2.2. Action of Groups.

In this subsection, we shall recall the notations of some matrix groups, the actions of the projective linear group and the projective semi-linear group on  $\bar{\mathbb{F}}_q = \mathbb{F}_q \cup \{\infty\}$ .

There are some matrix groups as follows:

(1) The affine group

$$AGL_2(\mathbb{F}_q) = \left\{ A = \begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} : a \in \mathbb{F}_q^*, b \in \mathbb{F}_q \right\}.$$

(2) The general linear group

$$GL_2(\mathbb{F}_q) = \left\{ A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{F}_q, ad - bc \neq 0 \right\}.$$

(3) The projective general linear group

$$PGL_2(\mathbb{F}_q) = GL_2(\mathbb{F}_q) / \{kE_2 : k \in \mathbb{F}_q^*\},$$

where  $E_2$  is the  $2 \times 2$  identity matrix.

(4) The projective semi-linear group

$$P\Gamma L_2(\mathbb{F}_q) = PGL_2(\mathbb{F}_q) \times Gal(\mathbb{F}_q/\mathbb{F}_2),$$

where  $Gal(\mathbb{F}_q/\mathbb{F}_2) = \langle \sigma \rangle$  is the Galois group.

Let  $\bar{\mathbb{F}}_q = \mathbb{F}_q \cup \{\infty\}$  be the projective line set and  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathbb{F}_q)$ . Then the projective general linear group  $PGL_2(\mathbb{F}_q)$  acts on  $\bar{\mathbb{F}}_q$  as follows:

$$PGL_2(\mathbb{F}_q) \times \bar{\mathbb{F}}_q \rightarrow \bar{\mathbb{F}}_q$$

$$\tilde{A}(\zeta) \mapsto \tilde{A}(\zeta) = \frac{a\zeta + b}{c\zeta + d}.$$

The projective semi-linear group  $P\Gamma L_2(\mathbb{F}_q)$  acts on  $\bar{\mathbb{F}}_q$  as follows:

$$P\Gamma L_2(\mathbb{F}_q) \times \bar{\mathbb{F}}_q \rightarrow \bar{\mathbb{F}}_q$$

$$(\tilde{A}, \sigma^j)(\zeta) \mapsto \tilde{A}(\sigma^j(\zeta)) = \frac{a\zeta^{2^j} + b}{c\zeta^{2^j} + d},$$

where  $\frac{1}{0} = \infty$ ,  $\frac{1}{\infty} = 0$  and  $\frac{a\infty+b}{c\infty+d} = \frac{a}{c}$ .

Let  $M = (A, \sigma^j) \in P\Gamma L_2(\mathbb{F}_q)$  be of order  $n$  and  $G = \langle M \rangle$  a cyclic subgroup of  $P\Gamma L_2(\mathbb{F}_q)$ , then for  $\alpha \in \bar{\mathbb{F}}_q$ ,

$$O_\alpha = G(\alpha) = \{M^i(\alpha) : 0 \leq i \leq n-1\}$$

is called an orbit of  $\alpha$  under the action of  $G$ .

### 2.3. Cyclic Expurgated or Extended Goppa Codes.

In this subsection, we shall recall the method to construct cyclic expurgated or extended Goppa codes. In fact, we can choose some special Goppa polynomials and support sets to obtain cyclic expurgated or extended Goppa codes. There are many articles ([3, 5, 6]) describing such results.

**Lemma 2.3.** *Let  $M = \left( \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \sigma^j \right) \in P\Gamma L_2(\mathbb{F}_q)$  with  $c \neq 0$ ,  $G = \langle M \rangle$  a subgroup of  $P\Gamma L_2(\mathbb{F}_q)$ ,  $O_\alpha = \{\alpha_1, \dots, \alpha_n\}$  an orbit of  $\alpha \in \overline{\mathbb{F}}_q$  under the action of  $G$ , and  $g(x) = \sum_{i=0}^r g_i x^i \in \mathbb{F}_q[x]$  a monic polynomial of degree  $r (< n)$  without roots in  $O_\alpha$  satisfying*

$$c^r g\left(\frac{a}{c}\right) \neq 0 \text{ and } (cx^{2^j} + d)^r g\left(\frac{ax^{2^j} + b}{cx^{2^j} + d}\right) = c^r g\left(\frac{a}{c}\right) g(x)^{2^j}. \quad (2.3)$$

- (1) If  $\infty \notin \mathcal{L} = O_\alpha$ , then the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g)$  is cyclic of length  $n$ .  
 (2) If  $\infty \in O_\alpha$ ,  $\mathcal{L} = O_\alpha \setminus \{\infty\}$ , then the extended Goppa code  $\bar{\Gamma}(\mathcal{L}, g)$  is cyclic of length  $n$ .

*Proof.* (1) Since  $O_\alpha = \{\alpha_1, \dots, \alpha_n\}$  is an orbit of  $\alpha \in \overline{\mathbb{F}}_q$  under the action of  $G$ ,

$$\alpha_{i+1} = M\alpha_i = \frac{a\alpha_i^{2^j} + b}{c\alpha_i^{2^j} + d}, i = 1, \dots, n,$$

where  $\alpha_{n+1} = \alpha_1$ .

If  $\tilde{H}$  is the parity-check matrix of  $\tilde{\Gamma}(\mathcal{L}, g)$  in (2.1), where  $\infty \notin \mathcal{L} = O_\alpha$ , then by (2.3),

$$\begin{aligned} \tilde{H} &= \begin{pmatrix} g(M\alpha_n)^{-1} & g(M\alpha_1)^{-1} & \cdots & g(M\alpha_{n-1})^{-1} \\ M\alpha_n g(M\alpha_n)^{-1} & M\alpha_1 g(M\alpha_1)^{-1} & \cdots & M\alpha_{n-1} g(M\alpha_{n-1})^{-1} \\ \vdots & \vdots & & \vdots \\ (M\alpha_n)^r g(M\alpha_n)^{-1} & (M\alpha_1)^r g(M\alpha_1)^{-1} & \cdots & (M\alpha_{n-1})^r g(M\alpha_{n-1})^{-1} \end{pmatrix} \\ &= \frac{1}{c^r g\left(\frac{a}{c}\right)} \begin{pmatrix} \frac{(c\alpha_n^{2^j} + d)^r}{g(\alpha_n)^{2^j}} & \frac{(c\alpha_1^{2^j} + d)^r}{g(\alpha_1)^{2^j}} & \cdots & \frac{(c\alpha_{n-1}^{2^j} + d)^r}{g(\alpha_{n-1})^{2^j}} \\ \frac{(a\alpha_n^{2^j} + b)(c\alpha_n^{2^j} + d)^{r-1}}{g(\alpha_n)^{2^j}} & \frac{(a\alpha_1^{2^j} + b)(c\alpha_1^{2^j} + d)^{r-1}}{g(\alpha_1)^{2^j}} & \cdots & \frac{(a\alpha_{n-1}^{2^j} + b)(c\alpha_{n-1}^{2^j} + d)^{r-1}}{g(\alpha_{n-1})^{2^j}} \\ \vdots & \vdots & & \vdots \\ \frac{(a\alpha_n^{2^j} + b)^r}{g(\alpha_n)^{2^j}} & \frac{(a\alpha_1^{2^j} + b)^r}{g(\alpha_1)^{2^j}} & \cdots & \frac{(a\alpha_{n-1}^{2^j} + b)^r}{g(\alpha_{n-1})^{2^j}} \end{pmatrix}. \end{aligned}$$

By some elementary row transformations,  $\tilde{H}$  can be changed into

$$\tilde{H}_1 = \begin{pmatrix} \frac{1}{g(\alpha_n)^{2^j}} & \frac{1}{g(\alpha_1)^{2^j}} & \cdots & \frac{1}{g(\alpha_{n-1})^{2^j}} \\ \frac{\alpha_n^{2^j}}{g(\alpha_n)^{2^j}} & \frac{\alpha_1^{2^j}}{g(\alpha_1)^{2^j}} & \cdots & \frac{\alpha_{n-1}^{2^j}}{g(\alpha_{n-1})^{2^j}} \\ \vdots & \vdots & & \vdots \\ \frac{(\alpha_n^r)^{2^j}}{g(\alpha_n)^{2^j}} & \frac{(\alpha_1^r)^{2^j}}{g(\alpha_1)^{2^j}} & \cdots & \frac{(\alpha_{n-1}^r)^{2^j}}{g(\alpha_{n-1})^{2^j}} \end{pmatrix}.$$

Then  $\mathbf{x} = (x_1, \dots, x_n) \in \tilde{\Gamma}(\mathcal{L}, g) \subseteq \mathbb{F}_2^n$  if and only if  $\tilde{H}\mathbf{x}^T = 0$  if and only if

$$\tilde{H}_1(x_n, x_1, \dots, x_{n-1})^T = 0$$

if and only if  $(x_n, x_1, \dots, x_{n-1}) \in \tilde{\Gamma}(\mathcal{L}, g)$ . Hence  $\tilde{\Gamma}(\mathcal{L}, g)$  is cyclic.

(2) For  $\infty \in O_\alpha$ , without loss of generality, let  $\alpha_n = \infty$  and  $O_\alpha = \{\alpha_1, \dots, \alpha_{n-1}, \alpha_n = \infty\}$ . Similarly, we know that  $\bar{\Gamma}(\mathcal{L}, g)$  is also cyclic.  $\square$

**Lemma 2.4.** (*[5, Theorem 6, Proposition 7]*)

(1) Let  $M = (A, \sigma^j) \in P\Gamma L_2(\mathbb{F}_q)$ ,  $\alpha$  be an element of an extension of  $\mathbb{F}_q$ , and  $O_\alpha(x) = \prod_{i=0}^{n-1} (x - M^i(\alpha))$ , where  $n$  is the length of the orbit of  $\alpha$  under  $G = \langle M \rangle$ .

Let  $g(x) = \sum_{i=0}^r g_i x^i \in \mathbb{F}_q[x]$  be a monic polynomial of degree  $r$ . Then  $g(x)$  satisfies the condition in (2.3) if and only if  $g(x)$  is a product of polynomials  $O_{\alpha_k}(x)$  for some  $\alpha_k$ .

(2) Let  $M \in P\Gamma L_2(\mathbb{F}_q)$ . Let  $g_1(x)$  and  $g_2(x)$  be two polynomials satisfying the condition in (2.3). If  $\deg g_1(x) + \deg g_2(x) < n$ , then  $g(x) = g_1(x)g_2(x)$  satisfies the condition in (2.3).

In Lemma 2.3 and 2.4, a method for constructing cyclic expurgated or extended Goppa codes is described, but their generator polynomials are not determined. Naturally, we shall consider it.

**Question:** How are their generator polynomials given in Lemmas 2.3 and 2.4?

In detail, for  $M = \left( \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \sigma^j \right) \in P\Gamma L_2(\mathbb{F}_q)$  and  $g(x) = \sum_{i=0}^r g_i x^i \in \mathbb{F}_q[x]$  a monic polynomial of degree  $r$ , there are two questions as follows:

(1) If  $\sigma^j = 1$ , i.e.,  $M = A \in PGL_2(\mathbb{F}_q)$ , can we give generator polynomials of cyclic expurgated or extended Goppa codes?

(2) If  $\sigma^j \neq 1$ , i.e.,  $M \notin PGL_2(\mathbb{F}_q)$ , can we obtain generator polynomials of cyclic expurgated or extended Goppa codes?

In this paper, we shall determine generator polynomials of cyclic expurgated or extended Goppa codes in the case of  $\sigma^j = 1$ .

### 3. GENERATOR POLYNOMIALS OF CYCLIC EXPURGATED OR EXTENDED GOPPA CODES

In this section, we shall give all the generator polynomials of cyclic expurgated or extended Goppa codes in the case of  $\sigma^j = 1$  of Section 2.3.

First of all, we shall describe the partitions of  $\overline{\mathbb{F}}_{q^2}$  and  $\overline{\mathbb{F}}_q$ , respectively. Thus by Lemma 2.4, we can obtain all Goppa polynomials  $g(x)$  satisfying the condition in (2.3). Besides, we could select the support set  $\mathcal{L}$  of a Goppa code such that its expurgated or extended code is cyclic. The results described in Lemmas 3.2 and 3.3 of [20] are important. For completeness, we include their proofs here.

Let  $PGL_2(\mathbb{F}_q) = GL_2(\mathbb{F}_q) / \{kE_2 : k \in \mathbb{F}_q^*\}$  be projective general linear group. If  $\tilde{A} \in PGL_2(\mathbb{F}_q)$  and  $A_1 = \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix}$  is a representative element of the class  $\tilde{A}$ , then  $|A_1| = a_1d_1 + b_1c_1 = \Delta = \delta^2 \neq 0$ ,  $\delta \in \mathbb{F}_q$ . For convenience, we always choose the representative element  $A = \frac{1}{\delta}A_1$  of the class  $\tilde{A}$  with  $|A| = 1$ .

**Lemma 3.1.** (*[20, Lemma 3.2]*) Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PGL_2(\mathbb{F}_q)$  and  $|A| = ad + bc =$

1. Then

(1)  $\text{ord}(A) = 2$  if and only if  $a = d$  and  $(b, c) \neq (0, 0)$ .

(2) If  $\text{ord}(A) = n > 2$  and  $|\lambda E_2 - A| = (\lambda - \rho)(\lambda - \rho^{-1})$ , where  $\rho, \rho^{-1} \in \mathbb{F}_{q^2}$ , then  $\text{ord}(A) = \text{ord}(\rho)$  and either  $n \mid (q - 1)$  or  $n \mid (q + 1)$ .

*Proof.* (1) For  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PGL_2(\mathbb{F}_q)$ , it is clear that  $\text{ord}(A) = 2$  if and only if  $A^2 = \begin{pmatrix} a^2 + bc & b(a + d) \\ c(a + d) & d^2 + bc \end{pmatrix} = kE_2$  and  $A \neq k'E_2$ ,  $k, k' \in \mathbb{F}_q^*$ , if and only if  $a = d$  and  $(b, c) \neq (0, 0)$ .

(2) If  $\text{ord}(A) = n > 2$ , then by  $|A| = 1$ ,

$$|\lambda E_2 - A| = \begin{vmatrix} \lambda + a & b \\ c & \lambda + d \end{vmatrix} = \lambda^2 + (a + d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1}),$$

where  $\rho, \rho^{-1} \in \mathbb{F}_{q^2}^*$  are two eigenvalues of  $A$  such that  $\rho + \rho^{-1} = a + d$ . In fact, if the above characteristic polynomial is reducible over  $\mathbb{F}_q$ , then  $\text{ord}(\rho) = \text{ord}(\rho^{-1}) = n$  and  $n \mid (q - 1)$ ; if it is irreducible over  $\mathbb{F}_q$ , then  $\text{ord}(\rho) = \text{ord}(\rho^{-1}) = n$  and  $n \mid (q + 1)$ .

By (1) and  $a \neq d$ ,  $\rho \neq \rho^{-1}$ . Then there is an invertible matrix  $P \in PGL_2(\mathbb{F}_{q^2})$  such that  $P^{-1}AP = \begin{pmatrix} \rho & 0 \\ 0 & \rho^{-1} \end{pmatrix}$ . It has that  $A^k = P \begin{pmatrix} \rho^k & 0 \\ 0 & \rho^{-k} \end{pmatrix} P^{-1} = cE_2$ ,  $c \in \mathbb{F}_q^*$ , if and only if  $\rho^k = \rho^{-k}$  if and only if  $\rho^{2k} = 1$ . Hence  $\text{ord}(A) = \text{ord}(\rho^2) = \text{ord}(\rho) = n$ .  $\square$

In the following, we always assume that  $c \neq 0$  and  $\text{ord}(A) = n > 2$ , which implies  $\rho \neq \rho^{-1}$ . It is worth mentioning that we just consider the case  $c \neq 0$  in the case of  $\sigma^j = 1$  of Section 2.3 because it is very simple if  $c = 0$ . So we omit it here.

**Lemma 3.2.** (*[20, Lemma 3.3]*) Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PGL_2(\mathbb{F}_q)$  be of order  $n > 2$  with  $|A| = ad + bc = 1$  and  $c \neq 0$ . Let  $\alpha$  be an element of an extension of  $\mathbb{F}_q$ . Then

$A(\alpha) = \alpha$  if and only if  $\alpha \in \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\}$ ; otherwise,  $n$  is the least positive integer such that  $A^n(\alpha) = \alpha$ , where  $\rho, \rho^{-1} \in \mathbb{F}_{q^2}^*$  are two eigenvalues of  $A$ .

*Proof.* By Lemma 3.1, there is an invertible matrix  $P$  such that  $A = P \begin{pmatrix} \rho & 0 \\ 0 & \rho^{-1} \end{pmatrix} P^{-1}$

and  $A^i(\alpha) = P \begin{pmatrix} \rho^i & 0 \\ 0 & \rho^{-i} \end{pmatrix} P^{-1}(\alpha)$ ,  $1 \leq i \leq n-1$ , where

$$P = \begin{pmatrix} a + \rho^{-1} & a + \rho \\ c & c \end{pmatrix}. \quad (3.1)$$

In fact, since  $\rho, \rho^{-1} \in \mathbb{F}_{q^2}^*$  are two eigenvalues of  $A$ ,

$$\begin{aligned} P^{-1}AP &= \begin{pmatrix} c & a + \rho \\ c & a + \rho^{-1} \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} a + \rho^{-1} & a + \rho \\ c & c \end{pmatrix} \\ &= \begin{pmatrix} c\rho & d\rho + 1 \\ c\rho^{-1} & d\rho^{-1} + 1 \end{pmatrix} \begin{pmatrix} a + \rho^{-1} & a + \rho \\ c & c \end{pmatrix} \\ &= \begin{pmatrix} c(a+d)\rho & c(\rho^2 + (a+d)\rho + 1) \\ c(\rho^{-2} + (a+d)\rho^{-1} + 1) & c(a+d)\rho^{-1} \end{pmatrix} \\ &= \begin{pmatrix} c(a+d)\rho & 0 \\ 0 & c(a+d)\rho^{-1} \end{pmatrix} = \begin{pmatrix} \rho & 0 \\ 0 & \rho^{-1} \end{pmatrix}. \end{aligned}$$

For  $1 \leq i \leq n-1$ ,  $A^i(\alpha) = \alpha$  if and only if  $\rho^{2i}P^{-1}(\alpha) = P^{-1}(\alpha)$ . By  $\text{ord}(\rho) = n$  odd,  $A^i(\alpha) = \alpha$  if and only if  $P^{-1}(\alpha) \in \{0, \infty\}$  if and only if  $\alpha \in \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\}$ .

Therefore,  $A(\alpha) = \alpha$  if and only if  $\alpha \in \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\}$ ; otherwise,  $n$  is the least positive integer such that  $A^n(\alpha) = \alpha$ .  $\square$

**Corollary 3.3.** *Let  $A$  be a matrix as in Lemma 3.2 and  $G = \langle A \rangle$  a subgroup of  $PGL_2(\mathbb{F}_q)$ . For each  $\alpha \in \overline{\mathbb{F}}_{q^2}$ , let  $O_\alpha = \{A^i(\alpha) : A^i \in G\}$  be the orbit of  $\alpha$  in  $\overline{\mathbb{F}}_{q^2}$  under the action of  $G$ . Then there is a partition of  $\overline{\mathbb{F}}_{q^2}$  as follows:*

$$\overline{\mathbb{F}}_{q^2} = \left\{ \frac{a+\rho}{c} \right\} \cup \left\{ \frac{a+\rho^{-1}}{c} \right\} \cup \bigcup_{i=1}^{\frac{q^2-1}{n}} O_{\alpha_i},$$

where  $\alpha_i \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\}$  and  $|O_{\alpha_i}| = \text{ord}(A) = n$ . Moreover,

$$O_\infty = \{P(\rho), P(\rho^2), \dots, P(\rho^{n-1}), P(1) = \infty\},$$

where  $P$  is defined as (3.1).

*Proof.* Let  $G = \langle A \rangle$  be a cyclic subgroup of  $PGL_2(\mathbb{F}_q)$  and  $\overline{\mathbb{F}}_{q^2}$  be the projective line set, then there is an action of  $G$  on  $\overline{\mathbb{F}}_{q^2}$ :

$$\begin{aligned} G \times \overline{\mathbb{F}}_{q^2} &\rightarrow \overline{\mathbb{F}}_{q^2}, \\ (A^i, \alpha) &\mapsto A^i(\alpha). \end{aligned}$$



Then  $\overline{\mathbb{F}}_{q^2}$  has a partition (in details, see [24, Proposition 272]). By Lemma 3.2,  $\frac{a+\rho}{c}$  and  $\frac{a+\rho^{-1}}{c}$  are the only two fixed points under the action of group  $G = \langle A \rangle$ ; since  $n$  is the least positive integer such that  $A^n(\alpha) = \alpha$  if  $\alpha \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\}$ ,  $O_\alpha = G(\alpha) = \{\alpha, A(\alpha), \dots, A^{n-1}(\alpha)\}$ . Hence there is a partition of  $\overline{\mathbb{F}}_{q^2}$  as follows:

$$\overline{\mathbb{F}}_{q^2} = \left\{ \frac{a+\rho}{c} \right\} \cup \left\{ \frac{a+\rho^{-1}}{c} \right\} \cup \bigcup_{i=1}^{\frac{q^2-1}{n}} O_{\alpha_i},$$

where  $\alpha_i \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\}$  and  $|O_{\alpha_i}| = \text{ord}(A) = n$ .

Let  $\Lambda = \begin{pmatrix} \rho & 0 \\ 0 & \rho^{-1} \end{pmatrix}$  and  $G' = \langle \Lambda \rangle$  a subgroup of  $PGL_2(\overline{\mathbb{F}}_{q^2})$ , then  $G'$  acts on the set  $\overline{\mathbb{F}}_{q^2}$ , so for  $\beta \notin \{0, \infty\}$ , there is an orbit of  $\beta$  under the action of  $G'$ :

$$O_\beta = \{\beta, \rho\beta, \dots, \rho^{n-1}\beta\}.$$

By  $A = P\Lambda P^{-1}$ ,  $P(0) = \frac{a+\rho}{c}$ ,  $P(\infty) = \frac{a+\rho^{-1}}{c}$ . Hence for  $\beta \notin \{0, \infty\}$ , there is an orbit of  $P(\beta)$  under the action of  $G = \langle A \rangle$ :

$$O_{P(\beta)} = \{P(\beta), P(\rho\beta), \dots, P(\rho^{n-1}\beta)\},$$

so

$$O_\infty = \{P(\rho), \dots, P(\rho^{n-1}), P(1) = \infty\}.$$

□

**Corollary 3.4.** *Let  $A$  be a matrix as in Lemma 3.2 and  $G = \langle A \rangle$  a subgroup of  $PGL_2(\mathbb{F}_q)$ . For each  $\alpha \in \overline{\mathbb{F}}_q$ , let  $O_\alpha = \{A^i(\alpha) : A^i \in G\}$  be the orbit of  $\alpha$  in  $\overline{\mathbb{F}}_q$  under the action of  $G$ .*

(1) *If  $\lambda^2 + (a+d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  is reducible over  $\mathbb{F}_q$ , then there is a partition of  $\overline{\mathbb{F}}_q$  as follows:*

$$\overline{\mathbb{F}}_q = \left\{ \frac{a+\rho}{c} \right\} \cup \left\{ \frac{a+\rho^{-1}}{c} \right\} \cup \bigcup_{i=1}^{\frac{q-1}{n}} O_{\alpha_i},$$

where  $\alpha_i \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\}$  and  $|O_{\alpha_i}| = \text{ord}(A) = n$ .

(2) *If  $\lambda^2 + (a+d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  is irreducible over  $\mathbb{F}_q$ , then there is a partition of  $\overline{\mathbb{F}}_q$  as follows:*

$$\overline{\mathbb{F}}_q = \bigcup_{i=1}^{\frac{q+1}{n}} O_{\alpha_i},$$

where  $|O_{\alpha_i}| = \text{ord}(A) = n$ .

**Proposition 3.5.** *Let  $A$  be a matrix as in Lemma 3.2,  $\lambda^2 + (a+d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$ ,  $\rho, \rho^{-1} \in \mathbb{F}_{q^2}$ ,  $G = \langle A \rangle$  a subgroup of  $PGL_2(\mathbb{F}_q)$ , and  $g(x) = \sum_{i=0}^r g_i x^i \in \mathbb{F}_{q^2}[x]$  a*

monic polynomial satisfying the condition in (2.3). Then both the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g)$  are cyclic of length  $n$ , where

$$\mathcal{L}' = O_\infty \setminus \{\infty\} = \{P(\rho), \dots, P(\rho^{n-1})\} = \{\gamma, A(\gamma), \dots, A^{n-2}(\gamma)\} \subseteq \mathbb{F}_{q^2},$$

and

$$\mathcal{L} = O_\alpha = \{\alpha_1 = \alpha, \alpha_2 = A(\alpha), \dots, \alpha_n = A^{n-1}(\alpha)\} \subseteq \mathbb{F}_{q^2}$$

is the orbit of  $\alpha$  under the action of  $G$  if  $\alpha \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\} \cup O_\infty$ .

(1) If the degree  $r$  of  $g(x)$  is not less than  $n-1$ , i.e.,  $r \geq n-1$ , then both the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g)$  are zero.

(2) If the degree  $r$  of  $g(x)$  is less than  $n-1$ , then

$$g(x) = g_1(x)^s g_2(x)^t, \quad (3.2)$$

where

$$g_1(x) = x + \frac{a+\rho}{c}, \quad g_2(x) = x + \frac{a+\rho^{-1}}{c}, \quad (3.3)$$

and  $s+t < n-1$ .

*Proof.* (1) If  $r \geq n-1$  and  $\tilde{H}$  is defined as (2.1), then  $\text{rank}_{\mathbb{F}_{q^2}}(\tilde{H}) = n$  and

$$\begin{aligned} \tilde{\Gamma}(\mathcal{L}, g) &= \left\{ \mathbf{c} = (c_1, \dots, c_n) \in \mathbb{F}_2^n : \tilde{H}\mathbf{c}^T = 0 \right\} \\ &\subseteq \left\{ \mathbf{c} = (c_1, \dots, c_n) \in \mathbb{F}_{q^2}^n : \tilde{H}\mathbf{c}^T = 0 \right\} = \{\mathbf{0}\}, \end{aligned}$$

Hence the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g)$  is zero. Similarly, we can prove that the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g)$  is also zero.

(2) If the degree  $r$  of  $g(x)$  is less than  $n-1$  and  $g(x)$  satisfies the condition in (2.3), then by Lemma 2.4, there is not any root of  $g(x)$  in an orbit  $O_{\alpha_i}$  with  $|O_{\alpha_i}| = n$  in Corollary 3.3. Hence  $g(x) = g_1(x)^s g_2(x)^t \in \mathbb{F}_{q^2}[x]$ , where  $g_1(x) = x + \frac{a+\rho}{c}$ ,  $g_2(x) = x + \frac{a+\rho^{-1}}{c}$ ,  $s, t \in \mathbb{N}$ , and  $s+t < n-1$ .  $\square$

**Remark 3.6.** In Proposition 3.5, if  $g(x) = \sum_{i=0}^r g_i x^i \in \mathbb{F}_q[x]$  is a monic polynomial satisfying the condition in (2.3), then

$$g(x) = \begin{cases} g_1(x)^s g_2(x)^t, & \text{if } \lambda^2 + (a+d)\lambda + 1 \text{ is reducible over } \mathbb{F}_q, \\ (g_1(x)g_2(x))^s, & \text{if } \lambda^2 + (a+d)\lambda + 1 \text{ is irreducible over } \mathbb{F}_q, \end{cases} \quad (3.4)$$

where  $s, t \in \mathbb{N}$ ,  $s+t < n-1$  if  $\lambda^2 + (a+d)\lambda + 1$  is reducible over  $\mathbb{F}_q$ , and  $2s < n-1$  if  $\lambda^2 + (a+d)\lambda + 1$  is irreducible over  $\mathbb{F}_q$ .

Indeed, if  $\lambda^2 + (a+d)\lambda + 1$  is reducible over  $\mathbb{F}_q$ , then  $g_1(x) = x + \frac{a+\rho}{c}$ ,  $g_2(x) = x + \frac{a+\rho^{-1}}{c} \in \mathbb{F}_q[x]$ , and  $g(x) = g_1(x)^s g_2(x)^t \in \mathbb{F}_q[x]$ . If  $\lambda^2 + (a+d)\lambda + 1$  is irreducible over  $\mathbb{F}_q$ , then  $\text{ord}(\rho) = \text{ord}(A) = n \mid (q+1)$ ,  $(\frac{a+\rho}{c})^q = \frac{a+\rho^{-1}}{c}$ ,  $g_1(x)^s g_2(x)^t = g(x) = (g(x))^q = g_2(x)^s g_1(x)^t$  by  $g(x) \in \mathbb{F}_q[x]$ , thus  $s = t$ .

In the following, we always assume that the degree  $r$  of  $g(x)$  is less than  $n - 1$ . We shall find generator polynomials of cyclic expurgated or extended Goppa codes with Goppa polynomial  $g(x) = g_1(x)^s g_2(x)^t$ ,  $s, t \in \mathbb{N}$  and  $s + t < n - 1$ . Specifically, we shall consider the Goppa polynomial  $g(x)$  in three cases:

- (1)  $g(x) = g_1(x)$  or  $g_2(x)$ ;
- (2)  $g(x) = g_1(x)^s$  or  $g_2(x)^t$ ,  $s, t \in \mathbb{N}$  and  $1 \leq s, t < n - 1$ ;
- (3)  $g(x) = g_1(x)^s g_2(x)^t$ ,  $s, t \in \mathbb{N}$  and  $s + t < n - 1$ ;

where  $g_1(x)$  and  $g_2(x)$  are just defined as (3.3).

### 3.1. $g(x) = g_1(x)$ or $g_2(x)$ .

In the subsection, we shall assume that  $g(x) = g_1(x)$  or  $g_2(x)$  and find generator polynomials of cyclic expurgated or extended Goppa codes in the case of  $\sigma^j = 1$  of Section 2.3.

**Theorem 3.7.** *Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PGL_2(\mathbb{F}_q)$  be of order  $n > 2$  with  $|A| = ad + bc = 1$  and  $c \neq 0$ ,  $G = \langle A \rangle$  a subgroup of  $PGL_2(\mathbb{F}_q)$ , and  $|\lambda E_2 - A| = \lambda^2 + (a + d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  reducible over  $\mathbb{F}_q$ . Let*

$$\mathcal{L}' = O_\infty \setminus \{\infty\} = \{P(\rho), \dots, P(\rho^{n-1})\} = \{\gamma, A(\gamma), \dots, A^{n-2}(\gamma)\} \subseteq \mathbb{F}_q$$

and

$$\mathcal{L} = O_\alpha = \{\alpha_1 = \alpha, \alpha_2 = A(\alpha), \dots, \alpha_n = A^{n-1}(\alpha)\} \subseteq \mathbb{F}_q$$

be the orbit of  $\alpha$  under the action of  $G$  if  $\alpha \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\} \cup O_\infty$ .

(1) If  $g_1(x) = x + \frac{a+\rho}{c} \in \mathbb{F}_q[x]$ , then both the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g_1)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g_1)$  are cyclic of length  $n$  and their generator polynomials are  $u_1(x) = (x+1)m_{\rho^{-1}}(x)$ , where  $m_{\rho^{-1}}(x)$  is the minimal polynomial of  $\rho^{-1}$  over  $\mathbb{F}_2$ .

(2) If  $g_2(x) = x + \frac{a+\rho^{-1}}{c} \in \mathbb{F}_q[x]$ , then both the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g_2)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g_2)$  are cyclic of length  $n$  and their generator polynomials are  $u_2(x) = (x+1)m_\rho(x)$ , where  $m_\rho(x)$  is the minimal polynomial of  $\rho$  over  $\mathbb{F}_2$ .

*Proof.* Let  $\lambda^2 + (a+d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  be reducible over  $\mathbb{F}_q$ , then by Proposition 3.5,

$$g_1(x) = x + \frac{a + \rho}{c}, \quad g_2(x) = x + \frac{a + \rho^{-1}}{c} \in \mathbb{F}_q[x]$$

satisfy the condition in (2.3). Hence by Lemma 2.3, the expurgated Goppa codes  $\tilde{\Gamma}(\mathcal{L}, g_1)$ ,  $\tilde{\Gamma}(\mathcal{L}, g_2)$  and the extended Goppa codes  $\bar{\Gamma}(\mathcal{L}', g_1)$ ,  $\bar{\Gamma}(\mathcal{L}', g_2)$  are all cyclic of length  $n$ . In the following, we only need to find the generator polynomial of  $\tilde{\Gamma}(\mathcal{L}, g_1)$ . Similarly, we can also obtain those of  $\bar{\Gamma}(\mathcal{L}', g_1)$ ,  $\tilde{\Gamma}(\mathcal{L}, g_2)$  and  $\bar{\Gamma}(\mathcal{L}', g_2)$ .

For the cyclic expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g_1)$ , we shall prove that the polynomial  $(x+1)m_{\rho^{-1}}(x)$  is its generator polynomial, where  $m_{\rho^{-1}}(x)$  is the minimal polynomial

of  $\rho^{-1}$  over  $\mathbb{F}_2$ . Now we consider the parity-check matrix of  $\tilde{\Gamma}(\mathcal{L}, g_1)$  as follows:

$$\tilde{H} = \begin{pmatrix} \frac{1}{\alpha_1 + \frac{a+\rho}{c}} & \cdots & \frac{1}{\alpha_n + \frac{a+\rho}{c}} \\ 1 & \cdots & 1 \end{pmatrix}.$$

Hence  $\mathbf{c} = (c_1, \dots, c_n) \in \tilde{\Gamma}(\mathcal{L}, g_1)$  if and only if  $\tilde{H}\mathbf{c}^T = 0$  if and only if  $\mathbf{c}$  is a solution of the system of equations

$$\begin{cases} \frac{c_1}{\alpha_1 + \frac{a+\rho}{c}} + \cdots + \frac{c_n}{\alpha_n + \frac{a+\rho}{c}} = 0, \\ c_1 + c_2 + \cdots + c_n = 0. \end{cases} \quad (3.5)$$

In the following, we shall prove that  $\mathbf{c} = (c_1, \dots, c_n) \in \tilde{\Gamma}(\mathcal{L}, g_1)$  if and only if  $\mathbf{c}$  is a solution of the system of equations

$$\begin{cases} c_1 + c_2e + \cdots + c_n e^{n-1} = 0, \\ c_1 + c_2 + \cdots + c_n = 0, \end{cases} \quad (3.6)$$

where  $e = \rho^{-2}$ . It means that the system in (3.5) is equivalent to the system in (3.6).

In (3.5),  $\frac{1}{\alpha_i + \frac{a+\rho}{c}} = B(\alpha_i)$ ,  $1 \leq i \leq n$ , where  $B = \begin{pmatrix} 0 & 1 \\ 1 & \frac{a+\rho}{c} \end{pmatrix}$ . On the other hand,

$$D = BAB^{-1} = \begin{pmatrix} a+d+\rho & c \\ \frac{\rho^2+(a+d)\rho+1}{c} & \rho \end{pmatrix} = \begin{pmatrix} \rho^{-2} & c\rho^{-1} \\ 0 & 1 \end{pmatrix} \triangleq \begin{pmatrix} e & f \\ 0 & 1 \end{pmatrix},$$

where  $e = \rho^{-2}$  and  $f = c\rho^{-1}$ . Then  $\text{ord}(D) = \text{ord}(A) = \text{ord}(e) = n$  and

$$D^i = BA^iB^{-1} = \begin{pmatrix} e^i & f(e^{i-1} + e^{i-2} + \cdots + 1) \\ 0 & 1 \end{pmatrix}, 1 \leq i \leq n-1.$$

By  $\alpha_{i+1} = A(\alpha_i)$ ,  $1 \leq i \leq n-1$ , the system (3.5) is equivalent to the system

$$\begin{cases} c_1B(\alpha_1) + c_2B(A(\alpha_1)) + \cdots + c_nB(A^{n-1}(\alpha_1)) = 0, \\ c_1 + c_2 + \cdots + c_n = 0. \end{cases}$$

Note that  $B(A^i(\alpha_1)) = BA^iB^{-1}B(\alpha_1) = D^iB(\alpha_1)$ ,  $1 \leq i \leq n-1$ . Hence the above system is equivalent to the system

$$\begin{cases} c_1B(\alpha_1) + c_2D(B(\alpha_1)) + \cdots + c_nD^{n-1}(B(\alpha_1)) = 0, \\ c_1 + c_2 + \cdots + c_n = 0. \end{cases}$$

Now we consider the first equation in the above system.

$$\begin{aligned} & c_1B(\alpha_1) + c_2D(B(\alpha_1)) + \cdots + c_nD^{n-1}(B(\alpha_1)) \\ &= (c_1 + c_2e + c_3e^2 + \cdots + c_n e^{n-1})B(\alpha_1) \\ &+ \frac{f}{e-1}(c_2(e-1) + c_3(e^2-1) + \cdots + c_n(e^{n-1}-1)) \\ &= (c_1 + c_2e + c_3e^2 + \cdots + c_n e^{n-1})(B(\alpha_1) + \frac{f}{e-1}) = 0. \end{aligned}$$

Suppose that  $B(\alpha_1) + \frac{f}{e-1} = 0$ , i.e.,  $\frac{1}{\alpha_1 + \frac{a+\rho}{c}} = \frac{c\rho^{-1}}{\rho^{-2}-1}$ . Then  $\alpha_1 = \frac{a+\rho^{-1}}{c}$ , which is contradictory with the choice of  $\alpha_1$ . Therefore, the system in (3.5) is equivalent to the system in (3.6).

Moreover, let  $m_{\rho^{-1}}(x)$  be the minimal polynomial of  $\rho^{-1}$  over  $\mathbb{F}_2$ , then it is also the minimal polynomial of  $e = \rho^{-2}$  over  $\mathbb{F}_2$ . Since  $\tilde{\Gamma}(\mathcal{L}, g_1)$  is a cyclic code of length  $n$ , whose elements satisfy the system (3.6), its generator polynomial is  $u_1(x) = (x-1)m_{\rho^{-1}}(x)$ .

Similarly, we can get generator polynomial of  $\bar{\Gamma}(\mathcal{L}', g_1)$  is  $u_1(x) = (x-1)m_{\rho^{-1}}(x)$  and generator polynomials of both  $\tilde{\Gamma}(\mathcal{L}, g_2)$  and  $\bar{\Gamma}(\mathcal{L}', g_2)$  are  $u_2(x) = (x-1)m_{\rho}(x)$ , where  $m_{\rho}(x)$  is the minimal polynomial of  $\rho$  over  $\mathbb{F}_2$ .  $\square$

**Remark 3.8.** In [33],  $m_{\rho}(x) = m_{\rho^{-1}}(x)$  is equivalent to the following conditions:

- (1) The polynomial  $m_{\rho}(x)$  is self-reciprocal.
- (2) There exists a positive integer  $\omega$  such that  $2^{\omega} \equiv -1 \pmod{n}$ .
- (3) Suppose  $n = p_1^{l_1} p_2^{l_2} \cdots p_t^{l_t}$  is the prime factorization, where  $p_1, \dots, p_t$  are distinct odd primes and  $l_1, \dots, l_t$  are positive integers. By this time,  $v_2(d_1) = \cdots = v_2(d_t) = \delta$ , where  $d_i$  is the order of 2 modulo  $p_i$ ,  $v_2(d_i)$  denotes the 2-adic valuation of  $d_i$ ,  $1 \leq i \leq t$ , and  $\delta$  is a positive integer.

**Remark 3.9.** In Theorem 3.7, if  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PGL_2(\mathbb{F}_q)$  and  $\lambda^2 + (a+d)\lambda + 1 = (\lambda-\rho)(\lambda-\rho^{-1})$  is reducible over  $\mathbb{F}_q$ , then the polynomial  $g(x)$  of degree 1 that satisfies the condition in (2.3) can only be  $g_1(x) = x + \frac{a+\rho}{c}$  or  $g_2(x) = x + \frac{a+\rho^{-1}}{c}$ . Hence all possible cyclic expurgated Goppa codes with Goppa polynomial of degree 1 are  $\tilde{\Gamma}(\mathcal{L}, g_1)$  and  $\tilde{\Gamma}(\mathcal{L}, g_2)$ ; all possible cyclic extended Goppa codes with Goppa polynomial of degree 1 are  $\bar{\Gamma}(\mathcal{L}', g_1)$  and  $\bar{\Gamma}(\mathcal{L}', g_2)$ .

**Corollary 3.10.** Let  $A \in PGL_2(\mathbb{F}_q)$  be the matrix,  $G = \langle A \rangle$  the subgroup in Theorem 3.7, and  $|\lambda E_2 - A| = \lambda^2 + (a+d)\lambda + 1 = (\lambda-\rho)(\lambda-\rho^{-1})$  irreducible over  $\mathbb{F}_q$ . Let

$$\mathcal{L}' = O_{\infty} \setminus \{\infty\} = \{P(\rho), \dots, P(\rho^{n-1})\} = \{\gamma, A(\gamma), \dots, A^{n-2}(\gamma)\} \subseteq \mathbb{F}_{q^2}$$

and

$$\mathcal{L} = O_{\alpha} = \{\alpha_1 = \alpha, \alpha_2 = A(\alpha), \dots, \alpha_n = A^{n-1}(\alpha)\} \subseteq \mathbb{F}_{q^2}$$

be the orbit of  $\alpha$  under the action of  $G$  if  $\alpha \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\} \cup O_{\infty}$ .

Then the conclusions of Theorem 3.7 still hold but with the Goppa polynomials  $g_1(x), g_2(x) \in \mathbb{F}_{q^2}[x]$ . Moreover, let  $\lambda^2 + (a+d)\lambda + 1$  be irreducible over  $\mathbb{F}_q$ ,  $\text{ord}(A) = n \mid (q+1)$ , then by Remark 3.8,  $m_{\rho}(x) = m_{\rho^{-1}}(x)$ .

By Lemma 2.2, we can get the following result.

**Corollary 3.11.** In Theorem 3.7,  $\tilde{\Gamma}(\mathcal{L}, g_i^2) = \tilde{\Gamma}(\mathcal{L}, g_i) = \langle h_i(x) \rangle = \bar{\Gamma}(\mathcal{L}', g_i^2) = \bar{\Gamma}(\mathcal{L}', g_i)$ ,  $i = 1, 2$ , are all cyclic codes of length  $n$ .

**Example 3.12.** Let  $\mathbb{F}_{26}^* = \langle \alpha \rangle$  and  $A = \begin{pmatrix} \alpha^5 & \alpha^{43} \\ \alpha^{13} & \alpha^{59} \end{pmatrix} \in PGL_2(\mathbb{F}_{26})$ , then  $|A| = 1$ ,  $\text{ord}(A) = 21 > 2$ , and  $\lambda^2 + (\alpha^5 + \alpha^{59})\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1}) \in \mathbb{F}_{26}[x]$ , where  $\rho = \alpha^3, \rho^{-1} = \alpha^{-3} = \alpha^{60} \in \mathbb{F}_{26}^*$  are two eigenvalues of  $A$ . The minimal polynomials of  $\rho^{-1}$  and  $\rho$  over  $\mathbb{F}_2$  are  $m_{\rho^{-1}}(x) = x^6 + x^4 + x^2 + x + 1$  and  $m_\rho(x) = x^6 + x^5 + x^4 + x^2 + 1$ , respectively. Let

$$\mathcal{L}' = O_\infty \setminus \{\infty\} = \{\alpha^{55}, \alpha^{59}, \alpha^{23}, \alpha^{15}, \alpha^7, \alpha^{34}, \alpha^{38}, 0, \alpha^{47}, \alpha^{18}, \alpha^{17}, \alpha^{53}, \alpha^{10}, \alpha^{42}, \alpha^{51}, \alpha^{20}, \alpha^{40}, \alpha^{13}, \alpha^{12}, \alpha^{46}\} \subseteq \mathbb{F}_{26}$$

and

$$\mathcal{L} = O_\alpha = \{\alpha, \alpha^{62}, \alpha^{11}, \alpha^{60}, \alpha^{14}, \alpha^5, \alpha^6, \alpha^{32}, \alpha^2, \alpha^8, \alpha^{27}, \alpha^{30}, \alpha^{48}, \alpha^{58}, \alpha^{52}, \alpha^{37}, \alpha^{21}, \alpha^{36}, \alpha^{44}, \alpha^{26}, \alpha^{50}\} \subseteq \mathbb{F}_{26},$$

where  $|\mathcal{L}'| = 20$  and  $|\mathcal{L}| = 21 = \text{ord}(A)$ .

(1) If  $g_1(x) = x + \frac{\alpha^5 + \alpha^3}{\alpha^{13}}$  and  $g_1(x)^2 = (x + \frac{\alpha^5 + \alpha^3}{\alpha^{13}})^2$ , we can obtain that the expurgated Goppa codes  $\tilde{\Gamma}(\mathcal{L}, g_1)$  and  $\tilde{\Gamma}(\mathcal{L}, g_1^2)$ , the extended Goppa codes  $\bar{\Gamma}(\mathcal{L}', g_1)$  and  $\bar{\Gamma}(\mathcal{L}', g_1^2)$  are all  $[21, 14, 4]$  cyclic codes and their generator polynomials are  $(x+1)(x^6 + x^4 + x^2 + x + 1) = (x+1)m_{\rho^{-1}}(x)$  by the Magma program.

(2) If  $g_2(x) = x + \frac{\alpha^5 + \alpha^{-3}}{\alpha^{13}}$  and  $g_2(x)^2 = (x + \frac{\alpha^5 + \alpha^{-3}}{\alpha^{13}})^2$ , we can obtain that the expurgated Goppa codes  $\tilde{\Gamma}(\mathcal{L}, g_2)$  and  $\tilde{\Gamma}(\mathcal{L}, g_2^2)$ , the extended Goppa codes  $\bar{\Gamma}(\mathcal{L}', g_2)$  and  $\bar{\Gamma}(\mathcal{L}', g_2^2)$  are all  $[21, 14, 4]$  cyclic codes and their generator polynomials are  $(x+1)(x^6 + x^5 + x^4 + x^2 + 1) = (x+1)m_\rho(x)$  by the Magma program.

**Example 3.13.** Let  $\mathbb{F}_{26}^* = \langle \alpha \rangle$  and  $A = \begin{pmatrix} \alpha^7 & 0 \\ 1 & \alpha^{-7} \end{pmatrix} \in PGL_2(\mathbb{F}_{26})$ , then  $|A| = 1$ ,  $\text{ord}(A) = 9 > 2$ , and  $\lambda^2 + (\alpha^7 + \alpha^{-7})\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1}) \in \mathbb{F}_{26}[x]$ , where  $\rho = \alpha^7, \rho^{-1} = \alpha^{-7} \in \mathbb{F}_{26}^*$  are two eigenvalues of  $A$ . Both the minimal polynomial of  $\rho$  and  $\rho^{-1}$  over  $\mathbb{F}_2$  are  $m_\rho(x) = m_{\rho^{-1}}(x) = x^6 + x^3 + 1$ . Let

$$\mathcal{L}' = O_\infty \setminus \{\infty\} = \{\alpha^7, \alpha^5, \alpha^{30}, \alpha, \alpha^8, \alpha^{51}, \alpha^{40}, \alpha^{56}\} \subseteq \mathbb{F}_{26}$$

and

$$\mathcal{L} = O_{\alpha^2} = \{\alpha^2, \alpha^{52}, \alpha^{35}, \alpha^{28}, \alpha^{38}, \alpha^{16}, \alpha^{19}, \alpha^{27}, \alpha^{26}\} \subseteq \mathbb{F}_{26},$$

where  $|\mathcal{L}'| = 8$  and  $|\mathcal{L}| = 9 = \text{ord}(A)$ .

(1) If  $g_1(x) = x$  and  $g_1^2(x) = x^2$ , we can obtain that the expurgated Goppa codes  $\tilde{\Gamma}(\mathcal{L}, g_1)$  and  $\tilde{\Gamma}(\mathcal{L}, g_1^2)$ , the extended Goppa codes  $\bar{\Gamma}(\mathcal{L}', g_1)$  and  $\bar{\Gamma}(\mathcal{L}', g_1^2)$  are all  $[9, 2, 6]$  cyclic codes and their generator polynomials are  $(x+1)(x^6 + x^3 + 1) = (x+1)m_{\rho^{-1}}(x)$  by the Magma program.

(2) If  $g_2(x) = x + \alpha^7 + \alpha^{-7}$  and  $g_2^2(x) = (x + \alpha^7 + \alpha^{-7})^2$ , we can obtain that the expurgated Goppa codes  $\tilde{\Gamma}(\mathcal{L}, g_2)$  and  $\tilde{\Gamma}(\mathcal{L}, g_2^2)$ , the extended Goppa codes  $\bar{\Gamma}(\mathcal{L}', g_2)$  and  $\bar{\Gamma}(\mathcal{L}', g_2^2)$  are all  $[9, 2, 6]$  cyclic codes and their generator polynomials are  $(x+1)(x^6 + x^3 + 1) = (x+1)m_\rho(x)$  by the Magma program.

**Example 3.14.** Let  $\mathbb{F}_{2^4}^* = \langle \alpha \rangle$  and  $\mathbb{F}_{2^8}^* = \langle \gamma \rangle$ , where  $\alpha = \gamma^{17}$ . Let  $A = \begin{pmatrix} \alpha^{11} & \alpha^5 \\ \alpha^3 & \alpha^6 \end{pmatrix} \in PGL_2(\mathbb{F}_{2^4})$ , then  $|A| = 1$ ,  $\text{ord}(A) = 17 > 2$ , and  $\lambda^2 + (\alpha^{11} + \alpha^6)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1}) \in \mathbb{F}_{2^8}[x]$ , where  $\rho = \gamma^{45}, \rho^{-1} = \gamma^{-45} = \gamma^{210} \in \mathbb{F}_{2^8}^*$  are two eigenvalues of  $A$ . Both the minimal polynomial of  $\rho$  and  $\rho^{-1}$  over  $\mathbb{F}_2$  are  $m_\rho(x) = m_{\rho^{-1}}(x) = x^8 + x^5 + x^4 + x^3 + 1$ . Let

$$\mathcal{L}' = O_\infty \setminus \{\infty\} = \{\gamma^{136}, \gamma^{119}, \gamma^{68}, 1, \gamma^{17}, \gamma^{34}, \gamma^{221}, \gamma^{170}, \gamma^{153}, \\ 0, \gamma^{238}, \gamma^{204}, \gamma^{102}, \gamma^{187}, \gamma^{85}, \gamma^{51}\} \subseteq \mathbb{F}_{2^8}$$

and

$$\mathcal{L} = O_{\gamma^3} = \{\gamma^3, \gamma^{26}, \gamma^{147}, \gamma^{172}, \gamma^{32}, \gamma^{87}, \gamma^{232}, \gamma^{128}, \gamma^{241}, \\ \gamma^{61}, \gamma^{144}, \gamma^{191}, \gamma^{39}, \gamma^{175}, \gamma^{38}, \gamma^{25}, \gamma^{78}\} \subseteq \mathbb{F}_{2^8}$$

where  $|\mathcal{L}'| = 16$  and  $|\mathcal{L}| = 17 = \text{ord}(A)$ .

(1) If  $g_1(x) = x + \frac{\alpha^{11} + \gamma^{45}}{\alpha^3}$  and  $g_1(x)^2 = (x + \frac{\alpha^{11} + \gamma^{45}}{\alpha^3})^2$ , we can obtain that the expurgated Goppa codes  $\tilde{\Gamma}(\mathcal{L}, g_1)$  and  $\tilde{\Gamma}(\mathcal{L}, g_1^2)$ , the extended Goppa codes  $\bar{\Gamma}(\mathcal{L}', g_1)$  and  $\bar{\Gamma}(\mathcal{L}', g_1^2)$  are all  $[17, 8, 6]$  cyclic codes and their generator polynomials are  $(x + 1)(x^8 + x^5 + x^4 + x^3 + 1) = (x + 1)m_{\rho^{-1}}(x)$  by the Magma program.

(2) If  $g_2(x) = x + \frac{\alpha^{11} + \gamma^{210}}{\alpha^3}$  and  $g_2(x)^2 = (x + \frac{\alpha^{11} + \gamma^{210}}{\alpha^3})^2$ , we can obtain that the expurgated Goppa codes  $\tilde{\Gamma}(\mathcal{L}, g_2)$  and  $\tilde{\Gamma}(\mathcal{L}, g_2^2)$ , the extended Goppa codes  $\bar{\Gamma}(\mathcal{L}', g_2)$  and  $\bar{\Gamma}(\mathcal{L}', g_2^2)$  are all  $[17, 8, 6]$  cyclic codes and their generator polynomials are  $(x + 1)(x^8 + x^5 + x^4 + x^3 + 1) = (x + 1)m_\rho(x)$  by the Magma program.

**3.2.  $g(x) = g_1(x)^s$  or  $g_2(x)^t$ ,  $s, t \in \mathbb{N}$  and  $s, t < n - 1$ .**

In the subsection, we shall assume that  $g(x) = g_1(x)^s$  or  $g_2(x)^t$ , where  $s, t \in \mathbb{N}$  and  $s, t < n - 1$ . We shall find generator polynomials of cyclic expurgated or extended Goppa codes in the case of  $\sigma^j = 1$  of Section 2.3. There is a surprising result that the expurgated or extended Goppa codes are all BCH codes, which are an important class of cyclic codes.

**Definition 3.15.** ([18, Definition 9.44]) Let  $a$  be a nonnegative integer and  $\alpha \in \mathbb{F}_{q^m}$  be a primitive  $n$ -th root of unity. A BCH code  $\mathcal{C} = \langle u(x) \rangle$  over  $\mathbb{F}_q$  of length  $n$  and designed distance  $\delta$ ,  $2 \leq \delta \leq n$ , is a cyclic code with the generator polynomial:

$$u(x) = \text{lcm}(m^{(a)}(x), m^{(a+1)}(x), \dots, m^{(a+\delta-2)}(x)),$$

where each  $m^{(i)}(x)$  is the minimal polynomial of  $\alpha^i$  over  $\mathbb{F}_q$ . Then the minimum distance of the BCH code  $\mathcal{C}$  is at least  $\delta$ .

**Theorem 3.16.** Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PGL_2(\mathbb{F}_q)$  be of order  $n > 2$  with  $|A| = ad + bc = 1$  and  $c \neq 0$ ,  $G = \langle A \rangle$  a subgroup of  $PGL_2(\mathbb{F}_q)$ , and  $|\lambda E_2 - A| = \lambda^2 + (a + d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  reducible over  $\mathbb{F}_q$ . Let

$$\mathcal{L}' = O_\infty \setminus \{\infty\} = \{P(\rho), \dots, P(\rho^{n-1})\} = \{\gamma, A(\gamma), \dots, A^{n-2}(\gamma)\} \subseteq \mathbb{F}_q$$

and

$$\mathcal{L} = O_\alpha = \{\alpha_1 = \alpha, \alpha_2 = A(\alpha), \dots, \alpha_n = A^{n-1}(\alpha)\} \subseteq \mathbb{F}_q$$

be the orbit of  $\alpha$  under the action of  $G$  if  $\alpha \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\} \cup O_\infty$ .

(1) If  $g_1(x)^s = (x + \frac{a+\rho}{c})^s \in \mathbb{F}_q[x]$ , then both the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g_1^s)$  are BCH codes of length  $n$  with designed distances  $\delta = s + 2$  and the generator polynomials:

$$u_1(x) = (x + 1)\text{lcm}\{m_{\rho^{-i}}(x) : i = 1, 2, \dots, s\}, \quad (3.7)$$

where each  $m_{\rho^{-i}}(x)$  is the minimal polynomial of  $\rho^{-i}$  over  $\mathbb{F}_2$ .

(2) If  $g_2(x)^t = (x + \frac{a+\rho^{-1}}{c})^t \in \mathbb{F}_q[x]$ , then both the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g_2^t)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g_2^t)$  are BCH codes of length  $n$  with designed distances  $\delta = t + 2$  and the generator polynomials:

$$u_2(x) = (x + 1)\text{lcm}\{m_{\rho^i}(x) : i = 1, 2, \dots, t\}, \quad (3.8)$$

where each  $m_{\rho^i}(x)$  is the minimal polynomial of  $\rho^i$  over  $\mathbb{F}_2$ .

Moreover, if  $\deg u_1(x) = n$ , then both  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$  and  $\bar{\Gamma}(\mathcal{L}', g_1^s)$  are zero; similarly, both  $\tilde{\Gamma}(\mathcal{L}, g_2^t)$  and  $\bar{\Gamma}(\mathcal{L}', g_2^t)$  are also zero if  $\deg u_2(x) = n$ .

*Proof.* Let  $\lambda^2 + (a+d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  be reducible over  $\mathbb{F}_q$ , then by Proposition 3.5,

$$g_1(x)^s = (x + \frac{a+\rho}{c})^s, \quad g_2(x)^t = (x + \frac{a+\rho^{-1}}{c})^t \in \mathbb{F}_q[x]$$

satisfy the condition in (2.3). Hence by Lemma 2.3, the expurgated Goppa codes  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$ ,  $\tilde{\Gamma}(\mathcal{L}, g_2^t)$  and the extended Goppa codes  $\bar{\Gamma}(\mathcal{L}', g_1^s)$ ,  $\bar{\Gamma}(\mathcal{L}', g_2^t)$  are all cyclic of length  $n$ . In the following, we only need to find the generator polynomial of  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$ . Similarly, we can also obtain those of  $\bar{\Gamma}(\mathcal{L}', g_1^s)$ ,  $\tilde{\Gamma}(\mathcal{L}, g_2^t)$  and  $\bar{\Gamma}(\mathcal{L}', g_2^t)$ .

For the cyclic expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$ , we shall prove that the polynomial  $u_1(x)$  in (3.7) is its generator polynomial. Now we consider the parity-check matrix of  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$  as follows:

$$\tilde{H} = \begin{pmatrix} \frac{1}{(\alpha_1 + \frac{a+\rho}{c})^s} & \frac{1}{(\alpha_2 + \frac{a+\rho}{c})^s} & \cdots & \frac{1}{(\alpha_n + \frac{a+\rho}{c})^s} \\ \frac{\alpha_1}{(\alpha_1 + \frac{a+\rho}{c})^s} & \frac{\alpha_2}{(\alpha_2 + \frac{a+\rho}{c})^s} & \cdots & \frac{\alpha_n}{(\alpha_n + \frac{a+\rho}{c})^s} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\alpha_1^s}{(\alpha_1 + \frac{a+\rho}{c})^s} & \frac{\alpha_2^s}{(\alpha_2 + \frac{a+\rho}{c})^s} & \cdots & \frac{\alpha_n^s}{(\alpha_n + \frac{a+\rho}{c})^s} \end{pmatrix}.$$

Since  $1, x, \dots, x^s$  and  $1, x + \frac{a+\rho}{c}, \dots, (x + \frac{a+\rho}{c})^s$  are two sets of bases of  $\mathbb{F}_q[x]_{\leq s}$  over  $\mathbb{F}_q$ , performing elementary row operations over  $\mathbb{F}_q$  on the above matrix, we can obtain



the equivalent matrix over  $\mathbb{F}_q$ :

$$\tilde{H}_1 = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ \frac{1}{\alpha_1 + \frac{a+\rho}{c}} & \frac{1}{\alpha_2 + \frac{a+\rho}{c}} & \cdots & \frac{1}{\alpha_n + \frac{a+\rho}{c}} \\ \vdots & \vdots & & \vdots \\ \frac{1}{(\alpha_1 + \frac{a+\rho}{c})^{s-1}} & \frac{1}{(\alpha_2 + \frac{a+\rho}{c})^{s-1}} & \cdots & \frac{1}{(\alpha_n + \frac{a+\rho}{c})^{s-1}} \\ \frac{1}{(\alpha_1 + \frac{a+\rho}{c})^s} & \frac{1}{(\alpha_2 + \frac{a+\rho}{c})^s} & \cdots & \frac{1}{(\alpha_n + \frac{a+\rho}{c})^s} \end{pmatrix}.$$

Then  $\tilde{H}_1$  is also the parity-check matrix of  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$ .

Hence  $\mathbf{c} = (c_1, \dots, c_n) \in \tilde{\Gamma}(\mathcal{L}, g_1^s)$  if and only if  $\tilde{H}_1 \mathbf{c}^T = 0$  if and only if  $\mathbf{c}$  is a solution of the system of equations

$$\begin{cases} c_1 + c_2 + \cdots + c_n = 0, \\ \frac{c_1}{\alpha_1 + \frac{a+\rho}{c}} + \cdots + \frac{c_n}{\alpha_n + \frac{a+\rho}{c}} = 0, \\ \cdots \\ \frac{c_1}{(\alpha_1 + \frac{a+\rho}{c})^{s-1}} + \cdots + \frac{c_n}{(\alpha_n + \frac{a+\rho}{c})^{s-1}} = 0, \\ \frac{c_1}{(\alpha_1 + \frac{a+\rho}{c})^s} + \cdots + \frac{c_n}{(\alpha_n + \frac{a+\rho}{c})^s} = 0. \end{cases} \quad (3.9)$$

Let  $B = \begin{pmatrix} 0 & 1 \\ 1 & \frac{a+\rho}{c} \end{pmatrix}$  and

$$D = BAB^{-1} = \begin{pmatrix} a+d+\rho & c \\ \frac{\rho^2 + (a+d)\rho + 1}{c} & \rho \end{pmatrix} = \begin{pmatrix} \rho^{-2} & c\rho^{-1} \\ 0 & 1 \end{pmatrix} \triangleq \begin{pmatrix} e & f \\ 0 & 1 \end{pmatrix},$$

where  $e = \rho^{-2}$  and  $f = c\rho^{-1}$ . Then  $\text{ord}(D) = \text{ord}(A) = \text{ord}(e) = n$  and

$$D^i = BA^iB^{-1} = \begin{pmatrix} e^i & f \frac{e^i - 1}{e - 1} \\ 0 & 1 \end{pmatrix}, 1 \leq i \leq n - 1.$$

In (3.9),  $\frac{1}{\alpha_i + \frac{a+\rho}{c}} = B(\alpha_i) = B(A^{i-1}\alpha_1) = BA^{i-1}B^{-1}B(\alpha_1) = D^{i-1}B(\alpha_1)$ ,  $1 \leq i \leq n$ . Denote  $\beta = B(\alpha_1) = \frac{1}{\alpha_1 + \frac{a+\rho}{c}}$ .

Hence the system (3.9) is equivalent to the system

$$\begin{cases} c_1 + c_2 + \cdots + c_n = 0, \\ c_1\beta + c_2D(\beta) + \cdots + c_nD^{n-1}(\beta) = 0, \\ \cdots \\ c_1\beta^{s-1} + c_2(D(\beta))^{s-1} + \cdots + c_n(D^{n-1}(\beta))^{s-1} = 0, \\ c_1\beta^s + c_2(D(\beta))^s + \cdots + c_n(D^{n-1}(\beta))^s = 0, \end{cases} \quad (3.10)$$

where  $\beta = B(\alpha_1) = \frac{1}{\alpha_1 + \frac{a+\rho}{c}}$ .

We shall prove that  $\mathbf{c} = (c_1, \dots, c_n) \in \tilde{\Gamma}(\mathcal{L}, g_1^s)$  if and only if  $\mathbf{c}$  is a solution of the system of equations

$$\begin{cases} c_1 + c_2 + \dots + c_n = 0, \\ c_1 + c_2e + \dots + c_n e^{n-1} = 0, \\ \dots \\ c_1 + c_2e^{s-1} + \dots + c_n(e^{s-1})^{n-1} = 0, \\ c_1 + c_2e^s + \dots + c_n(e^s)^{n-1} = 0, \end{cases} \quad (3.11)$$

where  $e = \rho^{-2}$ . It means that the system in (3.10) is equivalent to the system in (3.11). In the following, we shall prove that the system in (3.10) is equivalent to the system in (3.11) by induction on  $s = \deg g(x)$ .

If  $\deg g(x) = s = 1$ , then it has been proved in Theorem 3.7. We assume that  $s > 1$  and that the statement is shown for  $\deg g(x)$  equal to  $s - 1$ , i.e., the system

$$\begin{cases} c_1 + c_2 + \dots + c_n = 0, \\ c_1\beta + c_2D(\beta) + \dots + c_nD^{n-1}(\beta) = 0, \\ \dots \\ c_1\beta^{s-1} + c_2(D(\beta))^{s-1} + \dots + c_n(D^{n-1}(\beta))^{s-1} = 0, \end{cases} \quad (3.12)$$

is equivalent to the system

$$\begin{cases} c_1 + c_2 + \dots + c_n = 0, \\ c_1 + c_2e + \dots + c_n e^{n-1} = 0, \\ \dots \\ c_1 + c_2e^{s-1} + \dots + c_n(e^{s-1})^{n-1} = 0. \end{cases} \quad (3.13)$$

Now we consider the case  $\deg g(x) = s$ .

Let  $s = s_1 + 1$ , i.e.,  $s_1 = s - 1$ , then the last equation in (3.10) is that

$$\begin{aligned} 0 &= c_1\beta^s + c_2(D(\beta))^s + \dots + c_n(D^{n-1}(\beta))^s \\ &= c_1\beta^{s_1}\beta + c_2(D(\beta))^{s_1}(e\beta + f) + \dots + c_n(D^{n-1}(\beta))^{s_1}(e^{n-1}\beta + f\frac{e^{n-1}-1}{e-1}) \\ &= (c_1\beta^{s_1} + c_2(D(\beta))^{s_1}e + \dots + c_n(D^{n-1}(\beta))^{s_1}e^{n-1})\beta \\ &\quad + \frac{f}{e-1}(c_2(D(\beta))^{s_1}(e-1) + \dots + c_n(D^{n-1}(\beta))^{s_1}(e^{n-1}-1)) \\ &= (c_1\beta^{s_1} + c_2(D(\beta))^{s_1}e + \dots + c_n(D^{n-1}(\beta))^{s_1}e^{n-1})\beta \\ &\quad + \frac{f}{e-1}(c_2(D(\beta))^{s_1}e + \dots + c_n(D^{n-1}(\beta))^{s_1}e^{n-1} - c_2(D(\beta))^{s_1} - \dots - c_n(D^{n-1}(\beta))^{s_1}) \\ &= (c_1\beta^{s_1} + c_2(D(\beta))^{s_1}e + \dots + c_n(D^{n-1}(\beta))^{s_1}e^{n-1})(\beta + \frac{f}{e-1}). \end{aligned}$$

Since  $\beta + \frac{f}{e-1} = B(\alpha_1) + \frac{f}{e-1} \neq 0$  (otherwise,  $\frac{1}{\alpha_1 + \frac{a+\rho}{c}} = \frac{c\rho^{-1}}{\rho^{-2}-1}$ , i.e.,  $\alpha_1 = \frac{a+\rho^{-1}}{c}$ , which is contradictory with the choice of  $\alpha_1$ ), the last equation in (3.10) is equivalent to

$$0 = c_1\beta^{s_1} + c_2(D(\beta))^{s_1}e + \dots + c_n(D^{n-1}(\beta))^{s_1}e^{n-1}$$

$$\begin{aligned}
 &= c_1\beta^{s_1} + c_2(e\beta + f\frac{e-1}{e-1})^{s_1}e + \cdots + c_n(e^{n-1}\beta + f\frac{e^{n-1}-1}{e-1})^{s_1}e^{n-1} \\
 &= (c_1 + c_2e^s + \cdots + c_n(e^s)^{n-1})\beta^{s_1} \\
 &+ \sum_{i=1}^{s_1} \frac{C_{s_1}^i f^i \beta^{s_1-i}}{(e-1)^i} (c_2e^{s_1+1-i}(e-1)^i + \cdots + c_n(e^{n-1})^{s_1+1-i}(e^{n-1}-1)^i), \quad (3.14)
 \end{aligned}$$

where each  $C_{s_1}^i$  is a combinatorial number.

For  $1 \leq i \leq s_1$ ,  $i = \sum_{j=1}^k 2^{d_j}$  is called the 2-adic expansion of  $i$ , where  $d_1 < d_2 < \cdots < d_k$ , and  $k$  is called the algebraic degree of  $i$ . In (3.14),

$$\begin{aligned}
 &c_2e^{s_1+1-i}(e-1)^i + \cdots + c_n(e^{n-1})^{s_1+1-i}(e^{n-1}-1)^i \\
 &= c_2e^{s-i} \prod_{j=1}^k (e^{2^{d_j}} + 1) + \cdots + c_n(e^{n-1})^{s-i} \prod_{j=1}^k ((e^{n-1})^{2^{d_j}} + 1) \\
 &= c_2e^s + c_3(e^s)^2 + \cdots + c_n(e^s)^{n-1} \\
 &+ \sum_{j=1}^{2^k-1} (c_2e^{s-i+t_j} + c_3(e^{s-i+t_j})^2 + \cdots + c_n(e^{s-i+t_j})^{n-1}),
 \end{aligned}$$

where each  $t_j$  is a partial sum of  $2^{d_1}, 2^{d_2}, \dots, 2^{d_k}$  and  $t_j < i$ . By the induction hypothesis, each  $c_2e^{s-i+t_j} + c_3(e^{s-i+t_j})^2 + \cdots + c_n(e^{s-i+t_j})^{n-1} = c_1$ . Hence the equation in (3.14) is equivalent to

$$\begin{aligned}
 &(c_1 + c_2e^s + \cdots + c_n(e^s)^{n-1})(\beta^{s_1} + \sum_{i=1}^{s_1} \frac{C_{s_1}^i f^i \beta^{s_1-i}}{(e-1)^i}) \\
 &= (c_1 + c_2e^s + \cdots + c_n(e^s)^{n-1})(\beta + \frac{f}{e-1})^{s_1} = 0,
 \end{aligned}$$

where  $\beta = B(\alpha_1) = \frac{1}{\alpha_1 + \frac{a+p}{c}}$  and  $\beta + \frac{f}{e-1} \neq 0$ . Then the last equation in (3.10) holds if and only if  $c_1 + c_2e^s + \cdots + c_n(e^s)^{n-1} = 0$  under the induction hypothesis, which means the system in (3.10) is equivalent to the system in (3.11).

Therefore,  $\mathbf{c} = (c_1, \dots, c_n) \in \widetilde{\Gamma}(\mathcal{L}, g_1^s)$  if and only if  $\mathbf{c}$  is a solution of the system (3.11). Moreover, let  $m_{\rho^{-i}}(x)$  be the minimal polynomials of  $\rho^{-i}$  over  $\mathbb{F}_2$ , then they are also the minimal polynomials of  $e^i = \rho^{-2i}$  over  $\mathbb{F}_2$ ,  $i = 1, 2, \dots, s$ . Since  $\widetilde{\Gamma}(\mathcal{L}, g_1^s)$  is a cyclic code of length  $n$ , whose elements satisfy the system (3.11), its generator polynomial is

$$u_1(x) = (x+1)\text{lcm}\{m_{\rho^{-i}}(x) : i = 1, 2, \dots, s\}.$$

Similarly, we can get generator polynomial of  $\overline{\Gamma}(\mathcal{L}', g_1^s)$  is  $u_1(x)$  and generator polynomials of both  $\widetilde{\Gamma}(\mathcal{L}, g_2^t)$  and  $\overline{\Gamma}(\mathcal{L}', g_2^t)$  are

$$u_2(x) = (x+1)\text{lcm}\{m_{\rho^i}(x) : i = 1, 2, \dots, t\},$$

where  $m_{\rho^i}(x)$  are the minimal polynomials of  $\rho^i$  over  $\mathbb{F}_2$ .

If  $\deg u_1(x) = n$ , then the dimensions of both  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$  and  $\bar{\Gamma}(\mathcal{L}', g_1^s)$  are  $n - \deg u_1(x) = 0$ , so both  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$  and  $\bar{\Gamma}(\mathcal{L}', g_1^s)$  are zero; similarly, both  $\tilde{\Gamma}(\mathcal{L}, g_1^t)$  and  $\bar{\Gamma}(\mathcal{L}', g_1^t)$  are also zero if  $\deg u_2(x) = n$ .  $\square$

**Remark 3.17.** In Theorem 3.16, since  $\rho^j$  and  $\rho^{2j}$  have the same minimal polynomial over  $\mathbb{F}_2$  for every  $j \in \mathbb{Z}$ , (3.7) and (3.8) in Theorem 3.16 can be changed into

$$\begin{aligned} u_1(x) &= (x+1) \text{lcm}\{m_{\rho^{-i}}(x) : i = 1, 2, \dots, 2\lfloor \frac{s+1}{2} \rfloor\} \\ &= (x+1) \text{lcm}\{m_{\rho^{-i}}(x) : i = 1, 3, \dots, 2\lfloor \frac{s+1}{2} \rfloor - 1\} \end{aligned}$$

and

$$\begin{aligned} u_2(x) &= (x+1) \text{lcm}\{m_{\rho^i}(x) : i = 1, 2, \dots, 2\lfloor \frac{t+1}{2} \rfloor\} \\ &= (x+1) \text{lcm}\{m_{\rho^i}(x) : i = 1, 3, \dots, 2\lfloor \frac{t+1}{2} \rfloor - 1\}. \end{aligned}$$

By Definition 3.15, Theorem 3.16, and Remark 3.17, we can immediately obtain the following result.

**Corollary 3.18.** (1) In Theorem 3.16, the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g_1^s)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g_1^s)$  are BCH codes of length  $n$  with minimum distances  $d \geq 2\lfloor \frac{s+1}{2} \rfloor + 2$ .

(2) In Theorem 3.16, the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g_1^t)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g_1^t)$  are BCH codes of length  $n$  with minimum distances  $d \geq 2\lfloor \frac{t+1}{2} \rfloor + 2$ .

**Corollary 3.19.** Let  $A \in \text{PGL}_2(\mathbb{F}_q)$  be the matrix,  $G = \langle A \rangle$  the subgroup in Theorem 3.16, and  $|\lambda E_2 - A| = \lambda^2 + (a+d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  irreducible over  $\mathbb{F}_q$ . Let

$$\mathcal{L}' = O_\infty \setminus \{\infty\} = \{P(\rho), \dots, P(\rho^{n-1})\} = \{\gamma, A(\gamma), \dots, A^{n-2}(\gamma)\} \subseteq \mathbb{F}_{q^2}$$

and

$$\mathcal{L} = O_\alpha = \{\alpha_1 = \alpha, \alpha_2 = A(\alpha), \dots, \alpha_n = A^{n-1}(\alpha)\} \subseteq \mathbb{F}_{q^2}$$

be the orbit of  $\alpha$  under the action of  $G$  if  $\alpha \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\} \cup O_\infty$ .

Then the conclusions of Theorem 3.16 still hold but with the Goppa polynomials  $g_1^s(x), g_2^t(x) \in \mathbb{F}_{q^2}[x]$ . Moreover, let  $\lambda^2 + (a+d)\lambda + 1$  be irreducible over  $\mathbb{F}_q$ ,  $\text{ord}(A) = n \mid (q+1)$ , then by Remark 3.8,  $m_{\rho^i}(x) = m_{\rho^{-i}}(x), i = 1, 2, \dots, j, j = \max\{s, t\}$ .

**Example 3.20.** In Example 3.12, the minimal polynomials of  $\rho^{-3}, \rho^3, \rho^{-5}, \rho^5, \rho^{-7}, \rho^7$  over  $\mathbb{F}_2$  are  $m_{\rho^{-3}}(x) = x^3 + x^2 + 1$ ,  $m_{\rho^3}(x) = x^3 + x + 1$ ,  $m_{\rho^{-5}}(x) = m_\rho(x) = x^6 + x^5 + x^4 + x^2 + 1$ ,  $m_{\rho^5}(x) = m_{\rho^{-1}}(x) = x^6 + x^4 + x^2 + x + 1$ ,  $m_{\rho^{-7}}(x) = m_{\rho^7}(x) = x^2 + x + 1$ , respectively. Let  $g_1(x) = x + \frac{\alpha^5 + \alpha^3}{\alpha^{13}} \in \mathbb{F}_{2^6}[x]$ ,  $g_2(x) = x + \frac{\alpha^5 + \alpha^{-3}}{\alpha^{13}} \in \mathbb{F}_{2^6}[x]$ .

(1)

$$\begin{aligned} \tilde{\Gamma}(\mathcal{L}, g_1^3) &= \tilde{\Gamma}(\mathcal{L}, g_1^4) = \bar{\Gamma}(\mathcal{L}', g_1^3) = \bar{\Gamma}(\mathcal{L}', g_1^4) \\ &= \langle (x+1)(x^3 + x^2 + 1)(x^6 + x^4 + x^2 + x + 1) \rangle \\ &= \langle (x+1) \text{lcm}(m_{\rho^{-1}}(x), m_{\rho^{-3}}(x)) \rangle, \end{aligned}$$

$$\begin{aligned}
 \tilde{\Gamma}(\mathcal{L}, g_2^3) &= \tilde{\Gamma}(\mathcal{L}, g_2^4) = \bar{\Gamma}(\mathcal{L}', g_2^3) = \bar{\Gamma}(\mathcal{L}', g_2^4) \\
 &= \langle (x+1)(x^3+x+1)(x^6+x^5+x^4+x^2+x+1) \rangle \\
 &= \langle (x+1)\text{lcm}(m_\rho(x), m_{\rho^3}(x)) \rangle,
 \end{aligned}$$

and they are all [21, 11, 6] cyclic codes by the Magma program.

(2)

$$\begin{aligned}
 \tilde{\Gamma}(\mathcal{L}, g_1^5) &= \tilde{\Gamma}(\mathcal{L}, g_1^6) = \bar{\Gamma}(\mathcal{L}', g_1^5) = \bar{\Gamma}(\mathcal{L}', g_1^6) \\
 &= \langle (x+1)(x^3+x^2+1)(x^6+x^4+x^2+x+1)(x^6+x^5+x^4+x^2+1) \rangle \\
 &= \langle (x+1)\text{lcm}\{m_{\rho^{-i}}(x) : i = 1, 3, 5\} \rangle,
 \end{aligned}$$

$$\begin{aligned}
 \tilde{\Gamma}(\mathcal{L}, g_2^5) &= \tilde{\Gamma}(\mathcal{L}, g_2^6) = \bar{\Gamma}(\mathcal{L}', g_2^5) = \bar{\Gamma}(\mathcal{L}', g_2^6) \\
 &= \langle (x+1)(x^3+x+1)(x^6+x^4+x^2+x+1)(x^6+x^5+x^4+x^2+1) \rangle \\
 &= \langle (x+1)\text{lcm}\{m_{\rho^i}(x) : i = 1, 3, 5\} \rangle,
 \end{aligned}$$

and they are all [21, 5, 10] cyclic codes by the Magma program.

(3)

$$\begin{aligned}
 \tilde{\Gamma}(\mathcal{L}, g_1^7) &= \tilde{\Gamma}(\mathcal{L}, g_1^8) = \bar{\Gamma}(\mathcal{L}', g_1^7) = \bar{\Gamma}(\mathcal{L}', g_1^8) \\
 &= \langle (x+1)(x^2+x+1)(x^3+x^2+1)(x^6+x^4+x^2+x+1)(x^6+x^5+x^4+x^2+1) \rangle \\
 &= \langle (x+1)\text{lcm}\{m_{\rho^{-i}}(x) : i = 1, 3, 5, 7\} \rangle,
 \end{aligned}$$

$$\begin{aligned}
 \tilde{\Gamma}(\mathcal{L}, g_2^7) &= \tilde{\Gamma}(\mathcal{L}, g_2^8) = \bar{\Gamma}(\mathcal{L}', g_2^7) = \bar{\Gamma}(\mathcal{L}', g_2^8) \\
 &= \langle (x+1)(x^2+x+1)(x^3+x+1)(x^6+x^4+x^2+x+1)(x^6+x^5+x^4+x^2+1) \rangle \\
 &= \langle (x+1)\text{lcm}\{m_{\rho^i}(x) : i = 1, 3, 5, 7\} \rangle,
 \end{aligned}$$

and they are all [21, 3, 12] cyclic codes by the Magma program.

3.3.  $g(x) = g_1(x)^s g_2(x)^t$ ,  $s, t \in \mathbb{N}$  and  $s + t < n - 1$ .

In the subsection, we shall assume that  $g(x) = g_1(x)^s g_2(x)^t$ , where  $s, t \in \mathbb{N}$  and  $s + t < n - 1$ . We shall find generator polynomials of cyclic expurgated or extended Goppa codes in the case of  $\sigma^j = 1$  of Section 2.3.

**Theorem 3.21.** Let  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PGL_2(\mathbb{F}_q)$  be of order  $n > 2$  with  $|A| = ad + bc = 1$  and  $c \neq 0$ ,  $G = \langle A \rangle$  a subgroup of  $PGL_2(\mathbb{F}_q)$ , and  $|\lambda E_2 - A| = \lambda^2 + (a + d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  reducible over  $\mathbb{F}_q$ . Let

$$\mathcal{L}' = O_\infty \setminus \{\infty\} = \{P(\rho), \dots, P(\rho^{n-1})\} = \{\gamma, A(\gamma), \dots, A^{n-2}(\gamma)\} \subseteq \mathbb{F}_q$$

and

$$\mathcal{L} = O_\alpha = \{\alpha_1 = \alpha, \alpha_2 = A(\alpha), \dots, \alpha_n = A^{n-1}(\alpha)\} \subseteq \mathbb{F}_q$$

be the orbit of  $\alpha$  under the action of  $G$  if  $\alpha \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\} \cup O_\infty$ .

If  $g(x) = g_1(x)^s g_2(x)^t = (x + \frac{a+\rho}{c})^s (x + \frac{a+\rho^{-1}}{c})^t \in \mathbb{F}_q[x]$ , then both the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g)$  are cyclic of length  $n$  and their generator polynomials are

$$u(x) = \text{lcm}(u_1(x), u_2(x)), \quad (3.15)$$

where  $\tilde{\Gamma}(\mathcal{L}, g_1^s) = \bar{\Gamma}(\mathcal{L}', g_1^s) = \langle u_1(x) \rangle$  and  $\tilde{\Gamma}(\mathcal{L}, g_2^t) = \bar{\Gamma}(\mathcal{L}', g_2^t) = \langle u_2(x) \rangle$ ,  $u_1(x)$  and  $u_2(x)$  are defined as (3.7) and (3.8).

Moreover, if  $\deg u(x) = n$ , then both  $\tilde{\Gamma}(\mathcal{L}, g)$  and  $\bar{\Gamma}(\mathcal{L}', g)$  are zero.

*Proof.* Let  $\lambda^2 + (a+d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  be reducible over  $\mathbb{F}_q$ , then by Proposition 3.5,

$$g(x) = g_1(x)^s g_2(x)^t \in \mathbb{F}_q[x]$$

satisfies the condition in (2.3). Hence by Lemma 2.3, both the expurgated Goppa code  $\tilde{\Gamma}(\mathcal{L}, g)$  and the extended Goppa code  $\bar{\Gamma}(\mathcal{L}', g)$  are cyclic of length  $n$ .

Furthermore, by  $\gcd(g_1(x), g_2(x)) = 1$ ,  $\Gamma(\mathcal{L}, g) = \Gamma(\mathcal{L}, g_1^s) \cap \Gamma(\mathcal{L}, g_2^t)$ , then

$$\tilde{\Gamma}(\mathcal{L}, g) = \tilde{\Gamma}(\mathcal{L}, g_1^s g_2^t) = \tilde{\Gamma}(\mathcal{L}, g_1^s) \cap \tilde{\Gamma}(\mathcal{L}, g_2^t) = \langle \text{lcm}(u_1(x), u_2(x)) \rangle,$$

$$\bar{\Gamma}(\mathcal{L}', g) = \bar{\Gamma}(\mathcal{L}', g_1^s g_2^t) = \bar{\Gamma}(\mathcal{L}', g_1^s) \cap \bar{\Gamma}(\mathcal{L}', g_2^t) = \langle \text{lcm}(u_1(x), u_2(x)) \rangle,$$

where  $\tilde{\Gamma}(\mathcal{L}, g_1^s) = \bar{\Gamma}(\mathcal{L}, g_1^s) = \langle u_1(x) \rangle$  and  $\tilde{\Gamma}(\mathcal{L}, g_2^t) = \bar{\Gamma}(\mathcal{L}', g_2^t) = \langle u_2(x) \rangle$ .

If  $\deg u(x) = n$ , then the dimensions of both  $\tilde{\Gamma}(\mathcal{L}, g)$  and  $\bar{\Gamma}(\mathcal{L}', g)$  are  $n - \deg u(x) = 0$ , so both  $\tilde{\Gamma}(\mathcal{L}, g)$  and  $\bar{\Gamma}(\mathcal{L}', g)$  are zero.  $\square$

**Corollary 3.22.** Let  $A \in PGL_2(\mathbb{F}_q)$  be the matrix,  $G = \langle A \rangle$  the subgroup in Theorem 3.21, and  $|\lambda E_2 - A| = \lambda^2 + (a+d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  irreducible over  $\mathbb{F}_q$ . Let

$$\mathcal{L}' = O_\infty \setminus \{\infty\} = \{P(\rho), \dots, P(\rho^{n-1})\} = \{\gamma, A(\gamma), \dots, A^{n-2}(\gamma)\} \subseteq \mathbb{F}_{q^2}$$

and

$$\mathcal{L} = O_\alpha = \{\alpha_1 = \alpha, \alpha_2 = A(\alpha), \dots, \alpha_n = A^{n-1}(\alpha)\} \subseteq \mathbb{F}_{q^2}$$

be the orbit of  $\alpha$  under the action of  $G$  if  $\alpha \notin \{\frac{a+\rho}{c}, \frac{a+\rho^{-1}}{c}\} \cup O_\infty$ .

Then the conclusions of Theorem 3.21 still hold but with the Goppa polynomials  $g(x) = g_1^s(x)g_2^t(x) \in \mathbb{F}_{q^2}[x]$ . Moreover, let  $\lambda^2 + (a+d)\lambda + 1$  be irreducible over  $\mathbb{F}_q$ ,  $\text{ord}(A) = n \mid (q+1)$ , then by Remark 3.8,  $m_{\rho^i}(x) = m_{\rho^{-i}}(x)$ ,  $i = 1, 2, \dots, j$ ,  $j = \max\{s, t\}$ .

**Remark 3.23.** In Theorem 3.21, let  $\lambda^2 + (a+d)\lambda + 1 = (\lambda - \rho)(\lambda - \rho^{-1})$  be irreducible over  $\mathbb{F}_q$ ,  $g(x) = (g_1(x)g_2(x))^s \in \mathbb{F}_q[x]$  a monic polynomial satisfying the condition in (2.3), and  $\mathcal{L}, \mathcal{L}' \subseteq \mathbb{F}_q$ . Then the conclusions of Theorem 3.21 still hold.

So far, we have given all the answers to the case of  $\sigma^j = 1$  of Section 2.3.

**Example 3.24.** In Example 3.12, the minimal polynomials of  $\rho^{-3}$ ,  $\rho^3$ ,  $\rho^{-5}$ ,  $\rho^5$ ,  $\rho^{-7}$ , and  $\rho^7$  over  $\mathbb{F}_2$  are  $m_{\rho^{-3}}(x) = x^3 + x^2 + 1$ ,  $m_{\rho^3}(x) = x^3 + x + 1$ ,  $m_{\rho^{-5}}(x) = m_\rho(x) = x^6 + x^5 + x^4 + x^2 + 1$ ,  $m_{\rho^5}(x) = m_{\rho^{-1}}(x) = x^6 + x^4 + x^2 + x + 1$ ,  $m_{\rho^{-7}}(x) = m_{\rho^7}(x) = x^2 + x + 1$ , respectively. Let  $g_1(x) = x + \frac{\alpha^5 + \alpha^3}{\alpha^{13}} \in \mathbb{F}_{2^6}[x]$ ,  $g_2(x) = x + \frac{\alpha^5 + \alpha^{-3}}{\alpha^{13}} \in \mathbb{F}_{2^6}[x]$ .

(1) If  $g(x) = g_1(x)g_2(x)$ , then

$$\begin{aligned}\tilde{\Gamma}(\mathcal{L}, g) &= \bar{\Gamma}(\mathcal{L}', g) \\ &= \langle (x+1)(x^6 + x^4 + x^2 + x + 1)(x^6 + x^5 + x^4 + x^2 + x + 1) \rangle \\ &= \langle (x+1)\text{lcm}(m_{\rho^{-1}}(x), m_{\rho}(x)) \rangle,\end{aligned}$$

and they are [21, 8, 6] cyclic codes by the Magma program.

(2) If  $g(x) = g_1(x)g_2(x)^3$  or  $g(x) = g_1(x)g_2(x)^5$ , then

$$\begin{aligned}\tilde{\Gamma}(\mathcal{L}, g) &= \bar{\Gamma}(\mathcal{L}', g) \\ &= \langle (x+1)(x^3 + x + 1)(x^6 + x^4 + x^2 + x + 1)(x^6 + x^5 + x^4 + x^2 + x + 1) \rangle \\ &= \langle (x+1)\text{lcm}(m_{\rho^{-1}}(x), m_{\rho}(x), m_{\rho^3}(x)) \rangle,\end{aligned}$$

and they are [21, 5, 10] cyclic codes by the Magma program.

(3) If  $g(x) = g_1(x)^s g_2(x)^t$ , where  $(s, t) \in \{(3, 3), (3, 5), (5, 3), (5, 5)\}$ , then

$$\begin{aligned}\tilde{\Gamma}(\mathcal{L}, g) &= \bar{\Gamma}(\mathcal{L}', g) \\ &= \langle (x+1)(x^3 + x + 1)(x^3 + x^2 + 1)(x^6 + x^4 + x^2 + x + 1)(x^6 + x^5 + x^4 + x^2 + x + 1) \rangle \\ &= \langle (x+1)\text{lcm}(m_{\rho^{-1}}(x), m_{\rho^{-3}}(x), m_{\rho}(x), m_{\rho^3}(x)) \rangle,\end{aligned}$$

and they are [21, 2, 14] cyclic codes by the Magma program.

(4) If  $g(x) = g_1(x)g_2(x)^7$ , then

$$\begin{aligned}\tilde{\Gamma}(\mathcal{L}, g) &= \bar{\Gamma}(\mathcal{L}', g) \\ &= \langle (x+1)(x^2 + x + 1)(x^3 + x + 1)(x^6 + x^4 + x^2 + x + 1)(x^6 + x^5 + x^4 + x^2 + x + 1) \rangle \\ &= \langle (x+1)\text{lcm}(m_{\rho^{-1}}(x), m_{\rho}(x), m_{\rho^3}(x), m_{\rho^5}(x), m_{\rho^7}(x)) \rangle,\end{aligned}$$

and they are [21, 3, 12] cyclic codes by the Magma program.

(5) If  $g(x) = g_1(x)^7 g_2(x)$ , then

$$\begin{aligned}\tilde{\Gamma}(\mathcal{L}, g) &= \bar{\Gamma}(\mathcal{L}', g) \\ &= \langle (x+1)(x^2 + x + 1)(x^3 + x^2 + 1)(x^6 + x^4 + x^2 + x + 1)(x^6 + x^5 + x^4 + x^2 + x + 1) \rangle \\ &= \langle (x+1)\text{lcm}(m_{\rho^{-1}}(x), m_{\rho^{-3}}(x), m_{\rho^{-5}}(x), m_{\rho^{-7}}(x), m_{\rho}(x)) \rangle,\end{aligned}$$

and they are [21, 3, 12] cyclic codes by the Magma program.

Let  $M = \left( \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \sigma^j \right) \in P\Gamma L_2(\mathbb{F}_q)$  and  $g(x) = \sum_{i=0}^r g_i x^i \in \mathbb{F}_q[x]$  a monic polynomial of degree  $r$  satisfying the condition in (2.3). We gave generator polynomials of cyclic expurgated or extended Goppa codes if  $\sigma^j = 1$  in this paper. Based on the above discussion, now we have an open question:

If  $\sigma^j \neq 1$ , can you show generator polynomials of cyclic expurgated or extended Goppa codes?

## 4. CONCLUSION

Cyclic codes are an interesting type of linear codes and have wide applications in communication and storage systems due to their efficient encoding and decoding algorithms. Goppa codes are widely used in public-key cryptosystems. Cyclic Goppa codes, cyclic expurgated and extended Goppa codes were investigated (e.g., [2–6, 19]), but their generator polynomials have not shown so far.

This study aims to determine all the generator polynomials of cyclic expurgated or extended Goppa codes under some prescribed permutations induced by the projective general linear automorphism  $A \in PGL_2(\mathbb{F}_q)$ . Future research should be conducted to investigate the generator polynomials of cyclic expurgated or extended Goppa codes under some prescribed permutations induced by the projective semi-linear automorphism  $M \in P\Gamma L_2(\mathbb{F}_q)$ .

## ACKNOWLEDGEMENT

The authors would like to thank the editor and anonymous reviewers for their valuable comments that improved the quality of this paper.

## REFERENCES

- [1] M. Bardet, R. Mora, and J. P. Tillich, “Polynomial time key-recovery attack on high rate random alternant codes,” preprint 2023.
- [2] T. P. Berger, “Cyclic alternant codes induced by an automorphism of a GRS code,” in *Finite Field Theory, Applications and Algorithm (Contemporary Mathematics)*, R. Mullin and G. Mullen, Eds. Providence, RI, USA: American Mathematical Society, vol. 225, pp. 143-154, 1999.
- [3] T. P. Berger, “New classes of cyclic extended Goppa codes,” *IEEE Trans. Inf. Theory*, vol. 45, no. 4, pp. 1264-1266, 1999.
- [4] T. P. Berger, “Goppa and related codes invariant under a prescribed permutation,” *IEEE Trans. Inf. Theory*, vol. 46, no. 7, pp. 2628-2633, 2000.
- [5] T. P. Berger, “On the cyclicity of Goppa codes, parity-check subcodes of Goppa codes, and extended Goppa codes,” *Finite Fields Appl.*, vol. 6, no. 3, pp. 255-281, 2000.
- [6] T. P. Berger, “Quasi-cyclic Goppa codes,” In: *Proceedings of IEEE International Symposium on Information Theory*, 2000.
- [7] E. R. Berlekamp and O. Moreno, “Extended double-error-correcting binary Goppa codes are cyclic,” *IEEE Trans. Inform. Theory*, IT-19, pp. 817-818 1973.
- [8] A. Couvreur, I. M. Corbella, and R. Pellikaan, “A polynomial time attack against algebraic geometry code based public key cryptosystems,” In *2014 IEEE International Symposium on Information Theory*, pp. 1446-1450, 2014.
- [9] A. Couvreur, R. Mora, and J. P. Tillich, “A new approach based on quadratic forms to attack the McEliece cryptosystem,” preprint 2023.
- [10] B. Chen and G. Zhang, “Enumeration of Extended Irreducible Binary Goppa Codes,” *IEEE Trans. Inf. Theory*, vol. 68, no. 8, pp. 5145-5153, 2022.
- [11] B. Chen and G. Zhang, “A Tight Upper Bound on the Number of Non-Zero Weights of a Cyclic Code,” *IEEE Trans. Inf. Theory*, vol. 69, no. 2, pp. 995-1004, 2023.



- [12] G. I. Feng, K. K. Tzeng, "Further results about the cyclicity of generalized Goppa codes," *Discr. Math.* vol. 56, pp. 281-290, 1985.
- [13] J. C. Faugère, L. Perret, and F. D. Portzamparc, "Algebraic attack against variants of McEliece with Goppa polynomial of a special form," In *International Conference on the Theory and Application of Cryptology and Information Security*, pp. 21-41, Berlin, Heidelberg: Springer Berlin Heidelberg, 2014.
- [14] V. Goppa, "A new class of linear error correcting codes," *Probl. Peredachi Inf.*, vol. 6, no. 3, pp. 207-212, 1970.
- [15] V. Goppa, "Rational representation of codes and  $(L, g)$  codes," *Probl. Peredachi Inf.*, vol. 7, no. 3, pp. 41-49, 1971.
- [16] W. Huffman, V. Pless, "Fundamentals of Error Correcting codes," Cambridge University Press, Cambridge, 2003.
- [17] E. Kirshanova, and A. May, "Decoding McEliece with a Hint-Secret Goppa Key Parts Reveal Everything," *International Conference on Security and Cryptography for Networks*, Cham: Springer International Publishing, 2022.
- [18] R. Lidl and H. Niederreiter, "Finite Fields," Cambridge University Press, Cambridge, 2008.
- [19] X. Li, Q. Yue, "Non-binary irreducible quasi-cyclic parity-check subcodes of Goppa codes and extended Goppa codes," *Des. Codes Cryptogr.*, vol. 90, no. 7, pp. 1629-1647, 2022.
- [20] X. Li, Q. Yue, "Construction of expurgated and extended Goppa codes with dihedral automorphism groups," *IEEE Trans. Inf. Theory*, vol. 68, no. 10, pp. 6472-6480, 2022.
- [21] F. MacWilliams and N. Sloane, "The Theory of Error-Correcting codes," North Holland, Amsterdam, 1977.
- [22] R. J. McEliece, "A public-key cryptosystem based on algebraic coding theory," *DSN Progress Report 42-44*, pp. 114-116, 1978.
- [23] PQC Standardization Process: Announcing Four Candidates to be Standardized, Plus Fourth Round Candidates [EB/OL]. <https://csrc.nist.gov/News/2022/pqc-candidates-to-be-standardized-and-round-4>.
- [24] E. Richard. "Groups and Group Actions," Hilary and Trinity Terms, 2014.
- [25] R. M. Roth, "Introduction to Coding Theory," Cambridge University Press, Cambridge, 2006.
- [26] V. M. Sidelnikov and S. O. Shestakov, "On insecurity of cryptosystems based on generalized Reed-Solomon codes," *Discr. Math. Appl.*, vol. 2, no. 4, pp. 439-444, 1992.
- [27] H. Stichtenoth, "Which extended Goppa codes are cyclic?" *J. Comb. Theory*, vol. 51, pp. 205-220, 1989.
- [28] J. A. Thiong-Ly, "Automorphisms of two families of extended non binary cyclic Goppa codes," In *Lecture Notes in Computer Science*, vol. 229. New York/Berlin: Springer-Verlag, pp. 112-121, 1985.
- [29] K. K. Tzeng and C. Y. Yu, "Characterization theorems for extending Goppa codes to cyclic codes," *IEEE Trans. Inform. Theory*, vol. 25, pp. 246-250, 1979.
- [30] K. K. Tzeng and K. Zimmermann, "On extending Goppa codes to cyclic codes," *IEEE Trans. Inform. Theory*, vol. 21, pp. 712-716, 1975.
- [31] A. L. Vishnevetskii, "Cyclicity of extended Goppa codes," *Probl. Pered. Inform.*, vol. 18, pp. 171-175, 1982.
- [32] C. Wieschebrink, "Cryptanalysis of the Niederreiter public key scheme based on GRS subcodes," In *Post-Quantum Cryptography: Third International Workshop, PQCrypto 2010*, Darmstadt, Germany, May 25-28, 2010. *Proceedings 3* (pp. 61-72). Springer Berlin Heidelberg, 2010.

- [33] Y. Wu, Q. Yue, and S. Fan, “Self-reciprocal and self-conjugate-reciprocal irreducible factors of  $x^n - \lambda$  and their applications,” *Finite Fields Appl.*, vol. 63, pp. 101648, 2020.
- [34] C. Xing and S. Ling, “Coding Theory: A First Course,” Cambridge University Press, 2003.

School of Mathematics, Nanjing University of Aeronautics and Astronautics, Nanjing, 211100, P. R. China and State Key Laboratory of Cryptology, P.O. Box 5159, Beijing 100878, P. R. China  
*Email address:* `jiaxue0904@163.com`

College of Science, Nanjing University of Posts and Telecommunications, Nanjing, 210023, P. R. China  
*Email address:* `lfwzzu@126.com`

School of Mathematics, Nanjing University of Aeronautics and Astronautics, Nanjing, 211100, P. R. China and State Key Laboratory of Cryptology, P.O. Box 5159, Beijing 100878, P. R. China  
*Email address:* `sunhuan6558@163.com`

School of Mathematics, Nanjing University of Aeronautics and Astronautics, Nanjing, 211100, P. R. China and State Key Laboratory of Cryptology, P.O. Box 5159, Beijing 100878, P. R. China  
*Email address:* `yueqin@nuaa.edu.cn`