

FIRST EIGENVALUE OF JACOBI OPERATOR AND RIGIDITY RESULTS FOR CONSTANT MEAN CURVATURE HYPERSURFACES

MÁRCIO BATISTA, MARCOS P. CAVALCANTE, AND LUIZ R. MELO

ABSTRACT. In this paper, we obtain geometric upper bounds for the first eigenvalue λ_1^J of the Jacobi operator for both closed and compact with boundary hypersurfaces having constant mean curvature (CMC). As an application, we derive new rigidity results for the area of CMC hypersurfaces under suitable conditions on λ_1^J and the curvature of the ambient space. We also address the Jacobi-Steklov problem, proving geometric upper bounds for its first eigenvalue σ_1^J and deriving rigidity results related to the length of the boundary. Additionally, we present some results in higher dimensions related to the Yamabe invariants.

1. INTRODUCTION

In the context of the theory of minimal surfaces, stable surfaces or those of index one play a very special role. For instance, the well-known theorem by Schoen and Yau [29] states that if Σ is a closed orientable stable minimal surface in a Riemannian manifold M^3 with non-negative scalar curvature, then Σ is either a sphere or a torus. Moreover, infinitesimal rigidity is verified when Σ is a torus.

The stability index of a minimal surface can be defined as the number of negative eigenvalues of the Jacobi operator J associated with Σ . Geometrically, this index corresponds to the number of linearly independent normal directions in which infinitesimal variations result in a decrease in area. Indeed, controlling the index well often leads to a better understanding of the surface's geometry.

In this setting, it becomes natural to investigate geometric relationships involving the eigenvalues of J , in particular the first non-zero eigenvalue, denoted by λ_1^J . In [26], Perdomo obtained an estimate for λ_1^J for minimal surfaces in the sphere \mathbb{S}^3 , along with a rigidity result in the case of equality. More precisely, he demonstrated that if Σ is a compact, oriented minimal surface that is not totally geodesic in \mathbb{S}^3 , then the first eigenvalue of the Jacobi operator, λ_1^J , satisfies $\lambda_1^J \leq -4$. Furthermore, λ_1^J equals -4 if and only if Σ is isometric to the Clifford torus $\mathbb{S}^1\left(\frac{1}{\sqrt{2}}\right) \times \mathbb{S}^1\left(\frac{1}{\sqrt{2}}\right) \subset \mathbb{S}^3$.

Date: May 29, 2024.

2020 *Mathematics Subject Classification.* Primary: 53C42, 58J50. Secondary: 35P15, 49Q10.

From this work, several other estimates for λ_1^J have been obtained, but always in the case of closed submanifolds. We cite, for example, the works [1, 2, 10, 11, 12, 13, 24, 33], to name a few. We highlight the work of the first name author and Santos in [6], where they obtained general estimates for λ_1^J of closed constant mean curvature (CMC) surfaces, assuming only a bound on the curvature of the ambient manifold.

In this paper, we obtain new rigidity results for both closed and compact free boundary surfaces $\Sigma \subset M$ with constant mean curvature, and involving the first eigenvalue λ_1^J . Our results are related with some previous classical results, as for instance [4, 7, 25]. We recall that a CMC free-boundary surfaces are critical points for the area functional among volume-preserving variations that preserve the boundary $\partial\Sigma$ within ∂M . Notably, $\partial\Sigma$ intersects ∂M orthogonally. In this context, we obtain the following result.

Theorem 1.1. *Let M^3 be a Riemannian manifold with Ricci curvature $\text{Ric}_M \geq 2$ and convex boundary in the sense that its second fundamental form satisfies $\text{II}^{\partial M} \geq 0$. Let $\Sigma^2 \subset M^3$ be an immersed and orientable surface with constant mean curvature and free boundary in M . If $\lambda_1^J \geq -2$, then Σ has genus 0, a single connected boundary component, and $|\Sigma| \leq 2\pi$.*

If equality holds, then Σ is isometric to \mathbb{S}_+^2 . Moreover, if the Gaussian curvature of the boundary satisfies $K_{\partial M} \geq 1$, then M^3 is isometric to \mathbb{S}_+^3 .

This theorem follows from a general geometric inequality for the first eigenvalue λ_1^J , as proved in Theorems 2.3 and 2.4. In the rigidity part, we use the Hang-Wang Theorem [20].

In the case of surfaces without boundary, immersed in manifolds with nonnegative scalar curvature, we get:

Theorem 1.2. *Let M^3 be a Riemannian manifold with scalar curvature $S \geq 0$. Let $\Sigma^2 \subset M^3$ be a closed, immersed, and orientable surface with constant mean curvature $H \geq 2$ and $\lambda_1^J \geq -2$. Then Σ has genus 0 and $|\Sigma| \leq 4\pi$.*

Furthermore, if equality holds, then Σ is isometric to the unit sphere, and M^3 contains a subset isometric to the unit Euclidean ball whose boundary is Σ .

We obtain a similar result in the case of scalar curvature bounded from below by a negative constant:

Theorem 1.3. *Let M^3 be a Riemannian manifold with scalar curvature $S \geq -6$. Let $\Sigma^2 \subset M^3$ be an immersed, closed, and orientable surface with constant mean curvature $H \geq 2\sqrt{2}$ and $\lambda_1^J \geq -2$. Then Σ has genus 0 and $|\Sigma| \leq 4\pi$.*

Furthermore, if equality holds, then Σ is isometric to a sphere, and M^3 contains a subset isometric to the unit ball in hyperbolic space \mathbb{H}^3 .

We also derive theorems that holds in higher dimensions.

Theorem 1.4. *Let M^{n+1} be a closed orientable Riemannian manifold with scalar curvature $S \geq n(n+1)$, and let $\Sigma^n \subset M^{n+1}$ be a closed, immersed, orientable hypersurface with constant mean curvature H . If $\lambda_1^J \geq -n$, then*

$$(1.1) \quad n(n-1)|\Sigma| \leq \int_{\Sigma} S_{\Sigma} dA.$$

Moreover, the equality holds if, and only if, Σ is totally geodesic, $S|_{\Sigma} = n(n+1)$, and $\text{Ric}(N, N) = n$.

Furthermore, in the equality case with $K_M \geq 1$ and either

- (1) *n is even, or*
- (2) *n is odd and Σ is simply connected,*

then M^{n+1} is isometric to \mathbb{S}^{n+1} with the canonical metric.

In the case of hypersurfaces with boundary, we introduce a new eigenvalue problem, namely the Jacobi-Steklov problem, which utilizes the Jacobi operator instead of the Laplacian operator (see Section 4 for details). We denote the first eigenvalue of this problem by σ_1^J and provide upper estimates for σ_1^J in terms similar to those we obtained for λ_1^J (See Theorem 4.2).

As a consequence of our estimates and inspired by Mendes Theorem (see [23, Theorem 1.4]), we obtain the following rigidity theorem for the length of the boundary $\partial\Sigma$. In the rigidity part, we used a recent result by Mazet and Mendes [22].

Theorem 1.5. *Let M^3 be a Riemannian manifold with a non-empty boundary such that $\text{Ric}_M \geq 0$ and $\Pi^{\partial M} \geq g|_{\partial M}$. Let $\Sigma^2 \subset M^3$ be an immersed and orientable surface with constant mean curvature and free boundary such that $\sigma_1^J \geq -1$. Then, Σ has genus 0, a single connected boundary component, and $|\partial\Sigma| \leq 2\pi$.*

Furthermore, if equality holds, Σ is isometric to the disk \mathbb{D} and M^3 is isometric to the unit ball $\mathbb{B}^3 \subset \mathbb{R}^3$.

We also obtain the following estimate in higher dimensions. Here $H^{\partial\Sigma, \Sigma}$ denotes the mean curvature of $\partial\Sigma$ as a hypersurface of Σ .

Theorem 1.6. *Let (M^{n+1}, g) be a compact orientable Riemannian manifold with scalar curvature $S \geq 0$ and mean curvature of the boundary $H^{\partial M} \geq n$. Let $\Sigma^n \subset M^{n+1}$ be a compact, immersed, orientable hypersurface with constant mean curvature and free boundary. If $\sigma_1^J \geq -1$, then,*

$$(1.2) \quad 2(n-1)|\partial\Sigma| \leq \int_{\Sigma} S_{\Sigma} dA + 2 \int_{\partial\Sigma} H^{\partial\Sigma, \Sigma} da.$$

Moreover, equality holds if and only if Σ is totally geodesic, $\text{Ric}_M(N, N) = 0$, $\Pi^{\partial M}(N, N) = 1$, and $S|_{\Sigma} = 0$. In particular, $S_{\Sigma} = 0$ and $H^{\partial\Sigma, \Sigma} = n-1$. Furthermore, assuming $K_M \geq 0$, $\Pi^{\partial M} \geq g|_{\partial M}$, and the equality in (1.2), then Σ is isometric to \mathbb{B}^n and M is isometric to \mathbb{B}^{n+1} .

To conclude this paper, in Section 5 inspired in the work of Cai and Galloway [8] we present some geometric relations involving the eigenvalues λ_1^J and σ_1^J with the Yamabe invariants for manifolds with boundary introduced by Escobar in [16, 17].

2. THE FIRST EIGENVALUE OF THE JACOBI OPERATOR

Let M^{n+1} be an orientable Riemannian manifold with a (possibly empty) smooth boundary, and let Σ be a two-sided compact hypersurface with constant mean curvature, immersed in M^{n+1} . We denote by N the unit normal field along Σ .

If $\partial\Sigma \neq \emptyset$, we assume that $\partial\Sigma$ intersects ∂M orthogonally. In such case, Σ is a critical point of the area functional for variations that maintain the boundary of Σ within the boundary of M , and it is referred as a free-boundary hypersurface. Moreover, denoting ν as the unit conormal vector along $\partial\Sigma$, the formula for the second variation (see [27]) is given by the quadratic form

$$\mathcal{I}(\phi, \rho) = - \int_{\Sigma} \phi J \rho \, dv + \int_{\partial\Sigma} \phi \left\{ \frac{\partial \rho}{\partial \nu} - \Pi^{\partial M}(N, N) \rho \right\} da,$$

where $J = \Delta + \text{Ric}_M(N, N) + |A|^2$ is the Jacobi operator of Σ and $\Pi^{\partial M}$ denotes the second fundamental form of ∂M . Here Ric_M stands for the Ricci curvature of M , and A is the shape operator of Σ .

We say that $\rho \in C^\infty(\Sigma)$ is an *eigenfunction* of the Jacobi operator associated with the *eigenvalue* λ^J , if ρ is not identically zero and solves the Robin type boundary problem

$$(2.1) \quad \begin{cases} J\rho + \lambda^J \rho = 0, & \text{in } \Sigma, \\ \frac{\partial \rho}{\partial \nu} = \Pi^{\partial M}(N, N)\rho, & \text{on } \partial\Sigma. \end{cases}$$

Note that this is equivalent to saying that $\mathcal{I}(\rho, \phi) = \lambda^J \langle \rho, \phi \rangle_{L^2(\Sigma)}$, for every $\phi \in C^\infty(\Sigma)$. From standard elliptic theory, we know that the eigenvalues of the problem (2.1) are given by a diverging sequence denoted by

$$\lambda_1^J < \lambda_2^J \leq \lambda_3^J \leq \dots \nearrow \infty.$$

We recall that the index of Σ is defined as the number of negative eigenvalues, λ_k^J . In other words, it is the maximal dimension of a linear subspace V in $C^\infty(\Sigma)$ for which $\mathcal{I}(\rho, \rho) < 0$ for all $\rho \in V \setminus \{0\}$. This index is an important geometric invariant of Σ . For further details and significant results related to this topic, we refer to [3, 9, 27, 28].

The first eigenvalue λ_1^J can be obtained variationally by the Rayleigh formula

$$(2.2) \quad \lambda_1^J = \inf_{\phi \in C^\infty(\Sigma) \setminus \{0\}} \frac{\mathcal{I}(\phi, \phi)}{\int_{\Sigma} \phi^2 \, dv}.$$

In particular,

$$\lambda_1^J \leq \frac{\int_{\Sigma} |\nabla \phi|^2 - (\text{Ric}_M(N, N) + |A|^2) \phi^2 dv - \int_{\partial \Sigma} \text{II}^{\partial M}(N, N) \phi^2 da}{\int_{\Sigma} \phi^2 dv},$$

for every $\phi \in C^\infty(\Sigma) \setminus \{0\}$.

Remark 2.1. It is important to observe that all the above facts still hold if we are considering a hypersurface Σ without boundary, just omitting the boundary terms.

Tracing twice the Gauss equation we get the Schoen-Yau trick (cf. [29]), that is, the Ricci curvature in the normal direction can be written as

$$(2.3) \quad \text{Ric}_M(N, N) = \frac{1}{2}(S - S_\Sigma + H^2 - |A|^2),$$

where $H = \text{tr } A$ is the mean curvature of Σ , S is the scalar curvature of M and S_Σ is the scalar curvature of Σ .

Let us assume that Σ is a two dimensional surface immersed in M^3 . In this case, the Gauss curvature is given by $K_\Sigma = \frac{1}{2}S_\Sigma$. We will take the test function $\phi = 1$ in the variational characterization (2.2) of λ_1^J (see [29, Teorema 5.1]). Thus, using the equation (2.3) we have

$$\begin{aligned} \lambda_1^J &\leq \frac{\mathcal{I}(1, 1)}{\int_{\Sigma} 1 dv} = -\frac{\int_{\Sigma} (\text{Ric}_M(N, N) + |A|^2) dv + \int_{\partial \Sigma} \text{II}^{\partial M}(N, N) da}{|\Sigma|} \\ &= -\frac{1}{2|\Sigma|} \int_{\Sigma} (S - 2K_\Sigma + H^2 + |A|^2) dv - \frac{1}{|\Sigma|} \int_{\partial \Sigma} (H^{\partial M} - \kappa) da, \end{aligned}$$

where $H^{\partial M} = \text{tr } \text{II}^{\partial M}$ is mean curvature of ∂M in M , $\kappa = \langle \nabla_T \nu, T \rangle = \text{II}^{\partial M}(T, T)$ is the geodesic curvature of $\partial \Sigma$ in Σ and T is a unit vector tangent to $\partial \Sigma$.

Finally, using the Gauss-Bonnet Theorem, we have

$$(2.4) \quad \lambda_1^J \leq -\frac{1}{2|\Sigma|} \int_{\Sigma} (S + H^2 + |A|^2) dv - \frac{1}{|\Sigma|} \int_{\partial \Sigma} H^{\partial M} da + \frac{2\pi\chi(\Sigma)}{|\Sigma|}.$$

Note that if ρ is a first eigenfunction, using (2.1) and the Gauss equation (2.3), we have that

$$(2.5) \quad \begin{aligned} \Delta \rho &= -(\lambda_1^J + \text{Ric}_M(N, N) + |A|^2) \rho \\ &= -\left(\lambda_1^J + \frac{1}{2}S - \frac{1}{2}S_\Sigma + \frac{1}{2}H^2 + \frac{1}{2}|A|^2\right) \rho. \end{aligned}$$

The following lemma is well-known among experts, but we include a proof here for the completeness of our work. This result will be fundamental in characterizing the equality in our estimates.

Lemma 2.2. *Assume there exists $\rho \in C^\infty(\Sigma) \setminus \{0\}$ such that $\mathcal{I}(\rho, \rho) = \lambda_1^J \int_{\Sigma} \rho^2 dv$. Then, ρ is an eigenfunction of \mathcal{I} associated with the eigenvalue λ_1^J .*

Proof. Given $\phi \in C^\infty(\Sigma)$, consider the function

$$f(t) := \mathcal{I}(\rho + t\phi, \rho + t\phi) - \lambda_1^J \int_{\Sigma} (\rho + t\phi)^2 dv, \quad t \in \mathbb{R}.$$

From the definition of λ_1^J , it follows that $f(t) \geq 0$ for all $t \in \mathbb{R}$. Moreover, by hypothesis, $f(0) = 0$, and therefore, $t = 0$ is a global minimum point of f . Hence,

$$0 = f'(0) = 2\mathcal{I}(\rho, \phi) - 2\lambda_1^J \int_{\Sigma} \rho\phi dv.$$

That is, for every $\phi \in C^\infty(\Sigma)$, we have

$$- \int_{\Sigma} \phi J\rho dv + \int_{\partial\Sigma} \phi \left\{ \frac{\partial\rho}{\partial\nu} - \Pi^{\partial M}(N, N)\rho \right\} da = \lambda_1^J \int_{\Sigma} \rho\phi dv.$$

In particular, taking functions $\phi \in C^\infty(\Sigma)$ with compact support inside Σ , we conclude that $-J\rho = \lambda_1^J\rho$ and thus

$$\int_{\partial\Sigma} \phi \left\{ \frac{\partial\rho}{\partial\nu} - \Pi^{\partial M}(N, N)\rho \right\} da = 0,$$

for $\phi \in C^\infty(\Sigma)$. And once again, by the arbitrariness of $\phi \in C^\infty(\Sigma)$, we conclude that

$$\frac{\partial\rho}{\partial\nu} - \Pi^{\partial M}(N, N)\rho = 0,$$

and therefore ρ is an eigenfunction associated with the eigenvalue λ_1^J . \square

Using this terminology, we obtain the following estimate for λ_1^J with infinitesimal rigidity in the case of equality.

Theorem 2.3. *Let M^3 be a Riemannian manifold with boundary such that the scalar curvature and the mean curvature of the boundary satisfy $\inf S = a > -\infty$ and $\inf H^{\partial M} = b > -\infty$, respectively. Let $\Sigma^2 \subset M^3$ be a compact two-sided surface with constant mean curvature H and free boundary. Then*

$$\lambda_1^J \leq -\frac{1}{2} \left(\frac{3}{2}H^2 + a \right) - b \frac{|\partial\Sigma|}{|\Sigma|} + \frac{2\pi\chi(\Sigma)}{|\Sigma|}.$$

Moreover, equality holds if and only if the following conditions occur:

- (1) Σ is totally umbilical and the geodesic curvature of $\partial\Sigma$ in Σ is constant and equal to b .
- (2) $S|_{\Sigma} = a$, $H^{\partial M}|_{\partial\Sigma} = b$, and K is constant.
- (3) $\text{Ric}_M(N, N) = -\lambda_1^J - \frac{1}{2}H^2$ and $\Pi^{\partial M}(N, N) = 0$ along $\partial\Sigma$.

Proof. From the inequality (2.4), along with the assumptions $a \leq S$, $b \leq H^{\partial M}$ and the fact that $|A|^2 \geq \frac{1}{2}H^2$, we have

$$\begin{aligned} \lambda_1^J &\leq -\frac{1}{2|\Sigma|} \int_{\Sigma} (S + H^2 + |A|^2) dv - \frac{1}{|\Sigma|} \int_{\partial\Sigma} H^{\partial M} da + \frac{2\pi\chi(\Sigma)}{|\Sigma|} \\ &\leq -\frac{1}{2|\Sigma|} \int_{\Sigma} \left(a + \frac{3H^2}{2} \right) dv - b \frac{|\partial\Sigma|}{|\Sigma|} + \frac{2\pi\chi(\Sigma)}{|\Sigma|} \\ &\leq -\frac{a}{2} - \frac{3H^2}{4} - b \frac{|\partial\Sigma|}{|\Sigma|} + \frac{2\pi\chi(\Sigma)}{|\Sigma|}. \end{aligned}$$

Furthermore, if equality occurs, then equality holds in all the inequalities we used. In particular, $|A|^2 = \frac{1}{2}H^2$ and we have that Σ is totally umbilical. We also have that

$$\lambda_1^J = \frac{\mathcal{I}(1,1)}{\int_{\Sigma} 1 dv}$$

and thus, by Lemma 2.2, the constant function 1 is an eigenfunction associated with λ_1^J . Specifically, $S|_{\Sigma} = a$, $H^{\partial M}|_{\partial\Sigma} = b$, and by (2.5) we have that $\text{Ric}_M(N, N) = -\lambda_1^J - \frac{1}{2}H^2$ and K_{Σ} is constant equal to $a/2 + 3H^2/4 + \lambda_1^J$. Again, by (2.1) we have that $\text{II}(N, N) = 0$ and therefore $\text{II}(T, T) = H^{\partial M}|_{\partial\Sigma} = b$. The converse is immediate. \square

Next, by making a less refined estimate, we obtain a stricter constraint in the case of equality. More precisely:

Theorem 2.4. *Under the same conditions as Theorem 2.3, we have*

$$\lambda_1^J \leq -\frac{1}{2}(H^2 + a) - \frac{b}{|\Sigma|}|\partial\Sigma| + \frac{2\pi\chi(\Sigma)}{|\Sigma|}.$$

Moreover, equality holds if, and only if,

- (1) Σ is totally geodesic and $\partial\Sigma$ consists of geodesics of ∂M , and the geodesic curvature of $\partial\Sigma$ in Σ is constant and equal to b .
- (2) $S|_{\Sigma} = a$, $H^{\partial M}|_{\partial\Sigma} = b$, and K is constant.
- (3) $\text{Ric}_M(N, N) = -\lambda_1^J$ and $\text{II}^{\partial M}(N, N) = 0$ along $\partial\Sigma$.

Proof. The proof follows in the same manner as the previous theorem, but using the estimate $|A|^2 \geq 0$. In the case of equality, it follows that Σ is totally geodesic.

Furthermore, since Σ is totally geodesic and N is the normal vector of $\partial\Sigma$ in ∂M , it follows that the geodesic curvature of $\partial\Sigma$ in ∂M , given by $\langle -\nabla_T N, T \rangle$, is zero, where T is the unit tangent vector to $\partial\Sigma$. The converse is immediate. \square

We point out that the results in this present paper also hold in the case of manifolds with density (see [18] for a good introduction to this subject). In particular, in the case of surfaces without boundary (see Remark 2.1), we reobtain the main result in [6]. For constant density, the results read as follows.

Corollary 2.5 (Batista-Santos [6]). *Let (M^3, g) be a Riemannian manifold without boundary such that $\inf S \geq a > -\infty$. Let $\Sigma^2 \subset M^3$ be a closed surface with constant mean curvature H . Then, the first eigenvalue λ_1^J satisfies*

$$\lambda_1^J \leq -\frac{1}{2}(H^2 + a) + \frac{2\pi\chi(\Sigma)}{|\Sigma|}.$$

Moreover, equality holds if and only if Σ is totally geodesic, $S|_\Sigma = a$, and K_Σ is constant.

3. RIGIDITY RESULTS FOR THE AREA OF CMC SURFACES

In this section, as an application of our estimates for λ_1^J , we present the proofs of the area rigidity results for constant mean curvature surfaces, as cited in the Introduction.

We need the following characterization of the hemisphere of the 2-sphere. Although it is essentially proved in [20], we include the proof here for the sake of completeness.

Lemma 3.1. *Let (Σ^2, g) be a compact, orientable Riemannian surface with genus 0, constant Gaussian curvature equal to 1, and totally geodesic boundary. Then Σ is isometric to \mathbb{S}_+^2 .*

Proof. From the Gauss-Bonnet formula, we conclude that Σ is simply connected, in particular, $\partial\Sigma^2$ has only one component. Thus, using the Riemann Mapping Theorem, it follows that we can consider (Σ^2, g) as $(\overline{B}, g = e^{2u}|dz|^2)$, where $\overline{B} = \{z \in \mathbb{C}; |z| \leq 1\}$ and $|dz|^2$ is the canonical metric. Since $(\overline{B}, g = e^{2u}|dz|^2)$ has curvature 1 and the boundary is a convex circle, it can be isometrically embedded as a domain of the sphere \mathbb{S}^2 (see [20, Theorem 4]). As the boundary is totally geodesic, we conclude that $(\overline{B}, g = e^{2u}|dz|^2)$ is isometric to a hemisphere. \square

3.1. Proof of Theorem 1.1.

Proof. By assumption, we can apply Theorem 2.4 with $a \geq 6$ and $b \geq 0$. Writing $\chi(M) = 2 - 2g - r$, where g is the genus of Σ and r is the number of connected components of the boundary, we have:

$$\lambda_1^J \leq -\frac{1}{2}(H^2 + 6) + \frac{2\pi(2 - 2g - r)}{|\Sigma|},$$

and therefore,

$$-2 \leq \lambda_1^J \leq -3 + \frac{2\pi(2 - 2g - r)}{|\Sigma|} \leq -3 + \frac{2\pi}{|\Sigma|},$$

which proves our area estimate.

If equality holds, then by Theorem 2.4, Σ is totally geodesic, has constant Gaussian curvature equal to 1, and the geodesic curvature of $\partial\Sigma$ is zero. By Lemma 3.1, Σ is isometric to \mathbb{S}_+^2 and $\partial\Sigma$ is isometric to a circle of radius 1.

Furthermore, if the Gaussian curvature of ∂M is $K_{\partial M} \geq 1$ and contains geodesic which is a circle of radius 1, by Topogonov theorem [31] (see also

[20, Corollary 1]) we have that $\partial M = \mathbb{S}^2$. Now Hang-Wang theorem [20, Theorem 2] implies that M is isometric to the hemisphere \mathbb{S}_+^3 . \square

3.2. Proof of Theorem 1.2.

Proof. By Theorem 2.3, we have

$$\lambda_1^J \leq -\frac{1}{2} \left(\frac{3}{2}H^2 + a \right) + \frac{2\pi\chi(\Sigma)}{|\Sigma|}.$$

Given that $S \geq a = 0$, $H \geq 2$, and $\lambda_1^J \geq -2$, we obtain

$$(3.1) \quad -2 \leq \lambda_1^J \leq -3 + \frac{4\pi}{|\Sigma|}$$

and the estimate follows.

If equality occurs, then $H = 2$ and Σ is an immersed surface, totally umbilical with constant Gaussian curvature $K = 1$ and genus 0. Therefore, it is isometric to \mathbb{S}^2 . It follows from the Hang-Wang Theorem [19] that the region bounded by Σ is isometric to the unit Euclidean ball. \square

3.3. Proof of Theorem 1.3.

Proof. As in the previous theorem, we have

$$\lambda_1^J \leq -\frac{1}{2} \left(\frac{3}{2}H^2 + a \right) + \frac{2\pi(2-2g)}{|\Sigma|},$$

and given that $S \geq a = -6$, $H \geq 2\sqrt{2}$, and $\lambda_1^J \geq -2$, we obtain

$$(3.2) \quad -2 \leq \lambda_1^J \leq -3 + \frac{4\pi}{|\Sigma|}.$$

If equality occurs, then $H = 2\sqrt{2}$ and Σ is totally umbilical with constant Gaussian curvature $K = 1$ and genus 0. Therefore, it is isometric to \mathbb{S}^2 . It follows by Shi and Tam theorem, see [30, Theorem 3.8], that the region bounded by Σ is isometric to the unit hyperbolic ball. \square

3.4. Proof of Theorem 1.4.

Proof. Taking $\phi = 1$ in the variational characterization of λ_1^J and using the Gauss equation as given by (2.3), we obtain

$$\begin{aligned} \lambda_1^J &\leq -\frac{\int_{\Sigma} (\text{Ric}(N, N) + |A|^2) dv}{|\Sigma|} \\ &= -\frac{1}{2|\Sigma|} \int_{\Sigma} (S - S_{\Sigma} + H^2 + |A|^2) dv. \end{aligned}$$

Given the condition on the first eigenvalue and the hypothesis on the scalar curvature $S \geq n(n+1)$, we have

$$(3.3) \quad -n \leq \lambda_1 \leq -\frac{n(n+1)}{2} + \frac{1}{2} \frac{\int_{\Sigma} S_{\Sigma} dv}{|\Sigma|},$$

which proves the inequality.

The equality holds if, and only if, Σ is totally geodesic, the scalar curvature of M along Σ is constant and equal to $S|_{\Sigma} = n(n+1)$, and $\text{Ric}(N, N) = n$.

If we further assume that $K_M \geq 1$. For n even, we use Synge theorem and the orientability of Σ with the Gauss equation to deduce that Σ is isometric to the round sphere \mathbb{S}^n . For n odd and Σ simply connected, then the Gauss equation implies that Σ is also isometric to the round sphere \mathbb{S}^n . In particular, Σ is embedded.

Under the previous conclusions, we have that $M \setminus \Sigma$ consists of two connected components, say Ω_1 and Ω_2 (see [21]). Now, applying Hang-Wang theorem [20, Theorem 2] to each of these components, we conclude that $\Omega_1 = \Omega_2 = \mathbb{S}_+^{n+1}$, and the desired result follows. \square

4. THE JACOBI-STEKLOV PROBLEM

In this section we consider the Steklov problem associated with the Jacobi operator of a free boundary hypersurface Σ in M^{n+1} . More precisely, we say that $\rho \in C^\infty(\Sigma)$ is a *Steklov eigenfunction* associated with the Jacobi operator with eigenvalue σ if it solves the boundary value problem

$$(4.1) \quad \begin{cases} J\rho = 0, & \text{in } \Sigma, \\ B\rho = \sigma\rho, & \text{on } \partial\Sigma, \end{cases}$$

where J is the Jacobi operator and $B = \partial_\nu - \Pi^{\partial M}(N, N)$ is the boundary term operator in the second variation formula of the area (cf. Section 2).

Note that this is equivalent to saying that

$$\mathcal{I}(\rho, \phi) = \sigma \langle \rho, \phi \rangle_{L^2(\partial\Sigma)},$$

for all $\phi \in C^\infty(\Sigma)$.

Similarly to the eigenvalue problem for the Jacobi operator, the Steklov eigenvalues of problem (4.1) form a divergent sequence

$$\sigma_1^J \leq \sigma_2^J \leq \sigma_3^J \dots \nearrow \infty$$

and the first eigenvalue can be characterized variationally as

$$(4.2) \quad \sigma_1^J = \inf_{\phi \in C^\infty(\Sigma) \setminus \{0\}} \frac{\mathcal{I}(\phi, \phi)}{\int_{\partial\Sigma} \phi^2 da}.$$

We also have the characterization of the first eigenfunctions, analogous to what we did for the first eigenvalue of the Jacobi operator. The proof follows exactly the same scheme as Lemma 2.2, and will be omitted.

Lemma 4.1. *Suppose there exists $\rho \in C^\infty(\Sigma) \setminus \{0\}$ such that $\mathcal{I}(\rho, \rho) = \sigma_1^J \int_{\partial\Sigma} \rho^2 da$. Then, ρ is an eigenfunction of problem (4.1) associated with eigenvalue σ_1^J .*

We have the following estimate for the first eigenvalue σ_1^J :

Theorem 4.2. *Let M^3 be a Riemannian manifold with boundary such that $a = \inf S$ and $b = \inf H^{\partial M}$ are finite. Let $\Sigma^2 \subset M^3$ be a compact two-sided surface with constant mean curvature H and free boundary. Then*

$$(4.3) \quad \sigma_1^J \leq -\left(\frac{a}{2} + \frac{3}{4}H^2\right) \frac{|\Sigma|}{|\partial\Sigma|} + \frac{2\pi\chi(\Sigma)}{|\partial\Sigma|} - b.$$

Moreover, equality holds if, and only if,

- (1) Σ is totally umbilical and has constant Gaussian curvature.
- (2) $S|_\Sigma = a$, $H^{\partial M}|_{\partial\Sigma} = b$.
- (3) $\text{Ric}_M(N, N) = -\frac{1}{2}H^2$ along Σ and $\text{II}^{\partial M}(N, N) = -\sigma_1^J$ along $\partial\Sigma$.

Proof. Taking the function $\phi = 1$ in the variational characterization of σ_1^J and using the Gauss equation and Gauss-Bonnet theorem, we have

$$\begin{aligned} \sigma_1^J |\partial\Sigma| &\leq -\int_\Sigma \{(\text{Ric}_M(N, N) + |A|^2)\} dv - \int_{\partial\Sigma} \text{II}^{\partial M}(N, N) da \\ &= -\frac{1}{2} \int_\Sigma (S - 2K_\Sigma + H^2 + |A|^2) dv - \int_{\partial\Sigma} (H^{\partial M} - \kappa) da \\ &\leq -\frac{1}{2}(a + \frac{3}{2}H^2)|\Sigma| + 2\pi\chi(\Sigma) - b|\partial\Sigma|. \end{aligned}$$

If equality holds, then all the inequalities are equalities. In particular, Σ is totally umbilical, and by Lemma 4.1, the function $\phi = 1$ is an eigenfunction, and therefore satisfies the problem (4.1). Thus, we have $\text{Ric}_M(N, N) + H^2/2 = 0$ and $\sigma_1^J = -\text{II}^{\partial M}(N, N)$. The converse is immediate. \square

4.1. Proof of Theorem 1.5.

Proof. Since $\text{II}^{\partial M} \geq 1$, it follows that $b = \inf H^{\partial M} \geq 2$. Therefore, using inequality (4.3), we have

$$-1 \leq \sigma_1^J \leq \frac{2\pi\chi(\Sigma)}{|\partial\Sigma|} - 2,$$

which proves the first part.

If $|\partial\Sigma| = 2\pi$, $\chi(\Sigma) = 1$, that is, $r = 1$ and $g = 0$. Moreover, the geodesic curvature of the boundary $\partial\Sigma$ is $\kappa = \langle \nabla_T \nu, T \rangle = \text{II}^{\partial M}(T, T) = 1$, which implies that $K_\Sigma = 0$. Thus, Σ is isometric to the disk \mathbb{D} . Moreover, $\partial\Sigma$ is a simple closed geodesic of ∂M , since $\langle \nabla_T N, T \rangle = A(T, T) = 0$. We are in conditions to apply Mazet-Mendes Theorem [22, Theorem 5.1] and conclude the result. \square

4.2. Proof of Theorem 1.6.

Proof. We first observe that we can derive an estimate for σ_1^J in higher dimensions using the equation (2.3). Specifically,

$$(4.4) \quad \sigma_1^J |\partial\Sigma| \leq -\frac{1}{2} \int_\Sigma (S - S_\Sigma + H^2 + |A|^2) dv - \int_{\partial\Sigma} \text{II}^{\partial M}(N, N) da.$$

Now, recall that $H^{\partial\Sigma, \Sigma}$ denotes the mean curvature of $\partial\Sigma$ as a hypersurface of Σ . Since Σ is free-boundary, we have $H^{\partial M} = H^{\partial\Sigma, \Sigma} + \Pi^{\partial M}(N, N)$. So, inequality (1.2) follows directly from (4.4) and our assumptions.

If equality holds, then we have directly that Σ is totally geodesic, $S|_{\Sigma} = 0$, $H^{\partial M} = n$, and $\phi = 1$ is an eigenfunction. Thus, using (4.1), we have $\text{Ric}_M(N, N) = 0$ and $\Pi^{\partial M}(N, N) = 1$, and so $H^{\partial\Sigma, \Sigma} = n - 1$.

Furthermore, if $K_M \geq 0$ and $\Pi^{\partial M} \geq g|_{\partial\Sigma}$, then $K_{\Sigma} = 0$ and $\Pi^{\partial\Sigma, \Sigma} = g|_{\partial\Sigma}$. Therefore, we can use the exponential map to construct an isometry between a star-shaped domain in the tangent space and Σ . In particular, Σ and $\partial\Sigma$ are simply connected. By the Gauss equation, the sectional curvature of $\partial\Sigma$ is constant equal to one, and thus we conclude that $\partial\Sigma$ is isometric to \mathbb{S}^{n-1} and Σ is isometric to the ball \mathbb{B}^n by Xia's theorem [32].

Using again $K_M \geq 0$ and $\Pi^{\partial M} \geq g|_{\partial M}$, and the Gauss equation, we obtain that $\text{Ric}_{\partial M} \geq ng$. Given that Σ is a free boundary hypersurface and the previous conclusions, we have that $\partial\Sigma$ in ∂M is a totally geodesic round sphere, which divides ∂M into two components Ω_+ and Ω_- that are isometric to \mathbb{S}_+^n by the Hang-Wang theorem (see [20, Theorem 2]). Therefore, ∂M is isometric to \mathbb{S}^n and, applying again Xia's theorem, we obtain the desired result. \square

Remark 4.3. We observe that we can also apply the Mazet-Mendes theorem [22] to conclude the rigidity in the theorem above.

Remark 4.4. It is an interesting problem to obtain the rigidity part in Theorem 1.6 under a condition on the Ricci curvature.

5. ESTIMATES INVOLVING THE YAMABE INVARIANTS

It is noteworthy that the first eigenvalue of the Jacobi operator, λ_1^J , and the first eigenvalue of the Jacobi-Steklov problem, σ_1^J , for hypersurfaces with boundary are related to the Yamabe invariants. These geometric invariants were introduced by Escobar in [14] and [15], and play a crucial role in solving the Yamabe Problem with boundary.

Let (Σ^n, g) be a compact Riemannian manifold with boundary $\partial\Sigma$ and dimension $n \geq 3$. Following Escobar [15], we define the functional

$$Q_g(\varphi) = \frac{\int_{\Sigma} (a_n |\nabla \varphi|^2 + S_{\Sigma} \varphi^2) dv + 2 \int_{\partial\Sigma} H^{\partial\Sigma, \Sigma} \varphi^2 da}{\left(\int_{\Sigma} \varphi^{\frac{2n}{n-2}} dv \right)^{\frac{n-2}{n}}},$$

where φ is a non-zero smooth function on Σ , and $a_n = \frac{4(n-1)}{n-2}$.

The first *Yamabe constant for manifolds with boundary* is then defined as

$$Q_g(\Sigma) = \inf_{\varphi \in C^{\infty}(\Sigma) \setminus \{0\}} Q_g(\varphi).$$

It is not difficult to see that $Q_g(\Sigma)$ is invariant under conformal changes, and therefore, depends only on the conformal class $[g]$ of g (cf. [15]). Moreover, we have that $Q_g(\Sigma) \leq Q_g(\mathbb{S}_+^n)$.

The first *Yamabe invariant* of Σ is the supremum of the Yamabe constants over all conformal classes. That is,

$$\sigma(\Sigma) = \sup_{[g] \in \mathcal{C}(\Sigma)} Q_g(\Sigma),$$

where $\mathcal{C}(\Sigma)$ is the space of all conformal classes of Σ . Note that, by definition, $\sigma(\Sigma)$ is a differentiable invariant of Σ .

Armed with these new invariants, we have the following result, which can be viewed as a boundary version of [8, Theorem 6] due to Cai and Galloway.

Theorem 5.1. *Let (M^{n+1}, g) be an orientable Riemannian manifold with boundary, $n \geq 3$, with scalar curvature $S \geq a$, and mean-convex boundary, $H^{\partial M} \geq 0$. Let $\Sigma^n \subset M^{n+1}$ be a compact, orientable hypersurface with constant mean curvature H and free boundary.*

If $\sigma(\Sigma) < 0$, then $2\lambda_1^J + a + H^2 < 0$ and

$$(5.1) \quad |\Sigma| \geq \left(\frac{\sigma(\Sigma)}{2\lambda_1^J + a + H^2} \right)^{\frac{n}{2}}.$$

Moreover, the equality in (5.1) holds if and only if $\sigma(\Sigma) = Q_g(1)$, Σ is totally geodesic in M , $S|_{\Sigma} = a$, $H^{\partial M}|_{\partial\Sigma} = 0$, $\Pi^{\partial M}(N, N)|_{\partial\Sigma} = 0$, and $\text{Ric}_M(N, N) = -\lambda_1^J$. In particular, $S_{\Sigma} = a + 2\lambda_1^J < 0$, $H^{\partial\Sigma, \Sigma} = 0$, and $(\Sigma, g|_{\Sigma})$ is Einstein with totally geodesic boundary.

Proof. Using the variational characterization of λ_1^J given in (2.2), and noting that $a_n = \frac{4(n-1)}{n-2} > 2$, and using our assumptions about the curvatures, we have that

$$\begin{aligned} 2\lambda_1^J \int_{\Sigma} \varphi^2 dv &\leq 2 \int_{\Sigma} \{ |\nabla \varphi|^2 - (\text{Ric}_M(N, N) + |A|^2) \varphi^2 \} dv \\ &\quad - 2 \int_{\partial\Sigma} \Pi^{\partial M}(N, N) \varphi^2 da \\ &\leq \int_{\Sigma} \{ a_n |\nabla \varphi|^2 - (S - S_{\Sigma} + H^2 + |A|^2) \varphi^2 \} dv \\ &\quad - 2 \int_{\partial\Sigma} (H^{\partial M} - H^{\partial\Sigma, \Sigma}) \varphi^2 da \\ &\leq \int_{\Sigma} (a_n |\nabla \varphi|^2 + S_{\Sigma}) \varphi^2 dv - \int_{\Sigma} (a + H^2) \varphi^2 dv + 2 \int_{\partial\Sigma} H^{\partial\Sigma, \Sigma} \varphi^2 da. \end{aligned}$$

That is,

$$\begin{aligned}
(2\lambda_1^J + a + H^2) \int_{\Sigma} \varphi^2 dv &\leq \int_{\Sigma} (a_n |\nabla \varphi|^2 + S_{\Sigma}) \varphi^2 dv + 2 \int_{\partial \Sigma} H^{\partial \Sigma, \Sigma} \varphi^2 da \\
&= Q_g(\varphi) \left(\int_{\Sigma} \varphi^{\frac{2n}{n-2}} dv \right)^{\frac{n-2}{n}}.
\end{aligned}$$

Using that $Q_g(\Sigma) < 0$, we can take a sequence of smooth function $\{\varphi_l\}$ such that $Q_g(\varphi_l) < 0$ and $Q_g(\varphi_l) \rightarrow Q_g(\Sigma)$, when $l \rightarrow \infty$. In particular, $2\lambda_1^J + a + H^2 < 0$. On the other hand, using the Hölder inequality we have

$$\int_{\Sigma} \varphi_l^2 dv \leq |\Sigma|^{\frac{2}{n}} \left(\int_{\Sigma} \varphi_l^{\frac{2n}{n-2}} dv \right)^{\frac{n-2}{n}},$$

and using again $Q_g(\varphi_l) < 0$, we obtain

$$(2\lambda_1^J + a + H^2) |\Sigma|^{\frac{2}{n}} \leq Q_g(\varphi_l).$$

Making $l \rightarrow \infty$ and using the definition of $\sigma(\Sigma)$ we conclude the estimate.

If equality holds, all inequalities become equalities, and in particular, φ is constant. The remaining assertions follow straightforwardly. The converse is immediate.

In particular, $S_{\Sigma} = S|_{\Sigma} - 2\text{Ric}_M(N, N) = a + 2\lambda_1^J$, using the free boundary condition we have $H^{\partial \Sigma, \Sigma} = H^{\partial M}|_{\partial \Sigma} - \text{II}^{\partial M}(N, N) = 0$, and arguing as in the proof of Theorem 2.1 of [5] we conclude that $(\Sigma, g|_{\Sigma})$ is Einstein with totally geodesic boundary. \square

Escobar also introduced another invariant (see [14]) which we now describe. Consider the functional

$$(5.2) \quad Y(\varphi) = \frac{\int_{\Sigma} (a_n |\nabla \varphi|^2 + S_{\Sigma} \varphi^2) dv + 2 \int_{\partial \Sigma} H^{\partial \Sigma, \Sigma} \varphi^2 da}{\left(\int_{\partial \Sigma} \varphi^{\frac{2(n-1)}{n-2}} da \right)^{\frac{n-2}{n-1}}},$$

where φ is a smooth function not vanishing on the boundary.

The second *Yamabe constant for manifolds with boundary* is then defined as

$$Y(\Sigma, \partial \Sigma) = \inf_{\varphi \in C^{\infty}(\Sigma) \setminus \{0\}} Y(\varphi).$$

It is not difficult to see that $Y(\Sigma, \partial \Sigma)$ is invariant under conformal changes, and thus depends only on the conformal class $[g]$ of g (cf. [14]). Moreover, we have that $Y(\Sigma, \partial \Sigma) \leq Y(\mathbb{B}^n, \partial \mathbb{B}^n)$.

In this setting, the second *Yamabe invariant of Σ* is the supremum of these Yamabe constants over all conformal classes:

$$\tau(\Sigma, \partial \Sigma) = \sup_{[g] \in \mathcal{C}(\Sigma)} Y(\Sigma, \partial \Sigma).$$

Using this invariant we obtain the following estimate.

Theorem 5.2. *Let (M^{n+1}, g) be an orientable Riemannian manifold with boundary, $n \geq 3$, with scalar curvature $S \geq 0$, and $H^{\partial M} \geq b$. Let $\Sigma^n \subset M^{n+1}$ be a compact, orientable hypersurface with constant mean curvature H and free boundary. Assume that $\tau(\Sigma, \partial\Sigma) < 0$, then $\sigma_1^J + b < 0$ and*

$$(5.3) \quad |\partial\Sigma| \geq \left(\frac{\tau(\Sigma, \partial\Sigma)}{2(\sigma_1^J + b)} \right)^{n-1}.$$

Moreover, the equality in (5.3) holds if and only if $\tau(\Sigma, \partial\Sigma) = Y(1)$, Σ is totally geodesic in M , $S|_{\Sigma} = 0$, $H^{\partial M}|_{\partial\Sigma} = b$, $\Pi^{\partial M}(N, N)|_{\partial\Sigma} = -\sigma_1^J$, and $\text{Ric}_M(N, N) = 0$. In particular, $S_{\Sigma} = 0$, $H^{\partial\Sigma, \Sigma} = b + \sigma_1^J$, and $(\Sigma, g|_{\Sigma})$ is Ricci-flat with totally umbilical boundary.

Proof. Using Holder inequality and the hypothesis on the Yamabe invariant we obtain:

$$(5.4) \quad Y(\varphi) \int_{\partial\Sigma} \varphi^2 da \geq Y(\varphi) |\partial\Sigma|^{\frac{1}{n-1}} \left(\int_{\partial\Sigma} \varphi^{\frac{2(n-1)}{n-2}} da \right)^{\frac{n-2}{n-1}}.$$

By (4.2), we have:

$$\begin{aligned} 2\sigma_1^J \int_{\partial\Sigma} \varphi^2 da &\leq 2 \int_{\Sigma} \{ |\nabla\varphi|^2 - (\text{Ric}(N, N) + |A|^2) \varphi^2 \} dv \\ &\quad - 2 \int_{\partial\Sigma} \Pi^{\partial M}(N, N) \varphi^2 da \\ &\leq \int_{\Sigma} \{ a_n |\nabla\varphi|^2 - (S - S_{\Sigma} + H^2 + |A|^2) \varphi^2 \} dv \\ &\quad - 2 \int_{\partial\Sigma} (H^{\partial M} - H^{\partial\Sigma, \Sigma}) \varphi^2 da, \end{aligned}$$

where we used $a_n = \frac{4(n-1)}{n-2} > 2$. Thus,

$$\begin{aligned} (2\sigma_1^J + 2b) \int_{\partial\Sigma} \varphi^2 da &\leq \int_{\Sigma} [a_n |\nabla\varphi|^2 + S_{\Sigma} \varphi^2] dv + 2 \int_{\partial\Sigma} H^{\partial\Sigma, \Sigma} da \\ &= Y(\varphi) \left(\int_{\partial\Sigma} \varphi^{\frac{2(n-1)}{n-2}} da \right)^{\frac{n-2}{n-1}} \end{aligned}$$

By (5.4) we have

$$2(\sigma_1 + b) |\partial\Sigma|^{\frac{1}{n-1}} \leq Y(\varphi).$$

Using (5.2) and definitions after that, we get the estimate (5.3).

The analysis of the equality case is similar to the previous case. \square

Acknowledgments. The authors are grateful to Abraão Mendes for interesting discussions and suggestions on this paper. The authors were partially supported by the Brazilian National Council for Scientific and Technological Development - CNPq and the Foundation for Research Support of the State of Alagoas - FAPEAL. The first author was supported by

CNPq [Grants 405468/2021-0, 308440/2021-8 and 402463/2023-9] and FAPEAL [Grant 60030.0000001758/2022]. The second author was supported by CNPq [Grants 405468/2021-0 and 11136/2023-0] and FAPEAL [Grant 60030.0000000323/2023]. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

REFERENCES

1. Luis J. Alías, Abdénago Barros, and Aldir Brasil, Jr., *A spectral characterization of the $H(r)$ -torus by the first stability eigenvalue*, Proc. Amer. Math. Soc. **133** (2005), no. 3, 875–884. MR 2113939
2. Luis J. Alías, Miguel A. Meroño, and Irene Ortiz, *On the first stability eigenvalue of constant mean curvature surfaces into homogeneous 3-manifolds*, Mediterr. J. Math. **12** (2015), no. 1, 147–158. MR 3306032
3. Lucas Ambrozio, Alessandro Carlotto, and Ben Sharp, *Index estimates for free boundary minimal hypersurfaces*, Math. Ann. **370** (2018), no. 3-4, 1063–1078.
4. Lucas C. Ambrozio, *Rigidity of area-minimizing free boundary surfaces in mean convex three-manifolds*, J. Geom. Anal. **25** (2015), no. 2, 1001–1017.
5. Henrique Araujo, *Critical points of the total scalar curvature plus total mean curvature functional*, Indiana Univ. Math. J. **52** (2003), 85–108.
6. Márcio Batista and José I. Santos, *Upper bounds for the first stability eigenvalue of surfaces in 3-Riemannian manifolds*, Potential Anal. **49** (2018), no. 1, 91–103.
7. Hubert Bray, Simon Brendle, and Andre Neves, *Rigidity of area-minimizing two-spheres in three-manifolds*, Comm. Anal. Geom. **18** (2010), no. 4, 821–830. MR 2765731
8. Mingliang Cai and Gregory J. Galloway, *On the topology and area of higher-dimensional black holes*, Classical Quantum Gravity **18** (2001), no. 14, 2707–2718.
9. Marcos P. Cavalcante and Darlan F. de Oliveira, *Index estimates for free boundary constant mean curvature surfaces*, Pac. J. Math. **305** (2020), no. 1, 153–163.
10. Daguang Chen and Qing-Ming Cheng, *Estimates for the first eigenvalue of Jacobi operator on hypersurfaces with constant mean curvature in spheres*, Calc. Var. Partial Differential Equations **56** (2017), no. 2, Paper No. 50, 12. MR 3626321
11. Qing-Ming Cheng, *First eigenvalue of a Jacobi operator of hypersurfaces with a constant scalar curvature*, Proc. Amer. Math. Soc. **136** (2008), no. 9, 3309–3318. MR 2407097
12. Antonio W. Cunha, Henrique F. de Lima, and Fábio R. dos Santos, *On the first stability eigenvalue of closed submanifolds in the Euclidean and hyperbolic spaces*, Differential Geom. Appl. **52** (2017), 11–19. MR 3656418
13. Nguyen Thac Dung, Juncheol Pyo, and Hung Tran, *First stability eigenvalue of singular hypersurfaces with constant mean curvature in the unit sphere*, Proc. Amer. Math. Soc. **151** (2023), no. 2, 795–810. MR 4520028
14. José F. Escobar, *Conformal deformation of a Riemannian metric to a scalar flat metric with constant mean curvature on the boundary*, Ann. Math. (2) **136** (1992), no. 1, 1–50.
15. ———, *The Yamabe problem on manifolds with boundary*, J. Differential Geom. **35** (1992), no. 1, 21–84. MR 1152225
16. ———, *The geometry of the first non-zero Stekloff eigenvalue*, J. Funct. Anal. **150** (1997), no. 2, 544–556. MR 1479552
17. ———, *An isoperimetric inequality and the first Steklov eigenvalue*, J. Funct. Anal. **165** (1999), no. 1, 101–116. MR 1696453

18. José M. Espinar, *Gradient Schrödinger operators, manifolds with density and applications*, J. Math. Anal. Appl. **455** (2017), no. 2, 1505–1528.
19. Fengbo Hang and Xiadong Wang, *Rigidity and non-rigidity results on the sphere*, Commun. Anal. Geom. **14** (2006), no. 1, 91–106.
20. Fengbo Hang and Xiaodong Wang, *Rigidity theorems for compact manifolds with boundary and positive Ricci curvature*, J. Geom. Anal. **19** (2009), no. 3, 628–642.
21. H. Blaine Lawson, Jr., *The unknottedness of minimal embeddings*, Invent. Math. **11** (1970), 183–187. MR 287447
22. Laurent Mazet and Abraão Mendes, *Rigidity of min-max minimal disks in 3-balls with non-negative Ricci curvature*, arXiv 2307.03624 (2023), 1–27.
23. Abraão Mendes, *Rigidity of free boundary surfaces in compact 3-manifolds with strictly convex boundary*, J. Geom. Anal. **28** (2018), no. 2, 1245–1257.
24. Miguel A. Meroño and Irene Ortiz, *First stability eigenvalue characterization of CMC Hopf tori into Riemannian Killing submersions*, J. Math. Anal. Appl. **417** (2014), no. 1, 400–410. MR 3191435
25. Takashi Okayasu, *Area of minimal disks with free boundary*, Geometry and global analysis. Report of the 1st MSJ (Mathematical Society of Japan) International Research Institute, Sendai, Japan, July 12–23, 1993, Sendai: Tôhoku Univ., Mathematical Institute, 1993, pp. 447–449.
26. Oscar Perdomo, *First stability eigenvalue characterization of Clifford hypersurfaces*, Proc. Am. Math. Soc. **130** (2002), no. 11, 3379–3384.
27. Antonio Ros and Enaldo Vergasta, *Stability for hypersurfaces of constant mean curvature with free boundary*, Geom. Dedicata **56** (1995), no. 1, 19–33. MR 1338315
28. Pam Sargent, *Index bounds for free boundary minimal surfaces of convex bodies*, Proc. Am. Math. Soc. **145** (2017), no. 6, 2467–2480.
29. Richard Schoen and Shing-Tung Yau, *Existence of incompressible minimal surfaces and the topology of three dimensional manifolds with non-negative scalar curvature*, Ann. Math. (2) **110** (1979), 127–142.
30. Yuguang Shi and Luen-Fai Tam, *Rigidity of compact manifolds and positivity of quasi-local mass*, Classical Quantum Gravity **24** (2007), no. 9, 2357–2366.
31. V. A. Toponogov, *Evaluation of the length of a closed geodesic on a convex surface*, Dokl. Akad. Nauk SSSR **124** (1959), 282–284 (Russian).
32. Changyu Xia, *Rigidity of compact manifolds with boundary and nonnegative Ricci curvature*, Proc. Am. Math. Soc. **125** (1997), no. 6, 1801–1806.
33. Jonathan J. Zhu, *First stability eigenvalue of singular minimal hypersurfaces in spheres*, Calc. Var. Partial Differ. Equ. **57** (2018), no. 5, 13, Id/No 130.

CPMAT-IM, UNIVERSIDADE FEDERAL DE ALAGOAS, MACEIÓ - BRAZIL

Email address: mhbs@mat.ufal.br, marcos@pos.mat.ufal.br, luiz.melo@im.ufal.br