

Impacts of ALP on the Constraints of Dark Photon

Chuan-Ren Chen¹, Yuan-Feng Hsieh², and Chrisna Setyo Nugroho³

Department of Physics, National Taiwan Normal University, Taipei 116, Taiwan

Abstract

Dark sector may exist and interact with Standard Model (SM) through the $U(1)$ kinetic mixing. Through this portal-type interaction, dark photon from dark sector couples to SM fermions, and may explain the discrepancy between experimental data and SM calculations on muon anomalous magnetic moment, muon $g - 2$. However, current searches for dark photon impose stringent constraints on the mixing parameter ε for various dark photon masses, excluding the favorite parameter space for muon $g - 2$. In this paper, we study the case where a global $U(1)$ in dark sector is spontaneously broken, resulting a light pseudo-Goldstone, axion-like particle (ALP) a , which couples to dark photon and SM photon, $g_{a\gamma\gamma'}$. Through this interaction, dark photon may decay into photon and ALP when this channel is kinematically allowed. As a result, the experimental constraints on dark photon change significantly, and dark photon is able to explain the muon $g - 2$ anomaly when its mass is heavier than 10 GeV.

arXiv:2405.19087v1 [hep-ph] 29 May 2024

¹crchen@ntnu.edu.tw

²61041026S@ntnu.edu.tw

³setyo13nugros@ntnu.edu.tw

1 Introduction

Besides the discovery of Higgs boson, the Large Hadron Collider (LHC) is built for new physics signals. However, the null results after Run II at the LHC indicate that new physics may interact extremely weak with Standard Model (SM). The existence of dark sector that interacts with SM through portals naturally realizes this possibility. Depending on the structure of dark sector, the portal can be realized via scalars, sterile neutrinos or vectors. The simplest mechanism for vector portal is given by a kinetic mixing term between two $U(1)$ gauged symmetries from dark sector and SM, respectively. As a result, a new gauge boson γ' , referred to dark photon, interacts with the SM electromagnetic current J_{em}^μ as $\varepsilon A'_\mu J_{em}^\mu$, where A'_μ is the dark photon field and ε is the mixing parameter.

Several proposals have been made to detect dark photon, ranging from laboratory experiments to astronomical observations [1–3], and the ε is constrained to be tiny and less than about 10^{-3} for dark photon mass below about 100 GeV. On the other hand, the interactions between SM fermions and dark photon introduce additional one-loop diagrams to anomalous magnetic moments ($g - 2$). The long-standing tension between SM calculation and experimental measurements by BNL [4] and FermiLab [5] on muon $g - 2$ reaches 5σ level. It has been shown that such anomaly can be explained by a sub-GeV γ' [6]. However, the collider searches of γ' exclude this parameter space favored by muon $g - 2$.

Note that the constraints from collider searches mainly come from the null excesses on signals of e^+e^- or $\mu^+\mu^-$ pairs from γ' decays. If the structure of dark sector allows additional decay channels other than SM fermion final states, the total event rates of e^+e^- and $\mu^+\mu^-$ would be significantly changed. In this paper, we propose the existence of a pseudo-Goldstone boson when a global $U(1)$ in dark sector is broken, which is referred to axion-like particle (ALP). Similar to the QCD axion coupling to two photons, ALP interacts with dark photons and photons [7]. As a result, dark photon will decay into ALP and photon if ALP is lighter than dark photon¹. We will see later that the decay channel of $\gamma' \rightarrow a\gamma$, where a and γ are ALP and photon, respectively, could dominate the the decay channels for γ' heavier about 100 MeV and rescue the excluded parameter space for muon $g - 2$.

The paper is organized as follows: In section 2, we discuss the model containing the SM and dark sector with dark photon acting as a messenger between them. We provide the current experimental limits on dark photon coupling in Section 3. We recast the recent experimental constraints in the presence of new interaction in Section 4 and further discuss its consequences there. Our summary is presented in Section 5.

2 The Model

We focus on the dark sector where both global and gauged $U(1)$ symmetries are implemented. After the symmetries are spontaneously broken, a pseudo-Nambu-Goldstone boson dubbed axion-like particle (ALP), and a gauge boson dubbed dark photon are generated [7, 9]. The relevant Lagrangian are given as [7]

$$\mathcal{L} \supset \frac{1}{2} m_{\gamma'}^2 A'_\mu A'^\mu + \frac{1}{2} \varepsilon F_{\mu\nu} F'^{\mu\nu} - \frac{1}{2} g_{a\gamma\gamma'} a F_{\mu\nu} \tilde{F}'^{\mu\nu}, \quad (2.1)$$

¹For extremely light ALP, it could decay into two photons which is constrained by the survey of infrared photon spectrum by JWST [8].

where A'_μ is dark photon field, $F_{\mu\nu}$ and $F'_{\mu\nu}$ are field strength of photon and dark photon, respectively; $\tilde{F}'_{\mu\nu}$ is the dual tensor of $F'_{\mu\nu}$, while a represents the ALP. Because of the mixing term of photon and dark photon, the couplings between dark photon and fermions of the SM are introduced. As a result, the dark photons can be produced in the QED processes by simply replacing non-propagating photons by dark photons. Therefore, the production mechanism includes bremsstrahlung, annihilation, meson decays and Drell-Yan process. Regarding its decay, the dark photon γ' will decay into SM fermions if its mass is heavy enough. The partial decay width of dark photon into a pair of charged leptons ℓ is

$$\Gamma(\gamma' \rightarrow \ell^+\ell^-) = \frac{\varepsilon^2 e^2}{12\pi} m_{\gamma'} \left(1 + 2 \frac{m_\ell^2}{m_{\gamma'}^2}\right) \sqrt{1 - 4 \frac{m_\ell^2}{m_{\gamma'}^2}}, \quad (2.2)$$

where e and m_ℓ are the electric charge and mass of charged lepton ℓ . For quark channels of dark photon decay, the final state is represented by the hadrons. The partial decay width is therefore presented as

$$\Gamma(\gamma' \rightarrow \text{hadrons}) = \Gamma(\gamma' \rightarrow \mu^+\mu^-) R_t, \quad (2.3)$$

where R_t is defined as the ratio $R_t \equiv \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$, with σ being the cross section at the e^+e^- collider at a certain energy. For the numerical results shown later, we read out the values of R_t from PDG [10]. In addition, the dark photon may decay into an ALP and a photon due to the interaction $-\frac{1}{2}g_{a\gamma\gamma'}aF_{\mu\nu}\tilde{F}'^{\mu\nu}$ if it is kinematically allowed. In this case, we obtain the partial decay width as

$$\Gamma(\gamma' \rightarrow a\gamma) = \frac{g_{a\gamma\gamma'}^2}{96\pi} m_{\gamma'}^3 \left(1 - \frac{m_a^2}{m_{\gamma'}^2}\right)^3, \quad (2.4)$$

where m_a is the mass of ALP. Through this paper, we focus on the scenario where $m_{\gamma'} \gg m_a$, leading to a simple result $\Gamma(\gamma' \rightarrow a\gamma) \propto m_{\gamma'}^3$. The size of photon-dark-photon mixing parameter ε and strength of $g_{a\gamma\gamma'}$ depend on the details of the UV setup of the model. Here, we treat them as parameters for simplicity in our phenomenology study. We notice that the constraints on ε has been studied in details in the literature, see e.g. [2] and references therein, if the $g_{a\gamma\gamma'} \ll \varepsilon$. However, when $g_{a\gamma\gamma'}$ is of the same order of ε , the current constraints on ε change significantly and searching strategy of dark photon should be revisited, as we will discuss later in Section 4.

Fig. 1 shows the decay branching ratios of a dark photon and the ratio of branching ratio $\gamma' \rightarrow e^+e^-$ to branching ratio $\gamma' \rightarrow a\gamma$ for its mass $m_{\gamma'} > 1$ MeV, where benchmark values $\varepsilon = 10^{-2}$ (left) and $\varepsilon = 10^{-3}$ (right) are taken, while $g_{a\gamma\gamma'}$ is fixed to be 10^{-2} GeV $^{-1}$. When the value of $g_{a\gamma\gamma'}$ is as large as the mixing parameter ε , see the left panel, dark photon will mainly decay into γ and ALP when $m_{\gamma'}$ is larger than 1 GeV, while hadronic modes are subdominant. However, for a light dark photon lighter than about 200 MeV, e^+e^- mode is the main channel. The branching ratio of $\gamma' \rightarrow \mu^+\mu^-$ is also the same as e^+e^- mode and becomes the main decay channel up to $m_{\gamma'} \simeq 700$ MeV. For the window of dark photon mass between 700 MeV and 1 GeV, hadronic decay modes of dark photon is the leading channel. For $\varepsilon = 10^{-3}$ and $g_{a\gamma\gamma'} = 10^{-2}$ GeV $^{-1}$, e^+e^- mode is the dominant decay channel till $m_{\gamma'} \simeq 100$ MeV. Beyond 100 MeV, the $a\gamma$ channel takes over.

It is worth mentioning that the product of $\varepsilon g_{a\gamma\gamma'}$ can be bounded using the light shining through the wall (LSW) experiments [11–13] in the case of very light dark photon such that the decay channel to e^+e^- is absent. The reason can be easily understood since the converting




Figure 1: The decay branching ratios of dark photon. Two benchmark values, 10^{-2} and 10^{-3} are taken for mixing parameter ϵ , while effective dark photon-ALP-photon coupling $g_{a\gamma\gamma'}$ is fixed to be 10^{-2} GeV^{-1} .

probability of photon to dark photon is proportional to ϵ^2 while the decay probability inside the detector is approximately proportional to decay width which is proportional to $g_{a\gamma\gamma'}^2$. The conclusion is that, as the dark photon is of order of $O(10^4)$ eV, $\epsilon g_{a\gamma\gamma'}$ needs to be smaller than about 10^{-7} GeV^{-1} . The limit becomes weaker when mass of dark photon is even lighter. For $m_{\gamma'} \lesssim 10^{-1} \text{ eV}$, $\epsilon g_{a\gamma\gamma'}$ can be as large as 10^{-3} GeV^{-1} . Note that since we focus the case that $m_{\gamma'}$ is heavier than 1 MeV, the LSW experiment has no sensitivity on dark photon search. For the coupling strength $g_{a\gamma\gamma'}$ along, B-factories, e.g. BaBar, can impose stringent constraints through the search for mono-photon signal [14]. Due to limitation of the electron-beam energy, B-factories can only examine dark photon up to 10 GeV, and it is shown that the region of $g_{a\gamma\gamma'} \gtrsim 3 \times 10^{-3}$ is excluded [15].

3 Experimental Limits

Many experiments put efforts to search for dark photon, including colliders, fixed targets, and beam dumps. We focus on the one that searches for signals of electron and positron pairs from dark photon decay, since it is an inevitable decay channel of dark photon. We don't include invisible decay channel because this channel depends on dark matter assignment in dark sector. A summary of the current constraints on ϵ - $m_{\gamma'}$ plane is displayed in Fig. 2 which is adopted from Fig. 4 of [16].

For dark photon mass considered in this paper, $1 \text{ MeV} < m_{\gamma'} < 100 \text{ GeV}$, several experiments has been carried out to probe dark photon in the laboratories. In collider frontier, Babar collaboration puts stringent limit on dark photon parameter space based on visible search of γ' decay into e^+e^- and $\mu^+\mu^-$ [14]. Furthermore, NA48/2 experiment has performed dark photon search from pion decay $\pi^0 \rightarrow \gamma'\gamma$ followed by $\gamma' \rightarrow e^+e^-$ [17]. In addition, A1 and APEX collaborations employ electron bremsstrahlung in fixed target experiment $eZ \rightarrow eZ\gamma'$ with e^+e^- as final state of A' decay [18, 19]. Moreover, KLOE experiment has been looking for dark photon from $\phi \rightarrow \gamma'\eta$ followed by $\gamma' \rightarrow e^+e^-$ as well as $e^+e^- \rightarrow \gamma'\gamma$ with $\gamma' \rightarrow \pi^+\pi^-$ decay mode [20, 21]. Finally, the LHCb collaboration have performed dark photon search




Figure 2: The current constraints of mixing parameter ε for $m_{\gamma'}$ in the region of 1 MeV to 100 GeV. The shaded area is excluded [16]. The parameter space excluded by electron anomalous magnetic moment measurement (Δa_e) and the one favored by muon (Δa_μ) are indicated by gray region and red band, respectively.

using $\gamma' \rightarrow \mu^+ \mu^-$ decay mode from $pp \rightarrow \gamma'$ production channel [22].

On the other hand, beam dump experiments has put stringent limits on dark photon search, especially in sub-GeV dark photon mass. In both electron and proton beam dumps, dark photon search is inferred from its decay to e^+e^- final state. For electron beam dump, several experiments such as Orsay, E774, E137, E141, KEK, and NA64 has placed strict limits on dark photon [23–28]. Moreover, proton beam dump experiments like PS191, NOMAD, CHARM, ν -CAL I also exclude dark photon parameter space [29–33]. Furthermore, anomalous magnetic moment of both electron and muon put additional limits on dark photon parameter space [6, 34–37]. All these limits are presented in Fig.2 with exception of A1, APEX, and KLOE limits since they are already covered by BaBar experiment.

4 New Limits on Photon-Dark Photon Mixing

In all experiments mentioned above, the searches of dark photon focus on detecting electron and positron pairs from dark photon decay. The number of event (N) of these pairs in dark photon experiments is described by

$$N = \mathcal{L} \cdot \sigma_{\text{Prod}}(\gamma') \cdot \text{BR}(\gamma') \cdot \eta(\tau_{\gamma'}) . \quad (4.1)$$

Here, \mathcal{L} , $\sigma_{\text{Prod}}(\gamma')$, $\text{BR}(\gamma')$, and $\eta(\tau_{\gamma'})$ stand for the integrated luminosity, dark photon production cross section, branching ratio of dark photon decay into e^+e^- , and detector efficiency, respectively. In the presence of new photon-dark photon-axion interaction given by the last term of Eq.(2.1), one expects that, for $m_{\gamma'} > m_a$ considered in this work, dark photon could decay into the axion and photon. As a result, the e^+e^- branching ratio and lifetime of γ' will be different while leaving its production cross section intact. Therefore, limit on photon-dark photon mixing would be modified and should be revisited.

In term of the observed event number of e^+e^- , one have the relation

$$\mathcal{L}^{\text{New}} \cdot \sigma_{\text{Prod}}^{\text{New}}(\gamma') \cdot \text{BR}^{\text{New}}(\gamma') \cdot \eta^{\text{New}}(\tau_{\gamma'}) = \mathcal{L}^{\text{Old}} \cdot \sigma_{\text{Prod}}^{\text{Old}}(\gamma') \cdot \text{BR}^{\text{Old}}(\gamma') \cdot \eta^{\text{Old}}(\tau_{\gamma'}) , \quad (4.2)$$

where the superscript New (Old) denotes the photon-dark photon interaction with (without) axion. Since the integrated luminosity in an experiment is independent of the interaction in dark sector, one obtains $\mathcal{L}^{\text{New}} = \mathcal{L}^{\text{Old}}$. Furthermore, for $m_{\gamma'} > m_a$, the ratio of dark photon production cross section is

$$\frac{\sigma_{\text{Prod}}^{\text{New}}(\gamma')}{\sigma_{\text{Prod}}^{\text{Old}}(\gamma')} \approx \frac{\varepsilon_{\text{New}}^2}{\varepsilon_{\text{Old}}^2}, \quad (4.3)$$

where we have used the fact that the dark photon production cross section scales as ε^2 at colliders, and the dark photon from mesons decays is much less than the dominant decay channels. Thus, we can rewrite the number of event equation as

$$\varepsilon_{\text{New}}^2 \cdot \text{BR}^{\text{New}}(\gamma') \cdot \eta^{\text{New}}(\tau_{\gamma'}) \approx \varepsilon_{\text{Old}}^2 \cdot \text{BR}^{\text{Old}}(\gamma') \cdot \eta^{\text{Old}}(\tau_{\gamma'}). \quad (4.4)$$

In addition, the explicit expression of the new branching ratio $\text{BR}^{\text{New}}(\gamma')$ and the old one $\text{BR}^{\text{Old}}(\gamma')$ is given by

$$\text{BR}^{\text{New}}(\gamma') = \text{BR}^{\text{Old}}(\gamma') \left(1 + \frac{\Gamma(\gamma' \rightarrow a\gamma)}{\Gamma_{\text{tot}}^{\text{Old}}} \right)^{-1}, \quad (4.5)$$

where $\Gamma_{\text{tot}}^{\text{Old}}$ stands for the total decay width of dark photon without including ALP and $\Gamma(\gamma' \rightarrow a\gamma)$ is the partial decay width of dark photon decay into ALP plus a photon.

Finally, the detection efficiency η depends on the lifetime of dark photon, and varies according to the details of the experimental setups. The visible search for dark photon from BaBar relies on the prompt decay of dark photon into e^+e^- and $\mu^+\mu^-$ after being produced. Therefore, $\eta^{\text{New}}(\tau_{\gamma'})$ should be equal to $\eta^{\text{Old}}(\tau_{\gamma'})$. For NA48/2 and LHCb, one needs to consider the efficiency ratio [16]

$$\frac{\eta^{\text{New}}(\tau_{\gamma'})}{\eta^{\text{Old}}(\tau_{\gamma'})} \approx 1 - e^{-\tilde{t}/\tau_{\gamma'}^{\text{New}}}, \quad (4.6)$$

where \tilde{t} is the largest proper decay time that satisfied the experimental requirement. In beam dump experiments, the situation is more complicated. Following the analysis in [16], it is possible to construct the effective proper-time in a given decay region $[\tilde{t}_0, \tilde{t}_1]$. In this case, \tilde{t}_0 is the time dark photon spend inside the beam dump shield and \tilde{t}_1 denote the required time for dark photon to reach the detector from its production point. These two quantities are related by

$$\tilde{t}_1 = \tilde{t}_0 \left(1 + \frac{L_{\text{dec}}}{L_{\text{sh}}} \right), \quad (4.7)$$

where L_{dec} and L_{sh} stand for the length of decay volume and the length of the shield, respectively. Therefore, the corresponding efficiency ratio becomes

$$\frac{\eta^{\text{New}}(\tau_{\gamma'})}{\eta^{\text{Old}}(\tau_{\gamma'})} = \frac{e^{-\tilde{t}_0/\tau_{\gamma'}^{\text{New}}} - e^{-\tilde{t}_1/\tau_{\gamma'}^{\text{New}}}}{e^{-\tilde{t}_0/\tau_{\gamma'}^{\text{Old}}} - e^{-\tilde{t}_1/\tau_{\gamma'}^{\text{Old}}}}. \quad (4.8)$$

To obtain the values of \tilde{t}_0 at each dark photon mass $m_{\gamma'}$ inside the interval limits of $[\varepsilon_{\text{min}}, \varepsilon_{\text{max}}]$, one needs to solve the following equation [16]

$$\varepsilon_{\text{max}}^2 \eta \left[\tau_{\gamma'}(\varepsilon_{\text{max}}^2) \right] = \varepsilon_{\text{min}}^2 \eta \left[\tau_{\gamma'}(\varepsilon_{\text{min}}^2) \right], \quad (4.9)$$




Figure 3: The limits on ε - $m_{\gamma'}$ plane in the presence of ALP-photon-dark photon interaction. The light gray region below 10 GeV for $g_{a\gamma\gamma'} = 10^{-2}, 10^{-1} \text{ GeV}^{-1}$ is excluded [15].

which stems from the fact that the observed number of signals for setting the bound is the same.

The impacts of ALP in dark sector on dark photon searches are shown in Fig. 3. When the decay channel $\gamma' \rightarrow a\gamma$ is opened, even though the coupling strength $g_{a\gamma\gamma'} = 10^{-4} \text{ GeV}^{-1}$ is small, its effect shows up for mass of dark photon around 0.1 GeV in some beam dump experiments covering the region of $10^{-3} \gtrsim \varepsilon \gtrsim 10^{-8}$, e.g. E137 and ν -CAL I, see upper-left panel of Fig.3. This is because that these experiments have long decay volume, and efficiency is sensitive to change of decay lifetime. Also, the decay branching ratio of $\gamma' \rightarrow e^+e^-$ is comparable with that of $\gamma' \rightarrow a\gamma$ when $\varepsilon \lesssim 10^{-6}$ for $m_{\gamma'} \lesssim 0.1 \text{ GeV}$. As a result, the excluded ε is shifted to a larger value for a given dark photon mass. And the mass range been covered by beam-dump experiments becomes smaller. If $g_{a\gamma\gamma'}$ is increased to 10^{-3} GeV^{-1} , see the upper-right panel, the sensitivity for $m_{\gamma'}$ heavier than 30 MeV is lost. Furthermore, it is worth mentioning that part of the allowed parameter space, e.g. $20 \text{ MeV} \lesssim m_{\gamma'} \lesssim 30 \text{ MeV}$ and $10^{-5} \text{ GeV}^{-1} \lesssim g_{a\gamma\gamma'} \lesssim 10^{-3} \text{ GeV}^{-1}$ in Fig. 2 is excluded. For LHCb, BaBar and NA48/2, which are sensitive to heavy dark photon search with prompt decay with $\varepsilon \gtrsim 10^{-3}$, the branching of $\gamma' \rightarrow e^+e^-$ is still dominant. Therefore, the lower bounds of ε almost remain when $g_{a\gamma\gamma'} = 10^{-4}$ and 10^{-3} GeV^{-1} . In the case of large $g_{a\gamma\gamma'}$, e.g. 10^{-2} GeV^{-1} , the decay of dark photon is dominated by $\gamma' \rightarrow a\gamma$ channel for $m_{\gamma'} \gtrsim 0.1 \text{ GeV}$, even when $\varepsilon \simeq 10^{-2}$. Therefore, the lower limits of ε from LHCb and BaBar search become loose, especially for heavier dark photon, see lower-left panel of Fig.3. The allowed ε can even reach $\mathcal{O}(0.1)$ for $m_{\gamma'} \gtrsim 10 \text{ GeV}$. Interestingly, the excluded parameter space for dark photon to explain muon $g-2$ is now reopened due to the large $g_{a\gamma\gamma'}$. However, when we take into account the constraints on $g_{a\gamma\gamma'}$, the parameter space for dark photon lighter than 10 GeV is inconsistent

with the constraints of mono-photon searches at BaBar [15] shown by the vertical dashed line. If we take $g_{a\gamma\gamma'}$ to be larger, e.g. 0.1 GeV^{-1} , the LHCb constraints is getting more and more released, and a large parameter space is consistent with muon $g - 2$ result, see the red band in the lower-right of Fig. 3.

5 Summary

The existence of dark sector with portal-type coupling provides a natural and simple way for new physics that weakly interacts with the SM. Through the $U(1)$ kinetic mixing term between dark sector and SM, a massive dark photon from dark sector could couple to SM fermions, and the deviation of muon $g - 2$ from SM can be explained. However, experimental searches for dark photon impose stringent bounds on the mixing parameter ε and exclude the possibility for dark photon to solve the muon $g - 2$ anomaly. In this study, we propose that a light pseudo-Goldstone boson, axion-like particle, may exist in dark sector. Therefore, dark photon could decay into it with a photon, if this decay channel is kinematically allowed. Due to this additional decay channel, we found significant changes in the bounds on ε imposed by both beam-dump and collider experiments. When the coupling $g_{a\gamma\gamma'}$ between axion-like particle, dark photon and photon reaches 0.1, the dominant decay channel of dark photon will be $\gamma' \rightarrow a\gamma$ for mass of dark photon heavier than 0.1 GeV. As a result, the parameter space that is consistent with observation of muon $g - 2$, but previously excluded by LHCb, is now allowed.

Acknowledgment

We would like to acknowledge the support of National Center for Theoretical Sciences (NCTS). This work was supported in part by the National Science and Technology Council (NSTC) of Taiwan under Grant No.MOST 111-2112-M-003-006, 111-2811-M-003-025-, and 112-2811-M-003-004.

References

- [1] R. L. Workman [Particle Data Group], PTEP **2022** (2022), 083C01 doi:10.1093/ptep/ptac097
- [2] M. Fabbrichesi, E. Gabrielli and G. Lanfranchi, doi:10.1007/978-3-030-62519-1 [arXiv:2005.01515 [hep-ph]].
- [3] A. Caputo, A. J. Millar, C. A. J. O'Hare and E. Vitagliano, Phys. Rev. D **104** (2021) no.9, 095029 doi:10.1103/PhysRevD.104.095029 [arXiv:2105.04565 [hep-ph]].
- [4] G. W. Bennett *et al.* [Muon $g-2$], Phys. Rev. D **73**, 072003 (2006) doi:10.1103/PhysRevD.73.072003 [arXiv:hep-ex/0602035 [hep-ex]].
- [5] D. P. Aguillard *et al.* [Muon $g-2$], Phys. Rev. Lett. **131**, no.16, 161802 (2023) doi:10.1103/PhysRevLett.131.161802 [arXiv:2308.06230 [hep-ex]].
- [6] M. Pospelov, Phys. Rev. D **80**, 095002 (2009) doi:10.1103/PhysRevD.80.095002 [arXiv:0811.1030 [hep-ph]].

- [7] K. Kaneta, H. S. Lee and S. Yun, Phys. Rev. Lett. **118**, no.10, 101802 (2017) doi:10.1103/PhysRevLett.118.101802 [arXiv:1611.01466 [hep-ph]].
- [8] S. Roy, C. Blanco, C. Dessert, A. Prabhu and T. Temim, [arXiv:2311.04987 [hep-ph]].
- [9] H. S. Lee and W. Yin, Phys. Rev. D **99**, no.1, 015041 (2019) doi:10.1103/PhysRevD.99.015041 [arXiv:1811.04039 [hep-ph]].
- [10] <https://pdg.lbl.gov/2023/hadronic-xsections/>
- [11] T. Inada, T. Namba, S. Asai, T. Kobayashi, Y. Tanaka, K. Tamasaku, K. Sawada and T. Ishikawa, Phys. Lett. B **722**, 301-304 (2013) doi:10.1016/j.physletb.2013.04.033 [arXiv:1301.6557 [physics.ins-det]].
- [12] H. S. Lee, J. Lee and J. Yi, Phys. Rev. D **106**, no.1, 015011 (2022) doi:10.1103/PhysRevD.106.015011 [arXiv:2201.11906 [hep-ph]].
- [13] M. A. Ismail, C. S. Nugroho and H. T. K. Wong, Phys. Rev. D **107**, no.8, 082002 (2023) doi:10.1103/PhysRevD.107.082002 [arXiv:2211.13384 [hep-ph]].
- [14] J. P. Lees *et al.* [BaBar], Phys. Rev. Lett. **113**, no.20, 201801 (2014) doi:10.1103/PhysRevLett.113.201801 [arXiv:1406.2980 [hep-ex]].
- [15] P. deNiverville, H. S. Lee and M. S. Seo, Phys. Rev. D **98**, no.11, 115011 (2018) doi:10.1103/PhysRevD.98.115011 [arXiv:1806.00757 [hep-ph]].
- [16] P. Ilten, Y. Soreq, M. Williams and W. Xue, JHEP **06**, 004 (2018) doi:10.1007/JHEP06(2018)004 [arXiv:1801.04847 [hep-ph]].
- [17] J. R. Batley *et al.* [NA48/2], Phys. Lett. B **746**, 178-185 (2015) doi:10.1016/j.physletb.2015.04.068 [arXiv:1504.00607 [hep-ex]].
- [18] H. Merkel, P. Achenbach, C. Ayerbe Gayoso, T. Beranek, J. Bericic, J. C. Bernauer, R. Böhm, D. Bosnar, L. Correa and L. Debenjak, *et al.* Phys. Rev. Lett. **112**, no.22, 221802 (2014) doi:10.1103/PhysRevLett.112.221802 [arXiv:1404.5502 [hep-ex]].
- [19] S. Abrahamyan *et al.* [APEX], Phys. Rev. Lett. **107**, 191804 (2011) doi:10.1103/PhysRevLett.107.191804 [arXiv:1108.2750 [hep-ex]].
- [20] F. Archilli *et al.* [KLOE-2], Phys. Lett. B **706**, 251-255 (2012) doi:10.1016/j.physletb.2011.11.033 [arXiv:1110.0411 [hep-ex]].
- [21] A. Anastasi *et al.* [KLOE-2], Phys. Lett. B **757**, 356-361 (2016) doi:10.1016/j.physletb.2016.04.019 [arXiv:1603.06086 [hep-ex]].
- [22] Aaij, Roel and others Phys. Rev. Lett. **120**, 6, 061801 (2018) doi = 10.1103/PhysRevLett.120.061801 [arXiv:1710.02867 [hep-ph]]
- [23] M. Davier and H. Nguyen Ngoc, Phys. Lett. B **229**, 150-155 (1989) doi:10.1016/0370-2693(89)90174-3
- [24] A. Bross, M. Crisler, S. H. Pordes, J. Volk, S. Errede and J. Wrbanek, Phys. Rev. Lett. **67**, 2942-2945 (1991) doi:10.1103/PhysRevLett.67.2942

- [25] J. D. Bjorken, S. Ecklund, W. R. Nelson, A. Abashian, C. Church, B. Lu, L. W. Mo, T. A. Nunamaker and P. Rassmann, *Phys. Rev. D* **38**, 3375 (1988) doi:10.1103/PhysRevD.38.3375
- [26] E. M. Riordan, M. W. Krasny, K. Lang, P. De Barbaro, A. Bodek, S. Dasu, N. Varelas, X. Wang, R. G. Arnold and D. Benton, *et al. Phys. Rev. Lett.* **59**, 755 (1987) doi:10.1103/PhysRevLett.59.755
- [27] A. Konaka, K. Imai, H. Kobayashi, A. Masaike, K. Miyake, T. Nakamura, N. Nagamine, N. Sasao, A. Enomoto and Y. Fukushima, *et al. Phys. Rev. Lett.* **57**, 659 (1986) doi:10.1103/PhysRevLett.57.659
- [28] D. Banerjee *et al.* [NA64], *Phys. Rev. D* **101**, 7, 071101 (2020) doi:10.1103/PhysRevD.101.071101 [arXiv:1912.11389 [hep-ex]].
- [29] G. Bernardi, G. Carugno, J. Chauveau, F. Dicarolo, M. Dris, J. Dumarchez, M. Ferro-Luzzi, J. M. Levy, D. Lukas and J. M. Perreau, *et al. Phys. Lett. B* **166**, 479-483 (1986) doi:10.1016/0370-2693(86)91602-3
- [30] P. Astier *et al.* [NOMAD], *Phys. Lett. B* **506**, 27-38 (2001) doi:10.1016/S0370-2693(01)00362-8 [arXiv:hep-ex/0101041 [hep-ex]].
- [31] F. Bergsma *et al.* [CHARM], *Phys. Lett. B* **157**, 458-462 (1
- [32] J. Blumlein, J. Brunner, H. J. Grabosch, P. Lanius, S. Nowak, C. Rethfeldt, H. E. Ryseck, M. Walter, D. Kiss and Z. Jaki, *et al. Z. Phys. C* **51**, 341-350 (1991) doi:10.1007/BF01548556
- [33] J. Blumlein, J. Brunner, H. J. Grabosch, P. Lanius, S. Nowak, C. Rethfeldt, H. E. Ryseck, M. Walter, *Int. J. Mod. Phys. A* **7**, 3835-3850 (1992) doi:10.1142/S0217751X9200171X
- [34] M. Endo, K. Hamaguchi and G. Mishima, *Phys. Rev. D* **86**, 095029 (2012) doi:10.1103/PhysRevD.86.095029 [arXiv:1209.2558 [hep-ph]].
- [35] B. Abi *et al.* [Muon g-2], *Phys. Rev. Lett.* **126**, no.14, 141801 (2021) doi:10.1103/PhysRevLett.126.141801 [arXiv:2104.03281 [hep-ex]].
- [36] T. Aoyama, N. Asmussen, M. Benay [cite]oun, J. Bijnens, T. Blum, M. Bruno, I. Caprini, C. M. Carloni Calame, M. Cè and G. Colangelo, *et al. Phys. Rept.* **887**, 1-166 (2020) doi:10.1016/j.physrep.2020.07.006 [arXiv:2006.04822 [hep-ph]].
- [37] Bodas, Arushi and Coy, Rupert and King, Simon J. D. *Eur. Phys. J. C* **81**, 12 , 1065 (2021) doi = 10.1140/epjc/s10052-021-09850-x [arXiv:2102.07781 [hep-ph]]